

MSC 34D20

Global solvability, stability and oscillation criteria for systems of two first-order pseudo linear ordinary differential equations

G. A. Grigorian

0019 Armenia c. Yerevan, str. M. Bagramian 24/5

Institute of Mathematics of NAS of Armenia

E - mail: mathphys2@instmath.sci.am, phone: 098 62 03 05, 010 35 48 61

Abstract. In this paper we use the Riccati equation method with other ones to establish global solvability, stability and oscillation criteria for a class of two dimensional nonlinear systems of ordinary differential equations, which is a generalization of wide classes of second order nonlinear ordinary differential equations, studied by many authors. The applicability of some of these criteria are illustrated by examples.

Key words: the system of pseudo linear equations, the Riccati equation, global solvability, the van der Pol's equation, the Duffing's equation, the equation of Emden-Fowler.

1. Introduction. Let $P(t, u, v)$, $Q(t, u, v)$, $R(t, u, v)$, $S(t, u, v)$, $F(t, u, v)$ and $G(t, u, v)$ be real-valued locally integrable by t and continuous by u and v functions on $[t_0, +\infty) \times (\mathbb{R} \setminus \{0\}) \times (\mathbb{R} \setminus \{0\})$, where $\mathbb{R} \equiv (-\infty, +\infty)$. Consider the system of two first-order pseudo linear (i.e. like of linear) ordinary differential equations

$$\begin{cases} \phi' = P(t, \phi, \psi)\phi + Q(t, \phi, \psi)\psi + F(t, \phi, \psi), \\ \psi' = R(t, \phi, \psi)\phi + S(t, \phi, \psi)\psi + G(t, \phi, \psi), \quad t \geq t_0. \end{cases} \quad (1.1)$$

In the particular case, when $F(t, u, v) = G(t, u, v) \equiv 0$ we have a "pseudo homogeneous" case of (1.1):

$$\begin{cases} \phi' = P(t, \phi, \psi)\phi + Q(t, \phi, \psi)\psi, \\ \psi' = R(t, \phi, \psi)\phi + S(t, \phi, \psi)\psi, \quad t \geq t_0. \end{cases} \quad (1.2)$$

By a solution of the system (1.1) (of the system (1.2)) on $[t_1, t_2)$ ($t_0 \leq t_1 < t_2 \leq +\infty$) we mean an ordered pair $(\phi(t), \psi(t))$ of absolutely continuous on $[t_1, t_2)$ functions $\phi(t)$ and $\psi(t)$, satisfying (1.1) ((1.2)) almost everywhere on $[t_1, t_2)$. According to the general theory

of normal systems of ordinary differential equations for every $\alpha, \beta \in \mathbb{R} \setminus \{0\}$ and $t_1 \geq t_0$ there exists $t_2 > t_1$ ($t_2 \leq +\infty$) and a solution $(\phi(t), \psi(t))$ of the system (1.1) ((1.2)) on $[t_1, t_2)$ with $\phi(t_1) = \alpha$, $\psi(t_1) = \beta$. A great interest from the point of view of qualitative theory of differential equations represents the case $t_2 = +\infty$. In this case we say that the system (1.1) ((1.2)) is global solvable. Indicate some well (and not so well) known non linear second order ordinary differential equations, which are reducible to the systems of type (1.1) or (1.2):

1. the van der Pol equation with parametric excitation (see [1], p. 333):

$$\phi'' + \varepsilon(\phi^2 - 1)\phi' + (1 + \beta \cos t)\phi = 0; \quad (1.3)$$

2. the van der Pol equation (see [1], p. 234):

$$\phi'' + \varepsilon(\phi^2 - 1)\phi' + \phi = \Gamma \cos \omega t; \quad (1.4)$$

3. the equation of oscillation of an electronic contour with parametric excitation having an electronic lamp (see [2], p. 221):

$$\phi'' + 2(\lambda_1 + \lambda_2 \phi^2)\phi' + \omega^2(1 - h \cos \nu t)\phi = 0; \quad (1.5)$$

4. the van der Pol-Matheu equation for the dynamics of dust grain charges in dusty plasma (see [3]):

$$\phi'' - (\alpha - \beta \phi^2)\phi' + \omega_0^2(1 + h \cos \gamma t)\phi = 0; \quad (1.6)$$

5. the Lienard's equation (see [1], p. 387):

$$\phi'' + f(t, \phi')\phi' + g(\phi) = 0; \quad (1.7)$$

6. the equation of a pendulum with a light (see [1], p. 331):

$$\phi'' + \left(-\frac{g}{2} + \frac{\varepsilon}{a} \cos \omega t\right) \sin \phi = 0; \quad (1.8)$$

7. the equation of a pendulum (see [2], p. 34):

$$(ml^2(t)\phi')' + gl(t) \sin \phi = 0; \quad (1.9)$$

8. the equation of a pendulum with bob of mass m and rigid suspension of length a , hanging from a support, which is constrained to move with vertical and horizontal displacement $\zeta(t)$ and $\eta(t)$ respectively (see [1], p. 334);

$$a\phi'' + (g + \zeta''(t)) \sin \phi + \eta''(t) \cos \phi = 0; \quad (1.10)$$

9. the Duffing's equation (see [1], p 223):

$$\phi'' + k\phi' + \alpha\phi + \beta\phi^3 = \Gamma \cos \omega t; \quad (1.11)$$

10. the equation of the damped pendulum with periodic forcing of the pivot (see [1], p. 504):

$$\phi'' + \sin \phi = \varepsilon(\gamma \sin t \sin \phi - k\phi'); \quad (1.12)$$

11. the Reyligh's equation (see [1], p. 197):

$$\phi'' + \varepsilon \left(\frac{1}{3}(\phi')^3 - \phi' \right) + \phi = 0; \quad (1.13)$$

12. the equation for the relativistic perturbation of a planetary orbit (see [1], p. 218):

$$\phi'' + \phi = k(1 + \varepsilon\phi^2); \quad (1.14)$$

13. the nonlinear equation with the p -Laplacian (see [4])

$$(\Phi_p(x'))' + \frac{2(p-1)}{t}\Phi(x') + a(t)g(x) = 0, \quad t \geq t_0,$$

$p > 1$, $\Phi_p(y) = |y|^{p-2}y$, $a(t) > 0$, $xg(x) > 0$, $x \neq 0$;

14. (see [6])

$$(|y'|^{\alpha-1}y')' + q(t)|y|^{\beta-1}y = 0;$$

15. (see [7])

$$(a(t)\psi(x)K(x'))' + p(t)K(x') + q(t)f(x)g(x') = 0,$$

$a(t) > 0$, $xf(x) > 0$, $x \neq 0$, $K^2(y) < \gamma_1 yK(y)$, $q(t) > 0$, $0 < c \leq g(x')$, $\frac{f(x)}{x} > \gamma_2 > 0$, $x \neq 0$.

In this paper we prove global solvability, stability and oscillation criteria for the systems (1.1) and (1.2). We illustrate the applicability of some of these criteria by examples.

2. Auxiliary propositions. Let $(\phi(t), \psi(t))$ be a solution of the system (1.1) on $[t_0, t_1)$ ($t_1 \leq +\infty$). We can interpret $\phi(t)$ as a solution of the linear equation

$$\phi' = W(t)\phi + U(t), \quad t \in [t_0, t_1)$$

where $W(t) \equiv P(t, \phi(t), \psi(t))$, $U(t) = G(t, \phi(t), \psi(t))\psi(t) + F(t, \phi(t), \psi(t))\psi(t)$, $t \in [t_0, t_1)$. Then by Cauchy formula we have

$$\phi(t) = \exp \left\{ \int_{t_0}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} \left[\phi(t_0) + \int_{t_0}^t \exp \left\{ - \int_{t_0}^{\tau} P(s; \phi(s); \psi(s)) ds \right\} \times \right.$$

$$\times \left(Q(\tau, \phi(\tau), \psi(\tau))\psi(\tau) + F(\tau, \phi(\tau), \psi(\tau)) \right) d\tau \Big], \quad t \in [t_0, t_1]. \quad (2.1)$$

Analogously for $\psi(t)$ we obtain

$$\begin{aligned} \psi(t) = \exp \left\{ \int_{t_0}^t S(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} & \left[\psi(t_0) + \int_{t_0}^t \exp \left\{ - \int_{t_0}^{\tau} S(s, \phi(s), \psi(s)) ds \right\} \times \right. \\ & \left. \times \left(R(\tau, \phi(\tau), \psi(\tau))\phi(\tau) + G(\tau, \phi(\tau), \psi(\tau)) \right) d\tau \right], \quad t \in [t_0, t_1]. \quad (2.2) \end{aligned}$$

Substitute the right hand part of the last equality into (2.1). After some simplifications we obtain

$$\phi(t) = v_1(t) + \int_{t_0}^t K_1(t, \zeta) \phi(\zeta) d\zeta, \quad t \in [t_0, t_1], \quad (2.3)$$

$$\begin{aligned} \text{where } v_1(t) \equiv & \phi(t_0) \exp \left\{ \int_{t_0}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} + \\ & + \psi(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi(s), \psi(s)) ds + \int_{t_0}^{\tau} S(s, \phi(s), \psi(s)) ds \right\} + \\ & \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi(s), \psi(s)) ds \right\} \left[F(\tau, \phi(\tau), \psi(\tau)) + Q(\tau, \phi(t), \psi(\tau)) \int_{t_0}^{\tau} G(s, \phi(s), \psi(s)) ds \right] d\tau, \\ K_1(t, \zeta) \equiv & R(\zeta, \phi(\zeta), \psi(\zeta)) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t P(s, \phi(s), \psi(s)) ds + \int_{\zeta}^{\tau} S(s, \phi(s), \psi(s)) ds \right\} \times \\ & \times Q(\tau, \phi(\tau), \psi(\tau)) d\tau, \quad t, \zeta \in [t_0, t), \quad \zeta \leq t. \end{aligned}$$

By similar way from (2.1) and (2.2) we can obtain the equality

$$\psi(t) = v_2(t) + \int_{t_0}^t K_2(t, \zeta) \psi(\zeta) d\zeta, \quad t \in [t_0, t_1], \quad (2.4)$$

$$\begin{aligned}
\text{where } v_2(t) \equiv & \psi(t_0) \exp \left\{ \int_{t_0}^t S(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} + \\
& + \phi(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi(s), \psi(s)) ds + \int_{t_0}^{\tau} P(s, \phi(s), \psi(s)) ds \right\} + \\
& \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi(s), \psi(s)) ds \right\} \left[G(\tau, \phi(\tau), \psi(\tau)) + R(\tau, \phi(\tau), \psi(\tau)) \int_{t_0}^{\tau} F(s, \phi(s), \psi(s)) ds \right] d\tau, \\
K_2(t, \zeta) \equiv & Q(\zeta, \phi(\zeta), \psi(\zeta)) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t S(s, \phi(s), \psi(s)) ds + \int_{\zeta}^{\tau} P(s, \phi(s), \psi(s)) ds \right\} \times \\
& \times R(\tau, \phi(\tau), \psi(\tau)) d\tau, \quad t, \zeta \in [t_0, t], \quad \zeta \leq t.
\end{aligned}$$

Let $f_k(t)$, $g_k(t)$, $h_k(t)$, $k = 1, 2$ be real-valued locally integrable functions on $[t_0, +\infty)$. Consider the Riccati equations

$$y' + f_k(t)y^2 + g_k(t)y + h_k(t) = 0, \quad t \geq t_0. \quad (2.5_k)$$

$k = 1, 2$ and the differential inequalities

$$\eta + f_k(t)\eta^2 + g_k(t)\eta + h_k(t) \geq 0, \quad t \geq t_0. \quad (2.6_k)$$

$k = 1, 2$. By a solution of Eq. (2.5_k) (of inequality (2.6_k)) on $[t_1, t_2]$ ($t_0 \leq t_1 < t_2 \leq +\infty$) we mean an absolutely continuous function $y(t)$ ($\eta(t)$) on $[t_1, t_2]$, satisfying (2.5_k) ((2.6_k)) almost everywhere on $[t_1, t_2]$, $k = 1, 2$.

Remark 2.1. Every solution of Eq. (2.5₂) on $[t_0, t_1]$ is also a solution of the inequality (2.6₂) on $[t_0, t_1]$.

Remark 2.2. If $f_1(t) \geq 0$, $t \in [t_0, t_1]$, then every solution of the linear equation

$$\zeta' + g_1(t)\zeta + h_1(t) = 0, \quad t \in [t_0, t_1]$$

is also a solution of the inequality (2.6₁) on $[t_0, t_1]$.

Theorem 2.1. Let $y_2(t)$ be a solution of Eq. (2.5₂) on $[t_0, \tau_0]$ ($t_0 < \tau_0 \leq +\infty$) and let $\eta_1(t)$ and $\eta_2(t)$ be solutions of the inequalities (2.6₁) and (2.6₂) respectively on $[t_0, \tau_0]$ such that $y_2(t_0) \leq \eta_k(t_0)$ $k = 1, 2$. In addition let the following conditions be satisfied:

$$f_1(t) \geq 0, \quad \gamma - y_2(t_0) + \int_{t_0}^t \exp \left\{ \int_{t_0}^{\tau} [f_1(s)(\eta_1(s) + \eta_2(s)) + g_1(s)] ds \right\} \left[(f_2(\tau) - f_1(\tau))^2 y_2^2(\tau) + \right.$$

$(g_2(\tau) - g_1(\tau))y_2(\tau) + h_2(\tau) - h_1(\tau) \Big] d\tau \geq 0, \quad t \in [t_0, \tau_0)$ for some $\gamma \in [y_2(t_0), \eta_1(t_0)]$. Then Eq. (2.5₁) has a solution $y_1(t)$ on $[t_0, \tau_0)$ with $y_1(t_0) \geq \gamma$ and $y_1(t) \geq y_2(t), \quad t \in [t_0, \tau_0)$

Proof. By analogy of the proof of Theorem 3.1 from [7].

In the system (1.2) substitute

$$\psi = y\phi$$

We obtain

$$\begin{cases} \phi' = [P(t; \phi; \psi) + yQ(t, \phi, \psi)]\phi + F(t, \phi, \psi), \\ [y'\phi + y(t)\{[B(t, \phi, \psi) + y(t)Q(t, \phi, \psi)]\psi + F(t, \phi, \psi)\} = R(t, \phi, \psi)\phi + G(t, \phi, \psi), \quad t \geq t_0, \end{cases}$$

where $B(t, \phi, \psi) \equiv P(t, \phi, \psi) - S(t, \phi, \psi)$. It follows from here that if $(\phi(t), \psi(t))$ is a solution of the system (1.1) on $[t_0, t_1)$ with $\phi(t) \neq 0, \quad t \in [t_0, t_1)$ and $y(t)$ is the solution of the Riccati equation

$$\begin{aligned} y' + Q(t, \phi(t), \psi(t))y^2 + \left(B(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\phi(t)} \right) y - \\ - R(t, \phi(t), \psi(t)) - \frac{G(t, \phi(t), \psi(t))}{\phi(t)} = 0 \end{aligned} \quad (2.7)$$

on $[t_0, t_1)$ with $y(t_0) = \frac{\psi(t_0)}{\phi(t_0)}$ then

$$\begin{cases} \phi(t) = \phi(t_0) \exp \left\{ \int_{t_0}^t \left[P(\tau, \phi(\tau), \psi(\tau)) + \frac{F(\tau, \phi(\tau), \psi(\tau))}{\phi(\tau)} + \right. \right. \\ \left. \left. + y(\tau)Q(\tau; \phi(\tau); \psi(\tau)) \right] d\tau \right\}, \quad \text{or} \\ \phi(t) = \exp \left\{ \int_{t_0}^t [P(\tau; \phi(\tau), \psi(\tau)) + y(\tau)Q(\tau, \phi(\tau), \psi(\tau))] d\tau \right\} \left[\phi(t_0) + \right. \\ \left. + \int_{t_0}^t \exp \left\{ - \int_{t_0}^{\tau} [P(s, \phi(s), \psi(s)) + y(s)Q(s, \phi(s), \psi(s))] ds \right\} F(\tau, \phi(\tau), \psi(\tau)) d\tau \right] \\ \psi(t) = y(t)\phi(t), \quad t \in [t_0, t_1) \end{cases} \quad (2.8)$$

Analogously the substitution

$$\phi = z\psi$$

in (1.1) implies that if $(\phi(t), \psi(t))$ is a solution of the system (1.1) on $[t_0, t_1)$ with $\psi(t) \neq 0$, $t \in [t_0, t_1)$ and $z(t)$ is the solution of the Riccati equation

$$y' + R(t, \phi(t), \psi(t))y^2 - \left(B(t, \phi(t), \psi(t)) + \frac{G(t, \phi(t), \psi(t))}{\phi(t)} \right) y - Q(t, \phi(t), \psi(t)) - \frac{F(t, \phi(t), \psi(t))}{\phi(t)} = 0 \quad (2.9)$$

on $[t_0, t_1)$ with $z(t_0) = \frac{\phi(t_0)}{\psi(t_0)}$ then

$$\left\{ \begin{array}{l} \psi(t) = \psi(t_0) \exp \left\{ \int_{t_0}^t \left[S(\tau, \phi(\tau), \psi(\tau)) + \frac{G(\tau, \phi(\tau), \psi(\tau))}{\psi(\tau)} + z(\tau)R(\tau, \phi(\tau), \psi(\tau)) \right] d\tau \right\}, \text{ or} \\ \psi(t) = \exp \left\{ \int_{t_0}^t \left[S(\tau, \phi(\tau), \psi(\tau)) + z(\tau)QR(\tau, \phi(\tau), \psi(\tau)) \right] d\tau \right\} \left[\psi(t_0) + \int_{t_0}^t \exp \left\{ - \int_{t_0}^{\tau} \left[S(s, \phi(s), \psi(s)) + z(s)R(s, \phi(s), \psi(s)) \right] ds \right\} G(\tau, \phi(\tau), \psi(\tau)) d\tau \right] \\ \psi(t) = y(t)\phi(t), \quad t \in [t_0, t_1) \end{array} \right. \quad (2.10)$$

Note that we can interpret a solution $y(t)$ of Eq. (2.7) on $[t_0, t_1)$ as a solution of a linear equation

$$y' + H(t)y - R_1(t, \phi(t), \psi(t)) = 0$$

on $[t_0, t_1)$, where $H(t) \equiv Q(t, \phi(t), \psi(t))y(t) + B(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\phi(t)}$, $R_1(t) \equiv R(t, \phi(t), \psi(t)) + \frac{G(t, \phi(t), \psi(t))}{\phi(t)}$, $t \in [t_0, t_1)$. Then according to Cauchy formula we have

$$y(t) = \exp \left\{ - \int_{t_0}^t H(\tau) d\tau \right\} \left[y(t_0) + \int_{t_0}^t \exp \left\{ \int_{t_0}^{\tau} H(s) ds \right\} R_1(\tau, \phi(\tau), \psi(\tau)) d\tau \right], \quad (2.11)$$

$t \in [t_0, t_1)$. By analogy for a solution $z(t)$ of Eq. (2.9) we get

$$z(t) = \exp \left\{ - \int_{t_0}^t L(\tau) d\tau \right\} \left[z(t_0) + \int_{t_0}^t \exp \left\{ \int_{t_0}^{\tau} L(s) ds \right\} Q_1(\tau, \phi(\tau), \psi(\tau)) d\tau \right], \quad (2.12)$$

$t \in [t_0, t_1]$, where $L(t) \equiv R(t, \phi(t), \psi(t))z(t) - B(t, \phi(t), \psi(t)) + \frac{G(t, \phi(t), \psi(t))}{\psi(t)}$,
 $Q_1(t) \equiv Q(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\psi(t)}$, $t \in [t_0, t_1]$

Lemma 2.1. *Let $(\phi(t), \psi(t))$ be a solution of the system (1.2) on $[t_0, t_1]$. If $Q(t, \phi(t), \psi(t)) \geq 0$, $R(t, \phi(t), \psi(t)) \geq 0$, $t \in [t_0, t_1]$, then for every $\gamma \geq 0$ Eq. (2.7) (Eq. (2.9)) has a non negative solution $y(t)$ ($z(t)$) on $[t_0, t_1]$.*

Proof. Let us prove the existence of $y(t)$. Consider the Riccati equation

$$y' + Q(t, \phi(t), \psi(t))y^2 + B(t, \phi(t), \psi(t))y = 0, \quad t \in [t_0, t_1] \quad (2.13)$$

Obviously $y_2(t) \equiv 0$ is a solution of this equation on $[t_0, t_1]$. Then since $Q(t, \phi(t), \psi(t)) \geq 0$, $R(t, \phi(t), \psi(t)) \geq 0$, $t \in [t_0, t_1]$, using Theorem 2.1 to the pair of equations (2.7) and (2.13) we conclude that for every $\gamma \geq 0$ Eq. (2.7) has a non negative solution $y(t)$ on $[t_0, t_1]$ with $y(t_0) = \gamma$. The proof of existence of $z(t)$ can be made by analogy using Eq. (2.9) instead of Eq. (2.7). The lemma is proved.

Definition 2.1. *An interval $[t_0, T)$ is called the maximum existence interval for a solution $(\phi(t), \psi(t))$ of the system (1.1), if $(\phi(t), \psi(t))$ satisfies (1.1) almost everywhere on $[t_0, T)$ and cannot be continued to the right from T as a solution of Eq. (1.1).*

Lemma 2.2. *Assume $P(t, u, v)$, $Q(t, u, v)$, $R(t, u, v)$, $S(t, u, v)$, $F(t, u, v)$ and $G(t, u, v)$ are locally integrable and locally bounded by t and continuous by u and v on $[t_0, +\infty) \times \mathbb{R} \times \mathbb{R}$. If $(\phi(t), \psi(t))$ is a bounded solution of the system (1.1) on $[t_0, T)$ and $T < +\infty$, then $[t_0, T)$ cannot be the maximum existence interval for $(\phi(t), \psi(t))$.*

Proof. To prove the lemma it is enough to show that $(\phi(t), \psi(t))$ has a finite limit when $t \rightarrow T - 0$. By (1.1) we have

$$\phi(t) = \phi(t_0) + \int_{t_0}^t A_{\phi, \psi}(\tau) d\tau, \quad t \in [t_0, T), \quad (2.14)$$

where $A_{\phi, \psi}(\tau) \equiv P(\tau, \phi(\tau), \psi(\tau))\phi(\tau) + Q(\tau, \phi(\tau), \psi(\tau))\psi(\tau) + F(\tau, \phi(\tau), \psi(\tau))$, $\tau \in [t_0, T)$. Since $(\phi(t), \psi(t))$ is bounded on $[t_0, T)$ and $P(t, u, v)$, $Q(t, u, v)$, $R(t, u, v)$ are continuous by u , v on \mathbb{R} functions we have

$$|A_{\phi, \psi}(t)| \leq M, \quad t \in [t_0, T) \quad (2.15)$$

for some $M \geq 0$. Let $\{t_k\}_{k=1}^{+\infty}$ be a sequence such that $t_k < T$, $k = 1, 2, \dots$ and $\lim_{k \rightarrow +\infty} t_k = T$.

Then by (2.15) for any integers $m, n \geq 1$ we have $\left| \int_{t_m}^{t_n} A_{\phi, \psi}(\tau) d\tau \right| \leq M|t_m - t_n|$. Hence by (2.14) $|\phi(t_m) - \phi(t_n)| \leq M|t_m - t_n|$, i. e. the sequence $\{\phi(t_k)\}_{k=1}^{+\infty}$ is fundamental, which

proves that $\phi(t)$ has a finite limit when $t \rightarrow T - 0$. By analogy it can be proved the existence of the finite limit $\lim_{t \rightarrow T-0} \psi(t)$. The lemma is proved.

3. Global solvability, stability and oscillation criteria.

Definition 3.1. A solution $(\phi_0(t), \psi_0(t))$ of the system (1.1) ((1.2)) is called *Lyapunov (asymptotically) stable* if for every $\varepsilon > 0$ there exists $\delta > 0$ such that for every solution $(\phi(t), \psi(t))$ of the system (1.1) ((1.2)) with $|\phi(t_0) - \phi_0(t_0)| \leq \delta$, $|\psi(t_0) - \psi_0(t_0)| \leq \delta$ the inequalities

$$|\phi(t) - \phi_0(t)| \leq \varepsilon, \quad |\psi(t) - \psi_0(t)| \leq \varepsilon, \quad t \geq t_0$$

are valid (the relations

$$\lim_{t \rightarrow +\infty} (\phi(t) - \phi_0(t)) = \lim_{t \rightarrow +\infty} (\psi(t) - \psi_0(t)) = 0$$

are valid).

Definition 3.2. A solution $(\phi_0(t), \psi_0(t))$ of the system (1.1) ((1.2)) is called *global asymptotically stable* if for every its solution $(\phi(t), \psi(t))$ the functions $\phi(t) - \phi_0(t)$ and $\psi(t) - \psi_0(t)$ vanish at $+\infty$.

Let $P_0(t)$, $Q_0(t)$, $R_0(t)$, $S_0(t)$, $F_0(t)$ and $G_0(t)$ be real-valued locally integrable functions on $[t_0, +\infty)$. Consider the linear system of ordinary differential equations

$$\begin{cases} \phi' = P_0(t)\phi + Q_0(t)\psi + F_0(t), \\ \psi' = R_0(t)\phi + S_0(t)\psi + G_0(t), \quad t \geq t_0 \end{cases} \quad (3.1)$$

and the corresponding homogeneous one

$$\begin{cases} \phi' = P_0(t)\phi + Q_0(t)\psi, \\ \psi' = R_0(t)\phi + S_0(t)\psi, \quad t \geq t_0 \end{cases} \quad (3.2)$$

Theorem 3.1. Assume $P(t, u, v)$, $Q(t, u, v)$, $R(t, u, v)$, $S(t, u, v)$, $F(t, u, v)$ and $G(t, u, v)$ are locally integrable and locally bounded by t and continuous by u and v on $[t_0, +\infty) \times \mathbb{R} \times \mathbb{R}$. Moreover assume $P(t, u, v) \leq P_0(t)$, $|Q(t, u, v)| \leq Q_0(t)$, $|R(t, u, v)| \leq R_0(t)$, $S(t, u, v) \leq S_0(t)$, $|F(t, u, v)| \leq F_0(t)$, $|G(t, u, v)| \leq G_0(t)$, $t \geq t_0, u, v \in \mathbb{R}$. Then the following statements are valid.

- I) For every $\alpha, \beta \in \mathbb{R}$ the solution $(\phi(t), \psi(t))$ of the system (1.1) with $\phi(t_0) = \alpha$, $\psi(t_0) = \beta$ exists on $[t_0, +\infty)$.
- II) If a solution $(\phi_0(t), \psi_0(t))$ of the system (3.1) with $\phi_0(t_0) > 0$, $\psi_0(t_0) > 0$ is bounden on $[t_0, +\infty)$ (vanish at $+\infty$), then every solution $(\phi(t), \psi(t))$ of the system (1.1) with $|\phi(t_0)| \leq \phi_0(t_0)$, $|\psi(t_0)| \leq \psi_0(t_0)$ is also bounden on $[t_0, +\infty)$ (vanish at $+\infty$).

III) If the system (3.2) is Lyapunov (asymptotically) stable then the null solution $(0, 0)$ of the system (1.2) is Lyapunov (global asymptotically) stable.

Proof. Let us prove the statement I). Let $(\phi_0(t), \psi_0(t))$ be a solution of the system (1.1) with $\phi_0(t_0) = \alpha$, $\psi_0(t_0) = \beta$. Suppose this solutions does not exist on $[t_0, +\infty)$. Let then $[t_0, T)$ be the maximum existence interval for this solution. In virtue of (2.3) and (2.4) $\phi_0(t)$ and $\psi_0(t)$ are solutions of the Volterra equations

$$\phi(t) = \overset{o}{v}_1(t) + \int_{t_0}^t \overset{o}{K}_1(t, \zeta) \phi(\zeta) d\zeta, \quad t \in [t_0, T),$$

$$\psi(t) = \overset{o}{v}_2(t) + \int_{t_0}^t \overset{o}{K}_2(t, \zeta) \psi(\zeta) d\zeta, \quad t \in [t_0, T)$$

respectively, where

$$\begin{aligned} \overset{o}{v}_1(t) &\equiv \phi_0(t_0) \exp \left\{ \int_{t_0}^t P(\tau, \phi_0(\tau), \psi_0(\tau)) d\tau \right\} + \psi_0(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi_0(s), \psi_0(s)) ds \right. \\ &+ \left. \int_{t_0}^{\tau} S(s, \phi_0(s), \psi_0(s)) ds \right\} d\tau + \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi_0(s), \psi_0(s)) ds \right\} \times \\ &\times \left[F(\tau, \phi_0(\tau), \psi_0(\tau)) + Q(\tau, \phi_0(\tau), \psi_0(\tau)) \int_{t_0}^{\tau} G(s, \phi_0(s), \psi_0(s)) ds \right] d\tau \\ \overset{o}{v}_2(t) &\equiv \psi_0(t_0) \exp \left\{ \int_{t_0}^t S(\tau, \phi_0(\tau), \psi_0(\tau)) d\tau \right\} + \phi_0(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi_0(s), \psi_0(s)) ds \right. \\ &+ \left. \int_{t_0}^{\tau} P(s, \phi_0(s), \psi_0(s)) ds \right\} d\tau + \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi_0(s), \psi_0(s)) ds \right\} \times \\ &\times \left[G(\tau, \phi_0(\tau), \psi_0(\tau)) + R(\tau, \phi_0(\tau), \psi_0(\tau)) \int_{t_0}^{\tau} F(s, \phi_0(s), \psi_0(s)) ds \right] d\tau \end{aligned}$$

$$\begin{aligned}
\overset{\circ}{K}_1(t, \zeta) &\equiv R(t, \phi_0(t), \psi_0(t)) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t P(s, \phi_0(s), \psi_0(s)) ds + \int_{\zeta}^{\tau} S(s, \phi_0(s), \psi_0(s)) ds \right\} \times \\
&\quad \times Q(\tau, \phi_0(\tau), \psi_0(\tau)) d\tau, \\
\overset{\circ}{K}_2(t, \zeta) &\equiv Q(t, \phi_0(t), \psi_0(t)) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t S(s, \phi_0(s), \psi_0(s)) ds + \int_{\zeta}^{\tau} P(s, \phi_0(s), \psi_0(s)) ds \right\} \times \\
&\quad \times R(\tau, \phi_0(\tau), \psi_0(\tau)) d\tau,
\end{aligned}$$

It follows from the conditions of the theorem that

$$|\overset{\circ}{v}_j(t)| \leq \overset{\circ\circ}{v}_j(t), \quad |\overset{\circ}{K}_j(t, \zeta)| \leq \overset{\circ\circ}{K}_j(t, \zeta), \quad j = 1, 2, \quad t, \zeta \in [t_0, T), \quad \zeta \leq t, \quad (3.3)$$

where

$$\begin{aligned}
\overset{\circ\circ}{v}_1(t) &\equiv |\phi_0(t_0)| \exp \left\{ \int_{t_0}^t P_0(\tau) d\tau \right\} + |\psi_0(t_0)| \int_{t_0}^t \exp \left\{ \int_{\tau}^t P_0(s) ds + \int_{t_0}^{\tau} S_0(s) ds \right\} d\tau + \\
&\quad + \int_{t_0}^t \exp \left\{ \int_{\tau}^t P_0(s) ds \right\} \left[F_0(\tau) + Q_0(\tau) \int_{t_0}^{\tau} G_0(s) ds \right] d\tau \\
\overset{\circ\circ}{v}_2(t) &\equiv |\psi_0(t_0)| \exp \left\{ \int_{t_0}^t S_0(\tau) d\tau \right\} + |\phi_0(t_0)| \int_{t_0}^t \exp \left\{ \int_{\tau}^t S_0(s) ds + \int_{t_0}^{\tau} P_0(s) ds \right\} d\tau + \\
&\quad + \int_{t_0}^t \exp \left\{ \int_{\tau}^t S_0(s) ds \right\} \left[G_0(\tau) + R_0(\tau) \int_{t_0}^{\tau} F_0(s) ds \right] d\tau \\
\overset{\circ\circ}{K}_1(t, \zeta) &\equiv R_0(\zeta) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t P_0(s) ds + \int_{\zeta}^{\tau} S_0(s) ds \right\} Q_0(\tau) d\tau, \\
\overset{\circ\circ}{K}_2(t, \zeta) &\equiv Q_0(\zeta) \int_{\zeta}^t \exp \left\{ \int_{\tau}^t S_0(s) ds + \int_{\zeta}^{\tau} P_0(s) ds \right\} R_0(\tau) d\tau, \quad t, \zeta \in [t_0, T), \quad \zeta \leq t.
\end{aligned}$$

Let $\phi_{oo}(t)$ and $\psi_{oo}(t)$ be solutions of the Volterra equations

$$\phi(t) = {}^{oo}v_1(t) + \int_{t_0}^t {}^{oo}K_1(t, \zeta) \phi(\zeta) d\zeta, \quad t \in [t_0, T), \quad (3.4)$$

$$\psi(t) = {}^{oo}v_2(t) + \int_{t_0}^t {}^{oo}K_2(t, \zeta) \psi(\zeta) d\zeta, \quad t \in [t_0, T), \quad (3.5)$$

respectively. Show that $\phi_{oo}(t)$ and $\psi_{oo}(t)$ are bounded functions on $[t_0, T)$. By (3.4) and (3.5) for $\phi_{oo}(t)$ and $\psi_{oo}(t)$ we have the representations via series expansion

$$\phi_{oo}(t) = {}^{oo}v_1(t) + \int_{t_0}^t {}^{oo}K_1(t, \zeta) {}^{oo}v_1(\zeta) d\zeta + \int_{t_0}^t {}^{oo}K_1(t, \zeta) d\zeta \int_{t_0}^{\zeta} {}^{oo}K_1(t, \xi) {}^{oo}v_1(\xi) d\xi + \dots, \quad (3.6)$$

$$\psi_{oo}(t) = {}^{oo}v_2(t) + \int_{t_0}^t {}^{oo}K_2(t, \zeta) {}^{oo}v_2(\zeta) d\zeta + \int_{t_0}^t {}^{oo}K_2(t, \zeta) d\zeta \int_{t_0}^{\zeta} {}^{oo}K_2(t, \xi) {}^{oo}v_2(\xi) d\xi + \dots, \quad (3.7)$$

respectively. From the definitions of ${}^{oo}v_j(t)$, ${}^{oo}K_j(t, \zeta)$, $j = 1, 2$ is seen that they are bounded functions for $t, \zeta \in [t_0, T)$, $\zeta \leq t$. Then from (3.6) and (3.7) we obtain respectively:

$$|\phi_{oo}(t)| \leq m_1 \left(1 + M_1(T - t_0) + \frac{[M_1(T - t_0)]^2}{2!} + \dots \right) = m_1 \exp\{M_1(T - t_0)\}, \quad (3.8)$$

$$|\psi_{oo}(t)| \leq m_2 \left(1 + M_2(T - t_0) + \frac{[M_2(T - t_0)]^2}{2!} + \dots \right) = m_2 \exp\{M_2(T - t_0)\}, \quad (3.9)$$

where $m_j \equiv \sup_{t \in [t_0, T)} |{}^{oo}v_j(t)|$, $M_j \equiv \sup_{t, \zeta \in [t_0, T), \zeta \leq t} |{}^{oo}K_j(t, \zeta)|$, $j = 1, 2$. Hence $\phi_{oo}(t)$ and $\psi_{oo}(t)$ are bounded functions on $[t_0, T)$. This together with (3.3), (3.8) and (3.9) implies that $(\phi_0(t), \psi_0(t))$ is bounded on $[t_0, T)$. Then by virtue of Lemma 2.2 $[t_0, T)$ is not the maximum existence interval for $(\phi_0(t), \psi_0(t))$, which contradicts our assumption. The obtained contradiction completes the proof of the statement I). The statement III) follows immediately from the statement II). Prove the statement II). Let $(\phi_0(t), \psi_0(t))$ be a solution of the system (3.1) with $\phi_0(t_0) > 0$, $\psi_0(t_0) > 0$. Then $\phi_0(t)$ and $\psi_0(t)$ are

solutions of the Volterra equations (3.4) and (3.5) respectively on $[t_0, +\infty)$. Let $(\phi(t), \psi(t))$ be a solution of the system (1.1) with

$$|\phi(t_0)| \leq \phi_0(t_0), \quad |\psi(t_0)| \leq \psi_0(t_0) \quad (3.10)$$

(according to the statement I) $(\phi(t), \psi(t))$ exists on $[t_0, +\infty)$). In virtue of (2.3) and (2.4) for $\phi(t)$ and $\psi(t)$ the relations

$$\begin{aligned} \phi(t) &= v_1(t) + \int_{t_0}^t K_1(t, \zeta) \phi(\zeta) d\zeta, \quad t \geq t_0, \\ \psi(t) &= v_2(t) + \int_{t_0}^t K_2(t, \zeta) \psi(\zeta) d\zeta, \quad t \geq t_0 \end{aligned}$$

are valid respectively, where

$$\begin{aligned} v_1(t) &\equiv \phi(t_0) \exp \left\{ \int_{t_0}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} + \psi(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi(s), \psi(s)) ds + \right. \\ &+ \left. \int_{t_0}^{\tau} S(s, \phi(s), \psi(s)) ds \right\} d\tau + \int_{t_0}^t \exp \left\{ \int_{\tau}^t P(s, \phi(s), \psi(s)) ds \right\} \times \\ &\times \left[F(\tau, \phi(\tau), \psi(\tau)) + Q(\tau, \phi(\tau), \psi(\tau)) \int_{t_0}^{\tau} G(s, \phi(s), \psi(s)) ds \right] d\tau \\ v_2(t) &\equiv \psi(t_0) \exp \left\{ \int_{t_0}^t S(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} + \phi(t_0) \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi(s), \psi(s)) ds + \right. \\ &+ \left. \int_{t_0}^{\tau} P(s, \phi(s), \psi(s)) ds \right\} d\tau + \int_{t_0}^t \exp \left\{ \int_{\tau}^t S(s, \phi(s), \psi(s)) ds \right\} \times \\ &\times \left[G(\tau, \phi(\tau), \psi(\tau)) + R(\tau, \phi(\tau), \psi(\tau)) \int_{t_0}^{\tau} F(s, \phi(s), \psi(s)) ds \right] d\tau \end{aligned}$$

For $\phi_0(t)$ and $\psi_0(t)$ we have the relations

$$\phi_0(t) = {}^0u_1(t) + \int_{t_0}^t {}^0L_1(t, \zeta) \phi_0(\zeta) d\zeta, \quad t \geq t_0,$$

$$\psi_0(t) = {}^0u_2(t) + \int_{t_0}^t {}^0L_2(t, \zeta) \psi_0(\zeta) d\zeta, \quad t \geq t_0,$$

respectively, where

$$\begin{aligned} {}^0u_1(t) &\equiv \phi_0(t_0) \exp\left\{\int_{t_0}^t P_0(\tau) d\tau\right\} + \psi_0(t_0) \int_{t_0}^t \exp\left\{\int_{\tau}^t P_0(s) ds + \int_{t_0}^{\tau} S_0(s) ds\right\} d\tau + \\ &\quad + \int_{t_0}^t \exp\left\{\int_{\tau}^t P_0(s) ds\right\} \left[F_0(\tau) + Q_0(\tau) \int_{t_0}^{\tau} G_0(s) ds\right] d\tau \\ {}^0u_2(t) &\equiv \psi_0(t_0) \exp\left\{\int_{t_0}^t S_0(\tau) d\tau\right\} + \phi_0(t_0) \int_{t_0}^t \exp\left\{\int_{\tau}^t S_0(s) ds + \int_{t_0}^{\tau} P_0(s) ds\right\} d\tau + \\ &\quad + \int_{t_0}^t \exp\left\{\int_{\tau}^t S_0(s) ds\right\} \left[G_0(\tau) + R_0(\tau) \int_{t_0}^{\tau} F_0(s) ds\right] d\tau \\ {}^0L_1(t, \zeta) &\equiv R_0(\zeta) \int_{\zeta}^t \exp\left\{\int_{\tau}^t P_0(s) ds + \int_{\zeta}^{\tau} S_0(s) ds\right\} Q_0(\tau) d\tau, \\ {}^0L_2(t, \zeta) &\equiv Q_0(\zeta) \int_{\zeta}^t \exp\left\{\int_{\tau}^t S_0(s) ds + \int_{\zeta}^{\tau} P_0(s) ds\right\} R_0(\tau) d\tau, \quad t \geq \zeta \geq t_0. \end{aligned}$$

It follows from the conditions of the theorem that

$$|v_j(t)| \leq {}^0u_j(t), \quad |k_j(t, \zeta)| \leq |{}^0L_j(t, \zeta)|, \quad j = 1, 2, \quad t \geq \zeta \geq t_0.$$

Then from (3.10) it follows that

$$|\phi(t)| \leq \phi_0(t), \quad |\psi(t)| \leq \psi_0(t), \quad t \geq t_0. \quad (3.11)$$

Assume $(\phi_0(t), \psi_0(t))$ is bounded on $[t_0, +\infty)$ (vanish at $+\infty$). Then the relations (3.11) imply that $(\phi(t), \psi(t))$ is bounded on $[t_0, +\infty)$ (vanish at $+\infty$). The statement II) is proved. The proof of the theorem is complete.

Remark 3.1. From the proof of Theorem 3.1 is seen that it remains valid if we generalize its conditions assuming that $P(t, u, v)$, $Q(t, u, v)$, $R(t, u, v)$, $S(t, u, v)$, $F(t, u, v)$ and $G(t, u, v)$ are complex-valued locally integrable and locally bounded by t and continuous by u and v functions on $[t_0, +\infty) \times (\mathbb{C}) \times (\mathbb{C})$ and replace $\alpha, \beta \in \mathbb{R}$ by $\alpha, \beta \in \mathbb{C}$ and the conditions $P(t, u, v) \leq P_0(t)$, $S(t, u, v) \leq S_0(t)$ by $\operatorname{Re} P(t, u, v) \leq P_0(t)$, $\operatorname{Re} S(t, u, v) \leq S_0(t)$.

Example 3.1. Eq. (1.3) is equivalent to the system

$$\begin{cases} \phi' = \psi, \\ \psi' = -(1 + \beta \cos t)\phi - \varepsilon(\phi^2 - 1)\psi, \quad t \geq t_0. \end{cases}$$

Obviously for this system with $\varepsilon \geq 0$ the conditions of Theorem 3.1 are satisfied. Therefore for every $\alpha, \beta \in \mathbb{R}$ Eq. (1.3) has a solution $\phi(t)$ on $[t_0, +\infty)$ with $\phi(t_0) = \alpha$, $\psi(t_0) = \beta$. By similar way can be discussed the applicability of Theorem 3.1 to the equations (1.4) - (1.14). The applicability of Theorem 3.1 to the system of equations (see [1], p. 257):

$$\begin{cases} \phi' = -(\phi^2 + \psi^2 - 1)\phi + a \sin t \psi, \\ \psi' = -a \sin t \phi - (\phi^2 + \psi^2 - 1)\psi \end{cases}$$

is obvious.

Let $B_1(t)$ and $B_2(t)$ be real-valued continuous functions on $[t_0, +\infty)$. For any $c_1 \neq 0$ and $c_2 \neq 0$ set:

$$\begin{aligned} K(t, c_1, c_2) \equiv & c_1 \exp \left\{ \int_{t_0}^t P_0(s) ds + \int_{t_0}^t Q_0(\tau) \left[\frac{c_2}{c_1} \exp \left\{ - \int_{t_0}^{\tau} B_1(s) ds \right\} + \right. \right. \\ & \left. \left. + \int_{t_0}^{\tau} \exp \left\{ - \int_{\zeta}^{\tau} B_1(s) ds \right\} R_0(\zeta) d\zeta \right] d\tau \right\}, \\ L(t, c_1, c_2) \equiv & c_2 \exp \left\{ \int_{t_0}^t S_0(s) ds + \int_{t_0}^t R_0(\tau) \left[\frac{c_1}{c_2} \exp \left\{ \int_{t_0}^{\tau} B_2(s) ds \right\} + \right. \right. \end{aligned}$$

$$+ \int_{t_0}^{\tau} \exp \left\{ \int_{\zeta}^{\tau} B_2(s) ds \right\} Q_0(\zeta) d\zeta \Big] d\tau \Big\}, \quad t \geq t_0.$$

Theorem 3.2.. *Let for some $c_1 > 0$, $c_2 > 0$, $\varepsilon > 0$ and for every $t \geq t_0$, $u \in (0, K(t, c_1, c_2) + \varepsilon]$, $v \in (0, L(t, c_1, c_2) + \varepsilon]$ the inequalities $P(t, u, v) \leq P_0(t)$, $S(t, u, v) \leq S_0(t)$, $0 \leq Q(t, u, v) \leq Q_0(t)$, $0 \leq R(t, u, v) \leq R_0(t)$, $B_1(t) \leq B(t, u, v) \leq B_2(t)$ are valid. Then every solution $(\phi(t), \psi(t))$ of the system (1.2) with $\phi(t_0) = c_1$, $\psi(t_0) = c_2$ exists on $[t_0, +\infty)$ and*

$$0 < \phi(t) \leq K(t, c_1, c_2), \quad 0 < \psi(t) \leq L(t, c_1, c_2), \quad t \geq t_0. \quad (3.12)$$

Proof. Let $(\phi_0(t), \psi_0(t))$ be a solution of the system (1.2) with $\phi_0(t_0) = c_1$ and $\psi_0(t_0) = c_2$. Suppose this solution is not continuable on $[t_0, +\infty)$. Let then $[t_0, T)$ be the maximum existence interval for $(\phi_0(t), \psi_0(t))$. Show that the inequalities (3.12) are satisfied for all $t \in [t_0, T)$. By (2.1) and (2.2) we have

$$\phi_0(t) > 0, \quad \psi_0(t) > 0, \quad t \in [t_0, T). \quad (3.13)$$

Note that (3.12) is valid at least for $t = t_0$. Then we can set:

$$\bar{t}_1 \equiv \sup\{t \in [t_0, T) : \phi_0(\tau) \leq K(\tau, c_1, c_2), \quad \tau \in [t_0, t]\},$$

$$\bar{t}_2 \equiv \sup\{t \in [t_0, T) : \psi_0(\tau) \leq L(\tau, c_1, c_2), \quad \tau \in [t_0, t]\},$$

Assume $\bar{t}_1 \leq \bar{t}_2$ (the proof in the case $\bar{t}_2 \leq \bar{t}_1$ by analogy) and assume (3.12) is not valid for all $t \in [t_0, T)$. Then $\bar{t}_1 < T$ and, therefore, there exists $t_2 \in (t_0, T)$ such that

$$\phi_0(t_2) > K(t_2, c_1, c_2), \quad (3.14)$$

$$\phi_0(t) \leq K(t, c_1, c_2) + \varepsilon, \quad t \in [t_0, t_2], \quad (3.15)$$

$$\psi_0(t) \leq L(t, c_1, c_2) + \varepsilon, \quad t \in [t_0, t_2]. \quad (3.16)$$

Consider the Riccati equations

$$y' + Q(t, \phi_0(t), \psi_0(t))y^2 + B(t, \phi_0(t), \psi_0(t))y - R(t, \phi_0(t), \psi_0(t)) = 0, \quad (3.17)$$

$$z' + R(t, \phi_0(t), \psi_0(t))z^2 - B(t, \phi_0(t), \psi_0(t))z - Q(t, \phi_0(t), \psi_0(t)) = 0, \quad (3.18)$$

By Lemma 2.1 from the conditions of the theorem and from (3.15), (3.16) it follows that for every $\gamma \geq 0$, Eq. (3.17) (Eq. (3.18)) has a non negative solution $y_0(t)$ ($z_0(t)$) on $[t_0, t_2)$ with $y_0(t_0) = \gamma$ ($z_0(t_0) = \gamma$). By (2.11) we have

$$y_0(t) = y_0(t_0) \exp\left\{-\int_{t_0}^t H_0(\tau) d\tau\right\} + \int_{t_0}^t \exp\left\{-\int_{\tau}^t H_0(s) ds\right\} R(\tau, \phi_0(\tau), \psi_0(\tau)) d\tau,$$

$t \in [t_0, t_2)$, where $H_0(t) \equiv Q(t, \phi_0(t), \psi_0(t))y(t) + B(t, \phi_0(t), \psi_0(t))$, $t \in [t_0, t_2)$. Multiply both sides of this equality by $Q(t, \phi_0(t), \psi_0(t))$ and integrate from t_0 to t . We obtain

$$\begin{aligned} \int_{t_0}^t Q(\tau, \phi_0(\tau), \psi_0(\tau)) y_0(\tau) d\tau &= \int_{t_0}^t Q(\tau, \phi_0(\tau), \psi_0(\tau)) \times \\ &\left[y_0(t_0) \exp\left\{-\int_{t_0}^{\tau} H_0(s) ds\right\} + \int_{t_0}^{\tau} \exp\left\{-\int_{\zeta}^{\tau} H_0(s) ds\right\} R(\zeta, \phi_0(\zeta), \psi_0(\zeta)) d\zeta \right] d\tau, \end{aligned}$$

$t \in [t_0, t_2)$. By (2.8) this equality with conditions of the theorem implies

$$\phi_0(t_2) \leq K(t_2, c_1, c_2),$$

which contradicts (3.14). The obtained contradiction implies

$$0 < \phi(t) \leq K(t, c_1, c_2), \quad 0 < \psi(t) \leq L(t, c_1, c_2), \quad t \in [t_0, T). \quad (3.19)$$

From here and from (3.13) it follows that $(\phi_0(t), \psi_0(t))$ is bounded on $[t_0, T)$. Then by Lemma 2.2 $[t_0, T)$ is not the maximum existence interval for $(\phi_0(t), \psi_0(t))$, which contradicts our assumption. The obtained contradiction together with (3.13) and (3.19) completes the proof of the theorem.

Definition 3.3. A solution $(\phi(t), \psi(t))$ of the system (1.1) is called conditionally stable, if there exists an one dimensional manifold $J \ni (\phi(t_0), \psi(t_0))$ such that for every $\varepsilon > 0$ there exists $\delta > 0$ such that each solution $(\tilde{\phi}(t), \tilde{\psi}(t))$ of the system (1.1) with $(\tilde{\phi}(t_0), \tilde{\psi}(t_0)) \in J$ and $|\tilde{\phi}(t_0) - \phi(t_0)| + |\tilde{\psi}(t_0) - \psi(t_0)| < \delta$ exists on $[t_0, +\infty)$ and satisfies to the inequality $|\tilde{\phi}(t) - \phi(t)| + |\tilde{\psi}(t) - \psi(t)| < \varepsilon$, $t \geq t_0$.

Corollary 3.1. Let for some $c_1 > 0$, $c_2 > 0$, $\varepsilon > 0$ and for every $t \geq t_0$, $\pm u \in (0, K(t, c_1, c_2) + \varepsilon]$, $\pm v \in (0, L(t, c_1, c_2) + \varepsilon]$ the inequalities $P(t, \pm u, \pm v) \leq P_0(t)$, $S(t, \pm u, \pm v) \leq S_0(t)$, $0 \leq Q(t, \pm u, \pm v) \leq Q_0(t)$, $0 \leq R(t, \pm u, \pm v) \leq R_0(t)$, $B_1(t) \leq B(t, \pm u, \pm v) \leq B_2(t)$ are valid (the expression $(t, \pm u, \pm v)$ means (t, u, v) and $(t, -u, -v)$)

and let $\lim_{u \rightarrow 0, u \neq 0} P(t, u, v)u = 0$, $\lim_{v \rightarrow 0, v \neq 0} S(t, u, v)v = 0$. Then every solution $(\phi(t), \psi(t))$ of the system (1.2) with $\phi(t_0) = \lambda c_1$, $\psi(t_0) = \lambda c_2$, $|\lambda| \leq 1$ exists on $[t_0, +\infty)$ and

$$0 < |\phi(t)| \leq \lambda K(t, c_1, c_2), \quad 0 < |\psi(t)| \leq \lambda L(t, c_1, c_2), \quad t \geq t_0. \quad (3.20)$$

Moreover if $K(t, c_1, c_2)$ and $L(t, c_1, c_2)$ are bounded on $[t_0, +\infty)$ (c_1 and c_2 are fixed) then the null solution $(0, 0)$ of the system (1.2) is conditionally stable.

Proof. The substitution $\phi \rightarrow -\phi$, $\psi \rightarrow -\psi$ in the system (1.2) reduces it in the following one

$$\begin{cases} \phi' = P(t, -\phi, -\psi)\phi + Q(t, -\phi, -\psi)\psi, \\ \psi' = R(t, -\phi, -\psi)\phi + S(t, -\phi, -\psi)\psi, \quad t \geq t_0. \end{cases} \quad (3.21)$$

Moreover it is obvious that

$$K(t, \lambda c_1, \lambda c_2) = \lambda K(t, c_1, c_2), \quad L(t, \lambda c_1, \lambda c_2) = \lambda L(t, c_1, c_2), \quad t \geq t_0, \quad \lambda \in \mathbb{R}$$

By Theorem 3.2 it follows from here and from (3.21) that under the conditions of the corollary every solution $(\phi(t), \psi(t))$ of the system (1.2) with $\phi(t_0) = \lambda c_1$, $\psi(t_0) = \lambda c_2$, $|\lambda| \leq 1$ exists on $[t_0, +\infty)$ and the estimates (3.20) are valid. If as a manifold, indicated in Definition 3.3, we take $J \equiv \{(\lambda c_1, \lambda c_2)\}$ then from (3.20) and from the boundedness of $K(t, c_1, c_2)$ and $L(t, c_1, c_2)$ it follows the conditionally stable of the null solution $(0, 0)$ of the system (1.2). The corollary is proved.

Let $\alpha(t)$, $\beta(t)$, $P_\delta(t)$, $Q_\delta(t)$, $R_\delta(t)$, $S_\delta(t)$, $B_{1,\delta}(t)$, $B_{2,\delta}(t)$ be real-valued locally integrable functions on $[t_0, +\infty)$, $\delta > 0$. For any $c_1 \neq 0$, $c_2 \neq 0$ set:

$$\begin{aligned} K_\delta(t, c_1, c_2) &\equiv c_1 \exp \left\{ \int_{t_0}^t P_\delta(s) ds + \int_{t_0}^t Q_\delta(\tau) \left[\frac{c_2}{c_1} \exp \left\{ - \int_{t_0}^\tau B_{1,\delta}(s) ds \right\} + \right. \right. \\ &\quad \left. \left. + \int_{t_0}^\tau \exp \left\{ - \int_{\zeta}^\tau B_{1,\delta}(s) ds \right\} R_\delta(\zeta) d\zeta \right] d\tau \right\}, \\ L_\delta(t, c_1, c_2) &\equiv c_2 \exp \left\{ \int_{t_0}^t S_\delta(s) ds + \int_{t_0}^t R_\delta(\tau) \left[\frac{c_1}{c_2} \exp \left\{ \int_{t_0}^\tau B_{2,\delta}(s) ds \right\} + \right. \right. \\ &\quad \left. \left. + \int_{t_0}^\tau \exp \left\{ \int_{\zeta}^\tau B_{2,\delta}(s) ds \right\} Q_\delta(\zeta) d\zeta \right] d\tau \right\}, \quad t \geq t_0. \end{aligned}$$

Theorem 3.3. *Let for some $c_1 > 0$, $c_2 > 0$, $\varepsilon > 0$ and for every $\delta \in (0, \varepsilon)$, $t \geq t_0$, $u \in [\delta, K_\delta(t, c_1, c_2) + \varepsilon]$, $v \in [\delta, L_\delta(t, c_1, c_2) + \varepsilon]$ the inequalities*

$$\alpha(t) \leq P(t, u, v) \leq P_\delta(t), \quad \beta(t) \leq S(t, u, v) \leq S_\delta(t) \quad (3.22)$$

$0 \leq Q(t, u, v) \leq Q_\delta(t)$, $0 \leq R(t, u, v) \leq R_\delta(t)$, $B_{1,\delta}(t) \leq B(t, u, v) \leq B_{2,\delta}(t)$ are valid. Then every solution $(\phi(t), \psi(t))$ of the system (1.2) with $\phi(t_0) = c_1$, $\psi(t_0) = c_2$ exists on $[t_0, +\infty)$ and

$$\phi(t) > 0, \quad \psi(t) > 0, \quad t \geq t_0. \quad (3.23)$$

Proof. Let $(\phi_0(t), \psi_0(t))$ be a solution of the system (1.2) with $\phi_0(t_0) = c_1$, $\psi_0(t_0) = c_2$ and let $[t_0, T)$ be its maximum existence interval. We must show that

$$T = +\infty. \quad (3.24)$$

By (2.1) and (2.2) from the conditions (3.22) it follows that

$$\phi_0(t) \geq c_1 \exp \left\{ \int_{t_0}^t \alpha(\tau) d\tau \right\}, \quad \psi_0(t) \geq c_2 \exp \left\{ \int_{t_0}^t \beta(\tau) d\tau \right\}, \quad t \in [t_0, T);$$

Suppose $T < +\infty$. Then since $\alpha(t)$ and $\beta(t)$ are locally integrable from the last inequalities it follows that

$$\phi_0(t) \geq \delta_0, \quad \psi_0(t) \geq \delta_0, \quad t \in [t_0, T)$$

for some $\delta_0 > 0$. Show that

$$0 < \phi_0(t) \leq K_{\delta_0}(t, c_1, c_2), \quad 0 < \psi_0(t) \leq L_{\delta_0}(t, c_1, c_2), \quad t \in [t_0, T). \quad (3.25)$$

From the conditions of the theorem is seen that these inequalities are satisfied at least for $t = t_0$. Then we can set:

$$\bar{t}_1 \equiv \sup\{t \in [t_0, T) : \phi_0(\tau) \leq K_{\delta_0}(\tau, c_1, c_2), \tau \in [t_0, t]\},$$

$$\bar{t}_2 \equiv \sup\{t \in [t_0, T) : \psi_0(\tau) \leq L_{\delta_0}(\tau, c_1, c_2), \tau \in [t_0, t]\},$$

Assume $\bar{t}_1 < \bar{t}_2$ (the proof in the case $\bar{t}_2 \leq \bar{t}_1$ by analogy) and assume (δ) is not valid. Then $\bar{t}_t < T$ and there exists $t_2 \in (t_0, T)$ such that

$$\phi_0(t_2) > K_{\delta_0}(t_2, c_1, c_2), \quad (3.26)$$

$$\delta_0 \leq \phi_0(t) \leq K_{\delta_0}(t, c_1, c_2) + \varepsilon, \quad t \in [t_0, t_2],$$

$$\delta_0 \leq \psi_0(t) \leq L_{\delta_0}(t, c_1, c_2) + \varepsilon, \quad t \in [t_0, t_2].$$

Further similar to the proof of Theorem 3.2 from here we come to the conclusion that

$$\phi_0(t_2) \leq K_{\delta_0}(t, c_1, c_2),$$

which contradicts (3.26). The obtained contradiction proves (3.25). It follows from (3.25) that $(\phi_0(t), \psi_0(t))$ is bounded on $[t_0, T)$. Then by Lemma 2.2 (3.24) is valid. from (3.24) and from (3.25) it follows (3.23). The theorem is proved.

In contrast to the conditions of Theorem 3.2, the conditions of Theorem 3.3 admit that the functions Q and R have singularities for $u = 0$ and $v = 0$, which, at part, is illustrated in the following example.

Example 3.2. Consider the Emden-Fowler's equation (see [8], p. 171)

$$(t^\rho \phi')' - t^\sigma \phi^n = 0, \quad t \geq t_0 > 0, \quad \rho, \sigma, n \in \mathbb{R}. \quad (3.27)$$

This equation is equivalent to the following system

$$\begin{cases} \phi' = t^{-\rho} \psi, \\ \psi' = (t^\sigma \phi^{n-1}) \phi, \quad t \geq t_0. \end{cases} \quad (3.28)$$

Here $P(t, u, v) = S(t, u, v) \equiv 0$, $Q(t, u, v) = t^{-\rho}$, $R(t, u, v) = t^\sigma u^{n-1}$. Consider the case $\rho \neq 1$, $\sigma \neq -1$, $2 + \sigma - \rho \neq 0$. At first we consider the superlinear case $n > 1$. Let us take $P_0(t) = S_0(t) \equiv 0$, $Q_0(t) = t^{-\rho}$, $R_0(t) = t^\sigma$, $t \geq t_0 > 0$. Then it is not difficult to show that

$$L(t, c_1, c_2) = c_2 \exp \left\{ \left(\frac{c_1}{c_2} - \frac{t_0^{1-\rho}}{1-\rho} \right) \frac{t^{\sigma+1} - t_0^{\sigma+1}}{\sigma+1} + \frac{t^{2+\sigma-\rho} - t_0^{2+\sigma-\rho}}{(1-\rho)(2+\sigma-\rho)} \right\}, \quad (3.29)$$

$t \geq t_0 > 0$. Obviously the inequality $0 \leq Q(t, u, v) \leq Q_0(t)$, $t \geq t_0 > 0$ for the system (3.28) holds for all $u, v \in \mathbb{R}$. Then on the basis of (3.29) it is easy to verify that the conditions of Theorem 3.2 for the system (3.28) are satisfied provided:

$$1. \quad \rho < 1, \quad \sigma < -1, \quad 2 + \sigma - \rho < 0, \quad c_1 > 0, \quad 0 < c_2 < 1, \quad \frac{c_1}{c_2} \leq \frac{t_0^{1-\rho}}{1-\rho},$$

or

$$2. \quad \rho < 1, \quad \sigma < -1, \quad 2 + \sigma - \rho < 0, \quad c_1 > 0, \quad 0 < c_1 < \exp \left\{ \frac{t_0^{\sigma+1}}{\sigma+1} \right\}, \quad \frac{c_1}{c_2} > \frac{t_0^{1-\rho}}{1-\rho}.$$

In the case $\rho = 1$, $\sigma \neq -1$ we have

$$L(t, c_1, c_2) = c_2 \exp \left\{ \frac{c_1}{c_2} \frac{t^{\sigma+1} - t_0^{\sigma+1}}{\sigma+1} + \int_{t_0}^t \tau^\sigma \ln \frac{\tau}{t_0} d\tau \right\}.$$

Hence the conditions of Theorem 3.2 for the system (3.28) are satisfied provided

3. $\rho = 1$, $\sigma < -1$, $c_1 > 0$, $0 < c_2 < \exp\left\{\frac{c_1 t_0^{\sigma+1}}{c_2 \sigma+1} - \int_{t_0}^{+\infty} \tau^\sigma \ln \frac{\tau}{t_0} d\tau\right\}$. In the sublinear ($0 < n < 1$) and extra sublinear ($n < 0$) cases (the cases $n = 0$ and $n = 1$ are trivial) we can take $\alpha(t) = \beta(t) = P_\delta(t) = S_\delta(t) \equiv 0$, $Q_\delta(t) = t^{-\rho}$, $R_\delta(t) = t^\sigma \delta^{n-1}$ and apply Theorem 3.3.

Remark 3.2. The case $\rho > 1$, $n \neq 1$ is reducible to the considered case $\rho < 1$ (namely to its simpler case $\rho = 0$) by the transformation (see [8], p. 143)

$$s = \frac{t^{\rho-1}}{\rho-1}, \quad \phi = (\rho-1)^{(\rho-\sigma-2)/[(\rho-1)(n-1)]} \frac{\psi}{s} \quad (3.30)$$

which reduces Eq. (3.22) to the following one

$$\psi'' - s^{\sigma_1} \psi^n = 0, \quad (3.31)$$

where $\sigma_1 \equiv \frac{\sigma+\rho}{\rho-1} - (n+3)$. Note that in the case $\rho < 1$, $n \neq 1$ Eq. (3.27) is also reducible to its simpler form by the another transformation (see [8], p. 144)

$$\psi'' - s^{\sigma_2} \psi^n = 0, \quad (3.32)$$

where $\sigma_2 \equiv \frac{\sigma+\rho}{1-\rho}$. It was established in [8, p. 144] that for each pair of certain parameters ρ and n ($(\sigma+\rho)(\sigma+n-1) > 0$) the equation

$$\phi'' - t^\sigma \phi^n = 0$$

has a global solution of the form

$$\phi(t) \equiv c(\sigma, n) t^{w(\sigma, n)}$$

where $c(\sigma, n) \equiv \left[\frac{(\sigma+2)(\sigma+n+1)}{(n-1)^2} \right]^{1/(n-1)}$, $w(\sigma, n) \equiv -\frac{\sigma+2}{n-1}$. Due to (3.30) – (3.32) this means, that it is established the existence of a single global solution of Eq. (3.27) for each ordered triple (ρ, σ, n) of certain parameters ρ , σ , n . Whereas, in Example 3.2 has been established the existence of two-parameter families of global solutions of Eq. (3.27) for certain parameters ρ , σ , n .

Remark 3.3. The Wintner's theorem (see [9], pp. 29,30, Theorem 5.1) is not applicable neither to the van der Pol's nor to the Emden-Fowler's equations. Therefore on the basis of examples 3.1 and 3.2 we conclude that Theorem 3.1 and Theorem 3.2 are not consequences of the Wintner's theorem.

Theorem 3.4. *Let the following conditions be satisfied.*

- 1) $0 \leq Q(t, u, v) \leq Q_0(t), \quad t \geq t_0, \quad u > 0;$
- 2) $P(t, u, v) \leq P_0(t), \quad t \geq t_0, \quad u > 0;$
- 3) $0 \leq F(t, u, v) \leq F_0(t), \quad t \geq t_0, \quad u > 0;$
- 4) $B(t, u, v) + \frac{F(t, u, v)}{u} \leq L_0(t), \quad t \geq t_0, \quad u > 0;$
- 5) $0 \leq R(t, u, v) + \frac{G(t, u, v)}{u} \leq M_0(t), \quad t \geq t_0, \quad u > 0,$

where $Q_0(t), P_0(t), F_0(t), L_0(t), M_0(t)$ are some real-valued continuous functions on $[t_0, +\infty)$. Then every solution $(\phi(t), \psi(t))$ of the system (1.1) with $\phi(t_0) > 0, \psi(t_0) \geq 0$ exists on $[t_0, +\infty)$ and

$$\phi(t) > 0, \quad \psi(t) \geq 0, \quad t \geq t_0. \quad (3.33)$$

Proof. Let $(\phi(t), \psi(t))$ be a solution of the system (1.1) with $\phi(t_0) > 0, \psi(t_0) \geq 0$ and let $[t_0, T)$ be the maximum existence interval for it. We must show that

$$T = +\infty. \quad (3.34)$$

At first show that

$$\phi(t) > 0, \quad t \in [t_0, T). \quad (3.35)$$

Suppose this is not true. Then since $\phi(t_0) > 0$ we have $\phi(t) > 0, \quad t \in [t_0, T_1)$ and

$$\phi(T_1) = 0 \quad (3.36)$$

for some $T_1 \in (t_0, T)$. By the second and third relation of (2.8) we have

$$\begin{aligned} \phi(t) = \phi(t_0) \exp \left\{ \int_{t_0}^t \left[P(\tau, \phi(\tau), \psi(\tau)) + y(\tau) Q(\tau, \phi(\tau), \psi(\tau)) \right] d\tau \right\} + \\ \int_{t_0}^t \exp \left\{ \int_{\tau}^t \left[P(s, \phi(s), \psi(s)) + y(s) Q(s, \phi(s), \psi(s)) \right] ds \right\} F(\tau, \phi(\tau), \psi(\tau)) d\tau, \end{aligned} \quad (3.37)$$

$t \in [t_0, T_1),$

$$\psi(t) = y(t)\phi(t), \quad t \in [t_0, T_1), \quad (3.38)$$

where by the third relation of (2.8) $y(t)$ is the solution of the Riccati equation

$$y' + Q(t, \phi(t), \psi(t))y^2 + \left(B(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\phi(t)} \right) y -$$

$$-R(t, \phi(t), \psi(t)) - \frac{G(t, \phi(t), \psi(t))}{\phi(t)} = 0, \quad t \in [t_0, T_1) \quad (3.39)$$

with $y(t_0) = \frac{\psi(t_0)}{\phi(t_0)} \geq 0$. Consider the Riccati equation

$$y' + Q(t, \phi(t), \psi(t))y^2 + \left(B(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\phi(t)} \right) y = 0, \quad t \in [t_0, T_1). \quad (3.40)$$

Obviously $y_1(t) \equiv 0$ is a solution of this equation. Then applying Theorem 2.1 to (3.39) and (3.40) on the basis of the conditions 1) and 5) we conclude that

$$y(t) \geq 0, \quad t \in [t_0, T_1). \quad (3.41)$$

Consider the Riccati equation

$$y' + Q(t, \phi(t), \psi(t))y^2 + L_0(t)y - M_0(t) = 0, \quad t \in [t_0, T_1).$$

Let $y_2(t)$ be a solution of this equation with $y_2(t_0) \geq y(t_0)$. Using Theorem 2.1 to this equation and to Eq. (3.39) and taking into account 1), 4), 5) and (3.41) we conclude that $y_2(t)$ exists on $[t_0, T_1)$ and

$$y(t) \leq y_2(t), \quad t \in [t_0, T_1). \quad (3.42)$$

This together with (3.37), (3.39), (3.41), and the conditions 1), 2), 3) implies that $\phi(T_1) > 0$, which contradicts (3.36). The obtained contradiction proves (3.35). It follows from (3.35) that $y(t)$ is continuable on $[t_0, T)$ as a solution of the equation

$$\begin{aligned} y' + Q(t, \phi(t), \psi(t))y^2 + \left(B(t, \phi(t), \psi(t)) + \frac{F(t, \phi(t), \psi(t))}{\phi(t)} \right) y - \\ - R(t, \phi(t), \psi(t)) - \frac{G(t, \phi(t), \psi(t))}{\phi(t)} = 0, \quad t \in [t_0, T). \end{aligned}$$

and by analogy of the proofs of the inequalities (3.41) and (3.42) it can be proved that

$$0 \leq y(t) \leq y_2(t), \quad t \in [t_0, T), \quad (3.43)$$

where $y_2(t)$ is a solution of the Riccati equation

$$y' + Q(t, \phi(t), \psi(t))y^2 + L_0(t)y - M_0(t) = 0, \quad t \in [t_0, T).$$

with $y_2(t_0) \geq y(t_0)$. By (2.8) we have

$$\begin{aligned}
\phi(t) &= \phi(t_0) \exp \left\{ \int_{t_0}^t \left[P(\tau, \phi(\tau), \psi(\tau)) + y(\tau) Q(\tau, \phi(\tau), \psi(\tau)) \right] d\tau \right\} + \\
&\int_{t_0}^t \exp \left\{ \int_{\tau}^t \left[P(s, \phi(s), \psi(s)) + y(s) Q(s, \phi(s), \psi(s)) \right] ds \right\} F(\tau, \phi(\tau), \psi(\tau)) d\tau, \quad t \in [t_0, T]. \\
\psi(t) &= y(t) \phi(t), \quad t \in [t_0, T].
\end{aligned} \tag{3.44}$$

This together with the conditions 1), 2), 3) implies that $(\phi(t), \psi(t))$ is bounded on $[t_0, T]$ provided $T < +\infty$. Hence if $T < +\infty$ then by Lemma 2.2 $[t_0, T)$ cannot be the maximum existence interval for $(\phi(t), \psi(t))$. Therefore (3.34) is valid. From (3.34), (3.35), (3.43) and (3.44) it follows (3.33). The theorem is proved.

Remark 3.4. *Theorem 3.3 remains valid if we replace its conditions 2), 3) by the following one*

$$P(t, u, v) + \frac{F(t, u, v)}{u} \leq P_0(t), \quad t \geq t_0, \quad u > 0.$$

which can be substantiated by using in the proof of Theorem 3.3 the first relation of (2.8) instead of the second one of (2.8).

Theorem 3.5. *Let $q_0(t)$, $r_0(t)$, $\alpha_0(t)$, $\beta_0(t)$ be real-valued locally integrable on $[a, b]$ ($t_0 \leq a < b < +\infty$) functions such that*

$$Q(t, u, v) \geq q_0(t) \geq 0, \quad R(t, u, v) \leq r_0(t) \leq 0, \quad \alpha_0(t) \leq B(t, u, v) \leq \beta_0(t), \quad t \in [a, b],$$

$$\int_a^b \min \left[q_0(t) \exp \left\{ - \int_a^t \alpha_0(\tau) d\tau \right\}, -r_0(t) \exp \left\{ \int_a^t \beta_0(\tau) d\tau \right\} \right] dt \geq \pi. \tag{3.45}$$

Then the system (1.2) is oscillatory on the interval $[a, b]$.

Proof. Let $(\phi(t), \psi(t))$ be a nontrivial solution of the system (1.2) on $[a, b]$. Consider the transcendental system of equations

$$\begin{cases} \rho \sin \theta = \phi(t) \exp \left\{ - \int_a^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\}, \\ \rho \cos \theta = \psi(t) \exp \left\{ - \int_a^t S(\tau, \phi(\tau), \psi(\tau)) d\tau \right\}, \end{cases} \quad t \in [a, b] \tag{3.46}$$

We set

$$\rho(t) \equiv \sqrt{\phi^2(t) \exp\left\{-2 \int_a^t P(\tau, \phi(\tau), \psi(\tau)) d\tau\right\} + \psi^2(t) \exp\left\{-2 \int_a^t S(\tau, \phi(\tau), \psi(\tau)) d\tau\right\}},$$

$$\theta(t) \equiv \theta_0 + \int_a^t \frac{[\phi(\tau)E(\tau)]'\psi(\tau) - \phi(\tau)E(\tau)\psi'(\tau)}{\phi^2(\tau)E^2(\tau) + \psi^2(\tau)} d\tau, \quad t \in [a, b],$$

$$\text{where } \theta_0 \equiv \begin{cases} \arctan \frac{\phi(a)}{\psi(a)}, & \psi(a) \neq 0, \\ \pi/2 - \arctan \frac{\psi(a)}{\phi(a)}, & \phi(a) \neq 0, \end{cases} \quad E(t) \equiv \exp\left\{\int_a^t B(\tau, \phi(\tau), \psi(\tau)) d\tau\right\},$$

$t \in [a, b]$. It is not difficult to verify that $(\rho(t), \theta(t))$ is an absolutely continuously solution of the system (3.46) on $[a, b]$. Then by (1.2) we have

$$\rho'(t) \sin \theta(t) + \rho(t) \theta'(t) \cos \theta(t) = Q(t, \phi(t), \psi(t)) \exp\left\{-\int_a^t B(\tau, \phi(\tau), \psi(\tau)) d\tau\right\} \rho(t) \cos \theta(t),$$

$$\rho'(t) \cos \theta(t) - \rho(t) \theta'(t) \sin \theta(t) = R(t, \phi(t), \psi(t)) \exp\left\{\int_a^t B(\tau, \phi(\tau), \psi(\tau)) d\tau\right\} \rho(t) \sin \theta(t).$$

almost everywhere on $[a, b]$. Multiply both sides of these relations by $\cos \theta(t)$ and both sides of the second of the ones by $-\sin \theta(t)$ and summarize. We obtain

$$\begin{aligned} \rho(t) \theta'(t) &= \left[Q(t, \phi(t), \psi(t)) \exp\left\{-\int_a^t B(\tau, \phi(\tau), \psi(\tau)) d\tau\right\} \cos^2 \theta - \right. \\ &\quad \left. - R(t, \phi(t), \psi(t)) \exp\left\{\int_a^t B(\tau, \phi(\tau), \psi(\tau)) d\tau\right\} \sin^2 \theta \right] \rho(t), \quad t \in [a, b]. \end{aligned}$$

Since $\rho(t) \neq 0$, $t \in [a, b]$ from here and from the conditions of the theorem it follows

$$\theta(b) - \theta(a) \geq \int_a^b \min \left[q_0(t) \exp\left\{-\int_a^t \alpha_0(\tau) d\tau\right\}, -r_0(t) \exp\left\{\int_a^t \beta_0(\tau) d\tau\right\} \right] dt \geq \pi.$$

This together with (3.46) implies that the system (1.2) is oscillatory on the interval $[a, b]$. The theorem is proved.

Corollary 3.2. *Let the conditions of Theorem 3.5, except of (3.45), be satisfied for $b = +\infty$ and let*

$$\int_a^{+\infty} \min \left[q_0(t) \exp \left\{ - \int_a^t \alpha_0(\tau) d\tau \right\}, -r_0(t) \exp \left\{ \int_a^t \beta_0(\tau) d\tau \right\} \right] dt = +\infty. \quad (3.47)$$

Then the system (1.2) is oscillatory.

Proof. If (3.47) is satisfied, then there exists an infinitely large sequence $a_1 < b_1 < a_2 < b_2 < \dots < a_n < b_n < \dots$ such that

$$\int_{a_n}^{b_n} \min \left[q_0(t) \exp \left\{ - \int_{a_n}^t \alpha_0(\tau) d\tau \right\}, -r_0(t) \exp \left\{ \int_{a_n}^t \beta_0(\tau) d\tau \right\} \right] dt \geq \pi, \quad n = 1, 2, \dots$$

Then by Theorem 3.5 the system (1.2) is oscillatory on each of the intervals $[a_n, b_n]$, $n = 1, 2, \dots$. Hence, the system (1.2) is oscillatory. The corollary is proved.

Theorem 3.5 as well as Corollary 3.2 are hypothetic (conditional) in the sense that the existence of solutions of the system (1.2) on appropriate intervals $[a, b]$ and $[a, +\infty)$ respectively is only assumed but not proved. In the regular case, when the "coefficients" P, Q, R, S have no singularities, this conditionality is omitted by Theorem 3.1. The next theorem weakens the hypothetic character of Theorem 3.5 and Corollary 3.2 (to some extent) in the singular case (when the "coefficients" P, Q, R, S may have singularities).

Theorem 3.6. *Let the following conditions be satisfied*

$$P(t, u, v) < 0, \quad Q(t, u, v) \geq 0, \quad R(t, u, v) \leq 0, \quad S(t, u, v) < 0$$

$$\frac{Q(t, u, v)}{P(t, u, v)} \leq M, \quad \frac{R(t, u, v)}{S(t, u, v)} \leq \frac{1}{M}$$

for some $M > 0$. Then for every $\alpha, \beta \in \mathbb{R}$ any solution $(\phi(t), \psi(t))$ of the system (1.2) with $\phi(t_0) = \alpha, \psi(t_0) = \beta$ exists on $[t_0, +\infty)$.

Proof. Let $(\phi(t), \psi(t))$ be a solution of the system (1.2) with $\phi(t_0) = \alpha, \psi(t_0) = \beta$ and let $[t_0, T)$ be the maximum existence interval for $(\phi(t), \psi(t))$. We must show that

$$T = +\infty.$$

Assume $\alpha^2 + \beta^2 \neq 0$. Then by (2.1) and (2.2) two cases are possible

I^0) there exists $t_1 \in [t_0, T)$ such that $\phi(t) \neq 0, \psi(t) \neq 0, t \in [t_1, T)$

II^0) there exists infinite sequence $t_0 < t_1 < \dots < T$ such that

$$\begin{aligned} & \begin{cases} \phi(t) > 0, & t \in [t_1, t_2), \\ \psi(t_1) = 0, & \psi(t) \leq 0, & t \in [t_1, t_2); \end{cases} \\ & \begin{cases} \phi(t_2) = 0, & \phi(t) \leq 0, & t \in [t_2, t_3), \\ \psi(t) \leq 0, & t \in [t_2, t_3); \end{cases} \\ & \begin{cases} \phi(t) \leq 0, & t \in [t_3, t_4), \\ \psi(t_3) = 0, & \psi(t) \geq 0, & t \in [t_3, t_4); \end{cases} \\ & \begin{cases} \phi(t_4) = 0, & \phi(t) \geq 0, & t \in [t_4, t_5), \\ \psi(t) > 0, & t \in [t_4, t_5); \end{cases} \\ & \begin{cases} \phi(t) > 0, & t \in [t_5, t_6), \\ \psi(t_5) = 0, & \psi(t) \leq 0, & t \in [t_5, t_6); \end{cases} \end{aligned}$$

so on (roughly speaking the point $(\phi(t), \psi(t))$ rotates around of the origin of coordinate plane ϕ, ψ in the clockwise direction infinitely many times when t varies from t_1 to T). Assume the case I^0) takes place. and $\phi(t_1) = 0, \psi(t_1) > 0$. Then by (2.1) and (2.2) from the conditions of the theorem we obtain respectively

$$\begin{aligned} |\phi(t)| &= \left| \phi(t_1) \exp \left\{ \int_{t_1}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} - \right. \\ & \left. - \int_{t_1}^t \left(\exp \left\{ - \int_{t_1}^s P(s, \phi(s), \psi(s)) ds \right\} \right)' \frac{Q(\tau, \phi(\tau), \psi(\tau))}{P(\tau, \phi(\tau), \psi(\tau))} d\tau \exp \left\{ \int_{t_1}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} \right| \leq \\ & \leq M |\phi(t_1)| \exp \left\{ \int_{t_1}^t P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} \left(\exp \left\{ \int_{t_1}^t -P(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} - 1 \right) \leq M |\phi(t_1)|, \\ & |\psi(t)| \leq |\psi(t_1)|, \quad t \in [t_1, T). \end{aligned}$$

Hence $(\phi(t), \psi(t))$ is bounded on $[t_0, T)$ provided $T < +\infty$. By similar way it can be shown the boundedness of $(\phi(t), \psi(t))$ on $[t_0, T)$ for $T < +\infty$ in the remaining subcases:

$$\begin{aligned} \phi(t_1) &> 0, & \psi(t_1) &< 0; \\ \phi(t_1) &< 0, & \psi(t_1) &> 0; \\ \phi(t_1) &< 0, & \psi(t_1) &< 0. \end{aligned}$$

Thus by Lemma 2.2 in the case I^0) for $T < +\infty$ the interval $[t_0, T)$ cannot be the maximum existence interval for $(\phi(t), \psi(t))$. In the case II^0) using relations (2.1) and (2.2) on the basis of the conditions of the theorem we obtain

$$|\phi(t)| \leq |\phi(t_1)|, \quad t \in [t_1, t_2),$$

$$\begin{aligned} |\psi(t)| &\leq \left| \exp \left\{ \int_{t_1}^t S(\tau, \phi(\tau), \psi(\tau)) d\tau \right\} \times \right. \\ &\quad \left. \times \int_{t_1}^t \left(\exp \left\{ - \int_{t_1}^s S(s, \phi(s), \psi(s)) ds \right\} \right)' \frac{R(\tau, \phi(\tau), \psi(\tau))}{S(\tau, \phi(\tau), \psi(\tau))} \phi(\tau) d\tau \right| \leq \frac{|\phi(t_1)|}{M}, \quad t \in [t_1, t_2). \end{aligned}$$

Using these estimates by similar way we can obtain successively

$$|\phi(t)| \leq |\phi(t_1)|, \quad |\psi(t)| \leq \frac{|\phi(t_1)|}{M}, \quad t \in [t_k, t_{k+1}), \quad k = 2, 3, \dots$$

It follows from here that $(\phi(t), \psi(t))$ is bounded on $[t_0, T)$ provided $T < +\infty$. Therefore in the case II^0) for $T + \infty$ the interval $[t_0, T)$ cannot be the maximum existence interval of $(\phi(t), \psi(t))$. If $\alpha = \beta = 0$ then either $\phi(t) = \psi(t) \equiv 0$ form a solution of the system, or $\phi^2(t_1) + \psi^2(t_1) \neq 0$ for some $t_1 > t_0$. In the last case by already proven $(\phi(t), \psi(t))$ exists on $[t_1, +\infty)$. Hence $(\phi(t), \psi(t))$ exists on $[t_0, +\infty)$. The theorem is proved.

Remark 3.5. *Theorem 3.6 remains valid if we replace in its conditions the relations*

$$\frac{Q(t, u, v)}{P(t, u, v)} \leq M, \quad \frac{R(t, u, v)}{S(t, u, v)} \leq \frac{1}{M}$$

by the following ones

$$\frac{Q(t, u, v)}{S(t, u, v)} \leq M, \quad \frac{R(t, u, v)}{P(t, u, v)} \leq \frac{1}{M}.$$

This can be substantiated by using in the proof of Theorem 3.6 the relations (2.3) and (2.4) instead of (2.1) and (2.2) respectively.

References

1. D. W. Jordan and P. Smith, Nonlinear ordinary differential equations. Oxford University Press. Fourth Edition, 2007.
2. N. N. Bogolubov, Yu. A. Mitropolski, Asymptotic methods in the theory of nonlinear oscillation. Mskow, "Mir", 1974, pp. 1 - 410.
3. Momeni M., Kurakis I., Mosley - Fard M., Shukla P. K. A Van der Pol - Mathieu equation for the dynamics of dust grain charge in dusty plasmas. J. Phys, A: Math.Theor., v. 40, 2007, F473 - F481.
4. N. Yamaoka, Oscillation criteria for second-order damped nonlinear differential equations with p -Laplacian, J. Math. Anal. Appl. 322 (2007) 932-948.
5. M. Kitano and K. Takasi, On a class of second order quasilinear ordinary differential equations, Hiroshima Math. J. 25 (1995), 321-355.
6. X. Wang, G. Song, Oscillation Theorems for a Class of Nonlinear Second Order Differential Equations with Damping, Advances in Pure Mathematics, 2013, 3, 226-233.
7. 4. G. A. Grigorianm On two comparison tests for second-order linear ordinary differential equations, Diff. Urav. 47 (2011), 1225(1240 (in Russian), Diff. Eq. 47 (2011), 1237) 1252 (in English).
8. 6. Bellman R. Stability theory of differential equations, Moscow, Izdatelstvo inostrannoj literatury, 1954 (New York, Toronto, London, McGRAW-HILL BOOK COMPANY, INC, 1053).
9. 7. Ph. Hartman, Ordinary Differential Equations, SIAM, Classics in Applied Mathematics literature, 1954, 38, Philadelphia 2002.