

GENERALIZED QUASI YAMABE GRADIENT SOLITONS AND APPLICATIONS

SİNEM GÜLER AND BÜLENT ÜNAL

ABSTRACT. The purpose of this article is to study generalized quasi Yamabe gradient solitons on warped product manifolds. First, we obtain some necessary and sufficient conditions for the existence of generalized quasi Yamabe gradient solitons equipped with the warped product structure. Then we study three important applications in the Lorentzian and the neutral settings for the particular class, called as gradient Yamabe soliton: We proved the existence of the non-trivial gradient Yamabe soliton on generalized Robertson-Walker spacetimes, standard static spacetimes and Walker manifolds.

1. INTRODUCTION

In recent years, self-similar solutions, referred as soliton solutions of some geometric flow equations, have begun to be introduced and studied as they appear to be possible singularity models. The most famous and studied classes of them are the Ricci solitons, defined as fixed points of the Ricci flow and in this area, significant progress has been made. Then, in the late 1980's, Hamilton introduced the Yamabe flow to prove the Yamabe problem, [1]. Basically, the Yamabe problem consists of looking for a metric on an $n \geq 3$ dimensional manifold such that its scalar curvature is constant. Thus, the Yamabe flow is defined as the metric $g(t)$ on a Riemannian manifold (M^n, g) satisfying $\frac{\partial g(t)}{\partial t} = -\tau g(t)$, where τ is the scalar curvature of M . The solution of this problem for two-dimensional case is already guaranteed by the Uniformization Theorem. For the studies on this concept we refer to [2–4]. Gradient Yamabe solitons are the solutions of this flow and defined as follows:

A pseudo Riemannian manifold (M^n, g) is said to be a gradient Yamabe soliton if there exists a smooth function φ on M and a constant λ satisfying

$$(1.1) \quad \text{Hess}(\varphi) = (\tau - \lambda)g.$$

If $\lambda > 0$, $\lambda < 0$ or $\lambda = 0$, then (M^n, g) is called a shrinking, expanding or steady gradient Yamabe soliton respectively. One of the most important results recorded as the solution to the Yamabe problem is given in [4] and

2010 *Mathematics Subject Classification.* 53C25, and 53C40.

Key words and phrases. Generalized quasi gradient Yamabe solitons, warped product manifolds, generalized Robertson-Walker spacetimes, standard static spacetimes, Walker manifolds.

it is proved that the metric of any compact Yamabe gradient soliton is a metric of constant scalar curvature.

In [5] it was proved that gradient Yamabe soliton admits a warped product structure, and this result enables to make possible studies in both Riemannian and Lorentzian settings.

After the Ricci flow theory is introduced and some progress was made, to classify the Riemannian manifolds and generalize the Ricci solitons, almost gradient Ricci solitons, quasi Einstein manifolds and generalized quasi Einstein manifolds have been introduced and studied extensively. For these concepts, we refer to [6–11] and many others. Analogously, in literature some generalizations of the self-similar solutions of Yamabe flow are also described. First the notion of quasi Yamabe gradient soliton in [12, 13] and then notion of generalized quasi Yamabe gradient soliton were introduced in [15]:

An n -dimensional pseudo-Riemannian manifold (M, g) is said to be a generalized quasi Yamabe gradient soliton if there exist smooth functions φ and μ on M and also a constant λ satisfying

$$(1.2) \quad \text{Hess}(\varphi) = (\tau - \lambda)g + \mu d\varphi \otimes d\varphi$$

where $d\varphi$ is the dual 1-form of $\nabla\varphi$ and τ is the scalar curvature of M . Here, φ is called as a potential function and the underlying generalized quasi Yamabe gradient soliton (briefly GQY) is denoted by $(M, g, \varphi, \mu, \lambda)$.

If φ is a constant function, we say that (M, g) is a trivial generalized quasi Yamabe gradient soliton. Otherwise, it will be called non-trivial. The restricted case where μ is constant is called quasi Yamabe gradient soliton. Moreover, if $\mu = 0$, then the equation (1.2) reduces to the fundamental equation of gradient Yamabe soliton (1.1).

In [12] it is proved that compact quasi Yamabe gradient soliton has constant scalar curvature and in [14], Wang has studied the special generalized quasi Yamabe gradient soliton in which $\mu = \frac{1}{m}$ for some constant $m > 0$ and proved that m -quasi Yamabe gradient soliton also has warped product structure in the region $\|\nabla\varphi\| \neq 0$. Moreover, the warping function is completely defined by the potential function of the soliton. Then in [15], Neto and Oliveira have extended these results to the generalized quasi Yamabe gradient solitons.

Inspired by these studies, we investigate some necessary and sufficient conditions for the existence of generalized quasi Yamabe gradient solitons equipped with the warped product structure. Then we study three important applications in the Lorentzian and the neutral settings for the class of gradient Yamabe solitons. We proved the existence of the non-trivial gradient Yamabe soliton on generalized Robertson-Walker spacetimes, standard static spacetimes and Walker manifolds.

2. WARPED PRODUCT GENERALIZED QUASI YAMABE GRADIENT SOLITONS

Assume that (B, g_B) and (F, g_F) are two pseudo-Riemannian manifolds of dimensions r and s , respectively. Let $\pi : B \times F \rightarrow B$ and $\sigma : B \times F \rightarrow F$ be the natural projection maps of the Cartesian product $B \times F$ onto B and F , respectively. Also, let $b : B \rightarrow (0, \infty)$ be a positive real-valued smooth function. The warped product manifold $M = B \times_b F$ is the the product manifold $B \times F$ equipped with the metric tensor defined by

$$(2.1) \quad g = \pi^*(g_B) \oplus (b \circ \pi)^2 \sigma^*(g_F)$$

where $*$ denotes the pull-back operator on tensors [16–18]. The function b is called the warping function of the warped product manifold $B \times_b F$, and the manifolds B and F are called base and fiber, respectively. In particular, if $b = 1$, then $B \times_1 F = B \times F$ is the usual Cartesian product manifold. For the sake of simplicity, throughout this paper, all relations will be written, without involving the projection maps from $B \times F$ to each component B and F as in $g = g_B \oplus b^2 g_F$.

Proposition 1. *Let (M, g) be an n -dimensional pseudo-Riemannian manifold. Then $(M, g, \varphi, \mu, \lambda)$ is a generalized quasi Yamabe gradient soliton if and only if*

$$(2.2) \quad \text{Hess}(\theta) = -\frac{\theta}{m}(\tau - \lambda)g$$

where $\mu = 1/m$ and $\theta = e^{-\varphi/m}$.

Proof. Introduce $\mu = 1/m$ and $\theta = e^{-\varphi/m}$. Then $\nabla\theta = -\frac{\theta}{m}\nabla\varphi$ and $d\theta = -\frac{\theta}{m}d\varphi$. So, for vector fields X and Y on M , we have:

$$\begin{aligned} \text{Hess}(\varphi)(X, Y) &= g(\nabla_X \nabla \varphi, Y) \\ &= g\left(\frac{m}{\theta^2} X(\theta) \nabla \theta - \frac{m}{\theta} \nabla_X \nabla \theta, Y\right) \\ &= \frac{m}{\theta^2} X(\theta) Y(\theta) - \frac{m}{\theta} \text{Hess}(\theta) \\ &= \frac{1}{m} d\varphi \otimes d\varphi - \frac{m}{\theta} \text{Hess}(\theta) \end{aligned}$$

Thus equation (1.2) can be reduced to (2.2). \square

As a direct corollary of the above proposition, we have:

Corollary 1. *Every generalized quasi Yamabe gradient soliton is conformal gradient soliton.*

Now, we can refer to the theorem of Cheeger and Colding [19], where the authors characterized the warped product manifolds and proved that any conformal gradient soliton satisfying $\text{Hess}(\theta) = kg$, for some function k is isometric to a warped product on some open interval. Thus, in view of (2.2) we conclude that (M^n, g) is isometric to the warped product $I \times_l N$, for some positive function l , where $I \subseteq \mathbb{R}$ is an open interval. Thus:

Corollary 2. *Every generalized quasi Yamabe gradient soliton admits the warped product structure $I \times_l N$, for some positive function l , where $I \subseteq \mathbb{R}$ is an open interval.*

This section presents the main result:

Theorem 1. *Let $M = B \times_b F$ be a warped product manifold equipped with the metric $g = g_B \oplus b^2 g_F$. Then $(M, g, \varphi, \mu, \lambda)$ is a generalized quasi Yamabe gradient soliton if and only if the followings hold:*

- (1) *the potential function φ depends only on the base manifold B ,*
- (2) *the potential function φ and the warping function b cannot be orthogonal,*
- (3) *the base manifold B is also a conformal gradient soliton,*
- (4) *the scalar curvature τ_F of the fiber manifold (F, g_F) is constant.*

Proof. Assume that $(M, g, \varphi, \mu, \lambda)$ is a generalized quasi Yamabe gradient soliton which is also a warped product. If X, Y are vector fields on B and V, W are vector fields on F , then apply the last proposition and obtain: $\text{Hess}(\theta)(X, V) = 0$ since $g(X, V) = 0$. On the other hand, by decomposing $\nabla\theta$ on the base and fiber,

$$\begin{aligned} \text{Hess}(\theta)(X, V) &= g(\nabla_X \nabla\theta, V) \\ &= g(\nabla_X \tan(\nabla\theta), V) + g(\nabla_X \text{nor}(\nabla\theta), V) \\ &= bX(b)g_F(\text{nor}(\nabla\theta), V). \end{aligned}$$

Thus, $X(b)g_F(\text{nor}(\nabla\theta), V) = 0$. If b is not constant, the last equation implies that $\text{nor}(\nabla\theta) = 0$. So, $\theta = e^{-\varphi/m}$ depends only on the base manifold B , that is, φ is only defined on B , $\varphi \in C^\infty(B)$.

Moreover, $g(V, W) = b^2 g_F(V, W)$ and $\nabla\theta = \tan(\nabla\theta)$ since $\text{nor}(\nabla\theta) = 0$.

$$\begin{aligned} \text{Hess}(\theta)(V, W) &= g(\nabla_V \nabla\theta, W) \\ &= g(\nabla_V \tan(\nabla\theta), W) \\ &= b \tan(\nabla\theta) g_F(V, W). \end{aligned}$$

Thus, by the last proposition,

$$b \tan(\nabla\theta) g_F(V, W) = -\frac{\theta}{m}(\tau - \lambda) b^2 g_F(V, W)$$

Equivalently,

$$\left(b \tan(\nabla\theta)(b) + \frac{b^2 \theta}{m}(\tau - \lambda) \right) g_F(V, W) = 0.$$

Contracting the last equation over V and W , we obtain:

$$sb(\tan(\nabla\theta)(b) + \frac{b\theta}{m}(\tau - \lambda)) = 0.$$

Noting that $\tan(\nabla\theta) = \nabla^B(\theta)$ since $\theta \in \mathcal{C}^\infty(B)$, we have:

$$g_B(\nabla^B\theta, \nabla^B b) = (\lambda - \tau) \frac{b\theta}{m}.$$

Finally,

$$\begin{aligned} \text{Hess}(\theta)(X, Y) &= g(\nabla_X \nabla\theta, Y) \\ &= g_B(\nabla_X \tan(\nabla\theta), Y) \\ &= \text{Hess}^B(\theta)(X, Y) \end{aligned}$$

since $\theta \in \mathcal{C}^\infty(B)$, i.e, $\tan(\nabla\theta) = \nabla^B(\theta)$. By the last proposition, we can have:

$$\text{Hess}^B(\theta) = \frac{\theta}{m}(\lambda - \tau)g_B.$$

Hence, B is conformal gradient soliton and scalar curvature τ_F of the fiber manifold (F, g_F) is constant. The converse statement is straightforward. \square

Combining the above results, we can state that:

Corollary 3. *Let $(M, g, \varphi, \mu, \lambda)$ be a generalized quasi Yamabe gradient soliton satisfying the conditions (1)-(4) of Theorem 1. Then it admits the multiply warped product structure $I \times_l N \times_b F$, for some positive functions l and b , where $I \subseteq \mathbb{R}$ is an open interval.*

3. APPLICATIONS

3.1. Gradient Yamabe Soliton on Generalized Robertson-Walker Space-times. We first define generalized Robertson-Walker space-times. Let (F, g_F) be an s -dimensional Riemannian manifold and $b : I \rightarrow (0, \infty)$ be a smooth function. Then $(s + 1)$ -dimensional product manifold $I \times_b F$ furnished with the metric tensor

$$\bar{g} = -dt^2 \oplus b^2 g_F$$

is called a generalized Robertson-Walker space-time and is denoted by $M = I \times_b F$ where I is an open, connected subinterval of \mathbb{R} and dt^2 is the Euclidean metric tensor on I . This structure was introduced to extend Robertson-Walker space-times [20, 21] and have been studied by many authors, such as [22–24]. From now on, we will denote $\frac{\partial}{\partial t} \in \mathfrak{X}(I)$ by ∂_t to state our results in simpler forms.

We will apply our main result Theorem 1. Assume that $\varphi \in \mathcal{C}^\infty(I)$ is a potential function for a generalized Robertson-Walker space-time of the form $M = I \times_b F$.

The equation $(\tau - \lambda)g_{ij} = \text{Hess}_{ij}$ yields

$$(3.1) \quad \begin{cases} \varphi'' = -(\tau - \lambda), \\ b'\varphi' = (\tau - \lambda)b. \end{cases}$$

Thus $b\varphi'' = -b'\varphi'$. By solving the last ODE, as $b \neq 0$, we have:

$$\varphi(t) = \alpha \int_{t_0}^t \frac{1}{b(\bar{t})} d\bar{t} \quad \text{for some } \alpha \in \mathbb{R}.$$

Hence, we can state that:

Theorem 2. *A generalized Robertson-Walker space-time of the form $M = I \times_b F$ is a gradient Yamabe soliton with the potential function φ given by*

$$\varphi(t) = \alpha \int_{t_0}^t \frac{1}{b(\bar{t})} d\bar{t} \quad \text{for some } \alpha \in \mathbb{R}.$$

3.2. Gradient Yamabe Soliton on Standard Static Space-times. We begin by defining standard static space-times. Let (F, g_F) be an s -dimensional Riemannian manifold and $f : F \rightarrow (0, \infty)$ be a smooth function. Then $(s + 1)$ -dimensional product manifold $_f I \times F$ furnished with the metric tensor

$$g = -f^2 dt^2 \oplus g_F$$

is called a standard static space-time and is denoted by $M =_f I \times F$ where I is an open, connected subinterval of \mathbb{R} and dt^2 is the Euclidean metric tensor on I .

Note that standard static space-times can be considered as a generalization of the Einstein static universe [25–28] and many spacetime models that characterize the universe and the solutions of Einstein's field equations are known to have this structure.

Again we apply Theorem 1. Suppose that $\varphi \in C^\infty(F)$ is a potential function for a standard static space-time of the form $M =_f I \times F$.

The equation $(\tau - \lambda)g_{ij} = \text{Hess}_{ij}$ yields

$$(3.2) \quad \begin{cases} \nabla \varphi(f) = (\tau - \lambda)f, \\ \text{Hess}_F(\varphi) = (\tau - \lambda)g_F. \end{cases}$$

By contracting the last equation over F , we have $\Delta_F(f) = s(\tau - \lambda)$. Then we obtain:

$$\Delta_F(\varphi) = \frac{s}{f} \nabla \varphi(f).$$

Hence, we conclude that:

Theorem 3. *A standard static space-time of the form $M =_f I \times F$ is a gradient Yamabe soliton with the potential function φ given by*

$$\Delta_F(\varphi) = \frac{s}{f} \nabla \varphi(f).$$

3.3. Gradient Yamabe Soliton on 3-dimensional Walker Manifolds. Three-dimensional Lorentzian manifolds admitting a parallel degenerate line field is called Walker manifold, [29, 30]. Then there exist local coordinates (t, x, y) such that with respect to the local frame fields $\{\partial_t, \partial_x, \partial_y\}$ the Lorentzian metric tensor is of the form:

$$(3.3) \quad g = 2dt dy + dx^2 + \phi(t, x, y) dy^2,$$

for some function $\phi(t, x, y)$. The restricted case of Walker manifold where ϕ described as a function of only x and y is called strictly Walker manifold and these classes of Walker manifolds are geodesically complete. Also, it is known that a Walker manifold is Einstein if and only if it is flat, [29].

Now, we will investigate conditions on this particular class of manifolds to have gradient Yamabe solitons, that is,

$$(3.4) \quad \text{Hess}(f)_{ij} = (\tau - \lambda)g_{ij}$$

where f is a potential function.

Note that this equation implies that

$$\frac{\Delta(f)}{3} = \tau - \lambda.$$

By using the metric (3.3) and straightforward computations, we have:

$$(3.5) \quad \begin{cases} \text{Hess}(f)_{tt} = f_{tt}, \\ \text{Hess}(f)_{tx} = f_{tx}, \\ \text{Hess}(f)_{ty} = f_{ty} - \frac{1}{2}\phi_t f_t, \\ \text{Hess}(f)_{xx} = f_{xx}, \\ \text{Hess}(f)_{xy} = f_{xy} - \frac{1}{2}\phi_x f_t, \\ \text{Hess}(f)_{yy} = f_{yy} - \frac{1}{2}\phi\phi_t f_t + \frac{1}{2}\phi_t f_y. \end{cases}$$

Moreover,

$$(3.6) \quad \Delta(f) = -\phi f_{tt} + 2f_{ty} - \phi_t f_t + f_{xx}$$

By combining these, we get:

$$(3.7) \quad \tau - \lambda = \frac{1}{3} \left(-\phi f_{tt} + 2f_{ty} - \phi_t f_t + f_{xx} \right)$$

By applying equations (3.4), (3.5) and (3.7), we obtain the following system of PDEs:

$$(3.8) \quad \begin{cases} f_{tt} = 0, \\ f_{tx} = 0, \\ f_{xy} = \frac{1}{2}\phi_x f_t, \\ f_{xx} - f_{ty} = -\frac{1}{2}\phi_t f_t, \\ f_{yy} - f_{xx} = \frac{1}{2}\phi\phi_t f_t - \frac{1}{2}\phi_t f_y. \end{cases}$$

Notice first that $f_{tt} = 0$ and $f_{tx} = 0$ imply that $f(t, x, y) = tb(y) + c(x, y)$ for some functions b and c .

Thus $f_{xy} = \frac{1}{2}\phi_x f_t$ turns out to be $2c_{xy}(x, y) = \phi_x b(y)$. Substituting them in the last two equations of the system (3.8), we obtain

$$(3.9) \quad 2c_{xx}(x, y) + \phi_t b(y) = 2b'(y)$$

Differentiating the both sides of equation (3.9) with respect to t , we get $\phi_{tt}b(y) = 0$ this implies that $\phi_{tt} = 0$ or $b(y) = 0$.

• *Case 1:* $\phi_{tt} = 0$. Then the metric becomes Ricci-flat so we may assume that this case should be excluded.

- *Case 2:* If $\phi_{tt} \neq 0$, then $b(y) = 0$, i.e, $f(t, x, y) = c(x, y)$.

Thus $c_{xx} = 0$ and $c_{xy} = 0$ imply that $c(x, y) = \kappa x + \eta(y)$, for some $\kappa \in \mathbb{R}$. Hence $2\eta''(y) + \phi_t \eta'(y) = 0$. So, $\phi_t = -2\ln[\eta'(y)]$ and then $\phi(t, x, y) = -2t\ln[\eta'(y)] + \zeta(x, y)$. In this case, the potential function is given by: $f(x, y) = \kappa x + \eta(y)$. Hence we obtain the main result of this section:

Theorem 4. *Let (M, g) be a 3-dimensional Lorentzian Walker manifold equipped with metric:*

$$g = 2dt dy + dx^2 + \phi(t, x, y)dy^2.$$

Then (M, g) is a gradient Yamabe soliton if and only if

- (1) *the potential function of the soliton structure is given by*
 $f(x, y) = \kappa x + \eta(y)$ *for some $\kappa \in \mathbb{R}$.*
- (2) *the function defined in the Walker metric is given by*
 $\phi(t, x, y) = -2t\ln[\eta'(y)] + \zeta(x, y)$.

Note that by Equation (3.6), we can get: $\Delta(f) = 0$. In order to have a non-trivial potential function we may assume that M is non-compact due to Hopf's Lemma. Thus, we have:

Corollary 4. *Let (M, g) be a 3-dimensional Lorentzian Walker manifold equipped with metric (3.3). Then the gradient Yamabe soliton structure on (M, g) characterized by the conditions (1) and (2) of Theorem 4 is non-compact.*

3.4. Gradient Yamabe Soliton on 4-dimensional Walker Manifolds.

Before finishing the work, let's finally construct the gradient Yamabe soliton structure on the 4-dimensional Walker manifold.

A 4-dimensional Walker manifold is a triple (M, g, D) consisting of an indefinite metric g and an 2-dimensional parallel null plane D and in this case g has neutral signature $(-, -, +, +)$ and in suitable coordinates (x, y, z, t) such that with respect to the local frame fields $\{\partial_x, \partial_y, \partial_z, \partial_t\}$ it can be given by [29]

$$(3.10) \quad g = 2dx dz + 2dy dt + a(x, y, z, t)dz^2 + 2c(x, y, z, t)dz dt + b(x, y, z, t)dt^2,$$

for some functions $a(x, y, z, t), b(x, y, z, t), c(x, y, z, t)$ and $D = \langle \partial_x, \partial_y \rangle$. The case where $c(x, y, z, t) = 0$ was studied and locally conformally flatness of this special metric was investigated in [31].

Here, we consider the restricted case of Walker manifold where $a = c = 0$ and b described as a function of only t so the metric (3.10) reduces to the form

$$(3.11) \quad g = 2dx dz + 2dy dt + b(t)dt^2,$$

Now, we will investigate conditions on this particular class of manifolds to have gradient Yamabe solitons, that is,

$$(3.12) \quad \text{Hess}(f)_{ij} = (\tau - \lambda)g_{ij}$$

where f is a potential function.

Note that this equation implies that

$$\frac{\Delta(f)}{4} = \tau - \lambda.$$

By using the metric (3.11) and straightforward computations, we have:

$$(3.13) \quad \begin{cases} \text{Hess}(f)_{xx} = f_{xx}, & \text{Hess}(f)_{xy} = f_{xy}, \\ \text{Hess}(f)_{xz} = f_{xz}, & \text{Hess}(f)_{xt} = f_{xt}, \\ \text{Hess}(f)_{yy} = f_{yy}, & \text{Hess}(f)_{yz} = f_{yz}, \\ \text{Hess}(f)_{yt} = f_{yt}, & \text{Hess}(f)_{zz} = f_{zz}, \\ \text{Hess}(f)_{zt} = f_{zt}, & \text{Hess}(f)_{tt} = f_{tt} - \frac{1}{2}b_t f_y. \end{cases}$$

Moreover,

$$(3.14) \quad \Delta(f) = 2f_{xz} - bf_{yy} + 2f_{yt}.$$

By combining these, we get:

$$(3.15) \quad \tau - \lambda = \frac{1}{4} \left(2f_{xz} - bf_{yy} + 2f_{yt} \right)$$

By applying equations (3.12), (3.13) and (3.15), we obtain the following system of PDEs:

$$(3.16) \quad \begin{cases} f_{xx} = f_{xy} = f_{yy} = f_{yz} = f_{zz} = 0, \\ f_{xt} = f_{zt} = 0, \\ f_{xz} = f_{yt} = \frac{\Delta(f)}{4}, \\ f_{tt} - \frac{1}{2}b_t f_y = b \frac{\Delta(f)}{4}. \end{cases}$$

Notice first that $f_{xy} = 0$, $f_{xx} = 0$, $f_{yz} = 0$ and $f_{yy} = 0$ imply that $f(x, y, z, t) = x\beta(z, t) + yA(t) + B(z, t)$ for some functions β, A and B . Thus from $f_{xz} = f_{yt}$, we have $\beta_z(z, t) = A'(t)$. Since $f_{zz} = 0$ and $f_{xt} = 0$, $A'(t) = C'(z)$ which yields

$$(3.17) \quad A(t) = c_0 t + c_1, \quad C(z) = c_0 z + c_2, \quad \text{where } c_0, c_1, c_2 \in \mathbb{R}.$$

Also, from $f_{xt} = 0$ and $f_{zt} = 0$, we obtain $\beta(y, z) = \kappa z + \mu$ and $B(z, t) = c_3 z + E(t)$, respectively where $c_0 \in \mathbb{R}$. As a result, the potential function is given by

$$(3.18) \quad f(x, y, z, t) = x(c_0 z + c_2) + y(c_0 z + c_1) + c_3 z + E(t).$$

Substituting this in the last equation of the system (3.16), we obtain

$$(3.19) \quad 2E''(t) - b_t(c_0 t + c_1) = 2c_0 b.$$

Integrating (3.19) with respect to t and using integration by parts for the second term, we get $2E'(t) - b(c_0 t + c_1) = c_0 \int_{t_0}^t b(\bar{t}) d\bar{t}$.

Hence we obtain the main result of this section:

Theorem 5. *Let (M, g) be a 4-dimensional Walker manifold equipped with metric:*

$$(3.20) \quad g = 2dx dz + 2dy dt + b(t)dt^2.$$

Then (M, g) is a gradient Yamabe soliton if and only if its potential function is defined by

$$f(x, y, z, t) = x(c_0 z + c_2) + y(c_0 z + c_1) + c_3 z + E(t)$$

where $c_i \in \mathbb{R}$, $(i = 0, 1, 2, 3)$ and the metric function $b(t)$ and $E(t)$ are related by

$$2E'(t) - b(c_0 t + c_1) = c_0 \int_{t_0}^t b(\bar{t}) d\bar{t}.$$

Moreover, $\Delta(f) = 4c_0$.

If in additionally, M is compact, then by applying the Divergence Theorem, $c_0 = 0$ and so get conclude that:

Corollary 5. *Let (M, g) be a 4-dimensional Walker manifold equipped with metric (3.20). Then (M, g) is a compact gradient Yamabe soliton if and only if its potential function is defined by*

$$f(x, y, z, t) = c_2 x + c_1 y + c_3 z + \frac{c_1}{2} \int_{t_0}^t b(\bar{t}) d\bar{t}.$$

where $c_i \in \mathbb{R}$, $(i = 1, 2, 3)$.

REFERENCES

- [1] R.S. Hamilton, *The Ricci flow on surfaces*, in: *Mathematics and General Relativity*, Contemp. Math., **71** (1988) 237–262. [1](#)
- [2] S. Brendle, *Convergence of the Yamabe flow for arbitrary initial energy*, J. Differential Geom. **69** (2) (2005) 217–278. [1](#)
- [3] S. Brendle, *Convergence of the Yamabe flow in dimension 6 and higher*, Invent. Math., **170** (3) (2007) 541–576. [1](#)
- [4] P. Daskalopoulos, N. Sesum, *The classification of locally conformally flat Yamabe solitons*, Advances in Mathematics **240** (2013) 346–369. [1](#), [1](#)
- [5] H.-D. Cao, X. Sun, Y. Zhang, *On the structure of gradient Yamabe solitons*, Math. Res. Lett. **19** (2012) 767–774. [1](#)
- [6] H.-D. Cao, Q. Chen, *On locally conformally flat gradient steady Ricci solitons*, Trans. Am. Math. Soc., (2012) 23772391. [1](#)
- [7] J.S. Case, Y. Shu, G. Wei, *The non-existence of quasi Einstein metrics*, Pac. J. Math. **248**(2) (2010) 277284. [1](#)
- [8] J. S. Case, Y. Shu, G. Wei, *Rigidity of quasi Einstein metrics*, Differ. Geom. Appl. **29** (2011) 93100. [1](#)
- [9] C.-L. Bejan, S. Güler, *Laplace, Einstein and related equations*, Mediter. J.Math. **16**19, (2019) <https://doi.org/10.1007/s00009-018-1283-9>. [1](#)
- [10] G. Catino, *Generalized quasi-Einstein manifolds with harmonic Weyl tensor*, Math. Z. **271**(3) (2012) 751756. [1](#)
- [11] S. Altay Demirbağ, S. Güler, *Rigidity of (m, λ) -quasi Einstein manifolds*, Match. Nachr., **290**(1415) (2017) 21002110. [1](#)
- [12] G. Huang, H. Li, *On a classification of the quasi Yamabe gradient solitons*, Methods Appl. Anal. **21** (2014) 379390. [1](#), [1](#)

- [13] B. Leandro Neto, *A note on (anti-)self dual quasi Yamabe gradient solitons*, Results Math. (2015) 17. [1](#)
- [14] L.F. Wang, *On noncompact quasi Yamabe gradient solitons*, Differ. Geom. Appl. **31** (2013) 337348. [1](#)
- [15] B. Leandro Neto, H. P. Oliveira, *Generalized quasi Yamabe gradient solitons*, Diff. Geo. Appl., **49** (2016) 167–175. [1](#), [1](#)
- [16] R. L. Bishop and B. O’Neill, *Manifolds of negative curvature*, Trans. Amer. Math. Soc. **145** (1969) 1-49. [2](#)
- [17] B. O’Neill, *Semi-Riemannian Geometry with Applications to Relativity*, Academic Press Limited, London, 1983. [2](#)
- [18] S. Shenawy and B. Ünal, *2–Killing vector fields on warped product manifolds*, International Journal of Mathematics, **26** (2015) 1550065 [2](#)
- [19] J. Cheeger and T. Colding, *Lower bounds on Ricci curvature and the almost rigidity of warped products*, Ann. Math., **144** (1996), 189237. [2](#)
- [20] M. Sánchez, *On the Geometry of Generalized Robertson-Walker Spacetimes: geodesics*, Gen. Relativ. Gravitation **30** (1998) 915-932. [3.1](#)
- [21] M. Sánchez, *On the Geometry of Generalized Robertson-Walker Spacetimes: Curvature and Killing fields*, J. Geom. Phys. **31** (1999) 1-15. [3.1](#)
- [22] C. A. Mantica, L. G. Molinari, U. C. De, *A condition for a perfect fluid space-time to be a generalized Robertson-Walker space-time*, J. Math. Phys. 57 (2) (2016) 022508. [3.1](#)
- [23] C. A. Mantica, Y. J. Suh, U. C. De, *A note on generalized Robertson-Walker space-times*, Int. J. Geom. Meth. Mod. Phys. 13 (2016) 1650079. [3.1](#)
- [24] B.-Y. Chen, *A simple characterization of generalized Robertson-Walker space-times*, Gen. Relativ. Gravit. 46 (2014) 1833. [3.1](#)
- [25] D.E. Allison, *Energy conditions in standard static space-times*, General Relativity and Gravitation, **20** (1988) 115-122. [3.2](#)
- [26] D.E. Allison, *Geodesic Completeness in Static Space-times*, Geometriae Dedicata **26** (1988) 85-97. [3.2](#)
- [27] D.E. Allison and B. Ünal, *Geodesic Structure of Standard Static Space-times*, Journal of Geometry and Physics **46** (2003) 193-200. [3.2](#)
- [28] A. L. Besse, *Einstein Manifolds*, Classics in Mathematics, Springer-Verlag, Berlin, (2008). [3.2](#)
- [29] A. G.Walker, *Canonical form for a Riemannian space with a parallel field of null planes*, Quart. J. Math. Oxford, **21** (1950) 69-79. [3.3](#), [3.3](#), [3.4](#)
- [30] M. Chaichi, E. Garcia-Rio and M. E. Vazquez-Abal, *Three-dimensional Lorentz manifolds admitting a parallel null vector field*, J. Phys. A: Math. Gen., **38** (2005) 841–850. [3.3](#)
- [31] M. Chaichi, E. Garca-Ro, and Y. Matsushita, *Curvature properties of four-dimensional Walker metrics*, Classical and Quantum Gravity, **22** (2005) 559–577. [3.4](#)

(S. Güler) DEPARTMENT OF INDUSTRIAL ENGINEERING, ISTANBUL SABAHATTIN ZAIM UNIVERSITY, HALKALI, ISTANBUL, TURKEY.

Email address: sinem.guler@izu.edu.tr

(B. Ünal) DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, BILKENT, 06800 ANKARA, TURKEY

Email address: bulentunal@mail.com