# On extended 1-perfect bitrades<sup>\*</sup>

Evgeny A. Bespalov and Denis S. Krotov

#### Abstract

We prove the equivalence of several definitions of extended 1-perfect bitrades in the Hamming graph H(n,q) and prove the nonexistence of such bitrades for odd n.

### 1. Introduction

In this note, we discuss several different definitions of an extended 1-perfect bitrade in Hamming graphs H(n,q), q > 2, prove their equivalence, and prove the nonexistence of extended 1-perfect bitrades for odd n, generalizing some of the results of [3] on the nonexistence of extended 1-perfect codes. As usual, for different combinatorial objects like some classes of codes, designs, orthogonal arrays, etc., bitrades are defined as a generalization of the difference pair  $(C_0 \setminus C_1, C_1 \setminus C_0)$  of two objects  $C_0$  and  $C_1$  from the initial class. The main question is the following: which property of the initial class has to be generalized to form the definition of a bitrade. The five definitions of extended 1-perfect bitrades considered below reflect different definitions of an extended 1-perfect code as

- a code whose projection in any direction (puncturing in any coordinate) gives a 1-perfect code (this definition reflects a traditional meaning of the concept "extended code");
- a uniformly packed (in the wide sense [2]) code, with certain parameters;
- a completely regular code [5], with certain parameters;
- a diameter-3 perfect code [1];
- a distance-4 code with certain spectral constraints, namely, the characteristic function of code belongs to the sum of three eigenspaces of H(n,q), with the eigenvalues (q-1)n (the maximum eigenvalue of H(n,q)), q-2, and -n (the minimum eigenvalue of H(n,q)); these spectral restrictions can be also expressed in terms of the components of the dual distance distribution.

We note that our definitions are not in agree with known definition of extended 1perfect bitrades in the binary Hamming graph H(n, 2). The reason is that H(n, 2), or *n*-cube, is a bipartite graph, and it is natural to require that the components of

<sup>\*</sup>Funded by Russian Science Foundation, Grant 18-11-00136

the bitrade lie in one of its parts, either even or odd (which also allows to consider bitrades as objects in the halved *n*-cube). Relaxing this restriction leads to some degenerate cases, which are not interesting from theoretical point of view. Of course, H(n,q) can also be considered as a *q*-partite graph, but the parts are not uniquely defined, and we cannot guarantee for an extended 1-perfect code that it lies in one of them; neither it is natural to require this from a bitrade.

We finish the introduction by the observation that extended 1-perfect codes and extended 1-perfect bitrades do exist in  $H(2^k + 2, 2^k)$ , so the equivalences proved are about the objects from non-empty class.

### 2. Preliminaries

Given a graph G denote by vG its vertex set. A code in G is an arbitrary nonempty subset of vG.

Denote  $e_i^a = (0, \ldots, 0, a, 0, \ldots, 0)$ , where  $a \in \mathbb{Z}_q$  is placed in the *i*-th position. Denote by  $B(x) = \{z : d(x, z) \leq 1\}$  the radius-1 ball with center x. Denote by  $S_i(x)$  the radius-i sphere with center x, i.e.  $S_i(x) = \{y : d(x, y) = i\}$ . The set  $C_{x,i} = \bigcup_{a \in \mathbb{Z}_q} B(x + e_i^a)$  is called cylinder.

A pair  $(T_+, T_-)$  of disjoint subsets of vH(n, q) is called a *perfect bitrade* in H(n, q) if  $|B(x) \cap T_+| = |B(x) \cap T_-| \le 1$  for any  $x \in vH(n, q)$ .

**Lemma 1.** If  $(T_+, T_-)$  is a perfect bitrade in H(n, q), then  $T_+$  and  $T_-$  are codes with code distance 3.

For a vertex  $x = (x_1, \ldots, x_n) \in {}_{\vee}H(n, q)$  denote by  $x_i^a$  the vertex  $(x_1, \ldots, x_{i-1}, a, x_i, \ldots, x_n)$ of H(n + 1, q). Let C be a code in H(n + 1, q). The set  $\{x \in {}_{\vee}H(n, q) : x_i^a \in C$  for some  $a \in \mathbb{Z}_q\}$  is called the *i*-projection of C (or simply, projection). Let  $(T_+, T_-)$ be a pair of disjoint sets of vertices of H(n + 1, q), and let  $A_+$  and  $A_-$  be *i*-projections of  $T_+$  and  $T_-$  respectively. The pair  $(T'_+, T'_-)$ , where  $T'_+ = A_+ \setminus A_-$  and  $T'_- = A_- \setminus A_+$ , is called the *i*-projection (or simply, projection) of  $(T_+, T_-)$ .

A function  $f: vH(n,q) \to \mathbb{C}$  is called an *eigenfunction* of H(n,q) with eigenvalue  $\lambda$  if  $Af = \lambda f$ ,  $f \not\equiv 0$ , where A is the adjacency matrix of H(n,q). The Hamming graph H(n,q) has the following eigenvalues:  $\lambda_i = n(q-1) - iq$ ,  $i = 0, 1, \ldots, n$ . Denote by  $U_i(n,q) = \{f: Af = \lambda_i f\}$  the eigenspace of eigenfunctions of H(n,q) with eigenvalue  $\lambda_i$ . For functions  $f, g: vH(n,q) \to \mathbb{C}$  define  $(f,g) = \sum_{x \in vH(n,q)} f(x)g(x)$ .

### 3. Definitions of bitrade

**Definition 1.** A pair  $(T_+, T_-)$  of disjoint subsets of  $_{\vee}H(n, q)$  is called an extended perfect bitrade in H(n, q), if there is an  $(q^n, 3)$ -matrix F, whose columns are indexed by the vertices of H(n, q), such that:

- the first column is the characteristic  $(0, \pm 1)$ -function of  $(T_+, T_-)$
- in each row the sum of elements equals 0 and there are at most 2 nonzero elements

• there holds the matrix equation AF = FS, where A is the adjacency matrix of H(n,q) and

$$S = \begin{pmatrix} 0 & n(q-1) & 0 \\ 1 & q-2 & (n-1)(q-1) \\ 0 & n & n(q-2) \end{pmatrix}.$$

**Definition 2.** A pair  $(T_+, T_-)$  of disjoint subsets of  $_{\vee}H(n, q)$  is called an extended perfect bitrade in H(n + 1, q), if  $T_+$  and  $T_-$  are codes with code distance 4 and for every  $i \in \{1, \ldots, n\}$  the *i*-projection of  $(T_+, T_-)$  is a perfect bitrade in H(n, q).

Let A be an arbitrary subset of vH(n,q). For a vertex  $x \in vH(n,q)$ , define its weight  $w_A(x)$  with respect to A in the following way:  $w_A(x) = |S_0(x) \cap A| + |S_1(x) \cap A| + \frac{2}{n}|S_2(x) \cap A|$ .

**Definition 3.** A pair  $(T_+, T_-)$  of disjoint subsets of  ${}_{v}H(n,q)$  is called an extended perfect code in H(n,q) if for any vertex x we have  $w_{T_+}(x) = w_{T_-}(x) \leq 1$ .

**Lemma 2.** Let  $(T_+, T_-)$  be an extended perfect bitrade according to Definition 3. Then  $T_+$  and  $T_-$  are codes with code distance 4.

PROOF. Let there be vertices  $x \neq y$  in  $T_+$  such that  $d(x, y) \leq 3$ . Therefore, there is a path  $x = z_0, z_1, \ldots, z_l = y$ , where  $l \leq 3$ . So,  $w_+(z_1) \geq 1 + 2/n$  and we have contradiction Definition 3 of bitrade. For  $T_-$  the proof is similar.

For a pair  $(T_+, T_-)$  of subsets of vH(n,q) and  $x \in vH(n,q)$ , denote  $w_+(x) = w_{T_+}(x), w_-(x) = w_{T_-}(x)$ .

**Definition 4.** A pair  $(T_+, T_-)$  of disjoint subsets of vH(n, q) is called an extended perfect bitrade in H(n, q), if  $T_+$  and  $T_-$  are codes with code distance 4 and the characteristic  $(0, \pm 1)$ -function of pair belongs to the direct sum of eigenspaces, corresponding to eigenvalues -n and q - 2.

**Definition 5.** A pair  $(T_+, T_-)$  of disjoint subsets of vH(n, q) is called an extended perfect bitrade in H(n, q), if for any  $x \in vH(n, q)$  and  $i \in \{1, ..., n\}$  it holds  $|C_{x,i} \cap T_+| = |C_{x,i} \cap T_-| \le 1$ 

**Lemma 3.** Let A be a code in H(n,q) with code distance 4. Then for any vertex x of H(n,q)

- 1.  $w_A(x) \le 1$ .
- 2. If  $w_A(x) = 1$ , then either  $|S_0(x) \cap A| = 1$ ,  $|S_1(x) \cap A| = 0$ ,  $|S_2(x) \cap A| = 0$ , or  $|S_0(x) \cap A| = 0$ ,  $|S_1(x) \cap A| = 1$ ,  $|S_2(x) \cap A| = 0$ , or  $|S_0(x) \cap A| = 0$ ,  $|S_1(x) \cap A| = 0$ ,  $|S_2(x) \cap A| = n/2$ .

PROOF. If  $x \in A$ , there are no vertices from A in  $(S_1(x) \cup S_2(x))$ . If  $x \notin A$ and x is adjacent with some vertex from A, then  $|S_0(x) \cap A| = 0$ ,  $|S_1(x) \cap A| = 1$ ,  $|S_2(x) \cap A| = 0$ . In the other case, the statement follows from the fact that there are at most n/2 vertices in  $S_2(x)$  with pairwise distances not less than 4.

## 4. Equivalence of the definitions

Since all definitions of bitrade are symmetric with respect to  $T_+$  and  $T_-$ , we will omit one of symmetric cases in the proofs of this section.

#### **Proposition 1.** The definitions 2 and 3 are equivalent.

PROOF.  $3 \Longrightarrow 2$ . Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 3. Consider its *i*-projection for arbitrary *i*, a pair  $(T'_+, T'_-)$ . Let us prove that this pair is a perfect bitrade in H(n-1, q).  $T'_+$  and  $T'_-$  are codes with code distance not less than 3 (since the code distance of the projection of some code A decreases at most by 1 with respect to the code distance of A). For a vertex  $x \in H(n-1,q)$ , denote by  $c_x$  the clique  $\{x_i^a : a \in \mathbb{Z}_q\}$  in H(n,q) (pre-image of x in the projection) and denote by  $c_x(y)$  the vertex in the clique that has  $y \in \mathbb{Z}_q$  in position *i*. Let  $x \in T'_+$ . So,  $c_x(y) \in T_+$  for some  $y \in \mathbb{Z}_q$ . From the code distance we have  $|S_1(x) \cap T'_+| = 0$ . Suppose that  $|S_1(x) \cap T'_-| = 0$ . Then the vertex  $c_x(y)$  has no neighbours in  $T_-$ . Therefore, there are n/2 vertices from  $T_-$  in  $S_2(c_x(y))$  (by Lemma 3). Without loss of generality, assume that  $c_x(y)$  is the zero vertex. Since  $T_-$  has code distance 4, the supports of nonzero positions do not intersect. Hence, the cardinality of the union of the supports equals n. But i does not belong to this union (as in this case, one of the vertices belongs to a clique  $c_z$  such that d(z, x) = 1). Therefore, the union of the supports has cardinality at most n - 1, and we have contradiction.

Let  $x \notin (T'_+ \cup T'_-)$ . If there is a vertex from  $T_+$  and a vertex from  $T_-$  in the clique  $c_x$ , then from the code distance we have  $|S_1(x) \cap (T'_- \cup T'_+)| = 0$ . Assume that there are no vertices from  $(T_+ \cup T_-)$  in  $c_x$ . Suppose that there is a vertex z from  $T'_+$  and there are no vertices from  $T'_-$  in the neighbourhood of x. Then  $c_z(y) \in T_+$  for some  $y \in \mathbb{Z}_q$ ; hence,  $w_+(c_x(y)) = 1$ . Without loss of generality, we assume that  $c_x(y)$  is the zero vertex. So, there are n/2 vertices from  $T_-$  in  $S_2(c_x(y))$  and the supports of nonzero positions do not intersects. As before, we have that i does not belong to the union of these supports. Since n-1 positions remain, we have the contradiction.

 $2 \implies 3$ . Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 2. Consider an arbitrary vertex x. Without loss of generality, we assume that it is the zero vertex. Let us prove that  $w_+(x) = w_-(x)$ . Consider cases. Suppose  $x \in T_+$ . In this case,  $w_+(x) = 1$ . If there is a vertex from  $T_-$  in the neighbourhood of x, then  $w_{-}(x) = 1$ . Otherwise, let  $y_1, \ldots, y_k$  be vertices from  $T_{-}$  at the distance 2 from x and let  $I_i$  be the support of nonzero positions of  $y_i$ . The supports  $I_1, \ldots, I_i$ do not intersect. Let *i* does not belong to the union of their supports. Consider the *i*-projection. The projection of x has no neighbours from one part of perfect bitrade that contradicts the definition of perfect bitrade. Therefore, k = n/2 and  $w_{-}(x) = 1$ . Let  $x \notin (T_{+} \cup T_{-})$ . If there is a vertex from  $T_{+}$  and a vertex from  $T_{-}$  in the neighbourhood of x, then  $w_+(x) = w_-(y) = 1$ . Suppose that  $|S_1(x) \cap T_+| = 1$  and  $|S_1(x) \cap T_-| = 0$ . As before, there are n/2 vertices from  $T_-$  at the distance 2 from x, hence,  $w_{-}(x) = 1$ . If there are no vertices from  $(T_{+} \cup T_{-})$  in the neighbourhood of x, then analogously there is the same number of vertices from  $T_+$  and  $T_-$  at the distance 2 from x (as the union of the supports of nonzero positions is the same in both cases; otherwise, there is some position j such that x has a neighbour only in one part of the perfect bitrade in the j-projection).

Therefore, in all cases we have  $w_+(x) = w_-(x)$ , and  $(T_+, T_-)$  is an extended perfect bitrade according to Definition 3.

#### **Proposition 2.** The Definitions 1 and 3 are equivalent.

PROOF.  $3 \Longrightarrow 1$  Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 3. Let us construct a matrix F that satisfy Definition 1. The first column

of F is the characteristic  $(0, \pm 1)$  function of  $(T_+, T_-)$ . For a vertex v denote by f(v) the v-row of F. Let  $x \in T_+$ . If there is a vertex from  $T_-$  in the neighbourhood of x then define f(x) = (+1, -1, 0), otherwise, f(x) = (+1, 0, -1). Let  $x \in T_-$ . Analogously, if there is a vertex from  $T_+$  in the neighbourhood of x then define f(x) = (-1, +1, 0), otherwise, f(x) = (-1, 0, +1). Let  $x \notin (T_+ \cup T_-)$ . If  $|S_1(x) \cap T_+| = |S_1(x) \cap T_-|$ , then f(x) = (0, 0, 0). Otherwise, f(x) = (0, +1, -1) if  $|S_1(x) \cap T_+| = 1$  and f(x) = (0, -1, +1) if  $|S_1(x) \cap T_-| = 1$ .

Thus, we defined the matrix F. It remains to prove that AF = FS. Let x be an arbitrary vertex of H(n,q). Without loss of generality, we assume that x is the zero vertex. The neighbourhood of x is the union of n cliques of cardinality q - 1; denote them by  $N_1, \ldots, N_n$ . Consider some cases depending on f(x).

Let f(x) = (+1, -1, 0). By definition,  $x \in T_+$ , there is a vertex  $y \in T_-$  in  $S_1(x)$ , and there are no vertices from  $(T_- \cup T_+)$  in  $S_2(x)$ . The vertex y belongs to  $N_i$  for some i and f(y) = (-1, +1, 0). The remaining vertices of  $N_i$  have value (0, 0, 0). The remaining (n-1)(q-1) vertices from the neighbourhood have value (0, +1, -1). Hence,  $AF_x = (-1, +1, 0) + (n-1)(q-1)(0, +1, -1) = (-1, (n-1)(q-1) + 1, -(n-1)(q-1))$ . On the other hand, (+1, -1, 0)S = (-1, (n-1)(q-1) + 1, -(n-1)(q-1)).

Let f(x) = (+1, 0, -1). By definition,  $x \in T_+$ , there are no vertices from  $T_-$  in  $S_1(x)$ , and there are n/2 vertices from  $T_-$  in  $S_2(x)$ . So, in each clique  $N_i$  there is exactly one vertex adjacent to some vertex from  $T_-$ . Therefore, there are n vertices with value (0, 0, 0) and n(q-2) vertices with value (0, +1, -1) in the neighbourhood of x. Hence,  $AF_x = n(q-2)(0, +1, -1)$ . On the other hand, (+1, 0, -1)S equals (0, n(q-2), -n(q-2)).

Let f(x) = (0, +1, -1). By definition,  $x \notin (T_+ \cup T_-)$ , there is a vertex  $y \in T_+$  in  $S_1(x)$  and there are n/2 vertices from  $T_-$  in  $S_2(x)$ . The vertex y belongs to  $N_i$  for some i. If y is adjacent to some vertex from  $T_-$ , then there is a vertex y with value (+1, -1, 0), q-2 vertices in  $N_i$  with value (0, +1, -1) (not adjacent with vertices from  $T_-$ ), n-1 vertices with value (0, -1, +1) in the neighbourhood of x. All remaining vertices from the neighbourhood have value (0, 0, 0). Hence,  $AF_x = (+1, -1, 0) + (q-2)(0, +1, -1) + (n-1)(0, -1, +1) = (+1, -n+q-2, n-q+1)$ . If y is not adjacent with any vertex from  $T_-$ , then the neighbourhood of x containing a vertex y with value (1, 0, -1), q-3 vertices in  $N_i$  with value (0, +1, -1), and n-1 vertices with value (0, -1, +1). All remaining vertices from the neighbourhood have value (0, 0, -1, +1). Hence,  $AF_x = (+1, 0, -1) + (q-3)(0, +1, -1) + (n-1)(0, -1, +1) = (+1, q-n-2, n-q+1)$ . This value is also equal to (0, +1, -1)S.

Let f(x) = (0, 0, 0). In this case, by definition,  $x \notin (T_+ \cup T_-)$ . Suppose that there are  $y \in T_+$  and  $z \in T_-$  in the neighbourhood of x. Let y and z belong to the same clique  $N_i$  for some i. So, f(y) = (+1, -1, 0), f(z) = (-1, +1, 0) and all remaining vertices from the neighbourhood have value (0, 0, 0). Hence,  $AF_x = (0, 0, 0)$ . Let yand z belong to different cliques. So, the neighbourhood of x contains the vertex ywith value (+1, 0, -1), the vertex z with value (-1, 0, +1), q - 2 vertices with value (0, +1, -1), (q - 2) vertices with value (0, -1, +1). All remaining vertices from the neighbourhood have value (0, 0, 0). Hence,  $AF_x = (0, 0, 0)$ . It remains to consider the case when there are no vertices from  $(T_+ \cup T_-)$  in the neighbourhood of x. In this case, the number of vertices in  $S_1(x)$  that adjacent to some vertex from  $T_+$  and not adjacent with any vertex from  $T_-$  and not adjacent with any vertex from  $T_+$ . All remaining vertices from the neighbourhood have value (0, 0, 0). Hence,  $AF_x = (0, 0, 0)$ . This is also equal to (0, 0, 0)S = (0, 0, 0).

 $1 \Longrightarrow 3$  Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 1. Consider an arbitrary vertex x. Without loss of generality, we assume that x is the zero vertex. Let us prove that  $w_+(x) = w_-(x)$ . Remind that by definition  $T_+$  and  $T_-$  are codes with code distance 4, so, we can use Lemma 3. Consider some cases depending on the value of f(x).

Let f(x) = (+1, -1, 0). So,  $x \in T_+$  and  $w_+(x) = 1$ . From the matrix equation, in the vector  $AF_x$  the value in the first position equals -1. Therefore, there is a vertex from  $T_-$  in the neighbourhood of x, and hence  $w_-(x) = 1$ .

Let f(x) = (0, +1, -1) (f(x) = (0, -1, +1)). So, there is a vertex y from  $T_+$   $(T_-)$ and there are no vertices from  $T_ (T_+)$  in the neighbourhood of x. So,  $w_+(x) = 1$  $(w_-(x) = 1)$ . Let  $y \in N_i$ , and let  $z_1, \ldots, z_k$  be vertices from  $T_-$  in  $S_2(x)$ . Let f(y) =(+1, -1, 0). In this case, excluding y, the neighbourhood of x contains (q-2) vertices with value (0, +1, -1), 2k - 1 vertices with value (0, -1, +1), and the remaining vertices have value (0, 0, 0). Let f(y) = (+1, 0, -1). In this case, excluding y, the neighbourhood of x contains q-3 vertices with value (0, +1, -1), 2k - 1 vertices with value (0, -1, +1), and the remaining vertices have value (0, 0, 0). In both cases, from the matrix equation we have 2k = n and  $w_-(x) = 1$ .

Let f(x) = (+1, 0, -1). So,  $w_+(x) = 1$  and there are no vertices from  $T_-$  in the neighbourhood of x. Hence, the vertices from the neighbourhood have values (0, 0, 0) or (0, +1, -1). In the other hand, the sum of these values is equal to (0, n(q - 2), -n(q - 2)). Hence, the neighbourhood of x contains n vertices that has neighbour in  $T_-$ . Since each vertex from  $S_2(x)$  is adjacent with 2 vertices from  $S_1(x)$ , there are n/2 vertices from  $T_-$  in  $S_1(x)$ ; hence,  $w_-(x) = 1$ .

Let f(x) = (0, 0, 0). If there is a vertex from  $T_+$  and a vertex from  $T_-$  in the neighbourhood of x, then  $w_-(x) = w_+(x) = 1$ . Otherwise, the number of vertices with value (0, +1, -1) is equal to the number of vertices with value (0, -1, +1). All remaining vertices have value (0, 0, 0). Therefore,  $S_2(x)$  has the same number of vertices from  $T_+$  and  $T_-$ ; i.e.  $w_+(x) = w_-(x)$ .

Consider two functions  $f: {}_{\mathrm{v}}H(n,q) \to \mathbb{C}$  and  $\varphi: {}_{\mathrm{v}}H(1,q) \to \mathbb{C}$ . Define the function  $f *_i \varphi: {}_{\mathrm{v}}H(n-1,q) \to \mathbb{C}$  in the following way:  $f *_i \varphi(x) = \sum_{y=x_i^a:a \in \mathbb{Z}_q} f(y)\overline{\varphi(a)}$ .

**Lemma 4.** Let  $f \in U_j(n,q)$ , and let  $\psi \equiv 1$  be a constant function on H(1,q). Then

- 1. [4] If  $j \neq n$ , then  $f *_i \psi \in U_j(n-1,q)$  for any  $i \in \{1, ..., n\}$ .
- 2. [4] If j = n, then  $f *_i \equiv 0$  for any  $i \in \{1, ..., n\}$ .
- 3. If  $j \neq n$ , then there is i such that  $f *_i \psi \neq 0$

PROOF. Let  $j \neq n$ . Denote by  $c_{x,i}$  the following set of vertices  $\{x + e_i^a : a \in \mathbb{Z}_q\}$ , which form a clique in H(n,q). Denote by  $f_{x,i}$  the restriction of f on the set  $c_{x,i}$ . If  $f *_i \psi \equiv 0$  for any i, then  $f_{x,i}$  is an eigenfunction of H(1,q) with eigenvalue -1 for any x and i (indeed,  $(g,\psi) = 0$  for any  $g \in U_1(1,q)$  and  $(h,\psi) \neq 0$  for any  $h \in U_0(1,q)$ such that  $h \not\equiv 0$ ). In this case, f is eigenfunction of H(n,q) with eigenvalue  $\lambda_n = -n$ and we have a contradiction.

The following fact is well known and straightforward.

**Lemma 5.** A pair  $(T_+, T_-)$  of disjoint subsets of  $_{\vee}H(n, q)$  is a perfect bitrade in H(n, q) if and only if  $T_+$  and  $T_-$  are codes with code distance 3 and the characteristic  $(0, \pm 1)$ -function f belongs to  $U_l(n, q)$ , where  $\lambda_l = -1$  and, respectively,  $l = \frac{(n-1)(q-1)+1}{q}$ .

**Proposition 3.** The definitions 2 and 4 are equivalent.

PROOF.  $4 \implies 2$  Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 4. Let f be the characteristic  $(0, \pm 1)$ -function of the bitrade. By definition,  $f \in U_n(n,q) \oplus U_l(n,q)$ , where  $\lambda_l = q - 2$ , i.e.  $l = \frac{(n-1)(q-1)+1}{q}$ . Therefore, f = g + h, where  $g \in U_n(n,q)$ ,  $h \in U_l(n,q)$ . Let  $i \in \{1, \ldots, n\}$  be an arbitrary position, and let  $\psi \equiv 1$  be a constant function on H(1,q). We have,  $f *_i \psi = g *_i \psi + h *_i \psi = h *_i \psi \in U_l(n-1,q)$ . By Lemma 5,  $f *_i \psi$  is the characteristic function of some perfect bitrade in H(n-1,q). On the other hand, the pair  $(T'_+, T'_-)$  corresponding to  $f *_i \psi$  is a projection of  $(T_+, T_-)$ . Therefore,  $(T_+, T_-)$  is an extended perfect bitrade by Definition 2.

 $2 \Longrightarrow 4$  Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 2, and let f be its characteristic  $(0, \pm 1)$ -function. Let  $f \in U_{j_1}(n, q) \oplus \ldots \oplus U_{j_k}(n, q)$ . So,  $f = g_1 + \ldots + g_k$ , where  $g_t \in U_{j_t}$ . Denote  $l = \frac{(n-1)(q-1)+1}{q}$  ( $\lambda_l = -1$  in H(n-1,q) for this l). Suppose that there is  $m \in \{j_1, \ldots, j_k\} \setminus \{l, n\}$ . By Lemma 4, there is such r that  $g_m *_r \psi \neq 0$ . Hence,  $f *_r \psi = g_1 *_r \psi + \ldots + g_k *_r \psi = h_1 + \ldots + h_k$ , where  $h_t \in U_{j_t}(n-1,q)$  and  $h_m \neq 0$ . Therefore,  $f *_r \psi \notin U_l(n-1,q)$ , and we have a contradiction with the fact that  $f *_r \psi$  is a characteristic function of some bitrade.

#### **Proposition 4.** The Definitions 2, 3, and 5 are equivalent.

PROOF. 5  $\Longrightarrow$  2. Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 5, and let  $i \in \{1, \ldots, n\}$  be an arbitrary position. Consider the *i*projection of the bitrade; denote it by  $(T'_+, T'_-)$ . Let x be an arbitrary vertex of H(n-1,q) and B(x) be radius-1 ball in H(n-1,q). The pre-image of B(x) is a cylinder  $C_{y,i}$ , where  $y = x_i^0$ . If there are no vertices from  $(T_+ \cup T_-)$  in  $C_{y,i}$ , then there are no vertices from  $(T'_+ \cup T'_-)$  in B(x). Let  $C_{y,i} \cap T_+ = \{v\}, C_{y,i} \cap T_- = \{u\}$ . If the vertices u and v differ only in position i, then there are no vertices from  $(T'_+ \cup T'_-)$  in B(x); otherwise, we have  $|B(x) \cap T'_+| = |B(x) \cap T'_-| = 1$ .

 $3 \implies 5$ . Let  $(T_+, T_-)$  be an extended perfect bitrade in H(n, q) according to Definition 3. Consider  $C_{x,i}$  for arbitrary  $x \in {}_{v}H(n,q)$  and  $i \in \{1, \ldots, n\}$ . Since  $T_+$ and  $T_-$  are codes with code distance 4 by Lemma 2, there are at most one vertex from  $T_+$  and at most one vertex from  $T_-$  in  $C_{x,i}$ . It is sufficient to prove that if there is a vertex y from  $T_+$  in  $C_{x,i}$ , then there is a vertex from  $T_-$  in  $C_{x,i}$ . Since  $y \in T_+$ , there is a vertex z from  $C_{x,i}$  such that  $w_+(z) = 1$  and  $z - x = e_i^a$  for some  $a \in \mathbb{Z}_q$ . So, we have that either z is adjacent to some vertex u from  $T_-$ , or there are n/2 vertices from  $T_$ at the distance 2 from z. In the first case, u also belongs to  $C_{x,i}$ . In the second case, there is a vertex  $v \in T_-$  at the distance 2 from z such that z and v differ in position i and some other position (otherwise, there are vertices  $v_1 \bowtie v_2$  that differ from z in the same position, but in this case  $d(v_1, v_2) \leq 3$ ). So, v also belongs to  $C_{x,i}$ .

 $2 \Longrightarrow 3$ . By Proposition 1.

From Propositions 1 - 4 we have the following.

**Theorem 1.** The Definitions 1 – 5 are equivalent.

A pair  $(T_+, T_-)$  of disjoint subsets of  $_{v}H(n, q)$  is called an *extended perfect bitrade* in H(n, q) if it satisfies one of definitions 1 - 5.

**Theorem 2.** If there is an extended perfect bitrade  $(T_+, T_-)$  in H(n, q), then

- 1. n is even and
- 2. n = lq + 2 for some  $l \in \{0, 1, ...\}$ .

PROOF. 1) Let  $x \in T_+$ . Then there is a vertex  $y \in S_1(x)$  such that y is not adjacent to any vertex from  $T_-$ . Since,  $w_+(y) = 1$ , there are n/2 vertices from  $T_-$  in  $S_2(y)$ . Hence, n is even.

2) By definition, n(q-1) - iq equals q-2 for some *i*. Hence,  $n \equiv 2 \mod q$ .

## References

- R. Ahlswede, H. K. Aydinian, and L. H. Khachatrian. On perfect codes and related concepts. *Des. Codes Cryptography*, 22(3):221–237, 2001. DOI: 10.1023/A:1008394205999.
- [2] L. A. Bassalygo, G. V. Zaitsev, and V. A. Zinoviev. On uniformly packed codes. *Probl. Inf. Transm.*, 10(1):6–9, 1974. Translated from *Probl. Peredachi Inf.*, 10(1): 9-14, 1974.
- [3] Bespalov E. On the non-existence of extended perfect codes and some perfect colorings. E-print 2008.13260, arXiv.org, 2020. Available at http://arxiv.org/abs/2008.13260.
- [4] I. Yu. Mogilnykh and A. A. Valyuzhenich. Equitable 2-partitions of the Hamming graphs with the second eigenvalue. *Discrete Math.*, 343(11):112039(1–9), Nov. 2020. DOI: 10.1016/j.disc.2020.112039.
- [5] A. Neumaier. Completely regular codes. Discrete Math., 106-107:353–360, Sept. 1992. DOI: 10.1016/0012-365X(92)90565-W.