

A TRUNCATED MINIMAL FREE RESOLUTION OF THE RESIDUE FIELD

VAN C. NGUYEN AND OANA VELICHE

ABSTRACT. In a paper in 1962, Golod proved that the Betti sequence of the residue field of a local ring attains an upper bound given by Serre if and only if the homology algebra of the Koszul complex of the ring has trivial multiplications and trivial Massey operations. This is the origin of the notion of Golod ring. Using the Koszul complex components he also constructed a minimal free resolution of the residue field. In this article, we extend this construction up to degree five for *any* local ring. We describe how the multiplicative structure and the triple Massey products of the homology of the Koszul algebra are involved in this construction. As a consequence, we provide explicit formulas for the first six terms of a sequence that measures how far the ring is from being Golod.

1. INTRODUCTION

Throughout the paper, (R, \mathfrak{m}, k) is a local noetherian ring with maximal ideal \mathfrak{m} and residue field k , of embedding dimension n and codepth $c = n - \text{depth } R$. Let K^R be the Koszul complex of R on a minimal set of generators of \mathfrak{m} . Serre pointed out that there is always a coefficient-wise inequality between the Poincaré series of the R -module k and a series of rational form involving the ranks of the homologies of K^R :

$$(1.0.1) \quad P_k^R(t) := \sum_{i=0}^{\infty} \text{rank}_k \text{Tor}_i^R(k, k) t^i \preceq \frac{(1+t)^n}{1 - \sum_{i=1}^c \text{rank}_k H_i(K^R) t^{i+1}}.$$

In [8], Golod proved that a ring R attains this upper bound if and only if the graded-commutative algebra $H(K^R)$ has trivial multiplications and trivial Massey operations; such a ring is now called a *Golod ring*. In the same paper, he also constructed the minimal free resolution of the R -module k in terms of K^R .

In [2, Corollary 5.10], Avramov proved that the Poincaré series $P_k^R(t)$ is completely determined by the structure of the Koszul homology algebra $A := H(K^R)$ as an algebra with Massey operations. One can now ask the following question:

Question 1.1. For *any* local ring (R, \mathfrak{m}, k) , given the knowledge of its Koszul homology algebra A , with its products and Massey operations, how does one construct a *minimal* free resolution of the residue field k ?

In this paper, we answer this question explicitly up to degree five. For any local ring R , using the Koszul complex K^R as building blocks and the graded-commutative structure of its homology A , we construct a minimal free resolution of the R -module k , up to degree five, see Construction 3.1 and Theorem 3.2:

$$F : \quad F_5 \xrightarrow{\partial_5^F} F_4 \xrightarrow{\partial_4^F} F_3 \xrightarrow{\partial_3^F} F_2 \xrightarrow{\partial_2^F} F_1 \xrightarrow{\partial_1^F} F_0 \rightarrow k \rightarrow 0.$$

The higher degrees of the resolution can be extended similarly for some special cases of R , but in general it requires an understanding of higher Massey products, which remains elusive and is left for future projects. Indeed, Massey operations can be represented by using the A_∞ -algebra structure, which the Koszul homology algebra A possesses, see for example, [12] by Lu, Palmieri, Wu, and Zhang. Moreover, the A_∞ -structure was also used by Burke in [6] to construct certain projective resolutions. These connections suggest that perhaps the A_∞ -algebra structure of A may play a role in giving a complete answer to Question 1.1 from this perspective.

To prepare for Construction 3.1, in Section 2 we analyze in detail the multiplicative structure of the algebra A , up to degree four, in terms of bases of A_i and $A_i \cdot A_j$, and necessary maps. Moreover, in degree four the Massey products appear for the first time, as ternary Massey products $\langle A_1, A_1, A_1 \rangle$; we give a description of the elements of this set in Proposition 2.11.

Date: March 25, 2021.

2010 Mathematics Subject Classification. 13D02, 13D07, 13H10, 13C05.

Key words and phrases. Golod rings, minimal free resolution, Tor algebra, Massey products.

In Section 4, we obtain several direct applications of Theorem 3.2. In Corollary 4.1, we explicitly describe the Betti numbers $\beta_i := \text{rank}_k \text{Tor}_i^R(k, k)$, up to degree five, in terms of the multiplicative invariants of the Koszul homology algebra A . Let $a_i := \text{rank}_k A_i$ and consider the difference of series

$$\mathcal{P}(t) := \frac{(1+t)^n}{1 - \sum_{i=1}^c a_i t^{i+1}} - P_k^R(t)$$

that measures how far the ring R is from being Golod, that is, how far the Betti numbers of R are from their maximum possible values. In Proposition 4.2, we compute the first six coefficients of $\mathcal{P}(t)$ in terms of multiplicative invariants of A :

$$\mathcal{P}_0 = \mathcal{P}_1 = \mathcal{P}_2 = 0, \quad \mathcal{P}_3 = q_{11}, \quad \mathcal{P}_4 = (n+1)q_{11} + q_{12}, \quad \mathcal{P}_5 = \binom{n+1}{2} + 2a_1 q_{11} + (n+1)q_{12} + a - b,$$

where $q_{ij} = \text{rank}_k(A_i \cdot A_j)$, and a, b are described in Summary 2.13. For any local ring R of embedding dimension n with rational Poincaré series of the form $P_k^R(t) = (1+t)^n/d(t)$, we express the denominator, up to degree five, as follows:

$$d(t) = (1 - \sum_{i=1}^c a_i t^{i+1}) + q_{11}t^3 + (q_{11} + q_{12})t^4 + (q_{12} - b + a)t^5 + f(t)t^6,$$

for some $f(t) \in \mathbb{Z}[t]$, see Proposition 4.5. We consider a few classes of rings with such rational Poincaré series, see Corollary 4.10 and Examples 4.11 and 4.12, and using the coefficients of the denominator we obtain algebraic invariants of their Koszul homology algebra A . The Poincaré series $P_k^R(t)$ can be described by using the deviations ε_i 's, see for example [4]. In Corollary 4.13, we give a description of the first five deviations in terms of the algebraic invariants of A .

In Section 5, we illustrate Construction 3.1 through an example of a ring of codepth 4 examined by Avramov in [2]. In that example, Avramov provided a nontrivial indecomposable Massey product element in $\langle A_1, A_1, A_1 \rangle$ that does not come from multiplications of the homology. Using Proposition 2.11, we prove that, up to a scalar, this is the only element with this property, modulo the products in homology.

2. MULTIPLICATIVE STRUCTURE ON THE HOMOLOGY OF THE KOSZUL ALGEBRA

Let (R, \mathfrak{m}, k) be a local ring of embedding dimension n and codepth c . The differential graded algebra structure on the Koszul complex $K = K^R$ on a minimal set of generators of the maximal ideal \mathfrak{m} induces a graded-commutative algebra structure on the homology of K :

$$A := H(K) = A_0 \oplus A_1 \oplus A_2 \oplus \cdots \oplus A_c.$$

In this section we discuss this multiplicative structure up to degree four. We will use this structure extensively in constructing a truncated minimal free resolution of the residue field over R in the next section.

The following notation is used throughout the paper: the differential map on K is denoted by ∂^K , a homogeneous element of degree i in the Koszul complex K is denoted by π^i , a representative in K_i of an element in the homology A_i is denoted by p^i , and elements of R are denoted by α 's, β 's, and γ 's. The subscript of a homogeneous element indicates its index in a tuple and the superscript indicates its homological degree. Set

$$(2.0.1) \quad a_i := \text{rank}_k(A_i) \quad \text{and} \quad q_{ij} := \text{rank}_k(A_i \cdot A_j), \text{ for all } 0 \leq i, j \leq c.$$

It is clear that $A_0 = k$. For each $1 \leq i \leq a_1$ we consider $z_i^1 \in \text{Ker } \partial_1^K$ such that

$$(2.0.2) \quad \{[z_i^1]\}_{i=1, \dots, a_1} \text{ is a basis of } A_1.$$

In particular, every element in $\text{Ker } \partial_1^K$ can be written as a sum of elements in $\text{Im } \partial_2^K$ and a linear combination of $\{z_i^1\}_{i=1, \dots, a_1}$.

2.1. Products in degree two. Let $\overline{A_2}$ be a k -subspace of A_2 such that

$$A_2 = (A_1 \cdot A_1) \oplus \overline{A_2}.$$

For each $1 \leq \ell \leq a_2 - q_{11}$, we consider $z_\ell^2 \in \text{Ker } \partial_2^K$ such that

$$(2.1.1) \quad \{[z_\ell^2]\}_{\ell=1, \dots, a_2 - q_{11}} \text{ is a basis of } \overline{A_2}.$$

In particular, every element in $\text{Ker } \partial_2^K$ can be written as a sum of elements in $\text{Im } \partial_3^K$, a linear combination of $\{z_i^1 \wedge z_j^1\}_{i, j=1, \dots, a_1}$, and a linear combination of $\{z_\ell^2\}_{\ell=1, \dots, a_2 - q_{11}}$.

Consider the multiplication map:

$$(2.1.2) \quad \phi_1 : A_1 \otimes A_1 \rightarrow A_2, \quad \text{defined by} \quad \phi_1([x] \otimes [y]) = [x] \wedge [y].$$

As $\text{Im } \phi_1 = A_1 \cdot A_1$, we have $\text{rank}_k(\text{Ker } \phi_1) = a_1^2 - q_{11}$. For all $1 \leq i \leq a_1$ and $1 \leq s \leq a_1^2 - q_{11}$, we choose $\tilde{p}_{si}^1 \in \text{Ker } \partial_1^K$ such that

$$(2.1.3) \quad \left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \right\}_{s=1, \dots, a_1^2 - q_{11}} \quad \text{is a basis of } \text{Ker } \phi_1.$$

For each s , we have $\sum_{i=1}^{a_1} [z_i^1] \wedge [\tilde{p}_{si}^1] = 0$, so there exists $\tilde{\pi}_s^3 \in K_3$ such that

$$(2.1.4) \quad \partial_3^K(\tilde{\pi}_s^3) = \sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{si}^1.$$

Lemma 2.2. *Let $\tilde{p}_{si}^1 \in \text{Ker } \partial_1^K$ be as in (2.1.3). Then the vectors $\{([\tilde{p}_{s1}^1], \dots, [\tilde{p}_{sa_1}^1])\}_{s=1, \dots, a_1^2 - q_{11}}$ in $A_1^{a_1}$ are linearly independent.*

Proof. If α_s is in R such that $\sum_{s=1}^{a_1^2 - q_{11}} ([\tilde{p}_{s1}^1], \dots, [\tilde{p}_{sa_1}^1]) \wedge [\alpha_s] = 0$, then

$$\sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\alpha_s] = 0, \quad \text{for all } 1 \leq i \leq a_1.$$

By tensoring with $[z_i^1]$ and taking the sum over i we get

$$\sum_{i=1}^{a_1} [z_i^1] \otimes \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\alpha_s] = \sum_{s=1}^{a_1^2 - q_{11}} \left(\sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \right) \wedge [\alpha_s] = 0.$$

The desired conclusion now follows from (2.1.3). \square

Proposition 2.3. *Let $\{[z_i^1]\}_{i=1, \dots, a_1}$ and $\{[z_\ell^2]\}_{\ell=1, \dots, a_2 - q_{11}}$ be as in (2.0.2) and (2.1.1), respectively. If β_{ij} and α_ℓ are in R such that*

$$\sum_{i,j=1}^{a_1} [z_i^1] \wedge [z_j^1] \wedge [\beta_{ij}] + \sum_{\ell=1}^{a_2 - q_{11}} [z_\ell^2] \wedge [\alpha_\ell] = 0,$$

then the following hold:

- (a) $[\alpha_\ell] = 0$ for all $1 \leq \ell \leq a_2 - q_{11}$;
- (b) There exist γ_s in R such that for all $1 \leq i \leq a_1$

$$\sum_{j=1}^{a_1} [z_j^1] \wedge [\beta_{ij}] = \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\gamma_s],$$

where \tilde{p}_{si}^1 is as in (2.1.3).

Proof. (a): The first sum in the hypothesized equality is in $A_1 \cdot A_1$. The elements $\{[z_\ell^2]\}_{\ell=1, \dots, a_2 - q_{11}}$ form a basis in A_2 that completes a basis of $A_1 \cdot A_1$, thus $[\alpha_\ell] = 0$ for all $1 \leq \ell \leq a_2 - q_{11}$.

(b): By (a) it follows that $0 = \sum_{i,j=1}^{a_1} [z_i^1] \wedge [z_j^1] \wedge [\beta_{ij}] = \sum_{i=1}^{a_1} [z_i^1] \wedge \left(\sum_{j=1}^{a_1} [z_j^1] \wedge [\beta_{ij}] \right)$, which implies that $\sum_{i=1}^{a_1} [z_i^1] \otimes \left(\sum_{j=1}^{a_1} [z_j^1] \wedge [\beta_{ij}] \right)$ is in $\text{Ker } \phi_1$. The desired conclusion now follows from (2.1.3). \square

2.4. Products in degree three. Let $\overline{A_3}$ be a k -subspace of A_3 such that

$$A_3 = (A_1 \cdot A_2) \oplus \overline{A_3} = (A_1 \cdot A_1 \cdot A_1 + A_1 \cdot \overline{A_2}) \oplus \overline{A_3}.$$

For each $1 \leq t \leq a_3 - q_{12}$, we consider $z_t^3 \in \text{Ker } \partial_3^K$ such that

$$(2.4.1) \quad \{[z_t^3]\}_{t=1, \dots, a_3 - q_{12}} \quad \text{is basis of } \overline{A_3}.$$

In particular, every element in $\text{Ker } \partial_3^K$ can be written as a sum of elements in $\text{Im } \partial_4^K$, a linear combination of $\{z_i^1 \wedge z_j^1 \wedge z_k^1\}_{i,j,k=1, \dots, a_1}$, a linear combination of $\{z_i^1 \wedge z_\ell^2\}_{i=1, \dots, a_1, \ell=1, \dots, a_2 - q_{11}}$, and a linear combination of

$$\{z_t^3\}_{t=1,\dots,a_3-q_{12}}.$$

Consider the map:

$$(2.4.2) \quad \psi : A_1 \otimes A_1 \otimes A_1 \rightarrow ((A_1 \cdot A_1) \otimes A_1) \oplus (A_1 \otimes (A_1 \cdot A_1)),$$

defined by

$$\psi([x] \otimes [y] \otimes [z]) = ([x] \wedge [y] \otimes [z], [x] \otimes [y] \wedge [z])$$

for all $[x], [y], [z] \in A_1$ and set

$$(2.4.3) \quad b = \text{rank}_k(\text{Coker } \psi).$$

Remark that $b = 0$ if and only if ψ is surjective. If $q_{11} = 0$, then $A_1 \cdot A_1 = 0$ and hence $b = 0$.

Lemma 2.5. *Let a_i, q_{ij} be as in (2.0.1) and b be as in (2.4.3). The sequence*

$$A_1 \otimes A_1 \otimes A_1 \xrightarrow{\bar{\psi}} ((A_1 \cdot A_1) \otimes A_1) \oplus (A_1 \otimes A_2) \xrightarrow{\mu} A_3,$$

where

$$\begin{aligned} \bar{\psi}([x] \otimes [y] \otimes [z]) &= ([x] \wedge [y] \otimes [z], [x] \otimes [y] \wedge [z]), \text{ and} \\ \mu([u] \otimes [x], [y] \otimes [v]) &= [u] \wedge [x] - [y] \wedge [v], \end{aligned}$$

for all $[x], [y], [z] \in A_1$, $[u] \in (A_1 \cdot A_1)$ and $[v] \in A_2$, is a complex whose homology has rank

$$a_1 a_2 - a_1 q_{11} - q_{12} + b.$$

Proof. We first show that $\mu \circ \bar{\psi} = 0$.

$$\mu(\bar{\psi}([x] \otimes [y] \otimes [z])) = \mu([x] \wedge [y] \otimes [z], [x] \otimes [y] \wedge [z]) = [x] \wedge [y] \wedge [z] - [x] \wedge [y] \wedge [z] = 0.$$

Hence, the sequence in the statement of the lemma is a complex. The homology of this complex has

$$\begin{aligned} \text{rank}_k \left(\frac{\text{Ker } \mu}{\text{Im } \bar{\psi}} \right) &= \text{rank}_k(\text{Ker } \mu) - \text{rank}_k(\text{Im } \bar{\psi}) \\ &= \text{rank}_k \left(((A_1 \cdot A_1) \otimes A_1) \oplus (A_1 \otimes A_2) \right) - \text{rank}_k(\text{Im } \mu) - \text{rank}_k(\text{Im } \psi) \\ &= a_1 q_{11} + a_1 a_2 - q_{12} - \text{rank}_k \left(((A_1 \cdot A_1) \otimes A_1) \oplus (A_1 \otimes (A_1 \cdot A_1)) \right) + b \\ &= a_1 q_{11} + a_1 a_2 - q_{12} - 2a_1 q_{11} + b \\ &= a_1 a_2 - a_1 q_{11} - q_{12} + b. \end{aligned}$$

□

To describe a basis of the homology of the complex in Lemma 2.5, it is enough to observe that $\text{Im } \bar{\psi}$ is generated by elements of the form $([z_i^1] \wedge [z_j^1] \otimes [z_k^1], [z_i^1] \otimes [z_j^1] \wedge [z_k^1])$ for $1 \leq i, j, k \leq a_1$ and that in $\text{Ker } \mu / \text{Im } \bar{\psi}$ we have $[(z_i^1] \wedge [z_j^1] \otimes [z_k^1], 0) = -[(0, [z_i^1] \otimes [z_j^1] \wedge [z_k^1])]$, as $[(z_i^1] \wedge [z_j^1] \otimes [z_k^1], [z_i^1] \otimes [z_j^1] \wedge [z_k^1]) = 0$. We choose $\tilde{p}_{ui}^2 \in \text{Ker } \partial_2^K$ such that

$$(2.5.1) \quad \left\{ \left[0, \left(\sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{ui}^2] \right) \right] \right\}_{u=1,\dots,a_1 a_2 - a_1 q_{11} - q_{12} + b},$$

is a basis of the homology of the complex defined in Lemma 2.5. By definition of μ , for each u we have $\sum_{i=1}^{a_1} [z_i^1] \wedge [\tilde{p}_{ui}^2] = 0$ in A_3 . Therefore, for each u there exists $\tilde{\pi}_u^4 \in K_4$ such that

$$(2.5.2) \quad \partial_4^K(\tilde{\pi}_u^4) = \sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{ui}^2.$$

The following result gives a method for finding \tilde{p}_{ui}^2 from (2.5.1).

Proposition 2.6. Consider the map $\phi_2 : A_1 \otimes A_2 \rightarrow A_3$, defined by $\phi_2([x] \otimes [v]) = [x] \wedge [v]$, and set

$$\mathcal{A} = \left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \wedge [z_j^1] \mid 1 \leq j \leq a_1 \text{ and } 1 \leq s \leq a_1^2 - q_{11} \right\} \quad \text{and}$$

$$\mathcal{B} = \left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{ui}^2] \mid 1 \leq u \leq a_1 a_2 - a_1 q_{11} - q_{12} + b \right\},$$

where \tilde{p}_{si}^1 and \tilde{p}_{ui}^2 are as in (2.1.3) and (2.5.1) respectively. Then, the following hold:

(a) The set \mathcal{B} is linearly independent and

$$\text{Ker } \phi_2 = (\text{Span}_{\mathbf{k}} \mathcal{A}) \oplus (\text{Span}_{\mathbf{k}} \mathcal{B}).$$

(b) $\text{Span}_{\mathbf{k}} \mathcal{A} \subseteq A_1 \otimes (A_1 \cdot A_1)$ and for b defined in (2.4.3),

$$b = a_1 q_{11} - \text{rank}_{\mathbf{k}}(\text{Span}_{\mathbf{k}} \mathcal{A}).$$

(c) If B is a \mathbf{k} -subspace of $\text{Ker } \phi_2$ such that

$$\text{Ker } \phi_2 = (\text{Span}_{\mathbf{k}} \mathcal{A}) \oplus B,$$

then for every basis \mathcal{B}' of B the set $[(0, \mathcal{B}')] is a basis of the homology of the complex in Lemma 2.5.$

Proof. (a): The linear independence of \mathcal{B} follows from (2.5.1). Next, we show that the elements of \mathcal{A} and \mathcal{B} generate the kernel of ϕ_2 . By definitions of \tilde{p}_{si}^1 and \tilde{p}_{ui}^2 , it is clear that the listed elements are in the kernel of ϕ_2 . Let $\sum_{i=1}^{a_1} [z_i^1] \otimes [p_i^2] \in \text{Ker } \phi_2$, for some $[p_i^2] \in A_2$. Then $[(0, \sum_{i=1}^{a_1} [z_i^1] \otimes [p_i^2])] \in \text{Ker } \mu$, and thus by definition (2.5.1) for all u, i, j there exist $\delta_u \in R$ and $p_{ij}^1 \in \text{Ker } \partial_1^K$ such that

(2.6.1)

$$\left(0, \sum_{i=1}^{a_1} [z_i^1] \otimes [p_i^2]\right) = \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \left(0, \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{ui}^2] \wedge [\delta_u]\right) + \sum_{i,j=1}^{a_1} \left([z_i^1] \wedge [p_{ij}^1] \otimes [z_j^1], [z_i^1] \otimes [p_{ij}^1] \wedge [z_j^1]\right).$$

Comparing the first components of both sides of (2.6.1), for each j we have:

$$0 = \sum_{i=1}^{a_1} [z_i^1] \wedge [p_{ij}^1].$$

By (2.1.3), there exist ε_{js} in R such that for all $1 \leq i, j \leq a_1$ we can write:

$$[p_{ij}^1] = \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\varepsilon_{js}].$$

Comparing the second components of both sides of (2.6.1), and using the above expression for $[p_{ij}^1]$, for each i we have:

$$[p_i^2] = \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} [\tilde{p}_{ui}^2] \wedge [\delta_u] + \sum_{j=1}^{a_1} [p_{ij}^1] \wedge [z_j^1] = \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} [\tilde{p}_{ui}^2] \wedge [\delta_u] + \sum_{s=1}^{a_1^2 - q_{11}} \sum_{j=1}^{a_1} [\tilde{p}_{si}^1] \wedge [z_j^1] \wedge [\varepsilon_{js}].$$

Thus, $\sum_{i=1}^{a_1} [z_i^1] \otimes [p_i^2] \in \text{Ker } \phi_2$ is a linear combination of elements of \mathcal{A} and \mathcal{B} . Next, we show that $\text{Span}_{\mathbf{k}} \mathcal{A} \cap \text{Span}_{\mathbf{k}} \mathcal{B} = \{0\}$. By definition of \tilde{p}_{si}^1 we have $\sum_{i=1}^{a_1} [z_i^1] \wedge [\tilde{p}_{si}^1] = 0$ for all $1 \leq s \leq a_1^2 - q_{11}$. By definition of $\tilde{\psi}$, for all s and j we obtain

$$(2.6.2) \quad \tilde{\psi} \left(\sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \otimes [z_j^1] \right) = \left(0, \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \wedge [z_j^1]\right),$$

hence $(0, \mathcal{A}) \subseteq \text{Im } \tilde{\psi}$. On the other hand by (2.5.1), $[(0, \mathcal{B})]$ is a basis of the homology of the complex in Lemma 2.5. Therefore, $\text{Span}_{\mathbf{k}} \mathcal{A}$ and $\text{Span}_{\mathbf{k}} \mathcal{B}$ have no nontrivial elements in common, and part (a) holds.

(b): The inclusion follows from the definition of \mathcal{A} . Observe that $\text{rank}_k(\text{Ker } \phi_2) = a_1 a_2 - q_{12}$, and by the linear independence of \mathcal{B} from part (a), $\text{rank}_k(\text{Span}_k \mathcal{B}) = a_1 a_2 - q_{12} - (a_1 q_{11} - b)$. Therefore, by part (a), $\text{rank}_k(\text{Span}_k \mathcal{A}) = a_1 q_{11} - b$.

(c): By parts (a) and (b), every basis \mathcal{B}' of B has $a_1 a_2 - a_1 q_{11} - q_{12} + b$ elements. It is clear that $(0, \mathcal{B}') \subseteq \text{Ker } \mu$, with μ as in Lemma 2.5 and that the set $[(0, \mathcal{B}')]$ has at most $a_1 a_2 - a_1 q_{11} - q_{12} + b$ elements. It is enough to show that it is a generating set for the homology of the complex in Lemma 2.5 to conclude that $[(0, \mathcal{B}')]$ has exactly $a_1 a_2 - a_1 q_{11} - q_{12} + b$ elements, hence it forms a basis for the homology. Since $\mathcal{B} \subseteq \text{Ker } \phi_2$, every element of \mathcal{B} can be written as a linear combination of elements of \mathcal{A} and \mathcal{B}' . In particular, every basis element in $[(0, \mathcal{B})]$ can be written as a linear combination of elements in $[(0, \mathcal{A})]$ and $[(0, \mathcal{B}')]$. However, each element in $[(0, \mathcal{A})]$ is zero in the homology, as remarked in (2.6.2), thus $[(0, \mathcal{B}')] is a generating set. Part (c) now holds. $\square$$

Remark 2.7. In practice, one can apply Proposition 2.6(b) to compute the value b , instead of using definition (2.4.3). Similarly, instead of using definition (2.5.1) for \tilde{p}_{ui}^2 , by Proposition 2.6(c), one can find first \tilde{p}_{si}^1 as defined in (2.1.3), and then find a basis of a space complementary to

$$\text{Span}_k \mathcal{A} = \text{Span}_k \left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \wedge [z_j^1] \right\}_{1 \leq j \leq a_1, 1 \leq s \leq a_1^2 - q_{11}}$$

in the kernel of the multiplication map ϕ_2 , of the form

$$\left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{ui}^2] \right\}_{1 \leq u \leq a_1 a_2 - a_1 q_{11} - q_{12} + b}.$$

Proposition 2.8. Let $\{[z_i^1]\}_{i=1, \dots, a_1}$, $\{[z_\ell^2]\}_{\ell=1, \dots, a_2 - q_{11}}$, and $\{[z_t^3]\}_{t=1, \dots, a_3 - q_{12}}$ be as in (2.0.2), (2.1.1), and (2.4.1), respectively. If γ_{ijk} , $\beta_{i\ell}$, and α_t are in R such that

$$\sum_{i,j,k=1}^{a_1} [z_i^1] \wedge [z_j^1] \wedge [z_k^1] \wedge [\gamma_{ijk}] + \sum_{i=1}^{a_1} \sum_{\ell=1}^{a_2 - q_{11}} [z_i^1] \wedge [z_\ell^2] \wedge [\beta_{i\ell}] + \sum_{t=1}^{a_3 - q_{12}} [z_t^3] \wedge [\alpha_t] = 0,$$

then the following hold:

- (a) $[\alpha_t] = 0$ for all $1 \leq t \leq a_3 - q_{12}$;
- (b) There exist δ_u and ε_{js} in R such that

$$\sum_{j,k=1}^{a_1} [z_j^1] \wedge [z_k^1] \wedge [\gamma_{ijk}] + \sum_{\ell=1}^{a_2 - q_{11}} [z_\ell^2] \wedge [\beta_{i\ell}] = \sum_{j=1}^{a_1} \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [z_j^1] \wedge [\varepsilon_{js}] + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} [\tilde{p}_{ui}^2] \wedge [\delta_u],$$

where \tilde{p}_{si}^1 and \tilde{p}_{ui}^2 are as in (2.1.3) and (2.5.1) respectively.

Proof. (a): The first two sums in the hypothesized equality are in $A_1 \cdot A_2$. As $\{[z_t^3]\}_{t=1, \dots, a_3 - q_{12}}$ is a basis in A_3 that completes a basis of $A_1 \cdot A_2$, this implies $[\alpha_t] = 0$ for all $1 \leq t \leq a_3 - q_{12}$.

(b): By (a) it follows that

$$\sum_{i=1}^{a_1} [z_i^1] \wedge \left(\sum_{j,k=1}^{a_1} [z_j^1] \wedge [z_k^1] \wedge [\gamma_{ijk}] + \sum_{\ell=1}^{a_2 - q_{11}} [z_\ell^2] \wedge [\beta_{i\ell}] \right) = 0,$$

hence $\sum_{i=1}^{a_1} [z_i^1] \otimes \left(\sum_{j,k=1}^{a_1} [z_j^1] \wedge [z_k^1] \wedge [\gamma_{ijk}] + \sum_{\ell=1}^{a_2 - q_{11}} [z_\ell^2] \wedge [\beta_{i\ell}] \right)$ is in $\text{Ker } \phi_2$. The desired assertion now follows from Proposition 2.6(a). \square

2.9. Massey products in degree four. Massey products occur in degrees four and higher. In degree four, one may obtain only triple Massey products of elements of degree one. The *Massey product* of a triplet $[x], [y], [z] \in A_1$ satisfying

$$[x] \wedge [y] = 0 \quad \text{and} \quad [y] \wedge [z] = 0,$$

is a subset of A_4 , defined as follows:

$$\langle [x], [y], [z] \rangle = \{[\pi_{xy}^3 \wedge z + x \wedge \pi_{yz}^3] \mid \partial_3^K(\pi_{xy}^3) = x \wedge y \text{ and } \partial_3^K(\pi_{yz}^3) = y \wedge z\},$$

for some $\pi_{xy}^3, \pi_{yz}^3 \in K_3$.

Remark 2.10. Let $[x], [y], [z] \in A_1$ with $[x] \wedge [y] = 0$ and $[y] \wedge [z] = 0$. Choose π_{xy}^3 and π_{yz}^3 in K_3 such that $\partial_3^K(\pi_{xy}^3) = x \wedge y$ and $\partial_3^K(\pi_{yz}^3) = y \wedge z$. Then, every element of $\langle [x], [y], [z] \rangle$ is of the form

$$[(\pi_{xy}^3 + p_{xy}^3) \wedge z + x \wedge (\pi_{yz}^3 + p_{yz}^3)] = [\pi_{xy}^3 \wedge z + x \wedge \pi_{yz}^3] + [p_{xy}^3 \wedge z] + [x \wedge p_{yz}^3],$$

for some p_{xy}^3 and $p_{yz}^3 \in \text{Ker } \partial_3^K$. Therefore,

$$\langle [x], [y], [z] \rangle = [\pi_{xy}^3 \wedge z + x \wedge \pi_{yz}^3] + (A_3 \cdot [z]) + ([x] \cdot A_3).$$

The element $\pi_{xy}^3 \wedge z + x \wedge \pi_{yz}^3$ is called a *representative* of the triple Massey product $\langle [x], [y], [z] \rangle$. Here, $(A_3 \cdot [z] + [x] \cdot A_3)$ is called the *indeterminacy* of the Massey operation, see e.g., May [14] and [2, Section 7].

Let $\langle A_1, A_1, A_1 \rangle$ denote the set of elements in A_4 which are Massey products of triplets in A_1 . The next result describes the elements of this set.

Proposition 2.11. *Each element in $\langle A_1, A_1, A_1 \rangle \subseteq A_4$ has a representative*

$$\sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge p_s^1 \quad \text{such that} \quad \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1 \wedge p_s^1] = 0, \quad \text{for all } i = 1, \dots, a_1,$$

where $p_s^1 \in \text{Ker } \partial_1^K$, and \tilde{p}_{si}^1 and $\tilde{\pi}_s^3$ are defined in (2.1.3) and (2.1.4) respectively.

Proof. Let $[x], [y], [z] \in A_1$ such that $[x] \wedge [y] = 0$ and $[y] \wedge [z] = 0$. We write $[y] = \sum_{i=1}^{a_1} [z_i^1] \wedge [\alpha_i]$ for some

$\alpha_i \in R$. Since $[x] \wedge [y] = \sum_{i=1}^{a_1} [z_i^1] \wedge [x] \wedge [-\alpha_i] = 0$, we have $\sum_{i=1}^{a_1} [z_i^1] \otimes [x] \wedge [-\alpha_i] \in \text{Ker } \phi_1$. Hence, by (2.1.3)

there exist $\beta_s \in R$ such that $[x] \wedge [-\alpha_i] = \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\beta_s]$ for all i , so

$$[x] \wedge [y] = \sum_{s=1}^{a_1^2 - q_{11}} \left(\sum_{i=1}^{a_1} [z_i^1 \wedge \tilde{p}_{si}^1] \right) \wedge [\beta_s] = \sum_{s=1}^{a_1^2 - q_{11}} [\partial_3^K(\tilde{\pi}_s^3)] \wedge [\beta_s] = \left[\partial_3^K \left(\sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \beta_s \right) \right].$$

Therefore, we may choose $\pi_{xy}^3 = \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \beta_s$. Similarly, using the equality $[y] \wedge [z] = 0$, we may choose

$$\pi_{yz}^3 = \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \gamma_s, \quad \text{for some } \gamma_s \in R, \quad \text{where } [z] \wedge [\alpha_i] = \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [\gamma_s] \quad \text{for all } i.$$

It follows that any element in $\langle A_1, A_1, A_1 \rangle$ has a representative

$$\pi_{xy}^3 \wedge z + x \wedge \pi_{yz}^3 = \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge (z \wedge \beta_s - x \wedge \gamma_s).$$

If we set $p_s^1 = z \wedge \beta_s - x \wedge \gamma_s$, then for each $1 \leq i \leq a_1$ we have

$$\begin{aligned} \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1 \wedge p_s^1] &= \sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1] \wedge [z \wedge \beta_s - x \wedge \gamma_s] = \left(\sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1 \wedge \beta_s] \right) \wedge [z] + [x] \wedge \left(\sum_{s=1}^{a_1^2 - q_{11}} [\tilde{p}_{si}^1 \wedge \gamma_s] \right) \\ &= -[x] \wedge [z] \wedge [\alpha_i] + [x] \wedge [z] \wedge [\alpha_i] = 0. \end{aligned}$$

□

2.12. Products in degree four. Let $\overline{A_4}$ be a k -subspace of A_4 such that

$$A_4 = (A_1 \cdot A_3 + A_2 \cdot A_2 + \text{Span}_k \langle A_1, A_1, A_1 \rangle) \oplus \overline{A_4}$$

and set

$$(2.12.1) \quad a = \text{rank}_k (A_1 \cdot A_3 + A_2 \cdot A_2 + \text{Span}_k \langle A_1, A_1, A_1 \rangle).$$

For some choice of $z_r^4 \in \text{Ker } \partial_4^K$, let

$$(2.12.2) \quad \{z_r^4\}_{r=1, \dots, a_4 - a} \quad \text{be a basis of } \overline{A_4}.$$

In particular, every element in $\text{Ker } \partial_4^K$ can be written as a sum of elements in $\text{Im } \partial_5^K$, a linear combination of $\{z_i^1 \wedge p_i^3\}_{i=1, \dots, a_1}$, a linear combination of $\{z_\ell^2 \wedge p_\ell^2\}_{\ell=1, \dots, a_2 - q_{11}}$, a linear combination of $\{\tilde{\pi}_s^3 \wedge p_s^1\}_{s=1, \dots, a_1^2 - q_{11}}$, and a linear combination of $\{z_r^4\}_{r=1, \dots, a_4 - a}$, where $p_i^3 \in \text{Ker } \partial_3^K$, $p_\ell^2 \in \text{Ker } \partial_2^K$ and $p_s^1 \in \text{Ker } \partial_1^K$, such that p_s^1 is as in Proposition 2.11 and $\tilde{\pi}_s^3$ as in (2.1.4).

Summary 2.13. We summarize here all notations, introduced in this section, to be referred to throughout the rest of the paper:

$$\begin{aligned} A &= H(K^R) = A_0 \oplus A_1 \oplus A_2 \oplus \dots \\ A_2 &= (A_1 \cdot A_1) \oplus \overline{A_2} \\ A_3 &= (A_1 \cdot A_2) \oplus \overline{A_3} \\ A_4 &= (A_1 \cdot A_3 + A_2 \cdot A_2 + \text{Span}_k(\langle A_1, A_1, A_1 \rangle)) \oplus \overline{A_4} \end{aligned}$$

$$a_i = \text{rank}_k A_i, \text{ for } 1 \leq i \leq 4, \quad (2.0.1)$$

$$q_{ij} = \text{rank}_k A_i \cdot A_j, \text{ for } 1 \leq i \leq j \leq 4, \quad (2.0.1)$$

$$a = \text{rank}_k (A_1 \cdot A_3 + A_2 \cdot A_2 + \text{Span}_k \langle A_1, A_1, A_1 \rangle), \quad (2.12.1)$$

$$\psi: A_1 \otimes A_1 \otimes A_1 \rightarrow ((A_1 \cdot A_1) \otimes A_1) \oplus (A_1 \otimes (A_1 \cdot A_1)), \quad (2.4.2)$$

$$\psi([x] \otimes [y] \otimes [z]) = ([x] \wedge [y] \otimes [z], [x] \otimes [y] \wedge [z]) \text{ for all } [x], [y], [z] \in A_1$$

$$b = \text{rank}_k(\text{Coker } \psi), \quad (2.4.3), \text{ see also Proposition 2.6(b)}$$

$$\{[z_i^1]\}_{i=1, \dots, a_1} \text{ is a basis for } A_1, \quad (2.0.2)$$

$$\{[z_\ell^2]\}_{\ell=1, \dots, a_2 - q_{11}} \text{ is a basis for } \overline{A_2}, \quad (2.1.1)$$

$$\{[z_t^3]\}_{t=1, \dots, a_3 - q_{12}} \text{ is a basis for } \overline{A_3}, \quad (2.4.1)$$

$$\{[z_r^4]\}_{r=1, \dots, a_4 - a} \text{ is a basis for } \overline{A_4}, \quad (2.12.2)$$

$$\phi_1: A_1 \otimes A_1 \rightarrow A_2, \quad \phi_1([x] \otimes [y]) = [x] \wedge [y], \text{ for all } [x], [y] \in A_1, \quad (2.1.2)$$

$$\left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \right\}_{s=1, \dots, a_1^2 - q_{11}} \text{ is a basis of } \text{Ker } \phi_1, \quad (2.1.3)$$

$$\partial_3^K(\tilde{\pi}_s^3) = \sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{si}^1, \quad (2.1.4)$$

$$\phi_2: A_1 \otimes A_2 \rightarrow A_3, \quad \phi_2([x] \otimes [y]) = [x] \wedge [y], \text{ for all } [x] \in A_1, [y] \in A_2, \text{ Proposition 2.6}$$

$$\left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{ui}^2] \right\}_{1 \leq u \leq a_1 a_2 - a_1 q_{11} - q_{12} + b} \text{ is a basis of } B, \text{ where}$$

$$\text{Ker } \phi_2 = B \oplus \text{Span}_k \left\{ \sum_{i=1}^{a_1} [z_i^1] \otimes [\tilde{p}_{si}^1] \wedge [z_j^1] \right\}_{1 \leq j \leq a_1, 1 \leq s \leq a_1^2 - q_{11}}$$

$$\partial_4^K(\tilde{\pi}_u^4) = \sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{ui}^2, \quad (2.5.2).$$

Lemma 2.14. All the elements $z_i^1, z_\ell^2, z_t^3, z_r^4, \tilde{p}_{si}^1, \tilde{\pi}_s^3, \tilde{p}_{ui}^2$, and $\tilde{\pi}_u^4$, as in Summary 2.13, are in $\mathfrak{m}K$.

Proof. From the inclusions $\text{Ker } \partial_j^K \subseteq \mathfrak{m}K_j$ for all $j \geq 0$, we obtain that $z_i^1, z_\ell^2, z_t^3, z_r^4, \tilde{p}_{si}^1$, and \tilde{p}_{ui}^2 are in $\mathfrak{m}K$. The assertions for the preimages $\tilde{\pi}_s^3$ and $\tilde{\pi}_u^4$ follows from the inclusions $(\partial_j^K)^{-1}(\mathfrak{m}^2 K_{j-1}) \subseteq \mathfrak{m}K_j$ for $j = 3$ and 4 respectively. \square

3. TRUNCATED MINIMAL FREE RESOLUTION OF THE RESIDUE FIELD

In this section, we construct the beginning of the minimal free resolution of the residue field k over the local ring (R, \mathfrak{m}, k) , by using the Koszul complex K of R and the graded-commutative structure of the algebra $A = H(K)$ described in Section 2.

Construction 3.1. We consider the following sequence of free R -modules:

$$F : \quad F_5 \xrightarrow{\partial_5^F} F_4 \xrightarrow{\partial_4^F} F_3 \xrightarrow{\partial_3^F} F_2 \xrightarrow{\partial_2^F} F_1 \xrightarrow{\partial_1^F} F_0,$$

where

$$F_0 := K_0$$

$$F_1 := K_1$$

$$F_2 := K_2 \oplus K_0^{a_1}$$

$$F_3 := K_3 \oplus K_1^{a_1} \oplus K_0^{a_2 - q_{11}}$$

$$F_4 := K_4 \oplus K_2^{a_1} \oplus K_1^{a_2 - q_{11}} \oplus K_0^{a_3 - q_{12}} \oplus K_0^{a_1^2 - q_{11}}$$

$$F_5 := K_5 \oplus K_3^{a_1} \oplus K_2^{a_2 - q_{11}} \oplus K_1^{a_3 - q_{12}} \oplus K_0^{a_4 - a} \oplus K_1^{a_1^2 - q_{11}} \oplus K_0^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \oplus K_0^{a_1 a_2 - a_1 q_{11}}.$$

Using the elements described in Section 2, the differential maps of F are defined as follows.

$$(3.1.1) \quad \partial_1^F : K_1 \rightarrow K_0, \quad \text{is given by} \quad \partial_1^F := \partial_1^K.$$

$$(3.1.2) \quad \partial_2^F : K_2 \oplus K_0^{a_1} \rightarrow K_1, \quad \text{is given by} \quad \partial_2^F := (\partial_2^K \quad z^1 \wedge),$$

that is

$$\partial_2^F \left(\begin{pmatrix} \pi^2 \\ (\alpha_i)_{i=1, \dots, a_1} \end{pmatrix} \right) := \partial_2^K(\pi^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \alpha_i.$$

$$(3.1.3) \quad \partial_3^F : K_3 \oplus K_1^{a_1} \oplus K_0^{a_2 - q_{11}} \rightarrow K_2 \oplus K_0^{a_1}, \quad \text{is given by} \quad \partial_3^F := \begin{pmatrix} \partial_3^K & z^1 \wedge & -z^2 \wedge \\ 0 & (\partial_1^K)^{a_1} & 0 \end{pmatrix},$$

that is

$$\partial_3^F \left(\begin{pmatrix} \pi^3 \\ (\pi_i^1)_{i=1, \dots, a_1} \\ (\alpha_\ell)_{\ell=1, \dots, a_2 - q_{11}} \end{pmatrix} \right) := \begin{pmatrix} \partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 - \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \alpha_\ell \\ (\partial_1^K(\pi_i^1))_{i=1, \dots, a_1} \end{pmatrix}.$$

$$(3.1.4) \quad \partial_4^F : K_4 \oplus K_2^{a_1} \oplus K_1^{a_2 - q_{11}} \oplus K_0^{a_3 - q_{12}} \oplus K_0^{a_1^2 - q_{11}} \rightarrow K_3 \oplus K_1^{a_1} \oplus K_0^{a_2 - q_{11}}, \quad \text{is given by}$$

$$\partial_4^F := \begin{pmatrix} \partial_4^K & z^1 \wedge & z^2 \wedge & z^3 \wedge & -\tilde{\pi}^3 \wedge \\ 0 & (\partial_2^K)^{a_1} & 0 & 0 & (\tilde{p}^1 \wedge)^{a_1} \\ 0 & 0 & (\partial_1^K)^{a_2 - q_{11}} & 0 & 0 \end{pmatrix},$$

that is

$$\partial_4^F \left(\begin{pmatrix} \pi^4 \\ (\pi_i^2)_{i=1, \dots, a_1} \\ (\pi_\ell^1)_{\ell=1, \dots, a_2 - q_{11}} \\ (\alpha_t)_{t=1, \dots, a_3 - q_{12}} \\ (\beta_s)_{s=1, \dots, a_1^2 - q_{11}} \end{pmatrix} \right) := \begin{pmatrix} \partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^2 + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \pi_\ell^1 + \sum_{t=1}^{a_3 - q_{12}} z_t^3 \wedge \alpha_t - \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \beta_s \\ (\partial_2^K(\pi_i^2) + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{p}_{si}^1 \wedge \beta_s)_{i=1, \dots, a_1} \\ (\partial_1^K(\pi_\ell^1))_{\ell=1, \dots, a_2 - q_{11}} \end{pmatrix}.$$

$$(3.1.5) \quad \begin{aligned} \partial_5^F : K_5 \oplus K_3^{a_1} \oplus K_2^{a_2 - q_{11}} \oplus K_1^{a_3 - q_{12}} \oplus K_0^{a_4 - a} \oplus K_1^{a_1^2 - q_{11}} \oplus K_0^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \oplus K_0^{a_1 a_2 - a_1 q_{11}} \\ \rightarrow K_4 \oplus K_2^{a_1} \oplus K_1^{a_2 - q_{11}} \oplus K_0^{a_3 - q_{12}} \oplus K_0^{a_1^2 - q_{11}} \end{aligned}$$

is given by

$$\partial_5^F := \begin{pmatrix} \partial_5^K & z^1 \wedge & -z^2 \wedge & z^3 \wedge & z^4 \wedge & -\tilde{\pi}^3 \wedge & -\tilde{\pi}^4 \wedge & 0 \\ 0 & (\partial_3^K)^{a_1} & 0 & 0 & 0 & (\tilde{p}^1 \wedge)^{a_1} & (\tilde{p}^2 \wedge)^{a_1} & -(z^2 \wedge)^{a_1} \\ 0 & 0 & (\partial_2^K)^{a_2-q_{11}} & 0 & 0 & 0 & 0 & (z^1 \wedge)^{a_2-q_{11}} \\ 0 & 0 & 0 & (\partial_1^K)^{a_3-q_{12}} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & (\partial_1^K)^{a_1^2-q_{11}} & 0 & 0 & 0 \end{pmatrix},$$

that is

$$\partial_5^F \begin{pmatrix} \pi^5 \\ (\pi_i^3)_{i=1,\dots,a_1} \\ (\pi_\ell^2)_{\ell=1,\dots,a_2-q_{11}} \\ (\pi_t^1)_{t=1,\dots,a_3-q_{12}} \\ (\alpha_r)_{r=1,\dots,a_4-a} \\ (\pi_s'^1)_{s=1,\dots,a_1^2-q_{11}} \\ (\beta_u)_{u=1,\dots,a_1 a_2 - a_1 q_{11} - q_{12} + b} \\ (\gamma_{\ell i})_{\ell=1,\dots,a_2-q_{11}; i=1,\dots,a_1} \end{pmatrix} := \begin{pmatrix} \partial_5^K(\pi^5) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell^2 + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \pi_t^1 + \sum_{r=1}^{a_4-a} z_r^4 \wedge \alpha_r - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \pi_s'^1 - \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{\pi}_u^4 \wedge \beta_u \\ \left(\partial_3^K(\pi_i^3) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1 + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \beta_u - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \gamma_{\ell i} \right)_{i=1,\dots,a_1} \\ \left(\partial_2^K(\pi_\ell^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \gamma_{\ell i} \right)_{\ell=1,\dots,a_2-q_{11}} \\ \left(\partial_1^K(\pi_t^1) \right)_{t=1,\dots,a_3-q_{12}} \\ \left(\partial_1^K(\pi_s'^1) \right)_{s=1,\dots,a_1^2-q_{11}} \end{pmatrix}.$$

Theorem 3.2. *Let $(R, \mathfrak{m}, \mathfrak{k})$ be a local ring. The sequence F constructed in 3.1 is a truncated minimal free resolution of \mathfrak{k} over R , up to homological degree five.*

Proof. The minimality follows from Lemma 2.14. We show exactness at each degree by using the Koszul relations in the complex K , and the basis elements and maps defined in Section 2.

Exactness at degree one. $\text{Im } \partial_2^F \subseteq \text{Ker } \partial_1^F$:

$$\partial_1^F \circ \partial_2^F \begin{pmatrix} \pi^2 \\ (\alpha_i)_{i=1,\dots,a_1} \end{pmatrix} = \partial_1^K (\partial_2^K(\pi^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \alpha_i) = 0.$$

$\text{Im } \partial_2^F \supseteq \text{Ker } \partial_1^F$: If $\pi^1 \in \text{Ker } \partial_1^F$, then there exist $\pi^2 \in K_2$ and $\alpha_i \in R$ such that

$$\pi^1 = \partial_2^K(\pi^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \alpha_i, \quad \text{therefore,} \quad \pi^1 = \partial_2^F \begin{pmatrix} \pi^2 \\ (\alpha_i)_{i=1,\dots,a_1} \end{pmatrix}.$$

Exactness at degree two. $\text{Im } \partial_3^F \subseteq \text{Ker } \partial_2^F$:

$$\partial_2^F \circ \partial_3^F \begin{pmatrix} \pi^3 \\ (\pi_i^1)_{i=1,\dots,a_1} \\ (\alpha_\ell)_{\ell=1,\dots,a_2-q_{11}} \end{pmatrix} = \partial_2^F \begin{pmatrix} \partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \alpha_\ell \\ (\partial_1^K(\pi_i^1))_{i=1,\dots,a_1} \end{pmatrix}$$

$$\begin{aligned}
&= \partial_2^K \left(\partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \alpha_\ell \right) + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_1^K(\pi_i^1) \\
&= - \sum_{i=1}^{a_1} z_i^1 \wedge \partial_1^K(\pi_i^1) + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_1^K(\pi_i^1) = 0.
\end{aligned}$$

$\text{Im } \partial_3^F \supseteq \text{Ker } \partial_2^F$: If $\begin{pmatrix} \pi^2 \\ (\alpha_i)_{i=1, \dots, a_1} \end{pmatrix} \in \text{Ker } \partial_2^F$, then $\partial_2^K(\pi^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \alpha_i = 0$ in K_1 . In particular, we have $\sum_{i=1}^{a_1} [z_i^1] \wedge [\alpha_i] = 0$ in A_1 . Since $\{[z_i^1]\}_{i=1, \dots, a_1}$ is a basis of A_1 , we get $\alpha_i \in \mathfrak{m}$. It follows that for each i there exists $\pi_i^1 \in K_1$ such that

$$(3.2.1) \quad \alpha_i = \partial_1^K(\pi_i^1).$$

Hence we have

$$0 = \partial_2^K(\pi^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_1^K(\pi_i^1) = \partial_2^K \left(\pi^2 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 \right).$$

In particular, there exist $\pi^3 \in K_3$, $p_i^1 \in \text{Ker } \partial_1^K$, and $\beta_\ell \in R$ such that

$$\pi^2 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 = \partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge p_i^1 + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \beta_\ell, \quad \text{which implies}$$

$$(3.2.2) \quad \pi^2 = \partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^1 + p_i^1) + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \beta_\ell.$$

Combining (3.2.1) and (3.2.2) we get

$$\text{Ker } \partial_2^F \ni \begin{pmatrix} \pi^2 \\ (\alpha_i)_{i=1, \dots, a_1} \end{pmatrix} = \partial_3^F \begin{pmatrix} \pi^3 \\ (\pi_i^1 + p_i^1)_{i=1, \dots, a_1} \\ -(\beta_\ell)_{\ell=1, \dots, a_2-q_{11}} \end{pmatrix}.$$

Exactness at degree three. $\text{Im } \partial_4^F \subseteq \text{Ker } \partial_3^F$: We show that both components of the element

$$\partial_3^F \circ \partial_4^F \begin{pmatrix} \pi^4 \\ (\pi_i^2)_{i=1, \dots, a_1} \\ (\pi_\ell^1)_{\ell=1, \dots, a_2-q_{11}} \\ (\alpha_t)_{t=1, \dots, a_3-q_{12}} \\ (\beta_s)_{s=1, \dots, a_1^2-q_{11}} \end{pmatrix} \text{ are zero. The first component is:}$$

$$\begin{aligned}
&\partial_3^K \left(\partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^2 + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell^1 + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha_t - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \beta_s \right) \\
&\quad + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_2^K(\pi_i^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \left(\sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \beta_s \right) - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_1^K(\pi_\ell^1) \\
&= - \sum_{i=1}^{a_1} z_i^1 \wedge \partial_2^K(\pi_i^2) + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_1^K(\pi_\ell^1) - \sum_{s=1}^{a_1^2-q_{11}} \partial_3^K(\tilde{\pi}_s^3) \wedge \beta_s \\
&\quad + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_2^K(\pi_i^2) + \sum_{s=1}^{a_1^2-q_{11}} \left(\sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{si}^1 \right) \wedge \beta_s - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_1^K(\pi_\ell^1) = 0.
\end{aligned}$$

The last equality follows from the definition of $\tilde{\pi}_s^3$ in (2.1.4). As $\tilde{p}_{si}^1 \in \text{Ker } \partial_1^K$ for all i and s , we have $\partial_1^K \left(\partial_2^K(\pi_i^2) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \beta_s \right) = 0$, for each $1 \leq i \leq a_1$, so the second component is zero.

$\text{Im } \partial_4^F \supseteq \text{Ker } \partial_3^F$: If $\begin{pmatrix} \pi^3 \\ (\pi_i^1)_{i=1, \dots, a_1} \\ (\alpha_\ell)_{\ell=1, \dots, a_2-q_{11}} \end{pmatrix} \in \text{Ker } \partial_3^F$, then

$$(3.2.3) \quad \partial_3^K(\pi^3) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^1 - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \alpha_\ell = 0 \quad \text{in } K_2, \quad \text{and}$$

$$(3.2.4) \quad \partial_1^K(\pi_i^1) = 0 \quad \text{for all } 1 \leq i \leq a_1.$$

It follows that for each i , there exist $\pi_i^2 \in K_2$ and $\beta_{ij} \in R$ such that

$$(3.2.5) \quad \pi_i^1 = \partial_2^K(\pi_i^2) + \sum_{j=1}^{a_1} z_j^1 \wedge \beta_{ij}.$$

Thus (3.2.3) becomes:

$$(3.2.6) \quad \partial_3^K\left(\pi^3 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^2\right) + \sum_{i=1}^{a_1} z_i^1 \wedge \left(\sum_{j=1}^{a_1} z_j^1 \wedge \beta_{ij}\right) - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \alpha_\ell = 0.$$

In A_2 we obtain $\sum_{i,j=1}^{a_1} [z_i^1] \wedge [z_j^1] \wedge [\beta_{ij}] - \sum_{\ell=1}^{a_2-q_{11}} [z_\ell^2] \wedge [\alpha_\ell] = 0$. By Proposition 2.3, we get $[\alpha_\ell] = 0$ for all ℓ and $\sum_{j=1}^{a_1} [z_j^1] \wedge [\beta_{ij}] = \sum_{s=1}^{a_1^2-q_{11}} [\tilde{p}_{si}^1] \wedge [\beta'_s]$ for some β'_s in R . In particular, there exist $\pi_\ell'^1 \in K_1$ and $\pi_i'^2 \in K_2$ such that

$$(3.2.7) \quad \alpha_\ell = \partial_1^K(\pi_\ell'^1), \quad \text{and} \\ \sum_{j=1}^{a_1} z_j^1 \wedge \beta_{ij} = \partial_2^K(\pi_i'^2) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \beta'_s.$$

The equation (3.2.5) becomes:

$$(3.2.8) \quad \pi_i^1 = \partial_2^K(\pi_i^2 + \pi_i'^2) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \beta'_s,$$

and equation (3.2.6) now becomes:

$$\begin{aligned} 0 &= \partial_3^K\left(\pi^3 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^2\right) + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_2^K(\pi_i'^2) + \sum_{i=1}^{a_1} \sum_{s=1}^{a_1^2-q_{11}} z_i^1 \wedge \tilde{p}_{si}^1 \wedge \beta'_s - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_1^K(\pi_\ell'^1) \\ &= \partial_3^K\left(\pi^3 - \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^2 + \pi_i'^2) - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell'^1 + \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \beta'_s\right), \end{aligned}$$

where the second equality uses the definition of $\tilde{\pi}_s^3$ from (2.1.4). Hence, there exist $\pi^4 \in K_4$, $p_i^2 \in \text{Ker } \partial_2^K$, $p_\ell^1 \in \text{Ker } \partial_1^K$, and $\alpha'_t \in R$ such that

$$\begin{aligned} &\pi^3 - \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^2 + \pi_i'^2) - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell'^1 + \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \beta'_s \\ &= \partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge p_i^2 + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge p_\ell^1 + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha'_t. \end{aligned}$$

Thus,

$$(3.2.9) \quad \pi^3 = \partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^2 + \pi_i'^2 + p_i^2) + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge (\pi_\ell'^1 + p_\ell^1) + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha'_t - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \beta'_s.$$

The equations (3.2.7), (3.2.8), and (3.2.9), now yield

$$\text{Ker } \partial_3^F \ni \begin{pmatrix} \pi^3 \\ (\pi_i^1)_{i=1, \dots, a_1} \\ (\alpha_\ell)_{\ell=1, \dots, a_2-q_{11}} \end{pmatrix} = \partial_4^F \begin{pmatrix} \pi^4 \\ (\pi_i^2 + \pi_i'^2 + p_i^2)_{i=1, \dots, a_1} \\ (\pi_\ell'^1 + p_\ell^1)_{\ell=1, \dots, a_2-q_{11}} \\ (\alpha_t')_{t=1, \dots, a_3-q_{12}} \\ (\beta_s')_{s=1, \dots, a_1^2-q_{11}} \end{pmatrix}.$$

Exactness at degree four. $\text{Im } \partial_5^F \subseteq \text{Ker } \partial_4^F$: We show that all three components of the element

$$\partial_4^F \circ \partial_5^F \begin{pmatrix} \pi^5 \\ (\pi_i^3)_{i=1, \dots, a_1} \\ (\pi_\ell^2)_{\ell=1, \dots, a_2-q_{11}} \\ (\pi_t^1)_{t=1, \dots, a_3-q_{12}} \\ (\alpha_r)_{r=1, \dots, a_4-a} \\ (\pi_s'^1)_{s=1, \dots, a_1^2-q_{11}} \\ (\beta_u)_{u=1, \dots, a_1 a_2 - a_1 q_{11} - q_{12} + b} \\ (\gamma_{\ell i})_{\ell=1, \dots, a_2-q_{11}; i=1, \dots, a_1} \end{pmatrix} \text{ are zero. The first component is:}$$

$$\begin{aligned} & \partial_4^K \left(\partial_5^K(\pi^5) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell^2 + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \pi_t^1 + \sum_{r=1}^{a_4-a} z_r^4 \wedge \alpha_r - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \pi_s'^1 - \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{\pi}_u^4 \wedge \beta_u \right) \\ & + \sum_{i=1}^{a_1} z_i^1 \wedge \left(\partial_3^K(\pi_i^3) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1 + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \beta_u - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \gamma_{\ell i} \right) \\ & + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \left(\partial_2^K(\pi_\ell^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \gamma_{\ell i} \right) + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \partial_1^K(\pi_t^1) - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \partial_1^K(\pi_s') \\ & = - \sum_{i=1}^{a_1} z_i^1 \wedge \partial_3^K(\pi_i^3) - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_2^K(\pi_\ell^2) - \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \partial_1^K(\pi_t^1) - \sum_{s=1}^{a_1^2-q_{11}} \partial_3^K(\tilde{\pi}_s^3) \wedge \pi_s'^1 \\ & + \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \partial_1^K(\pi_s'^1) - \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \partial_4^K(\tilde{\pi}_u^4) \wedge \beta_u + \sum_{i=1}^{a_1} z_i^1 \wedge \partial_3^K(\pi_i^3) + \sum_{s=1}^{a_1^2-q_{11}} \left(\sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{si}^1 \right) \wedge \pi_s'^1 \\ & + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \left(\sum_{i=1}^{a_1} z_i^1 \wedge \tilde{p}_{ui}^2 \right) \wedge \beta_u - \sum_{\ell=1}^{a_2-q_{11}} \sum_{i=1}^{a_1} z_i^1 \wedge z_\ell^2 \wedge \gamma_{\ell i} + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \partial_2^K(\pi_\ell^2) \\ & + \sum_{\ell=1}^{a_2-q_{11}} \sum_{i=1}^{a_1} z_\ell^2 \wedge z_i^1 \wedge \gamma_{\ell i} + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \partial_1^K(\pi_t^1) - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \partial_1^K(\pi_s'^1) = 0, \end{aligned}$$

by definitions of $\tilde{\pi}_s^3$, \tilde{p}_{si}^1 , $\tilde{\pi}_u^4$, and \tilde{p}_{ui}^2 . For each $1 \leq i \leq a_1$, as \tilde{p}_{si}^1 is in $\text{Ker } \partial_1^K$ and \tilde{p}_{ui}^2 is in $\text{Ker } \partial_2^K$, we have

$$\begin{aligned} & \partial_2^K \left(\partial_3^K(\pi_i^3) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1 + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \beta_u - \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \gamma_{\ell i} \right) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \partial_1^K(\pi_s'^1) \\ & = - \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \partial_1^K(\pi_s'^1) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \partial_1^K(\pi_s'^1) = 0. \end{aligned}$$

Therefore, the second component is zero. For each $1 \leq \ell \leq a_2 - q_{11}$, $\partial_1^K(\partial_2^K(\pi_\ell^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \gamma_{\ell i}) = 0$. Thus, the third component is zero.

$$\text{Im } \partial_5^F \supseteq \text{Ker } \partial_4^F: \text{ If } \begin{pmatrix} \pi^4 \\ (\pi_i^2)_{i=1, \dots, a_1} \\ (\pi_\ell^1)_{\ell=1, \dots, a_2-q_{11}} \\ (\alpha_t)_{t=1, \dots, a_3-q_{12}} \\ (\beta_s)_{s=1, \dots, a_1^2-q_{11}} \end{pmatrix} \in \text{Ker } \partial_4^F, \text{ then}$$

$$(3.2.10) \quad \partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^2 + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell^1 + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha_t - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \beta_s = 0,$$

$$(3.2.11) \quad \partial_2^K(\pi_i^2) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \beta_s = 0 \quad \text{for all } 1 \leq i \leq a_1, \quad \text{and}$$

$$(3.2.12) \quad \partial_1^K(\pi_\ell^1) = 0 \quad \text{for all } 1 \leq \ell \leq a_2 - q_{11}.$$

The equality (3.2.12) implies that there exist $\pi_\ell'^2 \in K_2$ and $\gamma_{i\ell} \in R$ such that

$$(3.2.13) \quad \pi_\ell^1 = \partial_2^K(\pi_\ell'^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \gamma_{i\ell}.$$

In $A_1^{a_1}$ the equality (3.2.11) becomes $\sum_{s=1}^{a_1^2-q_{11}} ([\tilde{p}_{s1}^1, \dots, \tilde{p}_{sa_1}^1]) \wedge [\beta_s] = 0$. Applying Lemma 2.2 we obtain $[\beta_s] = 0$ for all s , thus

$$(3.2.14) \quad \beta_s = \partial_1^K(\pi_s'^1) \quad \text{for some } \pi_s'^1 \in K_1.$$

The equality (3.2.11) now becomes

$$\partial_2^K(\pi_i^2) + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \partial_1^K(\pi_s'^1) = \partial_2^K\left(\pi_i^2 - \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1\right) = 0.$$

Therefore, there exist π_i^3 in K_3 , δ_{ijk} and $\gamma'_{i\ell}$ in R such that for all i we have:

$$(3.2.15) \quad \pi_i^2 = \partial_3^K(\pi_i^3) + \sum_{j,k=1}^{a_1} z_j^1 \wedge z_k^1 \wedge \delta_{ijk} + \sum_{\ell=1}^{a_2-q_{12}} z_\ell^2 \wedge \gamma'_{i\ell} + \sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1.$$

Putting together (3.2.13), (3.2.14), and (3.2.15), into the equality (3.2.10) we get:

$$\begin{aligned} 0 &= \partial_4^K(\pi^4) + \sum_{i=1}^{a_1} z_i^1 \wedge \left(\sum_{s=1}^{a_1^2-q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1 + \partial_3^K(\pi_i^3) + \sum_{j,k=1}^{a_1} z_j^1 \wedge z_k^1 \wedge \delta_{ijk} + \sum_{\ell=1}^{a_2-q_{12}} z_\ell^2 \wedge \gamma'_{i\ell} \right) \\ &\quad + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \left(\partial_2^K(\pi_\ell'^2) + \sum_{i=1}^{a_1} z_i^1 \wedge \gamma_{i\ell} \right) + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha_t - \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \partial_1^K(\pi_s'^1) \\ &= \partial_4^K\left(\pi^4 + \sum_{s=1}^{a_1^2-q_{11}} \tilde{\pi}_s^3 \wedge \pi_s'^1 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge \pi_\ell'^2\right) \\ &\quad + \left(\sum_{i,j,k=1}^{a_1} z_i^1 \wedge z_j^1 \wedge z_k^1 \wedge \delta_{ijk} + \sum_{i=1}^{a_1} \sum_{\ell=1}^{a_2-q_{11}} z_i^1 \wedge z_\ell^2 \wedge (\gamma'_{i\ell} + \gamma_{i\ell}) + \sum_{t=1}^{a_3-q_{12}} z_t^3 \wedge \alpha_t \right). \end{aligned}$$

In A_3 this reduces to:

$$\sum_{i,j,k=1}^{a_1} [z_i^1] \wedge [z_j^1] \wedge [z_k^1] \wedge [\delta_{ijk}] + \sum_{i=1}^{a_1} \sum_{\ell=1}^{a_2-q_{11}} [z_i^1] \wedge [z_\ell^2] \wedge [\gamma'_{i\ell} + \gamma_{i\ell}] + \sum_{t=1}^{a_3-q_{12}} [z_t^3] \wedge [\alpha_t] = 0.$$

By Proposition 2.8, for each t there exists $\pi_t''^1 \in K_1$, and for each i there exist elements $\delta'_u, \varepsilon_{js}$ in R , and $\pi_i'^3 \in K_3$ such that

$$(3.2.16) \quad \alpha_t = \partial_1^K(\pi_t''^1), \quad \text{and}$$

$$\sum_{j,k=1}^{a_1} z_j^1 \wedge z_k^1 \wedge \delta_{ijk} + \sum_{\ell=1}^{a_2-q_{11}} z_\ell^2 \wedge (\gamma_{i\ell} + \gamma'_{i\ell}) = \partial_3^K(\pi_i^3) + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \delta'_u + \sum_{j=1}^{a_1} \sum_{s=1}^{a_1^2 - q_{11}} \tilde{p}_{si}^1 \wedge z_j^1 \wedge \varepsilon_{js}.$$

Thus, (3.2.10) further becomes:

$$\begin{aligned} 0 &= \partial_4^K \left(\pi^4 + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \pi_s'^1 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \pi_\ell'^2 \right) \\ &\quad + \sum_{i=1}^{a_1} z_i^1 \wedge \left(\partial_3^K(\pi_i^3) + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \delta'_u + \sum_{j=1}^{a_1} \sum_{s=1}^{a_1^2 - q_{11}} \tilde{p}_{si}^1 \wedge z_j^1 \wedge \varepsilon_{js} \right) + \sum_{t=1}^{a_3 - q_{12}} z_t^3 \wedge \partial_1^K(\pi_t''^1) \\ &= \partial_4^K \left(\pi^4 + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \pi_s'^1 - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \pi_\ell'^2 \right. \\ &\quad \left. - \sum_{i=1}^{a_1} z_i^1 \wedge \pi_i^3 + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{\pi}_u^4 \wedge \delta'_u + \sum_{j=1}^{a_1} \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge z_j^1 \wedge \varepsilon_{js} - \sum_{t=1}^{a_3 - q_{12}} z_t^3 \wedge \pi_t''^1 \right). \end{aligned}$$

The second equality above follows from definitions of $\tilde{\pi}_s^3$ and $\tilde{\pi}_u^4$ in (2.1.4) and (2.5.2). Therefore, there exist $\pi^5 \in K_5$, $\delta_r \in R$, $p_i^3 \in \text{Ker } \partial_3^K$, $p_\ell^2 \in \text{Ker } \partial_2^K$ and $p_s^1 \in \text{Ker } \partial_1^K$ with p_s^1 as in Proposition 2.11, such that

$$\begin{aligned} &\pi^4 - \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^3 + \pi_i'^3) + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \pi_\ell'^2 - \sum_{t=1}^{a_3 - q_{12}} z_t^3 \wedge \pi_t''^1 \\ &\quad + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge (\pi_s'^1 + \sum_{j=1}^{a_1} z_j^1 \wedge \varepsilon_{js}) + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{\pi}_u^4 \wedge \delta'_u \\ &= \partial_5^K(\pi^5) + \sum_{i=1}^{a_1} z_i^1 \wedge p_i^3 + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge p_\ell^2 + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge p_s^1 + \sum_{r=1}^{a_4 - a} z_r^4 \wedge \delta_r. \end{aligned}$$

This implies:

$$\begin{aligned} (3.2.17) \quad &\pi^4 = \partial_5^K(\pi^5) + \sum_{i=1}^{a_1} z_i^1 \wedge (\pi_i^3 + \pi_i'^3 + p_i^3) - \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge (\pi_\ell'^2 - p_\ell^2) + \sum_{t=1}^{a_3 - q_{12}} z_t^3 \wedge \pi_t''^1 + \sum_{r=1}^{a_4 - a} z_r^4 \wedge \delta_r \\ &\quad - \sum_{s=1}^{a_1^2 - q_{11}} \tilde{\pi}_s^3 \wedge \left(\pi_s'^1 + \sum_{j=1}^{a_1} z_j^1 \wedge \varepsilon_{js} - p_s^1 \right) - \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{\pi}_u^4 \wedge \delta'_u. \end{aligned}$$

By (3.2.16), the expression (3.2.15) becomes:

$$\begin{aligned} (3.2.18) \quad &\pi_i^2 = \partial_3^K(\pi_i^3) + \sum_{j,k=1}^{a_1} z_j^1 \wedge z_k^1 \wedge \delta_{ijk} + \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \gamma'_{i\ell} + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{p}_{si}^1 \wedge \pi_s'^1 \\ &= \partial_3^K(\pi_i^3 + \pi_i'^3) + \sum_{s=1}^{a_1^2 - q_{11}} \tilde{p}_{si}^1 \wedge \left(\pi_s'^1 + \sum_{j=1}^{a_1} z_j^1 \wedge \varepsilon_{js} \right) + \sum_{u=1}^{a_1 a_2 - a_1 q_{11} - q_{12} + b} \tilde{p}_{ui}^2 \wedge \delta'_u - \sum_{\ell=1}^{a_2 - q_{11}} z_\ell^2 \wedge \gamma_{i\ell}. \end{aligned}$$

We conclude that by using (3.2.13), (3.2.14), (3.2.16), (3.2.17), (3.2.18), and Proposition 2.11, the chosen kernel element is in the image of ∂_5^F :

$$\text{Ker } \partial_4^F \ni \begin{pmatrix} \pi^4 \\ (\pi_i^2)_{i=1, \dots, a_1} \\ (\pi_\ell^1)_{\ell=1, \dots, a_2 - q_{11}} \\ (\alpha_t)_{t=1, \dots, a_3 - q_{12}} \\ (\beta_s)_{s=1, \dots, a_1^2 - q_{11}} \end{pmatrix} = \partial_5^F \begin{pmatrix} \pi^5 \\ (\pi_i^3 + \pi_i'^3 + p_i^3)_{i=1, \dots, a_1} \\ (\pi_\ell'^2 - p_\ell^2)_{\ell=1, \dots, a_2 - q_{11}} \\ (\pi_t''^1)_{t=1, \dots, a_3 - q_{12}} \\ (\delta_r)_{r=1, \dots, a_4 - a} \\ \left(\pi_s'^1 + \sum_{j=1}^{a_1} z_j^1 \wedge \varepsilon_{js} - p_s^1 \right)_{s=1, \dots, a_1^2 - q_{11}} \\ (\delta'_u)_{u=1, \dots, a_1 a_2 - a_1 q_{11} - q_{12} + b} \\ (\gamma_{\ell i})_{\ell=1, \dots, a_2 - q_{11}; i=1, \dots, a_1} \end{pmatrix}.$$

□

Remark 3.3. If all multiplications on the algebra A are trivial, that is, $q_{ij} = 0$ for all $i, j \geq 1$, and all Massey operations are zero, Golod's construction in [8] gives a minimal free resolution of the residue field \mathbf{k} of a local ring R in terms of the Koszul complex K of R . Without these assumptions, our complex F in Construction 3.1 generalizes Golod's resolution up to degree five.

4. APPLICATIONS OF THE CONSTRUCTION

For a local ring R of embedding dimension n and codepth c , set $\beta_i := \text{rank}_{\mathbf{k}} \text{Tor}_i^R(\mathbf{k}, \mathbf{k})$ for $i \geq 0$ for the Betti numbers of \mathbf{k} over R . Let $\sum_{i=0}^{\infty} b_i t^i$ denote the series on the right hand side of the inequality (1.0.1). The sequence $\mathcal{P} := \{\mathcal{P}_i\}_{i \geq 0}$ defined by:

$$(4.0.1) \quad \mathcal{P}_i := b_i - \beta_i, \quad \text{for all } i \geq 0,$$

gives the coefficients of the series $\mathcal{P}(t)$ defined in the Introduction. The ring R is Golod if and only if $\mathcal{P}_i = 0$ for all $i \geq 0$. Our goal in this section is to give a description of the sequence $\{\mathcal{P}_i\}_{0 \leq i \leq 5}$ in terms of the invariants of the multiplicative structure of the algebra $A = H(K^R)$ and discuss various consequences of Theorem 3.2. First, using Theorem 3.2, we explicitly describe the Betti numbers β_i in terms of those invariants, up to degree five.

Corollary 4.1. *Let R be a local ring of embedding dimension n . Let a_i, q_{ij}, a and b be as in Summary 2.13. Then the following equalities hold:*

$$\begin{aligned} \beta_0 &= 1, & \beta_1 &= n, & \beta_2 &= \binom{n}{2} + a_1, \\ \beta_3 &= \binom{n}{3} + na_1 + a_2 - q_{11}, & \beta_4 &= \binom{n}{4} + \binom{n}{2} a_1 + na_2 + a_3 + a_1^2 - (n+1)q_{11} - q_{12}, \\ \beta_5 &= \binom{n}{5} + \binom{n}{3} a_1 + \binom{n}{2} a_2 + na_3 + a_4 + na_1^2 + 2a_1 a_2 - \left(\binom{n+1}{2} + 2a_1 \right) q_{11} - (n+1)q_{12} + b - a. \end{aligned}$$

Proof. This is a direct consequence of Theorem 3.2. The formulas for β_0, \dots, β_3 are clear, and we simplify the following expressions for β_4 and β_5 to obtain the ones in the statement:

$$\begin{aligned} \beta_4 &= \binom{n}{4} + \binom{n}{2} a_1 + a_1^2 - q_{11} + n(a_2 - q_{11}) + a_3 - q_{12} \\ \beta_5 &= \binom{n}{5} + \binom{n}{3} a_1 + n(a_1^2 - q_{11}) + a_1 a_2 - a_1 q_{11} - q_{12} + b \\ &\quad + \binom{n}{2} (a_2 - q_{11}) + a_1 a_2 - a_1 q_{11} + n(a_3 - q_{12}) + a_4 - a. \end{aligned}$$

□

Proposition 4.2. *Let R be a local ring of embedding dimension n and codepth c . Let a_i, q_{ij}, a , and b be as in Summary 2.13. The sequence \mathcal{P} from (4.0.1) satisfies the following equalities:*

$$\mathcal{P}_0 = \mathcal{P}_1 = \mathcal{P}_2 = 0, \quad \mathcal{P}_3 = q_{11}, \quad \mathcal{P}_4 = (n+1)q_{11} + q_{12}, \quad \text{and}$$

$$\mathcal{P}_5 = \left(\binom{n+1}{2} + 2a_1 \right) q_{11} + (n+1)q_{12} + a - b.$$

Proof. Recall that the right hand side of the inequality (1.0.1) is $\sum_{i=1}^{\infty} b_i t^i = \frac{(1+t)^n}{1 - \sum_{i=1}^c a_i t^{i+1}}$. Comparing the coefficients on both sides of (1.0.1) we have the following recursive formulas for b_i :

$$b_0 = 1, \quad b_1 = n, \quad b_i = \sum_{j=1}^{i-1} a_j b_{i-j-1} + \binom{n}{i}, \quad \text{for all } i \geq 2.$$

In particular, we obtain:

$$\begin{aligned} b_2 &= \binom{n}{2} + a_1, & b_4 &= \binom{n}{4} + \binom{n}{2} a_1 + n a_2 + a_3 + a_1^2, \\ b_3 &= \binom{n}{3} + n a_1 + a_2, & b_5 &= \binom{n}{5} + \binom{n}{3} a_1 + \binom{n}{2} a_2 + n a_3 + a_4 + n a_1^2 + 2 a_1 a_2. \end{aligned}$$

Now, the expression for $\mathcal{P}_i = b_i - \beta_i$ for $0 \leq i \leq 5$ follows from Corollary 4.1.

Alternatively, assume $q_{ij} = 0$ for all $1 \leq i+j \leq 4$, and Massey products are also trivial, we have $a = b = 0$. In Construction 3.1 we obtain the maximum possible values of the Betti numbers, so the differences between the actual Betti numbers and these maximum values are exactly the \mathcal{P}_i 's in the proposition. \square

Corollary 4.3. *Let R be a local ring, a_i , q_{ij} , a , and b be as in Summary 2.13 and \mathcal{P} as in (4.0.1).*

- (a) *If $q_{11} = q_{12} = 0$, then $\mathcal{P}_i = 0$ for all $0 \leq i \leq 4$ and $\mathcal{P}_5 = a$.*
- (b) *If $q_{11} = q_{12} = q_{13} = q_{22} = 0$, then $\mathcal{P}_5 = \text{rank}_k(\text{Span}_k\langle A_1, A_1, A_1 \rangle)$.*
- (c) *If $\text{codepth } R = 4$, then R is Golod if and only if $\mathcal{P}_i = 0$ for all $1 \leq i \leq 5$.*

Proof. (a) and (b) are straightforward from Proposition 4.2. For (c), result in [8] showed that R is Golod if and only if all products on A and all ternary Massey products are trivial. If $\text{codepth } R = 4$, then the r -ary Massey products are trivial for all $r \geq 4$. Thus, part (c) follows from Proposition 4.2. \square

Remark 4.4. A result of Burke [6, Corollary 6.10] implies that for a local ring R of codepth c the following implication holds:

$$\text{If } \mathcal{P}_i = 0 \text{ for all } 0 \leq i \leq c+1, \text{ then } \mathcal{P}_i = 0 \text{ for all } i \geq 0.$$

Thus, Corollary 4.3(c) is a consequence of this result as well.

We now examine some local rings with rational Poincaré series of certain forms and describe $P_k^R(t)$ in terms of the algebraic invariants of A .

Proposition 4.5. *Let (R, \mathfrak{m}, k) be a local ring of embedding dimension n and codepth c . Let a_i, q_{ij}, a , and b be as in Summary 2.13. If the Poincaré series of R is rational of the form $P_k^R(t) = (1+t)^n/d(t)$, then*

$$d(t) = \left(1 - \sum_{i=1}^c a_i t^{i+1} \right) + q_{11} t^3 + (q_{11} + q_{12}) t^4 + (q_{12} - b + a) t^5 + f(t) t^6,$$

for some $f(t) \in \mathbb{Z}[t]$.

Proof. Let \mathcal{P}_i be as in (4.0.1). Set

$$\mathcal{P}(t) = \sum_{i=0}^{\infty} \mathcal{P}_i t^i, \quad \alpha(t) = 1 - \sum_{i=1}^c a_i t^{i+1} \quad \text{and} \quad \gamma(t) = d(t) - \alpha(t).$$

Then

$$\begin{aligned} \mathcal{P}(t) &= \frac{(1+t)^n}{\alpha(t)} - \frac{(1+t)^n}{d(t)} = \frac{(1+t)^n \cdot \gamma(t)}{\alpha(t) \cdot (\alpha(t) + \gamma(t))} \iff \\ \frac{\gamma(t)}{\alpha(t) + \gamma(t)} &= \frac{\mathcal{P}(t) \cdot \alpha(t)}{(1+t)^n} \iff \frac{\gamma(t)}{\alpha(t)} = \frac{\mathcal{P}(t) \cdot \alpha(t)}{(1+t)^n - \mathcal{P}(t) \cdot \alpha(t)} \iff \gamma(t) = \frac{\mathcal{P}(t) \cdot (\alpha(t))^2}{(1+t)^n - \mathcal{P}(t) \cdot \alpha(t)}. \end{aligned}$$

We compare the coefficients of t^i for all $0 \leq i \leq 5$ on both sides of the following equality

$$\gamma(t) \cdot \left((1+t)^n - \mathcal{P}(t) \cdot \alpha(t) \right) = \mathcal{P}(t) \cdot (\alpha(t))^2.$$

By Proposition 4.2, $\mathcal{P}_0 = \mathcal{P}_1 = \mathcal{P}_2 = 0$, and thus the left and right hand sides of the above equation become:

$$\begin{aligned} \text{LHS} &= (\gamma_0 + \gamma_1 t + \gamma_2 t^2 + \gamma_3 t^3 + \gamma_4 t^4 + \gamma_5 t^5 + \dots) \\ &\quad \cdot \left(1 + nt + \binom{n}{2} t^2 + \left(\binom{n}{3} - \mathcal{P}_3 \right) t^3 + \left(\binom{n}{4} - \mathcal{P}_4 \right) t^4 + \left(\binom{n}{5} + \mathcal{P}_3 a_1 - \mathcal{P}_5 \right) t^5 + \dots \right), \\ \text{RHS} &= (\mathcal{P}_3 t^3 + \mathcal{P}_4 t^4 + \mathcal{P}_5 t^5 + \dots) \cdot (1 - 2a_1 t^2 - 2a_2 t^3 + (a_1^2 - 2a_4) t^4 + (a_1 a_2 - 2a_5) t^5 + \dots). \end{aligned}$$

It is clear that $\gamma_0 = \gamma_1 = \gamma_2 = 0$ and comparing the coefficients of t^3 we get: $\gamma_3 = \mathcal{P}_3 = q_{11}$. Comparing the coefficients of t^4 and by Proposition 4.2 we get:

$$\gamma_4 + n\gamma_3 = \mathcal{P}_4 \iff \gamma_4 = \mathcal{P}_4 - n\gamma_3 = (n+1)q_{11} + q_{12} - nq_{11} = q_{11} + q_{12}.$$

Finally, comparing the coefficients of t^5 and by Proposition 4.2 we get:

$$\begin{aligned} \gamma_5 + n\gamma_4 + \binom{n}{2} \gamma_3 &= \mathcal{P}_5 - 2a_1 \mathcal{P}_3 \iff \\ \gamma_5 &= \mathcal{P}_5 - 2a_1 \mathcal{P}_3 - n\gamma_4 - \binom{n}{2} \gamma_3 \\ &= \left(\binom{n+1}{2} + 2a_1 \right) q_{11} + (n+1)q_{12} - b + a - 2a_1 q_{11} - n(q_{11} + q_{12}) - \binom{n}{2} q_{11} \\ &= q_{12} - b + a. \end{aligned}$$

Therefore, the expression for $d(t) = \alpha(t) + \gamma(t)$ in the statement holds. \square

There are many classes of local rings for which the Poincaré series is rational of the form $P_k^R(t) = (1+t)^n/d(t)$. In light of Proposition 4.5, we write the coefficients of the polynomial $d(t)$ in terms of the invariants a_i, q_{ij}, b and a for some special cases and provide a *uniform expression* of the Poincaré series in these cases.

Corollary 4.6. *If (R, \mathfrak{m}, k) is a local ring of embedding dimension n and codepth at most 3, and a_i, q_{ij}, b are as in Summary 2.13, then*

$$P_k^R(t) = \frac{(1+t)^{n-1}}{1 - t - (a_1 - 1)t^2 - (a_3 - q_{11})t^3 + q_{12}t^4 - bt^5}.$$

Proof. By [3, Theorem 3.5] the Poincaré series of the ring R is rational given by $P_k^R(t) = (1+t)^n/d(t)$ with $\deg d(t) \leq 6$ and $d(-1) = 0$. Since $\text{codepth } R \leq 3$ we have $a_4 = 0$ and $a = 0$. Thus, Proposition 4.5 gives:

$$P_k^R(t) = \frac{(1+t)^n}{1 - a_1 t^2 - (a_2 - q_{11})t^3 - (a_3 - q_{11} - q_{12})t^4 - (b - q_{12})t^5 - bt^6}.$$

Simplifying the fraction by the common factor $(1+t)$, we get the desired conclusion. \square

Remark 4.7. In [5, Lemma 3.6] Avramov defined the invariant τ for non-Gorenstein ring R of codepth 3 as follows: $\tau = 1$ if R is of class **T**, and $\tau = 0$ otherwise. By comparing [5, (3.6.2)] and the Poincaré expression in Corollary 4.6 we see that τ is our $b = \text{rank}_k(\text{Coker } \psi)$ in (2.4.3), which is a multiplicative invariant of the homology algebra A . It follows from [3, Theorem 3.5] that $b = 1$ for a ring of class **C**(3) or **T**, and $b = 0$ for rings of class **C**(1), **C**(2), **S**, **B**, **G**(r), and **H**(q_{11}, q_{12}).

Corollary 4.8. *If (R, \mathfrak{m}, k) is a Gorenstein local ring, not a complete intersection, of embedding dimension n and codepth 4, and a_i, q_{ij}, a, b are as in Summary 2.13, then*

$$P_k^R(t) = \frac{(1+t)^{n-2}}{1 - 2t - (a_1 - 3)t^2 + (q_{11} - 2)t^3 + (-q_{11} + q_{12} - 1)t^4 - (q_{11} - q_{12} - b + a - 1)t^5}.$$

Proof. Since R is Gorenstein we have $a_2 = 2a_1 - 2, a_3 = a_1$, and $a_4 = 1$. The statement follows from [3, Theorem 3.5] and Proposition 4.5. \square

Remark 4.9. Using the Avramov's notation from [3] for the classification of Gorenstein local rings R of codepth 4 given by Kustin and Miller [11] and comparing the Poincaré expressions from [3, Theorem 3.5] and Corollary 4.8, we obtain the following table of algebra invariants of a Gorenstein local ring R :

Class	q_{11}	q_{12}	q_{22}	q_{13}	a	b
C(4)	6	4	1	1	1	4
GT	3	3	1	1	1	1
GS	0	0	1	1	1	0
GH(p)	p	$p+1$	1	1	1	0

We provide more examples of rings for which we can calculate the multiplicative invariants q_{ij}, a, b from the Poincaré series of the ring.

Corollary 4.10. *Let k be a field, I be an ideal of $k[x, y, z, w]$ such that $(x, y, z, w)^3 \subseteq I \subseteq (x, y, z, w)^2$, and set $R = k[x, y, z, w]/I$. Let a_i, q_{ij}, a, b be as in Summary 2.13. If R has $a_4 = 3$ and its Poincaré series is of the form $P_k^R(t) = \frac{1}{(1-t)(1-3t)}$, then*

$$q_{11} = a_2 - 8, \quad q_{12} = 8, \quad a = 3, \quad \text{and} \quad b = 0.$$

Proof. By hypothesis,

$$P_k^R(t) = \frac{(1+t)^4}{(1-t)(1-3t)(1+t)^4} = \frac{(1+t)^4}{1-7t^2-8t^3+3t^4+8t^5+3t^6}.$$

Proposition 4.5 thus gives:

$$(4.10.1) \quad a_1 = 7, \quad a_2 - q_{11} = 8, \quad a_3 - q_{11} - q_{12} = -3, \quad \text{and} \quad a_4 - q_{12} + b - a = -8.$$

From the second equality we obtain $q_{11} = a_2 - 8$ and from the third we get $q_{12} = a_3 - a_2 + 11$. Since $a_4 = 3$ and $a_1 = 7$, we get $a_2 = a_3 + 3$ and thus $q_{12} = 8$. The last equality in (4.10.1) becomes $a - b = 3$. By the definitions of a and b we have $0 \leq a \leq a_4 = 3$ and $b \geq 0$, thus $a = 3$ and $b = 0$. \square

Examples 4.11. Rings satisfying the hypotheses of Corollary 4.10 are discussed by Yoshino in [17] and Christensen and Veliche in [7]. For examples, consider the following ideals in $Q = \mathbb{Q}[x, y, z, w]$:

$$\begin{aligned} I_1 &= (yw, xw + zw + w^2, z^2 + w^2, xz + zw + w^2, y^2 + yz, xy + zw, x^2 + zw), \\ I_2 &= (zw + w^2, yw, z^2 + w^2, yz + xw + w^2, xz + w^2, xy + y^2 + xw + w^2, x^2 + xw + w^2), \\ I_3 &= (zw, yw, xw - w^2, yz, xz, xy - z^2, x^2 - y^2), \\ I_4 &= (w^2, yw + zw, xw, yz + z^2, y^2 + zw, xy + xz, x^2 + zw). \end{aligned}$$

The rings Q/I_i satisfy the hypotheses of Corollary 4.10 and have $a_2 = 10 + i$, for $1 \leq i \leq 4$.

Examples 4.12. In an unpublished note, Roos [16], inspired by a paper of Katthän [10], constructed several examples of non-Golod rings R of codepth 4 with trivial algebra multiplications on A . We provide here two of them. For each one, there exists a Golod homomorphism from a complete intersection ring, hence it has rational Poincaré series. The algebra multiplication on A was checked using the **DGAlgebras** package [15] of **Macaulay2** [13]. Proposition 4.5 confirms that indeed $q_{11} = q_{12} = 0$, and moreover it gives us the exact size of the space generated by the triple Massey products $\langle A_1, A_1, A_1 \rangle$, known to be nonzero.

Consider the following ideals in $Q = \mathbb{Q}[x, y, z, w]$:

$$\begin{aligned} J_1 &= (w^3, xy^2, xz^2 + yz^2, x^2w, x^2y + y^2w, y^2z + z^2w), \\ J_2 &= (w^3, xy^2, xz^2 + yz^2, x^2w + zw^2, y^2w + xzw, y^2z + yz^2). \end{aligned}$$

The rings Q/J_i with $i = 1, 2$ are non-Artinian of codepth 4 with

$$\begin{aligned} P_{Q/J_i}^Q(t) &= 1 + 6t + (10 + i)t^2 + (7 + i)t^3 + 2t^4, \\ P_k^{Q/J_i}(t) &= \frac{(1+t)^4}{1 - 6t^2 - (10 + i)t^3 - (7 + i)t^4 - t^5 + t^6}. \end{aligned}$$

For both rings Q/J_i , $q_{11} = 0 = q_{12}$ by Proposition 4.5, hence $b = 0$ and $a = 1$. Since $q_{22} = q_{13} = 0$, the space spanned by the ternary Massey products $\langle A_1, A_1, A_1 \rangle$ has rank one.

By [1, Example 7.1], not all local rings have rational Poincaré series. However, for every local ring R with residue field k , there exists a unique sequence of integers $\{\varepsilon_i\}_{i \geq 0}$ such that the Poincaré series of R can be expressed as

$$P_k^R(t) = \frac{\prod_{i=1}^{\infty} (1 + t^{2i-1})^{\varepsilon_{2i-1}}}{\prod_{i=1}^{\infty} (1 - t^{2i})^{\varepsilon_{2i}}},$$

and ε_i is called the i -th deviation of R ; see for example [4, Remark 7.1.1]. Theorem 3.2 allows us to describe the first five deviations in terms of the algebraic invariants of A .

Corollary 4.13. *Let R be a local ring of embedding dimension n and a_i, q_{ij} be as in Summary 2.13. Then the first five deviations of R are*

$$\begin{aligned} \varepsilon_1 &= n, & \varepsilon_2 &= a_1, & \varepsilon_3 &= a_2 - q_{11}, \\ \varepsilon_4 &= a_3 - q_{12} + \binom{a_1}{2} - q_{11}, \\ \varepsilon_5 &= a_4 + a_1 a_2 - a_1 q_{11} - q_{12} + b - a. \end{aligned}$$

Proof. By comparing the coefficients on the left and right sides of the equality

$$\prod_{i=1}^{\infty} (1 + t^{2i-1})^{\varepsilon_{2i-1}} = \prod_{i=1}^{\infty} (1 - t^{2i})^{\varepsilon_{2i}} \cdot (\beta_0 + \beta_1 t + \beta_2 t^2 + \beta_3 t^3 + \beta_4 t^4 + \beta_5 t^5 + \dots)$$

one obtains the following relations between the Betti numbers $\{\beta_i\}_{1 \leq i \leq 5}$ and the deviations $\{\varepsilon_i\}_{1 \leq i \leq 5}$:

$$\begin{aligned} \beta_1 &= \varepsilon_1, & \beta_2 &= \varepsilon_2 + \binom{\varepsilon_1}{2}, & \beta_3 &= \varepsilon_3 + \varepsilon_2 \varepsilon_1 + \binom{\varepsilon_1}{3}, \\ \beta_4 &= \varepsilon_4 + \varepsilon_3 \varepsilon_1 + \binom{1 + \varepsilon_2}{2} + \varepsilon_2 \binom{\varepsilon_1}{2} + \binom{\varepsilon_1}{4}, \\ \beta_5 &= \varepsilon_5 + \varepsilon_4 \varepsilon_1 + \varepsilon_3 \varepsilon_2 + \varepsilon_3 \binom{\varepsilon_1}{2} + \varepsilon_2^2 \varepsilon_1 - \varepsilon_1 \binom{\varepsilon_2}{2} + \varepsilon_2 \binom{\varepsilon_1}{3} + \binom{\varepsilon_1}{5}. \end{aligned}$$

The first four relations were also given by Avramov [4, page 62] and Gulliksen and Levin [9, Proposition 3.3.4, Theorem 4.4.3]. Note that we have corrected the expression for β_4 , compared to that given in [4, page 62]. The expressions for ε_i described in the statement follow from these relations and Corollary 4.1. \square

Remark 4.14. The formulas for $\varepsilon_2, \varepsilon_3, \varepsilon_4$ in Corollary 4.13 were previously given in [2, Corollary 6.2] and [9, Proposition 3.3.4]. The expression for ε_5 obtained by Avramov in [2, Corollary 6.2] is, in our notations:

$$\varepsilon_5 = a_4 + a_1 a_2 + a_1 q_{11} - q_{12} - a_1^3 + b' - a,$$

where $b' := \text{rank}_k(\text{Ker } \Gamma)$ with

$$\begin{aligned} \Gamma : A_1 \otimes A_1 \otimes A_1 &\rightarrow A_2 \otimes A_1 \oplus A_1 \otimes A_2, \quad \text{defined by} \\ \Gamma([x], [y], [z]) &= ([x] \wedge [y], [y] \wedge [z]), \quad \text{for all } [x], [y], [z] \in A_1. \end{aligned}$$

The map ψ in (2.4.2), that defines our invariant $b = \text{rank}_k(\text{Coker } \psi)$, differs from Γ just by its codomain. Thus, one can relate b' in [2] and our b by:

$$b' = \text{rank}_k(\text{Ker } \psi) = a_1^3 - 2a_1 q_{11} + b.$$

Note that in both references [2] and [9] there is a shift in the indexing of the deviations, their ε_i is our ε_{i+1} .

5. AN EXAMPLE ILLUSTRATING THE CONSTRUCTION

In this section we consider the codepth 4 artinian local ring

$$R = \mathbb{Q}[x, y, z, w] / (x^3, y^3, z^3 - xy^2, x^2 z^2, xyz^2, y^2 w, w^2)$$

from [2, Section 7]. This ring is of a particular interest to us, since its Koszul homology algebra $A = H(K^R)$ has nontrivial multiplication and a nontrivial ternary Massey product that does not come from this multiplication. The free modules $\{F_i\}_{i=0, \dots, 5}$ are given in terms of the Koszul algebra components $\{K_i\}_{i=0, \dots, 5}$, the ranks a_i of A_i , and the multiplicative invariants q_{ij}, a and b . The differential maps of the complex F in Construction 3.1 are given in terms of elements $z_i^1, z_\ell^2, z_t^3, z_r^4, \tilde{p}_{si}^1, \tilde{\pi}_s^3, \tilde{p}_{ui}^2$, and $\tilde{\pi}_u^4$, see

Summary 2.13. We explicitly describe them all for this ring. The bases of $A_i = H_i(K)$ are computed with **Macaulay2** [13]. We use our results from Section 2 to obtain the other elements needed in Construction 3.1.

5.1. The ring R is graded artinian with $R_i = 0$ for all $i \geq 6$. The other graded components have the following basis elements:

$$\begin{aligned} R_0 : & 1 \\ R_1 : & x, y, z, w \\ R_2 : & x^2, xy, xz, xw, y^2, yz, yw, z^2, zw \\ R_3 : & x^2y, x^2z, x^2w, xyz, xyw, xz^2, xzw, y^2z, yz^2, yzw, z^3, z^2w \\ R_4 : & x^2yz, x^2yw, x^2zw, xyzw, xz^3, xz^2w, y^2z^2, yz^2w, z^4 \\ R_5 : & x^2yzw, xz^4. \end{aligned}$$

5.2. The Koszul algebra of R has the form $K = \bigwedge(R^4) \cong R \oplus R^4 \oplus R^6 \oplus R^4 \oplus R$. Let $\{T_1, T_2, T_3, T_4\}$ be the standard ordered vector basis of the free module $K_1 = R^4$. Set $T_{ij} = T_i \wedge T_j$ and consider the ordered basis $\{T_{12}, T_{13}, T_{23}, T_{14}, T_{24}, T_{34}\}$ for the free module $K_2 = \bigwedge^2(R^4) \cong R^6$. Set $T_{ijk} = T_i \wedge T_j \wedge T_k$ and consider the ordered basis $\{T_{123}, T_{124}, T_{134}, T_{234}\}$ for the free module $K_3 = \bigwedge^3(R^4) \cong R^4$. Set $T_{1234} = T_1 \wedge T_2 \wedge T_3 \wedge T_4$ and consider the basis $\{T_{1234}\}$ for the free module $K_4 = \bigwedge^4(R^4) \cong R$. By **Macaulay2**, we obtain bases of $A_i = H_i(K)$ and ranks

$$a_1 = 7, \quad a_2 = 15, \quad a_3 = 14, \quad a_4 = 5.$$

5.3. A basis of A_1 is $\{[z_i^1]\}_{i=1,\dots,7}$ where:

$$\begin{aligned} z_1^1 &= wT_4, & z_2^1 &= x^2T_1, & z_3^1 &= ywT_2, & z_4^1 &= y^2T_2, \\ z_5^1 &= y^2T_1 - z^2T_3, & z_6^1 &= yz^2T_1, & z_7^1 &= xz^2T_1. \end{aligned}$$

5.4. Using Koszul and ring relations, we obtain that the space $A_1 \cdot A_1$ has rank $q_{11} = 7$ and its basis is given by the classes of

$$\begin{aligned} z_1^1 \wedge z_2^1 &= -x^2wT_{14}, & z_1^1 \wedge z_5^1 &= z^2wT_{34}, & z_1^1 \wedge z_6^1 &= -yz^2wT_{14}, & z_1^1 \wedge z_7^1 &= -xz^2wT_{14}, \\ z_2^1 \wedge z_3^1 &= x^2yT_{12}, & z_3^1 \wedge z_5^1 &= -yz^2wT_{23}, & z_4^1 \wedge z_5^1 &= -y^2z^2T_{23}. \end{aligned}$$

Therefore, a basis of $\overline{A_2}$ is $\{[z_\ell^2]\}_{\ell=1,\dots,8}$, where

$$\begin{aligned} z_1^2 &= ywT_{24}, & z_2^2 &= y^2T_{24}, & z_3^2 &= xz^2T_{12}, & z_4^2 &= yz^2T_{13}, \\ z_5^2 &= x^2yT_{12} - xz^2T_{13}, & z_6^2 &= x^2zT_{13}, & z_7^2 &= yz^2wT_{12}, & z_8^2 &= z^4T_{23}. \end{aligned}$$

5.5. All other products among the basis elements of A_1 are zero in A_2 , except for the Koszul relations of the nonzero products $A_1 \cdot A_1$ above. It follows that the kernel of the multiplication map $\phi_1 : A_1 \otimes A_1 \rightarrow A_2$ is generated by $a_1^2 - q_{11} = 42$ elements. We record the indices of basis elements of $A_1 \cdot A_1$ by the set

$$S = \{(1, 2), (1, 5), (1, 6), (1, 7), (2, 3), (3, 5), (4, 5)\}.$$

A basis of $\text{Ker } \phi_1$, as defined in (2.1.3), is given by elements of two types:

- (1) $[z_i^1] \otimes [z_j^1]$, for $1 \leq i, j \leq 7$, $(i, j) \notin S$ and $(j, i) \notin S$;
- (2) $[z_i^1] \otimes [z_j^1] + [z_j^1] \otimes [z_i^1]$, for $(i, j) \in S$.

Therefore, according to the types above, one defines $\tilde{p}_{s(i,j)k}^1 = (\tilde{p}_{s(i,j)k}^1)_{k=1,\dots,7}$ as:

$$\begin{aligned} (1) \quad \tilde{p}_{s(i,j)k}^1 &= \begin{cases} z_j^1 & \text{if } k = i \\ 0 & \text{if } k \neq i, \end{cases} \quad \text{for all } 1 \leq i, j \leq 7, \quad (i, j) \notin S \quad \text{and} \quad (j, i) \notin S; \\ (2) \quad \tilde{p}_{s(i,j)k}^1 &= \begin{cases} z_j^1 & \text{if } k = i \\ z_i^1 & \text{if } k = j \\ 0 & \text{if } k \neq i, j, \end{cases} \quad \text{for all } (i, j) \in S. \end{aligned}$$

5.6. Next, we find the nonzero elements $\tilde{\pi}_s^3 \in K_3$ defined in (2.1.4) as follows. For the elements coming from Koszul relations in K , we choose $\tilde{\pi}_s^3 = 0$ for those elements. The only nonzero products in K_2 come from elements of type (1) and they are

$$z_2^1 \wedge z_4^1 = x^2 y^2 T_{12} = -z_4^1 \wedge z_2^1 \quad \text{and} \quad z_5^1 \wedge z_7^1 = x z^4 T_{13} = -z_7^1 \wedge z_5^1.$$

Therefore, we choose the following nontrivial liftings in K_3 :

$$\tilde{\pi}_{s(2,4)}^3 = x z^2 T_{123} = -\tilde{\pi}_{s(4,2)}^3 \quad \text{and} \quad \tilde{\pi}_{s(5,7)}^3 = -x^2 y z T_{123} = -\tilde{\pi}_{s(7,5)}^3.$$

5.7. We describe next the elements of A_3 .

Claim 1. All the products in $A_1 \cdot A_1 \cdot A_1$ are zero.

Proof of Claim 1. First, all the products in $A_1 \cdot A_1 \cdot A_1$ not involving $[z_1^1]$ have the coefficients of $[T_{ijk}]$ of degree six, and $R_6 = 0$. Thus, all such products are zero. Second, the only pairs (i, j) with $1 < i < j \leq 7$ such that $[z_i^1] \wedge [z_j^1]$ is nonzero in $A_1 \cdot A_1$ are in the set $\{(2, 3), (3, 5), (4, 5)\}$, but $[z_1^1 \wedge z_3^1] = [z_1^1 \wedge z_4^1] = 0$. Hence, all products involving $[z_1^1]$ at least once are zero. The claim now follows. \square

Therefore, $A_1 \cdot A_2 = A_1 \cdot \overline{A_2}$, it has rank $q_{12} = 10$, and its basis is given by the classes of

$$\begin{aligned} z_1^1 \wedge z_3^2 &= x z^2 w T_{124}, & z_1^1 \wedge z_4^2 &= y z^2 w T_{134}, & z_1^1 \wedge z_5^2 &= x^2 y w T_{124} - x z^2 w T_{134}, \\ z_1^1 \wedge z_6^2 &= x^2 z w T_{134}, & z_2^1 \wedge z_1^2 &= x^2 y w T_{124}, & z_2^1 \wedge z_2^2 &= x^2 y^2 T_{124}, \\ z_3^1 \wedge z_6^2 &= -x^2 y z w T_{123}, & z_5^1 \wedge z_1^2 &= y z^2 w T_{234}, & z_5^1 \wedge z_2^2 &= y^2 z^2 T_{234}, \\ z_5^1 \wedge z_3^2 &= -x z^4 T_{123} = z_4^1 \wedge z_6^2. \end{aligned}$$

A basis of $\overline{A_3}$ is $\{[z_t^3]\}_{t=1, \dots, 4}$, where

$$z_1^3 = y z^2 w T_{123}, \quad z_2^3 = y^2 z^2 T_{123}, \quad z_3^3 = y z^2 w T_{124}, \quad z_4^3 = z^4 T_{234}.$$

5.8. Using Proposition 2.6, we describe the elements \tilde{p}_{ui}^2 , for $1 \leq u \leq a_1 a_2 - a_1 q_{11} - q_{12} + b = 46 + b$ and $1 \leq i \leq 7$, as defined in (2.5.1).

Claim 2. Let \mathcal{A}, \mathcal{B} be as in Proposition 2.6 and let b be as in Summary 2.13. Then

$$\text{Span}_{\mathbf{k}} \mathcal{A} = A_1 \otimes (A_1 \cdot A_1) \quad \text{and} \quad b = 0.$$

Proof of Claim 2. It is clear that $\text{Span}_{\mathbf{k}} \mathcal{A} \subseteq A_1 \otimes (A_1 \cdot A_1)$. For $1 \leq i, j, h \leq 7$, any nonzero element $[z_i^1] \otimes [z_j^1] \wedge [z_h^1]$ in $A_1 \otimes (A_1 \cdot A_1)$ has $(j, h) \in S$ or $(h, j) \in S$. We show that each such element is in $\text{Span}_{\mathbf{k}} \mathcal{A}$.

In the case $\{(i, j), (i, h)\} \not\subseteq S$ we have

$$[z_i^1] \otimes [z_j^1] \wedge [z_h^1] = \sum_{k=1}^7 [z_k^1] \otimes [\tilde{p}_{s(i,j)k}^1] \wedge [z_h^1],$$

as in case (1) of 5.5.

By Koszul relation, the case $(i, h) \in S$ reduces to the case $(i, j) \in S$ in which we have

$$[z_i^1] \otimes [z_j^1] \wedge [z_h^1] = [z_i^1] \otimes [z_j^1] \wedge [z_h^1] + [z_j^1] \otimes [z_i^1] \wedge [z_h^1] = \sum_{k=1}^7 [z_k^1] \otimes [\tilde{p}_{s(i,j)k}^1] \wedge [z_h^1],$$

as in case (2) of 5.5. The first equality above follows from the proof of Claim 1. This implies $\text{Span}_{\mathbf{k}} \mathcal{A} = A_1 \otimes (A_1 \cdot A_1)$, so $\text{rank}_{\mathbf{k}}(\text{Span}_{\mathbf{k}} \mathcal{A}) = a_1 q_{11}$. By Proposition 2.6(b) we get $b = 0$. \square

Remark that Claim 2 implies Claim 1, since $\text{Span}_{\mathbf{k}} \mathcal{A} \subseteq \text{Ker } \phi_2$.

By definition of $\overline{A_2}$ as in Summary 2.13 and respectively Claim 2, the following equalities hold:

$$A_1 \otimes A_2 = (A_1 \otimes (A_1 \cdot A_1)) \oplus (A_1 \otimes \overline{A_2}) = (\text{Span}_{\mathbf{k}} \mathcal{A}) \oplus (A_1 \otimes \overline{A_2}).$$

By Proposition 2.6, $\text{Ker } \phi_2 = (\text{Span}_{\mathbf{k}} \mathcal{A}) \oplus B$, where $B = (\text{Ker } \phi_2) \cap (A_1 \otimes \overline{A_2})$. In order to find a basis for B , we record the indices of basis elements of $A_1 \cdot \overline{A_2}$ by the set

$$U = \{(1, 3), (1, 4), (1, 5), (1, 6), (2, 1), (2, 2), (3, 6), (5, 1), (5, 2), (5, 3), (4, 6)\}.$$

A basis of B is given by elements of two types:

- (1') $[z_i^1] \otimes [z_\ell^2]$ for all $1 \leq i \leq 7$ and $1 \leq \ell \leq 8$ such that $(i, \ell) \notin U$;
 (2') $[z_4^1] \otimes [z_6^2] - [z_5^1] \otimes [z_3^2]$.

Therefore, according to the types above, one defines $\tilde{p}_{u(i,\ell)}^2 = (\tilde{p}_{u(i,\ell)k}^2)_{k=1,\dots,7}$ as:

$$(1') \quad \tilde{p}_{u(i,\ell)k}^2 = \begin{cases} z_\ell^2 & \text{if } k = i \\ 0 & \text{if } k \neq i, \end{cases} \text{ for all } 1 \leq i \leq 7 \text{ and } 1 \leq \ell \leq 8 \text{ such that } (i, \ell) \notin U;$$

$$(2') \quad \tilde{p}_{u(4,6)k}^2 = \begin{cases} z_6^2 & \text{if } k = 4 \\ -z_3^2 & \text{if } k = 5 \\ 0 & \text{if } k \neq 4, 5. \end{cases}$$

5.9. It is easy to check that in K_3 we have the equalities:

- (1') $z_i^1 \wedge z_\ell^2 = 0$ for all $1 \leq i \leq 7$ and $1 \leq \ell \leq 8$ such that $(i, \ell) \notin U$;
 (2') $z_4^1 \wedge z_6^2 - z_5^1 \wedge z_3^2 = 0$.

Therefore, we may choose $\tilde{\pi}_u^4 = 0$ for all $1 \leq u \leq 46$.

5.10. A similar argument as in the proof of Claim 1 gives $A_1 \cdot A_3 = 0$, and hence $q_{13} = 0$. Moreover, $A_2 \cdot A_2$ has rank $q_{22} = 2$ and its basis is given by the classes of

$$z_1^2 \wedge z_6^2 = -x^2 y z w T_{1234} \quad \text{and} \quad z_2^2 \wedge z_6^2 = -x z^4 T_{1234}.$$

As $A_1 \cdot A_3 = 0$, any element in $\langle A_1, A_1, A_1 \rangle \subseteq A_4$ has a representative in K_4 given by

$$(5.10.1) \quad \sum_{s=1}^{42} \tilde{\pi}_s^3 \wedge p_s^1 \quad \text{such that} \quad \sum_{s=1}^{42} [\tilde{p}_{sk}^1 \wedge p_s^1] = 0, \text{ for all } 1 \leq k \leq 7,$$

where $p_s^1 \in \text{Ker } \partial_1^K$ is as in Proposition 2.11, and \tilde{p}_{si}^1 and $\tilde{\pi}_s^3$ are defined in (2.1.3) and (2.1.4) respectively. As described above, the only nontrivial lifting elements $\tilde{\pi}_s^3 \in K_3$ are $\tilde{\pi}_{s(2,4)}^3 = -\tilde{\pi}_{s(4,2)}^3$ and $\tilde{\pi}_{s(5,7)}^3 = -\tilde{\pi}_{s(7,5)}^3$. Since all of them contain T_{123} , only the component of p_s^1 that contains $z_1^1 = w T_4$ contributes to a nonzero Massey product. Thus, there exist $\alpha, \beta \in R$ such that

$$\begin{aligned} \sum_{s=1}^{42} [\tilde{\pi}_s^3 \wedge p_s^1] &= [\tilde{\pi}_{s(2,4)}^3 \wedge p_{s(2,4)}^1] + [\tilde{\pi}_{s(5,7)}^3 \wedge p_{s(5,7)}^1] \\ &= [\tilde{\pi}_{s(2,4)}^3 \wedge z_1^1] \wedge [\alpha] + [\tilde{\pi}_{s(5,7)}^3 \wedge z_1^1] \wedge [\beta] \\ &= [x z^2 w T_{1234}] \wedge [\alpha] + [x^2 y z w T_{1234}] \wedge [\beta] \\ &= \langle [z_2^1], [z_4^1], [z_1^1] \rangle \wedge [\alpha] - [z_1^2 \wedge z_6^2] \wedge [\beta]. \end{aligned}$$

The last equality follows from the following computation:

$$\begin{aligned} \langle [z_2^1], [z_4^1], [z_1^1] \rangle &= \{[(\partial_3^K)^{-1}(z_2^1 \wedge z_4^1) \wedge z_1^1 + z_2^1 \wedge (\partial_3^K)^{-1}(z_4^1 \wedge z_1^1)]\} \\ &= \{[\tilde{\pi}_{s(2,4)}^3 \wedge z_1^1 + z_2^1 \wedge \tilde{\pi}_{s(4,1)}^3]\} \\ &= \{[x z^2 w T_{1234}]\}. \end{aligned}$$

By abusing notation, we write $\langle [z_2^1], [z_4^1], [z_1^1] \rangle = [x z^2 w T_{1234}]$, which is not in $A_1 \cdot A_3 + A_2 \cdot A_2$, as showed in [2, Section 7]. It is clear now that the rank of the k -vector space

$$A_1 \cdot A_3 + A_2 \cdot A_2 + \text{Span}_k \langle A_1 \cdot A_1 \cdot A_1 \rangle$$

is $a = 3$ and its basis is given by

$$[z_1^2] \wedge [z_6^2], \quad [z_2^2] \wedge [z_6^2], \quad \text{and} \quad \langle [z_2^1], [z_4^1], [z_1^1] \rangle.$$

Thus, a basis of $\overline{A_4}$ is $\{[z_r^4]\}_{r=1,2}$ where

$$z_1^4 = y z^2 w T_{1234} \quad \text{and} \quad z_2^4 = y^2 z^2 T_{1234}.$$

5.11. We conclude that for the ring R discussed in this section, and by Proposition 4.2 we have:

$$\begin{array}{llll} a_1 = 7, & a_2 = 15, & a_3 = 14, & a_4 = 5, \\ q_{11} = 7, & q_{12} = 10, & q_{13} = 0, & q_{22} = 2, \\ a = 3, & b = 0, & & \\ \mathcal{P}_3 = 7, & \mathcal{P}_4 = 45, & \mathcal{P}_5 = 221. & \end{array}$$

ACKNOWLEDGEMENT

The authors thank Hailong Dao for inspiring them to work on this project and for very fruitful discussions. They also thank Frank Moore for helping them with the `DGAlgebras` package [15] to check the computations, and thank the referee for helpful suggestions. The first author was supported by the Naval Academy Research Council in Summer 2020.

REFERENCES

- [1] D. J. Anick, *A counterexample to a conjecture of Serre*, Ann. of Math. (2) **115** (1982), no. 1, 1–33.
- [2] L. L. Avramov, *On the Hopf algebra of a local ring*, Izv. Akad. Nauk SSSR Ser. Mat. **38** (1974), 253–277.
- [3] L. L. Avramov, *Homological asymptotics of modules over local rings*, Commutative algebra (Berkeley, CA, 1987), Math. Sci. Res. Inst. Publ., Springer, New York, vol. 15, (1989), 33–62.
- [4] L. L. Avramov, *Infinite free resolutions*, Six Lectures on Commutative Algebra (Bellaterra, 1996), Progr. Math. **166** (1998), Birkhäuser, Basel, 1–118.
- [5] L. L. Avramov, *A cohomological study of local rings of embedding codepth 3*, J. Pure Appl. Algebra **216** (2012), no. 11, 2489–2506.
- [6] J. Burke, *Higher homotopies and Golod rings*, preprint, arXiv:1508.03782.
- [7] L. W. Christensen; O. Veliche, *Acyclicity over local rings with radical cube zero*, Illinois J. Math. **51** (2007), no. 4, 1439–1454.
- [8] E. S. Golod, *On the homology of some local rings*, Dokl. Akad. Nauk SSSR **144** (1962), 479–482 (in Russian); English translation: Soviet Math. Dokl. **3** (1962), 745–748.
- [9] T. H. Gulliksen; G. Levin, *Homology of local rings*, Queen’s Papers in Pure Appl. Math. **20** (1969), x+192 pp.
- [10] L. Katthän, *A non-Golod ring with a trivial product on its Koszul homology*, J. Algebra, **479** (2017), 244–262.
- [11] A. R. Kustin; M. Miller, *Classification of the Tor-Algebras of Codimension Four Gorenstein Local Rings*, Math. Z. **190** (1985), 341–355.
- [12] D.-M. Lu, J.H. Palmieri, Q.-S. Wu, and J.J. Zhang, *A-infinity structure on Ext-algebras*, J. Pure Appl. Alg. **213** (2009), 2017–2037.
- [13] D. R. Grayson; M. Stillman, *Macaulay2, a software system for research in algebraic geometry*, Available at <http://www.math.uiuc.edu/Macaulay2/>.
- [14] J. P. May, *Matric Massey products*, J. Algebra, **12**(4) (1969), 533–568.
- [15] F. Moore, *DGAlgebras: Data type for DG algebras. Version 1.0.1*, A Macaulay2 package available at <https://github.com/Macaulay2/M2/tree/master/M2/Macaulay2/packages>.
- [16] J.-E. Roos, *On some unexpected rings that are close to Golod rings*, (2016), unpublished notes.
- [17] Y. Yoshino, *Modules of G-dimension zero over local rings with the cube of maximal ideal being zero*, Commutative algebra, singularities and computer algebra (Sinaia, 2002), 255–273, NATO Sci. Ser. II Math. Phys. Chem., **115**, Kluwer Acad. Publ., Dordrecht, 2003.

DEPARTMENT OF MATHEMATICS, UNITED STATES NAVAL ACADEMY, ANNAPOLIS, MD 21402, U.S.A.

Email address: vnnguyen@usna.edu

URL: <https://sites.google.com/view/vcnguyen>

DEPARTMENT OF MATHEMATICS, NORTHEASTERN UNIVERSITY, BOSTON, MA 02115, U.S.A.

Email address: o.veliche@northeastern.edu

URL: <https://web.northeastern.edu/oveliche>