

Satisfiability problems on sums of Kripke frames

Ilya B. Shapirovsky

New Mexico State University, USA
Institute for Information Transmission Problems of Russian Academy of Sciences

July 6, 2022

Abstract

We consider the operation of sum on Kripke frames, where a family of frames-summands is indexed by elements of another frame. In many cases, the modal logic of sums inherits the finite model property and decidability from the modal logic of summands [BR10], [Sha18]. In this paper we show that, under a general condition, the satisfiability problem on sums is polynomial space Turing reducible to the satisfiability problem on summands. In particular, for many modal logics decidability in PSpace is an immediate corollary from the semantic characterization of the logic.

Keywords: sum of Kripke frames, finite model property, decidability, Turing reduction, PSpace, Japaridze’s polymodal logic, lexicographic product of modal logics, lexicographic sum of modal logics

1 Introduction

In classical model theory, there is a number of results (“composition theorems”) that reduce the theory (first-order, MSO) of a compound structure (e.g., sum or product) to the theories of its components, see, e.g., [Mos52, FV59, She75, Gur79] or [Gur85]. In this paper we use the composition method in the context of modal logic.

Given a family $(F_i \mid i \text{ in } I)$ of frames (structures with binary relations) indexed by elements of another frame I , the *sum of the frames F_i ’s over I* is obtained from their disjoint union by connecting elements of i -th and j -th distinct components according to the relations in I . Given a class \mathcal{F} of frames-summands and a class \mathcal{I} of frames-indices, $\sum_{\mathcal{I}} \mathcal{F}$ denotes the class of all sums of F_i ’s in \mathcal{F} over I in \mathcal{I} .

In many cases, this operation preserves the finite model property and decidability of the logic of summands [BR10], [Sha18]. In this paper we show that transferring results also hold for the complexity of the modal satisfiability problems on sums. In particular, it follows that for many logics PSpace-completeness is an immediate corollary of semantic characterization.

It is a classical result by R. Ladner that the decision problem for the logic of (finite) preorders S4 is in PSpace [Lad77]. In [Sha08], it was shown that the polymodal provability logic GLP is also decidable in PSpace. In spite of the significant difference between these two logics, there is a uniform proof for both these two particular systems, as well as for many other important modal logics: the general phenomenon is that the modal satisfiability problem on sums over Noetherian (in particular, finite) orders is polynomial space Turing reducible to the modal satisfiability problem on summands. In the case of S4 these summands are frames of the form $(W, W \times W)$ with the satisfiability problem being in NP, so in PSpace. And hence, S4 is in PSpace. In the case of GLP, the class of summands is even simpler: the only summand required

is an irreflexive singleton [Bek10]. (We will discuss these and other examples in Sections 4.5, 5.3, 5.4.)

The paper has the following structure. Section 2 contains preliminary material. Section 3 is about truth-preserving operations on sums of frames; it contains necessary tools for the complexity results, and also quotes recent results on the finite model property. This section is based on [Sha18]. The central Section 4 is about complexity. The reduction between sums and summands is described in Theorem 11 and in its more technical (but more tunable) version Theorem 17; these are the main results of the paper. This section elaborates earlier works [Sha08, Sha05] (in particular, our new results significantly generalize and simplify Theorems 22 and 35 in [Sha08]). In Section 5 we discuss modifications of the sum operation: (iterated) lexicographic sums of frames, which are important in the context of provability logics [Bek10]; lexicographic products of frames, earlier studied in [Bal09, Bal10, BM13, BFD16]; the operation of refinement of modal logics introduced in [BR10]. Further results and directions are discussed in Section 6.

2 Preparatory syntactical and semantical definitions

Let $A \leq \omega$. The set $ML(A)$ of *modal formulas over A* (or *A-formulas*, for short) is built from a countable set of *variables* $PV = \{p_0, p_1, \dots\}$ using Boolean connectives \perp, \rightarrow and unary connectives $\Diamond_a, a < A$ (*modalities*). The connectives $\vee, \wedge, \neg, \top, \Box_a$ are defined as abbreviations in the standard way, in particular $\Box_a \varphi$ is $\neg \Diamond_a \neg \varphi$.

An (A) -frame is a structure $F = (W, (R_a)_{a < A})$, where $W \neq \emptyset$ and $R_a \subseteq W \times W$ for $a < A$. A (*Kripke*) *model on F* is a pair $M = (F, \theta)$, where $\theta : PV \rightarrow 2^W$. We write $\text{dom}(F)$ for W , which is called the *domain* of F . We write $u \in F$ for $u \in \text{dom}(F)$. For $u \in W, V \subseteq W$, we put $R_a(u) = \{v \mid uR_av\}$, $R_a[V] = \cup_{v \in V} R_a(v)$.

The *truth relation* in a model is defined in the usual way, in particular $M, w \models \Diamond_a \varphi$ means that $M, v \models \varphi$ for some v in $R_a(w)$. A formula φ is *satisfiable in a model M* if $M, w \models \varphi$ for some w in M . A formula is *satisfiable in a frame F* (in a class \mathcal{F} of frames) if it is satisfiable in some model on F (in some model on a frame in \mathcal{F}). φ is *valid in a frame F* (in a class \mathcal{F} of frames) if $\neg \varphi$ is not satisfiable in F (in \mathcal{F}). Validity of a set of formulas means validity of every formula in this set.

The notions of p-morphism, generated subframe and submodel are defined in the standard way, see e.g. [GKWZ03, Section 1.4]. The notation $F \twoheadrightarrow G$ means that G is a p-morphic image of F ; $F \cong G$ means that F and G are isomorphic.

A (*propositional normal modal*) *logic* is a set L of formulas that contains all classical tautologies, the axioms $\neg \Diamond_a \perp$ and $\Diamond_a(p_0 \vee p_1) \rightarrow \Diamond_a p_0 \vee \Diamond_a p_1$ for each a in A , and is closed under the rules of modus ponens, substitution and monotonicity (if $\varphi \rightarrow \psi \in L$, then $\Diamond_a \varphi \rightarrow \Diamond_a \psi \in L$, for each a in A).

The set $\text{Log } \mathcal{F}$ of all formulas that are valid in a class \mathcal{F} of A -frames is a logic (see, e.g., [CZ97]); it is called the *logic of \mathcal{F}* ; such logics are called *Kripke complete*. A logic has the *finite model property* if it is the logic of a class of finite frames (a frame is finite, if its domain is). $\text{Fr } L$ denotes the class of all frames validating L .

Remark that for a Kripke complete logic, its decision problem is the validity problem on the class of all its frames (or on any other class \mathcal{F} such that $L = \text{Log}(\mathcal{F})$). The dual problem is the satisfiability: $\text{Sat}(\mathcal{F})$ is the set of all formulas satisfiable in \mathcal{F} (in the signature of \mathcal{F}). Remark that for the class PSpace (as well as for any other deterministic complexity class), $\text{Sat } \mathcal{F} \in \text{PSpace}$ iff $\text{Log } \mathcal{F} \in \text{PSpace}$.

Natural numbers are considered as finite ordinals. Given a sequence $\mathbf{v} = (v_0, v_1, \dots)$, we write $\mathbf{v}(i)$ for v_i .

3 Sums

We fix $A \leq \omega$ for the alphabet and consider the language $\text{ML}(A)$.

Consider a non-empty family $(F_i)_{i \in I}$ of A-frames $F_i = (W_i, (R_{i,a})_{a \in A})$. The *disjoint union* of these frames is the A-frame $\bigsqcup_{i \in I} F_i = (\bigsqcup_{i \in I} W_i, (R_a)_{a \in A})$, where $\bigsqcup_{i \in I} W_i = \bigcup_{i \in I} (\{i\} \times W_i)$ is the disjoint union of sets W_i , and

$$(i, w)R_a(j, v) \quad \text{iff} \quad i = j \ \& \ wR_{i,a}v.$$

Suppose that I is the domain of another A-frame $\mathbf{l} = (I, (S_a)_{a \in A})$.

Definition 1. The *sum of the family* $(F_i)_{i \in I}$ of A-frames over the A-frame $\mathbf{l} = (I, (S_a)_{a \in A})$ is the A-frame $\sum_{i \in \mathbf{l}} F_i = (\bigsqcup_{i \in I} W_i, (R_a^\Sigma)_{a \in A})$, where

$$(i, w)R_a^\Sigma(j, v) \quad \text{iff} \quad (i = j \ \& \ wR_{i,a}v) \text{ or } (i \neq j \ \& \ iS_a j).$$

The *sum of models* $\sum_{i \in \mathbf{l}} (F_i, \theta_i)$ is the model $(\sum_{i \in \mathbf{l}} F_i, \theta)$, where $(i, w) \in \theta(p)$ iff $w \in \theta_i(p)$.

For classes \mathcal{I}, \mathcal{F} of A-frames, let $\sum_{\mathcal{I}} \mathcal{F}$ be the class of all sums $\sum_{i \in \mathbf{l}} F_i$ such that $\mathbf{l} \in \mathcal{I}$ and $F_i \in \mathcal{F}$ for every i in \mathbf{l} .

Remark 1. We do not require that S_a 's are partial orders or even transitive relations. Also, we note that the relations R_a^Σ are independent of reflexivity of the relations S_a : if $\mathbf{l}' = (I, (S'_a)_{a \in A})$, where S'_a is the reflexive closure of S_a for each $a \in A$, then $\sum_{i \in \mathbf{l}} F_i = \sum_{i \in \mathbf{l}'} F_i$.

Example 1 (Skeleton and clusters). A *cluster* is a frame of the form $(W, W \times W)$.

Let $\mathbf{F} = (W, R)$ be a preorder. The *skeleton* of \mathbf{F} is the partial order $\text{sk}\mathbf{F} = (\overline{W}, \leq_R)$, where \overline{W} is the quotient set of W by the equivalence $R \cap R^{-1}$, and for $C, D \in \overline{W}$, $C \leq_R D$ iff $\exists w \in C \exists v \in D \ wRv$. The restriction of \mathbf{F} on an element of \overline{W} is called a *cluster in* \mathbf{F} .

It is easy to see that every preorder \mathbf{F} is isomorphic to the sum $\sum_{C \in \text{sk}\mathbf{F}} (C, C \times C)$ of its clusters over its skeleton.

Example 2. Suppose that $\mathbf{F} = (W, R)$ satisfies the property of *weak transitivity* $xRzRy \Rightarrow xRy \vee x = y$. Let \mathbf{l} be the skeleton of the preorder (W, R^*) , where R^* is the transitive reflexive closure of R . Then \mathbf{F} is isomorphic to a sum $\sum_{i \in \mathbf{l}} F_i$ such that every $F_i = (W_i, R_i)$ satisfies the property $x \neq y \Rightarrow xR_i y$.

We shall be mainly interested in the polymodal case where the indexing frame has only one non-empty relation.

Definition 2. Consider a unimodal frame $\mathbf{l} = (I, S)$ and a family $(F_i)_{i \in I}$ of A-frames (or A-models). For $a \in A$, the *a-sum* $\sum_{\mathbf{l}}^a F_i$ is the sum $\sum_{\mathbf{l}'} F_i$, where \mathbf{l}' is the A-frame whose domain is I , the a -th relation is S and other relations are empty. If \mathcal{F} is a class of A-frames, \mathcal{I} is a class of 1-frames, then $\sum_{\mathcal{I}}^a \mathcal{F}$ is the class of all sums $\sum_{\mathbf{l}}^a F_i$, where $\mathbf{l} \in \mathcal{I}$ and all F_i are in \mathcal{F} .

3.1 Basic truth-preserving operations

The theorem below is a collection of facts illustrating how sums interact with p-morphisms, generated subframes, and disjoint unions. Their proofs are straightforward from definitions, see [Sha18, Section 3] for the details.

Theorem 2 ([Sha18]).

1. Let I be an A-frame and $(F_i)_{i \in I}$ a family of A-frames. If J is a generated subframe of I , then $\sum_{i \in J} F_i$ is a generated subframe of $\sum_{i \in I} F_i$.
2. Let I, J be A-frames, $(F_i)_{i \in I}, (G_j)_{j \in J}$ families of A-frames. Suppose that all the relations in J are irreflexive.
 - (a) If $f : I \rightarrow J$ and $F_i \rightarrow G_{f(i)}$ for all i in I , then $\sum_{i \in I} F_i \rightarrow \sum_{j \in J} G_j$.
 - (b) If $I = J$ and $F_i \rightarrow G_i$ for all i in I , then $\sum_{i \in I} F_i \rightarrow \sum_{i \in I} G_i$.
 - (c) If $f : I \rightarrow J$, then $\sum_{i \in I} G_{f(i)} \rightarrow \sum_{j \in J} G_j$.
3. (a) Let I be an A-frame, $(J_i)_{i \in I}$ a family of A-frames, and $(F_{ij})_{i \in I, j \in J_i}$ a family of A-frames. Then

$$\sum_{i \in I} \sum_{j \in J_i} F_{ij} \cong \sum_{(i,j) \in \sum_{k \in I} J_k} F_{ij}.$$

- (b) Let I be a non-empty set, $(J_i)_{i \in I}$ a family of A-frames, and $(F_{ij})_{i \in I, j \in J_i}$ a family of A-frames. Then

$$\bigsqcup_{i \in I} \sum_{j \in J_i} F_{ij} \cong \sum_{(i,j) \in \bigsqcup_{k \in I} J_k} F_{ij}.$$

- (c) Let I be an A-frame, $(J_i)_{i \in I}$ a family of non-empty sets, and $(F_{ij})_{i \in I, j \in J_i}$ a family of A-frames. Then

$$\sum_{i \in I} \bigsqcup_{j \in J_i} F_{ij} \cong \sum_{(i,j) \in \sum_{k \in I} (J_k, (\emptyset)_A)} F_{ij},$$

where $(\emptyset)_A$ denotes the sequence of length A in which every element is the empty set.

3.2 Sums and universal modality

For an A-frame $F = (W, (R_0, R_1, \dots))$, let $F^{(\forall)}$ be the $(1 + A)$ -frame $(W, (W \times W, R_0, R_1, \dots))$. For a class \mathcal{F} of A-frames, $\mathcal{F}^{(\forall)} = \{F^{(\forall)} \mid F \in \mathcal{F}\}$.

In [Sha18], it was shown that if the classes $\mathcal{F}^{(\forall)}$ and $\mathcal{G}^{(\forall)}$ have the same logic, then for any class \mathcal{I} of A-frames, the logics of sums $\sum_{\mathcal{I}} \mathcal{F}$ and $\sum_{\mathcal{I}} \mathcal{G}$ are equal; moreover, the logics of the classes $\sum_{\mathcal{I}} \mathcal{F}^{(\forall)}$ and $\sum_{\mathcal{I}} \mathcal{G}^{(\forall)}$ are equal, thus we have $\text{Log } \sum_{\mathcal{J}} (\sum_{\mathcal{I}} \mathcal{F}) = \text{Log } \sum_{\mathcal{J}} (\sum_{\mathcal{I}} \mathcal{G})$ for any other class of frames-indices \mathcal{J} , and so on.

Theorem 3 ([Sha18], Theorem 4.11). *Let $\mathcal{I}, \mathcal{F}, \mathcal{G}$ be classes of A-frames. If $\text{Log } \mathcal{F}^{(\forall)} = \text{Log } \mathcal{G}^{(\forall)}$, then $\text{Log } \sum_{\mathcal{I}} \mathcal{F}^{(\forall)} = \text{Log } \sum_{\mathcal{I}} \mathcal{G}^{(\forall)}$, and hence $\text{Log } \sum_{\mathcal{I}} \mathcal{F} = \text{Log } \sum_{\mathcal{I}} \mathcal{G}$.*

In particular, it follows that if the logic of the class $\mathcal{F}^{(\forall)}$ has the finite model property, then the logic of the class of sums $\sum_{\mathcal{I}} \mathcal{F}$ is equal to the logic of the class of sums $\sum_{\mathcal{I}} \mathcal{G}$, where \mathcal{G} is a class of finite frames.

3.3 Decomposition of sums

To reduce satisfiability in sums to the satisfiability in summands, we will use an auxiliary notion: *satisfiability under conditions*.

Definition 3. A sequence $\Gamma = (\Gamma_a)_{a \in A}$, where Γ_a are sets of A-formulas, is called a *condition* (in the language $\text{ML}(A)$).

Consider a model $M = (W, (R_a)_{a \in A}, \theta)$, w in M . By induction on the length of an A-formula φ , we define the relation $M, w \models_{\Gamma} \varphi$ (“under the condition Γ , φ is true at w in M ”):

as usual, $M, w \not\models_{\Gamma} \perp$, $M, w \models_{\Gamma} p$ iff $M, w \models p$ for a variable p , $M, w \models_{\Gamma} \varphi \rightarrow \psi$ iff $M, w \not\models_{\Gamma} \varphi$ or $M, w \models_{\Gamma} \psi$; for $a \in A$,

$$M, w \models_{\Gamma} \Diamond_a \varphi \quad \text{iff} \quad \varphi \in \Gamma_a \text{ or } \exists v \in R_a(w) \ M, v \models_{\Gamma} \varphi.$$

In particular, if all Γ_a are empty, then we have the standard notion of truth in a Kripke model:

$$M, w \models_{(\emptyset)_A} \varphi \quad \text{iff} \quad M, w \models \varphi,$$

where $(\emptyset)_A$ denotes the condition consisting of empty sets. The truth under conditions is respected by the standard operations on Kripke models:

Proposition 4. *Let Γ be a condition and φ a formula.*

1. *If M' is a generated submodel of M , then $M', w \models_{\Gamma} \varphi$ iff $M, w \models_{\Gamma} \varphi$ for every w in M' .*
2. *If $M = \bigsqcup_{i \in I} M_i$, then $M, (i, w) \models_{\Gamma} \varphi$ iff $M_i, w \models_{\Gamma} \varphi$ for every i in I and every w in M_i .*
3. *If $f : M \twoheadrightarrow M'$, then $M, w \models_{\Gamma} \varphi$ iff $M', f(w) \models_{\Gamma} \varphi$ for every w in M .*

Proof. This proof completely reflects the proof of these facts for the standard truth relation in Kripke models and can be obtained by a straightforward induction on the length of φ . \square

Let $\text{sub}(\varphi)$ be the set of all subformulas of φ . We put

$$\varphi[M, \Gamma] = \{\psi \in \text{sub}(\varphi) \mid M, v \models_{\Gamma} \psi \text{ for some } v\}.$$

In particular, $\varphi[M, (\emptyset)_A]$ is the set of all subformulas of φ that are satisfiable in M .

A triple (φ, Φ, Γ) , where $\Phi \subseteq \text{sub}(\varphi)$, is called a *tie*. A tie (φ, Φ, Γ) is *satisfiable* in a frame F (in a class \mathcal{F} of frames) if there exists a model M on F (on a frame in \mathcal{F}) such that $\Phi = \varphi[M, \Gamma]$. Hence, we have:

Proposition 5. *A formula φ is satisfiable in a class \mathcal{F} of frames iff there exists $\Phi \subseteq \text{sub}(\varphi)$ such that $\varphi \in \Phi$ and the tie $(\varphi, \Phi, (\emptyset)_A)$ is satisfiable in \mathcal{F} .*

Classes of A-frames \mathcal{F} and \mathcal{G} are said to be *interchangeable*, in symbols $\mathcal{F} \equiv \mathcal{G}$, if the same ties are satisfiable in classes \mathcal{F} and \mathcal{G} . From the above proposition, it follows that if $\mathcal{F} \equiv \mathcal{G}$, then the logics of these classes are equal. Moreover, the following holds:

Proposition 6 ([Sha18], Proposition 4.10). *Let \mathcal{F} and \mathcal{G} be classes of A-frames. The following are equivalent:*

- *A tie is satisfiable in \mathcal{F} iff it is satisfiable in \mathcal{G} .*
- *A formula is satisfiable in $\mathcal{F}^{(\forall)}$ iff it is satisfiable in $\mathcal{G}^{(\forall)}$.*

The latter condition means that the logics of $\mathcal{F}^{(\forall)}$ and $\mathcal{G}^{(\forall)}$ are equal. The term ‘interchangeable’ is motivated by Theorem 3.

Hence, the language of ties is as expressible as the modal language with the universal modality. In Section 4.2, we will present explicit reductions between these languages.

Definition 4. Let V be a set of elements of a model $M = (W, (R_a)_{a \in A}, \theta)$. Given a formula φ and a condition Γ , let Δ be the condition defined as follows: for $a \in A$,

$$\Delta(a) = \Gamma(a) \cup \{\chi \in \text{sub}(\varphi) \mid \exists w \in R_a[V] \setminus V \ M, w \models_{\Gamma} \chi\}.$$

Δ is called the *external condition of V in M with respect to φ and Γ* .

Remark 7. If $\Gamma \subseteq \text{sub}(\varphi)^A$, then $\Delta \subseteq \text{sub}(\varphi)^A$ as well (this will be important in the next section, where we consider the *conditional satisfiability problem*).

Lemma 8 ([Sha18], Lemma 4.5). *Consider a sum of models $M = \sum_I M_i$, i in I , and the set $V = \{i\} \times \text{dom}(M_i)$. If Δ is the external condition of V in M with respect to some given φ , Γ , then for all v in M_i , χ in $\text{sub}(\varphi)$,*

$$M, (i, v) \models_{\Gamma} \chi \quad \text{iff} \quad M_i, v \models_{\Delta} \chi. \quad (1)$$

Lemma 9 below is a particular corollary of Lemma 8. It will be important for the proofs of our complexity results.

Consider $a \in A$ and models M_0, M_1 . The model $M_0 +^a M_1$ is obtained from the disjoint union of M_0 and M_1 by adding all the pairs of form $((0, w), (1, v))$ to the a -th relation; that is, in our general notation, $M_0 +^a M_1 = \sum_{(2, <)}^a M_i$.

For an A -condition $\Gamma = (\Gamma_0, \dots, \Gamma_{A-1})$ and a set of A -formulas Ψ , we put $\Gamma \cup^a \Psi = (\Gamma'_0, \dots, \Gamma'_{A-1})$, where $\Gamma'_a = \Gamma_a \cup \Psi$, and $\Gamma'_b = \Gamma_b$ for $b \neq a$.

Lemma 9. *For A -models M_0, M_1 , an A -formula φ , an A -condition Γ , and $a \in A$, we have*

$$\varphi[M_0 +^a M_1, \Gamma] = \varphi[M_0, \Gamma \cup^a \varphi[M_1, \Gamma]] \cup \varphi[M_1, \Gamma]. \quad (2)$$

Proof. Let V be the bottom part of the sum $M_0 +^a M_1$: $V = \{(0, w) \mid w \text{ is in } M_0\}$. Then

$$\Delta = \Gamma \cup^a \varphi[M_1, \Gamma]$$

is the external condition of V w.r.t. φ and Γ . The external condition of the top part $\{(1, v) \mid v \text{ is in } M_1\}$ w.r.t. φ and Γ is just Γ . By Lemma 8, we have for every w in M_0 , every v in M_1 , and every $\chi \in \text{sub}(\varphi)$:

$$\begin{aligned} M_0 +^a M_1, (0, w) \models_{\Gamma} \chi & \quad \text{iff} \quad M_0, w \models_{\Delta} \chi \\ M_0 +^a M_1, (1, v) \models_{\Gamma} \chi & \quad \text{iff} \quad M_1, v \models_{\Gamma} \chi. \end{aligned}$$

Now (2) follows. □

3.4 Sums over Noetherian orders

We say that I is a *Noetherian* (or *converse well-founded*) *order* if I is a strict partial order which has no infinite ascending chains. Let NPO be the class of all non-empty Noetherian orders (we say that a partial order is non-empty, if its domain is).

A strict partial order $(I, <)$ is called a (*transitive irreflexive*) *tree* if it has a least element (the *root*) and for all $i \in I$ the set $\{j \mid j < i\}$ is a finite chain. Let Tr_f be the class of all finite trees.

Consider a finite tree $I = (I, <)$. The *branching of i in I* , denoted by $\underline{(i, I)}$, is the number of immediate successors of i (j is an immediate successor of i , if $i < j$ and there is no k such that $i < k < j$); the *branching of I* , denoted by $\underline{(I)}$, is $\max \{\underline{(i, I)} \mid i \text{ in } I\}$. The *height of I* , denoted by $ht(I)$, is $\max \{|V| \mid V \text{ is a chain in } I\}$. For $h, b \in \omega$, let $\text{Tr}(h, b)$ be the class of all finite trees with height $\leq h$ and branching $\leq b$:

$$\text{Tr}(h, b) = \{I \in \text{Tr}_f \mid ht(I) \leq h \ \& \ \underline{(I)} \leq b\}.$$

Let $\bigsqcup \mathcal{F}$ be the class of frames of form $\bigsqcup_I F_i$, where I is a non-empty set, $F_i \in \mathcal{F}$ for all $i \in I$, and let $\bigsqcup_{\leq k} \mathcal{F}$ be the class of such frames with $0 < |I| \leq k$. Likewise for $\bigsqcup_{< k} \mathcal{F}$.

Let $\#\varphi$ denote the number of subformulas of a formula φ .

Theorem 10 ([Sha18]). *Let \mathcal{F} be a class of \mathbf{A} -frames, $a \in \mathbf{A}$, and \mathcal{I} a class of Noetherian orders containing all finite trees.*

1. *We have*

$$\text{Log } \sum_{\text{NPO}}^a \mathcal{F} = \text{Log } \sum_{\mathcal{I}}^a \mathcal{F} = \text{Log } \sum_{\text{Tr}_f}^a \mathcal{F}.$$

Moreover, for every \mathbf{A} -formula φ we have:

$$\varphi \text{ is satisfiable in } \sum_{\text{NPO}}^a \mathcal{F} \text{ iff } \varphi \text{ is satisfiable in } \sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}.$$

2. *Assume that \mathcal{I} is closed under finite disjoint unions. Then*

$$\sum_{\text{NPO}}^a \mathcal{F} \equiv \sum_{\mathcal{I}}^a \mathcal{F} \equiv \bigsqcup_{< \omega} \sum_{\text{Tr}_f}^a \mathcal{F}.$$

Moreover, for every \mathbf{A} -tie $\tau = (\varphi, \Phi, \mathbf{\Gamma})$ we have:

$$\tau \text{ is satisfiable in } \sum_{\text{NPO}}^a \mathcal{F} \text{ iff } \tau \text{ is satisfiable in } \bigsqcup_{\leq \#\varphi} \sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}. \quad (3)$$

In view of Proposition 5, the first statement of the theorem is a corollary of the second statement. For the key equivalence (3), see Theorem 5.2(i) in [Sha18].

Theorem 10 will be the crucial semantic tool for the complexity results.

4 Complexity

The main goal of this section is to show that the modal satisfiability problem on sums over Noetherian orders is polynomial space Turing reducible to the modal satisfiability problem on summands.

For problems A and B , we put $A \leq_{\text{T}}^{\text{PSpace}} B$ if there exists a polynomial space bounded oracle deterministic machine M with oracle B that decides A [SG77] (it is assumed that every tape of M , including the oracle tape, is polynomial space bounded).

Theorem 11. *Let $a < \mathbf{A} < \omega$, \mathcal{F} a class of \mathbf{A} -frames, and \mathcal{I} a class of Noetherian orders containing all finite trees. Then:*

$$1. \text{Sat } \sum_{\mathcal{I}}^a \mathcal{F} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}.$$

$$2. \text{ If also } \mathcal{I} \text{ is closed under finite disjoint unions, then } \text{Sat}(\sum_{\mathcal{I}}^a \mathcal{F})^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}.$$

This theorem will be proven in Section 4.3. For technical reasons, first we will address complexity of the *conditional satisfiability problem*. Let \mathbf{A} be finite, and let \mathcal{F} be a class of \mathbf{A} -frames. We shall be interested in whether a given tie $(\varphi, \Phi, \mathbf{\Gamma})$ is satisfiable in \mathcal{F} . The following simple observation shows that, w.l.o.g., we may assume that every $\mathbf{\Gamma}(a)$, $a < \mathbf{A}$, consists of subformulas of φ , and hence that $\mathbf{\Gamma}$ is a finite sequence of finite sets:

Proposition 12. *A tie $(\varphi, \Phi, \mathbf{\Gamma})$ is satisfiable in a class \mathcal{F} iff $(\varphi, \Phi, (\text{sub}(\varphi) \cap \mathbf{\Gamma}(a))_{a \in \mathbf{A}})$ is.*

Proof. It is immediate from Definition 3 that for any conditions $\mathbf{\Gamma}, \mathbf{\Delta}$ such that

$$\mathbf{\Gamma}(a) \cap \text{sub}(\varphi) = \mathbf{\Delta}(a) \cap \text{sub}(\varphi) \text{ for all } a \in \mathbf{A},$$

we have

$$M, w \models_{\Gamma} \chi \text{ iff } M, w \models_{\Delta} \chi$$

for every model M , every w in M , and every $\chi \in \text{sub}(\varphi)$.

The statement of the proposition is a particular case of this observation, where $\Delta(a) = \text{sub}(\varphi) \cap \Gamma(a)$ for all $a \in A$. \square

The *conditional satisfiability problem on \mathcal{F}* is to decide whether a given tie (φ, Φ, Γ) such that $\Gamma \subseteq \text{sub}(\varphi)^A$ is satisfiable in \mathcal{F} . In symbols, $\text{CSat } \mathcal{F}$ is the set

$$\{(\varphi, \Phi, \Gamma) \mid \varphi \in \text{ML}(A) \ \& \ \Phi \subseteq \text{sub}(\varphi) \ \& \ \Gamma \subseteq \text{sub}(\varphi)^A \ \& \ \text{the tie } (\varphi, \Phi, \Gamma) \text{ is satisfiable in } \mathcal{F}\}.$$

In Section 4.1 we will describe a decision procedure for the conditional satisfiability problem on sums over Noetherian orders $\sum_{\text{NPO}}^a \mathcal{F}$ with the oracle $\text{CSat } \mathcal{F}$. Next, in Section 4.2, we will describe reductions between $\text{CSat } \mathcal{F}$ and $\text{Sat } \mathcal{F}^{(\vee)}$, which will complete the proof of Theorem 11.

4.1 Arithmetic of conditional satisfiability

It will be convenient to encode subformulas of a given φ as Boolean vectors of length $\#\varphi$, considered as characteristic functions on $\text{sub}(\varphi)$. For $A \leq \omega$, the set of modal formulas is linearly ordered by a polynomial time computable relation \sqsupseteq such that if ψ is a subformula of φ , then $\varphi \sqsupseteq \psi$ (e.g., put $\varphi \sqsupseteq \psi$ if ψ is shorter than φ , and assume that formulas of the same length are ordered lexicographically). Let $(\psi_0, \dots, \psi_{\#\varphi-1})$ be the \sqsupseteq -chain of all subformulas of φ (hence, $\psi_0 = \varphi$); for $\mathbf{v} \in 2^{\#\varphi}$, we write $\varphi[\mathbf{v}]$ for $\{\psi_i \mid \mathbf{v}(i) = 1\}$; similarly, a sequence $\mathbf{U} = (\mathbf{u}_a)_{a \in A}$ of such vectors represents the condition $\Gamma = (\varphi[\mathbf{u}_a])_{a \in A}$. Hence, for a finite A , every tie $\tau = (\varphi, \Phi, \Gamma)$ with $\Gamma \subseteq \text{sub}(\varphi)^A$ is represented by a triple $\tau' = (\varphi, \mathbf{v}, (\mathbf{u}_a)_{a \in A})$, where $\mathbf{v}, \mathbf{u}_0, \dots, \mathbf{u}_{A-1} \in 2^{\#\varphi}$; this triple is also called a *tie*. In this case, by the satisfiability of τ' we mean the satisfiability of τ .

Let $\mathbf{0}_\varphi$ denote the sequence of length A of zero vectors of length $\#\varphi$ (that is, $\mathbf{0}_\varphi$ represents the condition, consisting of empty sets). In view of Proposition 12, we have the following reformulation of Proposition 5:

Proposition 13. *φ is satisfiable in \mathcal{F} iff there exists $\mathbf{v} \in 2^{\#\varphi}$ such that $\mathbf{v}(0) = 1$ and the tie $(\varphi, \mathbf{v}, \mathbf{0}_\varphi)$ is satisfiable in \mathcal{F} .*

For Boolean vectors $\mathbf{v} = (v_0, \dots, v_{l-1})$, $\mathbf{u} = (u_0, \dots, u_{l-1})$, let $\mathbf{v} + \mathbf{u}$ be their element-wise disjunction $(\max\{v_0, u_0\}, \dots, \max\{v_{l-1}, u_{l-1}\})$.

Lemma 14. *Let \mathcal{G} be a class of A -frames and $0 < b < \omega$. A tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\bigsqcup_{\leq b} \mathcal{G}$ iff there exist a positive $k \leq b$, $\mathbf{v}_0, \dots, \mathbf{v}_{k-1} \in 2^{\#\varphi}$ such that $\mathbf{v} = \sum_{i < k} \mathbf{v}_i$ and for every $i < k$ the tie $(\varphi, \mathbf{v}_i, \mathbf{U})$ is satisfiable in \mathcal{G} .*

Proof. This proposition is a corollary of Proposition 4 formulated in our new vector notation for ties. Indeed, by Proposition 4, we have

$$\varphi[\bigsqcup_{i < k} M_i, \Gamma] = \bigcup_{i < k} \varphi[M_i, \Gamma] \quad (4)$$

for every A -models M_0, \dots, M_{k-1} and every A -condition Γ . We are interested in the situation when $0 < k \leq b$ and frames of M_0, \dots, M_{k-1} are in \mathcal{G} . Assuming that Γ is the condition represented by \mathbf{U} , and vectors $\mathbf{v}, \mathbf{v}_1, \dots, \mathbf{v}_{k-1}$ are given by the identities $\varphi[\mathbf{v}] = \varphi[\bigsqcup_{i < k} M_i, \Gamma]$, $\varphi[\mathbf{v}_0] = \varphi[M_0, \Gamma]$, \dots , $\varphi[\mathbf{v}_{k-1}] = \varphi[M_{k-1}, \Gamma]$, we see that (4) takes the form $\mathbf{v} = \sum_{i < k} \mathbf{v}_i$. \square

For Boolean vectors $\mathbf{v}, \mathbf{u}_0, \dots, \mathbf{u}_{A-1}$ of the same length and $a \in A$, we put $(\mathbf{u}_0, \dots, \mathbf{u}_{A-1}) +^a \mathbf{v} = (\mathbf{u}'_0, \dots, \mathbf{u}'_{A-1})$, where $\mathbf{u}'_a = \mathbf{u}_a + \mathbf{v}$, and $\mathbf{u}'_b = \mathbf{u}_b$ for $b \neq a$.

Similarly to models, for A-frames \mathbf{F}_0 and \mathbf{F}_1 and $a \in A$ we define $\mathbf{F}_0 +^a \mathbf{F}_1$ as $\sum_{(2, <)}^a \mathbf{F}_i$; for classes \mathcal{F} and \mathcal{G} of A-frames, let $\mathcal{F} +^a \mathcal{G} = \{\mathbf{F} +^a \mathbf{G} \mid \mathbf{F} \in \mathcal{F} \ \& \ \mathbf{G} \in \mathcal{G}\}$.

Lemma 15. *Let \mathcal{F} and \mathcal{G} be classes of A-frames and $a \in A$. Then a tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\mathcal{F} +^a \mathcal{G}$ iff there exist $\mathbf{v}_0, \mathbf{v}_1 \in 2^{\# \varphi}$ such that*

1. $\mathbf{v} = \mathbf{v}_0 + \mathbf{v}_1$, and
2. $(\varphi, \mathbf{v}_0, \mathbf{U} +^a \mathbf{v}_1)$ is satisfiable in \mathcal{F} , and
3. $(\varphi, \mathbf{v}_1, \mathbf{U})$ is satisfiable in \mathcal{G} .

Proof. Let \mathbf{M}_0 be a model on a frame in \mathcal{F} , and let \mathbf{M}_1 be a model on a frame in \mathcal{G} . Let $\mathbf{\Gamma}$ be the condition represented by \mathbf{U} , i.e., $\mathbf{\Gamma} = (\varphi[\mathbf{U}(a)])_{a \in A}$, and let $\Psi = \varphi[\mathbf{M}_1, \mathbf{\Gamma}]$. By Lemma 9,

$$\varphi[\mathbf{M}_0 +^a \mathbf{M}_1, \mathbf{\Gamma}] = \varphi[\mathbf{M}_0, \mathbf{\Gamma} \cup^a \Psi] \cup \Psi. \quad (5)$$

For the “only if” part, assume that $\varphi[\mathbf{v}] = \varphi[\mathbf{M}_0 +^a \mathbf{M}_1, \mathbf{\Gamma}]$. Consider tuples $\mathbf{v}_0, \mathbf{v}_1 \in 2^{\# \varphi}$ such that

$$\varphi[\mathbf{v}_0] = \varphi[\mathbf{M}_0, \mathbf{\Gamma} \cup^a \Psi], \text{ and} \quad (6)$$

$$\varphi[\mathbf{v}_1] = \varphi[\mathbf{M}_1, \mathbf{\Gamma}]. \quad (7)$$

The identity (6) says that $(\varphi, \mathbf{v}_0, \mathbf{U} +^a \mathbf{v}_1)$ is satisfiable in \mathcal{F} , and the identity (7) says that $(\varphi, \mathbf{v}_1, \mathbf{U})$ is satisfiable in \mathcal{G} . Since $\Psi = \varphi[\mathbf{M}_1, \mathbf{\Gamma}]$, by (5) we obtain $\varphi[\mathbf{v}] = \varphi[\mathbf{v}_0] \cup \varphi[\mathbf{v}_1]$, that is $\mathbf{v} = \mathbf{v}_0 + \mathbf{v}_1$.

For the “if” part, assume that for some $\mathbf{v}_0, \mathbf{v}_1$ with $\mathbf{v} = \mathbf{v}_0 + \mathbf{v}_1$ we have (6) (this can be assumed since $(\varphi, \mathbf{v}_0, \mathbf{U} +^a \mathbf{v}_1)$ is satisfiable in \mathcal{F}), and (7) (this can be assumed since $(\varphi, \mathbf{v}_1, \mathbf{U})$ is satisfiable in \mathcal{G}). Since $\mathbf{v} = \mathbf{v}_0 + \mathbf{v}_1$, we have $\varphi[\mathbf{v}] = \varphi[\mathbf{v}_0] \cup \varphi[\mathbf{v}_1]$. Now by (5) we obtain that $\varphi[\mathbf{v}] = \varphi[\mathbf{M}_0 + \mathbf{M}_1, \mathbf{\Gamma}]$. The latter proves that $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\mathcal{F} +^a \mathcal{G}$. \square

Lemma 16. *Let \mathcal{F} be a class of A-frame, $a \in A$, and $0 < h, b < \omega$. A tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\sum_{\text{Tr}(h+1, b)}^a \mathcal{F}$ iff it is satisfiable in \mathcal{F} , or there exist a positive $k \leq b$ and $\mathbf{u}, \mathbf{v}_0, \dots, \mathbf{v}_{k-1} \in 2^{\# \varphi}$ such that*

1. $\mathbf{v} = \mathbf{u} + \sum_{i < k} \mathbf{v}_i$, and
2. the tie $(\varphi, \mathbf{u}, \mathbf{U} +^a \sum_{i < k} \mathbf{v}_i)$ is satisfiable in \mathcal{F} , and
3. for all $i < k$, the tie $(\varphi, \mathbf{v}_i, \mathbf{U})$ is satisfiable in $\sum_{\text{Tr}(h, b)}^a \mathcal{F}$.

Proof. By the definition of $\sum_{\text{Tr}(h+1, b)}^a \mathcal{F}$, the tie $\tau = (\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\sum_{\text{Tr}(h+1, b)}^a \mathcal{F}$ iff τ is satisfiable in \mathcal{F} or in $\mathcal{F} +^a \mathcal{G}$, where $\mathcal{G} = \bigsqcup_{\leq b} \sum_{\text{Tr}(h, b)}^a \mathcal{F}$.

By Lemma 15, τ is satisfiable in $\mathcal{F} +^a \mathcal{G}$ iff there exist $\mathbf{u}, \mathbf{u}' \in 2^{\# \varphi}$ such that $\mathbf{v} = \mathbf{u} + \mathbf{u}'$, $(\varphi, \mathbf{u}, \mathbf{U} +^a \mathbf{u}')$ is satisfiable in \mathcal{F} , and $(\varphi, \mathbf{u}', \mathbf{U})$ is satisfiable in \mathcal{G} . By Lemma 14, $(\varphi, \mathbf{u}', \mathbf{U})$ is satisfiable in \mathcal{G} iff there exist $0 < k \leq b$ and $\mathbf{v}_0, \dots, \mathbf{v}_{k-1} \in 2^{\# \varphi}$ such that $\mathbf{u}' = \sum_{i < k} \mathbf{v}_i$, and for all $i < k$ the tie $(\varphi, \mathbf{v}_i, \mathbf{U})$ is satisfiable in $\sum_{\text{Tr}(h, b)}^a \mathcal{F}$. \square

This lemma allows to describe the procedure $\text{CSatSum}_{\mathcal{F}}$ (Algorithm 1) which using an oracle for $\text{CSat } \mathcal{F}$ decides whether a given tie is satisfiable in $\sum_{\text{Tr}(h, b)}^a \mathcal{F}$. Namely, we have:

ALGORITHM 1: Decision procedure for $\text{CSat} \sum_{\text{Tr}(h,b)}^a \mathcal{F}$ with an oracle for $\text{CSat} \mathcal{F}$

$\text{CSatSum}_{\mathcal{F}}(\varphi, \mathbf{v}, \mathbf{U}, h, b) : \text{boolean}$

Input: A tie $(\varphi, \mathbf{v}, \mathbf{U})$; positive integers h, b

if $(\varphi, \mathbf{v}, \mathbf{U})$ *is satisfiable in* \mathcal{F} **then return** true;

if $h > 1$ **then**

for k *such that* $1 \leq k \leq b$ **do for** $\mathbf{u}, \mathbf{v}_0, \dots, \mathbf{v}_{k-1} \in 2^{\#\varphi}$ *such that* $\mathbf{v} = \mathbf{u} + \sum_{i < k} \mathbf{v}_i$ **do**

if $(\varphi, \mathbf{u}, \mathbf{U} + \sum_{i < k} \mathbf{v}_i)$ *is satisfiable in* \mathcal{F} **then**

if $\bigwedge_{i < k} \text{CSatSum}_{\mathcal{F}}(\varphi, \mathbf{v}_i, \mathbf{U}, h-1, b)$ **then return** true;

return false.

Theorem 17. *Let $a < A < \omega$, \mathcal{F} a class of A-frames, $(\varphi, \mathbf{v}, \mathbf{U})$ an A-tie, and $0 < h, b < \omega$. Then*

$$(\varphi, \mathbf{v}, \mathbf{U}) \text{ is satisfiable in } \sum_{\text{Tr}(h,b)}^a \mathcal{F} \text{ iff } \text{CSatSum}_{\mathcal{F}}(\varphi, \mathbf{v}, \mathbf{U}, h, b) \text{ returns true.}$$

This theorem will be our main technical tool for complexity results. For the first of its corollaries, we show how to reduce the conditional satisfiability on sums to the conditional satisfiability on summands.

Theorem 18. *Let $a < A < \omega$, \mathcal{F} a class of A-frames, and \mathcal{I} a class of Noetherian orders containing all finite trees. Then:*

$$1. \text{Sat} \sum_{\mathcal{I}}^a \mathcal{F} \leq_{\text{T}}^{\text{PSpace}} \text{CSat} \mathcal{F}.$$

$$2. \text{ If also } \mathcal{I} \text{ is closed under finite disjoint unions, then } \text{CSat} \sum_{\mathcal{I}}^a \mathcal{F} \leq_{\text{T}}^{\text{PSpace}} \text{CSat} \mathcal{F}.$$

Proof. By Theorem 10(1), a formula φ is satisfiable in $\sum_{\mathcal{I}}^a \mathcal{F}$ iff φ is satisfiable in $\sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}$, and by Proposition 13, this means that there exists a satisfiable in $\sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}$ tie $(\varphi, \mathbf{v}, \mathbf{0}_{\varphi})$ with $\mathbf{v}(0) = 1$. From Theorem 17 we obtain

Lemma 19. *A formula φ is satisfiable in $\sum_{\mathcal{I}}^a \mathcal{F}$ iff there exists $\mathbf{v} \in 2^{\#\varphi}$ such that $\mathbf{v}(0) = 1$ and $\text{CSatSum}_{\mathcal{F}}(\varphi, \mathbf{v}, \mathbf{0}_{\varphi}, \#\varphi, \#\varphi)$ returns true.*

Assume that \mathcal{I} is closed under finite disjoint unions. By Theorem 10(2), a tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\sum_{\mathcal{I}}^a \mathcal{F}$ iff it is satisfiable in $\bigsqcup_{\leq \#\varphi} \sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}$. By Lemma 14, this means that there exists $k \leq \#\varphi$ and tuples $\mathbf{v}^{(0)}, \dots, \mathbf{v}^{(k-1)} \in 2^{\#\varphi}$ such that $\mathbf{u} = \sum_{i < k} \mathbf{v}^{(i)}$ and $(\varphi, \mathbf{v}^{(i)}, \mathbf{U}, \#\varphi, \#\varphi)$ is satisfiable in $\sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}$ for every $i < k$. Using Theorem 17 again, we obtain

Lemma 20. *If \mathcal{I} is closed under finite disjoint unions, then a tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in $\sum_{\mathcal{I}}^a \mathcal{F}$ iff there exist $k \leq \#\varphi$ and tuples $\mathbf{v}^{(0)}, \dots, \mathbf{v}^{(k-1)} \in 2^{\#\varphi}$ such that $\mathbf{u} = \sum_{i < k} \mathbf{v}^{(i)}$ and $\text{CSatSum}_{\mathcal{F}}(\varphi, \mathbf{v}^{(i)}, \mathbf{U}, \#\varphi, \#\varphi)$ returns true for every $i < k$.*

Set $n = \#\varphi$. Let us estimate the amount of space used by $\text{CSatSum}_{\mathcal{F}}$ for the case $n = h = b$. At each call $\text{CSatSum}_{\mathcal{F}}$ needs $O(n^2)$ space to store new variables. The depth of recursion is bounded by n . Thus, we can reduce $\text{Sat} \sum_{\mathcal{I}}^a \mathcal{F}$ to the conditional satisfiability problem on \mathcal{F} in $O(n^3)$ space by Lemma 19. This proves the first statement of the theorem. In the case when \mathcal{I} is closed under finite disjoint unions, $\text{CSat} \sum_{\mathcal{I}}^a \mathcal{F}$ is also reducible to the conditional satisfiability problem on \mathcal{F} in $O(n^3)$ space by Lemma 20. This proves the second statement of the theorem. \square

4.2 Reductions between $\text{Sat } \mathcal{F}$, $\text{CSat } \mathcal{F}$, and $\text{Sat } \mathcal{F}^{(\forall)}$

By Proposition 13, the satisfiability problem on a class \mathcal{F} is polynomial space Turing reducible to the conditional satisfiability problem on \mathcal{F} . Let \leq_m^{PTime} denote the polynomial-time many-to-one reduction. Below we show that in many cases $\text{CSat } \mathcal{F}$ is polynomial time reducible to $\text{Sat } \mathcal{F}$, that is $\text{CSat } \mathcal{F} \leq_m^{\text{PTime}} \text{Sat } \mathcal{F}$; hence, in these cases, the above two problems are equivalent with respect to \leq_T^{PSpace} . This fact is close to a result obtained in [Hem96], where it was shown that in many situations there exists a (stronger than \leq_T^{PSpace} , but weaker¹ than \leq_m^{PTime}) reduction of $\text{Sat } \mathcal{F}^{(\forall)}$ to $\text{Sat } \mathcal{F}$. Let us discuss reductions between the above two problems in more details.

For a binary relation R on a set W , put $R^{\leq m} = \bigcup_{n \leq m} R^n$, where R^0 is the identity relation on W , $R^{n+1} = R \circ R^n$ (\circ denotes the composition of relations). Recall that R^* denotes the transitive reflexive closure of R : $R^* = \bigcup_{n < \omega} R^n$. R is said to be m -transitive if $R^{\leq m}$ includes R^{m+1} , or equivalently, $R^{\leq m} = R^*$ (e.g., every transitive relation is 1-transitive). For a frame $\mathbf{F} = (W, (R_a)_{a \in A})$, put $R_{\mathbf{F}} = \bigcup_{a \in A} R_a$. We say that \mathbf{F} is m -transitive if the relation $R_{\mathbf{F}}$ is. In particular, if one of the relations of \mathbf{F} is universal (i.e., is equal to $W \times W$), then \mathbf{F} is 1-transitive.

For w in \mathbf{F} , let $\mathbf{F}[w]$ denote the subframe of \mathbf{F} generated by the singleton $\{w\}$ (that is, $\mathbf{F}[w]$ is the restriction of \mathbf{F} on the set $\{v \mid w R_{\mathbf{F}}^* v\}$); such frames are called *cones*). A class \mathcal{F} of frames is *closed under taking cones* if for every \mathbf{F} in \mathcal{F} and for every w in \mathbf{F} , the cone $\mathbf{F}[w]$ is in \mathcal{F} .

A class \mathcal{F} of A-frames is said to be *preconical*, if

- \mathcal{F} is closed under taking cones, and
- there exists $m < \omega$ such that every frame in \mathcal{F} is m -transitive, and
- for every \mathbf{F} in \mathcal{F} , the relation $R_{\mathbf{F}}^*$ is downward directed (i.e., for every w, v in \mathbf{F} there exists u such that $u R_{\mathbf{F}}^* w$ and $u R_{\mathbf{F}}^* v$).

Proposition 21. *Let A be finite and \mathcal{F} a class of A-frames. If \mathcal{F} is preconical, then $\text{CSat } \mathcal{F} \leq_m^{\text{PTime}} \text{Sat } \mathcal{F}$.*

Proof. Given a condition $\mathbf{\Gamma}$ and a formula φ , we define the formula $[\varphi]^{\mathbf{\Gamma}}$ as follows: $[\perp]^{\mathbf{\Gamma}} = \perp$, $[p]^{\mathbf{\Gamma}} = p$ for variables, $[\varphi_1 \rightarrow \varphi_2]^{\mathbf{\Gamma}} = [\varphi_1]^{\mathbf{\Gamma}} \rightarrow [\varphi_2]^{\mathbf{\Gamma}}$, and

$$[\Diamond_a \varphi]^{\mathbf{\Gamma}} = \begin{cases} \top, & \text{if } \varphi \in \mathbf{\Gamma}(a), \\ \Diamond_a [\varphi]^{\mathbf{\Gamma}} & \text{otherwise.} \end{cases}$$

For every A-model \mathbf{M} , we have:

$$\mathbf{M}, w \models_{\mathbf{\Gamma}} \varphi \text{ iff } \mathbf{M}, w \models [\varphi]^{\mathbf{\Gamma}} \quad (8)$$

The proof is straightforward, see [Sha18, Lemma 4.7] for the details.

Let $\Diamond \varphi$ abbreviate the A-formula $\bigvee_{a \in A} \Diamond_a \varphi$, and let $\Diamond^0 \varphi = \varphi$, $\Diamond^{m+1} \varphi = \Diamond \Diamond^m \varphi$, $\Diamond^{\leq m} \varphi = \bigvee_{n \leq m} \Diamond^n \varphi$.

For a tie $(\varphi, \mathbf{v}, \mathbf{U})$ with $\mathbf{U} = (\mathbf{u}_a)_{a \in A}$, we put

$$\delta_m(\varphi, \mathbf{v}, \mathbf{U}) = \bigwedge_{\psi \in \varphi[\mathbf{v}]} \Diamond^{\leq m} [\psi]^{\mathbf{\Gamma}} \wedge \bigwedge_{\psi \in \text{sub}(\varphi) \setminus \varphi[\mathbf{v}]} \neg \Diamond^{\leq m} [\psi]^{\mathbf{\Gamma}}, \quad (9)$$

where $\mathbf{\Gamma}$ is the condition represented by \mathbf{U} , i.e., $\mathbf{\Gamma} = (\varphi[\mathbf{u}_a])_{a \in A}$.

Since \mathcal{F} is preconical, there exists a finite m such that every frame in \mathcal{F} is m -transitive. We claim that

$$(\varphi, \mathbf{v}, \mathbf{U}) \in \text{CSat } \mathcal{F} \text{ iff } \delta_m(\varphi, \mathbf{v}, \mathbf{U}) \in \text{Sat } \mathcal{F}. \quad (10)$$

¹I am using “stronger” and “weaker” in a non-strict sense.

First, assume that $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in a frame $\mathbf{F} \in \mathcal{F}$. This means that for a model \mathbf{M} based on \mathbf{F} we have $\varphi[\mathbf{v}] = \varphi[\mathbf{M}, \mathbf{\Gamma}]$, where $\mathbf{\Gamma}$ is the condition represented by \mathbf{U} . For every $\psi \in \varphi[\mathbf{v}]$ we choose a point w_ψ such that $\mathbf{M}, w_\psi \models_{\mathbf{\Gamma}} \psi$, and then put $V = \{w_\psi \mid \psi \in \varphi[\mathbf{v}]\}$. The relation $R_{\mathbf{F}}^*$ is downward directed, hence there exists a point w in \mathbf{M} such that $wR_{\mathbf{F}}^*v$ for all v in V ; by m -transitivity, $wR_{\mathbf{F}}^{\leq m}v$. It follows that if $\psi \in \varphi[\mathbf{v}]$, then $\mathbf{M}, w \models \Diamond^{\leq m}[\psi]^{\mathbf{\Gamma}}$: indeed, we have $\mathbf{M}, w_\psi \models [\psi]^{\mathbf{\Gamma}}$ by (8) and $wR_{\mathbf{F}}^{\leq m}w_\psi$. On the other hand, if a subformula ψ of φ is not in $\varphi[\mathbf{v}]$, then $[\psi]^{\mathbf{\Gamma}}$ is false at every point in \mathbf{M} by (8), and so $\mathbf{M}, w \models \neg \Diamond^{\leq m}[\psi]^{\mathbf{\Gamma}}$. It follows that the formula $\delta_m(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in \mathcal{F} .

Now assume that $\delta_m(\varphi, \mathbf{v}, \mathbf{U})$ is true at a point w in a model \mathbf{M} over a frame $\mathbf{F} \in \mathcal{F}$. Since \mathcal{F} is closed under taking cones, we may assume that $\mathbf{F} = \mathbf{F}[w]$ (recall that if a formula is true at a point in a model, then it is true at this point in the model generated by this point). Let ψ be a subformula of φ . Suppose that $\psi \in \varphi[\mathbf{v}]$. Then $\mathbf{M}, w \models \Diamond^{\leq m}[\psi]^{\mathbf{\Gamma}}$. Hence, the formula $[\psi]^{\mathbf{\Gamma}}$ is true at a point u of \mathbf{M} , which means that $\mathbf{M}, u \models_{\mathbf{\Gamma}} \psi$ by (8). Thus, $\psi \in \varphi[\mathbf{M}, \mathbf{\Gamma}]$. On the other hand, if $\psi \notin \varphi[\mathbf{v}]$, then $\mathbf{M}, w \models \neg \Diamond^{\leq m}[\psi]^{\mathbf{\Gamma}}$; by m -transitivity, we obtain that $[\psi]^{\mathbf{\Gamma}}$ is false at every point of \mathbf{M} ; using (8) again, we obtain $\psi \notin \varphi[\mathbf{M}, \mathbf{\Gamma}]$. Thus, $\varphi[\mathbf{v}] = \varphi[\mathbf{M}, \mathbf{\Gamma}]$. Since $\mathbf{\Gamma} = (\varphi[\mathbf{u}_a])_{a \in A}$, the tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in \mathcal{F} . \square

Proposition 22. *If A is finite and \mathcal{F} is a class of A -frames, then $\text{CSat } \mathcal{F} \leq_m^{\text{PTime}} \text{Sat } \mathcal{F}^{(\forall)}$.*

Proof. It is trivial that $\text{CSat } \mathcal{F} \leq_m^{\text{PTime}} \text{CSat } \mathcal{F}^{(\forall)}$ (the reduction increases by one the indexes of modalities occurring in formulas of a given tie).

The class $\mathcal{F}^{(\forall)}$ is preconical: if $\mathbf{F} \in \mathcal{F}$, then \mathbf{F} is 1-transitive, $\mathbf{F}[w] = \mathbf{F}$ for every $w \in \mathbf{F}$, and $R_{\mathbf{F}}^*$ is downward-directed, since $R_{\mathbf{F}}$ is the universal relation on \mathbf{F} . Hence, $\text{CSat } \mathcal{F}^{(\forall)} \leq_m^{\text{PTime}} \text{Sat } \mathcal{F}^{(\forall)}$ by Proposition 21. \square

Now let us describe a reduction of $\text{Sat } \mathcal{F}^{(\forall)}$ to $\text{CSat } \mathcal{F}$. This reduction is based on the following construction proposed in [Hem96]. Let φ be a formula in the language $\text{ML}(1 + A)$. For subformulas $\Diamond_0\psi$ of φ starting with \Diamond_0 , we choose distinct variables p_ψ not occurring in φ , and put for subformulas of φ : $\perp' = \perp$; $p' = p$ for variables; $(\varphi_1 \rightarrow \varphi_2)' = \varphi_1' \rightarrow \varphi_2'$; $(\Diamond_a\psi)' = \Diamond_a\psi'$ for $a > 0$; and $(\Diamond_0\psi)' = p_\psi$. We have for a class \mathcal{F} of A -frames:

$$\varphi \in \text{Sat } \mathcal{F}^{(\forall)} \text{ iff } \varphi' \wedge \bigwedge_{\Diamond_0\psi \in \text{sub}(\varphi)} ((\Diamond_0\psi' \leftrightarrow \Box_0 p_\psi) \wedge (\Box_0 p_\psi \vee \Box_0 \neg p_\psi)) \in \text{Sat } \mathcal{F}^{(\forall)}, \quad (11)$$

see [Hem96, Lemma 4.5] for details.²

Let η denote the formula in the right-hand side of the above equivalence. Let us show how the satisfiability of η in $\mathcal{F}^{(\forall)}$ can be expressed as the satisfiability of a tie in \mathcal{F} . Consider the formula

$$\xi_\varphi = \varphi' \wedge \bigwedge_{\Diamond_0\psi \in \text{sub}(\varphi)} (\psi' \wedge \neg p_\psi)$$

(the only role of the second conjunct is to have ψ' , $\neg p_\psi$, p_ψ that occur in η as subformulas of ξ_φ). Let \mathbf{M} be a model on a frame in $\mathcal{F}^{(\forall)}$, and let Φ be the set of subformulas of ξ_φ that are satisfiable in \mathbf{M} , that is, $\Phi = \xi_\varphi[\mathbf{M}, (\emptyset)_{1+A}]$. Then η is true at a point in \mathbf{M} iff

$$\varphi' \in \Phi \text{ and for every } \psi \text{ with } \Diamond_0\psi \in \text{sub}(\varphi), \psi' \in \Phi \text{ iff } p_\psi \in \Phi \text{ iff } \neg p_\psi \notin \Phi. \quad (12)$$

It follows that η is satisfiable in $\mathcal{F}^{(\forall)}$ iff there exists $\Phi \subseteq \text{sub}(\xi_\varphi)$ satisfying (12) and the tie $(\xi_\varphi, \Phi, (\emptyset)_{1+A})$ is satisfiable in $\mathcal{F}^{(\forall)}$. The formula ξ_φ and so, the formulas in Φ , do not contain \Diamond_0 . Hence, the satisfiability of $(\xi_\varphi, \Phi, (\emptyset)_{1+A})$ in $\mathcal{F}^{(\forall)}$ is equivalent to the satisfiability

²The formula in (11) is an equivalent form of the formula φ_{flat} used in [Hem96, Lemma 4.5].

of $(\dot{\xi}_\varphi, \dot{\Phi}, (\emptyset)_A)$ in \mathcal{F} , where $\dot{\xi}_\varphi$ and $\dot{\Phi}$ are obtained by decreasing indexes of modalities by 1. Putting everything together, we have shown that

$\varphi \in \text{Sat } \mathcal{F}^{(\forall)}$ iff there exists $\Phi \subseteq \text{sub}(\xi_\varphi)$ satisfying (12) s. t. $(\dot{\xi}, \dot{\Phi}, (\emptyset)_A)$ is satisfiable in \mathcal{F} .

This proves

Proposition 23. *If A is finite and \mathcal{F} is a class of A -frames, then $\text{Sat } \mathcal{F}^{(\forall)} \leq_T^{\text{PSpace}} \text{CSat } \mathcal{F}$.*

Remark 24. In fact, Proposition 23 provides a stronger reduction than \leq_T^{PSpace} .

4.3 Proof of Theorem 11

In view of the above propositions, Theorem 11 is an easy corollary of Theorem 18.

Proof of Theorem 11. We obtain $\text{Sat } \sum_{\mathcal{I}}^a \mathcal{F} \leq_T^{\text{PSpace}} \text{CSat } \mathcal{F}$ by Theorem 18(1), and then $\text{CSat } \mathcal{F} \leq_m^{\text{PTime}} \text{Sat } \mathcal{F}^{(\forall)}$ by Proposition 22. Since \leq_m^{PTime} is stronger than \leq_T^{PSpace} , we obtain $\text{CSat } \mathcal{F} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$. Hence, $\text{Sat } \sum_{\mathcal{I}}^a \mathcal{F} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$, which proves the first statement of the theorem.

To prove the second statement, we start with Proposition 23 and obtain $\text{Sat}(\sum_{\mathcal{I}}^a \mathcal{F})^{(\forall)} \leq_T^{\text{PSpace}} \text{CSat}(\sum_{\mathcal{I}}^a \mathcal{F})$. Now by Theorem 18(2) and Proposition 22, we obtain $\text{CSat } \sum_{\mathcal{I}}^a \mathcal{F} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$. So $\text{Sat}(\sum_{\mathcal{I}}^a \mathcal{F})^{(\forall)} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$, which completes the proof. \square

Recall that in the preconical case, the conditional satisfiability problem is reducible to the standard modal satisfiability problem (Proposition 21), and so $\text{CSat } \mathcal{F}$, $\text{Sat } \mathcal{F}$, and $\text{Sat } \mathcal{F}^{(\forall)}$ are \leq_T^{PSpace} -equivalent by Propositions 22, 23 (in general, the decidability of $\text{Sat } \mathcal{F}$ does not imply the decidability of $\text{Sat } \mathcal{F}^{(\forall)}$, see [Spa93, Theorem 4.2.1]). In this case, Theorem 11 can be reformulated in the following way:

Corollary 25. *Let $a < A < \omega$, \mathcal{F} a class of A -frames, and \mathcal{I} a class of Noetherian orders containing all finite trees. If \mathcal{F} is preconical, then:*

1. $\text{Sat } \sum_{\mathcal{I}}^a \mathcal{F} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}$.
2. *If also \mathcal{I} is closed under finite disjoint unions, then $\text{Sat}(\sum_{\mathcal{I}}^a \mathcal{F})^{(\forall)} \leq_T^{\text{PSpace}} \text{Sat } \mathcal{F}$.*

4.4 PSPACE-hardness: some corollaries of Ladner's construction

According to Ladner's theorem, every unimodal logic contained in S4 is PSpace-hard [Lad77]. In fact, Ladner's proof yields PSpace-hardness for a wider class of modal logics (e.g., for logics contained in the Gödel-Löb logic GL or in the Grzegorczyk logic GRZ), see [Spa93]. With minor modifications, Ladner's construction also works for logics contained in S4.1, GRZ.2 etc, for the polymodal case, and in particular – for sums.

To illustrate this, let us briefly discuss the proof. Consider a quantified Boolean formula $\eta = Q_1 p_1 \dots Q_m p_m \theta$, where $Q_1, \dots, Q_m \in \{\exists, \forall\}$, and θ is a propositional Boolean formula in variables p_1, \dots, p_m . Choose fresh variables q_0, \dots, q_m . Let $[\eta]_L$ be the following unimodal

formula:³

$$\begin{aligned}
& q_0 \wedge \bigwedge_{i < m} \Box^{\leq m}(q_i \rightarrow \Diamond q_{i+1}) \wedge \bigwedge_{\substack{i \neq j \\ i, j \leq m}} \Box^{\leq m}(q_i \rightarrow \neg q_j) \wedge \Box^m(q_m \rightarrow \theta) \wedge \\
& \bigwedge_{\{i < m \mid Q_{i+1} = \forall\}} \Box^i(q_i \rightarrow \Diamond(q_{i+1} \wedge p_{i+1}) \wedge \Diamond(q_{i+1} \wedge \neg p_{i+1})) \wedge \\
& \bigwedge_{1 \leq i \leq m-1} \Box^i(q_i \rightarrow \Box^{\leq m} p_i \vee \Box^{\leq m} \neg p_i)
\end{aligned}$$

where $\Box^m = \neg \Diamond^m \neg$; likewise for $\Box^{\leq m}$.

Let $\mathsf{T}_\eta = (W_\eta, R_\eta)$ be the *quantifier tree* of η : $W_\eta = \bigcup_{k \leq m} \{\sigma \in 2^k \mid \sigma(i) = 0 \text{ if } Q_{i+1} = \exists\}$; $\sigma_1 R_\eta \sigma_2$ iff $\sigma_1 \subset \sigma_2$ and $\text{dom}(\sigma_2) = \text{dom}(\sigma_1) + 1$. We remark that T_η is antitransitive (that is, $xRyRz \Rightarrow \neg(xRz)$ holds in T_η), and so is irreflexive. We have:

$$\eta \text{ is valid} \quad \Rightarrow \quad [\eta]_{\mathsf{L}} \text{ is satisfiable in } (W_\eta, R_\eta^*), \quad (13)$$

$$[\eta]_{\mathsf{L}} \text{ is satisfiable in a Kripke frame} \quad \Rightarrow \quad \eta \text{ is valid}; \quad (14)$$

see [BdRV02, Section 6.7] for the details.

Consider a unimodal logic $L \subseteq \mathsf{S4}$. Then (W_η, R_η^*) is an L -frame, hence every satisfiable in this frame formula is L -consistent. So we have:

$$\eta \text{ is valid} \quad \Rightarrow \quad [\eta]_{\mathsf{L}} \text{ is satisfiable in } (W_\eta, R_\eta^*) \quad \Rightarrow \quad [\eta]_{\mathsf{L}} \text{ is } L\text{-consistent} \quad \Rightarrow \quad \eta \text{ is valid},$$

that is, η is valid iff $[\eta]_{\mathsf{L}}$ is L -consistent. Thus, the L -consistency problem is PSpace-hard; synonymously, the (provability problem for the) logic L is PSpace-hard.

This proves Ladner's theorem, in its classical formulation. And, in fact, it proves more. Let GRZ.BIN be the logic of the class of all finite transitive reflexive trees with branching ≤ 2 . We have the following proper inclusions

$$\mathsf{S4} \subsetneq \mathsf{S4.1} \subsetneq \text{GRZ} \subsetneq \text{GRZ.BIN},$$

see, e.g., [CZ97], and [GDJ74] for the latter inclusion. Observe that (W_η, R_η^*) is a finite transitive reflexive tree. It immediately follows from the above reasonings that every logic contained in GRZ is PSpace-hard. Moreover, the branching of (W_η, R_η^*) is ≤ 2 ; thus, the result holds for logics contained in GRZ.BIN :

$$L \subseteq \text{GRZ.BIN} \quad \Rightarrow \quad L \text{ is PSpace-hard.}$$

This formulation does not include an important logic GL : its frames are irreflexive. However, we can reformulate (13) in the following way: if $R_\eta \subseteq R \subseteq R_\eta^*$, then

$$\eta \text{ is valid} \quad \Rightarrow \quad [\eta]_{\mathsf{L}} \text{ is satisfiable in } (W_\eta, R); \quad (15)$$

the proof is straightforward and is an immediate analog of the proof of (13) given in [BdRV02].

The condition $R_\eta \subseteq R \subseteq R_\eta^*$ can be reformulated as $R^* = R_\eta^*$, because T_η is a tree.

Definition 5. A class \mathcal{F} of unimodal frames is *thick* if for every finite transitive reflexive tree $\mathsf{T} = (T, \leq)$ whose branching is bounded by 2 there exist a relation R on T and a frame $\mathsf{F} \in \mathcal{F}$ such that $R^* = \leq$ and F is isomorphic to (T, R) .

³This variant of reduction is a slight modification of the one used in [BdRV02].

From (15) we obtain that if \mathcal{F} is thick then

$$\eta \text{ is valid} \quad \Rightarrow \quad [\eta]_L \text{ is satisfiable in } \mathcal{F}. \quad (16)$$

Thus, for a class \mathcal{F} of unimodal frames and a unimodal logic L , from (16) and (14) we have:

$$\mathcal{F} \text{ is thick and } L \subseteq \text{Log } \mathcal{F} \quad \Rightarrow \quad L \text{ is PSpace-hard.} \quad (17)$$

In particular, this formulation allows to apply Ladner's construction for logics below GL, since GL is the logic of a thick class (consisting of finite irreflexive transitive trees).

The logic S4.2 is not the logic of a thick class: it does not have trees of height > 1 within its frames. However, trees are contained as subframes in S4.2-frames. And this is also enough for PSpace-hardness due to the following relativization argument proposed by E. Spaan [Spa93].

Recall that a *subframe* of a frame $F = (W, (R_a)_{a \in A})$ is the restriction $F \upharpoonright V = (V, (R_a \cap (V \times V))_{a \in A})$, where $V \neq \emptyset$. For a class \mathcal{F} of frames, let $\text{Sub } \mathcal{F}$ be its closure under the subframe operation: $\text{Sub } \mathcal{F} = \{F \upharpoonright V \mid F \in \mathcal{F} \text{ and } \emptyset \neq V \subseteq \text{dom}(F)\}$. The satisfiability in $\text{Sub } \mathcal{F}$ can be reduced to the satisfiability in \mathcal{F} (see the proof of Theorem 2.2.1 in [Spa93]). Namely, for an A-formula φ and a variable q , let $[\varphi]_q$ be the A-formula inductively defined as follows: $[\perp]_q = \perp$, $[p]_q = p$ for variables, $[\psi_1 \rightarrow \psi_2]_q = [\psi_1]_q \rightarrow [\psi_2]_q$, and $[\Diamond_a \psi]_q = \Diamond_a([\psi]_q \wedge q)$ for $a \in A$. We put $[\varphi]_{\text{Rel}} = q \wedge [\varphi]_q$, where q is the first variable not occurring in φ . If $M = (F, \theta)$ is a model and $V = \theta(q)$, then for every $v \in V$ we have: $M \upharpoonright V, v \models \varphi$ iff $M, v \models [\varphi]_{\text{Rel}}$; the proof is by induction on φ . This immediately yields

Proposition 26 ([Spa93]). *φ is satisfiable in $\text{Sub } \mathcal{F}$ iff $[\varphi]_{\text{Rel}}$ is satisfiable in \mathcal{F} .*

Thus, $\text{Sat Sub } \mathcal{F}$ is polynomial time reducible to $\text{Sat } \mathcal{F}$.

It immediately follows from (17) and Proposition 26 that if $\text{Sub } \mathcal{F}$ is thick, then the logic of \mathcal{F} is PSpace-hard. Moreover, this holds for every $L \subseteq \text{Log } \mathcal{F}$: a quantified Boolean formula η is valid iff $[[\eta]_L]_{\text{Rel}}$ is L -consistent. Indeed, if η is valid, then $[\eta]_L$ is satisfiable in $\text{Sub } \mathcal{F}$ by (16), so $[[\eta]_L]_{\text{Rel}}$ is satisfiable in \mathcal{F} by Proposition 26, hence $[[\eta]_L]_{\text{Rel}}$ is L -consistent. The converse implication follows from (14). Thus, we have

Theorem 27 (A corollary of [Lad77] and [Spa93]). *Let \mathcal{F} be a class of unimodal frames. If $\text{Sub } \mathcal{F}$ is thick, then every unimodal $L \subseteq \text{Log } \mathcal{F}$ is PSpace-hard.*

For an A-frame $F = (W, (R_a)_{a \in A})$ and $a \in A$, let $F^{\upharpoonright \{a\}}$ be its reduct (W, R_a) . A class \mathcal{F} of A-frames is said to be *thick* if for some $a \in A$ the class $\{F^{\upharpoonright \{a\}} \mid F \in \mathcal{F}\}$ is thick.

Proposition 28. *Let \mathcal{F} and \mathcal{I} be classes of A-frames. If \mathcal{F} is non-empty and $\text{Sub } \mathcal{I}$ is thick, then $\text{Sub } \sum_{\mathcal{I}} \mathcal{F}$ is thick.*

Proof. Follows from Definitions 1 and 5. □

Corollary 29. *Let \mathcal{F} and \mathcal{I} be classes of A-frames. If \mathcal{F} is non-empty and $\text{Sub } \mathcal{I}$ is thick, then $\text{Sat } \sum_{\mathcal{I}} \mathcal{F}$ is PSpace-hard.*

4.5 Examples

For many logics, Theorem 11 gives a uniform proof of decidability in PSpace (PSpace-completeness in view of Corollary 29). Let us illustrate it with certain examples.

Example 3. It is well-known that the logic S4 as well as its expansion with the universal modality are PSpace-complete [Lad77],[Hem96].

Let us prove it via sums. Recall that clusters are frames of the form $(C, C \times C)$. Every preorder F is isomorphic to the sum $\sum_{C \in \text{sk}F} (C, C \times C)$ of its clusters over its skeleton $\text{sk}F$. The logic S4 has the finite model property, so it is enough to consider only finite indices, and hence S4 is the logic of the class

$$\sum_{\text{finite partial orders}} \text{clusters} \quad (18)$$

Thus, we have:

$$\text{Sat}(\text{preorders}) \leq_T^{\text{PSpace}} \text{Sat}(\text{clusters})$$

or dually

$$\text{S4} \leq_T^{\text{PSpace}} \text{S5}.$$

The satisfiability problem in clusters is in NP (one can easily check that a formula φ is satisfiable in a cluster iff it is satisfiable in a cluster of size $\#\varphi$; see, e.g., [CZ97, Section 18.3]). Now PSpace-completeness of S4 follows from (18): we have PSpace-upper bound by Theorem 11(1); PSpace-hardness is given by Corollary 29. Moreover, Theorem 11(2) gives PSpace-upper bound of the logic $\text{S4}^{[\vee]} = \text{Log}\{(W, W \times W, R) \mid (W, R) \text{ is a preorder}\}$.

Changing the class of summands, we obtain PSpace-completeness for other logics. Let S_0 and S_1 be an irreflexive and a reflexive singleton, respectively.

If we add S_0 to the class of summands-clusters, then we obtain PSpace-completeness of K4 and $\text{K4}^{[\vee]}$.

Letting the class of summands be $\{S_0\}$, we obtain PSpace-completeness for the logics GL. If the class of summands consists of a reflexive singleton S_1 , the above reasonings give PSpace-completeness of the Grzegorczyk logic, and if the class of summands is $\{S_0, S_1\}$ — for the weak Grzegorczyk logic (recall that the latter logic is characterized by frames whose reflexive closures are non-strict Noetherian orders [Lit07]; this class can be represented as $\sum_{\text{NPO}} \{S_0, S_1\}$).

The results discussed in Example 3 are well-known [Lad77],[Spa93],[Hem96]. To the best of our knowledge, the following result has never been published before.

Example 4. The *weak transitivity logic* wK4 is the logic of frames satisfying the condition $xRzRy \Rightarrow xRy \vee x = y$. The *difference logic* DL is the logic of the frames such that xRy whenever $x \neq y$; let \mathcal{F}_{\neq} be the class of such frames. The logic wK4 has the finite model property [Esa01], hence it is the logic of the class

$$\sum_{\text{finite partial orders}} \mathcal{F}_{\neq}, \quad (19)$$

see Example 2. It is not difficult to check that $\text{Sat } \mathcal{F}_{\neq}$ is in NP (like in the case of clusters, the size of a countermodel is linear in the length of a formula), and by Theorem 11 we obtain that

$$\text{wK4} \leq_T^{\text{PSpace}} \text{DL} \in \text{PSpace},$$

so wK4 is in PSpace (and PSpace-complete: PSpace-hardness follows from [Lad77], since $\text{wK4} \subseteq \text{S4}$, or from the representation (19) and Corollary 29). Moreover, since the class \mathcal{F}_{\neq} is preconical, $\text{wK4}^{[\vee]}$ is in PSpace.

Example 5. The logic wK4.2 is the logic of weakly transitive frames (considered in the above example) satisfying the Church-Rosser property $xRy_1 \& xRy_2 \Rightarrow \exists z (y_1Rz \& y_2Rz)$. Observe that every frame validating wK4.2 is either in \mathcal{F}_{\neq} , or is isomorphic to the sum of a frame validating wK4 and a frame in \mathcal{F}_{\neq} .

$$\text{Fr}(\text{wK4.2}) = \mathcal{F}_{\neq} \cup (\text{Fr}(\text{wK4}) +^0 \mathcal{F}_{\neq}).$$

Now that wK4.2 is decidable in PSpace follows from the previous example and the following theorem.

Theorem 30. *Let \mathcal{F} and \mathcal{G} be classes of \mathbf{A} -frames, and $a \in \mathbf{A}$. If $\text{Sat } \mathcal{G}^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$ (in particular, if $\text{Sat } \mathcal{G}^{(\forall)}$ is in PSpace), then $\text{Sat}(\mathcal{F} +^a \mathcal{G})^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$, and consequently $\text{Sat}(\mathcal{F} +^a \mathcal{G}) \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$.*

Proof. By Propositions 22 and 23, from $\text{Sat } \mathcal{G}^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$ we obtain $\text{CSat } \mathcal{G} \leq_{\text{T}}^{\text{PSpace}} \text{CSat } \mathcal{F}$. Using Lemma 15, from the latter reduction we obtain $\text{CSat}(\mathcal{F} +^a \mathcal{G}) \leq_{\text{T}}^{\text{PSpace}} \text{CSat } \mathcal{F}$. Using Propositions 22 and 23 again, we obtain that $\text{Sat}(\mathcal{F} +^a \mathcal{G})^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$. \square

Remark 31. This theorem can be strengthened in two aspects. First, it can be formulated for reductions stronger than $\leq_{\text{T}}^{\text{PSpace}}$. Another observation is that instead of the class $\mathcal{F} +^a \mathcal{G}$, i.e., a class of sums over $\mathbf{l} = (2, <)$, one can consider sums over an arbitrary finite indexing frame \mathbf{l} ; in this case, a reduction (with several oracles) from sums to summands would follow from Lemma 8.

We conclude this section with the following example. In topological semantics (\Diamond is for the derivation), wK4 is the logic of all topological spaces [Esa01]. In [BEG11] it was shown that the logic wK4T_0 of all T_0 -spaces is equal to the logic of all finite weakly transitive frames where clusters contain at most one irreflexive point. The satisfiability for such clusters is in NP, and we obtain

Corollary 32. *wK4T_0 is PSpace -complete.*

5 Variations

In this section we are interested in polymodal logics which can be characterized by models obtained via multiple applications of the sum operation. An important example of such logic is *Japaridze's polymodal logic* [Jap86, Bek10]; we also consider *lexicographic products of modal logics* introduced in [Bal09] and *refinements of logics* introduced in [BR10]; see Sections 5.3 and 5.4 below. We anticipate them with some general observations.

5.1 Iterated sums over unimodal indexes

Let \mathcal{F} be a class of \mathbf{A} -frames and \mathcal{I} a class of 1-frames. Let $\mathbf{a} = (a_0, \dots, a_{s-1})$ be a finite sequence of elements of \mathbf{A} . If \mathbf{a} is the empty sequence, let \mathbf{a} -iterated sums of \mathcal{F} be the elements of \mathcal{F} . For $0 < s < \omega$, let \mathbf{a} -iterated sums of \mathcal{F} be a_0 -sums of (a_1, \dots, a_{s-1}) -iterated sums: that is, an \mathbf{a} -iterated sum of frames in \mathcal{F} over frames in \mathcal{I} is a frame of form $\sum_{\mathbf{l}}^{a_0} H_i$, where $\mathbf{l} \in \mathcal{I}$ and every H_i is an (a_1, \dots, a_{s-1}) -iterated sum of frames in \mathcal{F} over frames in \mathcal{I} . The class of all such sums is denoted by $\sum_{\mathcal{I}}^{\mathbf{a}} \mathcal{F}$.

Since $\sum_{\mathcal{I}}^{\mathbf{a}} \mathcal{F}$ is $\sum_{\mathcal{I}}^{a_0} \sum_{\mathcal{I}}^{(a_1, \dots, a_{s-1})} \mathcal{F}$, by Theorem 11 we obtain

Corollary 33. *Let $\mathbf{A} < \omega$, \mathcal{F} a class of \mathbf{A} -frames, and \mathbf{a} a finite sequence of elements of \mathbf{A} . If \mathcal{I} is a class of Noetherian orders that contains all finite trees and is closed under finite disjoint unions, then $\text{Sat}(\sum_{\mathcal{I}}^{\mathbf{a}} \mathcal{F})^{(\forall)} \leq_{\text{T}}^{\text{PSpace}} \text{Sat } \mathcal{F}^{(\forall)}$.*

Theorem 10 reduces satisfiability on sums over Noetherian orders to sums over finite trees. This theorem can also be extended for the case of iterated sums [Sha18]. Namely, let $\mathbf{a} = (a_0, \dots, a_{s-1}) \in \mathbf{A}^s$, $0 < s < \omega$. Then for every \mathbf{A} -tie $\tau = (\varphi, \Phi, \Gamma)$ we have:

$$\tau \text{ is satisfiable in } \sum_{\text{NPO}}^{\mathbf{a}} \mathcal{F} \text{ iff } \tau \text{ is satisfiable in } \bigsqcup_{\leq \# \varphi} \sum_{\text{Tr}(\# \varphi, \# \varphi)}^{\mathbf{a}} \mathcal{F}. \quad (20)$$

This can be proven by induction of the length of \mathbf{a} with the help of the following lemma (see the proof of [Sha18, Theorem 5.2] for the details):

Lemma 34. *Let \mathcal{F} be a class of A-frames and $a \in A$. Then every frame in $\sum_{\text{NPO}}^a \sqcup \mathcal{F}$ is isomorphic to a frame in $\sum_{\text{NPO}}^a \mathcal{F}$.*

Proof. By Theorem 2(3c), a sum of form $\sum_{i \in I} \sqcup_{j \in J_i} F_{ij}$ is isomorphic to the sum $\sum_{(i,j) \in \sum_{k \in I} (J_k, \emptyset)} F_{ij}$. It remains to observe that if $\mathbf{l} = (I, <) \in \text{NPO}$ and $(J_i)_{i \in I}$ is a family of non-empty sets, then $\sum_{\mathbf{l}} (J_i, \emptyset) \in \text{NPO}$. \square

In view of Proposition 5, from (20) we obtain

Corollary 35. *Let \mathcal{F} be a class of A-frames, $s < \omega$, and $\mathbf{a} = (a_0, \dots, a_{s-1}) \in A^s$. Then for every A-formula φ we have:*

$$\varphi \text{ is satisfiable in } \sum_{\text{NPO}}^a \mathcal{F} \text{ iff } \varphi \text{ is satisfiable in } \sum_{\text{Tr}(\#\varphi, \#\varphi)}^a \mathcal{F}.$$

Remark 36. The operation of iterated sum can result in very tangled structures. In this remark, we expand the formal definition of this operation.

Let $0 \leq s < \omega$. Consider a tree $(T, <)$ such that every maximal chain in T has $s + 1$ elements. Let I be the set of maximal elements of $(T, <)$, $J = T \setminus I$, and for $i \in J$, let $\text{sc}(i)$ be the set of immediate successors of i . A structure $\mathbf{T} = (T, <, (S_i)_{i \in J})$ such that S_i is a binary relation on $\text{sc}(i)$ is called an *indexing tree*. If also $(\text{sc}(i), S_i) \in \mathcal{I}$ for every $i \in J$, then \mathbf{T} is an *indexing tree for \mathcal{I}* . For a sequence $\mathbf{a} = (a_0, \dots, a_{s-1}) \in A^s$, and a family $(F_i)_{i \in I}$ of A-frames, we define the \mathbf{a} -sum of $(F_i)_{i \in I}$ over the indexing tree \mathbf{T} , in symbols $\sum_{\mathbf{T}}^{\mathbf{a}} F_i$, as the following A-frame $(W, (R_a)_{a \in A})$. The domain of this structure is the disjoint union of the domains of F_i :

$$W = \{(i, w) \mid i \text{ is maximal in } \mathbf{T} \text{ and } w \text{ is in } F_i\}$$

To define the relations R_a , consider $(i, w), (j, v) \in W$. If $i = j$, then, as usual, we put $(i, w)R_a(j, v)$ iff $wR_{i,a}v$, where $R_{i,a}$ denotes the a -th relation in F_i . Assume that $i \neq j$. Let k be the infimum $\inf\{i, j\}$, and let h be the height of k in \mathbf{T} , that is the number of elements in $\{l \in T \mid l < k\}$. There are unique immediate successors i' and j' of k such that $i' \leq i$ and $j' \leq j$. Hence $0 \leq h < s$, $i' \neq j'$, and $i', j' \in \text{sc}(k)$. We put $(i, w)R_a(j, v)$ iff $i'S_k j'$ and a is the h -th element of \mathbf{a} .

The class $\sum_{\mathcal{I}}^* \mathcal{F}$ of *iterated sums of frames in \mathcal{F} over frames in \mathcal{I}* consists of such structures where \mathbf{T} is an indexing tree for \mathcal{I} and all F_i are in \mathcal{F} . One can see that for a fixed \mathbf{a} , the elements of $\sum_{\mathcal{I}}^* \mathcal{F}$ are (up to isomorphisms) \mathbf{a} -iterated sums $\sum_{\mathcal{I}}^{\mathbf{a}} \mathcal{F}$.

5.2 Lexicographic sums

The sum operation does not change the signature. In many cases (see below) it is convenient to characterize a polymodal logic via the following modification of \mathbf{a} -sums.

Definition 6. Let $\mathbf{l} = (I, S)$ be a unimodal frame, $(F_i)_{i \in I}$ a family of A-frames, and $F_i = (W_i, (R_{i,a})_{a \in A})$. The *lexicographic sum* $\sum_{\mathbf{l}}^{\text{lex}} F_i$ is the $(1 + A)$ -frame $(\sqcup_{i \in I} W_i, S^{\text{lex}}, (R_a)_{a \in A})$, where

$$\begin{aligned} (i, w)S^{\text{lex}}(j, u) & \quad \text{iff} \quad iSj, \\ (i, w)R_a(j, u) & \quad \text{iff} \quad i = j \ \& \ wR_{i,a}u. \end{aligned}$$

For a class \mathcal{F} of A-frames and a class \mathcal{I} of 1-frames, we define $\sum_{\mathcal{I}}^{\text{lex}} \mathcal{F}$ as the class of all sums $\sum_{\mathbf{l}}^{\text{lex}} F_i$, where $\mathbf{l} \in \mathcal{I}$ and all F_i are in \mathcal{F} .

Notice that lexicographic sums depend on reflexivity of the indexing frame (unlike sums considered in previous sections, see Remark 1.) This difference is explained in Propositions 37 and 38 below.

Recall that for an A -frame $F = (W, (R_i)_{i \in A})$, $F^{(\forall)}$ is the $(1+A)$ -frame $(W, W \times W, (R_i)_{i \in A})$. Let $F^{(\emptyset)}$ be the $(1+A)$ -frame $(W, \emptyset, (R_a)_{a < A})$; for a class \mathcal{F} of frames, let $\mathcal{F}^{(\emptyset)} = \{F^{(\emptyset)} \mid F \in \mathcal{F}\}$.

The following is immediate from the definitions:

Proposition 37. *For a unimodal frame $I = (I, S)$ and a family $(F_i)_{i \in I}$ of A -frames,*

$$\sum_I^{\text{lex}} F_i = \sum_{I'} F'_i = \sum_I^0 F'_i,$$

where I' is the $(1+A)$ -frame $(I, S, (\emptyset)_A)$, and for $i \in I$, $F'_i = F_i^{(\forall)}$ whenever i is reflexive in I , and $F'_i = F_i^{(\emptyset)}$ otherwise.

Proposition 38. *Let \mathcal{F} be a class of A -frames.*

1. *If \mathcal{I} is a class of irreflexive 1-frames, then $\sum_{\mathcal{I}}^{\text{lex}} \mathcal{F} = \sum_{\mathcal{I}}^0 \mathcal{F}^{(\emptyset)}$.*
2. *If \mathcal{I} is a class of reflexive 1-frames, then $\sum_{\mathcal{I}}^{\text{lex}} \mathcal{F} = \sum_{\mathcal{I}}^0 \mathcal{F}^{(\forall)}$.*

Proof. Immediate from Proposition 37. □

Given a class \mathcal{F} of frames in an alphabet A and a class of unimodal frames \mathcal{I} , let 0 -iterated lexicographic sums be elements of \mathcal{F} , and for $n < \omega$, let $(n+1)$ -iterated lexicographic sums be lexicographic sums of n -iterated sums. The class of n -iterated lexicographic sums of frames in \mathcal{F} over frames in \mathcal{I} is denoted by $\sum_{\mathcal{I}}^{\text{lex}^n} \mathcal{F}$.

The next fact is the iterated version of the above proposition. For a class \mathcal{F} of A -frames and $n < \omega$, let $\mathcal{F}^{(\emptyset)^n}$ be the class of $(n+A)$ -frames $(W, (\emptyset)_n, (R_a)_{a \in A})$ such that $(W, (R_a)_{a \in A}) \in \mathcal{F}$. Likewise, let $\mathcal{F}^{(\forall)^n}$ be the class of frames $(W, (S_a)_{a \in n+A})$ such that $S_a = W \times W$ for $a < n$, and $(W, (S_{n+a})_{a \in A}) \in \mathcal{F}$.

Proposition 39. *Let $A \leq \omega$, $0 < n < \omega$, and let \mathcal{F} be a class of A -frames.*

1. *If \mathcal{I} is a class of irreflexive 1-frames, then $\sum_{\mathcal{I}}^{\text{lex}^n} \mathcal{F} = \sum_{\mathcal{I}}^{(0, \dots, n-1)} \mathcal{F}^{(\emptyset)^n}$.*
2. *If \mathcal{I} is class of reflexive 1-frames, then $\sum_{\mathcal{I}}^{\text{lex}^n} \mathcal{F} = \sum_{\mathcal{I}}^{(0, \dots, n-1)} \mathcal{F}^{(\forall)^n}$.*

Proof. Follows from Proposition 37 by a straightforward induction on n . □

Propositions 38 and 39 allow to expand our results on the finite model property and complexity to the logics of (iterated) lexicographic sums.

5.3 Japaridze's polymodal logic

In this section we show that Japaridze's polymodal provability logic GLP is decidable in PSpace. First, this theorem was proven in [Sha08]. Here we provide a version of the proof based on Theorem 17.

GLP is a normal modal logic in the language $ML(\omega)$. This system was introduced in [Jap86] and plays an important role in proof theory (see, e.g., [Bek04]). GLP is known to be Kripke incomplete, so we cannot directly apply our tools to analyze it. However, in

[Bek10], L. Beklemishev introduced a modal logic J, a Kripke complete approximation of GLP. Semantically, J is characterised as the logic of frames called *stratified*, or *hereditary partial orderings* ([Bek10, Section 3]). They are defined as follows.

Definition 7. For $A \leq \omega$, let S_A be $(\{0\}, (\emptyset)_A)$, a singleton A-frame with empty relations. For $A < \omega$, let $\mathcal{J}(A)$ be the class of A-iterated lexicographic sums

$$\sum_{\text{NPO}}^{\text{lex}^A} \{S_\omega\} = \underbrace{\sum_{\text{NPO}}^{\text{lex}} \cdots \sum_{\text{NPO}}^{\text{lex}}}_{A \text{ times}} \{S_\omega\}.$$

The class \mathcal{J} of *hereditary partial orderings* is the class $\bigcup_{A < \omega} \mathcal{J}(A)$. The *logic* J is defined as the logic of the class \mathcal{J} .

In [Bek10], it was shown that GLP is polynomial time many-to-one reducible to J: there exists a polynomial time computable $f : \text{ML}(\omega) \rightarrow \text{ML}(\omega)$ such that $\varphi \in \text{GLP}$ iff $f(\varphi) \in \text{J}$ (for an explicit description of f , see [BFDJ14, Lemma 3.4]).

Our aim is to show that J is in PSpace. For our purposes, it is more convenient to work with a -sums. Since $S_\omega^{(\emptyset)^A} = S_\omega$, from Proposition 39(1) we have:

Proposition 40. *For every $A < \omega$, $\mathcal{J}(A) = \sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{A-1} \{S_\omega\}$.*

Formally, for every $A < \omega$, each frame in $\mathcal{J}(A)$ has infinitely many relations. Next facts allow us to consider frames of finite signatures. Let

$$\hat{\mathcal{J}}(A) = \sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{A-1} \{S_A\}.$$

For an alphabet C, a C-frame $F = (W, (R_a)_{a \in C})$, and $B \leq C$, let $F^{\upharpoonright B}$ be the reduct $(W, (R_a)_{a \in B})$; for a class \mathcal{F} of C-frames, $\mathcal{F}^{\upharpoonright B} = \{F^{\upharpoonright B} \mid F \in \mathcal{F}\}$.

Proposition 41. *For every $A < \omega$, we have:*

1. $\mathcal{J}(A)^{\upharpoonright A} = \hat{\mathcal{J}}(A)$;
2. *If $A > 0$, then every frame in $\mathcal{J}^{\upharpoonright A}$ is isomorphic to a frame in $\hat{\mathcal{J}}(A)$.*

Proof. From Definition 1 it is immediate that for alphabets $B \leq C$ and classes of C-frames \mathcal{F}, \mathcal{I} ,

$$\left(\sum_{\mathcal{I}} \cdots \sum_{\mathcal{I}} \mathcal{F} \right)^{\upharpoonright B} = \sum_{\mathcal{I}^{\upharpoonright B}} \cdots \sum_{\mathcal{I}^{\upharpoonright B}} (\mathcal{F}^{\upharpoonright B}). \quad (21)$$

Clearly, $S_\omega^{\upharpoonright A}$ is S_A . Now we obtain (1) from Proposition 40 and Definition 2.

To prove (2), let us fix $B < \omega$ and show that every frame in $\mathcal{J}(B)^{\upharpoonright A}$ is isomorphic to a frame in $\hat{\mathcal{J}}(A)$.

First, observe that $\mathcal{J}(B+1)$ contains copies of all frames in $\mathcal{J}(B)$. Indeed, every frame in the class $\sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{B-1} \{S_\omega\}$ is isomorphic to a frame in $\sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{B-1} \sum_{\text{NPO}}^B \{S_\omega\}$, since $\sum_{\text{NPO}}^B \{S_\omega\}$ contains a copy of S_ω . Hence, if $B < A < \omega$, $\mathcal{J}(A)$ contains copies of all frames in $\mathcal{J}(B)$.

The case $A < B$ is more interesting. Consider the class $\mathcal{G} = \sum_{\text{NPO}}^{A-1} \cdots \sum_{\text{NPO}}^{B-1} \{S_\omega\}$. By (21) and Proposition 40, we have:

$$\mathcal{J}(B)^{\upharpoonright A} = \left(\sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{B-1} \{S_\omega\} \right)^{\upharpoonright A} = \sum_{\text{NPO}}^0 \cdots \sum_{\text{NPO}}^{A-2} \mathcal{G}^{\upharpoonright A}.$$

By (21) again, $\mathcal{G}^{\uparrow A}$ can be represented as sums of disjoint unions of singletons:

$$\left(\sum_{\text{NPO}}^{A-1} \cdots \sum_{\text{NPO}}^{B-1} \{S_\omega\} \right)^{\uparrow A} = \sum_{\text{NPO}}^{A-1} \underbrace{\bigsqcup \cdots \bigsqcup}_{B-A \text{ times}} \{S_A\},$$

since a sum over the structure with all relations empty is just a disjoint sum. By Lemma 34, every frame in $\mathcal{G}^{\uparrow A}$ is isomorphic to a frame in the class $\sum_{\text{NPO}}^{A-1} \{S_A\}$. It follows that $\hat{\mathcal{J}}(A)$ contains copies of all frames in $\mathcal{J}(B)^{\uparrow A}$. \square

Proposition 42. *For every $A < \omega$, $J \cap \text{ML}(A) = \text{Log}(\hat{\mathcal{J}}(A))$.*

Proof. By Proposition 41(1), $\hat{\mathcal{J}}(A) \subseteq \mathcal{J}^{\uparrow A}$, so if φ is satisfiable in $\hat{\mathcal{J}}(A)$, then it is satisfiable in \mathcal{J} . Conversely, if $\varphi \in \text{ML}(A)$ is satisfiable in \mathcal{J} , then it is satisfiable in $\mathcal{J}^{\uparrow A}$, and hence — in $\hat{\mathcal{J}}(A)$ by Proposition 41(2). \square

It is trivial that the satisfiability problem on the singleton $S_A^{(\vee)}$ is in PSpace (in fact, it is in NP). From Corollary 33 and Proposition 41(1), we obtain

Corollary 43. *For every $A < \omega$, $\text{Sat } \hat{\mathcal{J}}(A)^{(\vee)} \in \text{PSpace}$.*

Hence, every fragment of J with finitely many modalities is in PSpace. The above corollary does not directly imply that $\text{Sat } \mathcal{J}$ is in PSpace. But the latter fact is a corollary of Theorem 17 and the observations below.

Lemma 44. *Let $A < \omega$. An A -formula φ is satisfiable in J iff φ is satisfiable in $\sum_{\text{Tr}(\#\varphi, \#\varphi)}^{(0, \dots, A-1)} \{S_A\}$.*

Proof. By Proposition 42, φ is satisfiable in J iff it is satisfiable in $\hat{\mathcal{J}}(A)$. By Corollary 35, the latter is equivalent to the satisfiability of φ in $\sum_{\text{Tr}(\#\varphi, \#\varphi)}^{(0, \dots, A-1)} \{S_A\}$. \square

Consider a formula φ in the language ML_ω . Let $a_0 < \cdots < a_{N-1}$ be the increasing sequence of indices of all occurring in φ modalities. Put $N(\varphi) = N$. Let $\hat{\varphi}$ be the result of replacing a_b -modalities by b -modalities in φ . Note that

$$\hat{\varphi} \text{ is an } N(\varphi)\text{-formula, and } N(\varphi) < \#\varphi = \#\hat{\varphi}. \quad (22)$$

Lemma 45. *[BFDJ14, Lemma 3.5]. $\varphi \in J$ iff $\hat{\varphi} \in J$.*

Theorem 46. *J is in PSpace.*

Proof. We describe a decision procedure for the conditional satisfiability on iterated sums of trees. For positive h, b and $a \leq A < \omega$, we define sums $\mathcal{S}(h, b, a, A)$ as follows: if $a = A$, let $\mathcal{S}(h, b, a, A)$ denote $\{S_A\}$ for all h, b ; if $a < A$, let $\mathcal{S}(h, b, a, A)$ denote $\sum_{\text{Tr}(h, b)}^a \sum_{\text{Tr}(A, A)}^{(a+1, \dots, A-1)} \{S_A\}$. In particular, $\mathcal{S}(h, b, A-1, A)$ is $\sum_{\text{Tr}(h, b)}^{A-1} \{S_A\}$, and $\mathcal{S}(A, A, 0, A) = \sum_{\text{Tr}(A, A)}^{(0, \dots, A-1)} \{S_A\}$. Using Theorem 17, we describe the procedure CSatJ (Algorithm 2) that decides whether a given A -tie $(\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable $\mathcal{S}(h, b, a, A)$:

Lemma 47. *Let $a \leq A < \omega$, $(\varphi, \mathbf{v}, \mathbf{U})$ an A -tie, and $0 < h, b < \omega$. Then*

$$(\varphi, \mathbf{v}, \mathbf{U}) \text{ is satisfiable in } \mathcal{S}(h, b, a, A) \text{ iff CSatJ}(\varphi, \mathbf{v}, \mathbf{U}, h, b, a, A) \text{ returns true.}$$

ALGORITHM 2: Decision procedure for $\text{CSat } \mathcal{S}(h, b, a, A)$

$\text{CSatJ}(\varphi, \mathbf{v}, \mathbf{U}, h, b, a, A) : \text{boolean}$

Input: An A -tie $(\varphi, \mathbf{v}, \mathbf{U})$; positive integers h, b ; $a \leq A < \omega$.
if $a = A$ **then return** $((\varphi, \mathbf{v}, \mathbf{U})$ is satisfiable in \mathcal{S}_A);
if $\text{CSatJ}(\varphi, \mathbf{v}, \mathbf{U}, A, A, a + 1, A)$ **then return** true;
if $h > 1$ **then**
 for k *such that* $1 \leq k \leq b$ **do for** $\mathbf{u}, \mathbf{v}_0, \dots, \mathbf{v}_{k-1} \in 2^{\#\varphi}$ *such that* $\mathbf{v} = \mathbf{u} + \sum_{i < k} \mathbf{v}_i$ **do**
 if $\text{CSatJ}(\varphi, \mathbf{u}, \mathbf{U} + \sum_{i < k} \mathbf{v}_i, A, A, a + 1, A)$ **then**
 if $\bigwedge_{i < k} \text{CSatJ}(\varphi, \mathbf{v}_i, \mathbf{U}, h - 1, b, a, A)$ **then return** true;
return false.

Proof. By induction on $A - a$. The case $A = a$ is trivial: $\mathcal{S}(h, b, A, A)$ consists of a single singleton \mathcal{S}_A . If $a < A$, then $\mathcal{S}(h, b, a, A)$ is $\sum_{\text{Tr}(h, b)}^a \mathcal{S}(A, A, a + 1, A)$. By induction hypothesis, $\text{CSatJ}(\tau, A, A, a + 1, A)$ decides whether a tie τ is satisfiable in $\mathcal{S}(A, A, a + 1, A)$. Observe that for $a < A$, the algorithm CSatJ is an instance of the algorithm $\text{CSatSum}_{\mathcal{F}}$ (Algorithm 1 in Section 4), where $\text{CSatJ}(\tau, A, A, a + 1, A)$ is the oracle for the summands $\sum_{\text{Tr}(A, A)}^{(a+1, \dots, A-1)} \{\mathcal{S}_A\}$. Now the induction step follows from Theorem 17. \square

Lemma 48. φ is satisfiable in \mathcal{J} iff $\hat{\varphi}$ is satisfiable in $\mathcal{S}(\#\varphi, \#\varphi, 0, \#\varphi)$.

Proof. We have from Lemmas 45 and 44: φ is satisfiable in \mathcal{J} iff $\hat{\varphi}$ is satisfiable in \mathcal{J} iff $\hat{\varphi}$ is satisfiable in $\sum_{\text{Tr}(A, A)}^{(0, \dots, A-1)} \{\mathcal{S}_A\}$, where A can be assumed equal to $\#\varphi$ by (22). The latter class is $\mathcal{S}(A, A, 0, A)$. \square

Hence, by Proposition 13, we obtain

$$\varphi \text{ is satisfiable in } \mathcal{J} \text{ iff } \exists \mathbf{v} \in 2^{\#\varphi} (\mathbf{v}(0) = 1 \ \& \ (\hat{\varphi}, \mathbf{v}, \mathbf{0}) \in \text{CSat } \mathcal{S}(\#\varphi, \#\varphi, 0, \#\varphi)),$$

where $\mathbf{0}$ represents the empty condition.

Let us estimate the amount of space used by CSatJ on the input $(\hat{\varphi}, \mathbf{v}, \mathbf{0}, \#\varphi, \#\varphi, 0, \#\varphi)$. On each recursive call, either the argument a increases by 1, or a does not change and the parameter h decreases by 1. Since $1 \leq h \leq \#\varphi$ and $0 \leq a \leq \#\varphi$, we obtain that the depth of recursion is bounded by $\#\varphi \cdot (\#\varphi + 1)$. For $n = \#\varphi$, at each call CSatJ needs $O(n^2)$ space to store new variables. Hence, to check the satisfiability of φ in \mathcal{J} we need $O(n^4)$ space. This completes the proof of the theorem. \square

Corollary 49. *Japaridze's polymodal provability logic GLP is decidable in PSpace.*

Remark 50. It is immediate that GLP and \mathcal{J} are PSpace-hard (e.g., it follows from the fact that the 1-modal fragments of these logics are the logic GL). From [CR03], it follows that one-variable fragment of GLP is PSpace-hard. In [Pak14], it was shown that even the constant (closed) fragment of GLP is PSpace-hard.

5.4 Refinements and lexicographic products of modal logics

In this paragraph we discuss algorithmic properties of logics obtained via the *refinement* and the *lexicographic product* operations.

The operation of *refinement of modal logics* was introduced in [BR10].

Definition 8. Let $F = (W, R)$ be a preorder and $\text{sk}F = (\overline{W}, \leq)$ its skeleton. Consider a family $(F_C)_{C \in \overline{W}}$ of A-frames such that $\text{dom}(F_C) = C$ for all $C \in \overline{W}$. The *refinement* of F by $(F_C)_{C \in \overline{W}}$ is the $(1 + A)$ -frame $(W, R, (R_a^\triangleright)_{a \in A})$, where

$$R_a^\triangleright \subseteq \bigcup_{C \in \overline{W}} C \times C \quad \text{for all } a \in A, \quad (23)$$

$$(W, (R_a^\triangleright)_{a \in A}) \upharpoonright C = F_C \quad \text{for all } C \in \overline{W}. \quad (24)$$

For a class \mathcal{I} of preorders and a class \mathcal{G} of A-frames let $\text{Ref}(\mathcal{I}, \mathcal{F})$ be the class of all refinements of frames from \mathcal{I} by frames in \mathcal{F} . For logics $L_1 \supseteq \text{S4}, L_2$, we put $\text{Ref}(L_1, L_2) = \text{Log Ref}(\text{Fr } L_1, \text{Fr } L_2)$.

In [BR10] it was shown that in many cases the refinement operation preserves the finite model property and decidability.

Refinements can be considered as sums according to the following fact:

Proposition 51. *The refinement of F by frames $(F_C)_{C \in \text{sk}F}$ is isomorphic to the sum $\sum_{C \in \text{sk}F}^0 F_C^{(\vee)}$.*

Proof. The isomorphism is given by $w \mapsto (C, w)$, where $w \in C$. \square

In view of Theorems 3 and 10, this observation provides another way to prove the finite model property of refinements (see [Sha18, Section 5.2] for more details). Moreover, representation of refinements as sums allows to obtain complexity results according to our theorems in Section 4.

Let us illustrate this with the logic $\text{Ref}(\text{S4}, \text{S4})$. In [BR10], it was proven that $\text{Ref}(\text{S4}, \text{S4})$ is decidable and that $\text{Ref}(\text{S4}, \text{S4}) = \text{Log Ref}(\text{QO}_f, \text{QO}_f)$, where QO_f is the class of all finite non-empty preorders. The latter fact in combination with above proposition yields

$$\text{Ref}(\text{S4}, \text{S4}) = \text{Log} \sum_{\text{Tr}_f}^0 \text{QO}_f^{(\vee)}.$$

Since the satisfiability problem for the class $\text{QO}_f^{(\vee)}$ is in PSpace ([Hem96]; see also Example 3), from Theorem 11 (and Corollary 29) we obtain

Corollary 52. $\text{Ref}(\text{S4}, \text{S4})$ is PSpace-complete.

A related operation is the *lexicographic product of modal logics* introduced in [Bal09] by Ph. Balbiani.

Definition 9. Consider frames $I = (I, S)$ and $F = (W, (R_a)_{a \in A})$. Their *lexicographic product* $I \bowtie F$ is the $(1 + A)$ -frame $(I \times W, S^\bowtie, (R_a^\bowtie)_{a \in A})$, where

$$\begin{aligned} (i, w) S^\bowtie (j, u) & \quad \text{iff} \quad i S j, \\ (i, w) R_a^\bowtie (j, u) & \quad \text{iff} \quad i = j \ \& \ w R_a u. \end{aligned}$$

In other words, $I \bowtie F$ is the lexicographic sum $\sum_{I \in \mathcal{I}}^{\text{lex}} F_i$, where $F_i = F$ for all i in I .

For a class \mathcal{I} of 1-frames and a class \mathcal{G} of A-frames, the class $\mathcal{I} \bowtie \mathcal{F}$ is the class of all products $I \bowtie F$ such that $I \in \mathcal{I}$ and $F \in \mathcal{F}$. For logics L_1, L_2 , we put $L_1 \bowtie L_2 = \text{Log}(\text{Fr } L_1 \bowtie \text{Fr } L_2)$.

As examples, consider the logics $\text{S4} \bowtie \text{S4}$ and $\text{GL} \bowtie \text{S4}$ and show that they are in PSpace (PSpace-complete). In [Sha18, Theorem 5.13], it was shown that $\text{S4} \bowtie \text{S4}$ is equal to $\text{Ref}(\text{S4}, \text{S4})$. Also, it was shown that

$$\text{GL} \bowtie \text{S4} = \text{Log}(\text{Tr}_f \bowtie \text{QO}_f) = \text{Log} \sum_{\text{Tr}_f}^0 \sum_{\text{Tr}_f}^1 C$$

where \mathcal{C} is the class of finite frames of form $(C, \emptyset, C \times C)$ (in terms of lexicographic sums, $\sum_{\text{Tr}_f}^0 \sum_{\text{Tr}_f}^1 \mathcal{C}$ is the class $\sum_{\text{Tr}_f}^{\text{lex}} \text{QO}_f$); see [Sha18, Theorem 5.14] for the proof. Hence, this logic is in PSpace by Corollary 33.

Corollary 53. *The lexicographic products $\text{S4} \bowtie \text{S4}$, $\text{GL} \bowtie \text{S4}$ are PSpace-complete.*

This result contrasts with the undecidability results for *modal products* of transitive logics [GKWZ05].

6 Conclusion

In many cases, sum-like operations preserve the finite model property and decidability [BR10], [Sha18]. In this paper we showed that transferring results can be obtained for the complexity of the modal satisfiability problems on sums. In particular, it follows that for many logics PSpace-completeness is immediate from their semantic characterizations.

Let us indicate some further results and directions.

- Sums and products with linear indices

In the linear case, *modal products* are typically (highly) undecidable [RZ01]. However, the modal satisfiability problem on the lexicographic squares of dense unbounded linear orders is in NP [BM13]. This positive result seems to be scalable according to the following observation: in many cases, φ is satisfiable in sums over linear (pre)orders iff φ is satisfiable in such sums that the length of indices is bounded by $\#\varphi$. In this situation there is a stronger than \leq_T^{PSpace} reduction between the sums and the summands.

- Further results on the finite model property and decidability of sums

In many cases, the finite model property of a modal logic L can be obtained by a filtration method; in this case we say that L *admits filtration*. It follows from [BR10] that filtrations of refinements can be reconstructed from filtrations of components. This result can be extended for a more general setting of lexicographic sums.

Also, the results obtained in [BR10] in a combination with Theorem 3 suggest the following conjecture: in the case of finitely many modalities, if $\text{Log } \mathcal{F}^{(\vee)}$ has the finite model property and $\text{Log } \mathcal{I}$ admits filtration, then the logic of $\sum_{\mathcal{I}} \mathcal{F}$ has the finite model property.

Another fact that we announce relates to the property of local finiteness of logics: if both the logics of indices and summands are locally finite, then the logic of lexicographic sums (a fortiori, of lexicographic products) is locally finite.

- Axiomatization of sums

For unimodal logics L_1, L_2 , let $\sum_{L_1}^{\text{lex}} L_2$ be the logic of the class $\sum_{\text{Fr } L_1}^{\text{lex}} \text{Fr } L_2$. Consider the following 2-modal formulas:

$$\alpha = \Diamond_1 \Diamond_0 p \rightarrow \Diamond_0 p, \quad \beta = \Diamond_0 \Diamond_1 p \rightarrow \Diamond_0 p, \quad \gamma = \Diamond_0 p \rightarrow \Box_1 \Diamond_0 p$$

One can see that these formulas are valid in every lexicographic sum $\sum_i^{\text{lex}} F_i$ (and hence, in every product $I \bowtie F$) of 1-frames. In many cases, these axioms provide a complete axiomatization of $\sum_{L_1}^{\text{lex}} L_2$. In particular, the logic $\sum_{\text{GL}}^{\text{lex}} \text{GL} = \text{Log}(\sum_{\text{NPO}}^{\text{lex}} \text{NPO})$, the bimodal fragment of the logic J considered in Section 5.3, is the logic $\text{GL} * \text{GL} + \{\alpha, \beta, \gamma\}$ [Bek10] ($L_1 * L_2$ denotes the *fusion* of L_1 and L_2 , $L + \Psi$ is for the least logic containing $L \cup \Psi$). Analogous

results hold for various lexicographic products [Bal09]; e.g., $S4 \succ S4 = S4 * S4 + \{\alpha, \beta, \gamma\}$. We announce the following results:

1. If $L_1 * L_2 + \{\alpha, \beta, \gamma\}$ is Kripke complete, and the class $\text{Fr } L_1$ (considered as a class of models in the classical model-theoretic sense) is first-order definable without equality, then

$$\sum_{L_1}^{\text{lex}} L_2 = L_1 * L_2 + \{\alpha, \beta, \gamma\}.$$

2. Assume that $\text{Fr } L_2$ is closed under direct products and validates the property $\forall x \exists y x R y$, and that the class $\text{Fr } L_1$ is first-order definable without equality. If the logic $L_1 * L_2 + \{\alpha, \beta, \gamma\}$ is Kripke complete, then

$$L_1 \succ L_2 = L_1 * L_2 + \{\alpha, \beta, \gamma\}.$$

At the same time, no general axiomatization results are known for the logics of sums in the sense of Definition 1.

- Sum-based operations in the Kripke-incomplete case

The operations we considered so far lead to Kripke complete logics. What could be definition of sums for modal algebras (models, general Kripke frames)? E.g., can we give a semantic characterization of an important Kripke-incomplete logic GLP by sums of form $\sum_{\text{NPO}} \dots \sum_{\text{NPO}} \mathcal{C}$?

7 Acknowledgements

I am sincerely grateful to the reviewers for multiple important suggestions on the earlier version of the manuscript.

References

- [Bal09] Philippe Balbiani. Axiomatization and completeness of lexicographic products of modal logics. In Silvio Ghilardi and Roberto Sebastiani, editors, *Frontiers of Combining Systems*, volume 5749 of *Lecture Notes in Computer Science*, pages 165–180. Springer, 2009.
- [Bal10] P. Balbiani. Axiomatizing the temporal logic defined over the class of all lexicographic products of dense linear orders without endpoints. In *2010 17th International Symposium on Temporal Representation and Reasoning*, pages 19–26, Sept 2010.
- [BdRV02] Patrick Blackburn, Maarten de Rijke, and Yde Venema. *Modal Logic*, volume 53 of *Cambridge Tracts in Theoretical Computer Science*. Cambridge University Press, 2002.
- [BEG11] Guram Bezhanishvili, Leo Esakia, and David Gabelaia. Spectral and t_0 -spaces in d -semantics. In Nick Bezhanishvili, Sebastian Löbner, Kerstin Schwabe, and Luca Spada, editors, *Logic, Language, and Computation*, pages 16–29, Berlin, Heidelberg, 2011. Springer Berlin Heidelberg.
- [Bek04] Lev D. Beklemishev. Provability algebras and proof-theoretic ordinals, i. *Annals of Pure and Applied Logic*, 128(1):103 – 123, 2004.

- [Bek10] Lev D. Beklemishev. Kripke semantics for provability logic GLP. *Annals of Pure and Applied Logic*, 161(6):756–774, 2010. The proceedings of the IPM 2007 Logic Conference.
- [BFD16] Philippe Balbiani and David Fernández-Duque. Axiomatizing the lexicographic products of modal logics with linear temporal logic. In Lev Beklemishev, Stéphane Demri, and András Máté, editors, *Advances in modal logic, vol. 11*, pages 78–96. College Publications, 2016.
- [BFDJ14] Lev D. Beklemishev, David Fernández-Duque, and Joost J. Joosten. On provability logics with linearly ordered modalities. *Studia Logica: An International Journal for Symbolic Logic*, 102(3):541–566, 2014.
- [BM13] Philippe Balbiani and Szabolcs Mikulás. Decidability and complexity via mosaics of the temporal logic of the lexicographic products of unbounded dense linear orders. In Pascal Fontaine, Christophe Ringeissen, and Renate A. Schmidt, editors, *Frontiers of Combining Systems*, pages 151–164, Berlin, Heidelberg, 2013. Springer Berlin Heidelberg.
- [BR10] Sergey Babenyshev and Vladimir Rybakov. Logics of Kripke meta-models. *Logic Journal of the IGPL*, 18(6):823–836, 2010.
- [CR03] A. V. Chagrov and M. N. Rybakov. How many variables does one need to prove pspace-hardness of modal logics. In Philippe Balbiani, Nobu-Yuki Suzuki, Frank Wolter, and Michael Zakharyashev, editors, *Advances in Modal Logic, Volume 4*, pages 71–82. CSLI Publications, 2003.
- [CZ97] Alexander Chagrov and Michael Zakharyashev. *Modal Logic*, volume 35 of *Oxford Logic Guides*. Oxford University Press, 1997.
- [Esa01] Leo Esakia. Weak transitivity-restitution. *Logical Studies*, 8:244–255, 2001.
- [FV59] S. Feferman and R. Vaught. The first order properties of products of algebraic systems. *Fundamenta Mathematicae*, 47(1):57–103, 1959.
- [GDJ74] D. M. Gabbay and D. H. J. De Jongh. A sequence of decidable finitely axiomatizable intermediate logics with the disjunction property. *Journal of Symbolic Logic*, 39(1):67–78, 1974.
- [GKWZ03] D. Gabbay, A. Kurucz, F. Wolter, and M. Zakharyashev. *Many-dimensional Modal Logics: Theory and Applications*. Studies in logic and the foundations of mathematics. North Holland Publishing Company, 2003.
- [GKWZ05] David Gabelaia, Agi Kurucz, Frank Wolter, and Michael Zakharyashev. Products of ‘transitive’ modal logics. *J. Symbolic Logic*, 70(3):993–1021, 09 2005.
- [Gur79] Yuri Gurevich. Modest theory of short chains. I. *J. Symb. Log.*, 44(4):481–490, 1979.
- [Gur85] Y. Gurevich. Monadic second-order theories. In J. Barwise and S. Feferman, editors, *Model-Theoretic Logics*, volume Volume 8 of *Perspectives in Mathematical Logic*, chapter XIII, pages 479–506. Springer-Verlag, New York, 1985.
- [Hem96] Edith Hemaspaandra. The price of universality. *Notre Dame J. Formal Logic*, 37(2):174–203, 04 1996.

- [Jap86] Giorgi K Japaridze. *The modal logical means of investigation of provability*. PhD thesis, Thesis in Philosophy, in Russian, Moscow, 1986.
- [Lad77] R. Ladner. The computational complexity of provability in systems of modal propositional logic. *SIAM Journal on Computing*, 6(3):467–480, 1977.
- [Lit07] Tadeusz Litak. The non-reflexive counterpart of Grz. *Bulletin of the Section of Logic*, 36, 01 2007.
- [Mos52] Andrzej Mostowski. On direct products of theories. *The Journal of Symbolic Logic*, 17(1):1–31, 1952.
- [Pak14] Fedor Pakhomov. On the complexity of the closed fragment of Japaridze’s provability logic. *Arch. Math. Log.*, 53(7-8):949–967, 2014.
- [RZ01] Mark Reynolds and Michael Zakharyashev. On the Products of Linear Modal Logics. *Journal of Logic and Computation*, 11(6):909–931, 2001.
- [SG77] Istvan Simon and John Gill. Polynomial reducibilities and upward diagonalizations. In *Proceedings of the Ninth Annual ACM Symposium on Theory of Computing*, STOC ’77, pages 186–194, New York, NY, USA, 1977. ACM.
- [Sha05] I. Shapirovsky. On PSPACE-decidability in transitive modal logic. In *Advances in Modal Logic*, volume 5, pages 269–287, London, 2005. College Publications. ISBN 1904987222.
- [Sha08] I. Shapirovsky. PSpace-decidability of Japaridze’s polymodal logic. In *Advances in Modal Logic*, volume 7 of *AiML*, pages 289–304, London, 2008. College Publications.
- [Sha18] I.B. Shapirovsky. Truth-preserving operations on sums of Kripke frames. In *Advances in Modal Logic*, volume 12, pages 541–558. College Publications, 2018. ISBN 978-1848902558.
- [She75] Saharon Shelah. The monadic theory of order. *Annals of Mathematics*, 102(3):379–419, 1975.
- [Spa93] E. Spaan. *Complexity of Modal Logics, PhD thesis*. University of Amsterdam, Institute for Logic, Language and Computation, 1993.