

A simple approach to proving the existence, uniqueness, and strong and weak convergence rates for a broad class of McKean–Vlasov equations

Abdul-Lateef Haji-Ali* Håkon Hoel† Raúl Tempone†‡

November 10, 2021

Abstract

By employing a system of interacting stochastic particles as an approximation of the McKean–Vlasov equation and utilizing classical stochastic analysis tools, namely Itô’s formula and Kolmogorov–Chentsov continuity theorem, we prove the existence and uniqueness of strong solutions for a broad class of McKean–Vlasov equations. Considering an increasing number of particles in the approximating stochastic particle system, we also prove the L^p strong convergence rate and derive the weak convergence rates using the Kolmogorov backward equation and variations of the stochastic particle system. Our convergence rates were verified by numerical experiments which also indicate that the assumptions made here and in the literature can be relaxed.

MSC 2010 subject classifications: Primary 65C05, 62P05

Keywords and phrases: Interacting stochastic particle systems, Stochastic mean-field limit, Weak and strong convergence rates.

1 Introduction

For $a, \sigma, \kappa_1, \kappa_2 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$, $t \geq 0$ and $\{W(s)\}_{s>0}$ being a one-dimensional Wiener process defined on some probability space with the natural filtration, we consider the following McKean–Vlasov equation:

$$\begin{aligned} Z(t) = \xi &+ \int_0^t a\left(Z(s), \int_{\mathbb{R}} \kappa_1(Z(s), z) \mu_s(dz)\right) ds \\ &+ \int_0^t \sigma\left(Z(s), \int_{\mathbb{R}} \kappa_2(Z(s), z) \mu_s(dz)\right) dW(s), \end{aligned} \quad (1)$$

where μ_s denotes the law of $Z(s)$ for all $s \geq 0$, and ξ denotes a random initial state whose law is μ_0 . For $a(x, y) = y$, with σ being constant and κ_1 being bounded and Lipschitz, [23, Theorem I.1.1] demonstrates the existence and uniqueness of a strong solution to (1). A recent analysis yielded the same result in [18] when the initial condition ξ has a finite fourth moment, $a(x, y) = \sigma(x, y) = y$, κ_2 does not change sign, and for all $x, x', y \in \mathbb{R}$, there exists a constant $C > 0$ such that

$$\begin{aligned} |\kappa_1(x, y)| + |\kappa_2(x, y)| &\leq C(1 + |x|), \\ |\kappa_2(x, y) - \kappa_2(x', y)| &\leq C(1 + |y|^2) |x - x'|. \end{aligned}$$

A typical method for obtaining strong approximations of Z is to solve the following d -dimensional system of Itô stochastic differential equations (SDEs), also known as an interacting stochastic

*a.hajiali@hw.ac.uk, Heriot-Watt University, Edinburgh, UK.

†{hoel,tempone}@uq.rwth-aachen.de, RWTH Aachen University, Germany.

‡raul.tempone@kaust.edu.sa, King Abdullah University of Science & Technology (KAUST), Thuwal, Saudi Arabia.

particle system with pairwise interaction kernels

$$\begin{aligned}
X_i^d(t) = \xi_i &+ \int_0^t a \left(X_i^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_i^d(s), X_j^d(s)) \right) ds \\
&+ \int_0^t \sigma \left(X_i^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_2(X_i^d(s), X_j^d(s)) \right) dW_i(s)
\end{aligned} \tag{2}$$

for $i \in \{1, 2, \dots, d\}$, ξ_i are i.i.d. and have the same law, μ_0 , and $\{W_i(s)\}_{s>0}$ are independent one-dimensional Wiener processes and independent of $\{\xi_i\}_{i=1}^d$. In other words, the law μ_t for $t \geq 0$ is approximated by an empirical measure based on the particles $\mathbf{X}^d := \{X_i^d(t)\}_{i=1}^d$. It should be noted that these particles are identically distributed but not independent.

The rate of strong convergence of X_i^d to a corresponding Z_i , which satisfies (1) with ξ_i and W_i , as the size of the particle system, d , increases when $a(x, y) = y$ and σ is a constant was shown to be $\mathcal{O}(1/\sqrt{d})$ in the classical reference [23, Theorem I.1.4]. See also the works [5, 6] and others by the same authors. For $\mathbf{Z} := \{Z_j^d\}_{j=1}^d$ being d independent samples of the solution to the McKean–Vlasov equation (1), we define the rate of weak convergence of as the rate of convergence of $\mathbb{E}[g(\mathbf{X}^d)]$ to $\mathbb{E}[g(\mathbf{Z})]$ for a certain class of functions, $g: \mathbb{R}^d \rightarrow \mathbb{R}$. The rate of weak convergence was established to be $\mathcal{O}(1/d)$ in, e.g., [16, Chapter 9] and [17, Theorem 6.1]. These works assume that $\kappa_2 \equiv 0$ and build upon semigroup theory in measure-valued function spaces to prove their results. Recently, there has been an increasing interest in extending proofs of strong and weak convergence in more general settings with nonlinear drift/diffusion coefficients. To that end, works such as [24, Theorem B.2] and [10, Theorem 2.17] use Lions-derivatives [8] to calculate derivatives with respect to measures and a master equation for probability measures [9]. For a more exhaustive literature review, see [10].

Our study proves the existence and uniqueness of solutions to (1) and derives, we believe more simply, the same rates of strong and weak convergence of $X_i^d \rightarrow Z_i$ as in the literature using the stochastic particle-based approximation (2) and classical stochastic analysis tools. In particular, we employ Itô’s formula, the classical Kolmogorov–Chentsov continuity theorem, the Kolmogorov backward equation and the first, second, and third variations of the SDE (2). [11]. We believe that this new analysis is simpler than the analysis in the literature and is fully presented in this manuscript. Moreover, it leads to clearer conditions on the class of admissible drift and diffusion coefficients. In some cases, such conditions are relaxed (cf. the main condition on strong existence (4) and those in [18] and the conditions on the $\mathcal{O}(1/d)$ weak convergence rate in Theorem 3.1 and [10, Theorem 2.17]). We emphasize that the tools and proofs we present here can be extended to more general settings with vector-valued $Z(t) \in \mathbb{R}^n$ for some integer $n > 1$ or time heterogeneous coefficients. However, we restrict ourselves to the above setting for a more transparent exposition.

In Section 2, we prove the existence and uniqueness of strong solutions to (1) by first bounding moments and strong errors of (2) as functions of the particle system d . We then prove that the conditional expectation of particles in (2) converges to a limit satisfying the McKean–Vlasov equation (1). We also establish the standard strong convergence rates in this section.

In Section 3, we derive the rate of weak convergence. The principal steps to achieve this result use the Kolmogorov backward equation to represent the weak error and the stochastic flows and the dual functions to bound the weights in the resulting dual weighted residual representation. Using the Kolmogorov backward equation to estimate the weak error in SDEs goes back to the ideas of Talay and Tubaro [25], who estimated the time discretization error for uniform deterministic time-steps. Later, the works [1, 2], extended the analysis to approximations with non-smooth observables and the probability density of the solution at a given time. Kloeden and Platen [15] generalized the analysis in [25] to weak approximations of a higher order. Later, in a series of works inspired by [25], the authors developed methods based on stochastic flows and dual functions to bound the weights in the resulting dual-weighted residual representation. This approach provided the analysis for the weak approximation of SDEs using nonuniform, possibly stochastic, time-steps, see [4, 19, 20, 22]. Furthermore, the same analysis line extended into the adaptive Multilevel Monte Carlo realm, [12, 13]. The closest inspiration for deriving the weak convergence rate goes back to the use of the mentioned techniques in the context of multiscale approximation. Those works derived macroscopic SDEs continuum models by choosing their drift and diffusion functions to

minimize the weak error in the given macroscopic observables when compared with a given base model. Particularly, [14] employed master equations with long-range interaction potentials as a base model as the stochastic Ising model with Glauber dynamics, whereas [26] determined the stochastic phase-field models from atomistic formulations by coarse-graining molecular dynamics to model the dendritic growth of a crystal in an under-cooled melt. Systems of coupled SDEs, with increasing size could be useful for approximating a non-Markovian behavior. For example, the recent work [3] investigated weak convergence rates for a rough stochastic volatility model emerging in mathematical finance, namely, the rough Bergomi model. As in this study, the analysis in [3] also employed a dual-weighted representation of the weak error, yielding an error expansion that characterizes the weak convergence rate.

Finally, we verify in Section 4 the strong and weak convergence rates that we obtained. In particular, we show numerically that the weak convergence rate is the same for an example stochastic particle system that does not satisfy regularity conditions of our theory or those of other theories in the literature cited above. Therefore, further work might be necessary to extend the current approaches. We then conclude the principal part of the manuscript and dedicate the appendix to prove the technical results and determine sufficient conditions to bound the derivatives of the value function for a generic multidimensional SDE. Given certain assumptions, we prove that the first, second, and third variations of an SDE have bounded moments. We then prove that the value function has bounded first, second and third derivatives based on these bounded moments.

In what follows, we will use the notation $A \lesssim B$ to denote that there is a constant $0 < c < \infty$ which is independent of d , the size of the particle system (2), such that $A \leq cB$. We will also denote the p -ary Cartesian power of a set \mathcal{I} by \mathcal{I}^p .

2 Strong Analysis

In this section, we prove the existence and uniqueness of a strong solution to (1) and establish the rates of strong convergence of X^d to such a solution. First observe that for any t , the particles $\{X_i^d(t)\}_{i=1}^d$ in (2) are *exchangeable*. Therefore, for any function $g : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\mathbb{E}[g(X_1^d(t), X_2^d(t), \dots, X_d^d(t))] = \mathbb{E}[g(X_{i_1}^d(t), X_{i_2}^d(t), \dots, X_{i_d}^d(t))]$$

for any permutation $\{i_1, i_2, \dots, i_d\}$ of $\{1, 2, \dots, d\}$. In particular, for any $i \in \{1, 2, \dots, d\}$ we have $\mathbb{E}[g(X_i^d(t))] = \mathbb{E}[g(X_1^d(t))]$. When such exchangeability is possible and the dependence on the index i is unimportant, we could write $\mathbb{E}[g(X^d)] := \mathbb{E}[g(X_i^d)]$. In the rest of the manuscript, for notational convenience, we will work under the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ where \mathcal{F} denotes the σ -algebra generated by $\{\xi_i, (W_i(s))_{0 \leq s} \}_{i \in \mathbb{N}}$. However, observe that our results rely only on probability spaces that involve finitely many Wiener processes and initial states. For some $p \geq 1$, we also consider the Lebesgue space

$$L^p(\Omega) = \{f : (\Omega, \mathcal{F}) \rightarrow (\mathbb{R}, \mathcal{B}(\mathbb{R})), \quad \text{where } f \text{ is measurable and } \mathbb{E}[|f|^p] < \infty\},$$

where $\mathcal{B}(\mathbb{R})$ denotes the Borel σ -algebra of \mathbb{R} , and the Bochner space

$$L^\infty([0, T], L^p(\Omega)) = \{f : [0, T] \rightarrow L^p(\Omega), \quad \text{where } f \text{ is Borel-measurable and } \operatorname{ess\,sup}_{0 \leq t \leq T} \mathbb{E}[|f(t, \cdot)|^p] < \infty\},$$

In this section, all lemmas and corollaries implicitly assume the following:

Assumption 2.1. *There exists a constant, $C > 0$, such that for all $x, y, z, w \in \mathbb{R}$,*

$$\begin{aligned} & |a(x, y) - a(w, z)| + |\sigma(x, y) - \sigma(w, z)| \\ & + |\kappa_1(x, y) - \kappa_1(w, z)| + |\kappa_2(x, y) - \kappa_2(w, z)| \leq C(|x - w| + |y - z|). \end{aligned} \quad (3)$$

and

$$|a(x, y)| + |\sigma(x, y)| + |\kappa_1(x, y)| + |\kappa_2(x, y)| \leq C(1 + |x| + |y|), \quad (4)$$

for all $x, y \in \mathbb{R}$. Note that (3) implies the linear growth condition (4) for sufficiently large C .

The existence and uniqueness of a strong solution to the system of particles (2) under the previous assumption are well established [21]. In what follows, for any integer i , let $\mathcal{F}_i(t)$ be the σ -algebra generated by $\{\xi_i, (W_i(s))_{0 \leq s \leq t}\}$ and define the $\mathcal{F}_i(t)$ -adapted process

$$Z_i^d(t) := \mathbb{E}[X_i^d(t) | \mathcal{F}_i(t)]. \quad (5)$$

Note that, given two independent Wiener processes W_i and W_j , the two corresponding processes, Z_i^d and Z_j^d , are similarly independent and have the same law by exchangeability.

Lemma 2.2 (Bounded moments of particles). *Let $\{X_i^d\}_{i=1}^d$ be the strong solution of (2); then, for any $p \geq 1$ and any $T > 0$, we have*

$$\sup_{0 \leq t \leq T} \mathbb{E}[|X^d(t)|^{2p}] \lesssim 1 + \mathbb{E}[|\xi|^{2p}]$$

where the hidden constant depends on p , T , and C only.

Proof. By Itô's formula and Tonelli's theorem

$$\begin{aligned} \mathbb{E}[|X_1^d(t)|^{2p}] &\leq \mathbb{E}[|\xi_1|^{2p}] + p \int_0^T \mathbb{E} \left[\left| a \left(X_1^d(t), \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s)) \right) \right| |X_1^d(s)|^{2p-1} \right] ds \\ &\quad + (2p-1) \int_0^T \mathbb{E} \left[\left| \sigma \left(X_1^d(t), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1(s), X_j(s)) \right) \right|^2 |X_1^d(s)|^{2p-2} \right] ds \end{aligned} \quad (6)$$

Using (4) and Young's inequality,

$$\begin{aligned} &\mathbb{E} \left[\left| a \left(X_1^d(t), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1(s), X_j(s)) \right) \right| |X_1^d(s)|^{2p-1} \right] \\ &\leq C \mathbb{E} \left[\left(1 + |X_1^d(s)| + C \left(1 + |X_1^d(s)| + \frac{1}{d} \sum_{j=1}^d |X_j(s)| \right) \right) |X_1^d(s)|^{2p-1} \right] \\ &\lesssim \mathbb{E}[|X_1^d(s)|^{2p-1}] + \mathbb{E}[|X_1^d(s)|^{2p}] + \frac{1}{d} \sum_{j=1}^d \mathbb{E}[|X_1^d(s)| |X_j^d(s)|^{2p-1}] \\ &\leq \frac{1}{2p} + \frac{2p-1}{2p} \mathbb{E}[|X_1^d(s)|^{2p}] + 3, \mathbb{E}[|X_1^d(s)|^{2p}] \end{aligned}$$

we used Hölder's and Young's inequalities and the fact that $\mathbb{E}[|X_1^d(s)|] = \mathbb{E}[|X_j^d(s)|]$ for all $j \in \{1, \dots, d\}$ by exchangeability. Similarly, we bound the second term in (6)

$$\mathbb{E} \left[\left| \sigma \left(X_1^d(t), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1(s), X_j(s)) \right) \right|^2 |X_1^d(s)|^{2p-2} \right] \lesssim \frac{1}{2p} + \frac{2p-1}{p} \mathbb{E}[|X_1^d(s)|^{2p}].$$

Substituting these bounds back and using Grönwall's inequality we arrive at the result. \square

Lemma 2.3 ($L^\infty([0, T], L^{2p}(\Omega))$ error of particle systems). *Let $\{X_i^d\}_{i=1}^d$ and $\{X_i^{2d}\}_{i=1}^{2d}$ be the strong solution of (2) with d and $2d$ particles, respectively. Then for any $T > 0$, any $i \in \{1, \dots, d\}$, and any integer $p \geq 1$, we have*

$$\sup_{0 \leq t \leq T} \mathbb{E}[|X_i^d(t) - X_i^{2d}(t)|^{2p}] \lesssim d^{-p} (1 + \mathbb{E}[|\xi|^{2p}]), \quad (7)$$

where the hidden constant depends only on p , T , and C .

Proof. By exchangeability, we consider $i = 1$ only. By Itô's formula, we can bound

$$\begin{aligned} \mathbb{E}[|X_1^d(t) - X_1^{2d}(t)|^{2p}] &\leq 2p \int_0^t \mathbb{E}[|\Delta a| |X_1^d(s) - X_1^{2d}(s)|^{2p-1}] ds \\ &\quad + p(2p-1) \int_0^t \mathbb{E}[(\Delta\sigma)^2 |X_1^d(s) - X_1^{2d}(s)|^{2p-2}] ds, \end{aligned} \quad (8)$$

where

$$\Delta a = a\left(X_1^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s))\right) - a\left(X_1^{2d}(s), \sum_{j=1}^{2d} \kappa_1(X_1^{2d}(s), X_j^{2d}(s))\right)$$

and $\Delta\sigma$ is defined similarly. Using Young's inequality yields

$$\mathbb{E}[|\Delta a| |X_1^d(s) - X_1^{2d}(s)|^{2p-1}] \leq \frac{1}{2p} \mathbb{E}[|\Delta a|^{2p}] + \frac{2p-1}{2p} \mathbb{E}[|X_1^d(s) - X_1^{2d}(s)|^{2p}]. \quad (9)$$

We then bound using (3)

$$|\Delta a| \leq C |X_1^d(s) - X_1^{2d}(s)| + C \left| \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s)) - \frac{1}{2d} \sum_{j=1}^{2d} \kappa_1(X_1^{2d}(s), X_j^{2d}(s)) \right|$$

Then, taking the $2p$ moment and using Jensen's inequality yield

$$\begin{aligned} \mathbb{E}[|\Delta a|^{2p}] &\leq 2^{2p-1} C^{2p} \mathbb{E}[|X_1^{2d}(s) - X_1^d(s)|^{2p}] \\ &\quad + 2^{2p-1} \left(\frac{C}{2}\right)^{2p} \mathbb{E}\left[\left(\frac{1}{d} \sum_{j=1}^d \Delta\kappa_j\right)^{2p}\right], \end{aligned} \quad (10)$$

where we denote

$$\Delta\kappa_j := \kappa_1(X_1^{2d}(s), X_j^{2d}(s)) - \kappa_1(X_1^d(s), X_{d+j}^{2d}(s)).$$

Here, for $\mathcal{I} = \{1, \dots, d\}$,

$$\mathbb{E}\left[\left(\frac{1}{d} \sum_{j=1}^d \Delta\kappa_j\right)^{2p}\right] = \frac{1}{d^{2p}} \sum_{\mathbf{k} \in \mathcal{I}^{2p}} \mathbb{E}\left[\prod_{j=1}^{2p} \Delta\kappa_{k_j}\right]. \quad (11)$$

Let $\mathcal{I}^{2p} = \mathcal{J}_{\mathcal{I}^{2p}} \cup \mathcal{U}_{1,1} \cup \mathcal{U}_{1,-1}$ with $\mathcal{J}_{\mathcal{I}^{2p}}$ is as defined in (55), i.e., the set of all multi-indices from \mathcal{I}^{2p} with no unique indices, $\mathcal{U}_{1,1}$ is the set of multi-indices with exactly one unique index equal to 1, and $\mathcal{U}_{1,-1}$ is the set of multi-indices with at least one unique index different from 1. We note that for $\mathbf{k} \in \mathcal{U}_{1,-1}$, i.e., there is an $\ell \in \mathcal{I} \setminus \{1\}$ such that $k_\ell \neq k_j$ for all $j \neq \ell$, we have

$$\mathbb{E}\left[\prod_{j=1}^{2p} \Delta\kappa_{k_j}\right] = \mathbb{E}\left[\Delta\kappa_{k_\ell} \times \prod_{j=1, j \neq \ell}^{2p} \Delta\kappa_{k_j}\right] = 0,$$

as the particles k_ℓ and $d + k_\ell$ inside $\Delta\kappa_{k_\ell}$ in the previous expectation are exchangeable. For other $\mathbf{k} \in \mathcal{J}_{\mathcal{I}^{2p}} \cup \mathcal{U}_{1,1}$, we bound using the Lipschitz condition (3) on κ_1 and Jensen's and Hölder's inequalities

$$\left| \mathbb{E}\left[\prod_{j=1}^{2p} \Delta\kappa_{k_j}\right] \right| \leq 2^{2p} C^{2p} \mathbb{E}[|X^{2d}(s)|^{2p}].$$

Finally noting, by Lemma A.1, that

$$\begin{aligned} |\mathcal{J}_{\mathcal{I}^{2p}}| &\leq c_{2p} d^p \\ |\mathcal{U}_{1,1}| = 2p \left| \mathcal{J}_{(\mathcal{I} \setminus \{1\})^{2p-1}} \right| &\leq c_{2p-1} (2p) (d-1)^{\lfloor (2p-1)/2 \rfloor} \leq 2 c_{2p-1} p d^p \end{aligned}$$

we conclude that

$$\left| \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta \kappa_j \right)^{2p} \right] \right| \lesssim d^{-p} \sup_{0 \leq t \leq T} \mathbb{E} \left[|X^{2d}(t)|^{2p} \right]. \quad (12)$$

We then substitute (12) into (10) and the result into (9) and then into (8). By similarly obtaining the corresponding bounds for the second term in (8) (involving $\Delta \sigma$), we arrive at

$$\begin{aligned} \mathbb{E} \left[|X_1^d(t) - X_1^{2d}(t)|^{2p} \right] &\lesssim \int_0^t \mathbb{E} \left[|X_1^d(s) - X_1^{2d}(s)|^{2p} \right] ds \\ &\quad + d^{-p} \sup_{0 \leq s \leq T} \mathbb{E} \left[|X^{2d}(s)|^{2p} \right]. \end{aligned} \quad (13)$$

Using Grönwall's inequality and Lemma 2.2, we arrive at the claimed result. \square

Corollary 2.4 ($L^\infty([0, T], L^{2p}(\Omega))$ distance of equally-sized particle systems). *Let $\{X_i^d\}_{i=1}^d$ be the strong solution of (2) with d particles, for a given $\{\xi_i, (W_i)_{t \geq 0}\}_{i=1}^d$. Define a copy of the system, (2), $\{\tilde{X}_i^d\}_{i=1}^d$, satisfying for all $i \in \{1, \dots, d\}$,*

$$\begin{aligned} \tilde{X}_i^d(t) = \tilde{\xi}_i &+ \int_0^t a \left(\tilde{X}_i^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(\tilde{X}_i^d(s), \tilde{X}_j^d(s)) \right) ds \\ &+ \int_0^t \sigma \left(\tilde{X}_i^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_2(\tilde{X}_i^d(s), \tilde{X}_j^d(s)) \right) d\tilde{W}_i(s), \end{aligned} \quad (14)$$

where $\{\tilde{\xi}_i\}_{i=1}^d$ are i.i.d. samples of ξ , and $\{(\tilde{W}_i(t))_{t \geq 0}\}_{i=1}^d$ are mutually independent Wiener processes and independent of $\{\tilde{\xi}_i\}_{i=1}^d$. Assume that for some $i \in \{1, \dots, d\}$, we have $\tilde{\xi}_i = \xi_i$ and, for all t , we have $\tilde{W}_i(t) = W_i(t)$ almost surely and $\{\tilde{\xi}_j, (\tilde{W}_j(t))_{t \geq 0}\}_{j=1}^d$ is independent of $\{\xi_j, (W_j(t))_{t \geq 0}\}_{j=1}^d$. Then, for any $T > 0$ and any integer $p \geq 1$, we have

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\left| \tilde{X}_i^d(t) - X_i^d(t) \right|^{2p} \right] \lesssim d^{-p} (1 + \mathbb{E}[|\xi|^{2p}]), \quad (15)$$

where the hidden constant depends only on p, T , and C .

Proof. By exchangeability, we only consider $i = 1$. We construct a particle system $\{X_i^{2d}\}_{i=1}^{2d}$, which satisfies (2) for $2d$ particles and where for $i \in \{2, \dots, d\}$ we set $\xi_{d+i} = \tilde{\xi}_i$ and $W_{d+i}(t) = \tilde{W}_i(t)$ for all t . Then, using Jensen's inequality

$$\mathbb{E} \left[|\tilde{X}_1^d(t) - X_1^d(t)|^{2p} \right] \lesssim 2^{2p-1} \mathbb{E} \left[|\tilde{X}_1^d(t) - X_1^{2d}(t)|^{2p} \right] + 2^{2p-1} \mathbb{E} \left[|X_1^d(t) - X_1^{2d}(t)|^{2p} \right],$$

we conclude using Lemma 2.3 and Lemma 2.2. \square

Lemma 2.5 ($L^\infty([0, T], L^{2p}(\Omega))$ error of conditional expectations). *For any integer $p \geq 1$, any $T > 0$ and any $i \in \{1, \dots, d\}$*

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[|Z_i^d(t) - Z_i^{2d}(t)|^{2p} \right] \lesssim d^{-p} (1 + \mathbb{E}[|\xi|^{2p}]), \quad (16)$$

where the hidden constant depends only on p, T and C .

Proof. The result follows by using Jensen inequality for conditional expectations in (15) and Lemma 2.2. \square

Lemma 2.6 (Moments of increments and pathwise Hölder regularity of conditional expectations). *For any $p \geq 1$ and $T, h \geq 0$ and any $i \in \{1, \dots, d\}$, we have*

$$\sup_{0 \leq t \leq T} \mathbb{E} \left[\left(Z_i^d(t) - Z_i^d(t+h) \right)^{2p} \right] \lesssim h^p (1 + \mathbb{E}[|\xi|^{2p}]), \quad (17)$$

where the hidden constant depends only on p, T and C . Furthermore, when $p > 1$ and $\mathbb{E}[|\xi|^{2p}]$ is finite, the process Z_i^d is a.s. γ -Hölder continuous in t , with $\gamma \in (0, 1/2 - 1/(2p))$.

Proof. By exchangeability, we only consider $i = 1$. Recall that $Z_1^d(t) = \mathbb{E}[X_1^d(t) | \mathcal{F}_1(t)]$; hence, we start by taking the conditional expectation of both sides of (2):

$$\begin{aligned} Z_1^d(t) &= \xi_1 + \mathbb{E}\left[\int_0^t a_1(s) ds \middle| \mathcal{F}_1(t)\right] + \mathbb{E}\left[\int_0^t \sigma_1(s) dW_1(s) \middle| \mathcal{F}_1(t)\right] \\ &= \xi_1 + \int_0^t \mathbb{E}[a_1(s) | \mathcal{F}_1(s)] ds + \int_0^t \mathbb{E}[\sigma_1(s) | \mathcal{F}_1(s)] dW_1(s), \end{aligned}$$

where

$$\begin{aligned} a_1(s) &= a\left(X_1(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1(s), X_j(s))\right) \\ \sigma_1(s) &= \sigma\left(X_1(s), \frac{1}{d} \sum_{j=1}^d \kappa_2(X_1(s), X_j(s))\right). \end{aligned}$$

Next, we bound

$$\begin{aligned} &\mathbb{E}\left[|Z_1^d(t+h) - Z_1^d(t)|^{2p}\right] \\ &\leq 2^{2p-1} \mathbb{E}\left[\left|\int_t^{t+h} \mathbb{E}[a_1(s) | \mathcal{F}_1(s)] ds\right|^{2p}\right] + 2^{2p-1} \mathbb{E}\left[\left|\int_t^{t+h} \mathbb{E}[\sigma_1(s) | \mathcal{F}_1(s)] dW_1(s)\right|^{2p}\right] \\ &\leq 2^{2p-1} h^{2p-1} \int_t^{t+h} \mathbb{E}[|a_1(s)|^{2p}] ds + 2^{2p-1} C_{\text{BDG}, 2p} \mathbb{E}\left[\left|\int_t^{t+h} (\mathbb{E}[\sigma_1(s) | \mathcal{F}_1(s)])^2 ds\right|^p\right] \\ &\leq 2^{2p-1} h^{2p-1} \int_t^{t+h} \mathbb{E}[|a_1(s)|^{2p}] ds + 2^{2p-1} C_{\text{BDG}, 2p} h^{p-1} \int_t^{t+h} \mathbb{E}[|\sigma_1(s)|^{2p}] ds, \end{aligned}$$

where we used Jensen's and Burkholder–Davis–Gundy's (BDG) inequalities ($C_{\text{BDG}, 2p}$ is the BDG constant [7]). Using (4) and Jensen's inequality, we can bound

$$\mathbb{E}[|a_1(s)|^{2p}] \leq 3^{2p-1}(1 + 3^{2p-1}) + 3^{2p-1}(2 + 3^{2p-1}) \sup_{0 \leq t \leq T} \mathbb{E}[|X_1(s)|^{2p}],$$

and similarly for $\mathbb{E}[|\sigma_1(s)|^{2p}]$. Substituting back and using Lemma 2.2 yield the claimed (17). Finally, when $p > 1$, we can use the Kolmogorov–Chentsov continuity theorem to conclude that there exists a modification of Z^d that is a.s. pathwise γ -Hölder continuous in t with $\gamma \in (0, 1/2 - 1/(2p))$. \square

Lemma 2.7 ($L^\infty([0, T], L^{2p}(\Omega))$ convergence of conditional expectations). *For an integer $p \geq 1$, assume that $\mathbb{E}[|\xi|^{2p}]$ is finite. Then, for any integer $i \leq 2^{k_0}$, the sequence $\{Z_i^{2^k}\}_{k=k_0}^\infty$ converges in $L^\infty([0, T], L^{2p}(\Omega))$ to the $\mathcal{F}_i(t)$ -adapted process Z_i^∞ . Furthermore, for any integers $i \neq j$, the processes Z_i^∞ and Z_j^∞ are independent and have the same law. Additionally, when $p > 1$, the process Z_i^∞ is γ -Hölder continuous in t , with $\gamma \in (0, 1/2 - 1/(2p))$.*

Proof. First, observe that Lemma 2.5 with $p = 2$ implies that $\{Z_i^{2^k}\}_{k=k_0}^\infty$ is a Cauchy sequence in the space $L^\infty([0, T], L^{2p}(\Omega))$ and hence converges to Z_i^∞ in that space. The limit is $\mathcal{F}_i(t)$ -adapted as Z_i^d is $\mathcal{F}_i(t)$ -adapted for all d . In addition, for any $i \neq j$, the limiting processes Z_i^∞ and Z_j^∞ are independent and have the same law as Z_i^d and Z_j^d are independent and have the same law for any d (recall that the filtrations $\mathcal{F}_i(t)$ and $\mathcal{F}_j(t)$ are independent). Then, consider the estimate

$$\begin{aligned} \mathbb{E}\left[|Z_i^\infty(t) - Z_i^\infty(t+h)|^{2p}\right] &\leq 3^{2p-1} \mathbb{E}\left[|Z_i^\infty(t) - Z_i^d(t)|^{2p}\right] + 3^{2p-1} \mathbb{E}\left[|Z_i^d(t) - Z_i^d(t+h)|^{2p}\right] \\ &\quad + 3^{2p-1} \mathbb{E}\left[|Z_i^d(t+h) - Z_i^\infty(t+h)|^{2p}\right]. \end{aligned} \tag{18}$$

We bound the first and third terms by a multiple of d^{-p} using (16), for example,

$$\begin{aligned} \mathbb{E}\left[|Z_i^\infty(t) - Z_i^d(t)|^{2p}\right]^{1/(2p)} &\leq \sum_{i=1}^{\infty} \mathbb{E}\left[|Z_i^{2^i d}(t) - Z_i^{2^{i-1}d}(t)|^{2p}\right]^{1/(2p)} \\ &\lesssim \sum_{i=1}^{\infty} (2^{i-1}d)^{-1/2} \lesssim d^{-1/2}. \end{aligned} \quad (19)$$

Then, using Lemma 2.6 to bound the second term in (18), substituting back in (18) and taking $d \rightarrow \infty$ yield

$$\mathbb{E}\left[(Z_i^\infty(t) - Z_i^\infty(t+h))^{2p}\right] \lesssim h^p,$$

where the constant is bounded for all $t \in [0, T]$ and h . Since $p > 1$, we can use the Kolmogorov–Chentsov continuity theorem to conclude that there exists a modification of Z_i^∞ that is a.s. pathwise γ -Hölder continuous with $\gamma \in (0, 1/2 - 1/(2p))$. \square

Corollary 2.8 (Rates of $L^\infty([0, T], L^{2p}(\Omega))$ convergence). *Let $\{X_i^d\}_{i=1}^d$ be the strong solution of (2) with d particles. For any $i \in \{1, \dots, d\}$, the following is true:*

$$\text{for any } p > 1, \quad \sup_{0 \leq t \leq T} \mathbb{E}[|Z_i^\infty(t)|^{2p}] \lesssim 1 + \mathbb{E}[|\xi|^{2p}], \quad (20a)$$

$$\sup_{0 \leq t \leq T} \mathbb{E}[|Z_i^d(t) - X_i^d(t)|^2] \lesssim d^{-1}(1 + \mathbb{E}[\xi^2]), \quad (20b)$$

$$\sup_{0 \leq t \leq T} \mathbb{E}[|Z_i^\infty(t) - X_i^d(t)|^2] \lesssim d^{-1}(1 + \mathbb{E}[\xi^2]). \quad (20c)$$

Here, the hidden constants depend only on p, T , and C .

Proof. Using (19), we bound

$$\begin{aligned} \mathbb{E}[|Z_i^\infty(t)|^{2p}] &\leq 2^{2p-1} \mathbb{E}[|Z_i^\infty(t) - Z_i^d(t)|^{2p}] + 2^{2p-1} \mathbb{E}[|Z_i^d(t)|^{2p}] \\ &\lesssim d^{-p} + 1 + \mathbb{E}[|\xi|^{2p}], \end{aligned}$$

taking the limit $d \rightarrow \infty$ yields (20a). For (20b), we know that for any $\mathcal{F}_i(t)$ -measurable $\tilde{Z}(t)$

$$\mathbb{E}[(Z_i^d(t) - X_i^d(t))^2] \leq \mathbb{E}[(\tilde{Z}(t) - X_i^d(t))^2],$$

because

$$Z_i^d(t) := \mathbb{E}[X_i^d(t) | \mathcal{F}_i(t)]$$

and the $L^2(\Omega)$ optimality of the conditional expectation. In particular, consider a particle system $\{\tilde{X}_j^d\}_{j=1}^d$ that satisfies (14) and set $\tilde{\xi}_i = \xi_i$ and $\tilde{W}_i(t) = W_i(t)$ for all t almost surely. Let $\mathcal{G}(t)$ be the σ -algebra generated by $\{\tilde{W}_j(s)\}_{0 \leq s \leq t}$ and $\tilde{\xi}_j$ for all $j \in \{1, \dots, d\} \setminus \{i\}$. Then

$$\mathbb{E}[(Z_i^d(t) - X_i^d(t))^2] \leq \mathbb{E}[(\tilde{X}_i^d(t) - X_i^d(t))^2 | \mathcal{G}(t)].$$

Hence,

$$\mathbb{E}[(Z_i^d(t) - X_i^d(t))^2] \leq \mathbb{E}[(\tilde{X}_i^d(t) - X_i^d(t))^2] \lesssim d^{-1},$$

using Corollary 2.4. Finally, for (20c),

$$\mathbb{E}[|Z_i^\infty(t) - X_i^d(t)|^2] \leq 2 \mathbb{E}[|Z_i^\infty(t) - Z_i^d(t)|^2] + 2 \mathbb{E}[|Z_i^d(t) - X_i^d(t)|^2],$$

and we conclude by bounding the first term using (19) and the second term using (20b). \square

Lemma 2.9 (McKean-Vlasov equation). *Assume that $\mathbb{E}[\xi^4]$ is bounded. Given the initial state ξ_i and the Wiener process $(W_i(t))_{t \geq 0}$ and letting $\mathcal{F}_i(t)$ be the σ -algebra generated by $\{\xi_i, (W_i(s))_{0 \leq s \leq t}\}$, the $L^\infty([0, T], L^2(\Omega))$ limit, Z_i^∞ , satisfies the McKean-Vlasov equation:*

$$\begin{aligned} Z_i^\infty(t) = \xi_i &+ \int_0^t a\left(Z_i^\infty(s), \int_{\mathbb{R}} \kappa_1(Z_i^\infty(s), z) \mu_s(dz)\right) ds \\ &+ \int_0^t \sigma\left(Z_i^\infty(s), \int_{\mathbb{R}} \kappa_2(Z_i^\infty(s), z) \mu_s(dz)\right) dW_i(s), \end{aligned} \quad (21)$$

where μ_s denotes the law of $Z_i^\infty(s)$.

Proof. Without loss of generality set $i = 1$. Denote the right hand side of (21) by $\bar{Z}_1^\infty(t)$, then

$$\mathbb{E}[|Z_1^\infty(t) - \bar{Z}_1(t)|^2] \leq 2\mathbb{E}[|Z_1^\infty(t) - X_1^d(t)|^2] + 2\mathbb{E}[|X_1^d(t) - \bar{Z}_1(t)|^2]. \quad (22)$$

We bound the first term using (20c). For the second term, by Itô's formula, we can bound (recall that $X_1^d(0) = Z_1^\infty(0)$)

$$\mathbb{E}[|X_1^d(T) - \bar{Z}_1^\infty(T)|^2] \leq 2\int_0^T \mathbb{E}[|\Delta a| |X_1^d(t) - Z_1^\infty(t)|] dt + \int_0^T \mathbb{E}[|\Delta\sigma|^2] dt, \quad (23)$$

where

$$\Delta a = a\left(X_1^d(t), \frac{1}{d}\sum_{j=1}^d \kappa_1(X_1^d(t), X_j^d(t))\right) - a\left(Z_1^\infty(t), \int_{\mathbb{R}} \kappa_1(Z_1^\infty(t), z)\mu_t(dz)\right),$$

and $\Delta\sigma$ is similarly defined. Using Young's inequality yields

$$\mathbb{E}[|\Delta a| |X_1^d(t) - Z_1^\infty(t)|] \leq \frac{1}{2}\mathbb{E}[|\Delta a|^2] + \frac{1}{2}\mathbb{E}[|X_1^d(t) - Z_1^\infty(t)|^2]. \quad (24)$$

We then use (3) to bound

$$\begin{aligned} |\Delta a| &\leq \left| a\left(Z_1^\infty(t), \int_{\mathbb{R}} \kappa_1(Z_1^\infty(t), z)\mu_t(dz)\right) - a\left(X_1^d(t), \frac{1}{d}\sum_{j=1}^d \kappa_1(X_1^d(t), X_j^d(t))\right) \right| \\ &\leq C|Z_1^\infty(t) - X_1^d(t)| + C\left| \int_{\mathbb{R}} \kappa_1(Z_1^\infty(t), z)\mu_t(dz) - \frac{1}{d}\sum_{j=1}^d \kappa_1(X_1^d(t), X_j^d(t)) \right| \\ &\leq C|Z_1^\infty(t) - X_1^d(t)| + C\frac{1}{d}\sum_{j=1}^d \left| \kappa_1(Z_1^\infty(t), Z_j^\infty(t)) - \kappa_1(X_1^d(t), X_j^d(t)) \right| \\ &\quad + C\left| \int_{\mathbb{R}} \kappa_1(Z_1^\infty(t), z)\mu_t(dz) - \frac{1}{d}\sum_{j=1}^d \kappa_1(Z_1^\infty(t), Z_j^\infty(t)) \right|. \end{aligned}$$

The second term can again be bounded using (3). Taking the second moment using Jensen's inequality and (20c) yield

$$\mathbb{E}[|\Delta a|^2] \leq (C^2 + 2C^4)d^{-1} + 3C^2\mathbb{E}\left[\left(\frac{1}{d}\sum_{j=1}^d \Delta\kappa_j\right)^2\right], \quad (25)$$

where we denote

$$\Delta\kappa_j := \int_{\mathbb{R}} \kappa_1(Z_1^\infty(t), z)\mu_t(dz) - \kappa_1(Z_1^\infty(t), Z_j^\infty(t)).$$

Here,

$$\mathbb{E}\left[\left(\frac{1}{d}\sum_{j=1}^d \Delta\kappa_j\right)^2\right] = \frac{1}{d^2}\sum_{j=1}^d \sum_{k=1}^d \mathbb{E}[\Delta\kappa_j \Delta\kappa_k].$$

When i, j and k are distinct, we have

$$\mathbb{E}[\Delta\kappa_j \Delta\kappa_k] = \mathbb{E}[\mathbb{E}[\Delta\kappa_j | Z_1^\infty(t)] \times \mathbb{E}[\Delta\kappa_k | Z_1^\infty(t)]] = 0,$$

since, for a given $Z_1^\infty(t)$, $\Delta\kappa_j$ and $\Delta\kappa_k$ are independent and $\mathbb{E}[\Delta\kappa_j | Z_1^\infty(t)] = 0$ when $j \neq i$. When i, j and k are not distinct, we bound using Hölder's inequality and the Lipschitz condition (3) on κ_1

$$|\mathbb{E}[\Delta\kappa_j \Delta\kappa_k]| \leq \mathbb{E}[(\Delta\kappa_j)^2] \leq 4C^2\mathbb{E}[|Z^\infty(t)|^2].$$

Substituting back, we have

$$\left| \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta \kappa_j \right)^2 \right] \right| \leq \frac{3 \times 4C^2}{d} \sup_{0 \leq s \leq T} \mathbb{E} \left[|Z^\infty(s)|^2 \right]. \quad (26)$$

We then substitute (26) into (25) and the result into (24) and then into (23). Similarly, obtaining the corresponding bounds for the second term in (23) (involving $\Delta\sigma$) and using Grönwall's inequality and Lemma 2.2, we arrive at

$$\mathbb{E}[|Z_1^\infty(t) - \bar{Z}_1(t)|^2] \lesssim d^{-1}.$$

Taking $d \rightarrow \infty$, we have that for some $T > 0$ and for all $t \in [0, T]$, $Z^\infty(t) = \bar{Z}(t)$ almost surely. Then, using the continuity of the map $t \rightarrow |Z^\infty(t) - \bar{Z}(t)|$, we conclude that

$$\mathbb{P}[|Z^\infty(t) - \bar{Z}(t)| = 0 \text{ for all } t \in [0, T]] = 1,$$

and hence, Z^∞ satisfies the McKean–Vlasov equation (21). \square

Theorem 2.10 (Existence and uniqueness). *Under Assumption 2.1 and assuming that $\mathbb{E}[\xi^4]$ is bounded, a strong solution to the McKean–Vlasov equation (1) exists and is unique.*

Proof. The existence of a solution is a direct implication of Lemma 2.9. The uniqueness follows in the usual way by assuming that two solutions to (1), Z^∞ and \bar{Z} , exist whose laws at time t are μ_t and $\bar{\mu}_t$, respectively. Then, using Itô's formula, we bound

$$\begin{aligned} \mathbb{E}[|Z^\infty(t) - \bar{Z}(t)|^2] &\leq \mathbb{E}[|Z^\infty(0) - \bar{Z}(0)|^2] \\ &\quad + \int_0^t \mathbb{E}[|\Delta a| \times |Z^\infty(s) - \bar{Z}(s)|] ds + \int_0^t \mathbb{E}[|\Delta\sigma|^2] ds, \end{aligned} \quad (27)$$

where

$$\Delta a := a\left(Z^\infty(t), \int_{\mathbb{R}} \kappa_1(Z^\infty(t), z) \mu_t(dz)\right) - a\left(\bar{Z}(t), \int_{\mathbb{R}} \kappa_1(\bar{Z}(t), z) \bar{\mu}_t(dz)\right),$$

and $\Delta\sigma$ is similarly defined. As before, using Young's inequality,

$$\mathbb{E}[|\Delta a| \times |Z^\infty(s) - \bar{Z}(s)|] \leq \frac{1}{2} \mathbb{E}[|\Delta a|^2] + \frac{1}{2} \mathbb{E}[|Z^\infty(s) - \bar{Z}(s)|^2].$$

Here,

$$|\Delta a|^2 \leq 2C^2 |Z^\infty(s) - \bar{Z}(s)|^2 + 2C^2 \left| \int_{\mathbb{R}^d} \kappa_1(Z^\infty(s), z) \mu_t(dz) - \int_{\mathbb{R}^d} \kappa_1(\bar{Z}(s), z) \bar{\mu}_s(dz) \right|^2,$$

using (3). We then recognize that for some random variables, $Y^\infty(s)$ and $\bar{Y}(s)$, being independent of $Z^\infty(s)$ and $\bar{Z}(s)$ and identically distributed, respectively, we have

$$\begin{aligned} &\left| \int_{\mathbb{R}^d} \kappa_1(Z^\infty(s), z) \mu_t(dz) - \int_{\mathbb{R}^d} \kappa_1(\bar{Z}(s), z) \bar{\mu}_s(dz) \right| \\ &= \left| \mathbb{E}[\kappa_1(Z^\infty(s), Y^\infty(s)) - \kappa_1(\bar{Z}(s), \bar{Y}(s)) \mid Z^\infty(s), \bar{Z}(s)] \right| \\ &\leq C |Z^\infty(s) - \bar{Z}(s)| + C \mathbb{E}[|Y^\infty(s) - \bar{Y}(s)|], \end{aligned}$$

using (3) again. Finally, since

$$\mathbb{E}[|Y^\infty(s) - \bar{Y}(s)|] = \mathbb{E}[|Z^\infty(s) - \bar{Z}(s)|],$$

we have, by further using Jensen's and Hölder's inequalities, that

$$\mathbb{E}[|\Delta a|^2] \leq 2C^2 (1 + 4C^2) \mathbb{E}[|Z^\infty(s) - \bar{Z}(s)|^2].$$

Similarly, we bound $\mathbb{E}[|\Delta\sigma|^2]$ and substitute into (27) and conclude using Grönwall's inequality that

$$\mathbb{E}[|Z^\infty(t) - \bar{Z}(t)|^2] \leq \exp(4C^2 (1 + 4C^2)) \mathbb{E}[|Z^\infty(0) - \bar{Z}(0)|^2].$$

Hence, when $Z^\infty(0) = \bar{Z}(0)$ we have that for some $T > 0$, for all $t \in [0, T]$, $Z^\infty(t) = \bar{Z}(t)$ almost surely. By using the continuity of the map $t \rightarrow |Z^\infty(t) - \bar{Z}(t)|$, we conclude that

$$\mathbb{P}[|Z^\infty(t) - \bar{Z}(t)| = 0 \text{ for all } t \in [0, T]] = 1,$$

and uniqueness is proven. \square

Theorem 2.11 (Rates of $L^\infty([0, T], L^{2p}(\Omega))$ convergence). *Under Assumption 2.1, let $\{X_i^p\}_{i=1}^d$ be the strong solution of (2) and $\{Z_i^\infty\}_{i=1}^d$ be the strong solution of (21); then, for any $i \in \{1, \dots, d\}$ and any integer $p \geq 1$ we have*

$$\sup_{0 \leq t \leq T} \mathbb{E}[|Z_i^\infty(t) - X_i^d(t)|^{2p}] \lesssim d^{-p}(1 + \mathbb{E}[|\xi|^{2p}]). \quad (28)$$

Here, the hidden constant depends only on p, T , and C .

Proof. This result is a generalization of (20c) for higher moments. The proof presented here has the same steps as the proofs of Lemmas 2.3 and 2.9 and is a simple extension of the proof in [23, Theorem I.1.4] using Itô's formula and Lemma A.1. By exchangeability, we only consider $i = 1$; then, we can bound using Itô's formula (recall that $X_1^d(0) = Z_1^\infty(0)$)

$$\begin{aligned} \mathbb{E}[|X_1^d(t) - Z_1^\infty(t)|^{2p}] &\leq p \int_0^t \mathbb{E}[|\Delta a| |X_1^d(s) - Z_1^\infty(s)|^{2p-1}] dt \\ &\quad + p(2p-1) \int_0^t \mathbb{E}[(\Delta \sigma)^2 |X_1^d(s) - Z_1^\infty(s)|^{2p-2}] dt, \end{aligned} \quad (29)$$

where

$$\Delta a := a\left(X_1^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s))\right) - a\left(Z_1^\infty(s), \int_{\mathbb{R}} \kappa_1(Z_1^\infty(s), z) \mu_s(dz)\right)$$

and $\Delta \sigma$ is similarly defined. Using Young's inequality yields

$$\mathbb{E}[|\Delta a| |X_1^d(s) - Z_1^\infty(s)|^{2p-1}] \leq \frac{1}{2p} \mathbb{E}[|\Delta a|^{2p}] + \frac{2p-1}{2p} \mathbb{E}[|X_1^d(s) - Z_1^\infty(s)|^{2p}]. \quad (30)$$

We then bound

$$\begin{aligned} |\Delta a| &\leq \left| a\left(Z_1^\infty(s), \int_{\mathbb{R}} \kappa_1(Z_1^\infty(s), z) \mu_s(dz)\right) - a\left(X_1^d(s), \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s))\right) \right| \\ &\leq C |Z_1^\infty(s) - X_1^d(s)| + C \left| \int_{\mathbb{R}} \kappa_1(Z_1^\infty(s), z) \mu_s(dz) - \frac{1}{d} \sum_{j=1}^d \kappa_1(X_1^d(s), X_j^d(s)) \right| \\ &\leq C |Z_1^\infty(s) - X_1^d(s)| + C \frac{1}{d} \sum_{j=1}^d \left| \kappa_1(Z_1^\infty(s), Z_j^\infty(s)) - \kappa_1(X_1^d(s), X_j^d(s)) \right| \\ &\quad + C \left| \int_{\mathbb{R}} \kappa_1(Z_1^\infty(s), z) \mu_s(dz) - \frac{1}{d} \sum_{j=1}^d \kappa_1(Z_1^\infty(s), Z_j^\infty(s)) \right|. \end{aligned}$$

The second term can be bounded using (3). Taking the p moment, using Jensen's inequality and the fact that for $i, j = 1, \dots, d$ we have

$$\mathbb{E}[|X_i^d(s) - Z_i^\infty(s)|^{2p}] = \mathbb{E}[|X_j^d(s) - Z_j^\infty(s)|^{2p}],$$

by exchangeability, yield

$$\begin{aligned} \mathbb{E}[|\Delta a|^{2p}] &\leq 3^{2p-1} (C^{2p} + 2^{2p-1} C^{4p}) \mathbb{E}[|X_i^d(s) - Z_i^\infty(s)|^{2p}] \\ &\quad + 3^{2p-1} C^{2p} \mathbb{E}\left[\left(\frac{1}{d} \sum_{j=1}^d \Delta \kappa_j\right)^{2p}\right], \end{aligned} \quad (31)$$

where we denote

$$\Delta\kappa_j := \Delta\kappa(Z_1(s), Z_j(s)) := \int_{\mathbb{R}} \kappa_1(Z_1(s), z) \mu_s(dz) - \kappa(Z_1(s), Z_j(s)) .$$

Here, for $\mathcal{I} = \{1, \dots, d\}$,

$$\mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta\kappa_j \right)^{2p} \right] = \frac{1}{d^{2p}} \sum_{\mathbf{k} \in \mathcal{I}^{2p}} \mathbb{E} \left[\prod_{j=1}^{2p} \Delta\kappa_{k_j} \right] .$$

This term is analogous to (11) and can be similarly bounded by noting that $\mathbb{E}[\Delta\kappa_j | Z_1(s)] = 0$ when $j \neq i$ and that for a multi-index $\mathbf{k} \in \mathcal{I}^{2p}$, such that there exists $\ell \in \mathcal{I} \setminus \{1\}$ for which $k_\ell \neq k_j$ for all $j \neq \ell$, we have

$$\mathbb{E} \left[\prod_{j=1}^{2p} \Delta\kappa_{k_j} \right] = \mathbb{E} \left[\mathbb{E}[\Delta\kappa_{k_\ell} | Z_1(s)] \mathbb{E} \left[\prod_{j=1, j \neq \ell}^{2p} \Delta\kappa_{k_j} \middle| Z_1(s) \right] \right] = 0 ,$$

since, for a given $Z_1(s)$, $\{\Delta\kappa_j\}_{j=1}^d$ are independent. Thus, we conclude, similarly to (12), that

$$\left| \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta\kappa_j \right)^{2p} \right] \right| \lesssim d^{-p} \sup_{0 \leq s \leq T} \mathbb{E} \left[|Z(s)|^{2p} \right] . \quad (32)$$

We then substitute (32) into (31) and the result into (30) and (29). By similarly obtaining the corresponding bounds for the second term in (29) (involving $\Delta\sigma$) and using Grönwall's inequality and (20a), we arrive at the result. \square

3 Rate of Weak Convergence

We now prove the rate of weak convergence of X^d , satisfying (2), to Z , the solution to (1). In this section, for a multi-index $\boldsymbol{\ell} \in \mathbb{N}^d$, we define the derivative

$$\frac{\partial^{|\boldsymbol{\ell}|}}{\partial \mathbf{x}^{\boldsymbol{\ell}}} := \frac{\partial^{|\boldsymbol{\ell}|}}{\prod_{j=1}^d \partial x_j^{\ell_j}} ,$$

where $|\boldsymbol{\ell}|$ is the ℓ^1 norm of $\boldsymbol{\ell}$. For $\mathbf{x} \in \mathbb{R}^d$, let $\|\mathbf{x}\|$ denote the Euclidean norm. Let $C^0(\mathbb{R}^d)$ denote the space of continuous functions and $C_b^0(\mathbb{R}^d)$ denote the space of continuous and bounded functions. Define the space $\overline{C}_b^k(\mathbb{R}^d)$ of continuous functions whose partial derivatives up to order k are continuous and bounded. That is

$$\overline{C}_b^k(\mathbb{R}^d) = \left\{ u \in C^0(\mathbb{R}^d) : \frac{\partial^{|\boldsymbol{\ell}|} u}{\partial \mathbf{x}^{\boldsymbol{\ell}}} \in C_b^0(\mathbb{R}^d) \quad \forall 1 \leq |\boldsymbol{\ell}| \leq k \right\}$$

with the associated semi-norm

$$|u|_{\overline{C}_b^k(\mathbb{R}^d)} = \sum_{\boldsymbol{\ell} \in \mathbb{N}^d, 1 \leq |\boldsymbol{\ell}| \leq k} \left\| \frac{\partial^{|\boldsymbol{\ell}|} u}{\partial \mathbf{x}^{\boldsymbol{\ell}}} \right\|_{\overline{C}_b^0(\mathbb{R}^d)} .$$

Theorem 3.1 (Weak convergence result). *Let $\mathbf{X}^d = \{X_i^d\}_{i=1}^d$ be the strong solution to (2) and let $\mathbf{Z} = \{Z_i\}_{i=1}^d$ be d independent samples of the strong solution to (1). Assume that*

$$a, \sigma, \kappa_1 \text{ and } \kappa_2 \in \overline{C}_b^3(\mathbb{R} \times \mathbb{R}) , \quad (33)$$

and that $\mathbb{E}[|\xi|^3]$ is bounded. Then for $g : \mathbb{R}^d \rightarrow \mathbb{R}$, we have

$$\left| \mathbb{E} \left[g(\mathbf{X}^d(T)) - g(\mathbf{Z}(T)) \right] \right| \lesssim d^{-1} |g|_{\overline{C}_b^3(\mathbb{R}^d)} . \quad (34)$$

Proof. Note that (33) implies (3) and hence (4), and, because $\mathbb{E}[|\xi|^3]$ is bounded, $\sup_{0 \leq t \leq T} \mathbb{E}[|Z(t)|^3]$ is also bounded by (20a). For convenience, let

$$C := \|a\|_{\overline{C}_b^3(\mathbb{R} \times \mathbb{R})} + \|\sigma\|_{\overline{C}_b^3(\mathbb{R} \times \mathbb{R})} + \|\kappa_1\|_{\overline{C}_b^3(\mathbb{R} \times \mathbb{R})} + \|\kappa_2\|_{\overline{C}_b^3(\mathbb{R} \times \mathbb{R})} + \mathbb{E}[|\xi|^3].$$

Define the operators

$$\begin{aligned} \mathcal{L}_d u &= \sum_{i=1}^d a \left(x_i, \frac{1}{d} \sum_{j=1}^d \kappa_1(x_i, x_j) \right) \frac{\partial u}{\partial x_i} + \frac{1}{2} \sigma^2 \left(x_i, \frac{1}{d} \sum_{j=1}^d \kappa_2(x_i, x_j) \right) \frac{\partial^2 u}{\partial x_i^2} \\ \text{and } \mathcal{L}_\infty u &= \sum_{i=1}^d a \left(x_i, \int \kappa_1(x_i, z) \mu_t(dz) \right) \frac{\partial u}{\partial x_i} + \frac{1}{2} \sigma^2 \left(x_i, \int \kappa_2(x_i, z) \mu_t(dz) \right) \frac{\partial^2 u}{\partial x_i^2}. \end{aligned}$$

Consider the value function $u(t, \mathbf{x})$ for $\mathbf{x} \in \mathbb{R}^d$, satisfying the PDE

$$\begin{aligned} \frac{\partial u}{\partial t} + \mathcal{L}_d u &= 0, \text{ for } 0 < t < T \\ \text{and } u(T, \mathbf{x}) &= g(\mathbf{x}). \end{aligned} \tag{35}$$

Under (33), u satisfies [11, Theorem 6.1 in Chapter 5]

$$u(t, \mathbf{x}) = \mathbb{E} \left[g(\mathbf{X}^d(T)) \mid X_i(t) = x_i \text{ for } i \in \{1, 2, \dots, d\} \right].$$

For $U(t) := u(t, \mathbf{Z}(t))$, Itô formula yields

$$\begin{aligned} \mathbb{E}[g(\mathbf{Z}(T))] - \mathbb{E}[g(\mathbf{X}^d(T))] &= \mathbb{E}[U(T) - U(0)] \\ &= \mathbb{E} \left[\int_0^T dU(t) \right] \\ &= \mathbb{E} \left[\int_0^T (\mathcal{L}_\infty - \mathcal{L}_d) u(t, \mathbf{Z}(t)) dt \right] \\ &= \int_0^T \mathbb{E}[(\mathcal{L}_\infty - \mathcal{L}_d) u(t, \mathbf{Z}(t))] dt. \end{aligned} \tag{36}$$

The last equality is satisfied under the boundedness conditions in (33), the integrability of Z , and the boundedness of the derivatives of u , which we will establish later. Dropping the u dependence on $\mathbf{Z}(t)$ and t , for brevity, we write

$$\mathbb{E}[(\mathcal{L}_\infty - \mathcal{L}_d)u] = \sum_{i=1}^d \left(\mathbb{E} \left[\Delta_i a \frac{\partial u}{\partial x_i} \right] + \frac{1}{2} \mathbb{E} \left[\Delta_i \sigma^2 \frac{\partial^2 u}{\partial x_i^2} \right] \right), \tag{37}$$

where

$$\begin{aligned} \Delta_i a &:= a \left(Z_i(t), \int \kappa_1(Z_i(t), z) \mu_t(dz) \right) - a \left(Z_i(t), \frac{1}{d} \sum_{j=1}^d \kappa_1(Z_i(t), Z_j(t)) \right) \\ \text{and } \Delta_i \sigma^2 &:= \sigma^2 \left(Z_i(t), \int \kappa_2(Z_i(t), z) \mu_t(dz) \right) - \sigma^2 \left(Z_i(t), \frac{1}{d} \sum_{j=1}^d \kappa_2(Z_i(t), Z_j(t)) \right). \end{aligned}$$

Focusing first on the term involving a , by Taylor expanding, we can bound

$$\begin{aligned} \left| \mathbb{E} \left[\Delta_i a \frac{\partial u}{\partial x_i} \right] \right| &\leq \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{1,j} \right) \frac{\partial u}{\partial x_i} \right] \right| \\ &\quad + \frac{1}{2} \|\bar{a}_2\|_{C_b^0(\mathbb{R})} \left\| \frac{\partial u}{\partial x_i} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{1,j} \right)^2 \right] \end{aligned} \tag{38}$$

where

$$\begin{aligned}\Delta_i \kappa_{1,j} &:= \Delta \kappa_1(Z_i(t), Z_j(t)) := \int \kappa_1(Z_i(t), z) \mu_t(dz) - \kappa_1(Z_i(t), Z_j(t)), \\ \bar{a}_1(x) &:= \frac{\partial a}{\partial y} \left(x, \int \kappa_1(x, z) \mu_t(dz) \right), \\ \text{and } \bar{a}_2(x) &:= \frac{\partial^2 a}{\partial y^2} \left(x, \int \kappa_1(x, z) \mu_t(dz) \right).\end{aligned}$$

By (12), we have

$$\mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{1,j} \right)^2 \right] \lesssim d^{-1} \sup_{0 \leq s \leq T} \mathbb{E} [|Z(s)|^2] \lesssim d^{-1}. \quad (39)$$

Furthermore,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{1,j} \right) \frac{\partial u}{\partial x_i} \right] \right| &\leq \frac{1}{d} \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) (\Delta_i \kappa_{1,i}) \frac{\partial u}{\partial x_i} \right] \right| \\ &+ \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \left(\frac{1}{d} \sum_{j=1, j \neq i}^d \Delta_i \kappa_{1,j} \right) \frac{\partial u}{\partial x_i} \right] \right|. \end{aligned} \quad (40)$$

Here,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) (\Delta_i \kappa_{1,i}) \frac{\partial u}{\partial x_i} \right] \right| &\leq \|\bar{a}_1\|_{C_b^0(\mathbb{R})} \times \left\| \frac{\partial u}{\partial x_i} \right\|_{C_b^0(\mathbb{R}^d)} \times \mathbb{E}[|\Delta_i \kappa_{1,i}|] \\ &\leq 2C^2 \left\| \frac{\partial u}{\partial x_i} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E}[|Z(t)|] \end{aligned} \quad (41)$$

and for $j \neq i$, using a Taylor expansion of $\frac{\partial u}{\partial x_i}$,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \Delta_i \kappa_{1,j} \frac{\partial u}{\partial x_i} \right] \right| &\leq \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \Delta_i \kappa_{1,j} \frac{\partial u}{\partial x_i}(t, \bar{Z}_j(t)) \right] \right| \\ &+ \left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \Delta_i \kappa_{1,j} \int_0^{Z_j(t)} \frac{\partial^2 u}{\partial x_i \partial x_j}(t, \tilde{Z}_j(t)) d\tilde{x} \right] \right|. \end{aligned} \quad (42)$$

Here, $\bar{Z}_j(t)$ and $\tilde{Z}_j(t)$ are the same as $Z(t)$ but with the j 'th entry replaced by 0 or \tilde{x} , respectively. Note that

$$\begin{aligned} \mathbb{E} \left[\bar{a}_1(Z_i(t)) \Delta_i \kappa_{1,j} \frac{\partial u}{\partial x_i}(t, \bar{Z}_j(t)) \right] &= \mathbb{E} \left[\bar{a}_1(Z_i(t)) \mathbb{E}[\Delta_i \kappa_{1,j} | Z_i(t)] \frac{\partial u}{\partial x_i}(t, \bar{Z}_j(t)) \right] \\ &= 0, \end{aligned} \quad (43)$$

because $Z_j(t)$ has law μ_t and is independent of $\bar{Z}_j(t)$. Using (33), we bound

$$\begin{aligned} &\left| \mathbb{E} \left[\bar{a}_1(Z_i(t)) \Delta_i \kappa_{1,j} \int_0^{Z_j(t)} \frac{\partial^2 u}{\partial x_i \partial x_j}(t, \tilde{Z}_j(t)) d\tilde{x} \right] \right| \\ &\leq \|\bar{a}_1\|_{C_b^0(\mathbb{R})} \left\| \frac{\partial^2 u(t, \cdot)}{\partial x_i \partial x_j} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E}[|\Delta \kappa(Z_i(t), Z_j(t))| \cdot |Z_j(t)|] \\ &\leq 2C^2 \left\| \frac{\partial^2 u(t, \cdot)}{\partial x_i \partial x_j} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E}[|Z(t)|^2]. \end{aligned} \quad (44)$$

Substituting (43) and (44) into (42), and the result and (41) into (40), and then substituting the result and (39) into (38), we arrive at

$$\left| \sum_{i=1}^d \mathbb{E} \left[\Delta_i a \frac{\partial u}{\partial x_i} \right] \right| \lesssim d^{-1} |u(t, \cdot)|_{\overline{C}_b^2(\mathbb{R}^d)}. \quad (45)$$

The term involving σ is treated similarly. By Taylor expanding σ^2 , we can bound

$$\begin{aligned} \left| \mathbb{E} \left[\Delta_i \sigma^2 \frac{\partial^2 u}{\partial x_i^2} \right] \right| &\leq \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i(t)) \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right) \frac{\partial^2 u}{\partial x_i^2} \right] \right| \\ &\quad + \left\| \frac{\partial^2 u}{\partial x_i^2} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E} \left[\left| \Sigma \right| \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right)^2 \right] \end{aligned} \quad (46)$$

where

$$\begin{aligned} \bar{\sigma}_1^2(x) &:= \frac{\partial \sigma^2}{\partial y} \left(x, \int \kappa_2(x, z) \mu_t(dz) \right), \\ \Sigma &:= \int_0^1 \bar{\sigma}_2^2 \left(s \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right) + \int \kappa_2(Z_i(t), z) \mu_t(dz) \right) (1-s) ds, \\ \text{and } \bar{\sigma}_2^2(x) &:= \frac{\partial^2 \sigma^2}{\partial y^2} \left(x, \int \kappa_2(x, z) \mu_t(dz) \right) \end{aligned}$$

for all $x \in \mathbb{R}$. Note that by (33), which implies (4), we have

$$\begin{aligned} \bar{\sigma}_1^2(x) &\leq 2C^2 (2 + 2|x| + \mathbb{E}[|Z(t)|]) \\ \bar{\sigma}_2^2(x) &\leq 2C^2 (3 + 2|x| + \mathbb{E}[|Z(t)|]) \\ |\Sigma| &\leq \int_0^1 \left| \bar{\sigma}_2^2 \left(s \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right) + \int \kappa_2(Z_i(t), z) \mu_t(dz) \right) \right| (1-s) ds \\ &\leq 2C^2 \int_0^1 \left(3 + \mathbb{E}[|Z(t)|] + 2 \left| s \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right) + \int \kappa_2(Z_i(t), z) \mu_t(dz) \right| \right) (1-s) ds \\ &\leq \frac{2}{3} C^2 \left(\frac{5}{2} (3 + \mathbb{E}[|Z(t)|]) + \left| \frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right| + |Z_i(t)| \right) \\ &\lesssim 1 + \left| \frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right| + |Z_i(t)| \end{aligned}$$

Next, by Hölder's inequality, (12) and (20a), we have

$$\begin{aligned} \mathbb{E} \left[\left| \Sigma \right| \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right)^2 \right] &\lesssim \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right)^2 \right] + \mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right)^3 \right] \\ &\quad + (\mathbb{E}[|Z_i(t)|^3])^{1/3} \left(\mathbb{E} \left[\left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right)^3 \right] \right)^{2/3} \\ &\lesssim d^{-1}. \end{aligned} \quad (47)$$

Furthermore,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \left(\frac{1}{d} \sum_{j=1}^d \Delta_i \kappa_{2,j} \right) \frac{\partial^2 u}{\partial x_i^2} \right] \right| &\leq \frac{1}{d} \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) (\Delta_i \kappa_{2,i}) \frac{\partial^2 u}{\partial x_i^2} \right] \right| \\ &\quad + \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \left(\frac{1}{d} \sum_{j=1, j \neq i}^d \Delta_i \kappa_{2,j} \right) \frac{\partial^2 u}{\partial x_i^2} \right] \right|. \end{aligned} \quad (48)$$

Here,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,i} \frac{\partial^2 u}{\partial x_i^2} \right] \right| &\leq \left\| \frac{\partial^2 u}{\partial x_i^2} \right\|_{C_b^0(\mathbb{R}^d)} \times \mathbb{E} [|\bar{\sigma}_1^2(Z_i)| \cdot |\Delta_i \kappa_{2,i}|] \\ &\leq 4C^3 \left\| \frac{\partial^2 u}{\partial x_i^2} \right\|_{C_b^0(\mathbb{R}^d)} (2\mathbb{E}[|Z(t)|] + 2(\mathbb{E}[|Z(t)|])^2 + \mathbb{E}[|Z(t)|^2]) \end{aligned} \quad (49)$$

and for $j \neq i$, using a Taylor expansion of $\frac{\partial^2 u}{\partial x_i^2}$,

$$\begin{aligned} \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,j} \frac{\partial^2 u}{\partial x_i^2} \right] \right| &\leq \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,j} \frac{\partial^2 u}{\partial x_i^2}(t, \bar{\mathbf{Z}}_j) \right] \right| \\ &\quad + \left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,j} \int_0^{Z_j} \frac{\partial^3 u}{\partial x_i^2 \partial x_j}(t, \tilde{\mathbf{Z}}_j) d\tilde{x} \right] \right|. \end{aligned} \quad (50)$$

Here, $\bar{\mathbf{Z}}_j, \tilde{\mathbf{Z}}_j$ are the same as \mathbf{Z} but with the j 'th entry replaced by 0 or \tilde{x} , respectively. Note that

$$\begin{aligned} \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,j} \frac{\partial^2 u}{\partial x_i^2}(t, \bar{\mathbf{Z}}_j) \right] &= \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \mathbb{E} [\Delta_i \kappa_{2,j} | Z_i] \frac{\partial^2 u}{\partial x_i^2}(t, \bar{\mathbf{Z}}_j) \right] \\ &= 0, \end{aligned} \quad (51)$$

because Z_j has law μ and is independent of $\bar{\mathbf{Z}}_j(t)$. Using (33), we bound

$$\begin{aligned} &\left| \mathbb{E} \left[\bar{\sigma}_1^2(Z_i) \Delta_i \kappa_{2,j} \int_0^{Z_j} \frac{\partial^3 u}{\partial x_i^2 \partial x_j}(t, \tilde{\mathbf{Z}}_j) d\tilde{x} \right] \right| \\ &\leq \left\| \frac{\partial^3 u(t, \cdot)}{\partial x_i^2 \partial x_j} \right\|_{C_b^0(\mathbb{R}^d)} \mathbb{E} [|\bar{\sigma}_1^2(Z_i)| \cdot |\Delta_i \kappa_{2,j}| \cdot |Z_j|] \\ &\leq 2C^3 \left\| \frac{\partial^3 u(t, \cdot)}{\partial x_i^2 \partial x_j} \right\|_{C_b^0(\mathbb{R}^d)} (2(\mathbb{E}[|Z(t)|])^2 + 3(\mathbb{E}[|Z(t)|])^3 + (2 + 3\mathbb{E}[|Z(t)|])\mathbb{E}[|Z(t)|^2]), \end{aligned} \quad (52)$$

Substituting (51) and (52) into (50), and the result and (49) into (48), and then substituting the result and (39) into (46), we arrive at

$$\left| \sum_{i=1}^d \mathbb{E} \left[\Delta_i \sigma \frac{\partial^2 u}{\partial x_i^2} \right] \right| \lesssim d^{-1} \left(\sum_{i=1}^d \left\| \frac{\partial^2 u(t, \cdot)}{\partial x_i^2} \right\|_{C_b^0(\mathbb{R}^d)} + \sum_{i=1}^d \sum_{j=1}^d \left\| \frac{\partial^3 u(t, \cdot)}{\partial x_i^2 \partial x_j} \right\|_{C_b^0(\mathbb{R}^d)} \right). \quad (53)$$

It is straightforward to see that Assumption B.1 is satisfied for $q = 3$ and

$$\begin{aligned} \nu_i(\mathbf{x}) &:= a \left(x_i, \frac{1}{d} \sum_{i=1}^d \kappa_1(x_i, x_j) \right) \\ \text{and } \varsigma_{i1}(\mathbf{x}) &:= \sigma \left(x_i, \frac{1}{d} \sum_{i=1}^d \kappa_2(x_i, x_j) \right), \end{aligned}$$

given (33). See the discussion after Assumption B.1. Hence, we use Proposition B.3 to bound $|u(t, \cdot)|_{\bar{C}_b^3(\mathbb{R}^d)} \lesssim |g|_{\bar{C}_b^3(\mathbb{R}^d)}$ and substitute this bound into (45) and (53). Finally, by substituting the results into (37) and combining them with (36), we arrive at the final result. \square

Corollary 3.2. *From the previous theorem, we can readily deduce that under the same condition and for an integer $k < d$ and $g : \mathbb{R}^k \rightarrow \mathbb{R}$, we have*

$$\begin{aligned} &\mathbb{E}[g(X_1^d(T), X_2^d(T), \dots, X_k^d(T)) - g(Z_1(T), Z_2(T), \dots, Z_k(T))] \\ &\lesssim \frac{1}{d} \binom{k}{3} \max_{\ell \in \mathbb{N}^k, 1 \leq |\ell| \leq 3} \left(\left\| \frac{\partial^{|\ell|} g}{\partial x^\ell} \right\|_{C_b^0(\mathbb{R}^k)} \right). \end{aligned}$$

Remark 3.3. In the special case when $\kappa_2 = 0$, we can relax (33) and only assume that a, σ and $\kappa_1 \in \overline{\mathcal{C}}_b^2(\mathbb{R} \times \mathbb{R})$. The result would then also involve only the first and second derivatives of g ,

$$\left| \mathbb{E} \left[g(\mathbf{X}^d(T)) - g(\mathbf{Z}(T)) \right] \right| \lesssim d^{-1} |g|_{\overline{\mathcal{C}}_b^2(\mathbb{R}^d)},$$

thus recovering a similar result to the one obtained, for example, in [16, Chapter 9].

4 Numerical Example

In this section, we verify the weak and strong convergence rates that we obtained using a simple McKean–Vlasov equation. For $r \in \mathbb{N}$ and $x \in \mathbb{R}$, consider the function

$$\phi_r(x) := \begin{cases} (1 - x^2)^r & |x| \leq 1 \\ 0 & |x| > 1 \end{cases}$$

and note that ϕ_0 is discontinuous while for $r > 0$ the k 'th derivative of ϕ_r exists and is uniformly bounded for all $k < r$, and the $(r - 1)$ 'th derivative is Lipschitz. Subsequently, consider the particle system (2) with $X_i(0)$ being uniformly distributed in $[-1, 1]$, and for $x, y \in \mathbb{R}$ set

$$\begin{aligned} a(x, y) &:= 2(x - 0.2) + y, \\ \sigma(x, y) &:= 0.2(1 + y), \\ \kappa_1(x, y) &:= \phi_1(10|x - y|) \\ \text{and } \kappa_2(x, y) &:= \phi_1(5|x - y|). \end{aligned} \tag{54}$$

Note that the previous a, σ, κ_1 , and κ_2 , the latter two being only Lipschitz, satisfy the conditions of Theorem 2.11 but not the conditions of Theorem 3.1. To approximate solutions to (2), we use an Euler–Maruyama time-stepping scheme with a fixed number of time-steps, N .

To illustrate the strong and weak convergence of X^d to Z , the corresponding solution of the McKean–Vlasov equation (1), we consider the sequence of systems, denoted by \mathbf{X}^d , satisfying (2) with an increasing number of particles, d . See Figure 1 for a histogram of the values of $X_i^d(1)$ for $d = 2,048$ and using $N = 64$. We also consider the discontinuous function

$$g(x) = \phi_0(10|x - 0.2|).$$

In Figure 2, we plot the quantities $(\mathbb{E}[(X_i^{2d} - X_i^d)^2])^{1/2}$ and $|\mathbb{E}[g(X_i^{2d}) - g(X_i^d)]|$, which verify the rates of Theorem 2.11 and Theorem 3.1, respectively. The same convergence rates were obtained with a different number of time-steps, $N = 32$. Finally, we note that even though κ_1 and κ_2 are only Lipschitz and g is discontinuous, the observed weak convergence rate is still $\mathcal{O}(d^{-1})$, as predicted by Theorem 3.1 when κ_1, κ_2 , and g were assumed to be three-times differentiable. Hence, the assumptions required by Theorem 3.1 and similar theories in the literature can be relaxed in certain cases.

5 Conclusions

We have proven the existence and uniqueness of solutions to (1) and derived the rates of strong and weak convergence of the stochastic particle system, satisfying (2), to the solution of the McKean–Vlasov equation (1). The rates are similar to those in the literature, but we derived them using simpler classical stochastic analysis techniques applied on a system of interacting, exchangeable stochastic particles. Extensions to the theory, such as having multiple kernels in the drift or diffusion coefficients or having n -way kernels, instead of the 2-way kernels presented here, are straightforward. The numerical results that we presented revealed that one could relax the regularity assumptions in the proof of the weak convergence rate on the coefficients and the functional, which could be the next step in this research.




Figure 1: A histogram of the values of $X_i^d(1)$ which follows the system of SDEs in (2) with (54). Here, the values were approximated using the Euler–Maruyama time-stepping scheme with $N = 64$ time-steps and $d = 2,048$.




Figure 2: Convergence rates $X_i^d(1)$ which follows the system of SDEs in (2) with (54). Here, the values were approximated using the Euler–Maruyama time-stepping scheme with $N = 64$ time-steps. Note that the rates are consistent with the predicted rates in Theorems 2.11 and 3.1 even though only the assumptions of the former are satisfied.

References

- [1] BALLY, V., AND TALAY, D. The Euler scheme for stochastic differential equations: error analysis with Malliavin calculus. *Mathematics and Computers in Simulation* 38, 1 (1995), 35 – 41.
- [2] BALLY, V., AND TALAY, D. The law of the Euler scheme for Stochastic Differential Equations: II. Convergence rate of the density. *Monte Carlo Methods and Applications* 2, 2 (1996), 93 – 128.
- [3] BAYER, C., HALL, E. J., AND TEMPONE, R. Weak error rates for option pricing under the rough Bergomi model, 2020.
- [4] BAYER, C., SZEPESSY, A., AND TEMPONE, R. Adaptive weak approximation of reflected and stopped diffusions. *Monte Carlo Methods and Applications* 16, 1 (2010), 1–67.
- [5] BOSSY, M., AND TALAY, D. Convergence rate for the approximation of the limit law of weakly interacting particles: application to the Burgers equation. *The Annals of Applied Probability* 6, 3 (1996), 818–861.
- [6] BOSSY, M., AND TALAY, D. A stochastic particle method for the McKean-Vlasov and the Burgers equation. *Mathematics of Computation of the American Mathematical Society* 66, 217 (1997), 157–192.
- [7] BURKHOLDER, D. L. Martingale transforms. *The Annals of Mathematical Statistics* 37, 6 (1966), 1494–1504.
- [8] CARDALIAGUET, P. Notes on mean field games. Tech. rep., Technical report, 2010.
- [9] CARDALIAGUET, P., DELARUE, F., LASRY, J.-M., AND LIONS, P.-L. The master equation and the convergence problem in mean field games.
- [10] CHASSAGNEUX, J.-F., SZPRUCH, L., AND TSE, A. Weak quantitative propagation of chaos via differential calculus on the space of measures.
- [11] FRIEDMAN, A. *Stochastic differential equations and applications*. Academic Press, 1975.
- [12] HOEL, H., HÄPPÖLÄ, J., AND TEMPONE, R. Construction of a mean square error adaptive euler–maruyama method with applications in multilevel monte carlo. In *Monte Carlo and Quasi-Monte Carlo Methods*, R. Cools and D. Nuyens, Eds. Springer International Publishing, 2016, pp. 29–86.
- [13] HOEL, H., VON SCHWERIN, E., SZEPESSY, A., AND TEMPONE, R. Implementation and analysis of an adaptive multilevel monte carlo algorithm. *Monte Carlo Methods and Applications* 20, 1 (2014), 1–41.
- [14] KATSOULAKIS, M. A., AND SZEPESSY, A. Stochastic hydrodynamical limits of particle systems. *Communications in Mathematical Sciences* 4, 3 (2006), 513–549.
- [15] KLOEDEN, P. E., AND PLATEN, E. *Numerical solution of stochastic differential equations*, vol. 23 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1992.
- [16] KOLOKOLTSOV, V. N. *Nonlinear Markov processes and kinetic equations*, vol. 182. Cambridge University Press, 2010.
- [17] MISCHLER, S., MOUHOT, C., AND WENNERBERG, B. A new approach to quantitative propagation of chaos for drift, diffusion and jump processes. *Probability Theory and Related Fields* 161, 1-2 (2015), 1–59.
- [18] MISHURA, Y. S., AND VERETENNIKOV, A. Y. Existence and uniqueness theorems for solutions of McKean–Vlasov stochastic equations.

- [19] MOON, K.-S., SZEPESSY, A., TEMPONE, R., AND ZOURARIS, G. E. Convergence rates for adaptive weak approximation of stochastic differential equations. *Stochastic analysis and applications* 23, 3 (2005), 511–558.
- [20] MORDECKI, E., SZEPESSY, A., TEMPONE, R., AND ZOURARIS, G. E. Adaptive weak approximation of diffusions with jumps. *SIAM Journal on Numerical Analysis* 46, 4 (2008), 1732–1768.
- [21] ØKSENDAL, B. *Stochastic differential equations: An introduction with applications*, sixth ed. Universitext. Springer-Verlag, Berlin, 2003.
- [22] SZEPESSY, A., TEMPONE, R., AND ZOURARIS, G. E. Adaptive weak approximation of stochastic differential equations. *Communications on Pure and Applied Mathematics: A Journal Issued by the Courant Institute of Mathematical Sciences* 54, 10 (2001), 1169–1214.
- [23] SZNITMAN, A.-S. Topics in propagation of chaos. In *Ecole d'été de probabilités de Saint-Flour XIX–1989*. Springer, 1991, pp. 165–251.
- [24] SZPRUCH, Ł., AND TSE, A. Antithetic multilevel particle system sampling method for McKean-Vlasov SDEs.
- [25] TALAY, D., AND TUBARO, L. Expansion of the global error for numerical schemes solving stochastic differential equations. *Stochastic Analysis and Applications* 8, 4 (1990), 483–509.
- [26] VON SCHWERIN, E., AND SZEPESSY, A. A stochastic phase-field model determined from molecular dynamics. *M2AN Math. Model. Numer. Anal.* 44, 4 (2010), 627–646.

Acknowledgements R. Tempone was partially supported by the KAUST Office of Sponsored Research (OSR) under Award numbers URF/1/2281-01-01, URF/1/2584-01-01 in the KAUST Competitive Research Grants Program Round 8, and the Alexander von Humboldt Foundation, through the Alexander von Humboldt Professorship award. A-L. Haji-Ali was supported by a Sabbatical Grant from the Royal Society of Edinburgh.

A Cardinality of Sets of Multi-indices that Do Not Contain Unique Indices

Lemma A.1 (Multi-indices with no unique indices). *Given a set \mathcal{I} , let \mathcal{I}^p be the p -ary Cartesian power of \mathcal{I} . Define the set of multi-indices with repeated indices as:*

$$\mathcal{J}_{\mathcal{I}^p} = \left\{ (k_1, k_2, \dots, k_p) \in \mathcal{I}^p : \forall j \in \{1, \dots, p\} \exists j' \in \{1, \dots, p\} \setminus \{j\} \text{ s.t. } k_j = k_{j'} \right\}. \quad (55)$$

Then, for any $p \geq 2$, there exists a constant c_p such that

$$|\mathcal{J}_{\mathcal{I}^p}| \leq c_p |\mathcal{I}|^{\lfloor p/2 \rfloor}. \quad (56)$$

Proof. We can directly compute that $|\mathcal{J}_{\mathcal{I}^2}| = |\mathcal{J}_{\mathcal{I}^3}| = |\mathcal{I}|$. Then, assuming that the result is true for p , we consider $\mathcal{J}_{\mathcal{I}^{p+2}}$

$$\begin{aligned} |\mathcal{J}_{\mathcal{I}^{p+2}}| &= |\mathcal{I}| + \sum_{i=1}^{|\mathcal{I}|} \sum_{j=2}^p \binom{p+2}{j} |\mathcal{J}_{(\mathcal{I} \setminus \{i\})^{p+2-j}}| \\ &= |\mathcal{I}| + \sum_{i=1}^{|\mathcal{I}|} \sum_{j=0}^{p-2} \binom{p+2}{j+2} |\mathcal{J}_{(\mathcal{I} \setminus \{i\})^{p-j}}| \\ &\leq |\mathcal{I}| + \sum_{i=1}^{|\mathcal{I}|} \sum_{j=0}^{p-2} \binom{p+2}{j+2} c_{p-j} (|\mathcal{I}| - 1)^{\frac{p-j}{2}} \\ &\leq |\mathcal{I}| + |\mathcal{I}|^{1+p/2} \sum_{j=0}^{p-2} \binom{p+2}{j+2} c_{p-j} (|\mathcal{I}| - 1)^{\frac{-j}{2}} =: c_{p+2} |\mathcal{I}|^{1+p/2}. \end{aligned}$$

□

B Moments Bounds for SDE Variations with Sobolev-Bounded Coefficients

In this section, for $T > 0$ and $(t, \mathbf{x}) \in [0, T] \times \mathbb{R}^d$, we consider a general SDE of the form

$$\mathbf{X}^{t, \mathbf{x}}(s) = \mathbf{x} + \int_t^s \boldsymbol{\nu}(\tau, \mathbf{X}^{t, \mathbf{x}}(\tau)) d\tau + \int_t^s \boldsymbol{\varsigma}(\tau, \mathbf{X}^{t, \mathbf{x}}(\tau)) d\mathbf{W}(\tau), \quad s \in [t, T], \quad (57)$$

with drift coefficient $\boldsymbol{\nu} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$, a diffusion coefficient $\boldsymbol{\varsigma} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^{d \times d'}$, and \mathbf{W} is a vector of d' independent standard Wiener processes over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ with the natural filtration. In what follows, for any $f : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}$, define the norm

$$\|f\|_\infty = \sup_{t \leq s \leq T} \|f(s, \cdot)\|_{C_b^q(\mathbb{R}^d)}.$$

Assumption B.1 (Bounded derivatives). *Given $T > 0$, there exists a constant $C_0 > 0$ such that for all $t \in [0, T]$ and $\mathbf{x} \in \mathbb{R}^d$*

$$(\|\boldsymbol{\nu}(t, \mathbf{x})\| + \|\boldsymbol{\varsigma}(t, \mathbf{x})\|) \leq C_0(1 + \|\mathbf{x}\|).$$

Moreover, for some strictly positive integer $q \geq 1$ and all $i \in \{1, \dots, d\}$, $m \in \{1, \dots, d'\}$ and $\boldsymbol{\ell} \in \mathbb{N}^d$ such that $|\boldsymbol{\ell}| \leq q$ the derivatives $\frac{\partial^{|\boldsymbol{\ell}|} \nu_i}{\partial \mathbf{x}^{\boldsymbol{\ell}}}$ and $\frac{\partial^{|\boldsymbol{\ell}|} \varsigma_{im}}{\partial \mathbf{x}^{\boldsymbol{\ell}}}$ exist and are continuous. Additionally, there exists a constant $C_q > 0$ such that

$$\sum_{k=1}^q \sum_{p=0}^k \sum_{\substack{\boldsymbol{\ell} \in \mathbb{N}^d \\ |\boldsymbol{\ell}|_1 = k-p}} \max_{\substack{i \in \{1, 2, \dots, d\} \\ \text{s.t. } \ell_i = 0}} \left\| \frac{\partial^k \nu_i}{\partial x_i^p \partial \mathbf{x}^{\boldsymbol{\ell}}} \right\|_\infty \leq C_q \quad (58a)$$

$$\text{and} \quad \sum_{k=1}^q \sum_{p=0}^k \sum_{\substack{\boldsymbol{\ell} \in \mathbb{N}^d \\ |\boldsymbol{\ell}|_1 = k-p}} \max_{\substack{i \in \{1, 2, \dots, d\} \\ \text{s.t. } \ell_i = 0}} \left(\sum_{m=1}^{d'} \left\| \frac{\partial^k \varsigma_{im}}{\partial x_i^p \partial \mathbf{x}^{\boldsymbol{\ell}}} \right\|_\infty^2 \right)^{1/2} \leq C_q. \quad (58b)$$

The conditions (58a) and (58b) deserve some explanation. Focusing on (58a), for $q = 1$, the left-hand side simplifies to

$$\max_i \left\| \frac{\partial \nu_i}{\partial x_i} \right\|_\infty + \sum_{\ell=1}^d \max_{i \neq \ell} \left\| \frac{\partial \nu_i}{\partial x_\ell} \right\|_\infty.$$

Hence, a sufficient condition is to bound the diagonal and off-diagonal elements of the Jacobian of $\boldsymbol{\nu}$, i.e.,

$$\left\| \frac{\partial \nu_i}{\partial x_\ell} \right\|_\infty = \begin{cases} \mathcal{O}(1) & i = \ell', \\ \mathcal{O}(d^{-1}) & i \neq \ell'. \end{cases}$$

For $q = 2$, the left-hand side would also include the terms

$$\max_i \left\| \frac{\partial^2 \nu_i}{\partial x_i^2} \right\|_\infty + \sum_{\ell=1}^d \max_{i \neq \ell} \left\| \frac{\partial^2 \nu_i}{\partial x_i \partial x_\ell} \right\|_\infty + \sum_{\ell=1}^d \sum_{\ell'=1}^d \max_{i \notin \{\ell, \ell'\}} \left\| \frac{\partial^2 \nu_i}{\partial x_\ell \partial x_{\ell'}} \right\|_\infty$$

and an additional condition on the second derivative would be required, for example

$$\left\| \frac{\partial^2 \nu_i}{\partial x_i \partial x_{\ell'}} \right\|_\infty = \begin{cases} \mathcal{O}(1) & i = \ell = \ell' \\ \mathcal{O}(d^{-1}) & i = \ell \neq \ell' \text{ or } i \neq \ell = \ell' \\ \mathcal{O}(d^{-2}) & i \neq \ell \neq \ell' \end{cases}$$

In general, a sufficient condition is

$$\left\| \frac{\partial^{|\boldsymbol{\ell}|} \nu_i}{\partial \mathbf{x}^{\boldsymbol{\ell}}} \right\|_\infty = \mathcal{O}(d^{1-|\boldsymbol{\ell}|+e_i|0}) \quad \text{for all } i \in \{1, \dots, d\} \text{ and } \boldsymbol{\ell} \in \mathbb{N}^d : |\boldsymbol{\ell}| \leq q,$$

where $|\ell|_0$ denotes the number of non-zero elements of ℓ and similarly

$$\left(\sum_{m=1}^{d'} \left\| \frac{\partial^{|\ell|} \zeta_{im}}{\partial \mathbf{x}^\ell} \right\|_\infty^2 \right)^{1/2} = \mathcal{O}(d^{1-|\ell+e_i|_0}) \quad \text{for all } i \in \{1, \dots, d\} \text{ and } \ell \in \mathbb{N}^d : |\ell| \leq q .$$

Proposition B.2 (L^p flux bound). *For $\mathbf{X}^{t,\mathbf{x}}$ satisfying (57) and Assumption B.1 being satisfied for $q \in \{1, 2, 3\}$, and for any $p \geq 2$ there exists a constant $K_{q,p}$, depending only on t, T, p , and C_q such that*

$$\sum_{k=1}^q \sum_{p=0}^k \sum_{\substack{\ell \in \mathbb{N}^d \\ |\ell|_1 = k-p}} \sup_{t \leq s \leq T} \max_{\substack{i \in \{1, 2, \dots, d\} \\ \text{s.t. } \ell_i = 0}} \mathbb{E} \left[\left| \frac{\partial^k X_i^{\mathbf{x}, t}}{\partial x_i^p \partial \mathbf{x}^\ell} (s) \right|^p \right]^{1/p} \leq K_{q,p} .$$

Proof. We will make repeated use of the following inequality

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} Y_i \right)^2 \right)^{p/2} \right] &\leq \left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} \right)^2 \right)^{p/2} \max_{i \in \mathcal{I}} \mathbb{E}[|Y_i|^p] \\ &\leq \left(\sum_{i \in \mathcal{I}} \left(\sum_{j \in \mathcal{J}} a_{i,j}^2 \right)^{1/2} \right)^p \max_{i \in \mathcal{I}} \mathbb{E}[|Y_i|^p] , \end{aligned} \quad (59)$$

for any random variable Y , any sequence of non-negative measurable functions $a_{i,j}$ and any index sets \mathcal{I} and \mathcal{J} . Both inequalities can be proven using Hölder's inequality,

$$\begin{aligned} \mathbb{E} \left[\left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} Y_i \right)^2 \right)^{p/2} \right] &\leq \mathbb{E} \left[\left(\sum_{j \in \mathcal{J}} \left(\sum_{i' \in \mathcal{I}} a_{i',j} \right) \sum_{i \in \mathcal{I}} a_{i,j} Y_i^2 \right)^{p/2} \right] \\ &= \mathbb{E} \left[\left(\sum_{i \in \mathcal{I}} \left(\sum_{j \in \mathcal{J}} \left(\sum_{i' \in \mathcal{I}} a_{i',j} \right) a_{i,j} \right) Y_i^2 \right)^{p/2} \right] \\ &\leq \left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} \right)^2 \right)^{p/2-1} \sum_{i \in \mathcal{I}} \left(\sum_{j \in \mathcal{J}} \left(\sum_{i' \in \mathcal{I}} a_{i',j} \right) a_{i,j} \right) \mathbb{E}[|Y_i|^p] \\ &\leq \left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} \right)^2 \right)^{p/2} \max_{i \in \mathcal{I}} \mathbb{E}[|Y_i|^p] , \\ \left(\sum_{j \in \mathcal{J}} \left(\sum_{i \in \mathcal{I}} a_{i,j} \right)^2 \right)^{1/2} &= \left(\sum_{i \in \mathcal{I}} \sum_{i' \in \mathcal{I}} \sum_{j \in \mathcal{J}} a_{i,j} a_{i',j} \right)^{1/2} \\ &\leq \left(\sum_{i \in \mathcal{I}} \sum_{i' \in \mathcal{I}} \left(\sum_{j \in \mathcal{J}} a_{i,j}^2 \right)^{1/2} \left(\sum_{j \in \mathcal{J}} a_{i',j}^2 \right)^{1/2} \right)^{1/2} \\ &= \sum_{i \in \mathcal{I}} \left(\sum_{j \in \mathcal{J}} a_{i,j}^2 \right)^{1/2} . \end{aligned}$$

In addition, note that

$$\max_{i \in \mathcal{I}} \sum_{j=1}^d a_{i,j} = \max_{i \in \mathcal{I}} a_{i,i} + \max_{i \in \mathcal{I}} \sum_{j=1, j \neq i}^d a_{i,j} \leq \max_{i \in \mathcal{I}} a_{i,i} + \sum_{j=1}^d \max_{i \in \mathcal{I}, i \neq j} a_{i,j}$$

First variation For brevity, denote $f_{x_j} = \frac{\partial f}{\partial x_j}$. First, note that the process $X_{i,x_j}^{t,x}$ exists under Assumption B.1 for $q \geq 1$ and satisfies for $s \geq t$ the SDE

$$\begin{aligned} X_{i,x_j}^{t,x}(s) &= \delta_{i,j} + \int_t^s \sum_{k=1}^d \nu_{i,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) d\tau \\ &\quad + \sum_{m=1}^{d'} \int_t^s \sum_{k=1}^d \varsigma_{im,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) dW_m(\tau), \end{aligned}$$

cf. [11, Theorem 5.3 in Chapter 5]. By Itô's formula

$$\begin{aligned} \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(s) \right|^p \right] &\leq \delta_{i,j} + p \int_t^s \mathbb{E} \left[\left| \sum_{k=1}^d \nu_{i,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) \right| \left| X_{i,x_j}^{t,x}(\tau) \right|^{p-1} \right] d\tau \\ &\quad + \frac{p(p-1)}{2} \int_t^s \sum_{m=1}^{d'} \mathbb{E} \left[\left(\sum_{k=1}^d \varsigma_{im,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) \right)^2 \left| X_{i,x_j}^{t,x}(\tau) \right|^{p-2} \right] d\tau. \end{aligned} \quad (60)$$

For the term involving ν , using Young's inequality, we can bound

$$\begin{aligned} &\mathbb{E} \left[\left| \sum_{k=1}^d \nu_{i,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) \right| \left| X_{i,x_j}^{t,x}(\tau) \right|^{p-1} \right] \\ &\leq \sum_{k=1}^d \mathbb{E} \left[\|\nu_{i,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \left| X_{i,x_j}^{t,x}(\tau) \right|^{p-1} \right] \\ &\leq \frac{1}{p} \mathbb{E} \left[\left(\sum_{k=1}^d \|\nu_{i,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \right)^p \right] + \frac{p-1}{p} \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(\tau) \right|^p \right]. \end{aligned} \quad (61)$$

By using Jensen's inequality and (59), we can further bound

$$\begin{aligned} &\mathbb{E} \left[\left(\sum_{k=1}^d \|\nu_{i,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \right)^p \right] \\ &\leq 2^{p-1} \|\nu_{i,x_j}\|_\infty^p \mathbb{E} \left[\left| X_{j,x_j}^{t,x}(\tau) \right|^p \right] + 2^{p-1} \mathbb{E} \left[\left(\sum_{k=1, k \neq j}^d \|\nu_{i,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \right)^p \right] \\ &\leq 2^{p-1} \|\nu_{i,x_j}\|_\infty^p \max_k \mathbb{E} \left[\left| X_{k,x_j}^{t,x}(\tau) \right|^p \right] + 2^{p-1} C_1^p \max_{k \neq j} \mathbb{E} \left[\left| X_{k,x_j}^{t,x}(\tau) \right|^p \right]. \end{aligned} \quad (62)$$

For the term involving ς , using Young's inequality,

$$\begin{aligned} &\mathbb{E} \left[\sum_{m=1}^{d'} \left(\sum_{k=1}^d \|\varsigma_{im,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \right)^2 \left| X_{i,x_j}^{t,x}(\tau) \right|^{p-2} \right] \\ &\leq \frac{2}{p} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1}^d \|\varsigma_{im,x_k}\|_\infty \left| X_{k,x_j}^{t,x}(\tau) \right| \right)^2 \right)^{p/2} \right] + \frac{p-2}{p} \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(\tau) \right|^p \right]. \end{aligned} \quad (63)$$

Using Jensen's inequality, we can further bound

$$\begin{aligned}
& \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1}^d \|\varsigma_{im, x_k}\|_\infty |X_{k, x_j}^{t, x}(\tau)| \right)^2 \right)^{p/2} \right] \\
& \leq 2^{p-1} \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j}\|_\infty^2 \right)^{p/2} \max_k \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right] \\
& \quad + 2^{p-1} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1, k \neq j}^d \|\varsigma_{im, x_k}\|_\infty |X_{k, x_j}^{t, x}(\tau)| \right)^2 \right)^{p/2} \right],
\end{aligned} \tag{64}$$

wherein using (59), we have

$$\begin{aligned}
& \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1, k \neq j}^d \|\varsigma_{im, x_k}\|_\infty |X_{k, x_j}^{t, x}(\tau)| \right)^2 \right)^{p/2} \right] \\
& \leq \left(\sum_{k=1, k \neq j}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_k}\|_\infty^2 \right)^{p/2} \right) \max_{k \neq j} \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right] \\
& \leq C_1^p \max_{k \neq j} \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right].
\end{aligned} \tag{65}$$

Combining (62) with (61) and (65) with (64) and (63) and substituting it into (60) and simplifying yield

$$\begin{aligned}
\mathbb{E} \left[|X_{i, x_j}^{t, x}(s)|^p \right] & \leq \delta_{ij} + c_1 \int_t^s \mathbb{E} \left[|X_{i, x_j}^{t, x}(\tau)|^p \right] d\tau \\
& \quad + c_2 \int_t^s \max_{k \neq j} \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right] d\tau \\
& \quad + 2^{p-1} (p-1) a_{ij}^p \int_t^s \max_k \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right] d\tau,
\end{aligned} \tag{66}$$

where

$$\begin{aligned}
c_1 & := \frac{p(p-1)}{2}, \\
c_2 & := p2^{p-1} C_1^p
\end{aligned}$$

$$\text{and } a_{ij}^p := \|\nu_{i, x_j}\|_\infty^p + \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j}\|_\infty^2 \right)^{p/2},$$

so that $a_{i, i} + \sum_{j=1}^d \max_{i \neq j} a_{i, j} \leq C_1$. Then, taking the maximum over all i in (66), we arrive at

$$\max_i \mathbb{E} \left[|X_{i, x_j}^{t, x}(s)|^p \right] \leq 1 + (c_1 + 2c_2) \int_t^s \max_i \mathbb{E} \left[|X_{i, x_j}^{t, x}(\tau)|^p \right] d\tau$$

and using Grönwall's inequality yields

$$\max_i \mathbb{E} \left[|X_{i, x_j}^{t, x}(s)|^p \right] \leq \exp((c_1 + 2c_2)(s-t)) =: D_1.$$

Taking the maximum over all $i \neq j$ in (66), we arrive at

$$\begin{aligned}
\max_{i \neq j} \mathbb{E} \left[|X_{i, x_j}^{t, x}(s)|^p \right] & \leq (c_1 + c_2) \int_t^s \max_{i \neq j} \mathbb{E} \left[|X_{i, x_j}^{t, x}(\tau)|^p \right] d\tau \\
& \quad + p2^{p-1} \left(\max_{i \neq j} (a_{ij}^p) \right) \int_t^s \max_k \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^p \right] d\tau
\end{aligned}$$

and using Grönwall's inequality yields

$$\begin{aligned}
\max_{i \neq j} \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(s) \right|^p \right] &\leq p 2^{p-1} \left(\max_{i \neq j} a_{ij}^p \right) \exp((c_1 + c_2)(s-t)) \int_t^s \max_k \mathbb{E} \left[\left| X_{k,x_j}^{t,x}(\tau) \right|^p \right] d\tau \\
&\leq p 2^{p-1} \left(\max_{i \neq j} a_{ij}^p \right) \exp((c_1 + c_2)(s-t)) \left(\int_t^s D_1 d\tau \right) \\
&=: \left(\max_{i \neq j} a_{ij}^p \right) \tilde{D}_1 .
\end{aligned}$$

Finally

$$\begin{aligned}
\max_i \mathbb{E} \left[\left| X_{i,x_i}^{t,x}(s) \right|^p \right]^{1/p} + \sum_{j=1}^d \max_{i \neq j} \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(s) \right|^p \right]^{1/p} &\leq D_1^{1/p} + \tilde{D}_1^{1/p} \sum_{j=1}^d \max_{i \neq j} a_{ij} \\
&\leq D_1^{1/p} + \tilde{D}_1^{1/p} C_1 \\
&=: K_{1,p} .
\end{aligned}$$

Second variation Because the proof steps were fully illustrated in the previous section, in this section, we simplify the presentation by using D_2 to denote constants depending only on t, T, p , and C_2 . Observe that these constants might change their values from one line to the next.

Again, denote $f_{x_j, x_{j'}} = \frac{\partial^2 f}{\partial x_j \partial x_{j'}}$ and note that the process $\left\{ X_{i,x_j, x_{j'}}^{t,x}(s) \right\}_{s \in [t, T]}$ exists under Assumption B.1 for $q \geq 2$ and satisfies for $s \in [t, T]$ the SDE

$$\begin{aligned}
X_{i,x_j, x_{j'}}^{t,x}(s) &= \int_t^s \sum_{k=1}^d \nu_{i,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j, x_{j'}}^{t,x}(\tau) d\tau \\
&\quad + \int_t^s \sum_{k=1}^d \sum_{k'=1}^d \nu_{i,x_k, x_{k'}}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) d\tau \\
&\quad + \sum_{m=1}^{d'} \int_t^s \sum_{k=1}^d \varsigma_{im, x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j, x_{j'}}^{t,x}(\tau) dW_m(\tau) \\
&\quad + \sum_{m=1}^{d'} \int_t^s \sum_{k=1}^d \sum_{k'=1}^d \varsigma_{im, x_k, x_{k'}}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) dW_m(\tau) ,
\end{aligned}$$

cf. [11, Theorem 5.4 in Chapter 5]. By Itô's formula,

$$\begin{aligned}
\mathbb{E} \left[\left| X_{i,x_j, x_{j'}}^{t,x}(s) \right|^p \right] &\leq p \int_t^s \mathbb{E} \left[(f_1 + f_3) \cdot \left| X_{i,x_j, x_{j'}}^{t,x}(\tau) \right|^{p-1} \right] d\tau \\
&\quad + \frac{p(p-1)}{2} \int_t^s \mathbb{E} \left[(f_2 + f_4) \cdot \left| X_{i,x_j, x_{j'}}^{t,x}(\tau) \right|^{p-2} \right] d\tau ,
\end{aligned} \tag{67}$$

where

$$\begin{aligned}
f_1 &:= \left| \sum_{k=1}^d \nu_{i,x_k}(\tau, X^{t,x}(\tau)) X_{k,x_j, x_{j'}}^{t,x}(\tau) \right| , \\
f_2 &:= \sum_{m=1}^{d'} \left(\sum_{k=1}^d (\varsigma_{im, x_k}(\tau, X^{t,x}(\tau))) X_{k,x_j, x_{j'}}^{t,x}(\tau) \right)^2 , \\
f_3 &:= \left| \sum_{k=1}^d \sum_{k'=1}^d \nu_{i,x_k, x_{k'}}(\tau, X^{t,x}(\tau)) X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) \right| \\
\text{and } f_4 &:= \sum_{m=1}^{d'} \left(\sum_{k=1}^d \sum_{k'=1}^d (\varsigma_{im, x_k, x_{k'}}(\tau, X^{t,x}(\tau))) X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) \right)^2 .
\end{aligned}$$

As before, the first step is to apply Young's inequality to each of the previous integrands. For integers $1 \leq q_u < p$ and $u \in \{1, 2, 3, 4\}$, we have

$$\mathbb{E} \left[f_u \cdot \left| X_{i,x_j,x_{j'}}^{t,x}(\tau) \right|^{p-q_u} \right] \leq \frac{q_u}{p} \mathbb{E} \left[f_u^{p/q_u} \right] + \frac{p-q_u}{p} \mathbb{E} \left[\left| X_{i,x_j,x_{j'}}^{t,x}(\tau) \right|^p \right].$$

We now turn our attention to bounding $\mathbb{E} [f_u^{p/q_u}]$ predominately using Jensen's inequality and (59). We begin by bounding $\mathbb{E} [f_2^{p/2}]$ and $\mathbb{E} [f_4^{p/2}]$. Bounding $\mathbb{E} [f_1^p]$ and $\mathbb{E} [f_3^p]$ is analogous.

$$\begin{aligned} \mathbb{E} [f_2^{p/2}] &\leq \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1}^d \|\zeta_{im,x_k}\|_\infty |X_{k,x_j,x_{j'}}^{t,x}(\tau)| \right)^2 \right)^{p/2} \right] \\ &\leq 4^{p-1} \sum_{k \in \{j,j'\}} \left(\sum_{m=1}^{d'} \|\zeta_{im,x_k}\|_\infty^2 \right)^{p/2} \mathbb{E} \left[|X_{k,x_j,x_{j'}}^{t,x}(\tau)|^p \right] \\ &\quad + 2^{p-1} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k \in \{1,2,\dots,d\} \setminus \{j,j'\}} \|\zeta_{im,x_k}\|_\infty |X_{k,x_j,x_{j'}}^{t,x}(\tau)| \right)^2 \right)^{p/2} \right] \\ &\leq 4^{p-1} \sum_{k \in \{j,j'\}} \left(\sum_{m=1}^{d'} \|\zeta_{im,x_k}\|_\infty^2 \right)^{p/2} \left(\mathbb{E} \left[|X_{k,x_j,x_{j'}}^{t,x}(\tau)|^p \right] \right) \\ &\quad + 2^{p-1} C_1^p \max_{k \in \{1,2,\dots,d\} \setminus \{j,j'\}} \mathbb{E} \left[|X_{k,x_j,x_{j'}}^{t,x}(\tau)|^p \right], \end{aligned}$$

where we used (59) in the last step. On the other hand,

$$\begin{aligned} \mathbb{E} [f_4^{p/2}] &\leq \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1}^d \sum_{k'=1}^d \|\zeta_{im,x_k,x_{k'}}\|_\infty X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) \right)^2 \right)^{p/2} \right] \\ &\leq 4^{p-1} \left(\sum_{m=1}^{d'} \|\zeta_{im,x_j,x_{j'}}\|_\infty^2 \right)^{p/2} \mathbb{E} \left[\left| X_{j,x_j}^{t,x}(\tau) X_{j',x_{j'}}^{t,x}(\tau) \right|^p \right] \\ &\quad + 4^{p-1} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k'=1, k' \neq j'}^d \|\zeta_{im,x_j,x_{k'}}\|_\infty X_{k',x_{j'}}^{t,x}(\tau) \right)^2 \right)^{p/2} \left| X_{j,x_j}^{t,x}(\tau) \right|^p \right] \\ &\quad + 4^{p-1} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1, k \neq j}^d \|\zeta_{im,x_k,x_{j'}}\|_\infty X_{j,x_k}^{t,x}(\tau) \right)^2 \right)^{p/2} \left| X_{j',x_{j'}}^{t,x}(\tau) \right|^p \right] \\ &\quad + 4^{p-1} \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1, k \neq j}^d \sum_{k'=1, k' \neq j'}^d \|\zeta_{im,x_k,x_{k'}}\|_\infty X_{k,x_j}^{t,x}(\tau) X_{k',x_{j'}}^{t,x}(\tau) \right)^2 \right)^{p/2} \right]. \end{aligned}$$

Looking at each term separately

$$\begin{aligned} \mathbb{E} \left[\left| X_{j,x_j}^{t,x}(\tau) X_{j',x_{j'}}^{t,x}(\tau) \right|^p \right] &\leq \left(\max_i \mathbb{E} \left[\left| X_{i,x_j}^{t,x}(\tau) \right|^{2p} \right] \right)^{1/2} \left(\max_i \mathbb{E} \left[\left| X_{i,x_{j'}}^{t,x}(\tau) \right|^{2p} \right] \right)^{1/2} \\ &\leq K_{1,2p}^{2p}. \end{aligned}$$

Moreover, using Hölder's inequality and (59)

$$\begin{aligned}
& \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k'=1, k' \neq j'}^d \|\varsigma_{im, x_j, x_{k'}}\|_{\infty} X_{k', x_{j'}}^{t, x}(\tau) \right)^2 \right)^{p/2} \left| X_{j, x_j}^{t, x}(\tau) \right|^p \right] \\
& \leq \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k'=1, k' \neq j'}^d \|\varsigma_{im, x_j, x_{k'}}\|_{\infty} X_{k', x_{j'}}^{t, x}(\tau) \right)^2 \right)^{2p} \right]^{1/2} \mathbb{E} \left[\left| X_{j, x_j}^{t, x}(\tau) \right|^{2p} \right]^{1/2} \\
& \leq \left(\sum_{k'=1}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j, x_{k'}}\|_{\infty}^2 \right)^{1/2} \right)^p \left(\max_{k' \neq j'} \mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2} \mathbb{E} \left[\left| X_{j, x_j}^{t, x}(\tau) \right|^{2p} \right]^{1/2} \\
& \leq K_{1,2p}^p \left(\sum_{k'=1}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j, x_{k'}}\|_{\infty}^2 \right)^{1/2} \right)^p \left(\max_{k' \neq j'} \mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2}
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E} \left[\left(\sum_{m=1}^{d'} \left(\sum_{k=1, k \neq j}^d \sum_{k'=1, k' \neq j'}^d \|\varsigma_{i, x_k, x_{k'}}\|_{\infty} X_{k, x_j}^{t, x}(\tau) X_{k', x_{j'}}^{t, x}(\tau) \right)^2 \right)^{p/2} \right] \\
& \leq \left(\sum_{k=1, k \neq j}^d \sum_{k'=1, k' \neq j'}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_k, x_{k'}}\|_{\infty}^2 \right)^{1/2} \right)^p \max_{k \neq j, k' \neq j'} \mathbb{E} \left[\left| X_{k, x_j}^{t, x}(\tau) X_{k', x_{j'}}^{t, x}(\tau) \right|^p \right] \\
& \leq C_2^p \left(\max_{k, k \neq j} \mathbb{E} \left[\left| X_{k, x_j}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2} \left(\max_{k', k' \neq j'} \mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2}.
\end{aligned}$$

Hence

$$\begin{aligned}
\mathbb{E}[f_4^{p/2}] & \leq 4^{p-1} \left(K_{1,2p}^{2p} F_{\varsigma_1, i, j, j'} + K_{1,2p}^p F_{\varsigma_2, i, j, j'} + K_{1,2p}^p F_{\varsigma_3, i, j, j'} + C_2^p F_{4, j, j'} \right) \\
& \leq D_2 (F_{\varsigma_1, i, j, j'} + F_{\varsigma_2, i, j, j'} + F_{\varsigma_3, i, j, j'} + F_{4, j, j'}),
\end{aligned}$$

where

$$\begin{aligned}
F_{\varsigma_1, i, j, j'} & := \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j, x_{j'}}\|_{\infty}^2 \right)^{p/2}, \\
F_{\varsigma_2, i, j, j'} & := \left(\sum_{k'=1}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j, x_{k'}}\|_{\infty}^2 \right)^{1/2} \right)^p \left(\max_{k' \neq j'} \mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2}, \\
F_{\varsigma_3, i, j, j'} & := \left(\sum_{k=1}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_k, x_{j'}}\|_{\infty}^2 \right)^{1/2} \right)^p \left(\max_{k \neq j} \mathbb{E} \left[\left| X_{k, x_j}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2} \\
\text{and } F_{4, j, j'} & := \left(\max_{k \neq j} \mathbb{E} \left[\left| X_{k, x_j}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2} \left(\max_{k' \neq j'} \mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2}.
\end{aligned}$$

Similarly, we bound $\mathbb{E}[f_3^p]$, to arrive at

$$\mathbb{E}[f_3^p] \leq D_2 (F_{\nu_1, i, j, j'} + F_{\nu_2, i, j, j'} + F_{\nu_3, i, j, j'} + F_{4, j, j'}),$$

where

$$\begin{aligned}
F_{\nu_1, i, j, j'} &:= \|\nu_{i, x_j, x_{j'}}\|_\infty^p, \\
F_{\nu_2, i, j, j'} &:= \left(\sum_{k'=1}^d \|\nu_{i, x_j, x_{k'}}\|_\infty \right)^p \left(\max_{k' \neq j'} \mathbb{E} \left[|X_{k', x_{j'}}^{t, x}(\tau)|^{2p} \right] \right)^{1/2}, \\
\text{and } F_{\nu_3, i, j, j'} &:= \left(\sum_{k=1}^d \|\nu_{i, x_k, x_{j'}}\|_\infty \right)^p \left(\max_{k \neq j} \mathbb{E} \left[|X_{k, x_j}^{t, x}(\tau)|^{2p} \right] \right)^{1/2}.
\end{aligned}$$

Using this and the bound on the first variation, we can find $b_{i, j, j'}$ such that

$$\int_t^s \mathbb{E}[f_3^p] + (p-1)\mathbb{E}[f_4^{p/2}] d\tau \leq b_{i, j, j'}^p \leq D_2.$$

We use the previous inequality in (67) and simplify to arrive at

$$\begin{aligned}
\mathbb{E} \left[|X_{i, x_j, x_{j'}}^{t, x}(s)|^p \right] &\leq 2c_1 \int_t^s \mathbb{E} \left[|X_{i, x_j, x_{j'}}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + c_2 \int_t^s \max_{k \in \{1, 2, \dots, d\} \setminus \{j, j'\}} \mathbb{E} \left[|X_{k, x_j, x_{j'}}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + 4^{p-1} (p-1) a_{i, j}^p \int_t^s \mathbb{E} \left[|X_{j, x_j, x_{j'}}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + 4^{p-1} (p-1) a_{i, j'}^p \int_t^s \mathbb{E} \left[|X_{j', x_j, x_{j'}}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + b_{i, j, j'}^p.
\end{aligned} \tag{68}$$

Recall that

$$a_{i, j}^p := \|\nu_{i, x_j}\|_\infty^p + \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j}\|_\infty^2 \right)^{p/2}$$

and that for all i we have $a_{i, i} + \sum_{j=1}^d \max_{i \neq j} a_{i, j} \leq C_1$. Then, taking the maximum over all i, j , and j' in (68) and using Grönwall's inequality then taking the p 'th root, we get

$$\max_{i, j, j'} \mathbb{E} \left[\left(X_{i, x_j, x_{j'}}^{t, x}(s) \right)^p \right]^{1/p} \leq D_2. \tag{69}$$

Setting $j' = i$ and taking the maximum over $i \neq j$ in (68) yield

$$\begin{aligned}
\max_{i \neq j} \mathbb{E} \left[|X_{i, x_j, x_i}^{t, x}(s)|^p \right] &\leq 2c_1 \int_t^s \max_{i \neq j} \mathbb{E} \left[|X_{i, x_j, x_i}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + c_2 \int_t^s \max_{i \neq j} \mathbb{E} \left[|X_{i, x_j, x_i}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + 4^{p-1} (p-1) \left(\max_{i \neq j} a_{i, j}^p \right) \int_t^s \max_{i, j'} \mathbb{E} \left[|X_{j', x_{j'}, x_i}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + 4^{p-1} (p-1) C_1^p \int_t^s \max_{i \neq j} \mathbb{E} \left[|X_{i, x_j, x_i}^{t, x}(\tau)|^p \right] d\tau \\
&\quad + \max_{i \neq j} b_{i, j, i}^p.
\end{aligned}$$

Using Grönwall's inequality, then taking the p 'th root and summing over j , we get

$$\begin{aligned}
\sum_{j=1}^d \max_{i \neq j} \mathbb{E} \left[|X_{i, x_i, x_j}^{t, x}(s)|^p \right]^{1/p} &\leq D_2 \sum_{j=1}^d \left(\max_{i \neq j} a_{i, j} \right) + D_2 \sum_{j=1}^d \left(\max_{i \neq j} b_{i, j, i} \right) \\
&\leq D_2 C_1 + D_2,
\end{aligned} \tag{70}$$

where we used (as we will later show) that

$$\sum_{j=1}^d \left(\max_{i \neq j} b_{i, j, i} \right) + \sum_{j=1}^d \sum_{j'=1}^d \left(\max_{i \notin \{j, j'\}} b_{i, j, j'} \right) \leq D_2 \tag{71}$$

Finally, taking the maximum over $i \notin \{j, j'\}$ in (68),

$$\begin{aligned}
\max_{i \notin \{j, j'\}} \mathbb{E} \left[\left| X_{i, x_j, x_{j'}}^{t, x}(s) \right|^p \right] &\leq 2c_1 \int_t^s \max_{i \notin \{j, j'\}} \mathbb{E} \left[\left| X_{i, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] d\tau \\
&+ c_2 \int_t^s \max_{k \notin \{j, j'\}} \mathbb{E} \left[\left| X_{k, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] d\tau \\
&+ 4^{p-1} (p-1) \max_{i \neq j} a_{i, j}^p \int_t^s \mathbb{E} \left[\left| X_{j, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] d\tau \\
&+ 4^{p-1} (p-1) \max_{i \neq j'} a_{i, j'}^p \int_t^s \mathbb{E} \left[\left| X_{j', x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] d\tau \\
&+ b_{i, j, j'}^p.
\end{aligned}$$

Then, using Grönwall's inequality, taking the p 'th root and summing over j and j' , we get

$$\begin{aligned}
&\sum_{j=1}^d \sum_{j'=1}^d \max_{i \notin \{j, j'\}} \mathbb{E} \left[\left| X_{i, x_j, x_{j'}}^{t, x}(s) \right|^p \right]^{1/p} \\
&\leq D_2 \left(\sum_{j=1}^d \max_{i \neq j} a_{i, j} \right) \left(\max_j \sum_{j'=1}^d \sup_{t \leq \tau \leq s} \mathbb{E} \left[\left| X_{j, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right]^{1/p} \right) \\
&+ D_2 \left(\sum_{j'=1}^d \max_{i \neq j'} a_{i, j'} \right) \left(\max_{j'} \sum_{j=1}^d \sup_{t \leq \tau \leq s} \mathbb{E} \left[\left| X_{j', x_j, x_{j'}}^{t, x}(\tau) \right|^p \right]^{1/p} \right) \\
&+ D_2 \sum_{j=1}^d \sum_{j'=1}^d \left(\max_{i \notin \{j, j'\}} b_{i, j, j'} \right).
\end{aligned}$$

The result follows by (71), and since $\sum_{j=1}^d \max_{i \neq j} a_{i, j} \leq C_1$ and by (69) and (70),

$$\begin{aligned}
&\max_j \sum_{j'=1}^d \sup_{t \leq \tau \leq s} \mathbb{E} \left[\left| X_{j, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] \\
&\leq \max_j \sup_{t \leq \tau \leq s} \mathbb{E} \left[\left| X_{j, x_j, x_j}^{t, x}(\tau) \right|^p \right] + \sum_{j'=1}^d \max_{j \neq j'} \sup_{t \leq \tau \leq s} \mathbb{E} \left[\left| X_{j, x_j, x_{j'}}^{t, x}(\tau) \right|^p \right] \\
&\leq D_2
\end{aligned}$$

It remains to show (71). We consider the terms $F_{\varsigma_u, i, j, j'}$ for $u \in \{1, 2, 3\}$ along with $F_{4, j, j'}$ and claim that $F_{\nu_u, i, j, j'}$ are treated similarly. First note that

$$\begin{aligned}
&\sum_{j=1}^d \max_{i \neq j} F_{\varsigma_1, i, i, j} + \sum_{j=1}^d \sum_{j'=1}^d \max_{i \notin \{j, j'\}} F_{\varsigma_1, i, j, j'} \\
&\leq \sum_{j=1}^d \max_{i \neq j} \left(\sum_{m=1}^{d'} \|Sim_{i, x_i, x_j}\|_\infty^2 \right)^{1/2} + \sum_{j=1}^d \sum_{j'=1}^d \max_{i \notin \{j, j'\}} \left(\sum_{m=1}^{d'} \|Sim_{x_j, x_{j'}}\|_\infty^2 \right)^{1/2} \\
&\leq C_2.
\end{aligned}$$

Next $F_{\varsigma_2, i, j, j'}$ and $F_{\varsigma_3, i, j, j'}$ are similar,

$$\begin{aligned}
\sum_{j=1}^d \max_{i \neq j} F_{\varsigma_2, i, i, j} &= \left(\max_{i \neq j} \sum_{k'=1}^d \left(\sum_{m=1}^{d'} \|Sim_{x_i, x_{k'}}\|_\infty^2 \right)^{1/2} \right) \\
&\times \left(\sum_{j=1}^d \max_{k' \neq j} \left(\mathbb{E} \left[\left| X_{k', x_j}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/(2p)} \right) \leq C_2 K_{1, 2p}
\end{aligned}$$

and

$$\begin{aligned} \sum_{j=1}^d \sum_{j'=1}^d \max_{i \notin \{j, j'\}} F_{\varsigma_2, i, j, j'} &= \left(\sum_{j=1}^d \max_{i \neq j} \sum_{k'=1}^d \left(\sum_{m=1}^{d'} \|\varsigma_{im, x_j, x_{k'}}\|_{\infty}^2 \right)^{1/2} \right) \\ &\quad \times \left(\sum_{j'=1}^d \max_{k' \neq j'} \left(\mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2p} \right) \leq C_2 K_{1, 2p} . \end{aligned}$$

Finally,

$$\begin{aligned} \sum_{j=1}^d \sum_{j'=1}^d F_{4, j, j'} &\leq \left(\sum_{j=1}^d \max_{k \neq j} \left(\mathbb{E} \left[\left| X_{k, x_j}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/2} \right) \\ &\quad \times \left(\sum_{j'=1}^d \max_{k' \neq j'} \left(\mathbb{E} \left[\left| X_{k', x_{j'}}^{t, x}(\tau) \right|^{2p} \right] \right)^{1/(2p)} \right) \leq K_{1, 2p}^2 . \end{aligned}$$

Third variation The result for $q = 3$ can be proven similarly. \square

Proposition B.3 (Bounds on derivatives of the value function). *Let u satisfy the Kolmogorov backward equation*

$$\begin{aligned} \frac{\partial u}{\partial t} + \mathcal{L}u &= 0 \\ \text{and } u(T, \mathbf{x}) &= g(\mathbf{x}) , \end{aligned} \tag{72}$$

where

$$\mathcal{L}u := \sum_{i=1}^d \nu_i \frac{\partial u}{\partial x_i} + \frac{1}{2} \sum_{i=1}^d \sum_{j=1}^d \left(\sum_{m=1}^{d'} \varsigma_{im} \varsigma_{jm} \right) \frac{\partial^2 u}{\partial x_i \partial x_j} .$$

and assume that the coefficients satisfy Assumption B.1 for some $q \in \{2, 3\}$. Then, for some constant D , depending only on C_q, t , and T , we have

$$|u(t, \cdot)|_{\overline{C}_b^q(\mathbb{R}^d)} \leq D |g|_{\overline{C}_b^q(\mathbb{R}^d)} .$$

Proof. First note that under Assumption B.1 for $q \geq 2$, u satisfies for all $t \in [0, T]$ and $\mathbf{x} \in \mathbb{R}^d$ [11, Theorem 6.1 in Chapter 5]

$$u(t, \mathbf{x}) = \mathbb{E} \left[g(\mathbf{X}^{t, \mathbf{x}}(T)) \right] . \tag{73}$$

Additionally, we can differentiate u with respect to the initial conditions by formally taking the derivative inside the expectation in (73) [11, Theorem 5.5 in Chapter 5]. Hence, we bound

$$\begin{aligned} \sum_{j=1}^d |u_{x_j}(t, \mathbf{x})| &= \sum_{j=1}^d \left| \mathbb{E} \left[\sum_{i=1}^d g_{x_i}(\mathbf{X}^{t, \mathbf{x}}(T)) \cdot X_{i, x_j}^{t, \mathbf{x}}(T) \right] \right| \\ &\leq \sum_{j=1}^d \sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} \cdot \mathbb{E} \left[\left| X_{i, x_j}^{t, \mathbf{x}}(T) \right| \right] \\ &\leq \sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} \cdot \left(\sum_{j=1}^d \mathbb{E} \left[\left| X_{i, x_j}^{t, \mathbf{x}}(T) \right| \right] \right) \\ &\leq \left(\sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} \right) \left(\max_i \sum_{j=1}^d \mathbb{E} \left[\left| X_{i, x_j}^{t, \mathbf{x}}(T) \right| \right] \right) \\ &\leq K_{1, 2} \left(\sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} \right) \end{aligned}$$

by Proposition B.2. Similarly,

$$\begin{aligned}
\sum_{j=1}^d \sum_{j'=1}^d |u_{x_j, x_{j'}}(t, \mathbf{x})| &\leq \sum_{j=1}^d \sum_{j'=1}^d \left| \mathbb{E} \left[\sum_{i=1}^d g_{x_i}(\mathbf{X}^{t, \mathbf{x}}(T)) \cdot X_{i, x_j, x_{j'}}^{t, \mathbf{x}}(T) \right] \right| \\
&+ \sum_{j=1}^d \sum_{j'=1}^d \left| \mathbb{E} \left[\sum_{i=1}^d \sum_{i'=1}^d g_{x_i, x_{i'}}(\mathbf{X}^{t, \mathbf{x}}(T)) \cdot X_{i, x_j}^{t, \mathbf{x}}(T) \cdot X_{i', x_{j'}}^{t, \mathbf{x}}(T) \right] \right| \\
&\leq \left(\sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} \right) \left(\max_i \sum_{j=1}^d \sum_{j'=1}^d \mathbb{E} \left[|X_{i, x_j, x_{j'}}^{t, \mathbf{x}}(T)| \right] \right) \\
&+ \left(\sum_{i=1}^d \sum_{i'=1}^d \|g_{x_i, x_{i'}}\|_{C_b^0(\mathbb{R}^d)} \right) \left(\max_i \sum_{j=1}^d \mathbb{E} \left[\left(X_{i, x_j}^{t, \mathbf{x}}(T) \right)^2 \right]^{1/2} \right)^2 \\
&\leq (K_{1,2}^2 + K_{2,2}) \left(\sum_{i=1}^d \|g_{x_i}\|_{C_b^0(\mathbb{R}^d)} + \sum_{i=1}^d \sum_{i'=1}^d \|g_{x_i, x_{i'}}\|_{C_b^0(\mathbb{R}^d)} \right).
\end{aligned}$$

It is easy to see that the previous proof extends to $q = 3$.

□