

# NEW INTEGRAL FORMULAS AND IDENTITIES INVOLVING SPECIAL NUMBERS AND FUNCTIONS DERIVED FROM CERTAIN CLASS OF SPECIAL COMBINATORIAL SUMS

YILMAZ SIMSEK

**ABSTRACT.** By applying  $p$ -adic integral on the set of  $p$ -adic integers in [27] (Interpolation Functions for New Classes Special Numbers and Polynomials via Applications of  $p$ -adic Integrals and Derivative Operator, Montes Taurus J. Pure Appl. Math. 3 (1), ...-..., 2021 Article ID: MTJPAM-D-20-00000), we constructed generating function for the special numbers and polynomials involving the following combinatorial sum and numbers:

$$y(n, \lambda) = \sum_{j=0}^n \frac{(-1)^n}{(j+1)\lambda^{j+1}(\lambda-1)^{n+1-j}}$$

The aim of this paper is to use the numbers  $y(n, \lambda)$  to derive some new and novel identities and formulas associated with the Bernstein basis functions, the Fibonacci numbers, the Harmonic numbers, the alternating Harmonic numbers, binomial coefficients and new integral formulas for the Riemann integral. We also investigate and study on open problems involving the numbers  $y(n, \lambda)$  in [27]. Moreover, we give relation among the numbers  $y(n, \frac{1}{2})$ , the Digamma function, and the Euler constant. Finally, we give conclusions for the results of this paper with some comments and observations.

2010 MATHEMATICS SUBJECT CLASSIFICATION. 005A15, 11B68, 11B73, 11B83, 26C05, 11S40, 11S80, 33B15.

**KEYWORDS AND PHRASES.** Generating function, Special numbers and polynomials, Bernoulli-type numbers and polynomials, Fibonacci numbers, Harmonic numbers, Stirling numbers, Daehee numbers, Digamma function.

## 1. INTRODUCTION

Due to the works in [1] and [17], we see that Daniel Bernoulli and Goldbach were studied not only on interpolating a sequence, but also on the partial sums of the harmonic series. But these studies were not without much success. That is, Bernoulli and Goldbach were working the following partial sums of the harmonic series

$$f(v) = \sum_{j=1}^v \frac{1}{j}$$

where  $v$  is a positive integer. In one of his studies, Euler found many formulas on these topics involving the Digamma function, and the Euler constant, interpolating a sequence, partial sums of the harmonic series. We can find these clues in Euler's letter to Christian Goldbach dated October 13, 1729. In this letter, Euler mentioned the formulas he found and sent the following formula, one of them, to Goldbach without any proof:

$$f\left(\frac{1}{2}\right) = 2 - 2\log(2)$$

(cf. [17, p.136], [1, pp. 283–284]). In [17, p.136], [1, pp. 283–284], the following formula was given:

$$f\left(\frac{1}{2}\right) = \int_0^1 \frac{1 - \sqrt{x}}{1 - x} dx.$$

The motivation of this paper is of come from the spirit of historical mathematical works of Bernoulli, Euler and Goldbach with the help of applications of the following combinatorial numbers which was defined in [27]:

$$(1) \quad y(n, \lambda) = \sum_{j=0}^n \frac{(-1)^n}{(j+1)\lambda^{j+1}(\lambda-1)^{n+1-j}}$$

It would be very appropriate to give a brief information about how the numbers  $y(n, \lambda)$  can be found. By applying  $p$ -adic integral on the set of  $p$ -adic integers in [27], we constructed generating function for the special numbers and polynomials involving combinatorial sums and many special numbers and polynomials. When we investigated and studied on interpolation functions for these new classes special numbers and polynomials with series algebraic operations, we performed on one of the these series multiplications with its inspiration, the numbers  $y(n, \lambda)$  given in equation (1) were defined. In [27], some of their properties for these numbers were discussed and given.

In [27], we have put forward the following open problems for the numbers  $y(n, \lambda)$ :

- (1) *One of the first questions that comes to mind what is generating function for the numbers  $y(n, -1)$  and the numbers  $y(n, \lambda)$ .*
- (2) *Some of the other questions are what are the special families of numbers the numbers  $y(n, -1)$  are related to.*
- (3) *What are the combinatorial applications of the numbers  $y(n, -1)$ .*
- (4) *Can we find a special arithmetic function representing this family of numbers?*

Partial solutions to some of the above open problems can be explored and studied. By using (1), we can prove the following theorems including formulas and identities related to the numbers  $y(n, \lambda)$ , the harmonic numbers, the Fibonacci numbers, the Bernstein basis functions, and the Riemann integral:

**Theorem 1.1.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(2) \quad y(n, \lambda) = \frac{-1}{\lambda} \sum_{j=0}^n \sum_{k=0}^j \binom{n+1}{j} \frac{B_k S_1(j, k)}{j! B_j^{n+1}(\lambda)}.$$

where  $B_j^n(\lambda)$ ,  $B_j$ , and  $S_1(j, k)$  denote the Bernstein basis function and the Bernoulli numbers, and the Stirling numbers of the first kind, respectively.

**Theorem 1.2.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$y(n, \lambda) = \frac{(-1)^n}{(\lambda - 1)^{n+2}} \int_0^{\frac{\lambda-1}{\lambda}} \frac{1 - x^{n+1}}{1 - x} dx.$$

**Theorem 1.3.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(3) \quad y\left(n, \frac{1}{2}\right) = 2^{n+2} \left( H_{\lfloor \frac{n}{2} \rfloor} - H_n + \frac{(-1)^{n+1}}{n+1} \right).$$

where  $H_n$  denotes the harmonic numbers and  $H_0 = 0$ .

**Theorem 1.4.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(4) \quad y\left(n, \frac{1}{2}\right) = \frac{1}{2^{n+2}} \int_0^1 \frac{1 + (-1)^n x^{n+1}}{1 + x} dx.$$

**Theorem 1.5.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(5) \quad \int_0^1 \frac{1 - x^{\lfloor \frac{n}{2} \rfloor}}{1 - x} dx - \int_0^1 \frac{1 - x^n}{1 - x} dx = \frac{y\left(n, \frac{1}{2}\right)}{2^{n+2}} + \frac{(-1)^n}{n+1},$$

where  $\lfloor x \rfloor$  denotes the integer part of  $x$ .

**Theorem 1.6.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(6) \quad y\left(n, \frac{1 + \sqrt{5}}{2}\right) = \sum_{j=0}^n \frac{(-1)^{j+1}}{(j+1) \left( \frac{1+\sqrt{5}}{2} F_{j+1} + F_j \right) \left( \frac{1-\sqrt{5}}{2} F_{n-j+1} + F_{n-j} \right)},$$

where  $F_n$  denotes the Fibonacci numbers.

Combinatorial sums and numbers, special numbers and polynomials are the main topics that can be used in almost all areas of mathematics. In addition, they are also one of the most frequently used topics in applied sciences involving mathematical modeling and algorithmic solutions for real world problems.

We give the following some basic standard notations, formulas, and definitions:

Let  $\mathbb{N}$ ,  $\mathbb{Z}$ ,  $\mathbb{R}$ , and  $\mathbb{C}$  denote the set of natural numbers, the set of integer numbers, the set of real numbers and the set of complex numbers, respectively, and also  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$ . Supposing that for  $z \in \mathbb{C}$  with  $z = x + iy$  ( $x, y \in \mathbb{R}$ );  $Re(z) = x$

and  $Im(z) = y$  and also  $\log z$  denotes the principal branch of the many-valued function  $Im(\log z)$  with the imaginary part of  $\log z$  constrained by

$$-\pi < Im(\log z) \leq \pi,$$

and  $\log e = 1$ .

$$0^n = \begin{cases} 1, & n = 0 \\ 0, & n \in \mathbb{N} \end{cases}$$

The Bernoulli polynomials,  $B_n(x)$ , are defined by the following generating function:

$$(7) \quad \frac{t}{e^t - 1} e^{xt} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!},$$

where  $|t| < 2\pi$  (cf. [1]-[29]; and references therein).

When  $x = 0$  in (7), we have

$$B_n(0) = B_n$$

where  $B_n$  denotes the Bernoulli numbers (cf. [1]-[29]; and references therein).

The Bernstein basis function  $B_j^n(\lambda)$  is defined by

$$B_j^n(\lambda) = \binom{n}{j} \lambda^j (1 - \lambda)^{n-j}$$

where  $j \in \{0, 1, 2, \dots, n\}$  (cf. [12]).

The Stirling numbers of the second kind,  $S_1(n, k)$ , are defined by the following generating function:

$$(8) \quad \frac{(\log(1+z))^k}{k!} = \sum_{n=0}^{\infty} S_1(n, k) \frac{z^n}{n!}$$

with  $S_1(n, k) = 0$  if  $k > n$ , and  $k \in \mathbb{N}_0$  (cf. [1]-[29]; and references therein).

In [5] and [6], with the aid of the Volkenborn integral on the set of  $p$ -adic integers, Kim defined the Daehee numbers  $D_n$  by the following generating function:

$$\frac{\log(1+z)}{z} = \sum_{n=0}^{\infty} D_n \frac{z^n}{n!},$$

where  $z \neq 0$  and  $|z| < 1$  (cf. [5], [6]).

By using the above equation, the following explicit formula for the Daehee numbers  $D_n$  is found:

$$(9) \quad D_n = (-1)^n \frac{n!}{n+1}$$

(cf. [5], [6]).

Substituting  $t = \log(1+z)$ ,  $x = 0$ , and  $z \neq 0$  into (7), by using Riordan method given in [14, p. 45, Exercise 19 (b)], after some elementary calculations, we have

$$(10) \quad \log(1+z) = z \sum_{n=0}^{\infty} B_n \frac{(\log(1+z))^n}{n!}.$$

Combining the above equation with (8), after some elementary calculations,  $|z| < 1$  we get

$$\sum_{m=1}^{\infty} (-1)^{m+1} \frac{z^{m-1}}{m} = \sum_{m=0}^{\infty} \sum_{n=0}^m B_n S_1(m, n) \frac{z^m}{m!}$$

here we use  $S_1(m, n) = 0$  if  $n > m$ . After some elementary calculations, equating the coefficients  $z^m$  on both sides of the previous equation, one has the following well known novel formula, which has been proven by other different methods:

$$(11) \quad \sum_{n=0}^m B_n S_1(m, n) = \frac{(-1)^m m!}{m+1}$$

(cf. [5], [14, p. 45, Exercise 19 (b)]).

There are many other proof of (11). One of them was given Kim [5] with the  $p$ -adic invariant integral on the set of  $p$ -adic integers. On the other hand, the relation is given by equation the (11), Riordan [14, p. 45, Exercise 19 (b)] represented it by the notation  $(b)_n$ .

By using (9) and (11), in the next section, we give relations among the numbers  $(b)_n$ , the Daehee numbers  $D_n$ , and the numbers  $y(n, \lambda)$ .

**1.1. Further remarks and observations on the numbers  $y(n, \lambda)$ .** Substituting  $\lambda = 2$  into (1), in [27] we defined the following combinatorial numbers:

$$(12) \quad y(n) := y(n, 2) = \sum_{j=0}^n \frac{(-1)^j}{(j+1)2^{j+1}}.$$

In [27], we also give a relationship between the numbers  $y(n)$  and the  $\lambda$ -Apostol-Daehee numbers.

Putting  $\lambda = 1$  in (1), we have

$$(13) \quad y(n, -1) = \frac{1}{2(n+1)} \sum_{j=0}^n \frac{1}{\binom{n}{j}}$$

(cf. [27]).

**Remark 1.** Recently many authors have studied on binomial coefficients involving the following well-known sums and their applications:

$$\begin{aligned} & \sum_{j=0}^n \frac{1}{\binom{n}{j}}, \\ & \sum_{j=0}^n \frac{(-1)^j}{\binom{n}{j}}, \\ & \sum_{j=0}^n \binom{n}{j} j^k \end{aligned}$$

and so on (cf. [2], [14], [4], [9]-[11], [11][16], [18]-[27], [30]).

**Remark 2.** Most recently, in [25, Equations. (3), (13), and (14)], we constructed the following generating function for the numbers  $y_6(m, n; \lambda, p)$  involving finite sums involving higher powers of binomial coefficients

$${}_pF_{p-1} \left[ \begin{matrix} -n, -n, \dots, -n \\ 1, 1, \dots, 1 \end{matrix} ; (-1)^p \lambda e^t \right] = n! \sum_{m=0}^{\infty} y_6(m, n; \lambda, p) \frac{t^m}{m!},$$

where

$$\begin{aligned} n!y_6(0, n; \lambda, p) &= {}_pF_{p-1} \left[ \begin{matrix} -n, -n, \dots, -n \\ 1, 1, \dots, 1 \end{matrix} ; (-1)^p \lambda \right] \\ &= \sum_{k=0}^n \binom{n}{k}^p k^m \lambda^k, \end{aligned}$$

and  ${}_pF_p$  denotes the well known generalized hypergeometric function which is defined by

$${}_pF_q \left[ \begin{matrix} \alpha_1, \dots, \alpha_p \\ \beta_1, \dots, \beta_q \end{matrix} ; z \right] = \sum_{m=0}^{\infty} \left( \frac{\prod_{j=1}^p (\alpha_j)^{\overline{m}}}{\prod_{j=1}^q (\beta_j)^{\overline{m}}} \right) \frac{z^m}{m!}.$$

where the above series converges for all  $z$  if  $p < q + 1$ , and for  $|z| < 1$  if  $p = q + 1$ . Assuming that all parameters have general values, real or complex, except for the  $\beta_j$ ,  $j = 1, 2, \dots, q$  none of which is equal to zero or a negative integer and also

$$(\alpha)^{\overline{v}} = \prod_{j=0}^{v-1} (\alpha + j),$$

and  $(\lambda)^{\overline{0}} = 1$  for  $\lambda \neq 1$ , where  $v \in \mathbb{N}$ ,  $\lambda \in \mathbb{C}$ . For the generalized hypergeometric function and their applications see for details (cf. [8], [25], [31]; and references therein).

Here, we noting that (in future studies) the relationships between the numbers  $y(n, \lambda)$  and the numbers  $y_6(0, n; \lambda, p)$ , which have potential use in not only in mathematics but also in other areas, may also be investigated.

## 2. PROOFS OF MAIN THEOREMS

In this chapter, the proofs of the theorems from Theorem 1.1 to Theorem 1.6, as well as the new novel formulas and relations derived in the light of the results of these theorems are given.

*Proof of Theorem 1.1.* From (1), we get

$$(14) \quad y(n, \lambda) = \frac{1}{\lambda} \sum_{j=0}^n (-1)^{j+1} \frac{\binom{n+1}{j}}{(j+1)!} \frac{1}{B_j^{n+1}(\lambda)}.$$

Combining (14) with (11) and (9), after some elementary calculations, we arrive at the desired result.  $\square$

*Proof of Theorem 1.2.* By using (1), we give integral representation for the numbers  $y(n, \lambda)$ . We set

$$(15) \quad y(n, \lambda) = \frac{(-1)^n}{(\lambda-1)^{n+2}} \sum_{j=0}^n \int_0^{\frac{\lambda-1}{\lambda}} x^j dx.$$

By using the above equation, we also get

$$(16) \quad y(n, \lambda) = \frac{(-1)^n}{(\lambda-1)^{n+2}} \int_0^{\frac{\lambda-1}{\lambda}} \frac{1-x^{n+1}}{1-x} dx.$$

Combining (15) and (16), proof is completed.  $\square$

*proof of Theorem 1.3.* Putting  $\lambda = \frac{1}{2}$  in (1), we have

$$(17) \quad y\left(n, \frac{1}{2}\right) = 2^{n+2} \sum_{j=0}^n \frac{(-1)^{j+1}}{j+1}.$$

We know from Sofo's work [28, Eq. (1.5)] that the right-hand side of the previous equation gives us a formula of the following well known alternating harmonic numbers, which are related to the Riemann zeta function, the Polylogarithm (or de Jonquiere's) function, the Psi (or Digamma) function and the other special functions:

$$(18) \quad \sum_{j=1}^n \frac{(-1)^j}{j} = H_{[\frac{n}{2}]} - H_n.$$

Combining (17) with (18), proof of Theorem 1.3 is completed.  $\square$

*Proof of Theorem 1.4.* Using (17), we get

$$(19) \quad y\left(n, \frac{1}{2}\right) = -2^{n+2} \sum_{j=0}^n (-1)^j \int_0^1 x^j dx.$$

Combining (19) with the following well known formula

$$\sum_{j=0}^n (-x)^j = \frac{1 + (-1)^n x^{n+1}}{1 + x},$$

we obtain

$$y\left(n, \frac{1}{2}\right) = \frac{1}{2^{n+2}} \int_0^1 \frac{1 + (-1)^n x^{n+1}}{1 + x} dx.$$

This last equation shows us that the proof of the theorem completed.  $\square$

*Proof of Theorem 1.5.* Combining (3) with (20), we get

$$y\left(n, \frac{1}{2}\right) = 2^{n+2} \left( \int_0^1 \frac{1 - x^{\lfloor \frac{n}{2} \rfloor}}{1 - x} dx - \int_0^1 \frac{1 - x^n}{1 - x} dx + \frac{(-1)^{n+1}}{n+1} \right).$$

After some elementary calculations in the above equation, we arrive at the desired result.  $\square$

*Proof of Theorem 1.6.* Putting  $\lambda = \frac{1+\sqrt{5}}{2}$  into (1), we get

$$y\left(n, \frac{1+\sqrt{5}}{2}\right) = \sum_{j=0}^n \frac{(-1)^n}{(j+1) \left(\frac{1+\sqrt{5}}{2}\right)^{j+1} \left(\frac{1+\sqrt{5}}{2} - 1\right)^{n+1-j}}.$$

Combining the previous equation with the following well known identities involving the Fibonacci numbers:

$$\left(\frac{1+\sqrt{5}}{2}\right)^{j+1} = \frac{1+\sqrt{5}}{2} F_{j+1} + F_j$$

(cf. [7]), after some elementary calculations, we arrive at the desired result.  $\square$

### 3. FORMULAS FOR THE NUMBERS $y\left(n, \frac{1}{2}\right)$ DERIVED FROM MAIN THEOREMS

In this section we give some formulas which are derived from main theorems.

In [1], [17], and [28], the Harmonic numbers are given by the following integral representation:

$$(20) \quad H_n = \int_0^1 \frac{1 - x^n}{1 - x} dx,$$



and

$$(21) \quad H_n = -n \int_0^1 x^{n-1} \ln(1-x) dx = -n \int_0^1 (1-x)^{n-1} \ln(x) dx,$$

where  $H_0 = 0$ .

Combining (3) with (21), we arrive at the following theorem:

**Theorem 3.1.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$y\left(n, \frac{1}{2}\right) = 2^{n+2} \left( n \int_0^1 (1-x)^{n-1} \ln(x) dx - \left[\frac{n}{2}\right] \int_0^1 (1-x)^{\left[\frac{n}{2}\right]-1} \ln(x) dx \right) + \frac{(-1)^{n+1} 2^{n+2}}{n+1}.$$

We now give relation among the numbers  $y\left(n, \frac{1}{2}\right)$ , the Psi (or Digamma) function and the Euler constant.

The Euler's constant (or Euler-Mascheroni) constant is given by

$$\gamma = \lim_{m \rightarrow \infty} \left( -\ln(m) + \sum_{j=1}^m \frac{1}{j} \right)$$

and the Psi (or Digamma) function

$$\psi(z) = \frac{d}{dz} \{\log \Gamma(z)\},$$

where  $\Gamma(z)$  denotes the Euler gamma function, which is defined by

$$\Gamma(z) = \int_0^\infty t^{z-1} e^{-t} dt$$

where  $z = x + iy$  with  $x > 0$ . For  $z = n \in \mathbb{N}$ ,  $\Gamma(n+1) = n!$  (cf. [1], [17], [29]).

Sofo [28] gave the following formula:

$$(22) \quad H_n = \gamma + \psi(n+1)$$

where  $H_0 = 0$ .

Combining (3) with (22), we arrive at the following theorem:

**Theorem 3.2.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$y\left(n, \frac{1}{2}\right) = 2^{n+2} \left( \frac{(-1)^{n+1}}{n+1} + H_{\left[\frac{n}{2}\right]} - \gamma - \psi(n+1) \right).$$

Combining (14) with (11), we also arrive at the following result.

**Corollary 3.3.** *Let  $n \in \mathbb{N}_0$ . Then we have*

$$(23) \quad y(n, \lambda) = \frac{-1}{\lambda} \sum_{j=0}^n \binom{n+1}{j} \frac{(b)_j}{j! B_j^{n+1}(\lambda)}$$

or

$$(24) \quad y(n, \lambda) = \frac{-1}{\lambda} \sum_{j=0}^n \binom{n+1}{j} \frac{D_j}{j! B_j^{n+1}(\lambda)}.$$

#### 4. CONCLUSION

In this study, some properties of the numbers  $y(n, \lambda)$  with special finite sums containing these numbers have been found. Many new novel formulas and relations blended with well known special functions including the Euler gamma function, the Euler-Mascheroni constant, the Psi function, special polynomials including Bernstein basis functions, and special numbers including the Fibonacci number, the Bernoulli numbers, the Euler numbers, the Stirling numbers, and the Daehee numbers, those of known by the names of well-known mathematicians.

On the other hand, new formulas and integral representations of the numbers  $y(n, \lambda)$  have been proved by using integral representations of the harmonic numbers and the alternating harmonic numbers, which are based on very old studies and have rich properties and relationships, and formulas related to other special functions, involving the Digamma function, the Euler gamma function, and Euler constant.

In [27], we gave some open problems for the numbers  $y(n, \lambda)$ . In section 1, we've made some efforts to find solutions some of them throughout this study.

In your next studies, it is planned to investigate the solutions of the above open questions, including the numbers  $y(n, \lambda)$ .

#### REFERENCES

- [1] R. E. Bradley, C. E. Sandifer, *Leonhard Euler Life, Work and Legacy*, Elsevier Science 2007.
- [2] L. Comtet, *Advanced Combinatorics*. Dordrecht-Holland/ Boston-U.S.A.: D. Reidel Publication Company, 1974.
- [3] G. B. Djordjevic, G. V. Milovanovic, *Special classes of polynomials*, University of Nis, Faculty of Technology Leskovac, 2014.
- [4] H. W. Gould, *Combinatorial Numbers and Associated Identities: Table 1: Stirling Numbers*, Edited and Compiled by Jocelyn Quaintance May 3, 2010, <https://math.wvu.edu/~hgould/Vol.7.PDF>
- [5] D.S. Kim, T. Kim, *Daehee numbers and polynomials*, Appl. Math. Sci. (Ruse), **7 (120)**, 5969–5976, 2013.
- [6] T. Kim, *An invariant  $p$ -adic integral associated with Daehee numbers*, Integral Transforms Spec. Funct. **13** (2002), 65–69.
- [7] T. Koshy, *Fibonacci and Lucas Numbers with Applications*, Wiley-Interscience, 2001.
- [8] W. Koepf, *Hypergeometric Summation, An Algorithmic Approach to Summation and Special Function Identities*, second edition, Springer-Verlag, London, 2014.

- [9] I. Kucukoglu, *Implementation of computation formulas for certain classes of Apostol-type polynomials and some properties associated with these polynomials*, arXiv:2012.09208v1.
- [10] I. Kucukoglu, B. Simsek and Y. Simsek, *An approach to negative hypergeometric distribution by generating function for special numbers and polynomials*, Turk. J. Math. **43** (2019), 2337–2353.
- [11] I. Kucukoglu and Y. Simsek, *On a family of special numbers and polynomials associated with Apostol-type numbers and polynomials and combinatorial numbers*, Appl. Anal. Discrete Math., **13** (2019), 478–494.
- [12] Lorentz, G.G: *Bernstein Polynomials*. Chelsea, New York (1986)
- [13] G. Ozdemir, Y. Simsek, and G. V. Milovanović, *Generating functions for special polynomials and numbers including Apostol-type and Humbert-type polynomials*. Mediterr. J. Math. **14** (2017), 1–17.
- [14] J. Riordan, *Introduction to Combinatorial Analysis*, Dover Publications, 2002.
- [15] S. Roman, *The Umbral Calculus*, New York, Dover Publications, 2005.
- [16] G.-C. Rota, *The number of partitions of a set*, American Math. Monthly **71** (5), 498–504, 1964.
- [17] C. E. Sandifer, *How Euler Did Even More*, The Mathematical Association of America, Washington, 2014.
- [18] Y. Simsek, *A new combinatorial approach to analysis: Bernstein basis functions, combinatorial identities and Catalan numbers*, Math. Meth. Appl. Sci. **38** (14) (2015), 3007–3021.
- [19] Y. Simsek, *Beta-Type Polynomials And Their Generating Functions*, Appl. Math. Comput. **254** (2015), 172–182.
- [20] Y. Simsek, *Combinatorial sums and binomial identities associated with the Beta-type polynomials*, Hacet. J. Math. Stat. **47** (5) (2018), 1144–1155.
- [21] Y. Simsek, *Computation methods for combinatorial sums and Euler-type numbers related to new families of numbers*, Math. Meth. Appl. Sci. **40** (7) (2017), 2347–2361.
- [22] Y. Simsek, *On generating functions for the special polynomials*, Filomat **31** (1) (2017), 9–16.
- [23] Y. Simsek, *New families of special numbers for computing negative order Euler numbers and related numbers and polynomials*, Appl. Anal. Discrete Math. **12** (2018), 1–35.
- [24] Y. Simsek, *Combinatorial identities and sums for special numbers and polynomials*, Filomat **32** (20) (2018), 6869–6877.
- [25] Simsek, Y. *Generating functions for finite sums involving higher powers of binomial coefficients: Analysis of hypergeometric functions including new families of polynomials and numbers*. J. Math. Anal. Appl. **477** (2019), 1328–1352.
- [26] Y. Simsek, *Explicit formulas for  $p$ -adic integrals: Approach to  $p$ -adic distributions and some families of special numbers and polynomials*, Montes Taurus J. Pure Appl. Math. Article ID: MTJPAM-D-19-00005, **1** (1) (2019), 1–76.
- [27] Y. Simsek, *Interpolation Functions for New Classes Special Numbers and Polynomials via Applications of  $p$ -adic Integrals and Derivative Operator*, Montes Taurus J. Pure Appl. Math. **3** (1) (2021), 1–24 Article ID: MTJPAM-D-20-00000
- [28] A. Sofo, *Quadratic alternating harmonic number sums*, J. Number Theory **154** (2015), 144–159.
- [29] H.M. Srivastava and J. Choi, *Zeta and  $q$ -Zeta Functions and Associated Series and Integrals*, Amsterdam, Elsevier Science Publishers, 2012.
- [30] B. Sury, T. Wang, and F.-Z. Zhao, *Some identities involving reciprocals of binomial coefficients*. J. Integer Sequences **7** (2004), Article 04.2.8.
- [31] R. Tremblay, B.J. Fugere, *Products of two restricted hypergeometric functions*, J. Math. Anal. Appl. **198**(3) (1996), 844–852.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE UNIVERSITY OF AKDENIZ TR-  
07058 ANTALYA-TURKEY

*Email address:* `ysimsek@akdeniz.edu.tr`