arXiv:2101.02248v2 [math.GM] 9 Apr 2021

Average Orders of the Euler Phi Function, The Dedekind Psi Function, The Sum of Divisors Function, And The Largest Integer Function

N. A. Carella

***Abstract*:** Let $x \geq 1$ be a large number, let $[x] = x - \{x\}$ be the largest integer function, and let $\varphi(n)$ be the Euler totient function. The result $\sum_{n\leq x} \varphi([x/n]) = (6/\pi^2)x\log x + O\left(x(\log x)^{2/3}(\log\log x)^{1/3}\right)$ was proved very recently. This note presents a short elementary proof, and sharpen the error term to $\sum_{n\leq x} \varphi([x/n]) = (6/\pi^2)x\log x + O(x)$. In addition, the first proofs of the asymptotics formulas for the finite sums $\sum_{n\leq x} \psi([x/n]) = (15/\pi^2)x\log x + O(x\log\log x)$, and $\sum_{n\leq x} \sigma([x/n]) = (\pi^2/6)x\log x + O(x\log\log x)$ are also evaluated here.

Contents

1 Introduction **2**

2 Euler Totient Function **2**

3 Auxiliary Results for the Phi Function **3**

4 Dedekind Psi Function **5**

5 Auxiliary Results for the Psi Function **6**

6 Sum of Divisors Function **8**

7 Auxiliary Results for the Sigma Function **8**

8 Numerical Data **10**

November 13, 2021
MSC2020: Primary 11N37, Secondary 11N05.
Keywords: Arithmetic function; Euler phi function; Dedekind psi function, Sum of divisors function; Average orders.

1 Introduction

Some new analytic techniques for evaluating the *fractional* finite sums $\sum_{n\leq x} f([x/n])$ for slow growing functions $f(n) \ll n^{\varepsilon}$, were recently introduced in [2]. Subsequently, the some analytic techniques for faster growing functions were introduced in the more recent literature as [4]. In this note, the standard analytic techniques originally developed for evaluating the average orders $\sum_{n\leq x} f(n)$ of arithmetic functions are modified to handle the fractional finite sums $\sum_{n\leq x} f([x/n])$ for multiplicative functions defined by Dirichlet convolutions $f(n) = \sum_{d|n} g(n/d)$, where $f, g : \mathbb{N} \longrightarrow \mathbb{C}$, and the fast rates of growth approximately $f(n) \gg n(\log n)^b$, for some $b \in \mathbb{Z}$. The modified standard techniques are simpler, more efficient and produce very short proofs.

As demonstrations, the fractional finite sum $\sum_{n\leq x} \varphi([x/n])$ of the Euler phi function φ in Theorem 2.1, the fractional finite sum $\sum_{n\leq x} \psi([x/n])$ of the Dedekind psi function ψ in Theorem 4.1, and the fractional finite sum $\sum_{n\leq x} \sigma([x/n])$ of the sum of divisors function σ in Theorem 6.1, are evaluated here. The three functions $\varphi(n) \leq \psi(n) \leq \sigma(n)$ share many similarities such as multiplicative structures, rates of growths, are defined by some Dirichlet convolutions $f(n) = \sum_{d|n} g(n/d)$, et cetera, and have similar proofs. Theorem 2.1 has a very short proof, and sharpen the error term of a very recent result $\sum_{n\leq x} \varphi([x/n]) = (6/\pi^2)x\log x + O\left(x(\log x)^{2/3}(\log\log x)^{1/3}\right)$ proved in [4] using a very complicated and lengthy proof. Further, Theorem 4.1, and Theorem 6.1 are new results in the literature.

2 Euler Totient Function

The Euler totient function $\varphi(n) = n\sum_{d|n} \mu(d)/d$ is multiplicative and satisfies the growth condition $\varphi(n) \gg n/\log\log n$. A very short proof for $\sum_{n\leq x} \varphi([x/n])$ is produced here. It is a modified version of the standard proof for the average order $\sum_{n\leq x} \varphi(n) = (3/\pi^2)x^2 + O(x\log x)$, which appears in [1, Theorem 3.7], and similar references.

Theorem 2.1. *If $x \geq 1$ is a large number, then,*

$$\sum_{n\leq x} \varphi\left(\left[\frac{x}{n}\right]\right) = \frac{6}{\pi^2} x\log x + O(x). \tag{1}$$

Proof. Use the identity $\varphi(n) = n\sum_{d|n} \mu(d)/d$ to rewrite the finite sum, and switch the order of summation:

$$\begin{aligned} \sum_{n\leq x} \varphi\left(\left[\frac{x}{n}\right]\right) &= \sum_{n\leq x} \left[\frac{x}{n}\right] \sum_{d\,|\,[x/n]} \frac{\mu(d)}{d} \\ &= \sum_{d\leq x} \frac{\mu(d)}{d} \sum_{\substack{n\leq x \\ d\,|\,[x/n]}} \left[\frac{x}{n}\right]. \end{aligned} \tag{2}$$

Apply Lemma 3.1 to remove the congruence on the inner sum index, and break it up into

two subsums. Specifically,

$$\begin{aligned}\sum_{d\le x}\frac{\mu(d)}{d}\sum_{\substack{n\le x\\ d\,|\,[x/n]}}\left[\frac{x}{n}\right] &= \sum_{d\le x}\frac{\mu(d)}{d}\sum_{n\le x}\left[\frac{x}{n}\right]\cdot\frac{1}{d}\sum_{0\le a\le d-1}e^{i2\pi a[x/n]/d} \\ &= \sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right]+\sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right]\sum_{0<a\le d-1}e^{i2\pi a[x/n]/d} \\ &= M(x) \quad + \quad E(x).\end{aligned} \tag{3}$$

The main term $M(x)$ is computed in Lemma 3.2 and the error term $E(x)$ is computed in Lemma 3.3. Summing these expressions complete the verification. ■

It is easy to verify that the expressions $M(x)$ and $E(x)$ imply the omega result

$$\sum_{n\le x}\varphi\left(\left[\frac{x}{n}\right]\right)-\frac{6}{\pi^2}x\log x=\Omega_{\pm}(x) \tag{4}$$

or a better result.

3 Auxiliary Results for the Phi Function

The detailed and elementary proofs of the preliminary results required in the proof of Theorem 2.1 concerning the Euler phi function $\varphi(n)=\sum_{d|n}\mu(d)d$ are recorded in this section.

Lemma 3.1. *Let $x\ge 1$ be a large number, and let $1\le d,n\le x$ be integers. Then,*

$$\frac{1}{d}\sum_{0\le a\le d-1}e^{i2\pi a[x/n]/d}=\begin{cases}1 & \text{if } d\mid[x/n],\\ 0 & \text{if } d\nmid[x/n],\end{cases} \tag{5}$$

Proof. The two cases $d\mid[x/n]$ and $d\nmid[x/n]$ are easily handled with the basic exponential sum

$$\sum_{0\le k\le q-1}e^{i2\pi km/q}=\begin{cases}q & \text{if } q\mid m,\\ 0 & \text{if } q\nmid m,\end{cases} \tag{6}$$

where $m\ne 0$, and $q\ge 1$ are integers. ■

Lemma 3.2. *Let $x\ge 1$ be a large number. Then,*

$$\sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right]=\frac{6}{\pi^2}x\log x+O\left(x\right). \tag{7}$$

Proof. Expand the bracket and evaluate the two subsums. Specifically,

$$\begin{aligned}\sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right] &= x\sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\frac{1}{n}-\sum_{d\le x}\frac{\mu(d)}{d^2}\sum_{n\le x}\left\{\frac{x}{n}\right\} \\ &= \frac{6}{\pi^2}x\log x+O\left(x\right)\end{aligned} \tag{8}$$

as claimed ■

Lemma 3.3. *Let $x \geq 1$ be a large number. Then,*

$$\sum_{d\leq x}\frac{\mu(d)}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{0<a\leq d-1}e^{i2\pi a[x/n]/d}=O\left(x\right). \tag{9}$$

Proof. As $n \leq x$ ranges over all its values, each value $m = [x/n] \leq x$ is repeated

$$\left[\frac{x}{n}\right]-\left[\frac{x}{n+1}\right]=\frac{x}{n(n+1)}+O(1) \tag{10}$$

times. Hence, reordering the triple sum $E(x)$, and substituting (10) yield

$$\begin{aligned} E(x) &= \sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{d\leq x}\frac{\mu(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi a[x/n]/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{d\leq x}\frac{\mu(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= S_0(x)+S_1(x). \end{aligned} \tag{11}$$

The finite subsums $S_0(x)$, and $S_1(x)$ correspond to the subsets of integers $n \leq x$ such that $d \mid [x/n]$, and $d \nmid [x/n]$, respectively.

Case I. The set of values $m = [x/n] \leq x$ such that $d \mid m$. Evaluating the indicator function returns

$$\begin{aligned} S_0(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d^2}\cdot(d-1) \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\left(-\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d^2}+\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d}\right). \end{aligned} \tag{12}$$

The first sum from right in (12) has the upper bound

$$\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d}=O(1), \tag{13}$$

this follows from [3, Theorem 5] and partial summation, or other summation methods. And the second sum from the right in (12) is represented by

$$\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d^2}=c_0(n)+O(1), \tag{14}$$

where $|c_0(n)|< 2$ depends on n. Replacing these expressions in (12) returns

$$\begin{aligned} S_0(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\left(c_0(n)+O(1)\right) \\ &= O\left(x\right). \end{aligned}$$

Case II. The set of values $m = [x/n] \leq x$ such that $d \nmid m$. Evaluating the indicator function returns

$$\begin{aligned} S_1(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\nmid m}}\frac{\mu(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\nmid m}}\frac{\mu(d)}{d^2}\cdot(-1) \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)(c_1(n)+O(1)) \\ &= O\left(x\right), \end{aligned} \tag{15}$$

where $|c_1(n)|< 2$ depends on n.

Summing the last two expressions yield $E(x) = S_0(x) + S_1(x) = O\left(x\right)$. ■

4 Dedekind Psi Function

The second result deals with the Dedekind function $\psi(n) = n\sum_{d|n}\mu(d)^2/d$. It is multiplicative and satisfies the growth condition $\psi(n) \gg n$. The first asymptotic formula for the fractional finite sum of the Dedekind function is given below.

Theorem 4.1. *If $x \geq 1$ is a large number, then,*

$$\sum_{n\leq x}\psi\left(\left[\frac{x}{n}\right]\right) = \frac{15}{\pi^2}x\log x + O\left(x\log\log x\right). \tag{16}$$

Proof. Use the identity $\psi(n) = n\sum_{d|n}\mu^2(d)/d$ to rewrite the finite sum, and switch the order of summation:

$$\begin{aligned} \sum_{n\leq x}\psi\left(\left[\frac{x}{n}\right]\right) &= \sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{d\,|\,[x/n]}\frac{\mu^2(d)}{d} \\ &= \sum_{d\leq x}\frac{\mu^2(d)}{d}\sum_{\substack{n\leq x\\ d\,|\,[x/n]}}\left[\frac{x}{n}\right]. \end{aligned} \tag{17}$$

Apply Lemma 3.1 to remove the congruence on the inner sum index, and break it up into two subsums. Specifically,

$$\begin{aligned} \sum_{d\leq x}\frac{\mu^2(d)}{d}\sum_{\substack{n\leq x\\ d\,|\,[x/n]}}\left[\frac{x}{n}\right] &= \sum_{d\leq x}\frac{\mu^2(d)}{d}\sum_{n\leq x}\left[\frac{x}{n}\right]\cdot\frac{1}{d}\sum_{0\leq a\leq d-1}e^{i2\pi a[x/n]/d} \\ &= \sum_{d\leq x}\frac{\mu^2(d)}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]+\sum_{d\leq x}\frac{\mu^2(d)}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{0<a\leq d-1}e^{i2\pi a[x/n]/d} \\ &= M_1(x) \quad + \quad E_1(x). \end{aligned} \tag{18}$$

The main term $M_1(x)$ is computed in Lemma 5.1 and the error term $E_1(x)$ is computed in Lemma 5.2. Summing these expressions complete the verification. ■

It is easy to verify that the subsums $M_1(x)$ and $E_1(x)$ imply the omega result

$$\sum_{n\le x}\psi\left(\left[\frac{x}{n}\right]\right)-\frac{15}{\pi^2}x\log x=\Omega_{\pm}(x) \tag{19}$$

or a better result. A sketch of the standard proof for the average order

$$\sum_{n\le x}\psi(n)=\frac{15}{\pi^2}x^2+O\left(x\log x\right), \tag{20}$$

appears in [1, Exercise 13, p. 71].

5 Auxiliary Results for the Psi Function

The detailed and elementary proofs of the preliminary results required in the proof of Theorem 4.1 concerning the Dedekind psi function $\psi(n)=\sum_{d|n}\mu^2(d)d$ are recorded in this section.

Lemma 5.1. *Let $x\ge 1$ be a large number. Then,*

$$\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right]=\frac{15}{\pi^2}x\log x+O\left(x\right). \tag{21}$$

Proof. Expand the bracket and evaluate the two subsums. Specifically,

$$\begin{aligned}\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right] &= x\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{n\le x}\frac{1}{n}-\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{n\le x}\left\{\frac{x}{n}\right\} \\ &= \frac{15}{\pi^2}x\log x+O\left(x\right)\end{aligned} \tag{22}$$

as claimed ■

Lemma 5.2. *Let $x\ge 1$ be a large number. Then,*

$$\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{n\le x}\left[\frac{x}{n}\right]\sum_{0<a\le d-1}e^{i2\pi a[x/n]/d}=O\left(x\log\log x\right). \tag{23}$$

Proof. As $n\le x$ ranges over all its values, each value $m=[x/n]\le x$ is repeated

$$\left[\frac{x}{n}\right]-\left[\frac{x}{n+1}\right]=\frac{x}{n(n+1)}+O(1) \tag{24}$$

times. Hence, reordering the triple sum $E(x)$, and substituting (24) yield

$$\begin{aligned}E_1(x) &= \sum_{n\le x}\left[\frac{x}{n}\right]\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{0<a\le d-1}e^{i2\pi a[x/n]/d} \\ &= \sum_{n\le x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{d\le x}\frac{\mu^2(d)}{d^2}\sum_{0<a\le d-1}e^{i2\pi am/d} \\ &= S_{10}(x)+S_{11}(x).\end{aligned} \tag{25}$$

The finite subsums $S_{10}(x)$, and $S_{11}(x)$ correspond to the subsets of integers $n\le x$ such that $d\mid[x/n]$, and $d\nmid[x/n]$, respectively.

Case I. The set of values $m = [x/n] \leq x$ such that $d \mid m$. Evaluating the indicator function returns

$$\begin{aligned} S_{10}(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu^2(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu^2(d)}{d^2}\cdot(d-1) \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\left(-\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu^2(d)}{d^2}+\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu^2(d)}{d}\right). \end{aligned} \tag{26}$$

The first sum from right in (26) has the upper bound

$$\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu^2(d)}{d}\leq\frac{\psi(m)}{m}=O(\log\log x), \tag{27}$$

since $m \leq x$. And the second sum from the right in (26) is represented by

$$\sum_{\substack{d\leq x\\ d\mid m}}\frac{\mu(d)}{d^2}=c_0(n)+O(1), \tag{28}$$

where $|c_{10}(n)|< 2$ depends on n. Replacing these expressions in (26) returns

$$\begin{aligned} S_{10}(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)(c_{10}(n)+O(\log\log x)) \\ &= O\left(x\log\log x\right). \end{aligned} \tag{29}$$

Case II. The set of values $m = [x/n] \leq x$ such that $d \nmid m$. Evaluating the indicator function returns

$$\begin{aligned} S_{11}(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\nmid m}}\frac{\mu^2(d)}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\nmid m}}\frac{\mu^2(d)}{d^2}\cdot(-1) \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)(c_{11}(n)+O(1)) \\ &= O\left(x\right), \end{aligned} \tag{30}$$

where $|c_{11}(n)|< 2$ depends on n.

Summing the last two expressions yield $E_1(x) = S_{10}(x) + S_{11}(x) = O\left(x\log\log x\right)$. ■

6 Sum of Divisors Function

The third result deals with the sum of divisors function $\sigma(n) = n\sum_{d|n} 1/d$. It is multiplicative and satisfies the growth condition $\sigma(n) \gg n$. The first asymptotic formula for the fractional sum of divisor function is given below.

Theorem 6.1. *If $x \geq 1$ is a large number, then,*

$$\sum_{n\leq x} \sigma\left(\left[\frac{x}{n}\right]\right) = \frac{\pi^2}{6} x\log x + O\left(x\log\log x\right). \tag{31}$$

Proof. Use the identity $\sigma(n) = n\sum_{d|n} 1/d$ to rewrite the finite sum, and switch the order of summation:

$$\begin{aligned}\sum_{n\leq x} \sigma\left(\left[\frac{x}{n}\right]\right) &= \sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{d\,|\,[x/n]}\frac{1}{d} \\ &= \sum_{d\leq x}\frac{1}{d}\sum_{\substack{n\leq x\\ d\,|\,[x/n]}}\left[\frac{x}{n}\right].\end{aligned} \tag{32}$$

Apply Lemma 3.1 to remove the congruence on the inner sum index, and break it up into two subsums. Specifically,

$$\begin{aligned}\sum_{d\leq x}\frac{1}{d}\sum_{\substack{n\leq x\\ d\,|\,[x/n]}}\left[\frac{x}{n}\right] &= \sum_{d\leq x}\frac{1}{d}\sum_{n\leq x}\left[\frac{x}{n}\right]\cdot\frac{1}{d}\sum_{0\leq a\leq d-1} e^{i2\pi a[x/n]/d} \\ &= \sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right] + \sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{0<a\leq d-1} e^{i2\pi a[x/n]/d} \\ &= M_2(x) \quad + \quad E_2(x).\end{aligned} \tag{33}$$

The main term $M_2(x)$ is computed in Lemma 7.1 and the error term $E_2(x)$ is computed in Lemma 7.2. Summing these expressions complete the verification. ■

It is easy to verify that the subsums $M_2(x)$ and $E_2(x)$ imply the omega result

$$\sum_{n\leq x} \sigma\left(\left[\frac{x}{n}\right]\right) - \frac{\pi^2}{6} x\log x = \Omega_{\pm}(x) \tag{34}$$

or a better result. The standard proof for the average order

$$\sum_{n\leq x}\sigma(n) = \frac{\pi^2}{12}x^2 + O\left(x\log x\right), \tag{35}$$

appears in [1, Theorem 3.4].

7 Auxiliary Results for the Sigma Function

The detailed and elementary proofs of the preliminary results required in the proof of Theorem 6.1 concerning the sum of divisor function $\sigma(n) = \sum_{d|n} d$ are recorded in this section.

Lemma 7.1. *Let $x \geq 1$ be a large number. Then,*

$$\sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]=\frac{\pi^2}{6}x\log x+O\left(x\right). \tag{36}$$

Proof. Expand the bracket and evaluate the two subsums. Specifically,

$$\begin{aligned}\sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right] &= x\sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\frac{1}{n}-\sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left\{\frac{x}{n}\right\} \\ &= \frac{\pi^2}{6}x\log x+O\left(x\right)\end{aligned} \tag{37}$$

as claimed ■

Lemma 7.2. *Let $x \geq 1$ be a large number. Then,*

$$\sum_{d\leq x}\frac{1}{d^2}\sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{0<a\leq d-1}e^{i2\pi a[x/n]/d}=O\left(x\log\log x\right). \tag{38}$$

Proof. As $n \leq x$ ranges over all its values, each value $m = [x/n] \leq x$ is repeated

$$\left[\frac{x}{n}\right]-\left[\frac{x}{n+1}\right]=\frac{x}{n(n+1)}+O(1) \tag{39}$$

times. Hence, reordering the triple sum $E_2(x)$, and substituting (39) yield

$$\begin{aligned}E_2(x) &= \sum_{n\leq x}\left[\frac{x}{n}\right]\sum_{d\leq x}\frac{1}{d^2}\sum_{0<a\leq d-1}e^{i2\pi a[x/n]/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{d\leq x}\frac{1}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= S_{20}(x)+S_{21}(x).\end{aligned} \tag{40}$$

The finite subsums $S_{20}(x)$, and $S_{21}(x)$ correspond to the subsets of integers $n \leq x$ such that $d \mid [x/n]$, and $d \nmid [x/n]$, respectively.

Case I. The set of values $m = [x/n] \leq x$ such that $d \mid m$. Evaluating the indicator function returns

$$\begin{aligned}S_{20}(x) &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{1}{d^2}\sum_{0<a\leq d-1}e^{i2\pi am/d} \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\sum_{\substack{d\leq x\\ d\mid m}}\frac{1}{d^2}\cdot(d-1) \\ &= \sum_{n\leq x}\left(\frac{x}{n(n+1)}+O(1)\right)\left(-\sum_{\substack{d\leq x\\ d\mid m}}\frac{1}{d^2}+\sum_{\substack{d\leq x\\ d\mid m}}\frac{1}{d}\right).\end{aligned} \tag{41}$$

The first sum from the right in (41) has the upper bound

$$\sum_{\substack{d\leq x\\ d\mid m}}\frac{1}{d}\leq\frac{\sigma(m)}{m}=O(\log\log x), \tag{42}$$

since $m \leq x$. And the second sum from the right in (41) is represented by

$$\sum_{\substack{d\leq x\\ d|m}} \frac{\mu(d)}{d^2} = c_0(n) + O(1), \tag{43}$$

where $|c_{20}(n)|< 2$ depends on n. Replacing these expressions in (41) returns

$$\begin{aligned} S_{20}(x) &= \sum_{n\leq x} \left(\frac{x}{n(n+1)} + O(1) \right) (c_{10}(n) + O(\log\log x)) \\ &= O\left(x \log\log x\right). \end{aligned} \tag{44}$$

Case II. The set of values $m = [x/n] \leq x$ such that $d \nmid m$. Evaluating the indicator function returns

$$\begin{aligned} S_{21}(x) &= \sum_{n\leq x} \left(\frac{x}{n(n+1)} + O(1) \right) \sum_{\substack{d\leq x\\ d\nmid m}} \frac{1}{d^2} \sum_{0<a\leq d-1} e^{i2\pi am/d} \\ &= \sum_{n\leq x} \left(\frac{x}{n(n+1)} + O(1) \right) \sum_{\substack{d\leq x\\ d\nmid m}} \frac{1}{d^2} \cdot (-1) \\ &= \sum_{n\leq x} \left(\frac{x}{n(n+1)} + O(1) \right) (c_{21}(n) + O(1)) \\ &= O\left(x\right), \end{aligned} \tag{45}$$

where $|c_{21}(n)|< 2$ depends on n.

Summing the last two expressions yield $E_2(x) = S_{20}(x) + S_{21}(x) = O\left(x \log\log x\right)$. ■

8 Numerical Data

Small numerical tables were generated by an online computer algebra system, the range of numbers $x \leq 10^5$ is limited by the wi-fi bandwidth. The error terms are defined by

$$E_1(x) = \sum_{n\leq x} \varphi\left(\left[\frac{x}{n}\right]\right) - \frac{6}{\pi^2} x \log x, \tag{46}$$

and

$$E_2(x) = \sum_{n\leq x} \sigma\left(\left[\frac{x}{n}\right]\right) - \frac{\pi^2}{6} x \log x, \tag{47}$$

respectively. All the calculations are within the predicted ranges $E_i(x) = O(x)$.

References

[1] Apostol, Tom M. *Introduction to analytic number theory*. Undergraduate Texts in Mathematics. Springer-Verlag, New York-Heidelberg, 1976.

[2] Olivier Bordelles, Randell Heyman, Igor E. Shparlinski. *On a sum involving the Euler function*, http://arXiv.org/abs/1808.00188.

Table 1: Numerical Data For $\sum_{n\leq x}\varphi([x/n])$.

x	$\sum_{n<x}\varphi([x/n])$	$6\pi^{-2}x\log x$	Error $E_1(x)$
10	17	14.00	3.00
100	275	279.96	−4.96
1000	4053	4199.41	146.41
10000	52201	55992.16	−3791.16
100000	673929	699901.94	−25972.94

Table 2: Numerical Data For $\sum_{n\leq x}\sigma([x/n])$.

x	$\sum_{n<x}\sigma([x/n])$	$6^{-1}\pi^2 x\log x$	Error $E_2(x)$
10	39	37.88	1.12
100	804	757.52	46.48
1000	12077	11362.80	714.20
10000	167617	151504.03	16112.97
100000	2033577	1893800.33	139776.67

[3] Montgomery, Hugh L.; Vaughan, Robert C. *Mean values of multiplicative functions.* Periodica Mathematica Hungarica volume 43, pages 199-214, (2002).

[4] Zhai, Wenguang. *On a sum involving the Euler function.* J. Number Theory 211 (2020), 199-219.

TotientAndDivisorsFunctions-04-09-21-553-arxiv.tex.