

No-boundary Wave Function, Wheeler-DeWitt Equation and Path Integral Analysis of the Bouncing ‘Quantum’ Cosmology

Karthik Rajeev^{*1}, Vikramaditya Mondal^{†1} and Sumanta Chakraborty^{‡1}

¹School of Physical Sciences

Indian Association for the Cultivation of Science, Kolkata-700032, India

Abstract

Bouncing models are alternatives to inflationary cosmology that replace the initial Big-Bang singularity by a ‘bouncing’ phase. A deeper understanding of the initial conditions of the universe, in these scenarios, requires knowledge of quantum aspects of bouncing models. In this work, we propose two classes of bouncing models that can be studied with great analytical ease and hence, provide test-bed for investigating more profound problems in quantum cosmology of bouncing universes. Our model’s two key ingredients enable us to do straightforward analytical calculations: (i) a convenient parametrization of the minisuperspace of FRLW spacetimes and (ii) two distinct choices of the effective perfect fluids that source the background geometry of the bouncing universe. We study the quantum cosmology of these models using both the Wheeler-de Witt equations and the path integral approach. In particular, we found a bouncing model analogue of the no-boundary wavefunction and presented a Lorentzian path integral representation for the same. We also discuss the introduction of real scalar perturbations.

^{*}krthkrajeev@gmail.com

[†]vikram.iisermail@gmail.com

[‡]sumantac.physics@gmail.com

Contents

1	Introduction	2
2	Mini-superspace model of cosmology: The general setup	3
3	Effective matter content of bouncing models: General analysis	6
4	The ground state wave function of the bouncing universe	10
4.1	The Wheeler-DeWitt equation and its solution	10
4.2	Path integral approach	12
4.2.1	The linear model	13
4.2.2	The quadratic model	15
4.3	Shear instability and its quantum analogue	19
5	Introducing matter field	22
5.1	Wheeler-DeWitt equation with matter and the WKB expansion	23
5.2	Classically allowed region	24
5.3	Classically forbidden region	26
6	Discussion	28

1 Introduction

The inflationary scenario gives rise to significant improvements to the Standard Big-Bang cosmology (SBC) [1–5]. Inflation is credited for solving several issues of SBC that include the horizon and flatness problems. The causal mechanism of structure formation that the theory of inflation offers has made it a proper science of precise and observationally verifiable predictions. Despite these desirable features, inflation is not devoid of serious conceptual challenges. For instance, if we consider the scalar field inflation in the context of Einstein gravity, it can be shown that there is an inevitable singularity before the onset of inflation [6, 7]. Another challenge to the inflationary models is the TransPlanckian issue [8, 9]. The term refers to the fact that if the universe expanded slightly over 65 e-folds during inflation, that is a little more than what is required to resolve certain issues of SBC, then one can show that the fluctuations which are within the Hubble radius today were originally at sub-Planckian length scales during the onset of inflation. The trans-Planckian problem challenges the validity of ‘quantum field theory in classical spacetime’ approximation which is at the heart of the analysis of perturbations in inflation. Deficiencies of the inflationary paradigm, such as these, have inspired the investigation of several viable alternative models for the early Universe.

‘Bouncing cosmology’ refers to a broad class of cosmological models characterized by a ‘bounce’, i.e., a smooth transition from a contracting to expanding phase. Bouncing models aim to resolve the fundamental problems of SBC without invoking inflation. For instance, by construction, bouncing cosmologies resolve the singularity problem as it replaces the singularity by a bounce. Bouncing models also avoids the TransPlanckian problem, since, in the bouncing scenario, a given length scale of fluctuation contracts to a non-zero minimum, thereby, assuring that all wavelengths of relevance to cosmology today have initially been far greater than the Planck length during the bounce. Nevertheless, bouncing models resolve the horizon problem, since, in a typical bouncing scenario, the sub-Hubble wavelengths of the seed fluctuation

modes exit the Hubble radius and then re-enter at a later time to emerge as a scale of current cosmological interest, just as in inflation (see, for instance, [10–12]). It is worth mentioning that there are challenges in bouncing models as well. For a recent review of progress and problems in bouncing models, the reader may consult [13].

Our aim, in this work, is to explore the quantum gravitational aspects of bouncing scenarios. There are, however, several models of bouncing cosmologies and most of them are conceptually disparate [14]. To see this, recall that the inevitable singularity that arises in homogeneous and isotropic cosmologies in Einstein gravity is a consequence of Hawking-Penrose singularity theorems. Hence, several ways of realizing a bounce can be viewed as essentially corresponding to the several ways of bypassing Hawking-Penrose singularity theorems’ assumptions. In order to realize a bouncing scenario, therefore, one might resort to any of the diverse models of unconventional physics which may include, among other things, concepts from modified gravity theories, string theory and loop quantum cosmology. Consequently, it is not easy to study the entire family of bouncing cosmologies in a single framework. Here, we adopt a simple phenomenological approach, which we can use to study a range of bouncing models of interest to theoretical cosmology.

Our starting point is the minisuperspace model with an effective potential $U_{eff}(a)$ for the scale factor. The functional form of $U_{eff}(a)$ is such that the scale factor’s classical dynamics correspond to a bouncing scenario. We introduce effective potential as a proxy for any of the unconventional physics that can accomplish a bounce. An obvious advantage of this approach is that we do not have to directly subscribe to any particular bounce model. It turns out that for several models a suitable $U_{eff}(a)$ may be chosen to capture its relevant features effectively. Our strategy then is to apply principles of quantum cosmology to study the quantum aspects of our model.

The paper’s structure is as follows: In Section 2, we briefly review the minisuperspace model of cosmology and introduce a convenient parametrization of the metric. In Section 3, we introduce two classes of perfect fluid densities that give rise to a wide range of bouncing cosmologies. We chose specific forms for these densities such that they generate effective potentials, that transform into a linear or quadratic form when rewritten in terms of a suitable variable. Quantum aspects of these models are then analyzed in Section 4 using the Wheeler-de Witt equation and a recent approach to quantum cosmology, for instance, prescribed in [15–17]. Followed by that, we introduce a massive scalar field conformally coupled to the background spacetime in Section 5. We conclude with the summary and discussion of our results in Section 6. (We shall henceforth work in units with $c = \hbar = 1$, unless otherwise specified.)

2 Mini-superspace model of cosmology: The general setup

We consider the mini-superspace of homogeneous and isotropic FLRW spacetimes with flat spatial sections. A convenient parametrization for this class of metrics is given by,

$$ds^2 = -\frac{\mathcal{N}^2(t)}{q(t)^p} dt^2 + q(t)^b |d\mathbf{x}|^2, \quad (1)$$

where p and b are both real numbers and are arbitrary at this point and shall later be fixed based on the necessity to have a quadratic action, to carry forward the analysis we hope to pursue here. It is easy to see that the scale factor is given by $a(t) = q(t)^{b/2}$. It must be emphasized that the time coordinate t , introduced above, is neither the cosmic time τ , nor the conformal time η . It is related to the cosmic time τ and the conformal time η through the following relations,

$$d\tau = \frac{\mathcal{N}(t)dt}{q(t)^{p/2}}; \quad d\eta = \frac{\mathcal{N}(t)dt}{q(t)^{(p+b)/2}}. \quad (2)$$

It we assume that the function $\mathcal{N}$ is a constant, then it is evident that as the parameters p and b take values such that $p + b = 0$ and $p = 0$, then the time coordinate t reduces to η and τ , respectively. For the metric ansatz presented in Eq. (1), the Einstein-Hilbert action takes the following form,

$$\mathcal{S}_{\text{EH}} = \int d^4x \sqrt{-g} R = V_3 \left(\frac{3b}{2} \right) \int dt q(t)^{\frac{3b+p-4}{2}} \left[\frac{1}{\mathcal{N}} (p + 2b - 2) \dot{q}^2 + 2 \frac{d}{dt} \left(\frac{1}{\mathcal{N}} q \dot{q} \right) - \frac{2}{\mathcal{N}} \dot{q}^2 \right], \quad (3)$$

where, the over-dot denotes derivative with respect to the time coordinate t and V_3 is the spatial volume¹. However, unless the action functional is quadratic in the dynamical variable, which corresponds to $q(t)$, it is difficult to employ the path-integral techniques. Thus, we demand that the following relation should hold: $3b + p = 4$. This condition means that the two independent parameters p and b in Eq. (1) boils down to a single parameter, which we choose to be b . For this choice, the Einstein-Hilbert action $\mathcal{S}_{\text{EH}}$, together with the Gibbons-Hawking-York boundary term, for a non-null boundary acquires the following form,

$$\begin{aligned} \mathcal{S}_{\text{EH}} &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} R - \frac{1}{\kappa} \int_{\text{Boundary}} d^3y \sqrt{h} K \\ &= \frac{V_3}{2\kappa} \int dt \left(-\frac{3b^2}{2\mathcal{N}} \right) \dot{q}^2 + 3bV_3 \left(\frac{1}{\mathcal{N}q\dot{q}} \right)_{\text{Boundary}} - \frac{1}{\kappa} \int_{\text{Boundary}} d^3y \sqrt{h} K \\ &= V_3 \int dt \left[-\frac{M}{2\mathcal{N}} \left(\frac{dq}{dt} \right)^2 \right], \end{aligned} \quad (4)$$

where, we have defined an “effective mass” $M \equiv \{(3b^2)/(2\kappa)\}$ with $\kappa = 8\pi G$. Note that, in the second line of the above equation, the extrinsic curvature terms, evaluated on the $t = \text{constant}$ boundary hypersurfaces, exactly cancels the total derivative term, thereby, reducing $\mathcal{S}_{\text{EH}}$ to the final expression. The parametrization of FRLW metric we presented in Eq. (1) contains several scenarios that have appeared in earlier works in the literature. These include the choice $b = 1/4$, which has been used to study de Sitter cosmology in [20] and more recently in [17] to describe the Lorentzian path integral approach to quantum cosmology. Additionally, for $b = 1/3$, the above parametrization has also appeared in [21] to rewrite the minisuperspace Lagrangian into a quadratic form. Finally, $b = 2$ corresponds to the standard representation of the metric in terms of the conformal time coordinate.

We have already described the Einstein-Hilbert part of the gravitational action along with the Gibbons-Hawking-York boundary term for the parametrization of the line element presented in Eq. (1). We will now introduce the matter sector in the picture. The most straightforward inclusion corresponds to the addition of a perfect fluid, with energy density ρ_{eff} and pressure $p_{\text{eff}} \propto \rho_{\text{eff}}$, for which the complete action describing gravity plus perfect fluid system is given by [22],

$$\begin{aligned} \mathcal{S}_{\text{perfect}} &= \frac{1}{2\kappa} \int d^4x \sqrt{-g} R - \frac{1}{\kappa} \int_{\text{Boundary}} d^3y \sqrt{h} K - \int d^4x \sqrt{-g} \rho_{\text{eff}}(q) \\ &= V_3 \int dt \left[-\frac{M}{2\mathcal{N}} \left(\frac{dq}{dt} \right)^2 + \mathcal{N} U_{\text{eff}}(q) \right], \end{aligned} \quad (5)$$

where, we have used Eq. (4) and have defined the effective ‘potential’ U_{eff} as,

$$U_{\text{eff}}(q) \equiv -q^{(3b-2)} \rho_{\text{eff}}(q). \quad (6)$$

¹For a non-compact flat FRLW universe, the spatial volume V_3 is infinite. We can, however, circumvent potential issues arising out of infinite V_3 by assuming that the spatial slices are locally flat but, *globally* compact, like a torus [18, 19]. The spacial volume V_3 , in such cases is finite.

As evident from Eq. (5), the Lagrangian resembles that of a point particle, with generalized coordinate q , moving in the effective potential U_{eff} that governs the dynamics of the background spacetime.²

Another possible addition to our ‘gravity-perfect fluid system’ is a non-minimally coupled scalar field of mass m , such that, the total action takes the following form,

$$\mathcal{S}_{\text{conformal}} = \frac{1}{2\kappa} \int d^4x \sqrt{-g} R - \frac{1}{\kappa} \int_{\text{Boundary}} d^3y \sqrt{h} K - \int d^4x \sqrt{-g} \rho_{\text{eff}}(q) + \int d^4x \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) - \frac{\xi}{2} R \phi^2 \right], \quad (7)$$

where, $\xi = 0$ corresponds to the minimally coupled scalar field and $\xi = (1/6)$ will yield the conformally coupled scalar field. Using the Fourier transform for the scalar field ϕ and using the fact that the scalar field is real, the scalar field action takes the following form,

$$\begin{aligned} \mathcal{S}_{\text{scalar}} &= \int d^4x \left[-\frac{1}{2} g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi - V(\phi) - \frac{\xi}{2} R \phi^2 \right] \\ &= \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \left[\frac{1}{2\mathcal{N}} q^2 |\dot{\phi}_{\mathbf{k}}|^2 - \frac{\mathcal{N}}{2} q^{2b-2} (|\mathbf{k}|^2 + m^2 q^b) |\phi_{\mathbf{k}}|^2 + \frac{3b^2\xi}{4\mathcal{N}} q^2 |\phi_{\mathbf{k}}|^2 + \frac{3b\xi}{2} \frac{1}{2\mathcal{N}} \frac{dq^2}{dt} \frac{d|\phi_{\mathbf{k}}|^2}{dt} \right] \\ &\quad - \frac{3b\xi}{2} \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{d}{dt} \left(\frac{1}{2\mathcal{N}} |\phi_{\mathbf{k}}|^2 \frac{dq^2}{dt} \right) \end{aligned} \quad (8)$$

Redefining, the scalar field as, $\phi_{\mathbf{k}} = \sqrt{V_3} q^c \tilde{\phi}_{\mathbf{k}}$, the above action can be expressed as,

$$\begin{aligned} \mathcal{S}_{\text{scalar}} &= V_3 \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \left[\frac{1}{2\mathcal{N}} q^{2+2c} |\dot{\tilde{\phi}}_{\mathbf{k}}|^2 + \frac{c+3b\xi}{4\mathcal{N}} q^{2c} \frac{dq^2}{dt} \frac{d|\tilde{\phi}_{\mathbf{k}}|^2}{dt} + \frac{c^2 + (3b^2\xi/2) + 6bc\xi}{2\mathcal{N}} q^{2c} q^2 |\tilde{\phi}_{\mathbf{k}}|^2 \right. \\ &\quad \left. - \frac{\mathcal{N}}{2} q^{2b-2+2c} (|\mathbf{k}|^2 + m^2 q^b) |\tilde{\phi}_{\mathbf{k}}|^2 \right] - \frac{3b\xi V_3}{2} \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{d}{dt} \left(\frac{1}{2\mathcal{N}} |\phi_{\mathbf{k}}|^2 \frac{dq^2}{dt} \right) \end{aligned} \quad (9)$$

Note that, if we wish to remove the second term in the matter action presented above then, the constant c must be chosen such that $c = -3b\xi$. In that case, the coefficient of the third term in the above action yields, $(b^2/2)(-18\xi^2 + 3\xi)$ and hence it can only vanish for $\xi = 0$ and $\xi = (1/6)$. Therefore, for conformally coupled scalar field, the action $\mathcal{S}_{\text{scalar}}$ simplifies considerably, such that the total action $\mathcal{S}_{\text{conformal}}$, presented in Eq. (7), takes the following form,

$$\begin{aligned} \mathcal{S}_{\text{conformal}} &= V_3 \int dt \left[-\frac{M}{2\mathcal{N}} \left(\frac{dq}{dt} \right)^2 + \mathcal{N} U_{\text{eff}}(q) \right] + V_3 \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \frac{d}{dt} \left(\frac{1}{2\mathcal{N}} \mathcal{S}_{\mathbf{k}} \right) \\ &\quad + V_3 \int dt \int \frac{d^3\mathbf{k}}{(2\pi)^3} \left[\frac{\mu(q)}{2\mathcal{N}} \left| \frac{d\tilde{\phi}_{\mathbf{k}}}{dt} \right|^2 - \frac{\mathcal{N}\mu(q)\omega_{\mathbf{k}}^2(q)}{2} |\tilde{\phi}_{\mathbf{k}}|^2 \right], \end{aligned} \quad (10)$$

where, the functions $\mu(q)$, $\omega_{\mathbf{k}}(q)$ and $\mathcal{S}_{\mathbf{k}}$ are defined as,

$$\mu(q) = q^{2-b}; \quad \omega_{\mathbf{k}}^2(q) = q^{2b-4} (k^2 + m^2 q^b); \quad \mathcal{S}_{\mathbf{k}} = -\frac{b}{4} |\phi_{\mathbf{k}}|^2 \frac{dq^2}{dt}. \quad (11)$$

²Although we have restricted ourselves to flat FRLW models, one can easily extend our analysis to closed as well as open FRLW cosmologies. In fact, one could imagine that $U_{\text{eff}}(q)$ contains a contribution of the form $-kq^{-2+2b} \propto \sqrt{-g} \frac{k}{a^2}$ (where $k = 0, \pm 1$ for flat, closed/open spatial slices), to account for the non-zero spatial curvature.

Thus, the action for the scalar field $\phi_{\mathbf{k}}$, as presented above, is also quadratic and each of the Fourier modes $\phi_{\mathbf{k}}$ depicts a harmonic oscillator with a time-dependent mass $\mu(q)$ and a time-dependent frequency $\omega_{\mathbf{k}}^2(q)$. Therefore, the equation of motion obtained by varying q and $\phi_{\mathbf{k}}$ are, respectively, given by:

$$M \frac{d^2 q}{dT^2} + \partial_q U_{\text{eff}}(q) = 0 \quad (12)$$

$$\frac{d}{dT} \left(\mu(q) \frac{d\phi_{\mathbf{k}}}{dT} \right) + \omega_{\mathbf{k}}^2(q) \phi_{\mathbf{k}} = 0 \quad (13)$$

where, $dT \equiv \mathcal{N}dt$. Furthermore, the momentum conjugate to q being $\pi_q = -(M/\mathcal{N})(dq/dt)$ and the momentum conjugate to $\tilde{\phi}_{\mathbf{k}}$ being $\pi_{\mathbf{k}} = (\mu(q)/2)(d\tilde{\phi}_{\mathbf{k}}/dt)$, which will be useful for later purposes. Out of the action functional, one can determine the Wheeler-DeWitt equation for the metric degree of freedom. The first step is to determine the Hamiltonian constraint, obtained by varying the action $\mathcal{S}_{\text{conformal}}$ with respect to the lapse function $\mathcal{N}$ and then setting $\mathcal{N} = 1$, which yields,

$$-\frac{MV_3}{2} \left(\frac{dq}{dt} \right)^2 - V_3 U_{\text{eff}}(q) + V_3 \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \left[\frac{\mu(q)}{2} \left(\frac{dQ_{\mathbf{k}}}{dt} \right)^2 + \frac{\mu(q)\omega_{\mathbf{k}}^2(q)}{2} Q_{\mathbf{k}}^2 \right] = 0. \quad (14)$$

where, we have defined:

$$Q_{\mathbf{k}} \equiv \begin{cases} \sqrt{2}\text{Re}[\tilde{\phi}_{\mathbf{k}}]; & k_z \geq 0 \\ \sqrt{2}\text{Im}[\tilde{\phi}_{\mathbf{k}}]; & k_z < 0 \end{cases} \quad (15)$$

Expressing $\dot{q}$ and $\dot{Q}_{\mathbf{k}}$ in terms of the conjugate momentum π_q and $\pi_{\mathbf{k}}$, along with promoting them to appropriate operators, such that $\pi_q \equiv -i\partial_q$ and $\pi_{\mathbf{k}} \equiv -i\partial_{Q_{\mathbf{k}}}$ (for ease of notation and since further confusion is unlikely to arise, we have henceforth renamed $Q_{\mathbf{k}}$ as $\phi_{\mathbf{k}}$), we obtain the following Wheeler-DeWitt equation,

$$\left\{ \frac{1}{2MV_3} \frac{\partial^2}{\partial q^2} - V_3 U_{\text{eff}}(q) + V_3 \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \left[-\frac{1}{2\mu(q)} \frac{\partial^2}{\partial \phi_{\mathbf{k}}^2} + \frac{\mu(q)\omega_{\mathbf{k}}^2(q)}{2} \phi_{\mathbf{k}}^2 \right] \right\} \Psi(q, \{\phi_{\mathbf{k}}\}) = 0, \quad (16)$$

where, $\Psi(q, \{\phi_{\mathbf{k}}\})$ is the wave function of the universe describing the evolution of the scale factor (through $q(t)$) and also the evolution of the matter field $\phi_{\mathbf{k}}$ in a coherent manner.

The first two terms of Eq. (16), which is just the purely gravitational part, governs the leading order quantum dynamics of the system considered here. The scalar field $\phi_{\mathbf{k}}$ is treated as a perturbation at the quantum level. As is evident from Eq. (16), the dynamics of the gravity sector is determined by the effective potential U_{eff} , which in turn is determined by the effective energy density ρ_{eff} of the perfect fluid system. In the next section, we will consider a family of perfect fluid models, with different ρ_{eff} , that can accomplish the bouncing scenario in the classical regime. Following that, we will discuss the associated quantum scenario.

3 Effective matter content of bouncing models: General analysis

In the previous section, we discussed the general aspects of minisuperspace model of cosmology consisting of a single dynamical variable $q(t)$. The ansatz for the spacetime metric was chosen such that the gravitational part of the action, including the Gibbons-Hawking-York boundary term, yields quadratic action for the

dynamical variable $q(t)$. In the matter sector, we considered a non-minimally coupled scalar field, and a perfect fluid with energy density ρ_{eff} . We also saw that the variable frequencies of the harmonic oscillator modes of the scalar field are functionals of q alone, if and only if, the coupling is either minimal or conformal. In this section, we will discuss the possible forms of the energy density ρ_{eff} , which is consistent with the quadratic nature of the action presented in Eq. (10).

For this purpose, consider the following behaviour for the effective energy density ρ_{eff} ,

$$\rho_{\text{eff}} = \rho_0 \left[\frac{c_1}{a^{n_1}} + \frac{c_2}{a^{n_2}} + \frac{c_3}{a^{n_3}} \right] , \quad (17)$$

where, ρ_0 is some energy density scale, c_1, c_2, c_3 are constants not necessarily positive, while n_1, n_2, n_3 are positive numbers, but not necessarily integers. Using the result, $a(t) = q(t)^{b/2}$, the effective energy density U_{eff} appearing in the action presented in Eq. (10), takes the following form,

$$U_{\text{eff}} = -q^{3b-2} \rho_{\text{eff}} = -\rho_0 \left[c_1 q^{3b-2-\frac{b}{2}n_1} + c_2 q^{3b-2-\frac{b}{2}n_2} + c_3 q^{3b-2-\frac{b}{2}n_3} \right] . \quad (18)$$

In order to work out the path integral arising out of the action dependent on U_{eff} , it is instructive to choose the powers n_1, n_2 and n_3 appropriately. Let us start by setting the coefficient of c_1 to be such that $3b - 2 - (b/2)n_1 = 0$, yielding,

$$b = \frac{4}{6 - n_1} ; \quad n_1 = \frac{2(3b - 2)}{b} . \quad (19)$$

Substituting the expression for b , presented above, in Eq. (18), the effective potential due to the matter field becomes

$$U_{\text{eff}} = -\rho_0 \left[c_1 + c_2 q^{\frac{2(n_1 - n_2)}{6 - n_1}} + c_3 q^{\frac{2(n_1 - n_3)}{6 - n_1}} \right] . \quad (20)$$

Setting the power of q in the term involving c_2 to identity and the power of q in the term involving c_3 to be quadratic, we obtain the powers n_2 and n_3 as a function of n_1 as,

$$U_{\text{eff}} = -\rho_0 [c_1 + c_2 q + c_3 q^2] ; \quad n_2 = \frac{3}{2}(n_1 - 2) ; \quad n_3 = 2n_1 - 6 . \quad (21)$$

On the other hand, we could have also taken n_2 to be fundamental and have expressed the powers n_1 and n_3 in terms of n_2 , which yield, $b = 6/(6 - n_2)$ and $n_1 = (2/3)(n_2 + 3)$ as well as $n_3 = (2/3)(2n_2 - 3)$. Similarly, choosing n_3 to be fundamental, one can express n_1 and n_2 in terms of it: $b = 8/(6 - n_3)$, $n_1 = (1/2)(n_3 + 6)$ and $n_2 = (3/2)\{1 + (n_3/2)\}$. Thus the above form for the energy density ρ_{eff} , presented in Eq. (17) with the choice of b given in Eq. (19) and the powers n_2 and n_3 given in Eq. (21), the action becomes quadratic in the variable $q(t)$ and thus path integral can be explicitly computed.

So far, our analysis has been fairly general. In what follows, we will be focussing on scenarios that give rise to ‘bounce’, by choosing the energy density ρ_{eff} appropriately. A generic bouncing scenario may be thought of as being sourced by two species of matter fields; one of them satisfying the energy condition, while the other violating the same. The latter is the responsible factor behind the bounce. Thus, in order to realise a bounce, $\rho_{\text{eff}}(a)$ should have components of both positive as well as negative energy densities which, in a generic situation, takes the following form,

$$\rho_{\text{eff}}(a) = \sum_n \frac{\rho_n^+}{a^n} - \sum_m \frac{\rho_m^-}{a^m} , \quad (22)$$

where ρ_n^+ and ρ_m^- as positive real numbers. The negative energy density terms could either be arising from an exotic matter field (as, for instance, in [23–25]) or emerging from the corrections to the Einstein-Hilbert action introduced by a UV-complete quantum gravity theory (as, for instance, in [26]).

If we assume that the bounce happens at sufficiently small values of the scale factor and keep in mind the fact that the leading order effects near the bounce are from the components violating the energy conditions, then in the simplest scenario, the effective energy density near the bounce is given by,

$$\rho_{\text{eff}}(a) \approx \frac{\rho_{n_+}^+}{a^{n_+}} - \frac{\rho_{n_-}^-}{a^{n_-}} , \quad (23)$$

with $n_- > n_+$. The powers n_+ and n_- are chosen such that the contribution to the action, arising out of the effective energy density ρ_{eff} , provides a quadratic action in the dynamical variable $q(t)$. Using the analysis leading to Eq. (21), the powers n_+ and n_- are chosen such that, we have the following two classes of effective energy densities:

$$\rho_{\text{eff}}^{(\text{I})}(a) = \rho_0 \left(\frac{1}{a^n} - \frac{1}{a^{\frac{(6+2n)}{3}}} \right) ; \quad b^{(\text{I})} = \frac{6}{6-n} , \quad (24)$$

$$\rho_{\text{eff}}^{(\text{II})}(a) = \rho_0 \left(\frac{1}{a^n} - \frac{1}{a^{\frac{(6+n)}{2}}} \right) ; \quad b^{(\text{II})} = \frac{8}{6-n} = \frac{4}{3}b^{(\text{I})} , \quad (25)$$

where, n and ρ_0 are real constants and, we have scaled the scale factor such that the bounce happens at $a = 1$, where ρ_{eff} identically vanishes. The necessary condition for both the effective energy densities to describe a bounce is $n < 6$, since then the power of a in the second term with negative coefficient will be less than that of the first term. It is worth mentioning that these two classes of energy densities together cover a wide range of physically relevant bouncing scenarios. For example, substituting $n = 3$ in the expression for $\rho_{\text{eff}}^{(\text{I})}(a)$, we obtain an effective energy density that describes a ‘matter bounce’ scenario (denoted by subscript ‘mb’) that may be realized, for instance, in the context of Horava-Lifshitz cosmology [26] or by invoking a non-canonical, ghost field [23]:

$$\rho_{\text{eff}}^{(\text{mb})}(a) = \rho_0 \left(\frac{1}{a^3} - \frac{1}{a^4} \right) . \quad (26)$$

Further, it is easy to see that the substitution of $n = 0$ in the first class of energy density, we obtain $\rho_{\text{eff}}(a) = \rho_0(1 - a^2)$, that is relevant for studying de Sitter spacetime in closed slicing [27–29]. As evident from Eq. (21), these two cases of energy densities correspond to $c_2 = 1 = -c_1$ with $c_3 = 0$ and $c_3 = 1 = -c_1$ with $c_2 = 0$, respectively. Thus in terms of the dynamical variable q , the corresponding effective potentials arising out of $\rho_{\text{eff}}^{(\text{I})}(a)$ and $\rho_{\text{eff}}^{(\text{II})}(a)$, respectively, reduce to the following simple forms:

$$U_{\text{eff}}^{(\text{I})}(q) = \rho_0 (1 - q) ; \quad (27)$$

$$U_{\text{eff}}^{(\text{II})}(q) = \rho_0 (1 - q^2) , \quad (28)$$

Keeping our later purposes in mind, it will be useful to introduce another constant h_n , which in terms of ρ_0 takes the following form,

$$h_n^2 \equiv \frac{1}{108} \kappa(n-6)^2 \rho_0 \quad (29)$$




Figure 1: The red graph shows the first kind of effective potential $U_{\text{eff}}^{(\text{I})}(q)$ and the brown dashed graph shows the second kind of effective potential $U_{\text{eff}}^{(\text{II})}(q)$. The shaded portion, with $q > 1$, denotes the classically allowed region, describing a bouncing universe.

Note that this constant h_n is very much like the Hubble constant associated with the constant energy density ρ_0 and will play a significant role in the subsequent discussion. Thus we have essentially brought down the analysis of a wide class of bouncing cosmologies to that of a particle in a one dimensional quadratic or linear potentials! For completeness, the effective potentials $U_{\text{eff}}^{(\text{I})}$ and $U_{\text{eff}}^{(\text{II})}$ have been plotted in Fig. 1. The point corresponding to $q = 1$ stands for the bounce and the region of parameter space with $q > 1$ depicts the classically allowed region.

The classical background with $q > 1$, that these energy densities give rise to, can be found by solving the Einstein's equations. If we choose our time coordinate such that the bounce happens at $t = 0$, then the background scale factors, corresponding to the energy densities $\rho_{\text{eff}}^{(\text{I})}(a)$ and $\rho_{\text{eff}}^{(\text{II})}(a)$, take the following forms, respectively:

$$a_{(\text{I})}(t) = (1 + h_n^2 t^2)^{\frac{3}{6-n}}, \quad (30)$$

$$a_{(\text{II})}(t) = \left[\cosh \left(\frac{3}{2} h_n t \right) \right]^{\frac{4}{6-n}}. \quad (31)$$

We caution the reader that the time coordinate t is *not* the cosmic time. Consequently, the apparent simple forms of the scale factors in Eq. (30) and Eq. (31) is rather deceptive, that is to say, the scale factors $a_{(\text{I})}(t)$ and $a_{(\text{II})}(t)$ could, in general, be complicated functions of the cosmic time τ . As an illustration, in the case of the ‘matter-bounce’ scenario described by the effective density in Eq. (26), it turns out that the time coordinate t coincides with the conformal time η and hence the background scale factor is given by:

$$a_{(\text{mb})}(t) = (1 + h_3^2 t^2) = (1 + h_3^2 \eta^2). \quad (32)$$

However, the same scale factor as a function of the cosmic time τ acquires a more complicated form:

$$a_{(\text{mb})}(\tau) = \left(\sqrt{\frac{9h_3^2 \tau^2}{4} + 1} - \frac{3h_3 \tau}{2} \right)^{2/3} + \left(\sqrt{\frac{9h_3^2 \tau^2}{4} + 1} + \frac{3h_3 \tau}{2} \right)^{2/3} - 1. \quad (33)$$

Generically, the functional dependence of the scale factor on the cosmic time and the conformal time is significantly complicated and is the prime reason to use the time coordinate t , rather than the conformal time η or the cosmic time τ .

The above discussion settles the issue of the matter content of the universe, in the bouncing scenario, which we will be interested in this work. There will be two components in the energy density of matter, near the bounce, with one of them violating the energy condition, in order to facilitate the bounce. The scaling of the energy densities with the scale factor is chosen, such that, the contribution of the effective potential to the Wheeler-DeWitt equation is at most quadratic in the dynamical variable $q(t)$. This, as we will see, will facilitate the path integral computation of the ground state wave function of the universe. In what follows, we will provide a quantum treatment of the bouncing universe, by first solving the Wheeler-DeWitt equation and then using the path-integral technique.

4 The ground state wave function of the bouncing universe

In this section, we analyze the quantum aspects of the bouncing model of the universe in order to determine the ground-state wave function and its properties. The analysis presented here is based on our description of the geometry and the matter content of the bouncing model, considered in the last sections. We shall take two different approaches for this purpose — (a) using the Wheeler-DeWitt equation and (2) the path integral approach. The quantum nature of the bouncing universe following both of these considerations and the connection between them will be elaborated in what follows.

4.1 The Wheeler-DeWitt equation and its solution

Concentrating on the gravity sector, described by the single dynamical variable $q(t)$, it is easy to see that the Wheeler-DeWitt equation reduces to one-dimensional time-independent Schrödinger equation for both the bouncing models described by energy densities $\rho_{\text{eff}}^{(\text{I})}$ and $\rho_{\text{eff}}^{(\text{II})}$, presented in Eq. (24) and Eq. (25), respectively. Taking a cue from Eq. (16), the associated Wheeler-DeWitt equation for these two models in terms of the dynamical variable $q(t)$, takes the following form,

$$\left[-\frac{\partial^2}{\partial q^2} + \alpha_n^2 (1 - q) \right] \Psi^{(\text{I})}(q) = 0 \quad (34)$$

$$\left[-\frac{\partial^2}{\partial q^2} + \frac{16}{9} \alpha_n^2 (1 - q^2) \right] \Psi^{(\text{II})}(q) = 0 \quad (35)$$

where, the factor (16/9) came from the relation between $b^{(\text{I})}$ and $b^{(\text{II})}$, as presented in Eq. (25). The constant α_n appearing in both the Wheeler-DeWitt equations can be expressed as $\beta_n V_3$, where using the expression for $b^{(\text{I})}$, from Eq. (24) and the expression for h_n from Eq. (29), β_n takes the following form,

$$\beta_n^2 \equiv 2M^{(\text{I})} \rho_0 = \frac{9h_n^2}{\kappa^2 \left(\frac{n}{6} - 1\right)^4} \equiv \frac{h_n^2}{\ell_{\text{p}}^4 \left(\frac{n}{6} - 1\right)^4} . \quad (36)$$

Here, we have introduced a new length scale $\ell_{\text{p}} = \sqrt{\kappa/3}$, which corresponds to the Planck length associated with the quantum nature of the gravitational interaction.

For the energy density $\rho_{\text{eff}}^{(\text{I})}(a)$, the effective potential $U_{\text{eff}}^{(\text{I})}(q)$ appearing in the Wheeler-DeWitt equation is linear in the dynamical variable q and hence the general solution to Eq. (34) can be written down in

terms of the Airy functions. The particular solution, which satisfy $\Psi^{(I)}(0) = 0$, i.e., the condition that the wave function must vanish at the singular point $a = 0$, is given by:

$$\Psi^{(I)}(q) = \mathcal{N}^{(I)} \left\{ \text{Bi} \left(\alpha_n^{2/3} \right) \text{Ai} \left[\alpha_n^{2/3} (1 - q) \right] - \text{Ai} \left(\alpha_n^{2/3} \right) \text{Bi} \left[\alpha_n^{2/3} (1 - q) \right] \right\}, \quad (37)$$

where, $\mathcal{N}^{(I)}$ is a normalisation constant. For $q > 1$, i.e., in the classical regime, one is interested in the limit $(\alpha_n/\ell_p^2) \gg 1$. In which case, we can use the following expansions of the Airy functions,

$$\begin{aligned} \text{Ai}(x) &\sim \frac{\sqrt{\pi}}{(-x)^{1/4}} \cos \left[-\frac{2}{3} (-x)^{3/2} + \frac{\pi}{4} \right]; \quad (x < 0, |x| \gg 1) \\ \text{Ai}(x) &\sim \frac{1}{\sqrt{\pi} x^{1/4}} \exp \left(-\frac{2}{3} x^{3/2} \right); \quad (x \gg 1) \end{aligned} \quad (38)$$

along with the fact that the expansion of $\text{Bi}(x)$ for negative, but large x is identical to that of $\text{Ai}(x)$, while for large positive x , $\text{Bi}(x)$ is exponentially growing, the exact opposite of the behaviour of $\text{Ai}(x)$, presented in Eq. (38). Thus for $q > 1$ and for large (α_n/ℓ_p^2) limit, the asymptotic expansion of the wave function $\Psi^{(I)}(q)$ takes the form:

$$\Psi^{(I)}(q) \approx \frac{\mathcal{N}^{(I)}}{\sqrt[4]{\alpha_n^{2/3} (q - 1)}} \exp \left(\frac{2}{3} \alpha_n \right) \cos \left[\frac{2}{3} \alpha_n (q - 1)^{3/2} - \frac{\pi}{4} \right]. \quad (39)$$

Notice that there is a noticeable similarity between the above solution of the Wheeler-DeWitt equation and that of the ground-state wave function of Hartle and Hawking (HH), associated with the ‘no-boundary’ proposal [6]. The precise connection of the above wave function with the no-boundary proposal will be made clear when we analyze the above problem using the path integral analysis.

On the other hand, the general solution to Eq. (35) with a quadratic effective potential $U_{\text{eff}}^{(\text{II})}$, as in Eq. (28) can be written down in terms of the parabolic cylinder functions $D_\nu(x)$. However, for our purpose it is convenient to work with another set of linearly independent solutions [30, p.314-p.317], namely, $W(a, -x)$ and $W(a, x)$. These functions are constructed from linear combinations of two linearly independent sets of standard parabolic cylinder functions. In general, the solution of Eq. (35) will involve two arbitrary constants, one of which can be eliminated by demanding that $\Psi^{(\text{II})}(0) = 0$, yielding:

$$\Psi^{(\text{II})}(q) = \mathcal{N}^{(\text{II})} \left[W \left(\frac{2}{3} \alpha_n, \frac{2\sqrt{2}}{\sqrt{3}} \sqrt{\alpha_n} q \right) - W \left(\frac{2}{3} \alpha_n, -\frac{2\sqrt{2}}{\sqrt{3}} \sqrt{\alpha_n} q \right) \right], \quad (40)$$

where, $\mathcal{N}^{(\text{II})}$ is a normalisation constant. For $q > 1$, i.e., in the classical regime, the asymptotic expansion of the above solution in the $(\alpha_n/\ell_p^2) \gg 1$ limit is given by [30],

$$\Psi^{(\text{II})}(q) \approx \frac{\mathcal{N}^{(\text{II})}}{\sqrt[4]{\frac{4}{3} \alpha_n (q^2 - 1)}} \exp \left(\frac{\pi}{3} \alpha_n \right) \cos \left[\frac{4}{3} \alpha_n \xi(q) - \frac{\pi}{4} \right], \quad (41)$$

where,

$$\xi(q) \equiv \frac{1}{2} q \sqrt{q^2 - 1} - \frac{1}{2} \log(q + \sqrt{q^2 - 1}) \quad (42)$$

It is worth emphasizing, that even if the effective potential $U_{\text{eff}}^{(\text{II})}$ was dependent on terms linear in the generalized coordinate q , the solution of the Wheeler-DeWitt equation would have an identical structure, as presented in Eq. (41). Thus the results obtained above is general enough to encompass all possible wave functions of the universe, with the matter degrees of freedom, yielding an effective potential, which is at most quadratic in the generalized degree of freedom q . Once again, we see that there is a close resemblance of the solution to the Wheeler-DeWitt equation to the Hartle-Hawking wave function compatible with the no-boundary proposal. As we shall shortly see, in the next subsection, through the path-integral prescription, that this resemblance is not accidental.

4.2 Path integral approach

Having described the solution to the Wheeler-DeWitt equation associated with linear and quadratic effective potentials, let us now try to determine the wave function of the universe, using the path-integral formalism. As we will see, the choice of contour for the evaluation of the path integral and the use of appropriate boundary conditions is extremely crucial in evaluating the path integral. The connection of the wave function obtained along these lines, with the no-boundary proposal of Hartle and Hawking will also be discussed. Prior to that, let us briefly review some fundamentals of the path integral approach to the mini-superspace model.

Following [20], it is convenient to use a gauge in which $\dot{\mathcal{N}} = 0$ and hence the ‘physical time’ T , defined through the following relation, $T = \int \mathcal{N} dt$, becomes $T = \mathcal{N}t$. For simplicity, but without losing generality, we will restrict ourselves within the time interval $0 \leq t \leq 1$, which translates into $0 \leq T \leq \mathcal{N}$. We can then construct a path integral kernel from the gravitational action $\mathcal{S}_{\text{perfect}}$, with the boundary condition that $q(T = \mathcal{N}) = q_1$ and $q(T = 0) = q_0$, as follows:

$$K(q_1, \mathcal{N}; q_0, 0) \equiv \int_{q(T=0)=q_0}^{q(T=\mathcal{N})=q_1} \mathcal{D}[q] \exp \left\{ i V_3 \int_0^{\mathcal{N}} dT' \left[-\frac{M}{2} \left(\frac{dq}{dT'} \right)^2 + U_{\text{eff}}(q) \right] \right\}, \quad (43)$$

where, Eq. (5) has been used. One can check that the Hamiltonian constraint indeed generates translation in the ‘physical time’ coordinate T . Furthermore, using the Kernel $K(q_1, \mathcal{N}; q_0, 0)$, it is indeed possible to construct the wave function of the universe $\Psi(q_1)$ at a later instant, from the wave function $\psi(q_0)$ at the initial instant, in the following manner:

$$\Psi(q_1) = \int_{\mathcal{C}} \left[\int_{-\infty}^{\infty} K(q_1, \mathcal{N}; q_0, 0) \psi(q_0) dq_0 \right] d\mathcal{N} \quad (44)$$

where, $\psi(q_0)$ is a square-integrable function in the appropriate domain and the integration over $\mathcal{N}$ is performed along a contour $\mathcal{C}$ in complex $\mathcal{N}$ -plane. The choice of an appropriate contour is not straightforward, since, if $\Psi(q)$ satisfies the Wheeler-DeWitt equation, we get the following condition,

$$\int_{-\infty}^{\infty} K(q_1, \mathcal{N}_f; q_0, 0) \psi(q_0) dq_0 - \int_{-\infty}^{\infty} K(q_1, \mathcal{N}_i; q_0, 0) \psi(q_0) dq_0 = 0, \quad (45)$$

where, $\mathcal{N}_i$ and $\mathcal{N}_f$ are the endpoints of the contour $\mathcal{C}$. One way of achieving this condition is to choose the endpoints such that, the kernel identically vanishes at these points. Another choice would be to choose the contour of T integration to be a closed one. Even though both of these possibilities are viable, for the purpose of our current discussion, we shall take the former choice and explore the structure of the path integral kernel for both linear and quadratic model.

4.2.1 The linear model

The path integral Kernel for the linear model is essentially the Kernel of a one dimensional particle, subject to a constant force and hence can be evaluated explicitly. The explicit form of the Kernel being,

$$K^{(I)}(q_1, \mathcal{N}; q_0, 0) = \sqrt{\frac{iM}{2\pi\mathcal{N}}} \exp \left[i\mathcal{S}_{\text{cl}}^{(I)}[q_1, q_0; \mathcal{N}] \right] \quad (46)$$

where, $\mathcal{S}_{\text{cl}}^{(I)}$ is just the action $\mathcal{S}_{\text{perfect}}[q]$ evaluated at the classical solution, which satisfies the boundary conditions $q(T=0) = q_0$ and $q(T=\mathcal{N}) = q_1 > 1$, such that the final state of the universe is described by a classical geometry. The explicit form of $\mathcal{S}_{\text{cl}}^{(I)}$, normalized with respect to the three-volume V_3 , in terms of q_0 , q_1 and the time interval $\mathcal{N}$ is given by,

$$\frac{\mathcal{S}_{\text{cl}}^{(I)}}{V_3} = \frac{1}{2\ell_p^2 \left(1 - \frac{n}{6}\right)^2} \left[-\frac{(q_1 - q_0)^2}{2\mathcal{N}} - h_n^2 (q_1 + q_0) \mathcal{N} + \frac{h_n^4}{6} \mathcal{N}^3 + 2h_n^2 \mathcal{N} \right]. \quad (47)$$

In the context of de Sitter spacetime, the original prescription of HH for defining a wave of the universe corresponds to choosing an initial wavefunction $\psi(q_0) \propto \delta(q_0)$. However, recent, more careful mathematical considerations have shown that this approach leads to perturbations that are unsuppressed [31, 32]. On the other hand, instead, if we choose an initial state $\psi(q_0)$ that corresponds to a well defined Euclidean momentum, the corresponding perturbations turn out to be Gaussian distributed [16, 17]. Motivated by this, we demand that the initial wave function $\psi(q_0)$ corresponds to that of a momentum eigenstate with momentum $p = -Mv$, where, $v \equiv (dq/dT)_{T=0} = \mathcal{N}^{-1}\dot{q}(0)$. It is then instructive to define the momentum space Kernel as the Fourier transform of the position space Kernel, yielding,

$$\begin{aligned} \mathcal{K}^{(I)}(q_1, \mathcal{N}; -Mv, 0) &\equiv \int K^{(I)}(q_1, \mathcal{N}; q_0, 0) e^{-iMvV_3q_0} dq_0 \\ &= e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(I)}[q_1, -Mv; \mathcal{N}]} \end{aligned} \quad (48)$$

where, $\tilde{\mathcal{S}}_{\text{cl}}^{(I)}$ is the action $\mathcal{S}_{\text{perfect}}[q] - \mathcal{S}_{\text{Boundary}}[q_0]$, evaluated at the classical solution $\tilde{q}_{\text{cl}}^{(I)}(t)$ which satisfies the boundary condition $\mathcal{N}^{-1}\dot{q}(0) = v$ and $q(T=\mathcal{N}) = q_1$, with the boundary term $\mathcal{S}_{\text{Boundary}}[q_0]$ having the form: $\mathcal{N}^{-1}MV_3q(0)\dot{q}(0)$ and, the dot, as usual, denotes a derivative with respect to t .

The classical solution to the equation of motion, for the dynamical variable $q(t)$, arising out of the action $\mathcal{S}_{\text{perfect}}$, with the above boundary conditions, takes the following explicit form:

$$\tilde{q}_{\text{cl}}^{(I)}(t) = h_n^2 \mathcal{N}^2 t^2 - \mathcal{N} (h_n^2 \mathcal{N} + v) + \mathcal{N} t v + q_1. \quad (49)$$

Motivated by the recent result of [17], in the context of de Sitter spacetime, where the no-boundary wavefunction was realised by choosing the initial momentum to be purely imaginary, we demand that the $v = 2ih_n$. With this choice of the initial velocity v and the classical solution $\tilde{q}_{\text{cl}}^{(I)}(t)$, presented above, the explicit form of the classical action $\tilde{\mathcal{S}}_{\text{cl}}^{(I)}$, which includes the boundary term as well, takes the following form,

$$\frac{\tilde{\mathcal{S}}_{\text{cl}}^{(I)}}{V_3} = \frac{1}{2\ell_p^2 \left(1 - \frac{n}{6}\right)^2} \left[\frac{2h_n^4}{3} \left(\mathcal{N} + \frac{i}{h_n} \right)^3 - 2h_n^2 \left(\mathcal{N} + \frac{i}{h_n} \right) (q_1 - 1) - \frac{4ih_n}{3} \right]. \quad (50)$$




Figure 2: The steepest descent/ascent curves in the complex $\mathcal{N}$ -plane, that pass through the saddle points $\bar{\mathcal{N}}_{\pm}^{(I)}$, for the linear model are represented by the black curves. The contour-plots for the function $\text{Re}[i2\ell_p^2\{(n/6)-1\}^2 V_3^{-1} \tilde{S}_{cl}^{(I)}]$ are also presented, using the colour coding scheme given by the right inset, to better visualise the descent/ascent directions (the plots are for the parameter values $h_n = 1$ and $q_1 = 3$). The dark red, horizontal line is the real line contour over which the integration of the lapse function needs to be performed to evaluate the no-boundary wavefunction. The real line contour can be smoothly deformed into the union of dashed red contours $\mathcal{C}_{+}^{(I)}$ and $\mathcal{C}_{-}^{(I)}$, thereby, enabling us to perform saddle point approximation. See text for more discussion.

In order to determine the wave function from the path integral Kernel, we need to integrate over the lapse function $\mathcal{N}$. The integral should be over a contour for which the Kernel vanishes sufficiently fast as one approaches the endpoints, i.e., the real part of $i\tilde{S}_{cl}^{(I)}$ should take a large and negative value as we approach the endpoints of the contour. Taking $\{\mathcal{N} + (i/h_n)\}$ as $re^{i\theta}$, we obtain from Eq. (50), that $\text{Re}(i\tilde{S}_{cl}^{(I)})$ takes large negative values, such that $\mathcal{K}^{(I)}(q_1, \mathcal{N}; -Mv, 0)$ vanishes, for large values of r , if $(\sin^3 \theta - 3 \sin \theta \cos^2 \theta) < 0$. Thus the momentum space Kernel identically vanishes in the asymptotic regions in the complex $\mathcal{N}$ -plane, provided we restrict ourselves to the following regions: $\text{Arg}[\{\mathcal{N} + i/(h_n)\}] \in (0, \pi/3) \cup (2\pi/3, \pi) \cup (4\pi/3, 5\pi/3)$ (These regions have been depicted in Fig. 2 in dark shades of blue). In addition, we also demand that the contours should pass through the saddle points, which for the action $\tilde{S}_{cl}^{(I)}$ turns out to be,

$$\bar{\mathcal{N}}_{\pm}^{(I)} = -\frac{i}{h_n} \pm \frac{\sqrt{q_1 - 1}}{h_n}, \quad (51)$$

The classical solutions $\tilde{q}_{cl}^{(I)}(t)$ with the above choice for the lapse function and with $v = 2ih_n$ is given by:

$$\tilde{q}_{cl,\pm}^{(I)}(t) = t^2 \left[(q_1 - 2) \mp 2i\sqrt{q_1 - 1} \right] + t \left(2 \pm 2i\sqrt{q_1 - 1} \right). \quad (52)$$

Therefore, we see that the saddle points describe geometries in which the scale factor $a(t) = q(t)^{3/(6-n)}$ vanishes at $t = 0$ (since $\tilde{q}_{cl,\pm}^{(I)}(t = 0) = 0$) and equals $q_1^{3/(6-n)}$ at $t = 1$ (since $\tilde{q}_{cl,\pm}^{(I)}(t = 1) = q_1$).

The Lorentzian contour, namely the real line, is the most natural choice for the $\mathcal{N}$ integration that evaluates the wavefunction. Fortunately, from Fig. 2, we see that this integral along the real line is convergent. In order to find the asymptotic expression for the wavefunction, however, we will employ the saddle point approximation. The steepest descent/ascent curves associated with the action $\tilde{\mathcal{S}}_{\text{cl}}^{(1)}$ are shown in Fig. 2 along with the saddle points $\bar{\mathcal{N}}_{\text{cl},\pm}^{(1)}$. The regions where the real part of $i\tilde{\mathcal{S}}_{\text{cl}}^{(1)}$ becomes large and negative are also presented. The relevant saddle points that contribute to the evaluation of the integral, according to Picard-Lefschetz theory [15, 17], are the ones that can be approached by flowing down (i.e., in view of Fig. 2, towards the direction of darker shades of blue) the original integration contour. This dictates that the appropriate contour, to which the real line should be deformed to, corresponds to $\mathcal{C}_+^{(1)} + \mathcal{C}_-^{(1)}$. With the relevant saddle points determined, we can now evaluate the saddle point approximation of the no-boundary wave function for the bouncing model, yielding,

$$\Psi^{(1)}(q_1) \sim \left(\frac{e^{i\frac{\pi}{4}}}{\sqrt{|\partial_{\mathcal{N}}^2 \tilde{\mathcal{S}}_{\text{cl}}^{(1)}|}} e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(1)}} \right)_{\bar{\mathcal{N}}_{\text{cl},+}^{(1)}} + \left(\frac{e^{-i\frac{\pi}{4}}}{\sqrt{|\partial_{\mathcal{N}}^2 \tilde{\mathcal{S}}_{\text{cl}}^{(1)}|}} e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(1)}} \right)_{\bar{\mathcal{N}}_{\text{cl},-}^{(1)}} \propto \exp\left(\frac{2\alpha_n}{3}\right) \frac{\cos\left[\frac{2}{3}(q-1)^{3/2}\alpha_n - \frac{\pi}{4}\right]}{\sqrt[4]{(\alpha_n)^2(q-1)}}, \quad (53)$$

which is clearly consistent with Eq. (39).

Having described the derivation of the no-boundary wave function from the path integral analysis with ‘mixed’ boundary condition, let us discuss its interpretation. We have seen that the saddle point solution $\tilde{q}_{\text{cl},\pm}^{(1)}(t)$ satisfies $\tilde{q}_{\text{cl},\pm}^{(1)}(0) = 0$ and $\tilde{q}_{\text{cl},\pm}^{(1)}(1) = q_1$. Therefore, the corresponding geometry may be imagined as describing an evolution from a point-sized to a finite-sized universe. This is reminiscent of the famous Hartle-Hawking saddle point geometry that appears in the context of Euclidean path integral approach to de Sitter cosmology [27]. Therein, the corresponding wavefunction of the universe is usually interpreted as describing tunnelling out of ‘nothing’. One might, therefore, be tempted to bestow an analogous interpretation on the wavefunction $\Psi^{(1)}(q_1)$, with the Hawking-Hartle geometry replaced by the appropriate generalization of it in the context of bounce (For representation purpose, in Fig. 3, we have presented the saddle point geometry corresponding to a ‘matter-bounce’ scenario). However, we argue that such an interpretation is questionable. A similar argument in the context of de Sitter spacetime can be found in [17], where the authors emphasize that the conventional interpretation of the no-boundary wavefunction, as describing ‘tunnelling from nothing’, is problematic. In order to appreciate this, note that, even though the dominant contribution to the path integral leading to $\Psi^{(1)}(q_1)$ is from the Hartle-Hawking like saddle point geometry, corresponding to a spacetime that emerges from zero size, the off-shell geometries can emerge from any size. This fact is evident from Eq. (48), where we have clearly performed a summation over *all* values of the ‘initial size’ q_0 to define the wavefunction. In the spirit of [17], and in the light of Eq. (48) it is more reasonable to interpret $\Psi^{(1)}(q_1)$ as describing a transition from the state of a specific Euclidean momentum.

4.2.2 The quadratic model

Having described the linear model in the previous section, we will now present the scenario in which the effective potential appearing in the Wheeler-DeWitt equation for the gravity and the perfect fluid system is quadratic in the dynamical variable q . In this model, the action is that of a one-dimensional particle moving in a quadratic potential, for which the path integral Kernel can be evaluated explicitly and takes




Figure 3: The saddle point geometry that would be relevant to Euclidean path integral approach to matter bounce. The surface denotes the $x - \eta$ plane, with periodic identification assumed along the x direction. The orange-coloured portion denotes the Lorentzian part of the geometry, while blue denotes the Euclidean portion. There is a true singularity at the centre of the Euclidean portion, unlike the Hawking-Hartle geometry which is smooth everywhere.

the following form:

$$K^{(\text{II})}(q_1, \mathcal{N}; q_0, 0) = \sqrt{\frac{3MiV_3}{2 \sinh\left(\frac{3}{2}h_n\mathcal{N}\right)}} e^{i\mathcal{S}_{\text{cl}}^{(\text{II})}[q_1, q_0; \mathcal{N}]} \quad (54)$$

where, $\mathcal{S}_{\text{cl}}^{(\text{II})}$ is the action $\mathcal{S}_{\text{perfect}}[q]$ (see Eq. (5)) evaluated at the classical solution satisfying the Dirichlet boundary conditions: $q(t=0) = q_0$ and $q(t=1) = q_1 > 1$. The explicit form of the action is given by the following expression,

$$\frac{\mathcal{S}_{\text{cl}}^{(\text{II})}}{V_3} = \frac{8}{9\ell_p^2 \left(\frac{n}{6} - 1\right)^2} \left\{ -\frac{3h_n}{4 \sinh\left(\frac{3}{2}h_n\mathcal{N}\right)} \left\{ \cosh\left(\frac{3}{2}h_n\mathcal{N}\right) (q_1^2 + q_0^2) - 2q_0q_1 \right\} + \frac{9}{8}h_n^2\mathcal{N} \right\}. \quad (55)$$

As in the case of linear model, we now demand that the initial wavefunction $\Psi(q_0)$ corresponds to a state of definite momentum $p = -Mv$. This implies that the relevant Kernel is the momentum space Kernel, with the following expression,

$$\begin{aligned} \mathcal{K}^{(\text{II})}(q_1, \mathcal{N}; -Mv, 0) &\equiv \int K^{(\text{II})}(q_1, \mathcal{N}; q_0, 0) e^{-iMvV_3q_0} dq_0 \\ &= \sqrt{\text{sech}\left(\frac{3}{2}h_n\mathcal{N}\right)} e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}[q_1, -Mv; \mathcal{N}]}, \end{aligned} \quad (56)$$

where, as in the linear case, $\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}$ is the action $\mathcal{S}_{\text{perfect}}[q] - \mathcal{S}_{\text{Boundary}}[q]$, evaluated at the classical solution $\hat{q}_{\text{cl}}^{(\text{II})}(t)$, which satisfies the boundary condition $\mathcal{N}^{-1}\dot{q}(0) = v$ and $q(1) = q_1$, with the boundary term

$\mathcal{S}_{\text{Boundary}}[q]$ having the form: $\mathcal{N}^{-1}MV_3q(0)\dot{q}(0)$ and, the dot denotes a derivative with respect to t . First of all, the explicit form of the solution $\tilde{q}_{\text{cl}}^{(\text{II})}(t)$, given the above boundary conditions, is given by:

$$\tilde{q}_{\text{cl}}^{(\text{II})}(t) = \frac{\text{sech}\left(\frac{3h_n\mathcal{N}}{2}\right)}{3h_n} \left[3h_nq_1 \cosh\left(\frac{3}{2}h_n\mathcal{N}t\right) + 2v \sinh\left(\frac{3}{2}h_n\mathcal{N}(t-1)\right) \right]. \quad (57)$$

Following [17] and the analysis in the case of the linear potential, for the quadratic model we have to choose $v = (3ih_n/2)$ in order to get the no-boundary wave function for the bouncing model under consideration. With this choice of the velocity v , the action on the classical trajectory $\tilde{q}_{\text{cl}}^{(\text{II})}(t)$ becomes:

$$\frac{\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}}{V_3} = \frac{8}{9\ell_p^2\left(\frac{n}{6}-1\right)^2} \left\{ \frac{9h_n^2\mathcal{N}}{8} - \frac{3}{4}h_n\text{sech}\left(\frac{3h_n\mathcal{N}}{2}\right) \left[(q_1^2+1) \sinh\left(\frac{3h_n\mathcal{N}}{2}\right) + 2iq_1 \right] \right\}. \quad (58)$$

Again, we need to find out the asymptotic regions in the complex $\mathcal{N}$ -plane, where the endpoints of the contours associated with the $\mathcal{N}$ integration must lie, so that the endpoint contribution to the kernel identically vanishes. This requires $\text{Re}[i\mathcal{S}_{\text{cl}}^{(\text{II})}] < 0$, which for large values of $|\mathcal{N}|$ demands: $\text{Im}[\mathcal{N}] > 0$. The immediate consequence of this condition is that, unlike in the case of the linear model, the $\mathcal{N}$ integral along real line contour and hence, the strictly Lorentzian path integral, is not convergent. Convergence of the $\mathcal{N}$ integration demands that we choose a slightly modified contour, namely, the continuous curve joining $-oe^{-i0^+}$ to oe^{i0^+} , represented by the red curve $\tilde{\mathcal{C}}^{(\text{II})}$ in Fig. 4.

In order to obtain the saddle points, first of all we need to compute the derivative of the classical action with respect to the lapse function $\mathcal{N}$, which yields,

$$\begin{aligned} \frac{1}{V_3} \frac{\partial \tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}}{\partial \mathcal{N}} &= \frac{9}{8}h_n^2 + \frac{9}{8}h_n^2\text{sech}\left(\frac{3h_n\mathcal{N}}{2}\right) \left[-(q_1^2+1) \text{sech}\left(\frac{3h_n\mathcal{N}}{2}\right) + 2iq_1 \tanh\left(\frac{3h_n\mathcal{N}}{2}\right) \right] \\ &= \frac{9}{8}h_n^2\text{sech}^2\left(\frac{3h_n\mathcal{N}}{2}\right) \left[iq_1 + \sinh\left(\frac{3h_n\mathcal{N}}{2}\right) \right]^2. \end{aligned} \quad (59)$$

Setting the above expression to zero, we obtain, $\sinh(3h_n\mathcal{N}/2) = -iq_1$, solving which we obtain an infinite number of saddle points for the action $\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}$ and they are given by:

$$\bar{\mathcal{N}}_{j,\pm}^{(\text{II})} = \frac{(4j-1)\pi i}{3h_n} \pm \frac{2}{3h_n} \log\left(q_1 + \sqrt{q_1^2-1}\right); \quad j \in \mathbb{Z}. \quad (60)$$

At these saddle points, substitution of the respective values of the lapse function $\mathcal{N}$ yields the following expression for the classical solution $\tilde{q}_{\text{cl}}^{(\text{II})}(t)$,

$$\tilde{q}_{\text{cl}(j,\pm)}^{(\text{II})}(t) = \sin \left[\left((4j-1)\pi \mp 2i \log(q_1 + \sqrt{q_1^2-1}) \right) \frac{t}{2} \right]; \quad j \in \mathbb{Z}. \quad (61)$$

It can be easily verified that the above solutions indeed satisfy the necessary boundary conditions, namely, $\tilde{q}_{\text{cl}(j,\pm)}^{(\text{II})}(0) = 0$ and $\tilde{q}_{\text{cl}(j,\pm)}^{(\text{II})}(1) = q_1$ and this holds for all $j \in \mathbb{Z}$.

The above analysis determines the saddle points of the classical action, $\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}$ in the complex $\mathcal{N}$ plane and the associated classical trajectory. In order to perform saddle point approximation of the no-boundary wave function, we need to first identify the relevant contour, which passes through the saddle points and




Figure 4: The steepest descent/ascent curves in the complex $\mathcal{N}$ -plane for the quadratic model are represented by the black curves. The contour-plots for $\text{Re} \left[i(9/8)\ell_p^2(n/6 - 1)^2 V_3^{-1} \tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})} \right]$ is also presented, with the colour coding scheme given in the right inset, to better visualise the descent/ascent directions (the parameter values for the above plot correspond to: $h_n = 1$ and $q_1 = 3$). The dashed purple lines are the branch-cuts of the pre-exponential factor $\sqrt{\text{sech}(3h_n\mathcal{N}/2)}$. The horizontal, green line is the real line contour. However, convergence of the $\mathcal{N}$ -integral demands that we choose a slightly modified contour, namely, the red continuous curve $\tilde{\mathcal{C}}^{(\text{II})}$, joining $-\infty e^{-i0^+}$ to ∞e^{i0^+} . This continuous red contour can be smoothly deformed into the dashed red curve $\mathcal{C}^{(\text{II})}$, that pass through the saddle points, thereby enabling us to evaluate the saddle point approximation. See text for more discussion.

have vanishing end-point contributions to the Kernel. The steepest descent/ascent curves for $i\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}(\mathcal{N})$ is depicted in Fig. 4. Besides, the contours associated with constant values of the $\text{Re}[i\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}]$ have also been depicted, to better visualize the descent/ascent directions. As prescribed by the Picard-Lefschetz theory, we can deform the original integration contour $\tilde{\mathcal{C}}^{(\text{II})}$ into the red dashed contour $\mathcal{C}^{(\text{II})}$ (see Fig. 4), without touching the branch-cuts of the pre-exponential factor $\text{sech}(3h_n\mathcal{N}/2)$ appearing in Eq. (59). Having identified the appropriate contour, we can now evaluate the $\mathcal{N}$ -integral using the saddle point approximation in the limit $(V_3 h_n / \ell_p^2) \gg 1$. The dominant contribution to the integral comes from the saddle points $\tilde{\mathcal{N}}_{0,+}^{(\text{II})}$ and $\tilde{\mathcal{N}}_{0,-}^{(\text{II})}$, since the contributions from $\mathcal{N}_{j,\pm}^{(\text{II})}$ are exponentially suppressed for $j > 1$. Therefore, the saddle point approximation of the no-boundary wave function for the quadratic class of bouncing models, is given by:

$$\Psi^{(\text{II})}(q_1) \sim \left(\frac{e^{i\frac{\pi}{4}} e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}}}{\sqrt{|\cosh(\frac{3}{2}h_n\mathcal{N})|}} \right)_{\mathcal{N}_{0,+}} + \left(\frac{e^{-i\frac{\pi}{4}} e^{i\tilde{\mathcal{S}}_{\text{cl}}^{(\text{II})}}}{\sqrt{|\cosh(\frac{3}{2}h_n\mathcal{N})|}} \right)_{\mathcal{N}_{0,-}} \propto e^{\frac{\pi}{3}\alpha_n \frac{\cos[\frac{4}{3}\alpha_n \xi(q) - \frac{\pi}{4}]}{\sqrt{(\alpha_n)^2(q^2 - 1)}}}, \quad (62)$$

where,

$$\xi(q) \equiv \frac{1}{2}q\sqrt{q^2 - 1} - \frac{1}{2}\log(q + \sqrt{q^2 - 1}). \quad (63)$$

Once again, we see that the result is consistent with that obtained from the Wheeler De-Witt equation.

Having derived the no-boundary wave function from the path-integral prescription with an appropriate choice of the integration contours, let us try to provide a physical interpretation to the same. Just as we have obtained in the case of the linear model, the saddle point solution $\tilde{q}_{\text{cl},\pm}^{(\text{II})}(t)$ indeed satisfies the relevant boundary conditions, given by: $\tilde{q}_{\text{cl},\pm}^{(\text{II})}(0) = 0$ and $\tilde{q}_{\text{cl},\pm}^{(\text{II})}(1) = q_1$. The corresponding geometry, therefore, acquires the interpretation of an evolution from a point-sized to a finite-sized universe. Once again, one is reminded of the quintessential saddle point geometry of the Euclidean path integral approach to de Sitter cosmology [27] and the corresponding wave function of the universe that is usually interpreted as describing tunnelling out of ‘nothing’. However, with the hindsight of our arguments in the case of the linear model, we maintain that such an interpretation is questionable. This is appreciated once we observe that even though the dominant contribution to the path integral leading to $\Psi^{(\text{II})}(q_1)$ is from the Hartle-Hawking like saddle point geometry, corresponding to a spacetime that emerges from zero size, the off-shell geometries can emerge from any size. This fact is evident from Eq. (56), where we have clearly performed a summation over *all* values of the ‘initial size’ q_0 to define the wavefunction. A more reasonable interpretation for $\Psi^{(\text{II})}(q_1)$, in the spirit of [17] and in the light of Eq. (56), seems to be that it describes a transition from the state of a specific Euclidean momentum.

4.3 Shear instability and its quantum analogue

In Section 2, we have discussed the metric ansatz, which is spatially homogeneous and isotropic. As a consequence, the geodesics comoving with the Hubble flow are free from any shear. In addition, the effect of any classical shear perturbations to the background FRW spacetime is to add an effective fluid with its density scaling as a^{-6} . Therefore, in the standard cosmology, the effects of shear can be safely ignored for sufficiently large values of the scale factor. On the contrary, for bouncing scenarios, the effective energy density arising from the classical shear will inevitably become dominant at sufficiently small values of the scale factor and can potentially affect the bouncing models [33]. We shall shortly discuss how this effect translates to in the quantum picture.

For a brief review of the classical picture along the lines discussed in this work, in particular following the metric ansatz presented in Section 2, we start with the following parametrization of the Bianchi I universe,

$$ds^2 = -\frac{\mathcal{N}(t)^2}{q(t)^{(4-3b)}} dt^2 + q(t)^b \left[\exp \left(\sqrt{\frac{2\kappa}{3}} \left\{ \theta_1(t) - \sqrt{3}\theta_2(t) \right\} \right) dx^2 \right. \\ \left. + \exp \left(\sqrt{\frac{2\kappa}{3}} \left\{ \theta_1(t) + \sqrt{3}\theta_2(t) \right\} \right) dy^2 + \exp \left(-2\sqrt{\frac{2\kappa}{3}} \theta_1(t) \right) dz^2 \right]. \quad (64)$$

It follows that $q(t)^{b/2}$ is the geometric mean of the ‘scale factors’ along the three independent spatial directions. Substitution of the above metric ansatz in the Einstein-Hilbert action with a perfect fluid source, i.e., in the action given by Eq. (5) yields:

$$\mathcal{S}_{\text{perfect}}^{(\text{shear})} = V_3 \int dt \left[-\frac{M}{2\mathcal{N}} \left(\frac{dq}{dt} \right)^2 + \mathcal{N} U_{\text{eff}}(q) \right] dt + V_3 \int dt \left[\frac{q^2}{2\mathcal{N}} \left(\frac{d\theta_1}{dt} \right)^2 + \frac{q^2}{2\mathcal{N}} \left(\frac{d\theta_2}{dt} \right)^2 \right]. \quad (65)$$

Variation of the above action with respect to the lapse function $\mathcal{N}$, and then setting the lapse function $\mathcal{N}$

to unity, we obtain the classical Hamiltonian constraint to read ³:

$$-\frac{M}{2} \left(\frac{dq}{dt} \right)^2 - U_{\text{eff}}(q) + \frac{q^2}{2} \left[\left(\frac{d\theta_1}{dt} \right)^2 + \left(\frac{d\theta_2}{dt} \right)^2 \right] = 0 . \quad (66)$$

Further, the variation of the action functional with respect to $\theta_{1,2}$ yields the classical equation of motion for θ_1 and θ_2 as: $\dot{\theta}_{1,2} \propto (1/q^2)$, where ‘dot’ denotes derivative with respect to t . Using this result in the constraint equation, we get:

$$-\frac{M}{2} \left(\frac{dq}{dt} \right)^2 - U_{\text{eff}}(q) + \frac{\rho_\theta}{q^2} = 0 \quad (67)$$

where, ρ_θ is a constant, related to the proportionality factor between $\dot{\theta}$ and $(1/q^2)$. Using the expression of the effective energy density U_{eff} , as given in Eq. (6), we see that the presence of the (ρ_θ/q^2) term in Eq. (67) can be interpreted as arising from an effective energy density ρ_θ , which scales as $q^{-3b} = a^{-6}$. If the value of ρ_θ is sufficiently large, such that (dq/dt) never vanishes, the bouncing scenario becomes classically forbidden. In particular, for the linear model, setting $(dq/dt) = 0$, we obtain, $\rho_0 q^3 - \rho_0 q^2 + \rho_\theta = 0$. Introducing, a new variable x through the following relation: $q = x + (1/3)$, the above algebraic equation translates into, $x^3 - (1/3)x + \{-(2/27) + (\rho_\theta/\rho_0)\} = 0$. The reduced algebraic equation would have positive real solution consistent with classical bounce, provided $\{-(2/27) + (\rho_\theta/\rho_0)\}^2 < 4(1/3)^6$. Thereby, we yield the following condition for a classical bounce to occur:

$$\frac{\rho_\theta}{\rho_0} < \frac{4}{27} . \quad (68)$$

On the other hand, for the quadratic model, the relevant algebraic equation becomes, $\rho_0 q^4 - \rho_0 q^2 + \rho_\theta = 0$ and hence real and positive solutions for q will exist, provided the following inequality holds,

$$\frac{\rho_\theta}{\rho_0} < \frac{1}{4} . \quad (69)$$

This finishes our discussion about the effect of shear on the classical FRW spacetime with a bouncing origin, where the contribution from the shear must satisfy the conditions presented in Eq. (68) and Eq. (69) respectively.

We shall now look at the quantum analysis of the problem and the starting point of the same is the Wheeler-DeWitt equation, which in the present context takes the following form,

$$\left[-\frac{1}{2MV_3} \left(\frac{\partial^2}{\partial q^2} \right) + V_3 U_{\text{eff}}(q) + \frac{1}{2q^2 V_3} \nabla^2 \right] \Psi(q, \theta_1, \theta_2) = 0 , \quad (70)$$

where, $\nabla^2 \equiv (\partial/\partial\theta_1)^2 + (\partial/\partial\theta_2)^2$. Given the above differential equation, it is evident that the (θ_1, θ_2) sector and the main gravitational degree of freedom q do not talk among each other. Hence, it is advisable to look for separable solutions of the Wheeler-DeWitt wave function $\Psi(q, \theta_1, \theta_2)$ of the following form,

$$\Psi(q, \theta_1, \theta_2) = \psi_{|\mathbf{k}|}(q) e^{iV_3(k_1\theta_1 + k_2\theta_2)} , \quad (71)$$

³If the spatial slices are closed, in addition to the terms in Eq. (66), we get an extra term given by $q^{2n/(6-n)}U(\theta_1, \theta_2)$ and $q^{(2n+4)/(6-n)}U(\theta_1, \theta_2)$, respectively, for the linear and the quadratic model, with $U(\theta_1, \theta_2)$ describing an effective potential for $\theta_{1,2}$. For sufficiently small values of the scale factor and $\theta_{1,2}$, one can reduce the constraint equation to Eq. (66), if we assume that we can neglect the extra term in comparison to U_{eff} . This is possible when $2 < n < 6$, for both the models.




Figure 5: The typical form of the total effective potential $U_{\text{total}} = U_{\text{eff}} - (\rho_\theta/q^2)$ has been demonstrated, when a classical bouncing scenario is allowed. The two classical turning points are marked $q_<$ and $q_>$, respectively and the shaded portions with $q > q_>$ and $q < q_<$ denote the classically allowed regions. The (blue) region to the left of the plot ($0 < q < q_<$) describes a big crunch scenario, wherein the classical domain the universe started from a zero size, reaches a maximum q value, and then reduces back to zero. While the (red) region to the right ($q_> < q < \infty$) describes the bouncing scenario, where the universe reaches a minimum value of q and then re-expands. Classically, a solution of the relevant Einstein's equations describes either of these two scenarios. However, quantum mechanically, tunnelling from one region to the other is allowed.

where, $|\mathbf{k}| = \sqrt{k_1^2 + k_2^2}$. Then $\psi_{|\mathbf{k}|}(q)$ can be shown to satisfy the following one-dimensional Schrödinger equation:

$$\left[-\frac{1}{2MV_3} \left(\frac{\partial^2}{\partial q^2} \right) + V_3 U_{\text{total}}(q) \right] \psi_{|\mathbf{k}|}(q) = 0, \quad (72)$$

where, the total effective potential $U_{\text{total}}(q)$ is given by,

$$U_{\text{total}}(q) \equiv U_{\text{eff}}(q) - \frac{\rho_\theta}{q^2}, \quad (73)$$

with the density $\rho_\theta \equiv (|\mathbf{k}|^2/2)$. We see that this is consistent with our classical analysis of the problem. Once again, we obtain the conditions Eq. (68) and Eq. (69), respectively, for the bounce to occur in the linear and quadratic models. The typical form of the total effective potential $U_{\text{total}}(q)$, when these conditions are met, is shown in Fig. 5.

Even though the bouncing scenario can be realized classically for the parameter ranges presented in Eq. (68) and Eq. (69), it is evident from Fig. 5 that there is also a region that corresponds to a classical crunch. Classically, however, a solution to Einstein's equations will correspond to either one of the two scenarios. Quantum mechanically, on the other hand, tunnelling is possible from one region to another. One can indeed compute the WKB probability $\mathcal{T}$ for tunnelling from the bouncing regime to the regime




Figure 6: The red graph shows $\mathcal{T}(\sigma_1)$ and the blue dashed graph shows $\mathcal{T}(\sigma_2)$, with $V_3\sqrt{2M\rho_0}$ fixed to a value of 50 in both the cases.

depicting classical crunch to get:

$$\mathcal{T} = \exp \left(-2 \int_{q_<}^{q_>} V_3 \sqrt{2MU_{\text{total}}(q)} dq \right) \equiv \exp \left[-2V_3 \sqrt{2M\rho_0} \Gamma \right], \quad (74)$$

where, $q_<$ and $q_>$ are classical turning points such that, $0 < q_< < q_>$ (see also Fig. 5). In the light of Eq. (68) and Eq. (69), it is convenient to define the following parameters: $\sigma_1 \equiv \sqrt{(27\rho_\theta/4\rho_0)}$ and $\sigma_2 \equiv \sqrt{(4\rho_\theta/\rho_0)}$, respectively, for the linear and quadratic models, such that $0 < \sigma_1, \sigma_2 < 1$. It is now straightforward to evaluate the defining integral in the exponential of Eq. (74) and hence determine $\Gamma(\sigma_1)$ and $\Gamma(\sigma_2)$ respectively. These can be expressed in terms of the Elliptic integrals with the following expressions:

$$\Gamma(\sigma_1) \approx (\alpha_1 - \alpha_2 \sigma_1) + \sigma_1^2 (\beta_1 + \beta_2 \ln \sigma_1) \quad (75)$$

$$\Gamma(\sigma_2) = \frac{\pi}{4} (1 - \sigma_2) \quad (76)$$

where, the coefficients appearing in the expression for $\Gamma(\sigma_1)$ has the following numerical expressions: $\alpha_1 = 0.666706$, $\alpha_2 = 0.607517$, $\beta_1 = -0.059189$ and $\beta_2 = 0.028686$, respectively. For illustration purpose, for a fixed value of $V_3\sqrt{2M\rho_0}$, the typical behaviour of the transition probability $\mathcal{T}$ as function of σ_1 and σ_2 , for the linear and the quadratic model are given in Fig. 6. As evident, when both σ_1 and σ_2 are near unity, i.e., when ρ_θ is comparable to ρ_0 , the tunnelling probability increases. However, normally we would like the contribution from the shear to be subdominant, and hence the tunnelling amplitude will be smaller.

5 Introducing matter field

So far, we have discussed the quantum cosmology of a universe with the matter source being a perfect fluid with energy density $\rho_{\text{eff}}(a)$, given by Eq. (24) and Eq. (25), respectively, such that the action functional becomes at most quadratic in the dynamical variable q . However, there can be further fundamental fields living in the spacetime, and for simplicity, we choose a single conformally coupled scalar field. This will

drive home the essential features arising out of addition of such matter fields in the quantum analysis of the bouncing model. Thus our starting point will be the Wheeler-DeWitt equation presented in Eq. (16), on which we will apply the WKB expansion and hence shall determine the behaviour of the matter fields in the classically allowed and in the classically forbidden regions explicitly. First of all, we will provide the key equations arising out of the WKB expansion of the Wheeler-DeWitt equation in the next section.

5.1 Wheeler-DeWitt equation with matter and the WKB expansion

As we have already remarked, we shall be treating the matter field ϕ , which is a conformally coupled scalar field appearing in Eq. (10), as a test field in a given background. The precise sense in which one can perform this analysis is through WKB expansion and it will be convenient for our purpose to follow the approach of [34, 35]. The starting point being the following ansatz for the Wheeler-DeWitt wave function $\Psi(q; \{\phi_k\})$ as a solution to Eq. (16), with the effective potentials of the form given in Eq. (27) or Eq. (28):

$$\Psi(q; \{\phi_k\}) = \exp \left[\frac{iV_3}{\ell_p^2} \sum_{j=0}^{\infty} \ell_p^{2j} S_j(q, \{\phi_k\}) \right] . \quad (77)$$

We now demand that $\Psi(q; \{\phi_k\})$ solves the Wheeler-DeWitt equation at all orders in ℓ_p^2 . At the first three leading orders, namely ℓ_p^{-4} , ℓ_p^{-2} and ℓ_p^0 , implies:

$$\partial_{\phi_k} S_0 = 0 ; \quad (78)$$

$$\left(\partial_q S_0^{(\text{I})} \right)^2 + \frac{h_n^2}{\left(\frac{n}{6} - 1 \right)^4} (1 - q) = 0 ; \quad (79)$$

$$\left(\partial_q S_0^{(\text{II})} \right)^2 + \left(\frac{16}{9} \right) \frac{h_n^2}{\left(\frac{n}{6} - 1 \right)^4} (1 - q^2) = 0 ; \quad (80)$$

$$-\frac{2i(\partial_q S_0)}{b^2} \partial_q \zeta(q, \{\phi_k\}) = V_3 \int \frac{d^3 p}{(2\pi)^3} \left(-\frac{1}{2\mu(q)} \partial_{\phi_p}^2 + \frac{1}{2} \mu(q) \omega_p(q)^2 \phi_p^2 \right) \zeta(q, \{\phi_k\}) , \quad (81)$$

where the function $\zeta(q, \{\phi_k\})$ is defined as,

$$\zeta(q, \{\phi_k\}) \equiv \frac{\sqrt{S'_0(q)}}{\sqrt{V_3}} e^{iV_3 S_1(q, \{\phi_k\})} . \quad (82)$$

The first equation $\partial_{\phi_k} S_0 = 0$, arising from setting the coefficient of ℓ_p^{-4} to zero, merely tells us that the zeroth order contribution to the action, i.e., S_0 is independent of the matter degrees of freedom ϕ_k . In addition, Eq. (79) and Eq. (80) are the Hamilton-Jacobi equations for the two scenarios of interest, as discussed in earlier sections. Finally, one can introduce the time coordinate t in the present formalism via the following relation:

$$-\frac{2(\partial_q S_0)}{b^2} = \frac{dq}{dt} . \quad (83)$$

With this definition, it is easy to see that Eq. (79) and Eq. (80) are equivalent to the classical constraint equation, given by Eq. (14), with the matter field neglected. Moreover, the third equation, namely Eq. (81),

takes form of the following time-dependent Schrödinger equation:

$$i\partial_t\zeta = V_3 \int \frac{d^3p}{(2\pi)^3} \left(-\frac{1}{2\mu(q(t))} \partial_{\phi_p}^2 + \frac{1}{2} \mu(q(t)) \omega_p^2(q(t)) \phi_p^2 \right) \zeta \quad (84)$$

where, $q(t)$ corresponds to the classical solutions of the purely gravitational part arising from Eq. (79) and Eq. (80), respectively. Therefore, Eq. (84) mathematically describes the paradigm of quantum field theory, with ϕ_k being the Fourier modes of the field in a time dependent background spacetime, which has been discussed extensively in the literature []. Following [], and keeping in mind the result that the function $\zeta(q(t); \{\phi_k\})$ satisfies the Schrödinger equation of a time-dependent harmonic oscillator, we have the following Gaussian solution for $\zeta(q(t); \{\phi_k\})$:

$$\zeta(q(t); \{\phi_k\}) \propto \exp \left[V_3 \int \frac{d^3k}{(2\pi)^3} \left(i\mu(t) \frac{\dot{u}_k}{2u_k} \phi_k^2 \right) \right]. \quad (85)$$

For the above solution for $\zeta(q(t); \{\phi_k\})$ to describe a coherent state of the scalar field and solve Eq. (84), the function $u_k(t)$ must satisfy the following differential equation:

$$\ddot{u}_k + \frac{\dot{\mu}}{\mu} \dot{u}_k + \omega_k^2 u_k = 0, \quad (86)$$

where, $\mu(t) \equiv \mu(q(t))$, $\omega_k^2(t) \equiv \omega_k^2(q(t))$ and the dot denotes a derivative with respect to the time coordinate t introduced in Eq. (83). Since, we want the solution for $\zeta(q(t); \{\phi_k\})$, presented in Eq. (85) to depict a stable situation, the real part of the argument of the exponential must be negative [28] and thus, in addition to Eq. (86), the solution u_k must satisfy the following condition:

$$\text{Re} \left[i\mu(t) \frac{\dot{u}_k}{u_k} \right] < 0. \quad (87)$$

In summary, we find that the leading order solution in ℓ_p of the Wheeler-DeWitt equation can be written as:

$$\Psi(q, \{\phi_k\}) \approx \frac{\sqrt{V_3}}{\sqrt{\partial_q S_0}} e^{i \frac{V_3}{\ell_p^2} S_0(q)} \zeta(q, \{\phi_k\}), \quad (88)$$

with $S_0(q)$ satisfying Eq. (79) or, Eq. (80) depending on the nature of the effective potential and ζ satisfies Eq. (84). This finishes our discussion on the WKB wave function for the bouncing models under consideration. In the subsequent sections we will analyze the nature of the solution u_k , which sits in the function $\zeta(q(t); \{\phi_k\})$ to completely understand the behaviour of $\Psi(q, \{\phi_k\})$. Since the effective potentials $U_{\text{eff}}(q)$ has classical turning points, pertaining to the bouncing models we are considering, it will be convenient to analyze the WKB solutions in the classically allowed and forbidden regions separately.

5.2 Classically allowed region

For the bouncing models under consideration, the classically allowed region corresponds to the following range of the dynamical variable: $1 < q < \infty$. As evident from Fig. 1, the effective potentials appearing in the Wheeler-DeWitt equation, i.e., $U_{\text{eff}}^{(\text{I})}(q)$ and $U_{\text{eff}}^{(\text{II})}(q)$, respectively take negative values and therefore, the solutions to Eq. (79) and Eq. (80), respectively, takes the form:

$$S_{0,\pm}^{(\text{I})}(q) = \pm \beta_n \ell_p^2 \int_1^{q(>1)} \sqrt{-(1-q')} dq'; \quad S_{0,\pm}^{(\text{II})}(q) = \pm \frac{4}{3} \beta_n \ell_p^2 \int_1^{q(>1)} \sqrt{-(1-q'^2)} dq'. \quad (89)$$

Here, we have used the definition of β_n , from Eq. (36). From Eq. (83), it follows that for $(\partial_q S_0) < 0$, we have $(dq/dt) > 0$, i.e., as the action decreases with q , the universe expands. Thus we notice that, $S_{0,-}^{(I,II)}(q)$ corresponds to the expanding phase, while, $S_{0,+}^{(I,II)}(q)$ corresponds to the contracting phase of the bouncing universe. Keeping this in mind, we integrate, Eq. (89) and then substituted the same in Eq. (88), to obtain the general solution to the Wheeler-DeWitt equation at $\mathcal{O}(\ell_p^0)$ as:

$$\Psi^{(I)}(q; \{\phi_k\}) \approx A_+^{(I)} \frac{e^{\frac{2i}{3}\alpha_n(q-1)^{\frac{3}{2}}}}{[\alpha_n^2(q-1)]^{\frac{1}{4}}} \zeta_+^{(I)}(q; \{\phi_k\}) + A_-^{(I)} \frac{e^{-\frac{2i}{3}\alpha_n(q-1)^{\frac{3}{2}}}}{[\alpha_n^2(q-1)]^{\frac{1}{4}}} \zeta_-^{(I)}(q; \{\phi_k\}) ; \quad (90)$$

$$\Psi^{(II)}(q; \{\phi_k\}) \approx A_+^{(II)} \frac{e^{\frac{4i}{3}\alpha_n\xi(q)}}{[\alpha_n^2(q^2-1)]^{\frac{1}{4}}} \zeta_+^{(II)}(q; \{\phi_k\}) + A_-^{(II)} \frac{e^{-\frac{4i}{3}\alpha_n\xi(q)}}{[\alpha_n^2(q^2-1)]^{\frac{1}{4}}} \zeta_-^{(II)}(q; \{\phi_k\}) . \quad (91)$$

In the above solutions of the Wheeler-DeWitt equation with matter fields, $A_{\pm}^{(I)}$ and $A_{\pm}^{(II)}$ are arbitrary constants, $\alpha_n \equiv V_3\beta_n$, with β_n defined in Eq. (36) and $\xi(q)$ is a function of the expansion parameter q , given by Eq. (42). In order to fix the unknown constants, we may impose the boundary condition that in the absence of the conformally coupled scalar field, the Wheeler-DeWitt wave function must take the form presented in Eq. (53) and Eq. (62), respectively for the two choices of the effective potential. This yields the following conditions on the unknown coefficients: $A_+^{(I)} = A_-^{(I)} \equiv A^{(I)}$ and $A_+^{(II)} = A_-^{(II)} \equiv A^{(II)}$. As emphasized above, these conditions ensure that the wave functions reduce to that of the Hawking-Hartle prescription when matter fields are neglected.

What remains, is to find out the appropriate solutions $\zeta(t)$ to Eq. (84), which appears in the solutions to the Wheeler-DeWitt equation directly. As remarked earlier, we shall take $\zeta(t)$ to be of the Gaussian form presented in Eq. (85) and hence the task of finding a solution for $\zeta(t)$ boils down to finding a solution for $u_k(t)$ from Eq. (86). Let us denote the independent solutions to Eq. (86) for the two effective potentials, as $u_{k,\pm}^{(I)}$ and $u_{k,\pm}^{(II)}$, respectively. These in turn will provide us the functions $\zeta_{\pm}^{(I)}$ and $\zeta_{\pm}^{(II)}$, used in Eq. (90) and Eq. (91), respectively. Thus the time dependent harmonic oscillator equation, Eq. (86) for the functions $u_{k,\pm}^{(I)}$ and $u_{k,\pm}^{(II)}$ take the following form:

$$\ddot{u}_{k,\pm}^{(I)} + \frac{\dot{\mu}^{(I)}}{\mu^{(I)}} \dot{u}_{k,\pm}^{(I)} + \left[\omega_{k,\pm}^{(I)}\right]^2 u_{k,\pm}^{(I)} = 0 , \quad (92)$$

$$\ddot{u}_{k,\pm}^{(II)} + \frac{\dot{\mu}^{(II)}}{\mu^{(II)}} \dot{u}_{k,\pm}^{(II)} + \left[\omega_{k,\pm}^{(II)}\right]^2 u_{k,\pm}^{(II)} = 0 , \quad (93)$$

where, the time dependent mass function $\mu(t)$ and the time dependent frequency function $\omega_k(t)$ has the following expressions, for the two choices of the effective potentials, appearing in the Wheeler-DeWitt equation:

$$\mu^{(I)}(t) = (1 + h_n^2 t^2)^{\frac{2(n-3)}{(n-6)}} ; \quad \omega_{k,\pm}^{(I)}(t)^2 = (1 + h_n^2 t^2)^{-\frac{4(n-3)}{(n-6)}} \left[k^2 + m^2 (1 + h_n^2 t^2)^{\frac{6}{6-n}} \right] . \quad (94)$$

$$\mu^{(II)}(t) = \left[\cosh\left(\frac{3}{2}h_n t\right) \right]^{\frac{2(n-2)}{(n-6)}} ; \quad \omega_{k,\pm}^{(II)}(t)^2 = \left[\cosh\left(\frac{3}{2}h_n t\right) \right]^{-\frac{4(n-2)}{(n-6)}} \left\{ k^2 + m^2 \left[\cosh\left(\frac{3}{2}h_n t\right) \right]^{\frac{8}{6-n}} \right\} . \quad (95)$$

The solutions to Eq. (92) and Eq. (93), which may lead to a stable wave function for the Wheeler-DeWitt equation are obtained by imposing the additional condition as in Eq. (87) on u_k . We shall now study various properties of the function u_k , leading to stable Wheeler-DeWitt wave functions.

For notational convenience, we shall momentarily ignore the subscripts ‘(I)’ and ‘(II)’ until absolutely necessary. The strategy will be as described in [], where solution generating technique of a time-dependent harmonic oscillator has been studied in great detail. The general solution to a time-dependent harmonic oscillator equation of the form Eq. (86) can be written in terms of a complex function $v_k(t)$ as follows:

$$u_k(t) = v_k^*(t) + B_k v_k(t) , \quad (96)$$

where, B_k is a constant and can be treated as the ratio between two Bogoliubov coefficients among the mode functions u_k and v_k . Additionally, we consider the functions v_k and v_k^* as linearly independent solutions of Eq. (86), that satisfies the following Wronskian condition:

$$i\mu(t) [v_k^* \dot{v}_k - v_k \dot{v}_k^*] = 1 . \quad (97)$$

For the solution u_k , presented above, we obtain: $\text{Re}[i\mu(t)(\dot{u}_k/u_k)] = \{(|B_k|^2 - 1)/2|u_k|^2\}$. Thus imposing the stability condition as presented in Eq. (87), on the solution u_k , we obtain,

$$|B_k|^2 < 1 . \quad (98)$$

It is worth mentioning that one can also express the functions $v_k(t)$ that satisfy Eq. (97) in terms of two real functions $\chi_k(t)$ and $\Theta_k(t)$ as follows: $v_k(t) = \chi_k(t)e^{-i\Theta_k(t)}$. Here, the phase factor $\Theta_k(t)$ is related to the amplitude $\chi_k(t)$ through the following relation: $\Theta_k(t) = \{1/2\mu(t)\chi_k^2(t)\}$ and the amplitude $\chi_k(t)$ satisfies the differential equation of a time-dependent harmonic oscillator with a source term, $\{1/4\mu(t)^2\chi_k^3(t)\}$. With this notation, general solutions $u_{k,\pm}^{(I)}$ and $u_{k,\pm}^{(II)}$ can be written as,

$$u_{k,\pm}^{(I,II)}(t) = \chi_k^{(I,II)}(t)e^{i\Theta_k^{(I,II)}(t)} + B_{k,\pm}^{(I,II)} \chi_k^{(I,II)}(t)e^{-i\Theta_k^{(I,II)}(t)} , \quad (99)$$

and the stability condition further implies the following restriction on the function $B_{k,\pm}^{(I,II)}$: $|B_{k,\pm}^{(I,II)}|^2 < 1$. Since the function $q(t)$, depicting expansion of the universe must be continuous at the bounce, i.e., at $q = 1$, it follows that: $\zeta_+^{(I,II)}(q = 1; \{\phi_k\}) = \zeta_-^{(I,II)}(q = 1; \{\phi_k\})$ [28, 36]. This condition translates to

$$B_{k,+}^{(I,II)} = B_{k,-}^{(I,II)} = B_k^{(I,II)} , \quad (100)$$

and all of these coefficients must have an absolute value less than unity. Thus a general solution to the Wheeler-DeWitt equation up to $\mathcal{O}(\ell_p^0)$, in the classical regime is given by Eq. (90) and Eq. (91), respectively, with the $\zeta(t)$ given by Eq. (85) and Eq. (99), keeping in mind the condition $|B_k|^2 < 1$, for stability. This completes our analysis of the solution to the Wheeler DeWitt equation in the classically allowed region. Next, we shall see how a similar analysis can be performed for the classically forbidden region as well.

5.3 Classically forbidden region

For completeness we present here an analysis of the wave function with matter field in the classically forbidden region. For the bouncing models we are considering, the classically forbidden region corresponds to the range $0 < q < 1$. In this range, the effective potentials $U_{\text{eff}}^{(I)}(q)$ and $U_{\text{eff}}^{(II)}(q)$ take positive values, as shown in Fig. 1. Thus using the same parametrizations as in the case of classically allowed region, the solutions to Eq. (79) and Eq. (80) can, therefore, be written as,

$$i\bar{S}_{0,\pm}^{(I)}(q) = \pm\beta_n\ell_p^2 \int_1^{q(<1)} dq' \sqrt{1-q'} ; \quad i\bar{S}_{0,\pm}^{(II)}(q) = \pm\frac{4}{3}\beta_n\ell_p^2 \int_1^{q(<1)} dq' \sqrt{1-q'^2} . \quad (101)$$

Note that the action $\bar{S}_{0,\pm}^{(I,II)}(q)$ is related to $S_{0,\pm}^{(I,II)}(q)$, derived in the previous section, through analytic continuation of the dynamical variable $q(t)$. Similarly, the time coordinate t must also be analytically continued, and hence following Eq. (83), it is convenient to define the Euclidean time coordinates $\tau_{\pm}$ in the following manner:

$$\frac{dq}{d\tau_{\pm}} = \mp i \frac{2(\partial_q \bar{S}_{0,\pm})}{b^2} . \quad (102)$$

It then follows that the Euclidean time coordinates $\tau_{\pm}$ defined above corresponds to the Wick rotations of the original Lorentzian time coordinate t , such that: $t \rightarrow i\tau_+$ and $t \rightarrow -i\tau_-$. Performing the Wick rotation and analytic continuation to the complex q plane, we can now write down the general solution to the Wheeler-DeWitt equation to $\mathcal{O}(\ell_p^0)$, in the classically forbidden region (i.e., $0 < q < 1$) as,

$$\Psi^{(I)}(q; \{\phi_k\}) \approx \bar{A}_+^{(I)} \frac{e^{-\frac{2}{3}\alpha_n(1-q)^{\frac{3}{2}}}}{[\alpha_n^2(1-q)]^{\frac{1}{4}}} \bar{\zeta}_+^{(I)}(q; \{\phi_k\}) + \bar{A}_-^{(I)} \frac{e^{\frac{2}{3}\alpha_n(1-q)^{\frac{3}{2}}}}{[\alpha_n^2(1-q)]^{\frac{1}{4}}} \bar{\zeta}_-^{(I)}(q; \{\phi_k\}) , \quad (103)$$

$$\Psi^{(II)}(q; \{\phi_k\}) \approx \bar{A}_+^{(II)} \frac{e^{-\frac{4}{3}\alpha_n \bar{\xi}(q)}}{[\alpha_n^2(1-q^2)]^{\frac{1}{4}}} \bar{\zeta}_+^{(II)}(q; \{\phi_k\}) + \bar{A}_-^{(II)} \frac{e^{\frac{4}{3}\alpha_n \bar{\xi}(q)}}{[\alpha_n^2(1-q^2)]^{\frac{1}{4}}} \bar{\zeta}_-^{(II)}(q; \{\phi_k\}) , \quad (104)$$

where, the function $\bar{\xi}(q)$ is defined as: $\bar{\xi}(q) = (1/2)[q\sqrt{1-q^2} - \cos^{-1}(q)]$. Further, the matter wave functions $\bar{\zeta}_{\pm}^{(I,II)}$ in the classically forbidden region in terms of the Euclidean time coordinate becomes,

$$\bar{\zeta}_{\pm}^{(I,II)}(\tau_{\pm}) \propto \exp \left[\int \frac{d^3k}{(2\pi)^3} \left(\pm \mu^{(I,II)} \frac{\bar{u}'_{k,\pm}{}^{(I,II)}}{2\bar{u}_{k,\pm}^{(I,II)}} \phi_k^2 \right) \right] , \quad (105)$$

where the prime denotes derivative with respect to the appropriate Euclidean time coordinates $\tau_{\pm}$ and, the functions $\bar{u}_{k,\pm}^{(I)}$ and $\bar{u}_{k,\pm}^{(II)}$ are solutions of the Euclidean versions of Eq. (92) and Eq. (93), respectively. For large values of α_n , we expect the wave functions $\Psi^{(I,II)}$ to exponentially decay to zero as $q \rightarrow 0$. Hence, we demand that the unknown coefficients should be such that, $\bar{A}_-^{(I)} = \bar{A}_-^{(II)} = 0$. This implies that the only Wick rotation relevant to the matter sector of the no-boundary wave function is $t \rightarrow i\tau_+$. Therefore, for notational convenience we shall henceforth avoid the subscript ‘+’ and express $\Psi^{(I,II)}$ in the classically forbidden region as:

$$\Psi^{(I)}(q; \{\phi_k\}) \approx \bar{A}^{(I)} \frac{e^{-\frac{2}{3}\alpha_n(1-q)^{\frac{3}{2}}}}{[\alpha_n^2(1-q)]^{\frac{1}{4}}} \bar{\zeta}^{(I)}(q; \{\phi_k\}) ; \quad (106)$$

$$\Psi^{(II)}(q; \{\phi_k\}) \approx \bar{A}^{(II)} \frac{e^{-\frac{4}{3}\alpha_n \bar{\xi}(q)}}{[\alpha_n^2(1-q^2)]^{\frac{1}{4}}} \bar{\zeta}^{(II)}(q; \{\phi_k\}) ; \quad (107)$$

where, $\bar{\zeta}^{(I,II)}(\tau)$ is given by Eq. (105), keeping in mind that only the ‘positive’ branch of the solution is allowed. Momentarily dropping the superscripts (I) and (II) for notational convenience, following our analysis for the classically allowed region, we can express the general solution $\bar{u}_k$ appearing in Eq. (105) in terms of two linearly independent solutions $\bar{v}_{k,1}$ and $\bar{v}_{k,2}$ of the time-dependent harmonic oscillator as:

$$\bar{u}_k = \bar{v}_{k,1} + \bar{B}_k \bar{v}_{k,2} . \quad (108)$$

Here, $\bar{B}_k$ is the ratio of the two Bogoliubov coefficients appearing from the rotation between the basis vectors $\bar{u}_k$ and $\bar{v}_k$. Following the analysis of the previous section, we may express the basis solutions $\bar{v}_{k,1}$ and $\bar{v}_{k,2}$ in terms of two real functions obtained by the Euclidean continuation of the solutions in the previous section, yielding,

$$\bar{v}_{k,1}(\tau) = \bar{\chi}_k(\tau)e^{-\bar{\Theta}_k(\tau)} ; \quad \bar{v}_{k,2}(\tau) = \bar{\chi}_k(\tau)e^{\bar{\Theta}_k(\tau)} , \quad (109)$$

where, $\bar{\chi}_k(\tau) \equiv \chi_k(i\tau)$ and $\bar{\Theta}'_k(\tau) \equiv \{1/2\mu(i\tau)\bar{\chi}_k^2(\tau)\}$. In terms of these two real functions along with the complex function $\bar{B}_k$, the stability condition presented in Eq. (87), for the classically forbidden region takes the following form:

$$\text{Re} \left[\mu(i\tau) \frac{\bar{u}'_k}{\bar{u}_k} \right] = \mu(i\tau) \left(\frac{\bar{\chi}'_k(\tau)}{\bar{\chi}_k(\tau)} + \frac{\bar{\Theta}'_k(\tau) (e^{4\bar{\Theta}_k(\tau)} |\bar{B}_k|^2 - 1)}{(e^{2\bar{\Theta}_k(\tau)} + \text{Re}[\bar{B}_k])^2 + \text{Im}[\bar{B}_k]^2} \right) < 0 \quad (110)$$

For the real function $\bar{\chi}_k(t)$, we may choose an initial condition, such that for some value of the Euclidean time τ_1 , $\{\bar{\chi}'_k(\tau_1)/\bar{\chi}_k(\tau_1)\} = 0$ and therefore, the stability condition presented in Eq. (110), at the Euclidean time τ_1 , translates to:

$$|\bar{B}_k^{(I,II)}|^2 < e^{-4\bar{\Theta}_k^{(I,II)}(\tau_1)} \quad (111)$$

where we have retained the superscripts for clarity. Additionally, the continuity at the classical turning point imposes the condition [28, 36]: $\bar{B}_k^{(I,II)} = B_k^{(I,II)}$. This implies that we cannot specify the matter-sector wave function uniquely, but only up to the constants $\bar{B}_k^{(I,II)}$. A special choice of $\bar{B}_k^{(I,II)}$ that corresponds to the condition that the solution $\bar{u}_k(\tau)$ vanish at $q = 0$ is given by

$$\bar{B}_k^{(I,II)} = -e^{-2\bar{\Theta}_k^{(I,II)}(\tau_0)} \quad (112)$$

where, τ_0 is defined via $q(\tau_0) = 0$. This choice may be considered as the generalization of the Euclidean vacuum introduced by Hartle and Hawking in the context of de Sitter spacetime. An interesting special case occurs when $\bar{\Theta}_k(\tau_0) \rightarrow \infty$ for all values of k . If this condition is satisfied, we can choose $\tau_1 = \tau_0$ and, therefore, Eq. (111) implies $\bar{B}_k^{(I,II)} = 0$, which in turn implies that a unique wavefunction exists.

6 Discussion

Bouncing cosmologies, generally, aim to solve the fundamental issues of the SBC without resorting to the inflation mechanism. Moreover, certain shortcomings of the inflationary paradigm, such as the ‘Trans-Planckian’ and the singularity problem, can be bypassed in a bouncing scenario. Consequently, cosmological models with a bounce, as against a singularity, are gaining interest as viable alternatives to the much-celebrated inflationary models. In light of this, we attempted to analyze the quantum aspects of bouncing cosmologies, which will be of relevance, especially, to the study of quantum seeds of structure formation.

Since there are several, conceptually disparate, routes to realize a bounce, it is practically difficult to adopt a single framework to study the set of *all* bouncing models in its entirety. Therefore, we employed a phenomenological approach, that accommodates the essential aspects of a wide class of bouncing models, while at the same time, render analytical calculations tractable. There are two key aspects to this approach;

one concerns the kinematics, while the other concerns the dynamics of the set of bouncing universes of our current interest. A convenient parametrization of the FRLW spacetime in terms of the variable $q(t)$, as given in Eq. (1), together with the condition $p = 4 - 3b$, is the kinematic ingredient of our scheme. Two classes of effective perfect-fluids that source the background FRLW spacetime, which are described by their respective energy densities $\rho_{\text{eff}}^{(\text{I})}$ and $\rho_{\text{eff}}^{(\text{II})}$, constitute the second, dynamical aspect. The aforementioned, specific parametrization of the FRLW metric was so chosen, such that it leaves the gravitational part of the action a quadratic functional of $\dot{q}(t)$. On the other hand, the specific form of the energy densities $\rho_{\text{eff}}^{(\text{I})}$ and $\rho_{\text{eff}}^{(\text{II})}$, as presented in Eq. (24) and Eq. (25), were so chosen such that: (1) the classical solutions of the corresponding equations of motion describe bouncing scenarios and (2) for an appropriate choice of the parameter b , the perfect-fluid part of the action is a linear/quadratic functional of $q(t)$, for densities $\rho_{\text{eff}}^{(\text{I})}$ and $\rho_{\text{eff}}^{(\text{II})}$, respectively. Hence, we refer to the class of bouncing universes described by $\rho_{\text{eff}}^{(\text{I})}$ as the ‘linear model’ and that by $\rho_{\text{eff}}^{(\text{II})}$ as the ‘quadratic model’. The kinematic and dynamic aspects of our approach, together, enable us to study two wide classes of bouncing cosmologies with considerable analytic comfort.

We explored the quantum aspects of our model using the framework of minisuperspaces. To this end, we have followed two different approaches: (1) using the appropriate minisuperspace Wheeler-de Witt equations and (2) using the minisuperspace path integral. Thanks to the specific choice of the dynamical variable $q(t)$ and the density of the effective fluid, we can find the solutions of the gravitational Wheeler-de Witt equation, for both classes of bouncing models we introduced, analytically. These solutions encode the initial conditions of the universe and hence, is of importance to early universe cosmology. In the context of de Sitter spacetime, a particular solution to the corresponding Wheeler-De Witt equation, namely, the no-boundary wavefunction, is widely utilized in cosmology for its several appealing features. On account of this, it is worthwhile to investigate the properties of the natural generalization of the no-boundary wavefunction for bouncing cosmologies. We presented the explicit expressions for the gravitational part of the bouncing-model counterpart of the no-boundary wavefunction, which are denoted by $\Psi^{(\text{I})}(q)$ and $\Psi^{(\text{II})}(q)$, respectively, for the linear and quadratic models.

The no-boundary wavefunction was originally envisioned as arising out of a Euclidean path integral over compact and regular metrics. Recently, it has become clear that, when one attempts a more rigorous calculation, such a Euclidean path integral is ill-defined and divergent. Moreover, even though a well defined, convergent Lorentzian path integral, with the initial condition corresponding to a zero-sized universe, can give rise to the no-boundary wavefunction, it leads to unstable perturbation. These issues can, however, be circumvented by imposing an initial condition corresponding to a well defined Euclidean momentum associated with the scale factor. Motivated by this, we investigated the extension of this new path integral approach, to the case of bouncing cosmologies. We found that the dominant contribution to the bouncing-model counterpart of the no-boundary wavefunction comes from a geometry that corresponds to a spacetime that evolves from zero size to a finite size. One is, thus, reminded of the Hawking-Hartle saddle geometry that appears in the context of Euclidean path integral approach to de Sitter cosmology. But, in view of the fact that one is fixing the initial momentum, as against setting the initial size to zero, in the new approach, the off-shell geometries can have *any* initial size. Consequently, one must not interpret the analogue of the no-boundary wavefunctions thus obtained, as the amplitudes for creation of the universe from ‘nothing’. A more sensible interpretation would be, that the wavefunctions are amplitudes for the transition from an initial state of well defined Euclidean momentum to a state of well-defined size.

Finally, we introduced a real scalar field conformally coupled to the background FRLW spacetime. The quantum theory of the gravity-scalar system can be studied using the corresponding Wheeler-de Witt equation. Since exact solutions could not be found in this case, we progressed by assuming that the scalar

fields are perturbations at the quantum level. To enforce this assumption mathematically, we started off by taking the solution to be exponential of a power series in ℓ_p^2 , with the lowest power being ℓ_p^{-2} . Demanding that this wavefunction solves the corresponding Wheeler-de Witt equation at all orders of ℓ_p^2 , implies that the coefficients of ℓ_p^2 in the power series that define the wavefunction satisfy a set of differential equations. At the first two leading orders, the corresponding differential equation satisfied by the coefficients can be shown to be equivalent to the Hamilton-Jacobi equation of the gravitational part. In the next order, we obtain a differential equation that essentially describes the paradigm of quantum field theory in curved spacetime. We then proceeded to find the bouncing-model analogue of the wavefunction that corresponds to the no-boundary proposal for the gravity-scalar system. Interestingly, we found that the wavefunction, in general, is not unique. Only when $\Theta_k(\tau)$, a certain function of the Euclidean time τ , diverges at the point of singularity, do we get a unique wavefunction. This has the consequence that the initial conditions of the universe have a certain level of arbitrariness that cannot, in general, be fixed without invoking further principles.

Acknowledgements

The authors gratefully acknowledge several helpful comments from L. Sriramkumar, which has improved the content of this work. Research of S.C. is funded by the INSPIRE Faculty fellowship from the DST, Government of India (Reg. No. DST/INSPIRE/04/2018/000893) and by the Start-Up Research Grant from SERB, DST, Government of India (Reg. No. SRG/2020/000409).

References

- [1] A. H. Guth, “The Inflationary Universe: A Possible Solution to the Horizon and Flatness Problems,” *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 139–148.
- [2] K. Sato, “First Order Phase Transition of a Vacuum and Expansion of the Universe,” *Mon. Not. Roy. Astron. Soc.* **195** (1981) 467–479.
- [3] A. D. Linde, “A New Inflationary Universe Scenario: A Possible Solution of the Horizon, Flatness, Homogeneity, Isotropy and Primordial Monopole Problems,” *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 149–153.
- [4] A. Albrecht and P. J. Steinhardt, “Cosmology for Grand Unified Theories with Radiatively Induced Symmetry Breaking,” *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 158–161.
- [5] A. A. Starobinsky, “A New Type of Isotropic Cosmological Models Without Singularity,” *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 130–133.
- [6] S. Hawking and R. Penrose, “The Singularities of gravitational collapse and cosmology,” *Proc. Roy. Soc. Lond. A* **314** (1970) 529–548.
- [7] A. Borde and A. Vilenkin, “Eternal inflation and the initial singularity,” *Phys. Rev. Lett.* **72** (1994) 3305–3309, [arXiv:gr-qc/9312022](#).
- [8] J. Martin and R. H. Brandenberger, “The TransPlanckian problem of inflationary cosmology,” *Phys. Rev. D* **63** (2001) 123501, [arXiv:hep-th/0005209](#).

- [9] R. H. Brandenberger and J. Martin, “Trans-planckian issues for inflationary cosmology,” *Classical and Quantum Gravity* **30** no. 11, (Apr, 2013) 113001.
<http://dx.doi.org/10.1088/0264-9381/30/11/113001>.
- [10] F. Finelli and R. Brandenberger, “On the generation of a scale invariant spectrum of adiabatic fluctuations in cosmological models with a contracting phase,” *Phys. Rev. D* **65** (2002) 103522, [arXiv:hep-th/0112249](#).
- [11] R. H. Brandenberger, “The Matter Bounce Alternative to Inflationary Cosmology,” [arXiv:1206.4196 \[astro-ph.CO\]](#).
- [12] M. Gasperini and G. Veneziano, “Pre - big bang in string cosmology,” *Astropart. Phys.* **1** (1993) 317–339, [arXiv:hep-th/9211021](#).
- [13] R. Brandenberger and P. Peter, “Bouncing Cosmologies: Progress and Problems,” *Found. Phys.* **47** no. 6, (2017) 797–850, [arXiv:1603.05834 \[hep-th\]](#).
- [14] D. Battefeld and P. Peter, “A Critical Review of Classical Bouncing Cosmologies,” *Phys. Rept.* **571** (2015) 1–66, [arXiv:1406.2790 \[astro-ph.CO\]](#).
- [15] J. Feldbrugge, J.-L. Lehners, and N. Turok, “Lorentzian Quantum Cosmology,” *Phys. Rev. D* **95** no. 10, (2017) 103508, [arXiv:1703.02076 \[hep-th\]](#).
- [16] A. Di Tucci and J.-L. Lehners, “No-Boundary Proposal as a Path Integral with Robin Boundary Conditions,” *Phys. Rev. Lett.* **122** no. 20, (2019) 201302, [arXiv:1903.06757 \[hep-th\]](#).
- [17] A. Di Tucci, J.-L. Lehners, and L. Sberna, “No-boundary prescriptions in lorentzian quantum cosmology,” *Phys. Rev. D* **100** (Dec, 2019) 123543.
<https://link.aps.org/doi/10.1103/PhysRevD.100.123543>.
- [18] J. Louko, “Semiclassical Path Measure and Factor Ordering in Quantum Cosmology,” *Annals Phys.* **181** (1988) 318–373.
- [19] J. Louko and P. J. Ruback, “SPATIALLY FLAT QUANTUM COSMOLOGY,” *Class. Quant. Grav.* **8** (1991) 91–122.
- [20] J. J. Halliwell, “Derivation of the wheeler-dewitt equation from a path integral for minisuperspace models,” *Phys. Rev. D* **38** (Oct, 1988) 2468–2481.
<https://link.aps.org/doi/10.1103/PhysRevD.38.2468>.
- [21] K. Rajeev, “Complex time route to quantum backreaction,” *Eur. Phys. J. C* **79** no. 11, (2019) 959, [arXiv:2001.02543 \[gr-qc\]](#).
- [22] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 1973.
- [23] R. N. Raveendran, D. Chowdhury, and L. Sriramkumar, “Viable tensor-to-scalar ratio in a symmetric matter bounce,” *JCAP* **01** (2018) 030, [arXiv:1703.10061 \[gr-qc\]](#).
- [24] R. N. Raveendran and L. Sriramkumar, “Viable scalar spectral tilt and tensor-to-scalar ratio in near-matter bounces,” *Phys. Rev. D* **100** no. 8, (2019) 083523, [arXiv:1812.06803 \[astro-ph.CO\]](#).

- [25] Y.-F. Cai, T. Qiu, Y.-S. Piao, M. Li, and X. Zhang, “Bouncing universe with quintom matter,” *JHEP* **10** (2007) 071, [arXiv:0704.1090 \[gr-qc\]](#).
- [26] R. Brandenberger, “Matter Bounce in Horava-Lifshitz Cosmology,” *Phys. Rev. D* **80** (2009) 043516, [arXiv:0904.2835 \[hep-th\]](#).
- [27] J. Hartle and S. Hawking, “Wave Function of the Universe,” *Adv. Ser. Astrophys. Cosmol.* **3** (1987) 174–189.
- [28] S. Wada, “Quantum cosmological perturbations in pure gravity,” *Nuclear Physics B* **276** no. 3, (1986) 729 – 743. <http://www.sciencedirect.com/science/article/pii/0550321386900738>.
- [29] A. Vilenkin, “Quantum Cosmology and the Initial State of the Universe,” *Phys. Rev. D* **37** (1988) 888.
- [30] F. W. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, *NIST handbook of mathematical functions hardback and CD-ROM*. Cambridge university press, 2010.
- [31] J. Feldbrugge, J.-L. Lehners, and N. Turok, “No smooth beginning for spacetime,” *Phys. Rev. Lett.* **119** no. 17, (2017) 171301, [arXiv:1705.00192 \[hep-th\]](#).
- [32] J. Feldbrugge, J.-L. Lehners, and N. Turok, “No rescue for the no boundary proposal: Pointers to the future of quantum cosmology,” *Phys. Rev. D* **97** no. 2, (2018) 023509, [arXiv:1708.05104 \[hep-th\]](#).
- [33] Y.-F. Cai, R. Brandenberger, and P. Peter, “Anisotropy in a non-singular bounce,” *Classical and Quantum Gravity* **30** no. 7, (Mar, 2013) 075019. <http://dx.doi.org/10.1088/0264-9381/30/7/075019>.
- [34] T. Singh and T. Padmanabhan, “NOTES ON SEMICLASSICAL GRAVITY,” *Annals Phys.* **196** (1989) 296–344.
- [35] T. Padmanabhan and T. Singh, “On the semiclassical limit of the Wheeler-DeWitt equation,” *Class. Quant. Grav.* **7** (1990) 411–426.
- [36] T. Vachaspati and A. Vilenkin, “On the Uniqueness of the Tunneling Wave Function of the Universe,” *Phys. Rev. D* **37** (1988) 898.