

On the trace anomaly of Chaudhuri-Choi-Rabinovici model

Yu Nakayama

Department of Physics, Rikkyo University, Toshima, Tokyo 171-8501, Japan

Abstract

Recently a non-supersymmetric conformal field theory with an exactly marginal deformation in the large N limit was constructed by Chaudhuri-Choi-Rabinovici. On a non-supersymmetric conformal manifold, c coefficient of the trace anomaly in four dimensions would generically change. In this model, we, however, find that it does not change at the first non-trivial order given by three-loop diagrams.

In four-dimensional conformal field theories, the trace anomaly has the form

$$T_{\mu}^{\mu} = c\text{Weyl}^2 - a\text{Euler} \quad (1)$$

and it is known that coefficient a cannot change under exactly marginal deformations, but coefficient c may [1][2][3][4][5][6][7]. However, there has been no explicit field theory example where c changes (except for the effective holographic constructions in [2]). The main obstruction has been that we have no good examples of non-supersymmetric conformal field theories with exactly marginal deformations; in superconformal field theories, while it is easier to realize exactly marginal deformations, c does not change [8].

Recently, Chaudhuri-Choi-Rabinovici have constructed a non-supersymmetric conformal field theory with an exactly marginal deformation in the large N limit [9].¹ This theory may serve as a first non-trivial check if c can really change under exactly marginal deformations. In this short note, we, however, show that it does not change at the first non-trivial order given by three-loop diagrams.

The model (called complex bifundamental model in [9]) is given by four $SU(N_c)$ gauge theories with names 1, 1', 2 and 2', each of which has N_f Dirac fermions in the fundamental representation. We have two complex scalars in the bifundamental representations Φ_1 (under gauge group 1 and 1') and Φ_2 (under gauge group 2 and 2'). It has no Yukawa interaction, absence of which is protected by chiral symmetry, but it has a scalar potential

$$V = \tilde{h}_1 \text{Tr}[\Phi_1^{\dagger} \Phi_1 \Phi_1^{\dagger} \Phi_1] + \tilde{h}_2 \text{Tr}[\Phi_2^{\dagger} \Phi_2 \Phi_2^{\dagger} \Phi_2] \\ + \tilde{f}_1 \text{Tr}[\Phi_1^{\dagger} \Phi_1] \text{Tr}[\Phi_1^{\dagger} \Phi_1] + \tilde{f}_2 \text{Tr}[\Phi_2^{\dagger} \Phi_2] \text{Tr}[\Phi_2^{\dagger} \Phi_2] + 2\tilde{\zeta} \text{Tr}[\Phi_1^{\dagger} \Phi_1] \text{Tr}[\Phi_2^{\dagger} \Phi_2]. \quad (2)$$

We take the Veneziano limit of $N_c, N_f \rightarrow \infty$ with fixed $x = \frac{N_f}{N_c}$ and consider the limit $x \rightarrow \frac{21}{4}$ to make the theory weakly coupled.

In terms of rescaled coupling constants ($i = 1, 2$)

$$\lambda_i = \frac{N_c g_i^2}{16\pi^2}, \quad h_i = \frac{N_c \tilde{h}_i}{16\pi^2}, \quad f_i = \frac{N_c^2 \tilde{f}_i}{16\pi^2}, \quad \zeta = \frac{N_c^2 \tilde{\zeta}}{16\pi^2}, \quad (3)$$

the renormalization group beta functions in the Veneziano limit are expressed as (no sum over i unless explicitly shown)

$$\beta_{\lambda_i} = -\frac{21 - 4x}{3} \lambda_i^2 + \frac{-54 + 26x}{3} \lambda_i^3$$

¹See also [10][11] for other recently constructed examples of non-supersymmetric field theories with exactly marginal deformations in different dimensions than four.

$$\begin{aligned}
\beta_{h_i} &= 8h_i^2 - 12\lambda_i h_i + \frac{3}{2}\lambda_i^2 \\
\beta_{f_i} &= 4f_i^2 + 16f_i h_i + 12h_i^2 + 4\zeta^2 - 12\lambda_i f_i + \frac{9}{2}\lambda_i^2 \\
\beta_\zeta &= \zeta \sum_{i=1}^2 (4f_i + 8h_i - 6\lambda_i) .
\end{aligned} \tag{4}$$

The zero of the beta functions was studied in [9] and they found that there exists a conformal manifold given by

$$\begin{aligned}
\lambda_1 = \lambda_2 = \lambda &= \frac{21 - 4x}{-54 + 26x} \\
h_1 = h_2 &= \frac{3 - \sqrt{6}}{4}\lambda \\
f_p \equiv \frac{f_1 + f_2}{2} &= \sqrt{\frac{3}{2}}\lambda \\
\zeta^2 + f_m^2 &= \frac{18\sqrt{6} - 39}{16}\lambda^2 ,
\end{aligned} \tag{5}$$

where $f_m \equiv \frac{f_1 - f_2}{2}$. From the last line of (5), we see that it has the topology of a circle. As long as λ is small, we may neglect higher order corrections.

We now ask if the coefficient c in the trace anomaly can change on this conformal manifold. In addition to the coupling constant independent contributions from the one-loop diagrams (that count a number of fields), the coupling constant dependent contributions to the trace anomaly that are relevant for us come from the three-loop diagrams shown in Fig 1. The detailed computation for diagram (A) (as well as other two-loop diagrams) can be found in [12][13][14],² but we only need the relative coefficient, so we can simply work on combinatorics.

Up to an overall proportionality factor, the result in the Veneziano limit is summarized as

$$c_{2,3\text{-loop}} = -4f_m^2 - 4\zeta^2 + c_\lambda \lambda^2 \tag{6}$$


on the conformal manifold, where c_λ is some numerical constant, which is unimportant for our discussions.³ Since the relative coefficient appearing here coincides what appears

²The three-loop diagrams of (B)(C)(D) are not evaluated in the literature, but we see that diagram (B) and (C) do not contribute to c . Diagram (D) may contribute in general, but the contributions to c in our theory do not depend on ζ or f_m from the symmetry of the diagrams.


³A typo in the two-loop gauge contribution [14] that could affect c_λ has been corrected in [15].

in the last line of (5), we conclude that c does not change on the conformal manifold although the value itself is perturbatively corrected. We also note that these two- and three-loop diagrams do not change the value of a as anticipated [1][16] (rather trivially without cancellation unlike c).


The result is surprising in the sense that we generically expect that c would change on non-supersymmetric conformal manifold. It is an interesting question to see if the higher loop corrections modify our conclusion. It may be possible to relate the all-loop argument for the existence of the exactly marginal deformation in [9] with the computation of c by closing all the external lines in beta functions to make vacuum diagrams.




(A)



(B)



(C)



(D)

Fig 1: Three-loop Feynman diagrams that could contribute to c .

Acknowledgements

This work is in part supported by JSPS KAKENHI Grant Number 17K14301. It is motivated from the online talk by Z. Komargodski at YITP workshop on Strings and Fields 2020, which the author watched on Youtube later.

References

- [1] H. Osborn, Nucl. Phys. B **363**, 486 (1991). doi:10.1016/0550-3213(91)80030-P
- [2] Y. Nakayama, JHEP **07**, 004 (2017) doi:10.1007/JHEP07(2017)004 [arXiv:1702.02324 [hep-th]].
- [3] D. Meltzer and E. Perlmutter, JHEP **07**, 157 (2018) doi:10.1007/JHEP07(2018)157 [arXiv:1712.04861 [hep-th]].
- [4] A. Bzowski, P. McFadden and K. Skenderis, JHEP **11**, 159 (2018) doi:10.1007/JHEP11(2018)159 [arXiv:1805.12100 [hep-th]].
- [5] S. N. Solodukhin, Phys. Lett. B **802**, 135235 (2020) doi:10.1016/j.physletb.2020.135235 [arXiv:1907.07916 [hep-th]].
- [6] V. Niarchos, C. Papageorgakis and E. Pomoni, JHEP **04**, 048 (2020) doi:10.1007/JHEP04(2020)048 [arXiv:1911.05827 [hep-th]].
- [7] V. Niarchos, C. Papageorgakis, A. Pini and E. Pomoni, [arXiv:2009.08375 [hep-th]].

- [8] D. Anselmi, D. Z. Freedman, M. T. Grisaru and A. A. Johansen, Nucl. Phys. B **526**, 543 (1998) doi:10.1016/S0550-3213(98)00278-8 [hep-th/9708042].
- [9] S. Chaudhuri, C. Choi and E. Rabinovici, [arXiv:2011.13981 [hep-th]].
- [10] N. Chai, S. Chaudhuri, C. Choi, Z. Komargodski, E. Rabinovici and M. Smolkin, Phys. Rev. Lett. **125**, no.13, 131603 (2020) doi:10.1103/PhysRevLett.125.131603
- [11] N. Chai, S. Chaudhuri, C. Choi, Z. Komargodski, E. Rabinovici and M. Smolkin, Phys. Rev. D **102**, no.6, 065014 (2020) doi:10.1103/PhysRevD.102.065014 [arXiv:2005.03676 [hep-th]].
- [12] I. Jack and H. Osborn, Nucl. Phys. B **234**, 331-364 (1984) doi:10.1016/0550-3213(84)90067-1
- [13] I. Jack, Nucl. Phys. B **234**, 365-378 (1984) doi:10.1016/0550-3213(84)90068-3
- [14] I. Jack, Nucl. Phys. B **253**, 323-352 (1985) doi:10.1016/0550-3213(85)90534-6
- [15] H. Osborn and A. Stergiou, JHEP **06**, 079 (2016) doi:10.1007/JHEP06(2016)079 [arXiv:1603.07307 [hep-th]].
- [16] Z. Komargodski and A. Schwimmer, JHEP **1112**, 099 (2011) doi:10.1007/JHEP12(2011)099 [arXiv:1107.3987 [hep-th]].