Existence results for a generalized mean field equation on a closed Riemann surface*

Linlin Sun^{a,b,*}, Yamin Wang^c, Yunyan Yang^c

^aSchool of Mathematics and Statistics, Wuhan University, Wuhan 430072, China
^bHubei Key Laboratory of Computational Science, Wuhan University, Wuhan, 430072, China
^cDepartment of Mathematics, Renmin University of China, Beijing 100872, China

Abstract

Let Σ be a closed Riemann surface, h a positive smooth function on Σ , ρ and α real numbers. In this paper, we study a generalized mean field equation

$$-\Delta u = \rho \left(\frac{he^{u}}{\int_{\Sigma} he^{u}} - \frac{1}{\operatorname{Area}(\Sigma)} \right) + \alpha \left(u - \int_{\Sigma} u \right),$$

where Δ denotes the Laplace-Beltrami operator. We first derive a uniform bound for solutions when $\rho \in (8k\pi, 8(k+1)\pi)$ for some non-negative integer number $k \in \mathbb{N}$ and $\alpha \notin \operatorname{Spec}(-\Delta) \setminus \{0\}$. Then we obtain existence results for $\alpha < \lambda_1(\Sigma)$ by using the Leray-Schauder degree theory and the minimax method, where $\lambda_1(\Sigma)$ is the first positive eigenvalue for $-\Delta$.

Keywords: generalized mean field equation, blow-up analysis, topological degree, minimax method 2020 MSC: 53C21, 58J05, 35J20, 35J61

1. Introduction

Let Σ be a closed Riemann surface with area one and Δ be the Laplace-Beltrami operator. The mean field equation is stated as follows

$$-\Delta u = \rho \left(\frac{he^u}{\int_{\Sigma} he^u} - 1 \right),\tag{1.1}$$

where ρ is a real number and h is a smooth function on Σ . It comes from the prescribed Gaussian curvature problem [4, 31, 9, 10, 8, 14, 27], and also appears in various context such as the abelian Chern-Simons-Higgs models [37, 33, 36, 35, 29, 28, 6, 39, 41].

The existence of solutions of the mean field equation has been widely studied in recent decades. Recall the strong Trudinger-Moser inequality [24, Theorem 1.7]

$$\sup_{u \in H^1(\Sigma), \int_{\Sigma} |\nabla u|^2 \le 1, \int_{\Sigma} u = 0} \int_{\Sigma} \exp\left(4\pi u^2\right) < \infty, \tag{1.2}$$

which implies the Trudinger-Moser inequality

$$\ln \int_{\Sigma} e^{u} \le \frac{1}{16\pi} \int_{\Sigma} |\nabla u|^{2} + \int_{\Sigma} u + c, \tag{1.3}$$

Email addresses: sunll@whu.edu.cn (Linlin Sun), 2017100918@ruc.edu.cn (Yamin Wang), yunyanyang@ruc.edu.cn (Yunyan Yang)

Preprint submitted to XXX

November 23, 2021

^{*}This work is partially supported by the National Natural Science Foundation of China (Grant Nos. 11971358, 11801420, 11721101), and by the National Key Research and Development Project SQ2020YFA070080.

^{*}Corresponding author.

where c is a uniform constant depends only on the geometry of Σ . Consequently, for $\rho < 8\pi$, the Trudinger-Moser inequality (1.3) gives a minimizer to the action functional

$$H^{1}(\Sigma) \ni u \mapsto \frac{1}{2} \int_{\Sigma} |\nabla u|^{2} + \rho \left(\int_{\Sigma} u - \ln \left| \int_{\Sigma} h e^{u} \right| \right).$$

Many partial existence results have been obtained for noncritical cases, see for examples Struwe and Tarantello [40], Ding, Jost, Li and Wang [21], Chen and Lin [12], Djadli [22] and the references therein. For the first critical case $\rho = 8\pi$ and h > 0, several sufficient conditions for the existence to the Nirenberg problem (i.e., $\rho = 8\pi$ and the genus is zero) were given by Moser [32], Aubin [1], Chang and Yang [9, 10], Ji [30] and others; a sufficient condition for the minimizing solution to the mean field equation on a closed Riemann surface Σ with positive genus was given by Ding, Jost, Li and Wang [20]; it was also independently proved by Nolasco and Tarantello [34] when Σ is a flat torus. Chen and Lin [11] obtained an existence result for general critical cases (i.e., $\rho = 8k\pi$) which generalized Ding, Jost, Li and Wang's result [20]. For sign-changing potential h, we refer the reader to [18, 19, 17, 27, 13, 14] and the references therein. For the uniqueness to (1.1) we refer the reader to Gui and Moradifam [26], Shi, Sun, Tian and Wei [38] and others. We refer to [42] for a nice survey on the mean field equation.

Among various improvements of (1.2), it was proved by Yang [43] that for all $\alpha < \lambda_1(\Sigma)$, the first positive eigenvalue of the negative Laplacian $-\Delta$, there holds

$$\sup_{u \in H^1(\Sigma), \int_{\Sigma} |\nabla u|^2 - \alpha \int_{\Sigma} u^2 \le 1, \int_{\Sigma} u = 0} \int_{\Sigma} e^{4\pi u^2} < \infty.$$
(1.4)

This leads to an analog of (1.3), namely for any $\alpha < \lambda_1(\Sigma)$, there exists some constant c depending only on the geometry of Σ such that for all $u \in H^1(\Sigma)$ with $\int_{\Sigma} u = 0$,

$$\ln \int_{\Sigma} e^{u} \le \frac{1}{16\pi} \int_{\Sigma} \left(|\nabla u|^{2} - \alpha u^{2} \right) + c. \tag{1.5}$$

In view of (1.4) and (1.5), it is natural to consider the following generalized mean field equation

$$-\Delta u = \rho \left(\frac{he^u}{\int_{\Sigma} he^u} - 1 \right) + \alpha \left(u - \int_{\Sigma} u \right), \tag{1.6}$$

where h is a smooth positive function on Σ and $\rho, \alpha \in \mathbb{R}$. The related functional would be written as

$$J_{\rho,\alpha}(u) = J_{\rho,\alpha,h}(u) = \frac{1}{2\rho} \int_{\Sigma} \left(|\nabla u|^2 - \alpha \left(u - \int_{\Sigma} u \right)^2 \right) + \int_{\Sigma} u - \ln \int_{\Sigma} h e^u, \quad u \in H^1(\Sigma).$$
 (1.7)

As an immediately consequence of (1.5), there exists a solution to (1.6) when $\rho < 8\pi$ and $\alpha < \lambda_1$ (Σ). In [44], Yang and Zhu gave a sufficient condition such that (1.6) has a solution when $\rho = 8\pi$ and $\alpha < \lambda_1$ (Σ).

Our aim is to study the existence problem for (1.6) when $\rho \neq 8k\pi$ with $k \in \mathbb{N}$ and $\alpha < \lambda_1$ (Σ). To achieve this goal, we begin by studying the blow-up phenomena for the generalized mean field equation (1.6). For the classical case (i.e., $\alpha = 0$), it is well known that the blow-up phenomena only occurs if the parameter ρ is a multiple of 8π . Therefore the set of solutions is compact when $\rho \in \mathbb{R} \setminus 8\pi\mathbb{N}^*$. Our first main theorem in the following can be viewed as an analogous conclusion.

Theorem 1.1. Let I, J, K be compact subsets with $I \subset \mathbb{R} \setminus 8\pi\mathbb{N}^*, J \subset (\mathbb{R} \setminus \text{Spec}(-\Delta)) \cup \{0\}$ and

$$K \subset \left\{ h \in C^{1,\tau}(\Sigma) : h \text{ is a positive function} \right\}$$

where $0 < \tau < 1$. Then there exists a constant C such that

$$||u||_{C^{1,\tau}(\Sigma)} \leq C$$

for all solutions u to (1.6) with

$$\rho \in I, \quad \alpha \in J, \quad h \in K.$$

Remark 1.2. We give an example to show that the blow-up happens for $0 < \rho < 8\pi$ and $\alpha = \lambda_1(\Sigma)$. Let $J_{\rho,\alpha}$ be defined as in (1.7). Set $\alpha_n = \lambda_1(\Sigma) - 1/n$ and let u_n be a minimizer of J_{ρ,α_n} by the Trudinger-Moser inequality (1.5). Then a straightforward calculation shows

$$\lim_{n\to\infty} J_{\rho,\alpha_n}(u_n) = \inf_{u\in H^1(\Sigma)} J_{\rho,\lambda_1(\Sigma)}(u) \le \lim_{t\to+\infty} J_{\rho,\lambda_1(\Sigma)}(t\xi) = -\lim_{t\to+\infty} \ln \int_{\Sigma} h e^{t\xi} = -\infty,$$

where ξ is a nonzero function which solves

$$-\Delta \xi = \lambda_1(\Sigma) \, \xi.$$

This implies that $\{u_n\}$ must be a blow-up sequence.

According to Theorem 1.1, one can define the Leray-Schauder degree $d_{\rho,\alpha,h}$ for (1.6) as follows (for more details about Leray-Schauder degree and its various properties we refer the reader to Chang [7, Chapter 3]). Let

$$X_{\tau} = \left\{ u \in C^{2,\tau}(\Sigma) : \int_{\Sigma} u = 0 \right\}.$$

Clearly X_{τ} , equipped with the $C^{2,\tau}(\Sigma)$ norm, is a Banach space. We introduce an operator $K_{\rho,\alpha,h}: X_{\tau} \longrightarrow X_{\tau}$ by

$$K_{\rho,\alpha,h}(u) = (-\Delta)^{-1} \left(\rho \left(\frac{he^u}{\int_{\Sigma} he^u} - 1 \right) + \alpha \left(u - \int_{\Sigma} u \right) \right).$$

The standard elliptic theory implies that $K_{\rho,\alpha,h}$ is a well defined compact operator. The generalized mean field equation (1.6) is equivalent to $(1 - K_{\rho,\alpha,h})u = 0$ in X_{τ} . For any bounded open set $B \subset X_{\tau}$, the Leray-Schauder degree $\deg(1 - K_{\rho,\alpha,h}, B, 0)$ is well defined provided $0 \notin (1 - K_{\rho,\alpha,h})(\partial B)$. Let

$$B_R^{X_\tau} = \left\{ u \in X_\tau : ||u||_{X_\tau} < R \right\}$$

be the ball in X_{τ} . Due to the compactness result in Theorem 1.1, we know that for all solutions u to (1.6)

$$\left\| u - \ln \int_{\Sigma} h e^{u} \right\|_{X_{\tau}} \le C,$$

which implies that

$$-C + \ln \int_{\Sigma} h e^{u} \le u \le C + \ln \int_{\Sigma} h e^{u}.$$

If $\oint_{\Sigma} u = 0$, then we obtain

$$\left| \ln \int_{\Sigma} h e^u \right| \leq C.$$

Consequently, the Leray-Schauder degree $\deg\left(1-K_{\rho,\alpha,h},B_R^{X_\tau},0\right)$ is well defined for R large and, in view of the homotopy invariance of the Leray-Schauder degree, is independent of R as R large. Thus

$$d_{\rho,\alpha} := \lim_{R \to \infty} \deg \left(1 - K_{\rho,\alpha,h}, B_R^{X_\tau}, 0 \right)$$

is well defined and is independent of h due to the homotopy invariance. In particular, according to Chen and Lin's result [12, Theorem 2], we have the following

Theorem 1.3. Let (Σ, g) be a closed Riemann surface, h a positive smooth function, $\lambda_1(\Sigma)$ the first positive eigenvalue of the negative Laplacian, ρ and α real numbers. Then we have for $\rho \in (8k\pi, 8(k+1)\pi)$, $k \in \mathbb{N}$ and $\alpha < \lambda_1(\Sigma)$,

$$d_{\rho,\alpha} = d_{\rho,0} = \begin{cases} \frac{\left(k - \chi\left(\Sigma\right)\right) \cdots \left(1 - \chi\left(\Sigma\right)\right)}{k!}, & k \in \mathbb{N}^*, \\ 1, & k = 0. \end{cases}$$

An obvious consequence of Theorem 1.3 yields an existence result. Namely,

Corollary 1.4. If $\chi(\Sigma) \leq 0$ or $\chi(\Sigma) = 2$ but $0 \leq k \leq 1$, then there exists at least $|d_{\rho,0}|$ solutions to (1.6) provided that $\rho \in \mathbb{R} \setminus 8\pi\mathbb{N}^*$ and $\alpha < \lambda_1(\Sigma)$.

We also consider the remaining cases of Corollary 1.4 and give the existence result for (1.6) on arbitrary closed Riemann surface. For this purpose, we employ a minimax scheme. This is a standard method now, which was used by Ding, Jost, Li and Wang [21] to study the mean field equation when $\rho \in (8\pi, 16\pi)$, and by Djadli [22] to solve the mean field equation when $\rho \in (8k\pi, 8(k+1)\pi)$ with $k \in \mathbb{N}^*$. Furthermore, Djadli and Malchiodi [23] employed it to discuss the constant Q-curvature equation; Battaglia, Jevnikar and Malchiodi [3] adopted it to study the Toda system.

Now we state our second main result as follows.

Theorem 1.5. Let (Σ, g) be a closed Riemann surface, h a positive smooth function, ρ and α real numbers. Assume that $\rho \in (8k\pi, 8(k+1)\pi)$ for some non-negative integer number $k \in \mathbb{N}$ and $\lambda_1(\Sigma)$ is the first positive eigenvalue of the negative Laplacian. Then there exists a solution to (1.6) provided $\alpha < \lambda_1(\Sigma)$.

Remark 1.6. For $\rho \in (8k\pi, 8(k+1)\pi)$ and $\alpha < \lambda_1(\Sigma)$, note that $d_{\rho,\alpha} = 0$ when Σ is a 2-sphere and $k \ge 2$. Consequently, there exist at least two solutions in this case.

Additionally, an analog of Theorem 1.5 reads

Remark 1.7. Let $0 = \lambda_0(\Sigma) < \lambda_1(\Sigma) < \cdots$ be all distinct eigenvalues of $-\Delta$, $E_{\lambda_k(\Sigma)}$ the eigenfunction space with respect to $\lambda_k(\Sigma)$, and $E_l = E_{\lambda_0(\Sigma)} \oplus \cdots \oplus E_{\lambda_l(\Sigma)}$. A similar argument above gives a critical point u of $J_{\rho,\alpha}$ in E_l^{\perp} provided $\rho \in \mathbb{R} \setminus 8\pi\mathbb{N}^*$ and $\alpha < \lambda_l(\Sigma)$, i.e.,

$$-\Delta \left[u - \sum_{j=0}^{L} \left\langle u, \phi_j \right\rangle \phi_j \right] = \rho \left[\frac{he^u}{\int_{\Sigma} he^u} - \sum_{j=0}^{L} \left\langle \frac{he^u}{\int_{\Sigma} he^u}, \phi_j \right\rangle \phi_j \right] + \alpha \left[u - \sum_{j=0}^{L} \left\langle u, \phi_j \right\rangle \phi_j \right],$$

where $\{\phi_0, \ldots, \phi_L\}$ is an orthonormal frame of E_l . This complements the results of Yang and Zhu [44].

The remaining part of this paper is organized as follows: firstly we study the compactness of the generalized mean field equation and prove Theorem 1.1 in section 2; secondly we give a new proof of a Trudinger-Moser inequality (cf. Theorem 3.1) in section 3; finally we obtain the existence result for (1.6) by using a minimax scheme and complete the proof of Theorem 1.5 in section 4. Hereafter we do not distinguish sequence and subsequence; moreover, we often denote various constants by the same C.

2. Compactness

In this section, we first review some facts about the blow-up analysis for the mean field equation with positive potential, and then we prove the compactness result for generalized mean field equation.

To study the general existence of the mean field equation (1.1) when $\rho \ge 8\pi$, we consider the blow-up analysis of a sequence u_n which solves

$$-\Delta u_n = \rho_n \left(\frac{h_n e^{u_n}}{\int_{\Sigma} h_n e^{u_n}} - 1 \right), \tag{2.1}$$

where

$$\rho_n \to \rho, \quad h_n \stackrel{C^1(\Sigma)}{\to} h.$$

Up to adding a constant, we may assume

$$\int_{\Sigma} h_n e^{u_n} = 1.$$

Since $\rho > 0$ and h is a positive smooth function, it yields that

$$\int_{\Sigma} e^{u_n} \le C. \tag{2.2}$$

Applying the Green representation formula (cf. [2, Theorem 4.13]) and the potential estimate (cf. [25, Lemma 7.12]), we obtain

$$\left\| u_n - \int_{\Sigma} u_n \right\|_{W^{1,p}(\Sigma)} \le C_p \left\| \Delta u_n \right\|_{L^1(\Sigma)} \le C_p, \quad \forall p \in (1,2).$$

If $\{u_n^+\}$ is bounded in $L^{\infty}(\Sigma)$, the standard elliptic estimate gives a uniform bound for $\{u_n\}$ in $L^{\infty}(\Sigma)$. In this case, $\{u_n\}$ is compact in $C^2(\Sigma)$. If $\{u_n\}$ is a blow-up sequence, i.e.,

$$\limsup_{n\to\infty} \max_{\Sigma} u_n = +\infty,$$

we may assume $\rho_n h_n e^{u_n} d\mu_{\Sigma}$ converges to a nonzero Radon measure μ . The singular set S of the blow-up sequence $\{u_n\}$ is defined by

$$S = \{x \in \Sigma : \mu(\{x\}) \ge 4\pi\}.$$

It is clear that S is a finite subset. According to Brezis-Merle's estimate ([5, Theorem 1]), for each smooth domain $\Omega \subset \Sigma$ and each solution u to

$$\begin{cases} -\Delta u = f, & \text{in } \Omega, \\ u = 0, & \text{in } \partial \Omega, \end{cases}$$

where $f \in L^1(\Sigma)$, we have for every $\delta \in (0, 4\pi)$

$$\int_{\Omega} \exp\left(\frac{(4\pi - \delta)|u|}{\|f\|_{L^{1}(\Sigma)}}\right) \le C_{\delta,\Omega}.$$

Consequently, for every compact subset $K \subset \Sigma \setminus S$, there is a constant $C = C_K$ such that (cf. [20, Lemma 2.8])

$$\left\|u_n - \int_{\Sigma} u_n\right\|_{L^{\infty}(K)} \le C_K,$$

which implies that S is nonempty. One can check (cf. [5, Page 1242-1243]) that $\lim_{n\to\infty} \oint_{\Sigma} u_n = -\infty$. Thus $\mu = \sum_{x\in\Sigma} \mu(\{x\}) \delta_x$. Moreover, the singular set can be characteristic of the blow-up set ([5, Page 1240-1241]), i.e.,

$$S = \left\{ x \in \Sigma : \exists \left\{ x_n \right\} \subset \Sigma, \ \lim_{n \to \infty} x_n = x, \ \lim_{n \to \infty} u_n(x_n) = +\infty \right\}.$$

For $x_0 \in S$, assume $S \cap B^{\Sigma}_{\delta}(x_0) = \{x_0\}$. Choose $B^{\Sigma}_{\delta}(x_0) \ni x_n \to x_0$ such that

$$\lambda_n := u_n(x_n) = \max_{B^{\Sigma}_{\delta}(x_0)} u_n \to +\infty.$$

Without loss of generality, assume $B_{\delta}^{\Sigma}(x_0)$ is a Euclidean ball. Consider a rescaling

$$\tilde{u}_n(x) = u_n \left(x_n + e^{-\lambda_n/2} x \right) - \lambda_n, \quad |x| \le e^{\lambda_n/2} \left(\delta - |x_n| \right).$$

From (2.1) and (2.2), it follows that

$$-\Delta_{\mathbb{R}^2}\tilde{u}_n(x) = \rho_n h_n \left(x_n + e^{-\lambda_n/2} x \right) e^{\tilde{u}_n(x)} - \rho_n e^{-\lambda_n},$$

and that

$$\int_{\mathbb{B}_R} e^{\tilde{u}_n} \le C_R.$$

The above argument implies that \tilde{u}_n converges strongly to u_∞ in $H^2_{loc}(\mathbb{R}^2)$ as $n \to +\infty$, and then Chen-Li's classification result ([15, Theorem 1]) leads to

$$u_{\infty}(x) = -2 \ln \left(1 + \frac{\rho h(x_0)}{8} |x|^2 \right).$$

By Fatou's Lemma, we conclude that

$$\mu(\{x_0\}) \ge 8\pi.$$

Recall the following Pohozaev identity (cf. [31, formula (8.1)]):

Assume u is a solution to

$$-\Delta u = fe^u - P$$
, in \mathbb{B} ,

where $f \in C^1(\mathbb{B})$ and $P \in L^{\infty}(\mathbb{B})$. Then for each $r \in (0,1)$ and every smooth function F,

$$\int_{\mathbf{B}_{r}} \left\langle \nabla^{2} F - \frac{1}{2} \Delta F g, \nabla u \otimes \nabla u \right\rangle + \frac{1}{2} \int_{\partial \mathbf{B}_{r}} |\nabla u|^{2} \left\langle \nabla F, \nu \right\rangle - \int_{\partial \mathbf{B}_{r}} \left\langle \nabla F, \nabla u \right\rangle \left\langle \nu, \nabla u \right\rangle \\
= \int_{\partial \mathbf{B}_{r}} f e^{u} \left\langle \nabla F, \nu \right\rangle - \int_{\mathbf{B}_{r}} e^{u} \left\langle \nabla f, \nabla F \right\rangle - \int_{\mathbf{B}_{r}} f e^{u} \Delta F - \int_{\mathbf{B}_{r}} P \left\langle \nabla F, \nabla u \right\rangle.$$
(2.3)

Take $F = \frac{1}{2} |x|^2$ in (2.3) to obtain

$$\frac{r}{2} \int_{\partial \mathbf{B}_{-}} |\nabla u|^{2} - r \int_{\partial \mathbf{B}_{-}} \langle v, \nabla u \rangle^{2} = r \int_{\partial \mathbf{B}_{-}} f e^{u} - \int_{\mathbf{B}_{-}} e^{u} \langle \nabla f, x \rangle - 2 \int_{\mathbf{B}_{-}} f e^{u} - \int_{\mathbf{B}_{-}} P \langle x, \nabla u \rangle. \tag{2.4}$$

Applying the Pohozaev identity (2.4) to the blow-up sequence u_n , we conclude that

$$\mu(\{x_0\}) = \lim_{r \to 0} \left(\frac{r}{2} \int_{\partial \mathbf{B}_r} \langle \nabla G, \nu \rangle^2 - \frac{r}{4} \int_{\mathbf{B}_r} |\nabla G|^2 \right) = \frac{\mu(\{x_0\})^2}{8\pi},$$

where G is the Green function satisfying

$$-\Delta G(\cdot, y) = \sum_{x \in S} \mu(\{x\}) \, \delta_x - \rho, \quad \int_{\Sigma} G(\cdot, y) = 0.$$

Therefore, if S has k points then $\rho = 8k\pi$. In other words, if $\rho \in \mathbb{R} \setminus 8\pi\mathbb{N}^*$, then there is a uniform constant C such that

$$||u_n||_{L^{\infty}(\Sigma)} \leq C.$$

We are in position to give a compactness result for the generalized mean field equation as below.

Proof of Theorem 1.1. We consider a sequence $u_n \in C^2(\Sigma)$ satisfying

$$-\Delta u_n = \rho_n \left(h_n e^{u_n} - 1 \right) + \alpha_n \left(u_n - \int_{\Sigma} u_n \right),\,$$

where

$$\rho_n \to \rho$$
, $\alpha_n \to \alpha$, $h_n \stackrel{C^{1,\tau}(\Sigma)}{\to} h$, as $n \to \infty$,

and $0 < \tau < 1$. It suffices to prove that

$$||u_n||_{L^{\infty}(\Sigma)} \leq C.$$

Arguing as in (2.2), one can check that

$$\int_{\Sigma} e^{u_n} \leq C.$$

We claim the following potential estimate:

$$\{u_n - \oint_{\Sigma} u_n\}$$
 is bounded in $W^{1,p}(\Sigma)$ for every $p \in (1,2)$.

In fact, for any fixed $p \in (0, 1)$, applying the classical potential estimate for Laplacian, one has

$$\left\|u_n - \int_{\Sigma} u_n\right\|_{W^{1,p}(\Sigma)} \le C \left\|\Delta u_n\right\|_{L^1(\Sigma)} \le C \left[\left\|\Delta u_n + \alpha_n \left(u_n - \int_{\Sigma} u_n\right)\right\|_{L^1(\Sigma)} + \left\|u_n - \int_{\Sigma} u_n\right\|_{L^1(\Sigma)}\right].$$

Assume for some $C_n \to +\infty$,

$$\left\| u_n - \int_{\Sigma} u_n \right\|_{W^{1,p}(\Sigma)} \ge C_n \left\| \Delta u_n + \alpha_n \left(u_n - \int_{\Sigma} u_n \right) \right\|_{L^1(\Sigma)}.$$

Set $w_n = \frac{u_n - \int_{\Sigma} u_n}{\|u_n - \int_{\Sigma} u_n\|_{W^{1,p}(\Sigma)}}$. Thus

$$1 = \|w_n\|_{W^{1,p}(\Sigma)} \ge C_n \|\Delta w_n + \alpha_n w_n\|_{L^1(\Sigma)}, \quad \int_{\Sigma} w_n = 0.$$

We may assume w_n converges to w weakly in $W^{1,p}(\Sigma)$ and strongly in $L^p(\Sigma)$. Then

$$\Delta w + \alpha w = 0, \quad \int_{\Sigma} w = 0.$$

The classical potential estimate yields that

$$1 = \|w_n\|_{W^{1,p}(\Sigma)} \leq C \left(\|\Delta w_n + \alpha_n w_n\|_{L^1(\Sigma)} + \|w_n\|_{L^1(\Sigma)} \right) \leq C \left(\|\Delta w_n + \alpha_n w_n\|_{L^1(\Sigma)} + \|w_n\|_{L^p(\Sigma)} \right).$$

Letting $n \to \infty$, one finds

$$1 \leq C ||w||_{L^p(\Sigma)}$$
.

In particular, α is a nonzero eigenvalue of Laplacian which is a contradiction. Consequently,

$$\left\| u_n - \int_{\Sigma} u_n \right\|_{W^{1,p}(\Sigma)} \le C \left\| \Delta u_n + \alpha_n \left(u_n - \int_{\Sigma} u_n \right) \right\|_{L^1(\Sigma)} \le C. \tag{2.5}$$

Let f_n be the solution of

$$-\Delta f_n = \alpha_n \left(u_n - \int_{\Sigma} u_n \right), \quad \int_{\Sigma} f_n = 0.$$
 (2.6)

Thanks to (2.5), a standard elliptic estimate for (2.6) implies that

$$||f_n||_{W^{3,p}(\Sigma)} \leq C, \quad \forall p \in (1,2).$$

Without loss of generality, we may assume f_n converges to f in $C^{1,\tau}(\Sigma)$ (0 < τ < 1). Define $\tilde{u}_n = u_n - f_n$ and $\tilde{h}_n = h_n e^{f_n}$. Then there holds

$$-\Delta \tilde{u}_n = \rho_n \left(\tilde{h}_n e^{\tilde{u}_n} - 1 \right),\,$$

and

$$\tilde{h}_n \stackrel{C^{1,\tau}(\Sigma)}{\to} \tilde{h} := he^f.$$

Therefore, we may assume $\alpha_n = 0$. The blow-up analysis for the mean field equation with positive potential function is applicable. In particular, we complete the proof.

3. An improved Trudinger-Moser inequality

From now on, we assume $\alpha < \alpha_1$. For $\rho \in (8k\pi, 8(k+1)\pi)(k \in \mathbb{N}^*)$, we shall adopt minimax arguments to derive the existence of solutions to (1.6) on arbitrary closed Riemann surface.

As an application of Theorem 1.1, we first prove the following Trudinger-Moser inequality, which was obtained by Yang [43].

Theorem 3.1. For $\alpha < \lambda_1$, there holds

$$I_{8\pi,\alpha}(u) := \frac{1}{16\pi} \int_{\Sigma} \left[|\nabla u|^2 - \alpha \left(u - \int_{\Sigma} u \right)^2 \right] + \int_{\Sigma} u - \ln \int_{\Sigma} e^u \ge -C, \quad \forall u \in H^1(\Sigma).$$

Proof. It is sufficient to consider the case $\alpha \in (0, \lambda_1)$. We claim that for every $0 < \varepsilon < 16\pi$, there is a positive constant C_{ε} such that

$$\frac{1}{16\pi - \varepsilon} \int_{\Sigma} \left[|\nabla u|^2 - \alpha \left(u - \int_{\Sigma} u \right)^2 \right] + \int_{\Sigma} u - \ln \int_{\Sigma} e^u \ge -C_{\varepsilon}, \quad \forall u \in H^1(\Sigma).$$
 (3.1)

In fact, according to the classical Truding-Moser inequality (1.3), we see that

$$\ln \int_{\Sigma} e^{u} \leq \int_{\Sigma} e^{u^{+}} \\
\leq \frac{1}{16\pi} \int_{\Sigma} \left| \nabla u^{+} \right|^{2} + \int_{\Sigma} u^{+} + C \\
\leq \frac{1}{16\pi} \int_{\Sigma} \left(\left| \nabla u^{+} \right|^{2} - \alpha \left(u^{+} - \int_{\Sigma} u^{+} \right)^{2} \right) + C \int_{\Sigma} \left| u^{+} \right|^{2} + C. \tag{3.2}$$

Without loss of generality, we may assume $\oint_{\Sigma} u = 0$. For each $\eta \in (0, 1)$, choose a > 0 such that

$$|\{x \in \Sigma : u(x) \ge a\}| = \eta.$$

Then it follows from (3.2) that

$$\ln \int_{\Sigma} e^{u} = a + \ln \int_{\Sigma} e^{u-a}
\leq \frac{1}{16\pi} \int_{\Sigma} \left(\left| \nabla (u-a)^{+} \right|^{2} - \alpha \left((u-a)^{+} - \int_{\Sigma} (u-a)^{+} \right)^{2} \right) + C \int_{\Sigma} \left| (u-a)^{+} \right|^{2} + C + a
= \frac{1}{16\pi} \int_{\Sigma} \left(\left| \nabla u \right|^{2} - \alpha u^{2} \right) - \frac{1}{16\pi} \int_{\Sigma} \left(\left| \nabla (u-a)^{-} \right|^{2} - \alpha \left((u-a)^{-} - \int_{\Sigma} (u-a)^{-} \right)^{2} \right)
+ \frac{\alpha}{8\pi} \int_{\Sigma} (u-a)^{+} \int_{\Sigma} (u-a)^{-} + C \int_{\Sigma} \left| (u-a)^{+} \right|^{2} + C + a
\leq \frac{1}{16\pi} \int_{\Sigma} \left(\left| \nabla u \right|^{2} - \alpha u^{2} \right) + \varepsilon \int_{\Sigma} \left| (u-a)^{-} \right|^{2} + C_{\varepsilon} \int_{\Sigma} \left| (u-a)^{+} \right|^{2} + C + a$$
(3.3)

On the other hand, Poincaré's inequality and Kato's inequality yield that

$$\left\| (u - a)^{+} \right\|_{L^{2}(\Sigma)} \le C \eta \left\| \nabla u \right\|_{L^{2}(\Sigma)} \le C \eta^{1/2} \left(\int_{\Sigma} \left(|\nabla u|^{2} - \alpha u^{2} \right) \right)^{1/2}, \tag{3.4}$$

and that

$$\left\| (u - a)^{-} \right\|_{L^{2}(\Sigma)} \le C \left(\int_{\Sigma} \left(|\nabla u|^{2} - \alpha u^{2} \right) \right)^{1/2}. \tag{3.5}$$

Insert (3.4) and (3.5) into (3.3), therefore

$$\ln \int_{\Sigma} e^{u} \leq \left(\frac{1}{16\pi} + \varepsilon + C_{\varepsilon}\eta\right) \int_{\Sigma} \left(|\nabla u|^{2} - \alpha u^{2}\right) + C + a.$$

Notice that

$$a\eta = a \int_{\{u \geq a\}} \leq \int_{\{u \geq a\}} u \leq \eta^{1/2} \|u\|_{L^2(\Sigma)} \leq C \eta^{1/2} \left(\int_{\Sigma} \left(|\nabla u|^2 - \alpha u^2 \right) \right)^{1/2}.$$

As a consequence,

$$\ln \int_{\Sigma} e^{u} \le \left(\frac{1}{16\pi} + \varepsilon + C_{\varepsilon} \eta \right) \int_{\Sigma} \left(|\nabla u|^{2} - \alpha u^{2} \right) + C + \frac{C_{\varepsilon}}{\eta}. \tag{3.6}$$

Let η small enough. Finally (3.1) follows from (3.6) immediately.

In view of (3.1), for each $\rho_n = 8\pi - 1/n$, there exits a minimizer $u_n \in H^1(\Sigma)$ to the functional

$$I_{\rho_{n},\alpha}(u) := \frac{1}{2\rho_{n}} \left(\int_{\Sigma} |\nabla u|^{2} - \alpha \left(u - \int_{\Sigma} u \right)^{2} \right) + \int_{\Sigma} u - \ln \int_{\Sigma} e^{u}, \quad u \in H^{1}(\Sigma).$$

$$(3.7)$$

It is obvious to see

$$\lim_{n\to\infty}I_{\rho_n,\alpha}(u_n)=\inf_{u\in H^1(\Sigma)}I_{8\pi,\alpha}(u).$$

One can check that if u_n blows up, then ([44, Theorem 1.1])

$$\inf_{u \in H^1(\Sigma)} I_{8\pi,\alpha}(u) = -1 - \ln \pi - \frac{1}{2} \max_{\Sigma} A,$$
(3.8)

where A is the regular part of the Green function G, i.e.,

$$-\Delta G(\cdot, y) = 8\pi \left(\delta_y - 1\right) + \alpha G(\cdot, y), \quad \int_{\Sigma} G(\cdot, y) = 0, \tag{3.9}$$

Here G takes the form

$$G(x, x_0) = -4 \ln r(x) + A(x_0) + \psi(x), \tag{3.10}$$

where r denotes the geodesic distance between x and x_0 , $\psi(x) \in C^1(\Sigma)$ and $\psi(x_0) = 0$. We next give a new proof of (3.8). Without loss of generality, we may assume u_n satisfies

$$-\Delta u_n = \rho_n \left(e^{u_n} - 1 \right) + \alpha \left(u_n - \int_{\Sigma} u_n \right)$$

and $u_n - \oint_{\Sigma} u_n$ converges to the Green function G weakly in $W^{1,p}(\Sigma)$ and strongly in $L^p(\Sigma)$. Let $\{x_0\}$ be the blow-up point. Let f_n be given by

$$-\Delta f_n = \alpha \left(u_n - \int_{\Sigma} u_n \right), \quad \int_{\Sigma} f_n = 0.$$

Due to the potential estimate (2.5), we obtain that f_n is compact in $C^1(\Sigma)$. Set $w_n = u_n - f_n$ and $k_n = e^{f_n}$. We may assume $k_n = e^{f_n}$ converges to $k := e^f$ in $C^1(\Sigma)$ as $n \to \infty$. And then, there holds

$$-\Delta \left(G(\cdot, x_0) - f\right) = 8\pi \left(\delta_{x_0} - 1\right).$$

Observe that

$$-\Delta w_n = \rho_n \left(k_n e^{w_n} - 1 \right).$$

It is easy to check that $\{w_n\}$ is a blow-up sequence. Since

$$I_{\rho_n,\alpha}(u_n) = \left[\frac{1}{2\rho_n} \int_{\Sigma} |\nabla w_n|^2 + \int_{\Sigma} w_n - \ln \int_{\Sigma} k_n e^{w_n}\right] + \left[\frac{\alpha}{2\rho_n} \int_{\Sigma} \left(u_n - \int_{\Sigma} u_n\right)^2 - \frac{1}{2\rho_n} \int_{\Sigma} |\nabla f_n|^2\right],$$

we conclude from the proof of [20, formula (1.4)] that

$$\lim_{n \to \infty} I_{\rho_n,\alpha}(u_n) = \lim_{n \to \infty} \left[\frac{1}{2\rho_n} \int_{\Sigma} |\nabla w_n|^2 + \int_{\Sigma} w_n - \ln \int_{\Sigma} k_n e^{w_n} \right] + \lim_{n \to \infty} \left[\frac{\alpha}{2\rho_n} \int_{\Sigma} \left(u_n - \int_{\Sigma} u_n \right)^2 - \frac{1}{2\rho_n} \int_{\Sigma} |\nabla f_n|^2 \right]$$

$$\geq -1 - \ln \pi - \left(\ln k(x_0) + \frac{1}{2} \tilde{A}(x_0) \right) + \frac{\alpha}{16\pi} \int_{\Sigma} G(\cdot, x_0)^2 - \frac{1}{16\pi} \int_{\Sigma} |\nabla f|^2 .$$
(3.11)

Here \tilde{A} is the regular part of the Green function \tilde{G} , i.e.,

$$-\Delta \tilde{G}(\cdot, y) = 8\pi \left(\delta_y - 1\right), \quad \int_{\Sigma} \tilde{G}(\cdot, y) = 0.$$

and in a local normal coordinates x centering at x_0 ,

$$\tilde{G}(x, x_0) = -4 \ln|x| + \tilde{A}(x_0) + \tilde{\psi}(x), \tag{3.12}$$

where $\tilde{\psi}$ is a smooth function with $\tilde{\psi}(x_0) = 0$. Note that $G(\cdot, x_0) = \tilde{G}(\cdot, x_0) + f$. Together with (3.10) and (3.12), it leads to $\tilde{A}(x_0) = A(x_0) - f(x_0)$ and

$$\frac{1}{16\pi} \int_{\Sigma} |\nabla f|^2 = -\frac{1}{16\pi} \int_{\Sigma} f \Delta f = \frac{\alpha}{16\pi} \int_{\Sigma} f G = \frac{1}{16\pi} \int_{\Sigma} f \left(-\Delta G + 8\pi \left(1 - \delta_{x_0}\right)\right) = -\frac{1}{16\pi} \int_{\Sigma} \Delta f G - \frac{f(x_0)}{2}. \tag{3.13}$$

Combing (3.11) and (3.13), one has

$$\lim_{n \to \infty} I_{\rho_n,\alpha}(u_n) \ge -1 - \ln \pi - \left(f(x_0) + \frac{1}{2} \tilde{A}(x_0) \right) + \frac{f(x_0)}{2} = -1 - \ln \pi - \frac{1}{2} A(x_0) \ge -1 - \ln \pi - \frac{1}{2} \max_{\Sigma} A.$$

On the other hand, we construct a sequence of functions $\{\phi_n\}$ (which is in [44]) satisfying

$$\lim_{n\to\infty} I_{8\pi,\alpha}\left(\phi_n - \oint_{\Sigma} \phi_n\right) = -1 - \ln \pi - \frac{1}{2} \max_{\Sigma} A.$$

Precisely, suppose that $A(x_0) = \max_{x \in \Sigma} A_x$. Let r = r(x) be the geodesic distance between x and x_0 . Set

$$\phi_n(x) = \begin{cases} c - 2\log\left(1 + \frac{r^2}{8r_n^2}\right), & x \in B_{Rr_n}^{\Sigma}(x_0), \\ G(x, x_0) - \eta(x)\psi(x), & x \in B_{2Rr_n}^{\Sigma}(x_0) \setminus B_{Rr_n}^{\Sigma}(x_0), \\ G(x, x_0), & x \in \Sigma \setminus B_{2Rr_n}^{\Sigma}(x_0), \end{cases}$$

where $r_n = e^{-\lambda_n/2}$, $\lambda_n = \max_{\Sigma} u_n$, $\eta \in C_0^{\infty}\left(B_{2Rr_n}^{\Sigma}(x_0)\right)$ is a cut-off function satisfying $\eta \equiv 1$ in $B_{Rr_n}^{\Sigma}(x_0)$ and $|\nabla \eta(x)| \leq \frac{4}{Rr_n}$ for $x \in B_{2Rr_n}^{\Sigma}(x_0)$, $G(\cdot, x_0)$ and ψ are defined in (3.9), (3.10) respectively. Moreover

$$c = 2\ln(1 + R^2/8) - 4\ln R - 4\ln r_n + A(x_0).$$

Then we estimate the three terms of (3.7) respectively. By a straightforward calculation as in the proof of ([44], Theorem 1.1.), we can get the desired result. Thus, we end the proof of (3.8) and complete the proof the theorem.

We next prove an improved Trudinger-Moser inequality which will be used in the next section. Namely,

Lemma 3.2. Assume $\alpha < \lambda_1(\Sigma)$. For positive integer number k, positive numbers δ_0, γ_0 , consider

$$X_{k,\delta_0,\gamma_0} := \left\{ u \in H^1(\Sigma) : \exists subdomains \ \Omega_i \subset \Sigma \ s.t. \ \min_{1 \le i < j \le k} \operatorname{dist} \left(\Omega_i, \Omega_j\right) \ge \delta_0 \ and \ \min_{1 \le i \le k} \int_{\Omega_i} e^u \ge \gamma_0 \int_{\Sigma} e^u \right\}.$$

For every $\varepsilon \in (0, 16k\pi)$, there exists a positive constant $C = C_{k,\delta_0,\gamma_0,\varepsilon}$ such that

$$\ln \int_{\Sigma} e^{u} \le \frac{1}{16k\pi - \varepsilon} \int_{\Sigma} \left(|\nabla u|^{2} - \alpha \left(u - \int_{\Sigma} u \right)^{2} \right) + \int_{\Sigma} u + C, \quad \forall u \in X_{k, \delta_{0}, \gamma_{0}}. \tag{3.14}$$

Proof. Given $u \in X_{k,\delta_0,\gamma_0}$ with $\oint_{\Sigma} u = 0$, applying an improved Trudinger-Moser inequality for $\alpha = 0$ (cf. [16, Theorem 2.1]), we obtain

$$\ln \int_{\Sigma} e^{u} \leq \frac{1}{16k\pi - \varepsilon} \int_{\Sigma} |\nabla u|^{2} + \int_{\Sigma} u + C, \quad \forall u \in X_{k,\delta_{0},\gamma_{0}}.$$

Using a similar method as in the proof of (3.1), we complete the proof.

4. Existence

The remaining part of this section is devoted to the proof of Theorem 1.5. We begin with the topological structure for the formal set of barycenters, which is to be used in the minimax argument. Denote $\mathcal{D}(\Sigma)$ by the distributions on Σ . We will use on $\mathcal{D}(\Sigma)$ the metric given by $C^1(\Sigma)^*$ and which will be denoted by $\mathbf{d}(\cdot, \cdot)$. In other words, for $\sigma, \zeta \in \mathcal{D}(\Sigma)$,

$$\mathbf{d}(\sigma,\zeta) = \sup_{\psi \in C^1(\Sigma): \|\psi\|_{C^1(\Sigma)} \le 1} \langle \sigma - \zeta, \psi \rangle.$$

We consider

$$\Sigma_{k} = \left\{ \sum_{i=1}^{k} t_{i} \delta_{x_{i}} : t_{i} \geq 0, x_{i} \in \Sigma, \sum_{i=1}^{k} t_{i} = 1 \right\} \subset \mathcal{D}(\Sigma)$$

which is known as the formal set of barycenters of Σ of order k. It is the fact that Σ_k is non-contractible for every $k \ge 1$ (cf. [22, Lemma 4.7]).

The first step is to construct a continuous projection from low sublevels of the functional to the k-th barycenters, precisely

Lemma 4.1. Let $\rho \in (8k\pi, 8(k+1)\pi)$, $k \in \mathbb{N}$ and $\alpha < \lambda_1(\Sigma)$. Then for L sufficiently large there exists a continuous projection

$$\Psi: J_{\rho,\alpha}^{-L} := \left\{ u \in H^1(\Sigma): J_{\rho,\alpha}(u) \le -L \right\} \longrightarrow \Sigma_k.$$

Moreover, if $\frac{e^{u_n}}{\int_{\Sigma} e^{u_n}} d\mu_{\Sigma}$ converges to $\sigma \in \Sigma_k$, then $\Psi(u_n) \to \sigma$.

Proof. The proof is similar to [19, Proposition 4.1]. There exists a large L such that for every $u_n \in J_{\rho,\alpha}^{-L}$, the following holds

If $J_{o,\alpha}(u_n) \to -\infty$, then up to a subsequence,

$$\sigma_n := \frac{e^{u_n}}{\int_{\Sigma} e^{u_n}} d\mu_{\Sigma} \to \sigma \in \Sigma_k.$$

One can prove this fact by contradiction that there exists k+1 points $\{x_1,\ldots,x_{k+1}\}\subset \operatorname{supp}(\sigma)$. Take r>0 such that $B_{2r}^{\Sigma}(x_i)\cap B_{2r}^{\Sigma}(x_i)=\emptyset$ for all $i\neq j$. There exists $\varepsilon>0$ such that $\sigma\left(B_r^{\Sigma}(x_i)\right)>2\varepsilon$ for all i. Thus we may assume

$$\frac{\int_{B_r^{\Sigma}(x_i)} e^{u_n}}{\int_{\Sigma} e^{u_n}} \ge \varepsilon, \quad i = 1, \dots, k+1.$$

The improved Trudinger-Moser inequality (3.14) implies that for any $\tilde{\varepsilon} \in (0, 16(k+1)\pi)$, there exists a constant $C = C(\tilde{\varepsilon}, \varepsilon, r)$ such that

$$\ln \int_{\Sigma} h e^{u_n} \le \ln \int_{\Sigma} e^{u_n} + C \le \frac{1}{16(k+1)\pi - \tilde{\varepsilon}} \int_{\Sigma} \left(|\nabla u_n|^2 - \alpha \left(u_n - \int_{\Sigma} u_n \right)^2 \right) + \int_{\Sigma} u_n + C. \tag{4.1}$$

Taking $\tilde{\varepsilon}$ small, we conclude from (4.1) that

$$J_{\rho,\alpha}(u_n) \geq -C$$
,

which yields a contradiction.

Therefore, for any $\varepsilon_0 > 0$, there exists L_0 large enough such that if $L > L_0$, then

$$\frac{e^{u}}{\int_{\Sigma} e^{u}} d\mu_{\Sigma} \in \{ \sigma \in \mathcal{D}(\Sigma) : \mathbf{d} (\sigma, \Sigma_{k}) < \varepsilon_{0} \}, \quad \forall u \in J_{\rho, \alpha}^{-L}.$$

Notice that for small ε_0 , there is a continuous retraction (cf. [3, Proposition 2.2])

$$\psi_k : \{ \sigma \in \mathcal{D}(\Sigma) : \mathbf{d}(\sigma, \Sigma_k) < \varepsilon_0 \} \longrightarrow \Sigma_k.$$

Now define Ψ by

$$\Psi: u \mapsto \psi_k \left(\frac{e^u}{\int_{\Sigma} e^u} d\mu_{\Sigma} \right)$$

to finish the proof.

Next, we aim to construct a continuous map $\Phi_{\lambda}: \Sigma_{k} \longrightarrow J_{\rho,\alpha}^{-L}$ such that $\Psi \circ \Phi_{\lambda}$ is homotopic to the identity map for λ large. For this purpose, we choose a smooth non-decreasing function $\eta: \mathbb{R} \longrightarrow \mathbb{R}$ such that

$$\eta(t) = \begin{cases} t, & t \le 1, \\ 2, & t \ge 2, \end{cases}$$

and denote $\eta_{\delta}(t) = \delta \eta (t/\delta)$ for small $\delta > 0$. Given $\sigma = \sum_{i=1}^{k} t_i \delta_{x_i} \in \Sigma_k$ and $\lambda > 0$, we define

$$\phi_{\lambda,\sigma}(x) = \ln\left(\sum_{i=1}^{k} t_i \frac{8\lambda^2}{\left(1 + \lambda^2 \eta_{\delta}^2 \left(\operatorname{dist}^{\Sigma}(x, x_i)\right)\right)^2}\right).$$

Then the following lemma holds.

Lemma 4.2. Let $\rho \in (8k\pi, 8(k+1)\pi)$, $k \in \mathbb{N}$ and $\alpha < \lambda_1(\Sigma)$. We can choose a small δ and large λ such that for any $\sigma \in \Sigma_k$,

$$J_{\rho,\alpha}\left(\phi_{\lambda,\sigma}\right) \leq -\left(1 - \frac{8k\pi}{\rho}\right)\ln\lambda,$$

and

$$\frac{e^{\phi_{\lambda,\sigma}}}{\int_{\Sigma} e^{\phi_{\lambda,\sigma}}} d\mu_{\Sigma} \to \sigma, \quad as \ \lambda \to +\infty.$$

Proof. Assume $\delta \ll 1$ and $\lambda \gg 1$. Set $\sigma = \sum_{i=1}^k t_i \delta_{x_i}, t_i \ge 0, \sum_{i=1}^k t_i = 1$. By definition

$$\phi_{\lambda,\sigma}(x) = \begin{cases} \ln\left(\frac{8\lambda^2}{(1+4\delta^2\lambda^2)^2}\right), & x \in \Sigma \setminus \left(\bigcup_{i=1}^k B_{2\delta}^{\Sigma}(x_i)\right), \\ \ln\left(\frac{8\lambda^2 t_i}{(1+\lambda^2\eta_{\delta}^2(\operatorname{dist}^{\Sigma}(x,x_i)))^2} + \frac{8\lambda^2(1-t_i)}{(1+4\delta^2\lambda^2)^2}\right), & x \in B_{2\delta}^{\Sigma}(x_i). \end{cases}$$

Choose local normal coordinates x around x_i . It is well known that

$$\det\left(g_{ij}(x)\right) = 1 + O\left(|x - x_i|^2\right).$$

For any fixed $\delta > 0$, and every smooth function ψ on Σ , it holds

$$\int_{\Sigma} e^{\phi_{\lambda,\sigma}} \psi = \sum_{i=1}^{k} t_{i} \int_{B_{r}^{\Sigma}(x_{i})} \frac{8\lambda^{2}}{\left(1 + \lambda^{2} \left(\operatorname{dist}^{\Sigma}(x, x_{i})\right)^{2}\right)^{2}} + O\left(r^{-4}\lambda^{-2}\right)$$

$$= \left(1 + O\left(r^{2}\right)\right) \sum_{i=1}^{k} t_{i} \int_{B_{r}(x_{i})} \frac{8\lambda^{2}}{\left(1 + \lambda^{2} |x - x_{i}|^{2}\right)^{2}} \psi(x) + O\left(r^{-4}\lambda^{-2}\right)$$

$$= \left(1 + O\left(r^{2}\right)\right) \sum_{i=1}^{k} t_{i} \int_{B_{r}(x_{i})} \frac{8\lambda^{2}}{\left(1 + \lambda^{2} |x - x_{i}|^{2}\right)^{2}} \left(\psi(x) - \psi(x_{i})\right) + 8\pi \sum_{i=1}^{k} t_{i} \psi(x_{i}) + O\left(r^{2}\right) + O\left(r^{-4}\lambda^{-2}\right)$$

$$= 8\pi \sum_{i=1}^{k} t_{i} \psi(x_{i}) + O\left(r\right) + O\left(r^{-4}\lambda^{-2}\right).$$
(4.2)

Passing to the limit $\lambda \to +\infty$ first, and then $r \to 0$, we have by (4.2) that

$$\lim_{\lambda \to +\infty} \int_{\Sigma} e^{\phi_{\lambda,\sigma}} = 8\pi$$

and that

$$\lim_{\lambda \to +\infty} \int_{\Sigma} \frac{e^{\phi_{\lambda,\sigma}}}{\int_{\Sigma} e^{\phi_{\lambda,\sigma}}} \psi = \sum_{i=1}^{k} t_i \psi(x_i) = \langle \sigma, \psi \rangle.$$

Thus we confirm the second claim.

For the first assertion, we compute

$$\int_{\Sigma} \left| \nabla \phi_{\lambda,\sigma} \right|^{2} = \sum_{i=1}^{k} \int_{B_{2\delta}^{\Sigma}(x_{i})} \left(\frac{\frac{8\lambda^{2}t_{i}}{(1+\lambda^{2}\eta_{\delta}^{2})^{2}}}{\frac{8\lambda^{2}t_{i}}{(1+\lambda^{2}\eta_{\delta}^{2})^{2}}} + \frac{8\lambda^{2}(1-t_{i})}{(1+4\delta^{2}\lambda^{2})^{2}} \right)^{2} \cdot \left(\frac{4\lambda^{2}\eta_{\delta}\eta_{\delta}'}{1+\lambda^{2}\eta_{\delta}^{2}} \right)^{2} \\
\leq \sum_{i=1}^{k} \int_{B_{\delta}^{\Sigma}(x_{i})} \left(\frac{4\lambda^{2}\operatorname{dist}^{\Sigma}(x, x_{i})}{1+\lambda^{2}\left(\operatorname{dist}^{\Sigma}(x, x_{i})\right)^{2}} \right)^{2} + C \\
= \sum_{i=1}^{k} \left(1 + O\left(\delta^{2}\right) \right) \int_{\mathbf{B}_{\delta}} \left(\frac{4\lambda^{2}|x|}{1+\lambda^{2}|x|^{2}} \right)^{2} + C \\
\leq 32k\pi \left(1 + O\left(\delta^{2}\right) \right) \ln \lambda + C. \tag{4.3}$$

This together with Poincaré's inequality leads to

$$\int_{\Sigma} \left(\phi_{\lambda,\sigma} - \int_{\Sigma} \phi_{\lambda,\sigma} \right)^{2} \leq \int_{\Sigma} \left(\phi_{\lambda,\sigma} - \ln \left(\frac{8\lambda^{2}}{(1 + 4\delta^{2}\lambda^{2})^{2}} \right) \right)^{2}$$

$$= \sum_{i=1}^{k} \int_{B_{2\delta}^{\Sigma}(x_{i})} \left(\phi_{\lambda,\sigma} - \ln \left(\frac{8\lambda^{2}}{(1 + 4\delta^{2}\lambda^{2})^{2}} \right) \right)^{2}$$

$$\leq C\delta^{2} \int_{B_{2\delta}^{\Sigma}(x_{i})} \left| \nabla \phi_{\lambda,\sigma} \right|^{2}$$

$$= O\left(\delta^{2}\right) \ln \lambda. \tag{4.4}$$

By (4.3) and (4.4), it follows from Jensen's inequality that

$$J_{\rho,\alpha}\left(\phi_{\lambda,\sigma}\right) \leq \frac{1}{2\rho} \int_{\Sigma} \left|\nabla\phi_{\lambda,\sigma}\right|^{2} + \int_{\Sigma} \phi_{\lambda,\sigma} - \ln\int_{\Sigma} e^{\phi_{\lambda,\sigma}} + C \int_{\Sigma} \left(\phi_{\lambda,\sigma} - \int_{\Sigma} \phi_{\lambda,\sigma}\right)^{2} + C$$

$$\leq \frac{1}{2\rho} \int_{\Sigma} \left|\nabla\phi_{\lambda,\sigma}\right|^{2} + C \int_{\Sigma} \left(\phi_{\lambda,\sigma} - \int_{\Sigma} \phi_{\lambda,\sigma}\right)^{2} + C$$

$$\leq \left(\frac{16k\pi}{\rho} - 2 + O\left(\delta^{2}\right)\right) \ln\lambda + C. \tag{4.5}$$

Since $\rho > 8k\pi$, choosing δ small and λ large, by (4.5), we obtain

$$J_{\rho,\alpha}\left(\phi_{\lambda,\sigma}\right) \leq -\left(1 - \frac{8k\pi}{\rho}\right) \ln \lambda.$$

This ends the proof of the lemma.

Define $\Phi_{\lambda}(\sigma) = \phi_{\lambda,\sigma}$. According to Lemma 4.1 and Lemma 4.2, we conclude that for large L and $\lambda \geq \lambda_L = e^{L/(1-8k\pi/\rho)}$

$$\Sigma_k \xrightarrow{\Phi_{\lambda}} J_{\rho,\alpha}^{-(1-8k\pi/\rho)\ln\lambda} \xrightarrow{\Psi} \Sigma_k$$

and $\lim_{\lambda \to +\infty} \Psi \circ \Phi_{\lambda} = \text{Id.}$ In particular, $\Psi \circ \Phi_{\lambda}$ is homotopic to the identity on Σ_k provided $\lambda \ge \lambda_L$.

Now we can prove the existence result by employing a minimax method.

Proof of Theorem 1.5. Let $\hat{\Sigma}_k = \Sigma_k \times [0, 1]/\Sigma \times \{0\}$ denote the cone over Σ_k . Choose *L* large and consider the following class

$$\Gamma_{\lambda} = \Gamma_{\lambda,\rho} = \left\{ \gamma \in C^0 \left(\hat{\Sigma}_k, H^1 \left(\Sigma \right) \right) : \gamma(\cdot \times \{1\}) = \Phi_{\lambda} \right\}, \quad \forall \lambda \geq e^{L/(1 - 8k\pi/\rho)}.$$

It is clear that Γ_{λ} is nonempty. Set

$$\bar{\Gamma}_{\lambda} = \bar{\Gamma}_{\lambda,\rho} = \inf_{\gamma \in \Gamma_{\lambda_M}} \sup_{z \in \hat{\Sigma}_k} J_{\rho,\alpha}\left(\gamma(z)\right) \le -\left(1 - \frac{8k\pi}{\rho}\right) \ln \lambda.$$

Using the fact that Σ_k is non-contractible and $\Psi \circ \Phi_{\lambda}$ is homotopic to the identity, we have

$$\bar{\Gamma}_{\lambda} > -2\left(1 - \frac{8k\pi}{\rho}\right) \ln \lambda.$$

For otherwise, there exists $\gamma \in \Gamma_{\lambda}$ with

$$\sup_{z \in \hat{\Sigma}_k} J_{\rho,\alpha}\left(\gamma(z)\right) \le -\frac{3}{2} \left(1 - \frac{8k\pi}{\rho}\right) \ln \lambda.$$

$$t \mapsto \Psi \circ \gamma(\cdot, t)$$

gives a homotopy in Σ_k between $\Psi \circ \Phi_{\lambda} = \Psi \circ \gamma(\cdot, 1)$ and a constant map $\Psi \circ \gamma(\cdot, 0)$. But this is impossible since Σ_k is non-contractible and $\Psi \circ \Phi_{\lambda}$ is homotopic to the identity.

By the monotonicity trick (cf. [3, Lemma 5.1]), there exists $\Lambda \subset (1 - \mu_0, 1 + \mu_0)$ such that Λ is dense in $[1 - \mu_0, 1 + \mu_0]$. Moreover, for any $\mu \in \Lambda$, the functional $J_{\mu\rho,\alpha}$ posses a bounded Palais-Smale sequence $\{u_n\}$ at level $\bar{\Gamma}_{\lambda,\mu\rho}$. Standard arguments show that there is a critical point u_n of $J_{\mu_n\rho,\alpha}$ for each $\mu_n \in \Lambda$. Then applying the compactness result in Theorem 1.1 and the denseness of Λ , we obtain a critical point of $J_{\rho,\alpha}$ and hence complete the proof.

References

- [1] Aubin, T., 1979. Meilleures constantes dans le théorème d'inclusion de Sobolev et un théorème de Fredholm non linéaire pour la transformation conforme de la courbure scalaire. J. Functional Analysis 32, 148–174. URL: https://doi.org/10.1016/0022-1236(79)90052-1, doi:10.1016/0022-1236(79)90052-1.
- [2] Aubin, T., 1998. Some nonlinear problems in Riemannian geometry. Springer Monographs in Mathematics, Springer-Verlag, Berlin. URL: https://doi.org/10.1007/978-3-662-13006-3, doi:10.1007/978-3-662-13006-3.
- [3] Battaglia, L., Jevnikar, A., Malchiodi, A., Ruiz, D., 2015. A general existence result for the Toda system on compact surfaces. Adv. Math. 285, 937–979. URL: https://doi.org/10.1016/j.aim.2015.07.036, doi:10.1016/j.aim.2015.07.036.
- [4] Berger, M.S., 1971. Riemannian structures of prescribed Gaussian curvature for compact 2-manifolds. J. Differential Geometry 5, 325–332. URL: http://projecteuclid.org/euclid.jdg/1214429996.
- [5] Brezis, H., Merle, F., 1991. Uniform estimates and blow-up behavior for solutions of $-\Delta u = V(x)e^u$ in two dimensions. Comm. Partial Differential Equations 16, 1223–1253. URL: https://doi.org/10.1080/03605309108820797, doi:10.1080/03605309108820797.
- [6] Caffarelli, L.A., Yang, Y.S., 1995. Vortex condensation in the Chern-Simons Higgs model: an existence theorem. Comm. Math. Phys. 168, 321–336. URL: http://projecteuclid.org/euclid.cmp/1104272361.
- [7] Chang, K.C., 2005. Methods in nonlinear analysis. Springer Monographs in Mathematics, Springer-Verlag, Berlin.
- [8] Chang, K.C., Liu, J.Q., 1993. On Nirenberg's problem. Internat. J. Math. 4, 35–58. URL: https://doi.org/10.1142/S0129167X93000042, doi:10.1142/S0129167X93000042.
- [9] Chang, S.Y.A., Yang, P.C., 1987. Prescribing Gaussian curvature on S². Acta Math. 159, 215–259. URL: https://doi.org/10.1007/BF02392560, doi:10.1007/BF02392560.
- [10] Chang, S.Y.A., Yang, P.C., 1988. Conformal deformation of metrics on S^2 . J. Differential Geom. 27, 259–296. URL: http://projecteuclid.org/euclid.jdg/1214441783.
- [11] Chen, C.C., Lin, C.S., 2002. Sharp estimates for solutions of multi-bubbles in compact Riemann surfaces. Comm. Pure Appl. Math. 55, 728–771. URL: https://doi.org/10.1002/cpa.3014, doi:10.1002/cpa.3014.
- [12] Chen, C.C., Lin, C.S., 2003. Topological degree for a mean field equation on Riemann surfaces. Comm. Pure Appl. Math. 56, 1667–1727. URL: https://doi.org/10.1002/cpa.10107, doi:10.1002/cpa.10107.
- [13] Chen, W., Li, C., 2008. A priori estimate for the Nirenberg problem. Discrete Contin. Dyn. Syst. Ser. S 1, 225–233. URL: https://doi.org/10.3934/dcdss.2008.1.225, doi:10.3934/dcdss.2008.1.225.
- [14] Chen, W.X., Ding, W.Y., 1987. Scalar curvatures on S². Trans. Amer. Math. Soc. 303, 365–382. URL: https://doi.org/10.2307/2000798, doi:10.2307/2000798.
- [15] Chen, W.X., Li, C., 1991a. Classification of solutions of some nonlinear elliptic equations. Duke Math. J. 63, 615–622. URL: https://doi.org/10.1215/S0012-7094-91-06325-8, doi:10.1215/S0012-7094-91-06325-8.
- [16] Chen, W.X., Li, C., 1991b. Prescribing Gaussian curvatures on surfaces with conical singularities. J. Geom. Anal. 1, 359–372. URL: https://doi.org/10.1007/BF02921311, doi:10.1007/BF02921311.
- [17] D'Aprile, T., De Marchis, F., Ianni, I., 2018. Prescribed Gauss curvature problem on singular surfaces. Calc. Var. Partial Differential Equations 57, Paper No. 99, 36. URL: https://doi.org/10.1007/s00526-018-1373-3, doi:10.1007/s00526-018-1373-3.
- [18] De Marchis, F., López-Soriano, R., 2016. Existence and non existence results for the singular Nirenberg problem. Calc. Var. Partial Differential Equations 55, Art. 36, 35. URL: https://doi.org/10.1007/s00526-016-0974-y, doi:10.1007/s00526-016-0974-y.
- [19] De Marchis, F., López-Soriano, R., Ruiz, D., 2018. Compactness, existence and multiplicity for the singular mean field problem with sign-changing potentials. J. Math. Pures Appl. (9) 115, 237-267. URL: https://doi.org/10.1016/j.matpur.2017.11.007, doi:10.1016/j.matpur.2017.11.007.
- [20] Ding, W., Jost, J., Li, J., Wang, G., 1997. The differential equation $\Delta u = 8\pi 8\pi he^u$ on a compact Riemann surface. Asian J. Math. 1, 230–248. URL: https://doi.org/10.4310/AJM.1997.v1.n2.a3, doi:10.4310/AJM.1997.v1.n2.a3.
- [21] Ding, W., Jost, J., Li, J., Wang, G., 1999. Existence results for mean field equations. Ann. Inst. H. Poincaré Anal. Non Linéaire 16, 653–666. URL: https://doi.org/10.1016/S0294-1449(99)80031-6, doi:10.1016/S0294-1449(99)80031-6.
- [22] Djadli, Z., 2008. Existence result for the mean field problem on Riemann surfaces of all genuses. Commun. Contemp. Math. 10, 205–220. URL: https://doi.org/10.1142/S0219199708002776, doi:10.1142/S0219199708002776.
- [23] Djadli, Z., Malchiodi, A., 2008. Existence of conformal metrics with constant *Q*-curvature. Ann. of Math. (2) 168, 813–858. URL: https://doi.org/10.4007/annals.2008.168.813, doi:10.4007/annals.2008.168.813.
- [24] Fontana, L., 1993. Sharp borderline Sobolev inequalities on compact Riemannian manifolds. Comment. Math. Helv. 68, 415–454. URL: https://doi.org/10.1007/BF02565828, doi:10.1007/BF02565828.

- [25] Gilbarg, D., Trudinger, N.S., 2001. Elliptic partial differential equations of second order. Classics in Mathematics, Springer-Verlag, Berlin. Reprint of the 1998 edition.
- [26] Gui, C., Moradifam, A., 2018. The sphere covering inequality and its applications. Invent. Math. 214, 1169–1204. URL: https://doi.org/10.1007/s00222-018-0820-2, doi:10.1007/s00222-018-0820-2.
- [27] Han, Z.C., 1990. Prescribing Gaussian curvature on S^2 . Duke Math. J. 61, 679–703. URL: https://doi.org/10.1215/S0012-7094-90-06125-3, doi:10.1215/S0012-7094-90-06125-3.
- [28] Hong, J., Kim, Y., Pac, P.Y., 1990. Multivortex solutions of the abelian Chern-Simons-Higgs theory. Phys. Rev. Lett. 64, 2230–2233. URL: https://doi.org/10.1103/PhysRevLett.64.2230, doi:10.1103/PhysRevLett.64.2230.
- [29] Jackiw, R., Weinberg, E.J., 1990. Self-dual Chern-Simons vortices. Phys. Rev. Lett. 64, 2234–2237. URL: https://doi.org/10.1103/PhysRevLett.64.2234, doi:10.1103/PhysRevLett.64.2234.
- [30] Ji, M., 2004. On positive scalar curvature on S^2 . Calc. Var. Partial Differential Equations 19, 165–182. URL: https://doi.org/10.1007/s00526-003-0214-0, doi:10.1007/s00526-003-0214-0.
- [31] Kazdan, J.L., Warner, F.W., 1974. Curvature functions for compact 2-manifolds. Ann. of Math. (2) 99, 14–47. URL: https://doi.org/10.2307/1971012, doi:10.2307/1971012.
- [32] Moser, J., 1973. On a nonlinear problem in differential geometry, in: Dynamical systems (Proc. Sympos., Univ. Bahia, Salvador, 1971), pp. 273–280.
- [33] Nolasco, M., 2003. Nontopological *N*-vortex condensates for the self-dual Chern-Simons theory. Comm. Pure Appl. Math. 56, 1752–1780. URL: https://doi.org/10.1002/cpa.10109, doi:10.1002/cpa.10109.
- [34] Nolasco, M., Tarantello, G., 1998. On a sharp Sobolev-type inequality on two-dimensional compact manifolds. Arch. Ration. Mech. Anal. 145, 161–195. URL: https://doi.org/10.1007/s002050050127, doi:10.1007/s002050050127.
- [35] Nolasco, M., Tarantello, G., 1999. Double vortex condensates in the Chern-Simons-Higgs theory. Calc. Var. Partial Differential Equations 9, 31–94. URL: https://doi.org/10.1007/s005260050132, doi:10.1007/s005260050132.
- [36] Nolasco, M., Tarantello, G., 2000. Vortex condensates for the SU(3) Chern-Simons theory. Comm. Math. Phys. 213, 599–639. URL: https://doi.org/10.1007/s002200000252, doi:10.1007/s002200000252.
- [37] Ricciardi, T., Tarantello, G., 2000. Vortices in the Maxwell-Chern-Simons theory. Comm. Pure Appl. Math. 53, 811-851. URL: https://doi.org/10.1002/(SICI)1097-0312(200007)53:7<811::AID-CPA2>3.0.C0;2-F, doi:10.1002/(SICI)1097-0312(200007)53:7<811::AID-CPA2>3.0.C0;2-F.
- [38] Shi, Y., Sun, J., Tian, G., Wei, D., 2019. Uniqueness of the mean field equation and rigidity of Hawking mass. Calc. Var. Partial Differential Equations 58, Paper No. 41, 16. URL: https://doi.org/10.1007/s00526-019-1496-1, doi:10.1007/s00526-019-1496-1.
- [39] Spruck, J., Yang, Y.S., 1995. Topological solutions in the self-dual Chern-Simons theory: existence and approximation. Ann. Inst. H. Poincaré Anal. Non Linéaire 12, 75–97. URL: https://doi.org/10.1016/S0294-1449(16)30168-8, doi:10.1016/S0294-1449(16)30168-8.
- [40] Struwe, M., Tarantello, G., 1998. On multivortex solutions in Chern-Simons gauge theory. Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 1, 109–121.
- [41] Tarantello, G., 1996. Multiple condensate solutions for the Chern-Simons-Higgs theory. J. Math. Phys. 37, 3769–3796. URL: https://doi.org/10.1063/1.531601.doi:10.1063/1.531601.
- [42] Tarantello, G., 2010. Analytical, geometrical and topological aspects of a class of mean field equations on surfaces. Discrete Contin. Dyn. Syst. 28, 931–973. URL: https://doi.org/10.3934/dcds.2010.28.931, doi:10.3934/dcds.2010.28.931.
- [43] Yang, Y., 2015. Extremal functions for Trudinger-Moser inequalities of Adimurthi-Druet type in dimension two. J. Differential Equations 258, 3161–3193. URL: https://doi.org/10.1016/j.jde.2015.01.004, doi:10.1016/j.jde.2015.01.004.
- [44] Yang, Y., Zhu, X., 2018. Existence of solutions to a class of Kazdan-Warner equations on compact Riemannian surface. Sci. China Math. 61, 1109–1128. URL: https://doi.org/10.1007/s11425-017-9086-6, doi:10.1007/s11425-017-9086-6.