

Complexity analysis of Bayesian learning of high-dimensional DAG models and their equivalence classes

Quan Zhou and Hyunwoong Chang

Department of Statistics, Texas A&M University

Abstract

Structure learning via MCMC sampling is known to be very challenging because of the enormous search space and the existence of Markov equivalent DAGs. Theoretical results on the mixing behavior are lacking. In this work, we prove the rapid mixing of a random walk Metropolis-Hastings algorithm, which reveals that the complexity of Bayesian learning of sparse equivalence classes grows only polynomially in n and p , under some high-dimensional assumptions. A series of high-dimensional consistency results is obtained, including the strong selection consistency of an empirical Bayes model for structure learning. Our proof is based on two new results. First, we derive a general mixing time bound on finite state spaces, which can be applied to local MCMC schemes for other model selection problems. Second, we construct high-probability search paths on the space of equivalence classes with node degree constraints by proving a combinatorial property of DAG comparisons. Simulation studies on the proposed MCMC sampler are conducted to illustrate the main theoretical findings.

Keywords: finite Markov chains; greedy equivalence search (GES); locally informed proposals; Poincaré inequality; random walk Metropolis-Hastings; rapid mixing; strong selection consistency.

1 Introduction

1.1 Gaussian DAG models and equivalence classes

A directed acyclic graph (DAG) encodes a set of conditional independence (CI) relations among node variables, which can be read off using the “d-separation” criterion [52]. Structure learning of DAG models from observational data plays a fundamental role in causal inference and has found many applications in machine learning and statistical data analysis [34]. In genomics, for example, DAG is a convenient device for conducting pathway analysis and inferring interactions among genes or proteins [41, 21].

Two DAGs with different edge sets can encode the same set of CI relations, in which case we say both belong to the same (Markov) equivalence class. For example, the DAGs $i \rightarrow j \rightarrow k$ and $i \leftarrow j \rightarrow k$ are Markov equivalent: both encode only one CI relation $i \perp\!\!\!\perp k \mid j$ (i.e., i, k are independent given j). But they are not Markov equivalent to $i \rightarrow j \leftarrow k$, since the latter encodes only one CI relation $i \perp\!\!\!\perp k$. A Gaussian DAG model represents a set

of multivariate normal distributions that satisfy the CI constraints encoded by the DAG. Due to normality, Markov equivalence further implies distributional equivalence [23], and thus observational data alone cannot distinguish between Markov equivalent DAGs; this is a main challenge in devising efficient structure learning algorithms [14].

This paper is chiefly concerned with the following problem: given n i.i.d. observations from a p -variate DAG-perfect normal distribution, estimate the equivalence class of the underlying DAG model. This is a model selection problem where the model space is a collection of p -vertex equivalence classes. We are most interested in high-dimensional settings where p grows much faster than n and the true DAG model is sparse.

The structure learning problem can be greatly simplified if the topological ordering of the variables is known. By ordering, we mean a permutation $\sigma \in \mathbb{S}^p$, where \mathbb{S}^p denotes the symmetric group on $\{1, \dots, p\}$, such that for any $i < j$, an edge connecting $\sigma(i)$ and $\sigma(j)$ is always directed as $\sigma(i) \rightarrow \sigma(j)$. Such a total ordering always exists, but may not be unique, for a DAG due to acyclicity. For example, for $1 \rightarrow 3 \leftarrow 2$, the ordering can be either $(1, 2, 3)$ or $(2, 1, 3)$. Any two different DAGs that share a same ordering cannot be Markov equivalent. This can be proved by contradiction: if the two DAGs are Markov equivalent, they must have the same skeleton [73], but the ordering uniquely determines the directions of all edges implying that the two DAGs must be the same. Henceforth, we refer to the problem as DAG selection when the ordering is known and reserve the term “structure learning” for learning equivalence classes when the ordering is unknown; the latter is the focus of this paper.

1.2 Algorithms for Bayesian structure learning

Most Bayesian structure learning methods aim to produce a posterior distribution of the DAG model or its equivalence class, which can be further used for making inference on quantities of interest via model averaging. To numerically approximate the posterior distribution, Markov chain Monte Carlo (MCMC) sampling is often invoked, and existing MCMC methods differ from each other mainly in three respects: the state space, the set of local operators, and the proposal scheme. The local operators decide which states the sampler may move to in the next iteration (i.e., they define the “neighborhood” of each state). The proposal scheme refers to how the proposal probabilities of these neighboring states are assigned. Most existing algorithms use either random walk Metropolis-Hastings (MH) or Gibbs schemes, but we note that informed proposal schemes recently proposed in Zanella [78] and Zhou et al. [82] can be applied as well.

There are three popular choices of the state space: states can be DAGs, equivalence classes or orderings. The famous “structure MCMC” sampler is a random walk MH algorithm that searches the DAG space using addition, deletion and reversal of single edges [42, 24]. It is straightforward to implement (one only needs to check acyclicity when proposing local moves) but may not be efficient since the sampler can spend a lot of time traversing large equivalence classes. Various methods have been proposed to improve the performance by using more complicated local operators [27, 68] or blocked Gibbs schemes [25]. Directly searching the space of equivalence classes seems more efficient, but a major challenge is to construct a proper set of local operators [4, 56, 47, 14, 54]; see Madigan et al. [43] and Castelletti

et al. [11] for MCMC samplers defined on the space of equivalence classes. Order MCMC methods [20, 19, 1] target a posterior distribution on the order space \mathbb{S}^p . They are motivated by the observation that given ordering, the conditional posterior distribution of DAGs can be evaluated relatively easily. More sophisticated MCMC schemes can be built by using partial orderings [50, 35]. It should be noted that the choice of the prior distribution typically depends on the state space, which results in essentially different posterior distributions on the three state spaces (see Section 7.1). We will focus on the space of equivalence classes.

In principle, by treating the logarithm of the posterior probability as a scoring criterion, deterministic score-based search algorithms can also be used to find the structure that maximizes the score (i.e., the maximum a posteriori estimate). This approach appears less popular in the Bayesian structure learning literature, probably because it cannot quantify the uncertainty in estimation. One of the most important score-based algorithms is the greedy equivalence search (GES) proposed by Meek [45] and Chickering [15], a two-stage greedy search algorithm defined on the space of equivalence classes. Nandy et al. [48] were the first to prove the high-dimensional consistency of GES (i.e., the search returns the true equivalence class with high probability for sufficiently large n) using an assumption called strong faithfulness. Though it is known that strong faithfulness is very restrictive [71], such conditions appear to be necessary for proving high-dimensional consistency results for many search methods [33]. We refer readers to Drton and Maathuis [16] and Scutari et al. [63] for other scored-based structure learning methods.

1.3 Overview of the paper

While many MCMC methods for structure learning have been proposed, to our knowledge, no theoretical result on the mixing time is available. This is probably because the structure of the state space is highly complicated. The primary goal of this paper is to fill the gap by deriving non-asymptotic mixing time bounds. We find that structure MCMC and order MCMC methods are, unfortunately, hard to analyze due to the technical difficulty in bounding sizes of equivalence classes. The equivalence class sampler of Castelletti et al. [11] uses six graph operators to move between equivalence classes [28], but we can explicitly construct slow mixing examples for this sampler with fixed p . This motivates us to propose our own equivalence class sampler, RW-GES, which uses a random walk proposal scheme that mimics and generalizes the local moves employed by GES. We prove a high-dimensional rapid mixing result for the RW-GES sampler, which essentially says that, under some conditions, the number of iterations needed to find the true equivalence class grows only polynomially in n and p with high probability. The proof consists of three steps, which we now explain separately.

In Section 2, we first develop a general theory on the complexity of local MCMC algorithms for model selection. This section is self-contained and of considerable independent interest. We build a weighted path argument and use a Poincaré-type inequality [32] to obtain a novel, generally applicable mixing time bound under a unimodal assumption; see Condition 1 and Theorem 2. This result can be applied to other model selection problems such as variable selection and stochastic block models. It sharpens the existing mixing time

bounds for random walk MH algorithms in the literature [77, 83] and can be utilized to derive theoretical guarantees for locally informed MH algorithms [78]. Our theory also reveals a link between optimization and sampling: if for some model selection problem there is a greedy local search with consistency guarantee, it is hopeful that, with some modifications, we may convert the greedy algorithm to a local MH sampler that has provable rapid mixing property. In general, rapid mixing is more difficult to prove and more informative than the consistency of a greedy search, since the former characterizes the overall complexity of the algorithm and requires an analysis of the local posterior landscape in the whole state space.

The RW-GES sampler for structure learning is formally introduced in Section 3. To impose sparsity, we define the state space to be the set of all equivalence classes that satisfy some node degree constraints. We do not explicitly define the target posterior distribution in this section (which will be done in Section 4); instead, we assume the posterior has some consistency property that typically holds for sufficiently large sample sizes. To use Theorem 2, we need to verify its assumption for the structure learning problem, which requires us to bound the neighborhood size (see Lemma 1) and construct “canonical paths” for the RW-GES sampler. We show by examples (see Examples 2 and 3) that a major and unique challenge in the path construction is to verify that the equivalence classes located on the “boundary” of the restricted search space cannot be local modes. To overcome this, we introduce a “swap” proposal move to RW-GES and prove a key combinatorial property of DAGs in Lemma 2. Combining it with the well-known Chickering algorithm [15], we obtain the canonical paths of RW-GES.

In Section 4, we propose an empirical Bayes model for structure learning and prove that it has the desired high-dimensional consistency property assumed in Section 3. Our model generalizes the DAG selection model of Lee et al. [39], and we show that it yields the same marginal fractional likelihood for Markov equivalent DAGs. The main result in this section is Theorem 5, which gives the strong selection consistency of our structure learning model. For Bayesian methods, such consistency results have only been established lately for the DAG selection problem with known ordering [10, 39]. Roughly speaking, in our consistency result, the maximum degree of searched DAGs is allowed to grow at rate $\sqrt{\log p}$; see Remark 10. The analogous high-dimensional consistency results for both variable selection and DAG selection are obtained as intermediate steps of our proof of Theorem 5.

The rapid mixing of RW-GES now follows from the mixing time bound given in Theorem 2 and the results of Sections 3 and 4. It is formally stated in Section 5. For comparison, we provide two slow mixing examples in the same section. The first one (see Example 4) shows why it is difficult to relax a key assumption used in our analysis, which is called the “strong beta-min condition” and is similar to the strong faithfulness assumption. The second (see Example 5) illustrates that the equivalence class sampler of Castelletti et al. [11] may mix slowly when p is small and n is large. We conduct simulation studies in Section 6 to show that our theoretical results hold “approximately” for moderately large sample sizes and provide useful guidance on the use of RW-GES in practice. Section 7 concludes the paper with discussions on why the structure MCMC is difficult to analyze and potential extensions of RW-GES. All proofs are relegated to the supplementary material [81]. For

readers' convenience, a notation table is given in Supplement A.

2 Mixing time bounds for model selection problems

2.1 A general setup

In this section, we use $\Theta = \Theta_p$ to denote a finite model space for a general model selection problem with p variables; for example, for the structure learning problem, each $\theta \in \Theta$ can be a unique equivalence class. Let $\mathcal{N}: \Theta \rightarrow 2^\Theta$ be given such that $\theta \notin \mathcal{N}(\theta)$ for each $\theta \in \Theta$; \mathcal{N} is called a neighborhood function. We say θ' is a neighbor of θ if and only if $\theta' \in \mathcal{N}(\theta)$. We say \mathcal{N} is “symmetric” if $\theta \in \mathcal{N}(\theta')$ always implies $\theta' \in \mathcal{N}(\theta)$. When we need to emphasize Θ is equipped with \mathcal{N} , we denote the space by (Θ, \mathcal{N}) . Let π denote a posterior distribution on Θ for a Bayesian procedure; assume it is known up to a normalizing constant and $\pi(\theta) > 0$ for each θ . Given a function $h: (0, \infty) \rightarrow (0, \infty)$, define a Markov chain \mathbf{K}^h on Θ by

$$\mathbf{K}^h(\theta, \theta') = \frac{h(\pi(\theta')/\pi(\theta))}{\sum_{\tilde{\theta} \in \mathcal{N}(\theta)} h(\pi(\tilde{\theta})/\pi(\theta))} \mathbb{1}_{\mathcal{N}(\theta)}(\theta') \quad (1)$$

where $\mathbb{1}$ is the indicator function. That is, given current state θ , \mathbf{K}^h moves to some $\theta' \in \mathcal{N}(\theta)$ with probability $\propto h(\pi(\theta')/\pi(\theta))$. Given \mathbf{K}^h , define another Markov chain \mathbf{P}^h by

$$\mathbf{P}^h(\theta, \theta') = \begin{cases} \mathbf{K}^h(\theta, \theta') \min \left\{ 1, \frac{\pi(\theta') \mathbf{K}^h(\theta', \theta)}{\pi(\theta) \mathbf{K}^h(\theta, \theta')} \right\}, & \text{if } \theta' \neq \theta, \\ 1 - \sum_{\tilde{\theta} \neq \theta} \mathbf{P}^h(\theta, \tilde{\theta}) & \text{if } \theta' = \theta. \end{cases} \quad (2)$$

If \mathbf{P}^h is irreducible, then π is the unique stationary distribution of \mathbf{P}^h . To avoid periodicity, we will often work with the lazy version $\mathbf{P}_{\text{lazy}}^h = (\mathbf{P}^h + \mathbf{I})/2$, where \mathbf{I} is the identity matrix.

Definition 1 (local Metropolis-Hastings algorithms). We say \mathbf{P}^h defined by (2) is a local MH algorithm with local proposal \mathbf{K}^h . If $h \equiv 1$, we say \mathbf{P}^h is the random walk MH algorithm. If h is non-constant and non-decreasing, we say \mathbf{K}^h is a (locally) informed proposal and \mathbf{P}^h is a (locally) informed MH algorithm.

The locally informed MH algorithm was proposed by Zanella [78]. The main idea is to assign larger proposal probabilities to those neighboring states with larger posterior so that the chain can quickly move to high-posterior regions. Let h in (1) be $h(u) = u^a$ for some $a \geq 0$. Observe that when $a = 0$, \mathbf{K}^h is reduced to the random walk proposal, and when $a \rightarrow \infty$ we obtain the greedy search (see definition below). So informed proposals are generally more aggressive than random walk but less aggressive than greedy search.

Definition 2 (greedy local search). A greedy (local) search on (Θ, \mathcal{N}) with initial state $\theta^{(0)}$ generates $\theta^{(1)}, \theta^{(2)}, \dots$, sequentially by letting $\theta^{(i)} = \arg \max_{\theta' \in \mathcal{N}(\theta^{(i-1)}) \cup \{\theta^{(i-1)}\}} \pi(\theta')$ for each $i \geq 1$. The search stops and returns $\theta^{(j)}$ if $\theta^{(j)} = \theta^{(j-1)}$.

The efficiency of both greedy search and MH algorithms largely depends on the choice of \mathcal{N} . For model selection problems, $\mathcal{N}(\cdot)$ is usually much smaller than Θ so that the algorithm is computationally affordable. But \mathcal{N} should also provide enough connectivity so that the algorithm cannot get trapped at sub-optimal local modes (θ is a local mode if $\pi(\theta) > \pi(\theta')$ for any $\theta' \in \mathcal{N}(\theta)$). We measure the convergence rate of MH algorithms using mixing time.

Definition 3 (mixing time). Let \mathbf{P} be an irreducible and aperiodic transition matrix defined on a finite state space Θ , with stationary distribution π . Define its mixing time by

$$T_{\text{mix}}(\mathbf{P}) = \max_{\theta \in \Theta} \min\{t \geq 0: \|\mathbf{P}^t(\theta, \cdot) - \pi(\cdot)\|_{\text{TV}} \leq 1/4\},$$

where $\|\cdot\|_{\text{TV}}$ denotes the total variation distance which takes value in $[0, 1]$.

Remark 1. We say an MCMC algorithm is rapidly mixing if its mixing time grows at most polynomially in the complexity parameters n (sample size) and p (number of variables). For most high-dimensional model selection problems, the size of Θ grows at least super-polynomially with p . For variable selection, which is probably the best-studied problem, Yang et al. [77] proved the rapid mixing of a random walk MH algorithm, and Zhou et al. [82] showed that an informed MH algorithm can converge much faster and obtain a mixing time independent of p .

2.2 A multi-purpose path method

We propose a general method for bounding the mixing time of \mathbf{P}^h defined in (2) and proving consistency properties of the posterior distribution π . The bounds to be derived in this section are non-asymptotic, and p is treated as a fixed constant. We begin by assuming that the triple $(\Theta, \mathcal{N}, \pi)$ satisfies the following condition, where $|\cdot|$ denotes the cardinality of a set.

Condition 1. $|\Theta| < \infty$, \mathcal{N} is symmetric, and $\pi > 0$. There exists a function $g: \Theta \rightarrow \Theta$, a state $\theta^* \in \Theta$ and constants $t_1, t_2 > 0, p > 1$ such that (i) $|\mathcal{N}(\theta)| \leq p^{t_1}$ for each $\theta \in \Theta$, and (ii) $g(\theta) \in \mathcal{N}(\theta)$ and $\pi(g(\theta))/\pi(\theta) \geq p^{t_2}$ for each $\theta \neq \theta^*$.

Remark 2. Part (ii) is equivalent to either of the following statements.

- (a) For any $\theta \neq \theta^*$, $\max_{\theta' \in \mathcal{N}(\theta)} \pi(\theta') \geq p^{t_2} \pi(\theta)$.
- (b) For any $\theta \neq \theta^*$, there exists $k < \infty$ and a sequence $(\theta_0 = \theta, \theta_1, \theta_2, \dots, \theta_k = \theta^*)$ such that $\theta_i \in \mathcal{N}(\theta_{i-1})$ and $\pi(\theta_i)/\pi(\theta_{i-1}) \geq p^{t_2}$ for each $i = 1, \dots, k$.

We introduce the function g because, for model selection problems, one often verifies Condition 1 by explicitly identifying some $g(\theta)$ for each θ . There may exist many choices of g so that Condition 1 holds. Without loss of generality, we always define $g(\theta^*) = \theta^*$. Then, part (ii) implies that for any θ there exists $k \leq |\Theta|$ such that $g^k(\theta) = \theta^*$, and θ^* is the only attracting fixed point of g . We will call g a canonical transition function and a sequence of the form $(\theta, g(\theta), \dots, g^k(\theta) = \theta^*)$ a canonical path. We can think of a canonical path as a candidate “greedy search path” since the posterior keeps increasing along the path, but note that a greedy search does not necessarily follow a canonical path since $g(\theta)$ may not be the maximizer of π in $\mathcal{N}(\theta)$.

Roughly speaking, in the model selection context, θ^* can be thought of as the “true” data-generating model, and Condition 1 can be interpreted as an algorithmic consistency property since it implies that θ^* is the unique mode of π and the greedy search always returns θ^* ; see part (i) of Theorem 1. For variable selection, Yang et al. [77] proved that

Condition 1 holds with high probability under some mild high-dimensional assumptions and then used the canonical path method of Sinclair [64] to bound the mixing time of the random walk MH algorithm. We generalize their result to our setup.

Theorem 1. *Let $\Theta, \mathcal{N}, \pi, g, \theta^*, t_1, t_2, p$ be as given in Condition 1. Let \mathbf{P}^h be given by (2) and $\mathbf{P}_{\text{lazy}}^h = (\mathbf{P}^h + \mathbf{I})/2$ be its lazy version. The following statements hold.*

(i) *The greedy search always returns θ^* regardless of the initial state.*

(ii) *If $t_2 > t_1$, then $\pi(\theta^*) \geq 1 - p^{-(t_2-t_1)}$.*

(iii) *If $t_2 > t_1$, then*

$$T_{\text{mix}}(\mathbf{P}_{\text{lazy}}^h) \leq \frac{4\ell_{\max}}{\{1 - p^{-(t_2-t_1)}\} \min_{\theta \neq \theta^*} \mathbf{P}^h(\theta, g(\theta))} \log \left(\frac{4}{\pi_{\min}} \right),$$

where $\ell_{\max} = \max_{\theta \neq \theta^} \min\{k \geq 1 : g^k(\theta) = \theta^*\}$ and $\pi_{\min} = \min_{\theta \in \Theta} \pi(\theta)$.*

(iv) *If $t_2 > t_1$ and $h \equiv 1$, then $\mathbf{P}^h(\theta, g(\theta)) \geq p^{-t_1}$.*

Proof. See Supplement B.4. □

Remark 3. Part (ii) of Theorem 1 shows that π concentrates on θ^* , which can be further used to show the strong selection consistency of a Bayesian model selection procedure (see Section 4.3). This is very useful since we only require polynomial (in p) bounds for the ratio $\pi(g(\theta))/\pi(\theta)$ in Condition 1, while $|\Theta|$ may be (super-)exponential in p . For a random walk MH algorithm, by parts (iii) and (iv), the order of mixing time is given by $p^{t_1} \ell_{\max} \log \pi_{\min}^{-1}$. For the greedy search, note that ℓ_{\max} is an upper bound for the steps needed to find θ^* , and in each step the search needs to evaluate π for at most p^{t_1} states. Hence, the greedy search and random walk MH algorithm have very similar complexity.

We now show that the mixing time bound in Theorem 1 can be improved. The new bound given in Theorem 2 has two major advantages. First, it does not involve ℓ_{\max} , which can be large. Second, it replaces $\min_{\theta \neq \theta^*} \mathbf{P}^h(\theta, g(\theta))$ in Theorem 1 with $\min_{\theta \neq \theta^*} \mathbf{P}^h(\theta, \mathcal{N}^*(\theta))$, where $\mathcal{N}^*(\theta)$ is the set of all “desirable moves” for \mathbf{P}^h at θ including $g(\theta)$. This is key to bounding the mixing times of informed MH algorithms. To prove Theorem 2, we use a novel path argument that may be of independent interest. For each $\theta \neq \theta^*$, we construct a set of paths from θ to θ^* using all desirable moves. By properly weighting these paths, we are able to bound the mixing time using a Poincaré-type inequality [32], which significantly generalizes the canonical path method. See Supplement B for details.

Theorem 2. *Let $\Theta, \mathcal{N}, \pi, g, \theta^*, t_1, t_2, p$ be as given in Condition 1, and \mathbf{P}^h be given by (2). For each $\theta \neq \theta^*$, define $\mathcal{N}^*(\theta) = \{\theta' \in \mathcal{N}(\theta) : \pi(\theta') \geq p^{t_2} \pi(\theta)\}$. Let $\pi_{\min} = \min_{\theta \in \Theta} \pi(\theta)$. If $t_2 > t_1$, then,*

$$T_{\text{mix}}(\mathbf{P}_{\text{lazy}}^h) \leq \frac{2C(p, t_1, t_2) \log \left(\frac{4}{\pi_{\min}} \right)}{\min_{\theta \neq \theta^*} \mathbf{P}^h(\theta, \mathcal{N}^*(\theta))}, \quad \text{where } C(p, t_1, t_2) = \frac{1 + (1 - p^{t_1-t_2})^{-1}}{[1 - p^{(t_1-t_2)/2}]^2}.$$

Proof. See Supplement B.6. □

Remark 4. Theorem 2 can be used to immediately improve some existing mixing time bounds in the literature. Both Yang et al. [77] and Zhuo and Gao [83] proved the rapid mixing of a random-walk MH algorithm for some high-dimensional discrete-state-space problem by showing Condition 1 holds for some g and using the canonical path method underlying Theorem 1. Theorem 2 shows that ℓ_{\max} can be dropped (in an asymptotic setting where $p \rightarrow \infty$ and $t_1 < t_2$ are fixed, $C(p, t_1, t_2) \rightarrow 2$).

Remark 5. Another important application of Theorem 2 is the mixing time analysis of informed MH algorithms. Define $\mathcal{N}^t(\theta) = \{\theta' \in \mathcal{N}(\theta) : \pi(\theta') \geq p^t \pi(\theta)\}$. If t_2 is sufficiently large and the function h in (1) is chosen properly, it is often possible to show that $\min_{\theta \neq \theta^*} \mathbf{P}^h(\theta, \mathcal{N}^t(\theta)) \geq c$ for some $t > t_1$ and fixed constant $c > 0$. Indeed, for the LIT-MH algorithm for variable selection considered in Zhou et al. [82], one can follow their calculations to verify that this holds for $c = 1/4$, and then by Theorem 2, the order of the mixing time is only $\log \pi_{\min}^{-1}$. This cannot be achieved by using Theorem 1, since $\mathbf{P}^h(\theta, g(\theta))$ can be as small as $O(p^{-t_1})$ (e.g., when all neighboring states have the same posterior probabilities).

The theory developed in this section relies on Condition 1, which is a property of the triple $(\Theta, \mathcal{N}, \pi)$. If Condition 1 holds, one can use Theorem 2 to study the mixing times of any local MH algorithm. For simplicity, for the structure learning problem to be studied in the rest of this paper, we will only consider the random walk proposal, and our main task is to construct a triple $(\Theta, \mathcal{N}, \pi)$ that satisfies Condition 1. We will often define \mathcal{N} on Θ and then use \mathcal{N} to refer to a neighborhood relation on a restricted space $\Theta_0 \subset \Theta$; this means that the neighborhood of $\theta \in \Theta_0$ is given by $\mathcal{N}(\theta) \cap \Theta_0$. Note that even if $(\Theta, \mathcal{N}, \pi)$ satisfies Condition 1, $(\Theta_0, \mathcal{N}, \pi)$ may not, which is one challenge in the sparse structure learning problem to be considered.

3 The RW-GES sampler and its canonical paths

3.1 Notation and terminology

We set up the notation to be used for the structure learning problem. Let $[p] = \{1, \dots, p\}$ and $|\cdot|$ denote the cardinality of a set. A subset of $[p]$ is typically denoted by S . The Hamming distance between two sets S, S' is denoted by $\text{Hd}(S, S') = |S \setminus S'| + |S' \setminus S|$.

A DAG G is a pair (V, E) where V is the vertex set and $E \subset V \times V$ is the set of directed edges. Throughout the paper, we assume $V = [p]$ for DAG models, representing random variables X_1, \dots, X_p . Note that $(i, i) \notin E$ for any $i \in [p]$. Let $|G|$ denote the number of edges in the DAG G ; thus, $|G| = |E|$. We use the notation $i \rightarrow j \in G$ to mean that $(i, j) \in E$ and $(j, i) \notin E$. The notation $i \rightarrow j \notin G$ means that $(i, j) \notin E$. For two DAGs $G = (V, E)$ and $G' = (V, E')$, we write $G' = G \cup \{i \rightarrow j\}$ if $E' = E \cup (i, j)$, and $G' = G \setminus \{i \rightarrow j\}$ if $E' = E \setminus (i, j)$. We write $G = G'$ if and only if G and G' have the same vertex set and edge set. Given a DAG G , we say node i is a parent of node j (and node j is a child of node i) if $i \rightarrow j \in G$. Let $\text{Pa}_j(G) = \{i \in [p] : i \rightarrow j \in G\}$ denote the set of parents of node j ; the in-degree of node j is $|\text{Pa}_j(G)|$. The maximum in-degree of G is $\max_j |\text{Pa}_j(G)|$. Similarly, let $\text{Ch}_j(G) = \{i \in [p] : j \rightarrow i \in G\}$, and $|\text{Ch}_j(G)|$ is called the out-degree of node j . The

degree of a node is the sum of its in-degree and out-degree, and the maximum degree of G is $\max_j |\text{Pa}_j(G) \cup \text{Ch}_j(G)|$. We may simply write Pa_j if we are not referring to a specific DAG or the underlying DAG is clear from context. The Hamming distance between two DAGs G, G' is defined by $\text{Hd}(G, G') = \sum_{j \in [p]} \text{Hd}(\text{Pa}_j(G), \text{Pa}_j(G'))$.

An equivalence class of DAGs is typically denoted by \mathcal{E} . We always interpret \mathcal{E} as a set of DAGs, and use $|\mathcal{E}|$ to denote the number of member DAGs in \mathcal{E} . The equivalence class of a DAG G is also denoted by $[G]$; thus, $\mathcal{E} = [G]$ if and only if $G \in \mathcal{E}$. The set of CI relations encoded by a DAG G or an equivalence class \mathcal{E} is denoted by $\mathcal{CI}(G)$ or $\mathcal{CI}(\mathcal{E})$, respectively. Note that we always have $\mathcal{CI}(G) = \mathcal{CI}([G])$.

We say a p -variate distribution μ is Markovian w.r.t. a p -vertex DAG G and G is an independence map (I-map) of μ if all CI relations encoded by G hold for μ . If the converse is also true, we say μ is faithful or perfectly Markovian w.r.t. G , and G is a perfect map of μ [66, 67]. We say μ is DAG-perfect if there exists some DAG that is a perfect map of μ . A DAG G is an I-map of a DAG G' and its equivalence class $[G']$ if $\mathcal{CI}(G) \subseteq \mathcal{CI}(G')$, and G is a minimal I-map (of G') if any sub-DAG of G (different from G) is not an I-map of G' . Given the set $\mathcal{CI}(G)$, a minimal I-map of G with ordering σ , which we denote by G_σ , can be uniquely defined as follows: for any $i < j$, $\sigma(i) \rightarrow \sigma(j) \in G_\sigma$ if and only if nodes $\sigma(i), \sigma(j)$ are not conditionally independent given nodes $\{\sigma(1), \dots, \sigma(j-1)\} \setminus \{\sigma(i)\}$ [65]. An example for $p = 3$ is given below. If μ is a p -variate positive measure, a unique minimal I-map of $\mathcal{CI}(\mu)$ with ordering σ can be constructed in an analogous manner [34, Chapter 3.4].

Example 1. Let $p = 3$ and G be the DAG $1 \rightarrow 3 \leftarrow 2$. Let G_σ denote the minimal I-map of G with ordering σ . If $\sigma = (1, 2, 3)$ or $(2, 1, 3)$, then $G_\sigma = G$ since σ is an ordering of G . If σ is any other ordering, then G_σ is the complete DAG (i.e., a DAG without missing edge). For example, if $\sigma = (1, 3, 2)$, G_σ has three edges $1 \rightarrow 3$, $1 \rightarrow 2$ and $3 \rightarrow 2$; in particular, the edge $1 \rightarrow 2$ is included since $1 \not\perp\!\!\!\perp 2 \mid 3$ in G .

3.2 Search spaces and posterior distributions

To apply the general theory developed in Section 2, it suffices to construct a triple $(\Theta, \mathcal{N}, \pi)$ that satisfies Condition 1. We will do this for both high-dimensional DAG selection and structure learning. Recall that for DAG selection, our goal is to estimate an underlying DAG model from the data when we know it has some ordering σ , and for structure learning, our goal is to estimate the equivalence class of the DAG model. We first define the search spaces (i.e., model space) for the two problems. Let \mathcal{G}_p denote the space of all p -vertex DAGs, which grows super-exponentially in p . We consider two sparsity constraints for DAGs, one for the maximum in-degree and the other for the maximum out-degree. For $d_{\text{in}}, d_{\text{out}} \in [p]$, define

$$\mathcal{G}_p(d_{\text{in}}, d_{\text{out}}) = \{G \in \mathcal{G}_p : \max_j |\text{Pa}_j(G)| \leq d_{\text{in}}, \text{ and } \max_j |\text{Ch}_j(G)| \leq d_{\text{out}}\}.$$

Since all Markov equivalent DAGs have the same skeleton, the two constraints ensure that the degree of any DAG $G' \in [G]$ for some $G \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$ is at most $d_{\text{in}} + d_{\text{out}}$. One may also use a single constraint for the maximum degree, but for the theoretical analysis to be

carried out in this paper, it is more convenient to specify $d_{\text{in}}, d_{\text{out}}$ separately. This setup is appealing to practitioners, since a DAG model with bounded degree is easier to visualize and interpret. Let $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ denote the space of “sparse equivalence classes” defined by

$$\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}) = \{[G]: G \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})\}.$$

Hence, $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ is the set of all equivalence classes that contain at least one member in $\mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$. We will use $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ as the model space for the sparse structure learning problem. The unrestricted space is denoted by $\mathcal{C}_p = \mathcal{C}_p(p, p)$.

Recall that \mathbb{S}^p is the space of all permutations of $[p]$. For each $\sigma \in \mathbb{S}^p$, let

$$\begin{aligned}\mathcal{G}_p^\sigma &= \{G \in \mathcal{G}_p: \sigma \text{ is a topological ordering of } G\} \\ &= \{G \in \mathcal{G}_p: \sigma(j) \rightarrow \sigma(i) \notin G \text{ for any } i < j\}.\end{aligned}$$

Note a DAG may have multiple orderings; in particular, the empty DAG belongs to \mathcal{G}_p^σ for any $\sigma \in \mathbb{S}^p$. Let $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}) = \mathcal{G}_p^\sigma \cap \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$ denote the space of sparse DAG models with ordering σ , which is the space we consider for the sparse DAG selection problem.

For our target posterior probability distributions, we assume they can be expressed by using a Bayesian scoring criterion $\psi: \mathcal{G}_p \rightarrow \mathbb{R}$ such that $\psi(G) = \psi(G')$ for any Markov equivalent G and G' , a property known as “score equivalence” [15]. For an equivalence class \mathcal{E} , define $\psi(\mathcal{E}) = \psi(G)$ using any $G \in \mathcal{E}$. Let the un-normalized posterior probability of a DAG G be given by $e^{\psi(G)}$, and that of an equivalence class \mathcal{E} be given by $e^{\psi(\mathcal{E})}$ (see Section 4.1 for more details). We further assume that ψ is decomposable: for each G ,

$$\psi(G) = \sum_{j \in [p]} \psi_j(\text{Pa}_j(G)),$$

where for each j , $\psi_j: 2^{[p]} \rightarrow \mathbb{R}$ gives the local score at node j .

3.3 Neighborhood functions and the RW-GES sampler

We define our neighborhood function on $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ by considering operations on all member DAGs of each equivalence class. To this end, we first define three neighborhoods on the unrestricted space \mathcal{G}_p for each DAG G , which correspond to three types of edge modification: addition, deletion and swap.

$$\begin{aligned}\mathcal{N}_{\text{add}}(G) &= \{G' \in \mathcal{G}_p: G' = G \cup \{i \rightarrow j\} \text{ for some } i \rightarrow j \notin G\}, \\ \mathcal{N}_{\text{del}}(G) &= \{G' \in \mathcal{G}_p: G' = G \setminus \{i \rightarrow j\} \text{ for some } i \rightarrow j \in G\}, \\ \mathcal{N}_{\text{swap}}(G) &= \{G' \in \mathcal{G}_p: G' = (G \cup \{k \rightarrow j\}) \setminus \{\ell \rightarrow j\} \text{ for some } k \rightarrow j \notin G, \ell \rightarrow j \in G\}.\end{aligned}$$

Note that a swap move consists of adding an incoming edge and deleting one at the same node, which is a straightforward extension of the swap proposal used in variable selection problems. Define the “add-delete-swap neighborhood” of G by

$$\mathcal{N}_{\text{ads}}(G) = \mathcal{N}_{\text{add}}(G) \cup \mathcal{N}_{\text{del}}(G) \cup \mathcal{N}_{\text{swap}}(G). \quad (3)$$

For each equivalence class $\mathcal{E} \in \mathcal{C}_p$, define

$$\mathcal{N}_{\text{ads}}(\mathcal{E}) = \{[G'] : G' \in \mathcal{N}_{\text{ads}}(G) \text{ for some } G \in \mathcal{E}\}, \quad (4)$$

and define the sets $\mathcal{N}_{\text{add}}(\mathcal{E})$, $\mathcal{N}_{\text{del}}(\mathcal{E})$ and $\mathcal{N}_{\text{swap}}(\mathcal{E})$ analogously; for example, $\mathcal{E}' \in \mathcal{N}_{\text{add}}(\mathcal{E})$ if and only if there exist $G \in \mathcal{E}$ and $G' \in \mathcal{E}'$ such that $G' \in \mathcal{N}_{\text{add}}(G)$. (The neighborhood notation is overloaded here, but the meaning should be clear from the argument.) Clearly, $\mathcal{N}_{\text{ads}}(\mathcal{E}) = \mathcal{N}_{\text{add}}(\mathcal{E}) \cup \mathcal{N}_{\text{del}}(\mathcal{E}) \cup \mathcal{N}_{\text{swap}}(\mathcal{E})$, and \mathcal{N}_{ads} is symmetric on both \mathcal{G}_p and \mathcal{C}_p . The following lemma gives a bound on the size of $\mathcal{N}_{\text{ads}}(\mathcal{E})$, which is needed later when we verify part (i) of Condition 1.

Lemma 1. *For any $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$,*

$$|\mathcal{N}_{\text{ads}}(\mathcal{E}) \cap \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})| \leq 3p(p-1)(d_{\text{in}} + d_{\text{out}})2^{d_{\text{in}} + d_{\text{out}}}.$$

Proof. See Supplement D.1. □

As explained in Section 2.1, we can construct a random walk MH algorithm on the restricted space $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ using \mathcal{N}_{ads} . The proposal distribution is given by $\mathbf{K}(\mathcal{E}, \mathcal{E}') = 1/|\mathcal{N}_{\text{ads}}(\mathcal{E})|$ for each $\mathcal{E}' \in \mathcal{N}_{\text{ads}}(\mathcal{E})$, where $\mathcal{N}_{\text{ads}}(\mathcal{E})$ is still defined on the unrestricted space (if we propose $\mathcal{E}' \notin \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, we simply reject the proposal). It should be noted that, in practice, there is no need to calculate the size of $\mathcal{N}_{\text{ads}}(\mathcal{E})$ or enumerate member DAGs in \mathcal{E} . States in $\mathcal{N}_{\text{ads}}(\mathcal{E})$ can be proposed very efficiently by using some local graph operators, which is explained in detail in Supplement H.1. We call this sampler random walk GES (RW-GES), since it uses a neighborhood function similar to that of the GES algorithm [15], which is a two-stage greedy search on the space \mathcal{C}_p that uses \mathcal{N}_{add} in the first stage and \mathcal{N}_{del} in the second. Swap moves are not used in GES, and we will use $\mathcal{N}_{\text{ges}}(\cdot) = \mathcal{N}_{\text{add}}(\cdot) \cup \mathcal{N}_{\text{del}}(\cdot)$ to denote the neighborhood relation used by GES.

3.4 Motivating examples

Assume the data-generating distribution is perfectly Markovian w.r.t. some DAG G^* (which henceforth is called the “true DAG”) and let $\mathcal{E}^* = [G^*]$ be the true equivalence class. In the classical asymptotic regime where p is fixed and sample size n tends to infinity, Chickering [15] proved that for a large class of Bayesian scoring criteria, GES and the greedy search on $(\mathcal{C}_p, \mathcal{N}_{\text{ges}})$ are consistent. According to our discussion following Condition 1, if we fix p and let $n \rightarrow \infty$, we can mimic the consistency proof of GES and use Theorem 1 to bound the mixing time of the random walk MH algorithm on $(\mathcal{C}_p, \mathcal{N}_{\text{ges}})$. The purpose of this subsection is to use examples to illustrate the technical challenges we encounter as we try to extend this argument to the space $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$.

To simplify the discussion, we assume the score ψ (i.e., log-posterior) satisfies the following condition, known as local consistency [15] (which is only used for making heuristic arguments in this section). It essentially says that all CI relations encoded by G^* can be correctly identified, which we expect to happen when $n = \infty$.

Condition 2. If distinct DAGs G, G' satisfy $G' = G \cup \{i \rightarrow j\}$, then (i) $\psi(G) > \psi(G')$ if $i \perp\!\!\!\perp j \mid \text{Pa}_j(G)$ in G^* , and (ii) $\psi(G') > \psi(G)$ if $i \not\perp\!\!\!\perp j \mid \text{Pa}_j(G)$ in G^* .

Under Condition 2, GES is consistent [15] and no equivalence class other than $[G^*]$ can be a local mode on $(\mathcal{C}_p, \mathcal{N}_{\text{ges}})$ (the reason will become clear in the next subsection). However, once we introduce the degree constraint (which is necessary for proving high-dimensional consistency results), local modes can arise on the boundary of the restricted space. To illustrate this, we construct two examples below. Example 2 explains why swap moves are useful and why in the consistency proof of GES we only consider edge removals when the current equivalence class is an I-map of \mathcal{E}^* . Example 3 shows that for the sparse DAG selection problem with degree constraints, local modes can also arise unexpectedly.

Example 2. Let $p = 3$ and DAGs G^*, G be given by

$$G^*: 1 \rightarrow 2 \rightarrow 3, \quad G: 2 \leftarrow 1 \rightarrow 3.$$

Consider how to increase the score of G by single-edge addition or deletion under Condition 2. Since $1 \not\perp\!\!\!\perp 2$ and $1 \not\perp\!\!\!\perp 3$ in G^* , both edges cannot be removed. However, since $2 \not\perp\!\!\!\perp 3 \mid 1$ in G^* , we can add the edge $2 \rightarrow 3$ to G to increase the score. The complete DAG $G \cup \{2 \rightarrow 3\}$ is an I-map of G^* , from which we should be able to remove the edge $1 \rightarrow 3$ since $1 \perp\!\!\!\perp 3 \mid 2$. One can apply the same argument to any other DAG in $\mathcal{E} = [G]$ and conclude that $\psi(\mathcal{E}) > \psi(\mathcal{E}')$ for any $\mathcal{E}' \in \mathcal{N}_{\text{del}}(\mathcal{E})$. In particular, we cannot remove the edge between nodes 1, 3 from any $G \in \mathcal{E}$, though the two nodes are not connected in G^* .

Next, we impose the constraint $d_{\text{in}} = 1$. Since G has two edges, we have $\mathcal{N}_{\text{add}}(\mathcal{E}) = \{\tilde{\mathcal{E}}\}$, where $\tilde{\mathcal{E}}$ is the equivalence class of all complete DAGs. But any complete DAG has maximum in-degree 2, which means that moving from \mathcal{E} to $\tilde{\mathcal{E}}$ is forbidden and \mathcal{E} is a local mode on $(\mathcal{C}_p(d_{\text{in}} = 1, d_{\text{out}} = p), \mathcal{N}_{\text{ges}})$. However, a swap move allows us to directly move from G to G^* by removing $1 \rightarrow 3$ and adding $2 \rightarrow 3$ simultaneously; that is, \mathcal{E} is not a local mode on $(\mathcal{C}_p(1, p), \mathcal{N}_{\text{ads}})$ where \mathcal{N}_{ads} is given by (4).

Example 3. Consider the DAG selection problem with $p = 5$ and $\sigma = (1, 2, 3, 4, 5)$. Let G^*, G be DAGs in \mathcal{G}_p^σ with edge sets

$$G^*: \{(1, 2), (1, 3), (2, 4), (2, 5)\}, \quad G: \{(1, 2), (1, 4), (2, 3), (2, 5)\}.$$

Under Condition 2, we can increase the score of G by adding $1 \rightarrow 3$ or $2 \rightarrow 4$, but deleting $1 \rightarrow 4$ or $2 \rightarrow 3$ will lower the score since $1 \not\perp\!\!\!\perp 4$ and $2 \not\perp\!\!\!\perp 3$ in G^* . Now let $d_{\text{in}} = d_{\text{out}} = 2$. Though $G^*, G \in \mathcal{G}_p^\sigma(2, 2)$, G is a local mode on $(\mathcal{G}_p^\sigma(2, 2), \mathcal{N}_{\text{ads}})$ because adding either $1 \rightarrow 3$ or $2 \rightarrow 4$ violates the out-degree constraint (note swap moves may not be helpful either).

3.5 Overview of the canonical path construction

Let the true DAG model $G^* \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$ and let $\mathcal{E}^* = [G^*]$. To verify Condition 1 for the triple $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$, we need to show that for any $\mathcal{E} \neq \mathcal{E}^*$, we can identify some $g(\mathcal{E}) \in \mathcal{N}_{\text{ads}}(\mathcal{E})$ such that $\psi(g(\mathcal{E})) > \psi(\mathcal{E})$. By Remark 2, this is equivalent to constructing a canonical path from any \mathcal{E} to \mathcal{E}^* . We briefly discuss the main idea behind our construction in this subsection. It will be helpful to think of the space $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ as the union of $\{\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}) : \sigma \in \mathbb{S}^p\}$ and think of structure learning as simultaneous DAG selection for all $p!$ orderings.

Suppose RW-GES starts at some \mathcal{E} which contains a member DAG $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ for an arbitrary $\sigma \in \mathbb{S}^p$. We will first construct a canonical path on $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, denoted by $(G_0 = G, G_1, G_2, \dots, G_k)$, where the terminal state G_k (if possible) is given by

$$\hat{G}(\sigma) = \arg \max_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})} \psi(G). \quad (5)$$

If Condition 2 holds and $G_\sigma^* \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, we claim $\hat{G}(\sigma) = G_\sigma^*$, where we recall G_σ^* is the minimal I-map of G^* with ordering σ . To show this, without loss of generality, assume $\sigma = (1, 2, \dots, p)$, and note that the following CI relations hold in G^* for each $j \in [p]$ by the definition of minimal I-maps (see Section 3.1):

$$j \perp\!\!\!\perp [j-1] \setminus \text{Pa}_j(G_\sigma^*) \mid \text{Pa}_j(G_\sigma^*), \text{ and } j \not\perp\!\!\!\perp i \mid [j-1] \setminus \{i\} \text{ for each } i \in \text{Pa}_j(G_\sigma^*). \quad (6)$$

Under Condition 2, the first property in (6) implies that if G is a DAG such that $\text{Pa}_j(G_\sigma^*) \subsetneq \text{Pa}_j(G)$, we can increase the score of G by removing some edge $\ell \rightarrow j$, and the second implies that if $\text{Pa}_j(G_\sigma^*) \not\subseteq \text{Pa}_j(G)$, we can add some edge $k \rightarrow j$ or perform a swap. This shows $\hat{G}(\sigma) = G_\sigma^*$ and suggests how we can construct the path from G to G_σ^* . However, as discussed in the previous subsection, the main challenge is to deal with the degree constraints.

Now suppose that RW-GES can move from \mathcal{E} to $[G_\sigma^*]$ following the path $(\mathcal{E}, \mathcal{E}_1, \mathcal{E}_2, \dots, \mathcal{E}_k)$ where $\mathcal{E}_i = [G_i]$. If $[G_\sigma^*] = \mathcal{E}^*$ (i.e., $G_\sigma^* = G^*$ or G_σ^* is Markov equivalent to G^*), we have obtained the path from \mathcal{E} from \mathcal{E}^* . If $[G_\sigma^*] \neq \mathcal{E}^*$, then one can use the famous Chickering algorithm [45, 15] to construct a path from $[G_\sigma^*]$ to \mathcal{E}^* (see Lemma D3 in Supplement D.4). Intuitively, since G_σ^* is an I-map of G^* , the skeleton of G^* must be a subset of the skeleton of G_σ^* (see Lemma C3), and we can remove edges from some other member DAG of $[G_\sigma^*]$.

Unfortunately, to rigorously prove that $\hat{G}(\sigma) = G_\sigma^*$ for all $\sigma \in \mathbb{S}^p$ in high-dimensional settings, one often needs to impose restrictive assumptions on the true data-generating mechanism, such as strong faithfulness [48]. To our knowledge, there is no fully satisfactory solution to this issue, and we will make a similar assumption in our theoretical analysis in Section 4 and assume $\hat{G}(\sigma) = G_\sigma^*$ in this section. Nevertheless, we will construct canonical paths of RW-GES using a flexible and finer argument, which, in some cases, can be used to show the rapid mixing of RW-GES under weaker assumptions; see Supplement I.

3.6 Canonical add-delete-swap paths of RW-GES

The discussion above suggests that we can construct the canonical paths of RW-GES by first constructing the canonical paths for all DAG selection problems. To this end, fix an arbitrary $\sigma \in \mathbb{S}^p$ first, and consider the sparse DAG selection problem with state space $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, neighborhood function \mathcal{N}_{ads} and posterior e^ψ . We treat G_σ^* as the true model, and we need to construct a candidate canonical transition function for this problem, $g^\sigma: \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}) \rightarrow \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, such that for any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$,

$$g^\sigma(G) \in \mathcal{N}_{\text{ads}}(G), \text{ and } (g^\sigma)^k(G) = G_\sigma^* \text{ for some } k < \infty.$$

For Condition 1 to hold we also need $\psi(g^\sigma(G)) > \psi(G)$. To overcome the out-degree constraint issue illustrated by Example 3, we will construct $g^\sigma(G)$ by first analyzing each node




Figure 1: An example for the operator g_j^σ . We consider four nodes with ordering $\sigma = (1, 2, 3, 4)$; assume $d_{\text{in}} = 3$. G_σ^* has three edges, $1 \rightarrow 2$, $2 \rightarrow 3$ and $3 \rightarrow 4$. Consider another DAG G with edges $1 \rightarrow 3$, $1 \rightarrow 4$ and $3 \rightarrow 4$. The DAGs $g_1^\sigma(G), g_2^\sigma(G), g_3^\sigma(G), g_4^\sigma(G)$ are shown above. For example, since $\text{Pa}_4(G_\sigma^*) = \{3\} \subset \text{Pa}_4(G) = \{1, 3\}$, node 4 is overfitted in G , and by part (ii) of Definition 4, $g_4^\sigma(G)$ is obtained by removing the edge $1 \rightarrow 4$ from G .

separately. Observe that if there is no out-degree constraint, the DAG selection problem is equivalent to p variable selection problems: for each j , we need to estimate the set Pa_j which takes value in the space $\mathcal{M}_p^\sigma(j, d_{\text{in}})$ defined by

$$\mathcal{M}_p^\sigma(j, d_{\text{in}}) = \{S \subseteq \mathcal{A}_p^\sigma(j) : |S| \leq d_{\text{in}}\}, \quad \mathcal{A}_p^\sigma(j) = \{k \in [p] : \sigma^{-1}(k) < \sigma^{-1}(j)\}, \quad (7)$$

where $\mathcal{A}_p^\sigma(j)$ is the set of variables that precede X_j in the ordering σ . Motivated by the discussion following (6), we construct a transition function on the space $\mathcal{M}_p^\sigma(j, d_{\text{in}})$ in Definition 4, which gives the “optimal” add-delete-swap move for Pa_j . Recall that we assume $\psi(G) = \sum_j \psi_j(\text{Pa}_j(G))$ for each G .

Definition 4. Assume $G_\sigma^* \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ and let $S_{\sigma,j}^* = \text{Pa}_j(G_\sigma^*)$. For each j , we construct $g_j^\sigma : \mathcal{M}_p^\sigma(j, d_{\text{in}}) \rightarrow \mathcal{M}_p^\sigma(j, d_{\text{in}})$ as follows. Fix an arbitrary $S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})$, and let $T = S_{\sigma,j}^* \setminus S$ and $R = S \setminus S_{\sigma,j}^*$.

- (i) If $S = S_{\sigma,j}^*$, let $g_j^\sigma(S) = S_{\sigma,j}^*$.
- (ii) If $S_{\sigma,j}^* \subset S$, let $g_j^\sigma(S) = S \setminus \{\tilde{\ell}\}$ where $\tilde{\ell} = \arg \max_{\ell \in R} \psi_j(S \setminus \{\ell\})$.
- (iii) If $S_{\sigma,j}^* \not\subseteq S$ and $|S| < d_{\text{in}}$, let $g_j^\sigma(S) = S \cup \{\tilde{k}\}$ where $\tilde{k} = \arg \max_{k \in T} \psi_j(S \cup \{k\})$.
- (iv) If $S_{\sigma,j}^* \not\subseteq S$ and $|S| = d_{\text{in}}$, let $g_j^\sigma(S) = (S \cup \{\tilde{k}\}) \setminus \{\tilde{\ell}\}$ where $(\tilde{k}, \tilde{\ell}) = \arg \max_{(k,\ell) \in T \times R} \psi_j((S \cup \{k\}) \setminus \{\ell\})$.

In case (ii), we say node j is (strictly) overfitted; in cases (iii) and (iv), we say it is underfitted. We use $g_j^\sigma(G)$ to denote the DAG obtained by replacing the parent set of j in G with $g_j^\sigma(\text{Pa}_j(G))$; that is, $\text{Pa}_j(g_j^\sigma(G)) = g_j^\sigma(\text{Pa}_j(G))$, and for any $i \neq j$, $\text{Pa}_i(g_j^\sigma(G)) = \text{Pa}_i(G)$.

Remark 6. It is clear from definition that $\text{Hd}(g_j^\sigma(S), S_{\sigma,j}^*) < \text{Hd}(S, S_{\sigma,j}^*)$ if $S \neq S_{\sigma,j}^*$. Further, $g_j^\sigma(G) \in \mathcal{N}_{\text{ads}}(G)$ and $\text{Hd}(g_j^\sigma(G), G_\sigma^*) < \text{Hd}(G, G_\sigma^*)$ if $\text{Pa}_j(G) \neq \text{Pa}_j(G_\sigma^*)$. In words, if node j is overfitted in G , $g_j^\sigma(G)$ is obtained by removing an incoming edge of node j . If node

j is underfitted, $g_j^\sigma(G)$ is obtained by adding an incoming edge of node j (if the in-degree constraint is violated, remove another incoming edge of node j). An example is provided in Figure 1. Note that this rationale is similar to that for GES and forward-backward stepwise regression. We always first transform an underfitted model to overfitted and then remove redundant variables or edges (recall Example 2).

Remark 7. Consider the variable selection problem with model space $\mathcal{M}_p^\sigma(j, d_{\text{in}})$ and true model $S_{\sigma,j}^*$. Yang et al. [77] proved that, under very mild high-dimensional assumptions, g_j^σ satisfies Condition 1 with high probability; that is,

$$\psi_j(g_j^\sigma(S)) - \psi_j(S) \geq t \log p, \quad \forall S \in \mathcal{M}_p^\sigma(j, d_{\text{in}}) \setminus \{\text{Pa}_j(G_\sigma^*)\}, \quad (8)$$

for some $t > 0$ (in their conclusion t is a universal constant).

Suppose that (8) holds for each j . Then, to show that the triple $(\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$ satisfies part (ii) of Condition 1, we only need to use the operators $\{g_j^\sigma : j \in [p]\}$ to construct a path from any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ to G_σ^* . At first glance, this seems trivial since we can use g_j^σ repeatedly to convert any $\text{Pa}_j(G)$ to $\text{Pa}_j(G_\sigma^*)$. However, the definition of g_j^σ only guarantees that $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, p)$, but the maximum out-degree of $g_j^\sigma(G)$ can be larger than that of G . Indeed, Example 3 in Section 3.4 shows that, in extreme cases, none of the operators $g_1^\sigma, \dots, g_p^\sigma$ yields a DAG that is different from G and belongs to $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. Fortunately, we are able to prove that, as long as d_{out} is chosen sufficiently large, there always exists some j such that g_j^σ yields a different DAG in $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. We define

$$d_\sigma^* = \max_{j \in [p]} |\text{Pa}_j(G_\sigma^*) \cup \text{Ch}_j(G_\sigma^*)|, \quad d^* = \max_{\sigma \in \mathbb{S}^p} d_\sigma^*, \quad (9)$$

where d^* will be used later in Theorem 3.

Lemma 2. Assume $d_\sigma^* \leq d_{\text{in}}$ and $\min\{d_\sigma^* d_{\text{in}} + 1, p\} \leq d_{\text{out}}$. For any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ such that $G \neq G_\sigma^*$, there exists some $j \in [p]$ such that $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ and $g_j^\sigma(G) \neq G$.

Proof. The key idea of the proof is to use the pigeonhole principle multiple times to derive the contradiction. See Supplement D.2. \square

Corollary 1. Let $\sigma \in \mathbb{S}^p$. Assume that $d_\sigma^* \leq d_{\text{in}}$ and $\min\{d_\sigma^* d_{\text{in}} + 1, p\} \leq d_{\text{out}}$.

- (i) There exists a function $g^\sigma : \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}) \rightarrow \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ such that for any $G \neq G_\sigma^*$, $g^\sigma(G) = g_j^\sigma(G) \neq G$ for some $j \in [p]$ and $(g^\sigma)^k(G) = G_\sigma^*$ for some $k \leq (d_\sigma^* + d_{\text{in}})p$.
- (ii) If (8) holds for each $j \in [p]$, Condition 1 holds for the triple $(\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$ with $t_1 = 3$ and $t_2 = t$.

Proof. See Supplement D.3. \square

We are now ready to construct a canonical transition function $g : \mathcal{C}_p(d_{\text{in}}, d_{\text{out}}) \rightarrow \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ for the structure learning problem using operators $\{g_j^\sigma : j \in [p], \sigma \in \mathbb{S}^p\}$. If \mathcal{E} contains a minimal I-map of G^* , we define $g(\mathcal{E})$ using Chickering algorithm [15]; see Lemma D3 in the supplement. If not, by the definition of $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, there exists $G \in \mathcal{E} \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ for

some $\sigma \in \mathbb{S}^p$, and we can define $g(\mathcal{E})$ using the function g^σ constructed for the DAG selection problem. But note that we need to fix the DAG representation of each \mathcal{E} so that $g(\mathcal{E})$ can be defined uniquely. We give an explicit construction of g in the proof of Theorem 3, the main result for this section.

Theorem 3. *Assume that $d^* \leq d_{\text{in}}$ and $\min\{d^*d_{\text{in}} + 1, p\} \leq d_{\text{out}}$. Then, $G_\sigma^* \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ for each $\sigma \in \mathbb{S}^p$. Further, there exists a function $g: \mathcal{C}_p(d_{\text{in}}, d_{\text{out}}) \rightarrow \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$ such that $g(\mathcal{E}^*) = \mathcal{E}^*$ and the following hold for any $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}}) \setminus \{\mathcal{E}^*\}$.*

- (i) $g(\mathcal{E}) = [g_j^\sigma(G)]$ for some $j \in [p]$, $\sigma \in \mathbb{S}^p$ and $G \in \mathcal{E} \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ such that $g_j^\sigma(G) \neq G$.
- (ii) There exist $k \leq (d^* + d_{\text{in}})p$ and $k \leq \ell \leq (2d^* + d_{\text{in}})p$ such that $g^k(\mathcal{E}) = G_\sigma^*$ for some $\sigma \in \mathbb{S}^p$ and $g^\ell(\mathcal{E}) = \mathcal{E}^*$.

Proof. See Supplement D.4. □

We conclude this section with the following corollary, which shows that to establish part (ii) of Condition 1 for the sparse structure learning problem, it only remains to prove that (8) holds for all j and σ simultaneously. This will be done rigorously in the next section.

Corollary 2. *Assume $d^* \leq d_{\text{in}}$, $\min\{d^*d_{\text{in}} + 1, p\} \leq d_{\text{out}}$ and ψ is score equivalent so that we can define $\psi(\mathcal{E}) = \psi(G)$ using any $G \in \mathcal{E}$. If (8) holds for each $\sigma \in \mathbb{S}^p$ and each $j \in [p]$, part (ii) of Condition 1 holds for the triple $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$ with $t_2 = t$.*

Proof. See Supplement D.5. □

4 High-dimensional consistency of an empirical Bayes model for structure learning

4.1 Model, prior and posterior distributions

Let X be an $n \times p$ data matrix where each row is an i.i.d. copy of a normal random vector $\mathbf{X} = (\mathbf{X}_1, \dots, \mathbf{X}_p)$. (The font for the random vector \mathbf{X} and that for the data matrix X are different.) Assume that, given a DAG G , the distribution of \mathbf{X} can be described by the structural equation model (SEM),

$$\mathbf{X} = B^\top \mathbf{X} + \mathbf{e}, \quad \mathbf{e} \sim N_p(0, \Omega), \quad (10)$$

for some $(B, \Omega) \in \mathcal{D}_p(G)$, where

$$\begin{aligned} \mathcal{D}_p(G) = \{ & (B, \Omega): B \in \mathbb{R}^{p \times p}, B_{ij} = 0 \text{ if } i \rightarrow j \notin G, \text{ for any } i, j \in [p]; \\ & \Omega = \text{diag}(\omega_1, \dots, \omega_p), \omega_i > 0 \text{ for any } i \in [p] \}. \end{aligned} \quad (11)$$

That is, each \mathbf{X}_j follows a linear regression model where explanatory variables with nonzero regression coefficients must be parents of node j in G . The matrix B is often called the weighted adjacency matrix. We can equivalently express (10) as

$$\mathbf{X} \sim N_p(0, \Sigma(B, \Omega)), \text{ where } \Sigma(B, \Omega) = (I - B^\top)^{-1} \Omega (I - B)^{-1} \quad (12)$$

is called the modified Cholesky decomposition (I denotes the identity matrix). The SEM representation of the Gaussian DAG model is used frequently in the literature [17, 72, 6].

Let $\pi_0(B, \Omega \mid G)$ denote the conditional prior distribution with support $\mathcal{D}_p(G)$. It suffices to specify it for $\{(\beta_j(G), \omega_j) : j = 1, \dots, p\}$, where $\beta_j(G)$ is the subvector of the j -th column of B with entries indexed by $\text{Pa}_j(G)$, and ω_j is the j -th diagonal element of Ω . We use the empirical prior proposed by Lee et al. [39], which is an extension of the empirical variable selection model of Martin et al. [44]. Our prior assumes that, given G , $(\beta_1(G), \omega_1), \dots, (\beta_p(G), \omega_p)$ are independently distributed according to

$$\pi_0(\omega_j \mid G) \propto \omega_j^{-\kappa/2-1},$$

$$\beta_j(G) \mid \text{Pa}_j(G) = S_j, \omega_j \sim N_{|S_j|} \left((X_{S_j}^\top X_{S_j})^{-1} X_{S_j}^\top X_j, \frac{\omega_j}{\gamma} (X_{S_j}^\top X_{S_j})^{-1} \right),$$

where $\gamma > 0, \kappa \geq 0$ are hyperparameters, X_j denotes the j -th column of the data matrix X , and X_S is the submatrix containing columns indexed by S . Next, we compute the marginal likelihood of G by integrating out (B, Ω) and using a fractional exponent $\alpha \in (0, 1)$ to offset the overuse of data caused by the empirical prior. The resulting fractional marginal likelihood is given by $f_\alpha(G) = \prod_{j=1}^p f_{\alpha,j}(\text{Pa}_j(G))$, where

$$f_{\alpha,j}(S) = (1 + \alpha\gamma^{-1})^{-|S|/2} \left\{ X_j^\top (I - X_S (X_S^\top X_S)^{-1} X_S^\top) X_j \right\}^{-(\alpha n + \kappa)/2}.$$

More details about this empirical prior are given in Supplement F.1.

For sparse DAG selection with ordering σ , the state space is $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. For each G on this space, we specify its prior probability by

$$\pi_0^\sigma(G) \propto (c_1 p^{c_2})^{-|G|}, \quad (13)$$

where $c_1 > 0, c_2 \geq 0$ are hyperparameters. We can then calculate the posterior distribution by $\pi_n^\sigma(G) \propto \pi_0^\sigma(G) f_\alpha(G)$. Using the fractional marginal likelihood f_α , we get

$$\pi_n^\sigma(G) \propto e^{\psi(G)} \mathbb{1}_{\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})}(G), \text{ where} \quad (14)$$

$$\psi(G) = \sum_{j=1}^p \psi_j(\text{Pa}_j(G)), \text{ and } e^{\psi_j(S)} = (c_1 p^{c_2})^{-|S|} f_{\alpha,j}(S). \quad (15)$$

For the sparse structure learning problem, we use the prior

$$\pi_0(\mathcal{E}, G) \propto (c_1 p^{c_2})^{-|G|} \pi_0(G \mid \mathcal{E}) \mathbb{1}_{\mathcal{E}}(G), \quad (16)$$

where $\pi_0(G \mid \mathcal{E})$ satisfies $\sum_{G \in \mathcal{E}} \pi_0(G \mid \mathcal{E}) = 1$. Denote the corresponding posterior distribution by π_n . Marginalizing out G from $\pi_n(\mathcal{E}, G)$, we get

$$\pi_n(\mathcal{E}) \propto \sum_{G \in \mathcal{E}} \pi_0(G \mid \mathcal{E}) e^{\psi(G)}.$$

In Lemma 3 below, we prove that ψ yields the same value for any Markov equivalent DAGs. Hence, we can define $\psi(\mathcal{E}) = \psi(G)$ using any $G \in \mathcal{E}$, and $\pi_n(\mathcal{E})$ can be expressed by

$$\pi_n(\mathcal{E}) \propto e^{\psi(\mathcal{E})} \mathbb{1}_{\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})}(\mathcal{E}). \quad (17)$$

The indicator function in (17) serves to remind us of the restricted search space. We do not consider estimating the DAG or ordering from π_n . Indeed, for $G \in \mathcal{E}$, $\pi_n(G)$ depends on the conditional prior probability $\pi_0(G \mid \mathcal{E})$, which we leave unspecified.

Lemma 3. *The function ψ defined by (15) satisfies that $\psi(G) = \psi(G')$ whenever G and G' are Markov equivalent DAGs.*

Proof. See Supplement F.2. □

We will refer to $\psi_j(\text{Pa}_j)$, $\psi(G)$, $\psi(\mathcal{E})$ as the scores of Pa_j , G and \mathcal{E} , respectively. Note that a scoring criterion derived from a nodewise normal-inverse-gamma prior for $(B, \Omega) \mid G$ does not necessarily have the property given in Lemma 3. For non-empirical prior distributions, see Geiger and Heckerman [23] and Peluso and Consonni [53] for related results.

4.2 High-dimensional setup

Let G^* denote the true DAG model and $\mathcal{E}^* = [G^*]$ be the true equivalence class that we want to recover from the data. Assume that each row of X is drawn independently from $N_p(0, \Sigma^*)$, a normal distribution perfectly Markovian w.r.t. G^* . We will show π_n defined in (17) concentrates on $[G^*]$ by first proving that for each σ , π_n^σ defined in (14) concentrates on the minimal I-map G_σ^* . Due to normality, G_σ^* can be equivalently defined by using the modified Cholesky decomposition.

Definition 5. Let Σ^* be positive definite and $N_p(0, \Sigma^*)$ be perfectly Markovian w.r.t. some DAG G^* . For each $\sigma \in \mathbb{S}^p$, let $\mathcal{D}_p(\sigma) = \cup_{G \in \mathcal{G}_p^\sigma} \mathcal{D}_p(G)$. By Lemma C6, we can define $(B_\sigma^*, \Omega_\sigma^*)$ to be the unique pair in $\mathcal{D}_p(\sigma)$ such that

$$(I - (B_\sigma^*)^\top)^{-1} \Omega_\sigma^* (I - B_\sigma^*)^{-1} = \Sigma^*.$$

Define G_σ^* to be the DAG such that $i \rightarrow j \in G_\sigma^*$ if and only if $(B_\sigma^*)_{ij} \neq 0$, which, by Lemma C5, is the minimal I-map of G^* with ordering σ .

Consider a high-dimensional setting with $p = p(n)$ tending to infinity. The true DAG model G^* , true covariance matrix Σ^* and prior parameters $c_1, c_2, \alpha, \gamma, d_{\text{in}}, d_{\text{out}}$ are all implicitly indexed by n . We say a constant is universal if it does not depend on n . To derive our consistency results, we need to make a few assumptions on the parameters and Σ^* .

(A1) There exist $\underline{\nu} = \underline{\nu}(n), \bar{\nu} = \bar{\nu}(n) > 0$ and a universal constant $\delta_0 > 0$ such that

$$0 < \frac{\underline{\nu}}{(1 - \delta_0)^2} \leq \lambda_{\min}(\Sigma^*) \leq \lambda_{\max}(\Sigma^*) \leq \frac{\bar{\nu}}{(1 + \delta_0)^2},$$

where $\lambda_{\min}, \lambda_{\max}$ denote the smallest and largest eigenvalues, respectively.

(A2) The sparsity parameter d_{in} and n, p satisfy that $d_{\text{in}} \log p = o(n)$.

(A3) Prior parameters satisfy that $\kappa \leq n$, $1 \leq c_1 \sqrt{1 + \alpha/\gamma} \leq p$, and

$$c_2 \geq (\alpha + 1)(4d_{\text{in}} + 6) + t$$

for some universal constant $t > 0$.

(A4) Assumption on the maximum in-degree of G_σ^* .

(A4.1) Let $\nu_0 = 4\bar{\nu}^2\underline{\nu}^{-4}(\bar{\nu} - \underline{\nu})^2$. For some $\sigma \in \mathbb{S}^p$, $(\nu_0 + 1) \max_{j \in [p]} |\text{Pa}_j(G_\sigma^*)| \leq d_{\text{in}}$.

(A4.2) Assumption (A4.1) holds for every $\sigma \in \mathbb{S}^p$.

(A5) Assumption on B_σ^*, B^* .

(A5.1) There exists a universal constant $C_\beta > 0$ such that for some $\sigma \in \mathbb{S}^p$,

$$\min \{ |(B_\sigma^*)_{ij}|^2 : (B_\sigma^*)_{ij} \neq 0 \} \geq 5(C_\beta + 4c_2) \frac{\bar{\nu}^2 \log p}{\alpha \underline{\nu}^2 n}, \quad (18)$$

where B_σ^* is given by Definition 5.

(A5.2) There exists a universal constant $C_\beta > 0$ such that (18) holds for every $\sigma \in \mathbb{S}^p$.

The first three assumptions are standard and commonly used in high-dimensional statistical theory. Assumption (A1) is the standard restricted eigenvalue condition [9]. Assumption (A2) controls the growth rates of p and d_{in} (which determines the maximum model size for nodewise variable selection), and, together with Assumption (A3), ensures that we cannot overfit the data; recall from (16) that the hyperparameter c_2 controls the penalty on the model size, so it plays the same role as the tuning parameter in the penalized likelihood methods. Such assumptions (especially a condition similar to $d_{\text{in}} \log p = o(n)$) are required for most high-dimensional problems including variable selection [75, 76, 77, 30], stochastic block model [22], covariance matrix estimation [36, 69, 51], undirected Gaussian graphical models [60, 7, 40] and DAG selection [10, 39]; see Banerjee et al. [8] for a recent review. Note that the numerical constants in our assumptions are very conservative. For example, Assumption (A3) suggests that c_2 should grow linearly with d_{in} , but in practice, one can use some c_2 much smaller than $4d_{\text{in}}$, which we will illustrate using a simulation study in Section 6.3.

Assumption (A4.1) requires that the maximum in-degree of the “true model” for DAG selection with ordering σ is sufficiently small compared with d_{in} . It is similar to Assumption D of Yang et al. [77] and is technically needed to show that an MH sampler using add-delete-swap moves cannot get stuck at DAG models with maximum in-degree equal to d_{in} . But unlike their setup, we assume both lower and upper restricted eigenvalues are available, which enables us to avoid imposing an irrepresentability condition as in Yang et al. [77, Assumption D]. Assumption (A4.2) restricts the maximum in-degree of all minimal I-maps of G^* , which is allowed to have the same order as d_{in} , if $\bar{\nu}, \underline{\nu}$ defined in Assumption (A1) can be bounded by universal constants.

Assumption (A5.1) is the well-known beta-min condition for DAG selection with ordering σ [10, 39]. According to Definition 5, the SEM representation (10) holds for $(B, \Omega) = (B_\sigma^*, \Omega_\sigma^*)$. Hence, Assumption (A5.1) just means that all nonzero regression coefficients (i.e., signal sizes) of the true SEM with ordering σ are sufficiently large. Assumption (A5.2) is for structure learning and assumes the beta-min condition holds uniformly over all $\sigma \in \mathbb{S}^p$; this is often known as the strong beta-min or permutation beta-min condition [71] and was used in Van de Geer and Bühlmann [72] and Aragam et al. [6]. If p and Σ^* are fixed, which implies

B_σ^* is fixed for all $\sigma \in \mathbb{S}^p$, then Assumption (A5.2) can always be satisfied by choosing some large n . We need the strong beta-min condition (or some similar assumption) since we want to first establish that with high probability, for every $\sigma \in \mathbb{S}^p$, the minimal I-map G_σ^* has the highest score among all DAGs in $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, which is needed for proving consistency results for structure learning. For methods based on CI tests, a similar assumption, known as “strong faithfulness”, is commonly used [48] (strong beta-min condition essentially replaces partial correlations in strong faithfulness with partial regression coefficients). Uhler et al. [71] showed that the volume of normal distributions that are strongly faithful is very small. Though strong faithfulness and strong beta-min condition are not directly comparable, both seem to be fairly restrictive [72]. Unfortunately, without them, we cannot preclude the possibility that GES or local MH algorithms get trapped at local modes; see Example 4 in Section 5.1. A discussion on how to overcome such limitations is given in Supplement I. We end this subsection with one more remark on Assumption (A1).

Remark 8. The restricted eigenvalue condition can be used to obtain some useful bounds related to B_σ^* and Ω_σ^* . Write $\Omega_\sigma^* = \text{diag}(\omega_{\sigma,1}^*, \dots, \omega_{\sigma,p}^*)$. The decomposition (12) implies that $\omega_{\sigma,k}^* \in (\underline{\nu}, \bar{\nu})$ for any $\sigma \in \mathbb{S}^p$ and $k \in [p]$ since the diagonal elements of Σ^* and $(\Sigma^*)^{-1}$ can be bounded by the extreme eigenvalues of Σ^* . Further, we can bound the ℓ^2 -norm of the true regression coefficients for node j by $\sum_{i \in [p]} (B_\sigma^*)_{ij}^2 \leq \omega_{\sigma,j}^* / \underline{\nu} - 1$, using the fact that the operator norm is no less than the ℓ^2 -norm of any column.

4.3 Strong selection consistency results

For a general model selection problem, we say a Bayesian procedure has strong selection consistency if the posterior probability of the true model converges to 1 in probability with respect to the true data-generating probability measure [31, 49, 10]. By part (ii) of Theorem 1, to prove the strong selection consistency, we only need to show that Condition 1 is satisfied for some universal $t_2 > t_1$.

We begin with the strong selection consistency for nodewise variable selection and DAG selection problems. It turns out that we only need (8) holds for any $j \in [p]$ and $\sigma \in \mathbb{S}^p$. By Corollary 2, this consistency property of $\{g_j^\sigma: j \in [p], \sigma \in \mathbb{S}^p\}$ is also key to the verification of Condition 1 for structure learning. The complete proof for Theorem 4 is highly technical, and the most involved step is to establish an analogous consistency result for a single variable selection problem using our empirical prior, which is treated in detail in Supplement E and may be of independent interest.

Theorem 4. *Let $X \in \mathbb{R}^{n \times p}$ have i.i.d. rows drawn from $N_p(0, \Sigma^*)$, which is perfectly Markovian w.r.t. G^* . Suppose Assumptions (A1), (A2), (A3), (A4.2) and (A5.2) hold. Let $t > 0$ be the universal constant given in Assumption (A3) and assume $C_\beta \geq 8t/3$. For sufficiently large n , with probability at least $1 - 3p^{-1}$, the following statements hold.*

(i) *Consistency of the operators $\{g_j^\sigma: j \in [p], \sigma \in \mathbb{S}^p\}$ given in Definition 4:*

$$\min \{ \psi_j(g_j^\sigma(S)) - \psi_j(S) : \sigma \in \mathbb{S}^p, j \in [p], S \in \mathcal{M}_p^\sigma(j, d_{\text{in}}) \setminus \{S_{\sigma,j}^*\} \} \geq t \log p,$$

where ψ_j is given in (15) and $S_{\sigma,j}^* = \text{Pa}_j(G_\sigma^*)$.

(ii) If $t > 2$, we have the strong selection consistency of nodewise variable selection,

$$\min_{\sigma \in \mathbb{S}^p} \min_{j \in [p]} \frac{\exp(\psi_j(S_{\sigma,j}^*))}{\sum_{S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} \exp(\psi_j(S))} \geq 1 - p^{-(t-2)},$$

where $\mathcal{M}_p^\sigma(j, d_{\text{in}})$ is defined in (7).

(iii) If $t > 3$, we have the strong selection consistency of sparse DAG selection,

$$\min_{\sigma \in \mathbb{S}^p} \frac{\exp(\psi(G_\sigma^*))}{\sum_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})} \exp(\psi(G))} \geq 1 - p^{-(t-3)},$$

where $\psi(G)$ is defined in (15).

Proof. See Supplement F.3. □

Remark 9. The universal constant t can be chosen arbitrarily large. Given any $t > 0$, in order that Theorem 4 holds, we can always choose some c_2 that has same order as d_{in} and assume that the universal constant C_β in Assumption (A5.2) is sufficiently large.

As a corollary, the strong selection consistency for a single DAG selection problem with ordering σ can be obtained by replacing Assumptions (A4.2) and (A5.2) with Assumptions (A4.1) and (A5.1). This result was also proved in Lee et al. [39] under similar assumptions, but the method we use is different (the primary goal of Lee et al. [39] was to derive minimax posterior convergence rates for the weighted adjacency matrix). Note that if σ is an ordering of G^* , then $G_\sigma^* = G^*$.

Corollary 3. Let $X \in \mathbb{R}^{n \times p}$ have i.i.d. rows drawn from the distribution $N_p(0, \Sigma^*)$, which is perfectly Markovian w.r.t. G^* . Suppose Assumptions (A1), (A2), (A3), (A4.1) and (A5.1) hold for some $t > 3$ and $C_\beta \geq 8t/3$. Let σ be as given in Assumptions (A4.1) and (A5.1). For sufficiently large n , with probability at least $1 - 3p^{-1}$,

$$\frac{\exp(\psi(G_\sigma^*))}{\sum_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})} \exp(\psi(G))} \geq 1 - p^{-(t-3)}$$

Proof. The proof is wholly analogous to that for Theorem 4. □

In order to show Condition 1 holds and use Theorem 1 to prove the strong selection consistency of sparse structure learning, it only remains to invoke Lemma 1 to bound the size of $\mathcal{N}_{\text{ads}}(\cdot)$ and then apply Corollary 2. Recall the definition of d_σ^* and d^* given in (9).

Theorem 5. Let $X \in \mathbb{R}^{n \times p}$ have i.i.d. rows drawn from $N_p(0, \Sigma^*)$, which is perfectly Markovian w.r.t. G^* . Suppose $d^* \leq d_{\text{in}}$, $d^* d_{\text{in}} + 1 \leq d_{\text{out}}$ and $d_{\text{in}} + d_{\text{out}} \leq t_0 \log_2 p$ for some universal constant $t_0 > 0$, and Assumptions (A1), (A2), (A3), (A4.2) and (A5.2) hold with $C_\beta \geq 8t/3$ and $t > t_0 + 3$. For sufficiently large n , with probability at least $1 - 3p^{-1}$,

$$\frac{\exp(\psi(\mathcal{E}^*))}{\sum_{\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})} \exp(\psi(\mathcal{E}))} \geq 1 - p^{-(t-t_0-3)},$$

where $\psi(\mathcal{E}) = \psi(G)$ for any $G \in \mathcal{E}$ and $\psi(G)$ is defined in (15). Further, the greedy search on $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$ returns \mathcal{E}^* regardless of the initial state.

Proof. See Supplement F.4. □

Remark 10. The assumption $d_{\text{in}} + d_{\text{out}} = O(\log p)$ is mild, since the total number of edges in the DAG may have order p even if $d_{\text{in}} + d_{\text{out}} = O(1)$. In light of Assumption (A4.2), we may assume d^*, d_{in} have approximately the same order. Thus, roughly speaking, the assumptions of Theorem 5 imply that d^*, d_{in} cannot grow faster than $\sqrt{\log p}$.

4.4 Consistency results for sub-Gaussian random matrices

The normality assumption on the true distribution of \mathbf{X} can be relaxed. We can extend the consistency result obtained in Theorem 4 to the case where X is a sub-Gaussian random matrix (we still consider the posterior distributions defined in Section 4.1). Let each row of X be an i.i.d. copy of a random vector \mathbf{X} which has mean zero, covariance matrix Σ^* and distribution μ . Assume that μ is sub-Gaussian with sub-Gaussian parameter bounded by a universal constant, and $N_p(0, \Sigma^*)$ is perfectly Markovian w.r.t. a DAG G^* (that is, μ is not necessarily perfectly Markovian w.r.t. G^*). This includes the case where some node variables are Gaussian and some are discrete and bounded [38]. Then, under a set of similar assumptions, we can prove a consistency result analogous to Theorem 4(i); see Theorem F1 in Supplement F.5. By Corollary 2, this proves part (ii) of Condition 1, and other strong selection consistency results in Theorem 4 follow.

The main idea of the proof of Theorem F1 is similar to the Gaussian case. We first generalize the variable selection results of Yang et al. [77] to random matrices, which is performed in Supplement E.3. However, the proof techniques are very different from the Gaussian case in that we need to use random matrix theory [74] and error propagation results to show that all minimal I-maps of G^* can be recovered from the empirical covariance matrix. The key distinction between the two scenarios is that in the sub-Gaussian case uncorrelatedness does not imply independence. Consequently, some calculations are more involved, and we need to require a slightly stronger assumption on c_2 : in the sub-Gaussian case, we require $d_{\text{in}} \bar{\nu}^4 / \underline{\nu}^6 = O(c_2)$, while in the Gaussian case we only need $d_{\text{in}} = O(c_2)$.

5 Mixing time results for Bayesian structure learning

5.1 Rapid mixing of the RW-GES sampler

Recall that RW-GES is simply the random walk MH algorithm defined by (2) with $h \equiv 1$ and the triple $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, \pi_n)$ where π_n is given by (17). In the proof of Theorem 5, we have verified that Condition 1 holds, and thus we can apply the mixing time bounds in Section 2.2 to obtain the main result of this work, rapid mixing of RW-GES.

Theorem 6. *Consider the setting of Theorem 5, and let $\pi_{\min} = \min_{\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})} \pi_n(\mathcal{E})$. Let \mathbf{P} denote the transition matrix of the RW-GES sampler and \mathbf{P}_{lazy} denote its lazy version. For sufficiently large n , with probability at least $1 - 3p^{-1}$, we have*

$$T_{\text{mix}}(\mathbf{P}_{\text{lazy}}) \leq Ct_0 p^{t_0+2} (\log p) \log \left(\frac{4}{\pi_{\min}} \right),$$

for some universal constant C , where t_0 is as given in Theorem 5.

Proof. See Supplement G.1. □

Corollary 4. Suppose Assumptions (A1) and (A2) hold. We have

$$\min_{\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})} \frac{\pi_n(\mathcal{E})}{\pi_n(\mathcal{E}^*)} \geq \left(c_1 p^{c_2} \sqrt{1 + \alpha/\gamma} \right)^{-p(d_{\text{in}} + d^*)} \left(\frac{2\bar{\nu}}{\underline{\nu}} \right)^{-p(\alpha n + \kappa)/2}.$$

Hence, under the setting of Theorem 6, the mixing time of the RW-GES sampler can be bounded by a polynomial of n and p .

Proof. See Supplement G.2. □

Remark 11. Corollary 4 implies that RW-GES is rapidly mixing with high probability. The term $\log \pi_{\min}$ in the mixing time bound is only used to handle the worst scenario where the chain starts from the state with minimum posterior probability. If the chain starts from some “good” estimate, the actual mixing rate of the chain can be much faster; see Sinclair [64, Proposition 1].

If in the beta-min condition, we only assume that the minimum edge weight of B^* (the weighted adjacency matrix of the true DAG G^*) is sufficiently large, the rapid mixing of RW-GES does not hold. It is not difficult to construct an explicit example where RW-GES is slowly mixing. In the following example, we let $p = 3$ be fixed and show that the mixing time grows exponentially in n . One can extend our example to the case $p = n$ by adding variables X_4, \dots, X_n such that, for any $j = 4, \dots, n$, the observed vector X_j is exactly orthogonal to all the other column vectors of the data matrix.

Example 4. Assume $p = 3$ and the true SEM is given by

$$X_1 = z_1, \quad X_2 = b_1 X_1 + z_2, \quad X_3 = b_2 X_2 + z_3,$$

where z_1, z_2, z_3 are vectors orthogonal to each other and $\|z_j\|_2^2 = n$ for each j . Thus, we can let the true DAG G^* be $1 \rightarrow 2 \rightarrow 3$. Suppose the prior parameters satisfy that $d_{\text{in}} = d_{\text{out}} = 2$, $c_2 = \sqrt{n}$, $\kappa = 0$, and c_1, α, γ are fixed constants such that $c_1 \sqrt{1 + \alpha/\gamma} = 1$. Assume the true regression coefficients $b_1, b_2 > 0$ are given by

$$b_1^2 = b_2^2 = \frac{K c_2 \log p}{\alpha n} = o(1),$$

where K is some large universal constant. So, b_1, b_2 satisfy the bound in (18). Consider the DAG \tilde{G} given by $1 \rightarrow 2 \leftarrow 3$, which has $[\tilde{G}] = \{\tilde{G}\}$. The topological ordering of \tilde{G} can be chosen to be $\sigma = (1, 3, 2)$, and the minimal I-map G_σ^* is a complete DAG. One can show that the edge weight of $1 \rightarrow 3$ in G_σ^* is $b_1 b_2$. It is easy to verify that $b_1^2 b_2^2 = o(c_2 n^{-1} \log p)$, so the true model fails to satisfy the strong beta-min condition. Indeed, we can prove that RW-GES is slowly mixing. See Supplement G.4.

5.2 Rapid mixing results for sparse DAG selection

Suppose the ordering is given and the search is restricted to $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. We can construct a random walk MH sampler using neighborhood function \mathcal{N}_{ads} defined in (3) and posterior distribution π_n^σ defined in (14), which is just the standard add-delete-swap MH sampler. Denote its transition matrix by \mathbf{P}^σ . If there is no out-degree constraint, by posterior modularity, one can perform sampling for the parent set of each node separately; thus, there is no need to directly draw DAG samples. However, when $d_{\text{out}} < p$, the posterior distributions of $\text{Pa}_1, \dots, \text{Pa}_p$ are not independent, and this add-delete-swap sampler provides a convenient solution. Since by Theorem 4(i) and Corollary 1, the triple $(\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, \pi_n^\sigma)$ satisfies Condition 1, the mixing time bound for \mathbf{P}^σ immediately follows from Theorem 2.

Theorem 7. *Suppose Assumptions (A1), (A2), (A3), (A4.1) and (A5.1) hold for some $\sigma \in \mathbb{S}^p$, $t > 3$ and $C_\beta \geq 8t/3$. Further, assume that $\min\{d_\sigma^* d_{\text{in}} + 1, p\} \leq d_{\text{out}}$. For sufficiently large n , with probability at least $1 - 3p^{-1}$, we have*

$$T_{\text{mix}}(\mathbf{P}_{\text{lazy}}^\sigma) \leq C d_{\text{in}} p^2 \log \left(\frac{4}{\pi_{\min}^\sigma} \right),$$

for some universal constant C , where $\pi_{\min}^\sigma = \min_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})} \pi_n^\sigma(G)$.

Proof. See Supplement G.3. □

Remark 12. The assumptions are much weaker than those used in Theorem 6. In particular, we can allow a much larger model size for each nodewise variable selection problem. This is mainly because for any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, we have $|\mathcal{N}_{\text{ads}}(G)| = O(d_{\text{in}} p^2)$. But for an equivalence class $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, the size of $\mathcal{N}_{\text{ads}}(\mathcal{E})$ may grow exponentially in $d_{\text{in}} + d_{\text{out}}$.

5.3 Slow mixing examples for a CPDAG sampler

The neighborhood $\mathcal{N}_{\text{ads}}(\mathcal{E})$ used in RW-GES can be very large for some \mathcal{E} , which seems to be undesirable. However, other choices of the neighborhood relation on \mathcal{C}_p (which may seem very reasonable) can cause the search algorithm to be trapped in sub-optimal local modes.

A popular approach to constructing sampling algorithms on \mathcal{C}_p is to use the CPDAG (completed partially directed acyclic graph) representations of equivalence classes. Any equivalence class \mathcal{E} can be uniquely represented by a CPDAG, a partially directed acyclic graph that satisfies two conditions: (i) it has the same skeleton as any $G \in \mathcal{E}$; (ii) an edge is directed if and only if the edge is directed in the same orientation in every $G \in \mathcal{E}$. A CPDAG is also called an essential graph [4]. One can define local proposal moves on \mathcal{C}_p by modifying CPDAGs. However, one can easily end up with a CPDAG sampler that is slowly mixing even when p is fixed and n goes to infinity.

Example 5. Let $p = 3$ and the true data-generating DAG G^* be $1 \rightarrow 3 \leftarrow 2$. Since G^* is the only member in $\mathcal{E}^* = [G^*]$, the CPDAG of \mathcal{E}^* is the same as G^* . Let $\tilde{\mathcal{E}}$ be the equivalence class that contains all complete DAGs. It is easy to verify that the CPDAG of $\tilde{\mathcal{E}}$ is a complete undirected graph. If we define the neighborhood of $\tilde{\mathcal{E}}$ as all the CPDAGs that can be obtained by adding or removing a directed or undirected edge from $\tilde{\mathcal{E}}$, then the

only CPDAGs we can move to from $\tilde{\mathcal{E}}$ are $1-2-3$, $1-3-2$ and $2-1-3$. However, given sufficiently large sample size, all these three CPDAGs should have much smaller score than $\tilde{\mathcal{E}}$. For example, the CPDAG $1-3-2$ encodes the CI relation $1 \perp\!\!\!\perp 2 \mid 3$, which does not exist in G^* , and thus connecting nodes 1 and 2 should increase the score. See Supplement G.5 for an explicit construction of this example and another 5-node example, where we further prove that the CPDAG sampler proposed by Castelletti et al. [11] is slowly mixing.

6 Simulation studies on the RW-GES sampler

6.1 A rapid mixing example

In this section, we present three simulation studies which illustrate the theoretical results we have proved. We first construct a rapid mixing example for $p = 100$ and $n = 800$. In order to approximately satisfy the strong beta-min condition, we randomly generate the true DAG G^* such that its maximum node degree is 2 and its largest connected sub-DAG only has 10 nodes, and then for each edge (i, j) in G^* , we sample B_{ij}^* from the uniform distribution on $(0.5, 1.5) \cup (-1.5, -0.5)$. The DAG G^* we obtain has 66 edges, among which 24 are directed in the CPDAG representation of $[G^*]$; see Supplement H.2 for the visualization. Each row of the data matrix X is drawn independently from $N_p(0, \Sigma^*)$ where $\Sigma^* = (I - (B^*)^\top)^{-1}(I - B^*)^{-1}$. We use $\alpha = 0.99, \gamma = 0.01, \kappa = 0, c_1 = 1, c_2 = 2$ and run 20 RW-GES chains, all initialized at the null model, for 5×10^4 iterations. All 20 runs are able to find the true equivalence class in about 10^5 iterations, which indicates a fast mixing rate; see the left panel of Figure 2. This example illustrates that though the strong beta-min condition is restrictive, if the true DAG is sufficiently sparse and has a “simple” structure, RW-GES can be rapidly mixing for a moderately large sample size (in Supplement H.2, we use this idea to explicitly construct toy examples with $p \gg n$ that satisfy all assumptions of Theorem 6). For comparison, we repeat the analysis by only using the first 200 observations, and we find that 11 chains fail to sample $[G^*]$. The right panel of Figure 2 suggests that these 11 chains get stuck at different local modes. Notice that $[G^*]$ still seems to have the largest posterior probability in this case, which implies that the beta-min condition at least holds for the true ordering (i.e., if the true ordering is known, we can recover the true DAG). However, since n is small, the strong beta-min condition is significantly violated, which makes the posterior distribution on the space of equivalence classes highly multimodal.

6.2 Performance in a high-dimensional scenario

The complexity of the structure learning problem largely depends on p and the sparsity level of the true DAG G^* . We can roughly measure the sparsity using the maximum degree of G^* , denoted by $\deg(G^*)$. Assumption (A2) and Remark 10 suggest that we consider $\deg(G^*) \log p = O(n)$ and $\deg(G^*) = O(\sqrt{\log p})$. In the second simulation study, we examine these asymptotic orders by using 7 simulation settings where n grows linearly and p grows exponentially. Given p , we generate G^* by first sampling a random ordering and then including each edge with probability $D/(p-1)$, where the parameter D gives the expected




Figure 2: Trajectories of 20 independent RW-GES runs for a simulated data set with $p = 100$. Left: $n = 800$; right: $n = 200$. The posterior probabilities are un-normalized and the log-posterior of $[G^*]$ is set to zero. Red crosses mark the times that RW-GES first collects $[G^*]$. Runs that never sample $[G^*]$ are shown in blue.

number of neighbors of each node. We let D grow at rate \sqrt{n} (so we actually let $\deg(G^*) \log p$ grow slightly faster than n). We generate X using the normal SEM associated with G^* and choose the hyperparameters in the same way as in Section 6.1. The number of RW-GES iterations is set to grow polynomially with p but slightly slower than p^2 . We always initiate the sampler at the null model and discard the first 80% iterations as burn-in. For each setting, we generate 20 replicates (G^* and X are re-sampled each time), and the results are shown in Table 1 (see Supplement H.2 for the definition of true/false positive rates). Observe that the true positive rate (for both skeleton and CPDAG estimation) becomes stable as p grows, while the false positive rate even decreases. Though for most real-world problems, the strong beta-min condition is unlikely to be satisfied and $[G^*]$ may not be correctly identified, this study shows that the theoretical insights on the MCMC complexity is useful. In particular, the performance of RW-GES seems stable under the asymptotic regime $\deg(G^*) \log p = O(n)$.

p	n	D	$N_{\text{mcmc}}/1000$	TPR (skeleton)	TPR	FPR
7	60	1.549	3	0.854 (0.03)	0.721 (0.06)	0.047 (0.02)
14	90	1.897	10	0.89 (0.02)	0.668 (0.06)	0.03 (0.006)
28	120	2.191	30	0.91 (0.01)	0.73 (0.03)	0.017 (0.003)
56	150	2.449	100	0.871 (0.01)	0.629 (0.03)	0.015 (0.002)
112	180	2.683	300	0.866 (0.01)	0.634 (0.02)	0.0091 (0.0005)
224	210	2.898	1000	0.86 (0.008)	0.634 (0.01)	0.0049 (0.0002)
448	240	3.098	3000	0.869 (0.004)	0.648 (0.008)	0.0027 (0.00007)

Table 1: Performance of RW-GES in 7 settings. For the k -th setting, $p = 7 \cdot 2^{k-1}$, $n = 30(k+1)$, $D = 0.2\sqrt{n}$ and the number of RW-GES iterations $N_{\text{mcmc}} \approx 300 \cdot (4p/7)^{1.66}$. TPR (skeleton): true positive rate with edge directions ignored; TPR: true positive rate (edge directions determined by the CPDAG); FPR: false positive rate. Results are averaged over 20 replicates, and the number in parentheses is the standard error.

6.3 On the choice of c_2

The third simulation study aims to investigate the optimal choice of c_2 , which is the most important prior hyperparameter of our model since it determines the order of the penalty on the graph size. We fix $p = 20$ and $n = 100$ and generate 50 true DAGs and data sets using the method described in Section 6.2 with $D = 4$ (recall this gives the expected degree of a single node). When implementing RW-GES, we impose the maximum degree constraint, denoted by d (i.e., the sampler only searches equivalence classes with maximum degree bounded by d); see Supplement H.1 for details. RW-GES is run for 40,000 iterations for each simulated data set. We first fix $c_2 = 1.3$ and try $d = 4, 5, \dots, 9$. The results are shown in the left column of Table 2. True positive rates increase with d , since some nodes in the true DAG may have large degrees and their incoming edges cannot all be detected if d is small. However, the false positive rate also increases because the search space quickly grows with d . Next, we repeat the experiment by setting $c_2 = 1.1 + 0.1d$, which, according to our tests, appears to yield close-to-optimal performance in this simulation setting. As can be seen from the right column of Table 2, the false positive rate remains roughly a constant and the true positive rates are comparable or even better than those for $c_2 = 1.3$. Recall that to prove posterior consistency, we assume c_2 is greater than $4(\alpha + 1)d_{\text{in}}$ plus some constant in Assumption (A3). This simulation study shows that, though the coefficient in Assumption (A3) is quite pessimistic, the linear growth rate (w.r.t. the maximum degree constraint) is a useful rule of thumb for tuning c_2 in practice.

	$c_2 = 1.3$			$c_2 = 1.1 + 0.1d$		
d	TPR (skeleton)	TPR	FPR	TPR (skeleton)	TPR	FPR
4	0.542 (0.02)	0.307 (0.02)	0.0824 (0.004)	0.539 (0.01)	0.316 (0.02)	0.0788 (0.004)
5	0.61 (0.01)	0.339 (0.02)	0.101 (0.004)	0.601 (0.01)	0.325 (0.02)	0.0977 (0.004)
6	0.665 (0.01)	0.383 (0.02)	0.115 (0.005)	0.657 (0.01)	0.393 (0.02)	0.101 (0.005)
7	0.706 (0.01)	0.412 (0.02)	0.123 (0.006)	0.699 (0.01)	0.419 (0.02)	0.096 (0.005)
8	0.72 (0.01)	0.413 (0.02)	0.132 (0.006)	0.694 (0.01)	0.421 (0.02)	0.0993 (0.006)
9	0.718 (0.01)	0.401 (0.02)	0.138 (0.007)	0.695 (0.01)	0.437 (0.02)	0.0932 (0.005)

Table 2: Simulation study with $p = 20$, $n = 100$ and expected node degree $D = 4$. Results are averaged over 50 data sets.

7 Discussion

7.1 Mixing of structure MCMC and order MCMC methods

In this work, we have only analyzed the mixing times of MCMC algorithms defined on the space of equivalence classes, but the same strategy can be pursued to study samplers defined on the DAG space and order space. Observe that the canonical paths we constructed in Section 3.6 for the RW-GES sampler can also be thought of as paths on the DAG space. Given an equivalence class \mathcal{E} , we first pick arbitrarily some $G \in \mathcal{E}$. If G has ordering σ , we

move from G to the minimal I-map G_σ^* by only add-delete-swap modifications of the DAG. To move from G_σ^* to G^* , we have to change the ordering. For RW-GES, the neighborhood function defined in (4) allows us to “switch” from G_σ^* to a Markov equivalent DAG \tilde{G} , which is still an I-map of G^* but no longer minimal, and then we can remove edges from \tilde{G} (the existence of such \tilde{G} is guaranteed by Chickering algorithm). Repeating this procedure, we obtain a path from G_σ^* to G^* .

Consider the classical structure MCMC sampler, a random walk MH algorithms defined on \mathcal{G}_p that use single-edge addition, deletion and reversal to propose local moves [42, 12]. Since any two Markov equivalent DAGs G, G' are connected by a sequence of covered edge reversals (see Supplement C.1), structure MCMC is able to traverse equivalence classes and move from G_σ^* to any other Markov equivalent DAG. Therefore, the canonical paths of RW-GES are also paths of structure MCMC (introduce swap moves if a restricted space is considered). The same argument can be applied to order MCMC samplers, since a covered edge reversal can be seen as an adjacent transposition on \mathbb{S}^p [65]. Unfortunately, the size of an equivalence class can easily be very large, and it is unclear whether structure MCMC can always quickly leave any equivalence class even if the maximum degree is bounded. In Supplement G.6, we construct an interesting example where G_0 is Markov equivalent to $G^* \cup \{2 \rightarrow 1\}$ but it is quite difficult for structure MCMC to remove the edge between nodes 1 and 2 from G_0 . Indeed, we show that on average it takes structure MCMC $O(p^4)$ iterations to move from G_0 to G^* , while it only takes RW-GES $O(p^2)$ iterations to move from $[G_0]$ to $[G^*]$. Nevertheless, we conjecture that structure MCMC is still rapidly mixing under the assumptions we used in Section 4 (recall “rapid mixing” only requires the mixing time to be polynomial in n and p), though the proof would probably require a skillful analysis of how the size of an equivalence class changes with single-edge modifications of its member DAGs.

One caveat is that the target posterior distributions on DAG and order spaces are typically different from our target π_n defined in (17). For example, for DAG MCMC methods, it is convenient to use the prior $\pi_0^{\text{dag}}(G) \propto (c_1 p^{c_2})^{-|G|}$ for $G \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$, which yields the posterior $\pi_n^{\text{dag}}(G) \propto e^{\psi(G)} \mathbb{1}_{\mathcal{G}_p(d_{\text{in}}, d_{\text{out}})}(G)$. Comparing them with (16) and (17), we see that $\pi_0^{\text{dag}}(\mathcal{E}) = \sum_{G \in \mathcal{E}} \pi_0^{\text{dag}}(G) \propto |\mathcal{E}| \pi_0(\mathcal{E})$ and $\pi_n^{\text{dag}}(\mathcal{E}) \propto |\mathcal{E}| \pi_n(\mathcal{E})$. Note that we do not use π_0^{dag} for equivalence class samplers [11] since calculating the size of \mathcal{E} can be extremely time-consuming. On the order space, the situation is more subtle since one DAG can be compatible with multiple orderings [18, 19]. If rapid mixing of structure MCMC can be established, we expect that the same argument can be used to show the strong selection consistency of π_n^{dag} .

7.2 Advantages and extensions of RW-GES

One advantage of RW-GES over GES is that RW-GES considers a restricted search space and is equipped with the swap proposal. This is particularly important to theoretical analysis. Non-sparse models can easily overfit the data (e.g. if node j has more than n parents, then X_j can be perfectly explained leading to an infinite score), which is why the sparsity constraint is necessary for proving high-dimensional consistency results. For GES, even if the maximum degree of G^* is bounded, there is still a possibility that GES visits non-sparse equivalence

classes along its search path and then its behavior becomes completely unpredictable. In the proof of Nandy et al. [48] on the high-dimensional consistency of GES, the authors directly assumed that the output of the first stage is not too large; see Assumption (A5) therein.

The main methodological difference between the two algorithms is that GES is essentially an optimization algorithm, while RW-GES is used for sampling. The general theory on the relation between optimization and sampling suggests that each has its own unique advantages [70]. In particular, when the sample size is not large, the posterior tends to be multimodal and MCMC sampling (if it converges) can yield better estimates via model averaging [29]. One can also use the output of GES as the initial state for RW-GES, which, to some extent, may achieve the benefits of both methods. In our theoretical analysis, we choose to focus on RW-GES just for its simplicity. One can generalize it in many ways to improve its performance in practice, for example, by using an informed proposal scheme or combining it with tempering techniques (i.e., running multiple RW-GES samplers at different temperatures). One simple modification that may significantly improve the sampler’s performance is to first estimate a large conditional independence graph [46, 60] and then use it to tune the proposal probabilities. This can be seen as a randomized extension of the method of Nandy et al. [48]. The canonical paths we construct in Section 3 can always be applied as long as the sampler proposes states from $\mathcal{N}_{\text{ads}}(\cdot)$ (or a superset of it). But one important takeaway from our theory is that using a neighborhood smaller than $\mathcal{N}_{\text{ads}}(\cdot)$ may lead to slow mixing even when the sample size is sufficiently large. A detailed investigation into more sophisticated local MCMC schemes using $\mathcal{N}_{\text{ads}}(\cdot)$ is left to future research.

Acknowledgements

The authors would like to thank all anonymous reviewers whose comments have helped improve the quality of the paper.

Supplementary material

Part A: a notation table. Part B: more results for mixing times of finite Markov chains and proofs for Section 2. Part C: preliminaries for graphical models. Part D: proofs for Section 3. Part E: auxiliary results for high-dimensional empirical variable selection. Part F: proofs for Section 4. Part G: proofs and examples for Sections 5 and 7. Part H: further details about RW-GES implementation and simulation studies. Part I: discussion on the case where the strong beta-min or faithfulness condition fails.

A Notation used in the main text

In the table below, we list the notation that is used frequently in Sections 3 to 7.

Notation	Description
$[p]$	$\{1, 2, \dots, p\}$
\mathbb{S}^p	set of all permutations of $[p]$
$ S $	cardinality of a set S
$N_p(\mu, \Sigma)$	p -variate normal distribution with covariance matrix Σ
\mathbf{X}	a random vector with components X_1, \dots, X_p
X, X_j, X_S	data matrix, column vector, submatrix with columns index by S
$ G $	number of edges in the DAG G
$\text{Hd}(S, S'), \text{Hd}(G, G')$	Hamming distance between two sets or DAGs
$\text{Pa}_j(G), \text{Ch}_j(G)$	set of parents/children of node j in the DAG G
$[G]$	the equivalence class that contains the DAG G
$\mathcal{CI}(G), \mathcal{CI}(\mathcal{E})$	set of CI relations encoded by a DAG G or an equivalence class \mathcal{E}
$\mathcal{G}_p, \mathcal{G}_p^\sigma$	set of p -vertex DAGs, set of p -vertex DAGs with ordering σ
$\mathcal{G}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$	sets of DAGs that satisfy the in-degree and out-degree constraints
$\mathcal{C}_p, \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$	sets of equivalence classes
$\mathcal{A}_p^\sigma(j)$	set of nodes that precede X_j in the ordering σ
$\mathcal{M}_p^\sigma(j, d_{\text{in}})$	set of possible values of $\text{Pa}_j(G)$ for $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, p)$; see (7)
$g_j^\sigma(S), g_j^\sigma(G)$	canonical transition functions on $\mathcal{M}_p^\sigma(j, d_{\text{in}})$ and $\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$
$\mathcal{N}_{\text{ads}}(\mathcal{E})$	add-delete-swap neighborhood of an equivalence class \mathcal{E} ; see (4)
$\mathcal{N}_{\text{add}}, \mathcal{N}_{\text{del}}, \mathcal{N}_{\text{swap}}$	addition/deletion/swap neighborhood of G or \mathcal{E}
$\Sigma(B, \Omega)$	Σ with a modified Cholesky decomposition given by (B, Ω)
$\mathcal{D}_p(G)$	set of pairs (B, Ω) such that $\Sigma(B, \Omega)$ is Markovian w.r.t. G ; see (11)
$\mathcal{D}_p(\sigma)$	set of pairs (B, Ω) compatible with ordering σ ; see (C.1)
π_0, π_n	prior and posterior distributions or density functions [†]
$d_{\text{in}}, d_{\text{out}}$	maximum in-degree/out-degree
c_1, c_2, κ, γ	hyperparameters for $\pi_0(\mathcal{E})$ and $\pi_0(B, \Omega \mid G)$
α	exponent for the fractional likelihood function
ψ_j, ψ	posterior scores; see (15)
$\Sigma^*, G^*, \mathcal{E}^*$	true covariance matrix, DAG model and equivalence class
\mathbb{P}^*	probability measure corresponding to the true model
G_σ^*	minimal I-map of G^* with ordering σ
$(B_\sigma^*, \Omega_\sigma^*)$	modified Cholesky decomposition of Σ^* in $\mathcal{D}_p(\sigma)$
$S_{\sigma, j}^*$	parent set of node j in G_σ^*
d_σ^*, d^*	maximum degree of the minimal I-map(s); see (9)
\mathbf{K}, \mathbf{P}	proposal and transition matrices of MH algorithms

[†]: when π_0 or π_n denotes a density function, its dominating measure depends on the context.

B Path methods and mixing times of Markov chains

B.1 On the equivalence between mixing time and hitting time

Loosely speaking, the mixing time $T_{\text{mix}}(\mathbf{P})$ gives the worst estimate for how many iterations it takes for a Markov chain to “enter stationarity”. Theorem 1 shows that if Condition 1 holds with $t_2 > t_1$, π concentrates on a single state θ^* . In this case, it turns out that entering stationarity essentially means to hit θ^* . Formally, we can prove that T_{mix} is equivalent to the expected hitting time of θ^* , up to constant factors, using the result of Peres and Sousi [55]; see Theorem B2 below. For an intuitive explanation, observe that if $\pi(\theta^*) \approx 1$, then $\mathbf{P}^t(\theta, \theta^*)$ needs to be sufficiently large so that $\|\mathbf{P}^t(\theta, \cdot) - \pi(\cdot)\|_{\text{TV}}$ is small, which suggests that hitting θ^* is necessary for the chain to “enter stationarity.” On the other hand, the chain regenerates each time it hits θ^* , and thus between two successive visits to θ^* , the chain has completed an independent cycle. So the length of each cycle gives an estimate for the mixing time.

Let Θ be finite and \mathbf{P} be the transition matrix of an irreducible Markov chain $(Y_t)_{t \in \mathbb{N}}$ such that \mathbf{P} is reversible with respect to π . Let \mathbb{Q}_θ denote the probability measure for $(Y_t)_{t \in \mathbb{N}}$ with initial value $Y_0 = \theta$, and let \mathbb{E}_θ be the corresponding expectation. For $t \in \mathbb{N}$, let $\mathbf{P}^t(\theta, \cdot) = \mathbb{Q}_\theta(Y_t \in \cdot)$ denote the t -step transition matrix. For any set $A \subseteq \Theta$, define the hitting time of A by $\text{hit}(A) = \min\{t \in \mathbb{N} : Y_t \in A\}$. Let $T_{\text{mix}}^{\text{L}}$ be the mixing time of the lazy chain with transition matrix $(\mathbf{P} + \mathbf{I})/2$. We have the following results.

Theorem B1 (Peres and Sousi [55]). *For some $a < 1/2$, define*

$$T_{\text{H}}^a = \max \{ \mathbb{E}_\theta[\text{hit}(A)] : \theta \in \Theta, A \subseteq \Theta, \pi(A) \geq a \}.$$

Then, $T_{\text{mix}}^{\text{L}}$ and T_{H}^a are equivalent up to constant factors.

Remark B1. This result was first proved by Aldous [2] for continuous-time Markov chains. Griffiths et al. [26] showed that the equivalence between $T_{\text{mix}}^{\text{L}}$ and T_{H}^a also holds for $a = 1/2$. “Up to constant factors” means that there exist constants $c_a, C_a > 0$ (which do not depend on \mathbf{P}) such that $c_a T_{\text{H}}^a \leq T_{\text{mix}}^{\text{L}} \leq C_a T_{\text{H}}^a$.

Theorem B2. *If there exists some state θ^* such that $\pi(\theta^*) > 1/2$, then $T_{\text{mix}}^{\text{L}}$ is equivalent, up to constant factors, to $T^* = \max_{\theta \in \Theta} \mathbb{E}_\theta[\text{hit}(\{\theta^*\})]$.*

Proof. Choose any $a \in (1 - \pi(\theta^*), 1/2)$. For any A with $\pi(A) \geq a$, we have $\theta^* \in A$ and thus $\text{hit}(A) \leq \text{hit}(\{\theta^*\})$. Hence, we have $T_{\text{H}}^a = \max \{ \mathbb{E}_\theta[\text{hit}(A)] : \theta \in \Theta, A = \{\theta^*\} \}$. The result then follows from Theorem B1. \square

Theorem B1 also suggests that rapid mixing is impossible if the chain can get stuck at some state with small stationary probability for exponentially many steps. This can be proved by an elementary calculation.

Theorem B3. *Consider an asymptotic setting where Θ, \mathbf{P}, π are implicitly indexed by n . For each n , assume there exists $\theta_0 \in \Theta$ such that $\pi(\theta_0) \leq 1/2$ and $\mathbf{P}(\theta_0, \theta_0) \geq 1 - e^{-cn}$, where $c > 0$ is a universal constant. Then $T_{\text{mix}}(\mathbf{P})$ cannot be bounded from above by any polynomial in n .*

Proof. Let $A_n = \Theta \setminus \{\theta_0\}$. By the property of total variation distance, $\|\mathbf{P}^t(\theta_0, \cdot) - \pi(\cdot)\|_{\text{TV}} \geq |\mathbf{P}^t(\theta_0, A_n) - \pi(A_n)|$. It then follows from Definition 3 that

$$\begin{aligned} T_{\text{mix}}(\mathbf{P}) &\geq \min \{t \in \mathbb{N}: |\mathbf{P}^t(\theta_0, A_n) - \pi(A_n)| \leq 1/4\} \\ &\geq \min \{t \in \mathbb{N}: \mathbf{P}^t(\theta_0, A_n) \geq 1/4\}, \end{aligned}$$

since $\pi(A_n) \geq 1/2$. Observe that $\mathbf{P}^t(\theta_0, \theta_0) \geq (1 - e^{-cn})^t \geq 1 - te^{-cn}$ for any $t \geq 1$. Hence, $\mathbf{P}^t(\theta_0, A_n) \leq te^{-cn}$, which yields the result. \square

B.2 Path methods for bounding mixing times

Let Θ be finite and \mathcal{N} be a symmetric neighborhood function. We set up some notation and definitions for describing edges and paths on the neighborhood graph (Θ, \mathcal{N}) . Let

$$\text{Edge}(\mathcal{N}) = \{(\theta, \eta) \in \Theta^2: \theta \in \mathcal{N}(\eta)\}$$

denote the set of all directed edges in (Θ, \mathcal{N}) ; in particular, (θ, η) and (η, θ) are treated as different edges for any $\theta \neq \eta$. Below is the definition of “paths” on (Θ, \mathcal{N}) . Note that we allow a path to contain repeated vertices (but not repeated edges), which is often known as a “trail” in the graph theory.

Definition B1. We say a finite sequence $\gamma = (\theta_0, \theta_1, \dots, \theta_{k-1}, \theta_k)$ is an \mathcal{N} -path (or simply path) from θ to θ' with length k if (i) $\theta_0 = \theta$, $\theta_k = \theta'$, (ii) for each $i = 1, \dots, k$, the edge $(\theta_{i-1}, \theta_i) \in \text{Edge}(\mathcal{N})$, and (iii) γ has no repeated edges. We will also denote such a path by $\gamma = (e_1, \dots, e_k)$ where $e_i = (\theta_{i-1}, \theta_i)$, and we write $e \in \gamma$ to mean that the path γ traverses the edge e . A path is assumed to contain at least one edge.

Throughout Supplement B, the letter e is reserved for denoting edges and γ for denoting paths, while elements of Θ are typically denoted by θ, η, z, w .¹ Given a path $\gamma = (\theta_0, \dots, \theta_k)$, its reversal is denoted by $\tilde{\gamma} = (\theta_k, \theta_{k-1}, \dots, \theta_0)$. Given $\gamma_1 = (\theta_0, \dots, \theta_k)$ and $\gamma_2 = (\eta_0, \dots, \eta_l)$ such that $\theta_k = \eta_0$, the concatenation of the two paths is denoted by $\gamma_1 \gamma_2 = (\theta_0, \dots, \theta_k, \eta_1, \dots, \eta_l)$. Let $\Gamma(\mathcal{N})$ denote the set of all \mathcal{N} -paths. For any $\theta \neq \eta$, let $\Gamma_{\theta\eta}(\mathcal{N})$ denote the set of all paths in $\Gamma(\mathcal{N})$ that start at θ and end at η ; sometimes we will also use the notation $\Gamma(\theta, \eta; \mathcal{N})$ as an alternative to $\Gamma_{\theta\eta}(\mathcal{N})$. We say (Θ, \mathcal{N}) is connected if $\Gamma_{\theta\eta}(\mathcal{N})$ is non-empty for any $\theta \neq \eta$.

We will prove the mixing time bounds in Theorems 1 and 2 for a larger class of Markov chains. All we need is the following assumption on \mathbf{P} .

Assumption B.1. Let \mathbf{P} be a Markov chain on Θ such that (i) \mathbf{P} is reversible with respect to some distribution $\pi > 0$, (ii) $\{\eta \neq \theta: \mathbf{P}(\theta, \eta) > 0\} = \mathcal{N}(\theta)$ for any $\theta \in \Theta$, and (iii) all eigenvalues of \mathbf{P} are non-negative.

Note that if (Θ, \mathcal{N}) is connected, then \mathbf{P} in Assumption B.1 is also irreducible. Our proofs of Theorems 1 and 2 rely on the following Poincaré-type inequality.

¹Unfortunately, we have to abuse some notation: γ is later used to denote a hyperparameter in our Bayesian structure learning model. But it should be clear that all notation used in Supplement B is not related to structure learning or DAG models, as we are considering a general state space Θ here.

Theorem B4. Let (Θ, \mathcal{N}) be connected and \mathbf{P} be given by Assumption B.1. For any $e = (\theta, \eta) \in \text{Edge}(\mathcal{N})$, define $\rho(e) = \pi(\theta)\mathbf{P}(\theta, \eta) = \pi(\eta)\mathbf{P}(\eta, \theta)$. Let $\phi: \Gamma(\mathcal{N}) \rightarrow [0, \infty)$ be such that

$$\sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N})} \phi(\gamma) = \pi(\theta)\pi(\eta), \quad \text{for any } \theta \neq \eta.$$

Then, for any function $\ell: \text{Edge}(\mathcal{N}) \rightarrow (0, \infty)$, we have

$$\text{Gap}(\mathbf{P})^{-1} \leq \max_{e \in \text{Edge}(\mathcal{N})} \left\{ \frac{1}{\rho(e)\ell(e)} \sum_{\theta \neq \eta} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_{\ell} \right\},$$

where $|\gamma|_{\ell} = \sum_{e \in \gamma} \ell(e)$ and $\text{Gap}(\mathbf{P})$ denotes the spectral gap of \mathbf{P} ,

Proof. See Saloff-Coste [62, Theorem 3.2.9] and Kahale [32]. \square

Remark B2. The function ℓ can be seen a generalized “length” function which assigns a weight to each $e \in \text{Edge}(\mathcal{N})$ [32]. If we let $\ell(e) = 1$ for each e , then $|\gamma|_{\ell}$ is just the length of γ and Theorem B4 reduces to the “multicommodity flow” method of Sinclair [64] (the function ϕ is called a “flow”). To bound the mixing time, we can apply Sinclair [64, Proposition 1] to get

$$T_{\text{mix}}(\mathbf{P}) \leq \frac{-\log[\min_{\theta \in \Theta} \pi(\theta)] + \log 4}{\text{Gap}(\mathbf{P})}, \quad (\text{B.1})$$

Note that the assumption that \mathbf{P} has non-negative spectrum is not needed for Theorem B4 but is necessary for (B.1).

Roughly speaking, to obtain good bounds on $\text{Gap}(\mathbf{P})$ using Theorem B4, we want to construct a flow ϕ by identifying at least one “high-probability” path between any $\theta \neq \eta$, where “high-probability” means that $\rho(e)$ is not too small for every edge e of the path. To prove Theorem 1, we only need to identify one such path between any $\theta \neq \eta$, which can be naturally constructed by using the function g in Condition 1; see Sections B.3 and B.4. To prove Theorem 2, we need a much finer construction of the flow ϕ which takes into account all “high-probability” moves at each state; see Sections B.5 and B.6.

B.3 A general method for constructing canonical paths

Before we prove Theorem 1, we first develop some general results for constructing “canonical path ensembles” using “canonical transition functions.”

Definition B2. A canonical path ensemble on (Θ, \mathcal{N}) is a set of \mathcal{N} -paths, one (and only one) for each ordered pair of two distinct states in Θ .

Definition B3. We say $g: \Theta \rightarrow \Theta$ is a canonical transition function on (Θ, \mathcal{N}) with a unique fixed point θ^* if (i) $g(\theta^*) = \theta^*$; (ii) for any $\theta \neq \theta^*$, $g(\theta) \in \mathcal{N}(\theta)$ and there exists some finite k such that $g^k(\theta) = \theta^*$.

Lemma B1. Suppose (Θ, \mathcal{N}) is connected, and fix some $\theta^* \in \Theta$. There exists a canonical transition function g on (Θ, \mathcal{N}) with fixed point θ^* . Further, g induces a canonical path ensemble on (Θ, \mathcal{N}) such that each canonical path is an \mathcal{N}_g -path, where $\mathcal{N}_g(\theta) = \{\theta' \in \Theta: g(\theta') = \theta, \text{ or } g(\theta) = \theta'\}$.

Proof. First, we show that such a function g exists. Since (Θ, \mathcal{N}) is connected, for any $\theta \neq \theta^*$, there exists a shortest \mathcal{N} -path from θ to θ^* , which we denote by $(\theta_0 = \theta, \theta_1, \dots, \theta_k = \theta^*)$. Define $\tilde{g}(\theta)$ to be the state θ_1 on this path. Clearly, \tilde{g} is a canonical transition function.

Next, we explicitly construct a canonical path ensemble \mathcal{T} using an arbitrary canonical transition function g . The path from θ to η in \mathcal{T} will be denoted by $\gamma_{\mathcal{T}}(\theta, \eta)$, which is unique. Let

$$\mathbb{N} = \{0, 1, \dots\}, \quad \mathbb{N}^+ = \{1, 2, \dots\}, \quad k(\theta) = \min\{i \in \mathbb{N} : g^i(\theta) = \theta^*\} < \infty.$$

For $\theta \neq \theta^*$, define $\gamma_{\mathcal{T}}(\theta, \theta^*) = (\theta, g(\theta), \dots, g^{k(\theta)}(\theta) = \theta^*)$ (note that it cannot contain any duplicate state since otherwise $k(\theta)$ does not exist). Since \mathcal{N} is symmetric, we have $\theta \in \mathcal{N}(g(\theta))$ for each $\theta \neq \theta^*$, and thus we can define $\gamma_{\mathcal{T}}(\theta^*, \theta) = \tilde{\gamma}_{\mathcal{T}}(\theta, \eta)$. The construction of $\gamma_{\mathcal{T}}(\theta, \eta)$ for $\eta \neq \theta^*$ is divided into three cases.

Case 1. $\eta = g^j(\theta)$ for some $j \in \mathbb{N}^+$.

Case 2. $\theta = g^i(\eta)$ for some $i \in \mathbb{N}^+$.

Case 3. Neither Case 1 nor Case 2 holds.

For Case 1, since $\gamma_{\mathcal{T}}(\theta, \theta^*) = (\theta, g(\theta), \dots, g^{j-1}(\theta), \eta, g^{j+1}(\theta), \dots, g^k(\theta) = \theta^*)$, we can simply define $\gamma_{\mathcal{T}}(\theta, \eta) = (\theta, g(\theta), \dots, g^{j-1}(\theta), \eta)$, which is a sub-path of $\gamma_{\mathcal{T}}(\theta, \theta^*)$. Case 2 can be handled similarly. For Case 3, we define

$$\gamma_{\mathcal{T}}(\theta, \eta) = \gamma_{\mathcal{T}}(\theta, \theta^*) \gamma_{\mathcal{T}}(\theta^*, \eta).$$

To prove $\gamma_{\mathcal{T}}(\theta, \eta)$ has no duplicate edges, it suffices to show that paths $\gamma_{\mathcal{T}}(\theta, \theta^*)$ and $\gamma_{\mathcal{T}}(\theta^*, \eta)$ do not share any states except θ^* . We prove it by contradiction. Suppose $w \neq \theta^*$ exists in both paths. Then $w = g^s(\theta) = g^t(\eta)$ for some $s, t \in \mathbb{N}^+$. Without loss of generality, assume $s > t$. But this implies that $\eta = g^{s-t}(\theta)$, which yields the contradiction. \square

B.4 Proof of Theorem 1

Proof of Theorem 1(i). We say θ is a local maximum if $\pi(\theta) \geq \pi(\theta')$ for any $\theta' \in \mathcal{N}(\theta)$. Part (i) follows upon observing that any $\theta \neq \theta^*$ cannot be a local maximum, and thus θ^* is the only local and also global maximum. \square

Proof of Theorem 1(ii). For each $\theta \in \Theta$ and $k \in \mathbb{N}$, let

$$g^{-k}(\theta) = \{\theta' \in \Theta : g^k(\theta') = \theta, g^{k-1}(\theta') \neq \theta\}.$$

Since \mathcal{N} is symmetric, we have $|g^{-1}(\theta)| \leq |\mathcal{N}(\theta)|$. Further, note that $\Theta = \bigcup_{k \geq 0} g^{-k}(\theta^*)$, and for each $k \geq 1$, we can write

$$g^{-k}(\theta^*) = \{g^{-1}(\theta') : \theta' \in g^{-(k-1)}(\theta^*)\}.$$

A recursive calculation using Condition 1 then shows that $\pi(\theta)/\pi(\theta^*) \leq p^{-kt_2}$ for any $\theta \in g^{-k}(\theta^*)$, and $|g^{-k}(\theta^*)| \leq p^{kt_1}$. Hence,

$$\frac{\sum_{\theta \in \Theta} \pi(\theta)}{\pi(\theta^*)} \leq \sum_{k=0}^{\infty} \frac{\pi(g^{-k}(\theta^*))}{\pi(\theta^*)} \leq \sum_{k=0}^{\infty} p^{-k(t_2-t_1)} = \frac{1}{1 - p^{-(t_2-t_1)}},$$

from which the result follows. \square

Proof of Theorem 1(iii). We prove the claim for any Markov chain \mathbf{P} that satisfies Assumption B.1, which includes $\mathbf{P}_{\text{lazy}}^h$ as a special case (note that $\mathbf{P}^h(\theta, g(\theta)) = 2\mathbf{P}_{\text{lazy}}^h(\theta, g(\theta))$ for $\theta \neq \theta^*$). The existence of g implies that (Θ, \mathcal{N}) is connected and \mathbf{P} is irreducible.

Let $\mathcal{T} = \{\gamma_{\mathcal{T}}(\theta, \eta) : \theta, \eta \in \Theta, \text{ and } \theta \neq \eta\}$ be the canonical path ensemble induced by g , as constructed in the proof of Lemma B.1. It is clear from construction that for any $\theta \neq \theta'$, we have $|\gamma_{\mathcal{T}}(\theta, \theta')| \leq |\gamma_{\mathcal{T}}(\theta, \theta^*)| + |\gamma_{\mathcal{T}}(\theta', \theta^*)| \leq 2\ell_{\max}$, where ℓ_{\max} is as defined in Theorem 1. By Sinclair [64, Corollary 6], which is a special case of Theorem B.4, we have

$$\frac{1}{\text{Gap}(\mathbf{P})} \leq \max_{e \in \text{Edge}(\mathcal{N})} \frac{2\ell_{\max}}{\rho(e)} \sum_{(\theta, \eta) : e \in \gamma_{\mathcal{T}}(\theta, \eta)} \pi(\theta)\pi(\eta). \quad (\text{B.2})$$

Consider an arbitrary $e = (z, w) \in \text{Edge}(\mathcal{N})$. Observe that for \mathcal{T} constructed in Lemma B.1,

$$\{(\theta, \eta) : e \in \gamma_{\mathcal{T}}(\theta, \eta)\} \neq \emptyset \text{ only if } w = g(z) \text{ or } z = g(w).$$

So, to bound the maximum term in (B.2), we can assume $e = (z, w)$ for $w = g(z)$, which implies that $z \neq \theta^*$ (the case $e = (w, z)$ can be analyzed similarly). Define

$$\Lambda(z) = \{\theta \in \Theta : z = g^k(\theta), k \in \mathbb{N}\}$$

as the ancestor set of z w.r.t. the transition function g , where we define $g^0(z) = z$. If $(z, w) \in \gamma_{\mathcal{T}}(\theta, \eta)$ for some $\theta \neq \eta$, according to our construction of \mathcal{T} , it is straightforward to verify that $\theta \in \Lambda(z)$. Therefore, $\{(\theta, \eta) : (z, w) \in \gamma_{\mathcal{T}}(\theta, \eta)\} \subseteq \Lambda(z) \times \Theta$. It follows that

$$\begin{aligned} \frac{1}{\rho(e)} \sum_{(\theta, \eta) : e \in \gamma_{\mathcal{T}}(\theta, \eta)} \pi(\theta)\pi(\eta) &\leq \frac{1}{\rho(e)} \sum_{(\theta, \eta) \in \Lambda(z) \times \Theta} \pi(\theta)\pi(\eta) \\ &= \frac{1}{\rho(e)} \sum_{\theta \in \Lambda(z)} \pi(\theta) \sum_{\eta \in \Theta} \pi(\eta) \\ &= \frac{\pi(\Lambda(z))}{\pi(z)\mathbf{P}(z, w)}. \end{aligned} \quad (\text{B.3})$$

Analogously to the proof of part (ii), we can write $\Lambda(z) = \bigcup_{k \in \mathbb{N}} g^{-k}(z)$, and by Condition 1, we have $|g^{-k}(z)| \leq p^{kt_1}$ and $\pi(\theta)/\pi(z) \leq p^{-kt_2}$ for any $\theta \in g^{-k}(z)$. Since $t_2 > t_1$,

$$\frac{\pi(\Lambda(z))}{\pi(z)} = \sum_{k \in \mathbb{N}} \frac{\pi(g^{-k}(z))}{\pi(z)} \leq \sum_{k \in \mathbb{N}} p^{-k(t_2 - t_1)} = \frac{1}{1 - p^{-(t_2 - t_1)}}. \quad (\text{B.4})$$

Combining (B.2), (B.3) and (B.4), we get

$$\frac{1}{\text{Gap}(\mathbf{P})} \leq \frac{2\ell_{\max}}{\{1 - p^{-(t_2 - t_1)}\} \min_{\theta \neq \theta^*} \mathbf{P}(\theta, g(\theta))}.$$

The proof is then completed by invoking (B.1). \square

Proof of Theorem 1(iv). By definition of \mathbf{P}^h , if $h \equiv 1$, we have $\mathbf{K}^h(\theta, g(\theta)) = 1/|\mathcal{N}(\theta)|$. Using Condition 1, we find that

$$\begin{aligned} \mathbf{P}^h(\theta, g(\theta)) &= \frac{1}{|\mathcal{N}(\theta)|} \min \left\{ 1, \frac{\pi(g(\theta))}{\pi(\theta)} \frac{1/|\mathcal{N}(g(\theta))|}{1/|\mathcal{N}(\theta)|} \right\} \\ &\geq \frac{1}{|\mathcal{N}(\theta)|} \min \{1, p^{t_2 - t_1}\} = \frac{1}{|\mathcal{N}(\theta)|} \geq p^{-t_1}, \end{aligned}$$

which yields the claim. \square

B.5 A general method for constructing flows

We propose a general method for constructing a flow ϕ , which is motivated by Condition 1. We say $f: \Gamma(\mathcal{N}) \rightarrow [0, 1]$ is a unit flow if $\sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N})} f(\gamma) = 1$ for any $\theta \neq \eta$. To construct a flow ϕ as described in Theorem B4, it suffices to find a unit flow f first and then let $\phi(\gamma) = f(\gamma)\pi(\theta)\pi(\eta)$ for each $\gamma \in \Gamma_{\theta\eta}(\mathcal{N})$. Let $R > 1$ be a fixed constant and define

$$\mathcal{N}_R(\theta) = \left\{ \theta' \in \mathcal{N}(\theta) : \frac{\pi(\theta')}{\pi(\theta)} \geq R \right\}, \quad \text{for } \theta \neq \theta^*,$$

$$\text{Edge}(\mathcal{N}_R) = \{(\theta, \eta) : \theta \neq \theta^*, \eta \in \mathcal{N}_R(\theta)\}.$$

Assume that there exists some $\theta^* \in \Theta$ such that $\mathcal{N}_R(\theta)$ is not empty for any $\theta \neq \theta^*$ (this would be true if Condition 1 holds); note \mathcal{N}_R is “asymmetric”. Now we construct a unit flow f_R by only using edges in $\text{Edge}(\mathcal{N}_R)$ and their reversals.

Let \mathbf{P} be given by Assumption B.1, and define another transition matrix \mathbf{P}_R by letting $\mathbf{P}_R(\theta^*, \theta^*) = 1$ and

$$\mathbf{P}_R(\theta, \theta') = \frac{\mathbf{P}(\theta, \theta')}{\mathbf{P}(\theta, \mathcal{N}_R(\theta))} \mathbb{1}_{\mathcal{N}_R(\theta)}(\theta'), \quad \text{for any } \theta \neq \theta^*,$$

Clearly, \mathbf{P}_R has one (and only one) absorbing state, θ^* . Thus, for any $\theta \neq \theta^*$, we can construct a unit flow from θ to θ^* by running the Markov chain \mathbf{P}_R with initial state θ ; the weight of a path from θ to θ^* is simply given by the probability of the chain \mathbf{P}_R moving along the path. Reversing the paths from θ to θ^* and keeping the weights unchanged, we obtain a unit flow from θ^* to θ . To construct a unit flow from θ to η for some distinct $\theta, \eta \in \Theta \setminus \{\theta^*\}$, we treat θ^* as the “hub” and concatenate each path of \mathbf{P}_R from θ to θ^* and each path (after reversal) of \mathbf{P}_R from η to θ^* ; that is, we only consider paths in $\Gamma_{\theta\eta}(\mathcal{N})$ that visit θ^* once. A formal definition of f_R is given below, followed by a toy example illustrating the construction of f_R . We prove a useful inequality for the “load” of an edge in f_R in Lemma B2.

Definition B4. For $k \geq 1$ and $\gamma = (\theta_0, \theta_1, \dots, \theta_k) \in \Gamma(\mathcal{N})$, define $f_R(\gamma)$ as follows.

- (i) If θ^* does not occur in γ or θ^* occurs at least twice in γ , let $f_R(\gamma) = 0$.
- (ii) If $\theta_k = \theta^*$, let $f_R(\gamma) = \prod_{i=1}^k \mathbf{P}_R(\theta_{i-1}, \theta_i)$.
- (iii) If $\theta_0 = \theta^*$, let $f_R(\gamma) = f_R(\tilde{\gamma})$.
- (iv) If $\theta_j = \theta^*$ for some $1 \leq j \leq k-1$, let $\gamma_1 = (\theta_0, \dots, \theta_{j-1}, \theta^*)$ and $\gamma_2 = (\theta^*, \theta_{j+1}, \dots, \theta_k)$ and define $f_R(\gamma) = f_R(\gamma_1)f_R(\gamma_2)$.

Example B1. Let $\Theta = \{\theta_1, \dots, \theta_9, \theta^*\}$ and \mathbf{P}_R be given by Figure B1. Then, from θ_1 to θ^* there are three paths, and the weights are given by

$$f_R((\theta_1, \theta_2, \theta_4, \theta^*)) = \frac{1}{3}, \quad f_R((\theta_1, \theta_2, \theta_5, \theta^*)) = \frac{1}{6}, \quad f_R((\theta_1, \theta_3, \theta^*)) = \frac{1}{2}.$$

From θ_3 to θ^* there is only one path (θ_3, θ^*) . Hence, from θ_1 to θ_3 there are $3 \times 1 = 3$ paths; for example, the weight of the path $\gamma = (\theta_1, \theta_3, \theta^*, \theta_3)$ is given by

$$f_R(\gamma) = f_R((\theta_1, \theta_3, \theta^*))f_R((\theta^*, \theta_3)) = 1/2.$$




Figure B1: An example of the Markov chain \mathbf{P}_R . If $\mathbf{P}_R(\theta, \theta') > 0$, then we draw an arrow from θ to θ' and denote the value of $\mathbf{P}_R(\theta, \theta')$ near the arrow.

Note that γ does not contain repeated edges, though the state θ_3 occur twice. It is also clear that $\mathbf{P}_R^3(\theta, \theta^*) = 1$ for any $\theta \in \Theta$; that is, from any θ we can move to θ^* in at most 3 steps.

Lemma B2. *Let $\theta^* \in \Theta$ and assume $\mathcal{N}_R(\theta)$ is non-empty for every $\theta \neq \theta^*$. The function $f_R: \Gamma(\mathcal{N}) \rightarrow [0, 1]$ constructed in Definition B4 is a unit flow. Further, for any $e = (z, w)$ such that $e \in \text{Edge}(\mathcal{N}_R)$ or $\bar{e} \in \text{Edge}(\mathcal{N}_R)$, we have*

$$\sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} f_R(\gamma) \leq \mathbf{P}_R(z, w), \quad \forall \theta, \eta \in \Theta, \theta \neq \eta.$$

Proof. Since the Markov chain \mathbf{P}_R can only stay at the state θ^* , we can use

$$\Gamma(\mathcal{N}_R) = \{\gamma = (\theta_0, \theta_1, \dots, \theta_k): k \geq 1, \theta_i \in \mathcal{N}_R(\theta_{i-1}) \text{ for } i = 1, \dots, k\} \quad (\text{B.5})$$

to denote all possible paths of \mathbf{P}_R . Let $\Gamma(\theta, \eta; \mathcal{N}_R)$ denote the paths in $\Gamma(\mathcal{N}_R)$ that start at θ and end at η . Fix some $e = (z, w) \in \text{Edge}(\mathcal{N}_R)$. We prove that

$$\sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N})} f_R(\gamma) = 1, \quad \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} f_R(\gamma) \leq \mathbf{P}_R(z, w)$$

by considering three cases for the choice of (θ, η) . For $\bar{e} = (w, z)$, the second inequality can be proved by a symmetry argument.

First, let $\theta \neq \theta^*$ and $\eta = \theta^*$. It suffices to consider paths in $\Gamma(\theta, \theta^*; \mathcal{N}_R)$ since any other γ has $f_R(\gamma) = 0$. Fix an arbitrary integer $j \geq |\Theta|$. Observe that $\mathbf{P}_R^j(\theta, \theta^*) = 1$. Hence, by our definition of f_R , $\sum_{\gamma \in \Gamma(\theta, \theta^*; \mathcal{N}_R)} f_R(\gamma)$ is just the sum of the probabilities of all j -step paths of the Markov chain \mathbf{P}_R with initial state θ , which must be one. Similarly, $\sum_{\gamma \in \Gamma(\theta, \theta^*; \mathcal{N}): e \in \gamma} f_R(\gamma)$ is the sum of the probabilities of all j -step paths that traverse the edge e . Hence, it is less than or equal to $\mathbf{P}_R(z, w)$, where the equality is attained if all j -step paths visit the state z .

Second, let $\theta = \theta^*$ and $\eta \neq \theta^*$. By our definition of f_R , we have

$$\sum_{\gamma \in \Gamma(\theta^*, \eta; \mathcal{N})} f_R(\gamma) = \sum_{\gamma \in \Gamma(\theta^*, \eta; \mathcal{N})} f_R(\tilde{\gamma}) = \sum_{\gamma \in \Gamma(\eta, \theta^*; \mathcal{N})} f_R(\gamma) = 1.$$

The edge e is not traversed by any γ if $\tilde{\gamma} \in \Gamma(\eta, \theta^*; \mathcal{N})$.

Third, consider distinct $\theta, \eta \in \Theta \setminus \{\theta^*\}$. For any $\gamma \in \Gamma_{\theta\eta}(\mathcal{N})$, note that $f_R(\gamma) = 0$ unless we can write $\gamma = \gamma_1\gamma_2$ for some $\gamma_1 \in \Gamma(\theta, \theta^*; \mathcal{N}_R)$ and $\gamma_2 \in \Gamma(\theta^*, \eta; \mathcal{N}_R)$. Therefore,

$$\begin{aligned} \sum_{\gamma \in \Gamma(\theta, \eta; \mathcal{N})} f_R(\gamma) &= \sum_{\gamma_1\gamma_2: \gamma_1 \in \Gamma(\theta, \theta^*; \mathcal{N}_R), \gamma_2 \in \Gamma(\theta^*, \eta; \mathcal{N}_R)} f_R(\gamma) \\ &= \sum_{\gamma_1 \in \Gamma(\theta, \theta^*; \mathcal{N}_R)} \sum_{\gamma_2 \in \Gamma(\theta^*, \eta; \mathcal{N}_R)} f_R(\gamma_1)f_R(\gamma_2) = 1. \end{aligned}$$

The edge e can only be traversed by the γ_1 segment of γ , and thus

$$\sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} f_R(\gamma_1) \leq \sum_{\gamma_1 \in \Gamma(\theta, \theta^*; \mathcal{N}_R): e \in \gamma} f_R(\gamma) \leq \mathbf{P}_R(z, w),$$

by what we have proved for the first case. \square

B.6 Proof of Theorem 2

Proof. We prove the claim for any Markov chain \mathbf{P} that satisfies Assumption B.1. Let $R = p^{t_2}$. By Condition 1, $\mathcal{N}_R(\theta)$ is non-empty for every $\theta \neq \theta^*$. So, we can use the unit flow f_R constructed in Definition B.4 to define a flow ϕ by letting $\phi(\gamma) = f_R(\gamma)\pi(\theta)\pi(\eta)$ for $\gamma \in \Gamma_{\theta\eta}(\mathcal{N})$. To apply Theorem B.4, it only remains to specify the length function. Observe that $\sum_{\gamma \in \Gamma(\mathcal{N}): e \in \gamma} f_R(\gamma) > 0$ if and only if $e \in \text{Edge}(\mathcal{N}_R)$ or $\tilde{e} \in \text{Edge}(\mathcal{N}_R)$. Define

$$\ell(e) = \ell(\tilde{e}) = \pi(z)^{-q}, \quad \text{if } e = (z, w), w \in \mathcal{N}_R(z),$$

for some $q \in (0, 1)$. If $\gamma = (\theta_0 = \theta, \theta_1, \dots, \theta_k = \theta^*) \in \Gamma(\theta, \theta^*; \mathcal{N}_R)$, then

$$|\gamma|_\ell = \sum_{i=0}^{k-1} \pi(\theta_i)^{-q} \leq \sum_{i=0}^{k-1} \pi(\theta)^{-q} p^{-it_2q} \leq \frac{\pi(\theta)^{-q}}{1 - p^{-t_2q}}.$$

For any $\gamma \in \Gamma_{\theta\eta}(\mathcal{N})$ such that $f_R(\gamma) > 0$, we can write $\gamma = \gamma_1\gamma_2$ where $\gamma_1 \in \Gamma(\theta, \theta^*; \mathcal{N}_R)$ and $\tilde{\gamma}_2 \in \Gamma(\eta, \theta^*; \mathcal{N}_R)$, which yields

$$|\gamma|_\ell = |\gamma_1|_\ell + |\gamma_2|_\ell \leq \frac{\pi(\theta)^{-q} + \pi(\eta)^{-q}}{1 - p^{-t_2q}}.$$

Fix some $e = (z, w) \in \text{Edge}(\mathcal{N}_R)$. (By symmetry, the case $\tilde{e} \in \text{Edge}(\mathcal{N}_R)$ can be analyzed by the same argument.) By Lemma B.2 and the above bound on $|\gamma|_\ell$, we obtain that

$$\begin{aligned} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_\ell &= \pi(\theta)\pi(\eta) \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} f_R(\gamma)|\gamma|_\ell \\ &\leq \frac{\pi(\theta)^{1-q}\pi(\eta) + \pi(\theta)\pi(\eta)^{1-q}}{1 - p^{-t_2q}} \mathbf{P}_R(z, w). \end{aligned}$$

Recall $\Gamma(\theta, \eta; \mathcal{N}_R)$ defined in (B.5). Let

$$\Lambda(z) = \{\theta \in \Theta: \Gamma(\theta, z; \mathcal{N}_R) \neq \emptyset\} \cup \{z\}.$$

Note that if a path $\gamma = (\theta_0 = \theta, \theta_1, \dots, \theta_k)$ with $f_R(\gamma) \geq 0$ traverses the edge e , we must have $\theta \in \Lambda(z)$. Hence,

$$\begin{aligned} \sum_{\theta \neq \eta} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_\ell &\leq \sum_{\theta \in \Lambda(z)} \sum_{\eta \in \Theta} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_\ell \\ &\leq \frac{\mathbf{P}_R(z, w)}{1 - p^{-t_2q}} \sum_{\theta \in \Lambda(z)} \sum_{\eta \in \Theta} [\pi(\theta)^{1-q}\pi(\eta) + \pi(\theta)\pi(\eta)^{1-q}]. \end{aligned} \quad (\text{B.6})$$

Since \mathcal{N} is symmetric, for any θ ,

$$|\{\eta \in \Theta: \theta \in \mathcal{N}_R(\eta)\}| \leq |\mathcal{N}(\theta)| \leq p^{t_1}.$$

Applying this argument recursively, we find that for any $z \neq \theta^*$,

$$\sum_{\theta \in \Lambda(z)} \pi(\theta)^c \leq \sum_{k=0}^{\infty} \pi(z)^c p^{(t_1-t_2c)k} \leq \frac{\pi(z)^c}{1 - p^{t_1-t_2c}},$$

for any $c > 0$ such that the series converges. Similarly,

$$\sum_{\eta \in \Theta} \pi(\eta)^c = \sum_{\eta \in \Lambda(\theta^*)} \pi(\eta)^c \leq \frac{\pi(\theta^*)^c}{1 - p^{t_1-t_2c}} \leq \frac{1}{1 - p^{t_1-t_2c}}.$$

Plugging these bounds into (B.6), we obtain that

$$\begin{aligned} \sum_{\theta \neq \eta} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_\ell &\leq \frac{\mathbf{P}_R(z, w)}{1 - p^{-t_2q}} \left\{ \sum_{\theta \in \Lambda(z)} \pi(\theta)^{1-q} + \sum_{\theta \in \Lambda(z)} \sum_{\eta \in \Theta} \pi(\theta)\pi(\eta)^{1-q} \right\} \\ &\leq \frac{\mathbf{P}_R(z, w)}{1 - p^{-t_2q}} \left\{ \frac{\pi(z)^{1-q}}{1 - p^{t_1-t_2(1-q)}} + \frac{\pi(z)}{(1 - p^{t_1-t_2})(1 - p^{t_1-t_2(1-q)})} \right\} \\ &\leq \frac{\mathbf{P}_R(z, w)}{1 - p^{-t_2q}} \frac{\pi(z)^{1-q}}{1 - p^{t_1-t_2(1-q)}} \left(1 + \frac{1}{1 - p^{t_1-t_2}} \right). \end{aligned}$$

Choose $q = (t_2 - t_1)/(2t_2)$ such that

$$t_1 - t_2(1 - q) = \frac{t_1 - t_2}{2} = -t_2q.$$

Using $\mathbf{P}_R(z, w) = \mathbf{P}(z, w)/\mathbf{P}(z, \mathcal{N}_R(z))$ and $\rho(e)\ell(e) = \pi(z)^{1-q}\mathbf{P}(z, w)$, we find that

$$\frac{1}{\rho(e)\ell(e)} \sum_{\theta \neq \eta} \sum_{\gamma \in \Gamma_{\theta\eta}(\mathcal{N}): e \in \gamma} \phi(\gamma)|\gamma|_\ell \leq \frac{1 + (1 - p^{t_1-t_2})^{-1}}{[1 - p^{(t_1-t_2)/2}]^2} \mathbf{P}(z, \mathcal{N}_R(z))^{-1}.$$

The proof is then completed by applying (B.1) and Theorem B4. We have the coefficient 2 on the numerator of the bound given in the theorem because we consider the lazy version. \square

C Preliminaries for graphical models

C.1 Markov equivalence and independence maps

Below are some useful results for checking whether one DAG is Markov equivalent or an I-map of another DAG. The skeleton of a DAG is the unique undirected graph obtained by replacing all edges in the DAG with undirected ones. A v-structure is a triple $i \rightarrow j \leftarrow k$ (note that there is no edge between i and k). By an edge reversal, we mean to change an existing edge $i \rightarrow j$ to $j \rightarrow i$. We say the edge $i \rightarrow j$ is covered if and only if $\text{Pa}_i = \text{Pa}_j \setminus \{i\}$. Two nodes i, j are said to be adjacent in G if $i \rightarrow j \in G$ or $j \rightarrow i \in G$.

Lemma C1. *Two DAGs are Markov equivalent if and only if they have the same skeleton and v-structures.*

Proof. See Verma and Pearl [73]. □

Lemma C2. *Two DAGs G_1, G_2 are Markov equivalent if and only if we can transform G_1 to G_2 by a sequence of covered edge reversals.*

Proof. See Chickering [13, Theorem 2]. □

Lemma C3. *Let the DAG H be an I-map of the DAG $G \neq H$. We can create a DAG G' from G by at least one of the following two operations such that H is an I-map of G' .*

(i) *Reverse a covered edge $i \rightarrow j$ in G such that $j \rightarrow i \in H$.*

(ii) *Add an edge $i \rightarrow j$ to G such that i, j are adjacent in H .*

Consequently, there exists a sequence of DAGs $(G_0 = G, G_1, \dots, G_{m-1}, G_m = H)$ such that for each $i = 1, \dots, m$, H is an I-map of G_i , and G_i is obtained from G_{i-1} by either a covered edge reversal or a single-edge addition.

Proof. See Theorem 4 and the remark following it in Chickering [15]. □

C.2 Gaussian DAG models

We provide some background on the decomposition of $\Sigma(B, \Omega)$ given in (12). It is called the modified Cholesky decomposition of Σ (up to permutation of rows and columns), and B is the modified Cholesky factor (after scaling). Let $\mathcal{D}_p(\sigma) = \bigcup_{G \in \mathcal{G}_p^\sigma} \mathcal{D}_p(G)$ where $\mathcal{D}_p(G)$ is as given in (11). Observe that, equivalently, $\mathcal{D}_p(\sigma)$ can be defined by

$$\begin{aligned} \mathcal{D}_p(\sigma) = \{ (B, \Omega) : B \in \mathbb{R}^{p \times p}, B_{ij} = 0 \text{ if } \sigma^{-1}(i) \geq \sigma^{-1}(j), \text{ for any } i, j \in [p]; \\ \Omega = \text{diag}(\omega_1, \dots, \omega_p), \omega_i > 0 \text{ for any } i \in [p] \}. \end{aligned} \quad (\text{C.1})$$

Lemma C4. *For any positive definite matrix $\Sigma \in \mathbb{R}^{p \times p}$ and $\sigma \in \mathbb{S}^p$, the decomposition $\Sigma = (I - B^\top)^{-1} \Omega (I - B)^{-1}$ for $(B, \Omega) \in \mathcal{D}_p(\sigma)$ exists and is unique.*

Proof. First, permute the rows and columns of Σ using σ so that B becomes upper triangular after permutation. The result then follows from the existence and uniqueness of Cholesky decomposition for positive definite matrices. See also Aragam et al. [5, Lemma 2.1]. □

Lemma C5. Let $\Sigma = (I - B^\top)^{-1}\Omega(I - B)^{-1}$ for some $(B, \Omega) \in \cup_{\sigma \in \mathbb{S}^p} \mathcal{D}_p(\sigma)$. Let G be a DAG such that $i \rightarrow j \in G$ if and only if $B_{ij} > 0$. Then G is a minimal I-map of $N_p(0, \Sigma)$; that is, $N_p(0, \Sigma)$ is Markovian w.r.t. G but not Markovian w.r.t. any sub-DAG of G .

Proof. See Peters and Bühlmann [57, Theorem 1]. \square

Lemma C6. Let $N_p(0, \Sigma)$ be a non-degenerate multivariate normal distribution Markovian w.r.t. a p -vertex DAG G . Then the decomposition $\Sigma = (I - B^\top)^{-1}\Omega(I - B)^{-1}$ for $(B, \Omega) \in \mathcal{D}_p(G)$ exists and is unique.

Proof. Let $\mathbf{X} = (X_1, \dots, X_p) \sim N_p(0, \Sigma)$. The Markovian assumption implies that the density of \mathbf{X} factorizes according to G . Further, for any $S \subseteq [p] \setminus \{j\}$, $X_j \mid \mathbf{X}_S$ follows a normal distribution and the conditional expectation is linear in \mathbf{X}_S , which shows that $(B, \Omega) \in \mathcal{D}_p(G)$ exists. Let σ be a topological ordering of G . Since $\mathcal{D}_p(G) \subseteq \mathcal{D}_p(\sigma)$, the uniqueness follows from Lemma C4. \square

Lemma C7. Let G be a p -vertex DAG and with edge set E . Let B be a $p \times p$ matrix such that $B_E = \{B_{ij} : (i, j) \in E\}$ is sampled from an absolutely continuous distribution on $\mathbb{R}^{|E|}$ and other entries of B are zeroes. Let $\Omega = \text{diag}(\omega_1, \dots, \omega_p)$ be a diagonal matrix where $(\omega_1, \dots, \omega_p)$ is sampled from an absolutely continuous distribution on $(0, \infty)^p$. Let $\Sigma = (I - B^\top)^{-1}\Omega(I - B)^{-1}$. Then $N_p(0, \Sigma)$ is perfectly Markovian w.r.t. G almost surely.

Proof. See Spirtes et al. [66, Theorem 3.2]. \square

Example C1. We give an example for minimal I-maps of Gaussian DAG models. Let Z_1, Z_2, Z_3 be independent standard normal random variables. Define

$$\mathbf{X}_1 = Z_1, \quad \mathbf{X}_2 = b_1 X_1 + Z_2, \quad \mathbf{X}_3 = b_2 X_2 + Z_3. \quad (\text{C.2})$$

The DAG model G corresponding to this SEM is $1 \rightarrow 2 \rightarrow 3$, which only has one ordering $\sigma = c(1, 2, 3)$. Denote the weighted adjacency matrix of G by B , which has $B_{12} = b_1$ and $B_{23} = b_2$ (and all other entries are zero). Normality of (Z_1, Z_2, Z_3) implies that $\mathbf{X} = (X_1, X_2, X_3) \sim N_3(0, \Sigma)$ where

$$\Sigma = \begin{bmatrix} 1 & b_1 & b_1 b_2 \\ b_1 & b_1^2 + 1 & (b_1^2 + 1)b_2 \\ b_1 b_2 & (b_1^2 + 1)b_2 & b_1^2 b_2^2 + b_2^2 + 1 \end{bmatrix} = (I - B^\top)^{-1}(I - B)^{-1}.$$

A routine calculation shows that (C.2) can be equivalently written as

$$\begin{aligned} \mathbf{X}_1 &= Z_1, \\ \mathbf{X}_3 &= b_1 b_2 \mathbf{X}_1 + b_2 Z_2 + Z_3, \\ \mathbf{X}_2 &= \frac{b_1}{b_2^2 + 1} \mathbf{X}_1 + \frac{b_2}{b_2^2 + 1} \mathbf{X}_3 + \frac{1}{b_2^2 + 1} Z_2 - \frac{b_2}{b_2^2 + 1} Z_3. \end{aligned} \quad (\text{C.3})$$

This SEM clearly corresponds to the ordering $\tau = (1, 3, 2)$, from which one obtains the modified Cholesky decomposition factor $(B_\sigma, \Omega_\sigma)$; that is, $\Sigma = (I - B_\sigma^\top)^{-1}\Omega_\sigma(I - B_\sigma)^{-1}$

where

$$B_\sigma = \begin{bmatrix} 0 & b_1/(b_2^2 + 1) & b_1 b_2 \\ 0 & 0 & 0 \\ 0 & b_2/(b_2^2 + 1) & 0 \end{bmatrix}, \quad \Omega_\sigma = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1/(b_2^2 + 1) & 0 \\ 0 & 0 & b_2^2 + 1 \end{bmatrix}.$$

In particular, the i -th diagonal element of Ω_σ is the error variance of \mathbf{X}_i in the SEM (C.3).

C.3 CPDAGs and CPDAG operators

A partially directed acyclic graph (PDAG) contains directed and/or undirected edges but no directed cycles (a cycle is directed if it has at least one directed edge). For a PDAG H , we use $\text{Adj}(H)$ to denote the set of all nodes that are connected to node j by either a directed or undirected edge, in which case we say the two nodes are adjacent, and $\text{Un}_j(H) = \text{Adj}(H) \setminus (\text{Pa}_j(H) \cup \text{Ch}_j(H))$ to denote the set of nodes that are connected to j by an undirected edge.

For an equivalence class \mathcal{E} , its CPDAG or essential graph, denoted by $\text{EG}(\mathcal{E})$, is the unique PDAG with the same skeleton as any $G \in \mathcal{E}$ such that an edge in $\text{EG}(\mathcal{E})$ is directed if and only if the edge is directed in the same orientation in every $G \in \mathcal{E}$. For convenience, we will also denote $\text{EG}(\mathcal{E})$ by $\text{EG}(G)$ for any $G \in \mathcal{E}$. Given a CPDAG $\text{EG}(\mathcal{E})$, a CPDAG operator defines some specific modification of $\text{EG}(\mathcal{E})$, such as adding or deleting edges. After we apply such an operator to $\text{EG}(\mathcal{E})$, we may get a PDAG that is not a CPDAG, but it may be converted to a CPDAG in a unique way.

Definition C1 (consistent extension). A DAG G is a consistent extension of a PDAG H if (i) G and H have the same skeleton and v-structures, and (ii) each directed edge in H has the same orientation as in G .

Definition C2. A CPDAG operator is valid if the modified graph is a PDAG and has a consistent extension.

Note that He et al. [28] used a slight different definition of “valid” operators: they further required that a valid operator must also be “effective”, which means that all modified edges occur in the resulting CPDAG [28, Definition 3]. By Definition C1, all consistent extensions of a given PDAG are Markov equivalent. Hence, a valid CPDAG operator always results in a unique CPDAG (i.e., a unique equivalence class).

He et al. [28] considered the following six types of CPDAG operators: insert/delete an undirected edge, insert/delete a direct edge, and make/remove a v-structure. “Make a v-structure” means to convert a subgraph $i - j - k$ in the CPDAG to $i \rightarrow j \leftarrow k$; similarly, “remove a v-structure” means to convert a v-structure $i \rightarrow j \leftarrow k$ to $i - j - k$. However, there exist CPDAGs $\text{EG}(\mathcal{E})$ and $\text{EG}(\mathcal{E}')$ such that we can get $\text{EG}(\mathcal{E}')$ from $\text{EG}(\mathcal{E})$ by one of these operators, but not vice versa. To overcome this problem, He et al. [28, Definition 9] constructed a set of rules defining, for each CPDAG, which of the above six operators are allowed. The resulting set of operators is both “irreducible” and “reversible” on the unrestricted CPDAG space.

Chickering [15] introduced two types of CPDAG operators for implementing the GES algorithm, *Insert* and *Delete*, which can be defined as follows.

Definition C3 ($\text{Insert}(i, j, S)$). Given a CPDAG H , “ $\text{Insert}(i, j, S)$ ” operator is defined for non-adjacent nodes i, j and a set $S \subseteq \text{Un}_j(H) \setminus \text{Ad}_i(H)$. It modifies H by (i) inserting the edge $i \rightarrow j$, and (ii) for each $k \in S$, directing the edge between k and j as $k \rightarrow j$.

Definition C4 ($\text{Delete}(i, j, S)$). Given a CPDAG H , “ $\text{Delete}(i, j, S)$ ” operator is defined for adjacent nodes i, j connected as either $i - j$ or $i \rightarrow j$ and a set $S \subseteq \text{Un}_j(H) \cap \text{Ad}_i(H)$. It modifies H by (i) deleting the edge between i and j , and (ii) for each $k \in S$, directing the edge between k and j as $j \rightarrow k$ and any undirected edge between k and i as $i \rightarrow k$.

There are easily testable conditions for checking whether some *Insert* or *Delete* operator is valid; see Chickering [15, Theorems 15 and 17]. Given a CPDAG H with p nodes, let $\mathcal{O}_{\text{ges}}(H)$ denote the set of all *Insert* and *Delete* operators defined for H (which may or may not be valid). Let d denote the degree of H . Then, we always have $|\mathcal{O}_{\text{ges}}(H)| \leq p^2 2^d$. Further, the time complexity of converting the resulting PDAG to CPDAG is $O(pd^3)$ [15]. For each $k \in S$, the operator $\text{Insert}(i, j, S)$ creates a new v-structure $i \rightarrow j \leftarrow k$. Similarly, for each $k \in S$, the operator $\text{Delete}(i, j, S)$ creates a new v-structure $i \rightarrow k \leftarrow j$. Hence, given a CPDAG H , the only *Insert* operators that are not distinguishable (i.e., two operators result in the same CPDAG) are $\text{Insert}(i, j, \emptyset)$ and $\text{Insert}(j, i, \emptyset)$ when $\text{Pa}_i(H) = \text{Pa}_j(H)$ by Chickering [14, Theorem 34], and the only *Delete* operators that are not distinguishable are $\text{Delete}(i, j, S)$ and $\text{Delete}(j, i, S)$ when i, j are connected by an undirected edge in H . This observation reveals that the number of operators in $\mathcal{O}_{\text{ges}}(H_1)$ that can convert a CPDAG H_1 to another CPDAG H_2 is always the same as the number of operators in $\mathcal{O}_{\text{ges}}(H_2)$ that can convert H_2 to H_1 . To see this, fix an arbitrary CPDAG H_2 which can be obtained by applying $\text{Insert}(i, j, S)$ to some CPDAG H_1 (which implies i, j are not adjacent in H_1). If the edge $i \rightarrow j$ is directed in H_2 , then there is only one *Delete* operator that can transform H_2 back to H_1 . If i, j are connected by an undirected edge in H_2 , it implies that $S = \emptyset$ and $\text{Pa}_i(H_1) = \text{Pa}_j(H_1)$ by Chickering [14, Theorem 34]. Thus, both $\text{Insert}(i, j, \emptyset)$ and $\text{Insert}(j, i, \emptyset)$ can convert H_1 to H_2 , and there are two *Delete* operators that can transform H_2 back to H_1 .

D Proofs for Section 3

D.1 Proof of Lemma 1

We need two auxiliary lemmas.

Lemma D1. *Let G_1 and G_2 be two Markov equivalent DAGs such that $\text{Pa}_j(G_1) = \text{Pa}_j(G_2)$ and $i \notin \text{Pa}_j(G_1)$. Suppose that $G'_1 = G_1 \cup \{i \rightarrow j\}$ and $G'_2 = G_2 \cup \{i \rightarrow j\}$ are both DAGs. Then G'_1, G'_2 are Markov equivalent.*

Proof. By Lemma C1, G'_1 and G'_2 have the same skeleton. It suffices to show that G'_1 and G'_2 share the same v-structures. One can see that a v-structure remains unchanged unless it involves both i and j . There are only two cases where adding the edge $i \rightarrow j$ affects some v-structure:

1. For some node k , a new v-structure $i \rightarrow j \leftarrow k$ is formed.
2. For some node k , an existing v-structure $i \rightarrow k \leftarrow j$ is shielded.

We show that if Case 1 or Case 2 happens in G'_1 , it also happens in G'_2 . If $i \rightarrow j \leftarrow k$ is a v-structure in G'_1 for $k \in \text{Pa}_j(G_1)$, i, k are not adjacent in G'_1 (“adjacent” means two nodes are connected by an edge). Then i, k must be non-adjacent in G'_2 as well since G'_1 and G'_2 have the same skeleton. By the assumption $\text{Pa}_j(G_1) = \text{Pa}_j(G_2)$, we find that $k \in \text{Pa}_j(G_2)$ and thus G'_2 contains the v-structure $i \rightarrow j \leftarrow k$. If G_1 has a v-structure $i \rightarrow k \leftarrow j$ for some node k , so does G_2 by Lemma C1. Adding $i \rightarrow j$ shields the v-structure both in G'_1 and G'_2 . \square

Lemma D2. *Let G_1 and G_2 be two Markov equivalent DAGs such that $\text{Pa}_j(G_1) = \text{Pa}_j(G_2)$ and $i \in \text{Pa}_j(G_1)$. Let $G'_1 = G_1 \setminus \{i \rightarrow j\}$ and $G'_2 = G_2 \setminus \{i \rightarrow j\}$. Then G'_1, G'_2 are Markov equivalent.*

Proof. The proof is similar to that of Lemma D1. we show that G'_1 and G'_2 share the same v-structures. There are only two cases where deleting the edge $i \rightarrow j$ affects the v-structure.

1. For some node k , a new v-structure $i \rightarrow k \leftarrow j$ is formed.
2. For some node k , an existing v-structure $i \rightarrow j \leftarrow k$ is broken up.

Let $i \rightarrow k \leftarrow j$ be a v-structure in G'_1 . Observe that the assumptions $\text{Pa}_j(G_1) = \text{Pa}_j(G_2)$ and that G_1 and G_2 are Markov equivalent imply $\text{Ch}_j(G_1) = \text{Ch}_j(G_2)$. Since $k \in \text{Ch}_j(G_1)$, G'_2 also has the edge $k \leftarrow j$. Further, we must have $i \rightarrow k \in G'_2$, since G_2 is acyclic. This shows that $i \rightarrow k \leftarrow j$ is also a v-structure in G'_2 . If G_1 has a v-structure $i \rightarrow j \leftarrow k$ for some node k , so does G_2 . Removing $i \rightarrow j$ destroys the v-structure in both G'_1 and G'_2 . \square

Proof of Lemma 1. Fix an arbitrary $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$. By the definition of $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, there exists some $G_0 \in \mathcal{E}$ such that $G_0 \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}})$. Since any Markov equivalent DAGs must have the same skeleton, for any $G \in \mathcal{E}$, the degree of any node is bounded by $d_{\text{in}} + d_{\text{out}}$ and the set of adjacent nodes is fixed. Define $\text{Pa}_j(\mathcal{E}) = \{\text{Pa}_j(G) : G \in \mathcal{E}\}$, the collection of all possible parent sets of node j . It then follows that $|\text{Pa}_j(\mathcal{E})| \leq 2^{d_{\text{in}} + d_{\text{out}}}$.

Consider $\mathcal{N}_{\text{add}}(\mathcal{E})$. Suppose that \mathcal{E}' is obtained by adding the edge $i \rightarrow j$ to some $G \in \mathcal{E}$. By Lemma D1, for $G_1, G_2 \in \mathcal{E}$, $G_1 \cup \{i \rightarrow j\}$ and $G_2 \cup \{i \rightarrow j\}$ are still Markov equivalent if the resulting graphs are DAGs and $\text{Pa}_j(G) = \text{Pa}_j(G')$. But $|\text{Pa}_j(\mathcal{E})| \leq 2^{d_{\text{in}}+d_{\text{out}}}$ implies that adding the edge $i \rightarrow j$ can yield at most $2^{d_{\text{in}}+d_{\text{out}}}$ different equivalence classes. Hence, $|\mathcal{N}_{\text{add}}(\mathcal{E})| \leq p(p-1)2^{d_{\text{in}}+d_{\text{out}}}$ since there are at most $p(p-1)$ directed edges we can add.

A similar argument can be applied to $\mathcal{N}_{\text{del}}(\mathcal{E})$. For any $G \in \mathcal{E}$, we have $|G| \leq pd_{\text{in}}$, but these edges may be directed in either way. Using Lemma D2, we then find that $|\mathcal{N}_{\text{del}}(\mathcal{E})| \leq 2pd_{\text{in}}2^{d_{\text{in}}+d_{\text{out}}}$. Finally, for any $G \in \mathcal{E}$, $|\mathcal{N}_{\text{swap}}(G)| \leq p(p-1)(d_{\text{in}}+d_{\text{out}})$. Hence, $|\mathcal{N}_{\text{swap}}(\mathcal{E})| \leq p(p-1)(d_{\text{in}}+d_{\text{out}})2^{d_{\text{in}}+d_{\text{out}}}$. Combining the results for the three cases, we obtain the asserted upper bound on $|\mathcal{N}_{\text{ads}}(\mathcal{E})|$. \square

D.2 Proof of Lemma 2

Proof. We prove by contradiction. Assume that for any $j \in [p]$ such that $g_j^\sigma(G) \neq G$, we have $g_j^\sigma(G) \notin \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. It follows that there exists no j such that $\text{Pa}_j(G_\sigma^*) \subsetneq \text{Pa}_j(G)$. Otherwise, $g_j^\sigma(G)$ can be obtained from G by removing some incoming edge of node j and thus $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$.

Since there is no strictly overfitted node, there must exist some underfitted node. Let

$$U = \{u \in [p] : \text{Pa}_u(G_\sigma^*) \not\subseteq \text{Pa}_u(G)\}$$

be the set of all underfitted nodes. Recall that for any $u \in U$, $g_u^\sigma(G)$ is constructed by adding an incoming edge of node u to G . Define

$$A = \{a \in [p] : g_u^\sigma(G) = G \cup \{a \rightarrow u\} \text{ for some } u \in U\}.$$

Fix an arbitrary $a \in A$ and suppose that $g_j^\sigma(G) = G \cup \{a \rightarrow j\}$ for some j . If $|\text{Ch}_a(G)| < d_{\text{out}}$, one can verify that $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ (if $\text{Pa}_j(G) < d_{\text{in}}$, we can simply add the edge $a \rightarrow j$; if $\text{Pa}_j(G) = d_{\text{in}}$, we perform a swap). Thus, $|\text{Ch}_a(G)| = d_{\text{out}}$ for every $a \in A$.

For any node $u \in U$, there exists some node $a(u) \in A$ such that $a(u) \rightarrow u$ is in G_σ^* but not in G . But any node a can have at most d_σ^* outgoing edges in G_σ^* , which implies that

$$|A| \geq \frac{|U|}{d_\sigma^*}. \quad (\text{D.1})$$

For each $a \in A$, define

$$c_a = |\{u \in U : g_u^\sigma(G) = G \cup \{a \rightarrow u\}\}|, \quad F_a = \text{Ch}_a(G) \setminus \text{Ch}_a(G_\sigma^*).$$

So c_a is the number of nodes in U which we want to connect with the parent node a . Observe that $c_a \leq |\text{Ch}_a(G_\sigma^*) \setminus \text{Ch}_a(G)|$ and $\sum_{a \in A} c_a = |U|$. Using $|\text{Ch}_a(G)| = d_{\text{out}}$ and $|\text{Ch}_a(G_\sigma^*)| \leq d_\sigma^*$, we find that $|F_a| \geq d_{\text{out}} - d_\sigma^* + c_a$. Hence,

$$\sum_{a \in A} |F_a| \geq \sum_{a \in A} (d_{\text{out}} - d_\sigma^* + c_a) \geq \frac{|U|d_{\text{out}}}{d_\sigma^*} \quad (\text{D.2})$$

where the last step follows from (D.1) and $\sum_{a \in A} c_a = |U|$.

For any node $f_a \in F_a$, the edge $a \rightarrow f_a$ is in G but not in G_σ^* . Define

$$E = \{(a, f) : a \in A, f \in F_a\}.$$

Clearly, $|E| = \sum_{a \in A} |F_a|$. Since the maximum in-degree of G is at most d_{in} , for any $f \in [p]$, we have $|\{a \in A : (a, f) \in E\}| \leq d_{\text{in}}$. Consider the set $\bar{F} = \bigcup_{a \in A} F_a$. By (D.2) we have that

$$|\bar{F}| \geq \frac{|E|}{d_{\text{in}}} \geq \frac{|U|d_{\text{out}}}{d_\sigma^* d_{\text{in}}} > |U|,$$

since we assume $d_{\text{out}} > d_{\text{in}} d_\sigma^*$. For any node $f \in \bar{F}$, we have $\text{Pa}_f(G) \neq \text{Pa}_f(G_\sigma^*)$. Because we have already shown that no node in G can be strictly overfitted, all nodes in \bar{F} must be underfitted; thus, we must have $|\bar{F}| \leq |U|$, which yields the contradiction. \square

D.3 Proof of Corollary 1

Proof. Consider part (i) first. By Lemma 2, for any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ such that $G \neq G_\sigma^*$, there exists some $j \in [p]$ such that $G' = g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ and $G' \neq G$. Define $g^\sigma(G) = g_j^\sigma(G)$. Then, by the definition of g_j^σ , we always have $\text{Hd}(g^\sigma(G), G_\sigma^*) < \text{Hd}(G, G_\sigma^*)$ and

$$\begin{aligned} \psi(g^\sigma(G)) - \psi(G) &= \psi(g_j^\sigma(G)) - \psi(G) \\ &= \sum_i \{\psi_i(g_j^\sigma(G)) - \psi_i(G)\} \\ &= \psi_j(g_j^\sigma(G)) - \psi_j(G) \\ &= \psi_j(g_j^\sigma(\text{Pa}_j(G))) - \psi_j(\text{Pa}_j(G)). \end{aligned} \tag{D.3}$$

Since $\text{Hd}(G, G_\sigma^*) \leq (d_\sigma^* + d_{\text{in}})p$ for any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, the claim then follows.

For part (ii) of Corollary 1, observe that for any $G \neq G_\sigma^*$, if (8) holds, then it follows from (D.3) that $e^{\psi(g^\sigma(G))}/e^{\psi(G)} \geq p^t$. It is also clear from construction that $g^\sigma(G) \in \mathcal{N}_{\text{ads}}(G)$. Hence, g^σ is the canonical transition function that satisfies part (ii) of Condition 1 with $t_2 = t$. It only remains to bound the neighborhood size. Clearly, $|\mathcal{N}_{\text{add}}(G) \cup \mathcal{N}_{\text{del}}(G)| \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}) \leq p(p-1)/2$ since there are only $p(p-1)/2$ directed edges compatible with ordering σ . For the swap moves, observe that if $\sigma(j) = k$, then there are at most $k-1$ incoming edges of node j we can add and d_{in} incoming edges of node j we can remove. Hence, $|\mathcal{N}_{\text{swap}}(G)| \leq \sum_{k=1}^p (k-1)d_{\text{in}} \leq d_{\text{in}}p(p-1)/2$. Without loss of generality, we can assume $d_{\text{in}} \leq p-1$, which gives

$$|\mathcal{N}_{\text{ads}}(G) \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})| \leq \frac{(d_{\text{in}}+1)p(p-1)}{2} \leq \frac{d_{\text{in}}p^2}{2}. \tag{D.4}$$

This shows that part (ii) of Condition 1 holds with $t_1 = 3$. \square

D.4 Proof of Theorem 3

Consider the claim $G_\sigma^* \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ first. Without loss of generality, assume $d_{\text{in}} \leq p, d_{\text{out}} \leq p$. If $d^* = 0$, then G^* is the empty DAG and the claim holds trivially. So, we can assume $d^* \geq 1$. But then $\min\{d^*d_{\text{in}} + 1, p\} \leq d_{\text{out}}$ implies that $d_{\text{out}} \geq d_{\text{in}} \geq d^*$. Since d^* is the maximum degree of all minimal I-maps, this shows that $G_\sigma^* \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ for each σ .

To prove the existence of function g , we first derive an auxiliary lemma.

Lemma D3. Assume $d^* \leq d_{\text{in}} \leq d_{\text{out}}$. Consider G_σ^* for some $\sigma \in \mathbb{S}^p$ such that $\mathcal{E} = [G_\sigma^*] \neq \mathcal{E}^*$. There exist $\mathcal{E}' \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, $G \in \mathcal{E}$ and $\tau \in \mathbb{S}^p$ such that (i) $\mathcal{E}' \in \mathcal{N}_{\text{del}}(\mathcal{E})$ and $\mathcal{CI}(\mathcal{E}) \subset \mathcal{CI}(\mathcal{E}') \subseteq \mathcal{CI}(\mathcal{E}^*)$, and (ii) $G \in \mathcal{G}_p^\tau(d_{\text{in}}, d_{\text{out}})$ and $\mathcal{E}' = [g_j^\tau(G)]$ for some $j \in [p]$.

Proof. The first conclusion of the lemma is essentially the same as Chickering [15, Lemma 10]. To prove it, note that since $\mathcal{E} = [G_\sigma^*]$, we have $\mathcal{CI}(\mathcal{E}) = \mathcal{CI}(G_\sigma^*) \subset \mathcal{CI}(\mathcal{E}^*)$. By Lemma C3, there exist some finite m and a sequence of DAGs, $(G_0 = G^*, G_1, \dots, G_{m-1}, G_m = G_\sigma^*)$, such that, for each $k \in [m]$, $\mathcal{CI}(G_k) \subseteq \mathcal{CI}(G_{k-1})$ and G_k is obtained from G_{k-1} by either a covered edge reversal or an edge addition. Let $\ell = \max\{j \leq m: |G_j| = |G_\sigma^*| - 1\}$, which clearly exists. Then, $G_\ell \in \mathcal{N}_{\text{del}}(G_{\ell+1})$, $G_{\ell+1} \in \mathcal{E}$ by Lemma C2, and $\mathcal{CI}(G_{\ell+1}) \subset \mathcal{CI}(G_\ell) \subseteq \mathcal{CI}(G^*)$.

Let τ be a topological ordering of $G_{\ell+1}$. For each $i \in [p]$, we have $\text{Pa}_i(G_\tau^*) \subseteq \text{Pa}_i(G_{\ell+1})$, since $G_{\ell+1}$ is an I-map of G^* but G_τ^* is the unique minimal I-map with ordering τ . Meanwhile, $\mathcal{CI}(G_\ell) \subset \mathcal{CI}(G^*)$ and $|G_\ell| < |G_{\ell+1}|$ imply that there exists some node j such that $\text{Pa}_j(G_\tau^*) \subsetneq \text{Pa}_j(G_{\ell+1})$. Consider the DAG $G' = g_j^\tau(G_{\ell+1})$, which by definition satisfies that $G' = G_{\ell+1} \setminus \{i \rightarrow j\}$ for some $i \neq \text{Pa}_j(G_\tau^*)$. Hence, $G' \in \mathcal{N}_{\text{del}}(G)$, $\mathcal{E}' = [G'] \in \mathcal{N}_{\text{del}}(\mathcal{E})$ and $\mathcal{CI}(\mathcal{E}) = \mathcal{CI}(G_{\ell+1}) \subset \mathcal{CI}(\mathcal{E}') \subseteq \mathcal{CI}(\mathcal{E}^*)$. To conclude the proof, notice that the maximum degree of $G_{\ell+1}$ is bounded by d^* since $G_{\ell+1}$ and G_σ^* are Markov equivalent. It then follows from the assumption $d^* \leq d_{\text{in}} \leq d_{\text{out}}$ that $G_{\ell+1} \in \mathcal{G}_p^\tau(d_{\text{in}}, d_{\text{out}})$ and thus $\mathcal{E}' \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$. \square

Proof of Theorem 3. We give an explicit construction of g that satisfies the required conditions. For each $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, define the “distance” from \mathcal{E} to \mathcal{E}^* by

$$\Delta^*(\mathcal{E}) = \min \left\{ \text{Hd}(G, G_\sigma^*) + |G_\sigma^*| - |G^*| : G \in \mathcal{E} \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \sigma \in \mathbb{S}^p \right\}, \quad (\text{D.5})$$

where we recall Hd denotes the Hamming distance. Note that for any σ , $|G_\sigma^*| \geq |G^*|$ since G_σ^* is an I-map of G^* ; recall Lemma C3. Thus, $\Delta^*(\mathcal{E}) = 0$ if and only if $\mathcal{E} = \mathcal{E}^*$. We will construct g on $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}})$ such that $\Delta^*(g(\mathcal{E})) < \Delta^*(\mathcal{E})$ for any $\mathcal{E} \neq \mathcal{E}^*$.

Let $(\bar{G}(\mathcal{E}), \bar{\sigma}(\mathcal{E}))$ be the pair that attains the minimum in the definition of $\Delta^*(\mathcal{E})$ (if there are multiple such pairs, fix one of them). The pair $(\bar{G}(\mathcal{E}), \bar{\sigma}(\mathcal{E}))$ can be seen as a “canonical” representation of the equivalence class \mathcal{E} . Fix an arbitrary $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, and we define $g(\mathcal{E})$ as follows.

1. Let $(G, \sigma) = (\bar{G}(\mathcal{E}), \bar{\sigma}(\mathcal{E}))$ be its canonical representation.
2. If $G \neq G_\sigma^*$, by Lemma 2, there exists some j such that $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ and $g_j^\sigma(G) \neq G$. Let $G' = g_j^\sigma(G)$ and $g(\mathcal{E}) = [G'] \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$.
3. If $G = G_\sigma^*$ but $\mathcal{E} \neq \mathcal{E}^*$, by Lemma D3, there exist some $G_0 \in \mathcal{E}, \tau \in \mathbb{S}^p, j \in [p]$ such that $G_0 \in \mathcal{G}_p^\tau(d_{\text{in}}, d_{\text{out}})$, $G' = g_j^\tau(G_0) \in \mathcal{N}_{\text{del}}(G_0)$, and G' is still an I-map of G^* . Let $g(\mathcal{E}) = [G'] \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$.
4. If $\mathcal{E} = \mathcal{E}^*$, let $g(\mathcal{E}) = \mathcal{E}$.

It follows from the definition of \mathcal{N}_{ads} that $g(\mathcal{E}) \in \mathcal{N}_{\text{ads}}(\mathcal{E})$ for every $\mathcal{E} \neq \mathcal{E}^*$. Consider $g_j^\sigma(G)$ found in step (2). By Lemma 2 and the definition of g_j^σ , we have

$$\Delta^*([g_j^\sigma(G)]) \leq \text{Hd}(g_j^\sigma(G), G_\sigma^*) + |G_\sigma^*| - |G^*| < \text{Hd}(G, G_\sigma^*) + |G_\sigma^*| - |G^*| = \Delta^*(\mathcal{E})$$

since $g_j^\sigma(G) \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ and (G, σ) is chosen to be the canonical representation of \mathcal{E} . If $g(\mathcal{E})$ is defined in step (3) as $g_j^\tau(G_0)$ for some $G_0 \in \mathcal{E}$ (and $G = G_\sigma^* \in \mathcal{E}$), we claim

$$\Delta^*([g_j^\tau(G_0)]) \leq \text{Hd}(g_j^\tau(G_0), G_\tau^*) + |G_\tau^*| - |G^*| < |G_\sigma^*| - |G^*| = \Delta^*(\mathcal{E}).$$

This is because $g_j^\tau(G_0)$ is an I-map of G^* with ordering τ , and thus $\text{Hd}(g_j^\tau(G_0), G_\tau^*) = |g_j^\tau(G_0)| - |G_\tau^*|$. The above inequality then follows upon noticing that $|G_\sigma^*| = |G_0| > |g_j^\tau(G_0)|$.

Hence, for any $\mathcal{E} \neq \mathcal{E}^*$, we have $\Delta^*(g(\mathcal{E})) < \Delta^*(\mathcal{E})$. Since $\Delta^*(\mathcal{E}) = 0$ if and only if $\mathcal{E} = \mathcal{E}^*$, g induces a canonical path from \mathcal{E} to \mathcal{E}^* for every $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$. To bound the length of such paths, it suffices to bound $\Delta^*(\mathcal{E})$. Observe that

$$\begin{aligned} \Delta^*(\mathcal{E}) &\leq \max_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \sigma \in \mathbb{S}^p} \text{Hd}(G, G_\sigma^*) + \max_{\sigma \in \mathbb{S}^p} (|G_\sigma^*| - |G^*|) \\ &\leq \max_{G \in \mathcal{G}_p(d_{\text{in}}, d_{\text{out}}), \sigma \in \mathbb{S}^p} (|G| + |G_\sigma^*|) + \max_{\sigma \in \mathbb{S}^p} |G_\sigma^*| \\ &\leq (d^* + d_{\text{in}})p + d^*p = (2d^* + d_{\text{in}})p, \end{aligned}$$

which concludes the proof. \square

D.5 Proof of Corollary 2

Proof. Fix an arbitrary $\mathcal{E} \neq \mathcal{E}^*$, and let $g(\mathcal{E}) = [g_j^\sigma(G)]$ for $j \in [p], \sigma \in \mathbb{S}^p$ and $G \in \mathcal{E} \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$ as given in Theorem 3. The proof of Theorem 3 also shows that $g_j^\sigma(G) \in \mathcal{N}_{\text{ads}}(G) \cap \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, and thus $g(\mathcal{E}) \in \mathcal{N}_{\text{ads}}(\mathcal{E}) \cap \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$. By the score equivalence and (D.3),

$$\psi(g(\mathcal{E})) - \psi(\mathcal{E}) = \psi(g_j^\sigma(G)) - \psi(G) = \psi_j(g_j^\sigma(\text{Pa}_j(G))) - \psi_j(\text{Pa}_j(G)). \quad (\text{D.6})$$

So, if (8) holds, $e^{\psi(g(\mathcal{E}))}/e^{\psi(\mathcal{E})} \geq p^t$, which completes the proof. \square

E High-dimensional empirical variable selection

By the SEM representation, the DAG selection problem can be seen as a series of variable selection problems. In this supplement, we prove some high-dimensional consistency results for a single variable selection problem using the empirical normal-inverse-gamma prior. We still use X to denote an $n \times p$ data matrix and we use y to denote a response vector. Let

$$\mathcal{S}(m, d) = \{S \subseteq [m]: |S| \leq d\} \quad (\text{E.1})$$

denote the set of candidate models for some $m, d \in [p]$; that is, we want to select at most d variables from X_1, \dots, X_m to explain y . We allow m to be less than p , which is different from standard variable selection setups, so that the results can be easily applied to DAG selection problems. In particular, without loss of generality, one may assume $y = X_{m+1}$ if $m < p$.

E.1 Model, prior and posterior distributions

Let $y \in \mathbb{R}^n$, $X \in \mathbb{R}^{n \times p}$ and $m, d \in [p]$ be given, and $\mathcal{S}(m, d)$ be as defined in (E.1). Consider the following empirical Bayes model for variable selection [44],

$$\begin{aligned} y &= X_S \beta_S + \varepsilon, \quad \varepsilon \sim N_n(0, \omega I), \\ \pi_0(S) &\propto (c_1 p^{c_2})^{-|S|} \mathbb{1}_{\mathcal{S}(m, d)}(S), \\ \pi_0(\omega) &\propto \omega^{-\kappa/2-1}, \\ \beta_S \mid S, \omega &\sim N_{|S|} \left\{ (X_S^\top X_S)^{-1} X_S^\top y, \frac{\omega}{\gamma} (X_S^\top X_S)^{-1} \right\}, \end{aligned}$$

where I denotes the identity matrix and $c_1 > 0, c_2 \geq 0, \kappa \geq 0, \gamma > 0$ are hyperparameters. Using a fractional likelihood with exponent $\alpha \in (0, 1)$, we find that

$$\pi_n(S) \propto c_1^{-|S|} p^{-c_2|S|} \left(1 + \frac{\alpha}{\gamma}\right)^{-|S|/2} (y^\top \Phi_S^\perp y)^{-(\alpha n + \kappa)/2} \mathbb{1}_{\mathcal{S}(m, d)}(S), \quad (\text{E.2})$$

where Φ_S^\perp (and Φ_S) denotes the projection matrix,

$$\Phi_S = X_S (X_S^\top X_S)^{-1} X_S^\top, \quad \Phi_S^\perp = I - \Phi_S. \quad (\text{E.3})$$

The exponentiation of the posterior score function ψ_j defined in (15) has exactly the same form as (E.2). Thus, the analysis of the DAG selection and structure learning problem can be reduced to that of all possible nodewise variable selection problems.

As in Definition 4, given a true model $S^* \subseteq [m]$ (which will be defined later), we define a transition function $g: \mathcal{S}(m, d) \rightarrow \mathcal{S}(m, d)$ by

$$g(S) = \begin{cases} S, & \text{if } S = S^*, \\ \arg \max_{S' \in \mathcal{N}_{\text{del}}^*(S)} \pi_n(S'), & \text{if } S^* \subset S, \\ \arg \max_{S' \in \mathcal{N}_{\text{add}}^*(S)} \pi_n(S'), & \text{if } S^* \not\subseteq S, |S| < d \\ \arg \max_{S' \in \mathcal{N}_{\text{swap}}^*(S)} \pi_n(S'), & \text{if } S^* \not\subseteq S, |S| = d, \end{cases} \quad (\text{E.4})$$

where

$$\begin{aligned} \mathcal{N}_{\text{add}}^*(S) &= \{S \cup \{k\}: k \in S^* \setminus S\}, \quad \mathcal{N}_{\text{del}}^*(S) = \{S \setminus \{\ell\}: \ell \in S \setminus S^*\}, \\ \mathcal{N}_{\text{swap}}^*(S) &= \{(S \cup \{k\}) \setminus \{\ell\}: k \in S^* \setminus S, \ell \in S \setminus S^*\}. \end{aligned}$$

The goal of this supplement is to obtain a lower bound on $\pi_n(g(S))/\pi_n(S)$ for all $S \in \mathcal{S}(m, d) \setminus S^*$. If the bound is sufficiently large, it then yields the strong consistency of the variable selection procedure. We will consider two sets of conditions. In the first scenario (see Section E.2), we treat $X_{[m]}$ as fixed and impose conditions on the noise part of y , while in the second (see Section E.3), we treat both $X_{[m]}$ and y as random and, assuming $y = X_{m+1}$, start by estimating the covariance matrix for the joint distribution of $(X_1, \dots, X_m, X_{m+1})$.

E.2 High-dimensional consistency: first scenario

Suppose that y can be written as

$$y = y_0 + \epsilon, \quad \text{where } y_0 = X_{S^*} \beta_{S^*}^* \quad (\text{E.5})$$

for some $S^* \subseteq [m]$, $\beta_{S^*}^* \in \mathbb{R}^{|S^*|}$. The vector y_0 represents the signal part of y . We make the following assumptions. The first one controls the multicollinearity of the data, and the second assumes the prior parameters are properly chosen. The last three assumptions concern the behavior of the true model, where we treat the predictors (i.e., $X_{[m]}$) as fixed and require the signal size (true regression coefficient) be sufficiently large and the errors be “well behaved”.

Theorem E1. *Consider the distribution π_n given in (E.2) defined on $\mathcal{S}(m, d)$ for some $m \leq p$ and $d \geq 1$, where $X \in \mathbb{R}^{n \times p}$ and y is given by (E.5). Suppose the following hold.*

(E1.1) *There exist $\underline{\nu}, \bar{\nu} > 0$ such that $y^\top y \leq n\bar{\nu}$ and*

$$n\underline{\nu} \leq \lambda_{\min}(X_S^\top X_S) \leq \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu},$$

for any $S \subseteq [m]$ with $|S| \leq 2d + 1$.

(E1.2) *The vector ϵ in (E.5) satisfies the following for some $\omega^* \in [\underline{\nu}, \bar{\nu}]$ and $\rho \geq 2$.*

$$\begin{aligned} \min_{S: |S| \leq d} \epsilon^\top \Phi_S^\perp \epsilon &\geq n\omega^*/2, \\ \max_{S: |S| \leq d} \max_{j \notin S} \epsilon^\top (\Phi_{S \cup \{j\}} - \Phi_S) \epsilon &\leq \rho\omega^* \log p. \end{aligned}$$

(E1.3) *Prior parameters satisfy that $\kappa \leq n$, $c_1 \sqrt{1 + \alpha/\gamma} \in [1, p]$, and $c_2 \geq (\alpha + 1)\rho + t$ for some constant $t > 0$.*

(E1.4) *The model space parameter d and true model S^* in (E.5) satisfy*

$$\left\{ \frac{4\bar{\nu}^2(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^4} + 1 \right\} |S^*| \leq d.$$

(E1.5) *There exists a constant $C_\beta \geq 8t/3$ such that*

$$\beta_{\min}^2 = \min\{|\beta_j^*|^2: \beta_j^* \neq 0\} \geq 5(C_\beta + 4c_2) \frac{\bar{\nu}^2 \log p}{\alpha \underline{\nu}^2 n}.$$

Then, for the function g defined in (E.4), we have

$$\frac{\pi_n(g(S))}{\pi_n(S)} \geq p^t, \quad \forall S \in \mathcal{S}(m, d) \setminus \{S^*\}.$$

Proof. It follows from Lemmas E1, E2 and E3 to be proved below, each for one subcase in the definition of g (except the case $S = S^*$). \square

First, consider an overfitted model S such that $S^* \subset S$. To show that the posterior probability increases when we remove some covariate in $S \setminus S^*$, we only need Assumption (E1.2), which controls the behavior of ϵ , and Assumption (E1.3), which requires the penalty on model size be sufficiently large.

Lemma E1. *Suppose Assumptions (E1.2) and (E1.3) hold. For any $S \in \mathcal{S}(m, d)$ such that $S^* \subset S$ and any $j \in S \setminus S^*$, we have*

$$\frac{\pi_n(S)}{\pi_n(S \setminus \{j\})} \leq p^{-c_2 + (\alpha+1)\rho} \leq p^{-t}.$$

Proof. Let $S' = S \setminus \{j\}$ for any $j \in S \setminus S^*$. Then, it follows from (E.2) and the inequality $1 + x \leq e^x$ that

$$\begin{aligned} \frac{\pi_n(S)}{\pi_n(S')} &\leq c_1^{-1} (1 + \alpha/\gamma)^{-1/2} p^{-c_2} \exp \left\{ \frac{\alpha n + \kappa}{2} \frac{y^\top (\Phi_S - \Phi_{S'}) y}{y^\top \Phi_S^\perp y} \right\} \\ &= c_1^{-1} (1 + \alpha/\gamma)^{-1/2} p^{-c_2} \exp \left\{ \frac{\alpha n + \kappa}{2} \frac{\epsilon^\top (\Phi_S - \Phi_{S'}) \epsilon}{\epsilon^\top \Phi_S^\perp \epsilon} \right\}. \end{aligned}$$

The second step follows from the observations $S^* \subseteq S$ and $S^* \subseteq S'$. A routine calculation using Assumptions (E1.2) and (E1.3) then yields the result. \square

For an underfitted model $S \in \mathcal{S}(m, d)$ (underfitted means $S^* \setminus S \neq \emptyset$), bounding $\pi_n(g(S))/\pi_n(S)$ is much more difficult due to the collinearity in the design matrix. We need to use Corollary E1 proved in Section E.4. To show that any underfitted model S has a neighboring model with much larger posterior probability, we consider two subcases according as S is saturated (we say S is saturated if $|S| = d$). Note that an underfitted and saturated model exists only if $d < m$. If S is unsaturated, we can add some covariate in $S^* \setminus S$ so that the reduction in residual sum of squares would be significant. If S is saturated, we perform a swap move: add some covariate in $S^* \setminus S$ and remove another in $S \setminus S^*$.

Lemma E2. *Suppose Assumptions (E1.1), (E1.2), (E1.3) and (E1.5) hold. Let $S \in \mathcal{S}(m, d)$ be an underfitted model. There exists some $k \in S^* \setminus S$ such that*

$$\frac{\pi_n(S)}{\pi_n(S \cup \{k\})} \leq p^{-3c_2 - C_\beta/2}.$$

Proof. Let $S' = S \cup \{k\}$ for some $k \in S^* \setminus S$. Then,

$$\frac{\pi_n(S)}{\pi_n(S')} \leq c_1 (1 + \alpha/\gamma)^{1/2} p^{c_2} \exp \left\{ -\frac{\alpha n + \kappa}{2} \frac{y^\top (\Phi_{S'} - \Phi_S) y}{y^\top \Phi_S^\perp y} \right\}. \quad (\text{E.6})$$

By Assumption (E1.1), $y^\top \Phi_S^\perp y \leq y^\top y \leq n\bar{\nu}$. By Corollary E1 and Assumption (E1.5), k can be chosen such that

$$\|(\Phi_{S'} - \Phi_S) y_0\|_2^2 \geq \frac{n\nu^2}{\bar{\nu}} \beta_{\min}^2 \geq 5(C_\beta + 4c_2) \frac{\bar{\nu} \log p}{\alpha} \geq (C_\beta^{1/2} + 4c_2^{1/2})^2 \frac{\bar{\nu} \log p}{\alpha}. \quad (\text{E.7})$$

By Assumptions (E1.2) and (E1.3),

$$\|(\Phi_{S'} - \Phi_S)\epsilon\|_2^2 \leq \rho\omega^* \log p < \frac{c_2\omega^* \log p}{\alpha} \leq \frac{c_2\bar{\nu} \log p}{\alpha}. \quad (\text{E.8})$$

The reverse triangle inequality then yields that

$$\|(\Phi_{S'} - \Phi_S)y\|_2^2 \geq \left\{ \sqrt{C_\beta} + 3\sqrt{c_2} \right\}^2 \frac{\bar{\nu} \log p}{\alpha} \geq (C_\beta + 9c_2) \frac{\bar{\nu} \log p}{\alpha}.$$

Thus, the exponent in (E.6) can be bounded by

$$\frac{\alpha n + \kappa y^\top (\Phi_{S'} - \Phi_S)y}{2 y^\top \Phi_S^\perp y} \geq \frac{C_\beta + 9c_2}{2} \log p.$$

By Assumptions (E1.2) and (E1.3), $c_2 \geq 2$ and thus $9c_2/2 \geq 4c_2 + 1$. The proof is completed upon recalling that $c_1(1 + \alpha/\gamma)^{1/2}p^{c_2} \leq p$. \square

Lemma E3. Suppose Assumptions (E1.1), (E1.2), (E1.3), (E1.4) and (E1.5) hold. Let $S \in \mathcal{S}(m, d)$ be an underfitted model with $|S| = d$. There exist some $k \in S^* \setminus S$ and $j \in S \setminus S^*$ such that

$$\frac{\pi_n(S)}{\pi_n((S \cup \{k\}) \setminus \{j\})} \leq p^{-3C_\beta/8}.$$

Proof. Let $S' = (S \cup \{k\}) \setminus \{j\}$ for some $k \in S^* \setminus S$ and $j \in S \setminus S^*$. Then,

$$\frac{\pi_n(S)}{\pi_n(S')} \leq \exp \left\{ -\frac{\alpha n + \kappa y^\top (\Phi_{S'} - \Phi_S)y}{2 y^\top \Phi_S^\perp y} \right\}.$$

By Corollary E1, we can pick k and j such that

$$\|(\Phi_{S \cup \{k\}} - \Phi_S)y_0\|_2^2 \geq \frac{n\underline{\nu}^2 \|\beta_{S^* \setminus S}^*\|_2^2}{\bar{\nu}|S^* \setminus S|}, \quad \|(\Phi_{S' \cup \{j\}} - \Phi_{S'})y_0\|_2^2 \leq \frac{n\bar{\nu}(\bar{\nu} - \underline{\nu})^2 \|\beta_{S^* \setminus S}^*\|_2^2}{\underline{\nu}^2|S \setminus S^*|}.$$

By Assumption (E1.4),

$$\frac{\|(\Phi_{S \cup \{k\}} - \Phi_S)y_0\|_2^2}{\|(\Phi_{S' \cup \{j\}} - \Phi_{S'})y_0\|_2^2} \geq \frac{\underline{\nu}^4|S \setminus S^*|}{\bar{\nu}^2(\bar{\nu} - \underline{\nu})^2|S^* \setminus S|} \geq \frac{\underline{\nu}^4(d - |S^*|)}{\bar{\nu}^2(\bar{\nu} - \underline{\nu})^2|S^*|} \geq 4.$$

Then, using (E.7), (E.8) and triangle inequalities, we find that

$$\begin{aligned} \|(\Phi_{S \cup \{k\}} - \Phi_S)y\|_2^2 &\geq \left\{ \sqrt{C_\beta} + 3\sqrt{c_2} \right\}^2 \frac{\bar{\nu} \log p}{\alpha}, \\ \|(\Phi_{S' \cup \{j\}} - \Phi_{S'})y\|_2^2 &\leq \left\{ \frac{\sqrt{C_\beta}}{2} + 3\sqrt{c_2} \right\}^2 \frac{\bar{\nu} \log p}{\alpha}. \end{aligned}$$

Hence,

$$\|(\Phi_{S'} - \Phi_S)y\|_2^2 = \|(\Phi_{S \cup \{k\}} - \Phi_S)y\|_2^2 - \|(\Phi_{S' \cup \{j\}} - \Phi_{S'})y\|_2^2 \geq \frac{3C_\beta\bar{\nu} \log p}{4\alpha}.$$

The result then follows by a calculation similar to the proof of Lemma E2. \square

E.3 High-dimensional consistency: second scenario

We now re-derive the high-dimensional consistency result for the model described in Section E.1 by treating both X and y as random and viewing linear regression as orthogonal projection. This result is useful when one wants to use random matrix theory to study DAG selection or structure learning problems. We will use the following notation. Given a matrix M , let $\|M\|_{\text{op}} = \sup_{\|b\|_2=1} \|Mb\|_2$ denote its operator norm. Given a positive definite matrix Σ and sets (or integers) A, B , we use $\Sigma_{A,B}$ to denote the submatrix of Σ with rows indexed by A and columns indexed by B , and, by an abuse of notation, we simply write $\Sigma_A = \Sigma_{A,A}$ ². Further, we use

$$\Sigma_{A,B|C} = \Sigma_{A,B} - \Sigma_{A,C}(\Sigma_C)^{-1}\Sigma_{C,B} \quad (\text{E.9})$$

to denote the “partial covariance” and write $\Sigma_{A|C} = \Sigma_{A,A|C}$.

The following theorem has the same conclusion as Theorem E1, but we replace Assumption (E1.2) in Theorem E1 with $\zeta = O(\sqrt{n^{-1}d \log p})$ where ζ measures the difference between Σ^* and the empirical covariance matrix; all the other four assumptions are very similar.

Theorem E2. *Consider the distribution π_n given in (E.2) defined on $\mathcal{S}(m, d)$ for some $m \in [p-1]$ and $d \geq 1$, where $X \in \mathbb{R}^{n \times p}$, $y = X_{m+1}$ and each row of $X_{[m+1]}$ is an i.i.d. copy of a random vector $\mathbf{X} = (X_1, \dots, X_{m+1})$ with mean zero and covariance matrix Σ^* . Define*

$$\beta^* = (\Sigma_{[m]}^*)^{-1}\Sigma_{[m],m+1}^*, \quad S^* = \{j: \beta_j^* \neq 0\}, \quad \hat{\Sigma} = n^{-1}X^\top X.$$

Assume the following hold.

(E2.1) *There exist constants $\bar{\nu} \geq 1 \geq \underline{\nu}$ such that $\underline{\nu} \leq \lambda_{\min}(\Sigma^*) \leq \lambda_{\max}(\Sigma^*) \leq \bar{\nu}$ and, for any $S \in \mathcal{S}(m+1, 2d+1)$, $\underline{\nu} \leq \lambda_{\min}(\hat{\Sigma}_S) \leq \lambda_{\max}(\hat{\Sigma}_S) \leq \bar{\nu}$.*

(E2.2) *There exists a constant $C_{\text{err}} > 0$ such that $\zeta^2 \leq C_{\text{err}} n^{-1} d \log p$ where*

$$\zeta = \max_{S \in \mathcal{S}(m+1, 2d)} \|\hat{\Sigma}_S - \Sigma_S^*\|_{\text{op}}.$$

(E2.3) *Prior parameters satisfy that $\alpha \in (0, 1]$, $\kappa \leq n$, $c_1 \sqrt{1 + \alpha/\gamma} \in [1, p]$, and*

$$c_2 \geq C_{\text{pen}} \frac{d\bar{\nu}^4}{\underline{\nu}^6},$$

for some constant $C_{\text{pen}} > 0$.

(E2.4) *The model space parameter d and true model size $|S^*|$ satisfy*

$$\left\{ \frac{20\bar{\nu}^2(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^4} + 1 \right\} |S^*| \leq d.$$

²This notation is only used for covariance matrices and only used in Supplements E and F. Note that its meaning is different from X_S , which denotes the submatrix of X with columns indexed by S .

(E2.5) There exists a constant C_β such that

$$\beta_{\min}^2 = \min\{|\beta_j^*|^2 : \beta_j^* \neq 0\} \geq C_\beta \frac{c_2 \bar{\nu}^2 \log p}{\alpha \underline{\nu}^2 n}.$$

Given any $t > 0$, we can choose sufficiently large C_{pen} and C_β , which only depend on t and C_{err} , such that

$$\frac{\pi_n(g(S))}{\pi_n(S)} \geq p^t, \quad \forall S \in \mathcal{S}(m, d) \setminus \{S^*\}.$$

Remark E1. There is no loss of generality in assuming $\bar{\nu} \geq 1 \geq \underline{\nu}$ since one can always scale Σ^* . By the modified Cholesky decomposition, there exist a strictly upper triangular matrix B^* and a diagonal matrix Ω^* such that $\Sigma^* = (I - (B^*)^\top)^{-1} \Omega^* (I - B^*)^{-1}$. By the block matrix inversion formula, one can show that $\beta^* = B_{[m], m+1}^*$. Further, for any $j \in [m]$, the block matrix inversion formula yields that

$$\Sigma_{j, m+1}^* = \Sigma_{j, [m]}^* (\Sigma_{[m]}^*)^{-1} \Sigma_{[m], m+1}^* = \Sigma_{j, [m]}^* \beta^* = \Sigma_{j, S^*}^* \beta_{S^*}^*, \quad (\text{E.10})$$

which will be useful in the proof of Theorem E2.

Before we prove Theorem E2, we derive a lemma for controlling the error propagation via matrix operations, which is the main technical difference between the proofs of Theorem E1 and Theorem E2. Note that for any matrix M , we have $\|M\|_{\text{op}} \geq \max_{i,j} |M_{ij}|$, so Lemma E4 also yields a bound on the entrywise maximum error.

Lemma E4. Let $\Sigma^*, \hat{\Sigma} \in \mathbb{R}^{(m+1) \times (m+1)}$ be positive definite matrices that satisfy Assumption (E2.1) in Theorem E2. For any $S, T \in \mathcal{S}(m+1, d)$ such that $S \cap T = \emptyset$,

$$\begin{aligned} \|\hat{\Sigma}_{T|S} - \Sigma_{T|S}^*\|_{\text{op}} &\leq \zeta \bar{\nu}^2 / \underline{\nu}^2, \\ \|(\hat{\Sigma}_S)^{-1} \hat{\Sigma}_{S,T} - (\Sigma_S^*)^{-1} \Sigma_{S,T}^*\|_{\text{op}} &\leq 2\zeta \bar{\nu} / \underline{\nu}^2, \end{aligned}$$

where $\zeta = \max_{S \in \mathcal{S}(m+1, 2d)} \|\hat{\Sigma}_S - \Sigma_S^*\|_{\text{op}}$.

Proof. Let $\Sigma = \hat{\Sigma}$ or Σ^* . Since $(\Sigma_{T|S})^{-1}$ is a principal submatrix of $(\Sigma_{S \cup T})^{-1}$,

$$\bar{\nu}^{-1} \leq \lambda_{\min}((\Sigma_{S \cup T})^{-1}) \leq \lambda_{\min}((\Sigma_{T|S})^{-1}) \leq \lambda_{\max}((\Sigma_{T|S})^{-1}) \leq \lambda_{\max}((\Sigma_{S \cup T})^{-1}) \leq \underline{\nu}^{-1}.$$

Given invertible matrices A, B of the same dimension, $A^{-1} - B^{-1} = A^{-1}(B - A)B^{-1}$. Hence, by the sub-multiplicative property of operator norm,

$$\|(\hat{\Sigma}_{T|S})^{-1} - (\Sigma_{T|S}^*)^{-1}\|_{\text{op}} \leq \|(\hat{\Sigma}_{S \cup T})^{-1} - (\Sigma_{S \cup T}^*)^{-1}\|_{\text{op}} \leq \zeta / \underline{\nu}^2.$$

Applying the same argument again, we get $\|\hat{\Sigma}_{T|S} - \Sigma_{T|S}^*\|_{\text{op}} \leq \zeta \bar{\nu}^2 / \underline{\nu}^2$.

To prove the second inequality, we use the identity $AB - A'B' = A(B - B') + (A - A')B'$ for matrices A, A', B, B' . Since $\hat{\Sigma}_{S,T} - \Sigma_{S,T}^*$ is a submatrix of $\hat{\Sigma}_{S \cup T} - \Sigma_{S \cup T}^*$, we have $\|\hat{\Sigma}_{S,T} - \Sigma_{S,T}^*\|_{\text{op}} \leq \zeta$. It follows that

$$\|(\hat{\Sigma}_S)^{-1} \hat{\Sigma}_{S,T} - (\Sigma_S^*)^{-1} \Sigma_{S,T}^*\|_{\text{op}} \leq \zeta / \underline{\nu} + \zeta \bar{\nu} / \underline{\nu}^2,$$

which yields the asserted bound since $\bar{\nu} \geq \underline{\nu}$. \square

Proof of Theorem E2. Since $y = X_{m+1}$, we can write

$$n^{-1}y^\top \Phi_{S'}^\perp y = \hat{\Sigma}_{m+1|S}, \quad n^{-1}y^\top (\Phi_{S \cup \{j\}} - \Phi_S)y = (\hat{\Sigma}_{j|S})^{-1}(\hat{\Sigma}_{j,m+1|S})^2,$$

for any $S \subset [m]$ and $j \in [m] \setminus S$. As in the proof of Theorem E1, we split the proof into three cases.

Case 1: overfitted Suppose $S^* \subset S \in \mathcal{S}(m, d)$ and let $S' = S \setminus \{j\}$ for some $j \in S \setminus S^*$. By Assumption (E2.3), we have

$$\frac{\pi_n(S)}{\pi_n(S')} \leq p^{-c_2} \exp \left\{ \frac{n(\hat{\Sigma}_{j|S'})^{-1}(\hat{\Sigma}_{j,m+1|S'})^2}{\hat{\Sigma}_{m+1|S}} \right\}.$$

Assumption (E2.1) implies that $\hat{\Sigma}_{m+1|S} \geq \underline{\nu}$ and $(\hat{\Sigma}_{j|S'})^{-1} \leq \underline{\nu}^{-1}$. By (E.10), we have $\Sigma_{j,m+1|S'}^* = \Sigma_{j,S^*|S'}^* \beta_{S^*}^* = 0$ since $S^* \subseteq S'$. Apply Lemma E4 to get $|\hat{\Sigma}_{j,m+1|S'}| \leq \zeta \bar{\nu}^2 / \underline{\nu}^2$. It follows by Assumption (E2.2) that

$$\frac{\pi_n(S)}{\pi_n(S')} \leq p^{-c_2} \exp \left\{ \frac{n \zeta^2 \bar{\nu}^4}{\underline{\nu}^6} \right\} \leq \exp \left\{ \left(-c_2 + \frac{C_{\text{err}} d \bar{\nu}^4}{\underline{\nu}^6} \right) \log p \right\},$$

Clearly, $d \bar{\nu}^4 / \underline{\nu}^6 \geq 1$ since $\underline{\nu} \leq \bar{\nu}$ and we assume $\underline{\nu} \leq 1$. Hence, it suffices to choose $C_{\text{pen}} \geq C_{\text{err}} + t$ so that $\pi_n(S) / \pi_n(S') \leq p^{-t}$.

Case 2: underfitted and unsaturated Let $S \in \mathcal{S}(m, d-1)$ be such that $S^* \setminus S \neq \emptyset$. By Assumptions (E2.1) and (E2.3), for any $k \in S^* \setminus S$,

$$\frac{\pi_n(S)}{\pi_n(S \cup \{k\})} \leq p^{c_2+1} \exp \left\{ -\frac{\alpha n}{2} \frac{(\hat{\Sigma}_{k,m+1|S})^2}{\bar{\nu}^2} \right\}. \quad (\text{E.11})$$

Define

$$\hat{b} = (\hat{\Sigma}_{S \cup S^*})^{-1} \hat{\Sigma}_{S \cup S^*, m+1}, \quad b^* = (\Sigma_{S \cup S^*}^*)^{-1} \Sigma_{S \cup S^*, m+1}^*. \quad (\text{E.12})$$

Applying the first part of Lemma E5 to $\hat{\Sigma}_{S \cup S^* \cup \{m+1\}}$ with $T = S^* \setminus S$, $V = S$ and $l = |S \cup S^*|$, we get

$$\sum_{k \in T} (\hat{\Sigma}_{k,m+1|S})^2 \geq \lambda_{\min}(\hat{\Sigma}_{T|S})^2 \|\hat{b}_T\|_2^2 \geq \underline{\nu}^2 \|\hat{b}_T\|_2^2.$$

Observe that $b_T^* = \beta_T^*$; this can be proved by applying the block matrix inversion formula to $(\Sigma_{[m]}^*)^{-1}$. Then, by Lemma E4, for any $k \in S^* \setminus S$, we have $|\hat{b}_k - \beta_k^*| \leq \|\hat{b} - b^*\|_{\text{op}} \leq 2\zeta \bar{\nu} / \underline{\nu}^2$, and thus

$$(\hat{b}_k)^2 \geq (|\beta_k^*| - 2\zeta \bar{\nu} / \underline{\nu}^2)^2.$$

By Assumptions (E2.2), (E2.3) and (E2.5),

$$|\beta_k^*|^2 \geq C_\beta \frac{c_2 \bar{\nu}^2 \log p}{\alpha \underline{\nu}^2 n} \geq C_\beta C_{\text{pen}} \frac{d \bar{\nu}^6 \log p}{\alpha \underline{\nu}^8 n} \geq \frac{C_\beta C_{\text{pen}}}{C_{\text{err}}} \frac{\zeta^2 \bar{\nu}^6}{\underline{\nu}^8} \geq \frac{C_\beta C_{\text{pen}}}{C_{\text{err}}} \left(\frac{\zeta \bar{\nu}}{\underline{\nu}^2} \right)^2.$$

As long as $C_\beta C_{\text{pen}} / C_{\text{err}}$ is sufficiently large, we have $(\hat{b}_k)^2 \geq (\beta_k^*)^2 / 2$. It follows that

$$\sum_{k \in S^* \setminus S} (\hat{\Sigma}_{k,m+1|S})^2 \geq \frac{\underline{\nu}^2}{2} \|\beta_{S^* \setminus S}^*\|_2^2. \quad (\text{E.13})$$

Choosing an optimal k , we get $(\hat{\Sigma}_{k,m+1|S})^2 \geq \underline{\nu}^2 \beta_{\min}^2 / 2$. Using (E.11) and Assumption (E2.5), we find that

$$\frac{\pi_n(S)}{\pi_n(S \cup \{k\})} \leq p^{-(C_\beta/4-1)c_2+1}.$$

Consequently, we only need to choose sufficiently large $C_\beta = C_\beta(t, C_{\text{err}}, C_{\text{pen}})$ so that $\pi_n(S)/\pi_n(S') \leq p^{-t}$.

Case 3: underfitted and saturated Let $S \subseteq [m]$ be such that $S^* \setminus S \neq \emptyset$ and $|S| = d$. For any $k \in S^* \setminus S$ and $j \in S \setminus S^*$, by Assumption (E2.1),

$$\frac{\pi_n(S)}{\pi_n(S')} \leq \exp \left\{ -\frac{\alpha n(R_1 - R_2)}{2\bar{\nu}} \right\},$$

where $R_1 = (\hat{\Sigma}_{k|S})^{-1}(\hat{\Sigma}_{k,m+1|S})^2$, $R_2 = (\hat{\Sigma}_{j|S'})^{-1}(\hat{\Sigma}_{j,m+1|S'})^2$,

and $S' = (S \cup \{k\}) \setminus \{j\}$. By (E.13), there exists some $k \in S^* \setminus S$ such that

$$R_1 \geq \frac{\underline{\nu}^2}{2\bar{\nu}} \frac{\|\beta_{S^* \setminus S}^*\|_2^2}{|S^* \setminus S|} \geq \frac{\underline{\nu}^2}{2\bar{\nu}} \beta_{\min}^2 \geq \frac{C_\beta C_{\text{pen}}}{2C_{\text{err}}} \frac{\zeta^2 \bar{\nu}^5}{\underline{\nu}^6}. \quad (\text{E.14})$$

Let \hat{b} and b^* be as defined in (E.12). To bound R_2 , we apply the second part of Lemma E5 to $\hat{\Sigma}_{S \cup S^* \cup \{m+1\}}$. Letting $U = S \cup \{k\}$, $V = U \cap S^*$, $T = (U \setminus S^*) \cup (S^* \setminus U)$ and $W = U \setminus S^*$, we get that

$$\sum_{j \in W} \left\{ (\hat{\Sigma}_{j|U \setminus \{j\}})^{-1} \hat{\Sigma}_{j,m+1|U \setminus \{j\}} \right\}^2 \leq \|(\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,T} \hat{b}_T\|_2^2. \quad (\text{E.15})$$

Since $S^* \subset S \cup S^*$, we have $b^* = \beta_{S \cup S^*}^*$, which implies $b_T^* = \beta_T^*$. By Lemma E4 and Assumptions (E2.2), (E2.3) and (E2.5), for any $i \in S^* \setminus U$, we have $(\hat{b}_i)^2 \leq 2(\beta_i^*)^2$ as long as $C_\beta C_{\text{pen}}/C_{\text{err}}$ is sufficiently large. Hence, applying Lemma E4 again, we obtain

$$\|\hat{b}_{S^* \setminus U}\|_2^2 \leq 2\|\beta_{S^* \setminus U}^*\|_2^2, \quad \|\hat{b}_{U \setminus S^*}\|_2^2 \leq 2|U \setminus S^*| \zeta^2 \bar{\nu}^2 / \underline{\nu}^4.$$

Since $U \setminus S^*$ and $S^* \setminus U$ are disjoint, it follows that

$$\begin{aligned} \|(\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,T} \hat{b}_T\|_2^2 &= \|(\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,U \setminus S^*} \hat{b}_{U \setminus S^*} + (\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,S^* \setminus U} \hat{b}_{S^* \setminus U}\|_2^2 \\ &\leq 2\|\hat{b}_{U \setminus S^*}\|_2^2 + 2\|(\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,S^* \setminus U} \hat{b}_{S^* \setminus U}\|_2^2 \\ &\leq 4|U \setminus S^*| \zeta^2 \bar{\nu}^2 / \underline{\nu}^4 + 2\|(\hat{\Sigma}_U)^{-1} \hat{\Sigma}_{U,S^* \setminus U}\|_{\text{op}}^2 \|\hat{b}_{S^* \setminus U}\|_2^2 \\ &\leq 4|U \setminus S^*| \frac{\zeta^2 \bar{\nu}^2}{\underline{\nu}^4} + 4 \frac{(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^2} \|\beta_{S^* \setminus U}^*\|_2^2, \end{aligned}$$

where in the last step we have used Lemma E6 to bound the operator norm. Choosing an optimal j in (E.15), we get

$$\begin{aligned} R_2 &= (\hat{\Sigma}_{j|S'}) \left\{ (\hat{\Sigma}_{j|S'})^{-1} (\hat{\Sigma}_{j,m+1|S'}) \right\}^2 \leq 4\bar{\nu} \left(\frac{\zeta^2 \bar{\nu}^2}{\underline{\nu}^4} + \frac{(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^2 |U \setminus S^*|} \|\beta_{S^* \setminus U}^*\|_2^2 \right) \\ &\leq 8 \max \left\{ \frac{\zeta^2 \bar{\nu}^3}{\underline{\nu}^4}, \frac{\bar{\nu}(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^2 |U \setminus S^*|} \|\beta_{S^* \setminus U}^*\|_2^2 \right\}. \end{aligned}$$

If the first term in the maximum is larger, by (E.14), we only need to let $C_\beta C_{\text{pen}}/C_{\text{err}}$ be sufficiently large so that $R_1/R_2 \geq 5/4$. If the second term is larger (note $S^* \setminus U = S^* \setminus S$), one can use (E.14) and Assumption (E2.4) to show that $R_1/R_2 \geq 5/4$. To summarize, we can choose $k \in S^* \setminus S$ and $j \in S \setminus S^*$ such that

$$\frac{\pi_n(S)}{\pi_n((S \cup \{k\}) \setminus \{j\})} \leq \exp \left\{ -\frac{\alpha n R_1}{8\bar{\nu}} \right\} \leq \exp \left\{ -\frac{\alpha n \nu^2}{16\bar{\nu}^2} \beta_{\min}^2 \right\}.$$

A routine calculation then completes the proof. \square

E.4 Auxiliary lemmas

In this section, we prove some useful results for bounding the change in residual sum of squares for optimal addition or deletion moves. We first prove a general linear algebra result in Lemma E5 and then use it to obtain Corollary E1, which is similar to Lemma 8 of Yang et al. [77] [cf. 3, Lemma 1]. Note that Lemma 8 of Yang et al. [77] requires an irrepresentability assumption and involves the constant $\max_{S \in \mathcal{S}(m,d)} \|(\hat{\Sigma}_S)^{-1} \hat{\Sigma}_{S, S^* \setminus S}\|_{\text{op}}$. But we directly bound this constant using Lemma E6.

Lemma E5. *Let $\Sigma \in \mathbb{R}^{(l+1) \times (l+1)}$ be positive definite for some $l \in \mathbb{N}$ and define $\beta = (\Sigma[l])^{-1} \Sigma[l, l+1]$. Let V, U be nonempty sets such that $V \subset U \subseteq [l]$, and let $T = [l] \setminus V$ and $W = U \setminus V$. Then,*

$$\begin{aligned} \sum_{k \in T} (\Sigma_{k, l+1|V})^2 &\geq (\lambda_{\min}(\Sigma_{T|V}))^2 \|\beta_T\|_2^2, \\ \sum_{j \in W} \{(\Sigma_{j|U \setminus \{j\}})^{-1} \Sigma_{j, l+1|U \setminus \{j\}}\}^2 &\leq \|(\Sigma_U)^{-1} \Sigma_{U, T} \beta_T\|_2^2. \end{aligned}$$

Proof. Using $\Sigma_{k, l+1} = \Sigma_{k, [l]} \beta$ for any $k \in [l]$, we find that

$$\Sigma_{k, l+1|V} = \Sigma_{k, [l]|V} \beta = \Sigma_{k, 1|V} \beta_1 + \cdots + \Sigma_{k, l|V} \beta_l.$$

Observe that $\Sigma_{k, i|V} = 0$ if $i \in V$. Hence, $\Sigma_{k, l+1|V} = \Sigma_{k, T|V} \beta_T$. It follows that

$$\sum_{k \in T} (\Sigma_{k, l+1|V})^2 = \|\Sigma_{T|V} \beta_T\|_2^2 \geq (\lambda_{\min}(\Sigma_{T|V}))^2 \|\beta_T\|_2^2.$$

For the second claim, without loss of generality, assume $U = \{1, 2, \dots, |U|\}$ and define

$$b = (\Sigma_U)^{-1} \Sigma_{U, l+1} = (\Sigma_U)^{-1} \Sigma_{U, [l]} \beta.$$

We have $b_j = (\Sigma_{j|U \setminus \{j\}})^{-1} \Sigma_{j, l+1|U \setminus \{j\}}$ for each $j \in U$; this can be proved by applying the block matrix inversion formula to $(\Sigma_U)^{-1}$ with blocks $U \setminus \{j\}$ and $\{j\}$. Note that we can write $b = b^{(1)} + b^{(2)}$, where

$$b^{(1)} = (\Sigma_U)^{-1} \Sigma_{U, V} \beta_V, \quad b^{(2)} = (\Sigma_U)^{-1} \Sigma_{U, T} \beta_T,$$

and $b^{(1)}$ satisfies $b_W^{(1)} = 0$. Therefore, for any $j \in W$, $b_j = b_j^{(2)}$. Summing over $j \in W$, we get

$$\sum_{j \in W} \{(\Sigma_{j|U \setminus \{j\}})^{-1} \Sigma_{j, l+1|U \setminus \{j\}}\}^2 = \sum_{j \in W} (b_j^{(2)})^2 \leq \|(\Sigma_U)^{-1} \Sigma_{U, T} \beta_T\|_2^2,$$

which yields the asserted inequality. \square

Corollary E1. Let y_0 be as defined in (E.5) for some $S^* \in \mathcal{S}(m, d)$. Suppose

$$n\underline{\nu} \leq \min_{S \in \mathcal{S}(m, 2d+1)} \lambda_{\min}(X_S^\top X_S) \leq \max_{S \in \mathcal{S}(m, 2d+1)} \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu},$$

for some $\underline{\nu}, \bar{\nu} \in (0, \infty)$. Let Φ_S be as given in (E.3). For any $S \in \mathcal{S}(m, d)$,

$$\begin{aligned} \sum_{k \in S^* \setminus S} \|(\Phi_{S \cup \{k\}} - \Phi_S)y_0\|_2^2 &\geq \frac{n\underline{\nu}^2}{\bar{\nu}} \|\beta_{S^* \setminus S}^*\|_2^2, \\ \sum_{j \in S \setminus S^*} \|(\Phi_S - \Phi_{S \setminus \{j\}})y_0\|_2^2 &\leq \frac{n\bar{\nu}(\bar{\nu} - \underline{\nu})^2}{\underline{\nu}^2} \|\beta_{S^* \setminus S}^*\|_2^2. \end{aligned}$$

Proof. Let $\hat{\Sigma} = n^{-1}X^\top X$. Using $y_0 = X_{S^*}\beta_{S^*}^*$, the formula

$$\Phi_{S \cup \{k\}} - \Phi_S = \Phi_S^\perp X_k (X_k^\top \Phi_S^\perp X_k)^{-1} X_k^\top \Phi_S^\perp, \quad (\text{E.16})$$

and the eigenvalue assumption, we find that

$$\begin{aligned} n^{-1} \|(\Phi_{S \cup \{k\}} - \Phi_S)y_0\|_2^2 &= (\hat{\Sigma}_{k|S})^{-1} (\hat{\Sigma}_{k, S^*|S} \beta_{S^*}^*)^2 \geq \bar{\nu}^{-1} (\hat{\Sigma}_{k, S^*|S} \beta_{S^*}^*)^2, \\ n^{-1} \|(\Phi_S - \Phi_{S \setminus \{j\}})y_0\|_2^2 &= (\hat{\Sigma}_{j|S \setminus \{j\}})^{-1} (\hat{\Sigma}_{j, S^*|S \setminus \{j\}} \beta_{S^*}^*)^2 \leq \bar{\nu} \left\{ (\hat{\Sigma}_{j|S \setminus \{j\}})^{-1} \hat{\Sigma}_{j, S^*|S \setminus \{j\}} \beta_{S^*}^* \right\}^2. \end{aligned}$$

Let $S' = S^* \setminus S$. By an argument analogous to the proof of Lemma E5, one can show that

$$\begin{aligned} \sum_{k \in S^* \setminus S} (\hat{\Sigma}_{k, S^*|S} \beta_{S^*}^*)^2 &= \|\hat{\Sigma}_{k, S'|S} \beta_{S'}^*\|_2^2, \\ \sum_{j \in S \setminus S^*} \left\{ (\hat{\Sigma}_{j|S \setminus \{j\}})^{-1} \hat{\Sigma}_{j, S^*|S \setminus \{j\}} \beta_{S^*}^* \right\}^2 &\leq \|(\hat{\Sigma}_S)^{-1} \hat{\Sigma}_{S, S'} \beta_{S'}^*\|_2^2. \end{aligned}$$

The asserted inequalities follow from the eigenvalue assumption and Lemma E6 below. \square

Lemma E6. Let $A = [A_1 \ A_2]$ be an $n \times k$ matrix for some $k \leq n$, $\lambda_{\max}(A^\top A) = \nu_{\max}$ and $\lambda_{\min}(A^\top A) = \nu_{\min}$. Then, $\|A_1^\top A_2\|_{\text{op}} \leq \nu_{\max} - \nu_{\min}$.

Proof. Suppose the dimension of A_i is $n \times k_i$ for $i = 1, 2$. By the definition of operator norm,

$$\begin{aligned} \|A_1^\top A_2\|_{\text{op}} &= \max_{b_2 \in \mathbb{R}^{k_2}: \|b_2\|=1} \|A_1^\top A_2 b_2\| \\ &= \max \left\{ b_1^\top A_1^\top A_2 b_2 : b_1 \in \mathbb{R}^{k_1}, b_2 \in \mathbb{R}^{k_2}, \|b_1\| = \|b_2\| = 1 \right\}. \end{aligned}$$

Since $\|A_1\|_{\text{op}} \leq \|A\|_{\text{op}} = \sqrt{\nu_{\max}}$ and $\sigma_{\min}(A) = \sqrt{\nu_{\min}}$, we have

$$\begin{aligned} 2b_1^\top A_1^\top A_2 b_2 &= \|A_1 b_1\|_2^2 + \|A_2 b_2\|_2^2 - \|(A_1 b_1 - A_2 b_2)\|_2^2 \\ &\leq \nu_{\max} \|b_1\|_2^2 + \nu_{\max} \|b_2\|_2^2 - \nu_{\min} (\|b_1\|_2^2 + \|b_2\|_2^2). \end{aligned}$$

Hence, if $\|b_1\| = \|b_2\| = 1$, $b_1^\top A_1^\top A_2 b_2 \leq \nu_{\max} - \nu_{\min}$. A similar argument yields that $b_1^\top A_1^\top A_2 b_2 \geq \nu_{\min} - \nu_{\max}$, which completes the proof. \square

F Proofs for Section 4

F.1 Empirical Bayes Gaussian DAG model

Let $X_{(i)}$ denote the i -th row of the data matrix X . We model the conditional distribution of X given G by

$$\begin{aligned} X_{(1)}, \dots, X_{(n)} &| B, \Omega \stackrel{\text{i.i.d.}}{\sim} N_p(0, \Sigma(B, \Omega)), \\ \Sigma(B, \Omega) &= (I - B^\top)^{-1} \Omega (I - B)^{-1}, \\ (B, \Omega) &| G \sim \pi_0(B, \Omega | G), \quad \forall (B, \Omega) \in \mathcal{D}_p(G). \end{aligned} \tag{F.1}$$

Since a linear transformation of a normal random vector is still normal, the conditional distribution of X given (B, Ω) can also be expressed by the SEM,

$$X_j = \sum_{i \neq j} B_{ij} X_i + \varepsilon_j, \quad \varepsilon_j \sim N_n(0, \omega_j I),$$

for $j = 1, \dots, p$, where $\varepsilon_1, \dots, \varepsilon_p$ are independent error vectors. We use the model (F.1) for two reasons. First, by Lemma C6, if Σ is positive definite and $N_p(0, \Sigma)$ is Markovian w.r.t. G , then there exists a unique pair $(B, \Omega) \in \mathcal{D}_p(G)$ such that $\Sigma = \Sigma(B, \Omega)$. Second, by Lemma C7, if the edge weights (entries B_{ij} for $i \rightarrow j \in G$) are sampled from an absolutely continuous distribution (which is true for our empirical prior), the resulting distribution $N_p(0, \Sigma)$ is almost surely perfectly Markovian w.r.t. G . There is little loss of generality in assuming that X has mean zero, since the normality implies that any CI statement about X_1, \dots, X_p can be determined by using Σ alone.

Recalling $\Omega = \text{diag}(\omega_1, \dots, \omega_p)$ and using the notation $\beta_j(G)$ defined in Section 4.1, we can express our empirical prior $\pi_0(B, \Omega | G)$ by

$$\begin{aligned} &\pi_0(B, \Omega | \text{Pa}_1(G) = S_1, \dots, \text{Pa}_p(G) = S_p) \\ &\propto \prod_{j=1}^p \omega_j^{-\kappa/2-1} N_{|S_j|} \left(\beta_j(G); (X_{S_j}^\top X_{S_j})^{-1} X_{S_j}^\top X_j, \frac{\omega_j}{\gamma} (X_{S_j}^\top X_{S_j})^{-1} \right), \end{aligned} \tag{F.2}$$

where $N_q(b; \mu, \Sigma)$ denotes the density function of $N_q(\mu, \Sigma)$ evaluated at b . Note that since (B, Ω) takes value in $\mathcal{D}_p(G)$, $\beta_j(G)$ contains all nonzero regression coefficients for the response vector X_j . The prior mean for $\beta_j(G)$ is simply the ordinary-least-squares estimator. Let $L(B, \Omega)$ denote the likelihood function (the dependency on X is omitted). Since the empirical prior relies on the observed data, to counteract its effect we use a fractional likelihood with exponent $\alpha \in (0, 1)$, which yields the conditional posterior distribution,

$$\pi_n(B, \Omega | G) \propto \pi_0(B, \Omega | G) L(B, \Omega)^\alpha = \frac{\pi_0(B, \Omega | G)}{L(B, \Omega)^{1-\alpha}} L(B, \Omega).$$

This shows that the effective prior distribution for $(B, \Omega) | G$ is $\pi_0(B, \Omega | G) / L^{1-\alpha}(B, \Omega)$; see Martin et al. [44], Lee et al. [39] for more discussion. By a routine calculation using the conjugacy of normal-inverse-gamma prior, we obtain the fractional marginal likelihood

function f_α given in the main text,

$$\begin{aligned} f_\alpha(G) &= \int \pi_0(B, \Omega \mid G) L(B, \Omega)^\alpha d(B, \Omega) \\ &= \left(1 + \frac{\alpha}{\gamma}\right)^{-|G|/2} \prod_{j=1}^p (X_j^\top \Phi_{\text{Pa}_j}^\perp X_j)^{-(\alpha n + \kappa)/2}, \end{aligned} \quad (\text{F.3})$$

where we recall Φ_S^\perp is defined by

$$\Phi_S = X_S (X_S^\top X_S)^{-1} X_S^\top, \quad \Phi_S^\perp = I - \Phi_S, \quad \forall S \subseteq [p].$$

We can use f_α to derive posterior distributions of DAGs or equivalence classes. For example, given a prior distribution of DAGs, $\tilde{\pi}_0(G)$, we can calculate the posterior by

$$\tilde{\pi}_n(G) = \int \tilde{\pi}_n(G, B, \Omega) d(B, \Omega) \propto \int \tilde{\pi}_0(G) \pi_0(B, \Omega \mid G) L(B, \Omega)^\alpha d(B, \Omega) = \tilde{\pi}_0(G) f_\alpha(G).$$

F.2 Proof of Lemma 3

Proof. This is equivalent to proving that the marginal fractional likelihood defined in (F.3) is the same for Markov equivalent DAGs (since they have the same skeleton and thus the same number of edges). By Lemma C2, if two DAGs are Markov equivalent, then there exists a sequence of covered edge reversals that can transform one to the other. So it suffices to show that any covered edge reversal does not change the marginal likelihood. Let G, G' be two DAGs that differ by a covered edge reversal. Thus, there exist $i \neq j$ such that $i \rightarrow j \in G$, $j \rightarrow i \in G'$, $\text{Pa}_i(G) = \text{Pa}_j(G) \setminus \{i\}$, and all the other edges are exactly the same in the two DAGs. By (15), for $S = \text{Pa}_i(G)$ we have

$$\frac{\exp(\psi(G))}{\exp(\psi(G'))} = \left(\frac{X_i^\top \Phi_S^\perp X_i X_j^\top \Phi_{S \cup \{i\}}^\perp X_j}{X_i^\top \Phi_{S \cup \{j\}}^\perp X_i X_j^\top \Phi_S^\perp X_j} \right)^{-(\alpha n + \kappa)/2}.$$

It then follows from (E.16) that

$$\begin{aligned} (X_i^\top \Phi_S^\perp X_i)(X_j^\top \Phi_{S \cup \{i\}}^\perp X_j) &= (X_i^\top \Phi_S^\perp X_i) X_j^\top \left(\Phi_S^\perp - \frac{\Phi_S^\perp X_i X_i^\top \Phi_S^\perp}{X_i^\top \Phi_S^\perp X_i} \right) X_j \\ &= (X_i^\top \Phi_S^\perp X_i)(X_j^\top \Phi_S^\perp X_j) - (X_j^\top \Phi_S^\perp X_i)^2. \end{aligned}$$

By symmetry, we conclude that $\psi(G) = \psi(G')$. \square

F.3 Proof of Theorem 4

Proof of Theorem 4(i). We will use Theorem E1 to prove Theorem 4. The main challenge is to show the assumptions of Theorem E1 are satisfied for all the $p!p$ variable selection problems (there are $p!$ orderings and each corresponds to p variable selection problems).

For every $\sigma \in \mathbb{S}^p$, we have an SEM representation for the distribution $N_p(0, \Sigma^*)$ given by

$$\mathbf{X} = (B_\sigma^*)^\top \mathbf{X} + \mathbf{e}_\sigma, \quad \mathbf{e}_\sigma \sim N_p(0, \Omega_\sigma^*).$$

where $(B_\sigma^*, \Omega_\sigma^*)$ is the modified Cholesky decomposition of Σ^* given in Definition 5. Denote the diagonal elements of Ω_σ^* by $\omega_{\sigma,1}^*, \dots, \omega_{\sigma,p}^*$. Using the data matrix, we can rewrite the SEM model as

$$X_j = \sum_{i \neq j} (B_\sigma^*)_{ij} X_i + \varepsilon_{\sigma,j}, \quad \varepsilon_{\sigma,j} \sim N_n(0, \omega_{\sigma,j}^* I). \quad (\text{F.4})$$

for $j = 1, \dots, p$. Note that the error vector $\varepsilon_{\sigma,j}$ depends on the permutation σ . Define the standardized error vector by

$$z_{\sigma,j} = (\omega_{\sigma,j}^*)^{-1/2} \varepsilon_{\sigma,j}, \quad \mathcal{Z} = \{z_{\sigma,j} : \sigma \in \mathbb{S}^p, j \in [p]\}. \quad (\text{F.5})$$

Let $\beta_{\sigma,j}^* = (B_\sigma^*)_{\text{Pa}_j(G_\sigma^*),j}$ be the subvector of the j -th column of (B_σ^*) with entries indexed by $S_{\sigma,j}^* = \text{Pa}_j(G_\sigma^*)$. As observed in Van de Geer and Bühlmann [72, Section 7.4.1], $\beta_{\sigma,j}^*$ and $\varepsilon_{\sigma,j}$ only depend on the set $S_{\sigma,j}^*$; see also Aragam et al. [5, Proposition 8.5]. Since the maximum degree of G_σ^* is bounded by d^* , the number of possible parent sets for any node is at most p^{d^*} and thus

$$|\mathcal{Z}| \leq p \cdot p^{d^*} = p^{d^*+1}. \quad (\text{F.6})$$

Let $\mathbb{V}_{\sigma,j}$ denote the variable selection problem with response variable X_j , set of candidate predictor variables $\{X_i : i \in \mathcal{A}_p^\sigma(j)\}$ and true data-generating model given in (F.4) parameterized by $(S_{\sigma,j}^*, \beta_{\sigma,j}^*, \omega_{\sigma,j}^*, \varepsilon_{\sigma,j})$. Recall that $\mathcal{A}_p^\sigma(j)$ is the index set of variables that precede X_j in the permutation σ and thus $S_{\sigma,j}^* \subseteq \mathcal{A}_p^\sigma(j)$. For $\mathbb{V}_{\sigma,j}$ with arbitrary $\sigma \in \mathbb{S}^p$ and $j \in [p]$, Assumptions (E1.3), (E1.4) and (E1.5) used in Theorem E1 directly follow from Assumptions (A3), (A4.2) and (A5.2), respectively. Further, by Remark 8, we always have $\omega_{\sigma,j}^* \in (\underline{\nu}, \bar{\nu})$ when Assumption (A1) holds. For sufficiently large n , by Lemmas F1 and F2 that we prove below using Assumptions (A1) and (A2), Assumptions (E1.1) and (E1.2) in Theorem E1 hold for all $p!p$ variable selection problems in the set $\{\mathbb{V}_{\sigma,j} : \sigma \in \mathbb{S}^p, j \in [p]\}$ with $\rho = 4d_{\text{in}} + 6$ and probability at least $1 - 3p^{-1}$, where ρ is as given in Assumption (E1.2). The claim then follows from Theorem E1. \square

Lemma F1. *If Assumption (A1) holds and $d_{\text{in}} \log p = o(n)$, then for sufficiently large n ,*

$$\mathbb{P}^* \left\{ n\underline{\nu} \leq \min_{S \in \mathcal{M}_p(2d_{\text{in}})} \lambda_{\min}(X_S^\top X_S) \leq \max_{S \in \mathcal{M}_p(2d_{\text{in}})} \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu} \right\} \geq 1 - 2e^{-n\delta_0^2/16},$$

where $\mathcal{M}_p(2d_{\text{in}}) = \{S \subseteq [p] : |S| \leq 2d_{\text{in}}\}$.

Proof. For any $S \subseteq [p]$, let $A_S = (\Sigma_S^*)^{-1/2} X_S^\top X_S (\Sigma_S^*)^{-1/2}$ where Σ_S^* denotes the $|S| \times |S|$ submatrix of Σ^* with rows/columns indexed by S . By Rudelson and Vershynin [61, Theorem 2.6], for any S and $a > 0$, we have

$$\mathbb{P}^* \left\{ (\sqrt{n} - \sqrt{|S|} - a)^2 \leq \lambda_{\min}(A_S) \leq \lambda_{\max}(A_S) \leq (\sqrt{n} + \sqrt{|S|} + a)^2 \right\} \geq 1 - 2e^{-a^2/2}.$$

By the submultiplicative property of operator norms,

$$\lambda_{\min}(A_S) \lambda_{\min}(\Sigma_S^*) \leq \lambda_{\min}(X_S^\top X_S), \quad \lambda_{\max}(A_S) \lambda_{\max}(\Sigma_S^*) \geq \lambda_{\max}(X_S^\top X_S).$$

Choose $a = \delta_0 \sqrt{n}/2$ and $n \geq 8d_{\text{in}}/\delta_0^2$; the latter is allowed since $d_{\text{in}} = o(n)$. Using Assumption (A1), we find that

$$\mathbb{P}^* \left\{ n\underline{\nu} \leq \lambda_{\min}(X_S^\top X_S) \leq \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu} \right\} \geq 1 - 2e^{-a^2/2}.$$

Observe that $\{S \subseteq [p]: |S| \leq 2d_{\text{in}}\}$ contains less than $p^{2d_{\text{in}}}$ elements. By the union bound and the assumption that $d_{\text{in}} \log p = o(n)$, $n\underline{\nu} \leq \lambda_{\min}(X_S^\top X_S) \leq \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu}$ holds for all $S \in \mathcal{M}_p(2d_{\text{in}})$ with probability at least $1 - 2e^{-n\delta_0^2/8+o(n)}$, from which the claim follows. \square

Lemma F2. Suppose Assumption (A2) holds and $d^* \leq d_{\text{in}}$. Let $z_{\sigma,j}$ be as defined in (F.5). Then, for sufficiently large n ,

$$\begin{aligned} \mathbb{P}^* \left\{ \min_{\sigma \in \mathbb{S}^p} \min_{j \in [p]} \min_{S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} z_{\sigma,j}^\top \Phi_S^\perp z_{\sigma,j} \leq n/2 \right\} &\leq e^{-n/96}, \\ \mathbb{P}^* \left\{ \max_{\sigma \in \mathbb{S}^p} \max_{j \in [p]} \max_{S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} \max_{k \in \mathcal{A}_p^\sigma(j)} z_{\sigma,j}^\top (\Phi_{S \cup \{k\}} - \Phi_S) z_{\sigma,j} \geq (4d_{\text{in}} + 6) \log p \right\} &\leq 2p^{-1}. \end{aligned}$$

Proof. Let $z \sim N_n(0, I)$. Given a fixed projection matrix Φ_S^\perp , we have $z^\top \Phi_S^\perp z \sim \chi_{n-|S|}^2$. By Laurent and Massart [37, Lemma 1], for any $a > 0$,

$$\mathbb{P} \left\{ \frac{z^\top \Phi_S^\perp z}{(n - |S|)} \leq 1 - a \right\} \leq e^{-(n-|S|)a^2/4}.$$

Choosing $a = 1/3$ and sufficiently large n so that $|S| \leq d_{\text{in}} \leq n/4$, we obtain that

$$\mathbb{P} \left\{ z^\top \Phi_S^\perp z \leq n/2 \right\} \leq e^{-n/48}.$$

Observe that for any $S \subseteq \mathcal{A}_p^\sigma(j)$, $z_{\sigma,j}$ is independent of X_S under the probability measure \mathbb{P}^* . Hence, we can treat Φ_S^\perp as fixed and apply the union bound and (F.6) to get

$$\mathbb{P}^* \left\{ \min_{\sigma \in \mathbb{S}^p} \min_{j \in [p]} \min_{S \subseteq \mathcal{A}_p^\sigma(j): |S| \leq d_{\text{in}}} z_{\sigma,j}^\top \Phi_S^\perp z_{\sigma,j} \leq \frac{n}{2} \right\} \leq p^{d_{\text{in}}+d^*+1} e^{-n/48}.$$

The first asserted inequality then follows from the assumptions $d_{\text{in}} \log p = o(n)$ and $d^* \leq d_{\text{in}}$.

For any $S \subseteq [p]$ and $k \notin S$, $\Phi_{S \cup \{k\}} - \Phi_S$ is another projection matrix with rank 1. Hence, using a standard tail bound for Gaussian distribution, for any $a > 0$, we find that

$$\mathbb{P} \left\{ z^\top (\Phi_{S \cup \{k\}} - \Phi_S) z \geq a \log p \right\} \leq 2e^{-a \log p/2},$$

if $z \sim N_n(0, I)$ independently of $X_{S \cup \{k\}}$. Another application of the union bound then yields the second inequality. \square

Proof of Theorem 4(ii). For any $S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})$, define

$$\mathcal{N}_j^\sigma(S) = \{S' \in \mathcal{M}_p^\sigma(j, d_{\text{in}}): \exists k, l \in [p] \text{ s.t. } S' = (S \cup \{k\}) \setminus \{l\}\}.$$

Note that we allow $k \in S$ and $l \in [p] \setminus S$ so that $\mathcal{N}_j^\sigma(S)$ includes the models that can be obtained from S by an addition, deletion or swap. Observe that the neighborhood relation defined by \mathcal{N}_j^σ is symmetric, and $|\mathcal{N}_j^\sigma(S)| \leq 1 + p + (p - d_{\text{in}})d_{\text{in}} \leq p^2$ (if $p \geq 2$) for each $S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})$. By part (i), the triple $(\mathcal{M}_p^\sigma(j, d_{\text{in}}), \mathcal{N}_j^\sigma, e^{\psi_j})$ satisfies Condition 1 with $t_1 = 2$, $t_2 = t$ and g_j^σ being the canonical transition function. The result then follows from Theorem 1(ii). \square

Proof of Theorem 4(iii). It suffices to prove the claim for $d_{\text{out}} = p$. Observe that

$$\sum_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, p)} e^{\psi(G)} = \sum_{G: \text{Pa}_j(G) \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} \prod_{j=1}^p e^{\psi_j(\text{Pa}_j(G))} = \prod_{j=1}^p \sum_{S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} e^{\psi_j(S)}.$$

Thus, using part (ii), we find that

$$\frac{\sum_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, p)} e^{\psi(G)}}{e^{\psi(G_\sigma^*)}} = \prod_{j=1}^p \frac{\sum_{S \in \mathcal{M}_p^\sigma(j, d_{\text{in}})} e^{\psi_j(S)}}{e^{\psi_j(S_{\sigma, j}^*)}} \leq (1 - p^{-(t-2)})^{-p} \leq \frac{1}{1 - p^{-(t-3)}},$$

for every $\sigma \in \mathbb{S}^p$, from which the result follows. \square

F.4 Proof of Theorem 5

Proof. By Lemma 1, for any $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$,

$$|\mathcal{N}_{\text{ads}}(\mathcal{E})| \leq 3p(p-1)(d_{\text{in}} + d_{\text{out}})p^{t_0} \leq 3t_0p^{t_0+2} \log_2 p = o(p^{t_0+3}).$$

Let \mathcal{F} denote the event on which the conclusion of Theorem 4 holds, which happens with probability at least $1 - 3p^{-1}$. By Theorem 4(i) and Corollary 2, on \mathcal{F} , the triple $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, \pi_n)$ satisfies Condition 1 with $t_1 = t_0 + 3$ and $t_2 = t$. The results then follow from Theorem 1(i) and (ii). \square

F.5 Consistency results for sub-Gaussian random matrices

Now we generalize our consistency results by considering sub-Gaussian random matrices.

Theorem F1. *Let X be an $n \times p$ random matrix, of which each row is an i.i.d. copy of a p -dimensional sub-Gaussian random vector with mean zero, covariance matrix Σ^* and sub-Gaussian parameter bounded by a universal constant $C_{\text{sub}} \in (0, \infty)$. Suppose $N_p(0, \Sigma^*)$ is perfectly Markovian w.r.t. a DAG G^* . Let $(B_\sigma^*, \Omega_\sigma^*)$ be as given in Definition 5.*

(F1) *There exist $\underline{\nu}, \bar{\nu} > 0$ and a universal constant $\delta_0 > 0$ such that*

$$\frac{\underline{\nu}}{(1 - \delta_0)^2} \leq \lambda_{\min}(\Sigma^*) \leq \lambda_{\max}(\Sigma^*) \leq \frac{\bar{\nu}}{(1 + \delta_0)^2}.$$

(F2) *The in-degree parameter d_{in} and n, p satisfy that $d_{\text{in}} \log p = o(n)$.*

(F3) *Prior parameters satisfy that $\alpha \in (0, 1]$, $\kappa \leq n$, $c_1 \sqrt{1 + \alpha/\gamma} \in [1, p]$, and $c_2 \geq C_{\text{pen}} d_{\text{in}} \bar{\nu}^4 / \underline{\nu}^6$ for some universal constant $C_{\text{pen}} > 0$.*

(F4) *$(\nu_0 + 1) \max_{\sigma \in \mathbb{S}^p} \max_{j \in [p]} |\text{Pa}_j(G_\sigma^*)| \leq d_{\text{in}}$ where $\nu_0 = 20\bar{\nu}^2 \underline{\nu}^{-4} (\bar{\nu} - \underline{\nu})^2$.*

(F5) *There exists a universal constant $C_\beta > 0$ such that*

$$\min_{\sigma \in \mathbb{S}^p} \min_{j \in [p]} |(B_\sigma^*)_{ij} : (B_\sigma^*)_{ij} \neq 0| \geq C_\beta \frac{c_2 \bar{\nu}^2 \log p}{\alpha \underline{\nu}^2 n}.$$

Let $t > 0$ be an arbitrary universal constant, and choose sufficiently large C_β, C_{pen} . For sufficiently large n , with probability at least $1 - 3p^{-d_{\text{in}}}$,

$$\min \{ \psi_j(g_j^\sigma(S)) - \psi_j(S) : \sigma \in \mathbb{S}^p, j \in [p], S \in \mathcal{M}_p^\sigma(j, d_{\text{in}}) \setminus \{S_{\sigma,j}^*\} \} \geq t \log p,$$

where $S_{\sigma,j}^* = \{i : (B_\sigma^*)_{ij} \neq 0\}$, ψ_j is given by (15) and $\{g_j^\sigma : j \in [p], \sigma \in \mathbb{S}^p\}$ is given by Definition 4. That is, the conclusion of part (i) of Theorem 4 holds.

Remark F1. The assumption that $N_p(0, \Sigma^*)$ is perfectly Markovian w.r.t. G^* implies that B_σ^* is the weighted adjacency matrix of G_σ^* , the minimal I-map of G^* . Observe that the scoring criterion we use only depends on the data via the sample covariance matrix $n^{-1}X^\top X$, which explains why we impose the assumption on Σ^* instead of the actual CI relations among the variables. One may also consider a DAG-perfect SEM model with sub-Gaussian errors as in Nandy et al. [48], in which case the distribution of \mathbf{X} is perfectly Markovian w.r.t. G^* .

Proof. We will use Theorem E2. Let Σ_σ^* denote the permuted true covariance matrix such that $(\Sigma_\sigma^*)_{ij} = \text{Cov}(\mathbf{X}_{\sigma(i)}, \mathbf{X}_{\sigma(j)})$. As in the proof of Theorem 4, let $\mathbb{V}_{\sigma,j}$ denote the variable selection problem with response variable \mathbf{X}_j and set of candidate predictor variables $\{\mathbf{X}_i : i \in \mathcal{A}_p^\sigma(j)\}$. The true covariance matrix associated with the problem $\mathbb{V}_{\sigma,j}$ is given by $(\Sigma_\sigma^*)_{[m+1]}$ where $m = |\mathcal{A}_p^\sigma(j)|$, which enables us to express \mathbf{X}_j by

$$\mathbf{X}_j = \sum_{i \neq j} (B_\sigma^*)_{ij} \mathbf{X}_i + \mathbf{e}_{\sigma,j},$$

where $\mathbf{e}_{\sigma,j}$ is a random variable uncorrelated (but not necessarily independent) with $\{\mathbf{X}_i : i \in \mathcal{A}_p^\sigma(j)\}$. It only remains to verify the assumptions used in Theorem E2.

By Lemma F3 we prove below, under Assumptions (F1) and (F2), for sufficiently large n , Assumptions (E2.1) and (E2.2) hold for all problems in $\{\mathbb{V}_{\sigma,j} : \sigma \in \mathbb{S}^p, j \in [p]\}$ with probability at least $1 - 3p^{-d_{\text{in}}}$, and the constant C_{err} in Assumption (E2.2) only depends on C_{sub} (and does not depend on σ or j); denote this event set by \mathcal{F}_n . Assumptions (E2.3), (E2.4) and (E2.5) directly follow from Assumptions (F3), (F4) and (F5), respectively. Therefore, given any $t > 0$, on the event \mathcal{F}_n we can apply Theorem E2 to all problems in $\{\mathbb{V}_{\sigma,j} : \sigma \in \mathbb{S}^p, j \in [p]\}$, which means that for any σ, j , there exist sufficiently large C_{pen}, C_β such that

$$\min \{ \psi_j(g_j^\sigma(S)) - \psi_j(S) : S \in \mathcal{M}_p^\sigma(j, d_{\text{in}}) \setminus \{S_{\sigma,j}^*\} \} \geq t \log p.$$

Further, by Lemma F3 and Theorem E2, the choice of C_{pen}, C_β only depends on universal constants t and C_{sub} . Hence, we can choose C_{pen}, C_β such that the above bound holds simultaneously for all $\sigma \in \mathbb{S}^p$ and $j \in [p]$. \square

Lemma F3. Suppose Assumption (F1) holds and $d_{\text{in}} \log p = o(n)$. Then there exist universal constants $K_0, C_0 \in (0, \infty)$, which only depend on C_{sub} , such that, for sufficiently large n , with probability at least $1 - 2e^{-C_0 n \delta_0^2 / 16} - 2p^{-d_{\text{in}}}$,

$$\begin{aligned} \max_{S \in \mathcal{M}_p(2d_{\text{in}})} \|n^{-1}X_S^\top X_S - \Sigma_S^*\|_{\text{op}} &\leq K_0 \sqrt{\frac{d_{\text{in}} \log p}{n}}, \text{ and} \\ n\underline{\nu} &\leq \min_{S \in \mathcal{M}_p(2d_{\text{in}})} \lambda_{\min}(X_S^\top X_S) \leq \max_{S \in \mathcal{M}_p(2d_{\text{in}})} \lambda_{\max}(X_S^\top X_S) \leq n\bar{\nu}, \end{aligned}$$

where $\mathcal{M}_p(2d_{\text{in}}) = \{S \subseteq [p] : |S| \leq 2d_{\text{in}}\}$.

Proof. By Vershynin [74, Remark 5.40], for any $a > 0$ and $S \subseteq [p]$, the following inequality holds with probability at least $1 - 2e^{-C_0 a^2}$,

$$\|n^{-1}X_S^\top X_S - \Sigma_S^*\|_{\text{op}} \leq \max\{u, u^2\}, \quad \text{where } u = C_1 \sqrt{\frac{|S|}{n}} + \frac{a}{\sqrt{n}},$$

and $C_0, C_1 > 0$ are universal constants. Choose $a = (K_0/2)\sqrt{d_{\text{in}} \log p}$ for some sufficiently large universal constant K_0 . Observe that $|\{S \subseteq [p] : |S| \leq 2d_{\text{in}}\}| < p^{2d_{\text{in}}}$. Hence, the assumption $d_{\text{in}} \log p = o(n)$ and a union bound yields that, for sufficiently large n ,

$$\mathbb{P}^* \left\{ \max_{S \in \mathcal{M}_p(2d_{\text{in}})} \|n^{-1}X_S^\top X_S - \Sigma_S^*\|_{\text{op}} \leq K_0 \sqrt{\frac{d_{\text{in}} \log p}{n}} \right\} \geq 1 - 2p^{-d_{\text{in}}},$$

where we have assumed again K_0 is sufficiently large.

To prove the second part, define $A_S = (\Sigma_S^*)^{-1/2} X_S^\top X_S (\Sigma_S^*)^{-1/2}$. By Vershynin [74, Theorem 5.39], for any $a > 0$, with probability at least $1 - 2e^{-C_0 a^2}$,

$$(\sqrt{n} - C_1 \sqrt{|S|} - a)^2 \leq \lambda_{\min}(A_S) \leq \lambda_{\max}(A_S) \leq (\sqrt{n} + C_1 \sqrt{|S|} + a)^2.$$

The rest is analogous to the proof of Lemma F1. □

G Proofs and more examples for Sections 5 and 7

G.1 Proof of Theorem 6

Proof of Theorem 6. In the proof of Theorem 5, we have shown that, with probability at least $1 - 3p^{-1}$, the triple $(\mathcal{C}_p(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, e^\psi)$ satisfies Condition 1 for

$$p^{t_1} = 3t_0 p^{t_0+2} \log_2 p, \quad t_2 = t.$$

The mixing time bound for RW-GES then follows from Theorem 1(iv) and Theorem 2. \square

G.2 Proof of Corollary 4

Proof. Define $c_3 = c_1 \sqrt{1 + \alpha/\gamma}$. By (15), for any $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$, we have

$$\begin{aligned} \frac{e^{\psi(G)}}{e^{\psi(G_\sigma^*)}} &= \prod_{j=1}^p \frac{e^{\psi_j(\text{Pa}_j(G))}}{e^{\psi_j(\text{Pa}_j(G_\sigma^*))}} = (c_3 p^{c_2})^{|G_\sigma^*| - |G|} \prod_{j=1}^p \left(\frac{X_j^\top \Phi_{\text{Pa}_j(G)}^\perp X_j}{\varepsilon_{\sigma,j}^\top \Phi_{\text{Pa}_j(G_\sigma^*)}^\perp \varepsilon_{\sigma,j}} \right)^{-(\alpha n + \kappa)/2} \\ &\geq (c_3 p^{c_2})^{-pd_{\text{in}}} \prod_{j=1}^p \left(\frac{X_j^\top X_j}{\varepsilon_{\sigma,j}^\top \Phi_{\text{Pa}_j(G_\sigma^*)}^\perp \varepsilon_{\sigma,j}} \right)^{-(\alpha n + \kappa)/2} \\ &\geq (c_3 p^{c_2})^{-pd_{\text{in}}} \prod_{j=1}^p \left(\frac{n\bar{\nu}}{n\omega_{\sigma,j}^*/2} \right)^{-(\alpha n + \kappa)/2}, \end{aligned}$$

where the last step follows from Lemmas F1 and F2. By Remark 8, $\omega_{\sigma,j}^* \geq \underline{\nu}$, which yields

$$\min_{G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})} \frac{\exp(\psi(G))}{\exp(\psi(G_\sigma^*))} \geq (c_3 p^{c_2})^{-pd_{\text{in}}} \left(\frac{2\bar{\nu}}{\underline{\nu}} \right)^{-p(\alpha n + \kappa)/2}, \quad \forall \sigma \in \mathbb{S}^p. \quad (\text{G.1})$$

By Lemma D3, there exists a Chickering sequence $(G_0 = G_\sigma^*, G_1, \dots, G_k = G^*)$ for some $k \leq pd^*$ such that, for $i \in [k]$, G_i is obtained from G_{i-1} by covered edge reversals and a single edge deletion. Since by removing any single edge from a DAG in \mathcal{G}_p^τ , its posterior score can increase by at most $c_3 p^{c_2}$, we have

$$\frac{\exp(\psi([G_\sigma^*]))}{\exp(\psi(\mathcal{E}^*))} \geq (c_3 p^{c_2})^{-pd^*}. \quad (\text{G.2})$$

For any $\mathcal{E} \in \mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, there exists a pair (G, σ) such that $G \in \mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}})$. Thus,

$$\frac{\pi_n(\mathcal{E})}{\pi_n(\mathcal{E}^*)} = \frac{e^{\psi([G_\sigma^*])}}{e^{\psi(\mathcal{E}^*)}} \frac{e^{\psi(G)}}{e^{\psi(G_\sigma^*)}},$$

Combining (G.1) and (G.2), we obtain the asserted bound on $\pi_n(\mathcal{E})/\pi_n(\mathcal{E}^*)$. Under the setting of Theorem 5, we have the strong selection consistency, which implies that $\pi_n(\mathcal{E}^*) \geq 1/2$ for all sufficiently large n . Hence, $\log \pi_n(\mathcal{E})$ can be bounded a polynomial of n and p . This completes the rapid mixing proof for RW-GES. \square

G.3 Proof of Theorem 7

Proof. By Theorem 4(i) and Corollary 1, with probability at least $1 - 3p^{-1}$, the triple $(\mathcal{G}_p^\sigma(d_{\text{in}}, d_{\text{out}}), \mathcal{N}_{\text{ads}}, \pi_n^\sigma)$ satisfies Condition 1 for $p^{t_1} = d_{\text{in}} p^2/2$ and $t_2 = t$. (Note that though Corollary 1 used $t_1 = 3$, by (D.4) the neighborhood size actually can be bounded by $d_{\text{in}} p^2/2$.) The mixing time bound then follows from Theorem 1(iv) and Theorem 2. \square




Figure G1: DAG models with three vertices. Each DAG represents a unique equivalence class. Any other 3-vertex DAG model is Markov equivalent to one of these DAGs. For Example 4, the true DAG is $G^* = G_4$ and the local mode is the equivalence class of $\tilde{G} = G_7$. For Example G1, $G^* = G_9$ and $\tilde{G} = G_{10}$.

G.4 Proof of Example 4

Proof. Consider the space $\mathcal{G}_p(2, 2)$ for $p = 3$, the collection of all 3-vertex DAG models. By Andersson et al. [4], there are 11 labeled equivalence classes, which we show in Figure G1 (for each equivalence class we plot one DAG member). The true DAG is given by $G^* = G_4$ and the local mode is the equivalence class that contains $\tilde{G} = G_7$. Consider $\sigma = (1, 3, 2)$, which is a topological ordering of \tilde{G} . The corresponding SEM representation can be written as

$$\begin{aligned} X_1 &= z_1, \\ X_3 &= b_1 b_2 X_1 + b_2 z_2 + z_3, \\ X_2 &= \frac{b_1}{b_2^2 + 1} X_1 + \frac{b_2}{b_2^2 + 1} X_3 + \frac{1}{b_2^2 + 1} z_2 - \frac{b_2}{b_2^2 + 1} z_3. \end{aligned}$$

We prove the slow mixing by verifying the two conditions in Theorem B3:

1. $\mathbf{P}([G_7], [G_7]) \geq 1 - e^{-c\sqrt{n}}$ for some universal constant c .
2. $\pi_n([G_7]) \leq 1/2$ for all sufficiently large n .

Consider (i) first. Since $[G_7] = \{G_7\}$, the set $\mathcal{N}_{\text{ads}}([G_7])$ is determined by all the neighbors of G_7 . Observe that $G_1, G_2 \in \mathcal{N}_{\text{del}}(G_7)$ and $G_{10} \in \mathcal{N}_{\text{add}}(G_7)$. It can be shown that for $K \geq 4$ and sufficiently large n ,

$$\begin{aligned} \frac{\pi_n([G_7])}{\pi_n([G_1])} &= \exp \left\{ -c_2 \log p + \frac{\alpha n}{2} \log(1 + b_2^2) \right\} \geq p^{c_2/2}, \\ \frac{\pi_n([G_7])}{\pi_n([G_2])} &= \exp \left\{ -c_2 \log p + \frac{\alpha n}{2} \log \frac{(b_1^2 + 1)(b_2^2 + 1)}{b_1^2 b_2^2 + b_2^2 + 1} \right\} \geq p^{c_2/2}, \\ \frac{\pi_n([G_7])}{\pi_n([G_{10}])} &= \exp \left\{ c_2 \log p - \frac{\alpha n}{2} \log \frac{b_1^2 b_2^2 + b_2^2 + 1}{b_2^2 + 1} \right\} \geq p^{c_2/2}. \end{aligned}$$

To prove the first, we use $b_1^2 = b_2^2 = o(1)$ and $\log(1 + x) \sim x$ as $x \downarrow 0$ to get $\alpha n \log(1 + b_2^2) \sim K c_2 \log p$. Hence, if $K \geq 4$ and n is sufficiently large, $\alpha n \log(1 + b_2^2) \geq 3 c_2 \log p$. The proof

of the second inequality is similar. For the last one, we use $b_1^2 b_2^2 = o(b_1^2)$ to show that $c_2 \log p$ has a larger order than the other term in the exponent. Since $c_2 = \sqrt{n}$, by the Metropolis rule, for any $\mathcal{E}' \neq [G_7]$,

$$\mathbf{P}([G_7], \mathcal{E}') = \mathbf{K}([G_7], \mathcal{E}') \min \left\{ 1, \frac{\pi_n(\mathcal{E}') \mathbf{K}(\mathcal{E}', [G_7])}{\pi_n([G_7]) \mathbf{K}([G_7], \mathcal{E}')} \right\} \leq \frac{\pi_n(\mathcal{E}')}{\pi_n([G_7])} < p^{-\sqrt{n}/2}.$$

It then follows that (i) holds since $|\mathcal{N}_{\text{ads}}([G_7])|$ is bounded. Note that if p goes to infinity, we can still use $|\mathcal{N}_{\text{ads}}([G_7])| \leq 3p^2$ to show (i).

To prove (ii), we only need to compare G_7 with the true model G_4 . Another routine calculation yields that, for large n ,

$$\begin{aligned} \frac{\pi_n([G_4])}{\pi_n([G_7])} &= \exp \left\{ \frac{\alpha n}{2} \log \frac{b_1^2 b_2^2 + b_2^2 + 1}{b_2^2 + 1} \right\} \\ &\geq \exp \left\{ \frac{\alpha n}{4} b_1^2 b_2^2 \right\} = \exp \left\{ \frac{K^2 (\log p)^2}{4\alpha} \right\}. \end{aligned}$$

Since $p = 3$, as long as we choose K such that $K^2 \geq 4/\alpha$, we have $\pi_n([G_4])/\pi_n([G_7]) \geq 3^{\log 3} \approx 3.34$, from which (ii) follows. \square

G.5 Slow mixing examples for a CPDAG sampler

He et al. [28, Definition 9] constructed a “perfect” set of CPDAG operators: insert/delete an undirected edge, insert/delete a direct edge, and make/remove a v-structure; see Supplement C.3 for details. This set of CPDAG operators is said to be reversible and irreducible, which means that the induced neighborhood relation on \mathcal{C}_p is symmetric and neighborhood graph is connected (i.e., a random walk using this set of operators is irreducible). Let $\mathcal{N}_{\mathcal{C}}$ denote the corresponding neighborhood function. A random walk MH algorithm can be constructed by proposing a state in $\mathcal{N}_{\mathcal{C}}(\cdot)$ uniformly at random. This algorithm was implemented in Castelletti et al. [11]. Hence, compared with RW-GES, the only difference is that the neighborhood $\mathcal{N}_{\text{ads}}(\cdot)$ is replaced by $\mathcal{N}_{\mathcal{C}}(\cdot)$. Unfortunately, this “reversible” CPDAG sampler can be slowly mixing even if the strong beta-min condition holds.

Example G1. This is a more explicit construction of Example 5. Let $p = 3$ and n tend to infinity. The extension to the case $p = n$ is straightforward. Let the true SEM model be given by

$$X_1 = z_1, \quad X_2 = z_2, \quad X_3 = a_1 X_1 + a_2 X_2 + z_3,$$

where z_1, z_2, z_3 are as given in Example 4 but the coefficients $a_1, a_2 > 0$ are assumed to be fixed. Hence, the true DAG model G^* is $1 \rightarrow 3 \leftarrow 2$, which is a CPDAG itself. Choose prior parameters $c_1, c_2, \kappa, \alpha, \gamma$ as in Example 4. Let $\tilde{\mathcal{E}}$ be the equivalence class that contains all complete DAGs; $\text{EG}(\tilde{\mathcal{E}})$ is a complete undirected graph. The set $\mathcal{N}_{\mathcal{C}}(\tilde{\mathcal{E}})$ contains three equivalence classes represented by CPDAGs of the form $i - j - k$, but $\mathcal{E}^* \notin \mathcal{N}_{\mathcal{C}}(\tilde{\mathcal{E}})$. Below we prove that the random walk MH algorithm equipped with the neighborhood function $\mathcal{N}_{\mathcal{C}}$ is slowly mixing since it can get stuck at $\tilde{\mathcal{E}}$ for exponentially many steps.

Proof. We still use the numbering in Figure G1. The true DAG is $G^* = G_9$, and the local mode we consider is the equivalence class generated by $\tilde{G} = G_{10}$. By He et al. [28, Definition 9], $\mathcal{N}_C([G_{10}]) = \{[G_4], [G_5], [G_6]\}$. Since p, a_1, a_2, α are fixed constants and $c_2 = \sqrt{n}$, for sufficiently large n , we find that

$$\begin{aligned}\frac{\pi_n([G_{10}])}{\pi_n([G_9])} &= p^{-c_2} = p^{-\sqrt{n}}, \\ \frac{\pi_n([G_{10}])}{\pi_n([G_4])} &= \exp \left\{ -c_2 \log p + \frac{\alpha n}{2} \log(a_1^2 + 1) \right\} \geq e^{cn}, \\ \frac{\pi_n([G_{10}])}{\pi_n([G_5])} &= \exp \left\{ -c_2 \log p + \frac{\alpha n}{2} \log(a_2^2 + 1) \right\} \geq e^{cn}, \\ \frac{\pi_n([G_{10}])}{\pi_n([G_6])} &= \exp \left\{ -c_2 \log p + \frac{\alpha n}{2} \log \frac{(a_1^2 + 1)(a_2^2 + 1)}{a_1^2 + a_2^2 + 1} \right\} \geq e^{cn},\end{aligned}$$

for some universal constant $c > 0$. Thus, for the random walk MH algorithm using neighborhood relation \mathcal{N}_C , we have $\mathbf{P}([G_{10}], [G_{10}]) \geq 1 - 3e^{-cn}$. Since $[G_{10}]$ has negligible posterior probability, the chain is slowly mixing by Theorem B3. \square

We give another more complicated example with 5 nodes.

Example G2. Let $p = 5$ and the true DAG G^* be as given in Figure G2. Consider the DAG $H = G \cup \{1 \rightarrow 4\}$. By (dd2) in Definition 9 of He et al. [28], $[G^*] \notin \mathcal{N}_C([H])$. Note that since \mathcal{N}_C defines a symmetric relation, this is equivalent to claiming that we cannot move from $\text{EG}(G^*)$ to $\text{EG}(H)$, which is easy to prove: if we add $1 \rightarrow 4$ to $\text{EG}(G^*)$, all consistent extensions of the resulting PDAG cannot have the v-structures $1 \rightarrow 4 \leftarrow 3$ and $1 \rightarrow 4 \leftarrow 2$, which exist in $\text{EG}(H)$. Actually, according to He et al. [28, Definition 9], there are only 8 possible operations that we may apply to the CPDAG of H , which we list in Table G1. Each operation uniquely defines a resulting CPDAG and thus $|\mathcal{N}_C([H])| = 8$. In Figure G3, we plot a member DAG for each equivalence class in $\mathcal{N}_C([H])$.

If ψ satisfies Condition 2 (local consistency), one can verify that $[H]$ has a larger posterior probability than all 8 neighboring equivalence classes. First, since H is an I-map of G^* , H_1, H_2, H_3 should all have smaller scores. For $j = 4, \dots, 8$, observe that H_j is obtained from H by removing an edge between two nodes that are not conditionally independent given any set of other nodes. Hence, $[H]$ is a local mode on $(\mathcal{C}_p, \mathcal{N}_C)$.

On the restricted space $\mathcal{C}_p(d_{\text{in}}, d_{\text{out}})$, the maximum size of $\mathcal{N}_C(\cdot)$ tends to be much smaller than that of $\mathcal{N}_{\text{ads}}(\cdot)$ since the former can only grow polynomially in $d_{\text{in}} + d_{\text{out}}$. However, for two different DAGs G, G' such that $G' = G \cup \{i \rightarrow j\}$, we may have $[G] \notin \mathcal{N}_C([G'])$ because removing the edge between i and j from $\text{EG}(G')$ does not result in a unique CPDAG (more precisely, the modified graph has no consistent extension; see Supplement C.3 for details). If G happens to be the true DAG model, then G' is likely to be a local mode that can trap the algorithm for an enormous number of iterations, since other modifications of G' either yield a DAG that is not an I-map of G or a DAG with more “redundant” edges. Nevertheless, as originally proposed in He et al. [28], one can define a random walk on $(\mathcal{C}_p, \mathcal{N}_C)$ for efficiently sampling CPDAGs from the uniform distribution.




Figure G2: A slow mixing example for the random walk MH algorithm using neighborhood function $\mathcal{N}_{\mathcal{C}}$. G^* : the true DAG; $[G^*]$: the CPDAG of G^* ; H : a DAG representing a local mode; $[H]$: the CPDAG of H .

Table G1: Edge operations that may be applied to the CPDAG of H in Figure G2.

Operator	Related edge(s)
InsertU (insert an undirected edge)	$1 - 2, 1 - 3, 4 - 5$
DeleteU (delete an undirected edge)	$2 - 3$
InsertD (insert a directed edge)	None
DeleteD (delete a directed edge)	$3 \rightarrow 4, 3 \rightarrow 5, 2 \rightarrow 4, 2 \rightarrow 5$
MakeV (make a v-structure)	None
RemoveV (remove a v-structure)	None




Figure G3: Characterization of $\mathcal{N}_C([H])$ where H is as given in Figure G2. For each $\mathcal{E} \in \mathcal{N}_C([H])$, we plot a member DAG of \mathcal{E} . The 8 DAGs correspond to the 8 operations listed in Table G1. H_1 : insert $1 - 2$; H_2 : insert $1 - 3$; H_3 : insert $4-5$; H_4 : delete $2 - 3$; H_5 : delete $3 \rightarrow 4$; H_6 : delete $3 \rightarrow 5$; H_7 : delete $2 \rightarrow 4$; H_8 : delete $2 \rightarrow 5$.

G.6 An example for comparing RW-GES and structure MCMC

Consider the true data-generating DAG G^* and another DAG G_0 shown in Figure G4 and assume $n = \infty$. Observe that G_0 is a minimal I-map of G^* . Define $G_1 = G^* \cup \{2 \rightarrow 1\}$, which is Markov equivalent to G_0 . Let $\mathcal{E}^* = [G^*]$ and $\mathcal{E}_0 = [G_0]$. For ease of presentation, we assume the following condition on ψ holds, which is similar to but stronger than Condition 2: for any distinct DAGs G, G' such that $G' = G \cup \{i \rightarrow j\}$,

- (i) $\psi(G) - \psi(G') = \infty$ if $i \perp\!\!\!\perp j \mid \text{Pa}_j(G)$ in G^* ;
- (ii) $\psi(G') - \psi(G) = \infty$ if $i \not\perp\!\!\!\perp j \mid \text{Pa}_j(G)$ in G^* .

If p and the true data-generating model (perfectly Markovian w.r.t. G) are fixed and we let $n \rightarrow \infty$, this condition should hold for a large class of scoring criteria [15]. We make this assumption so that if an MH algorithm proposes to add $i \rightarrow j$ to G but $i \perp\!\!\!\perp j \mid \text{Pa}_j(G)$ in




Figure G4: The DAGs considered in Supplement G.6. G^* : the true DAG. G_0 : a minimal I-map of G^* .

G^* , the proposal will be rejected with probability one. Further, we assume the search spaces are unrestricted, i.e., \mathcal{C}_p for RW-GES and \mathcal{G}_p for structure MCMC, and do not consider swap moves of RW-GES (but the same conclusion still holds if swap moves are used).

Analysis of RW-GES By the definition of minimal I-maps given in Section A, one can verify that G_0 is a minimal I-map of G^* with ordering $(p, p-1, \dots, 2, 1)$. Hence, it is straightforward to check that the only proposal in $\mathcal{N}_{\text{add}}(\mathcal{E}_0) \cup \mathcal{N}_{\text{del}}(\mathcal{E}_0)$ that will be accepted is \mathcal{E}^* . Since the maximum degree of G_0 is always 3 (which does not depend on p), analogously to the proof of Lemma 1, one can show that $|\mathcal{N}_{\text{add}}(\mathcal{E}_0) \cup \mathcal{N}_{\text{del}}(\mathcal{E}_0)| \leq 2^3 p^2 = 8p^2$. That is, on average it takes RW-GES less than $8p^2$ iterations to propose \mathcal{E}^* , which will always be accepted by our assumption on ψ .

Analysis of structure MCMC Structure MCMC is defined on the DAG space which uses single-edge addition, deletion and reversal as the proposal moves. For structure MCMC, to move from G_0 to G^* , we have to move to G_1 first, which requires the sampler to reverse all edges of G_0 except the edge $2 \rightarrow 1$. Moreover, these edges have to be reversed from right to left.

To show this, note that since G_0 is a minimal I-map, we cannot add or remove any edge of G_0 . We can only move from G_0 to DAGs in $\mathcal{N}_{\text{rev}}(G_0)$, the set of all DAGs that can be obtained from G_0 by reversing one edge. Reversing $2 \rightarrow 1$ yields a Markov equivalent DAG but has no effect (the resulting DAG is essentially the same as G_0 compared to G^*), so we will ignore the reversal of this edge henceforth. Reversing $3 \rightarrow 1$ results in a cycle and thus is not allowed. Reversing $3 \rightarrow 2$ results in the v-structure $2 \rightarrow 3 \leftarrow 4$, which does not exist in G^* ; thus, the resulting DAG is not an I-map of G^* and will be rejected by our assumption on ψ . Similarly, reversing $4 \rightarrow 3$ will also be rejected since it results in the v-structure $3 \rightarrow 4 \leftarrow 5$ (assuming $p \geq 5$). So the only edge we can reverse other than $2 \rightarrow 1$ is the edge $p \rightarrow (p-1)$. Repeating this argument, we see that structure MCMC has to reverse the edges $p \rightarrow (p-1)$, $(p-1) \rightarrow (p-2)$, \dots , $4 \rightarrow 3$ in order; denote the resulting DAG by G_2 (see Figure G5). Note that after the edge $p \rightarrow (p-1)$ is reversed, structure MCMC may reverse the edge $(p-1) \rightarrow p$ again and return to G_0 . Hence, moving from G_0 to G_2 usually takes much more than $p-3$ successful edge reversals. Indeed, this can be seen as a reflected symmetric random walk on $\{4, 5, \dots, p\}$, and by a standard result from random walk theory, the expected number of successful edge reversals needed to move from G_0 to G_2

is $(p - 3)^2$. Since in structure MCMC the neighborhood size of each DAG has order p^2 , we conclude that on average it takes structure MCMC $O(p^4)$ iterations to move from G_0 to G^* .




Figure G5: A likely path of structure MCMC from G_0 to G^* in the example considered in Supplement G.6. Edges that have just been reversed are shown in red.

H Implementation and simulation studies on RW-GES

H.1 Implementation of RW-GES

To implement the RW-GES sampler, we need to propose new states from $\mathcal{N}_{\text{ads}}(\mathcal{E})$ for each \mathcal{E} . Though \mathcal{N}_{ads} is constructed by applying add-delete-swap moves to all member DAGs in \mathcal{E} , we do not need to enumerate these DAGs to sample from $\mathcal{N}_{\text{ads}}(\mathcal{E})$. Instead, recall the “Insert” and “Delete” operators of the GES algorithm defined in Supplement C.3. They can be used to generate the sets $\mathcal{N}_{\text{add}}(\mathcal{E})$ and $\mathcal{N}_{\text{del}}(\mathcal{E})$, which only involve local modifications of the CPDAG $\text{EG}(\mathcal{E})$. Define

$$\begin{aligned}\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E}) &= \{\text{Insert}(i, j, S) : S \subseteq \text{Un}_j(\text{EG}(\mathcal{E})) \setminus \text{Ad}_i(\text{EG}(\mathcal{E}))\}, \\ \mathcal{O}_{ij}^{\text{del}}(\mathcal{E}) &= \{\text{Delete}(i, j, S) : S \subseteq \text{Un}_j(\text{EG}(\mathcal{E})) \cap \text{Ad}_i(\text{EG}(\mathcal{E}))\}.\end{aligned}$$

By Definition C3, $\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E})$ is the set of all insertion operators that add the edge $i \rightarrow j$ to $\text{EG}(\mathcal{E})$. By Definition C4, $\mathcal{O}_{ij}^{\text{del}}(\mathcal{E})$ is the set of all deletion operators that remove the edge $i \rightarrow j$ or $i - j$ from $\text{EG}(\mathcal{E})$. Further, as explained in Supplement C.3, if O is a valid operator in $\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E})$ that converts \mathcal{E} to \mathcal{E}' , then it can be matched with an operator $O' \in \mathcal{O}_{ij}^{\text{del}}(\mathcal{E}')$ that can convert \mathcal{E}' back to \mathcal{E} . Now we describe an efficient implementation of RW-GES.

Algorithm H1. Let \mathcal{E} denote the current equivalence class and $\text{EG}(\mathcal{E})$ be its CPDAG. Sample an ordered pair (i, j) uniformly from $[p]^2$ such that $i \neq j$.

(i) If i, j are adjacent in $\text{EG}(\mathcal{E})$, sample O uniformly from $\mathcal{O}_{ij}^{\text{del}}(\mathcal{E})$.

(ii) If i, j are not adjacent in $\text{EG}(\mathcal{E})$, sample O uniformly from $\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E})$.

Apply the operator O to $\text{EG}(\mathcal{E})$. If the resulting PDAG has no consistent extension, stay at \mathcal{E} . If the resulting PDAG has a consistent extension G' , propose moving to $\mathcal{E}' = [G']$ and accept it with probability

$$\text{case (i): } \min \left\{ 1, \frac{\pi_n(\mathcal{E}')}{\pi_n(\mathcal{E})} \frac{|\mathcal{O}_{ij}^{\text{del}}(\mathcal{E})|}{|\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E}')|} \right\}, \quad \text{case (ii): } \min \left\{ 1, \frac{\pi_n(\mathcal{E}')}{\pi_n(\mathcal{E})} \frac{|\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E})|}{|\mathcal{O}_{ij}^{\text{del}}(\mathcal{E}')|} \right\}.$$

To implement the node degree constraints, for simplicity, we require that the maximum degree $\leq d$ for some positive integer d (i.e., any node is adjacent to at most d nodes). Then, in case (ii) of Algorithm H1, if the degree of node j is equal to d but the degree of node i is less than d , we replace the addition with a swap proposal. Note that there is no need to check the degree constraint in case (i) of Algorithm H1. Under our degree constraint, it is easy to check that $|\mathcal{O}_{ij}^{\text{ins}}(\mathcal{E})| \leq 2^{d-1}$ and $|\mathcal{O}_{ij}^{\text{del}}(\mathcal{E})| \leq 2^{d-1}$. Hence, the probability of proposing any insertion or deletion operator is at least $p^{-2}2^{1-d}$, which is greater than or equal to $2p^{-(2+t_0)}$ if we assume $d \leq t_0 \log_2 p$. We refer readers to Chickering [15] for how to convert DAGs to CPDAGs and tricks that can further expedite CPDAG operations.

If we run the RW-GES sampler for N iterations (after burn-in), we get N equivalence classes $\mathcal{E}^{(1)}, \dots, \mathcal{E}^{(N)}$. For each $\mathcal{E}^{(k)}$, we convert it to the corresponding CPDAG $H^{(k)} = \text{EG}(\mathcal{E}^{(k)})$, and we check which edges are in $H^{(k)}$; $i \rightarrow j, i \leftarrow j$ and $i - j$ are treated as different edges. For each $i < j$, RW-GES outputs the frequencies of the edge $i \rightarrow j, j \rightarrow i$ and $i - j$ being in $H^{(1)}, H^{(2)}, \dots, H^{(N)}$; denote them by $\hat{P}(i \rightarrow j), \hat{P}(j \rightarrow i)$ and $\hat{P}(i - j)$, respectively.

H.2 Further details about simulation studies

First simulation study Figure H1 shows the CPDAG representing the equivalence class of the DAG G^* used in the simulation study presented in Section 6.1. The true covariance matrix Σ^* has $\lambda_{\max}(\Sigma^*) = 9.6$ and $\lambda_{\min}(\Sigma^*) = 0.15$. The simulated sample covariance matrix $\hat{\Sigma}_n = n^{-1}X^\top X$ has $\lambda_{\max}(\hat{\Sigma}_n) = 12.3$ and $\lambda_{\min}(\hat{\Sigma}_n) = 0.04$ for $n = 200$ and $\lambda_{\max}(\hat{\Sigma}_n) = 10.7$ and $\lambda_{\min}(\hat{\Sigma}_n) = 0.11$ for $n = 800$.

This simulation study also illustrates a typical way to construct a high-dimensional data set that satisfies the strong beta-min condition. Recall that this condition is quite restrictive mainly because we need (18) to hold for all $\sigma \in \mathbb{S}^p$ but \mathbb{S}^p grows super-exponentially with p . However, if the true DAG G^* can be partitioned into disconnected sub-DAGs, then the true covariance matrix Σ^* and its inverse are block diagonal (after row/column permutation), and the strong beta-min condition is much easier to satisfy. In particular, if nodes i, j belong to a maximal connected sub-DAG with k nodes, the mapping $\sigma \mapsto (B_\sigma^*)_{ij}$ can take at most 2^{k-1} distinct values (since the parent set of node j has at most 2^{k-1} possibilities). For the CPDAG shown in Figure H1, there is one maximal connected subDAG with 10 nodes, and the other connected subDAGs have at most 3 nodes. A more general and simpler way to construct examples that satisfy the assumptions of Theorem 6 is given below.

Example H1. Let q be a fixed positive integer and Σ_0 be a fixed $q \times q$ positive definite covariance matrix with $\lambda_{\min}(\Sigma_0) = 2\underline{\nu}$ and $\lambda_{\max}(\Sigma_0) = \bar{\nu}/2$, where $\underline{\nu}, \bar{\nu}$ are nonzero universal constants. Let $p = mq$ where $m = \lfloor \exp(n^\xi) \rfloor$ for some $\xi \in (0, 1)$. Let the true covariance matrix $\Sigma^* = \text{diag}(\Sigma_0, \dots, \Sigma_0)$ (i.e., a block diagonal matrix). Then, we have $\lambda_{\min}(\Sigma^*) = 2\underline{\nu}$ and $\lambda_{\max}(\Sigma^*) = \bar{\nu}/2$, which satisfies the restricted eigenvalue condition (A1). Further, the block diagonal structure of Σ^* implies that $d^* \leq q$ and

$$\beta_{\min} = \{|(B_\sigma^*)_{ij}| : (B_\sigma^*)_{ij} \neq 0, i, j \in [p], \sigma \in \mathbb{S}^p\}$$

is a universal nonzero constant (since it can be determined using Σ_0 .) Choose $d_{\text{in}} = \nu_0 q$ where $\nu_0 = 4\bar{\nu}^2(\bar{\nu} - \underline{\nu})^2/\underline{\nu}^4 + 1$ and $d_{\text{out}} = 2\nu_0 q^2$ so that $d^* d_{\text{in}} + 1 \leq d_{\text{out}}$ and Assumptions (A2) and (A4.2) hold. Since $d_{\text{in}}, d_{\text{out}}$ are fixed constants, we can assume that $d_{\text{in}} + d_{\text{out}} \leq \log_2 p$ for sufficiently large n . For the prior parameters, we choose $\alpha = 1/2$ and $c_2 = 6d_{\text{in}} + 9 + t$ with $t > 4$ so that Assumption (A3) holds. Finally, the strong beta-min condition is satisfied for sufficiently large n since β_{\min} is a fixed constant but the lower bound in (18) goes to zero. Note that the true underlying DAG model G^* can have at most $p(q-1)/2$ edges.

Second simulation study For the simulation study presented in Section 6.2, the median extreme eigenvalues of the true covariance matrix and sample covariance matrix are reported in Table H1.

The TPR and FPR statistics reported in Tables 1 and 2 are calculated as follows. Let $H^* = \text{EG}(G^*)$ be the CPDAG of the true equivalence class. Recall that we say i, j are




Figure H1: The true CPDAG used to generate the data set used in the first simulation study.

adjacent in H^* if they are connected by either a directed or undirected edge. Define

$$\begin{aligned} \text{TPs} &= \sum_{i < j} \mathbb{1}\{i, j \text{ are adjacent in } H^*\} \left\{ \hat{P}(i \rightarrow j) + \hat{P}(j \rightarrow i) + \hat{P}(i - j) \right\}, \\ \text{TP} &= \sum_{i < j} \left[\mathbb{1}\{i \rightarrow j \in H^*\} \hat{P}(i \rightarrow j) + \mathbb{1}\{j \rightarrow i \in H^*\} \hat{P}(j \rightarrow i) + \mathbb{1}\{i - j \in H^*\} \hat{P}(i - j) \right], \\ \text{FP} &= \sum_{i < j} \mathbb{1}\{i, j \text{ are not adjacent in } H^*\} \left\{ \hat{P}(i \rightarrow j) + \hat{P}(j \rightarrow i) + \hat{P}(i - j) \right\}. \end{aligned}$$

Then, “TPR (skeleton)” is defined as $\text{TPs}/|G^*|$, where $|G^*|$ is also the number of edges in H^* . TPR is defined as $\text{TP}/|G^*|$, and FPR is defined as $\text{FP}/\{p(p-1)/2 - |G^*|\}$.

p	n	$\lambda_{\max}(\Sigma^*)$	$\lambda_{\min}(\Sigma^*)$	$\lambda_{\max}(\hat{\Sigma}_n)$	$\lambda_{\min}(\hat{\Sigma}_n)$
7	60	7.7 (2.9–15)	0.17 (0.12–0.3)	6.8 (3.6–19)	0.14 (0.09–0.28)
14	90	13 (5.6–35)	0.15 (0.11–0.23)	14 (6.8–43)	0.11 (0.07–0.2)
28	120	30 (16–102)	0.10 (0.08–0.13)	33 (16–113)	0.06 (0.03–0.09)
56	150	75 (43–188)	0.09 (0.07–0.11)	78 (41–173)	0.04 (0.04–0.05)
112	180	149 (73–348)	0.08 (0.06–0.09)	152 (75–353)	0.02 (0.01–0.02)
224	210	252 (146–631)	0.07 (0.06–0.08)	239 (148–633)	0
448	240	411 (251–844)	0.06 (0.06–0.07)	436 (264–814)	0

Table H1: Extreme eigenvalues of Σ^* and $\hat{\Sigma}_n = n^{-1}X^\top X$ for the simulation study presented in Section 6.2. Each cell gives the median value out of the 20 replicates with 0.05 and 0.95 quantiles given in parentheses.

Third simulation study For the simulation study conducted in Section 6.3, results for RW-GES without swap moves are shown in Table H2. Comparing Table 2 with Table H2,

one can see that using swap moves does improve the performance of RW-GES, especially when d is small.

	$c_2 = 1.3$			$c_2 = 1.1 + 0.1d$		
d	TPR (skeleton)	TPR	FPR	TPR (skeleton)	TPR	FPR
4	0.484 (0.01)	0.271 (0.01)	0.1 (0.004)	0.495 (0.01)	0.299 (0.02)	0.0961 (0.004)
5	0.569 (0.01)	0.304 (0.02)	0.118 (0.004)	0.575 (0.02)	0.338 (0.02)	0.108 (0.005)
6	0.649 (0.01)	0.37 (0.02)	0.123 (0.005)	0.647 (0.01)	0.385 (0.02)	0.107 (0.005)
7	0.696 (0.01)	0.404 (0.02)	0.125 (0.006)	0.675 (0.01)	0.398 (0.02)	0.104 (0.006)
8	0.707 (0.01)	0.403 (0.02)	0.136 (0.006)	0.691 (0.01)	0.419 (0.02)	0.0981 (0.006)
9	0.718 (0.01)	0.407 (0.02)	0.14 (0.007)	0.692 (0.01)	0.433 (0.021)	0.0962 (0.006)

Table H2: Simulation study shown in Table 2 without the swap proposal.

I Discussion on the strong beta-min or faithfulness condition

In our theoretical results for structure learning, we assume that the true distribution of \mathbf{X} , denoted by μ , is faithful w.r.t. some DAG G^* , largely because we want to use the “Chickering sequence” argument from the consistency proof for GES [15]. While for PC algorithms, it is known that the faithfulness assumption can be relaxed [80, 58], there seems no existing result on the consistency of GES beyond faithfulness; we can only find some related discussion in Zhalama et al. [79]. The reason is that, without faithfulness, even if $n \rightarrow \infty$, GES and RW-GES can get trapped at some “sub-optimal” minimal I-map of μ ; that is, there may exist $\sigma, \tau \in \mathbb{S}^p$ such that $|G_\sigma^\mu| > |G_\tau^\mu|$ but any $\mathcal{E}' \in \mathcal{N}_{\text{del}}([G_\sigma^\mu])$ is not an I-map of μ , where G_σ^μ denotes the minimal I-map of μ with ordering σ . An example is given in the proof of Theorem 2.4 in Raskutti and Uhler [59] (though it was constructed for some other purpose). From a Bayesian perspective, faithfulness is probably not a restrictive assumption: if the true distribution of \mathbf{X} is generated randomly according to the SEM (10) with $G = G^*$ and $\pi_0(B, \Omega \mid G)$ being continuous, then the distribution of \mathbf{X} is almost surely faithful to G^* ; see Lemma C7 in the supplement.

In high-dimensional settings, much more of a concern is the strong beta-min condition or the strong faithfulness assumption, which can often fail to hold in reality [71]. Here we describe a more flexible approach which may help overcome this limitation. Let \underline{b}^2 be the lower bound given in (18) (the constant in the bound can be adjusted if necessary). For each $\sigma \in \mathbb{S}^p$, define a DAG $\tilde{G}(\sigma)$ such that

$$i \rightarrow j \in \tilde{G}(\sigma) \text{ if and only if } (B_\sigma^*)_{ij}^2 \geq \underline{b}^2. \quad (\text{I.1})$$

As long as the other nonzero entries of B_σ^* are sufficiently small (so that their summed effects for each node have the same order as the noise), $\tilde{G}(\sigma)$ should coincide with the DAG $\hat{G}(\sigma)$ defined in (5), and we can prove consistency and rapid mixing results for the DAG selection problem with ordering σ by treating $\tilde{G}(\sigma)$ as the “true” model [cf. 77]. All we need is just to replace G_σ^* with $\tilde{G}(\sigma)$ in the construction of canonical paths. In some cases, we can prove the rapid mixing of RW-GES using this method. One extreme example is when all coefficients in the generating SEM are sufficiently close to zero and then RW-GES always quickly moves to the null model. Below is a more interesting example.

Example I1. Let $p = 3$ and consider the following SEM for the random vector \mathbf{X} ,

$$\mathbf{X}_1 = Z_1, \quad \mathbf{X}_2 = (1 + \epsilon)\mathbf{X}_1 + Z_2, \quad \mathbf{X}_3 = \mathbf{X}_1 - \mathbf{X}_2 + Z_3,$$

where Z_1, Z_2, Z_3 are standard normal random variables. Denote the distribution of \mathbf{X} by μ . If $\epsilon = 0$, one can easily show $1 \perp\!\!\!\perp 3$ and μ is faithful to the DAG \tilde{G} given by $1 \rightarrow 2 \leftarrow 3$. Now let ϵ be nonzero but very small. Then, μ is faithful to the complete DAG, but given a moderate sample size, the effective “true” DAG is still \tilde{G} since the evidence for $1 \not\perp\!\!\!\perp 3$ is too weak to be detected (the posterior mass will not concentrate on the equivalence class of complete DAGs either). However, our path method can be used to prove the rapid mixing of RW-GES, $\tilde{G}(\sigma)$ defined in (I.1) just becomes the minimal I-map of \tilde{G} in this case.

Nevertheless, in general, we cannot say much about the local posterior landscape at $[\tilde{G}(\sigma)]$, if $\tilde{G}(\sigma)$ is not a minimal I-map of G^* . It is likely that for some $\sigma \in \mathbb{S}^p$, $[\tilde{G}(\sigma)]$ becomes a sub-optimal local mode. But the simulation study in Section 6.2 suggests that our theory still holds “approximately” in the sense that the chain can quickly find models that are not too different from the true one (i.e., the chain mixes rapidly in most parts of the space). In addition to parallel tempering, another strategy for overcoming multimodality is to choose a larger neighborhood than the one used in RW-GES so that when the strong beta-min condition fails, we can modify the construction of the canonical path from $\tilde{G}(\sigma)$ to some “best” DAG \tilde{G} . In this regard, the DAG MCMC samplers proposed in Grzegorzcyk and Husmeier [27] and Su and Borsuk [68] may be very useful. They are equipped with proposal moves that are much more complex than single-edge modifications, which enable the samplers to jump between DAGs that encode very different CI relations. Similar ideas might be used to improve other sampling methods, including those defined on the space of equivalence classes.

References

- [1] Raj Agrawal, Caroline Uhler, and Tamara Broderick. Minimal I-MAP MCMC for scalable structure discovery in causal dag models. In *International Conference on Machine Learning*, pages 89–98, 2018.
- [2] David J Aldous. Some inequalities for reversible Markov chains. *Journal of the London Mathematical Society*, 2(3):564–576, 1982.
- [3] Hongzhi An, Da Huang, Qiwei Yao, and Cun-Hui Zhang. Stepwise searching for feature variables in high-dimensional linear regression. *Technical report*, 2008.
- [4] Steen A Andersson, David Madigan, and Michael D Perlman. A characterization of Markov equivalence classes for acyclic digraphs. *The Annals of Statistics*, 25(2):505–541, 1997.
- [5] Bryon Aragam, Arash A Amini, and Qing Zhou. Learning directed acyclic graphs with penalized neighbourhood regression. *arXiv preprint arXiv:1511.08963*, 2015.
- [6] Bryon Aragam, Arash Amini, and Qing Zhou. Globally optimal score-based learning of directed acyclic graphs in high-dimensions. In *Advances in Neural Information Processing Systems*, pages 4450–4462, 2019.
- [7] Sayantan Banerjee and Subhashis Ghosal. Bayesian structure learning in graphical models. *Journal of Multivariate Analysis*, 136:147–162, 2015.
- [8] Sayantan Banerjee, Ismaël Castillo, and Subhashis Ghosal. Bayesian inference in high-dimensional models. *arXiv preprint arXiv:2101.04491*, 2021.
- [9] Peter J Bickel and Elizaveta Levina. Regularized estimation of large covariance matrices. *The Annals of Statistics*, 36(1):199–227, 2008.

- [10] Xuan Cao, Kshitij Khare, and Malay Ghosh. Posterior graph selection and estimation consistency for high-dimensional Bayesian DAG models. *The Annals of Statistics*, 47(1):319–348, 2019.
- [11] Federico Castelletti, Guido Consonni, Marco L Della Vedova, and Stefano Peluso. Learning Markov equivalence classes of directed acyclic graphs: An objective Bayes approach. *Bayesian Analysis*, 13(4):1235–1260, 2018.
- [12] Robert Castelo and Tomás Kocka. On inclusion-driven learning of Bayesian networks. *Journal of Machine Learning Research*, 4(Sep):527–574, 2003.
- [13] David Maxwell Chickering. A transformational characterization of equivalent bayesian network structures. *Preceedings of the Eleventh Conference on Uncertainty in Artificial Intelligence*, 1995.
- [14] David Maxwell Chickering. Learning equivalence classes of Bayesian-network structures. *Journal of machine learning research*, 2(Feb):445–498, 2002.
- [15] David Maxwell Chickering. Optimal structure identification with greedy search. *Journal of machine learning research*, 3(Nov):507–554, 2002.
- [16] Mathias Drton and Marloes H Maathuis. Structure learning in graphical modeling. *Annual Review of Statistics and Its Application*, 4:365–393, 2017.
- [17] Mathias Drton, Rina Foygel, and Seth Sullivant. Global identifiability of linear structural equation models. *The Annals of Statistics*, 39(2):865–886, 2011.
- [18] Daniel Eaton and Kevin Murphy. Bayesian structure learning using dynamic programming and MCMC. pages 101–108, 2007.
- [19] Byron Ellis and Wing Hung Wong. Learning causal Bayesian network structures from experimental data. *Journal of the American Statistical Association*, 103(482):778–789, 2008.
- [20] Nir Friedman and Daphne Koller. Being Bayesian about network structure: A Bayesian approach to structure discovery in Bayesian networks. *Machine learning*, 50(1-2):95–125, 2003.
- [21] Bin Gao and Yuehua Cui. Learning directed acyclic graphical structures with genetical genomics data. *Bioinformatics*, 31(24):3953–3960, 2015.
- [22] Chao Gao, Aad W van der Vaart, and Harrison H Zhou. A general framework for Bayes structured linear models. *The Annals of Statistics*, 48(5):2848–2878, 2020.
- [23] Dan Geiger and David Heckerman. Parameter priors for directed acyclic graphical models and the characterization of several probability distributions. *The Annals of Statistics*, 30(5):1412–1440, 2002.

- [24] Paolo Giudici and Robert Castelo. Improving Markov chain Monte Carlo model search for data mining. *Machine learning*, 50(1-2):127–158, 2003.
- [25] Robert JB Goudie and Sach Mukherjee. A gibbs sampler for learning dags. *The Journal of Machine Learning Research*, 17(1):1032–1070, 2016.
- [26] Simon Griffiths, Ross Kang, Roberto Oliveira, and Viresh Patel. Tight inequalities among set hitting times in Markov chains. *Proceedings of the American Mathematical Society*, 142(9):3285–3298, 2014.
- [27] Marco Grzegorczyk and Dirk Husmeier. Improving the structure MCMC sampler for Bayesian networks by introducing a new edge reversal move. *Machine Learning*, 71(2-3):265, 2008.
- [28] Yangbo He, Jinzhu Jia, and Bin Yu. Reversible MCMC on Markov equivalence classes of sparse directed acyclic graphs. *The Annals of Statistics*, 41(4):1742–1779, 2013.
- [29] Jennifer A Hoeting, David Madigan, Adrian E Raftery, and Chris T Volinsky. Bayesian model averaging: a tutorial (with comments by M. Clyde, David Draper and EI George, and a rejoinder by the authors. *Statistical science*, 14(4):382–417, 1999.
- [30] Seonghyun Jeong and Subhashis Ghosal. Posterior contraction in sparse generalized linear models. *Biometrika*, 108(2):367–379, 2021.
- [31] Valen E Johnson and David Rossell. Bayesian model selection in high-dimensional settings. *Journal of the American Statistical Association*, 107(498):649–660, 2012.
- [32] Nabil Kahale. A semidefinite bound for mixing rates of Markov chains. *Random Structures & Algorithms*, 11(4):299–313, 1997.
- [33] Markus Kalisch and Peter Bühlmann. Estimating high-dimensional directed acyclic graphs with the PC-algorithm. *Journal of Machine Learning Research*, 8(Mar):613–636, 2007.
- [34] Daphne Koller and Nir Friedman. *Probabilistic graphical models: principles and techniques*. MIT press, 2009.
- [35] Jack Kuipers and Giusi Moffa. Partition MCMC for inference on acyclic digraphs. *Journal of the American Statistical Association*, 112(517):282–299, 2017.
- [36] Clifford Lam and Jianqing Fan. Sparsistency and rates of convergence in large covariance matrix estimation. *Annals of statistics*, 37(6B):4254, 2009.
- [37] Beatrice Laurent and Pascal Massart. Adaptive estimation of a quadratic functional by model selection. *Annals of Statistics*, pages 1302–1338, 2000.
- [38] Steffen L Lauritzen. Propagation of probabilities, means, and variances in mixed graphical association models. *Journal of the American Statistical Association*, 87(420):1098–1108, 1992.

- [39] Kyoungjae Lee, Jaeyong Lee, and Lizhen Lin. Minimax posterior convergence rates and model selection consistency in high-dimensional DAG models based on sparse cholesky factors. *The Annals of Statistics*, 47(6):3413–3437, 2019.
- [40] Song Liu, Taiji Suzuki, Raissa Relator, Jun Sese, Masashi Sugiyama, and Kenji Fukumizu. Support consistency of direct sparse-change learning in Markov networks. *The Annals of Statistics*, 45(3):959–990, 2017.
- [41] Marloes H Maathuis, Diego Colombo, Markus Kalisch, and Peter Bühlmann. Predicting causal effects in large-scale systems from observational data. *Nature Methods*, 7(4):247–248, 2010.
- [42] David Madigan, Jeremy York, and Denis Allard. Bayesian graphical models for discrete data. *International Statistical Review/Revue Internationale de Statistique*, pages 215–232, 1995.
- [43] David Madigan, Steen A Andersson, Michael D Perlman, and Chris T Volinsky. Bayesian model averaging and model selection for Markov equivalence classes of acyclic digraphs. *Communications in Statistics–Theory and Methods*, 25(11):2493–2519, 1996.
- [44] Ryan Martin, Raymond Mess, and Stephen G Walker. Empirical Bayes posterior concentration in sparse high-dimensional linear models. *Bernoulli*, 23(3):1822–1847, 2017.
- [45] Christopher Meek. *Graphical Models: Selecting causal and statistical models*. PhD thesis, PhD thesis, Carnegie Mellon University, 1997.
- [46] Nicolai Meinshausen and Peter Bühlmann. High-dimensional graphs and variable selection with the lasso. *The annals of statistics*, 34(3):1436–1462, 2006.
- [47] Paul Munteanu and Mohamed Bendou. The EQ framework for learning equivalence classes of Bayesian networks. In *Proceedings 2001 IEEE International Conference on Data Mining*, pages 417–424. IEEE, 2001.
- [48] Preetam Nandy, Alain Hauser, and Marloes H Maathuis. High-dimensional consistency in score-based and hybrid structure learning. *The Annals of Statistics*, 46(6A):3151–3183, 2018.
- [49] Naveen Naidu Narisetty and Xuming He. Bayesian variable selection with shrinking and diffusing priors. *The Annals of Statistics*, 42(2):789–817, 2014.
- [50] Teppo Mikael Niinimäki, Pekka Parviainen, and Mikko Koivisto. Partial order MCMC for structure discovery in Bayesian networks. In *Proceedings of the Twenty-Seventh Conference Conference on Uncertainty in Artificial Intelligence (UAI-11)*. AUAI Press, 2011.
- [51] Debdeep Pati, Anirban Bhattacharya, Natesh S Pillai, and David Dunson. Posterior contraction in sparse Bayesian factor models for massive covariance matrices. *The Annals of Statistics*, 42(3):1102–1130, 2014.

- [52] Judea Pearl. *Probabilistic reasoning in intelligent systems: networks of plausible inference*. Morgan Kaufmann, 1988.
- [53] Stefano Peluso and Guido Consonni. Compatible priors for model selection of high-dimensional gaussian dags. *Electronic Journal of Statistics*, 14(2):4110–4132, 2020.
- [54] Jose M Pena. Approximate counting of graphical models via MCMC. In *AISTATS*, pages 355–362, 2007.
- [55] Yuval Peres and Perla Sousi. Mixing times are hitting times of large sets. *Journal of Theoretical Probability*, 28(2):488–519, 2015.
- [56] Michael D Perlman. Graphical model search via essential graphs. *Contemporary Mathematics*, 287:255–266, 2001.
- [57] Jonas Peters and Peter Bühlmann. Identifiability of gaussian structural equation models with equal error variances. *Biometrika*, 101(1):219–228, 2014.
- [58] Joseph Ramsey, Peter Spirtes, and Jiji Zhang. Adjacency-faithfulness and conservative causal inference. UAI’06, page 401–408, Arlington, Virginia, USA, 2006. AUAI Press. ISBN 0974903922.
- [59] Garvesh Raskutti and Caroline Uhler. Learning directed acyclic graph models based on sparsest permutations. *Stat*, 7(1):e183, 2018.
- [60] Garvesh Raskutti, Bin Yu, Martin J Wainwright, and Pradeep Ravikumar. Model selection in Gaussian graphical models: High-dimensional consistency of ℓ_1 -regularized MLE. *Advances in Neural Information Processing Systems*, 21, 2008.
- [61] Mark Rudelson and Roman Vershynin. Non-asymptotic theory of random matrices: extreme singular values. In *Proceedings of the International Congress of Mathematicians 2010 (ICM 2010) (In 4 Volumes) Vol. I: Plenary Lectures and Ceremonies Vols. II–IV: Invited Lectures*, pages 1576–1602. World Scientific, 2010.
- [62] Laurent Saloff-Coste. Lectures on finite Markov chains. In *Lectures on probability theory and statistics*, pages 301–413. Springer, 1997.
- [63] Marco Scutari, Catharina Elisabeth Graafland, and José Manuel Gutiérrez. Who learns better Bayesian network structures: Accuracy and speed of structure learning algorithms. *International Journal of Approximate Reasoning*, 115:235–253, 2019.
- [64] Alistair Sinclair. Improved bounds for mixing rates of Markov chains and multicommodity flow. *Combinatorics, probability and Computing*, 1(4):351–370, 1992.
- [65] Liam Solus, Yuhao Wang, and Caroline Uhler. Consistency guarantees for greedy permutation-based causal inference algorithms. *arXiv preprint arXiv:1702.03530*, 2017.
- [66] Peter Spirtes, Clark N Glymour, Richard Scheines, and David Heckerman. *Causation, prediction, and search*. MIT press, 2000.

- [67] Milan Studený. *Probabilistic conditional independence structures*. Springer Science & Business Media, 2006.
- [68] Chengwei Su and Mark E Borsuk. Improving structure mcmc for bayesian networks through markov blanket resampling. *The Journal of Machine Learning Research*, 17(1): 4042–4061, 2016.
- [69] Tingni Sun and Cun-Hui Zhang. Sparse matrix inversion with scaled Lasso. *The Journal of Machine Learning Research*, 14(1):3385–3418, 2013.
- [70] Kunal Talwar. Computational separations between sampling and optimization. *Advances in Neural Information Processing Systems*, 32, 2019.
- [71] Caroline Uhler, Garvesh Raskutti, Peter Bühlmann, and Bin Yu. Geometry of the faithfulness assumption in causal inference. *The Annals of Statistics*, pages 436–463, 2013.
- [72] Sara Van de Geer and Peter Bühlmann. ℓ^0 -penalized maximum likelihood for sparse directed acyclic graphs. *The Annals of Statistics*, 41(2):536–567, 2013.
- [73] Thomas Verma and Judea Pearl. *Equivalence and synthesis of causal models*. UCLA, Computer Science Department, 1991.
- [74] Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.
- [75] Martin J Wainwright. Sharp thresholds for high-dimensional and noisy sparsity recovery using ℓ_1 -constrained quadratic programming (lasso). *IEEE transactions on information theory*, 55(5):2183–2202, 2009.
- [76] Yun Yang and Surya T Tokdar. Minimax-optimal nonparametric regression in high dimensions. *The Annals of Statistics*, 43(2):652–674, 2015.
- [77] Yun Yang, Martin J Wainwright, and Michael I Jordan. On the computational complexity of high-dimensional Bayesian variable selection. *The Annals of Statistics*, 44(6): 2497–2532, 2016.
- [78] Giacomo Zanella. Informed proposals for local MCMC in discrete spaces. *Journal of the American Statistical Association*, 115(530):852–865, 2020.
- [79] Zhalama, Jiji Zhang, and Wolfgang Mayer. Weakening faithfulness: some heuristic causal discovery algorithms. *International journal of data science and analytics*, 3(2): 93–104, 2017.
- [80] Jiji Zhang and Peter Spirtes. Detection of unfaithfulness and robust causal inference. *Minds and Machines*, 18(2):239–271, 2008.
- [81] Quan Zhou and Hyunwoong Chang. Supplementary material for “complexity analysis of Bayesian learning of high-dimensional DAG models and their equivalence classes”.

- [82] Quan Zhou, Jun Yang, Dootika Vats, Gareth O Roberts, and Jeffrey S Rosenthal. Dimension-free mixing for high-dimensional Bayesian variable selection. *arXiv preprint arXiv:2105.05719*, 2021.
- [83] Bumeng Zhuo and Chao Gao. Mixing time of Metropolis-Hastings for Bayesian community detection. *Journal of Machine Learning Research*, 22:10–1, 2021.