

EMERGENT BEHAVIORS OF HIGH-DIMENSIONAL KURAMOTO MODELS ON STIEFEL MANIFOLDS

SEUNG-YEAL HA, MYEONGJU KANG, AND DOHYUN KIM

ABSTRACT. We study emergent asymptotic dynamics for the first and second-order high-dimensional Kuramoto models on Stiefel manifolds which extend the previous consensus models on Riemannian manifolds including several matrix Lie groups. For the first-order consensus model on the Stiefel manifold proposed in [Markdahl et al, 2018], we show that the homogeneous ensemble relaxes the complete consensus state exponentially fast. On the other hand for a heterogeneous ensemble, we provide a sufficient condition leading to the phase-locked state in which relative distances between two states converge to definite values in a large coupling strength regime. We also propose a second-order extension of the first-order one by adding an inertial effect, and study emergent behaviors using Lyapunov functionals such as an energy functional and an averaged distance functional.

1. INTRODUCTION

Collective behaviors of biological and chemical oscillators have been widely studied not only in applied mathematics but also in other scientific disciplines, for instance, flocking of drones [30, 46] and passivity-based distributed optimization [11, 29, 51] in control theory, social dynamics [2, 3, 32, 44] and swarming behavior in quantitative biology [5, 14, 15, 23, 50]. Despite of its crucial roles in biological processes, the mathematical study of such collective dynamics has been started only after the seminal work of Winfree [61] and Kuramoto [33, 34] a half century ago. Among other models describing collective oscillatory behavior, to name a few, the Cucker-Smale model [13], the Kuramoto model [12, 33, 34] and the Vicsek model [58], our main interest lies in consensus models on Riemannian manifolds. So far, there has been much available literature dealing with emergent dynamics on Riemannian manifolds, for instance, on the unit sphere \mathbb{S}^{d-1} in [45, 62], on the hyperboloid \mathbb{H}^{d-1} in [49] and on the matrix Lie groups including special orthogonal group $\text{SO}(d)$ in [16, 53, 56], the unitary group $\text{U}(d)$ in [27, 28, 39], the *Lohe group* in [26] and even for the quaternions \mathbb{H}_1 in [17]. In particular, we are concerned with the Stiefel manifold [55] in $\mathbb{R}^{n \times p}$ for $p \leq n$ consisting of all rectangular matrices satisfying the relations:

$$\text{St}(p, n) := \{S \in M_{n,p}(\mathbb{R}) : S^\top S = I_p\}, \quad \|S\|_{\mathbb{F}}^2 := \text{tr}(S^\top S) = p,$$

where S^\top is defined as the transpose of a matrix S and $M_{n,p}(\mathbb{R})$ denotes the set of all $n \times p$ matrices with real entries.

Note that the unit sphere and (special) orthogonal group can be recovered from the specific choices of (p, n) . Thus, the Stiefel manifold is a manifold including \mathbb{S}^d , $\text{SO}(d)$ and

Date: November 13, 2021.

2010 Mathematics Subject Classification. 34D06, 34C15, 35B35.

Key words and phrases. Consensus, emergence, Kuramoto model, Lohe model, second-order extension, Stiefel manifold, synchronization.

Acknowledgment. The work of S.-Y. Ha was supported by NRF grant (2017R1A2B2001864).

$O(d)$ as special cases. We will briefly review basic properties of the Stiefel manifold in Section 2.1. It is worthwhile to mention that an optimization problem $\min_{S \in \text{St}(p,n)} f(S)$ for an objective function $f : \text{St}(p,n) \rightarrow \mathbb{R}$ has been extensively studied due to its computational difficulty and broad applications, e.g., the linear eigenvalue problem [21, 59], the nearest low-rank correlation matrix problem [35], singular value decomposition [38, 54], Riemmanian optimization [63] and applications to computer vision [40, 57]. We refer the reader to [18, 52, 60] and reference therein for introductions to optimization problem on the Stiefel manifold and applications.

In this paper, we mainly consider the consensus model in [41, 42] in which particles interact with neighboring ones on the Stiefel manifold and for the consistency of the paper, we briefly introduce the model below. Let $(\text{St}(p,n), \|\cdot\|)$ be the Stiefel manifold canonically embedded into the Euclidean space $\mathbb{R}^{n \times p}$ with its Frobenius (or Euclidean) norm and $\mathcal{S} := (S_1, \dots, S_N)$ denote state ensemble of particles. Next, we define the potential function \mathcal{V} as the total misfit functional with a symmetric connectivity matrix $\mathcal{A} := (a_{ik})$:

$$\mathcal{V}(\mathcal{S}) := \frac{\kappa}{N^2} \sum_{i,k=1}^N a_{ik} \|S_i - S_k\|_F^2, \quad a_{ik} = a_{ki} > 0.$$

Then, the gradient flow with the potential function \mathcal{V} reads as

$$\dot{S}_i = -\nabla_{S_i} \mathcal{V} = \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left[S_k - \frac{1}{2}(S_i S_i^\top S_k + S_i S_k^\top S_i) \right], \quad i = 1, \dots, N. \quad (1.1)$$

See Section 2.2 for detailed description. We finally add generalized natural frequencies $\Xi_i \in \mathfrak{so}(p)$ into (1.1) to find the desired model so that the Stiefel manifold is still positively invariant:

$$\dot{S}_i = S_i \Xi_i + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left[S_k - \frac{1}{2}(S_i S_i^\top S_k + S_i S_k^\top S_i) \right], \quad i = 1, \dots, N. \quad (1.2)$$

Since the Stiefel manifold is compact, the global existence of a unique solution to (1.2) directly follows from the standard Cauchy-Lipschitz theory. Detailed description will be presented in Section 2.1.

For the perspective of optimization, a gradient flow (1.1) with a total distance as a potential can be regarded an optimization problem of a given target (or objective) function defined on the Stiefel manifold. Thus, our model with a consensus estimate provides a method which tackles an optimization problem on the Stiefel manifold by means of dynamical systems approach. It should be mentioned that such optimization problems on manifolds have extensively studied in literature [4, 7, 37, 48, 52]. In particular, consensus-based optimization (CBO for brevity) algorithm toward a global optimization [6, 19, 20, 22, 47] has been recently proposed (see [31] for CBO on the Stiefel manifold). On the other hand for a control (or consensus protocol) perspective, it is mentioned in [43] that *ad-hoc* control algorithm would be employed in a specific situation. In this manner, one can consider (1.1) as a control problem by regarding the network structure a_{ik} as an external control (or parameter) to obtain a desired pattern formation (see [36] on the unit sphere and *an Olympic ring* in [10] for a distributed control approach for the Cucker-Smale flocking model [13]). For instance, if we set a_{ik} to be negative (or competitive), one can expect deployment where all agents tend to splay state, whereas complete consensus would be achieved for a positive

(or cooperative) a_{ik} . However in this paper, we restrict ourselves to a cooperative network so that emergence of complete consensus is our primary concern.

We analyze system (1.2) with all-to-all network topology and left-translation invariance. More precisely, we choose all-to-all network $a_{ij} = \kappa/N$ where κ denotes the coupling strength so that all particles communicate with neighbors through the same weight. In addition, we do not consider the effect of Ω_i by setting $\Omega_i \equiv O$ due to the left-translation property (see Lemma 2.2 for details). Then, our first-order consensus model is governed by the following Cauchy problem:

$$\begin{cases} \dot{S}_i = S_i \Xi_i + \frac{\kappa}{N} \sum_{k=1}^N \left[S_k - \frac{1}{2} (S_i S_i^\top S_k + S_i S_k^\top S_i) \right], & t > 0, \\ S_i(0) = S_i^{\text{in}} \in \text{St}(p, n), & i = 1, \dots, N. \end{cases} \quad (1.3)$$

Although the gradient flow structure guarantees that the Stiefel manifold is positively invariant along the flow (1.2), we provide its alternative and direct proof in Lemma 2.1.

Next, we turn to the second-order extension of the first-order model (1.3). In [25], the (first-order) Lohe matrix model on the unitary group has been extended to a second-order one by incorporating inertial effect. Similarly, we can also extend the first-order model (1.3) into a second-order one whose dynamics is governed by the following Cauchy problem:

$$\begin{cases} m \ddot{S}_i = -m S_i \dot{S}_i^\top \dot{S}_i - \gamma \dot{S}_i + S_i \Xi_i + \frac{m}{\gamma} (2 \dot{S}_i \Xi_i - S_i \Xi_i S_i^\top \dot{S}_i + S_i \dot{S}_i^\top S_i \Xi_i) \\ \quad + \frac{\kappa}{N} \sum_{k=1}^N \left(S_k - \frac{1}{2} (S_i S_i^\top S_k + S_i S_k^\top S_i) \right), & t > 0, \\ S_i(0) = S_i^{\text{in}} \in \text{St}(p, n), \quad \dot{S}_i^{t, \text{in}} S_i^{\text{in}} + S_i^{t, \text{in}} \dot{S}_i^{\text{in}} = O, & i = 1, \dots, N, \end{cases} \quad (1.4)$$

where m and γ represent mass and friction, respectively. Although it seems that the model looks quite complicated, we show the the Stiefel manifold is still positively invariant along system (1.4). Of course, if we turn off the inertial effect, that is, $m = 0$, then the first-order model (1.3) can be recovered from (1.4) straightforwardly. We also note that the Cauchy problems (1.3) and (1.4) have a unique global solution due to the compactness of Stiefel manifold and standard Cauchy-Lipschitz theory. Here, a global solution is referred as a solution which exists for all time (or globally). In other words, a solution does not blow up in finite time. For more detailed description of (1.4), we refer the reader to Section 2.2. Next, we recall several concepts for consensus as follows:

Definition 1.1. Let $\mathcal{S} = (S_1, \dots, S_N)$ be a global solution to (1.3) or (1.4).

(1) System (1.3) or (1.4) exhibits complete consensus if the following convergence holds:

$$\lim_{t \rightarrow \infty} \|S_i(t) - S_j(t)\|_F = 0, \quad \text{for all } i, j = 1, \dots, N.$$

(2) System (1.3) or (1.4) exhibits practical consensus if the following convergence holds:

$$\lim_{\kappa \rightarrow \infty} \limsup_{t \rightarrow \infty} \|S_i(t) - S_j(t)\|_F = 0, \quad \text{for all } i, j = 1, \dots, N.$$

(3) A global solution to system (1.3) or (1.4) tends to a phase-locked state if the following relation holds:

$$\lim_{t \rightarrow \infty} S_i^\top S_j \quad \text{exists for all } i, j = 1, \dots, N.$$

The main results of this paper deal with the emergent collective behaviors for the first-order model (1.3) and the second-order model (1.4). First, we consider the first-order model (1.3) with a homogeneous ensemble. The corresponding proof can be found in Section 3.1

Theorem 1.1. *Suppose that initial data and system parameters satisfy*

$$\mathcal{D}(\mathcal{S}^{\text{in}}) < \sqrt{2}, \quad \Xi \equiv O, \quad a_{ik} : \text{undirected and connected graph},$$

and let \mathcal{S} be a global solution to (1.2). Then, system (1.2) exhibits complete consensus exponentially fast.

The results in [41, 42] deal with almost global consensus without any explicit decay estimate. On the other hand, for a heterogeneous ensemble, we establish the emergence of the locked state exponentially fast in a large coupling regime.

Theorem 1.2. *Suppose that the coupling strength and initial data satisfy (3.11) and let \mathcal{S} be a global solution to (1.2). Then, \mathcal{S} tends to a locked state.*

For a detailed initial framework and proof, we refer to Section 3.2.

Next, we turn to the second-order model (1.4). As in the first-order one, we consider both homogeneous and heterogeneous ensembles. For the desired results, we first derive an energy estimate (Proposition 4.1):

$$\begin{aligned} \frac{d}{dt} \left(\frac{m}{N} \sum_{i=1}^N \|\dot{S}_i\|_{\mathbb{F}}^2 + \frac{\kappa}{2N^2} \sum_{i,j=1}^N \|S_i - S_j\|_{\mathbb{F}}^2 \right) \\ = -\frac{2\gamma}{N} \sum_{i=1}^N \|\dot{S}_i\|_{\mathbb{F}}^2 + \frac{1}{N} \sum_{i=1}^N \text{tr}(\dot{S}_i^{\top} S_i \Xi_i - \Xi_i S_i^{\top} \dot{S}_i), \quad t > 0. \end{aligned} \quad (1.5)$$

In what follows, we assume that the network topology satisfies

$$\begin{aligned} a_m := \min a_{ik}, \quad a_M := \max a_{ik}, \quad d(\mathcal{A}) := \max |a_{ik} - a_{jk}|, \\ \frac{1}{N} \sum_{k=1}^N a_{ik} \equiv \xi, \quad i = 1, \dots, N, \quad 0 < \Lambda := a_m - \frac{N-1}{N}(a_M + d(\mathcal{A})) < 8pa_M^2. \end{aligned}$$

Furthermore, for the averaged relative distances $\mathcal{G} := \frac{1}{N^2} \sum_{i,j=1}^N \|S_i - S_j\|_{\mathbb{F}}^2$, we derive a second-order differential inequality (Lemma 4.3) for \mathcal{G} :

$$m\ddot{\mathcal{G}} + \gamma\dot{\mathcal{G}} + 2\kappa\xi\mathcal{G} \leq 16mD(\dot{\mathcal{S}})^2 + 8\|\Xi\|_{\infty} + \frac{16m\sqrt{p}\|\Xi\|_{\infty}}{\gamma}D(\dot{\mathcal{S}}), \quad (1.6)$$

where $D(\dot{\mathcal{S}})$ and $\|\Xi\|_{\infty}$ are defined as follows:

$$D(\dot{\mathcal{S}}) := \max_{1 \leq i \leq N} \|\dot{S}_i\|_{\mathbb{F}}, \quad \|\Xi\|_{\infty} := \max_{1 \leq i \leq N} \|\Xi_i\|_{\mathbb{F}}.$$

Based on two estimates (1.5) and (1.6), for a homogeneous ensemble, we show that the complete consensus occurs for some admissible initial data.

Theorem 1.3. *Suppose that system parameters and initial data satisfy*

$$m > 0, \quad \gamma > 0, \quad \kappa > 0, \quad \Xi_i \equiv O \quad \text{for } i = 1, \dots, N,$$

$$\mathcal{E}(0) = \frac{m}{N} \sum_{i=1}^N \|\dot{S}_i^0\|_F^2 + \frac{\kappa}{2N^2} \sum_{i,j=1}^N a_{ij} \|S_i^0 - S_j^0\|_F^2 < \infty,$$

and let \mathcal{S} be a global solution to (1.4). Then, we have

$$\lim_{t \rightarrow \infty} \|\dot{S}_i(t)\|_F = 0 \quad \text{for } i = 1, \dots, N.$$

Moreover, system (1.4) exhibits the complete consensus:

$$\lim_{t \rightarrow \infty} \mathcal{G}(t) = 0.$$

We refer the reader to Section 4.1 for the proof. In contrast, for a heterogeneous ensemble, we assume that the inertia and the coupling satisfy the following relation:

$$m\kappa^{1+\eta} = \mathcal{O}(1) \quad \text{for some } \eta > 0.$$

Then, under this setting, we arrive at the following result.

Theorem 1.4. *Suppose that system parameters and initial data satisfy*

$$D(\dot{S}^{\text{in}}) < \frac{1}{\gamma} (\|\Xi\|_\infty + \kappa a_M \sqrt{p}), \quad m = \frac{m_0}{\kappa^{1+\eta}}, \quad (1.7)$$

and let \mathcal{S} be a global solution to (1.4). Then, system (1.4) exhibits practical consensus:

$$\lim_{\kappa \rightarrow \infty} \limsup_{t \rightarrow \infty} \mathcal{G}(t) = 0.$$

The proof can be found in Section 4.2.

The rest of the paper is organized as follows. In Section 2, we briefly discuss properties of the Stiefel manifold to be used later and provide descriptions of the first-order and second-order models. In Section 3, we present proofs of Theorem 1.1 and Theorem 1.2 which deal with the first-order model. In Section 4, rigorous justification of Theorem 1.3 and Theorem 1.4 for the second-order model is provided. Finally, Section 5 is devoted to a brief summary of our main results and discussion for a future work.

Notation: We denote by $M_{n,p}(\mathbb{R})$ as the set of all $n \times p$ real matrices and for notational simplicity, we set $M_n(\mathbb{R}) := M_{n,n}(\mathbb{R})$. In addition, O is the zero matrix regardless of its size.

2. PRELIMINARIES

In this section, we briefly discuss the Stiefel manifold and present detailed description and properties of the first-order and second-order consensus models on the Stiefel manifold to be used later in later sections.

2.1. The Stiefel manifold. We define Stiefel manifold and Frobenius norm:

$$\text{St}(p, n) := \{S \in M_{n,p}(\mathbb{R}) : S^\top S = I_p\}, \quad \|S\|_{\mathbb{F}}^2 := \text{tr}(S^\top S) = p \quad \text{for } S \in \text{St}(p, n).$$

Alternatively, it can be defined as the set of all p -tuples (x_1, \dots, x_p) of orthonormal vectors in \mathbb{R}^n or it is isomorphic to a homogeneous space:

$$\text{St}(p, n) \simeq \text{O}(n)/\text{O}(n-p).$$

In addition, if p is strictly less than n , then one also finds

$$\text{St}(p, n) \simeq \text{SO}(n)/\text{SO}(n-p).$$

Thus, the Stiefel manifold $\text{St}(p, n)$ is a compact set whose dimension is $pn - p(p+1)/2$. Furthermore, it is well known that $\text{St}(p, n)$ reduces to several well-known manifolds, for instance,

$$\text{St}(1, n) = \mathbb{S}^{n-1} \subseteq \mathbb{R}^n, \quad \text{St}(n-1, n) = \text{SO}(n), \quad \text{St}(n, n) = \text{O}(n). \quad (2.1)$$

We set $\mathfrak{so}(n)$ to be the special orthogonal Lie algebra associated with $\text{SO}(n)$. Then, we define two maps $\text{skew} : M_n(\mathbb{R}) \rightarrow \mathfrak{so}(n)$ and $\text{sym} : M_n(\mathbb{R}) \rightarrow \mathfrak{so}(n)^\perp$ as

$$\text{skew}(X) := \frac{1}{2}(X - X^\top), \quad \text{sym}(X) := \frac{1}{2}(X + X^\top).$$

Then, it is easy to see that the tangent space and the normal space of $\text{St}(p, n)$ at a point S are defined by

$$\begin{aligned} \text{T}_S \text{St}(p, n) &:= \{A \in M_{n,p}(\mathbb{R}) : \text{sym}(S^\top A) = O\} = \{A \in M_{n,p}(\mathbb{R}) : S^\top A + A^\top S = O\}, \\ \text{N}_S \text{St}(p, n) &:= \{SV : V \text{ is a } p \times p \text{ symmetric matrix}\}, \end{aligned}$$

and the projection of X onto $\text{N}_S \text{St}(p, n)$ is given by $S\text{sym}(S^\top X)$. Thus, the projection by $\Pi : M_{n,p}(\mathbb{R}) \times \text{St}(p, n) \rightarrow \text{T}_S \text{St}(p, n)$ is written as

$$\Pi(X, S) = X - S\text{sym}(S^\top X) = S\text{skew}(S^\top X) + (I_n - SS^\top)X.$$

For further details for the Stiefel manifold, we refer to [18].

2.2. A first-order consensus model on $\text{St}(p, n)$. In this subsection, we review the first-order model proposed in [41, 42] and study its basic property. First, we state the positive invariance of $\text{St}(p, n)$ for (1.2) which can be guaranteed from the gradient flow structure.

Lemma 2.1 (Positive invariance of the Stiefel manifold). *Let \mathcal{S} be a global solution to (1.2) with the initial data $\mathcal{S}^{\text{in}} := (S_1^{\text{in}}, \dots, S_N^{\text{in}})$. Then, we have*

$$S_i^{\text{in}} \in \text{St}(p, n) \implies S_i(t) \in \text{St}(p, n), \quad t > 0.$$

Next, we consider the left-translation invariance property whose proof directly follows from straightforward calculations.

Lemma 2.2 (Left-translation invariance). *For all $L \in \text{O}(n)$, system (1.2) is invariant under left-translation by an $n \times n$ orthogonal matrix in the sense that a transformed variable $V_i := LS_i$ satisfies*

$$\dot{V}_i = \Omega_i V_i + V_i \Xi_i + \sum_{k=1}^N a_{ik} \left(V_k - \frac{1}{2}(V_i V_i^\top V_k + V_i V_k^\top V_i) \right).$$

The model (1.2) on the Stiefel manifold in fact includes several first-order models on the Riemannian manifolds such as \mathbb{S}^{n-1} in [45], $\text{SO}(n)$ in [53] and \mathbb{S}^1 in [33]:

$$\begin{aligned}
\text{(i)} \quad \dot{R}_i &= \Omega_i R_i + \sum_{j=1}^N a_{ij} R_i \text{skew}(R_i^\top R_j), \quad R_i \in \text{SO}(n). \\
\text{(ii)} \quad \dot{x}_i &= \Omega_i x_i + (I_n - x_i \otimes x_i) \sum_{j=1}^N a_{ij} x_j, \quad x_i \in \mathbb{S}^{n-1}. \\
\text{(iii)} \quad \dot{\theta}_i &= \nu_i + \sum_{j=1}^N a_{ij} \sin(\theta_j - \theta_i), \quad \theta \in \mathbb{R}.
\end{aligned} \tag{2.2}$$

Reduction basically follows from the property (2.1) and the projection operator, since the models (1.2) and (2.2) share the gradient flow structure. Moreover, system (1.2) satisfies the following splitting property for a homogeneous ensemble $\Xi_i \equiv \Xi$ for $i = 1, \dots, N$. Then, the rotated variable $Y_i := S_i e^{-t\Xi}$ satisfies (1.2) with $\Xi \equiv O$. For a proof on the reduction and splitting property, we refer to Proposition 1 of [42]. Thus, when we consider a homogeneous ensemble, we set $\Xi \equiv O$ without loss of generality.

Remark 2.1. *If we perform left-multiplication by S_i^\top in (1.2), then we obtain the following reduced dynamics:*

$$S_i^\top \dot{S}_i = \Xi_i + \frac{\kappa}{2N} \sum_{k=1}^N (S_i^\top S_k - S_k^\top S_i), \tag{2.3}$$

which means that (2.3) can be uniquely derived from (1.2). In other words, if $\{S_i\}_{i=1}^N$ is a solution to (1.2), then it also becomes a solution to (2.3). However, the converse statement might not hold, since there does not exist an inverse matrix of S_i^\top . Hence, a solution set for (1.2) is a proper subset of that for (2.3).

Below, we briefly recall previous results in [41, 42] on the first-order model on the Stiefel manifolds. As far as the authors know, there has been only two literatures which concern with the consensus model (1.3) on the Stiefel manifolds. A graph \mathcal{G} is a pair (V, E) where $V = \{1, \dots, N\}$ and E is a subset of V where each subset is of cardinality two. Suppose that $\mathcal{S} = (S_1, \dots, S_N)$ is the state of particles interacting through the graph \mathcal{G} .

Theorem 2.1. [41, 42] *Suppose the pair (p, n) satisfy*

$$p \leq \frac{2}{3}n - 1,$$

and \mathcal{G} is connected. Let \mathcal{S} be a global solution to (1.1). Then, the consensus manifold \mathcal{C} defined by

$$\mathcal{C} := \{(S_i)_{i=1}^N \in \text{St}(p, n)^N : S_i = S_j, \quad \forall \{i, j\} \in E\}$$

is almost globally asymptotically stable.

2.3. A second-order extension. In this subsection, we introduce a second-order extension (1.4) of (1.3) by adding suitable inertia like terms:

$$\begin{aligned} m\ddot{S}_i &= -mS_i\dot{S}_i^\top\dot{S}_i - \gamma\dot{S}_i + S_i\Xi_i + \frac{m}{\gamma}(2\dot{S}_i\Xi_i - S_i\Xi_iS_i^\top\dot{S}_i + S_i\dot{S}_i^\top S_i\Xi_i) \\ &\quad + \frac{\kappa}{N} \sum_{k=1}^N \left(S_k - \frac{1}{2}(S_iS_i^\top S_k + S_iS_k^\top S_i) \right). \end{aligned} \quad (2.4)$$

Note that in a formal zero inertia limit $m \rightarrow 0$, system (2.4) reduces to the first-order model (1.2). In order to show that the proposed model (2.4) is a suitable extension on the Stiefel manifold, we need to make sure that the governing manifold $\text{St}(p, n)$ is still positively invariant along (2.4).

Lemma 2.3 (Positive invariance of the Stiefel manifold). *Suppose the initial data $(S^{\text{in}}, \dot{S}^{\text{in}})$ satisfy*

$$S_i^{\text{in}} \in \text{St}(p, n), \quad \dot{S}_i^{t, \text{in}} S_i^{\text{in}} + S_i^{t, \text{in}} \dot{S}_i^{\text{in}} = O, \quad i = 1, \dots, N, \quad (2.5)$$

and let \mathcal{S} be a global solution to (2.4). Then, we have

$$S_i(t) \in \text{St}(p, n), \quad i = 1, \dots, N, \quad t \geq 0.$$

Proof. We take the left multiplication of S_i^\top to (2.4) to obtain

$$\begin{aligned} mS_i^\top\ddot{S}_i &= -mS_i^\top S_i\dot{S}_i^\top\dot{S}_i - \gamma S_i^\top\dot{S}_i + S_i^\top S_i\Xi_i + \frac{m}{\gamma} \left(2S_i^\top\dot{S}_i\Xi_i - S_i^\top S_i\Xi_iS_i^\top\dot{S}_i + S_i^\top S_i\dot{S}_i^\top S_i\Xi_i \right) \\ &\quad + \frac{\kappa}{2N} \sum_{k=1}^N \left[S_i^\top S_k - \frac{1}{2}(S_i^\top S_iS_i^\top S_k + S_i^\top S_iS_k^\top S_i) \right]. \end{aligned} \quad (2.6)$$

We transpose (2.6) and use the skew-symmetry of Ξ_i to find

$$\begin{aligned} m\ddot{S}_i^\top S_i &= -m\dot{S}_i^\top\dot{S}_iS_i^\top S_i - \gamma\dot{S}_i^\top S_i - \Xi_iS_i^\top S_i + \frac{m}{\gamma} \left(-2\Xi_i\dot{S}_i^\top S_i + \dot{S}_i^\top S_i\Xi_iS_i^\top S_i - \Xi_iS_i^\top\dot{S}_iS_i^\top S_i \right) \\ &\quad + \frac{\kappa}{2N} \sum_{k=1}^N \left[S_k^\top S_i - \frac{1}{2}(S_k^\top S_iS_i^\top S_i + S_i^\top S_kS_i^\top S_i) \right]. \end{aligned} \quad (2.7)$$

We add (2.6) and (2.7) to get

$$\begin{aligned} m(S_i^\top\ddot{S}_i + \ddot{S}_i^\top S_i) &= -m(S_i^\top S_i\dot{S}_i^\top\dot{S}_i + \dot{S}_i^\top\dot{S}_iS_i^\top S_i) - \gamma(S_i^\top\dot{S}_i + \dot{S}_i^\top S_i) + [S_i^\top S_i, \Xi_i] \\ &\quad + \frac{m}{\gamma}\mathcal{J}_1 + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{J}_{2k}, \end{aligned} \quad (2.8)$$

where \mathcal{J}_1 and \mathcal{J}_{2k} are defined as

$$\begin{aligned} \mathcal{J}_1 &:= 2S_i^\top\dot{S}_i\Xi_i - S_i^\top S_i\Xi_iS_i^\top\dot{S}_i + S_i^\top S_i\dot{S}_i^\top S_i\Xi_i - 2\Xi_i\dot{S}_i^\top S_i + \dot{S}_i^\top S_i\Xi_iS_i^\top S_i - \Xi_iS_i^\top\dot{S}_iS_i^\top S_i, \\ \mathcal{J}_{2k} &:= S_i^\top S_k + S_k^\top S_i - \frac{1}{2}(S_i^\top S_iS_i^\top S_k + S_i^\top S_iS_k^\top S_i + S_k^\top S_iS_i^\top S_i + S_i^\top S_kS_i^\top S_i). \end{aligned}$$

We recall the notation:

$$H_i = I_p - S_i^\top S_i, \quad \dot{H}_i = -(\dot{S}_i^\top S_i + S_i^\top \dot{S}_i) \quad \text{and} \quad \ddot{H}_i = -(\ddot{S}_i^\top S_i + 2\dot{S}_i^\top \dot{S}_i + S_i^\top \ddot{S}_i).$$

Then, system (2.8) can be rewritten in terms of H_i and its derivatives:

$$m\ddot{H}_i + \gamma\dot{H}_i = -m(H_i\dot{S}_i^\top\dot{S}_i + \dot{S}_i^\top\dot{S}_iH_i) + [H_i, \Xi_i] - \frac{m}{\gamma}\mathcal{J}_1 - \frac{\kappa}{2N}\sum_{k=1}^N\mathcal{J}_{2k}. \quad (2.9)$$

In what follows, we present the estimates for \mathcal{J}_1 and \mathcal{J}_2 , respectively.

- (Estimate on \mathcal{J}_1): By direct calculation, one has

$$\begin{aligned} \mathcal{J}_1 &= 2S_i^\top\dot{S}_i\Xi_i - \Xi_iS_i^\top\dot{S}_i + \dot{S}_i^\top S_i\Xi_i + H_i(\dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i) \\ &\quad - 2\Xi_i\dot{S}_i^\top S_i + \dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i - (\dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i)H_i \\ &= 2\Xi_i\dot{H}_i - 2\dot{H}_i\Xi_i + H_i(\dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i) - (\dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i)H_i \\ &= 2[\Xi_i, \dot{H}_i] + [H_i, \dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i]. \end{aligned}$$

- (Estimate on \mathcal{J}_{2k}): Since the communication term of first-order and second-order models are same, it follows from Lemma 2.1 that

$$\mathcal{J}_{2k} = H_i(S_i^\top S_k + S_k^\top S_i) + (S_i^\top S_k + S_k^\top S_i)H_i.$$

In (2.9), we use the calculation of \mathcal{J}_1 and \mathcal{J}_2 to find the second-order (autonomous) matrix-valued equation for H_i :

$$\begin{aligned} m\ddot{H}_i + \gamma\dot{H}_i + mH_i(\dot{S}_i^\top\dot{S}_i) + m(\dot{S}_i^\top\dot{S}_i)H_i - [H_i, \Xi_i] + \frac{m}{\gamma}\left(2[\Xi_i, \dot{H}_i] + [H_i, \dot{S}_i^\top S_i\Xi_i - \Xi_iS_i^\top\dot{S}_i]\right) \\ + \frac{\kappa}{2N}\sum_{k=1}^N\left[H_i(S_i^\top S_k + S_k^\top S_i) + (S_i^\top S_k + S_k^\top S_i)H_i\right] = O. \end{aligned} \quad (2.10)$$

One can check that $H_i = O$ becomes a solution to (2.10) satisfying the initial assumption (2.5). Since a solution to the Cauchy problem (2.10) with the initial assumption (2.5) is unique, we conclude that

$$H_i(t) = O, \quad t > 0.$$

This yields the desired result. \square

Next, we focus on the situation in which solution operators for (2.4) can be expressed as a composition of two operators. In the proof of Lemma 2.3, we note that positive invariance of $\text{St}(p, n)$ is also valid, when the fourth term in the right-hand side of (2.4) is absent. However, the second-order model (2.4) satisfies such property for a homogeneous ensemble when the fourth term in the right-hand side of (2.4) is included.

Lemma 2.4. *Suppose that the initial data and frequency matrices satisfy*

$$S_i^{\text{in}} \in \text{St}(p, n), \quad \dot{S}_i^{t, \text{in}} S_i^{\text{in}} + S_i^{t, \text{in}} \dot{S}_i^{\text{in}} = O, \quad \Xi_i \equiv \Xi, \quad i = 1, \dots, N,$$

and let \mathcal{S} be a global solution to (2.4). Then, $Y_i := S_i e^{-\frac{\Xi}{\gamma}t}$ satisfies

$$m\ddot{Y}_i = -mY_i\dot{Y}_i^\top\dot{Y}_i - \gamma\dot{Y}_i + \frac{\kappa}{N}\sum_{k=1}^N\left(Y_k - \frac{1}{2}(Y_iY_i^\top Y_k + Y_iY_k^\top Y_i)\right).$$

Proof. By direct calculation, we use the ansatz for Y_i to find its derivatives:

$$\begin{aligned}\dot{Y}_i &= \left(\dot{S}_i - \frac{S_i \Xi_i}{\gamma} \right) e^{-\frac{\Xi_i}{\gamma} t}, \quad \ddot{Y}_i = \left(\ddot{S}_i - \frac{2\dot{S}_i \Xi_i}{\gamma} + \frac{S_i \Xi_i^2}{\gamma^2} \right) e^{-\frac{\Xi_i}{\gamma} t}, \\ Y_i^\top &= e^{\frac{\Xi_i}{\gamma} t} S_i^\top, \quad \dot{Y}_i^\top = e^{\frac{\Xi_i}{\gamma} t} \left(\dot{S}_i^\top + \frac{\Xi_i S_i^\top}{\gamma} \right).\end{aligned}$$

In what follows, we consider the three terms:

$$m\ddot{Y}_i + mY_i\dot{Y}_i^\top\dot{Y}_i, \quad \gamma\dot{Y}_i, \quad Y_k - \frac{1}{2}(Y_i Y_i^\top Y_k + Y_i Y_k^\top Y_i).$$

- (Estimate of $m\ddot{Y}_i + mY_i\dot{Y}_i^\top\dot{Y}_i$): we observe

$$\begin{aligned}m\ddot{Y}_i + mY_i\dot{Y}_i^\top\dot{Y}_i &= m \left(\ddot{S}_i - \frac{2\dot{S}_i \Xi_i}{\gamma} + \frac{S_i \Xi_i^2}{\gamma^2} \right) e^{-\frac{\Xi_i}{\gamma} t} + mS_i \left(\dot{S}_i^\top \dot{S}_i + \frac{\Xi_i S_i^\top \dot{S}_i}{\gamma} - \frac{\dot{S}_i^\top S_i \Xi_i}{\gamma} - \frac{\Xi_i^2}{\gamma^2} \right) e^{-\frac{\Xi_i}{\gamma} t} \\ &= \left(m\ddot{S}_i + mS_i \dot{S}_i^\top \dot{S}_i - \frac{2m\dot{S}_i \Xi_i}{\gamma} + \frac{mS_i \Xi_i S_i^\top \dot{S}_i}{\gamma} - \frac{mS_i \dot{S}_i^\top S_i \Xi_i}{\gamma} \right) e^{-\frac{\Xi_i}{\gamma} t}.\end{aligned}$$

- (Estimate of $\gamma\dot{Y}_i$): By direct calculation,

$$\gamma\dot{Y}_i = (\gamma\dot{S}_i - S_i \Xi_i) e^{-\frac{\Xi_i}{\gamma} t}.$$

- (Estimate of $Y_k - \frac{1}{2}(Y_i Y_i^\top Y_k + Y_i Y_k^\top Y_i)$): we use the ansatz of Y_i to find

$$Y_k - \frac{1}{2}(Y_i Y_i^\top Y_k + Y_i Y_k^\top Y_i) = \left(S_k - \frac{1}{2}(S_i S_i^\top S_k + S_i S_k^\top S_i) \right) e^{-\frac{\Xi_i}{\gamma} t}.$$

Finally, we combine the calculations above to find our desired result:

$$\begin{aligned}m\ddot{Y}_i + mY_i\dot{Y}_i^\top\dot{Y}_i + \gamma\dot{Y}_i - \frac{\kappa}{N} \sum_{k=1}^N \left(Y_k - \frac{1}{2}(Y_i Y_i^\top Y_k + Y_i Y_k^\top Y_i) \right) \\ = \left(m\ddot{S}_i + mS_i \dot{S}_i^\top \dot{S}_i + \gamma\dot{S}_i - S_i \Xi_i - \frac{m}{\gamma}(2\dot{S}_i \Xi_i - S_i \Xi_i S_i^\top \dot{S}_i + S_i \dot{S}_i^\top S_i \Xi_i) \right. \\ \left. - \frac{\kappa}{N} \sum_{k=1}^N \left(S_k - \frac{1}{2}(S_i S_i^\top S_k + S_i S_k^\top S_i) \right) \right) e^{-\frac{\Xi_i}{\gamma} t} = O.\end{aligned}$$

□

We finally close this section by introducing second-order Grönwall-type inequalities.

Lemma 2.5. [9, 24] *Let $y = y(t)$ be a nonnegative C^2 -function satisfying the following differential inequality:*

$$a\ddot{y} + b\dot{y} + cy \leq \varepsilon(t), \quad t > 0.$$

Then, the following estimates hold:

(1) Suppose that $b^2 - 4ac > 0$ and $\varepsilon(t) \equiv \varepsilon_0$ is a given positive constant. Then, we have

$$y(t) \leq \frac{\varepsilon_0}{c} + \left(y(0) + \frac{d}{c}\right) e^{-\nu_1 t} + \frac{a}{\sqrt{b^2 - 4ac}} \left(y'(0) + \nu_1 y(0) - \frac{2\varepsilon_0}{b - \sqrt{b^2 - 4ac}}\right) (e^{-\nu_2 t} - e^{-\nu_1 t}).$$

where ν_1 and ν_2 are given as follows:

$$\nu_1 := \frac{b + \sqrt{b^2 - 4ac}}{2a}, \quad \nu_2 := \frac{b - \sqrt{b^2 - 4ac}}{2a}.$$

Then, one has

$$\limsup_{t \rightarrow \infty} y(t) \leq \frac{\varepsilon_0}{c}.$$

(2) Suppose that $b^2 - 4ac < 0$. Then, we have

$$y(t) \leq \frac{4a\varepsilon_0}{b^2} + \left(y(0) - \frac{4a\varepsilon_0}{b^2} + \left(\frac{b}{2a}y(0) + y'(0) - \frac{2\varepsilon_0}{b}\right)t\right) e^{-\frac{b}{2a}t}.$$

Moreover,

$$\limsup_{t \rightarrow \infty} y(t) \leq \frac{4a\varepsilon_0}{b^2}.$$

(3) We furthermore assume that $\varepsilon(t)$ is a nonnegative continuously differentiable function decaying to zero as $t \rightarrow \infty$. Then, we have

$$\lim_{t \rightarrow \infty} y(t) = 0.$$

Proof. For the first two assertions, we refer the reader to [9, Lemma 3.1]. On the other hand for the last assertion, we refer the reader to [24, Lemma 4.9]. \square

3. FIRST-ORDER CONSENSUS MODEL

In this section, we study the asymptotic behavior of the first-order model. In Section 3.1, we consider the homogeneous ensemble in which all natural frequencies are the same so that the complete consensus can be achieved. In Section 3.2, we are concerned with the heterogeneous ensemble where that phase-locked states can arise under some framework with a large coupling strength regime.

Recall the synchronization quantity $H_{ij} \in M_{p,p}(\mathbb{R})$:

$$H_{ij} := I_p - S_i^\top S_j, \quad i, j = 1, \dots, N.$$

Then, we observe from the definition of $\text{St}(p, n)$

$$\|S_i - S_j\|_{\mathbb{F}}^2 = \text{tr}(I_p - S_i^\top S_j + I_p - S_j^\top S_i) = \text{tr}(H_{ij} + H_{ji}) \leq \sqrt{p} \|H_{ij} + H_{ji}\|_{\mathbb{F}}, \quad (3.1)$$

where we used the inequality:

$$\text{tr}(AB) \leq \|A\|_{\mathbb{F}} \|B\|_{\mathbb{F}} \quad \text{for } A = I_p \text{ and } B = H_{ij} + H_{ji}.$$

On the other hand, we see

$$\begin{aligned} \|H_{ij} + H_{ji}\|_{\mathbb{F}} &= \|I_p - S_i^\top S_j + I_p - S_j^\top S_i\|_{\mathbb{F}} = \|S_i^\top S_i - S_i^\top S_j + S_j^\top S_j - S_j^\top S_i\|_{\mathbb{F}} \\ &\leq \|(S_i^\top - S_j^\top)(S_i - S_j)\|_{\mathbb{F}} \leq \|S_i - S_j\|_{\mathbb{F}}^2. \end{aligned} \quad (3.2)$$

In (3.1) and (3.2), one has

$$\frac{1}{\sqrt{p}} \|S_i - S_j\|_F^2 \leq \|H_{ij} + H_{ji}\|_F \leq \|S_i - S_j\|_F^2. \quad (3.3)$$

Thus, the following equivalence holds:

$$\lim_{t \rightarrow \infty} \|S_i - S_j\|_F = 0 \iff \lim_{t \rightarrow \infty} \|H_{ij} + H_{ji}\|_F = 0.$$

3.1. A homogeneous ensemble. In this subsection, we consider a homogeneous ensemble where all frequency matrices are same:

$$\Xi_i \equiv O, \quad i = 1, \dots, N.$$

Lemma 3.1. *Let S be a solution to (1.3) with $\Xi_i \equiv \Xi$. Then, $\|S_i - S_j\|_F^2$ satisfies*

$$\begin{aligned} \frac{d}{dt} \|S_i - S_j\|^2 &\leq -\frac{\kappa}{N} \sum_{k=1}^N a_{ik} (\|S_i - S_j\|^2 - \|S_j - S_k\|^2) - \frac{\kappa}{N} \sum_{k=1}^N a_{jk} (\|S_i - S_j\|^2 - \|S_i - S_k\|^2) \\ &\quad - (2 - \|S_i - S_j\|^2) \cdot \frac{\kappa}{2N} \sum_{k=1}^N (a_{ik} \|S_i - S_k\|^2 + a_{jk} \|S_j - S_k\|^2). \end{aligned} \quad (3.4)$$

Moreover, for the lower bound estimate, one has

$$\frac{d}{dt} d_{ij}^2 \geq -4\kappa a_M d_{ij}^2, \quad t > 0.$$

Proof. First, we derive the dynamics for $\frac{d}{dt}(S_j^\top S_i) = \dot{S}_j^\top S_i + S_j^\top \dot{S}_i$. For this, we observe

$$\begin{aligned} S_j^\top \dot{S}_i &= S_j^\top S_i \Xi_i + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left[S_j^\top S_k - \frac{1}{2} (S_j^\top S_i S_i^\top S_k + S_j^\top S_i S_k^\top S_i) \right], \\ \dot{S}_j^\top S_i &= -\Xi_j S_j^\top S_i + \frac{\kappa}{N} \sum_{k=1}^N a_{jk} \left[S_k^\top S_i - \frac{1}{2} (S_k^\top S_j S_j^\top S_i + S_j^\top S_k S_j^\top S_i) \right]. \end{aligned} \quad (3.5)$$

Then, we calculate (3.5)₁ + (3.5)₂ to find the dynamics for $A_{ji} := S_j^\top S_i$:

$$\begin{aligned} \frac{d}{dt} A_{ji} &= A_{ji} \Xi_i - \Xi_j A_{ji} \\ &\quad + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left[A_{jk} - \frac{1}{2} (A_{ji} A_{ik} + A_{ji} A_{ki}) \right] + a_{jk} \left[A_{ki} - \frac{1}{2} (A_{kj} A_{ji} + A_{jk} A_{ji}) \right]. \end{aligned} \quad (3.6)$$

Again, if we recall the notation:

$$H_{ji} = I_p - A_{ji} = I_p - S_j^\top S_i,$$

then (3.6) directly yields the dynamics for H_{ji} :

$$\begin{aligned} \frac{d}{dt}H_{ji} &= (\Xi_j - \Xi_i) + (H_{ji}\Xi_i - \Xi_j H_{ji}) - \frac{\kappa}{N} \sum_{k=1}^N (a_{ik} + a_{jk})H_{ji} \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N a_{ik}(2H_{jk} - H_{ik} - H_{ki}) + a_{ik}(H_{ji}H_{ik} + H_{ji}H_{ki}) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N a_{jk}(2H_{ki} - H_{kj} - H_{jk}) + a_{jk}(H_{kj}H_{ji} + H_{jk}H_{ji}). \end{aligned} \quad (3.7)$$

We interchange the index $i \leftrightarrow j$ in (3.7) to get

$$\begin{aligned} \frac{d}{dt}H_{ij} &= (\Xi_i - \Xi_j) + (H_{ij}\Xi_j - \Xi_i H_{ij}) - \frac{\kappa}{N} \sum_{k=1}^N (a_{jk} + a_{ik})H_{ij} \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N a_{jk}(2H_{ik} - H_{jk} - H_{kj}) + a_{jk}(H_{ij}H_{jk} + H_{ij}H_{kj}) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N a_{ik}(2H_{kj} - H_{ki} - H_{ik}) + a_{ik}(H_{ki}H_{ij} + H_{ik}H_{ij}). \end{aligned} \quad (3.8)$$

We add (3.7) and (3.8) to find

$$\begin{aligned} \frac{d}{dt}(H_{ij} + H_{ji}) &= -\frac{\kappa}{N} \sum_{k=1}^N (a_{ik} + a_{jk})(H_{ij} + H_{ji}) + (H_{ji}\Xi_i - \Xi_j H_{ji} + H_{ij}\Xi_j - \Xi_i H_{ij}) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N (a_{jk} - a_{ik})(H_{ik} + H_{ki} - H_{jk} - H_{kj}) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N a_{ik}(H_{ji}(H_{ik} + H_{ki}) + (H_{ik} + H_{ki})H_{ij}) \\ &\quad + a_{jk}((H_{jk} + H_{kj})H_{ji} + H_{ij}(H_{jk} + H_{kj})). \end{aligned} \quad (3.9)$$

We take the trace in (3.9) to find

$$\begin{aligned} \frac{d}{dt}\|S_i - S_j\|^2 &= -\frac{\kappa}{N} \sum_{k=1}^N (a_{ik} + a_{jk})\|S_i - S_j\|^2 + \text{tr}((H_{ij} - H_{ji})(\Xi_i - \Xi_j)) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N (a_{jk} - a_{ik})(\|S_i - S_k\|^2 - \|S_j - S_k\|^2) \\ &+ \frac{\kappa}{2N} \sum_{k=1}^N \text{tr} \left[(a_{ik}(H_{ik} + H_{ki}) + a_{jk}(H_{jk} + H_{kj}))(H_{ij} + H_{ji}) \right]. \end{aligned} \quad (3.10)$$

Finally, we use $\Xi_i \equiv O$ and the inequality (3.3) to estimate the last term in (3.10):

$$\begin{aligned} \frac{d}{dt} \|S_i - S_j\|^2 &\leq -\frac{\kappa}{N} \sum_{k=1}^N a_{ik} (\|S_i - S_j\|^2 - \|S_j - S_k\|^2) - \frac{\kappa}{N} \sum_{k=1}^N a_{jk} (\|S_i - S_j\|^2 - \|S_i - S_k\|^2) \\ &\quad - (2 - \|S_i - S_j\|^2) \cdot \frac{\kappa}{2N} \sum_{k=1}^N (a_{ik} \|S_i - S_k\|^2 + a_{jk} \|S_j - S_k\|^2). \end{aligned}$$

□

We are now ready to prove a proof of Theorem 1.1. For this, we define the maximal diameter:

$$\mathcal{D}(\mathcal{S}(t)) := \max_{1 \leq i, j \leq N} \|S_i(t) - S_j(t)\|_F, \quad t \geq 0.$$

(Proof of Theorem 1.1): For each $t > 0$, we choose the maximal index (i_t, j_t) :

$$D(\mathcal{S}(t)) = \|S_{i_t} - S_{j_t}\|_F.$$

Then, it follows from (3.4) in Lemma 3.1 that $D(\mathcal{S})$ satisfies

$$\frac{d}{dt} \mathcal{D}(\mathcal{S})^2 \leq -(2 - \mathcal{D}(\mathcal{S})^2) \cdot \frac{\kappa}{2N} \sum_{k=1}^N (a_{ik} d_{i_t k}^2 + a_{jk} d_{j_t k}^2) \leq -\frac{\kappa a_m}{4} (2 - \mathcal{D}(\mathcal{S})^2) \mathcal{D}(\mathcal{S})^2.$$

Thus, the region $\{t \geq 0 : \mathcal{D}(\mathcal{S}(t)) < \sqrt{2}\}$ is positively invariant and the desired assertion directly follows from the dynamical systems theory. □

Remark 3.1. In Theorem 2.1, their initial framework leading to the complete consensus only depends on the relation of (p, n) (in fact, $p \leq \frac{2n}{3} - 1$), and the initial data \mathcal{S}^{in} is not restricted except for a measure zero set, so that the global consensus is achieved. In contrast, for our result, any (p, n) can be chosen; however, as a trade-off, smallness on the initial data would be imposed, for instance, $\mathcal{D}(\mathcal{S}^{\text{in}}) < \sqrt{2}$.

3.2. A heterogeneous ensemble. In this subsection, we deal with the heterogeneous ensemble, and provide a sufficient framework leading to the phase-locked state. In order to capture the phase-locked state, we briefly present our strategy introduced in [28] consisting of several steps:

- Step A: let \mathcal{S} and $\tilde{\mathcal{S}}$ two solutions to (1.3). Derive a temporal evolution of the maximal quantity (inter-diameter) denoted by $d(\mathcal{S}, \tilde{\mathcal{S}})$:

$$d(\mathcal{S}, \tilde{\mathcal{S}}) := \max_{1 \leq i, j \leq N} \|S_i^\top S_j - \tilde{S}_i^\top \tilde{S}_j\|_F = \max_{1 \leq i, j \leq N} \|A_{ij} - \tilde{A}_{ij}\|_F,$$

which measures the relative distance between two solution configurations \mathcal{S} and $\tilde{\mathcal{S}}$ (see Lemma 3.2).

- Step B: in Step A, since we need a smallness on the diameter $D(\mathcal{S})$, we find a positively invariant region of $\mathcal{D}(\mathcal{S})$ (see Lemma 3.3).

- Step C: together with a temporal evolution of $d(\mathcal{S}, \tilde{\mathcal{S}})$ and a positively invariant region of $\mathcal{D}(\mathcal{S})$, we show that $d(\mathcal{S}, \tilde{\mathcal{S}})$ converges to zero. Then, since our system is autonomous, $\tilde{S}_i(t) = S_i(t+T)$ also becomes a solution to the system for any T . By discretizing the time $t \in \mathbb{R}_+$ as $n \in \mathbb{Z}_+$ and setting $T = m \in \mathbb{Z}_+$, we deduce that $\{S_j^\top(n) S_i(n)\}_{n \in \mathbb{Z}_+}$ is indeed a

Cauchy sequence in the radius p -ball of $M_p(\mathbb{R})$ and hence each $S_j^\top S_i$ converges to a constant $p \times p$ matrix.

In what follows, we provide the detailed justification of Step A, Step B and Step C. For this, we assume

$$\Lambda := a_m - \frac{N-1}{N}(a_M + d(\mathcal{A})) \in (0, 8pa_M^2).$$

In other words, difference between the maximum and minimum would be small. Below, we study the temporal evolution of $d(\mathcal{S}, \tilde{\mathcal{S}})$ in the following lemma.

Lemma 3.2. *Let \mathcal{S} and $\tilde{\mathcal{S}}$ be two solutions to (1.3). Then, the inter-diameter $d(\mathcal{S}, \tilde{\mathcal{S}})$ satisfies*

$$\frac{d}{dt}d(\mathcal{S}, \tilde{\mathcal{S}}) \leq -2\kappa(\Lambda - a_M\sqrt{p}(\mathcal{D}(\mathcal{S}) + \mathcal{D}(\tilde{\mathcal{S}})))d(\mathcal{S}, \tilde{\mathcal{S}}), \quad t > 0.$$

Proof. Since the proof is lengthy, we leave its proof in Appendix A. \square

Next, we find a positively invariant region of $\mathcal{D}(\mathcal{S})$. For this, we define $0 < \alpha < \beta$ through the relation:

$$\alpha \text{ and } \beta \text{ are two positive roots of } r^3 - 2r + \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa_* a_m} = 0,$$

where κ_* is a positive constant defined by

$$\kappa_* := \frac{16p^2 a_M^3 \|\Xi\|_\infty}{a_m(8pa_M^2 \Lambda - \Lambda^3)}.$$

Lemma 3.3. *Suppose that the coupling strength and the initial data satisfy*

$$\kappa > \max \left\{ \frac{\sqrt{6p}\|\Xi\|_\infty}{9 a_m}, \frac{16p^2 a_M^3 \|\Xi\|_\infty}{a_m(8pa_M^2 \Lambda - \Lambda^3)} \right\}, \quad \mathcal{D}(\mathcal{S}^{\text{in}}) < \beta, \quad (3.11)$$

and let \mathcal{S} a global solution to (1.3). Then, there exists a finite entrance time $T_2 > 0$ such that

$$\mathcal{D}(\mathcal{S}(t)) < \alpha < \frac{\Lambda}{2a_M\sqrt{p}}, \quad t > T_2.$$

Proof. Recall the estimate (3.4) in Lemma 3.1 and find the differential inequality of $\mathcal{D}(\mathcal{S})$ for a heterogeneous ensemble:

$$\begin{aligned} \frac{d}{dt}\mathcal{D}(\mathcal{S}) &\leq -\frac{\kappa a_m}{2}\mathcal{D}(\mathcal{S}) + \frac{\kappa a_m}{4}\mathcal{D}(\mathcal{S})^3 + \frac{\sqrt{p}}{2}\|\Xi\|_\infty \\ &= \frac{\kappa a_m}{4} \left(-2\mathcal{D}(\mathcal{S}) + \mathcal{D}(\mathcal{S})^3 + \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa a_m} \right) =: f(\mathcal{D}(\mathcal{S})), \end{aligned}$$

where we introduce an auxiliary cubic function f :

$$f(r) = r^3 - 2r + \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa a_m}, \quad r \geq 0.$$

Then, it follows from the simple calculus that

$$f \text{ attains the minimum at } r_* := \sqrt{\frac{2}{3}} \text{ with } f(r_*) = \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa a_m} - \frac{2}{3}\sqrt{\frac{2}{3}}.$$

Thus, since $f(r_*) < 0$ is due to (3.11)₁, we see that f has two positive roots r_1 and r_2 such that

$$0 < r_1 < \sqrt{\frac{2}{3}}, \quad \sqrt{\frac{2}{3}} < r_2 < \sqrt{2}.$$

Note that (3.11)₁ indeed gives $f(r_*) < 0$. In addition, we observe

$$\lim_{\kappa \rightarrow \infty} r_1 = 0 \quad \lim_{t \rightarrow \infty} r_2 = \sqrt{2}.$$

We use the dynamical systems theory to show that if $\mathcal{D}(S^{\text{in}}) < r_2$, then there exists a finite entrance time $T_3 > 0$ such that

$$\mathcal{D}(S(t)) < r_1, \quad t > T_3.$$

More precisely, we split the case into two parts. First, suppose initial data satisfy

$$\mathcal{D}(S^{\text{in}}) \leq r_1.$$

Then, at the time $t = t_*$ when $\mathcal{D}(S(t_*)) = r_1$, we have

$$\frac{d}{dt}\mathcal{D}(S) \leq 0.$$

Thus, $\mathcal{D}(S)$ does not increase at $t = t_*$ and hence is restricted in the interval $[0, r_1]$. Second, suppose that the initial data satisfy $r_1 < \mathcal{D}(S^{\text{in}}) < r_2$. Again, at the instant time $t = t_*$ when $r_1 < \mathcal{D}(S(t_*)) < r_2$,

$$\frac{d}{dt}\mathcal{D}(S) < 0.$$

Thus, $\mathcal{D}(S)$ starts to strictly decrease. Then, by applying same argument in Proposition 3.1 in [8], we find such finite entrance time $T_3 > 0$.

In order to find the desired invariant region, we have to assume

$$f\left(\frac{\Lambda}{2a_M\sqrt{p}}\right) = \left(\frac{\Lambda}{2a_M\sqrt{p}}\right)^3 - 2 \cdot \left(\frac{\Lambda}{2a_M\sqrt{p}}\right) + \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa a_m} < 0, \quad (3.12)$$

and one can check from algebraic manipulation that the relation (3.12) is equivalent to (3.11)₁. Then, the cubic equation

$$r^3 - 2r + \frac{2\sqrt{p}\|\Xi\|_\infty}{\kappa_* a_m} = 0$$

has two positive roots: α and β . Hence, under the assumption (3.11), there exists such finite entrance time $T_2 > 0$ so that

$$\mathcal{D}(S(t)) < \alpha < \frac{\Lambda}{2a_M\sqrt{p}}, \quad t > T_2.$$

□

Finally, we use Lemma 3.2 and Lemma 3.3 to present a proof of Theorem 1.2.

(Proof of Theorem 1.2): Since we assume (3.11), we use Lemmas 3.2 and 3.3 to show that

$$\frac{d}{dt}d(S, \tilde{S}) \leq -2\kappa(\Lambda - 2a_M\sqrt{p}\alpha)d(S, \tilde{S}), \quad \text{a.e. } t > T_2.$$

This implies

$$d(\mathcal{S}, \tilde{\mathcal{S}})(t) \leq d(\mathcal{S}, \tilde{\mathcal{S}})(T_2) e^{-2\kappa(\Lambda - 2a_M \sqrt{p\alpha})(t - T_2)}, \quad \text{a.e. } t > T_2. \quad (3.13)$$

Since we are only interested in large-time behavior, without loss of generality, we would set $T_2 = 0$. On the other hand for any $T \geq 0$, a shifted solution $\{S_i(t + T)\}$ also becomes a global solution to (1.3) with a shifted initial data $\{S_i(T)\}$, as (1.3) is autonomous. Then, (3.13) yields

$$\|S_j^\top S_i(t + T) - S_j^\top S_i(t)\| \leq \max_{1 \leq i, j \leq N} \|S_j^\top S_i(T) - S_j^\top S_i(0)\| e^{-2\kappa(\Lambda - 2a_M \sqrt{p\alpha})t}, \quad t \geq 0. \quad (3.14)$$

Especially for $T = 1$ and $t = n \in \mathbb{Z}_+$ in (3.14), we have

$$\|S_j^\top S_i(n+1) - S_j^\top S_i(n)\| \leq \max_{1 \leq i, j \leq N} \|S_j^\top S_i(1) - S_j^\top S_i(0)\| e^{-2\kappa(\Lambda - 2a_M \sqrt{p\alpha})n}, \quad n \in \mathbb{Z}_+. \quad (3.15)$$

By induction argument, (3.15) gives for $m \in \mathbb{Z}_+$,

$$\|S_j^\top S_i(n+m) - S_j^\top S_i(n)\| \leq \max_{1 \leq i, j \leq N} \|S_j^\top S_i(1) - S_j^\top S_i(0)\| \frac{e^{-2\kappa(\Lambda - 2a_M \sqrt{p\alpha})n}}{1 - e^{-2\kappa(\Lambda - 2a_M \sqrt{p\alpha})}}, \quad n \in \mathbb{Z}_+.$$

Hence, the discretized sequence $\{S_j^\top(n)S_i(n)\}_{n \in \mathbb{Z}_+}$ is indeed Cauchy in the p -ball $B_p(O) := \{X \in M_p(\mathbb{R}) : \|X\|_F \leq p\}$. Consequently, for each i, j , it converges to a constant $p \times p$ matrix $\Gamma_{ji}^\infty \in B_p(O)$. \square

As a corollary, we show that normalized velocities synchronize.

Corollary 3.1. *Suppose that the coupling strength and initial data satisfy (3.11) and let \mathcal{S} be a global solution to (1.3). Then, the normalized velocities $S_i^\top \dot{S}_i$ and $S_j^\top \dot{S}_j$ synchronize:*

$$\lim_{t \rightarrow \infty} \|S_i^\top \dot{S}_i - S_j^\top \dot{S}_j\|_F = 0.$$

Proof. For two solutions \mathcal{S} and $\tilde{\mathcal{S}}$, we observe from (2.3)

$$\begin{aligned} S_i^\top \dot{S}_i - \tilde{S}_i^\top \dot{\tilde{S}}_i &= \left[\Xi_i + \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} (S_i^\top S_k - S_k^\top S_i) \right] - \left[\Xi_i + \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} (\tilde{S}_i^\top \tilde{S}_k - \tilde{S}_k^\top \tilde{S}_i) \right] \\ &= \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} \left[(S_i^\top S_k - \tilde{S}_i^\top \tilde{S}_k) - (S_k^\top S_i - \tilde{S}_k^\top \tilde{S}_i) \right]. \end{aligned}$$

Thus, we find

$$\|S_i^\top \dot{S}_i - \tilde{S}_i^\top \dot{\tilde{S}}_i\|_F \leq \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} (\|S_i^\top S_k - \tilde{S}_i^\top \tilde{S}_k\| + \|S_k^\top S_i - \tilde{S}_k^\top \tilde{S}_i\|) \leq \kappa a_M d(\mathcal{S}, \tilde{\mathcal{S}}).$$

Finally, we use the same argument in Theorem 1.2 to show that $\{S_i^\top \dot{S}_i\}$ becomes a Cauchy sequence and hence it converges to the same constant matrix. This shows the desired result. \square

4. A SECOND-ORDER CONSENSUS MODEL

In this section, we study the emergent dynamics of the second-order model (1.4). Before we present the estimates, energy functionals are defined and their temporal evolutions are derived as a crucial step. After that, the homogeneous ensemble are considered in Section 4.1 and we provide a sufficient framework leading to the complete consensus. On the other hand, we deal with the heterogeneous ensemble in Section 4.2 and present a sufficient framework for the practical consensus in the large coupling strength and small inertia regime.

As in [25], we define a total energy functional associated to (1.4):

$$\mathcal{E} =: \underbrace{\frac{m}{N} \sum_{i=1}^N \|\dot{S}_i\|_{\mathbb{F}}^2}_{=: \mathcal{K}} + \underbrace{\frac{\kappa}{2N^2} \sum_{i,j=1}^N a_{ij} \|S_i - S_j\|_{\mathbb{F}}^2}_{=: \mathcal{L}},$$

where \mathcal{K} represents a (total) kinetic energy and \mathcal{L} describes the interaction energy between the states of particles $\{S_i\}$ measuring the degree of consensus.

Next, we study the temporal-evolution of the total energy \mathcal{E} . First, we differentiate $\|\dot{S}_i\|_{\mathbb{F}}^2$ with respect to t to find

$$m \frac{d}{dt} \|\dot{S}_i\|_{\mathbb{F}}^2 = m \frac{d}{dt} \text{tr}(\dot{S}_i^\top \dot{S}_i) = m \text{tr}(\ddot{S}_i^\top \dot{S}_i + \dot{S}_i^\top \ddot{S}_i) = m \text{tr}(\dot{S}_i^\top \ddot{S}_i + (\dot{S}_i^\top \ddot{S}_i)^\top).$$

First, we observe the term $m \dot{S}_i^\top \ddot{S}_i$:

$$\begin{aligned} m \dot{S}_i^\top \ddot{S}_i &= -m \dot{S}_i^\top S_i \dot{S}_i^\top \dot{S}_i - \gamma \dot{S}_i^\top \dot{S}_i + \dot{S}_i^\top S_i \Xi_i + \frac{m}{\gamma} (2 \dot{S}_i^\top \dot{S}_i \Xi_i - \dot{S}_i^\top S_i \Xi_i S_i^\top \dot{S}_i + \dot{S}_i^\top S_i \dot{S}_i^\top S_i \Xi_i) \\ &\quad + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left(\dot{S}_i^\top S_k - \frac{1}{2} (\dot{S}_i^\top S_i S_i^\top S_k + \dot{S}_i^\top S_i S_k^\top S_i) \right) \\ &=: m \mathcal{I}_1 - \gamma \mathcal{I}_2 + \mathcal{I}_3 + \frac{m}{\gamma} \mathcal{I}_4 + \frac{\kappa}{N} \sum_{k=1}^N \mathcal{I}_5. \end{aligned} \tag{4.1}$$

In next lemma, we provide estimates for \mathcal{I}_k , $k = 1, \dots, 5$.

Lemma 4.1. *Let \mathcal{I}_k , $k = 1, \dots, 5$ be the terms defined in (4.1). Then, one has*

$$\begin{aligned} \text{tr}(\mathcal{I}_1 + \mathcal{I}_1^\top) &= 0, \quad \text{tr}(\mathcal{I}_2 + \mathcal{I}_2^\top) = 2 \|\dot{S}_i\|_{\mathbb{F}}^2, \quad \text{tr}(\mathcal{I}_3 + \mathcal{I}_3^\top) = \text{tr}(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i) \\ \text{tr}(\mathcal{I}_4 + \mathcal{I}_4^\top) &= 0, \quad \text{tr}(\mathcal{I}_5 + \mathcal{I}_5^\top) = a_{ik} \text{tr}(S_i^\top S_k + S_k^\top S_i). \end{aligned}$$

Proof. Below, we present estimates of $\text{tr}(\mathcal{I}_k + \mathcal{I}_k^\top)$, $k = 1, \dots, 5$, respectively.

- (Estimate of $\text{tr}(\mathcal{I}_1 + \mathcal{I}_1^\top)$): we recall the identity $\dot{S}_i^\top S_i + S_i^\top \dot{S}_i = O_p$ to see

$$\text{tr}(\mathcal{I}_1 + \mathcal{I}_1^\top) = \text{tr}(\dot{S}_i^\top S_i \dot{S}_i^\top \dot{S}_i + \dot{S}_i^\top \dot{S}_i S_i^\top \dot{S}_i) = \text{tr} \left[\dot{S}_i^\top \dot{S}_i \left(\dot{S}_i^\top S_i + S_i^\top \dot{S}_i \right) \right] = 0.$$

- (Estimate of $\text{tr}(\mathcal{I}_2 + \mathcal{I}_2^\top)$): by direct observation,

$$\text{tr}(\mathcal{I}_2 + \mathcal{I}_2^\top) = 2 \text{tr}(\mathcal{I}_2) = 2 \|\dot{S}_i\|_{\mathbb{F}}^2.$$

- (Estimate of $\text{tr}(\mathcal{I}_3 + \mathcal{I}_3^\top)$): we use the property of the trace $\text{tr}(AB) = \text{tr}(BA)$ to find

$$\text{tr}(\mathcal{I}_3 + \mathcal{I}_3^\top) = \text{tr}\left(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i\right).$$

- (Estimate of $\text{tr}(\mathcal{I}_4 + \mathcal{I}_4^\top)$): we first see

$$\begin{aligned} \text{tr}(\mathcal{I}_4) &= \text{tr}\left(\left(2\dot{S}_i^\top \dot{S}_i - S_i^\top \dot{S}_i \dot{S}_i^\top S_i + \dot{S}_i^\top S_i \dot{S}_i^\top S_i\right)\Xi_i\right) \\ &= \text{tr}\left(\left(2\dot{S}_i^\top \dot{S}_i - S_i^\top \dot{S}_i \dot{S}_i^\top S_i - \dot{S}_i^\top S_i S_i^\top \dot{S}_i\right)\Xi_i\right). \end{aligned} \quad (4.2)$$

Since the matrix $2\dot{S}_i^\top \dot{S}_i - S_i^\top \dot{S}_i \dot{S}_i^\top S_i - \dot{S}_i^\top S_i S_i^\top \dot{S}_i$ in (4.2) is symmetric and Ξ_i is skew-symmetric, we obtain

$$\text{tr}(\mathcal{I}_4 + \mathcal{I}_4^\top) = 0.$$

- (Estimate of $\text{tr}(\mathcal{I}_5 + \mathcal{I}_5^\top)$): we again use the identity $\dot{S}_i^\top S_i + S_i^\top \dot{S}_i = 0$ to see

$$\begin{aligned} \text{tr}(\mathcal{I}_5 + \mathcal{I}_5^\top) &= a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i) - \frac{a_{ik}}{2} \text{tr}(\dot{S}_i^\top S_i S_i^\top S_k + \dot{S}_i^\top S_i S_k^\top S_i + S_k^\top S_i S_i^\top \dot{S}_i + S_i^\top S_k S_i^\top \dot{S}_i) \\ &= a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i) - \frac{a_{ik}}{2} \text{tr}[\dot{S}_i^\top S_i (S_i^\top S_k + S_k^\top S_i) + (S_k^\top S_i + S_i^\top S_k) S_i^\top \dot{S}_i] \\ &= a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i) - \frac{a_{ik}}{2} \text{tr}[(\dot{S}_i^\top S_i + S_i^\top \dot{S}_i)(S_i^\top S_k + S_k^\top S_i)] = a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i). \end{aligned}$$

□

Next, we provide an estimate on the time-rate of change of the total energy.

Proposition 4.1. *Let S be a global solution to (1.4). Then, the total energy functional \mathcal{E} satisfies*

$$\frac{d\mathcal{E}}{dt} = -\frac{2\gamma}{N} \sum_{i=1}^N \|\dot{S}_i\|_F^2 + \frac{1}{N} \sum_{i=1}^N \text{tr}(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i). \quad (4.3)$$

Proof. We collect all the estimates of $\text{tr}(\mathcal{I}_i + \mathcal{I}_i^\top)$ for $i = 1, \dots, 5$ to find

$$m \frac{d}{dt} \|\dot{S}_i\|_F^2 = -2\gamma \|\dot{S}_i\|_F^2 + \text{tr}(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i) + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i), \quad (4.4)$$

and sum (4.4) over all i and divide the resulting relation by N to obtain

$$\begin{aligned} \frac{m}{N} \frac{d}{dt} \sum_{i=1}^N \|\dot{S}_i\|_F^2 &= -2\frac{\gamma}{N} \sum_{i=1}^N \|\dot{S}_i\|_F^2 + \frac{1}{N} \sum_{i=1}^N \text{tr}(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i) + \kappa \text{tr}(\dot{S}_c^\top S_c + S_c^\top \dot{S}_c) \\ &= -2\frac{\gamma}{N} \sum_{i=1}^N \|\dot{S}_i\|_F^2 + \frac{1}{N} \sum_{i=1}^N \text{tr}(\dot{S}_i^\top S_i \Xi_i - \Xi_i S_i^\top \dot{S}_i) + \frac{\kappa}{N^2} \sum_{i,k=1}^N a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i). \end{aligned} \quad (4.5)$$

On the other hand, we observe

$$\begin{aligned} \frac{\kappa}{N^2} \sum_{i,k=1}^N a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i) &= \frac{\kappa}{2N^2} \sum_{i,k=1}^N a_{ik} \text{tr}(\dot{S}_i^\top S_k + S_k^\top \dot{S}_i + \dot{S}_k^\top S_i + S_i^\top \dot{S}_k) \\ &= \frac{\kappa}{2N^2} \frac{d}{dt} \left(\sum_{i,k=1}^N a_{ik} \text{tr}(S_i^\top S_k + S_k^\top S_i) \right) = -\frac{\kappa}{2N^2} \frac{d}{dt} \left(\sum_{k=1}^N a_{ik} \|S_i - S_k\|^2 \right). \end{aligned}$$

Finally in (4.5), we obtain the desired dissipation energy estimate. \square

Remark 4.1. *It follows from (4.4) that*

$$m \frac{d}{dt} \|\dot{S}_i\|_{\mathbb{F}}^2 \leq -2\gamma \|\dot{S}_i\|_{\mathbb{F}}^2 + 2\|\dot{S}_i\|_{\mathbb{F}} \cdot \|S_i \Xi_i\|_{\mathbb{F}} + 2\kappa a_M \sqrt{p} \|\dot{S}_i\|_{\mathbb{F}}, \quad t > 0,$$

or equivalently

$$m \frac{d}{dt} \|\dot{S}_i\|_{\mathbb{F}} \leq -\gamma \|\dot{S}_i\|_{\mathbb{F}} + \|\Xi_i\|_{\mathbb{F}} + \kappa a_M \sqrt{p}, \quad t > 0.$$

We set the maximal value of $\|\Xi_i\|_{\mathbb{F}}$:

$$\|\Xi\|_{\infty} := \max_{1 \leq i \leq N} \|\Xi_i\|_{\mathbb{F}}.$$

Then, Grönwall's lemma yields a uniform boundedness and uniform continuity of $\|\dot{S}_i\|_{\mathbb{F}}$:

$$\sup_{0 \leq t < \infty} \|\dot{S}_i(t)\|_{\mathbb{F}} \leq \max \left\{ \max_{1 \leq i \leq N} \|\dot{S}_i^{\text{in}}\|_{\mathbb{F}}, \frac{1}{\gamma} (\|\Xi\|_{\infty} + \kappa a_M \sqrt{p}) \right\}. \quad (4.6)$$

In what follows, we impose the condition on the network topology:

$$\frac{1}{N} \sum_{k=1}^N a_{ik} \equiv \xi > 0, \quad i = 1, \dots, N.$$

In other words, every columns (or rows) have a common average ξ . We denote several quantities:

$$\mathcal{G}(t) := \frac{1}{N^2} \sum_{i,j=1}^N \|S_i(t) - S_j(t)\|_{\mathbb{F}}^2, \quad \mathcal{D}(\dot{S}(t)) := \max_{1 \leq i \leq N} \|\dot{S}_i(t)\|_{\mathbb{F}}, \quad \|\Xi_i\|_{\infty} := \max_{1 \leq i \leq N} \|\Xi_i\|_{\mathbb{F}}.$$

Below, we derive a second-order differential inequality for \mathcal{G} .

Lemma 4.2. *Let S be a solution to (1.4). Then, \mathcal{G} satisfies*

$$m\ddot{\mathcal{G}} + \gamma\dot{\mathcal{G}} + 2\kappa\xi\mathcal{G} \leq 16m\mathcal{D}(\dot{S})^2 + 8\|\Xi\|_{\infty} + \frac{16m\sqrt{p}\|\Xi\|_{\infty}}{\gamma}\mathcal{D}(\dot{S}). \quad (4.7)$$

Proof. Since the proof is lengthy, we postpone the justification of (4.7) in Appendix B. \square

Below, we quote Barbalat's lemma without proofs.

Lemma 4.3. [1] (i) *Suppose that a real-valued function $f : [0, \infty) \rightarrow \mathbb{R}$ is uniformly continuous and satisfies*

$$\lim_{t \rightarrow \infty} \int_0^t f(s) \, ds \quad \text{exists.}$$

Then, f tends to zero as $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} f(t) = 0.$$

(ii) Suppose that a real-valued function $f : [0, \infty) \rightarrow \mathbb{R}$ is continuously differentiable, and $\lim_{t \rightarrow \infty} f(t) = \alpha \in \mathbb{R}$. If f' is uniformly continuous, then

$$\lim_{t \rightarrow \infty} f'(t) = 0.$$

4.1. A homogeneous ensemble. In this part, we consider a homogeneous ensemble whose frequency matrices are the same, i.e., $\Xi_i \equiv \Xi$. Due to the property in Lemma 2.4 for a homogeneous ensemble, without loss of generality, we may set

$$\Xi_i \equiv O, \quad i = 1, \dots, N.$$

We are now ready to provide a proof of Theorem 1.3.

(Proof of Theorem 1.3) For the zero convergence of $\|\dot{S}_i\|_{\mathbb{F}}$, we use in (4.3) the condition $\Xi_i \equiv O$ to get

$$\frac{d\mathcal{E}}{dt} = -\frac{2\gamma}{N} \sum_{i=1}^N \|\dot{S}_i\|_{\mathbb{F}}^2, \quad t > 0.$$

We integrate the above equation on the interval $[0, t]$ to obtain

$$\frac{2\gamma}{N} \int_0^t \sum_{i=1}^N \|\dot{S}_i(s)\|_{\mathbb{F}}^2 ds = \mathcal{E}(0) - \mathcal{E}(t) \leq \mathcal{E}(0).$$

Hence, one has

$$\int_0^\infty \|\dot{S}_i(t)\|_{\mathbb{F}}^2 dt < \infty.$$

It follows from Remark 4.1 that $\left| \frac{d}{dt} \|\dot{S}_i\|_{\mathbb{F}} \right|$ is uniformly bounded, which implies the uniform continuity of $\|\dot{S}_i\|_{\mathbb{F}}$. Hence, we can apply Barbalat's lemma [1] to derive the desired zero convergence of $\|\dot{S}_i\|_{\mathbb{F}}$. On the other hand for the zero convergence of \mathcal{G} , we recall the differential inequality (4.7). Then, since we have $\|\Xi\|_\infty = 0$ and $\mathcal{D}(\dot{S})$ converges to zero, we apply Lemma 2.5 to obtain the desired result. \square

4.2. A heterogeneous ensemble. In this part, we study a heterogeneous ensemble and present a proof of Theorem 1.4. For this, we impose a small inertia assumption such that there exist $m_0 > 0$ which does not depend on κ and $\eta > 0$ satisfying

$$m = \frac{m_0}{\kappa^{1+\eta}}.$$

Since κ will tend to infinity, the relation above is referred as a small inertial assumption. The reason why we need such assumption is clarified throughout the proof.

(Proof of Theorem 1.4): since we assume (1.7)₁, it follows from (4.6) that

$$\mathcal{D}(\dot{S}(t)) < \frac{1}{\gamma} (\|\Xi\|_\infty + \kappa a_M \sqrt{p}).$$

Then, the right-hand side of (4.7) becomes

$$\begin{aligned}
& 16mD(\dot{\mathcal{S}})^2 + 8\|\Xi\|_\infty + \frac{16m\sqrt{p}\|\Xi\|_\infty}{\gamma}\mathcal{D}(\dot{\mathcal{S}}) \\
& \leq \frac{16m}{\gamma^2}(\|\Xi\|_\infty + \kappa a_M\sqrt{p})^2 + \frac{16m\sqrt{p}\|\Xi\|_\infty}{\gamma^2}(\|\Xi\|_\infty + \kappa a_M\sqrt{p}) + 8\|\Xi\|_\infty \\
& = \frac{16m}{\gamma^2}(\|\Xi\|_\infty + \kappa a_M\sqrt{p})(\|\Xi\|_\infty + \kappa a_M\sqrt{p} + \sqrt{p}\|\Xi\|_\infty) + 8\|\Xi\|_\infty \\
& = \mathcal{O}(m\kappa^2) + 8\|\Xi\|_\infty = \mathcal{O}(m\kappa^2) + \mathcal{O}(1).
\end{aligned}$$

In order to apply second-order Grönwall's inequality in Lemma 2.5 is used, we consider two cases:

$$(i) \quad \gamma^2 - 8\xi m\kappa > 0. \quad (ii) \quad \gamma^2 - 8\xi m\kappa < 0. \quad (4.8)$$

For the first case, we have

$$\limsup_{t \rightarrow \infty} \mathcal{G}(t) \leq \frac{1}{2\xi\kappa}(\mathcal{O}(m\kappa^2) + \mathcal{O}(1)) = \mathcal{O}(m\kappa) + \mathcal{O}(\kappa^{-1}) = \mathcal{O}(\kappa^{-\eta}) + \mathcal{O}(\kappa^{-1}), \quad (4.9)$$

where we used the relation (1.7)₂ in the last equality. On the other hand for the second case, we have

$$\limsup_{t \rightarrow \infty} \mathcal{G}(t) \leq \frac{4m}{\gamma^2}(\mathcal{O}(m\kappa^2) + \mathcal{O}(1)) = \mathcal{O}(m^2\kappa^2) + \mathcal{O}(m) = \mathcal{O}(\kappa^{-2\eta}) + \mathcal{O}(\kappa^{-(1+\eta)}), \quad (4.10)$$

where (1.7)₂ is used. By combining (4.9) and (4.10) and letting κ to infinity, we derive the desired estimate. \square

Remark 4.2. *Then, two conditions in (4.8) can be rewritten in terms of m_0 and η :*

$$(i) \quad \kappa > \left(\frac{8\xi m_0}{\gamma^2}\right)^{\frac{1}{\eta}}. \quad (ii) \quad \gamma < \left(\frac{8\xi m_0}{\kappa^\eta}\right)^{\frac{1}{2}}.$$

In (i), we only require a large coupling strength κ as $\kappa \rightarrow \infty$ and there is no assumption on γ . However for (ii), γ depends on κ . In fact, γ also tends to zero when $\kappa \rightarrow \infty$ as m does. To thi

5. CONCLUSION

In this paper, we have studied emergent dynamics of the first-order and second-order consensus models with both homogeneous and heterogeneous ensembles in which all states of the particles are restricted to the Stiefel manifold $\text{St}(p, n)$ consisting of all orthonormal k -frames in \mathbb{R}^n . As mentioned in Section 1, due to the structural property of Stiefel manifolds, it has been widely used for optimization problems especially in the engineering community. Our first-order model was first suggested in [41, 42] based on the gradient flow of the total distance between all particles. For a homogeneous ensemble in which frequency matrices are the same, we show that the relaxation process towards the complete consensus state is always achieved with an exponential rate. In addition, we provide a subset of initial data leading to the complete consensus independent of the choice (p, n) , whereas the results in [41, 42] show that the complete consensus is globally stable if (p, n) satisfies the specific relation and hence, the global consensus is achieved. On the other hand, for the heterogeneous ensemble, the phase-locked state can emerge under the large coupling strength regime, when the smallness assumption on the initial data is imposed. We also study the asymptotic

emergent dynamics of a second-order consensus model which naturally extends the first-order model by incorporating the inertial force. For a homogeneous ensemble, we present a sufficient framework leading to the complete consensus state based on a second-order Grönwall's inequality and energy estimate. In contrast, for a heterogeneous ensemble, a sufficient framework for the practical consensus state is provided under the large coupling and small inertia regime. In fact, the phase-locked state for the second-order model is not considered in this work. Thus, we leave this problem for a future work.

APPENDIX A. PROOF OF LEMMA 3.2

In this appendix, we provide a proof of Lemma 3.2 in which the differential inequality for $d(\mathcal{S}, \tilde{\mathcal{S}})$ is derived.

Recall the relation (3.6) in Lemma 3.1

$$\begin{aligned}
\frac{d}{dt}A_{ji} &= A_{ji}\Xi_i - \Xi_j A_{ji} \\
&\quad + \frac{\kappa}{N} \sum_{k=1}^N a_{ik} \left[A_{jk} - \frac{1}{2}(A_{ji}A_{ik} + A_{ji}A_{ki}) \right] + a_{jk} \left[A_{ki} - \frac{1}{2}(A_{kj}A_{ji} + A_{jk}A_{ji}) \right] \\
&= A_{ji}\Xi_i - \Xi_j A_{ji} + \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} (A_{jk} - A_{kj}) + a_{jk} (A_{ki} - A_{ik}) \\
&\quad + \frac{\kappa}{2N} \sum_{k=1}^N \left(a_{ik} (A_{jk} + A_{kj}) + a_{jk} (A_{ki} + A_{ik}) \right. \\
&\quad \quad \left. - a_{ik} A_{ji} (A_{ik} + A_{ki}) - a_{jk} (A_{jk} + A_{kj}) A_{ji} \right).
\end{aligned}$$

Similarly, we find

$$\begin{aligned}
\frac{d}{dt}\tilde{A}_{ji} &= \tilde{A}_{ji}\Xi_i - \Xi_j \tilde{A}_{ji} + \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} (\tilde{A}_{jk} - \tilde{A}_{kj}) + a_{jk} (\tilde{A}_{ki} - \tilde{A}_{ik}) \\
&\quad + \frac{\kappa}{2N} \sum_{k=1}^N \left(a_{ik} (\tilde{A}_{jk} + \tilde{A}_{kj}) + a_{jk} (\tilde{A}_{ki} + \tilde{A}_{ik}) \right. \\
&\quad \quad \left. - a_{ik} \tilde{A}_{ji} (\tilde{A}_{ik} + \tilde{A}_{ki}) - a_{jk} (\tilde{A}_{jk} + \tilde{A}_{kj}) \tilde{A}_{ji} \right).
\end{aligned}$$

Below, we consider the dynamics of $A_{ji} - \tilde{A}_{ji}$:

$$\begin{aligned}
\frac{d}{dt}(A_{ji} - \tilde{A}_{ji}) &= (A_{ji} - \tilde{A}_{ji})\Xi_i - \Xi_j (A_{ji} - \tilde{A}_{ji}) + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{J}_{3k} \\
&\quad + \frac{\kappa}{2N} \sum_{k=1}^N \left(a_{ik} ((A_{jk} - \tilde{A}_{jk}) - (A_{kj} - \tilde{A}_{kj})) \right. \\
&\quad \quad \left. + a_{jk} ((A_{ki} - \tilde{A}_{ki}) - (A_{ik} - \tilde{A}_{ik})) \right), \tag{A.1}
\end{aligned}$$

where \mathcal{J}_3 is defined as

$$\begin{aligned}
\mathcal{J}_{3k} &:= (a_{ik}(A_{jk} - \tilde{A}_{jk}) - a_{jk}(A_{jk}A_{ji} - \tilde{A}_{jk}\tilde{A}_{ji})) \\
&\quad + (a_{ik}(A_{kj} - \tilde{A}_{kj}) - a_{jk}(A_{kj}A_{ji} - \tilde{A}_{kj}\tilde{A}_{ji})) \\
&\quad + (a_{jk}(A_{ki} - \tilde{A}_{ki}) - a_{ik}(A_{ji}A_{ki} - \tilde{A}_{ji}\tilde{A}_{ji})) \\
&\quad + (a_{jk}(A_{ik} - \tilde{A}_{ik}) - a_{jk}(A_{ji}A_{ik} - \tilde{A}_{ji}\tilde{A}_{ik})) \\
&=: \mathcal{J}_{3k,1} + \mathcal{J}_{3k,2} + \mathcal{J}_{3k,3} + \mathcal{J}_{3k,4}.
\end{aligned} \tag{A.2}$$

Since $\mathcal{J}_{3k,j}$, $j = 1, \dots, 4$ has the same form, it suffices to provide the estimate for $\mathcal{J}_{3k,1}$:

$$\begin{aligned}
\mathcal{J}_{3k,1} &= a_{ik}(A_{jk} - \tilde{A}_{jk}) - a_{jk}(A_{jk}A_{ji} - \tilde{A}_{jk}\tilde{A}_{ji}) \\
&= -a_{jk}(A_{ji} - \tilde{A}_{ji}) + a_{jk}(I_p - \tilde{A}_{jk})(A_{ji} - \tilde{A}_{ji}) + a_{jk}(A_{jk} - \tilde{A}_{jk})(I_p - A_{ji}) \\
&\quad + (a_{ik} - a_{jk})(A_{jk} - \tilde{A}_{jk}).
\end{aligned} \tag{A.3}$$

Thus, in (A.1), we multiply $(A_{ji} - \tilde{A}_{ji})^\top$ and take the trace to find

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \|A_{ji} - \tilde{A}_{ji}\|_{\mathbb{F}}^2 \\
&= \text{tr}[\{(A_{ji} - \tilde{A}_{ji})\Xi_i - \Xi_j(A_{ji} - \tilde{A}_{ji})\}(A_{ji} - \tilde{A}_{ji})^\top] + \frac{\kappa}{2N} \sum_{k=1}^N \text{tr}[\mathcal{J}_{3k}(A_{ji} - \tilde{A}_{ji})^\top] \\
&\quad + \frac{\kappa}{2N} \sum_{k=1}^N a_{jk} \text{tr}[\{(A_{ki} - \tilde{A}_{ki}) - (A_{ik} - \tilde{A}_{ik})\}(A_{ji} - \tilde{A}_{ji})^\top] \\
&\quad + \frac{\kappa}{2N} \sum_{k=1}^N a_{ik} \text{tr}[\{(A_{jk} - \tilde{A}_{jk}) - (A_{kj} - \tilde{A}_{kj})\}(A_{ji} - \tilde{A}_{ji})^\top] \\
&=: \mathcal{J}_4 + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{J}_{5k} + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{J}_{6k} + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{J}_{7k}.
\end{aligned} \tag{A.4}$$

Below, we estimate the terms \mathcal{J}_4 and \mathcal{J}_{jk} , $j = 5, 6, 7$, separately.

• (Estimate on \mathcal{J}_4): for a skew-symmetric matrix Y and a matrix B , we observe the following identity:

$$\text{tr}[YBB^\top] = 0.$$

Then, we use the skew-symmetry of Ξ_i to see

$$\begin{aligned}
\mathcal{J}_4 &= \text{tr}[\{(A_{ji} - \tilde{A}_{ji})\Xi_i - \Xi_j(A_{ji} - \tilde{A}_{ji})\}(A_{ji} - \tilde{A}_{ji})^\top] \\
&= \text{tr}[\Xi_i(A_{ji} - \tilde{A}_{ji})^\top(A_{ji} - \tilde{A}_{ji})] - \text{tr}[\Xi_j(A_{ji} - \tilde{A}_{ji})(A_{ji} - \tilde{A}_{ji})^\top] \\
&= 0.
\end{aligned}$$

• (Estimate on \mathcal{J}_{5k}): Due to (A.2) and (A.3), it suffices to estimate the term involving $\mathcal{J}_{3k,1}$. Note that

$$\begin{aligned}
\text{tr}[\mathcal{J}_{3k,1}(A_{ji} - \tilde{A}_{ji})^\top] &\leq -a_{jk}\|A_{ji} - \tilde{A}_{ji}\|^2 + a_{jk}\sqrt{p}[D(\mathcal{S}) + D(\tilde{\mathcal{S}})]d(\mathcal{S}, \tilde{\mathcal{S}})^2 \\
&\quad + (a_{ik} - a_{jk})\text{tr}[(A_{jk} - \tilde{A}_{jk})(A_{ji} - \tilde{A}_{ji})^\top],
\end{aligned}$$

where we used the inequality:

$$\|I_p - A_{ji}\|_F = \|S_j^\top S_j - S_j^\top S_i\|_F \leq \sqrt{p}D(\mathcal{S}).$$

Hence, \mathcal{J}_{5k} can be estimated as follows:

$$\begin{aligned} \mathcal{J}_{5k} &\leq \text{tr}[(\mathcal{J}_{3k,1} + \dots + \mathcal{J}_{3k,4})(A_{ji} - \tilde{A}_{ji})^\top] \\ &= -2(a_{ik} + a_{jk})\|A_{ji} - \tilde{A}_{ji}\|^2 + 2(a_{jk} + a_{ik})\sqrt{p}[D(\mathcal{S}) + D(\tilde{\mathcal{S}})]d(\mathcal{S}, \tilde{\mathcal{S}})^2 \\ &\quad + (a_{ik} - a_{jk})\text{tr}((A_{jk} - \tilde{A}_{jk})(A_{ji} - \tilde{A}_{ji})^\top) + (a_{ik} - a_{jk})\text{tr}((A_{kj} - \tilde{A}_{kj})(A_{ji} - \tilde{A}_{ji})^\top) \\ &\quad + (a_{jk} - a_{ik})\text{tr}((A_{ki} - \tilde{A}_{ki})(A_{ji} - \tilde{A}_{ji})^\top) + (a_{ij} - a_{ik})\text{tr}((A_{ik} - \tilde{A}_{ik})(A_{ji} - \tilde{A}_{ji})^\top) \\ &\leq -2\kappa a_m \|A_{ji} - \tilde{A}_{ji}\|_F^2 + 2\kappa a_M \sqrt{p}[D(\mathcal{S}) + D(\tilde{\mathcal{S}})]d(\mathcal{S}, \tilde{\mathcal{S}})^2 + \mathcal{J}_{5k,1}, \end{aligned}$$

where sum of the last four terms are denoted by $\mathcal{J}_{5k,1}$. Then, we observe

$$\begin{aligned} \mathcal{J}_{5k,1} + \mathcal{J}_{6k} + \mathcal{J}_{7k} &= a_{jk}\text{tr}(((A_{ki} - \tilde{A}_{ki}) - (A_{ik} - \tilde{A}_{ik}))(A_{ji} - \tilde{A}_{ji})^\top) \\ &\quad + a_{ik}\text{tr}(((A_{jk} - \tilde{A}_{jk}) - (A_{kj} - \tilde{A}_{kj}))(A_{ji} - \tilde{A}_{ji})^\top) \\ &\quad + (a_{ik} - a_{jk})\text{tr}((A_{jk} - \tilde{A}_{jk})(A_{ji} - \tilde{A}_{ji})^\top) + (a_{ik} - a_{jk})\text{tr}((A_{kj} - \tilde{A}_{kj})(A_{ji} - \tilde{A}_{ji})^\top) \\ &\quad + (a_{jk} - a_{ik})\text{tr}((A_{ki} - \tilde{A}_{ki})(A_{ji} - \tilde{A}_{ji})^\top) + (a_{ij} - a_{ik})\text{tr}((A_{ik} - \tilde{A}_{ik})(A_{ji} - \tilde{A}_{ji})^\top) \\ &\leq 4(N-1)(a_M + d(\mathcal{A}))d(\mathcal{S}, \tilde{\mathcal{S}})^2. \end{aligned}$$

In (A.4), we collect all estimates to obtain the desired inequality:

$$\frac{d}{dt}d(\mathcal{S}, \tilde{\mathcal{S}}) \leq -2\kappa(\Lambda - a_M \sqrt{p}(D(\mathcal{S}) + D(\tilde{\mathcal{S}})))d(\mathcal{S}, \tilde{\mathcal{S}}), \quad \Lambda = \left(a_m - \frac{N-1}{N}(a_M + d(\mathcal{A})) \right).$$

APPENDIX B. PROOF OF LEMMA 4.3

In this appendix, we present a proof of Lemma 4.3 in which the differential inequality for $\|\mathcal{D}_{ij}\|_F^2$ is derived. First, we observe the following second-order ODE of \mathcal{D}_{ij} :

$$\begin{aligned} m\ddot{\mathcal{D}}_{ij} + \gamma\dot{\mathcal{D}}_{ij} + \kappa\xi_i\mathcal{D}_{ij} + \kappa(\xi_i - \xi_j)S_j &= -m \left(S_i\dot{S}_i^\top\dot{S}_i - S_j\dot{S}_j^\top\dot{S}_j \right) + (S_i\Xi_i - S_j\Xi_j) \\ &\quad - \frac{m}{\gamma} \left(2\dot{S}_i\Xi_i - S_i\Xi_iS_i^\top\dot{S}_i + S_i\dot{S}_i^\top S_i\Xi_i - 2\dot{S}_j\Xi_j + S_j\Xi_jS_j^\top\dot{S}_j - S_j\dot{S}_j^\top S_j\Xi_j \right) \\ &\quad + \frac{\kappa}{N} \sum_{k=1}^N a_{jk} \left(\frac{1}{2}S_jS_j^\top S_k + \frac{1}{2}S_jS_k^\top S_j - S_j \right) - a_{ik} \left(\frac{1}{2}S_iS_i^\top S_k + \frac{1}{2}S_iS_k^\top S_i - S_i \right). \end{aligned} \tag{B.1}$$

It follows from the definition of Stiefel manifold that

$$\mathcal{D}_{ik}^\top \mathcal{D}_{ik} = (S_i - S_k)^\top (S_i - S_k) = 2I_p - S_i^\top S_k - S_k^\top S_i,$$

or equivalently,

$$S_i - \frac{1}{2}S_iS_i^\top S_k - \frac{1}{2}S_iS_k^\top S_i = \frac{1}{2}S_i\mathcal{D}_{ik}^\top \mathcal{D}_{ik}. \tag{B.2}$$

In (B.1), we use the relation (B.2) to see

$$\begin{aligned}
& m\ddot{\mathcal{D}}_{ij} + \gamma\dot{\mathcal{D}}_{ij} + \kappa\xi_i\mathcal{D}_{ij} + \kappa(\xi_i - \xi_j)S_j \\
&= -m(S_i\dot{S}_i^\top\dot{S}_i - S_j\dot{S}_j^\top\dot{S}_j) + (S_i\Xi_i - S_j\Xi_j) + \frac{\kappa}{2N} \sum_{k=1}^N \left(a_{ik}S_i\mathcal{D}_{ik}^\top\mathcal{D}_{ik} - a_{jk}S_j\mathcal{D}_{jk}^\top\mathcal{D}_{jk} \right) \\
&\quad + \frac{m}{\gamma} \left(2\dot{S}_i\Xi_i - S_i\Xi_i S_i^\top\dot{S}_i + S_i\dot{S}_i^\top S_i\Xi_i - 2\dot{S}_j\Xi_j + S_j\Xi_j S_j^\top\dot{S}_j - S_j\dot{S}_j^\top S_j\Xi_j \right) \\
&=: -m\mathcal{I}_6 + \mathcal{I}_7 + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{I}_{8k} + \frac{m}{\gamma}\mathcal{I}_9.
\end{aligned} \tag{B.3}$$

By using (B.3), we derive a second-order ODE of $\|\mathcal{D}_{ij}\|_{\mathbb{F}}^2$:

$$\begin{aligned}
& m\frac{d^2}{dt^2} \|\mathcal{D}_{ij}\|_{\mathbb{F}}^2 + \gamma\frac{d}{dt} \|\mathcal{D}_{ij}\|_{\mathbb{F}}^2 + 2\kappa\xi_i \|\mathcal{D}_{ij}\|_{\mathbb{F}}^2 + 2\kappa(\xi_i - \xi_j)\text{tr}(I_p - S_j^\top S_i) \\
&= \text{tr} \left[m \left(\ddot{\mathcal{D}}_{ij}^\top\mathcal{D}_{ij} + 2\dot{\mathcal{D}}_{ij}^\top\dot{\mathcal{D}}_{ij} + \mathcal{D}_{ij}^\top\ddot{\mathcal{D}}_{ij} \right) + \gamma \left(\dot{\mathcal{D}}_{ij}^\top\mathcal{D}_{ij} + \mathcal{D}_{ij}^\top\dot{\mathcal{D}}_{ij} \right) + 2\kappa\mathcal{D}_{ij}^\top\mathcal{D}_{ij} \right] \\
&= \text{tr} \left[\left(m\ddot{\mathcal{D}}_{ij} + \gamma\dot{\mathcal{D}}_{ij} + \kappa\mathcal{D}_{ij} \right)^\top \mathcal{D}_{ij} \right] + \text{tr} \left[\mathcal{D}_{ij}^\top \left(m\ddot{\mathcal{D}}_{ij} + \gamma\dot{\mathcal{D}}_{ij} + \kappa\mathcal{D}_{ij} \right) \right] + 2m\|\dot{\mathcal{D}}_{ij}\|_{\mathbb{F}}^2 \\
&= 2\text{tr} \left[\mathcal{D}_{ij}^\top \left(-m\mathcal{I}_6 + \mathcal{I}_7 + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{I}_{8k} + \frac{m}{\gamma}\mathcal{I}_9 \right) \right] + 2m\|\dot{\mathcal{D}}_{ij}\|_{\mathbb{F}}^2.
\end{aligned} \tag{B.4}$$

We now assume

$$\xi_i = \xi_j \equiv \xi, \quad i, j = 1, \dots, N.$$

Then, we sum the relation (B.4) with respect to $i, j = 1, \dots, N$ to find temporal evolutions

for $\mathcal{G} = \frac{1}{N^2} \sum_{i,j=1}^N \|\mathcal{D}_{ij}\|_{\mathbb{F}}^2$:

$$\begin{aligned}
& m\ddot{\mathcal{G}} + \gamma\dot{\mathcal{G}} + 2\kappa\xi\mathcal{G} \\
&= \frac{2}{N^2} \sum_{i,j=1}^N \text{tr} \left[\mathcal{D}_{ij}^\top \left(-m\mathcal{I}_6 + \mathcal{I}_7 + \frac{\kappa}{2N} \sum_{k=1}^N \mathcal{I}_{8k} + \frac{m}{\gamma}\mathcal{I}_9 \right) \right] + \frac{2m}{N^2} \sum_{i,j=1}^N \|\dot{\mathcal{D}}_{ij}\|_{\mathbb{F}}^2.
\end{aligned} \tag{B.5}$$

Below, we present estimates for \mathcal{I}_k , $k = 6, \dots, 9$, respectively.

- (Estimate of \mathcal{I}_6): we use the fact

$$\|PAQ^\top\|_{\mathbb{F}} = \|A\|_{\mathbb{F}} \quad \text{for } P, Q \in \text{St}(p, n) \quad \text{and } A \in \mathcal{M}_{p,p}(\mathbb{R}), \tag{B.6}$$

which can be proved as follows:

$$\|PAQ^\top\|_{\mathbb{F}}^2 = \text{tr}(QA^\top P^\top PAQ^\top) = \text{tr}(QA^\top AQ^\top) = \text{tr}(A^\top AQ^\top Q) = \text{tr}(A^\top A) = \|A\|_{\mathbb{F}}^2.$$

Then, we use (B.6) to see

$$\begin{aligned}
\operatorname{tr} \left(\mathcal{D}_{ij}^\top \mathcal{I}_6 \right) &= \operatorname{tr} \left((S_i^\top - S_j^\top) (S_i \dot{S}_i^\top \dot{S}_i - S_j \dot{S}_j^\top \dot{S}_j) \right) \\
&= \operatorname{tr} \left(\dot{S}_i^\top \dot{S}_i - S_i^\top S_j \dot{S}_j^\top \dot{S}_j - S_j^\top S_i \dot{S}_i^\top \dot{S}_i + \dot{S}_j^\top \dot{S}_j \right) \\
&= \|\dot{S}_i\|_{\mathbb{F}}^2 + \|\dot{S}_j\|_{\mathbb{F}}^2 - \operatorname{tr} \left(S_j \dot{S}_j^\top \dot{S}_j S_i^\top + S_i \dot{S}_i^\top \dot{S}_i S_j^\top \right) \\
&\leq 2(\|\dot{S}_i\|_{\mathbb{F}}^2 + \|\dot{S}_j\|_{\mathbb{F}}^2) \leq 4D(\dot{S})^2.
\end{aligned}$$

- (Estimate of \mathcal{I}_7): we use the maximality of $\|\Xi\|_\infty$ to find

$$\begin{aligned}
\operatorname{tr} \left(\mathcal{D}_{ij}^\top \mathcal{I}_7 \right) &= \operatorname{tr} \left[\left(S_i^\top - S_j^\top \right) (S_i \Xi_i - S_j \Xi_j) \right] = \operatorname{tr} \left(\Xi_i - S_i^\top S_j \Xi_j - S_j^\top S_i \Xi_i + \Xi_j \right) \\
&\leq 2 \|\Xi_i\|_{\mathbb{F}} + 2 \|\Xi_j\|_{\mathbb{F}} \leq 4 \|\Xi\|_\infty.
\end{aligned}$$

- (Estimate of \mathcal{I}_8): by straightforward calculation,

$$\begin{aligned}
\frac{1}{N^3} \sum_{i,j,k} \operatorname{tr} (\mathcal{D}_{ij}^\top \mathcal{I}_{8k}) &= \frac{1}{N^3} \sum_{i,j,k} a_{ik} \operatorname{tr} ((S_i^\top - S_j^\top) S_i \mathcal{D}_{ik}^\top \mathcal{D}_{ik}) - a_{jk} \operatorname{tr} ((S_i^\top - S_j^\top) S_j \mathcal{D}_{jk}^\top \mathcal{D}_{jk}) \\
&= \frac{1}{2N^3} \sum_{i,j,k} \left(a_{ik} \operatorname{tr} (\mathcal{D}_{ik}^\top \mathcal{D}_{ik}) - a_{jk} \operatorname{tr} (\mathcal{D}_{jk}^\top \mathcal{D}_{jk}) \right. \\
&\quad \left. + a_{jk} S_i^\top S_j \mathcal{D}_{jk}^\top \mathcal{D}_{jk} - a_{ik} S_j^\top S_i \mathcal{D}_{ik}^\top \mathcal{D}_{ik} \right) = 0,
\end{aligned}$$

where we used the interchange of the index $i \leftrightarrow j$.

- (Estimate of \mathcal{I}_9): we use $\|S\|_{\mathbb{F}} = \sqrt{p}$ for $S \in \operatorname{St}(p, n)$ to see

$$\begin{aligned}
\operatorname{tr} \left(\mathcal{D}_{ij}^\top \mathcal{I}_9 \right) &= \operatorname{tr} \left[\left(S_i^\top - S_j^\top \right) \left(2\dot{S}_i \Xi_i - S_i \Xi_i S_i^\top \dot{S}_i + S_i \dot{S}_i^\top S_i \Xi_i \right. \right. \\
&\quad \left. \left. - 2\dot{S}_j \Xi_j + S_j \Xi_j S_j^\top \dot{S}_j - S_j \dot{S}_j^\top S_j \Xi_j \right) \right] \\
&= \operatorname{tr} (S_i^\top \dot{S}_i \Xi_i - \Xi_i S_i^\top \dot{S}_i - 2S_i^\top \dot{S}_j \Xi_j + S_i^\top S_j \Xi_j S_j^\top \dot{S}_j - S_i^\top S_j \dot{S}_j^\top S_j \Xi_j \\
&\quad + S_j^\top \dot{S}_j \Xi_j - \Xi_j S_j^\top \dot{S}_j - 2S_j^\top \dot{S}_i \Xi_i + S_j^\top S_i \Xi_i S_i^\top \dot{S}_i - S_j^\top S_i \dot{S}_i^\top S_i \Xi_i) \\
&\leq 8\sqrt{p}D(\dot{S})\|\Xi\|_\infty.
\end{aligned}$$

In (B.5), we combine all estimates for $\operatorname{tr}(\mathcal{D}_{ij}^\top \mathcal{I}_k)$, $k = 6, \dots, 9$ and the fact that $\|\dot{\mathcal{D}}_{ij}\|_{\mathbb{F}}^2 \leq 4D(\dot{S})^2$ to obtain the desired differential inequality for \mathcal{G} :

$$m\ddot{\mathcal{G}} + \gamma\dot{\mathcal{G}} + 2\kappa\xi\mathcal{G} \leq 16mD(\dot{S})^2 + 8\|\Xi\|_\infty + \frac{16m\sqrt{p}\|\Xi\|_\infty}{\gamma}D(\dot{S}).$$

REFERENCES

- [1] Barbălat, I.: *Systèmes d'équations différentielles d'oscillations non Linéaires*. Rev. Math. Pures Appl. **4** (1959), 267-270.
- [2] Bellomo, N. and Dogbé, C.: *On the modeling of traffic and crowds: A survey of models, speculations, and perspectives*. SIAM Rev. **53** (2011), 409-463.
- [3] Bellomo, N. and Gibelli, L.: *Behavioral crowds: Modeling and Monte Carlo simulations toward validation*. Comput. Fluids **141** (2016), 13-21.
- [4] Bloch, A. M., Crouch, P. E. and Sanyal, A. K.: *A variational problem on Stiefel manifolds*. Nonlinearity **19** (2006), 2247-2276.
- [5] Buck, J. and Buck, E.: *Biology of synchronous flashing of fireflies*. Nature **211** (1966), 562-564.
- [6] Carrillo, J. A., Choi, Y.-P., Totzeck, C. and Tse, O.: *An analytical framework for consensus-based global optimization method*. Math. Models Methods Appl. Sci. **28** (2018), 1037-1066.
- [7] Chen, S., Ma, S., So, A. M.-C. and Zhang, T.: *Proximal gradient method for nonsmooth optimization over the Stiefel manifold*. SIAM J. Optim. **30** (2020), 210-239.
- [8] Choi, Y.-P., Ha, S.-Y., Jung, S. and Kim, Y.: *Asymptotic formation and orbital stability of phase-locked states for the Kuramoto model*. Phys. D **241**, 735-754 (2012).
- [9] Choi, Y.-P., Ha, S.-Y. and Yun, S.-B.: *Complete synchronization of Kuramoto oscillators with finite inertia*. Phys. D **240** (2011), 32-44.
- [10] Choi, Y.-P., Kalsie, D., Peszek, J. and Peters, A.: *A collisionless singular Cucker-Smale model with decentralized formation control*. SIAM J. Appl. Dyn. Syst. **18** (2019), 1954-1981.
- [11] Chopra, N. and Spong, M. W.: *Passivity-based control of multi-agent systems*. In: Kawamura, S., Svinin, M. (eds) Advances in robot control. Springer, Berlin, Heidelberg.
- [12] Chopra, N. and Spong, M. W.: *On Exponential Synchronization of Kuramoto Oscillators* IEEE Trans. Automat. Control **54** (2009), 353-357.
- [13] Cucker, F. and Smale, S.: *Emergent behavior in flocks*. IEEE Trans. Automat. Control **52** (2007), 852-862.
- [14] Cumin, D. and Unsworth, C. P.: *Generalizing the Kuramoto model for the study of neuronal synchronization in the brain*. Phys. D **226** (2007), 181-196.
- [15] Danino, T., Mondragon-Palomino, O., Tsimring, L. and Hasty, J.: *A synchronized quorum of genetic clocks*. Nature **463** (2010), 326-330.
- [16] Degond, P., Frouvelle, A. and Merino-Aceituno, S.: *A new flocking model through body attitude coordination*. Math. Models Methods Appl. Sci. **27** (2017), 1005-1049.
- [17] Degond, P., Frouvelle, A., Merino-Aceituno, S. and Trescases, A.: *Quaternions in collective dynamics*. Multiscale Model. Simul. **16** (2018), 28-77.
- [18] Edelman, A., Arias, T. A. and Smith, S.: *The geometry of algorithms with orthogonality constraints*. SIAM J. Matrix Anal. Appl. **20** (1998), 303-353.
- [19] Fornasier, M., Huang, H., Pareschi, L. and Sünnen, P.: *Consensus-based optimization on hypersurfaces: well-posedness and mean-field limit*. To appear in Math. Models Method Appl. Sci.
- [20] Fornasier, M., Huang, H., Pareschi, L. and Sünnen, P.: *Consensus-based optimization on the sphere II: convergence to global minimizers and machine learning*. archived as arXiv:2001.11988v3.
- [21] Golub, G.H. and Van Loan, C.F.: *Matrix computations*, 4th edn. The Johns Hopkins University Press, Baltimore (2013)
- [22] Grassi, S. and Pareschi, L.: *From particle swarm optimization to consensus based optimization: stochastic modeling and mean-field limit*. archived as arXiv:2012.05613v1.
- [23] Gregor, T., Fujimoto, K., Masaki, N. and Sawai, S.: *The onset of collective behavior in social amoebae*. Science **328** (2010), 1021-1025.
- [24] Ha, S.-Y. and Kim, D.: *A second-order particle swarm model on a sphere and emergent dynamics*. SIAM J. Appl. Dyn. Syst. **18** (2019), 80-116.
- [25] Ha, S.-Y. and Kim, D.: *Emergent behavior of a second-order Lohe matrix model on the unitary group*. J. Stat. Phys. **175** (2019), 904-931.
- [26] Ha, S.-Y., Ko, D. and Ryoo, S. W.: *Emergent dynamics of a generalized Lohe model on some class of Lie groups*. J. Stat. Phys. **168** (2017), 171-207.
- [27] Ha, S.-Y., Ko, D. and Ryoo, S. W.: *On the relaxation dynamics of Lohe oscillators on some Riemannian manifolds*. J. Stat. Phys. **172** (2018), 1427-1478.

- [28] Ha, S.-Y. and Ryoo S.-W.: *On the emergence and orbital stability of phase-locked states for the Lohe model*. J. Stat. Phys. **163** (2016), 411-439.
- [29] Hatanaka, T., Chopra, N., Ishizaki, T. and Li, N.: *Passivity-based distributed optimization with communication delays using PI consensus algorithm*. IEEE Trans. Automat. Control **63** (2018), 4421-4428.
- [30] Hung, S. M. and Givigi, S. N.: *A q-learning approach to flocking with UAVs in a stochastic environment*. IEEE Trans. Cybern. **47** (2017), 186-197.
- [31] Kim, J., Kang, M., Kim, D., Ha, S.-Y. and Yang, I.: *A stochastic consensus method for nonconvex optimization on the Stiefel manifold*. To appear in Proceedings of the 59th IEEE Conference on Decision and Control (2020).
- [32] Klar, A. and Wegener, R.: *Enskog-like kinetic models for vehicular traffic*. J. Stat. Phys. **87** (1997), 91-114.
- [33] Kuramoto, Y.: *Chemical Oscillations, Waves and Turbulence*. Springer-Verlag, Berlin, 1984.
- [34] Kuramoto, Y.: *Self-entrainment of a population of coupled non-linear oscillators*. In *International Symposium on Mathematical Problems in Mathematical Physics*. Lecture Notes in Theoretical Physics **30** (1975), 420-422.
- [35] Li, Q. and Qi, H.: *A sequential semismooth Newton method for the nearest low-rank correlation matrix problem*. SIAM J. Optim. **21** (2011), 1641-1666.
- [36] Li, W. and Spong, M. W.: *Unified cooperative control of multiple agents on a sphere for different spherical patterns*. IEEE Trans. Automat. Control **59** (2014), 1283-1289.
- [37] Liu, C. and Boumal, N.: *Simple algorithms for optimization on Riemannian manifolds with constraints*. Appl. Math. Optim **82** (2020), 949-981.
- [38] Liu, X., Wen, Z. and Zhang, Y.: *Limited memory block Krylov subspace optimization for computing dominant singular value decompositions*. SIAM J. Sci. Comput. **35** (2013), 1641-1668.
- [39] Lohe, M. A.: *Non-Abelian Kuramoto model and synchronization*. J. Phys. A **42** (2009), 395101.
- [40] Lui, Y. M.: *Advances in matrix manifolds for computer vision*. Image Vision Comput. **30** (2012), 380-388.
- [41] Markdahl, J., Thunberg, J. and Gonçalves, J.: *Towards almost global synchronization on the Stiefel manifold*. Proc. of the 57th IEEE Conference on Decision and Control (2018).
- [42] Markdahl, J., Thunberg, J. and Gonçalves, J.: *High-dimensional Kuramoto models on Stiefel manifolds synchronize complex networks almost globally*. Automatica **113** (2020), 108736.
- [43] Markdahl, J.: *Synchronization on Riemannian manifolds: Multiply connected implies multistable*. IEEE Trans. Automat. Control (2020), <https://10.1109/TAC.2020.3030849>.
- [44] Nelson, P.: *A kinetic model of vehicular traffic and its associated bimodal equilibrium solution*. Transp. Theory Statist. Phys. **24** (1995), 383-409.
- [45] Olfati-Saber, R.: *Swarms on sphere: A programmable swarm with synchronous behaviors like oscillator networks*. Proc. of the 45th IEEE conference on Decision and Control (2006), 5060-5066.
- [46] Perea, L., Elosegui, P. and Gomez, G.: *Extension of the Cucker-Smale control law to space flight formations*. J. Guid. Control **32** (2009), 527-537.
- [47] Pinnau, R., Totzeck, C., Tse, O. and Martin, S.: *A consensus-based model for global optimization and its mean-field limit*. Math. Models Methods Appl. Sci. **27** (2017), 183-204.
- [48] Pölitz, C., Duivesteyn, W. and Morik, K.: *Interpretable domain adaptation via optimization over the Stiefel manifold*. Mach. Learn. **104** (2016), 315-336.
- [49] Ritchie, L. M. Lohe, M.A. and Williams, A.G.: *Synchronization of relativistic particles in the hyperbolic Kuramoto model*. Chaos **28** (2018), 053116.
- [50] Rubenstein, M., Cornejo, A. and Nagapal, R.: *Programmable self-assembly in a thousand-robot swarm*. Science **345** (2014), 795-799.
- [51] Sabattini, L., Secchi, C., Chopra, N. and Gasparri, A.: *Distributed control of multirobot systems with global connectivity maintenance*. IEEE Trans. Robot. **29** (2013), 1326-1332.
- [52] Sarlette, A. and Sepulchre, R.: *Consensus optimization on manifolds*. SIAM J. Control Optim. **48** (2009), 56-76.
- [53] Sarlette, A., Sepulchre, R., Leonard, N. E.: *Autonomous rigid body attitude synchronization*. Automatica **45** (2009), 572-577.
- [54] Sato, H. and Iwai, T.: *A Riemannian optimization approach to the matrix singular value decomposition*. SIAM J. Optim. **23** (2013), 188-212.

- [55] Stiefel, E.: *Richtungsfelder und fernparallelismus in n-dimensionalem mannig faltigkeiten*. Commentarii Math. Helvetici **8** (1935), 305-353.
- [56] Tron, R., Afsari, B. and Vidal, R.: *Intrinsic consensus on $SO(3)$ with almost-global convergence*. 2012 IEEE Conference on Decision and Control (2012), 2052-2058.
- [57] Turaga, P., Veeraraghavan, A., Srivastava, A. and Chellappa, R.: *Statistical Computations on Grassmann and Stiefel Manifolds for Image and Video-Based Recognition*. IEEE T. Pattern Anal. **33** (2011), 2273-2286.
- [58] Vicsek, T., Czirók, A., Ben-Jacob, E., Cohen, I. and Shochet, O.: *Novel type of phase transition in a system of self-driven particles*. Phys. Rev. Lett. **75** (1995), 1226-1229.
- [59] Wen, Z., Yang, C., Liu, X. and Zhang, Y.: *Trace-penalty minimization for large-scale eigenspace computation*. J. Sci. Comput. **66** (2016), 1175-1203.
- [60] Wen, Z. and Yin, W.: *A feasible method for optimization with orthogonality constraints*. Math. Program. **142** (2013), 397-434.
- [61] Winfree, A. T.: *Biological rhythms and the behavior of populations of coupled oscillators*. J. Theor. Biol. **16** (1967), 15-42.
- [62] Zhang, J. and Zhu, J.: *Exponential synchronization of the high-dimensional Kuramoto model with identical oscillators under digraphs*. Automatica **102** (2019), 122-128.
- [63] Zhu, X.: *A Riemannian conjugate gradient method for optimization on the Stiefel manifold*. Comput. Optim. Appl. **67** (2017), 73-110.

(S.-Y. Ha)

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS
 SEOUL NATIONAL UNIVERSITY, SEOUL 08826 AND
 KOREA INSTITUTE FOR ADVANCED STUDY, HOEGIRO 85, SEOUL, 02455, REPUBLIC OF KOREA
Email address: syha@snu.ac.kr

(M. Kang)

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS
 SEOUL NATIONAL UNIVERSITY, SEOUL 08826, REPUBLIC OF KOREA
Email address: bear0117@snu.ac.kr

(D. Kim)

DEPARTMENT OF MATHEMATICS,
 SUNGSHIN WOMEN'S UNIVERSITY, SEOUL 02844, REPUBLIC OF KOREA
Email address: dohyunkim@sungshin.ac.kr