NUMERICAL RADIUS IN HILBERT C*-MODULES

ALI ZAMANI

ABSTRACT. Utilizing the linking algebra of a Hilbert C^* -module $(\mathcal{V}, \|\cdot\|)$, we introduce $\Omega(x)$ as a definition of numerical radius for an element $x \in \mathcal{V}$ and then show that $\Omega(\cdot)$ is a norm on \mathcal{V} such that $\frac{1}{2}||x|| \leq \Omega(x) \leq ||x||$. In addition, we obtain an equivalent condition for $\Omega(x) = \frac{1}{2}||x||$. Moreover, we present a refinement of the triangle inequality for the norm $\Omega(\cdot)$. Some other related results are also discussed.

1. INTRODUCTION AND PRELIMINARIES

The notion of Hilbert C^* -module is a natural generalization of that of Hilbert space arising under replacement of the field of scalars \mathbb{C} by a C^* -algebra. This concept plays a significant role in the theory of operator algebras, quantum groups, noncommutative geometry and K-theory; see [10, 11].

Let us give that some necessary background and set up our notation. An element a in a C^* -algebra \mathscr{A} is called positive (we write $0 \leq a$) if $a = b^*b$ for some $b \in \mathscr{A}$. For an element a of \mathscr{A} , we denote by

Re
$$a = \frac{1}{2}(a + a^*)$$
, Im $a = \frac{1}{2i}(a - a^*)$

the real and the imaginary part of a. By \mathscr{A}' we denote the dual space of \mathscr{A} . A positive linear functional of \mathscr{A} is a map $\varphi \in \mathscr{A}'$ such that $0 \leq \varphi(a)$ whenever $0 \leq a$. The set of all states of \mathscr{A} , that is, the set of all positive linear functionals of \mathscr{A} of norm 1, is denoted by $\mathscr{S}(\mathscr{A})$. An inner product module over \mathscr{A} is a (left) \mathscr{A} -module \mathscr{V} equipped with an \mathscr{A} -valued inner product $\langle \cdot, \cdot \rangle$, which is \mathbb{C} -linear and \mathscr{A} -linear in the first variable and has the properties $\langle x, y \rangle^* = \langle y, x \rangle$ as well as $0 \leq \langle x, x \rangle$ with equality if and only if x = 0. The \mathscr{A} -module \mathscr{V} is called a Hilbert \mathscr{A} -module if it is complete with respect to the norm $||x|| = ||\langle x, x \rangle||^{\frac{1}{2}}$. In a Hilbert \mathscr{A} -module \mathscr{V} we have the following version of the Cauchy-Schwarz inequality:

$$\langle y, x \rangle \langle x, y \rangle \le \|x\|^2 \langle y, y \rangle, \qquad (x, y \in \mathscr{V}).$$
 (1.1)

Every C^* -algebra \mathscr{A} can be regarded as a Hilbert C^* -module over itself where the inner product is defined by $\langle a, b \rangle = a^*b$. Let \mathscr{V} and \mathscr{W} be two Hilbert \mathscr{A} modules. A mapping $T : \mathscr{V} \longrightarrow \mathscr{W}$ is called adjointable if there exists a mapping $S : \mathscr{W} \longrightarrow \mathscr{V}$ such that $\langle Tx, y \rangle = \langle x, Sy \rangle$ for all $x \in \mathscr{V}, y \in \mathscr{W}$. The unique mapping S is denoted by T^* and is called the adjoint operator of T. The space

²⁰¹⁰ Mathematics Subject Classification. Primary 46L05; Secondary 47A30, 47A12, 46B20.

Key words and phrases. C^* -algebra, Hilbert C^* -module, linking algebra, numerical range, numerical radius, inequality.

A. ZAMANI

 $\mathbb{B}(\mathscr{V},\mathscr{W})$ of all adjointable maps between Hilbert \mathscr{A} -modules \mathscr{V} and \mathscr{W} is a Banach space, while $\mathbb{B}(\mathscr{V}) := \mathbb{B}(\mathscr{V},\mathscr{V})$ is a C^* -algebra. By $\mathbb{K}(\mathscr{V},\mathscr{W})$ we denote the closed linear subspace of $\mathbb{B}(\mathscr{V},\mathscr{W})$ spanned by $\{\theta_{x,y} : x \in \mathscr{W}, y \in \mathscr{V}\}$, where $\theta_{x,y}$ is defined by $\theta_{x,y}(z) = x \langle y, z \rangle$. Elements of $\mathbb{K}(\mathscr{V},\mathscr{W})$ are often referred to as "compact" operators. We write $\mathbb{K}(\mathscr{V})$ for $\mathbb{K}(\mathscr{V},\mathscr{V})$. Given a Hilbert \mathscr{A} -module \mathscr{V} , the linking algebra $\mathbb{L}(\mathscr{V})$ is defined as the matrix algebra of the form

$$\mathbb{L}(\mathscr{V}) = \begin{bmatrix} \mathbb{K}(\mathscr{A}) & \mathbb{K}(\mathscr{V}, \mathscr{A}) \\ \mathbb{K}(\mathscr{A}, \mathscr{V}) & \mathbb{K}(\mathscr{V}) \end{bmatrix}.$$

Then $\mathbb{L}(\mathscr{V})$ has a canonical embedding as a closed subalgebra of the adjointable operators on the Hilbert \mathscr{A} -module $\mathscr{A} \oplus \mathscr{V}$ via

$$\begin{bmatrix} X & Y \\ Z & W \end{bmatrix} \begin{bmatrix} a \\ x \end{bmatrix} = \begin{bmatrix} Xa + Yx \\ Za + Wx \end{bmatrix}$$

which makes $\mathbb{L}(\mathscr{V})$ a C^* -algebra (cf. [15], Lemma 2.32 and Corollary 3.21). Each $x \in \mathscr{V}$ induces the maps $r_x \in \mathbb{B}(\mathscr{A}, \mathscr{V})$ and $l_x \in \mathbb{B}(\mathscr{V}, \mathscr{A})$ given by $r_x(a) = xa$ and $l_x(y) = \langle x, y \rangle$, respectively, such that $r_x^* = l_x$. The map $x \mapsto r_x$ is an isometric linear isomorphism of \mathscr{V} to $\mathbb{K}(\mathscr{A}, \mathscr{V})$ and $x \mapsto l_x$ is an isometric conjugate linear isomorphism of \mathscr{V} to $\mathbb{K}(\mathscr{A}, \mathscr{N})$. Further, every $a \in \mathscr{A}$ induces the map $T_a \in \mathbb{K}(\mathscr{A})$ given by $T_a(b) = ab$. The map $a \mapsto T_a$ defines an isomorphism between C^* -algebras \mathscr{A} and $\mathbb{K}(\mathscr{A})$. Therefore, we may write

$$\mathbb{L}(\mathscr{V}) = \left\{ \begin{bmatrix} T_a & l_y \\ r_x & T \end{bmatrix} : a \in \mathscr{A}, \, x, y \in \mathscr{V}, \, T \in \mathbb{K}(\mathscr{V}) \right\},\,$$

and identify the C^* -subalgebras of compact operators with the corresponding corners in the linking algebra: $\mathbb{K}(\mathscr{A}) = \mathbb{K}(\mathscr{A} \oplus 0) \subseteq \mathbb{K}(\mathscr{A} \oplus \mathscr{V}) = \mathbb{L}(\mathscr{V})$ and $\mathbb{K}(\mathscr{V}) = \mathbb{K}(0 \oplus \mathscr{V}) \subseteq \mathbb{K}(\mathscr{A} \oplus \mathscr{V}) = \mathbb{L}(\mathscr{V})$. We refer the reader to [10, 11] for more information on Hilbert C^* -modules and linking algebras.

Now, let $\mathbb{B}(\mathscr{H})$ denote the C^* -algebra of all bounded linear operators on a complex Hilbert space \mathscr{H} with inner product $[\cdot, \cdot]$. The numerical range of an element $A \in \mathbb{B}(\mathscr{H})$ is defined

$$W(A) := \{ [A\xi, \xi] : \xi \in \mathscr{H}, \|\xi\| = 1 \}.$$

It is known that W(A) is a nonempty bounded convex subset of \mathbb{C} (not necessarily closed). This concept is useful in studying linear operators and have attracted the attention of many authors in the last few decades (e.g., see [8], and references therein). The numerical radius of A is given by

$$w(A) = \sup \{ |[A\xi, \xi]| : \xi \in \mathscr{H}, ||\xi|| = 1 \}.$$

It is known that $w(\cdot)$ is a norm on $\mathbb{B}(\mathscr{H})$ and satisfies

$$\frac{1}{2} \|A\| \le w(A) \le \|A\|$$

for each $A \in \mathbb{B}(\mathcal{H})$. Some generalizations of the numerical radius $A \in \mathbb{B}(\mathcal{H})$ can be found in [2, 22].

In the next section, we first utilize the linking algebra $\mathbb{L}(\mathscr{V})$ of a Hilbert \mathscr{A} -module \mathscr{V} to introduce $\Phi(x)$ as a definition of numerical range for an arbitrary

element $x \in \mathscr{V}$. We then use this set to define numerical radius of x and denote it by $\Omega(x)$. In particular, we show that $\Omega(\cdot)$ is a norm on \mathscr{V} , which is equivalent to the norm $\|\cdot\|$ and the following inequalities hold for every $x \in \mathscr{V}$:

$$\frac{1}{2} \|x\| \le \Omega(x) \le \|x\|.$$
(1.2)

We also establish an inequality that refines the first inequality in (1.2). In addition, we prove that $\Omega(x) = \frac{1}{2} ||x||$ if and only if $||x|| = \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|$ for all complex unit λ . Furthermore, for $x \in \mathscr{V}$ and $a \in \mathscr{A}$ we prove that

$$\Omega(xa \pm xa^*) \le 2 \|a \pm a^*\| \Omega(x).$$

We finally present a refinement of the triangle inequality for the norm $\Omega(\cdot)$.

2. Main results

We start our work with the following definition.

Definition 2.1. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . The numerical range of $x \in \mathscr{V}$ is defined as the set

$$\Phi(x) := \left\{ \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) : \varphi \in \mathcal{S} \big(\mathbb{L}(\mathscr{V}) \big) \right\}.$$

Next, we present some properties of the numerical range in Hilbert C^* -modules.

Theorem 2.2. Let x and y be elements of a Hilbert \mathscr{A} -module \mathscr{V} and let $\alpha \in \mathbb{C}$. Then

- (i) $\Phi(\alpha x) = \alpha \Phi(x)$ (homogeneous).
- (ii) $\Phi(x+y) \subseteq \Phi(x) + \Phi(y)$ (subadditive).
- (iii) $\Phi(x)$ is a nonempty compact convex subset of \mathbb{C} .

Proof. Let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . For every $a \in \mathscr{A}$, we have

$$r_{\alpha x}(a) = (\alpha x)a = \alpha(xa) = (\alpha r_x)(a)$$

and

$$r_{x+y}(a) = (x+y)a = xa + ya = (r_x + r_y)(a).$$

Hence $r_{\alpha x} = \alpha r_x$ and $r_{x+y} = r_x + r_y$. Thus (i) and (ii) follow easily from the definition.

We now prove (iii). Since the existence of states on $\mathbb{L}(\mathscr{V})$ is guaranteed by the Hahn–Banach theorem, we have $\Phi(x) \neq \emptyset$. The convexity of $\Phi(x)$ is an easy consequence of the fact that a convex combination of two states is also a state. As for the compactness, note that the set $\mathcal{S}(\mathbb{L}(\mathscr{V}))$ is a weak*-closed subset of the unit ball $\{\varphi \in \mathbb{L}'(\mathscr{V}) : \|\varphi\| \leq 1\}$ of $\mathbb{L}'(\mathscr{V})$. Since, by the Banach–Alaoglu theorem, the latter is weak*-compact, the same is true for $\mathcal{S}(\mathbb{L}(\mathscr{V}))$. Hence $\Phi(x)$, the image of the weak*-continuous mapping $\varphi \mapsto \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right)$ for $\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$, is compact in \mathbb{C} . Remark 2.3. It is known that the set of all states of a unital C^* -algebra $\mathscr{A} \subseteq \mathbb{B}(\mathscr{H})$ is a weak*-closed convex hull of the set of all vector states of \mathscr{A} , i.e., the states of \mathscr{A} of the form $A \to [A\xi, \xi]$ for some unit vector ξ in \mathscr{H} . Also, for the Hilbert module $\mathscr{V} = \mathbb{B}(\mathscr{H})$ over the C^* -algebra $\mathbb{B}(\mathscr{H})$ is well known to be valid $\mathbb{K}(\mathbb{B}(\mathscr{H})) = \mathbb{K}(\mathscr{V}, \mathbb{B}(\mathscr{H})) = \mathbb{K}(\mathbb{B}(\mathscr{H}), \mathscr{V}) = \mathbb{K}(\mathscr{V}) = \mathbb{B}(\mathscr{H})$ (see [5, Remark 1.13]), so all corners in the linking algebra $\mathbb{L}(\mathscr{V})$ are equal to $\mathbb{B}(\mathscr{H})$. Hence, for $A \in \mathbb{B}(\mathscr{H})$, we have $\Phi(A) = \overline{W(A)}$.

Now, we are in a position to introduce numerical radius for elements of a Hilbert C^* -module. Some other related topics can be found in [3, 6, 12, 16, 17, 19].

Definition 2.4. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . The numerical radius of an element $x \in \mathscr{V}$ is defined as

$$\Omega(x) := \sup \left\{ \left| \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right| : \varphi \in \mathcal{S} \big(\mathbb{L}(\mathscr{V}) \big) \right\}.$$

In the following theorem, we prove that $\Omega(\cdot)$ is a norm on Hilbert C^* -module \mathscr{V} , which is equivalent to the norm $\|\cdot\|$.

Theorem 2.5. Let \mathscr{V} be a Hilbert \mathscr{A} -module. Then $\Omega(\cdot)$ is a norm on \mathscr{V} and the following inequalities hold for every $x \in \mathscr{V}$:

$$\frac{1}{2}\|x\| \le \Omega(x) \le \|x\|.$$

Proof. Let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . Let $x \in \mathscr{V}$. Clearly, $\Omega(x) \geq 0$. Let us now suppose $\Omega(x) = 0$. Then, by Definition 2.4, $\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} = 0$. Since $\left\| \begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right\| = \|x\|$, we get $\|x\| = 0$ and therefore, x = 0. Further, by Theorem 2.2 (i)-(ii), for $y, z \in \mathscr{V}$ and $\alpha \in \mathbb{C}$ we have $\Omega(\alpha y) = |\alpha|\Omega(y)$ and $\Omega(y + z) \leq \Omega(y) + \Omega(z)$. Thus $\Omega(\cdot)$ is a norm on \mathscr{V} .

On the other hands, for every $\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$, we have

$$\left|\varphi\left(\begin{bmatrix}0 & 0\\r_x & 0\end{bmatrix}\right)\right| \le \left\|\begin{bmatrix}0 & 0\\r_x & 0\end{bmatrix}\right\| = \|x\|.$$

So, by taking the supremum over $\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$ in the above inequality, we deduce that

$$\Omega(x) \le \|x\|. \tag{2.1}$$

Now let $\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} = \operatorname{Re}\left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) + i \operatorname{Im}\left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right)$ be the Cartesian decomposition of $\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix}$. By [13, Theorem 3.3.6], there exist $\varphi_1, \varphi_2 \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$ such that

$$\left|\varphi_1\left(\operatorname{Re}\left(\begin{bmatrix}0&0\\r_x&0\end{bmatrix}\right)\right)\right| = \left\|\operatorname{Re}\left(\begin{bmatrix}0&0\\r_x&0\end{bmatrix}\right)\right\|$$
(2.2)

and

$$\left|\varphi_{2}\left(\operatorname{Im}\left(\begin{bmatrix}0 & 0\\r_{x} & 0\end{bmatrix}\right)\right)\right| = \left\|\operatorname{Im}\left(\begin{bmatrix}0 & 0\\r_{x} & 0\end{bmatrix}\right)\right\|.$$
(2.3)

Therefore, by (2.2) and (2.3), we have

$$\begin{split} \frac{1}{2} \|x\| &= \frac{1}{2} \left\| \begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right\| \\ &\leq \frac{1}{2} \left\| \operatorname{Re} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right\| + \frac{1}{2} \left\| \operatorname{Im} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right\| \\ &= \frac{1}{2} \left| \varphi_1 \left(\operatorname{Re} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right| + \frac{1}{2} \left| \varphi_2 \left(\operatorname{Im} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right| \\ &= \frac{1}{4} \left| \varphi_1 \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) + \overline{\varphi_1} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right| + \frac{1}{4} \left| \varphi_2 \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) - \overline{\varphi_2} \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right| \\ &\leq \frac{1}{2} \left| \varphi_1 \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right| + \frac{1}{2} \left| \varphi_2 \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right| \leq \frac{1}{2} \Omega(x) + \frac{1}{2} \Omega(x) = \Omega(x), \end{split}$$

whence

$$\frac{1}{2}\|x\| \le \Omega(x). \tag{2.4}$$

From (2.1) and (2.4), we deduce the desired result.

For $A \in \mathbb{B}(\mathscr{H})$, we note that (see [20]) $w(A) = \sup_{\lambda \in \mathbb{T}} ||\operatorname{Re}(\lambda A)||$. Here, as usual, \mathbb{T} is the unit circle of the complex plane \mathbb{C} . This motivates the following result.

Theorem 2.6. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . Then

$$\Omega(x) = \frac{1}{2} \sup_{\lambda \in \mathbb{T}} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|,$$

for every $x \in \mathscr{V}$.

Proof. Let $x \in \mathscr{V}$. First, we show that

$$\sup_{\lambda \in \mathbb{T}} \left| \operatorname{Re} \left(\lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right| = \left| \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right|$$
(2.5)

for every $\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$.

Let $\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))$. We may assume that $\varphi\left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix}\right) \neq 0$, otherwise (2.5) trivially holds. Put

$$\lambda_0 = \frac{\overline{\varphi}\left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right)}{\left| \varphi\left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right|}.$$

Then we have

$$\begin{vmatrix} \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \end{vmatrix} = \begin{vmatrix} \operatorname{Re} \left(\lambda_0 \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \end{vmatrix}$$
$$\leq \sup_{\lambda \in \mathbb{T}} \begin{vmatrix} \operatorname{Re} \left(\lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \end{vmatrix}$$
$$\leq \sup_{\lambda \in \mathbb{T}} \begin{vmatrix} \lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \end{vmatrix} = \begin{vmatrix} \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \end{vmatrix},$$

and hence (2.5) holds. Now, since $\begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix}$ is self adjoint for any $\lambda \in \mathbb{T}$, by [13, Theorem 3.3.6], we obtain

$$\left\| \begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| = \sup_{\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \left| \varphi \left(\begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix} \right) \right|.$$
(2.6)

Therefore,

$$\sup_{\lambda \in \mathbb{T}} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| \stackrel{(2.6)}{=} \sup_{\lambda \in \mathbb{T} \varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \sup_{\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \left| \varphi \left(\operatorname{Re} \left(\lambda \begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right|$$
$$= 2 \sup_{\lambda \in \mathbb{T} \varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \sup_{\lambda \in \mathbb{T} \varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \left| \operatorname{Re} \left(\lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right|$$
$$= 2 \sup_{\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \sup_{\lambda \in \mathbb{T}} \left| \operatorname{Re} \left(\lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right|$$
$$= 2 \sup_{\varphi \in \mathcal{S}(\mathbb{L}(\mathscr{V}))} \sup_{\lambda \in \mathbb{T}} \left| \operatorname{Re} \left(\lambda \varphi \left(\begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right) \right) \right|$$

Thus

$$\frac{1}{2} \sup_{\lambda \in \mathbb{T}} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| = \Omega(x).$$

We can obtain a refinement of inequality (2.4) as follows.

Theorem 2.7. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . For $x \in \mathscr{V}$ the following inequality holds:

$$\frac{1}{8} \left(4\|x\| + 2|\Gamma - \Gamma'| + \Delta + \Delta' \right) \le \Omega(x),$$

where $\Gamma = \max\left\{ \|x\|, \left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} \right\| \right\}, \Gamma' = \max\left\{ \|x\|, \left\| \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\| \right\}, \Delta = \left\| \|x\| - \left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} \right\|$
and $\Delta' = \left\| \|x\| - \left\| \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\| \right|.$

Proof. Since
$$\Omega(x) = \frac{1}{2} \sup_{\lambda \in \mathbb{T}} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|$$
, by taking $\lambda = 1$ and $\lambda = i$, we have
 $\Omega(x) \ge \frac{1}{2} \left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} \right\|$ and $\Omega(x) \ge \frac{1}{2} \left\| \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\|$. (2.7)

So, by (2.4) and (2.7) we have $\Omega(x) \ge \frac{1}{2} \max\{\Gamma, \Gamma'\}$. Therefore,

$$\begin{split} \Omega(x) &\geq \frac{\Gamma + \Gamma'}{4} + \frac{|\Gamma - \Gamma'|}{4} \\ &= \frac{1}{4} \left(\frac{1}{2} \left(\|x\| + \left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} \right\| \right) + \frac{1}{2} \Delta \right) \\ &\quad + \frac{1}{4} \left(\frac{1}{2} \left(\|x\| + \left\| \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\| \right) + \frac{1}{2} \Delta' \right) + \frac{|\Gamma - \Gamma'|}{4} \\ &= \frac{1}{8} \left(\left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} \right\| + \left\| \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\| \right) + \frac{1}{4} \|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} \\ &\geq \frac{1}{8} \left\| \begin{bmatrix} 0 & l_x \\ r_x & 0 \end{bmatrix} + \begin{bmatrix} 0 & -l_x \\ r_x & 0 \end{bmatrix} \right\| + \frac{1}{4} \|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} \\ &= \frac{1}{4} \left\| \left\| \begin{bmatrix} 0 & 0 \\ r_x & 0 \end{bmatrix} \right\| + \frac{1}{4} \|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} \\ &= \frac{1}{4} \|x\| + \frac{1}{4} \|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} \\ &= \frac{1}{2} \|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} . \end{split}$$

Thus

$$\frac{1}{2}\|x\| + \frac{\Delta + \Delta'}{8} + \frac{|\Gamma - \Gamma'|}{4} \le \Omega(x).$$

In the following result, we state a necessary and sufficient condition for the equality case in the inequality (2.4).

Corollary 2.8. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . Let $x \in \mathscr{V}$. Then $\Omega(x) = \frac{1}{2} ||x||$ if and only if $||x|| = \left\| \begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|$ for all $\lambda \in \mathbb{T}$.

Proof. Let us first suppose that $\Omega(x) = \frac{1}{2} ||x||$. For every $\lambda \in \mathbb{T}$ then we have $\Omega(\lambda x) = \frac{1}{2} ||\lambda x||$. Therefore, by Theorem 2.7, we obtain

$$\Delta = \left| \|\lambda x\| - \left\| \begin{bmatrix} 0 & l_{\lambda x} \\ r_{\lambda x} & 0 \end{bmatrix} \right\| = 0.$$

From this it follows that $||x|| = \left\| \begin{bmatrix} 0 & \lambda l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|.$

Conversely, if
$$||x|| = \left\| \begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix} \right\|$$
 for all $\lambda \in \mathbb{T}$, then
$$\frac{1}{2} \sup_{\lambda \in \mathbb{T}} \left\| \begin{bmatrix} 0 & \overline{\lambda}l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| = \frac{1}{2} ||x||,$$

and so, by Theorem 2.6, $\Omega(x) = \frac{1}{2} ||x||$.

For every $a \in \mathscr{A}$ and $x \in \mathscr{V}$, by the inequalities (2.1) and (2.4), we have

$$\Omega(xa + xa^*) \le ||xa + xa^*|| \le 2||a|| ||x|| \le 4||a||\Omega(x),$$

and hence

$$\Omega(xa + xa^*) \le 4 \|a\| \Omega(x). \tag{2.8}$$

In the following theorem, we improve the inequality (2.8).

Theorem 2.9. Let \mathscr{V} be a Hilbert \mathscr{A} -module. Let $a \in \mathscr{A}$ and $x \in \mathscr{V}$. Then

 $\Omega(xa + xa^*) \le 2||a + a^*||\Omega(x).$

Proof. Let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . For every $b \in \mathscr{A}$ and $y \in \mathscr{V}$, we have

$$r_{xa}(b) = (xa)b = x(ab) = x(T_a(b)) = r_x T_a(b)$$

and

$$l_{xa}(y) = \langle xa, y \rangle = a^* \langle x, y \rangle = a^* (l_x(y)) = T_{a^*} l_x(y).$$

Hence $r_{xa} = r_x T_a$ and $l_{xa} = T_{a^*} l_x$. Now, let $\lambda \in \mathbb{T}$. Therefore,

$$\begin{split} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_{(xa+xa^*)} \\ \lambda r_{(xa+xa^*)} & 0 \end{bmatrix} \right\| &= \left\| \begin{bmatrix} 0 & \overline{\lambda} (T_a * l_x + T_a l_x) \\ \lambda (r_x T_a + r_x T_{a^*}) & 0 \end{bmatrix} \right\| \\ &= \left\| \begin{bmatrix} 0 & \overline{\lambda} T_{a+a^*} & l_x \\ \lambda r_x T_{a+a^*} & 0 \end{bmatrix} \right\| \\ &= \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \begin{bmatrix} T_{a+a^*} & 0 \\ 0 & 0 \end{bmatrix} + \begin{bmatrix} T_{a+a^*} & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| \\ &\leq 2 \left\| \begin{bmatrix} T_{a+a^*} & 0 \\ 0 & 0 \end{bmatrix} \right\| \left\| \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix} \right\| \\ &\leq 4 \| a + a^* \| \Omega(x), \end{split}$$

and so

$$\frac{1}{2} \left\| \begin{bmatrix} 0 & \overline{\lambda} l_{(xa+xa^*)} \\ \lambda r_{(xa+xa^*)} & 0 \end{bmatrix} \right\| \le 2 \|a+a^*\|\Omega(x).$$

Taking the supremum over $\lambda \in \mathbb{T}$ in the above inequality, we deduce that

$$\Omega(xa + xa^*) \le 2||a + a^*||\Omega(x).$$

As an immediate consequence of Theorem 2.9, we have the following result.

Corollary 2.10. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $a \in \mathscr{A}$ and $x \in \mathscr{V}$. If $xa = xa^*$, then

$$\Omega(xa) \le ||a+a^*||\Omega(x).$$

Remark 2.11. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $a \in \mathscr{A}$ and $x \in \mathscr{V}$. Replace a by ia in Theorem 2.9, to obtain $\Omega(xa - xa^*) \leq 2||a - a^*||\Omega(x)$. Thus

$$\Omega(xa \pm xa^*) \le 2 \|a \pm a^*\| \Omega(x).$$

In what follows, r(a) stands for the spectral radius of an arbitrary element a in a C^* -algebra \mathscr{A} . It is well known that for every $a \in \mathscr{A}$, we have $r(a) \leq ||a||$ and that equality holds in this inequality if a is normal. The following lemma gives us a spectral radius inequality for sums of elements in C^* -algebras.

Lemma 2.12. [21, Lemma 3.5] Let \mathscr{A} be a C^* -algebra and let $a, b \in \mathscr{A}$. Then

$$r(a+b) \le \left\| \begin{bmatrix} \|a\| & \|ab\|^{1/2} \\ \|ab\|^{1/2} & \|b\| \end{bmatrix} \right\|.$$

Now, we present a refinement of the triangle inequality for the numerical radius in Hilbert C^* -modules. We use some ideas of [1, Theorem 3.4]. We refer the reader to [4, 7, 14, 18] for more information on the triangle inequality.

Theorem 2.13. Let \mathscr{V} be a Hilbert \mathscr{A} -module and let $\mathbb{L}(\mathscr{V})$ be the linking algebra of \mathscr{V} . Let $x, y \in \mathscr{V}$. Then

$$\Omega(x+y) \leq \left\| \begin{bmatrix} \Omega(x) & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & \Omega(y) \end{bmatrix} \right\| \leq \Omega(x) + \Omega(y).$$
Proof. Let $\lambda \in \mathbb{T}$. Put $a = \begin{bmatrix} 0 & \overline{\lambda} l_x \\ \lambda r_x & 0 \end{bmatrix}$ and $b = \begin{bmatrix} 0 & \overline{\lambda} l_y \\ \lambda r_y & 0 \end{bmatrix}$. Then
$$\|a\| \leq 2\Omega(x) \quad \text{and} \quad \|b\| \leq 2\Omega(y).$$

Also, for every $c \in \mathscr{A}$ and $z \in \mathscr{V}$, we have

$$l_x r_y(c) = l_x(yc) = \langle x, yc \rangle = \langle x, y \rangle c = T_{\langle x, y \rangle}(c)$$

and

$$r_{x}l_{y}(z) = r_{x}(\langle y, z \rangle) = x \langle y, z \rangle = \theta_{x,y}(z).$$

Thus $l_{x}r_{y} = T_{\langle x,y \rangle}$ and $r_{x}l_{y} = \theta_{x,y}$. Therefore, $ab = \begin{bmatrix} T_{\langle x,y \rangle} & 0\\ 0 & \theta_{x,y} \end{bmatrix}$ and hence,

$$\begin{aligned} & \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0\\ 0 & \theta_{x,y} \end{bmatrix} \right\| = \|ab\| \le \|a\| \|b\| \le 4\Omega(x)\Omega(y). \tag{2.9} \end{aligned}$$

Since $\begin{bmatrix} 0 & \overline{\lambda}l_{\langle x+y \rangle} \\ \lambda r_{\langle x+y \rangle} & 0 \end{bmatrix}$ is a self adjoint element of C^{*} -algebra $\mathbb{L}(\mathcal{V})$, we have

$$\begin{aligned} & \left\| \begin{bmatrix} 0 & \overline{\lambda}l_{\langle x+y \rangle} \\ \lambda r_{\langle x+y \rangle} & 0 \end{bmatrix} \right\| = r \left(\begin{bmatrix} 0 & \overline{\lambda}l_{\langle x+y \rangle} \\ \lambda r_{\langle x+y \rangle} & 0 \end{bmatrix} \right). \end{aligned}$$

Therefore, by Lemma 2.12, we obtain

$$\begin{vmatrix} 0 & \overline{\lambda}l_{(x+y)} \\ \lambda r_{(x+y)} & 0 \end{vmatrix} \end{vmatrix} = r \left(\begin{bmatrix} 0 & \overline{\lambda}l_{(x+y)} \\ \lambda r_{(x+y)} & 0 \end{vmatrix} \right)$$
$$= r(a+b)$$
$$\leq \left\| \begin{bmatrix} \|a\| & \|ab\|^{1/2} \\ \|ab\|^{1/2} & \|b\| \end{bmatrix} \right\|.$$

So, by the norm monotonicity of matrices with nonnegative entries (see, e.g., [9, p. 491]), we get

$$\begin{split} \left\| \begin{bmatrix} 0 & \overline{\lambda}l_{(x+y)} \\ \lambda r_{(x+y)} & 0 \end{bmatrix} \right\| &\leq \left\| \begin{bmatrix} \sup_{\lambda \in \mathbb{T}} & \sup_{\lambda \in \mathbb{T}} \|a\| & \sup_{\lambda \in \mathbb{T}} \|ab\|^{1/2} \\ \sup_{\lambda \in \mathbb{T}} \|ab\|^{1/2} & \sup_{\lambda \in \mathbb{T}} \|b\| \end{bmatrix} \right\| \\ &= \left\| \begin{bmatrix} 2\Omega(x) & \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & 2\Omega(y) \end{bmatrix} \right\|. \end{split}$$

Therefore, for every $\lambda \in \mathbb{T}$ we have

$$\frac{1}{2} \left\| \begin{bmatrix} 0 & \overline{\lambda}l_{(x+y)} \\ \lambda r_{(x+y)} & 0 \end{bmatrix} \right\| \leq \left\| \begin{bmatrix} \Omega(x) & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & \Omega(y) \end{bmatrix} \right\|,$$

whence

$$\Omega(x+y) \le \left\| \begin{bmatrix} \Omega(x) & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & \Omega(y) \end{bmatrix} \right\|.$$
(2.10)

On the other hand, by (2.9), we have

$$\begin{aligned} & \left\| \begin{bmatrix} \Omega(x) & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & \Omega(y) \end{bmatrix} \right\| \\ & = \frac{1}{2} \left(\Omega(x) + \Omega(y) + \sqrt{(\Omega(x) - \Omega(y))^2 + \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|} \right) \\ & \leq \frac{1}{2} \left(\Omega(x) + \Omega(y) + \sqrt{(\Omega(x) - \Omega(y))^2 + 4\Omega(x)\Omega(y)} \right) = \Omega(x) + \Omega(y). \end{aligned}$$
(2.11)

Thus

$$\left\| \begin{bmatrix} \Omega(x) & \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} \\ \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,y \rangle} & 0 \\ 0 & \theta_{x,y} \end{bmatrix} \right\|^{1/2} & \Omega(y) \end{bmatrix} \right\| \leq \Omega(x) + \Omega(y),$$

and the proof is completed.

As a consequence of Theorem 2.13, we have the following result.

Corollary 2.14. Let \mathscr{V} be a Hilbert \mathscr{A} -module, and $x, y \in \mathscr{V}$. If $\Omega(x + y) = \Omega(x) + \Omega(y)$, then

$$\Omega(x)\Omega(y) = \frac{1}{4} \left\| \begin{bmatrix} T_{\langle x,y\rangle} & 0\\ 0 & \theta_{x,y} \end{bmatrix} \right\|$$
$$\frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,x\rangle} & 0\\ 0 & \theta \end{bmatrix} \right\|^{1/2}.$$

In particular, $\Omega(x) = \frac{1}{2} \left\| \begin{bmatrix} T_{\langle x,x \rangle} & 0\\ 0 & \theta_{x,x} \end{bmatrix} \right\|^{1/2}$

The following lemma must be known to specialists. For the sake of completeness we include the proof.

Lemma 2.15. Let \mathscr{V} be a Hilbert \mathscr{A} -module, and $x, y \in \mathscr{V}$. Then

$$\|\theta_{x,y}\| = \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \right\|.$$

Proof. We may assume that $x, y \neq 0$ otherwise the identity trivially holds. We have

$$\begin{aligned} \left\| \theta_{x,y} \left(\frac{y \langle x, x \rangle^{1/2}}{\left\| y \langle x, x \rangle^{1/2} \right\|} \right) \right\|^2 &= \frac{\left\| x \langle y, y \rangle \langle x, x \rangle^{1/2} \right\|^2}{\left\| y \langle x, x \rangle^{1/2} \right\|^2} \\ &= \frac{\left\| \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle \langle y, y \rangle \langle x, x \rangle^{1/2} \right\|}{\left\| \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle^{1/2} \right\|} \\ &= \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle^{1/2} \right\| \\ &= \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle^{1/2} \right\| = \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \right\|^2, \end{aligned}$$

and so

$$\left\|\theta_{x,y}\left(\frac{y\langle x,x\rangle^{1/2}}{\left\|y\langle x,x\rangle^{1/2}\right\|}\right)\right\| = \left\|\langle x,x\rangle^{1/2}\langle y,y\rangle^{1/2}\right\|.$$

Hence

$$\|\theta_{x,y}\| \ge \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \right\|.$$
(2.12)

On the other hand, let $z \in \mathscr{V}$ with ||z|| = 1. By (1.1) we have $\langle y, z \rangle \langle z, y \rangle \leq \langle y, y \rangle$ and hence by Theorem 2.2.5(2) of [13] it follows that

$$\langle x, x \rangle^{1/2} \langle y, z \rangle \langle z, y \rangle \langle x, x \rangle^{1/2} \le \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle^{1/2}.$$

So, [13, Theorem 2.2.5(3)] implies

$$\left\| \langle x, x \rangle^{1/2} \langle y, z \rangle \langle z, y \rangle \langle x, x \rangle^{1/2} \right\| \le \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle \langle x, x \rangle^{1/2} \right\|.$$
(2.13)

Therefore,

$$\begin{aligned} \|\theta_{x,y}(z)\| &= \|x\langle y, z\rangle\| \\ &= \|\langle z, y\rangle \langle x, x\rangle \langle y, z\rangle\|^{1/2} \\ &= \left\|\langle x, x\rangle^{1/2} \langle y, z\rangle \langle z, y\rangle \langle x, x\rangle^{1/2}\right\|^{1/2} \\ &\stackrel{(2.13)}{\leq} \left\|\langle x, x\rangle^{1/2} \langle y, y\rangle \langle x, x\rangle^{1/2}\right\|^{1/2} = \left\|\langle x, x\rangle^{1/2} \langle y, y\rangle^{1/2}\right\|, \end{aligned}$$

whence

$$\|\theta_{x,y}\| \le \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \right\|.$$
(2.14)

Utilizing (2.12) and (2.14), we conclude that $\|\theta_{x,y}\| = \|\langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \|$. \Box

We close this paper with the following result.

Corollary 2.16. Let \mathscr{V} be a Hilbert \mathscr{A} -module, and $x, y \in \mathscr{V}$. If $\langle x, y \rangle = 0$, then

$$\Omega(x+y) \le \frac{1}{2} \left(\Omega(x) + \Omega(y) + \sqrt{(\Omega(x) - \Omega(y))^2 + \left\| \langle x, x \rangle^{1/2} \langle y, y \rangle^{1/2} \right\|} \right) \le \Omega(x) + \Omega(y).$$

Proof. Since $\langle x, y \rangle = 0$, we have $T_{\langle x, y \rangle} = 0$. Hence from (2.10), (2.11) and Lemma 2.15, we deduce the desired result.

References

- A. Abu-Omar and F. Kittaneh, Notes on some spectral radius and numerical radius inequalities, Studia Math. 227 (2015), no. 2, 97–109.
- A. Abu-Omar and F. Kittaneh, A generalization of the numerical radius, Linear Algebra Appl. 569 (2019), 323–334.
- A. F. Albideewi and M. Mabruk, On maps compressing the numerical range between C^{*}algebras, Adv. Oper. Theory 2 (2) (2017), 108–113.
- A. Al-Natoor and W. Audeh, Refinement of triangle inequality for the Schatten p-norm, Adv. Oper. Theory 5 (2020), no. 4, 1635–1645.
- D. Bakić and B. Guljaš, On a class of module maps of Hilbert C*-modules, Math. Commun. 7(2) (2002), 177–192.
- L. Carvalho, C. Diogo and S. Mendes, The star-center of the quaternionic numerical range, Linear Algebra Appl. 603 (2020), 166–185.
- R. Eskandari, M. S. Moslehian and D. Popovici, Operator equalities and characterizations of orthogonality in pre-Hilbert C^{*}-modules, Proc. Edinburgh Math. Soc. (2021), doi: 10.1017/S0013091521000341.
- K. E. Gustafson and D. K. M. Rao, Numerical range. The field of values of linear operators and matrices, Universitext. Springer-Verlag, New York, 1997.
- R. A. Horn and C. R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- E. C. Lance, Hilbert C^{*}-modules. A Toolkit for Operator Algebraists, London Mathematical Society Lecture Note Series, vol. 210, Cambridge University Press, Cambridge, 1995.
- V. M. Manuilov and E. V. Troitsky, *Hilbert C^{*}-modules*, In: Translations of Mathematical Monographs. 226, American Mathematical Society, Providence, RI, 2005.

- M. Mehrazin, M. Amyari and M. E. Omidvar, A new type of numerical radius of operators on Hilbert C^{*}-module, Rend. Circ. Mat. Palermo (2) 69 (2020), no. 1, 29–37.
- 13. G. J. Murphy, C*-Algebras and Operator Theory, Academic Press, New York, 1990.
- D. Popovici, Norm equalities in pre-Hilbert C^{*}-modules, Linear Algebra Appl. 436 (2012), no. 1, 59–70.
- I. Raeburn and D. P. Williams, Morita equivalence and continuous-trace C^{*}-algebras, Mathematical Surveys and Monographs 60, AMS, Philadelphia, 1998.
- R. Rajić, On the algebra range of an operator on a Hilbert C*-module over compact operators, Proc. Amer. Math. Soc. 131 (2003), no. 10, 3043–3051.
- 17. R. Rajić, A generalized q-numerical range, Math. Commun. 10 (2005), no. 1, 31–45.
- R. Rajić, Characterization of the norm triangle equality in pre-Hilbert C^{*}-modules and applications, J. Math. Inequal. 3 (2009), no. 3, 347–355.
- 19. D. Thaghizadeh, M. Zahraei, A. Peperko and N. Haj Aboutalebi, On the numerical ranges of matrices in max algebra, Banach J. Math. Anal. 14 (2020), 1773–1792.
- T. Yamazaki, On upper and lower bounds of the numerical radius and an equality condition, Studia Math. 178 (2007), no. 1, 83–89.
- A. Zamani, Characterization of numerical radius parallelism in C*-algebras, Positivity 23 (2019), no. 2, 397–411.
- 22. A. Zamani and P. Wójcik, Another generalization of the numerical radius for Hilbert space operators, Linear Algebra Appl. 609 (2021), 114–128.

School of Mathematics and Computer Sciences, Damghan University, Damghan, P. O. BOX 36715-364, Iran

Email address: zamani.ali85@yahoo.com