

# Distance 4 Curves on Closed Surfaces of Arbitrary Genus

Kuwari Mahanta and Sreekrishna Palaparthi\*

**ABSTRACT.** Let  $S_g$  denote a closed, orientable surface of genus  $g \geq 2$  and  $\mathcal{C}(S_g)$  be the associated curve complex. The mapping class group of  $S_g$ ,  $Mod(S_g)$  acts on  $\mathcal{C}(S_g)$  by isometries. Since Dehn twists about certain curves generate  $Mod(S_g)$ , one can ask how Dehn twists move specific vertices in  $\mathcal{C}(S_g)$  away from themselves. We show that if two curves represent vertices at a distance 3 in  $\mathcal{C}(S_g)$  then the Dehn twist of one curve about another yields two vertices at distance 4. This produces many tractable examples of distance 4 vertices in  $\mathcal{C}(S_g)$ . We also show that the minimum intersection number of any two curves at a distance 4 on  $S_g$  is at most  $(2g - 1)^2$ .

**Keywords :** Curve complex, Minimal intersection number, Distance 4 curves, Filling pairs of curves

**Mathematics Subject Classification :** 27M60, 20F65, 20F67


## 1. Introduction

Let  $S_g$  denote a closed, orientable surface of genus  $g \geq 2$ . Throughout this article, a curve on  $S_g$  will mean an essential simple closed curve on it. For two curves,  $\alpha, \beta$  in  $S_g$ ,  $i(\alpha, \beta)$  denotes their geometric intersection number and  $T_\alpha(\beta)$  denotes the Dehn twist of the curve  $\beta$  about the curve  $\alpha$ . In [9], Harvey associated with  $S_g$  a simplicial complex called the complex of curves which is denoted by  $\mathcal{C}(S_g)$  and defined as follows. The 0-skeleton,  $\mathcal{C}^0(S_g)$  of this complex is in one-to-one correspondence with isotopy classes of essential simple closed curves on  $S_g$ . Two vertices span an edge in  $\mathcal{C}(S_g)$  if and only if these vertices have mutually disjoint representatives.  $\mathcal{C}^0(S_g)$  can be equipped with a metric,  $d$  by defining the distance between any two vertices to be the minimum number of edges in any edge path between them in  $\mathcal{C}(S_g)$ . By the distance between two curves on  $S_g$ , we mean the distance between the corresponding vertices in  $\mathcal{C}(S_g)$ . The minimal intersection number between any two curves on  $S_g$  which are at a distance  $n$  is denoted by  $i_{min}(g, n)$ .


Masur and Minsky, in their seminal paper [12], proved that  $\mathcal{C}(S)$  is a  $\delta$ -hyperbolic space with the metric  $d$ . Later, it was shown that the  $\delta$  can be chosen to be independent of the surface  $S_g$ , see [1], [5], [6], [10], [14]. The coarse geometry of the curve complex has various applications to 3-manifolds, Teichmüller theory and mapping class groups. One can see [13] for many such applications.

[16], [18], [17] and [4] gave algorithms to compute distances between two vertices in  $\mathcal{C}(S_g)$ . The algorithm in [4], although for closed surfaces is by far the most effective in calculating distances in  $\mathcal{C}(S_g)$ . Given vertices,  $\nu, \mu$  in  $\mathcal{C}(S_g)$ , the algorithm works by giving an initially efficient geodesic,  $\nu_0 = \nu, \nu_1, \dots, \nu_n = \mu$  where  $\nu_1$  is chosen from a list of  $n^{6g-6}$  possible vertices.

In [3], Aougab and Huang gave all  $Mod(S_g)$  orbits of minimally intersecting distance 3 curves in  $\mathcal{C}(S_g)$  by giving the curves as permutations which are solutions



**Figure 1** Distance 3 vertices in  $\mathcal{C}(S_4)$




**Figure 2** Distance 4 vertices in  $\mathcal{C}(S_4)$

to a system of equation in the permutation group of order  $8g - 4$ . They further showed that for genus,  $g \geq 3$ ,  $i_{\min}(g, 3) = 2g - 1$  and that all pairs of curves which intersect at  $2g - 1$  points and cuts up  $S_{g \geq 3}$  into a disc are at distance 3.

For vertices at distance 4 in  $\mathcal{C}(S_g)$ , we found only limited pictures (like, Figure 2) of such curves in  $S_{g \leq 3}$  and none for  $S_{g > 3}$ . In [8], the authors gave a test to determine when two vertices in  $\mathcal{C}(S_g)$  are at a distance  $\geq 4$  using the efficient geodesic algorithm of [4]. Using the MICC software, they showed that  $i_{\min}(2, 4) = 12$  by giving all minimally intersecting pairs of curves at distance 4. They also gave examples of curves which are at a distance 4 on a surface of genus 3 and concluded that  $i_{\min}(3, 4) \leq 29$  (refer [8] Theorem 1.8).

Although  $i_{\min}(g, 4)$  is still not known, in [2], Aougab and Taylor proved that  $i_{\min}(g, 4) = O(g^2)$  by answering a more general question by Dan Margalit that  $i_{\min}(g, n) = O(g^{n-2})$ .

In this article, we give a method to construct examples of vertices at distance 4 in  $\mathcal{C}(S_g)$  and improve the known upper bound of  $i_{\min}(g, 4)$  to  $(2g - 1)^2$ . In particular, using any of the minimally intersecting filling pairs of curves on  $S_3$ , as described in [3], one can get examples of curves at distance 4 in  $\mathcal{C}(S_3)$  which intersect at 25 points. This implies that  $i_{\min}(3, 4) \leq 25$ . Using a minimally intersecting pair



**Figure 3**  $Q_1, Q_2, Q_3, Q_4$  is an example of an initially efficient geodesic in  $\mathcal{C}(S_2)$

of curves (Figure 1) described in [3] as a permutation of 28 symbols, we use our method to construct a pair of distance 4 curves in  $\mathcal{C}(S_4)$  as in Figure 2.

Such workable examples of distance 4 curves were a result of trying to investigate the effect of certain Dehn twists on distances in  $\mathcal{C}(S_g)$  as follows:

REMARK 1. *If  $d(\alpha, \gamma) = 1$  or  $2$ , then  $d(\alpha, T_\gamma(\alpha)) = d(\alpha, \gamma)$ .*

In general, one can ask the following question:

QUESTION 1. *If  $d(\alpha, \gamma) > 2$ , then what is the relation between  $d(\alpha, \gamma)$  and  $d(\gamma, T_\gamma(\alpha))$ ?*

In this article, we prove that if  $d(\alpha, \gamma) \geq 3$  then  $d(\alpha, T_\gamma(\alpha)) \geq 3$ . We then answer the question 1 for  $d(\alpha, \gamma) = 3$  and show that  $d(\alpha, T_\gamma(\alpha)) = d(\alpha, \gamma) + 1$ .

## 2. Setup

For any ordered index in this work, we follow cyclical ordering. For instance, if  $i \in \{1, 2, \dots, k\}$ ,  $i = k + 1$  will indicate  $i = 1$ .

Two curves,  $\nu_1$  and  $\nu_2$  on  $S_g$  are said to be a filling pair if every component of  $S_g \setminus \{\nu_1, \nu_2\}$  is a disk. A component,  $D$  of  $S_g \setminus \{\nu_1, \nu_2\}$  is said to be an  $2n$ -gon if its boundary comprises of  $n$  arcs of  $\nu_1$  and  $\nu_2$ . Consider a pair,  $\alpha, \beta$ , of filling curves on  $S_g$  with geometric intersection number,  $i(\alpha, \beta) = k$ . Let their set of intersection points be  $\{w_1, \dots, w_k\}$ . Define a triangulation,  $G$  of  $S$  as follows:

Let  $\{w'_1, \dots, w'_k\}$  be the 0-skeleton of  $G$ . There is an edge between  $w'_i$  and  $w'_j$  in  $G$  if and only if there is an arc of  $\alpha$  or,  $\beta$  between the two intersection points,  $w_i, w_j$  in  $S$ . As  $\alpha$  and  $\beta$  are filling pairs, for each disk component,  $D$  of  $S_g \setminus \{\alpha, \beta\}$  attach a disk to the cycle formed by the edges of  $G$  corresponding to the arcs of  $\alpha, \beta$  that form the boundary of  $D$ . Using the Euler characteristic of a 2-dimensional complex, the number of faces,  $f$  of  $G$  is given by  $f = k + 2 - 2g$ .

Consider a geodesic,  $\nu_0, \dots, \nu_N$  of length  $N$  in  $\mathcal{C}^0(S)$ . An arc,  $\omega$  in  $S$  is a *reference arc* for the triple  $\nu_0, \nu_1, \nu_N$  if  $\omega$  and  $\nu_1$  are in minimal position and the interior of  $\omega$  is disjoint from  $\nu_0 \cup \nu_N$ . The oriented geodesic  $\nu_0, \dots, \nu_N$  is said to be *initially efficient* if  $i(\nu_1, \omega) \leq N - 1$  for all choices of reference arc,  $\omega$ . The authors of [4] prove that there exists an initially efficient geodesic between any two vertices of  $\mathcal{C}(S)$ .

The following theorem from [8] gives a criterion for detecting vertices in  $\mathcal{C}(S)$  at distance at-least 4.

THEOREM 1 (Theorem 1.3, [8]). *For the filling pair,  $\kappa, \omega$ , let  $\Gamma \subset \mathcal{C}^0(S)$  be the collection of all vertices such that the following hold :*

- (1) for  $\bar{\gamma} \in \Gamma$ ,  $d(\kappa, \bar{\gamma}) = 1$ ; and
- (2) for  $\bar{\gamma} \in \Gamma$ ; for each segment,  $b \subset \omega \setminus \kappa$ ,  $i(\bar{\gamma}, b) \leq 1$ .

Then  $d(\kappa, \omega) \geq 4$  if and only if  $d(\bar{\gamma}, \omega) \geq 3$  for all  $\bar{\gamma} \in \Gamma$ .

Let  $\lambda$  and  $\mu$  be two simple closed curves on  $S_g$  and let  $R_\lambda$  and  $R_\mu$  be closed regular neighborhoods of  $\lambda$  and  $\mu$  respectively. We say that the 4-tuple  $(\lambda, \mu, R_\lambda, R_\mu)$  is *amenable to Dehn twist in special position* if the following hold:

- (1)  $\lambda$  and  $\mu$  intersect transversely and minimally on  $S_g$ ,
- (2)  $\lambda$  and  $\mu$  fill  $S_g$ ,
- (3) the number of components of  $R_\lambda \cap R_\mu$  is equal to the number of components of  $S \setminus (\lambda \cup \mu)$  and each of these components is a disc.

When  $\lambda$  and  $\mu$  fill  $S_g$ , by considering a Euclidean model of  $S_g$ , it is easy to see that a 4-tuple  $(\lambda, \mu, R_\lambda, R_\mu)$  amenable to Dehn twist in special position always exists.


Consider a 4-tuple  $(\lambda, \mu, R_\lambda, R_\mu)$  which is amenable to Dehn twist in special position. Let  $i(\lambda, \mu) = k$  and  $K := \{1, 2, \dots, k\}$ . We construct a curve in the isotopy class of  $T_\lambda(\mu)$  which we call  $T_\lambda(\mu)$  in special position w.r.t. the 4-tuple  $(\lambda, \mu, R_\lambda, R_\mu)$ . Start at any one of the components of  $R_\lambda \cap R_\mu$  and label it as  $A_1$ . Since  $\mu$  intersects  $\lambda$  transversely, the arc  $\mu_1$  of  $\mu$  contained in  $A_1$  which has its endpoints  $X$  and  $Y$  on boundary arcs of  $R_\lambda$  is such that  $X$  and  $Y$  lie on distinct boundary component of  $\partial R_\lambda$ . We call the component of  $\partial R_\lambda$  containing  $X$  to be  $\partial_+ R_\lambda$  and the other component containing  $Y$  to be  $\partial_- R_\lambda$ . Equip  $A_1$  with the Euclidean metric such that it is a square in the  $xy$ -plane. Two opposite sides of  $A_1$  are formed from the arcs of  $\partial R_\lambda$  and the two remaining sides formed from arcs of  $\partial R_\mu$  and the  $x$ -axis lies along  $\mu_1$  and the value of the  $x$ -coordinate increases from  $X$  to  $Y$ . Orient  $\mu_1$  from  $X$  to  $Y$ . This induces an orientation on  $\mu$ . Next we pick  $k$  distinct points  $\{q_1, q_2, \dots, q_k\}$  in the interior of  $\mu_1$  such that the  $x$ -coordinate of  $q_i$  is greater than the  $x$  coordinate of  $q_j$  whenever  $i > j$  and  $i, j \in K$ . For each  $i \in K$ , let  $\lambda_i$  be a curve in  $R_\lambda$  which is isotopic to  $\lambda$  and passes through  $q_i$ . Further for each  $i, j \in K, i \neq j$  let  $\lambda_i$  and  $\lambda_j$  be disjoint.

Orient  $\lambda_1$  such that the  $y$ -coordinate on  $\lambda_1$  increases when following this orientation in the disk  $A_1$ . Starting with  $A_1$ , label the subsequent disk components,  $R_\lambda \cap R_\mu$ , as  $A_2, A_3, \dots, A_k$ , in the orientation of  $\lambda_1$ . For each  $i \in K$ ,  $A_i$  contains a unique arc of  $\mu$  which we label as  $\mu_i$ .  $\mu_i$  gets an induced orientation from  $\mu$ . For each  $i \in K$ , equip  $A_i$  with Euclidean metric and assume it to be a square in the  $xy$ -plane where  $\mu_i$  lies along the  $x$ -axis with the  $x$  coordinate increasing along the orientation of  $\mu_i$ . Assume  $A_i$  to be positioned such that  $\mu_i$  is the line segment joining the mid-points of the left and right sides of the square. In this orientation, call the component of  $\partial R_\mu$  which appears above  $\mu_i$  as  $\partial_+ R_\mu$  and the component of  $\partial R_\mu$  below  $\mu_i$  as  $\partial_- R_\mu$ . However, note that the side of  $A_i$  which is formed of the arcs of  $\partial_+ R_\lambda$  could either be to the right or to the left of this square. Accordingly, the side of  $A_i$  which is formed of the arcs of  $\partial_- R_\lambda$  could either be to the left or to the right of this square. For  $i, j \in K$ , by an isotopy inside  $A_i$ , we can assume that all the arcs of  $\lambda_j$  in  $A_i$  are straight lines.


For each  $i, j \in K$ , let  $u_{i,j} := A_i \cap \lambda_j \cap \partial_+ R_\mu$  and  $v_{i,j} := A_i \cap \lambda_j \cap \partial_- R_\mu$ . Also for each  $i \in K$  let the left end point of  $\mu_i$  in the square  $A_i$  be  $v_{i,0}$  and the right end point of  $\mu_i$  in the square  $A_i$  be  $u_{i,k+1}$ . Construct the Dehn twist of  $\mu$  about  $\lambda$  as follows: For each  $j \in K \cup \{0\}$  draw line segments,  $\theta_{i,j}$ , connecting  $v_{i,j}$  to  $u_{i,j+1}$ .  $T_\lambda(\mu)$  is the curve

$$((\mu \cup (\cup_{i \in K} \lambda_i)) \cap (S \setminus (\cup_{i \in K} A_i))) \cup (\cup_{i,j \in K} \theta_{i,j}).$$

The schematic, figure 4, shows  $A_i$  before and after this transformation. In the complement of  $A_i$ 's the transformation described above does not disturb the curves  $\lambda_i$ 's and  $\mu$ . In [7], an algorithm to obtain the Dehn twist,  $T_\lambda(\mu)$  has been described



**Figure 4** Disk of transformation before (figure on the left) and after (figure on the right) the Dehn twist



**Figure 5** Surgery of the curves to obtain  $T_\lambda(\mu)$

such that the curves in the discs of transformation are as in figure 5. The line segments in figure 4 are isotopic to the corresponding curves in 5 which shows that the above transformation indeed results in  $T_\lambda(\mu)$ . When  $T_\lambda(\mu)$  is constructed as above and as shown in figure 4, we say that  $T_\lambda(\mu)$  is in *special position w.r.t.  $\lambda$  and  $\mu$* . We call the  $k$  copies of  $\lambda$ ,  $\lambda_i$ ,  $i \in K$ , and  $\mu$  to be the *scaffolding* for  $T_\lambda(\mu)$ . We call the Euclidean disks  $A_i$ ,  $i \in K$ , along with the line segments  $\theta_{i,j}$ 's for  $j \in K$  to be the *disks of transformation* for  $T_\lambda(\mu)$ . The points  $u_{i,j}$ 's,  $v_{i,j}$ 's,  $u_{i,k+1}$  and  $v_{i,0}$  for  $i, j \in K$  shall hold their meaning as defined in the context of the disks of transformations. So, using these phrases, when  $T_\lambda(\mu)$  is in special position w.r.t.  $\lambda$  and  $\mu$ , the scaffolding of  $T_\lambda(\mu)$  remains unchanged outside its disks of transformation. Inside the disks of transformation for  $T_\lambda(\mu)$ , the schematic in figure 4 describes the changes to its scaffolding.


### 3. Distance 4 curves in $\mathcal{C}(S_{g \geq 2})$

**THEOREM 2.** *Let  $S$  be a surface of genus  $g \geq 2$ . Let  $\alpha$  and  $\gamma$  be two curves on  $S$  with  $d(\alpha, \gamma) = 3$ . Then,  $d(T_\gamma(\alpha), \alpha) = 4$ .*

**PROOF.** Let  $\nu_0, \nu_1, \nu_2, \nu_3$  be a geodesic from the vertex  $\nu_0$  corresponding to  $\alpha$  to the vertex  $\nu_3$  corresponding to  $\gamma$  in  $\mathcal{C}(S)$ . Let  $T_\gamma(\nu_0)$  be the vertex in  $\mathcal{C}(S)$  corresponding to  $T_\gamma(\alpha)$ . The existence of the path  $T_\gamma(\nu_0), T_\gamma(\nu_1), T_\gamma(\nu_2) = \nu_2, \nu_1, \nu_0$  gives that  $d(T_\gamma(\alpha), \alpha) \leq 4$ . We prove that  $d(T_\gamma(\alpha), \alpha) \geq 4$  by using Theorem 1 with  $\kappa = T_\gamma(\alpha)$  and  $\omega = \alpha$ , hence showing that  $d(T_\gamma(\alpha), \alpha) = 4$ .

**CLAIM 1.**  $T_\gamma(\alpha)$  and  $\alpha$  fill  $S$ .

*Proof of claim 1:* Let  $i(\gamma, \alpha) = k$ ,  $K := \{1, 2, \dots, k\}$ ,  $K_{-1} := \{1, 2, \dots, k-1\}$  and  $K_{2-2g} := \{1, 2, \dots, k+2-2g\}$ . We refer to section 2 for the terminology used here. Since  $\alpha$  and  $\gamma$  fill  $S$ , there is a 4-tuple  $(\alpha, \gamma, R_\alpha, R_\gamma)$  which is amenable to



**Figure 6** The scaffolding for  $T_{Q_1}(Q_4)$ , where  $Q_1$  and  $Q_4$  are from example in figure 3 and the shaded region is a rectangle of the scaffolding

Dehn twist in special position. Let  $T_\gamma(\alpha)$  be in special position w.r.t to  $\alpha$  and  $\gamma$ . We denote the disks of transformation of  $T_\gamma(\alpha)$  by  $A_i$ , for  $i \in K$ . By an isotopy we assume the curve  $\alpha$  to be disjoint from  $T_\gamma(\alpha) \setminus A_i$  for  $i \in K$  and in each  $A_i$  we further assume the arc  $\alpha_i := \alpha \cap A_i$  to be a straight line segment below the segment connecting  $v_{i,0}$  and  $u_{i,k+1}$ .


For  $i \in K$ , let  $g_i$  along with  $\alpha$  be the scaffolding for  $T_\gamma(\alpha)$ . For  $j \in K_{-1}$ , one of the components of  $S \setminus \{g_j, g_{j+1}\}$  is an annulus,  $G_j$ . Any component of  $G_j \setminus \alpha$  is a 4-gon which we call as a *rectangle of the scaffolding for  $T_\gamma(\alpha)$* . Figure 6 shows an example of such a rectangle of the scaffolding. The disks  $A_i$ ,  $i \in K$ , further divide each rectangle of the scaffolding into three components. There is a unique  $i \in K$  such that  $A_i$  and  $A_{i+1}$  intersect a given rectangle of the scaffolding. Denote a rectangle of the scaffolding formed out of  $G_j$  with its arcs of  $\alpha$  lying in  $A_i$  and  $A_{i+1}$  by  $B_{i,j}$ . Denote the sub-rectangles  $B_{i,j} \cap A_i$ , by  $C'_{i,j}$  and  $B_{i,j} \cap A_{i+1}$ , by  $C''_{i+1,j}$ . Also let  $B'_{i,j} := B_{i,j} \setminus (C'_{i,j} \cup C''_{i+1,j})$ . Let

$$B = \cup_{i=1}^k \cup_{j=1}^{k-1} B_{i,j}.$$


$S \setminus (\alpha \cup \gamma)$  has  $k + 2 - 2g$  disk components by Euler characteristic considerations. If  $F_p$  is a disk component of  $S \setminus (\alpha \cup \gamma)$ , for some  $p \in K_{2-2g}$ , then  $F'_p := F_p \setminus B$  is a single disk as  $B$  intersects any  $F_p$  only in disks which contain a boundary arc of  $F_p$ , namely arcs of  $\gamma$ . The components of  $S \setminus (\alpha \cup g_1 \cup \dots \cup g_k)$  comprise of  $k(k-1)$  rectangles of the scaffolding for  $T_\gamma(\alpha)$ , namely  $B_{i,j}$  where  $i \in K$ ,  $j \in K_{-1}$ , and  $k + 2 - 2g$  even sided polygonal discs, namely  $F'_p$ , where  $p \in K_{2-2g}$ . Let  $F''_p$  denote  $F'_p \setminus R_\alpha$  for  $p \in K_{2-2g}$ .

For each  $j \in K$  let  $w_{i,j} := \theta_{i,j} \cap \alpha_i$ . For each  $i \in K$  and  $j \in K_{-1}$ , let  $D''_{i,j}$  be the parallelogram with vertices  $v_{i,j}, v_{i,j+1}, w_{i,j}$  and  $w_{i,j+1}$  and  $D'_{i,j+1}$  be the parallelogram with vertices  $w_{i,j}, w_{i,j+1}, u_{i,j+1}, u_{i,j+2}$ . In each disk  $A_i$ , for  $i \in K$ , there is a pentagon,  $P_{i,1}$ , which is above  $\alpha_i$  and bounded by the lines  $\theta_{i,0}$ ,  $\partial R_\gamma$ ,  $\alpha_i$ ,  $\theta_{i,1}$  and the line segment of  $\partial_+ R_\alpha$  between  $u_{i,1}$  and  $u_{i,2}$ . Likewise, in each disk  $A_i$ , for  $i \in K$ , there is a triangle,  $T_{i,k+1}$ , which is bounded by the lines  $\alpha_i$ ,  $\theta_{i,k}$  and  $\partial R_\gamma$ . Figure 7 shows a schematic before and after the transformation to the disk  $A_i$ ; the figure to the left shows the rectangles  $C'_{i,1}$  and  $C''_{i,k}$  and the figure on the right shows  $P_{i,1}$  and  $T_{i,k+1}$ .


Figure 8 shows a schematic of  $R_\alpha$  before and after the transformation to the scaffolding of  $T_\gamma(\alpha)$ . The shaded region in the figure on the left shows  $C'_{i,j}$  and  $C''_{i,j-1}$  for some indices  $i, j$ . The shaded region in the figure on the right shows  $D'_{i,j}$  and  $D''_{i,j-1}$  for some indices  $i, j$ .



**Figure 7** The disk of transformation for  $T_\gamma(\alpha)$ : the figure on the left shows the portion of the scaffolding for  $T_\gamma(\alpha)$ ; the figure on the right shows the pentagon  $P_{i,1}$ , the triangle  $T_{i,k+1}$  and the parallelograms formed due to  $\alpha_i$  and  $T_\gamma(\alpha)$



**Figure 8** A schematic of  $R_\alpha$  (figure on the left) and after (figure on the right) the Dehn twist



**Figure 9** Two adjacent disks of transformation in  $R_\alpha$

For  $i \in K$ , note that all the disks  $A_i$ , occur in some sequence in the annulus  $R_\alpha$  when moving along  $\alpha$ . So, a disk  $A_i$  is connected to some disk  $A_j$  on the left and to some other disk  $A_p$  on the right by a single arc of  $\alpha \setminus R_\gamma$ , for some distinct indices  $i, j, p \in K$ . The schematic for two disks  $A_i$  and  $A_j$ , for some  $i, j \in K$ , which are connected via a single arc of  $\alpha \setminus R_\gamma$  and an arc of  $T_\gamma(\alpha) \setminus R_\gamma$  is as shown in the figure 9. Note that this schematic is generic since for every  $j \in K$ , there is a distinct  $i \in K$  such that  $A_j$  occurs to the left of  $A_i$ , in the sense mentioned above.

Figure 9 is a schematic of a portion of figure 8. In any of the cases, *viz.*  $\partial_+ R_\gamma$  and  $\partial_+ R_\gamma$  face each other,  $\partial_+ R_\gamma$  and  $\partial_- R_\gamma$  face each other or  $\partial_- R_\gamma$  and  $\partial_- R_\gamma$  face each other, In this schematic, we see that the pentagon  $P_{i,1}$  of the disk  $A_i$  is connected to the triangle  $T_{j,k+1}$  of  $A_j$  via an arc of  $\alpha \setminus R_\gamma$ ,  $\omega_{i,j}$ , and an arc of  $T_\gamma(\alpha)$ ,  $\eta_{i,j}$ . The disk,  $R_{i,j}$  outside  $R_\gamma$  bounded by  $\omega_{i,j}$ ,  $\eta_{i,j}$  and two arcs of  $\partial R_\gamma$ , will be called a *conduit*. Equip the conduit with the Euclidean metric and assume that  $R_{i,j}$  is a rectangle with two opposite sides  $\omega_{i,j}$  and  $\eta_{i,j}$ . Now  $P_{i,1} \cup R_{i,j} \cup T_{j,k+1}$  is

a 4-gon bounded by four arcs *viz.* (i)  $\theta_{i,0} \cup \eta_{i,j} \cup \theta_{j,k}$ , (ii)  $\alpha_j \cup \omega_{i,j} \cup \alpha_i$ , (iii)  $\theta_{i,1}$  and (iv) the arc of  $\partial_+ R_\alpha$  between  $u_{i,1}$  and  $u_{i,2}$ . This protracted 4-gon will be denoted by  $D'_{i,1}$ .

Let  $S' = S \setminus R_\alpha$ . The components of  $S \setminus (\alpha \cup T_\gamma(\alpha))$  are the components of  $S' \setminus T_\gamma(\alpha)$  and the components of  $R_\alpha \setminus (\alpha \cup T_\gamma(\alpha))$  glued at the boundary of  $R_\alpha$ . Since the changes to the scaffolding of  $T_\gamma(\alpha)$  is restricted to  $R_\alpha$ , the components of  $S' \setminus T_\gamma(\alpha)$  are precisely the disc components of  $S' \setminus (g_1 \cup \dots \cup g_k)$ .

The components of  $S' \setminus (g_1 \cup \dots \cup g_k)$  are  $B'_{i,j}$ ,  $i \in \{1, 2 \dots k\}$ ,  $j \in K_{-1}$ , along with disks  $F''_p$ ,  $p \in K_{2-2g}$ , as explained above. The components of  $R_\alpha \setminus (\alpha \cup T_\gamma(\alpha))$  will be examined using the schematic figure 9 of a portion of  $R_\alpha$ . There are four kinds of regions in  $R_\alpha$ . The upper disk regions, like  $R_1$  in the schematic figure 9, the lower disk regions, like  $R_2$  in the schematic figure 9, and the disks  $D'_{i,j}$ ,  $D''_{i,j}$ ,  $i \in K$ ,  $j \in K_{-1}$ . Figure 9 shows how the upper and lower disk regions are glued to disks  $F''_p$  for  $p \in K_{2-2g}$ . For each  $p \in K_{2-2g}$ , after gluing the lower disk regions and the upper disk regions to the respective disks  $F''_p$ , we get disks which we denote by  $F'''_p$ . We know that  $F'''_p$  is a disk because the upper and the lower disk regions are disjoint, except for the points  $w_{i,j}$  on the boundary and share a single arc of  $\partial R_\alpha$  with a unique  $F''_p$ . For each  $p \in K_{2-2g}$ , we call  $F'''_p$  to be the *modified disk* corresponding to the initial disk  $F_p$ .


For each  $i \in K$  and  $j \in K_{-1}$ , the line segment of  $\partial_+ R_\alpha$  between  $u_{i,j}$   $u_{i,j+1}$  is the common boundary of  $C'_{i,j}$  and  $D'_{i,j}$ . Likewise, for each such  $i, j$ , the line segment of  $\partial_- R_\alpha$  between  $v_{i,j}$   $v_{i,j+1}$  is the common boundary of  $C''_{i,j}$  and  $D''_{i,j}$ . So, for such  $i, j$ , when considering the components of  $S \setminus (\alpha \cup g_1 \cup \dots \cup g_k)$  the rectangular core  $B'_{i,j}$  is connected to  $C'_{i,j}$  along the boundary segment  $u_{i,j}$   $u_{i,j+1}$  and to  $C''_{i+1,j}$  along the boundary segment  $v_{i+1,j}$   $v_{i+1,j+1}$ , whereas when considering the components of  $S \setminus (\alpha \cup T_\gamma(\alpha))$ , the rectangular core  $B'_{i,j}$  is connected to  $D'_{i,j}$  along the boundary segment  $u_{i,j}$   $u_{i,j+1}$  and  $D''_{i+1,j}$  along the boundary segment  $v_{i+1,j}$   $v_{i+1,j+1}$ . So the rectangles of the scaffolding for  $T_\gamma(\alpha)$ ,  $B_{i,j}$ , which are components of  $S \setminus (\alpha \cup g_1 \cup \dots \cup g_k)$ , after the transformation in the disks of transformation for  $T_\gamma(\alpha)$  result in disks  $E_{i,j} := B'_{i,j} \cup D'_{i,j} \cup D''_{i+1,j}$  which now are components of  $S \setminus (\alpha \cup T_\gamma(\alpha))$ . For each  $p \in K_{2-2g}$ ,  $F'''_p$  is a disk as seen earlier. The components of  $S \setminus (\alpha \cup T_\gamma(\alpha))$  are precisely the disks  $F'''_p$  and  $E_{i,j}$  where  $p \in K_{2-2g}$ ,  $i \in K$  and  $j \in K_{-1}$ . This proves that the components of  $S \setminus (\alpha \cup T_\gamma(\alpha))$  are all disks and hence proving Claim 1.

The components of  $R_\gamma \setminus T_\gamma(\alpha)$  are disks and their boundary consists of two arc segments of  $T_\gamma(\alpha)$  and one each of  $\partial_+ R_\gamma$  and  $\partial_- R_\gamma$ . We call these disks as *rectangular tracks*. The word tracks derives its motivation from how these tracks appear in  $R_\gamma$ . Figure 10 shows  $R_\gamma$  and rectangular tracks inside  $R_\gamma$ .


Since  $i(\alpha, \gamma) = k$ , there are  $k$  components of  $\alpha \cap R_\gamma$ . Every component of  $\alpha \setminus T_\gamma(\alpha)$  is either contained in  $R_\gamma$  or, has a sub-arc which is contained in  $R_\gamma$ . For any  $i \in K$ ,  $\alpha_i$  intersects the rectangular tracks.

Let  $i_0 \in K$ . In the schematic figure 7,  $A_{i_0}$  has exactly  $k + 1$  arcs of  $T_\gamma(\alpha)$ . Call  $\theta_{i_0,0}$  to be the *leftmost arc* of  $A_{i_0}$  and  $\theta_{i_0,k}$  to be the *rightmost arc* of  $A_{i_0}$ . Let us consider one component of  $T_\gamma(\alpha) \cap R_\gamma$ , call it  $\rho_{i_0}$ , which intersects  $A_{i_0}$  in its leftmost arc. This  $\rho_{i_0}$  intersects  $A_{i_0}$  precisely in the arcs  $\theta_{i_0,0}$  and  $\theta_{i_0,k}$  and it intersects  $A_j$  for every  $j \in K \setminus \{i_0\}$  in the arcs  $\theta_{j,m}$  where  $m = (j - i_0)(\text{mod } k)$ . This is easily seen from the construction of  $T_\gamma(\alpha)$  in special position w.r.t.  $\alpha$  and  $\gamma$ . From this discussion it is clear that  $\rho_{i_0}$  intersects each  $\alpha_j$ , for  $j \in K$ , exactly once. It is also clear that, for  $j \in K$ , the points of  $\rho_{i_0} \cap \alpha_j$  lie on  $\rho_{i_0}$  in the order  $\alpha_{i_0+1}, \dots, \alpha_k, \alpha_1, \dots, \alpha_{i_0-1}, \alpha_{i_0}$  when  $\rho_{i_0}$  is traversed from  $\partial_+ R_\gamma$  to  $\partial_- R_\gamma$ . We now consider two arc components,  $\rho_{i_0}$  and  $\rho_{i_0+1}$ , of  $T_\gamma(\alpha) \cap R_\gamma$  and the rectangular track,  $T_{i_0}$ , which is enclosed by these two components in  $R_\gamma$ . We equip this rectangular






**Figure 10** The rectangular tracks shown inside the annulus  $R_\gamma$



**Figure 11** A rectangular track  $T_i$  along with arcs of  $\alpha_i$  in it



**Figure 12**  $A_i$  shown inside  $R_\gamma$  in the two possible ways : the figure on the left shows  $\alpha_i$  oriented from top to bottom; the figure on the right shows  $\alpha_i$  oriented from bottom to top

tracks  $T_{i_0}$  with the Euclidean metric so that the boundary arcs  $\rho_{i_0}$ ,  $\rho_{i_0+1}$ , and the arcs of  $T_{i_0} \cap \partial R_\gamma$  are all straight lines and so that  $T_{i_0}$  is a rectangle. We refer to  $T_{i_0} \cap \partial_+ R_\gamma$  as the left end of the rectangle and  $T_{i_0} \cap \partial_- R_\gamma$  as the right end of this rectangular track. We can draw the arcs of  $\alpha_j$ , for  $j \in K$ , as straight line segments in the rectangular tracks  $T_{i_0}$ . Figure 11 shows a schematic of  $T_i$  where  $i \in K$ .

From this schematic, at both the left and right end of this rectangular track  $T_i$ ,  $\alpha_i$  is a common boundary to a triangle and a pentagon. We call  $\alpha_i$  as the *starting arc* of this rectangular track  $T_i$ .

Figure 12 shows two possible schematics when  $A_i$  is pictured in  $R_\gamma$ .

For any of the two possible cases observed in figure 12, a portion of one of the two pentagons of  $T_i$  appears in the  $A_i$  which is between  $\alpha_i$  and  $\partial_+ R_\alpha$ , where  $\alpha_i$  is the starting arc of this track. We call this pentagon *the upper pentagon of the*

rectangular track  $T_i$ , owing to the viewpoint that  $\partial_+ R_\alpha$  is the upper boundary of  $R_\alpha$ . A portion of the other pentagon of  $T_i$  appears in  $A_i$  which is between  $\alpha_i$  and  $\partial_- R_\alpha$ . We call this pentagon *the lower pentagon of the rectangular track*. Likewise, we define the *upper triangle* and the *lower triangle* of a rectangular track  $T_i$ .

Let  $\bar{\gamma} \in \Gamma$  as in the statement of the Theorem 1. We prove that  $d(\bar{\gamma}, \alpha) \geq 3$  by showing that  $\bar{\gamma}$  and  $\alpha$  fill  $S$ . By Theorem 1, this will imply that  $d(T_\gamma(\alpha), \alpha) \geq 4$ .

It can be observed that  $i(\bar{\gamma} \cap \alpha) \neq 0$  because if  $\bar{\gamma}$  is disjoint from both  $\alpha$  and  $T_\gamma(\alpha)$  then  $\bar{\gamma}$  is non-essential as it will lie completely in one of the disc components of  $S \setminus (T_\gamma(\alpha) \cup \alpha)$ . Since  $d(\bar{\gamma}, \gamma) \geq d(T_\gamma(\alpha), \gamma) - d(\bar{\gamma}, T_\gamma(\alpha)) = d(T_\gamma(\alpha), T_\gamma(\gamma)) - d(\bar{\gamma}, T_\gamma(\alpha)) = 3 - 1 = 2$ , we also conclude that  $i(\bar{\gamma} \cap \gamma) \neq 0$ . Since  $\bar{\gamma}$  intersects  $\gamma$ , it intersects  $R_\gamma$ . It cannot be completely contained in  $R_\gamma$  because every simple closed curve contained in an annulus bounds a disk or is isotopic to the core curve of the annulus. Since neither of these is true, it follows that  $\bar{\gamma}$  intersects  $R_\gamma$  in arcs. Since  $i(\bar{\gamma}, T_\gamma(\alpha)) = \phi$ , each component of  $\bar{\gamma} \cap R_\gamma$  has to be completely contained in one of the rectangular tracks described by  $T_\gamma(\alpha)$ . Such a component arc of  $\bar{\gamma}$  could either be boundary reducible or essential in  $R_\gamma$ .


We consider an isotopy  $I_1$  of  $\bar{\gamma}$ , as follows: In the case that a component arc of  $\bar{\gamma}$  in  $R_\gamma$  is boundary reducible in  $R_\gamma$ , we can perform the boundary reduction of  $\bar{\gamma}$  preserving its minimal intersection position with  $\alpha$  and  $T_\gamma(\alpha)$ . This is possible because an arc of  $\bar{\gamma}$  which is boundary reducible in  $R_\gamma$  and is contained in the disk  $T_i$  will bound a bigon with one boundary arc of  $R_\gamma$  in  $T_i$ . Also, since  $\bar{\gamma}$  was already in minimal intersection position with  $\alpha$ , it does not bound bigons with the arcs  $\alpha_j$  inside  $T_i$ . Call the isotopy of  $\bar{\gamma}$  which reduces all the boundary-reducible arcs of  $\bar{\gamma} \cap R_\gamma$  as  $I_1$ . After the isotopy  $I_1$ , we can assume that all the arcs of  $\bar{\gamma}$  in  $R_\gamma$  are essential. We know that there is at-least one component of  $\bar{\gamma} \cap R_\gamma$  which is an essential arc of  $R_\gamma$  as  $\bar{\gamma}$  cannot be disjoint from  $R_\gamma$ . By the hypothesis that  $i(\bar{\gamma}, b) \leq 1$  for  $b \subset \alpha \setminus T_\gamma(\alpha)$  each rectangular track can contain at-most one component of  $\bar{\gamma} \cap R_\gamma$ .

Next, we describe an isotopy  $I_2$  of  $\bar{\gamma}$  such that all the points of  $\bar{\gamma} \cap \alpha$  will lie inside  $R_\gamma$  and so that no new boundary reducible arc components of  $\bar{\gamma} \cap R_\gamma$  are introduced and  $\bar{\gamma}$ 's minimal intersection position with  $\alpha$  and  $T_\gamma(\alpha)$  is retained. To this end, suppose that a point of  $\bar{\gamma} \cap \alpha$  lies outside  $R_\gamma$ .


Following the construction of the disk  $D'_{i,1}$  described above using figure 9, we see that the upper pentagon of the rectangular track  $T_i$  is connected to the upper triangle of the rectangular track  $T_j$  via a conduit  $R_{i,j}$  where  $i, j \in K$  are such that  $A_j$  is to the left of  $A_i$  in  $R_\alpha$  as in schematic 9.

If a point of  $\bar{\gamma} \cap \alpha$ ,  $x_0$ , lies outside  $R_\gamma$ , then it has to lie on  $\omega_{i,j}$  for some  $i$  and  $j$  such that  $i, j \in K, i \neq j$ . We now refer to the dotted line in figure 13. Since the intersection of  $\bar{\gamma}$  and  $\alpha$  is transverse, an arc of  $\bar{\gamma}$ , call it  $\delta$  lies on the two sides of the conduit  $R_{i,j}$ , one inside and one outside  $R_{i,j}$ . The endpoint  $P$  of the arc  $\delta$  inside  $R_{i,j}$  is also the endpoint of some other arc of  $\bar{\gamma}$  as  $\bar{\gamma}$  is a closed curve. If  $P$  connects to an arc of  $\bar{\gamma}$  lying in the upper triangular region of the track  $T_j$ , then an essential arc  $\delta_1$  of  $\bar{\gamma} \cap R_\gamma$  lies in  $T_j$  with its endpoint  $Q$  on  $\partial R_\gamma$  in the upper triangle of  $T_j$  so that  $\delta$ , the arc  $PQ$  and  $\delta_1$  together form a bigon with  $\alpha$  contradicting the minimal intersection position of  $\bar{\gamma}$  with  $\alpha$ . So,  $P$  connects to an arc of  $\bar{\gamma}$  in the upper pentagon in the track  $T_i$  as is the dotted line in figure 13. Consider an isotopy  $I_2$  which slides the point  $x_0$  onto  $\alpha_i$ . The image of the arc component of  $\bar{\gamma} \cap R_\gamma$  which is in  $T_i$ , under  $I_2$  has its endpoint in the lower triangle of  $T_i$  and the image of  $x_0$  lies in  $R_\gamma$ . A schematic for this isotopy  $I_2$  is shown in figure 13.

After finitely many such isotopies, we can now assume that all the points of  $\bar{\gamma} \cap \alpha$  lie inside  $R_\gamma$ . Now consider an isotopy  $I_3$  of  $\bar{\gamma}$  as follows: If any of the



**Figure 13** The isotopy  $I_2$  moving points of  $\bar{\gamma} \cap \alpha$  into  $R_\gamma$



**Figure 14** A schematic showing the normalization move, the isotopy  $I_3$


components of  $\bar{\gamma} \cap R_\gamma$  has its endpoint on the boundary of the upper triangle of  $T_j$ , for some  $j \in K$ , then by the above discussion,  $\bar{\gamma}$  cannot intersect  $\omega_{i,j}$  or  $\eta_{i,j}$ , for some  $i \in K$  such that the arcs of  $T_i$  and  $T_j$  forms the opposite sides of a conduit  $R_{i,j}$ . So  $\bar{\gamma} \cap R_{i,j}$  is an arc  $MN$  which has its endpoints  $M \in T_j$  and  $N \in T_i$  on  $\partial R_\gamma$ . Further, since  $\bar{\gamma}$  is a closed curve,  $\bar{\gamma} \cap T_i$  is an arc with its endpoint as  $N$  such that  $N$  necessarily lies in the upper pentagon of  $T_i$ . Conversely, if any of the components of  $\bar{\gamma} \cap R_\gamma$  has its endpoint,  $z_0$ , on the boundary of the upper pentagon of  $T_i$ , then it should be connected to an arc,  $g$ , of  $\bar{\gamma}$  in the conduit  $R_{i,j}$ . Note that the endpoints,  $z_0, z'_0$  of  $g$  are on  $\partial R_\gamma$ . There exists an arc component of  $\bar{\gamma} \cap R_\gamma$  lying in  $T_j$  such that  $z'_0$  is on the boundary of the upper triangle of  $T_j$ , as the dotted line in figure 14 shows. If any such arc  $g$  of  $\bar{\gamma}$  exists, consider an isotopy,  $I_3$ , of  $g$  such that the image,  $I_3(g)$ , lies outside  $R_{i,j}$ . A schematic of this is figure 14.

The component of  $\bar{\gamma} \cap R_\gamma$  in  $T_j$  now has an endpoint on the boundary of the lower pentagon of  $T_j$  and the component of  $\bar{\gamma} \cap R_\gamma$  in  $T_i$  has an endpoint on the boundary of the lower triangle of  $T_i$ . Also the image of  $\bar{\gamma} \cap \alpha$  under  $I_3$  moves a point of  $\bar{\gamma} \cap \alpha$  from the boundary of the upper triangle of  $T_j$  to the boundary of the lower pentagon of  $T_i$ . We call  $I_3$  to be a *normalization* move on  $\bar{\gamma}$ . After finitely many normalization moves performed on  $\bar{\gamma}$ , wherever applicable, we can assume that every component of  $\bar{\gamma} \cap R_\gamma$  is contained in a rectangular track  $T_i$  for some  $i \in K$  such that the endpoints of that component lie on the boundary of the lower triangle and the lower pentagon of  $T_i$ . So a schematic of every component of  $\bar{\gamma} \cap R_\gamma$  inside  $T_i$  is as in figure 15.


After these isotopies  $I_1, I_2, I_3$  of  $\bar{\gamma}$ , we say that  $\bar{\gamma}$  is in a *rectified position*. We now prove that  $\bar{\gamma}$  in rectified position and  $\alpha$  fill  $S$ . From now on we assume that  $\bar{\gamma}$  is in a rectified position.

For  $i \in K$ , let  $H_i$  be the rectangular component of  $R_\gamma \setminus (\cup_{i \in K} \alpha_i)$  containing the arcs  $a_i$  and  $a_{i+1}$  on its boundary. Each of these  $H_i$  contains a unique segment,  $\gamma_i$ , of the core curve  $\gamma$ . The schematic 16 shows  $H_1$  and  $\gamma_1$  for instance.

We say that an arc,  $g$  of  $\bar{\gamma}$  *covers*  $\gamma_i$  if  $g \subset H_i$  has its end points on  $\alpha_i$  and  $\alpha_{i+1}$  and  $g$  is isotopic in  $H_i$  to  $\gamma_i$  through arcs whose end points stay on  $\alpha_i, \alpha_{i+1}$ .



**Figure 15** The portion of  $\bar{\gamma}$  in rectified position inside  $T_i$




**Figure 16** Schematic showing  $H_1$  and  $\gamma_1$  in  $R_\gamma$

Since  $\gamma$  and  $\alpha$  form a filling pair, the set of essential arcs,  $\{\gamma_1, \dots, \gamma_k\}$  fill  $S \setminus \alpha$ . It follows that  $\bar{\gamma}$  fills  $S$  along with  $\alpha$  if segments of  $\bar{\gamma} \setminus \alpha$  cover  $\gamma_i$  for all  $i$  with  $i \in K$ .


Since  $\bar{\gamma}$  is in rectified position, each component of  $\bar{\gamma} \cap R_\gamma$  already covers all  $\gamma_i$  except one as in figure 15. More precisely, if a component of  $\bar{\gamma} \cap R_\gamma$  is in a rectangular track  $T_i$ , then  $\bar{\gamma}$  covers every  $\gamma_j$  where  $j$  is such that  $1 \leq j \leq k$  and  $j \neq i - 1$ . So, if  $\bar{\gamma} \cap R_\gamma$  has two distinct components, then each component has to lie in  $T_i$  for distinct  $i$  and hence  $\bar{\gamma}$  covers  $\gamma_j$  for  $j \in \{1, 2, \dots, k\}$ . We conclude that  $\bar{\gamma}$  and  $\alpha$  fill  $S$  in this case. Now it remains to show that if there is a single component of  $\bar{\gamma} \cap R_\gamma$ , which is an essential arc of  $R_\gamma$  and is contained in some rectangular track  $T_i$ , then  $\bar{\gamma}$  and  $\alpha$  fill  $S$ . As in the previous case,  $\bar{\gamma}$  covers every  $\gamma_j$  where  $j$  is such that  $1 \leq j \leq k$  and  $j \neq i - 1$ . The components of  $S \setminus (\alpha \cup_{1 \leq j \leq k, j \neq i-1} \gamma_j)$  will be disks except possibly one which could be a cylinder. This can be seen as follows. Since  $\alpha$  and  $\gamma$  fill  $S$ , the components of  $S \setminus \alpha \cup_{1 \leq j \leq k} \gamma_j$  are disks. Each segment of  $\gamma_j \setminus \alpha$  for  $j \in \{1, 2, \dots, k\}$  contributes to two distinct edges of a component  $J_0$  or two separate components  $J, J'$  of  $S \setminus \alpha \cup_{1 \leq j \leq k} \gamma_j$ .

Let  $P_1 := \bar{\gamma} \cap \alpha_i$  and  $P_2 := \bar{\gamma} \cap \alpha_{i-1}$  be points in  $T_i$  which appear on the unique component of  $\bar{\gamma} \cap R_\gamma$ . Let  $[P_1, P_2]$  represent the arc of  $\bar{\gamma}$  in  $R_\gamma$  with endpoints  $P_1$  and  $P_2$  and  $\bar{\gamma}_1 := \bar{\gamma} \setminus [P_1, P_2]$ .  $\bar{\gamma}_1$  is contained in all the components of  $S \setminus \alpha \cup_{1 \leq j \leq k, j \neq i-1} \gamma_j$  which contain the arcs  $\alpha_{i-1}$  and  $\alpha_i$  on their boundary. We know that there is at-least one such component because  $\gamma_{i-1}$  is also such an arc which joins  $\alpha_{i-1}$  to  $\alpha_i$ . If  $\gamma_{i-1}$  is the boundary of  $J, J'$ , then it would have been an arc which connected  $\alpha_{i-1}$  on one disk to  $\alpha_i$  on another disk. Note that both  $\alpha_i$  and  $\alpha_{i-1}$  are also boundary arcs of both  $J$  and  $J'$ . So, we would find  $P_1$  on the disk containing  $\alpha_i$  and  $P_2$  on the disk containing  $\alpha_{i-1}$ . When we join  $J$  and  $J'$  along  $\gamma_{i-1}$  we get a disk where  $\bar{\gamma}_1$  is an arc from  $P_1$  to  $P_2$  intersecting  $\gamma_{i-1}$ . Cutting along  $\bar{\gamma}_1$  still yields two different disks. The schematic, figure 17 shows this situation.

If  $\gamma_{i-1}$  were on the boundary of  $J_0$  representing two edges of  $J_0$  then it would have been an arc which connected  $\alpha_{i-1}$  to  $\alpha_i$ . When we glue  $J_0$  to itself along



**Figure 17** The figure on the left shows disks  $J$  and  $J'$  formed by cutting along  $\gamma_{i-1}$ ; the figure on the right shows the new disks formed when  $J \cup J'$  are cut along  $\bar{\gamma}_1$



**Figure 18** The disk  $J_0$  glued to itself along  $\gamma_{i-1}$  and cut along  $\bar{\gamma}_1$

$\gamma_{i-1}$ , we get a cylinder,  $A$ , where  $\alpha_i$  and  $\alpha_{i-1}$  will be arcs on different boundary components of  $A$ . So we would find  $P_1$  and  $P_2$  on distinct boundaries of  $A$  and hence  $\bar{\gamma}_1$  would be an essential arc on  $A$ . So cutting  $A$  along this arc  $\bar{\gamma}_1$  would yield a disk as shown in the schematic, figure 18.

In any case, we get disks by cutting  $S \setminus \alpha$  along the arcs of  $\bar{\gamma} \setminus \alpha$ . So this proves the theorem.  $\square$

Claim 1 of Theorem 2 doesn't use the hypothesis that  $d(\alpha, \gamma) = 3$ . So we have the following:

**COROLLARY 1.** *If  $\alpha$  and  $\gamma$  are a pair of curves which fill  $S$ , then  $\alpha$  and  $T_\gamma(\alpha)$  also fill  $S$ .*


**COROLLARY 2.** *For a surface of genus  $g \geq 3$ ,  $i_{\min}(g, 4) \leq (2g - 1)^2$ .*

**PROOF.** Aougab and Huang [3] proved that  $i_{\min}(g, 3) = 2g - 1$  for  $g \geq 3$ . Now, on  $S_g$ , for  $g \geq 3$ , suppose that  $\alpha$  and  $\beta$  are two such minimally intersecting curves with  $d(\alpha, \beta) = 3$ . Then  $i(\alpha, T_\beta(\alpha)) = (2g - 1)^2$  and by Theorem 2,  $d(\alpha, T_\beta(\alpha)) = 4$ . So  $i_{\min}(g, 4) \leq (2g - 1)^2$ .  $\square$

#### 4. An initially efficient geodesic

**LEMMA 1.** *If  $\alpha = \nu_0, \nu_1, \nu_2, \nu_3 = \gamma$  is an initially efficient geodesic then so is  $T_\gamma(\alpha), T_\gamma(\nu_1), \nu_2, \nu_1, \alpha$ .*

**PROOF.** For  $p \in K_{2-2g}$ , let  $F_p''$  be the components of  $S \setminus \{\alpha, R_\gamma\}$  as in the proof of Theorem 2. Since the geodesic  $\alpha, \nu_1, \nu_2, \gamma$  is an initially efficient one, each segment of  $\nu_1$  intersects every reference arc in  $E_i$  at most twice. In particular, arcs of  $\partial(R_\gamma)$  that form the edges of  $E_i$  intersect  $\nu_1$  at most twice. It follows from here that there are at the most two segments of  $\nu_1$  in each rectangular track  $T_i$  as




**Figure 19** There can be at-most two distinct segments of  $T_\gamma(\nu_1)$  in any rectangular component of  $S \setminus (\alpha \cup T_\gamma(\alpha))$  in  $R_\gamma$

defined in section 3. A schematic of this is shown in figure 19. Further, since the interior of a reference arc is disjoint from  $\alpha \cup T_\gamma(\alpha)$ , it is sufficient to check for the initial efficiency of the geodesic,  $T_\gamma(\alpha), T_\gamma(\nu_1), \nu_2, \nu_1, \alpha$  in the modified disks  $F_p'''$ , abbreviated  $F$ , corresponding to  $F_p$ , abbreviated  $E$ .

Since  $E$  and  $F$  are homeomorphic to a  $2g$ -gon. Without loss of generality assume  $E$  and  $F$  to be a regular Euclidean regular polygon with  $2g$  sides. Starting at any segment of  $\alpha$  in  $E$ , we label the edge as  $\alpha_1$ . Label the edges of  $E$  in a clockwise direction, starting at  $\alpha_1$  as  $\gamma_1, \alpha_2, \gamma_2, \dots, \gamma_g$ . Let  $S' = S \setminus R_\gamma$ . Since the components of  $S' \setminus \{\alpha, \gamma\}$  and  $S' \setminus \{\alpha, T_\gamma(\alpha)\}$  are the same, it follows that for every edge,  $a_{j_0}$  in  $F$  corresponding to  $\alpha$ , there exists a unique  $i_0 \in \{1, \dots, g\}$  such that  $a_{j_0} \subset \alpha_{i_0}$ . Index the edges,  $a_{j_0}$  of  $F$  such that  $j_0 = i_0$ . Label the edge of  $T_\gamma(\alpha)$  in  $F$  between  $a_i$  and  $a_{i+1}$  as  $t_i$ . Let  $\omega$  be a reference arc in  $F$  with end points on  $t_p$  and  $t_q$  for some  $p, q \in \{1, \dots, g\}$ . Suppose to the contrary that  $\omega \cap T_\gamma(\nu_1) \geq 3$ . Then there exists three segments,  $z_1, z_2, z_3$  of  $T_\gamma(\nu_1)$  in  $F$  such that  $z_j \cap \omega \neq \emptyset$ . For  $j \in \{1, 2, 3\}$ , let the end points of  $z_j$  lie on  $a_{j_1}$  and  $a_{j_2}$ . From our previous discussion on Dehn twist and figure 20, there exists arcs of  $\nu_1$  in  $E$  with end points on  $\gamma_{j_1}$  and  $\gamma_{j_2}$  for all  $j \in \{1, 2, 3\}$ . Consider a line segment,  $\omega'$  in  $E$  from an interior point of  $a_p$  to an interior point of  $a_q$ . Then  $\omega'$  is a reference arc for the triple,  $\alpha, \nu_1, \gamma$  and  $\omega' \cap \nu_1 \geq 3$ . This contradicts that  $\alpha, \nu_1, \nu_2, \gamma$  is an initially efficient geodesic. Hence,  $\omega \cap T_\gamma(\nu_1) \leq 2$  for any choice of reference arc,  $\omega$  for the triple  $T_\gamma(\alpha), T_\gamma(\nu_1), \alpha$ .

Since  $T_\gamma(\alpha), T_\gamma(\nu_1), \nu_2, \nu_1, \alpha$  is already a geodesic we have that  $d(T_\gamma(\nu_1), \alpha) = 3$ . This gives that  $T_\gamma(\nu_1)$  is an initially efficient geodesic of distance 4 from  $T_\gamma(\alpha)$  to  $\alpha$ .

□



**Figure 20** Initial efficiency of  $T_\gamma(a_1)$  follows from the initial efficiency of  $a_1$

## References

- [1] Aougab, Tarik . Uniform hyperbolicity of the graphs of curves. *Geom. Topol.* 17 (2013), no. 5, 2855–2875.
- [2] Aougab, Tarik ; Taylor, Samuel J. Small intersection numbers in the curve graph. *Bull. Lond. Math. Soc.* 46 (2014), no. 5, 989–1002.
- [3] Aougab, Tarik ; Huang, Shinniyih . Minimally intersecting filling pairs on surfaces. *Algebr. Geom. Topol.* 15 (2015), no. 2, 903–932.
- [4] Birman, Joan ; Margalit, Dan ; Menasco, William . Efficient geodesics and an effective algorithm for distance in the complex of curves. *Math. Ann.* 366 (2016), no. 3-4, 1253–1279.
- [5] Bowditch, Brian H. Uniform hyperbolicity of the curve graphs. *Pacific J. Math.* 269 (2014), no. 2, 269–280.
- [6] Clay, Matt ; Rafi, Kasra ; Schleimer, Saul . Uniform hyperbolicity of the curve graph via surgery sequences. *Algebr. Geom. Topol.* 14 (2014), no. 6, 3325–3344.
- [7] Farb, Benson ; Margalit, Dan . A primer on mapping class groups. Princeton Mathematical Series, 49. Princeton University Press, Princeton, NJ, 2012. xiv+472 pp. ISBN: 978-0-691-14794-9
- [8] Glenn, Paul ; Menasco, William W. ; Morrell, Kayla ; Morse, Matthew J. MICC: a tool for computing short distances in the curve complex. *J. Symbolic Comput.* 78 (2017), 115–132.
- [9] Harvey, W. J. Boundary structure of the modular group. *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, pp. 245–251, *Ann. of Math. Stud.*, 97, Princeton Univ. Press, Princeton, N.J., 1981.
- [10] Hensel, Sebastian ; Przytycki, Piotr ; Webb, Richard C. H. 1-slim triangles and uniform hyperbolicity for arc graphs and curve graphs. *J. Eur. Math. Soc. (JEMS)* 17 (2015), no. 4, 755–762.
- [11] Leasure, Jason Paige . Geodesics in the complex of curves of a surface. Thesis (Ph.D.)—The University of Texas at Austin. ProQuest LLC, Ann Arbor, MI, 2002. 56 pp. ISBN: 978-0496-62255-9
- [12] Masur, H. A. ; Minsky, Y. N. Geometry of the complex of curves. II. Hierarchical structure. *Geom. Funct. Anal.* 10 (2000), no. 4, 902–974.
- [13] Minsky, Yair N. Curve complexes, surfaces and 3-manifolds. *International Congress of Mathematicians. Vol. II*, 1001–1033, *Eur. Math. Soc.*, Zürich, 2006.
- [14] Przytycki, Piotr ; Sisto, Alessandro . A note on acylindrical hyperbolicity of mapping class groups. *Hyperbolic geometry and geometric group theory*, 255–264, *Adv. Stud. Pure Math.*, 73, Math. Soc. Japan, Tokyo, 2017.
- [15] Schleimer, Saul. Notes on the Complex of Curves. <http://homepages.warwick.ac.uk/~masgar/Maths/notes.pdf>, 2006

- [16] Shackleton, Kenneth J. Tightness and computing distances in the curve complex. *Geom. Dedicata* 160 (2012), 243–259.
- [17] Watanabe, Yohsuke . Intersection numbers in the curve graph with a uniform constant. *Topology Appl.* 204 (2016), 157–167.
- [18] Webb, Richard C. H. Combinatorics of tight geodesics and stable lengths. *Trans. Amer. Math. Soc.* 367 (2015), no. 10, 7323–7342.

KUWARI MAHANTA, Department of Mathematics, Indian Institute of Technology Guwahati, Assam 781039, India, email : kuwari.mahanta@iitg.ac.in

SREEKRISHNA PALAPARTHI (\*corresponding author), Department of Mathematics, Indian Institute of Technology Guwahati, Assam 781039, India, email : passkrishna@iitg.ac.in