

Optimal Clustering in Anisotropic Gaussian Mixture Models

Xin Chen¹ and Anderson Y. Zhang²

¹ *University of Washington*

² *University of Pennsylvania*

Abstract

We study the clustering task under anisotropic Gaussian Mixture Models where the covariance matrices from different clusters are unknown and are not necessarily the identical matrix. We characterize the dependence of signal-to-noise ratios on the cluster centers and covariance matrices and obtain the minimax lower bound for the clustering problem. In addition, we propose a computationally feasible procedure and prove it achieves the optimal rate within a few iterations. The proposed procedure is a hard EM type algorithm, and it can also be seen as a variant of the Lloyd's algorithm that is adjusted to the anisotropic covariance matrices.

1 Introduction

Clustering is a fundamentally important task in statistics and machine learning [7, 2]. The most popular and studied model for clustering is the Gaussian Mixture Model (GMM) [18, 20] which can be written as

$$Y_j = \theta_{z_j^*}^* + \epsilon_j, \text{ where } \epsilon_j \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \Sigma_{z_j^*}^*), \forall j \in [n].$$

Here $Y = (Y_1, \dots, Y_n)$ are the observations with n being the sample size. Let k be the number of clusters that is assumed to be known. Denote $\{\theta_a^*\}_{a \in [k]}$ to be unknown centers and $\{\Sigma_a^*\}$ to be unknown covariance matrices for the k clusters. Let $z^* \in [k]^n$ be the cluster structure such that for each index $j \in [n]$, the value of z_j^* indicates which cluster the j th data point belongs to. The goal is to recover z^* from Y . For any estimator \hat{z} , its clustering performance is measured by a misclustering error rate $h(\hat{z}, z^*)$ which will be introduced later in (4).

There have been increasing interests in the theoretical and algorithmic analysis of clustering under GMMs. When the GMM is isotropic (that is, all the covariance matrices $\{\Sigma_a^*\}_{a \in [k]}$ are equal to the same identity matrix), [15] obtains the minimax rate for clustering which takes a form of $\exp(-(1 + o(1))(\min_{a \neq b} \|\theta_a^* - \theta_b^*\|)^2/8)$ under the loss $h(\hat{z}, z^*)$. Various methods have been studied in the isotropic setting. k -means clustering [16] might be the most natural choice but it is NP-hard [4]. As a local approach to optimize the k -mean objects, Lloyd's algorithm [13] is one of the most popular clustering algorithms and has achieved many successes in different disciplines [24]. [15, 8] establishes computational and statistical guarantees for the Lloyd's algorithm by showing it achieves the optimal rates after a few iterations provided with some decent initialization. Another popular approach to clustering especially for high-dimensional data is spectral clustering [22, 19, 21], which is an umbrella term for clustering after a dimension reduction through a spectral decomposition. [14, 17, 1] proves the spectral clustering also achieves the optimality under the isotropic GMM. Another line of work is to consider semidefinite programming (SDP) as a convex relaxation of the k -means objective. Its statistical properties have been studied in [6, 9].

In spite of all the exciting results, most of the existing literature focuses on isotropic GMMs, and clustering under the anisotropic case where the covariance matrices are not necessarily the identity matrix is not well-understood. The results of some papers [15, 6] hold under sub-Gaussian mixture models, where the errors ϵ_j are assumed to follow some sub-Gaussian distribution with variance proxy σ^2 . It seems that their result already covers the anisotropic case, as $\{\mathcal{N}(0, \Sigma_a^*)\}_{a \in [k]}$ are indeed sub-Gaussian distributions. However, from a minimax point of view, among all the sub-Gaussian distributions with variance proxy σ^2 , the least favorable case (the case where clustering is the most difficult) is when the errors are $\mathcal{N}(0, \sigma^2)$. Therefore, the minimax rates for clustering under the sub-Gaussian mixture model is essentially the one under isotropic GMMs, and methods such as the Lloyd’s algorithm that requires no covariance matrix information can be rate-optimal. As a result, the aforementioned results are all for isotropic GMMs.

A few papers have explored the direction of clustering under anisotropic GMMs. [3] gives a polynomial-time clustering algorithm that provably works well when the Gaussian distributions are well separated by hyperplanes. Their idea is further developed in [11] which allows the Gaussians to be overlapped with each other but only for two-cluster cases. A recent paper [23] proposes another method for clustering under a balanced mixture of two elliptical distributions. They give a provable upper bound of their clustering performance with respect to an excess risk. Nevertheless, it remains unknown what is the fundamental limit of clustering under the anisotropic GMMs and whether there is any polynomial-time procedure that achieves it.

In this paper, we will investigate the optimal rates of the clustering task under two anisotropic GMMs. Model 1 is when the covariance matrices are all equal to each other (i.e., homogeneous) and are equal to some unknown matrix Σ^* . Model 2 is more flexible, where the covariance matrices are unknown and are not necessarily equal to each other (i.e., heterogeneous). The contribution of this paper is two-fold, summarized as follows.

Our first contribution is on the fundamental limits. We obtain the minimax lower bound for clustering under the anisotropic GMMs with respect to the loss $h(\hat{z}, z^*)$. We show it takes the form

$$\inf_{\hat{z}} \sup_{z^* \in [k]^n} \mathbb{E}h(z, z^*) \geq \exp\left(-\left(1 + o(1)\right) \frac{(\text{signal-to-noise ratio})^2}{8}\right),$$

where the signal-to-noise ratio under Model 1 is equal to $\min_{a, b \in [k]: a \neq b} \|(\theta_a^* - \theta_b^*)^T \Sigma^{*-1/2}\|$ and the one for Model 2 is more complicated. For both models, we can see the minimax rates depend not only on the centers but also the covariance matrices. This is different from the isotropic case whose signal-to-noise ratio is $\min_{a \neq b} \|\theta_a^* - \theta_b^*\|$. Our results precisely capture the role the covariance matrices play in the clustering problem. It shows covariance matrices impact the fundamental limits of the clustering problem through complicated interaction with the centers especially in Model 2. The minimax lower bounds are obtained by establishing connections with Linear Discriminant Analysis (LDA) and Quadratic Discriminant Analysis (QDA).

Our second and more important contribution is on the computational side. We propose a computationally feasible and rate-optimal algorithm for the anisotropic GMM. Popular methods including the Lloyd’s algorithm and the spectral clustering no longer work well as they are developed under the isotropic case and only consider the distances among the centers [3]. We study an *adjusted Lloyd’s algorithm* which estimates the covariance matrices in each iteration and recovers the clusters using the covariance matrix information. It can also be seen as a hard EM algorithm [5]. As an iterative algorithm, we give a statistical and computational guarantee and guidance to practitioners by showing that it obtains the minimax lower bound within $\log n$ iterations. That is, let $z^{(t)}$ be the

output of the algorithm after t iterations, we have with high probability,

$$h(z^{(t)}, z^*) \leq \exp\left(-\frac{(1+o(1))(\text{signal-to-noise ratio})^2}{8}\right),$$

holds for all $t \geq \log n$. The algorithm can be initialized by popular methods such as the spectral clustering or the Lloyd's algorithm. In numeric studies, we show the proposed algorithm improves greatly from the two aforementioned methods under anisotropic GMMs, and matches the optimal exponent given in the minimax lower bound.

Paper Organization. The remaining paper is organized as follows. In Section 2, we study Model 1 where the covariance matrices are unknown but homogeneous. In Section 3, we consider Model 2 where covariance matrices are unknown and heterogeneous. For both cases, we obtain the minimax lower bound for clustering and study the adjusted Lloyd's algorithm. In Section 4, we provide a numeric comparison with other popular methods. The proofs of theorems in Section 2 are given in Section 5 and the proofs for Section 3 are included in Section 6. All the technical lemmas are included in Section 7.

Notation. Let $[m] = \{1, 2, \dots, m\}$ for any positive integer m . For any set S , we denote $|S|$ for its cardinality. For any matrix $X \in \mathbb{R}^{d \times d}$, we denote $\lambda_1(X)$ to be its smallest eigenvalue and $\lambda_d(X)$ to be its largest eigenvalue. In addition, we denote $\|X\|$ to be its operator norm. For any two vectors u, v of the same dimension, we denote $\langle u, v \rangle = u^T v$ to be its inner product. For any positive integer d , we denote I_d to be the $d \times d$ identity matrix. We denote $\mathcal{N}(\mu, \Sigma)$ to be the normal distribution with mean μ and covariance matrix Σ . We denote $\mathbb{I}\{\cdot\}$ to be the indicator function. Given two positive sequences a_n, b_n , we denote $a_n = o(b_n)$ if $a_n/b_n = o(1)$ when $n \rightarrow \infty$. We write $a_n \lesssim b_n$ if there exists a constant $C > 0$ independent of n such that $a_n \leq C b_n$ for all n .

2 GMM with Unknown but Homogeneous Covariance Matrices

2.1 Model

We first consider a GMM where covariance matrices of different clusters are unknown but are assumed to be equal to each other. The data generating process can be displayed as follow:

Model 1:
$$Y_j = \theta_{z_j^*}^* + \epsilon_j, \text{ where } \epsilon_j \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \Sigma^*), \forall j \in [n]. \quad (1)$$

It is called Stretched Mixture Model in [23] as the density of Y_j is elliptical. Throughout the paper, we call it *Model 1* for simplicity and to distinguish it from a more complicated model that will be introduced in Section 3. The goal is to recover the underlying cluster assignment vector z^* from Y .

Signal-to-noise Ratio. Define the signal-to-noise ratio

$$\text{SNR} = \min_{a, b \in [k]: a \neq b} \|(\theta_a^* - \theta_b^*)^T \Sigma^{*-1/2}\|, \quad (2)$$

which is a function of all the centers $\{\theta_a^*\}_{a \in [k]}$ and the covariance matrix Σ^* . As we will show later in Theorem 2.1, SNR captures the difficulty of the clustering problem and determines the minimax rate. For the geometric interpretation of SNR, we defer it after presenting Theorem 2.2.

A quantity closely related to SNR is the minimum distance among the centers. Define Δ as

$$\Delta = \min_{a,b \in [k]: a \neq b} \|\theta_a^* - \theta_b^*\|. \quad (3)$$

Then we can see SNR and Δ are in the same order if all eigenvalues of the covariance matrix Σ^* is assumed to be constants. If Σ^* is further assumed to be an identical matrix, then we have SNR equal to Δ . As a result, in [15, 8, 14] where isotropic GMMs are studied, Δ plays the role of signal-to-noise ratio and appears in the minimax rates.

Loss Function. To measure the clustering performance, we consider the misclustering error rate defined as follows. For any $z, z^* \in [k]^n$, we define

$$h(z, z^*) = \min_{\psi \in \Psi} \frac{1}{n} \sum_{j=1}^n \mathbb{I}\{\psi(z_j) \neq z_j^*\}, \quad (4)$$

where $\Psi = \{\psi : \psi \text{ is a bijection from } [k] \text{ to } [k]\}$. Here the minimum is over all the permutations over $[k]$ due to the identifiability issue of the labels $1, 2, \dots, k$. Another loss that will be used is $\ell(z, z^*)$ defined as

$$\ell(z, z^*) = \sum_{j=1}^n \left\| \theta_{z_j}^* - \theta_{z_j^*}^* \right\|^2. \quad (5)$$

It also measures the clustering performance of z considering the distances among the true centers. It is related to $h(z, z^*)$ as $h(z, z^*) \leq \ell(z, z^*) / (n\Delta^2)$ and hence provides more information than $h(z, z^*)$. We will mainly use $\ell(z, z^*)$ in the technical analysis but will eventually present the results using $h(z, z^*)$ which is more interpretable.

2.2 Minimax Lower Bound

We first establish the minimax lower bound for the clustering problem under Model 1.

Theorem 2.1. *Under the assumption $\frac{\text{SNR}}{\sqrt{\log k}} \rightarrow \infty$, we have*

$$\inf_{\hat{z}} \sup_{z^* \in [k]^n} \mathbb{E}h(z, z^*) \geq \exp\left(-\frac{(1+o(1))\text{SNR}^2}{8}\right). \quad (6)$$

If $\text{SNR} = O(1)$ instead, we have $\inf_{\hat{z}} \sup_{z^ \in [k]^n} \mathbb{E}h(z, z^*) \geq c$ for some constant $c > 0$.*

Theorem 2.1 allows the cluster numbers k to grow with n and shows that $\text{SNR} \rightarrow \infty$ is a necessary condition to have a consistent clustering if k is a constant. Theorem 2.1 holds for any arbitrary $\{\theta_a^*\}_{a \in [k]}$ and Σ^* , and the minimax lower bound depend on them through SNR. The parameter space is only for z^* while $\{\theta_a^*\}_{a \in [k]}$ and Σ^* are fixed. Hence, (6) can be interpreted as a pointwise result, and it captures precisely the explicit dependence of the minimaxity on $\{\theta_a^*\}_{a \in [k]}$ and Σ^* .

Theorem 2.1 is closely related to the Linear Discriminant Analysis (LDA). If there are only two clusters, and if the centers and the covariance matrix are known, then estimating each z_j^* is exactly the task of LDA: we want to figure out which normal distribution an observation Y_j is generated from, where the two normal distributions have different means but the same covariance matrix. In fact, this is also how Theorem 2.1 is proved: we will first reduce the estimation problem of z^* into

a two-point hypothesis testing problem for each individual z_j^* , the error of which is given in Lemma 2.1 by the analysis of LDA, and then aggregate all the testing errors together.

In the following lemma, we give a sharp and explicit formula for the testing error of the LDA. Here we have two normal distributions $\mathcal{N}(\theta_1^*, \Sigma^*)$ and $\mathcal{N}(\theta_2^*, \Sigma^*)$ and an observation X that is generated from one of them. We are interested in estimating which distribution it is from. By Neyman-Pearson lemma, it is known that the likelihood ratio test $\mathbb{I} \{2(\theta_2^* - \theta_1^*)^T (\Sigma^*)^{-1} X \geq \theta_2^{*T} (\Sigma^*)^{-1} \theta_2^* - \theta_1^{*T} (\Sigma^*)^{-1} \theta_1^*\}$ is the optimal testing procedure. Then by using the Gaussian tail probability, we are able to obtain the optimal testing error, the lower bound of which is given in Lemma 2.1.

Lemma 2.1 (Testing Error for Linear Discriminant Analysis). *Consider two hypotheses $\mathbb{H}_0 : X \sim \mathcal{N}(\theta_1^*, \Sigma^*)$ and $\mathbb{H}_1 : X \sim \mathcal{N}(\theta_2^*, \Sigma^*)$. Define a testing procedure*

$$\phi = \mathbb{I} \{2(\theta_2^* - \theta_1^*)^T (\Sigma^*)^{-1} X \geq \theta_2^{*T} (\Sigma^*)^{-1} \theta_2^* - \theta_1^{*T} (\Sigma^*)^{-1} \theta_1^*\}.$$

Then we have $\inf_{\hat{\phi}} (\mathbb{P}_{\mathbb{H}_0}(\hat{\phi} = 1) + \mathbb{P}_{\mathbb{H}_1}(\hat{\phi} = 0)) = \mathbb{P}_{\mathbb{H}_0}(\phi = 1) + \mathbb{P}_{\mathbb{H}_1}(\phi = 0)$. If $\|(\theta_2^* - \theta_1^*)^T (\Sigma^*)^{-\frac{1}{2}}\| \rightarrow \infty$, we have

$$\inf_{\hat{\phi}} (\mathbb{P}_{\mathbb{H}_0}(\hat{\phi} = 1) + \mathbb{P}_{\mathbb{H}_1}(\hat{\phi} = 0)) \geq \exp \left(-(1 + o(1)) \frac{\|(\theta_2^* - \theta_1^*)^T (\Sigma^*)^{-\frac{1}{2}}\|^2}{8} \right).$$

Otherwise, $\inf_{\hat{\phi}} (\mathbb{P}_{\mathbb{H}_0}(\hat{\phi} = 1) + \mathbb{P}_{\mathbb{H}_1}(\hat{\phi} = 0)) \geq c$ for some constant $c > 0$.




Figure 1: A geometric interpretation of SNR.

With the help of Lemma 2.1, we have a geometric interpretation of SNR. In the left panel of Figure 1, we have two normal distributions $\mathcal{N}(\theta_1^*, \Sigma^*)$ and $\mathcal{N}(\theta_2^*, \Sigma^*)$ for X to be generated from. The black line represents the optimal testing procedure ϕ displayed in Lemma 2.1 that divides the space into two half-spaces. To calculate the testing error, we can make a transformation $X' = (\Sigma^*)^{-\frac{1}{2}}(X - \theta_1^*)$ so that the two normal distributions become isotropic: $\mathcal{N}(0, I_d)$ and $\mathcal{N}((\Sigma^*)^{-\frac{1}{2}}(\theta_2^* - \theta_1^*), I_d)$ as displayed in the right panel. Then the distance between the two centers are $\|(\Sigma^*)^{-\frac{1}{2}}(\theta_2^* - \theta_1^*)\|$, and the distance between a center and the black curve is half of it. Then $\mathbb{P}_{\mathbb{H}_0}(\hat{\phi} = 1)$ is the probability of $\mathcal{N}(0, I_d)$ in the grayed area, which is equal to $\exp(-(1 + o(1))\|(\Sigma^*)^{-\frac{1}{2}}(\theta_2^* - \theta_1^*)\|^2/8)$ by Gaussian tail probability. As a result, $\|(\Sigma^*)^{-\frac{1}{2}}(\theta_2^* - \theta_1^*)\|$ is the effective distance between the two centers of $\mathcal{N}(\theta_1^*, \Sigma^*)$ and $\mathcal{N}(\theta_2^*, \Sigma^*)$ for the clustering problem, considering the geometry of the covariance matrix. Since we have multiple clusters, SNR defined in (2) can be interpreted as the minimum effective distances among the centers $\{\theta_a^*\}_{a \in [k]}$ considering the anisotropic structure of Σ^* and it captures the intrinsic difficulty of the clustering problem.

2.3 Rate-Optimal Adaptive Procedure

In this section, we will propose a computationally feasible and rate-optimal procedure for clustering under Model 1. Summarized in Algorithm 1, the proposed algorithm is a variant of the Lloyd algorithm. Starting from some initialization, it updates the estimation of the centers $\{\theta_a^*\}_{a \in [k]}$ (in (7)), the covariance matrix Σ^* (in (8)), and the cluster assignment vector z^* (in (9)) iteratively. It differs from the Lloyd's algorithm in the sense that Lloyd's algorithm is for isotropic GMMs without the covariance matrix update (8). In addition, in (9) it updates the estimation of z_j^* by $\operatorname{argmin}_{a \in [k]} (Y_j - \theta_a^{(t)})^T (Y_j - \theta_a^{(t)})$ instead. To distinguish them from each other, we call the classical Lloyd's algorithm as the *vanilla Lloyd's algorithm*, and name Algorithm 1 as the *adjusted Lloyd's algorithm*, as it is adjusted to the unknown and anisotropic covariance matrix.

Algorithm 1 can also be interpreted as a hard EM algorithm. If we apply the Expectation Maximization (EM) for Model 1, we will have an M step for estimating parameters $\{\theta_a^*\}_{a \in [k]}$ and Σ^* and an E step for estimating z^* . It turns out the updates on the parameters (7) - (8) are exactly the same as the updates of EM (M step). However, the update on z^* in Algorithm 1 is different from that in the EM. Instead of taking a conditional expectation (E step), we also take a maximization in (9). As a result, Algorithm 1 consists solely of M steps for both the parameters and z^* , which is known as a hard EM algorithm.

Algorithm 1: Adjusted Lloyd's Algorithm for Model 1 (1).

Input: Data Y , number of clusters k , an initialization $z^{(0)}$, number of iterations T

Output: $z^{(T)}$

1 **for** $t = 1, \dots, T$ **do**

2 Update the centers:

$$\theta_a^{(t)} = \frac{\sum_{j \in [n]} Y_j \mathbb{I}\{z_j^{(t-1)} = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j^{(t-1)} = a\}}, \quad \forall a \in [k]. \quad (7)$$

3 Update the covariance matrix:

$$\Sigma^{(t)} = \frac{\sum_{a \in [k]} \sum_{j \in [n]} (Y_j - \theta_a^{(t)})(Y_j - \theta_a^{(t)})^T \mathbb{I}\{z_j^{(t-1)} = a\}}{n}. \quad (8)$$

4 Update the cluster estimations:

$$z_j^{(t)} = \operatorname{argmin}_{a \in [k]} (Y_j - \theta_a^{(t)})^T (\Sigma^{(t)})^{-1} (Y_j - \theta_a^{(t)}), \quad j \in [n]. \quad (9)$$

In Theorem 2.2, we give a computational and statistical guarantee of the proposed Algorithm 1. We show that starting from a decent initialization, within $\log n$ iterations, Algorithm 1 achieves an error rate $\exp(-(1 + o(1))\text{SNR}^2/8)$ which matches with the minimax lower bound given in Theorem 2.1. As a result, Algorithm 1 is a rate-optimal procedure. In addition, the algorithm is fully adaptive to the unknown $\{\theta_a^*\}_{a \in [k]}$ and Σ^* . The only information assumed to be known is k the number of clusters, which is commonly assumed to be known in clustering literature [15, 8, 14]. The theorem also shows that the number of iterations to achieve the optimal rate is at most $\log n$,

which provides implementation guidance to practitioners.

Theorem 2.2. *Assume $kd = O(\sqrt{n})$ and $\min_{a \in k} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ for some constant $\alpha > 0$. Assume $\frac{\text{SNR}}{k} \rightarrow \infty$ and $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$. For Algorithm 1, suppose $z^{(0)}$ satisfies $\ell(z^{(0)}, z^*) = o(n/k)$ with probability at least $1 - \eta$. Then with probability at least $1 - \eta - n^{-1} - \exp(-\text{SNR})$, we have*

$$h(z^{(t)}, z^*) \leq \exp\left(-\left(1 + o(1)\right)\frac{\text{SNR}^2}{8}\right), \quad \text{for all } t \geq \log n.$$

We have remarks on the assumptions of Theorem 2.2. We allow the number of clusters k to grow with n . When k is a constant, the assumption on $\text{SNR} \rightarrow \infty$ is the necessary condition to have a consistent recovery of z^* according to the minimax lower bound presented in Theorem 2.1. The assumption on Σ^* is to make sure the covariance matrix is well-conditioned. The dimensionality d is assumed to be at most $O(\sqrt{n})$, an assumption that is stronger than that in [15, 8, 14] which only needs $d = O(n)$. This is due to that compared to these papers, we need to estimate the covariance matrix Σ^* and to have a control on the estimation error $\|\Sigma^{(t)} - \Sigma^*\|$.

The requirement for the initialization $\ell(z^{(0)}, z^*) = o(n/k)$ can be fulfilled by simple procedures. A popular choice is the vanilla Lloyd's algorithm the performance of which is studied in [15, 8]. Since ϵ_j are sub-Gaussian random variables with proxy variance λ_{\max} , [8] implies the vanilla Lloyd's algorithm output \hat{z} satisfies $\ell(\hat{z}, z^*) \leq n \exp(-\left(1 + o(1)\right)\Delta^2/(8\lambda_{\max}))$ with probability at least $1 - \exp(-\Delta) - n^{-1}$, under the assumption that $\text{SNR}/k \rightarrow \infty$. Note that [8] is for isotropic GMMs, but its results can be extended to sub-Gaussian mixture models with nearly identical proof. Then we have $\ell(\hat{z}, z^*) = o(n/k)$, as Δ^2/λ_{\max} and SNR^2 are both in the same order under the assumption $\text{SNR}/k \rightarrow \infty$. As a result, we immediately have the following corollary.

Corollary 2.1. *Assume $kd = O(\sqrt{n})$ and $\min_{a \in k} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ for some constant $\alpha > 0$. Assume $\frac{\text{SNR}}{k} \rightarrow \infty$ and $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$. Using the vanilla Lloyd's algorithm as the initialization $z^{(0)}$ in Algorithm 1, we have with probability at least $1 - n^{-1} - \exp(-\text{SNR}) - \exp(-\Delta)$,*

$$h(z^{(t)}, z^*) \leq \exp\left(-\left(1 + o(1)\right)\frac{\text{SNR}^2}{8}\right), \quad \text{for all } t \geq \log n.$$

3 GMM with Unknown and Heterogeneous Covariance Matrices

3.1 Model

In this section, we study the GMM with covariance matrices from each cluster unknown and not necessarily equal to each other. The data generation process can be displayed as follow,

Model 2:
$$Y_j = \theta_{z_j^*}^* + \epsilon_j, \quad \text{where } \epsilon_j \stackrel{\text{ind}}{\sim} \mathcal{N}(0, \Sigma_{z_j^*}^*), \forall j \in [n]. \quad (10)$$

We call it *Model 2* throughout the paper to distinguish it from Model 1 studied in Section 2. The difference between (10) and (1) is that we now have $\{\Sigma_a^*\}_{a \in [k]}$ instead of a shared Σ^* . We consider the same loss functions as in (4) and (5).

Signal-to-noise Ratio. The signal-to-noise ratio for Model 2 is defined as follows. We use the notation SNR' to distinguish it from SNR for Model 1. Compared to SNR , SNR' is much more

complicated and does not have an explicit formula. We first define a space $\mathcal{B}_{a,b} \in \mathbb{R}^d$ for any $a, b \in [k]$ such that $a \neq b$:

$$\mathcal{B}_{a,b} = \left\{ x \in \mathbb{R}^d : x^T \Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} (\theta_a^* - \theta_b^*) + \frac{1}{2} x^T \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I_d \right) x \leq -\frac{1}{2} (\theta_a^* - \theta_b^*)^T \Sigma_b^{*-1} (\theta_a^* - \theta_b^*) + \frac{1}{2} \log |\Sigma_a^*| - \frac{1}{2} \log |\Sigma_b^*| \right\}.$$

We then define $\text{SNR}'_{a,b} = 2 \min_{x \in \mathcal{B}_{a,b}} \|x\|$ and

$$\text{SNR}' = \min_{a,b \in [k]: a \neq b} \text{SNR}'_{a,b}. \quad (11)$$

The form of SNR' is closely connected to the testing error of the Quadratic Discriminant Analysis (QDA), which we will give in Lemma 3.1. For the interpretation of the SNR' (especially from a geometric point of view), we defer it after presenting Lemma 3.1. Here let us consider a few special cases where we are able to simplify SNR' : (1) When $\Sigma_a^* = \Sigma^*$ for all $a \in [k]$, by simple algebra, we have $\text{SNR}'_{a,b} = \|(\theta_a^* - \theta_b^*)^T \Sigma^{*-1}\|$ for any $a, b \in [k]$ such that $a \neq b$. Hence, $\text{SNR}' = \text{SNR}$ and Model 2 is reduced to the Model 1. (2) When $\Sigma^* = \sigma_a^2 I_d$ for any $a \in [k]$ where $\sigma_1, \dots, \sigma_k > 0$ are large constants, we have $\text{SNR}'_{a,b}, \text{SNR}'_{b,a}$ both close to $2\|\theta_a^* - \theta_b^*\|/(\sigma_a + \sigma_b)$. From these examples, we can see SNR' is determined by both the centers $\{\theta_a^*\}_{a \in [k]}$ and the covariance matrices $\{\Sigma_a^*\}_{a \in [k]}$.

3.2 Minimax Lower Bound

We first establish the minimax lower bound for the clustering problem under Model 2.

Theorem 3.1. *Under the assumption $\frac{\text{SNR}'}{\sqrt{\log k}} \rightarrow \infty$, we have*

$$\inf_{\hat{z}} \sup_{z^* \in [k]^n} \mathbb{E} h(z, z^*) \geq \exp \left(-(1 + o(1)) \frac{\text{SNR}'^2}{8} \right).$$

If $\text{SNR}' = O(1)$ instead, we have $\inf_{\hat{z}} \sup_{z^ \in [k]^n} \mathbb{E} h(z, z^*) \geq c$ for some constant $c > 0$.*

Despite that the statement of Theorem 3.1 looks similar to that of Theorem 2.1, the two minimax lower bounds are different from each other due to the discrepancy in the dependence of the centers and the covariance matrices in SNR' and SNR . By the same argument as in Section 2.2, the minimax lower bound established in Theorem 3.1 is closely related to the Quadratic Discriminant Analysis (QDA) between two normal distributions with different means and different covariance matrices.

Lemma 3.1 (Testing Error for Quadratic Discriminant Analysis). *Consider two hypotheses $\mathbb{H}_0 : X \sim \mathcal{N}(\theta_1^*, \Sigma_1^*)$ and $\mathbb{H}_1 : X \sim \mathcal{N}(\theta_2^*, \Sigma_2^*)$. Define a testing procedure*

$$\phi = \mathbb{I} \left\{ \log |\Sigma_1^*| + (x - \theta_1^*)^T \Sigma_1^* (x - \theta_1^*) \geq \log |\Sigma_2^*| + (x - \theta_2^*)^T \Sigma_2^* (x - \theta_2^*) \right\}.$$

Then we have $\inf_{\hat{\phi}} (\mathbb{P}_{\mathbb{H}_0}(\hat{\phi} = 1) + \mathbb{P}_{\mathbb{H}_1}(\hat{\phi} = 0)) = \mathbb{P}_{\mathbb{H}_0}(\phi = 1) + \mathbb{P}_{\mathbb{H}_1}(\phi = 0)$. If $\min \{\text{SNR}'_{1,2}, \text{SNR}'_{2,1}\} \rightarrow \infty$, we have

$$\inf_{\hat{\phi}} (\mathbb{P}_{H_0}(\hat{\phi} = 1) + \mathbb{P}_{H_1}(\hat{\phi} = 0)) \geq \exp \left(-(1 + o(1)) \frac{\min \{\text{SNR}'_{1,2}, \text{SNR}'_{2,1}\}^2}{8} \right).$$

Otherwise, $\inf_{\hat{\phi}} (\mathbb{P}_{H_0}(\hat{\phi} = 1) + \mathbb{P}_{H_1}(\hat{\phi} = 0)) \geq c$ for some constant $c > 0$.




Figure 2: A geometric interpretation of SNR' .

From Lemma 3.1, we can have a geometric interpretation of SNR' . In the left panel of Figure 2, we have two normal distributions $\mathcal{N}(\theta_1^*, \Sigma_1^*)$ and $\mathcal{N}(\theta_2^*, \Sigma_2^*)$ where X can be generated from. The black curve represents the optimal testing procedure ϕ displayed in Lemma 3.1. Since Σ_1^* is not necessarily equal to Σ_2^* , the black curve is not necessarily a straight line. If \mathbb{H}_0 is true, then the probability for X to be incorrectly classified is when X falls in the gray area, which is $\mathbb{P}_{H_0}(\hat{\phi} = 1)$. To calculate it, we can make a transformation $X' = (\Sigma_1^*)^{-\frac{1}{2}}(X - \theta_1^*)$. Then displayed in the right panel of Figure 2, the two distributions become $\mathcal{N}(0, I_d)$ and $\mathcal{N}((\Sigma_1^*)^{-\frac{1}{2}}(\theta_2^* - \theta_1^*), (\Sigma_1^*)^{-\frac{1}{2}}\Sigma_2^*(\Sigma_1^*)^{-\frac{1}{2}})$, and the optimal testing procedure $\mathbb{I}\{X' \in \mathcal{B}_{1,2}\}$. As a result, in the right panel of Figure 2, $\mathcal{B}_{1,2}$ represents the space colored by gray, and the black curve is its boundary. Then $\mathbb{P}_{H_0}(\hat{\phi} = 1)$ is equal to $\mathbb{P}(\mathcal{N}(0, I_d) \in \mathcal{B}_{1,2})$, which can be shown to be determined by the minimum distance between the center of $\mathcal{N}(0, I_d)$ and the space $\mathcal{B}_{1,2}$. Denote the minimum distance by $\text{SNR}'_{1,2}/2$, by Lemmas 7.10 and Lemma 7.11, we can show $\mathbb{P}(\mathcal{N}(0, I_d) \in \mathcal{B}_{1,2}) = \exp(-(1 + o(1))\text{SNR}'_{1,2}/8)$. As a result, the SNR' can be interpreted as the minimum effective distance among the centers $\{\theta_a^*\}_{a \in [k]}$ considering the anisotropic and heterogeneous structure of $\{\Sigma_a^*\}_{a \in [k]}$ and it captures the intrinsic difficulty of the clustering problem under Model 2.

3.3 Optimal Adaptive Procedure

In this section, we will propose a computationally feasible and rate-optimal procedure for clustering under Model 2. Similar to Algorithm 1, the proposed Algorithm 2 can be seen as a variant of the Lloyd's algorithm that is adjusted to the unknown and heterogeneous covariance matrices. It can also be interpreted as a hard EM algorithm under Model 2. Algorithm 2 differs from Algorithm 1 in (13) and (14), as now there are k covariance matrices to be estimated.

In Theorem 3.2, we give a computational and statistical guarantee of the proposed Algorithm 2. We show that provided with some decent initialization, Algorithm 2 is able to achieve the minimax lower bound within $\log n$ iterations. The assumptions needed in Theorem 3.2 are similar to those in Theorem 3.2, except that we require stronger assumptions on k and the dimensionality d since now we have k (instead of one) covariance matrices to be estimated. In addition, by $\max_{a,b \in [k]} \lambda_d(\Sigma_a^*)/\lambda_1(\Sigma_b^*) = O(1)$ we not only assume each of the k covariance matrices is well-conditioned, but also assume they are comparable to each other.

Theorem 3.2. *Assume $k, d = O(1)$ and $\min_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ for some constant $\alpha > 0$. Assume $\text{SNR}' \rightarrow \infty$ and $\max_{a,b \in [k]} \lambda_d(\Sigma_a^*)/\lambda_1(\Sigma_b^*) = O(1)$. For Algorithm 2, suppose $z^{(0)}$ satisfies*

Algorithm 2: Adjusted Lloyd’s Algorithm for Model 2 (10).

Input: Data Y , number of clusters k , an initialization $z^{(0)}$, number of iterations T

Output: $z^{(T)}$

1 **for** $t = 1, \dots, T$ **do**

2 Update the centers:

$$\theta_a^{(t)} = \frac{\sum_{j \in [n]} Y_j \mathbb{I}\{z_j^{(t-1)} = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j^{(t-1)} = a\}}, \quad \forall a \in [k]. \quad (12)$$

3 Update the covariance matrices:

$$\Sigma_a^{(t)} = \frac{\sum_{j \in [n]} (Y_j - \theta_a^{(t)})(Y_j - \theta_a^{(t)})^T \mathbb{I}\{z_j^{(t-1)} = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j^{(t-1)} = a\}}, \quad \forall a \in [k]. \quad (13)$$

4 Update the cluster estimations:

$$z_j^{(t)} = \operatorname{argmin}_{a \in [k]} (Y_j - \theta_a^{(t)})^T (\Sigma_a^{(t)})^{-1} (Y_j - \theta_a^{(t)}) + \log |\Sigma_a^{(t)}|, \quad j \in [n]. \quad (14)$$

5 **end**

$\ell(z^{(0)}, z^*) = o(n/k)$ with probability at least $1 - \eta$. Then with probability at least $1 - \eta - n^{-1} - \exp(-SNR')$, we have

$$h(z^{(t)}, z^*) \leq \exp\left(-\left(1 + o(1)\right) \frac{SNR'^2}{8}\right), \quad \text{for all } t \geq \log n.$$

The vanilla Lloyd’s algorithm can be used as the initialization for Algorithm 2. Under the assumption that $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$, Model 2 is also a sub-Gaussian mixture model. By the same argument as in Section 2.3 we have the following corollary.

Corollary 3.1. *Assume $k, d = O(1)$ and $\min_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ for some constant $\alpha > 0$. Assume $SNR' \rightarrow \infty$ and $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$. Using the vanilla Lloyd’s algorithm as the initialization $z^{(0)}$ in Algorithm 2, we have with probability at least $1 - n^{-1} - \exp(-SNR)$,*

$$h(z^{(t)}, z^*) \leq \exp\left(-\left(1 + o(1)\right) \frac{SNR'^2}{8}\right), \quad \text{for all } t \geq \log n.$$

4 Numerical Studies

In this section, we compare the performance of the proposed methods with other popular clustering methods on synthetic datasets under different settings.

Model 1. The first simulation is designed for the GMM with unknown but homogeneous covariance matrices (i.e., Model 1). We independently generate $n = 1200$ samples with dimension




Figure 3: Left: Performance of Algorithm 1 compared with other methods under Model 1. Right: Performance of Algorithm 2 compared with other methods under Model 2.

$d = 50$ from $k = 30$ clusters. Each cluster has 40 samples. We set $\Sigma^* = U^T \Lambda U$, where Λ is a 50×50 diagonal matrix with diagonal elements selected from 0.5 to 8 with equal space and U is a randomly generated orthogonal matrix. The centers $\{\theta_a^*\}_{a \in [n]}$ are orthogonal or each other with $\|\theta_1^*\| = \dots = \|\theta_{30}^*\| = 9$. We consider four popular clustering methods: (1) the spectral clustering method [14] (denoted as “spectral”), (2) the vanilla Lloyd’s algorithm [15] (denoted as “vanilla Lloyd”), (3) the proposed Algorithm 1 initialized by the spectral clustering (denoted as “spectral + Alg 1”), and (4) Algorithm 1 initialized by the vanilla Lloyd (denoted as “vanilla Lloyd + Alg 1”). The comparison is presented in the left panel of Figure 3.

In the plot, the x -axis is the number of iterations and the y -axis is the logarithm of the mis-clustering error rate, i.e., $\log(h)$. Each of the curves plotted is an average of 100 independent trials. We can see the proposed Algorithm 1 outperforms the spectral clustering and the vanilla Lloyd’s algorithm significantly. What is more, the dashed line represents the optimal exponent $-\text{SNR}^2/8$ of the minimax lower bound given in Section 2.1. Then we can see Algorithm 1 achieves it after 3 iterations. This justifies the conclusion established in Theorem 2.2 that Algorithm 1 is rate-optimal.

Model 2. We also compare the performances of four methods (spectral, vanilla Lloyd, spectral + Alg 2, and vanilla Lloyd + Alg 2) for the GMM with unknown and heterogeneous covariance matrices (i.e., Model 2). In this case, we take $n = 1200$, $k = 3$ and $d = 5$. We set $\Sigma_1^* = I$, $\Sigma_2^* = \Lambda_2$ which is a 5×5 diagonal matrix with elements generated from 0.5 to 8 with equal space and $\Sigma_3^* = U^T \Lambda_3 U$, where Λ_3 is a diagonal matrix with elements selected uniformly from 0.5 to 2 and U is a randomly generated orthogonal matrix. To simplify the calculation of SNR' , we take θ_1^* as a randomly selected unit vector, $\theta_2^* = \theta_1^* + 5e_1$ with e_1 denoting the vector with a 1 in the first coordinate and 0’s elsewhere and $\theta_3^* = \theta_2^* + v_1$ with v_1 randomly selected satisfying $\|v_1\| = 10$. The comparison is presented in the right panel of Figure 3 where each curve plotted is an average of 100 independent trials.

From the plot, we can clearly see the proposed Algorithm 2 improves greatly the spectral clustering and the vanilla Lloyd algorithm. The dashed line represents the optimal exponent $-\text{SNR}'^2/8$ of the minimax lower bound given in Section 3.1, which is achieved by Algorithm 2. Hence, this numerically justifies Theorem 3.2 that Algorithm 1 is rate-optimal for clustering under Model 2.

5 Proofs in Section 2

5.1 Proofs for The Lower Bound

Proof of Lemma 2.1. Note that ϕ is the likelihood ratio test. Hence by the Neyman-Pearson lemma, it is the optimal procedure. Let $\epsilon \sim \mathcal{N}(0, I_d)$. By Gaussian tail probability, we have

$$\begin{aligned}
\mathbb{P}_{\mathbb{H}_0}(\phi = 1) + \mathbb{P}_{\mathbb{H}_1}(\phi = 0) &= \mathbb{P}\left(2(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-1}(\theta_1 + \epsilon) \geq \theta_2^{*T}(\Sigma^*)^{-1}\theta_2^* - \theta_1^{*T}(\Sigma^*)^{-1}\theta_1^*\right) \\
&\quad + \mathbb{P}\left(2(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-1}(\theta_2 + \epsilon) < \theta_2^{*T}(\Sigma^*)^{-1}\theta_2^* - \theta_1^{*T}(\Sigma^*)^{-1}\theta_1^*\right) \\
&= 2\mathbb{P}\left(2(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-1}(\theta_1 + \epsilon) \geq \theta_2^{*T}(\Sigma^*)^{-1}\theta_2^* - \theta_1^{*T}(\Sigma^*)^{-1}\theta_1^*\right) \\
&= 2\mathbb{P}\left(\epsilon > \frac{1}{2}\|(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-\frac{1}{2}}\|\right) \\
&\geq C \min\left\{1, \frac{1}{\|(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-\frac{1}{2}}\|} \exp\left(-\frac{\|(\theta_2^* - \theta_1^*)^T(\Sigma^*)^{-\frac{1}{2}}\|^2}{8}\right)\right\},
\end{aligned}$$

for some constant $C > 0$. The proof is complete. \square

Proof of Theorem 2.1. We adopt the idea from [15]. Without loss of generality, assume the minimum in (2) is achieved at $a = 1, b = 2$ so that $\text{SNR} = (\theta_1^* - \theta_2^*)^T(\Sigma^*)^{-1}(\theta_1^* - \theta_2^*)$. Consider an arbitrary $\bar{z} \in [k]^n$ such that $|\{i \in [n] : \bar{z}_i = a\}| \geq \lceil \frac{n}{k} - \frac{n}{8k^2} \rceil$ for any $a \in [k]$. Then for each $a \in [k]$, we can choose a subset of $\{i \in [n] : \bar{z}_i = a\}$ with cardinality $\lceil \frac{n}{k} - \frac{n}{8k^2} \rceil$, denoted by T_a . Let $T = \cup_{a \in [k]} T_a$. Then we can define a parameter space

$$\mathcal{Z} = \{z \in [k]^n : z_i = \bar{z}_i \text{ for all } i \in T \text{ and } z_i \in \{1, 2\} \text{ if } i \in T^c\}.$$

Notice that for any $z \neq \tilde{z} \in \mathcal{Z}$, we have $\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{z_i \neq \tilde{z}_i\} \leq \frac{k}{n} \frac{n}{8k^2} = \frac{1}{8k}$ and $\frac{1}{n} \sum_{i=1}^n \mathbb{I}\{\psi(z_i) \neq \tilde{z}_i\} \geq \frac{1}{n}(\frac{n}{2k} - \frac{n}{8k^2}) \geq \frac{1}{4k}$ for any permutation ψ on $[k]$. Thus we can conclude

$$h(z, \tilde{z}) = \frac{1}{n} \sum_{i=1}^n \mathbb{I}\{z_i \neq \tilde{z}_i\}, \quad \text{for all } z, \tilde{z} \in \mathcal{Z}.$$

We notice that

$$\begin{aligned}
\inf_{\hat{z}} \sup_{z^* \in [k]^n} \mathbb{E}h(\hat{z}, z^*) &\geq \inf_{\hat{z}} \sup_{z^* \in \mathcal{Z}} \mathbb{E}h(\hat{z}, z^*) \\
&\geq \inf_{\hat{z}} \frac{1}{|\mathcal{Z}|} \sum_{z^* \in \mathcal{Z}} \mathbb{E}h(\hat{z}, z^*) \\
&\geq \frac{1}{n} \sum_{i \in T^c} \inf_{\hat{z}_i} \frac{1}{|\mathcal{Z}|} \sum_{z^* \in \mathcal{Z}} \mathbb{P}_{z^*}(\hat{z}_i \neq z_i).
\end{aligned}$$

Now consider a fixed $i \in T^c$. Define $\mathcal{Z}_a = \{z \in \mathcal{Z} : z_i = a\}$ for $a = 1, 2$. Then we can see $\mathcal{Z} = \mathcal{Z}_1 \cup \mathcal{Z}_2$ and $\mathcal{Z}_1 \cap \mathcal{Z}_2 = \emptyset$. What is more, there exists a one-to-one mapping $f(\cdot)$ between \mathcal{Z}_1 and \mathcal{Z}_2 , such that for any $z \in \mathcal{Z}_1$, we have $f(z) \in \mathcal{Z}_2$ with $[f(z)]_j = z_j$ for any $j \neq i$ and $[f(z)]_i = 2$. Hence, we can reduce the problem into a two-point testing probe and then apply Lemma 2.1. We

first consider the case that $\text{SNR} \rightarrow \infty$. We have

$$\begin{aligned}
\inf_{\hat{z}_i} \frac{1}{|\mathcal{Z}|} \sum_{z^* \in \mathcal{Z}} \mathbb{P}_{z^*}(\hat{z}_i \neq z_i) &= \inf_{\hat{z}_i} \frac{1}{|\mathcal{Z}|} \sum_{z^* \in \mathcal{Z}_1} (\mathbb{P}_{z^*}(\hat{z}_i \neq 1) + \mathbb{P}_{f(z^*)}(\hat{z}_i \neq 2)) \\
&\geq \frac{1}{|\mathcal{Z}|} \sum_{z^* \in \mathcal{Z}_1} \inf_{\hat{z}_i} (\mathbb{P}_{z^*}(\hat{z}_i \neq 1) + \mathbb{P}_{f(z^*)}(\hat{z}_i \neq 2)) \\
&\geq \frac{|\mathcal{Z}_1|}{|\mathcal{Z}|} \exp\left(- (1 + \eta) \frac{\text{SNR}^2}{8}\right) \\
&\geq \frac{1}{2} \exp\left(- (1 + \eta) \frac{\text{SNR}^2}{8}\right),
\end{aligned}$$

for some $\eta = o(1)$. Here the second inequality is due to Lemma 2.1. Then,

$$\begin{aligned}
\inf_{\hat{z}} \sup_{z^* \in [k]^n} \mathbb{E}h(\hat{z}, z^*) &\geq \frac{|T^c|}{2n} \exp\left(- (1 + \eta) \frac{\text{SNR}^2}{8}\right) = \frac{1}{16k} \exp\left(- (1 + \eta) \frac{\text{SNR}^2}{8}\right) \\
&= \exp\left(- (1 + \eta') \frac{\text{SNR}^2}{8}\right),
\end{aligned}$$

for some other $\eta' = o(1)$, where we use $\text{SNR}^2 / \log k \rightarrow \infty$.

The proof for the case $\text{SNR} = O(1)$ is similar and hence is omitted here. \square

5.2 Proofs for The Upper Bound

In this section, we will prove Theorem 2.2 using the framework developed in [8] for analyzing iterative algorithms. The key idea to establish statistical guarantees of the proposed iterative algorithm (i.e., Algorithm 1) is to perform a ‘‘one-step’’ analysis. That is, assume we have an estimation z for z^* . Then we can apply (7), (8), and (9) on z to obtain $\{\hat{\theta}_a(z)\}_{a \in [k]}$, $\hat{\Sigma}(z)$, and $\hat{z}(z)$ sequentially, which all depend on z . Then $\hat{z}(z)$ can be seen as a refined estimation of z^* . We will first build the connection between $\ell(z, z^*)$ with $\ell(\hat{z}(z), z^*)$ as in Lemma 5.1. To establish the connection, we will decompose the loss $\ell(\hat{z}(z), z^*)$ into several errors according to the difference in their behaviors. Then we will give conditions (Condition 5.2.1 - 5.2.3), under which we will show these errors are either negligible or well controlled by $\ell(z, z^*)$. With Lemma 5.1 established, in Lemma 5.2 we will show the connection can be extended to multiple iterations, under two more conditions (Condition 5.2.4 - 5.2.5). Last, we will show all these conditions hold with high probability, and hence prove Theorem 2.2.

In the statement of Theorem 2.2, the covariance matrix Σ^* is assumed to satisfy $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$. Without loss of generality, we can replace it by assuming Σ^* satisfies

$$\lambda_{\min} \leq \lambda_1(\Sigma^*) \leq \lambda_d(\Sigma^*) \leq \lambda_{\max} \quad (15)$$

where $\lambda_{\min}, \lambda_{\max} > 0$ are two constants. This is due to the following simple argument using the scaling properties of normal distributions. Let $\{Y_j\}$ be some dataset generated according to Model 1 with parameters $\{\theta_a^*\}_{a \in [k]}$, Σ^* , and z^* . The assumption $\lambda_d(\Sigma^*)/\lambda_1(\Sigma^*) = O(1)$ is equivalent to assume there exist some constants $\lambda_{\min}, \lambda_{\max} > 0$ and some quantity $\sigma > 0$ that may depend on n such that $\lambda_{\min}\sigma^2 \leq \lambda_1(\Sigma^*) \leq \lambda_d(\Sigma^*) \leq \lambda_{\max}\sigma^2$. Then performing a scaling transformation we obtain another dataset $Y'_j = Y_j/\sigma$. Note that: 1) $\{Y'_j\}$ can be seen to be generated from Model 1 with parameters $\{\theta_a^*/\sigma\}_{a \in [k]}$, Σ^*/σ^2 , and z^* , 2) clustering on $\{Y_j\}$ is equivalent to clustering on $\{Y'_j\}$, 3) by the definition in (2), the SNRs that are associated with the data generating processes of $\{Y'_j\}$

and $\{Y_j\}$ are exactly equal to each other, and 4) we have $\lambda_{\min} \leq \lambda_1(\Sigma^*/\sigma^2) \leq \lambda_d(\Sigma^*/\sigma^2) \leq \lambda_{\max}$. Hence, in this section, we will assume (15) holds and it will not lose any generality.

In the proof, we will mainly use the loss $\ell(\cdot, \cdot)$ for convenience. Recall Δ is defined as the minimum distance among centers in (3). We have

$$h(z, z^*) \leq \frac{\ell(z, z^*)}{n\Delta^2}. \quad (16)$$

The algorithmic guarantees Lemma 5.1 and Lemma 5.2 are established with respect to the $\ell(\cdot, \cdot)$ loss. But eventually we will use (16) to convert it into a result with respect to $h(\cdot, \cdot)$ in the proof of Theorem 2.2.

Error Decomposition for the One-step Analysis: Consider an arbitrary $z \in [k]^n$. Apply (7), (8), and (9) on z to obtain $\{\hat{\theta}_a(z)\}_{a \in [k]}$, $\hat{\Sigma}(z)$, and $\hat{z}(z)$:

$$\begin{aligned} \hat{\theta}_a(z) &= \frac{\sum_{j \in [n]} Y_j \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}, \quad a \in [k] \\ \hat{\Sigma}(z) &= \frac{\sum_{a \in [k]} \sum_{j \in [n]} (Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T \mathbb{I}\{z_j = a\}}{n}, \\ \hat{z}_j(z) &= \underset{a \in [k]}{\operatorname{argmin}} (Y_j - \hat{\theta}_a(z))^T (\hat{\Sigma}(z))^{-1} (Y_j - \hat{\theta}_a(z)), \quad j \in [n]. \end{aligned}$$

For simplicity we use \hat{z} that is short for $\hat{z}(z)$. Let $j \in [n]$ be an arbitrary index with $z_j^* = a$. According to (9), z_j^* will be incorrectly estimated after on iteration in \hat{z}_j if $a \neq \operatorname{argmin}_{a \in [k]} (Y_j - \hat{\theta}_a(z))^T (\hat{\Sigma}(z))^{-1} (Y_j - \hat{\theta}_a(z))$. That is, it is important to analyze the event

$$\langle Y_j - \hat{\theta}_b(z), (\hat{\Sigma}(z))^{-1} (Y_j - \hat{\theta}_b(z)) \rangle \leq \langle Y_j - \hat{\theta}_a(z), (\hat{\Sigma}(z))^{-1} (Y_j - \hat{\theta}_a(z)) \rangle, \quad (17)$$

for any $b \in [k] \setminus \{a\}$. Note that $Y_j = \theta_a^* + \epsilon_j$. After some rearrangements, we can see (17) is equivalent to,

$$\begin{aligned} &\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1} (\hat{\theta}_a(z^*) - \hat{\theta}_b(z^*)) \rangle \\ &\leq -\frac{1}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1} (\theta_a^* - \theta_b^*) \rangle + F_j(a, b, z) + G_j(a, b, z) + H_j(a, b, z), \end{aligned}$$

where

$$\begin{aligned} F_j(a, b, z) &= \langle \epsilon_j, (\hat{\Sigma}(z))^{-1} (\hat{\theta}_b(z) - \hat{\theta}_b(z^*)) \rangle - \langle \epsilon_j, (\hat{\Sigma}(z))^{-1} (\hat{\theta}_a(z) - \hat{\theta}_a(z^*)) \rangle \\ &\quad + \langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1}) (\hat{\theta}_b(z^*) - \hat{\theta}_a(z^*)) \rangle, \\ G_j(a, b, z) &= \left(\frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_a(z)) \rangle - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle \right) \\ &\quad + \left(\frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}(z^*))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle \right) \\ &\quad + \left(-\frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_b(z)) \rangle + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle \right) \\ &\quad + \left(-\frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle \right), \end{aligned}$$

and

$$\begin{aligned}
H_j(a, b, z) = & \left(-\frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \frac{1}{2} \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \theta_b^*) \rangle \right) \\
& + \left(-\frac{1}{2} \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \theta_b^*) \rangle + \frac{1}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle \right) \\
& + \left(\frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \hat{\theta}_a(z^*)) \rangle \right).
\end{aligned}$$

Here $\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_a(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle$ is the main term that will lead to the optimal rate. Among all the remaining terms, $F_j(a, b, z)$ includes all terms involving ϵ_j . $G_j(a, b, z)$ includes all terms related to z and $H_j(a, b, z)$ consists of terms that only involves z^* . Readers can refer [8] for more information about the decomposition.

Conditions and Guarantees for One-step Analysis. To establish the guarantee for the one-step analysis, we first give several conditions on the error terms $F_j(a, b, z)$, $G_j(a, b, z)$ and $H_j(a, b, z)$.

Condition 5.2.1. Assume that

$$\max_{\{z:l(z,z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|H_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq \frac{\delta}{4}$$

holds with probability with at least $1 - \eta_1$ for some $\tau, \delta, \eta_1 > 0$.

Condition 5.2.2. Assume that

$$\max_{\{z:l(z,z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 l(z, z^*)} \leq \frac{\delta^2}{256}$$

holds with probability with at least $1 - \eta_2$ for some $\tau, \delta, \eta_2 > 0$.

Condition 5.2.3. Assume that

$$\max_{\{z:l(z,z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|G_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq \frac{\delta}{8}$$

holds with probability with at least $1 - \eta_3$ for some $\tau, \delta, \eta_3 > 0$.

We next define a quantity that we refer it as the ideal error,

$$\xi_{\text{ideal}}(\delta) = \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I}\{\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_a(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1-\delta}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle\}.$$

Lemma 5.1. Assumes Conditions 5.2.1 - 5.2.3 hold for some $\tau, \delta, \eta_1, \eta_2, \eta_3, > 0$. We then have

$$\mathbb{P} \left(\ell(\hat{z}, z^*) \leq 2\xi_{\text{ideal}}(\delta) + \frac{1}{4} \ell(z, z^*) \text{ for any } z \in [k]^n \text{ such that } \ell(z, z^*) \leq \tau \right) \geq 1 - \eta,$$

where $\eta = \sum_{i=1}^3 \eta_i$.

Proof. We notice that

$$\begin{aligned}
\mathbb{I}\{\hat{z}_j = b\} &\leq \mathbb{I}\left\{\langle Y_j - \hat{\theta}_b(z), (\hat{\Sigma}(z))^{-1}(Y_j - \hat{\theta}_b(z)) \rangle \leq \langle Y_j - \hat{\theta}_a(z), (\hat{\Sigma}(z))^{-1}(Y_j - \hat{\theta}_a(z)) \rangle\right\} \\
&= \mathbb{I}\left\{\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \hat{\theta}_b(z^*)) \rangle\right. \\
&\quad \left. \leq -\frac{1}{2}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle + F_j(z_j^*, b, z) + G_j(z_j^*, b, z) + H_j(z_j^*, b, z)\right\} \\
&\leq \mathbb{I}\left\{\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1-\delta}{2}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle\right\} \\
&\quad + \mathbb{I}\left\{\frac{\delta}{2}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \leq F_j(z_j^*, b, z) + G_j(z_j^*, b, z) + H_j(z_j^*, b, z)\right\} \\
&\leq \mathbb{I}\left\{\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1-\delta}{2}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle\right\} \\
&\quad + \mathbb{I}\left\{\frac{\delta}{8}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \leq F_j(z_j^*, b, z)\right\} \\
&\leq \mathbb{I}\left\{\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1-\delta}{2}\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle\right\} \\
&\quad + \frac{64F_j(z_j^*, b, z)^2}{\delta^2\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2},
\end{aligned}$$

where the third inequality comes from Condition 5.2.1. Thus, we have

$$\begin{aligned}
&\ell(\hat{z}, z^*) \\
&= \sum_{j=1}^n \sum_{b \in [k] \setminus \{a\}} \left\| \theta_b^* - \theta_{z_j^*}^* \right\|^2 \mathbb{I}\{\hat{z}_j = b\} \\
&\leq \xi_{\text{ideal}}(\delta) + \sum_{j=1}^n \sum_{b \in [k] \setminus \{a\}} \left\| \theta_b^* - \theta_{z_j^*}^* \right\|^2 \mathbb{I}\{\hat{z}_j = b\} \frac{64F_j(z_j^*, b, z)^2}{\delta^2\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2} \\
&\leq \xi_{\text{ideal}}(\delta) + \frac{\ell(z, z^*)}{4},
\end{aligned}$$

which implies Lemma 5.1. Here the last inequality uses Conditions 5.2.2 and 5.2.3. \square

Conditions and Guarantees for Multiple Iterations. In the above we establish a statistical guarantee for the one-step analysis. Now we will extend the result to multiple iterations. That is, starting from some initialization $z^{(0)}$, we will characterize how the losses $\ell(z^{(0)}, z^*)$, $\ell(z^{(1)}, z^*)$, $\ell(z^{(2)}, z^*)$, \dots , decay. We impose a condition on $\xi_{\text{ideal}}(\delta)$ and a condition for $z^{(0)}$.

Condition 5.2.4. *Assume that*

$$\xi_{\text{ideal}}(\delta) \leq \frac{3\tau}{8}$$

holds with probability with at least $1 - \eta_4$ for some $\tau, \delta, \eta_4 > 0$.

Finally, we need a condition on the initialization.

Condition 5.2.5. Assume that

$$\ell(z^{(0)}, z^*) \leq \tau$$

holds with probability with at least $1 - \eta_5$ for some $\tau, \eta_5 > 0$.

With these conditions satisfied, we can give a lemma that shows the linear convergence guarantee for our algorithm.

Lemma 5.2. Assumes Conditions 5.2.1 - 5.2.5 hold for some $\tau, \delta, \eta_1, \eta_2, \eta_3, \eta_4, \eta_5 > 0$. We then have

$$\ell(z^{(t)}, z^*) \leq 2\xi_{ideal}(\delta) + \frac{1}{4}\ell(z^{(t-1)}, z^*)$$

for all $t \geq 1$, with probability at least $1 - \eta$, where $\eta = \sum_{i=1}^5 \eta_i$.

Proof. By Condition 5.2.4, 5.2.5, and a mathematical induction argument, we can easily know $\ell(z^{(t)}, z^*) \leq \tau$ for any $t \geq 0$. Thus, Lemma 5.2 is a direct extension of Lemma 5.1. \square

With-high-probability Results for the Conditions and The Proof of The Main Theorem. Recall the definition of Δ in (3). Recall that in (15) we assume $\lambda_{\min} \leq \lambda_1(\Sigma^*) \leq \lambda_d(\Sigma^*) \leq \lambda_{\max}$ for two constants $\lambda_{\min}, \lambda_{\max} > 0$. Hence we have Δ is in the same order of SNR. Specifically, we have

$$\frac{1}{\lambda_{\max}}\Delta \leq \text{SNR} \leq \frac{1}{\lambda_{\min}}\Delta. \quad (18)$$

Hence the assumption $\text{SNR}/k \rightarrow \infty$ in the statement of Theorem 2.2 is equivalently $\Delta/k \rightarrow \infty$. Next, we give two lemmas to clarify the Conditions. The first lemma shows that δ can be taken for some $o(1)$ term and the second lemma shows for any $\delta = o(1)$, $\xi_{ideal}(\delta)$ is upper bounded by the desired minimax rate multiply by the sample size n .

Lemma 5.3. Under the same conditions as in Theorem 2.2, for any constant $C' > 0$, there exists some constant $C > 0$ only depending on α and C' such that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|H_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq C \sqrt{\frac{k(d + \log n)}{n}} \quad (19)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq Ck^3 \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right) \quad (20)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|G_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq Ck \left(\frac{\tau}{n} + \frac{1}{\Delta} \sqrt{\frac{\tau}{n}} + \frac{d\sqrt{\tau}}{n\Delta} \right) \quad (21)$$

with probability at least $1 - n^{-C'}$.

Proof. Under the conditions of Theorem 2.2, the inequalities (33)-(38) hold with probability at least $1 - n^{-C'}$. In the remaining proof, we will work on the event these inequalities hold. Denote $\hat{\Sigma}_a(z) = \frac{\sum_{j \in [n]} (Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}$ and $\Sigma_a^* = \Sigma^*$ for any $a \in [k]$. Then we have the equivalence

$$\hat{\Sigma}(z^*) - \Sigma^* = \sum_{a=1}^k \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{n} (\hat{\Sigma}_a(z^*) - \Sigma_a^*).$$

Hence, we can use the results from Lemma 7.7 and Lemma 7.8.

By (43) and (44), we have

$$\|\hat{\Sigma}(z^*) - \Sigma^*\| \lesssim \sqrt{\frac{k(d + \log n)}{n}},$$

and

$$\begin{aligned} \|\hat{\Sigma}(z) - \hat{\Sigma}(z^*)\| &= \left\| \sum_{a=1}^k \frac{\sum_{j=1}^n \mathbb{I}\{z_j = a\}}{n} \hat{\Sigma}_a(z) - \sum_{a=1}^k \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{n} \hat{\Sigma}_a(z^*) \right\| \\ &\lesssim \left\| \sum_{a=1}^k \frac{\sum_{j=1}^n \mathbb{I}\{z_j = a\}}{n} (\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)) \right\| + \left\| \sum_{a=1}^k \frac{\sum_{j=1}^n (\mathbb{I}\{z_j = a\} - \mathbb{I}\{z_j^* = a\})}{n} \hat{\Sigma}_a(z^*) \right\| \\ &\lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)} + \frac{k}{n\Delta^2}\ell(z, z^*) \\ &\lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}. \end{aligned}$$

By the assumption that $kd = O(\sqrt{n})$, $\frac{\Delta}{k} \rightarrow \infty$ and $\tau = o(n/k)$, we have $\|\hat{\Sigma}(z^*) - \Sigma^*\|, \|\hat{\Sigma}(z) - \hat{\Sigma}(z^*)\| = o(1)$, which implies $\|(\hat{\Sigma}(z^*))^{-1}\|, \|(\hat{\Sigma}(z))^{-1}\| \lesssim 1$. Thus, we have

$$\|(\hat{\Sigma}(z^*))^{-1} - (\Sigma^*)^{-1}\| \leq \|(\hat{\Sigma}(z^*))^{-1}\| \|\hat{\Sigma}(z^*) - \Sigma^*\| \|(\Sigma^*)^{-1}\| \lesssim \sqrt{\frac{k(d + \log n)}{n}}, \quad (22)$$

and similarly

$$\|(\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1}\| \lesssim \frac{k}{n}\ell(z, z^*) + \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}. \quad (23)$$

Now we start to prove (19)-(21). Let $F_j(a, b, z) = F_j^{(1)}(a, b, z) + F_j^{(2)}(a, b, z) + F_j^{(3)}(a, b, z)$ where

$$F_j^{(1)}(a, b, z) := \langle \epsilon_j, (\hat{\Sigma}(z))^{-1}(\hat{\theta}_b(z) - \hat{\theta}_b(z^*)) \rangle - \langle \epsilon_j, (\hat{\Sigma}(z))^{-1}(\hat{\theta}_a(z) - \hat{\theta}_a(z^*)) \rangle,$$

$$F_j^{(2)}(a, b, z) := -\langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_a^* - \theta_b^*) \rangle,$$

$$F_j^{(3)}(a, b, z) := -\langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_b^* - \hat{\theta}_b(z^*)) \rangle + \langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_a^* - \hat{\theta}_a(z^*)) \rangle.$$

Notice that

$$\begin{aligned} &\sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j^{(2)}(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \\ &\lesssim \sum_{j=1}^n \sum_{b=1}^k \frac{\left| \langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_{z_j^*}^* - \theta_b^*) \rangle \right|^2}{\|\theta_{z_j^*}^* - \theta_b^*\|^2 \ell(z, z^*)} \\ &\leq \sum_{b=1}^k \sum_{a \in [k] \setminus \{b\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \frac{\left| \langle \epsilon_j, ((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_a^* - \theta_b^*) \rangle \right|^2}{\|\theta_a^* - \theta_b^*\|^2 \ell(z, z^*)} \\ &\leq \sum_{b=1}^k \sum_{a \in [k] \setminus \{b\}} \frac{\|((\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1})(\theta_a^* - \theta_b^*)\|^2}{\|\theta_a^* - \theta_b^*\|^2 \ell(z, z^*)} \left\| \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j \epsilon_j^T \right\| \\ &\lesssim k^3 \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right), \end{aligned}$$

where we use (34), (23), and the fact that $\ell(z, z^*) \leq \tau$ and $kd = O(\sqrt{n})$ for the last inequality. From (41) we have $\max_{a \in [k]} \|\theta_a^* - \hat{\theta}_a(z^*)\| = o(1)$ under the assumption $kd = O(\sqrt{n})$. By the similar analysis as in $F_j^{(2)}(a, b, z)$, we have

$$\sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j^{(3)}(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \lesssim k^3 \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right).$$

Similarly, we have

$$\begin{aligned} & \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j^{(1)}(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \\ & \lesssim \sum_{b=1}^k \sum_{a \in [k] \setminus \{b\}} \frac{\|(\hat{\Sigma}(z))^{-1}(\hat{\theta}_a(z) - \hat{\theta}_a(z^*))\|^2}{\|\theta_a^* - \theta_b^*\|^2 \ell(z, z^*)} \left\| \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j \epsilon_j^T \right\| \\ & \lesssim \frac{k^3}{\Delta^4}, \end{aligned}$$

where we use (42) and the fact that $(\hat{\Sigma}(z))^{-1}$ has bounded operator norm. Combining these terms together, we obtain (20).

Next, for (19), by (41) we have

$$\begin{aligned} & | - \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \theta_b^*) \rangle | \\ & \leq | \langle \theta_b^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_b^* - \hat{\theta}_b(z^*)) \rangle | + 2 | \langle \theta_b^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \theta_b^*) \rangle | \\ & \lesssim \frac{k(d + \log n)}{n} + \sqrt{\frac{k(d + \log n)}{n}} \|\theta_a^* - \theta_b^*\|, \end{aligned}$$

and

$$\langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \hat{\theta}_a(z^*)) \rangle \lesssim \frac{k(d + \log n)}{n}.$$

By (22) we have

$$- \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \theta_b^*) \rangle + \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle \lesssim \sqrt{\frac{k(d + \log n)}{n}} \|\theta_a^* - \theta_b^*\|^2.$$

Using the results above we can get (19).

Finally we are going to establish (21). Recall the definition of $G_j(a, b, z)$ which has four terms. For the third and fourth terms, we have

$$\begin{aligned} & - \langle \theta_a^* - \hat{\theta}_b(z), (\hat{\Sigma}(z))^{-1}(\theta_a^* - \hat{\theta}_b(z)) \rangle + \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle \\ & \lesssim \|\hat{\theta}_b(z) - \hat{\theta}_b(z^*)\|^2 + \|\hat{\theta}_b(z) - \hat{\theta}_b(z^*)\| \|\theta_a^* - \theta_b^*\|, \end{aligned}$$

and

$$\begin{aligned} & - \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}(z^*))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle \\ & \lesssim \|\theta_a^* - \theta_b^*\|^2 \|(\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1}\|. \end{aligned}$$

We can easily verify that the other two terms are smaller than the above two terms. Then, by using (42) and (23), we have

$$\begin{aligned}
& \frac{|G_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \\
& \lesssim \frac{\|\hat{\theta}_b(z) - \hat{\theta}_b(z^*)\|^2 + \|\hat{\theta}_b(z) - \hat{\theta}_b(z^*)\| \|\theta_{z_j^*}^* - \theta_b^*\| + \|\theta_{z_j^*}^* - \theta_b^*\|^2 \|(\hat{\Sigma}(z))^{-1} - (\hat{\Sigma}(z^*))^{-1}\|}{\|\theta_{z_j^*}^* - \theta_b^*\|^2} \\
& \lesssim \frac{k\tau}{n} + \frac{k}{\Delta} \sqrt{\frac{\tau}{n}} + \frac{kd\sqrt{\tau}}{n\Delta}.
\end{aligned}$$

□

Lemma 5.4. *With the same conditions as Theorem 2.2, for any sequence $\delta_n = o(1)$, we have*

$$\xi_{ideal}(\delta_n) \leq n \exp\left(- (1 + o(1)) \frac{SNR^2}{8}\right).$$

with probability at least $1 - n^{-C'} - \exp(-SNR)$.

Proof. Under the conditions of Theorem 2.2, the inequalities (33)-(38) hold with probability at least $1 - n^{-C'}$. In the remaining proof, we will work on the event these inequalities hold. Recall the definition of ξ_{ideal} , we can write

$$\begin{aligned}
\xi_{ideal}(\delta) &= \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ \langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \hat{\theta}_b(z^*)) \rangle \leq -\frac{1-\delta}{2} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \\
&\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ \langle \epsilon_j, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \leq -\frac{1-\delta-\bar{\delta}}{2} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \\
&+ \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ \langle \epsilon_j, ((\hat{\Sigma}(z^*))^{-1} - (\Sigma^*)^{-1})(\theta_{z_j^*}^* - \theta_b^*) \rangle \leq -\frac{\bar{\delta}}{6} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \\
&+ \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ \langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_{z_j^*}(z^*) - \theta_{z_j^*}^*) \rangle \leq -\frac{\bar{\delta}}{6} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \\
&+ \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ -\langle \epsilon_j, (\hat{\Sigma}(z^*))^{-1}(\hat{\theta}_b(z^*) - \theta_b^*) \rangle \leq -\frac{\bar{\delta}}{6} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \\
&=: M_1 + M_2 + M_3 + M_4.
\end{aligned}$$

where $\bar{\delta} = \bar{\delta}_n$ is some sequence to be chosen later. We bound the four terms respectively. Suppose

$\epsilon_j = (\Sigma^*)^{1/2} \omega_j$, where $w_j \stackrel{iid}{\sim} \mathcal{N}(0, I_d)$. By (22), we know

$$\begin{aligned}
M_2 &\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ \frac{\bar{\delta}}{6\lambda_{\max}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \leq \lambda_{\max} \|w_j\| \|(\hat{\Sigma}(z^*))^{-1} - (\Sigma^*)^{-1}\| \|\theta_{z_j^*}^* - \theta_b^*\| \right\} \\
&\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ C\bar{\delta} \|\theta_{z_j^*}^* - \theta_b^*\| \sqrt{\frac{n}{d + \log n}} \leq \|w_j\| \right\} \\
&\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I} \left\{ C\bar{\delta}^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2 \frac{n}{d + \log n} - 2d \leq \|w_j\|^2 - 2d \right\},
\end{aligned}$$

where C is a constant which may differ line by line. Recall that $kd = O(\sqrt{n})$, $\min_{a \neq b} \|\theta_a^* - \theta_b^*\| \rightarrow \infty$, and $\Delta/k \rightarrow \infty$ by assumption. Let $n^{-\frac{1}{4}} = o(\bar{\delta})$. Using the χ^2 tail probability in Lemma 7.1, we have for any $a \neq b \in [k]$,

$$\mathbb{E}M_2 \leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \exp\left(-C\bar{\delta}^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2 \sqrt{n}\right) \leq n \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right).$$

We can obtain similar bounds on M_3 and M_4 by using (41). For M_1 , the Gaussian tail bound leads to the inequality

$$\begin{aligned}
&\mathbb{P} \left\{ \langle \epsilon_j, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle \leq -\frac{1 - \delta - \bar{\delta}}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle \right\} \\
&= \mathbb{P} \left\{ \langle w_j, (\Sigma^*)^{-1/2}(\theta_a^* - \theta_b^*) \rangle \leq -\frac{1 - \delta - \bar{\delta}}{2} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle \right\} \\
&\leq \exp\left(-\frac{(1 - \delta - \bar{\delta})^2}{8} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle\right).
\end{aligned}$$

Thus,

$$\begin{aligned}
\mathbb{E}M_1 &\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \exp\left(-\frac{(1 - \delta - \bar{\delta})^2}{8} \langle \theta_a^* - \theta_b^*, (\Sigma^*)^{-1}(\theta_a^* - \theta_b^*) \rangle\right) \\
&\leq n \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right).
\end{aligned}$$

Overall, we have $\mathbb{E}\xi_{\text{ideal}} \lesssim n \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right)$. By the Markov's inequality, we have

$$\mathbb{P}(\xi_{\text{ideal}}(\delta_n) \geq \mathbb{E}\xi_{\text{ideal}} \exp(\text{SNR})) \leq \exp(-\text{SNR}).$$

In other words, with probability at least $1 - \exp(-\text{SNR})$, we have

$$\xi_{\text{ideal}}(\delta_n) \leq \mathbb{E}\xi_{\text{ideal}}(\delta_n) \exp(\text{SNR}) \leq n \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right).$$

□

Proof of Theorem 2.2. By Lemmas 5.2 - 5.4, we have Conditions 5.2.1 - 5.2.5 satisfied with probability at least $1 - \eta - n^{-1} - \exp(-\text{SNR})$. Then applying Lemma 5.2, we have

$$\ell(z^{(t)}, z^*) \leq n \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right) + \frac{1}{4} \ell(z^{(t-1)}, z^*), \quad \text{for all } t \geq 1.$$

By (16) and there exists a constant C such that $\Delta \leq C\text{SNR}$, we can conclude

$$h(z^{(t)}, z^*) \leq \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right) + 4^{-t}, \quad \text{for all } t \geq 1.$$

Notice that $h(z, z^*)$ takes value in the set $\{j/n : j \in [n] \cup \{0\}\}$, the term 4^{-t} in the above inequality should be negligible as long as $4^{-t} = o(n^{-1})$. Thus, we can claim

$$h(z^{(t)}, z^*) \leq \exp\left(- (1 + o(1)) \frac{\text{SNR}^2}{8}\right), \quad \text{for all } t \geq \log n.$$

□

6 Proofs in Section 3

6.1 Proofs for The Lower Bound

Proof of Lemma 3.1. The Neyman-Pearson lemma tells us the likelihood ratio test ϕ is the optimal procedure. Following the proof of Lemma 2.1, we have

$$\begin{aligned} \mathbb{P}_{\mathbb{H}_0}(\phi = 1) + \mathbb{P}_{\mathbb{H}_1}(\phi = 0) &= \mathbb{P}(\epsilon \in \mathcal{B}_{1,2}) + \mathbb{P}(\epsilon \in \mathcal{B}_{2,1}) \\ &\geq \exp\left(- \frac{1 + o(1)}{8} \text{SNR}'_{1,2}\right) + \exp\left(- \frac{1 + o(1)}{8} \text{SNR}'_{2,1}\right), \end{aligned}$$

where the last inequality is by Lemma 7.11. □

Proof of Theorem 3.1. The proof is identical to the proof of Theorem 2.1 and is omitted here. □

6.2 Proofs for The Upper Bound

We adopt a similar proof idea as in Section 5.2. We first present an error decomposition for one-step analysis for Algorithm 2. In Lemma 6.1, we show the loss decays after a one-step iteration under Conditions 6.2.1 - 6.2.6. Then in Lemma 6.2 we extend the result to multiple iterations, under two extra Conditions 6.2.7 - 6.2.8. Last we show all the conditions are satisfied with high probability and thus prove Theorem 3.2.

In the statement of Theorem 3.2, we assume $\max_{a,b \in [k]} \lambda_d(\Sigma_a^*) / \lambda_1(\Sigma_b^*) = O(1)$ for the covariance matrix $\{\Sigma_a^*\}_{a \in [k]}$. Without loss of generality, we can replace it by assuming Σ^* satisfies

$$\lambda_{\min} \leq \min_{a \in [k]} \lambda_1(\Sigma_a^*) \leq \max_{a \in [k]} \lambda_d(\Sigma_a^*) \leq \lambda_{\max} \quad (24)$$

where $\lambda_{\min}, \lambda_{\max} > 0$ are two constants. This is due to the scaling properties of the normal distributions. The reasoning is the same as that of (15) for Model 1 and hence is omitted here. In the remaining of this section, we will assume (24) holds for the covariance matrices.

Error Decomposition for the One-step Analysis: Consider an arbitrary $z \in [k]^n$. Apply (12), (13), and (14) on z to obtain $\{\hat{\theta}_a(z)\}_{a \in [k]}$, $\{\hat{\Sigma}_a(z)\}_{a \in [k]}$, and $\hat{z}(z)$:

$$\begin{aligned}\hat{\theta}_a(z) &= \frac{\sum_{j \in [n]} Y_j \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}, \\ \hat{\Sigma}_a(z) &= \frac{\sum_{j \in [n]} (Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}, \\ \hat{z}_j(z) &= \operatorname{argmin}_{a \in [k]} (Y_j - \hat{\theta}_a(z))^T (\hat{\Sigma}_a(z))^{-1} (Y_j - \hat{\theta}_a(z)) + \log |\hat{\Sigma}_a(z)|, \quad j \in [n].\end{aligned}$$

For simplicity we denote \hat{z} short for $\hat{z}(z)$. Let $j \in [n]$ be an arbitrary index with $z_j^* = a$. According to (9), z_j^* will be incorrectly estimated after on iteration in \hat{z} if $a \neq \operatorname{argmin}_{a \in [k]} (Y_j - \hat{\theta}_a(z))^T (\hat{\Sigma}_a(z))^{-1} (Y_j - \hat{\theta}_a(z)) + \log |\hat{\Sigma}_a(z)|$. That is, it is important to analyze the event

$$\langle Y_j - \hat{\theta}_b(z), (\hat{\Sigma}_b(z))^{-1} (Y_j - \hat{\theta}_b(z)) \rangle + \log |\hat{\Sigma}_b(z)| \leq \langle Y_j - \hat{\theta}_a(z), (\hat{\Sigma}_a(z))^{-1} (Y_j - \hat{\theta}_a(z)) \rangle + \log |\hat{\Sigma}_a(z)|, \quad (25)$$

for any $b \in [k] \setminus \{a\}$. After some rearrangements, we can see (25) is equivalent to,

$$\begin{aligned}& \langle \epsilon_j, (\hat{\Sigma}_b(z^*))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle - \langle \epsilon_j, (\hat{\Sigma}_a(z^*))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle \\ & + \frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_b(z^*))^{-1} - (\hat{\Sigma}_a(z^*))^{-1}) \epsilon_j \rangle - \frac{1}{2} \log |\Sigma_a^*| + \frac{1}{2} \log |\Sigma_b^*| \\ & \leq -\frac{1}{2} \langle \theta_a^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_a^* - \theta_b^*) \rangle \\ & + F_j(a, b, z) + Q_j(a, b, z) + G_j(a, b, z) + H_j(a, b, z) + K_j(a, b, z) + L_j(a, b, z),\end{aligned}$$

where

$$\begin{aligned}F_j(a, b, z) &= \langle \epsilon_j, (\hat{\Sigma}_b(z))^{-1} (\hat{\theta}_b(z) - \hat{\theta}_b(z^*)) \rangle - \langle \epsilon_j, (\hat{\Sigma}_a(z))^{-1} (\hat{\theta}_a(z) - \hat{\theta}_a(z^*)) \rangle \\ & - \langle \epsilon_j, ((\hat{\Sigma}_b(z))^{-1} - (\hat{\Sigma}_b(z^*))^{-1}) (\theta_a^* - \hat{\theta}_b(z^*)) \rangle \\ & + \langle \epsilon_j, ((\hat{\Sigma}_a(z))^{-1} - (\hat{\Sigma}_a(z^*))^{-1}) (\theta_a^* - \hat{\theta}_a(z^*)) \rangle,\end{aligned}$$

$$Q_j(a, b, z) = -\frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_b(z))^{-1} - (\hat{\Sigma}_b(z^*))^{-1}) \epsilon_j \rangle + \frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_a(z))^{-1} - (\hat{\Sigma}_a(z^*))^{-1}) \epsilon_j \rangle,$$

$$\begin{aligned}G_j(a, b, z) &= \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z), (\hat{\Sigma}_a(z))^{-1} (\theta_a^* - \hat{\theta}_a(z)) \rangle - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}_a(z))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle \\ & + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}_a(z))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}_a(z^*))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle \\ & - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z), (\hat{\Sigma}_b(z))^{-1} (\theta_a^* - \hat{\theta}_b(z)) \rangle + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}_b(z))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle \\ & - \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}_b(z))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}_b(z^*))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle,\end{aligned}$$

$$\begin{aligned}H_j(a, b, z) &= -\frac{1}{2} \langle \theta_a^* - \hat{\theta}_b(z^*), (\hat{\Sigma}_b(z^*))^{-1} (\theta_a^* - \hat{\theta}_b(z^*)) \rangle + \frac{1}{2} \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}_b(z^*))^{-1} (\theta_a^* - \theta_b^*) \rangle \\ & - \frac{1}{2} \langle \theta_a^* - \theta_b^*, (\hat{\Sigma}_b(z^*))^{-1} (\theta_a^* - \theta_b^*) \rangle + \frac{1}{2} \langle \theta_a^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_a^* - \theta_b^*) \rangle \\ & + \frac{1}{2} \langle \theta_a^* - \hat{\theta}_a(z^*), (\hat{\Sigma}_a(z^*))^{-1} (\theta_a^* - \hat{\theta}_a(z^*)) \rangle,\end{aligned}$$

$$K_j(a, b, z) := \frac{1}{2}(\log |\hat{\Sigma}_a(z)| - \log |\hat{\Sigma}_a(z^*)|) - \frac{1}{2}(\log |\hat{\Sigma}_b(z)| - \log |\hat{\Sigma}_b(z^*)|),$$

and

$$L_j(a, b, z) := \frac{1}{2}(\log |\hat{\Sigma}_a(z^*)| - \log |\Sigma_a^*|) - \frac{1}{2}(\log |\hat{\Sigma}_b(z^*)| - \log |\Sigma_b^*|).$$

Among these terms, F_j, G_j, H_j are nearly identical to their counterparts in Section 5.2 with $\hat{\Sigma}(z)$ replaced by $\hat{\Sigma}_a(z)$ or $\hat{\Sigma}_b(z)$. There are three extra terms not appearing in Section 5.2: Q_j is a quadratic term of ϵ_j and K_j, L_j are terms involving matrix determinants.

Conditions and Guarantees for One-step Analysis. To establish the guarantee for the one-step analysis, we first give several conditions on the error terms.

Condition 6.2.1. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|H_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq \frac{\delta}{12}$$

holds with probability with at least $1 - \eta_1$ for some $\tau, \delta, \eta_1 > 0$.

Condition 6.2.2. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq \frac{\delta^2}{288}$$

holds with probability with at least $1 - \eta_2$ for some $\tau, \delta, \eta_2 > 0$.

Condition 6.2.3. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|G_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq \frac{\delta}{12}$$

holds with probability with at least $1 - \eta_3$ for some $\tau, \delta, \eta_3 > 0$.

Condition 6.2.4. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{Q_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq \frac{\delta^2}{288}$$

holds with probability with at least $1 - \eta_4$ for some $\tau, \delta, \eta_4 > 0$.

Condition 6.2.5. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{K_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq \frac{\delta^2}{288}$$

holds with probability with at least $1 - \eta_5$ for some $\tau, \delta, \eta_5 > 0$.

Condition 6.2.6. Assume that

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|L_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq \frac{\delta}{12}$$

holds with probability with at least $1 - \eta_6$ for some $\tau, \delta, \eta_6 > 0$.

We next define a quantity that refers to as the ideal error,

$$\begin{aligned} \xi_{\text{ideal}}(\delta) &= \sum_{j=1}^p \sum_{b \in [k] \setminus \{z_j^*\}} \|\theta_{z_j^*}^* - \theta_b^*\|^2 \mathbb{I}\{\langle \epsilon_j, (\hat{\Sigma}_b(z^*))^{-1}(\theta_a^* - \hat{\theta}_b(z^*)) \rangle - \langle \epsilon_j, (\hat{\Sigma}_a(z^*))^{-1}(\theta_a^* - \hat{\theta}_a(z^*)) \rangle\} \\ &+ \frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_b(z^*))^{-1} - (\hat{\Sigma}_a(z^*))^{-1}) \epsilon_j \rangle - \frac{1}{2} \log |\Sigma_a^*| + \frac{1}{2} \log |\Sigma_b^*| \leq -\frac{1-\delta}{2} \langle \theta_a^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_a^* - \theta_b^*) \rangle. \end{aligned}$$

Lemma 6.1. *Assumes Conditions 6.2.1 - 6.2.6 hold for some $\tau, \delta, \eta_1, \dots, \eta_6 > 0$. We then have*

$$\mathbb{P} \left(\ell(\hat{z}, z^*) \leq 2\xi_{\text{ideal}}(\delta) + \frac{1}{4} \ell(z, z^*) \text{ for any } z \in [k]^n \text{ such that } \ell(z, z^*) \leq \tau \right) \geq 1 - \eta,$$

where $\eta = \sum_{i=1}^6 \eta_i$.

Proof. The proof of this lemma is quite similar to the proof of Lemma 5.1. The additional terms Q_j and K_j can be dealt with the same way as F_j while L_j can be dealt with the same way as H_j . We omit the details here. \square

Conditions and Guarantees for Multiple Iterations. In the above we establish a statistical guarantee for the one-step analysis. Now we will extend the result to multiple iterations. That is, starting from some initialization $z^{(0)}$, we will characterize how the losses $\ell(z^{(0)}, z^*)$, $\ell(z^{(1)}, z^*)$, $\ell(z^{(2)}, z^*)$, \dots , decay. We impose a condition on $\xi_{\text{ideal}}(\delta)$ and a condition for $z^{(0)}$.

Condition 6.2.7. *Assume that*

$$\xi_{\text{ideal}}(\delta) \leq \frac{\tau}{2}$$

holds with probability with at least $1 - \eta_7$ for some $\tau, \delta, \eta_7 > 0$.

Finally, we need a condition on the initialization.

Condition 6.2.8. *Assume that*

$$\ell(z^{(0)}, z^*) \leq \tau$$

holds with probability with at least $1 - \eta_8$ for some $\tau, \eta_8 > 0$.

With these conditions satisfied, we can give a lemma that shows the linear convergence guarantee for our algorithm.

Lemma 6.2. *Assumes Conditions 6.2.1 - 6.2.8 hold for some $\tau, \delta, \eta_1, \dots, \eta_8 > 0$. We then have*

$$\ell(z^{(t)}, z^*) \leq \xi_{\text{ideal}}(\delta) + \frac{1}{2} \ell(z^{(t-1)}, z^*)$$

for all $t \geq 1$, with probability at least $1 - \eta$, where $\eta = \sum_{i=1}^8 \eta_i$.

Proof. The proof of this lemma is the same as the proof of Lemma 5.2. \square

With-high-probability Results for the Conditions and The Proof of The Main Theorem. Recall the definition of Δ in (3). Lemma 6.3 shows SNR' is in the same order with Δ , which will play a similar role as (18) in Section 5.2. It immediately implies the assumption $\text{SNR}' \rightarrow \infty$ in the statement of Theorem 3.2 is equivalently $\Delta \rightarrow \infty$. The proof of Lemma 6.3 is deferred to Section 7.

Lemma 6.3. *Assume $\text{SNR}' \rightarrow \infty$ and $d = O(1)$. Further assume there exist constants $\lambda_{\min}, \lambda_{\max} > 0$ such that $\lambda_{\min} \leq \lambda_1(\Sigma_a^*) \leq \lambda_d(\Sigma_a^*) \leq \lambda_{\max}$ for any $a \in [k]$. Then, there exist constants $C_1, C_0 > 0$ only depending on $\lambda_{\min}, \lambda_{\max}, d$ such that*

$$C_1 \|\theta_a^* - \theta_b^*\| \leq \text{SNR}'_{a,b} \leq C_2 \|\theta_a^* - \theta_b^*\|,$$

for any $a \neq b \in [k]$. As a result, SNR' is in the same order of Δ .

Lemma 6.4 and Lemma 6.5 are counterparts of Lemmas 5.3 and 5.4 in Section 5.2.

Lemma 6.4. *Under the same conditions as in Theorem 3.2, for any constant $C' > 0$, there exists some constant $C > 0$ only depending on $\alpha, C', \lambda_{\min}, \lambda_{\max}$ such that*

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|H_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq C \sqrt{\frac{k(d + \log n)}{n}} \quad (26)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{F_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq C k^3 \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right) \quad (27)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|G_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq C k \left(\frac{\tau}{n} + \frac{1}{\Delta} \sqrt{\frac{\tau}{n}} + \frac{d\sqrt{\tau}}{n\Delta} \right) \quad (28)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{Q_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq C \frac{k^3 d^2}{\Delta^2} \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right) \quad (29)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{K_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} \leq C \frac{k^3 d^2}{\Delta^2} \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right) \quad (30)$$

$$\max_{\{z: \ell(z, z^*) \leq \tau\}} \max_{j \in [n]} \max_{b \in [k] \setminus \{z_j^*\}} \frac{|L_j(z_j^*, b, z)|}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle} \leq C \frac{d}{\Delta^2} \sqrt{\frac{k(d + \log n)}{n}} \quad (31)$$

with probability at least $1 - n^{-C'} - \frac{4}{nd}$.

Proof. Under the conditions of Theorem 3.2, the inequalities (33)-(38) hold with probability at least $1 - n^{-C'}$. In the remaining proof, we will work on the event these inequalities hold. Hence, we can use the results from Lemma 7.7 and 7.8. Using the same arguments as in the proof of Lemma 5.3, we can get (26), (27) and (28).

As for (29), we first use Lemma 7.9 to have $\sum_{j=1}^n \|\epsilon_j\|^4 \leq 3nd$ with probability at least $1 - 4/(nd)$. Then, we have

$$\begin{aligned} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{Q_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1} (\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} &\lesssim \sum_{j=1}^n \sum_{b=1}^k \frac{Q_j(z_j^*, b, z)^2}{\Delta^2 \ell(z, z^*)} \\ &\leq k \sum_{j=1}^n \|\epsilon_j\|^4 \frac{\max_{a \in [k]} \|(\hat{\Sigma}_a(z))^{-1} - (\hat{\Sigma}_a(z^*))^{-1}\|^2}{\Delta^2 \ell(z, z^*)} \\ &\lesssim \frac{k^3 d^2}{\Delta^2} \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n\Delta^2} \right), \end{aligned}$$

where the last inequality is due to (53) and the fact that $\ell(z, z^*) \leq \tau$.

Next for (30), notice that by (43), (44), and $\text{SNR}' \rightarrow \infty$, we have for any $1 \leq i \leq d$, $\frac{\lambda_{\min}}{2} \leq \lambda_i(\hat{\Sigma}_a(z^*)) \leq 2\lambda_{\max}$ and

$$\left| \log\left(1 + \max_{a \in [k]} \frac{\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2}{\lambda_i(\hat{\Sigma}_a(z^*))}\right) \right| \leq \left| \log\left(1 - \max_{a \in [k]} \frac{\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2}{\lambda_i(\hat{\Sigma}_a(z^*))}\right) \right|.$$

Thus by Lemma 7.6, we know

$$\begin{aligned} & \max_{a \in [k]} \left| \log |\hat{\Sigma}_a(z)| - \log |\hat{\Sigma}_a(z^*)| \right| \\ &= \max_{a \in [k]} \left| \log \frac{|\hat{\Sigma}_a(z)|}{|\hat{\Sigma}_a(z^*)|} \right| \\ &\leq \left| \sum_{i=1}^d \log\left(1 - \frac{\max_{a \in [k]} \|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2}{\lambda_i(\hat{\Sigma}_a(z^*))}\right) \right| \\ &\leq \sum_{i=1}^d \log \left(1 + \max_{a \in [k]} \frac{\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2}{\lambda_i(\hat{\Sigma}_a(z^*))} + \frac{\max_{a \in [k]} \frac{\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2^2}{\lambda_i^2(\hat{\Sigma}_a(z^*))}}{1 - \max_{a \in [k]} \frac{\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2}{\lambda_i(\hat{\Sigma}_a(z^*))}} \right) \\ &\lesssim d \|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2, \end{aligned} \tag{32}$$

where the last inequality is due to the fact that $\lambda_i(\hat{\Sigma}_a(z^*))$ is at the constant rate, $\|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|_2 = o(1)$ and the inequality $\log(1+x) \leq x$ for any $x > 0$. (32) yields to the inequality

$$\begin{aligned} \sum_{j=1}^n \max_{b \in [k] \setminus \{z_j^*\}} \frac{K_j(z_j^*, b, z)^2 \|\theta_{z_j^*}^* - \theta_b^*\|^2}{\langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle^2 \ell(z, z^*)} &\lesssim \sum_{j=1}^n \frac{d^2 \max_{a \in [k]} \|\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)\|^2}{\Delta^2 \ell(z, z^*)} \\ &\lesssim \frac{k^2 d^2}{\Delta^2} \left(\frac{\tau}{n} + \frac{1}{\Delta^2} + \frac{d^2}{n \Delta^2} \right). \end{aligned}$$

Finally for (31), by (43) and the similar argument as (32), we can get

$$\max_{a \in [k]} \left| \log |\hat{\Sigma}_a(z^*)| - \log |\Sigma_a^*| \right| \lesssim d \sqrt{\frac{k(d + \log n)}{n}}$$

which implies (31). We complete the proof. \square

Lemma 6.5. *With the same conditions as Theorem 3.2, for any sequence $\delta_n = o(1)$, we have*

$$\xi_{\text{ideal}}(\delta_n) \leq n \exp\left(-\frac{(1 + o(1)) \text{SNR}'^2}{8}\right).$$

with probability at least $1 - n^{-C'} - \exp(-\text{SNR}')$.

Proof. Under the conditions of Theorem 3.2, the inequalities (33)-(38) hold with probability at least $1 - n^{-C'}$. In the remaining proof, we will work on the event these inequalities hold. Similar to the proof of Lemma 5.4, we have a decomposition $\xi_{\text{ideal}} \leq \sum_{i=1}^6 M_i$ where

$$\begin{aligned} M_1 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 &\mathbb{I} \left\{ \langle \epsilon_j, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle + \frac{1}{2} \langle \epsilon_j, ((\Sigma_b^*)^{-1} - (\Sigma_{z_j^*}^*)^{-1}) \epsilon_j \rangle \right. \\ &\left. - \frac{1}{2} \log |\Sigma_{z_j^*}^*| + \frac{1}{2} \log |\Sigma_b^*| \leq -\frac{1 - \delta - \bar{\delta}}{2} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\} \end{aligned}$$

is the main term and

$$M_2 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ \langle \epsilon_j, ((\hat{\Sigma}_b(z^*))^{-1} - (\Sigma_b^*)^{-1})(\theta_{z_j^*}^* - \theta_b^*) \rangle \leq -\frac{\bar{\delta}}{10} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\}$$

$$M_3 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ -\langle \epsilon_j, (\hat{\Sigma}_{z_j^*}(z^*))^{-1}(\theta_{z_j^*}^* - \hat{\theta}_{z_j^*}(z^*)) \rangle \leq -\frac{\bar{\delta}}{10} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\}$$

$$M_4 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ -\langle \epsilon_j, (\hat{\Sigma}_b(z^*))^{-1}(\hat{\theta}_b(z^*) - \theta_b^*) \rangle \leq -\frac{\bar{\delta}}{10} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\}$$

$$M_5 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ \frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_b(z^*))^{-1} - (\Sigma_b^*)^{-1})\epsilon_j \rangle \leq -\frac{\bar{\delta}}{10} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\}$$

$$M_6 := \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ -\frac{1}{2} \langle \epsilon_j, ((\hat{\Sigma}_{z_j^*}(z^*))^{-1} - (\Sigma_{z_j^*}^*)^{-1})\epsilon_j \rangle \leq -\frac{\bar{\delta}}{10} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right\}.$$

Using the same arguments as the proof of Lemma 5.4, we can choose some $\bar{\delta} = \bar{\delta}_n = o(1)$ which is slowly diverging to zero satisfying

$$\mathbb{E}M_i \leq n \exp \left(-(1 + o(1)) \frac{\text{SNR}'^2}{2} \right) \quad \text{for } i = 2, 3, 4.$$

As for M_5 , by (43) we have

$$M_5 \leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{I} \left\{ C\bar{\delta} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \leq \|w_j\|^2 \sqrt{\frac{\log n}{n}} \right\},$$

where C is a constant and $w_j \stackrel{iid}{\sim} \mathcal{N}(0, I_d)$. Since there exists some constant C' such that $\text{SNR}' \leq C'\Delta$, we can choose appropriate $\bar{\delta} = o(1)$ such that

$$\begin{aligned} \mathbb{E}M_5 &\leq \sum_{j=1}^n \sum_{b \in [k] \setminus \{z_j^*\}} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \mathbb{P} \left\{ C\bar{\delta} \left\| \theta_{z_j^*}^* - \theta_b^* \right\|^2 \sqrt{\frac{n}{\log n}} \leq \|w_j\|^2 \right\} \\ &\leq n \exp \left(-(1 + o(1)) \frac{\text{SNR}'^2}{8} \right). \end{aligned}$$

M_6 is essentially the same with M_5 . Finally for M_1 , using Lemma 7.10, we have

$$\begin{aligned}
& \mathbb{P}\left(\langle \epsilon_j, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle + \frac{1}{2} \langle \epsilon_j, ((\Sigma_b^*)^{-1} - (\Sigma_{z_j^*}^*)^{-1}) \epsilon_j \rangle \right. \\
& \quad \left. - \frac{1}{2} \log |\Sigma_{z_j^*}^*| + \frac{1}{2} \log |\Sigma_b^*| \leq -\frac{1 - \delta - \bar{\delta}}{2} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right) \\
& = \mathbb{P}\left(\langle w_j, (\Sigma_{z_j^*}^*)^{\frac{1}{2}} (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle + \frac{1}{2} \langle w_j, ((\Sigma_{z_j^*}^*)^{\frac{1}{2}} (\Sigma_b^*)^{-1} (\Sigma_{z_j^*}^*)^{\frac{1}{2}} - I_d) w_j \rangle \right. \\
& \quad \left. - \frac{1}{2} \log |\Sigma_{z_j^*}^*| + \frac{1}{2} \log |\Sigma_b^*| \leq -\frac{1 - \delta - \bar{\delta}}{2} \langle \theta_{z_j^*}^* - \theta_b^*, (\Sigma_b^*)^{-1}(\theta_{z_j^*}^* - \theta_b^*) \rangle \right) \\
& \leq \exp\left(- (1 - o(1)) \frac{\text{SNR}'_{z_j^*, b}}{8}\right).
\end{aligned}$$

Then we have

$$\mathbb{E}M_1 \leq n \exp\left(- (1 + o(1)) \frac{\text{SNR}'^2}{8}\right).$$

Using the Markov's inequality we complete the proof of Lemma 6.5. \square

Proof of Theorem 3.2. By Lemmas 6.2-6.5, we can obtain the result by arguments used in the proof of Theorem 2.2 and hence is omitted here. \square

7 Technical Lemmas

Here are the technical lemmas.

Lemma 7.1. *For any $x > 0$, we have*

$$\begin{aligned}
\mathbb{P}(\chi_d^2 \geq d + 2\sqrt{dx} + 2x) &\leq e^{-x}, \\
\mathbb{P}(\chi_d^2 \leq d - 2\sqrt{dx}) &\leq e^{-x}.
\end{aligned}$$

Proof. These results are Lemma 1 of [12]. \square

Lemma 7.2. *For any $z^* \in [k]^n$ and $k \in [n]$, consider independent vectors $\epsilon_j \sim \mathcal{N}(0, \Sigma_{z_j^*}^*)$ for any $j \in [n]$. Assume there exists a constant $\lambda_{\max} > 0$ such that $\|\Sigma_a^*\| \leq \lambda_{\max}$ for any $a \in [k]$. Then, for any constant $C' > 0$, there exists some constant $C > 0$ only depending on C', λ_{\max} such that*

$$\max_{a \in [k]} \left\| \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j}{\sqrt{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}} \right\| \leq C \sqrt{d + \log n}, \quad (33)$$

$$\max_{a \in [k]} \frac{1}{d + \sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \left\| \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j \epsilon_j^T \right\| \leq C, \quad (34)$$

$$\max_{T \subset [n]} \left\| \frac{1}{\sqrt{|T|}} \sum_{j \in T} \epsilon_j \right\| \leq C \sqrt{d + n}, \quad (35)$$

$$\max_{a \in [k]} \max_{T \subset \{j: z_j^* = a\}} \left\| \frac{1}{\sqrt{|T|(d + \sum_{j=1}^n \mathbb{I}\{z_j^* = a\})}} \sum_{j \in T} \epsilon_j \right\| \leq C, \quad (36)$$

with probability at least $1 - n^{-C'}$. We have used the convention that $0/0 = 0$.

Proof. Note that ϵ_j is sub-Gaussian with parameter λ_{\max} which is a constant. The inequalities (33) and (35) are respectively Lemmas A.4, A.1 in [15]. The inequality (34) is a slight extension of Lemma A.2 in [15]. This extension can be done by a standard union bound argument. The proof of (36) is identical to that of (35). \square

Lemma 7.3. *Consider the same assumptions as in Lemma 7.2. Assume additionally $\min_{a \in k} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ for some constant $\alpha > 0$ and $\frac{k(d+\log n)}{n} = o(1)$. Then, for any constant $C' > 0$, there exists some constant $C > 0$ only depending on $\alpha, C', \lambda_{\max}$ such that*

$$\max_{a \in [k]} \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j \epsilon_j^T - \Sigma_a^* \right\| \leq C \sqrt{\frac{k(d+\log n)}{n}}, \quad (37)$$

with probability at least $1 - n^{-C'}$.

Proof. Note that we have $\epsilon_j = \Sigma_{z_j^*}^{*\frac{1}{2}} \eta_j$ where $\eta_j \stackrel{iid}{\sim} \mathcal{N}(0, I_d)$ for any $j \in [n]$. Since $\max_a \|\Sigma_a^*\| \leq \lambda_{\max}$, we have

$$\max_{a \in [k]} \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \epsilon_j \epsilon_j^T - \Sigma_a^* \right\| \leq \lambda_{\max} \max_{a \in [k]} \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \eta_j \eta_j^T - I_d \right\|.$$

Define

$$Q_a = \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \eta_j \eta_j^T - I_d.$$

Take $S^{d-1} = \{y \in \mathbb{R}^d : \|y\| = 1\}$ and $N_\epsilon = \{v_1, \dots, v_{|N_\epsilon|}\}$ is an ϵ -covering of S^{d-1} . In particular, we pick $\epsilon < \frac{1}{4}$, then $|N_\epsilon| \leq 9^d$. By the definition of the ϵ -covering, we have

$$\|Q_a\| \leq \frac{1}{1 - 2\epsilon} \max_{i=1, \dots, |N_\epsilon|} |v_i^T Q_a v_i| \leq 2 \max_{i=1, \dots, |N_\epsilon|} |v_i^T Q_a v_i|.$$

For any $v \in N_\epsilon$,

$$v^T Q_a v = \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} (v^T \eta_j \eta_j^T v - 1).$$

Denote $n_a = \sum_{j=1}^n \mathbb{I}\{z_j^* = a\}$. Then $\sum_{j=1}^n \mathbb{I}\{z_j^* = a\} v^T \eta_j \eta_j^T v \sim \chi_{n_a}^2$. Using Lemma 7.1, we have

$$\begin{aligned} P(\max_{a \in [k]} \|Q_a\| \geq t) &\leq \sum_{a=1}^k P(\|Q_a\| \geq t) \\ &\leq \sum_{a=1}^k \sum_{i=1}^{|N_\epsilon|} P(|v_i^T Q_a v_i| \geq t/2) \\ &\leq \sum_{a=1}^k 2 \exp \left\{ -\frac{n_a}{8} \min\{t, t^2\} + d \log 9 \right\}. \end{aligned}$$

Since $\frac{k(d+\log n)}{n} = o(1)$ and $n_a \geq \alpha n/k$ where α is a constant, we can take $t = C'' \sqrt{\frac{k(d+\log n)}{n}}$ for some large constant C'' and the proof is complete. \square

Lemma 7.4. *Consider the same assumptions as in Lemma 7.2. Then, for any $s = o(n)$ and for any constant $C' > 0$, there exists some constant $C > 0$ only depending on C', λ_{\max} such that*

$$\max_{T \subset [n]: |T| \leq s} \frac{1}{|T| \log \frac{n}{|T|} + \min\{1, \sqrt{|T|}\}d} \left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| \leq C, \quad (38)$$

with probability at least $1 - n^{-C'}$. We have used the convention that $0/0 = 0$.

Proof. Consider any $a \in [s]$ and a fixed $T \subset [n]$ such that $|T| = a$. Similar to the proof of Lemma 7.3, we can take $S^{d-1} = \{y \in \mathbb{R}^d : \|y\| = 1\}$ and its ϵ -covering N_ϵ with $\epsilon < \frac{1}{4}$ and $|N_\epsilon| \leq 9^d$. Then we have

$$\left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| = \sup_{\|w\|=1} \sum_{j \in T} (w^T \epsilon_j)^2 \leq 2 \max_{w \in N_\epsilon} \sum_{j \in T} (w^T \epsilon_j)^2.$$

Note that $w^T \epsilon_j / \sqrt{\lambda_{\max}}$ is a sub-Gaussian random variable with parameter 1. By [10], for any fixed $w \in N_\epsilon$, we have

$$\mathbb{P} \left(\sum_{j \in T} (w^T \epsilon_j)^2 \geq \lambda_{\max} (a + 2\sqrt{at} + 2t) \right) \leq \exp(-t).$$

Since $a = o(n)$, there exists a constant C_0 such that $2a \leq C_0 a \log \frac{n}{a}$. We can take $t = \tilde{C}(a \log \frac{n}{a} + d)$ with $\tilde{C} = \frac{C}{16} - \frac{C_0}{4}$, then $a + 2\sqrt{at} + 2t \leq \frac{C}{4}(a \log \frac{n}{a} + d)$. Thus,

$$\mathbb{P} \left(\sum_{j \in T} (w^T \epsilon_j)^2 \geq \frac{C}{4}(a \log \frac{n}{a} + d) \right) \leq \exp \left(-\tilde{C}(a \log \frac{n}{a} + d) \right).$$

Hence, we have

$$\mathbb{P} \left(\left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| \geq \frac{C}{2}(a \log \frac{n}{a} + d) \right) \leq 9^d \exp \left(-\tilde{C}(a \log \frac{n}{a} + d) \right).$$

As a result,

$$\begin{aligned} \mathbb{P} \left\{ \max_{T \subset [n], 1 \leq |T| \leq s} \frac{1}{|T| \log \frac{n}{|T|} + d} \left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| \geq C \right\} &\leq \sum_{a=1}^s \mathbb{P} \left\{ \max_{|T|=a} \left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| \geq C(a \log \frac{n}{a} + d) \right\} \\ &\leq \sum_{a=1}^s \binom{n}{a} \max_{|T|=a} \mathbb{P} \left\{ \left\| \sum_{j \in T} \epsilon_j \epsilon_j^T \right\| \geq C(a \log \frac{n}{a} + d) \right\} \\ &\leq \sum_{a=1}^s \binom{n}{a} 9^d \exp \left(-\tilde{C}(a \log \frac{n}{a} + d) \right). \end{aligned}$$

Since $a \log \frac{n}{a}$ is an increasing function when $a \in [1, s]$ and $a \log \frac{n}{a} \geq \log n \geq \log s$, a choice of $\tilde{C} = 3 + C'$, that is $C = 16C' + 4C_0 + 48$, can yield the desired result.

Finally, to allow $|T| = 0$, we note that $d \leq \min\{1, \sqrt{|T|}\}d$. The proof is complete. \square

Lemma 7.5. For any $z^* \in [k]^n$ and $k \in [n]$, assume $\min_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$ and $\ell(z, z^*) = o(\frac{n\Delta^2}{k})$, then

$$\max_{a \in [k]} \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \leq 2. \quad (39)$$

Proof. For any $z \in [k]^n$ such that $\ell(z, z^*) = o(n)$ and any $a \in [k]$, we have

$$\begin{aligned} \sum_{j=1}^n \mathbb{I}\{z_j = a\} &\geq \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} - \sum_{j=1}^n \mathbb{I}\{z_j \neq z_j^*\} \\ &\geq \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} - \frac{\ell(z, z^*)}{\Delta^2} \\ &\geq \frac{\alpha n}{2k}, \end{aligned} \quad (40)$$

which implies

$$\begin{aligned} \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} &\leq \frac{\sum_{j=1}^n \mathbb{I}\{z_j = a\} + \sum_{j=1}^n \mathbb{I}\{z_j \neq z_j^*\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \\ &\leq 1 + \frac{\alpha n / 2k}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \\ &\leq 2. \end{aligned}$$

Thus, we obtain (39). □

The next lemma is the famous Weyl's Theorem and we omit the proof here.

Lemma 7.6 (Weyl's Theorem). Let A and B be any two $d \times d$ symmetric real matrix. Then for any $1 \leq i \leq d$, we have

$$\lambda_i(A + B) \leq \lambda_d(A) + \lambda_i(B).$$

In the following lemma, we are going to analyze estimation errors of $\{\Sigma_a^*\}_{a \in [k]}$ under the anisotropic GMMs. For any $z \in [k]^n$ and for any $a \in [k]$, recall the definitions

$$\begin{aligned} \hat{\theta}_a(z) &= \frac{\sum_{j \in [n]} Y_j \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}, \\ \hat{\Sigma}_a(z) &= \frac{\sum_{j \in [n]} (Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T \mathbb{I}\{z_j = a\}}{\sum_{j \in [n]} \mathbb{I}\{z_j = a\}}. \end{aligned}$$

Lemma 7.7. For any $z^* \in [k]^n$ and $k \in [n]$, consider independent vectors $Y_j = \theta_{z_j^*}^* + \epsilon_j$ where $\epsilon_j \sim \mathcal{N}(0, \Sigma_{z_j^*}^*)$ for any $j \in [n]$. Assume there exist constants $\lambda_{\min}, \lambda_{\max} > 0$ such that $\lambda_{\min} \leq \lambda_1(\Sigma_a^*) \leq \lambda_d(\Sigma_a^*) \leq \lambda_{\max}$ for any $a \in [k]$, and a constant $\alpha > 0$ such that $\min_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} \geq \frac{\alpha n}{k}$. Assume $\frac{k(d+\log n)}{n} = o(1)$ and $\frac{\Delta}{k} \rightarrow \infty$. Assume (33)-(38) hold. Then for any $\tau = o(n)$ and for any

constant $C' > 0$, there exists some constant $C > 0$ only depending on $\alpha, \lambda_{\max}, C'$ such that

$$\max_{a \in [k]} \left\| \hat{\theta}_a(z^*) - \theta_a^* \right\| \leq C \sqrt{\frac{k(d + \log n)}{n}}, \quad (41)$$

$$\max_{a \in [k]} \left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\| \leq C \left(\frac{k}{n\Delta} \ell(z, z^*) + \frac{k\sqrt{d+n}}{n\Delta} \sqrt{\ell(z, z^*)} \right), \quad (42)$$

$$\max_{a \in [k]} \left\| \hat{\Sigma}_a(z^*) - \Sigma_a^* \right\| \leq C \sqrt{\frac{k(d + \log n)}{n}}, \quad (43)$$

$$\max_{a \in [k]} \left\| \hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*) \right\| \leq C \left(\frac{k}{n} \ell(z, z^*) + \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{kd}{n\Delta} \sqrt{\ell(z, z^*)} \right), \quad (44)$$

for all z such that $\ell(z, z^*) \leq \tau$.

Proof. Using (33) we obtain (41). By the same argument of (118) in [8], we can obtain (42). By (33) and (37) and (41), we can obtain (43). In the remaining of the proof, we will establish (53).

Since $\frac{k(d+\log n)}{n} = o(1)$, we have $\|\hat{\Sigma}_a(z^*)\| \lesssim 1$ for any $a \in [k]$. The difference $\hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*)$ will be decomposed into several terms. We notice that

$$\left\| \hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*) \right\| \leq S_1 + S_2, \quad (45)$$

where

$$S_1 := \left\| \frac{1}{\sum \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a\} \left((Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T - (Y_j - \hat{\theta}_a(z^*))(Y_j - \hat{\theta}_a(z^*))^T \right) \right\|,$$

and

$$S_2 := \left\| \left(\frac{1}{\sum \mathbb{I}\{z_j = a\}} - \frac{1}{\sum \mathbb{I}\{z_j^* = a\}} \right) \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} (Y_j - \hat{\theta}_a(z^*))(Y_j - \hat{\theta}_a(z^*))^T \right\|.$$

Also, we notice that

$$S_1 \leq L_1 + L_2 + L_3, \quad (46)$$

where

$$L_1 := \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\} \left((Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T - (Y_j - \hat{\theta}_a(z^*))(Y_j - \hat{\theta}_a(z^*))^T \right) \right\|,$$

$$L_2 := \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} (Y_j - \hat{\theta}_a(z))(Y_j - \hat{\theta}_a(z))^T \right\|,$$

$$L_3 := \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j \neq a, z_j^* = a\} (Y_j - \hat{\theta}_a(z^*))(Y_j - \hat{\theta}_a(z^*))^T \right\|.$$

For L_1 , we have

$$\begin{aligned}
L_1 &\leq \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\} (\hat{\theta}_a(z) - \hat{\theta}_a(z^*)) (\hat{\theta}_a(z) - \hat{\theta}_a(z^*))^T \right\| \\
&\quad + 2 \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\} (Y_j - \hat{\theta}_a(z^*)) (\hat{\theta}_a(z) - \hat{\theta}_a(z^*))^T \right\| \\
&\lesssim \left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\|^2 \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} + \left\| \theta_a^* - \hat{\theta}_a(z^*) \right\| \left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\| \frac{\sum_{j=1}^n \mathbb{I}\{z_j^* = a\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \\
&\quad + \left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\| \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\} \epsilon_j \right\|. \tag{47}
\end{aligned}$$

By (36), (39), (40), we have uniformly for any $a \in [k]$,

$$\begin{aligned}
\left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\} \epsilon_j \right\| &\lesssim \frac{\sqrt{\sum_{j=1}^n \mathbb{I}\{z_j = z_j^* = a\}}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sqrt{d + \sum_{j=1}^n \mathbb{I}\{z_j^* = a\}} \\
&\lesssim 1. \tag{48}
\end{aligned}$$

Since $\max_{a \in [k]} \left\| \hat{\theta}_a(z^*) - \theta_a^* \right\| = o(1)$, by (39), (42), (41), (47), and (48), we have uniformly for any $a \in [k]$,

$$L_1 \lesssim \left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\| \lesssim \frac{k}{n\Delta} \ell(z, z^*) + \frac{k\sqrt{d+n}}{n\Delta} \sqrt{\ell(z, z^*)}. \tag{49}$$

To bound L_2 , we first give the following simple fact. For any positive integer m and any $\{u_j\}_{j \in [m]}, \{v_j\}_{j \in [m]} \in \mathbb{R}^d$, we have $\left\| \sum_{j \in [m]} (u_j + v_j)(u_j + v_j)^T \right\| \leq 2 \left\| \sum_{j \in [m]} u_j u_j^T \right\| + 2 \left\| \sum_{j \in [m]} v_j v_j^T \right\|$. Hence, for L_2 , we have the following decomposition

$$L_2 \leq 2R_1 + 2R_2, \tag{50}$$

where

$$\begin{aligned}
R_1 &:= \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} (Y_j - \theta_a^*) (Y_j - \theta_a^*)^T \right\|, \\
R_2 &:= \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} (\theta_a^* - \hat{\theta}_a(z)) (\theta_a^* - \hat{\theta}_a(z))^T \right\|.
\end{aligned}$$

Since $\max_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} \leq \frac{\ell(z, z^*)}{\Delta^2}$, we have

$$\begin{aligned}
R_2 &\leq \left\| \theta_a^* - \hat{\theta}_a(z) \right\|^2 \frac{\sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\}}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \\
&\lesssim \left(\left\| \hat{\theta}_a(z) - \hat{\theta}_a(z^*) \right\|^2 + \left\| \hat{\theta}_a(z^*) - \theta_a^* \right\|^2 \right) \frac{k\ell(z, z^*)}{n\Delta^2}. \tag{51}
\end{aligned}$$

By (38) and the fact that $\max_{a \in [k]} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} \leq \frac{\ell(z, z^*)}{\Delta^2}$, we also have

$$\begin{aligned}
R_1 &\leq 2 \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} (\theta_{z_j^*}^* - \theta_{z_j}^*) (\theta_{z_j^*}^* - \theta_{z_j}^*)^T \right\| \\
&\quad + 2 \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} \epsilon_j \epsilon_j^T \right\| \\
&\leq 2 \frac{\sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} \|\theta_{z_j^*}^* - \theta_{z_j}^*\|^2}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} + 2 \left\| \frac{1}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \sum_{j=1}^n \mathbb{I}\{z_j = a, z_j^* \neq a\} \epsilon_j \epsilon_j^T \right\| \\
&\lesssim \frac{k\ell(z, z^*)}{n} + \frac{\frac{\ell(z, z^*)}{\Delta^2} \log \frac{n\Delta^2}{\ell(z, z^*)} + d\sqrt{\frac{\ell(z, z^*)}{\Delta^2}}}{n/k}.
\end{aligned}$$

We are going to simplify the above bounds for R_1, R_2 . Under the assumption that $\frac{k(d+\log n)}{n} = o(1)$, $\Delta/k \rightarrow \infty$, and $\ell(z, z^*) \leq \tau = o(n)$, we have $\max_{a \in [k]} \|\hat{\theta}_a(z) - \hat{\theta}_a(z^*)\| = o(1)$, $\max_{a \in [k]} \|\hat{\theta}_a(z^*) - \theta_a^*\| = o(1)$, and $\frac{k\ell(z, z^*)}{n\Delta^2} = o(1)$. Hence $R_2 \lesssim \frac{k\ell(z, z^*)}{n\Delta^2}$. Also we have

$$\frac{k\ell(z, z^*)}{n\Delta^2} \log \frac{n\Delta^2}{\ell(z, z^*)} = \frac{k\sqrt{\ell(z, z^*)}}{n\Delta} \sqrt{\frac{\ell(z, z^*)}{\Delta^2} \left(\log \frac{n\Delta^2}{\ell(z, z^*)} \right)^2} \leq \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta}.$$

where in the last inequality, we use the fact that $x(\log(n/x))^2$ is an increasing function of x when $0 < x = o(n)$. Then,

$$L_2 \lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}.$$

Since L_3 is similar to L_2 , by (46) we have uniformly for any $a \in [k]$

$$S_1 \lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}. \quad (52)$$

To bound S_2 , by (70) in [8], we have uniformly for any $a \in [k]$,

$$S_2 = \frac{\left| \sum_{j=1}^n \mathbb{I}\{z_j^* = a\} - \sum_{j=1}^n \mathbb{I}\{z_j = a\} \right|}{\sum_{j=1}^n \mathbb{I}\{z_j = a\}} \left\| \hat{\Sigma}_a(z^*) \right\|^2 \lesssim \frac{k}{n} \frac{\ell(z, z^*)}{\Delta^2},$$

where we use (43). Since $\frac{k}{n} \frac{\ell(z, z^*)}{\Delta^2} \lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta}$, by (45) and the facts that $\ell(z, z^*) \leq \tau = o(n)$ we have

$$\max_{a \in [k]} \left\| \hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*) \right\| \lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}.$$

□

Lemma 7.8. *Under the same assumption as in Lemma 7.7, if additional we assume $kd = O(\sqrt{n})$ and $\tau = o(n/k)$, there exists some constant $C > 0$ only depending on $\alpha, \lambda_{\min}, \lambda_{\max}, C'$ such that*

$$\max_{a \in [k]} \left\| (\hat{\Sigma}_a(z))^{-1} - (\hat{\Sigma}_a(z^*))^{-1} \right\| \leq C \left(\frac{k}{n}\ell(z, z^*) + \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)} \right). \quad (53)$$

Proof. By (43) we have $\max_{a \in [k]} \|\hat{\Sigma}_a(z^*)\|, \max_{a \in [k]} \|(\hat{\Sigma}_a(z^*))^{-1}\| \lesssim 1$. By (44) we also have $\max_{a \in [k]} \|\hat{\Sigma}_a(z)\|, \max_{a \in [k]} \|(\hat{\Sigma}_a(z))^{-1}\| \lesssim 1$. Hence,

$$\begin{aligned} \max_{a \in [k]} \left\| (\hat{\Sigma}_a(z))^{-1} - (\hat{\Sigma}_a(z^*))^{-1} \right\| &\leq \max_{a \in [k]} \left\| (\hat{\Sigma}_a(z^*))^{-1} \right\| \left\| \hat{\Sigma}_a(z) - \hat{\Sigma}_a(z^*) \right\| \left\| (\hat{\Sigma}_a(z))^{-1} \right\| \\ &\lesssim \frac{k\sqrt{n\ell(z, z^*)}}{n\Delta} + \frac{k}{n}\ell(z, z^*) + \frac{kd}{n\Delta}\sqrt{\ell(z, z^*)}. \end{aligned} \quad (54)$$

□

Lemma 7.9. *Let $W_i \stackrel{iid}{\sim} \chi_d^2$ for any $i \in [n]$ where n, d are positive integers. Then we have*

$$\mathbb{P} \left(\sum_{i=1}^n W_i^2 \geq 3nd^2 \right) \leq \frac{4}{nd}.$$

Proof. We have $\mathbb{E} \sum_{i=1}^n W_i^2 = nd(d+2)$ and $\mathbb{E} \sum_{i=1}^n W_i^4 = nd(d+2)(d+4)(d+6)$. Then we have $\text{Var} \left(\sum_{i=1}^n W_i^2 \right) = 8nd(d+2)(d+3)$. Then we obtain the desired result by Chebyshev's inequality. □

Proof of Lemma 6.3. Consider any $a \neq b \in [k]$. We are going to prove

$$\frac{-\sqrt{\lambda_{\max}} + \sqrt{\lambda_{\max} + \frac{\lambda_{\min}(\lambda_{\min} + \lambda_{\max})}{2\lambda_{\max}}}}{\lambda_{\min} + \lambda_{\max}} \|\theta_a^* - \theta_b^*\| \leq \text{SNR}'_{a,b} \leq \lambda_{\min}^{-1/2} \|\theta_a^* - \theta_b^*\| + \sqrt{\frac{3}{2}d} + \sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}}. \quad (55)$$

We first prove the upper bound. Denote $\Xi_{a,b} = \theta_a^* - \theta_b^*$. Since we have assumed $\text{SNR}' \rightarrow \infty$, we have that $0 \notin \mathcal{B}_{a,b}$. Note that we have an equivalent expression of $\mathcal{B}_{a,b}$:

$$\mathcal{B}_{a,b} = \left\{ -(\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} + y \in \mathbb{R}^d : 2y^T (\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} + y^T \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) y - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \leq 0 \right\}.$$

We consider the following scenarios.

(1). If $\lambda_1 \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) \geq 0$, we have $|\Sigma_a^*| \geq |\Sigma_b^*|$. Let $y = 0$, we can know $-(\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} \in \mathcal{B}_{a,b}$. This tells us $\text{SNR}'_{a,b} \leq \left\| -(\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} \right\| \leq \lambda_{\min}^{-1/2} \|\Xi_{a,b}\|$.

(2). If $\lambda_1 \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) \leq -1$, let $A := \Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I$ and assume $U^T A U = V$, where U is an orthogonal matrix and $V := \text{diag}\{v_1, \dots, v_d\}$ is a diagonal matrix with diagonal elements $v_1 \leq v_2 \leq \dots \leq v_d$ and $v_1 \leq -1$. We can rewrite $y = Uz$ with $z = (z_1, \dots, z_d)^T$ and $U^T (\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} = (\tau_1, \dots, \tau_d)^T$. Since $-\log |\Sigma_a^*| + \log |\Sigma_b^*| \leq d \log \frac{\lambda_{\max}}{\lambda_{\min}}$, we can take $z_1 = -\text{sign}(\tau_1) \sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}}$ and $z_i = 0$ for $i \geq 2$. Then, we have

$$\begin{aligned} &2y^T (\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} + y^T \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) y - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &= 2z_1 \tau_1 + v_1 z_1^2 - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &\leq -2\sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}} |\tau_1| - d \log \frac{\lambda_{\max}}{\lambda_{\min}} - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &\leq 0. \end{aligned}$$

It means $-(\Sigma_a^*)^{-\frac{1}{2}}\Xi_{a,b} + y \notin \mathcal{B}_{a,b}$ and hence $\text{SNR}'_{a,b} \leq \left\| -(\Sigma_a^*)^{-\frac{1}{2}}\Xi_{a,b} \right\| + \|y\|$. Then Thus we have $\text{SNR}'_{a,b} \leq \left\| -(\Sigma_a^*)^{-\frac{1}{2}}\Xi_{a,b} \right\| + \sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}} \leq \lambda_{\min}^{-1/2} \|\Xi_{a,b}\| + \sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}}$.

(3). If $-1 < \lambda_1 \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) < 0$, we still use the notations in scenario (2). Notice that $\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} = A + I$, we have

$$\begin{aligned} \log \frac{|\Sigma_a^*|}{|\Sigma_b^*|} &= \log(1 + v_1) \cdots \cdots (1 + v_d) \\ &\geq d \log(1 + v_1) \\ &\geq \frac{3}{2} d v_1. \end{aligned}$$

Now we take $z_1 = -\text{sign}(\tau_1) \sqrt{\frac{3}{2}d}$ and $z_i = 0$ for $i \geq 2$, then we have

$$\begin{aligned} &2y^T (\Sigma_a^*)^{-\frac{1}{2}} \Xi_{a,b} + y^T \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) y - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &= 2z_1 \tau_1 + v_1 z_1^2 - \log |\Sigma_a^*| + \log |\Sigma_b^*| - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &\leq -2|\tau_1| \sqrt{\frac{3}{2}d} - \Xi_{a,b}^T (\Sigma_a^*)^{-1} \Xi_{a,b} \\ &\leq 0. \end{aligned}$$

Thus we have $\text{SNR}'_{a,b} \leq \left\| -(\Sigma_a^*)^{-\frac{1}{2}}\Xi_{a,b} \right\| + \sqrt{\frac{3}{2}d} \leq \lambda_{\min}^{-1/2} \|\Xi_{a,b}\| + \sqrt{\frac{3}{2}d}$.

Overall, we have $\text{SNR}'_{a,b} \leq \lambda_{\min}^{-1/2} \|\Xi_{a,b}\| + \sqrt{\frac{3}{2}d} + \sqrt{d \log \frac{\lambda_{\max}}{\lambda_{\min}}}$ for all the three scenarios.

To prove the lower bound, we have

$$\begin{aligned} x^T \Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} (\theta_a^* - \theta_b^*) + \frac{1}{2} x^T \left(\Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right) x &\geq - \left\| \Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \right\| \|x\| \|\Xi_{a,b}\| - \frac{1}{2} \|x\|^2 \left\| \Sigma_a^{*\frac{1}{2}} \Sigma_b^{*-1} \Sigma_a^{*\frac{1}{2}} - I \right\| \\ &\geq - \frac{\sqrt{\lambda_{\max}}}{\lambda_{\min}} \|x\| \|\Xi_{a,b}\| - \frac{1}{2} \left(\frac{\lambda_{\max}}{\lambda_{\min}} + 1 \right) \|x\|^2. \end{aligned}$$

By the upper bound we know for any $a \neq b \in [k]$, $\|\Xi_{a,b}\| \rightarrow \infty$ when $\text{SNR}' \rightarrow \infty$. Thus, we have

$$\begin{aligned} \frac{\sqrt{\lambda_{\max}}}{\lambda_{\min}} \|x\| \|\Xi_{a,b}\| + \frac{1}{2} \left(\frac{\lambda_{\max}}{\lambda_{\min}} + 1 \right) \|x\|^2 &\geq \frac{1}{2} \Xi_{a,b}^T \Sigma_b^{*-1} \Xi_{a,b} - \log |\Sigma_a^*| + \log |\Sigma_b^*| \\ &\geq \frac{1}{4\lambda_{\max}} \|\Xi_{a,b}\|^2. \end{aligned}$$

Hence,

$$\|x\| \geq \frac{-\sqrt{\lambda_{\max}} + \sqrt{\lambda_{\max} + \frac{\lambda_{\min}(\lambda_{\min} + \lambda_{\max})}{2\lambda_{\max}}}}{\lambda_{\min} + \lambda_{\max}} \|\Xi_{a,b}\|.$$

□

In the following lemmas, we are going to establish connections between testing errors and $\{\text{SNR}'_{a,b}\}_{a \neq b}$. Consider any $a, b \in [k]$ such that $a \neq b$. Let $\eta \sim \mathcal{N}(0, I_d)$, $\Xi_{a,b} = \theta_a^* - \theta_b^*$, and $\Delta_{a,b} = \|\Xi_{a,b}\|$. Define

$$\begin{aligned} \mathcal{B}_{a,b}(\delta) &= \left\{ x \in \mathbb{R}^d : x^T \Sigma_a^{*\frac{1}{2}} (\Sigma_b^*)^{-1} \Xi_{a,b} + \frac{1}{2} x^T \left(\Sigma_a^{*\frac{1}{2}} (\Sigma_b^*)^{-1} \Sigma_a^{*\frac{1}{2}} - I_d \right) x \right. \\ &\quad \left. \leq -\frac{1-\delta}{2} \Xi_{a,b}^T (\Sigma_b^*)^{-1} \Xi_{a,b} + \frac{1}{2} \log |\Sigma_a^*| - \frac{1}{2} \log |\Sigma_b^*| \right\}, \end{aligned}$$

for any $\delta \in \mathbb{R}$. In addition, we define

$$\text{SNR}'_{a,b}(\delta) = \min_{x \in \mathcal{B}_{a,b}(\delta)} 2 \|x\|,$$

$$\text{and } P_{a,b}(\delta) = \mathbb{P}(\eta \in \mathcal{B}_{a,b}(\delta)).$$

Recall the definitions of $\mathcal{B}_{a,b}$ and $\text{SNR}'_{a,b}$ in Section 3. Then they are a special case of $\mathcal{B}_{a,b}(\delta)$ and $\text{SNR}'_{a,b}(\delta)$ with $\delta = 0$. That is, we have $\mathcal{B}_{a,b} = \mathcal{B}_{a,b}(0)$ and $\text{SNR}'_{a,b} = \text{SNR}'_{a,b}(0)$.

Lemma 7.10. *Assume $d = O(1)$ and $\lambda_{\min} \leq \lambda_1(\Sigma_a^*), \lambda_1(\Sigma_b^*) \leq \lambda_d(\Sigma_a^*), \lambda_d(\Sigma_b^*) \leq \lambda_{\max}$ where $\lambda_{\min}, \lambda_{\max} > 0$ are constants. Under the condition $\text{SNR}'_{a,b} \rightarrow \infty$, for any positive sequence $\delta = o(1)$, there exists a $\tilde{\delta} = o(1)$ that depends on $\delta, d, \Delta_{a,b}, \lambda_{\min}, \lambda_{\max}$ such that*

$$P_{a,b}(\delta) \leq \exp\left(-\frac{1-\tilde{\delta}}{8} \text{SNR}'_{a,b}{}^2\right)$$

Proof. For convenience and conciseness, we will use the notation $\theta_a, \theta_b, \Sigma_a, \Sigma_b$ instead of $\theta_a^*, \theta_b^*, \Sigma_a^*, \Sigma_b^*$ throughout the proof. By Lemma 6.3, we have $\text{SNR}'_{a,b}$ in the same order of $\Delta_{a,b}$, which means $\Delta_{a,b} \rightarrow \infty$.

Assume we had obtained $\text{SNR}'_{a,b}(\delta) \geq (1-o(1))\text{SNR}'_{a,b}$. Then by Lemma 6.3, we have $\text{SNR}'_{a,b}(\delta)$ in the same order of $\Delta_{a,b}$ which is far bigger than d by assumption. Since $\|\eta\|^2 \sim \chi_d^2$, using Lemma 7.1, we have

$$\begin{aligned} P_{a,b}(\delta) &\leq P\left(\|\eta\|^2 \geq \frac{\text{SNR}'_{a,b}{}^2(\delta)}{4}\right) \leq \exp\left(-\left(1 - O\left(\frac{d}{\Delta_{a,b}^2}\right)\right) \frac{\text{SNR}'_{a,b}{}^2(\delta)}{8}\right) \\ &\leq \left(1 - O\left(\frac{d}{\Delta_{a,b}^2}\right) \frac{(1-o(1))\text{SNR}'_{a,b}{}^2}{8}\right) \end{aligned}$$

which is the desired result. Hence, the proof of this lemma is all about establishing $\text{SNR}'_{a,b}(\delta) \geq (1-o(1))\text{SNR}'_{a,b}$.

To prove it, we first simplify $\text{SNR}'_{a,b}(\delta)$. In spite of some abuse of notation, denote $\lambda_1 \leq \dots \leq \lambda_d$ to be the eigenvalues of $\Sigma_a^{\frac{1}{2}} (\Sigma_b)^{-1} \Sigma_a^{\frac{1}{2}} - I_d$ such that its eigen-decomposition can be written as $\Sigma_a^{\frac{1}{2}} (\Sigma_b)^{-1} \Sigma_a^{\frac{1}{2}} - I_d = \sum_{i=1}^d \lambda_i u_i u_i^T$, where $\{u_i\}$ are orthogonal vectors. Denote $U = (u_1, \dots, u_d)$ and

$$v = U^T \Sigma_a^{\frac{1}{2}} \Sigma_b^{-1} \Xi_{a,b}$$

and

$$\mathcal{B}'_{a,b}(\delta) = \left\{ y \in \mathbb{R}^d : \sum_i y_i v_i + \frac{1}{2} \sum_i \lambda_i y_i^2 \leq -\frac{1-\delta}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|} \right\}.$$

Then $\mathcal{B}'_{a,b}(\delta)$ can be seen a reflection-rotation of $\mathcal{B}_{a,b}(\delta)$ by the transformation $y = U^t x$. Hence we have $\text{SNR}'_{a,b}(\delta) = \min_{y \in \mathcal{B}'_{a,b}(\delta)} 2 \|y\|$ for any δ . What is more, let $\bar{\mathcal{B}}'_{a,b}(\delta)$ to be its boundary, i.e.,

$$\bar{\mathcal{B}}'_{a,b}(\delta) = \left\{ y \in \mathbb{R}^d : \sum_i y_i v_i + \frac{1}{2} \sum_i \lambda_i y_i^2 = -\frac{1-\delta}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|} \right\}.$$

Since $0 \notin \mathcal{B}'_{a,b}(\delta)$, we have $\text{SNR}'_{a,b}(\delta) = 2 \min_{y \in \bar{\mathcal{B}}'_{a,b}(\delta)} \|y\|$. As a result, we only need to work on $\bar{\mathcal{B}}'_{a,b}(\delta)$ instead of $\mathcal{B}_{a,b}(\delta)$. Denote $\bar{\mathcal{B}}'_{a,b}$ to be $\bar{\mathcal{B}}'_{a,b}(0)$ for simplicity.

We then give an equivalent expression of $\mathcal{B}'_{a,b}(\delta)$. From (55), we have an upper bound of $\text{SNR}'_{a,b}$: $\text{SNR}'_{a,b} \leq 2\lambda_{\min}^{-1/2} \Delta_{a,b}$ where we use $\Delta_{a,b} \gg \lambda_{\min}, \lambda_{\max}, d$. The same upper bound actually holds for $\mathcal{B}'_{a,b}(\delta)$ for any $\delta = o(1)$ following the same proof. Define $S = \{y \in \mathbb{R}^d : \|y\| \leq 2\lambda_{\min}^{-1/2} \Delta_{a,b}\}$. We then have

$$\text{SNR}'_{a,b}(\delta) = 2 \min_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y\|.$$

We have the following inequality. Let $g(y) : \bar{\mathcal{B}}'_{a,b}(\delta) \rightarrow \bar{\mathcal{B}}'_{a,b}(0)$ be any mapping. By the triangle inequality, we have $\|y\| \geq \|g(y)\| - \|y - g(y)\|$. We have

$$\begin{aligned} 2^{-1} \text{SNR}'_{a,b}(\delta) &= \min_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y\| \\ &\geq \min_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} (\|g(y)\| - \|y - g(y)\|) \\ &\geq \min_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|g(y)\| - \max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\| \\ &\geq \min_{y \in \bar{\mathcal{B}}'_{a,b}(0)} \|y\| - \max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\| \\ &\geq \text{SNR}'_{a,b} - \max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\|. \end{aligned} \quad (56)$$

As a result, if we are able to find some g such that $\max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\| = o(1) \text{SNR}'_{a,b}$, we will immediately have $\text{SNR}'_{a,b}(\delta) \geq (1 - o(1)) \text{SNR}'_{a,b}$ and the proof will be complete.

Let $w \in \mathbb{R}^d$ be some vector. Define $g(y) = y + w \operatorname{argmin}_{t \in \mathbb{R}: y+tw \in \bar{\mathcal{B}}'_{a,b}} |t|$. If $g(y)$ is a well-defined mapping, we have

$$\max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\| = \max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \min_{t \in \mathbb{R}: y+tw \in \bar{\mathcal{B}}'_{a,b}} \|w\| |t|, \quad (57)$$

which can be used to derive an upper bound. However, to make $g(y)$ well-defined, we need for any $y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S$, there exists some $t \in \mathbb{R}$ such that $y + tw \in \bar{\mathcal{B}}'_{a,b}$. This means we have the following two equations:

$$\begin{aligned} \sum_i y_i v_i + \frac{1}{2} \sum_i \lambda_i y_i^2 &= -\frac{1-\delta}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|}, \\ \text{and } \sum_i (y_i + tw_i) v_i + \frac{1}{2} \sum_i \lambda_i (y_i + tw_i)^2 &= -\frac{1}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|}. \end{aligned} \quad (58)$$

It is equivalent to require t to satisfy

$$t \sum_i (w_i v_i + \lambda_i y_i w_i) + \frac{t^2}{2} \sum_i \lambda_i w_i^2 = -\frac{\delta}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}. \quad (59)$$

Hence, all we need is to find a decent vector w such that: for any $y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S$ there exists a t satisfying (59), and we can obtain the desired upper bound for (57).

In the following, we will consider four different scenarios according to the spectral $\{\lambda_i\}$. For each scenario, we will construct a w with decent bounds for (57). Denote $\delta' = \sqrt{\delta}$.

Scenario 1: $|\lambda_1|, |\lambda_d| \leq \delta'$. We choose $w = v/\|v\|$. Note that we have $\|v\|$ in the same order of $\Delta_{a,b}$ and $\|\Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}\|$ in the same order of $\Delta_{a,b}^2$. Note that we have

$$\begin{aligned} t \sum_i (w_i v_i + \lambda_i y_i w_i) + \frac{t^2}{2} \sum_i \lambda_i w_i^2 &\leq t \|v\| + |t| \|y\| \sqrt{\sum_i \lambda_i^2 w_i^2} + \frac{t^2}{2} \sum_i \lambda_i w_i^2 \\ &\leq t \|v\| + |t| \delta' \|y\| + \frac{t^2 \delta'}{2} \\ &\leq t \|v\| + 2 |t| \delta' \lambda_{\min}^{-1/2} \Delta_{a,b} + \frac{t^2 \delta'}{2}, \end{aligned}$$

where in the last inequality we use $y \in S$. Define $t_0 = -\delta^{1/2} \Delta_{a,b}$. Then we have

$$t_0 \sum_i (w_i v_i + \lambda_i y_i w_i) + \frac{t_0^2}{2} \sum_i \lambda_i w_i^2 \lesssim -\delta^{1/2} \Delta_{a,b}^2 + \delta^{1/2} \delta' \Delta_{a,b}^2 + \delta \delta' \Delta_{a,b}^2 \ll -\delta \Delta_{a,b}^2 \lesssim -\frac{\delta}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}.$$

Hence for any $y \in S$ there exists a $t \in (t_0, 0)$ such that (59) is satisfied. Hence, $|t_0| = \delta^{1/2} \Delta_{a,b}$ is an upper bound for (57).

Scenario 2: $\lambda_1 < -\delta'$. We choose $w = e_1$ which is the first standard basis of \mathbb{R}^d . Then, (59) can be written as

$$\lambda_1 t^2 + 2(v_1 + \lambda_1 y_1)t + \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} = 0.$$

Since $\lambda_1 < 0$, the above equation has two different solutions $t_1, t_2 \in \mathbb{R}$. Simple algebra leads to

$$\min\{|t_1|, |t_2|\} \leq \sqrt{\frac{\delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}}{-\lambda_1}} \leq \sqrt{\frac{\delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}}{\delta'}} \lesssim \delta^{1/4} \Delta_{a,b}.$$

Hence, an upper bound for (57) is $O(\delta^{1/4} \Delta_{a,b})$.

Scenario 3: $\lambda_1 \geq -\delta'$ and there exists a $j \in [d]$ such that $\lambda_j \leq \delta'$ and $|v_j| \geq \sqrt{\delta'} \Delta_{a,b}$. We choose $w = e_j$. Then (59) can be written as

$$\lambda_j t^2 + 2(v_j + \lambda_j y_j)t + \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} = 0.$$

Note that for any $y \in S$, we have $|v_j + \lambda_j y_j| \geq |v_j| - |\lambda_j y_j| \geq \sqrt{\delta'} \Delta_{a,b} - \delta'(2\lambda_{\min}^{-1/2} \Delta_{a,b}) \geq \sqrt{\delta'} \Delta_{a,b}/2$. Denote $t_0 = -\text{sign}(v_j + \lambda_j y_j) \sqrt{\delta'} \Delta_{a,b}$. Then we have

$$\begin{aligned} \lambda_j t_0^2 + 2(v_j + \lambda_j y_j)t_0 + \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} &= \lambda_j \delta' \Delta_{a,b}^2 - |v_j + \lambda_j y_j| \sqrt{\delta'} \Delta_{a,b} + \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} \\ &\leq -\left(\frac{\delta'}{2} - \delta'^2\right) \Delta_{a,b}^2 + \delta O(\Delta_{a,b}^2) \\ &\leq 0. \end{aligned}$$

As a result, there exists some $t \in (t_0, 0)$ satisfying (59). Hence, $|t_0| = \delta^{1/2} \Delta_{a,b}$ is an upper bound for (57).

Scenario 4: $\lambda_1 \geq -\delta'$ and $|v_j| < \sqrt{\delta'} \Delta_{a,b}$ for all $j \in [d]$ such that $\lambda_j \leq \delta'$. This scenario is slightly more complicated as we need w to be dependent on y . Denote it as $w(y)$. Then (56) still holds and (57) can be changed into

$$\max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \|y - g(y)\| = \max_{y \in \bar{\mathcal{B}}'_{a,b}(\delta) \cap S} \min_{t \in \mathbb{R}: y+tw \in \bar{\mathcal{B}}'_{a,b}} \|w(y)\| |t|. \quad (60)$$

Denote $m \in [d]$ to be the integer such that $\lambda_j \leq \delta'$ for all $j \leq m$ and $\lambda_j > \delta'$ for all $j > m$. We can have $m < d$ otherwise this scenario can be reduced to Scenario 1. Define

$$[w(y)]_i = - \left(y_i + \frac{v_i}{\lambda_i} \right) \mathbb{I}\{i > m\}.$$

for any $i \in [d]$. Instead of using (59), we will analyze it slightly differently.

For $y \in \bar{\mathcal{B}}'_{a,b}(\delta)$, (58) can be rewritten as

$$\sum_{i>m} \lambda_i \left(y_i + \frac{v_i}{\lambda_i} \right)^2 = \sum_{i>m} \frac{v_i^2}{\lambda_i} - (1 - \delta) \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right). \quad (61)$$

On the other hand, if $g(y)$ is well-defined, we need $g(y) \in \bar{\mathcal{B}}'_{a,b}$ which means

$$\sum_{i>m} \lambda_i \left([g(y)]_i + \frac{v_i}{\lambda_i} \right)^2 = \sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right).$$

Note that we have $(y_i + v_i/\lambda_i)(1-t) = [g(y)]_i + v_i/\lambda_i$ for $i > m$ and $[g(y)]_i = y_i$ for $i \leq m$. Then the above display can be written as

$$(1-t)^2 \sum_{i>m} \lambda_i \left(y_i + \frac{v_i}{\lambda_i} \right)^2 = \sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right).$$

Together with (61) multiplied, the above equation leads to

$$(1-t)^2 \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} = (2t - t^2) \left(\sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right) \right). \quad (62)$$

It is sufficient to find some $0 < t_0 < 1$ such that

$$\frac{(1-t_0)^2}{t_0(2-t_0)} \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} \leq \sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right), \quad (63)$$

then there definitely exists some $0 < t < t_0$ satisfying (62).

We are going to give a lower bound for the right hand side of (63). Particularly, we need to lower bound $\sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}$. Denote $\tilde{y} = U^T(-\Sigma_a^{-1/2} \Xi_{a,b})$. Then using the definition of v and $\{\lambda_i\}$, we have

$$\begin{aligned} 2 \sum_{i \in [k]} \tilde{y}_i v_i + \sum_{i \in [k]} \lambda_i \tilde{y}_i^2 &= 2(-\Sigma_a^{-\frac{1}{2}} \Xi_{a,b})^T \Sigma_a^{\frac{1}{2}} \Sigma_b^{-1} \Xi_{a,b} + (-\Sigma_a^{-\frac{1}{2}} \Xi_{a,b})^T \left(\Sigma_a^{\frac{1}{2}} \Sigma_b^{-1} \Sigma_a^{\frac{1}{2}} - I_d \right) (-\Sigma_a^{-\frac{1}{2}} \Xi_{a,b}) \\ &= -\Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} - \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b}. \end{aligned}$$

Then we have

$$\begin{aligned} \sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} &= \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \sum_{i>m} \lambda_i \left(\tilde{y}_i + \frac{v_i}{\lambda_i} \right)^2 + \left(2 \sum_{i \leq m} \tilde{y}_i v_i + \sum_{i \leq m} \lambda_i \tilde{y}_i^2 \right) \\ &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \left(2 \sum_{i \leq m} \tilde{y}_i v_i + \sum_{i \leq m} \lambda_i \tilde{y}_i^2 \right). \end{aligned} \quad (64)$$

Hence, the right hand side of (63) can be lower bounded by

$$\begin{aligned} &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} + \left(2 \sum_{i \leq m} \tilde{y}_i v_i + \sum_{i \leq m} \lambda_i \tilde{y}_i^2 \right) - \left(2 \sum_{i \leq m} y_i v_i + \sum_{i \leq m} \lambda_i y_i^2 \right) \\ &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - 8 \left(\sqrt{\delta'} \sqrt{d} \lambda_{\min}^{-\frac{1}{2}} \Delta_{a,b}^2 + \delta' \lambda_{\min}^{-1} \Delta_{a,b}^2 \right) \\ &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - 16 \sqrt{\delta'} \sqrt{d} \lambda_{\min}^{-\frac{1}{2}} \Delta_{a,b}^2, \end{aligned} \quad (65)$$

where we use both $\tilde{y}, y \in S$ and the assumption that $|v_i| \leq \sqrt{\delta'} \Delta_{a,b}$ and $|\lambda_i| \leq \delta'$ for any $i \leq m$. Then a sufficient condition for (63) is t_0 satisfies

$$\frac{(1-t_0)^2}{t_0(2-t_0)} \delta \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} \leq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} + 16 \sqrt{\delta'} \sqrt{d} \lambda_{\min}^{-\frac{1}{2}} \Delta_{a,b}^2.$$

Since $\Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b}$ is in the same order of $\Delta_{a,b}^2$ and $\log \frac{|\Sigma_a|}{|\Sigma_b|} \lesssim d = O(1)$, it can be achieved by $t_0 = \sqrt{\delta}$.

As a result, from (60) we have

$$\begin{aligned} \max_{y \in \mathcal{B}'_{a,b}(\delta) \cap S} \|y - g(y)\| &\leq \max_{y \in \mathcal{B}'_{a,b}(\delta) \cap S} \|w(y)\| |t_0| \leq |t_0| \max_{y \in \mathcal{B}'_{a,b}(\delta) \cap S} 2 \sqrt{\|y\|^2 + \frac{\|v\|^2}{\delta'}} \\ &\lesssim \sqrt{\delta} \left(\Delta_{a,b} + \delta'^{-\frac{1}{2}} \Delta_{a,b} \right) \leq 2\delta^{\frac{1}{4}} \Delta_{a,b}. \end{aligned}$$

Combining the above four scenarios, we can see we all have $\max_{y \in \mathcal{B}'_{a,b}(\delta) \cap S} \|y - g(y)\| \lesssim \delta^{\frac{1}{4}} \Delta_{a,b}$ which is $o(1) \text{SNR}'_{a,b}$. By the argument before the discussion of the four scenarios, we have $\text{SNR}'_{a,b}(\delta) \geq (1 - o(1)) \text{SNR}'_{a,b}$ and the proof is complete. \square

Lemma 7.11. Assume $d = O(1)$ and $\lambda_{\min} \leq \lambda_1(\Sigma_a^*), \lambda_1(\Sigma_b^*) \leq \lambda_d(\Sigma_a^*), \lambda_d(\Sigma_b^*) \leq \lambda_{\max}$ where $\lambda_{\min}, \lambda_{\max} > 0$ are constants. Under the condition $\text{SNR}'_{a,b} \rightarrow \infty$, for any positive sequence $\delta = o(1)$, there exists a $\tilde{\delta} = o(1)$ that depends on $d, \Delta_{a,b}, \lambda_{\min}, \lambda_{\max}$ such that

$$P_{1,2}(0) \geq \exp\left(-\frac{1+\tilde{\delta}}{8}\text{SNR}'_{a,b}\right).$$

Proof. For convenience and conciseness, we will use the notation $\theta_a, \theta_b, \Sigma_a, \Sigma_b$ instead of $\theta_a^*, \theta_b^*, \Sigma_a^*, \Sigma_b^*$ throughout the proof. From Lemma 6.3, we know $\text{SNR}'_{a,b}$ is in the same order of $\Delta_{a,b}$, which means $\Delta_{a,b} \rightarrow \infty$. Similar to the proof of Lemma 7.10, denote $\lambda_1 \leq \dots \leq \lambda_d$ to be the eigenvalues of $\Sigma_a^{\frac{1}{2}}(\Sigma_b)^{-1}\Sigma_a^{\frac{1}{2}} - I_d$ such that its eigen-decomposition can be written as $\Sigma_a^{\frac{1}{2}}(\Sigma_b)^{-1}\Sigma_a^{\frac{1}{2}} - I_d = \sum_{i=1}^d \lambda_i u_i u_i^T$, where $\{u_i\}$ are orthogonal vectors. Denote $U = (u_1, \dots, u_d)$, $v = U^T \Sigma_a^{\frac{1}{2}} \Sigma_b^{-1} \Xi_{a,b}$. Then denote

$$\mathcal{B}'_{a,b} = \left\{ y \in \mathbb{R}^d : \sum_i y_i v_i + \frac{1}{2} \sum_i \lambda_i y_i^2 \leq -\frac{1}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|} \right\},$$

and its boundary

$$\bar{\mathcal{B}}'_{a,b} = \left\{ y \in \mathbb{R}^d : \sum_i y_i v_i + \frac{1}{2} \sum_i \lambda_i y_i^2 = -\frac{1}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|} \right\}.$$

By the same argument as in the proof of Lemma 7.10, $\mathcal{B}'_{a,b}$ can be seen a reflection-rotation of $\mathcal{B}_{a,b}$ by the transformation $y = U^t x$. Hence we have $\text{SNR}'_{a,b} = \min_{y \in \mathcal{B}'_{a,b}} 2 \|y\|$ and we can work on $\mathcal{B}'_{a,b}$ instead of $\mathcal{B}_{a,b}$. Denote $\bar{y} \in \mathcal{B}'_{a,b}$ to be the one such that $2 \|\bar{y}\| = \text{SNR}'_{a,b}$. From the proof of Lemma 7.10 we also know $\bar{y} \in S$ which is defined as $S = \{y \in \mathbb{R}^d : \|y\| \leq 2\lambda_{\min}^{-1/2} \Delta_{a,b}\}$. In addition, we know $\bar{y} \in \bar{\mathcal{B}}'_{a,b}$.

We first give the main idea of the remaining proof. Denote $p(y)$ to be the density function of $y \sim \mathcal{N}(0, I_d)$. We will construct a set $T \subset \mathbb{R}^d$ around \bar{y} such that for any $y \in T$ we have $y \in \mathcal{B}'_{a,b}$ and $\|y - \bar{y}\| = o(\Delta_{a,b})$. Then we have

$$\begin{aligned} P_{1,2}(0) &\geq |T| \inf_{y \in T} p(y) = |T| \frac{1}{(2\pi)^{\frac{d}{2}}} \exp\left(-\frac{1}{2} \max_{y \in T} \|y\|^2\right) \\ &= |T| \frac{1}{(2\pi)^{\frac{d}{2}}} \exp\left(-(1+o(1)) \frac{\text{SNR}'_{a,b}}{8}\right). \end{aligned} \quad (66)$$

Hence if $\log |T| = o(\text{SNR}'_{a,b})$ then the proof will be complete. So it is all about constructing such T . We will consider four scenarios same as in the proof of Lemma 7.10. Let $\delta = o(1)$ be some positive sequence going to 0 very slowly and denote $\delta' = \sqrt{\delta}$.

Scenario 1: $|\lambda_1|, |\lambda_d| \leq \delta'$. Define $w = v/\|v\|$. We define T as follows:

$$T = \{y = \bar{y} + s : \|(I_d - w w^T)s\| \leq \delta' |w^T s|, w^T s \in [-\delta \Delta_{a,b}, 0]\}.$$

Since $\bar{y} \in \bar{\mathcal{B}}'_{a,b}$, we have

$$\sum_i \bar{y}_i v_i + \frac{1}{2} \sum_i \lambda_i \bar{y}_i^2 = -\frac{1}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|}. \quad (67)$$

It is obvious $\max_{y \in T} \|y - \bar{y}\| \leq 2\delta\Delta_{a,b}$. Hence we only need to show that for any $y \in T$, $y \in \mathcal{B}'_{a,b}$, i.e.,

$$\sum_i (\bar{y}_i + s_i)v_i + \frac{1}{2} \sum_i \lambda_i (\bar{y}_i + s_i)^2 \leq -\frac{1}{2} \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \frac{1}{2} \log \frac{|\Sigma_a|}{|\Sigma_b|}. \quad (68)$$

From the above two displays, we need to show

$$2 \sum_i s_i v_i + \sum_i \lambda_i s_i^2 + 2 \sum_i \lambda_i \bar{y}_i s_i \leq 0. \quad (69)$$

Note that s satisfies $\|s\| \leq 2|w^T s|$.

$$\begin{aligned} 2 \sum_i s_i v_i + \sum_i \lambda_i s_i^2 + 2 \sum_i \lambda_i \bar{y}_i s_i &\leq 2\|v\| |w^T s| + \delta' \|s\|^2 + \delta' \|\bar{y}\| \|s\| \\ &\leq 2\|v\| |w^T s| + \delta' |w^T s|^2 + \delta' \|\bar{y}\| |w^T s| \\ &= |w^T s| (-2\|v\| + \delta' |w^T s| + \delta' \|\bar{y}\|) \\ &\leq 0, \end{aligned}$$

where we use the fact that $\|v\|, \|\bar{y}\|$ are in the order of $\Delta_{a,b}$. Hence, for any $y \in T$, we have shown $y \in \mathcal{B}'_{a,b}$. From Lemma 7.12, we have $|T| \geq \exp\left(d \log \frac{\delta \delta' \Delta_{a,b}}{4} - \frac{d}{2} \log d\right)$. Since $d = O(1)$, $\Delta_{a,b} \rightarrow \infty$, and δ goes to 0 slowly, (66) leads to $P_{1,2}(0) \geq \exp\left(- (1 + o(1)) \frac{\text{SNR}'_{a,b}}{8}\right)$.

Scenario 2: $\lambda_1 < -\delta'$. Denote e_1 the first standard basis in \mathbb{R}^d . Define T as

$$T = \left\{ y = \bar{y} + s : \|(I_d - e_1 e_1^T) s\| \leq \delta |e_1^T s|, \text{sign}(v_1 + \lambda_1 \bar{y}_1) e_1^T s \in [-2\delta^{\frac{1}{4}} \Delta_{a,b}, -\delta^{\frac{1}{4}} \Delta_{a,b}] \right\}.$$

Here for the sign function we define $\text{sign}(0) = 1$. It is obvious $\max_{y \in T} \|y - \bar{y}\| \leq 2\delta\Delta_{a,b}$. Hence we only need to establish (69) to show that for any $y \in T$, $y \in \mathcal{B}'_{a,b}$. Note that

$$\begin{aligned} 2 \sum_i s_i v_i + \sum_i \lambda_i s_i^2 + 2 \sum_i \lambda_i \bar{y}_i s_i &= 2s_1(v_1 + \lambda_1 \bar{y}_1) + \lambda_1 s^2 + 2 \sum_{i \geq 2} s_i (v_i + \lambda_i \bar{y}_i) + 2 \sum_{i \geq 2} \lambda_i s_i^2 \\ &\leq 2s_1(v_1 + \lambda_1 \bar{y}_1) + \lambda_1 s^2 + 2\|(I - e_1 e_1^T) s\| \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) + 2 \max_j |\lambda_j| \|(I - e_1 e_1^T) s\|^2 \\ &\leq \left(\lambda_1 + \delta^2 \max_j |\lambda_j| \right) s_1^2 - 2|v_1 + \lambda_1 \bar{y}_1| |s_1| + 2 \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \delta |s_1| \\ &\leq -\frac{\delta'}{2} s_1^2 + O(\Delta_{a,b}) \delta |s_1|, \end{aligned}$$

where we use $\max_j |\lambda_j| = O(1)$ and $\|v\|, \|\bar{y}\|$ are in the order of $\Delta_{a,b}$. It is easy to verify the right hand side is negative when $s_1 \in [-2\delta^{\frac{1}{4}} \Delta_{a,b}, -\delta^{\frac{1}{4}} \Delta_{a,b}]$. From Lemma 7.12, we have $|T| \geq \exp\left(d \log \frac{\delta^{\frac{5}{4}} \Delta_{a,b}}{4} - \frac{d}{2} \log d\right)$. Then (66) leads to the desired result.

Scenario 3: $\lambda_1 \geq -\delta'$ and there exists a $j \in [d]$ such that $\lambda_j \leq \delta'$ and $|v_j| \geq \sqrt{\delta'} \Delta_{a,b}$. Denote e_j the j th standard basis in \mathbb{R}^d . Define T as

$$T = \left\{ y = \bar{y} + s : \|(I_d - e_j e_j^T) s\| \leq \delta' |e_j^T s|, \text{sign}(v_j + \lambda_j \bar{y}_j) e_j^T s \in [-\delta \Delta_{a,b}, 0] \right\}.$$

Again define $\text{sign}(0) = 1$ and it is obvious $\max_{y \in T} \|y - \bar{y}\| \leq 2\delta\Delta_{a,b}$. Now we are going to verify (69), i.e., to show $\lambda_j s_j^2 + 2s_j(v_j + \lambda_j \bar{y}_j) \leq -\sum_{i \neq j} \lambda_i s_i^2 - 2\sum_{i \neq j} s_i(v_i + \lambda_i \bar{y}_i)$. On one hand, we have

$$\begin{aligned} \lambda_j s_j^2 + 2s_j(v_j + \lambda_j \bar{y}_j) &= \lambda_j s_j^2 - 2|s_j| |v_j + \lambda_j \bar{y}_j| \\ &\leq -\delta' \delta \Delta_{a,b} |s_j| - 2(\sqrt{\delta'} \Delta_{a,b} - \delta' O(\Delta_{a,b})) |s_j| \\ &\leq -\sqrt{\delta'} \Delta_{a,b} |s_j|. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} -\sum_{i \neq j} \lambda_i s_i^2 - 2\sum_{i \neq j} s_i(v_i + \lambda_i \bar{y}_i) &\geq -\max_j |\lambda_j| \|(I - e_j e_j^T) s\|^2 - 2\|(I - e_j e_j^T) s\| \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \\ &\geq -\max_j |\lambda_j| \delta' 2 |s_j|^2 - 2\delta' |s_j| \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \\ &\geq -2\delta' \left(\delta \Delta_{a,b} \max_j |\lambda_j| + \|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) |s_j| \\ &\geq -2\delta' O(\Delta_{a,b}) |s_j| \\ &\geq -\sqrt{\delta'} \Delta_{a,b} |s_j|, \end{aligned}$$

we use $\max_j |\lambda_j| = O(1)$ and $\|v\|, \|\bar{y}\|$ are in the order of $\Delta_{a,b}$. Hence (69) is established. From Lemma 7.12, we have $|T| \geq \exp\left(d \log \frac{\delta \delta' \Delta_{a,b}}{4} - \frac{d}{2} \log d\right)$. Then (66) leads to the desired result.

Scenario 4: $\lambda_1 \geq -\delta'$ and $|v_j| < \sqrt{\delta'} \Delta_{a,b}$ for all $j \in [d]$ such that $\lambda_j \leq \delta'$. Denote $m \in [d]$ to be the integer such that $\lambda_j \leq \delta'$ for all $j \leq m$ and $\lambda_j > \delta'$ for all $j > m$. We can have $m < k$ otherwise this scenario can be reduced to Scenario 1.

Define $w \in \mathbb{R}^d$ to be unit vector such that

$$w_i = \begin{cases} \frac{\lambda_i \bar{y}_i + v_i}{\sqrt{\sum_{j>m} (\lambda_j \bar{y}_j + v_j)^2}}, & \text{for all } i > m, \\ 0, & \text{o.w..} \end{cases}$$

Define

$$T = \{y = \bar{y} + s : \|(I_d - w w^T) s\| \leq \delta' |w^T s|, w^T s \in [-\delta \Delta_{a,b}, 0]\}.$$

Now we are going to verify (69), i.e., to show $2\sum_{i>m} s_i(v_i + \lambda_i \bar{y}_i) + \sum_i \lambda_i s_i^2 + 2\sum_{i \leq m} s_i(v_i + \lambda_i \bar{y}_i) \leq 0$. On one hand, we have

$$\begin{aligned} 2\sum_{i>m} s_i(v_i + \lambda_i \bar{y}_i) &= 2\sqrt{\sum_{j>m} (v_j + \lambda_j \bar{y}_j)^2} \sum_{i>m} s_i w_i = -2\sqrt{\sum_{j>m} (v_j + \lambda_j \bar{y}_j)^2} |w^T s| \\ &\leq -2\sqrt{\delta'} \sqrt{\sum_{j>m} \left(\bar{y}_j + \frac{v_j}{\lambda_j}\right)^2} |w^T s|. \end{aligned}$$

We are going to give a lower bound for $\sum_{j>m} \left(\bar{y}_j + \frac{v_j}{\lambda_j}\right)^2$. Note that (67) can be written as

$$\sum_{i>m} \lambda_i \left(\bar{y}_i + \frac{v_i}{\lambda_i}\right)^2 = \sum_{i>m} \frac{v_i^2}{\lambda_i} - \Xi_{a,b}^T \Sigma_b^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2\sum_{i \leq m} \bar{y}_i v_i + \sum_{i \leq m} \lambda_i \bar{y}_i^2\right).$$

Denote $\tilde{y} = U^T(-\Sigma_a^{-1/2}\Xi_{a,b})$. Using (64), we have

$$\begin{aligned} \sum_{i>m} \lambda_i \left(\bar{y}_i + \frac{v_i}{\lambda_i} \right)^2 &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \left(2 \sum_{i \leq m} \tilde{y}_i v_i + \sum_{i \leq m} \lambda_i \tilde{y}_i^2 \right) + \log \frac{|\Sigma_a|}{|\Sigma_b|} - \left(2 \sum_{i \leq m} \bar{y}_i v_i + \sum_{i \leq m} \lambda_i \bar{y}_i^2 \right) \\ &\geq \Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b} + \log \frac{|\Sigma_a|}{|\Sigma_b|} - 16\sqrt{\delta' d} \lambda_{\min}^{-\frac{1}{2}} \Delta_{a,b}^2, \\ &\geq C \Delta_{a,b}^2, \end{aligned}$$

for some constant $C > 0$. Here the second inequality is by the same argument as (65) and the last inequality uses the fact that $\Xi_{a,b}^T \Sigma_a^{-1} \Xi_{a,b}$ is in the order of $\Delta_{a,b}^2$ and $d = O(1)$. Hence,

$$\sum_{i>m} \lambda_i \left(\bar{y}_i + \frac{v_i}{\lambda_i} \right)^2 \leq -2\sqrt{C\delta'} \Delta_{a,b} |w^T s|.$$

On the other hand, we have

$$\begin{aligned} \sum_i \lambda_i s_i^2 + 2 \sum_{i \leq m} s_i (v_i + \lambda_i \bar{y}_i) &\leq \max_j |\lambda_j| \|s\|^2 + 2 \sqrt{\sum_{i \leq m} s_i^2} \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \\ &\leq 2 \max_j |\lambda_j| |w^T s|^2 + 2 \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \|(I - ww^T)s\| \\ &\leq 2 \left(\max_j |\lambda_j| \delta \Delta_{a,b} + \delta' \left(\|v\| + \max_j |\lambda_j| \|\bar{y}\| \right) \right) |w^T s| \\ &\leq O(\delta' \Delta_{a,b}) |w^T s|, \end{aligned}$$

where we use the properties of s as $y \in T$. Summing the above two displays together, we have (69) satisfied. From Lemma 7.12, we have $|T| \geq \exp\left(d \log \frac{\delta \delta' \Delta_{a,b}}{4} - \frac{d}{2} \log d\right)$. Then (66) leads to the desired result. \square

Lemma 7.12. *Consider any positive integer d and any $0 < r < 1$, $t > 0$. Define a set $T = \left\{ y \in \mathbb{R}^d : \left(\sum_{i \geq 2} y_i^2 \right)^{1/2} \leq r |y_1|, y_1 \in [-2t, -t] \right\}$. Then we have*

$$|T| \geq \exp\left(d \log \frac{rt}{2} - \frac{d}{2} \log d\right).$$

Proof. Define a d -dimensional ball $B = \left\{ y \in \mathbb{R}^d : (y_1 + 1.5t)^2 + \sum_{i \geq 2} y_i^2 \leq (rt/2)^2 \right\}$. We can easily verify that $B \in T$. First, for all $y \in B$, we have $y_1 \in [-2t, -t]$ as $r \in (0, 1)$. Then, we have $\left(\sum_{i \geq 2} y_i^2 \right)^{1/2} \leq rt/2 \leq r |y_1|$. As a result, by the expression of the volume of a d -dimensional ball, we have

$$|T| \geq |B| = \frac{\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2} + 1)} \left(\frac{rt}{2} \right)^d \geq \frac{1}{d^{\frac{d}{2}}} \left(\frac{rt}{2} \right)^d = \exp\left(d \log \frac{rt}{2} - \frac{d}{2} \log d\right),$$

where $\Gamma(\cdot)$ is the Gamma function. \square

References

- [1] E. Abbe, J. Fan, and K. Wang. An ℓ_p -theory of PCA and spectral clustering. *arxiv preprint*, 2020.
- [2] Christopher M Bishop. *Pattern recognition and machine learning*. springer, 2006.
- [3] S Charles Brubaker and Santosh S Vempala. Isotropic pca and affine-invariant clustering. In *Building Bridges*, pages 241–281. Springer, 2008.
- [4] Sanjoy Dasgupta. *The hardness of k-means clustering*. Department of Computer Science and Engineering, University of California . . . , 2008.
- [5] Arthur P Dempster, Nan M Laird, and Donald B Rubin. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society: Series B (Methodological)*, 39(1):1–22, 1977.
- [6] Yingjie Fei and Yudong Chen. Hidden integrality of sdp relaxation for sub-gaussian mixture models. *arXiv preprint arXiv:1803.06510*, 2018.
- [7] Jerome Friedman, Trevor Hastie, and Robert Tibshirani. *The elements of statistical learning*, volume 1. Springer series in statistics New York, 2001.
- [8] Chao Gao and Anderson Y Zhang. Iterative algorithm for discrete structure recovery. *arXiv preprint arXiv:1911.01018*, 2019.
- [9] Christophe Giraud and Nicolas Verzelen. Partial recovery bounds for clustering with the relaxed k means. *arXiv preprint arXiv:1807.07547*, 2018.
- [10] Daniel Hsu, Sham Kakade, Tong Zhang, et al. A tail inequality for quadratic forms of sub-gaussian random vectors. *Electronic Communications in Probability*, 17, 2012.
- [11] Adam Tauman Kalai, Ankur Moitra, and Gregory Valiant. Efficiently learning mixtures of two gaussians. In *Proceedings of the forty-second ACM symposium on Theory of computing*, pages 553–562, 2010.
- [12] Beatrice Laurent and Pascal Massart. Adaptive estimation of a quadratic functional by model selection. *Annals of Statistics*, pages 1302–1338, 2000.
- [13] Stuart Lloyd. Least squares quantization in pcm. *IEEE transactions on information theory*, 28(2):129–137, 1982.
- [14] Matthias Löffler, Anderson Y Zhang, and Harrison H Zhou. Optimality of spectral clustering for gaussian mixture model. *arXiv preprint arXiv:1911.00538*, 2019.
- [15] Yu Lu and Harrison H Zhou. Statistical and computational guarantees of lloyd’s algorithm and its variants. *arXiv preprint arXiv:1612.02099*, 2016.
- [16] James MacQueen et al. Some methods for classification and analysis of multivariate observations. 1967.
- [17] M. Ndaoud. Sharp optimal recovery in the two component gaussian mixture model. *arXiv preprint*, 2019.

-
- [18] Karl Pearson. Contributions to the mathematical theory of evolution. *Philosophical Transactions of the Royal Society of London. A*, 185:71–110, 1894.
- [19] Daniel A Spielman and Shang-Hua Teng. Spectral partitioning works: Planar graphs and finite element meshes. In *Proceedings of 37th Conference on Foundations of Computer Science*, pages 96–105. IEEE, 1996.
- [20] D Michael Titterton, Adrian FM Smith, and Udi E Makov. *Statistical analysis of finite mixture distributions*. Wiley,, 1985.
- [21] S. Vempala and G. Wang. A spectral algorithm for learning mixture models. *J. Comput. Syst. Sci.*, 68(4):841–860, 2004.
- [22] Ulrike Von Luxburg. A tutorial on spectral clustering. *Statistics and computing*, 17(4):395–416, 2007.
- [23] Kaizheng Wang, Yuling Yan, and Mateo Diaz. Efficient clustering for stretched mixtures: Landscape and optimality. *arXiv preprint arXiv:2003.09960*, 2020.
- [24] Xindong Wu, Vipin Kumar, J Ross Quinlan, Joydeep Ghosh, Qiang Yang, Hiroshi Motoda, Geoffrey J McLachlan, Angus Ng, Bing Liu, S Yu Philip, et al. Top 10 algorithms in data mining. *Knowledge and information systems*, 14(1):1–37, 2008.