

# THE ULTRAMETRIC GROMOV-WASSERSTEIN DISTANCE

FACUNDO MÉMOLI, AXEL MUNK, ZHENGCHAO WAN, AND CHRISTOPH WEITKAMP

**ABSTRACT.** In this paper, we investigate compact ultrametric measure spaces which form a subset  $\mathcal{U}^w$  of the collection of all metric measure spaces  $\mathcal{M}^w$ . In analogy with the notion of the ultrametric Gromov-Hausdorff distance on the collection of ultrametric spaces  $\mathcal{U}$ , we define ultrametric versions of two metrics on  $\mathcal{U}^w$ , namely of Sturm's Gromov-Wasserstein distance of order  $p$  and of the Gromov-Wasserstein distance of order  $p$ . We study the basic topological and geometric properties of these distances as well as their relation and derive for  $p = \infty$  a polynomial time algorithm for their calculation. Further, several lower bounds for both distances are derived and some of our results are generalized to the case of finite ultra-dissimilarity spaces. Finally, we study the relation between the Gromov-Wasserstein distance and its ultrametric version (as well as the relation between the corresponding lower bounds) in simulations and apply our findings for phylogenetic tree shape comparisons.

## 1. INTRODUCTION

Over the last decade the acquisition of ever more complex data, structures and shapes has increased dramatically. Consequently, the need to develop meaningful methods for comparing general objects has become more and more apparent. In numerous applications, e.g. in molecular biology [17, 43, 54], computer vision [45, 61] and electrical engineering [55, 77], it is important to distinguish between different objects in a pose invariant manner: two instances of the a given object in *different* spatial orientations are deemed to be equal. Furthermore, also the comparisons of graphs, trees, ultrametric spaces and networks, where mainly the underlying connectivity structure matters, have grown in importance [21, 29]. One possibility to compare two general objects in a pose invariant manner is to model them as metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  and regard them as elements of the collection of isometry classes of compact metric spaces denoted by  $\mathcal{M}$  (i.e. two compact metric spaces  $(X, d_X)$  and  $(Y, d_Y)$  are in the same class if and only if they are isometric to each other which we denote by  $X \cong Y$ ). It is possible to compare  $(X, d_X)$  and  $(Y, d_Y)$  via the *Gromov-Hausdorff distance* [32, 41], which is a metric on  $\mathcal{M}$ . It is defined as

$$d_{\text{GH}}(X, Y) := \inf_{Z, \phi, \psi} d_{\text{H}}^{(Z, d_Z)}(\phi(X), \psi(Y)), \quad (1)$$

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings into a metric space  $(Z, d_Z)$  and  $d_{\text{H}}^{(Z, d_Z)}$  denotes the *Hausdorff distance in  $Z$* . The Hausdorff distance is a metric on the collection of compact subsets of a metric space  $(Z, d_Z)$ , which is denoted by  $\mathcal{S}(Z)$ , and for  $A, B \in \mathcal{S}(Z)$  defined as follows

$$d_{\text{H}}^{(Z, d_Z)}(A, B) := \max \left( \sup_{a \in A} \inf_{b \in B} d_Z(a, b), \sup_{b \in B} \inf_{a \in A} d_Z(a, b) \right). \quad (2)$$

While the Gromov-Hausdorff distance has been applied successfully for various shape and data analysis tasks (see e.g. [12, 13, 14, 15, 16, 19, 20, 69]), it turns out that it is generally convenient to equip the modelled objects with more structure and to model them as

*metric measure spaces* [66, 67]. A metric measure space  $\mathcal{X} = (X, d_X, \mu_X)$  is a triple, where  $(X, d_X)$  denotes a metric space and  $\mu_X$  stands for a Borel probability measure on  $X$  with full support. This additional probability measure can be thought of as signalling the importance of different regions in the modelled object. Moreover, two metric measure spaces  $\mathcal{X} = (X, d_X, \mu_X)$  and  $\mathcal{Y} = (Y, d_Y, \mu_Y)$  are considered as isomorphic (denoted by  $\mathcal{X} \cong_w \mathcal{Y}$ ) if and only if there exists an isometry  $\varphi : (X, d_X) \rightarrow (Y, d_Y)$  such that  $\varphi_{\#}\mu_X = \mu_Y$ . Here,  $\varphi_{\#}$  denotes the pushforward map induced by  $\varphi$ . From now on,  $\mathcal{M}^w$  denotes the collection of all (isomorphism classes of) compact metric measure spaces.

The additional structure of the metric measure spaces allows to regard the modelled objects as probability measures instead of compact sets. Hence, it is possible to substitute the Hausdorff component in Equation (1) by a relaxed notion of proximity, namely the *Wasserstein distance*. This distance is fundamental to a variety of mathematical developments and is also known as Kantorovich distance [47], Kantorovich-Rubinstein distance [48], Mallows distance [63] or as the Earth Mover's distance [85]. Given a compact metric space  $(Z, d_Z)$ , let  $\mathcal{P}(Z)$  denote the space of probability measures on  $Z$  and let  $\alpha, \beta \in \mathcal{P}(Z)$ . Then, the Wasserstein distance of order  $p$ , for  $1 \leq p < \infty$ , between  $\alpha$  and  $\beta$  is defined as

$$d_{W,p}^{(Z,d_Z)}(\alpha, \beta) := \left( \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \int_{Z \times Z} d_Z^p(x, y) \mu(dx \times dy) \right)^{\frac{1}{p}}, \quad (3)$$

and for  $p = \infty$  as

$$d_{W,\infty}^{(Z,d_Z)}(\alpha, \beta) := \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \sup_{(x,y) \in \text{supp}(\mu)} d_Z(x, y), \quad (4)$$

where  $\text{supp}(\mu)$  stands for the support of  $\mu$  and  $\mathcal{C}(\alpha, \beta)$  denotes the set of all couplings of  $\alpha$  and  $\beta$ , i.e., the set of all probability measures  $\mu$  on the product space  $Z \times Z$  such that

$$\mu(A \times Z) = \alpha(A) \quad \text{and} \quad \mu(Z \times B) = \beta(B)$$

for all Borel measurable sets  $A$  and  $B$  of  $Z$ . It is worth noting that the Wasserstein distance between probability measures on the real line admits a closed form solution (see [99] and Remark 2.12).

Sturm [92] has shown that replacing the Hausdorff distance in Equation (1) with the Wasserstein distance indeed yields a meaningful metric on  $\mathcal{M}^w$ . Let  $\mathcal{X} = (X, d_X, \mu_X)$  and  $\mathcal{Y} = (Y, d_Y, \mu_Y)$  be two metric measure spaces. Then, *Sturm's Gromov-Wasserstein distance* of order  $p$ ,  $1 \leq p \leq \infty$ , is defined as

$$d_{GW,p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) := \inf_{Z, \phi, \psi} d_{W,p}^{(Z,d_Z)}(\phi_{\#}\mu_X, \psi_{\#}\mu_Y), \quad (5)$$

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings into the metric space  $(Z, d_Z)$ .

Based on similar ideas but starting from a different representation of the Gromov-Hausdorff distance, Mémoli [66, 67] derived a computationally more tractable and topologically equivalent metric on  $\mathcal{M}^w$ , namely the *Gromov-Wasserstein distance*: For  $1 \leq p < \infty$ , the *p-distortion* of a coupling  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  is defined as

$$\text{dis}_p(\mu) := \left( \iint_{X \times Y \times X \times Y} |d_X(x, x') - d_Y(y, y')|^p \mu(dx \times dy) \mu(dx' \times dy') \right)^{1/p} \quad (6)$$

and for  $p = \infty$  it is given as

$$\text{dis}_\infty(\mu) := \sup_{\substack{x, x' \in \mathcal{X}, y, y' \in \mathcal{Y} \\ \text{s.t. } (x, y), (x', y') \in \text{supp}(\mu)}} |d_X(x, x') - d_Y(y, y')|.$$

The *Gromov-Wasserstein distance* of order  $p$ ,  $1 \leq p \leq \infty$ , is defined as

$$d_{\text{GW}, p}(\mathcal{X}, \mathcal{Y}) := \frac{1}{2} \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \text{dis}_p(\mu). \quad (7)$$

It is known that in general  $d_{\text{GW}, p} \leq d_{\text{GW}, p}^{\text{sturm}}$  and that the inequality can be strict [67]. Although both  $d_{\text{GW}, p}^{\text{sturm}}$  and  $d_{\text{GW}, p}$ ,  $1 \leq p \leq \infty$ , are in general NP-hard to compute [67], it is possible to efficiently approximate  $d_{\text{GW}, p}$  via conditional gradient descent [67, 79]. This has led to numerous applications and extensions of this distance [4, 18, 24, 87, 95].

In many cases, since the direct computation of either of these distances can be onerous, the determination of the degree of similarity between two datasets is performed via firstly computing *invariant features* out of each dataset (e.g. global distance distributions [75]) and secondly by suitably comparing these features. This point of view has motivated the exploration of inverse problems arising from the study of such features [11, 67, 68, 93].

Clearly,  $\mathcal{M}^w$  contains various, extremely general spaces. However, in many applications it is possible to have prior knowledge about the metric measure spaces under consideration and it is often reasonable to restrict oneself to work on a specific sub-collections  $\mathcal{O}^w \subseteq \mathcal{M}^w$ . For instance, it could be known that the metrics of the spaces considered are induced by the shortest path metric on some underlying trees and hence it is unnecessary to consider the calculation of  $d_{\text{GW}, p}^{\text{sturm}}$  and  $d_{\text{GW}, p}$ ,  $1 \leq p \leq \infty$ , for all of  $\mathcal{M}^w$ . The potential advantages of focusing on a specific sub-collection  $\mathcal{O}^w$  are twofold. On the one hand, it might be possible to use the features of  $\mathcal{O}^w$  to gain computational benefits. On the other hand, it might be possible to refine the definition  $d_{\text{GW}, p}^{\text{sturm}}$  and  $d_{\text{GW}, p}$ ,  $1 \leq p \leq \infty$ , to obtain more informative comparisons on  $\mathcal{O}^w$ . Naturally, it is of interest to identify and study these subclasses and the corresponding refinements. This approach has been pursued to study (variants of) the Gromov-Hausdorff distance on compact *ultrametric spaces* by Zarichnyi [105] and Qiu [80], and on compact *p-metric spaces* by Mémoli et al. [70]. Here, the metric space  $(X, d_X)$  is called a *p-metric space* ( $1 \leq p < \infty$ ), if for all  $x, x', x'' \in X$  it holds

$$d_X(x, x'') \leq (d_X(x, x')^p + d_X(x', x'')^p)^{1/p}.$$

Further, the metric space  $(X, u_X)$  is called an *ultrametric space*, if  $u_X$  fulfills for all  $x, x', x'' \in X$  that

$$u_X(x', x'') \leq \max(u_X(x, x'), u_X(x', x'')). \quad (8)$$

In particular, note that ultrametrics can be considered as the limiting case of *p-metrics* as  $p \rightarrow \infty$ . In particular, Mémoli et al. [70] derived a polynomial time algorithm for the calculation of the *ultrametric Gromov-Hausdorff distance*  $u_{\text{GH}}$  between two compact ultrametric spaces  $(X, u_X)$  and  $(Y, u_Y)$  (see Section 2.2), which is defined as

$$u_{\text{GH}}(X, Y) := \inf_{Z, \phi, \psi} d_{\text{H}}^{(Z, u_Z)}(\phi(X), \psi(Y)), \quad (9)$$

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings into a common *ultrametric space*  $(Z, u_Z)$  and  $d_{\text{H}}^{(Z, u_Z)}$  denotes the Hausdorff distance on  $Z$ .

A further motivation to study (surrogates of) the distances  $d_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}$  restricted on a subset  $\mathcal{O}^w$  comes from the idea of *slicing* which originated as a method to efficiently estimate the Wasserstein distance  $d_{\text{W},p}^{\mathbb{R}^d}(\alpha, \beta)$  between probability measures  $\alpha$  and  $\beta$  supported in a high dimensional euclidean space  $\mathbb{R}^d$  [85]. The original idea is that given any line  $\ell$  in  $\mathbb{R}^d$  one first obtains  $\alpha_\ell$  and  $\beta_\ell$ , the respective pushforwards of  $\alpha$  and  $\beta$  under the orthogonal projection map  $\pi_\ell : \mathbb{R}^d \rightarrow \ell$ , and then one invokes the explicit formula for the Wasserstein distance for probability measures on  $\mathbb{R}$  (see Remark 2.12) to obtain a lower bound to  $d_{\text{W},p}^{\mathbb{R}^d}(\alpha, \beta)$  without incurring the possibly high computational cost associated to solving an optimal transportation problem. This lower bound is improved via repeated (often random) selections of the line  $\ell$  [9, 53, 85].

Recently, Le et al. [58] pointed out that, thanks to the fact that the 1-Wasserstein distance also admits an explicit formula when the underlying metric space is a tree [28, 34, 65], one can also devise *tree slicing* estimates of the distance between two given probability measures by suitably projecting them onto tree-like structures. Most likely, the same strategy is successful for suitable projections on random ultrametric spaces, as on these there is also an explicit formula for the Wasserstein distance [50]. The same line of work has also recently been explored in the Gromov-Wasserstein scenario [57, 98] and could be extended based on efficiently computable restrictions (or surrogates of)  $d_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}$ . Inspired by the results of Mémoli et al. [70] on the ultrametric Gromov-Hausdorff distance and the results of Kloeckner [50], who derived an explicit representation of the Wasserstein distance on ultrametric spaces, we study the collection of compact *ultrametric measure spaces*  $\mathcal{U}^w \subseteq \mathcal{M}^w$ , where  $\mathcal{X} = (X, u_X, \mu_X) \in \mathcal{U}^w$ , whenever the underlying metric space  $(X, u_X)$  is a compact ultrametric space.

In terms of applications, ultrametric spaces (and thus also ultrametric *measure spaces*) arise naturally in statistics as metric encodings of dendrograms [19, 46] which is a graph theoretical representations of ultrametric spaces, in the context of phylogenetic trees [90], in theoretical computer science in the probabilistic approximation of finite metric spaces [5, 35], and in physics in the context of a mean-field theory of spin glasses [71, 81].

Especially for phylogenetic trees (and dendrograms), where one tries to characterize the structure of an underlying evolutionary process or the difference between two such processes, it is important to have a meaningful method of comparison, i.e., to have a meaningful metric on  $\mathcal{U}^w$ . However, it is evident from the definition of  $d_{\text{GW},p}^{\text{sturm}}$  and the relationship between  $d_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}$  (see [67]), that the ultrametric structure of  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  is not taken into account in the computation of either  $d_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  or  $d_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$ ,  $1 \leq p \leq \infty$ . Hence, we suggest, just as for the ultrametric Gromov-Hausdorff distance, to adapt the definition of  $d_{\text{GW},p}^{\text{sturm}}$  (see Equation (5)) as well as the one of  $d_{\text{GW},p}$  (see Equation (7)) and verify in the following that this makes the comparisons of ultrametric measure spaces more sensitive and leads for  $p = \infty$  to a *polynomial time* algorithm for the derivation of the proposed metrics.

**1.1. The proposed approach.** Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be ultrametric measure spaces. Reconsidering the definition of Sturm's Gromov-Wasserstein distance in Equation (5), we propose to only infimize over ultrametric spaces  $(Z, u_Z)$  in Equation (5). Thus, we define for  $p \in [1, \infty]$  *Sturm's ultrametric Gromov-Wasserstein distance* of order  $p$

as

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) := \inf_{Z, \phi, \psi} d_{\text{W},p}^{(Z, u_Z)}(\phi_{\#}\mu_X, \psi_{\#}\mu_Y), \quad (10)$$

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings into an ultrametric space  $(Z, u_Z)$ .

In the subsequent sections of this paper, we will establish many theoretically appealing properties of  $u_{\text{GW},p}^{\text{sturm}}$ . Unfortunately, we will verify that, although an explicit formula for the Wasserstein distance of order  $p$  on ultrametric spaces exists [50], for  $p \in [1, \infty)$  the calculation of  $u_{\text{GW},p}^{\text{sturm}}$  yields a highly non-trivial combinatorial optimization problem (see Section 3.1.1). Therefore, we demonstrate that an adaption of the Gromov-Wasserstein distance defined in Equation (7) yields a topologically equivalent and easily approximable distance on  $\mathcal{U}^w$ . In order to define this adaption, we need to introduce some notation. For  $a, b \geq 0$  and  $1 \leq q < \infty$  let

$$\Lambda_q(a, b) := |a^q - b^q|^{1/q}.$$

Further define  $\Lambda_\infty(a, b) := \max(a, b)$  whenever  $a \neq b$  and  $\Lambda_\infty(a, b) = 0$  if  $a = b$ .

Now, we can rewrite  $d_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , as follows

$$d_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = \frac{1}{2} \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left( \iint_{X \times Y \times X \times Y} (\Lambda_1(d_X(x, x'), d_Y(y, y')))^p \mu(dx \times dy) \mu(dx' \times dy') \right)^{1/p}. \quad (11)$$

Considering the derivation of  $d_{\text{GW},p}$  in [67] and the results on the closely related ultrametric Gromov-Hausdorff distance studied in [70], this suggests to replace  $\Lambda_1$  in Equation (11) with  $\Lambda_\infty$  in order to incorporate the ultrametric structures of  $(X, u_X, \mu_X)$  and  $(Y, u_Y, \mu_Y)$  into the comparison. Hence, we define the *p-ultra-distortion* of a coupling  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  for  $1 \leq p < \infty$  as

$$\text{dis}_p^{\text{ult}}(\mu) := \left( \iint_{X \times Y \times X \times Y} (\Lambda_\infty(u_X(x, x'), u_Y(y, y')))^p \mu(dx \times dy) \mu(dx' \times dy') \right)^{1/p}. \quad (12)$$

and for  $p = \infty$  as

$$\text{dis}_\infty^{\text{ult}}(\mu) := \sup_{\substack{x, x' \in \mathcal{X}, y, y' \in \mathcal{Y} \\ \text{s.t. } (x, y), (x', y') \in \text{supp}(\mu)}} \Lambda_\infty(u_X(x, x'), u_Y(y, y')).$$

The *ultrametric Gromov-Wasserstein distance* of order  $p \in [1, \infty]$ , is given as

$$u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) := \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \text{dis}_p^{\text{ult}}(\mu). \quad (13)$$

Due to the structural similarity between  $d_{\text{GW},p}$  and  $u_{\text{GW},p}$ , we can expect (and later verify) that many properties of  $d_{\text{GW},p}$  extend to  $u_{\text{GW},p}$ . In particular, we will establish that also  $u_{\text{GW},p}$  can be approximated<sup>1</sup> via conditional gradient descent and admits several polynomial time computable lower bounds which are useful in applications.

<sup>1</sup>Here “approximation” is meant in the sense that one can write code which will locally minimize the functional. There are in general no theoretical guarantees that these algorithms will converge to a global minimum.

It is worth mentioning that Sturm [93] studied the family of so-called  $L^{p,q}$ -distortion distances similar to our construction of  $u_{\text{GW},p}$ . In our language, for any  $p, q \in [1, \infty)$ , the  $L^{p,q}$ -distortion distance is constructed by infimizing over the  $(p, q)$ -distortion defined by replacing  $\Lambda_\infty$  with  $(\Lambda_q)^q$  in Equation (12). This distance shares many properties with  $d_{\text{GW},p}$ .

## 1.2. Overview of our results.

We give a brief overview of our results.

**Section 2.** We generalize the results of Carlsson and Mémoli [19] on the relation between ultrametric spaces and dendrograms and establish a bijection between compact ultrametric spaces and *proper dendrograms* (see Definition 2.1). After recalling some results on the ultrametric Gromov-Hausdorff distance (see Equation (9)), we use the connection between compact ultrametric spaces and dendrograms to reformulate the explicit formula for the  $p$ -Wasserstein distance ( $1 \leq p < \infty$ ) on ultrametric spaces derived by Kloeckner [50] in terms of proper dendrograms. This allows us to derive a formulation of the  $\infty$ -Wasserstein distance on ultrametric spaces and to study the Wasserstein distance on compact subspaces of the ultrametric space  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ , which will be relevant when studying lower bounds of  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ .

**Section 3.** We demonstrate that  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , are  $p$ -metrics on the collection of ultrametric measure spaces  $\mathcal{U}^w$ . We derive several alternative representations for  $u_{\text{GW},p}^{\text{sturm}}$  and study the relation between the metrics  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$ . In particular, we show that, while for  $1 \leq p < \infty$  it holds in general that  $u_{\text{GW},p} \leq 2^{\frac{1}{p}} u_{\text{GW},p}^{\text{sturm}}$ , both metrics coincide for  $p = \infty$ , i.e.,  $u_{\text{GW},\infty} = u_{\text{GW},\infty}^{\text{sturm}}$ . Furthermore, we show how this equality in combination with an alternative representation of  $u_{\text{GW},\infty}$  leads to a *polynomial time algorithm* for the calculation of  $u_{\text{GW},\infty}^{\text{sturm}} = u_{\text{GW},\infty}$ . Moreover, we study the topological properties of  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  and  $(\mathcal{U}^w, u_{\text{GW},p})$ ,  $1 \leq p \leq \infty$ . Most importantly, we show that  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$  induce the same topology on  $\mathcal{U}^w$  which is also different from the one induced by  $d_{\text{GW},p}^{\text{sturm}}/d_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ . While we further prove that the metric spaces  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  and  $(\mathcal{U}^w, u_{\text{GW},p})$ ,  $1 \leq p < \infty$ , are neither complete nor separable metric space, we demonstrate that the ultrametric space  $(\mathcal{U}^w, u_{\text{GW},\infty}^{\text{sturm}})$ , which coincides with  $(\mathcal{U}^w, u_{\text{GW},\infty})$ , is complete. Finally, we establish that  $(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$  is a geodesic space.

**Section 4.** Unfortunately, it does not seem to be possible to derive a polynomial time algorithm for the calculation of  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$ ,  $1 \leq p < \infty$ . Consequently, based on easily computable invariant features, in Section 4 we derive several polynomial time computable lower bounds for  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ . Due to the structural similarity between  $d_{\text{GW},p}$  and  $u_{\text{GW},p}$ , these are in a certain sense analogue to those derived in [66, 67] for  $d_{\text{GW},p}$ . Among other things, we show that

$$u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) := \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \|\Lambda_\infty(u_X, u_Y)\|_{L^p(\gamma)}. \quad (14)$$

We verify that the lower bound  $\mathbf{SLB}_p^{\text{ult}}$  can be reformulated in terms of the Wasserstein distance on the ultrametric space  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  (we derive an explicit formula for  $d_{\text{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  in Section 2.3). This allows us to efficiently calculate  $\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y})$  in  $O((m \vee n)^2)$ , where  $m$  stands for the cardinality of  $X$  and  $n$  for the one of  $Y$ .

**Section 5.** As the ultrametric space assumption is somewhat restrictive (especially in the context of phylogenetic trees, see [90]), we prove in Section 5 that the results on  $u_{\text{GW},p}$  can be extended to the more general *ultra-dissimilarity spaces* (see Definition 5.1). In particular, we prove that  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , is a metric on the *isomorphism classes* of ultra-dissimilarity spaces (see Definition 5.5).

**Section 6.** We illustrate the behaviour and relation between  $u_{\text{GW},1}$  (which can be approximated via conditional gradient descent) and  $\text{SLB}_1^{\text{ult}}$  in a set of illustrative examples. Additionally, we carefully illustrate the differences between  $u_{\text{GW},1}$  and  $\text{SLB}_1^{\text{ult}}$ , and  $d_{\text{GW},1}$  and  $\text{SLB}_1$  (see Section 4 for a definition), respectively.

**Section 7.** Finally, we apply our ideas to *phylogenetic tree shape comparison*. To this end, we compare two sets of phylogenetic tree shapes based on the HA protein sequences from human influenza collected in different regions with the lower bound  $\text{SLB}_1^{\text{ult}}$ . In particular, we contrast our results in both settings to the ones obtained with the tree shape metric introduced in Equation (4) of Colijn and Plazzotta [25].

**1.3. Related work.** In order to better contextualize our contribution, we now describe related work, both in applied and computational geometry, and in phylogenetics (where notions of distance between trees have arisen naturally).

*Metrics between trees: the phylogenetics perspective.* In phylogenetics, where one chief objective is to infer the evolutionary relationship between species via methods that evaluate observable traits, such as DNA sequences, the need to be able to measure dissimilarity between different trees arises from the fact that the process of reconstruction of a phylogenetic tree may depend on the set of genes being considered. At the same time, even for the same set of genes, different reconstruction methods could be applied which would result in different trees. As such, this has led to the development of many different metrics for measuring distance between phylogenetic trees. Examples include the Robinson-Foulds metric [84], the subtree-prune and regraft distance [42], and the nearest-neighbor interchange distance [83].

As pointed out in [76], many of these distances tend to quantify differences between tree topologies and often do not take into account edge lengths. A certain phylogenetic tree metric space which encodes for edge lengths was proposed in [6] and studied algorithmically in [76]. This tree space assumes that the all trees have the same set of taxa. An extension to the case of trees over different underlying sets is given in [40]. Lafond et al. [56] considered one type of metrics on possibly *multilabeled* phylogenetic trees with a fixed number of leafs. As the authors pointed out, a multilabeled phylogenetic tree in which no leafs are repeated is just a standard phylogenetic tree, whereas a multilabeled phylogenetic tree in which all labels are equal defines a *tree shape*. The authors then proceeded to study the computational complexity associated to generalizations of some of the usual metrics for phylogenetic trees (such as the Robinson-Foulds distance) to the multilabeled case. Colijn and Plazzotta [25] studied a metric between (binary) phylogenetic tree shapes based on a bottom to top enumeration of specific connectivity structures. The authors applied their metric to compare evolutionary trees based on the HA protein sequences from human influenza collected in different regions.

*Metrics between trees: the applied geometry perspective.* From a different perspective, ideas from applied geometry and applied and computational topology have been applied to the

comparison of tree shapes in applications in probability, clustering and applied and computational topology.

Metric trees are also considered in probability theory in the study of models for random trees together with the need to quantify their distance; Evans [33] described some variants of the Gromov-Hausdorff distance between metric trees. See also [39] for the case of metric measure space representations of trees and a certain Gromov-Prokhorov type of metric on the collection thereof.

Trees, in the form of dendrograms, are abundant in the realm of hierarchical clustering methods. In their study of the *stability* of hierarchical clustering methods, Carlsson and Mémoli [19] utilized the Gromov-Hausdorff distance between the ultrametric representation of dendrograms. Schmiedl [88] proved that computing the Gromov-Hausdorff distance between tree metric spaces is NP-hard. Liebscher [59] suggested some variants of the Gromov-Hausdorff distance which are applicable in the context of phylogenetic trees. As mentioned before, Zarichnyi [105] introduced the ultrametric Gromov-Hausdorff distance  $u_{\text{GH}}$  between compact ultrametric spaces (a special type of tree metric spaces). Certain theoretical properties such as precompactness of  $u_{\text{GH}}$  has been studied in [80]. In contrast with the NP-hardness of computing  $d_{\text{GH}}$ , Mémoli et al. [70] devised a polynomial time algorithm for computing  $u_{\text{GH}}$ .

In computational topology *merge trees* arise through the study of the sublevel sets of a given function [1, 82] with the goal of shape simplification. Morozov et al. [74] developed the notion of *interleaving distance* between merge trees which is related to the Gromov-Hausdorff distance between trees through bi-Lipschitz bounds. In [2], exploiting the connection between the interleaving distance and the Gromov-Hausdorff between metric trees, the authors approached the computation of the Gromov-Hausdorff distance between metric trees in general and provide certain approximation algorithms. Touli and Wang [96] devised fixed-parameter tractable (FPT) algorithms for computing the interleaving distance between metric trees. One can imply from their methods an FPT algorithm to compute a 2-approximation of the Gromov-Hausdorff distance between ultrametric spaces. Mémoli et al. [70] devised an FPT algorithm for computing the exact value of the Gromov-Hausdorff distances between ultrametric spaces.

## 2. PRELIMINARIES

In this section we briefly summarize the basic notions and concepts required throughout the paper.

**2.1. Ultrametric spaces and dendrograms.** We begin by describing compact ultrametric spaces in terms of *proper dendrograms*. To this end, we introduce some definitions and some notation. Given a set  $X$ , a *partition* of  $X$  is a set  $P_X = \{X_i\}_{i \in I}$  where  $I$  is any index set,  $\emptyset \neq X_i \subseteq X$ ,  $X_i \cap X_j = \emptyset$  for all  $i \neq j \in I$  and  $\bigcup_{i \in I} X_i = X$ . We call each element  $X_i$  a *block* of the given partition  $P_X$  and denote by  $\mathbf{Part}(X)$  the collection of all partitions of  $X$ . For two partitions  $P_X$  and  $P'_X$  we say that  $P_X$  is *finer* than  $P'_X$ , if for every block  $X_i \in P_X$  there exists a block  $X'_j \in P'_X$  such that  $X_i \subseteq X'_j$ .

**Definition 2.1** (Proper dendrogram). Given a set  $X$  (not necessarily finite), a *proper dendrogram*  $\theta_X : [0, \infty) \rightarrow \mathbf{Part}(X)$  is a map satisfying the following conditions:

- (1)  $\theta_X(s)$  is finer than  $\theta_X(t)$  for any  $0 \leq s < t < \infty$ ;
- (2)  $\theta_X(0)$  is the finest partition consisting only singleton sets;
- (3) There exists  $T > 0$  such that for any  $t \geq T$ ,  $\theta_X(t) = \{X\}$  is the trivial partition;
- (4) For each  $t > 0$ , there exists  $\varepsilon > 0$  such that  $\theta_X(t) = \theta_X(t')$  for all  $t' \in [t, t + \varepsilon]$ .
- (5) For any distinct points  $x, x' \in X$ , there exists  $T_{xx'} > 0$  such that  $x$  and  $x'$  belong to different blocks in  $\theta_X(T_{xx'})$ .
- (6) For each  $t > 0$ ,  $\theta_X(t)$  consists of only finitely many blocks.
- (7) Let  $\{t_n\}_{n \in \mathbb{N}}$  be a decreasing sequence such that  $\lim_{n \rightarrow \infty} t_n = 0$  and let  $X_n \in \theta_X(t_n)$ . If for any  $1 \leq n < m$ ,  $X_m \subseteq X_n$ , then  $\bigcap_{n \in \mathbb{N}} X_n \neq \emptyset$ .

When  $X$  is finite, a function  $\theta_X : [0, \infty) \rightarrow \mathbf{Part}(X)$  satisfying conditions (1) to (4) will satisfy conditions (5), (6) and (7) automatically, and thus a proper dendrogram reduces to the usual dendrogram (see [19, Sec. 3.1] for a formal definition). Let  $\theta_X$  be a proper dendrogram over a set  $X$ . For any  $x \in X$  and  $t \geq 0$ , we denote by  $[x]_t^X$  the block in  $\theta(t)$  that contains  $x \in X$  and abbreviate  $[x]_t^X$  to  $[x]_t$  when the underlying set  $X$  is clear from the context. Similar to [19], who considered the relation between finite ultrametric spaces and dendrograms, we will prove that there is a bijection between compact ultrametric spaces and proper dendrograms. In particular, one can show that the subsequent theorem generalizes [19, Theorem 9]. Since its proof depends on several concepts not yet introduced, we postpone it to Appendix A.1.1.

**Theorem 2.2.** *Given a set  $X$ , denote by  $\mathcal{U}(X)$  the collection of all compact ultrametrics on  $X$  and  $\mathcal{D}(X)$  the collection of all proper dendrograms over  $X$ . For any  $\theta \in \mathcal{D}(X)$ , consider  $u_\theta$  defined as follows:*

$$\forall x, x' \in X, \quad u_\theta(x, x') := \inf\{t \geq 0 \mid x, x' \text{ belong to the same block of } \theta(t)\}.$$

*Then,  $u_\theta \in \mathcal{U}(X)$  and the map  $\Delta_X : \mathcal{D}(X) \rightarrow \mathcal{U}(X)$  sending  $\theta$  to  $u_\theta$  is a bijection.*

**Remark 2.3.** From now on, we denote by  $\theta_X$  the proper dendrogram corresponding to a given compact ultrametric  $u_X$  on  $X$  under the bijection given above. Note that a block  $[x]_t$  in  $\theta_X(t)$  is actually the closed ball  $B_t(x)$  in  $X$  centered at  $x$  with radius  $t$ . So for each  $t \geq 0$ ,  $\theta_X(t)$  partitions  $X$  into a union of several closed balls in  $X$  with respect to  $u_X$ .

**2.2. The ultrametric Gromov-Hausdorff distance.** Both  $d_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , are by construction closely related to the Gromov-Hausdorff distance. In a recent paper, Mémoli et al. [70] studied an ultrametric version of this distance, namely the *ultrametric Gromov-Hausdorff distance* (denoted as  $u_{\text{GH}}$ ). Since we will demonstrate several connections between  $u_{\text{GW},p}^{\text{sturm}}$ ,  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , and this distance, we briefly summarize some of the results in [70]. We start by recalling the formal definition of  $u_{\text{GH}}$ .

**Definition 2.4.** Let  $(X, u_X)$  and  $(Y, u_Y)$  be two compact ultrametric spaces. Then, the *ultrametric Gromov-Hausdorff* between  $X$  and  $Y$  is defined as

$$u_{\text{GH}}(X, Y) = \inf_{Z, \phi, \psi} d_{\text{H}}^Z(\phi(X), \psi(Y)),$$




FIGURE 1. **Metric quotient:** An ultrametric space (black) and its quotient at level  $t$  (red).

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings (distance preserving transformations) into the ultrametric space  $(Z, u_Z)$ .

Zarichnyi [105] has shown that  $u_{\text{GH}}$  is an ultrametric on the isometry classes of compact ultrametric spaces, which are denoted by  $\mathcal{U}$ , and Mémoli et al. [70] identified a structural theorem (cf. Theorem 2.5) that gives rise to a polynomial time algorithm for the calculation of  $u_{\text{GH}}$ . More precisely, it was proven in [70] that  $u_{\text{GH}}$  can be calculated via so-called *quotient ultrametric spaces*, which we define next. Let  $(X, u_X)$  be an ultrametric space and let  $t \geq 0$ . We define an equivalence relation  $\sim_t$  on  $X$  as follows:  $x \sim_t x'$  if and only if  $u_X(x, x') \leq t$ . We denote by  $[x]_t^X$  (resp.  $[x]_t$ ) the equivalence class of  $x$  under  $\sim_t$  and by  $X_t$  the set of all such equivalence classes. In fact,  $[x]_t^X = \{x' \in X \mid u(x, x') \leq t\}$  is exactly the closed ball centered at  $x$  with radius  $t$  and corresponds to a block in the corresponding proper dendrogram  $\theta_X(t)$  (see Remark 2.3). Thus, one can think of  $X_t$  as a “set representation” of  $\theta_X(t)$ . We define an ultrametric  $u_{X_t}$  on  $X_t$  as follows:

$$u_{X_t}([x]_t, [x']_t) := \begin{cases} u_X(x, x'), & [x]_t \neq [x']_t \\ 0, & [x]_t = [x']_t. \end{cases}$$

Then,  $(X_t, u_{X_t})$  is an ultrametric space and we call  $(X_t, u_{X_t})$  the *quotient* of  $(X, u_X)$  at level  $t$  (see Figure 1 for an illustration). It is straightforward to prove that the quotient of a compact ultrametric space at level  $t > 0$  is a finite ultrametric space (cf. [102, Lemma 2.3]). Furthermore, the quotient spaces characterize  $u_{\text{GH}}$  as follows.

**Theorem 2.5** (Structural theorem for  $u_{\text{GH}}$ , [70, Theorem 5.7]). *Let  $(X, u_X)$  and  $(Y, u_Y)$  be two compact ultrametric spaces. Then,*

$$u_{\text{GH}}(X, Y) = \inf \{t \geq 0 \mid X_t \cong Y_t\}.$$

**Remark 2.6.** Let  $(X, u_X)$  and  $(Y, u_Y)$  denote two finite ultrametric spaces and let  $t \geq 0$ . The quotient spaces  $X_t$  and  $Y_t$  can be considered as vertex weighted, rooted trees [70]. Hence, it is possible to check whether  $X_t \cong Y_t$  in polynomial time [3]. Consequently, Theorem 2.5 induces a simple, polynomial time algorithm to calculate  $u_{\text{GH}}$  between two finite ultrametric spaces.

**2.3. Wasserstein distance on ultrametric spaces.** Kloeckner [50] uses the representation of ultrametric spaces as so called *synchronized rooted trees* to derive an explicit formula for the Wasserstein distance on ultrametric spaces. By the constructions of the dendrograms and of the synchronized rooted trees (see Appendix A.2.1), it is immediately clear how to




FIGURE 2. **Illustration of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ :** This is the dendrogram for a subspace of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  consisting of 5 arbitrary distinct points of  $\mathbb{R}_+$ .

reformulate the results of Kloeckner [50] on compact ultrametric spaces in terms of proper dendrograms. To this end, we need to introduce some notation. For a compact ultrametric space  $X$ , let  $\theta_X$  be the associated proper dendrogram and let  $V(X) := \bigcup_{t>0} \theta_X(t) = \{[x]_t \mid x \in X, t > 0\}$ . It can be shown that  $V(X)$  is the collection of all closed balls in  $X$  except for singletons  $\{x\}$  such that  $x$  is a cluster point<sup>2</sup> (see Lemma A.8). For  $B \in V(X)$ , we denote by  $B^*$  the smallest (under inclusion) element in  $V(X)$  such that  $B \subsetneq B^*$  (for the existence and uniqueness of  $B^*$  see Lemma A.1).

**Theorem 2.7** (The Wasserstein distance on ultrametric spaces, [50, Theorem 3.1]). *Let  $(X, u_X)$  be a compact ultrametric space. For all  $\alpha, \beta \in \mathcal{P}(X)$  and  $1 \leq p < \infty$ , we have*

$$(d_{W,p}^X)^p(\alpha, \beta) = 2^{-1} \sum_{B \in V(X) \setminus \{X\}} (\text{diam}(B^*)^p - \text{diam}(B)^p) |\alpha(B) - \beta(B)|. \quad (15)$$

While Theorem 2.7 is only valid for  $p < \infty$ , it can be extended to the case  $p = \infty$ .

**Lemma 2.8.** *Let  $X$  be a compact ultrametric space. Then, for any  $\alpha, \beta \in \mathcal{P}(X)$ , we have*

$$d_{W,\infty}^X(\alpha, \beta) = \max_{B \in V(X) \setminus \{X\} \text{ and } \alpha(B) \neq \beta(B)} \text{diam}(B^*). \quad (16)$$

The proof of Lemma 2.8 is technical and we postpone it to Appendix A.1.2.

**2.3.1. Wasserstein distance on  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ .** The non-negative half real line  $\mathbb{R}_{\geq 0}$  endowed with  $\Lambda_\infty$  turns out to be an ultrametric space (cf. [70, Remark 1.14]). Finite subspaces of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  are of particular interest in this paper. These spaces possess a particular structure (see Figure 2) and the computation of the Wasserstein distance on them can be further simplified.

**Theorem 2.9** ( $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  between finitely supported measures). *Suppose  $\alpha, \beta$  are two probability measures supported on a finite subset  $\{x_0, \dots, x_n\}$  of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  such that  $0 \leq x_0 < x_1 < \dots < x_n$ . Denote  $\alpha_i := \alpha(\{x_i\})$  and  $\beta_i := \beta(\{x_i\})$ . Then, we have for  $p \in [1, \infty)$  that*

$$d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(\alpha, \beta) = 2^{-\frac{1}{p}} \left( \sum_{i=0}^{n-1} \left| \sum_{j=0}^i (\alpha_j - \beta_j) \right| \cdot |x_{i+1}^p - x_i^p| + \sum_{i=0}^n |\alpha_i - \beta_i| \cdot x_i^p \right)^{\frac{1}{p}}. \quad (17)$$

<sup>2</sup>A cluster point  $x$  in a topological space  $X$  is such that any neighborhood of  $x$  contains countably many points in  $X$ .

Let  $F_\alpha$  and  $F_\beta$  denote the cumulative distribution functions of  $\alpha$  and  $\beta$ , respectively. Then, for the case  $p = \infty$  we obtain

$$d_{W,\infty}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(\alpha, \beta) = \max \left( \max_{0 \leq i \leq n-1, F_\alpha(x_i) \neq F_\beta(x_i)} x_{i+1}, \max_{0 \leq i \leq n, \alpha_i \neq \beta_i} x_i \right).$$

*Proof.* Clearly,  $V(X) = \{\{x_0, x_1, \dots, x_i\} \mid i = 1, \dots, n\} \cup \{\{x_i\} \mid i = 1, \dots, n\}$  (recall that each set corresponds to a closed ball). Thus, we conclude the proof by applying Theorem 2.7 and Lemma 2.8.  $\square$

**Remark 2.10** (The case  $p = 1$ ). Note that when  $p = 1$ , for any finitely supported probability measures  $\alpha, \beta \in \mathcal{P}(\mathbb{R}_{\geq 0})$ ,

$$d_{W,1}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(\alpha, \beta) = \frac{1}{2} \left( d_{W,1}^{(\mathbb{R}, \Lambda_1)}(\alpha, \beta) + \int_{\mathbb{R}} x |\alpha - \beta|(dx) \right).$$

The formula indicates that the 1-Wasserstein distance on  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  is the average of the usual 1-Wasserstein distance on  $(\mathbb{R}_{\geq 0}, \Lambda_1)$  and a “weighted total variation distance”. The weighted total variation like distance term is sensitive to difference of supports. For example, let  $\alpha = \delta_{x_1}$  and  $\beta = \delta_{x_2}$ , then  $\int_{\mathbb{R}} x |\alpha - \beta|(dx) = x_1 + x_2$  if  $x_1 \neq x_2$ .

**Remark 2.11** (Extension to compactly supported measures). In fact,  $X \subseteq (\mathbb{R}_{\geq 0}, \Lambda_\infty)$  is compact if and only if it is either a finite set or countable with 0 being the unique cluster point (w.r.t. the usual Euclidean distance  $\Lambda_1$ ) (see Lemma A.2). Hence, it is straightforward to extend Theorem 2.9 to compactly supported measures and we refer to Appendix A.3 for the missing details.

**Remark 2.12** (Closed-form solution for  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}$ ). We know that there is a closed-form solution for Wasserstein distance on  $\mathbb{R}$  with the usual Euclidean distance  $\Lambda_1$ :

$$d_{W,p}^{(\mathbb{R}, \Lambda_1)}(\alpha, \beta) = \left( \int_0^1 |F_\alpha^{-1}(t) - F_\beta^{-1}(t)|^p dt \right)^{\frac{1}{p}},$$

where  $F_\alpha$  and  $F_\beta$  are cumulative distribution functions of  $\alpha$  and  $\beta$ , respectively. We have also obtained a closed-form solution for  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  in Theorem 2.9. We generalize these formulas to the case  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}$  when  $q \in (1, \infty)$  and  $q \leq p$  in Appendix A.3.1.

### 3. ULTRAMETRIC GROMOV-WASSERSTEIN DISTANCES

In this section we investigate the properties of  $u_{GW,p}^{\text{sturm}}$  as well as  $u_{GW,p}$ ,  $1 \leq p \leq \infty$ , and study the relation between them.

**3.1. Sturm’s ultrametric Gromov-Wasserstein distance.** We begin by establishing several basic properties of  $u_{GW,p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , including a proof that  $u_{GW,p}^{\text{sturm}}$  is indeed a metric (or more precisely a  $p$ -metric) on the collection of compact ultrametric measure spaces  $\mathcal{U}^w$ .

The definition of  $u_{GW,p}^{\text{sturm}}$  given in Equation (10) is clunky, technical and in general not easy to work with. Hence, the first observation to make is the fact that  $u_{GW,p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , shares a further property with  $d_{GW,p}^{\text{sturm}}$ :  $u_{GW,p}^{\text{sturm}}$  can be calculated by minimizing over pseudo-ultrametrics instead of isometric embeddings.

**Lemma 3.1.** *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be two ultrametric measure spaces. Let  $\mathcal{D}^{\text{ult}}(u_X, u_Y)$  denote the collection of all pseudo-ultrametrics  $u$  on the disjoint union  $X \sqcup Y$  such that  $u|_{X \times X} = u_X$  and  $u|_{Y \times Y} = u_Y$ . Let  $p \in [1, \infty]$ . Then, it holds that*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)} d_{\text{W},p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y), \quad (18)$$

where  $d_{\text{W},p}^{(X \sqcup Y, u)}$  denotes the Wasserstein pseudometric of order  $p$  defined in Equation (34) (resp. in Equation (35) for  $p = \infty$ ) in Appendix B.5.1 of the supplement.

*Proof.* The above lemma follows by the same arguments as Lemma 3.3 (iii) in [92].  $\square$

**Remark 3.2** (Wasserstein pseudometric). The *Wasserstein pseudometric* is a natural extension of the Wasserstein distance to pseudometric spaces and has for example been studied in Thorsley and Klavins [94]. In Appendix B.5.1 we carefully show that it is closely related to the Wasserstein distance on a canonically induced metric space. We further establish that the Wasserstein distance and the Wasserstein pseudometric share many relevant properties. Hence, we do not notationally distinguish between these two concepts.

The representation of  $u_{\text{GW},p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , given by the above lemma is much more accessible and we first use it to establish the subsequent basic properties of  $u_{\text{GW},p}^{\text{sturm}}$  (see Appendix B.1.1 for a full proof).

**Proposition 3.3.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, the following holds:*

- (1) *For any  $p \in [1, \infty]$ , we always have that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \geq d_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .*
- (2) *For any  $1 \leq p \leq q \leq \infty$ , we have that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq u_{\text{GW},q}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .*
- (3) *It holds that  $\lim_{p \rightarrow \infty} u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .*

Moreover, we use Lemma 3.1 to prove that  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is indeed a metric space.

**Theorem 3.4.**  *$u_{\text{GW},p}^{\text{sturm}}$  is a  $p$ -metric on the collection  $\mathcal{U}^w$  of compact ultrametric measure spaces. In particular, when  $p = \infty$ ,  $u_{\text{GW},\infty}^{\text{sturm}}$  is an ultrametric.*

In order to increase the readability of this section we postpone the proof of Theorem 3.4 to Appendix B.1.2. In the course of the proof, we will, among other things, verify the existence of optimal metrics and optimal couplings in Equation (18) (see Proposition B.1). Furthermore, it is important to note that the topology induced on  $\mathcal{U}^w$  by  $u_{\text{GW},p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , is different from the one induced by  $d_{\text{GW},p}^{\text{sturm}}$ . This is well illustrated in the following example.

**Example 3.5** ( $u_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}^{\text{sturm}}$  induce different topologies). This example is an adaptation from Mémoli et al. [70, Example 3.14]. For each  $a > 0$ , denote by  $\Delta_2(a)$  the two-point metric space with interpoint distance  $a$ . Endow with  $\Delta_2(a)$  the uniform probability measure  $\mu_a$  and denote the corresponding ultrametric measure space  $\hat{\Delta}_2(a)$ . Now, let  $\mathcal{X} := \hat{\Delta}_2(1)$  and let  $\mathcal{X}_n := \hat{\Delta}_2(1 + \frac{1}{n})$  for  $n \in \mathbb{N}$ . It is easy to check that for any  $1 \leq p \leq \infty$ ,  $d_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{X}_n) = \frac{1}{2n}$  and  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{X}_n) = 2^{-\frac{1}{p}}(1 + \frac{1}{n})$  where we adopt the convention that  $1/\infty = 0$ . Hence, as  $n$




FIGURE 3. **Common ultrametric spaces:** Representation of the two kinds of ultrametric spaces  $Z$  (middle and right) into which we can isometrically embed the spaces  $X$  and  $Y$  (left).

goes to infinity  $\mathcal{X}_n$  will converge to  $\mathcal{X}$  in the sense of  $d_{\text{GW},p}^{\text{sturm}}$ , but not in the sense of  $u_{\text{GW},p}^{\text{sturm}}$ , for any  $1 \leq p \leq \infty$ .

**3.1.1. Alternative representations of  $u_{\text{GW},p}^{\text{sturm}}$ .** In this subsection, we derive an alternative representation for  $u_{\text{GW},p}^{\text{sturm}}$  defined in Equation (10). We mainly focus on the case  $p < \infty$ , however it turns out that the results also hold for  $p = \infty$  (see Section 3.3).

Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  and recall the original definition of  $u_{\text{GW},p}^{\text{sturm}}$ ,  $p \in [1, \infty]$ , given in Equation (10), i.e.,

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{Z, \phi, \psi} d_{\text{W},p}^{(Z, u_Z)}(\phi_{\#} \mu_X, \psi_{\#} \mu_Y),$$

where  $\phi : X \rightarrow Z$  and  $\psi : Y \rightarrow Z$  are isometric embeddings into an ultrametric space  $(Z, u_Z)$ . It turns out that we only need to consider relatively few possibilities of mapping two ultrametric spaces into a common ultrametric space. Exemplarily, this is shown in Figure 3, where we see two finite ultrametric spaces and two possibilities for a common ultrametric space  $Z$ . Indeed, it is straightforward to write down all reasonable embeddings and target spaces. We define the set

$$\mathcal{A} := \{(A, \varphi) \mid \emptyset \neq A \subseteq X \text{ is closed and } \varphi : A \hookrightarrow Y \text{ is an isometric embedding}\}. \quad (19)$$

Clearly,  $\mathcal{A} \neq \emptyset$ , as it holds for each  $x \in X$  that  $\{(\{x\}, \varphi_y)\}_{y \in Y} \subseteq \mathcal{A}$ , where  $\varphi_y$  is the map sending  $x$  to  $y \in Y$ . Another possibility to construct elements in  $\mathcal{A}$  is illustrated in the subsequent example.

**Example 3.6.** Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  be finite spaces and let  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$ . If  $u^{-1}(0) \neq \emptyset$ , we define  $A := \pi_X(u^{-1}(0)) \subseteq X$ , where  $\pi_X : X \times Y \rightarrow X$  is the canonical projection. Then, the map  $\varphi : A \rightarrow Y$  defined by sending  $x \in A$  to  $y \in Y$  such that  $u(x, y) = 0$  is an isometric embedding and in particular,  $(A, \varphi) \in \mathcal{A}$ .

Now, fix two compact spaces  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Let  $(A, \varphi) \in \mathcal{A}$  and let  $Z_A = X \sqcup (Y \setminus \varphi(A)) \subseteq X \sqcup Y$ . Furthermore, define  $u_{Z_A} : Z_A \times Z_A \rightarrow \mathbb{R}_{\geq 0}$  as follows:

- (1)  $u_{Z_A}|_{X \times X} := u_X$  and  $u_{Z_A}|_{Y \setminus \varphi(A) \times Y \setminus \varphi(A)} := u_Y|_{Y \setminus \varphi(A) \times Y \setminus \varphi(A)}$ ;
- (2) For any  $x \in A$  and  $y \in Y \setminus \varphi(A)$  define  $u_{Z_A}(x, y) := u_Y(y, \varphi(x))$ ;
- (3) For  $x \in X \setminus A$  and  $y \in Y \setminus \varphi(A)$  let  $u_{Z_A}(x, y) := \inf\{\max(u_X(x, a), u_Y(\varphi(a), y)) \mid a \in A\}$ ;
- (4) For any  $x \in X$  and  $y \in Y \setminus \varphi(A)$ ,  $u_{Z_A}(y, x) := u_{Z_A}(x, y)$ .

Then,  $(Z_A, u_{Z_A})$  is an ultrametric space such that  $X$  and  $Y$  can be mapped isometrically into  $Z_A$  (see [105, Lemma 1.1]). Let  $\phi_{(A,\varphi)}^X$  and  $\psi_{(A,\varphi)}^Y$  denote the corresponding isometric embeddings of  $X$  and  $Y$ , respectively. This allows us to derive the following statement, whose proof is postponed to Appendix B.1.3.

**Theorem 3.7.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, we have for each  $p \in [1, \infty)$  that*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{(A,\varphi) \in \mathcal{A}} d_{\text{W},p}^{Z_A} \left( (\phi_{(A,\varphi)}^X)_{\#} \mu_X, (\psi_{(A,\varphi)}^Y)_{\#} \mu_Y \right). \quad (20)$$

**Remark 3.8.** Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two finite ultrametric measure spaces. The representation of  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$ ,  $1 \leq p \leq \infty$  given by Theorem 3.7 is very explicit and recasts the computation of  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$ ,  $1 \leq p \leq \infty$ , as a combinatorial problem. In fact, as  $\mathcal{X}$  and  $\mathcal{Y}$  are finite, the set  $\mathcal{A}$  in Equation (20) can be further reduced. More precisely, we demonstrate in Appendix B.1.3 (see Corollary B.7) that it is sufficient to infimize over the set of all *maximal pairs*, denoted by  $\mathcal{A}^*$ . Here, a pair  $(A, \varphi_1) \in \mathcal{A}$  is denoted as *maximal*, if for all pairs  $(B, \varphi_2) \in \mathcal{A}$  with  $A \subseteq B$  and  $\varphi_2|_A = \varphi_1$  it holds  $A = B$ . Using the ultrametric Gromov-Hausdorff distance (see Equation (9)) it is possible to determine if two ultrametric spaces are isometric in polynomial time [70, Theorem 5.7]. However, this is clearly not sufficient to identify all  $(A, \varphi) \in \mathcal{A}^*$  in polynomial time. Especially, for a given, viable  $A \subseteq X$ , there are usually multiple ways to define the corresponding map  $\varphi$ . Furthermore, we have for  $1 \leq p < \infty$  neither been able to further restrict the set  $\mathcal{A}^*$  nor to identify the optimal  $(A^*, \varphi^*)$ . This just leaves a brute force approach which is computationally not feasible. On the other hand, for  $p = \infty$  we are able to explicitly construct the optimal pair  $(A^*, \varphi^*)$  (see Theorem 3.22).

**3.2. The ultrametric Gromov-Wasserstein distance.** In the following, we consider basic properties of  $u_{\text{GW},p}$  and prove the analogue of Theorem 3.4, i.e., we verify that also  $u_{\text{GW},p}$  is a  $p$ -metric,  $1 \leq p \leq \infty$ , on the collection of ultrametric measure spaces.

The subsequent proposition collects three basic properties of  $u_{\text{GW},p}$  which are also shared by  $u_{\text{GW},p}^{\text{sturm}}$  (cf. Proposition 3.3). We refer to Appendix B.2.1 for its proof.

**Proposition 3.9.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, the following holds:*

- (1) *For any  $p \in [1, \infty]$ , we always have that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \geq d_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$ .*
- (2) *For any  $1 \leq p \leq q \leq \infty$ , it holds  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \leq u_{\text{GW},q}(\mathcal{X}, \mathcal{Y})$ ;*
- (3) *We have that  $\lim_{p \rightarrow \infty} u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ .*

Next, we verify that  $u_{\text{GW},p}$  is indeed a metric on the collection of ultrametric measure spaces.

**Theorem 3.10.** *The ultrametric Gromov-Wasserstein distance  $u_{\text{GW},p}$  is a  $p$ -metric on the collection  $\mathcal{U}^w$  of compact ultrametric measure spaces. In particular, when  $p = \infty$ ,  $u_{\text{GW},\infty}$  is an ultrametric.*

The full proof of Theorem 3.10, which is based on the existence of optimal couplings in Equation (13) (see Proposition B.10), is postponed to Appendix B.2.2.




FIGURE 4. **Weighted Quotient:** An ultrametric measure space (black) and its weighted quotient at level  $t$  (red).

**Remark 3.11** ( $u_{\text{GW},p}$  and  $d_{\text{GW},p}$  induce different topologies). Reconsidering Example 3.5, it is easy to verify that in this setting  $u_{\text{GW},p}(\mathcal{X}, \mathcal{X}_n) = 2^{-\frac{1}{p}}(1 + \frac{1}{n})$  while  $d_{\text{GW},p}(\mathcal{X}, \mathcal{X}_n) = \frac{1}{2^{1/pn}}$ ,  $1 \leq p \leq \infty$ . Hence, just like  $u_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}^{\text{sturm}}$ ,  $u_{\text{GW},p}$  and  $d_{\text{GW},p}$  do not induce the same topology on  $\mathcal{U}^w$ . This result can also be obtained from Section 3.4 where we derive that  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$  give rise to the same topology.

**Remark 3.12.** By the same arguments as for  $d_{\text{GW},p}$ ,  $1 \leq p < \infty$ , [67, Sec. 7], it follows that for two finite ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$  the computation of  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$ ,  $1 \leq p < \infty$ , boils down to solving a (non-convex) quadratic program. This is in general NP-hard [78]. On the other hand, for  $p = \infty$ , we will derive a polynomial time algorithm to determine  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$  (cf. Section 3.2.1).

**3.2.1. Alternative representations of  $u_{\text{GW},\infty}$ .** In the following, we will derive an alternative representation of  $u_{\text{GW},\infty}$  that resembles the one of  $u_{\text{GH}}$  derived in [70, Theorem 5.7]. It also leads to a polynomial time algorithm for the computation of  $u_{\text{GW},\infty}$ . For this purpose, we define the *weighted quotient* of an ultrametric measure space. Let  $\mathcal{X} = (X, u_X, \mu_X) \in \mathcal{U}^w$  and let  $t \geq 0$ . Then, the *weighted quotient* of  $\mathcal{X}$  at level  $t$ , is given as  $\mathcal{X}_t = (X_t, u_{X_t}, \mu_{X_t})$ , where  $(X_t, u_{X_t})$  is the quotient of the ultrametric space  $(X, u_X)$  at level  $t$  (see Section 2.2) and  $\mu_{X_t} \in \mathcal{P}(X_t)$  is the push forward of  $\mu_X$  under the canonical quotient map  $Q_t : (X, u_X) \rightarrow (X_t, u_{X_t})$  sending  $x$  to  $[x]_t$  for  $x \in X$ . Figure 4 illustrates the weighted quotient in a simple example. Based on this definition, we show the following theorem, whose proof is postponed to Appendix B.2.3.

**Theorem 3.13.** *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be two compact ultrametric measure spaces. Then, it holds that*

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = \min \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}.$$

**Remark 3.14.** The weighted quotients  $\mathcal{X}_t$  and  $\mathcal{Y}_t$  can be considered as vertex weighted, rooted trees and thus it is possible to verify whether  $\mathcal{X}_t \cong_w \mathcal{Y}_t$  in polynomial time [3]. In consequence, we obtain an polynomial time algorithm for the calculation of  $u_{\text{GW},\infty}$ . See Section 6.1.2 for details.

The representations of  $u_{\text{GH}}$  in Theorem 2.5 and  $u_{\text{GW},\infty}$  in Theorem 3.13 strongly resemble themselves. As a direct consequence of both Theorem 2.5 and Theorem 3.13, we obtain the following comparison between the two metrics

**Corollary 3.15.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, it holds that*

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) \geq u_{\text{GH}}(X, Y). \quad (21)$$

The inequality in Equation (21) is sharp and we illustrate this as follows. By Mémoli et al. [70, Corollary 5.8] we know that if the considered ultrametric spaces  $(X, u_X)$  and  $(Y, u_Y)$  have different diameters (w.l.o.g.  $\text{diam}(X) < \text{diam}(Y)$ ), then  $u_{\text{GH}}(X, Y) = \text{diam}(Y)$ . The same statement also holds for  $u_{\text{GW},\infty}$ .

**Corollary 3.16.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  be such that  $\text{diam}(X) < \text{diam}(Y)$ . Then,*

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = \text{diam}(Y) = u_{\text{GH}}(X, Y).$$

*Proof.* The rightmost equality follows directly from Corollary 5.8 of Mémoli et al. [70]. As for the leftmost equality, let  $t := \text{diam}(Y)$ , then it is obvious that  $\mathcal{X}_t \cong_w * \cong_w \mathcal{Y}_t$ , where  $*$  denotes the one point ultrametric measure space. Let  $s \in (\text{diam}(X), \text{diam}(Y))$ , then  $\mathcal{X}_t \cong_w *$  whereas  $\mathcal{Y} \not\cong_w *$ . By Theorem 3.13,  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = t = \text{diam}(Y)$ .  $\square$

**3.3. The relation between  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ .** In this section, we study the relation of  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$  and establish the topological equivalence between the two metrics.

**3.3.1. Lipschitz relation.** We first study the Lipschitz relation between  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$ . For this purpose, we have to distinguish the cases  $p < \infty$  and  $p = \infty$ .

*The case  $p < \infty$ .* We start the consideration of this case by proving that it is essentially enough to consider the case  $p = 1$  (see Theorem 3.17). To this end, we need to introduce some notation. For each  $\alpha > 0$ , we define a function  $S_\alpha : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  by  $x \mapsto x^\alpha$ . Given an ultrametric space  $(X, u_X)$  and  $\alpha > 0$ , we abuse the notation and denote by  $S_\alpha(X)$  the new space  $(X, S_\alpha \circ u_X)$ . It is obvious that  $S_\alpha(X)$  is still an ultrametric space. This transformation of metric spaces is also known as the *snowflake transform* [26]. Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  denote two ultrametric measure spaces. Let  $1 \leq p < \infty$ . We denote by  $S_p(\mathcal{X})$  the ultrametric measure space  $(X, S_p \circ u_X, \mu_X)$ . The snowflake transform can be used to relate  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  as well as  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  with  $u_{\text{GW},1}(S_p(\mathcal{X}), S_p(\mathcal{Y}))$  and  $u_{\text{GW},1}^{\text{sturm}}(S_p(\mathcal{X}), S_p(\mathcal{Y}))$ , respectively.

**Theorem 3.17.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  and let  $p \in [1, \infty)$ . Then, we obtain*

$$(u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}))^p = u_{\text{GW},1}(S_p(\mathcal{X}), S_p(\mathcal{Y})) \quad \text{and} \quad (u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}))^p = u_{\text{GW},1}^{\text{sturm}}(S_p(\mathcal{X}), S_p(\mathcal{Y})).$$

We give full proof of Theorem 3.17 in Appendix B.2.4. Based on this result, we can directly relate the metrics  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$  by only considering the case  $p = 1$  and prove the following Theorem 3.18 (see Appendix B.3.1 for a detailed proof).

**Theorem 3.18.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, we have for  $p \in [1, \infty)$  that*

$$u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \leq 2^{\frac{1}{p}} u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}).$$

The subsequent example verifies that the coefficient in Theorem 3.18 is tight.

**Example 3.19.** For each  $n \in \mathbb{N}$ , let  $\mathcal{X}_n$  be the three-point space  $\Delta_3(1)$  (i.e. the 3-point metric labeled by  $\{x_1, x_2, x_3\}$  where all distances are 1) with a probability measure  $\mu_X^n$  such that  $\mu_X^n(x_1) = \mu_X^n(x_2) = \frac{1}{2n}$  and  $\mu_X^n(x_3) = 1 - \frac{1}{n}$ . Let  $Y = *$  and  $\mu_Y$  be the only probability measure on  $Y$ . Then, it is routine (using Proposition B.23 from Appendix B.5.3) to check that  $u_{\text{GW},1}(\mathcal{X}_n, \mathcal{Y}) = \frac{2}{n} \left(1 - \frac{3}{4n}\right)$  and  $u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}_n, \mathcal{Y}) = \frac{1}{n}$ . Therefore, we have

$$\lim_{n \rightarrow \infty} \frac{u_{\text{GW},1}(\mathcal{X}_n, \mathcal{Y})}{u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}_n, \mathcal{Y})} = 2.$$

**Example 3.20** ( $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$  are not bi-Lipschitz equivalent). Following [67, Remark 5.17], we verify in Appendix B.3.2 that for any positive integer  $n$

$$u_{\text{GW},p}^{\text{sturm}}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \geq \frac{1}{4} \text{ and } u_{\text{GW},p}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \leq \left(\frac{3}{2n}\right)^{\frac{1}{p}}.$$

Here,  $\hat{\Delta}_n(1)$  denotes the  $n$ -point metric measure space with interpoint distance 1 and the uniform probability measure. Thus, there exists no constant  $C > 0$  such that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq C \cdot u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  holds for every input spaces  $\mathcal{X}$  and  $\mathcal{Y}$ . Hence,  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$  are not bi-Lipschitz equivalent.

*The case  $p = \infty$ .* Next, we consider the relation between  $u_{\text{GW},\infty}^{\text{sturm}}$  and  $u_{\text{GW},\infty}$ . By taking the limit  $p \rightarrow \infty$  in Theorem 3.18, one might expect that  $u_{\text{GW},\infty}^{\text{sturm}} \geq u_{\text{GW},\infty}$ . In fact, we prove that the equality holds (for the full proof see Appendix B.3.3).

**Theorem 3.21.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Then, it holds that*

$$u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}).$$

One application of Theorem 3.21 is to explicitly derive the minimizing pair  $(A, \phi) \in \mathcal{A}^*$  in Equation (31) for  $p = \infty$  (see Appendix B.3.4 for an explicit construction):

**Theorem 3.22.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Let  $s := u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  and assume that  $s > 0$ . Then, there exists  $(A, \phi) \in \mathcal{A}$  defined in Equation (19) such that*

$$u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = d_{\text{W},\infty}^{Z_A}(\mu_X, \mu_Y),$$

where  $Z_A$  denotes the ultrametric space defined in Section 3.1.1.

**3.3.2. Topological equivalence between  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ .** Mémoli [67] proved the topological equivalence between  $d_{\text{GW},p}$  and  $d_{\text{GW},p}^{\text{sturm}}$ . We establish an analogous result for  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ . To this end, we recall the *modulus of mass distribution*.

**Definition 3.23** (Greven et al. [39, Def. 2.9]). Given  $\delta > 0$  we define the *modulus of mass distribution* of  $\mathcal{X} \in \mathcal{U}^w$  as

$$v_\delta(\mathcal{X}) := \inf \{ \varepsilon > 0 \mid \mu_X(\{x : \mu_X(B_\varepsilon^\circ(x)) \leq \delta\}) \leq \varepsilon \}, \quad (22)$$

where  $B_\varepsilon^\circ(x)$  denotes the open ball centered at  $x$  with radius  $\varepsilon$ .

We note that  $v_\delta(\mathcal{X})$  is non-decreasing, right-continuous and bounded above by 1. Furthermore, it holds that  $\lim_{\delta \searrow 0} v_\delta(\mathcal{X}) = 0$  [39, Lemma 6.5]. With Definition 3.23 at hand, we derive the following theorem.

**Theorem 3.24.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ ,  $p \in [1, \infty)$  and  $\delta \in (0, \frac{1}{2})$ . Then, whenever  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) < \delta^5$  we have*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq (4 \cdot \min(v_\delta(\mathcal{X}), v_\delta(\mathcal{Y})) + \delta)^{\frac{1}{p}} \cdot M,$$

where  $M := 2 \cdot \max(\text{diam}(X), \text{diam}(Y)) + 54$ .

**Remark 3.25.** Since it holds that  $\lim_{\delta \searrow 0} v_\delta(\mathcal{X}) = 0$  and that  $2^{-1/p} u_{\text{GW},p}^{\text{sturm}} \geq u_{\text{GW},p}$  (see Theorem 3.18), the above theorem gives the topological equivalence between  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ ,  $1 \leq p < \infty$  (the topological equivalence between  $u_{\text{GW},\infty}^{\text{sturm}}$  and  $u_{\text{GW},\infty}$  holds trivially thanks to Theorem 3.21).

The proof of the Theorem 3.24 follows the same strategy used for proving Proposition 5.3 in [67] and we refer to Appendix B.3.5 for the details.

**3.4. Topological and geodesic properties.** In this section, we consider the topology induced by  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$  on  $\mathcal{U}^w$  and discuss the geodesic properties of both  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$  for  $1 \leq p \leq \infty$ .

**3.4.1. Completeness and separability.** We study completeness and separability of the two metrics  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$ ,  $1 \leq p \leq \infty$ , on  $\mathcal{U}^w$ . To this end, we derive the subsequent theorem whose proof is postponed to Appendix B.4.1.

**Theorem 3.26.** (1) *For  $p \in [1, \infty)$ , the metric space  $(\mathcal{U}^w, u_{\text{GW},p})$  is neither complete nor separable.*

(2) *For  $p \in [1, \infty)$ , the metric space  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is neither complete nor separable.*

(3)  *$(\mathcal{U}^w, u_{\text{GW},\infty}) = (\mathcal{U}^w, u_{\text{GW},\infty}^{\text{sturm}})$  is complete but not separable.*

**3.4.2. Geodesic property.** A *geodesic* in a metric space  $(X, d_X)$  is a continuous function  $\gamma : [0, 1] \rightarrow X$  such that for each  $s, t \in [0, 1]$ ,  $d_X(\gamma(s), \gamma(t)) = |s - t| \cdot d_X(\gamma(0), \gamma(1))$ . We say a metric space is geodesic if for any two distinct points  $x, x' \in X$ , there exists a geodesic  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = x'$ . For any  $p \in [1, \infty)$ , the notion of  $p$ -geodesic is introduced in [70]: A  $p$ -geodesic in a metric space  $(X, d_X)$  is a continuous function  $\gamma : [0, 1] \rightarrow X$  such that for each  $s, t \in [0, 1]$ ,  $d_X(\gamma(s), \gamma(t)) = |s - t|^{1/p} \cdot d_X(\gamma(0), \gamma(1))$ . Similarly, we say a metric space is  $p$ -geodesic if for any two distinct points  $x, x' \in X$ , there exists a  $p$ -geodesic  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = x'$ . Note that a 1-geodesic is a usual geodesic and a 1-geodesic space is a usual geodesic space. The subsequent theorem establishes  $(p)$ -geodesic properties of  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  for  $p \in [1, \infty)$ . A full proof is given in Appendix B.4.2.

**Theorem 3.27.** *For any  $p \in [1, \infty)$ , the space  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is  $p$ -geodesic.*

**Remark 3.28.** Due to the fact that a  $p$ -geodesic space cannot be geodesic when  $p > 1$  (cf. Lemma B.15),  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is not geodesic for all  $p > 1$ .

**Remark 3.29.** Though the geodesic properties of  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$ ,  $1 \leq p < \infty$  are clear, we remark that geodesic properties of  $(\mathcal{U}^w, u_{\text{GW},p})$ ,  $1 \leq p < \infty$ , still remain unknown to us.

**Remark 3.30** (The case  $p = \infty$ ). Being an ultrametric space itself (cf. Theorem 3.10),  $(\mathcal{U}^w, u_{\text{GW},\infty})$  ( $= (\mathcal{U}^w, u_{\text{GW},\infty}^{\text{sturm}})$ ) is *totally disconnected*, i.e., any subspace with at least two elements is disconnected [89]. This in turn implies that each continuous curve in  $(\mathcal{U}^w, u_{\text{GW},\infty})$  is constant. Therefore,  $(\mathcal{U}^w, u_{\text{GW},\infty})$  is not a  $p$ -geodesic space for any  $p \in [1, \infty)$ .

#### 4. LOWER BOUNDS FOR $u_{\text{GW},p}$

Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be two ultrametric measure spaces. The metrics  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$  respect the ultrametric structure of the spaces  $\mathcal{X}$  and  $\mathcal{Y}$ . Thus, one would hope that comparing ultrametric measure spaces with  $u_{\text{GW},p}^{\text{sturm}}$  or  $u_{\text{GW},p}$  is more meaningful than doing it with the usual Gromov-Wasserstein distance or Sturm's distance. Unfortunately, for  $p < \infty$ , the computation of both  $u_{\text{GW},p}^{\text{sturm}}$  and  $u_{\text{GW},p}$  is complicated and for  $p = \infty$  both metrics are extremely sensitive to differences in the diameters of the considered spaces (see Corollary 3.16). Thus, it is not feasible to use these metrics in many applications. However, we can derive meaningful lower bounds for  $u_{\text{GW},p}$  (and hence also for  $u_{\text{GW},p}^{\text{sturm}}$ ) that resemble those of the Gromov-Wasserstein distance. Naturally, the question arises whether these lower bounds are better/sharper than the ones of the usual Gromov-Wasserstein distance in this setting. This question is addressed throughout this section and will be readdressed in Section 6 as well as Section 7.

In [67], the author introduced three lower bounds for  $d_{\text{GW},p}$  that are computationally less expensive than the calculation of  $d_{\text{GW},p}$ . We will briefly review these three lower bounds and then define candidates for the corresponding lower bounds for  $u_{\text{GW},p}$ . In the following, we always assume  $p \in [1, \infty]$ .

**First lower bound.** Let  $s_{X,p} : X \rightarrow \mathbb{R}_{\geq 0}$ ,  $x \mapsto \|u_X(x, \cdot)\|_{L^p(\mu_X)}$ . Then, the first lower bound  $\text{FLB}_p(\mathcal{X}, \mathcal{Y})$  for  $d_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  is defined as follows

$$\text{FLB}_p(\mathcal{X}, \mathcal{Y}) := \frac{1}{2} \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \|\Lambda_1(s_{X,p}(\cdot), s_{Y,p}(\cdot))\|_{L^p(\mu)}.$$

Following our intuition of replacing  $\Lambda_1$  with  $\Lambda_\infty$ , we define the ultrametric version of **FLB** as

$$\text{FLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) := \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \|\Lambda_\infty(s_{X,p}(\cdot), s_{Y,p}(\cdot))\|_{L^p(\mu)}.$$

**Second lower bound.** The second lower bound  $\text{SLB}_p(\mathcal{X}, \mathcal{Y})$  for  $d_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  is given as

$$\text{SLB}_p(\mathcal{X}, \mathcal{Y}) := \frac{1}{2} \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \|\Lambda_1(u_X, u_Y)\|_{L^p(\gamma)}.$$

Thus, we define the ultrametric second lower bound between two ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$  as follows:

$$\text{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) := \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \|\Lambda_\infty(u_X, u_Y)\|_{L^p(\gamma)}.$$

**Third lower bound.** Before we introduce the final lower bound, we have to define several functions. First, let  $\Gamma_{X,Y}^1 : X \times Y \times X \times Y \rightarrow \mathbb{R}_{\geq 0}$ ,  $(x, y, x', y') \mapsto \Lambda_1(u_X(x, x'), u_Y(y, y'))$  and let  $\Omega_p^1 : X \times Y \rightarrow \mathbb{R}_{\geq 0}$ ,  $p \in [1, \infty]$ , be given by

$$\Omega_p^1(x, y) := \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| \Gamma_{X,Y}^1(x, y, \cdot, \cdot) \right\|_{L^p(\mu)}.$$

Then, the third lower bound  $\mathbf{TLB}_p$  is given as

$$\mathbf{TLB}_p(\mathcal{X}, \mathcal{Y}) := \frac{1}{2} \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| \Omega_p^1(\cdot, \cdot) \right\|_{L^p(\mu)}.$$

Analogously to the definition of previous ultrametric versions, we define  $\Gamma_{X,Y}^\infty : X \times Y \times X \times Y \rightarrow \mathbb{R}_{\geq 0}$ ,  $(x, y, x', y') \mapsto \Lambda_\infty(u_X(x, x'), u_Y(y, y'))$ . Further, for  $p \in [1, \infty]$ , let  $\Omega_p^\infty : X \times Y \rightarrow \mathbb{R}_{\geq 0}$  be given by

$$\Omega_p^\infty(x, y) := \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| \Gamma_{X,Y}^\infty(x, y, \cdot, \cdot) \right\|_{L^p(\mu)}.$$

Then, the ultrametric third lower bound between two ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$  is defined as

$$\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) := \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| \Omega_p^\infty(\cdot, \cdot) \right\|_{L^p(\mu)}.$$

**4.1. Properties and computation of the lower bounds.** Next, we examine the quantities  $\mathbf{FLB}^{\text{ult}}$ ,  $\mathbf{SLB}^{\text{ult}}$  and  $\mathbf{TLB}^{\text{ult}}$  more closely. Since  $\Lambda_\infty(a, b) \geq \Lambda_1(a, b) = |a - b|$  for any  $a, b \geq 0$ , it is easy to conclude that  $\mathbf{FLB}_p^{\text{ult}} \geq \mathbf{FLB}_p$ ,  $\mathbf{SLB}_p^{\text{ult}} \geq \mathbf{SLB}_p$  and  $\mathbf{TLB}_p^{\text{ult}} \geq \mathbf{TLB}_p$ . Moreover, the three ultrametric lower bounds satisfy the following theorem (for a complete proof see Appendix C.1.1).

**Theorem 4.1.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  and let  $p \in [1, \infty]$ .*

- (1)  $u_{\text{GW}, \infty}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{FLB}_\infty^{\text{ult}}(\mathcal{X}, \mathcal{Y})$ .
- (2)  $u_{\text{GW}, p}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y})$ .

**Remark 4.2.** Interestingly, it turns out that  $\mathbf{FLB}_p^{\text{ult}}$  is not a lower bound of  $u_{\text{GW}, p}$  in general when  $p < \infty$ . For example, let  $X = \{x_1, x_2, \dots, x_n\}$  and  $Y = \{y_1, \dots, y_n\}$  and define  $u_X$  such that  $u_X(x_1, x_2) = 1$  and  $u_X(x_i, x_j) = 2\delta_{i \neq j}$  for  $(i, j) \neq (1, 2)$ ,  $(i, j) \neq (2, 1)$  and  $i, j = 1, \dots, n$ . Let  $u_Y(y_i, y_j) = 2\delta_{i \neq j}$ ,  $i, j = 1, \dots, n$ , and let  $\mu_X$  and  $\mu_Y$  be uniform measures on  $X$  and  $Y$ , respectively. Then,  $u_{\text{GW}, 1}(\mathcal{X}, \mathcal{Y}) \leq \frac{4}{n^2}$  whereas  $\mathbf{FLB}_1^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \frac{4n-4}{n^2}$  which is greater than  $u_{\text{GW}, 1}(\mathcal{X}, \mathcal{Y})$  as long as  $n > 2$ . Moreover, we have in this case that  $\mathbf{FLB}_1^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = O\left(\frac{1}{n}\right)$  whereas  $u_{\text{GW}, 1}(\mathcal{X}, \mathcal{Y}) = O\left(\frac{1}{n^2}\right)$ . Hence, there exists no constant  $C > 0$  such that  $\mathbf{FLB}_1^{\text{ult}} \leq C \cdot u_{\text{GW}, 1}$  in general.

**Remark 4.3.** There exist ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$  such that  $\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = 0$  whereas  $u_{\text{GW}, p}(\mathcal{X}, \mathcal{Y}) > 0$  (examples described in [67, Figure 8] will serve the purpose). Furthermore, there are spaces  $\mathcal{X}$  and  $\mathcal{Y}$  such that  $\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = 0$  whereas  $\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) > 0$  (see Appendix C.1.3). The analogous statement holds true for  $\mathbf{TLB}_p$  and  $\mathbf{SLB}_p$ , which are nevertheless useful in various applications (see e.g. [37]).

From the structure of  $\mathbf{SLB}_p^{\text{ult}}$  and  $\mathbf{TLB}_p^{\text{ult}}$  it is obvious that their computations leads to different optimal transport problems (see e.g. [99]). However, in analogy to Chowdhury and Mémoli [23, Theorem 3.1] we can rewrite  $\mathbf{SLB}_p^{\text{ult}}$  and  $\mathbf{TLB}_p^{\text{ult}}$  in order to further simplify their computation. The full proof of the subsequent proposition is given in Appendix C.1.2.

**Proposition 4.4.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  and let  $p \in [1, \infty]$ . Then, we find that*

- (1)  $\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = d_{\mathbf{W}, p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}((u_X)_\#(\mu_X \otimes \mu_X), (u_Y)_\#(\mu_Y \otimes \mu_Y));$
- (2) *For each  $x, y \in X \times Y$ ,  $\Omega_p^\infty(x, y) = d_{\mathbf{W}, p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(u_X(x, \cdot)_\# \mu_X, u_Y(y, \cdot)_\# \mu_Y).$*

**Remark 4.5.** Since we have by Theorem 2.9 an explicit formula for the Wasserstein distance on  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  between finitely supported probability measures, these alternative representations of the lower bound  $\mathbf{SLB}_p^{\text{ult}}$  and the cost functional  $\Omega_p^\infty$  drastically reduce the computation time of  $\mathbf{SLB}_p^{\text{ult}}$  and  $\mathbf{TLB}_p^{\text{ult}}$ , respectively. In particular, we note that this allows us to compute  $\mathbf{SLB}_p^{\text{ult}}$ ,  $1 \leq p \leq \infty$ , between finite ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$  with  $|X| = m$  and  $|Y| = n$  in  $O((m \vee n)^2)$  steps.

Proposition 4.4 allows us to directly compare the two lower bounds  $\mathbf{SLB}_1^{\text{ult}}$  and  $\mathbf{SLB}_1$ .

**Corollary 4.6.** *For any finite ultrametric measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$ , we have that*

$$\mathbf{SLB}_1^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \mathbf{SLB}_1(\mathcal{X}, \mathcal{Y}) + \frac{1}{2} \int_{\mathbb{R}} t |(u_X)_\#(\mu_X \otimes \mu_X) - (u_Y)_\#(\mu_Y \otimes \mu_Y)| (dt). \quad (23)$$

*Proof.* The claim follows directly from Proposition 4.4 and Remark 2.10.  $\square$

This corollary implies that  $\mathbf{SLB}_p^{\text{ult}}$  is more rigid than  $\mathbf{SLB}_p$ , since the second summand on the right hand side of Equation (23) is sensitive to distance perturbations. This is also illustrated very well in the subsequent example.

**Example 4.7.** Recall notations from Example 3.5. For any  $d, d' > 0$ , we let  $X := \Delta_2(d)$  and let  $Y := \Delta_2(d')$ . Assume that  $X$  and  $Y$  have underlying sets  $\{x_1, x_2\}$  and  $\{y_1, y_2\}$ , respectively. Define  $\mu_X \in \mathcal{P}(X)$  and  $\mu_Y \in \mathcal{P}(Y)$  as follows. Let  $\alpha_1, \alpha_2 \geq 0$  be such that  $\alpha_1 + \alpha_2 = 1$ . Let  $\mu_X(x_1) = \mu_Y(y_1) := \alpha_1$  and let  $\mu_X(x_2) = \mu_Y(y_2) := \alpha_2$ . Then, it is easy to verify that

- (1)  $u_{\mathbf{GW}, 1}(\mathcal{X}, \mathcal{Y}) = \mathbf{SLB}_1^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = 2\alpha_1\alpha_2\Lambda_\infty(d, d').$
- (2)  $d_{\mathbf{GW}, 1}(\mathcal{X}, \mathcal{Y}) = \mathbf{SLB}_1(\mathcal{X}, \mathcal{Y}) = \alpha_1\alpha_2\Lambda_1(d, d') = \alpha_1\alpha_2|d - d'|.$
- (3)  $\frac{1}{2} \int_{\mathbb{R}} t |(u_X)_\#(\mu_X \otimes \mu_X) - (u_Y)_\#(\mu_Y \otimes \mu_Y)| (dt) = \alpha_1\alpha_2(d + d')\delta_{d \neq d'}.$

From 1 and 2 we observe that both second lower bounds are tight. Moreover, since we obviously have that  $(d + d')\delta_{d \neq d'} + |d - d'| = 2\Lambda_\infty(d, d')$ , we have also verified Equation (23) through this example. Unlike  $\mathbf{SLB}_1(\mathcal{X}, \mathcal{Y})$  being proportional to  $|d - d'|$ , as long as  $d \neq d'$ , even if  $|d - d'|$  is small,  $\Lambda_\infty(d, d') = \max(d, d')$  which results in a large value of  $\mathbf{SLB}_1^{\text{ult}}(\mathcal{X}, \mathcal{Y})$  when  $d$  and  $d'$  are large numbers. This example illustrates that  $\mathbf{SLB}_1^{\text{ult}}$  (and hence  $u_{\mathbf{GW}, 1}$ ) is rigid with respect to distance perturbation.

5.  $u_{\text{GW},p}$  ON ULTRA-DISSIMILARITY SPACES

A natural generalization of ultrametric spaces is provided by *ultra-dissimilarity spaces*. These spaces naturally occur when working with symmetric ultranetworks (see [91]) or phylogenetic tree data (see [90]). In this section, we will introduce these spaces and briefly illustrate to what extent the results for  $u_{\text{GW},p}$  can be adapted for ultra-dissimilarity measure spaces. We start by formally introducing *ultra-dissimilarity spaces*.

**Definition 5.1** (Ultra-dissimilarity spaces). An *ultra-dissimilarity space* is a couple  $(X, u_X)$  consisting of a set  $X$  and a function  $u_X : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying the following conditions for any  $x, y, z \in X$ :

- (1)  $u_X(x, y) = u_X(y, x)$ ;
- (2)  $u_X(x, y) \leq \max(u_X(x, z), u_X(z, y))$ ;
- (3)  $\max(u_X(x, x), u_X(y, y)) \leq u_X(x, y)$  and the equality holds if and only if  $x = y$ .

**Remark 5.2.** Note that when  $(X, u_X)$  is an ultrametric space the third condition is trivially satisfied.

In the following, we restrict ourselves to *finite* ultra-dissimilarity spaces to avoid technical issues in topology (see [22, 23] for a more complete treatment of infinite spaces). One important aspect of ultra-dissimilarity spaces is the connection with the so-called *treegrams* [70, 91], which can be regarded as generalized dendrograms. For a finite set  $X$ , let  $\mathbf{SubPart}(X)$  denote the collection of all *subpartitions* of  $X$ : Any partition  $P'$  of a non-empty subset  $X' \subseteq X$  is called a subpartition of  $X$ . Given two subpartitions  $P_1, P_2$ , we say  $P_1$  is coarser than  $P_2$  if each block in  $P_2$  is contained in some block in  $P_1$ .

**Definition 5.3** (Treegrams). A *treegram*  $T_X : [0, \infty) \rightarrow \mathbf{SubPart}(X)$  is a map parametrizing a nested family of subpartitions over the same set  $X$  and satisfying the following conditions:

- (1) For any  $0 \leq s < t < \infty$ ,  $T_X(t)$  is coarser than  $T_X(s)$ ;
- (2) There exists  $t_X > 0$  such that for any  $t \geq t_X$ ,  $T_X(t) = \{X\}$ ;
- (3) For each  $t \geq 0$ , there exists  $\varepsilon > 0$  such that  $T_X(t) = T_X(t')$  for all  $t' \in [t, t + \varepsilon]$ ;
- (4) For each  $x \in X$ , there exists  $t_x \geq 0$  such that  $\{x\}$  is a block in  $T_X(t_x)$ .

Similar to Theorem 2.2, which correlates ultrametrics to dendrograms, there exists an equivalence relation between ultra-dissimilarity functions and treegrams on a finite set (see Figure 5 for an illustration).

**Proposition 5.4** (Smith et al. [91]). Given a finite set  $X$ , denote by  $\mathcal{U}_{\text{dis}}(X)$  the collection of all ultrametric dissimilarity functions on  $X$  and by  $\mathcal{T}(X)$  the collection of all treegrams over  $X$ . Then, there exists a bijection  $\Delta_X : \mathcal{T}(X) \rightarrow \mathcal{U}_{\text{dis}}(X)$ .

An *ultra-dissimilarity measure space* is a triple  $\mathcal{X} = (X, u_X, \mu_X)$  where  $(X, u_X)$  is an ultra-dissimilarity space and  $\mu_X$  is a probability measure fully supported on  $X$ . Just as for metric




FIGURE 5. **Treegrams:** Relation between ultra-dissimilarity functions and treegrams

spaces or metric measure spaces, it is important to have a notion of isomorphism between ultra-dissimilarity spaces.

**Definition 5.5** (Isomorphism). Given two ultra-dissimilarity measure spaces  $\mathcal{X}$  and  $\mathcal{Y}$ , we say they are *isomorphic*, denoted  $\mathcal{X} \cong_w \mathcal{Y}$ , if there is a bijective function  $f : X \rightarrow Y$  such that  $f_{\#}\mu_X = \mu_Y$  and for any  $x, x' \in X$  it holds  $u_Y(f(x), f(x')) = u_X(x, x')$ . The collection of all isomorphism classes of ultra-dissimilarity spaces is denoted by  $\mathcal{U}_{\text{dis}}^w$ .

Given the previous results it is straightforward to show that  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , is a metric on the isomorphism classes of  $\mathcal{U}_{\text{dis}}^w$ . For the complete proof of the subsequent statement, we refer to Appendix D.1.1.

**Theorem 5.6.** *The ultrametric Gromov-Wasserstein distance  $u_{\text{GW},p}$  is a  $p$ -metric on  $\mathcal{U}_{\text{dis}}^w$ .*

**Remark 5.7.** Since  $u_{\text{GW},p}$  translates to a metric on  $\mathcal{U}_{\text{dis}}^w$ , it is clear that it admits the lower bounds introduced in Section 4.

## 6. COMPUTATIONAL ASPECTS

In this section, we investigate algorithms for approximating/calculating  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ . Furthermore, we evaluate for  $p < \infty$  the performance of the computationally efficient lower bound  $\text{SLB}^{\text{ult}}$  introduced in Section 4 and compare our findings to the results of the classical Gromov-Wasserstein distance  $d_{\text{GW},p}$  (see Equation (7)). Matlab implementations of the presented algorithms and comparisons are available at <https://github.com/ndag/uGW>.

**6.1. Algorithms.** Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be two finite ultrametric measure spaces with cardinalities  $m$  and  $n$ , respectively.

**6.1.1. The case  $p < \infty$ .** We have already noted in Remark 3.12 that calculating  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  for  $p < \infty$  yields a non-convex quadratic program (which is an NP-hard problem in general [78]). Solving this is not feasible in practice. However, in many practical applications it is sufficient to work with good approximations. Therefore, we propose to approximate  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  for  $p < \infty$  via conditional gradient descent. To this end, we note that the

gradient  $G$  that arises from Equation (12) can in the present setting be expressed with the following partial derivative with respect to  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$

$$G_{i,j} = 2 \sum_{k=1}^m \sum_{l=1}^n (\Lambda_\infty(u_X(x_i, x_k), u_Y(y_j, y_l)))^p \mu_{kl}, \quad \forall 1 \leq i \leq m, 1 \leq j \leq n. \quad (24)$$

As we deal with a non-convex minimization problem, the performance of the gradient descent strongly depends on the starting coupling  $\mu^{(0)}$ . Therefore, we follow the suggestion of Chowdhury and Needham [24] and employ a Markov Chain Monte Carlo Hit-And-Run sampler to obtain multiple random start couplings. Running the gradient descent from each point in this ensemble greatly improves the approximation in many cases. For a precise description of the proposed procedure, we refer to Algorithm 1.

---

**Algorithm 1**  $u_{\text{GW},p}(X, Y, p, N, L)$

---

```

//Create a list of random couplings
couplings = CreateRandomCouplings(N)
stat_points = cell(N)
for i=1:N do
     $\mu^{(0)} = \text{couplings}\{i\}$ 
    for j=1:L do
         $G = \text{Gradient from Equation (24) w.r.t. } \mu^{(j-1)}$ 
         $\tilde{\mu}^{(j)} = \text{Solve OT with ground loss } G$ 
         $\gamma^{(j)} = \frac{2}{j+2}$ 
        //Alt. find  $\gamma \in [0, 1]$  that minimizes  $\text{dis}_p^{\text{ult}}(\mu^{(j-1)} + \gamma(\tilde{\mu}^{(j)} - \mu^{(j-1)}))$ 
         $\mu^{(j)} = (1 - \gamma^{(j)})\mu^{(j-1)} + \gamma^{(j)}\tilde{\mu}^{(j)}$ 
    end for
    stat_points{i} =  $\mu^{(L)}$ 
end for
Find  $\mu^*$  in stat_points that minimizes  $\text{dis}_p^{\text{ult}}(\mu)$ 
result =  $\text{dis}_p^{\text{ult}}(\mu^*)$ 

```

---

6.1.2. *The case  $p = \infty$ .* For  $p = \infty$ , it follows by Theorem 3.13 that

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}. \quad (25)$$

This identity allows us to construct a polynomial time algorithm for  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$  based on the ideas of Mémoli et al. [70, Sec. 8.2.2]. More precisely, let  $\text{spec}(X) := \{u_X(x, x') \mid x, x' \in X\}$  denote the spectrum of  $X$ . Then, it is evident that in order to find the infimum in Equation (25), we only have to check  $\mathcal{X}_t \cong_w \mathcal{Y}_t$  for each  $t \in \text{spec}(X) \cup \text{spec}(Y)$ , starting from the largest to the smallest and  $u_{\text{GW},\infty}$  is given as the smallest  $t$  such that  $\mathcal{X}_t \cong_w \mathcal{Y}_t$ . This can be done in polynomial time by considering  $\mathcal{X}_t$  and  $\mathcal{Y}_t$  as labeled, weighted trees (e.g. by using a slight modification of the algorithm in Example 3.2 of [3]). This gives rise to a simple algorithm (see Algorithm 2) to calculate  $u_{\text{GW},\infty}$ .




FIGURE 6. **Ultrametric measure spaces:** Four non-isomorphic ultrametric measure spaces denoted (from left to right) as  $\mathcal{X}_i = (X_i, d_{X_i}, \mu_{X_i})$ ,  $1 \leq i \leq 4$ .

---

**Algorithm 2**  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$

---

```

spec = sort(spec(X) ∪ spec(Y), 'descent')
for i = 1 : length(spec) do
    t = spec(i)
    if  $\mathcal{X}_t \not\equiv_w \mathcal{Y}_t$  then
        return spec(i - 1)
    end if
end for
return 0

```

---

**6.2. The relation between  $u_{\text{GW},1}$ ,  $u_{\text{GW},\infty}$  and  $\text{SLB}_1^{\text{ult}}$ .** In order to understand how  $u_{\text{GW},p}$  (or at least its approximation),  $u_{\text{GW},\infty}$  and  $\text{SLB}_p^{\text{ult}}$  are influenced by small changes in the structure of the considered ultrametric measure spaces, we exemplarily consider the ultrametric measure spaces  $\mathcal{X}_i = (X_i, d_{X_i}, \mu_{X_i})$ ,  $1 \leq i \leq 4$ , displayed in Figure 6. These ultrametric measure spaces differ only by one characteristic (e.g. one side length or the equipped measure). Exemplarily, we calculate  $u_{\text{GW},1}(\mathcal{X}_i, \mathcal{X}_j)$  (approximated with Algorithm 1, where  $L = 5000$  and  $N = 40$ ),  $\text{SLB}_1^{\text{ult}}(\mathcal{X}_i, \mathcal{X}_j)$  and  $u_{\text{GW},\infty}(\mathcal{X}_i, \mathcal{X}_j)$ ,  $1 \leq i, j \leq 4$ . The results suggest that  $\text{SLB}_1^{\text{ult}}$  and  $u_{\text{GW},1}$  are influenced by the change in the diameter of the spaces the most (see Table 2 and Table 3 in Appendix E.1 for the complete results). Changes in the metric influence  $\text{SLB}_1^{\text{ult}}$  in a similar fashion as  $u_{\text{GW},1}$ , while changes in the measure have less impact on  $\text{SLB}_1^{\text{ult}}$ . Further, we observe that  $u_{\text{GW},\infty}$  attains for almost all comparisons the maximal possible value. Only the comparison of  $\mathcal{X}_1$  with  $\mathcal{X}_3$ , where the only small scale structure of the space was changed, yields a value that is smaller than the maximum of the diameters of the considered spaces.

**6.3. Comparison of  $u_{\text{GW},1}$ ,  $\text{SLB}_1^{\text{ult}}$ ,  $d_{\text{GW},1}$  and  $\text{SLB}_1$ .** In the remainder of this section, we will demonstrate the differences between  $u_{\text{GW},1}$ ,  $\text{SLB}_1^{\text{ult}}$ ,  $d_{\text{GW},1}$  and  $\text{SLB}_1$ . To this end, we first compare the metric measure spaces in Figure 6 based on  $d_{\text{GW},1}$  and  $\text{SLB}_1$ . We observe that  $d_{\text{GW},1}$  (approximated in the same manner as  $u_{\text{GW},1}$ ) and  $\text{SLB}_1$  are hardly influenced by the differences between the ultrametric measure spaces  $\mathcal{X}_i$ ,  $1 \leq i \leq 4$ . In particular, it is remarkable that  $d_{\text{GW},1}$  is affected the most by the changes made to the measure and not the metric structure (see Table 4 in Appendix E.2 for the complete results).

Next, we consider the differences between the aforementioned quantities more generally. For this purpose, we generate 4 ultrametric spaces  $Z_k$ ,  $1 \leq k \leq 4$ , with totally different dendrogram structures, whose diameters are between 0.5 and 0.6 (for the precise construction of these spaces see Appendix E.2). For each  $t = 0, 0.2, 0.4, 0.6$ , we perturb each  $Z_k$  independently to generate 15 ultrametric spaces  $Z_{k,t}^i$ ,  $1 \leq i \leq 15$ , such that  $(Z_{k,t}^i)_t \equiv (Z_k)_t$  for all  $i$ . The spaces  $Z_{k,t}^i$  are called *perturbations of  $Z_k$  at level  $t$*  (see Figure 7 for an illustration and see Appendix E.2 for more details). The spaces  $Z_{k,t}^i$  are endowed with the uniform probability measure and we obtain a collection of ultrametric measure spaces  $\mathcal{Z}_{k,t}^i$ . Naturally, we refer to  $k$  as the class of the ultrametric measure space  $\mathcal{Z}_{k,t}^i$ . We compute for each  $t$  the quantities  $u_{\text{GW},1}$ ,  $\mathbf{SLB}_1^{\text{ult}}$ ,  $d_{\text{GW},1}$  and  $\mathbf{SLB}_1$  among the resulting 60 ultrametric measure spaces. The results, where the spaces have been ordered lexicographically by  $(k, i)$ , are visualized in Figure 8. As previously, we observe that  $u_{\text{GW},1}$  and  $\mathbf{SLB}_1^{\text{ult}}$  as well as  $d_{\text{GW},1}$  and  $\mathbf{SLB}_1$  behave in a similar manner. More precisely, we see that both  $d_{\text{GW},1}$  and  $\mathbf{SLB}_1$  discriminate well between the different classes and that their behavior does not change too much for an increasing level of perturbation. On the other hand,  $u_{\text{GW},1}$  and  $\mathbf{SLB}_1^{\text{ult}}$  are very sensitive to the level of perturbation. For small  $t$  they discriminate better than  $d_{\text{GW},1}$  and  $\mathbf{SLB}_1$  between the different classes and pick up clearly that the perturbed spaces differ. However, if the level of perturbation becomes too large both quantities start to discriminate between spaces from the same class (see Figure 8).




FIGURE 7. Randomly sampled ultrametric measure spaces: Illustration of  $Z_k$  for  $k = 2, 3, 4, 5$  (top row) and instances for perturbations of  $Z_4$  with respect to perturbation level  $t \in \{0, 0.2, 0.4, 0.6\}$  (bottom row).

In conclusion,  $u_{\text{GW},1}$  and  $\mathbf{SLB}_1^{\text{ult}}$  are sensitive to differences in the large scales of the considered ultrametric measure spaces. While this leads (from small  $t$ ) to good discrimination in the above example, it also highlights that they are (different from  $d_{\text{GW},1}$  and  $\mathbf{SLB}_1$ ) susceptible to large scale noise.




FIGURE 8.  $u_{\text{GW},1}/\text{SLB}_1^{\text{ult}}$  and  $d_{\text{GW},1}/\text{SLB}_1$  among randomly generated ultrametric measure spaces: Heatmap representations of  $\text{SLB}_1^{\text{ult}}(\mathcal{Z}_{n,t}^i, \mathcal{Z}_{n',t}^{i'})$  (top row),  $u_{\text{GW},1}(\mathcal{Z}_{n,t}^i, \mathcal{Z}_{n',t}^{i'})$  (second row),  $\text{SLB}_1(\mathcal{Z}_{n,t}^i, \mathcal{Z}_{n',t}^{i'})$  (third row) and  $d_{\text{GW},1}(\mathcal{Z}_{n,t}^i, \mathcal{Z}_{n',t}^{i'})$  (bottom row),  $k, k' \in \{2, \dots, 5\}$  and  $i, i' \in \{1, \dots, 15\}$ .

## 7. PHYLOGENETIC TREE SHAPES

Rooted phylogenetic trees (for a formal definition see e.g., [90]) are a common tool to visualize and analyze the evolutionary relationship between different organisms. In combination with DNA sequencing, they are an important tool to study the rapid evolution of different pathogens. It is well known that the (unweighted) shape of a phylogenetic tree, i.e., the tree's connectivity structure without referring to its labels or the length of its branches, carries important information about macroevolutionary processes (see e.g., [8, 27, 72, 104]). In order to study the evolution of and the relation between different pathogens, it is of great interest to compare the shapes of phylogenetic trees created on the basis of different data sets. Currently, the number of tools for performing phylogenetic tree shape comparison is quite limited and the development of new methods for this is an active field of research

[25, 49, 60, 73]. It is well known that certain classes of phylogenetic trees (as well as their respective tree shapes) can be identified as ultrametric spaces [90, Sec. 7]. On the other hand, general phylogenetic trees are closely related to treegrams (see Definition 5.3). In the following, we will use this connection and demonstrate exemplarily that the computationally efficient lower bound  $\mathbf{SLB}_1^{\text{ult}}$  has some potential for comparing phylogenetic tree shapes. In particular, we contrast it to the metric defined for this application in Equation (4) of Colijn and Plazzotta [25], in the following denoted as  $d_{\text{CP},2}$ , and study the behavior of  $\mathbf{SLB}_1$  in this framework.




FIGURE 9. **Transforming a phylogenetic tree shape into an ultra-dissimilarity space:** In this figure, we illustrate the treegram corresponding to the ultra-dissimilarity space generated by Equation (26) with respect to the phylogenetic tree shape on the left. Note that the treegram preserves the tree structure and the smallest birth time of points is exactly 0.

In this section, we reconsider phylogenetic tree shape comparisons from Colijn and Plazzotta [25] and thereby study HA protein sequences from human influenza A (H3N2) (data downloaded from NCBI on 22 January 2016). More precisely, we investigate the relation between two samples of size 200 of phylogenetic tree shapes with 500 tips. Phylogenetic trees from the first sample are based on a random subsample of size 500 of 2168 HA-sequences that were collected in the USA between March 2010 and September 2015, while trees from the second sample are based on a random subsample of size 500 of 1388 HA-sequences gathered in the tropics between January 2000 and October 2015 (for the exact construction of the trees see [25]). Although both samples of phylogenetic trees are based on HA protein sequences from human influenza A, we expect them to be quite different. On the one hand, influenza A is highly seasonal outside the tropics (where this seasonal variation is absent) with the majority of cases occurring in the winter [86]. On the other hand, it is well known that the undergoing evolution of the HA protein causes a ‘ladder-like’ shape of long-term influenza phylogenetic trees [51, 62, 101, 103] that is typically less developed in short term data sets. Thus, also the different collection period of the two data sets will most likely influence the respective phylogenetic tree shapes.

In order to compare the phylogenetic tree shapes of the resulting 400 trees, we have to transform the phylogenetic tree shapes into ultra-dissimilarity measure spaces  $\mathcal{X}_i = (X_i, u_{X_i}, \mu_{X_i})$ ,  $1 \leq i \leq 400$ . To this end, we discard all the labels, denote by  $X_i$  the tips of the  $i$ ’th phylogenetic tree and refer to the corresponding tree shape as  $\mathcal{T}_i$ . Next, we define the ultra-dissimilarities  $u_{X_i}$  on  $X_i$ ,  $1 \leq i \leq 400$ . For this purpose, we set all edge length in the

considered phylogenetic trees to one and construct  $u_{X_i}$  as follows: let  $x_1^i, x_2^i \in X_i$  and let  $a_{1,2}^i$  be the most recent common ancestor of  $x_1^i$  and  $x_2^i$ . Let  $d_{a_{1,2}}^i$  be the length of the shortest path from  $a_{1,2}^i$  to the root, let  $d_1^i$  be the length of the shortest path from  $x_1^i$  to the root and let  $d^i$  be the length of the longest shortest path from any tip to the root. Then, we define for any  $x_1^i, x_2^i \in X_i$

$$u_{X_i}(x_1^i, x_2^i) = \begin{cases} d^i - d_{a_{1,2}}^i & \text{if } x_1^i \neq x_2^i \\ d^i - d_1^i & \text{if } x_1^i = x_2^i, \end{cases} \quad (26)$$

and weight all tips in  $X_i$  equally (i.e.  $\mu_{X_i}$  is the uniform measure on  $X_i$ ). This naturally transforms the collection of phylogenetic tree shapes  $\mathcal{T}_i$ ,  $1 \leq i \leq 400$ , into a collection of ultra-dissimilarity spaces (see Figure 9 for an illustration), which allows us to directly apply  $\mathbf{SLB}_1^{\text{ult}}$  to compare them (once again we exemplarily choose  $p = 1$ ).

In Figure 10 we contrast our findings for the comparisons of the shapes  $\mathcal{T}_i$ ,  $1 \leq i \leq 400$ , to those obtained by computing the metric  $d_{\text{CP},2}$  described in [25]. The top row of Figure 10 visualizes the dissimilarity matrix for the comparisons of all 400 phylogenetic tree shapes (the first 200 entries correspond to the tree shapes from the US-influenza and the second 200 correspond to the ones from the tropic influenza) obtained by applying  $\mathbf{SLB}_1^{\text{ult}}$  as heat map (left) and as multidimensional scaling plot (right). The heat map shows that the collection of US trees is divided into a large group  $\mathcal{G}_1 := (\mathcal{T}_i)_{1 \leq i \leq 161}$ , that is well separated from the phylogenetic tree shapes based on tropical data  $\mathcal{G}_3 := (\mathcal{T}_i)_{201 \leq i \leq 400}$ , and a smaller subgroup  $\mathcal{G}_2 := (\mathcal{T}_i)_{162 \leq i \leq 200}$ , that seems to be more similar (in the sense of  $\mathbf{SLB}_1^{\text{ult}}$ ) to the tropical phylogenetic tree shapes. In the following  $\mathcal{G}_1$  and  $\mathcal{G}_2$  are referred to as *US main* and *US secondary group*, respectively. This division is even more evident in the MDS-plot on the right (black points represent trees shapes from the US main group, blue points trees shapes from the US secondary group and red points trees shapes based on the tropical data).

We remark that in order to highlight the subgroups the US tree shapes have been reordered according to the output permutation of a single linkage dendrogram (w.r.t.  $\mathbf{SLB}_1^{\text{ult}}$ ) based on the US tree submatrix created by MATLAB [64] and that the tropical tree shapes have been reordered analogously.

The second row of Figure 10 displays the analogous plots for  $d_{\text{CP},2}$ . It is noteworthy, that the coloring in the MDS-plot of the left is the same, i.e.,  $T_1 \in \mathcal{G}_1$  is represented by a black point,  $T_2 \in \mathcal{G}_2$  by a blue one and  $T_3 \in \mathcal{G}_3$  by a red one. Interestingly, the analysis based on these plots differs from the previous one. Using  $d_{\text{CP},2}$  to compare the phylogenetic tree shapes at hand, we can split the data into two clusters, where one corresponds to the US data and the other one to the tropical data, with only a small overlap (see the MDS-plot in the second row of Figure 10 on the right). In particular, we notice that  $d_{\text{CP},2}$  does not clearly distinguish between the US groups  $\mathcal{G}_1$  and  $\mathcal{G}_2$ .

In order to analyze the different findings of  $\mathbf{SLB}_1^{\text{ult}}$  and  $d_{\text{CP},2}$ , we collect and compare different characteristics of the tree shapes in the groups  $\mathcal{G}_i$ ,  $1 \leq i \leq 3$ . More precisely, we concentrate on various “metric” properties of the considered ultra-dissimilarity spaces like  $\frac{1}{5002|\mathcal{G}_i|} \sum_{\mathcal{T}_i \in \mathcal{G}_i} \sum_{x, x' \in X_i} u_{X_i}(x, x')$  (“mean average distance”) or  $\frac{1}{|\mathcal{G}_i|} \sum_{\mathcal{T}_i \in \mathcal{G}_i} \max\{u_{X_i}(x, x') | x, x' \in X_i\}$  (“mean maximal distance”),  $1 \leq i \leq 3$ , (these influence  $\mathbf{SLB}_1^{\text{ult}}$  strongly) as well as the




FIGURE 10. **Phylogenetic tree shape comparison:** Visualization of the dissimilarity matrices for the comparison of the phylogenetic tree shapes  $\mathcal{T}_i$ ,  $1 \leq i \leq 400$ , based on  $\mathbf{SLB}_1^{\text{ult}}$  (top row) and  $d_{\text{CP},2}$  (bottom row) as heat maps (left) and MDS-plots (right).

mean numbers of certain connectivity structures, like the 4- and 5-structures (these influence  $d_{\text{CP},2}$ , for a formal definition see [25]). These values (see Table 1) show that the mean average distance and the mean maximal distance differ drastically between the two groups of the US tree shapes. The tree shapes in these two groups are completely different from a metric perspective and the values for the secondary US group strongly resemble those of the tropic tree shapes. On the other hand, the connectivity characteristics do not change too much between the US main and secondary group. Hence, the metric  $d_{\text{CP},2}$  does not clearly divide the US trees into two groups, although the differences are certainly present. When carefully checking the phylogenetic trees, the reasons for the differences between trees in the US main group and US secondary group are not immediately apparent. Nevertheless, it is remarkable that trees from the secondary US cluster generally contain more samples from California and Florida (on average 1.92 and 0.88 more) and less from Maryland, Kentucky and Washington (on average 0.73, 0.83 and 0.72 less).

To conclude this section, we remark that using  $\mathbf{SLB}_1$  instead of  $\mathbf{SLB}_1^{\text{ult}}$  for comparing the ultra-dissimilarity spaces  $\mathcal{X}_i$ ,  $1 \leq i \leq 400$ , gives comparable results (cf. Figure 11, coloring and ordering as previously). Nevertheless, we observe (as we already have in Section 6) that  $\mathbf{SLB}_1^{\text{ult}}$  is more discriminating than  $\mathbf{SLB}_1$ . Furthermore, we mention that so far we have only considered unweighted phylogenetic tree shapes. However, the branch lengths of the considered phylogenetic trees are relevant in many examples, because they can for instance reflect the (inferred) genetic distance between evolutionary events [25]. While the branch lengths cannot easily be included in the metric  $d_{\text{CP},2}$ , the modeling of phylogenetic tree shapes as ultra-dissimilarity spaces is extremely flexible. It is straightforward to include

	USA (main group)	USA (secondary group)	Tropics
Mean Avg. Dist.	36.16	61.88	53.45
Mean Max. Dist.	56.12	86.13	94.26
Mean Num. of 4-Struc.	15.61	14.08	7.81
Mean Num. of 5-Struc.	28.04	27.97	35.82

TABLE 1. **Tree shape characteristics:** The means of several metric and connectivity characteristics of the ultra-dissimilarity spaces  $\mathcal{X}_i$  and the corresponding phylogenetic tree shapes  $\mathcal{T}_i$ ,  $1 \leq i \leq 400$ , for the three groups  $\mathcal{G}_i$ ,  $1 \leq i \leq 3$ .

branch lengths into the comparisons or to put emphasis on specific features (via weights on the corresponding tips). However, this is beyond the scope of this illustrative data analysis.




FIGURE 11. **Phylogenetic tree shape comparison based on  $\text{SLB}_1$ :** Representation of the dissimilarity matrices for the comparisons of the ultra-dissimilarity spaces  $\mathcal{X}_i$ ,  $1 \leq i \leq 400$ , based on  $\text{SLB}_1$  as heat maps (left) and MDS-plots (right).

## 8. CONCLUDING REMARKS

Since we suspect that computing  $u_{\text{GW},p}$  and  $u_{\text{GW},p}^{\text{sturm}}$  for finite  $p$  leads to NP-hard problems, it seems interesting to identify suitable collections of ultrametric measure spaces where these distances can be computed in polynomial time as done for the Gromov-Hausdorff distance in [70].

**Acknowledgements.** We are grateful to Prof. Colijn for sharing the data from [25] with us. F.M. and Z.W. acknowledge funding from the NSF under grants NSF CCF 1740761, NSF DMS 1723003, and NSF RI 1901360. A.M. and C.W. gratefully acknowledge support by the DFG Research Training Group 2088 and Cluster of Excellence MBExC 2067. F.M. and A.M. thank the Mathematisches Forschungsinstitut Oberwolfach. Conversations which eventually led to this project were initiated during the 2019 workshop “Statistical and Computational Aspects of Learning with Complex Structure”.

## REFERENCES

- [1] Georgy M. Adelson-Velskii and Aleksandr S. Kronrod. About level sets of continuous functions with partial derivatives. In *Dokl. Akad. Nauk SSSR*, volume 49, pages 239–241, 1945.

- [2] Pankaj K. Agarwal, Kyle Fox, Abhinandan Nath, Anastasios Sidiropoulos, and Yusu Wang. Computing the Gromov-Hausdorff distance for metric trees. *ACM Transactions on Algorithms (TALG)*, 14(2): 1–20, 2018.
- [3] Alfred V. Aho and John E. Hopcroft. *The design and analysis of computer algorithms*. Pearson Education India, 1974.
- [4] David Alvarez-Melis and Tommi S. Jaakkola. Gromov-Wasserstein alignment of word embedding spaces. *arXiv preprint arXiv:1809.00013*, 2018.
- [5] Yair Bartal. Probabilistic approximation of metric spaces and its algorithmic applications. In *Proceedings of 37th Conference on Foundations of Computer Science*, pages 184–193. IEEE, 1996.
- [6] Louis J. Billera, Susan P. Holmes, and Karen Vogtmann. Geometry of the space of phylogenetic trees. *Advances in Applied Mathematics*, 27(4):733–767, 2001.
- [7] Patrick Billingsley. *Convergence of Probability Measures*. John Wiley & Sons, 2013.
- [8] Michael G.B. Blum and Olivier François. Which random processes describe the tree of life? A large-scale study of phylogenetic tree imbalance. *Systematic Biology*, 55(4):685–691, 2006.
- [9] Nicolas Bonneel, Julien Rabin, Gabriel Peyré, and Hanspeter Pfister. Sliced and radon Wasserstein barycenters of measures. *Journal of Mathematical Imaging and Vision*, 51(1):22–45, 2015.
- [10] Leon Bottou, Martin Arjovsky, David Lopez-Paz, and Maxime Oquab. Geometrical insights for implicit generative modeling. In *Braverman Readings in Machine Learning. Key Ideas from Inception to Current State*, pages 229–268. Springer, 2018.
- [11] Daniel Brinkman and Peter J. Olver. Invariant histograms. *The American Mathematical Monthly*, 119(1):4–24, 2012.
- [12] Alexander M. Bronstein, Michael M. Bronstein, and Ron Kimmel. Efficient computation of isometry-invariant distances between surfaces. *SIAM Journal on Scientific Computing*, 28(5):1812–1836, 2006.
- [13] Alexander M. Bronstein, Michael M. Bronstein, and Ron Kimmel. Generalized multidimensional scaling: A framework for isometry-invariant partial surface matching. *Proceedings of the National Academy of Sciences*, 103(5):1168–1172, 2006.
- [14] Alexander M. Bronstein, Michael M. Bronstein, Alfred M. Bruckstein, and Ron Kimmel. Partial similarity of objects, or how to compare a centaur to a horse. *International Journal of Computer Vision*, 84(2):163, 2009.
- [15] Alexander M. Bronstein, Michael M. Bronstein, and Ron Kimmel. Topology-invariant similarity of nonrigid shapes. *International journal of computer vision*, 81(3):281, 2009.
- [16] Alexander M. Bronstein, Michael M. Bronstein, Ron Kimmel, Mona Mahmoudi, and Guillermo Sapiro. A Gromov-Hausdorff framework with diffusion geometry for topologically-robust non-rigid shape matching. *International Journal of Computer Vision*, 89(2-3):266–286, 2010.
- [17] Peter Brown, Wayne Pullan, Yuedong Yang, and Yaoqi Zhou. Fast and accurate non-sequential protein structure alignment using a new asymmetric linear sum assignment heuristic. *Bioinformatics*, 32(3): 370–377, 2016.
- [18] Charlotte Bunne, David Alvarez-Melis, Andreas Krause, and Stefanie Jegelka. Learning generative models across incomparable spaces. *arXiv preprint arXiv:1905.05461*, 2019.
- [19] Gunnar Carlsson and Facundo Mémoli. Characterization, stability and convergence of hierarchical clustering methods. *Journal of machine learning research*, 11(Apr):1425–1470, 2010.
- [20] Frédéric Chazal, David Cohen-Steiner, Leonidas J Guibas, Facundo Mémoli, and Steve Y Oudot. Gromov-hausdorff stable signatures for shapes using persistence. In *Proceedings of the Symposium on Geometry Processing*, pages 1393–1403, 2009.
- [21] Jie Chen and Ilya Safro. Algebraic distance on graphs. *SIAM Journal on Scientific Computing*, 33(6): 3468–3490, 2011.
- [22] Samir Chowdhury. *Metric and Topological Approaches to Network Data Analysis*. PhD thesis, The Ohio State University, 2019.
- [23] Samir Chowdhury and Facundo Mémoli. The Gromov-Wasserstein distance between networks and stable network invariants. *Information and Inference: A Journal of the IMA*, 8(4):757–787, 2019.
- [24] Samir Chowdhury and Tom Needham. Generalized spectral clustering via Gromov-Wasserstein learning. *arXiv preprint arXiv:2006.04163*, 2020.
- [25] Caroline Colijn and Giacomo Plazzotta. A metric on phylogenetic tree shapes. *Systematic biology*, 67(1):113–126, 2018.

- [26] Guy David, Stephen W. Semmes, Stephen Semmes, and Guy Rene Pierre Pierre. *Fractured fractals and broken dreams: Self-similar geometry through metric and measure*, volume 7. Oxford University Press, 1997.
- [27] Adel Dayarian and Boris I. Shraiman. How to infer relative fitness from a sample of genomic sequences. *Genetics*, 197(3):913–923, 2014.
- [28] Khanh Do Ba, Huy L. Nguyen, Huy N. Nguyen, and Ronitt Rubinfeld. Sublinear time algorithms for Earth Mover’s distance. *Theory of Computing Systems*, 48(2):428–442, 2011.
- [29] Yihe Dong and Will Sawin. COPT: Coordinated optimal transport on graphs. *arXiv preprint arXiv:2003.03892*, 2020.
- [30] Dmitry Dordovskiy, Oleksiy Dovgoshey, and Eugeny Petrov. Diameter and diametrical pairs of points in ultrametric spaces. *P-Adic Numbers, Ultrametric Analysis, and Applications*, 3(4):253–262, 2011.
- [31] Richard M. Dudley. *Real analysis and probability*. CRC Press, 2018.
- [32] David A. Edwards. The structure of superspace. In *Studies in topology*, pages 121–133. Elsevier, 1975.
- [33] Steven N. Evans. *Probability and Real Trees: École D’Été de Probabilités de Saint-Flour XXXV-2005*. Springer, 2007.
- [34] Steven N. Evans and Frederick A. Matsen. The phylogenetic Kantorovich–Rubinstein metric for environmental sequence samples. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 74(3):569–592, 2012.
- [35] Jittat Fakcharoenphol, Satish Rao, and Kunal Talwar. A tight bound on approximating arbitrary metrics by tree metrics. *Journal of Computer and System Sciences*, 69(3):485–497, 2004.
- [36] Gerald B Folland. *Real analysis: modern techniques and their applications*, volume 40. John Wiley & Sons, 1999.
- [37] Manuela Gellert, Md Faruq Hossain, Felix Jacob Ferdinand Berens, Lukas Willy Bruhn, Claudia Urbainsky, Volkmar Liebscher, and Christopher Horst Lillig. Substrate specificity of thioredoxins and glutaredoxins—towards a functional classification. *Heliyon*, 5(12):e02943, 2019.
- [38] Clark R. Givens and Rae Michael Shortt. A class of Wasserstein metrics for probability distributions. *Michigan Math. J.*, 31(2):231–240, 1984. doi: 10.1307/mmj/1029003026. URL <https://doi.org/10.1307/mmj/1029003026>.
- [39] Andreas Greven, Peter Pfaffelhuber, and Anita Winter. Convergence in distribution of random metric measure spaces ( $\lambda$ -coalescent measure trees). *Probability Theory and Related Fields*, 145(1-2):285–322, 2009.
- [40] Gillian Grindstaff and Megan Owen. Geometric comparison of phylogenetic trees with different leaf sets. *arXiv preprint arXiv:1807.04235*, 2018.
- [41] M. Gromov. Groups of polynomial growth and expanding maps (with an appendix by Jacques Tits). *Publications Mathématiques de l’IHÉS*, 53:53–78, 1981.
- [42] Jotun Hein. Reconstructing evolution of sequences subject to recombination using parsimony. *Mathematical biosciences*, 98(2):185–200, 1990.
- [43] Liisa Holm and Chris Sander. Protein structure comparison by alignment of distance matrices. *Journal of molecular biology*, 233(1):123–138, 1993.
- [44] Norman R. Howes. *Modern analysis and topology*. Springer Science & Business Media, 2012.
- [45] Anil K. Jain and Chitra Dorai. 3d object recognition: Representation and matching. *Statistics and Computing*, 10(2):167–182, 2000.
- [46] Nicholas Jardine and Robin Sibson. *Mathematical taxonomy*. John Wiley & Sons, 1971.
- [47] Leonid V. Kantorovich. On the translocation of masses, cr (dokl.) acad. *Sci. URSS (NS)*, 37:199, 1942.
- [48] Leonid V. Kantorovich and G Rubinstein. On a space of completely additive functions (russ.). *Vestnik Leningrad Univ*, 13:52–59, 1958.
- [49] Jaehee Kim, Noah A. Rosenberg, and Julia A. Palacios. A metric space of ranked tree shapes and ranked genealogies. *bioRxiv*, 2019.
- [50] Benoît R. Kloeckner. A geometric study of Wasserstein spaces: Ultrametrics. *Mathematika*, 61(1):162–178, 2015.
- [51] Katia Koelle, Priya Khatri, Meredith Kamradt, and Thomas B. Kepler. A two-tiered model for simulating the ecological and evolutionary dynamics of rapidly evolving viruses, with an application to influenza. *Journal of The Royal Society Interface*, 7(50):1257–1274, 2010.

- [52] Andreĭ N. Kolmogorov and Sergei V. Fomin. *Elements of the theory of functions and functional analysis*, volume 1. Courier Corporation, 1957.
- [53] Soheil Kolouri, Kimia Nadjahi, Umut Simsekli, Roland Badeau, and Gustavo Rohde. Generalized sliced Wasserstein distances. In *Advances in Neural Information Processing Systems*, pages 261–272, 2019.
- [54] Irina Kufareva and Ruben Abagyan. Methods of protein structure comparison. In *Homology Modeling*, pages 231–257. Springer, 2011.
- [55] Hao-Yuan Kuo, Hong-Ren Su, Shang-Hong Lai, and Chin-Chia Wu. 3D object detection and pose estimation from depth image for robotic bin picking. In *2014 IEEE international conference on automation science and engineering (CASE)*, pages 1264–1269. IEEE, 2014.
- [56] Manuel Lafond, Nadia El-Mabrouk, Katharina T. Huber, and Vincent Moulton. The complexity of comparing multiply-labelled trees by extending phylogenetic-tree metrics. *Theoretical Computer Science*, 760:15–34, 2019.
- [57] Tam Le, Nhat Ho, and Makoto Yamada. Fast tree variants of Gromov-Wasserstein. *arXiv preprint arXiv:1910.04462*, 2019.
- [58] Tam Le, Makoto Yamada, Kenji Fukumizu, and Marco Cuturi. Tree-sliced variants of Wasserstein distances. In *Advances in neural information processing systems*, pages 12304–12315, 2019.
- [59] Volkmar Liebscher. New Gromov-inspired metrics on phylogenetic tree space. *Bulletin of mathematical biology*, 80(3):493–518, 2018.
- [60] Pengyu Liu, Matthew Gould, and Caroline Colijn. Polynomial phylogenetic analysis of tree shapes. *BioRxiv*, 2020.
- [61] David G. Lowe. Local feature view clustering for 3D object recognition. In *Proceedings of the 2001 IEEE Computer Society Conference on Computer Vision and Pattern Recognition. CVPR 2001*, volume 1, pages I–I. IEEE, 2001.
- [62] Marta Luksza and Michael Lässig. A predictive fitness model for influenza. *Nature*, 507(7490):57–61, 2014.
- [63] Colin L. Mallows. A note on asymptotic joint normality. *The Annals of Mathematical Statistics*, pages 508–515, 1972.
- [64] MATLAB. *MATLAB: Accelerating the pace of engineering and science*. The MathWorks, Inc., 2019. URL <https://www.mathworks.com>.
- [65] Andrew McGregor and Daniel Stubbs. Sketching Earth-Mover distance on graph metrics. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 274–286. Springer, 2013.
- [66] Facundo Mémoli. On the use of Gromov-Hausdorff distances for shape comparison. In M. Botsch, R. Pajarola, B. Chen, and M. Zwicker, editors, *Eurographics Symposium on Point-Based Graphics*. The Eurographics Association, 2007. ISBN 978-3-905673-51-7. doi: 10.2312/SPBG/SPBG07/081-090.
- [67] Facundo Mémoli. Gromov-Wasserstein distances and the metric approach to object matching. *Foundations of computational mathematics*, 11(4):417–487, 2011.
- [68] Facundo Mémoli and Tom Needham. Distance distributions and inverse problems for metric measure spaces. *arXiv preprint arXiv:1810.09646*, 2021.
- [69] Facundo Mémoli and Guillermo Sapiro. Comparing point clouds. In *Proceedings of the 2004 Eurographics/ACM SIGGRAPH symposium on Geometry processing*, pages 32–40, 2004.
- [70] Facundo Mémoli, Zane Smith, and Zhengchao Wan. Gromov-Hausdorff distances on  $p$ -metric spaces and ultrametric spaces. *arXiv preprint arXiv:1912.00564*, 2019.
- [71] Marc Mézard, Giorgio Parisi, and Miguel Angel Virasoro. *Spin glass theory and beyond: An Introduction to the Replica Method and Its Applications*, volume 9. World Scientific Publishing Company, 1987.
- [72] Arne O. Mooers and Stephen B. Heard. Inferring evolutionary process from phylogenetic tree shape. *The quarterly review of Biology*, 72(1):31–54, 1997.
- [73] Alexey Anatolievich Morozov. Extension of Colijn-Plazotta tree shape distance metric to unrooted trees. *BioRxiv*, page 506022, 2018.
- [74] Dmitriy Morozov, Kenes Beketayev, and Gunther Weber. Interleaving distance between merge trees. *Discrete and Computational Geometry*, 49(22-45):52, 2013.

- [75] Robert Osada, Thomas Funkhouser, Bernard Chazelle, and David Dobkin. Shape distributions. *ACM Transactions on Graphics (TOG)*, 21(4):807–832, 2002.
- [76] Megan Owen and J. Scott Provan. A fast algorithm for computing geodesic distances in tree space. *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 8(1):2–13, 2010.
- [77] Chavdar Papazov, Sami Haddadin, Sven Parusel, Kai Krieger, and Darius Burschka. Rigid 3D geometry matching for grasping of known objects in cluttered scenes. *The International Journal of Robotics Research*, 31(4):538–553, 2012.
- [78] Panos M. Pardalos and Stephen A. Vavasis. Quadratic programming with one negative eigenvalue is NP-hard. *Journal of Global optimization*, 1(1):15–22, 1991.
- [79] Gabriel Peyré, Marco Cuturi, and Justin Solomon. Gromov-Wasserstein averaging of kernel and distance matrices. In *International Conference on Machine Learning*, pages 2664–2672, 2016.
- [80] Derong Qiu. Geometry of non-archimedean Gromov-Hausdorff distance. *P-Adic Numbers, Ultrametric Analysis, and Applications*, 1(4):317, 2009.
- [81] Rammal Rammal, Gérard Toulouse, and Miguel A. Virasoro. Ultrametricity for physicists. *Reviews of Modern Physics*, 58:765–788, 1986.
- [82] Georges Reeb. Sur les points singuliers d’une forme de pfaff complètement intégrable ou d’une fonction numérique [on the singular points of a completely integrable pfaff form or of a numerical function]. *Comptes Rendus Acad. Sciences Paris*, 222:847–849, 1946.
- [83] David F. Robinson. Comparison of labeled trees with valency three. *Journal of combinatorial theory, Series B*, 11(2):105–119, 1971.
- [84] David F. Robinson and Leslie R. Foulds. Comparison of phylogenetic trees. *Mathematical biosciences*, 53(1-2):131–147, 1981.
- [85] Yossi Rubner, Carlo Tomasi, and Leonidas J. Guibas. The Earth Mover’s distance as a metric for image retrieval. *International journal of computer vision*, 40(2):99–121, 2000.
- [86] Colin A. Russell, Terry C. Jones, Ian G. Barr, Nancy J. Cox, Rebecca J. Garten, Vicky Gregory, Ian D. Gust, Alan W. Hampson, Alan J. Hay, Aeron C. Hurt, et al. The global circulation of seasonal influenza a (H3N2) viruses. *Science*, 320(5874):340–346, 2008.
- [87] Meyer Scetbon, Gabriel Peyré, and Marco Cuturi. Linear-time gromov wasserstein distances using low rank couplings and costs. *arXiv preprint arXiv:2106.01128*, 2021.
- [88] Felix Schmiedl. Computational aspects of the Gromov-Hausdorff distance and its application in non-rigid shape matching. *Discret. Comput. Geom.*, 57(4):854–880, 2017. doi: 10.1007/s00454-017-9889-4. URL <https://doi.org/10.1007/s00454-017-9889-4>.
- [89] Stephen Semmes. An introduction to the geometry of ultrametric spaces. *arXiv preprint arXiv:0711.0709*, 2007.
- [90] Charles Semple, Mike Steel, et al. *Phylogenetics*, volume 24. Oxford University Press on Demand, 2003.
- [91] Zane Smith, Samir Chowdhury, and Facundo Mémoli. Hierarchical representations of network data with optimal distortion bounds. In *2016 50th Asilomar Conference on Signals, Systems and Computers*, pages 1834–1838. IEEE, 2016.
- [92] Karl-Theodor Sturm. On the geometry of metric measure spaces. *Acta mathematica*, 196(1):65–131, 2006.
- [93] Karl-Theodor Sturm. The space of spaces: Curvature bounds and gradient flows on the space of metric measure spaces. *arXiv preprint arXiv:1208.0434*, 2012.
- [94] David Thorsley and Eric Klavins. Model reduction of stochastic processes using Wasserstein pseudometrics. In *2008 American Control Conference*, pages 1374–1381. IEEE, 2008.
- [95] Vayer Titouan, Nicolas Courty, Romain Tavenard, and Rémi Flamary. Optimal transport for structured data with application on graphs. In *International Conference on Machine Learning*, pages 6275–6284, 2019.
- [96] Elena Farahbakhsh Touli and Yusu Wang. FPT-algorithms for computing Gromov-Hausdorff and interleaving distances between trees. *arXiv preprint arXiv:1811.02425*, 2018.
- [97] Sergei S. Vallender. Calculation of the Wasserstein distance between probability distributions on the line. *Theory of Probability & Its Applications*, 18(4):784–786, 1974.
- [98] Titouan Vayer, Rémi Flamary, Romain Tavenard, Laetitia Chapel, and Nicolas Courty. Sliced Gromov-Wasserstein. *arXiv preprint arXiv:1905.10124*, 2019.

- [99] Cédric Villani. *Topics in optimal transportation*. American Mathematical Soc., 2003.
- [100] Cédric Villani. *Optimal transport: Old and new*, volume 338. Springer Science & Business Media, 2008.
- [101] Erik M. Volz, Katia Koelle, and Trevor Bedford. Viral phylodynamics. *PLoS Comput Biol*, 9(3): e1002947, 2013.
- [102] Zhengchao Wan. A novel construction of Urysohn universal ultrametric space via the Gromov-Hausdorff ultrametric. *arXiv preprint arXiv:2007.08105*, 2020.
- [103] Kim B. Westgeest, Miranda de Graaf, Mathieu Fourment, Theo M. Bestebroer, Ruud van Beek, Monique I.J. Spronken, Jan C. de Jong, Guus F Rimmelzwaan, Colin A. Russell, Albert D.M.E. Osterhaus, et al. Genetic evolution of the neuraminidase of influenza a (H3N2) viruses from 1968 to 2009 and its correspondence to haemagglutinin evolution. *The Journal of general virology*, 93(Pt 9): 1996, 2012.
- [104] Taoyang Wu and Kwok Pui Choi. On joint subtree distributions under two evolutionary models. *Theoretical population biology*, 108:13–23, 2016.
- [105] Ihor Zarichnyi. Gromov-Hausdorff ultrametric. *arXiv preprint math/0511437*, 2005.

## APPENDIX A. MISSING DETAILS FROM SECTION 2

**A.1. Proofs from Section 2.** In this section we give the proofs of various results from Section 2.

**A.1.1. Proof of Theorem 2.2.** Recall that for a given  $\theta \in \mathcal{D}(X)$ , we define  $u_\theta : X \times X \rightarrow \mathbb{R}_{\geq 0}$  as follows

$$u_\theta(x, x') := \inf\{t \geq 0 \mid x \text{ and } x' \text{ belong to the same block of } \theta(t)\}.$$

It is straightforward to verify that  $u_\theta$  is an ultrametric. For any Cauchy sequence  $\{x_n\}_{n \in \mathbb{N}}$  in  $(X, u_\theta)$ , let  $D_i := \sup_{m, n \geq i} u_\theta(x_m, x_n)$  for each  $i \in \mathbb{N}$ . Then, each  $D_i < \infty$  and  $\lim_{i \rightarrow \infty} D_i = 0$ . By definition of  $u_\theta$ , we have that for each  $i \in \mathbb{N}$  the set  $\{x_n\}_{n=i}^\infty$  is contained in the block  $[x_i]_{D_i} \in \theta(D_i)$ . Let  $X_i := [x_i]_{D_i}$  for each  $i \in \mathbb{N}$ . Then, obviously we have that  $X_j \subseteq X_i$  for any  $1 \leq i < j$ . By condition (7) in Definition 2.1, we have that  $\bigcap_{i \in \mathbb{N}} X_i \neq \emptyset$ . Choose  $x_* \in \bigcap_{i \in \mathbb{N}} X_i$ , then it is easy to verify that  $x_* = \lim_{n \rightarrow \infty} x_n$  and thus  $(X, u_\theta)$  is a complete space. To prove that  $(X, u_\theta)$  is a compact space, we need to verify that for each  $t > 0$ ,  $X_t$  is a finite space (cf. Lemma A.7). Since  $\theta(t)$  is finite by condition (6) in Definition 2.1, we have that  $X_t = \{[x]_t \mid x \in X\} = \theta(t)$  is finite and thus  $X$  is compact. Therefore, we have proved that  $u_\theta \in \mathcal{U}(X)$ . Based on this, the map  $\Delta_X : \mathcal{D}(X) \rightarrow \mathcal{U}(X)$  by  $\theta \mapsto u_\theta$  is well-defined.

Now given  $u \in \mathcal{U}(X)$ , we define a map  $\theta_u : [0, \infty) \rightarrow \mathbf{Part}(X)$  as follows: for each  $t \geq 0$ , consider the equivalence relation  $\sim_t$  with respect to  $u$ , i.e.,  $x \sim_t x'$  if and only if  $u(x, x') \leq t$ . This is actually the same equivalence relation defined in Section 2.2 for introducing quotient ultrametric spaces. We then let  $\theta_u(t)$  to be the partition induced by  $\sim_t$ , i.e.,  $\theta_u(t) = X_t$ . It is not hard to show that  $\theta_u$  satisfies conditions (1)–(5) in Definition 2.1. Since  $X$  is compact, then  $\theta_u(t) = X_t$  is finite for each  $t > 0$  and thus  $\theta_u$  satisfies condition (6) in Definition 2.1. Now, let  $\{t_n\}_{n \in \mathbb{N}}$  be a decreasing sequence such that  $\lim_{n \rightarrow \infty} t_n = 0$  and let  $X_n \in \theta_X(t_n)$  such that for any  $1 \leq n < m$ ,  $X_m \subseteq X_n$ . Since each  $X_n = [x_n]_{t_n}$  for some  $x_n \in X$ ,  $X_n$  is a compact subset of  $X$ . Since  $X$  is also complete, we have that  $\bigcap_{n \in \mathbb{N}} X_n \neq \emptyset$ . Therefore,  $\theta_u$  satisfies condition (7) in Definition 2.1 and thus  $\theta_u \in \mathcal{D}(X)$ . Then, we define the map  $\Upsilon_X : \mathcal{U}(X) \rightarrow \mathcal{D}(X)$  by  $u \mapsto \theta_u$ .

It is easy to check that  $\Upsilon_X$  is the inverse of  $\Delta_X$  and thus we have established that  $\Delta_X : \mathcal{D}(X) \rightarrow \mathcal{U}(X)$  is bijective.

A.1.2. *Proof of Lemma 2.8.* First of all, we show that the right hand side of Equation (16) is well defined. More precisely, we employ Lemma A.7 to prove that the supremum

$$\sup_{B \in V(X) \setminus \{X\} \text{ and } \alpha(B) \neq \beta(B)} \text{diam}(B^*)$$

is attained. For arbitrary  $B_0 \in V(X) \setminus \{X\}$  such that  $\alpha(B_0) \neq \beta(B_0)$ , we have that  $\text{diam}(B_0^*) > 0$ . By Lemma A.7 the spaces  $X_t$  are finite for  $t > 0$ . Since  $V(X) = \{[x]_t \mid x \in X, t > 0\} = \bigcup_{t>0} X_t$ , there are only finitely many  $B \in V(X) \setminus \{X\}$  such that  $\text{diam}(B) \geq \text{diam}(B_0^*)$  and thus  $\text{diam}(B^*) \geq \text{diam}(B_0^*)$ . This implies that the supremum is attained and thus

$$\sup_{B \in V(X) \setminus \{X\} \text{ and } \alpha(B) \neq \beta(B)} \text{diam}(B^*) = \max_{B \in V(X) \setminus \{X\} \text{ and } \alpha(B) \neq \beta(B)} \text{diam}(B^*). \quad (27)$$

Let  $B_1$  denote the maximizer in Equation (27) and let  $\delta := \text{diam}(B_1^*)$ . It is easy to see that for any  $x \in X$ ,  $\alpha([x]_\delta) = \beta([x]_\delta)$ .

By Strassen's theorem (see for example [31, Theorem 11.6.2]),

$$d_{W,\infty}(\alpha, \beta) = \inf\{r \geq 0 \mid \text{for any closed subset } A \subseteq X, \alpha(A) \leq \beta(A^r)\}, \quad (28)$$

where  $A^r := \{x \in X \mid u_X(x, A) \leq r\}$ .

Since  $\alpha(B_1) \neq \beta(B_1)$ , we assume without loss of generality that  $\alpha(B_1) > \beta(B_1)$ . By definition of  $B_1^*$ , it is obvious that  $(B_1)^\delta = B_1^*$  (recall:  $\delta := \text{diam}(B_1^*)$ ) and  $(B_1)^r = B_1$  for all  $0 \leq r < \delta$ . Therefore,  $\alpha(B_1) \leq \beta((B_1)^r)$  only when  $r \geq \delta$ . By Equation (28), this implies that  $d_{W,\infty}(\alpha, \beta) \geq \delta$ . Conversely, for any closed set  $A$ , we have that  $A^\delta = \bigcup_{x \in A} [x]_\delta$ . For two closed balls in ultrametric spaces, either one includes the other or they have no intersection. Therefore, there exists a subset  $S \subseteq A$  such that  $[x]_\delta \cap [x']_\delta = \emptyset$  for all  $x, x' \in S$  and  $x \neq x'$ , and that  $A^\delta = \bigsqcup_{x \in S} [x]_\delta$ . Then,  $\alpha(A) \leq \alpha(A^\delta) = \sum_{x \in S} \alpha([x]_\delta) = \sum_{x \in S} \beta([x]_\delta) = \beta(A^\delta)$ . Hence,  $d_{W,\infty}(\alpha, \beta) \leq \delta$  and thus

$$d_{W,\infty}(\alpha, \beta) = \max_{B \in V(X) \setminus \{X\} \text{ and } \alpha(B) \neq \beta(B)} \text{diam}(B^*).$$

**A.2. Technical issues from Section 2.** In the following, we address various technical issues from Section 2.

A.2.1. *Synchronized rooted trees.* A *synchronized rooted tree*, is a combinatorial tree  $T = (V, E)$  with a root  $o \in V$  and a height function  $h : V \rightarrow [0, \infty)$  such that  $h^{-1}(0)$  coincides with the leaf set and  $h(v) < h(v^*)$  for each  $v \in V \setminus \{o\}$ , where  $v^*$  is the parent of  $v$ . Similar as in Theorem 2.2 that there exists a correspondence between ultrametric spaces and dendrograms, an ultrametric space  $X$  uniquely determines a synchronized rooted tree  $T_X$  [50].

Now given a compact ultrametric space  $(X, u_X)$ , we construct the corresponding synchronized rooted tree  $T_X$  via the dendrogram  $\theta_X$  associated with  $u_X$ . Recall from Section 2.3 that  $V(X) := \bigcup_{t>0} \theta_X(t)$ . For each  $B \in V(X) \setminus \{X\}$ , denote by  $B^*$  the smallest element in  $V(X)$  such that  $B \subsetneq B^*$ , whose existence is guaranteed by the following lemma:

**Lemma A.1.** *Let  $X$  be a compact ultrametric space and let  $V(X) = \bigcup_{t>0} \theta_X(t)$ , where  $\theta_X$  is as defined in Remark 2.3. For each  $B \in V(X)$  such that  $B \neq X$ , there exists  $B^* \in V(X)$  such that  $B^* \neq B$  and  $B^* \subseteq B'$  for all  $B' \in V(X)$  with  $B \subsetneq B'$ .*

*Proof.* Let  $\delta := \text{diam}(B)$ . Let  $x \in B$ , then  $B = [x]_\delta$ . By Lemma A.7,  $X_\delta$  is a finite set. Consider  $\delta^* := \min\{u_{X_\delta}([x]_\delta, [x']_\delta) \mid [x']_\delta \neq [x]_\delta\}$ . Let  $B^* := [x]_{\delta^*}$ , then  $B^*$  is the smallest element in  $V(X)$  containing  $B$  under inclusion. Indeed,  $B^* \neq B$  and if  $B \subseteq B'$  for some  $B' \in V(X)$ , then  $B' = [x]_r$  for some  $r > \delta$ . It is easy to see that for all  $\delta < r < \delta^*$ ,  $[x]_r = [x]_\delta$ . Therefore, if  $B' \neq B$ , we must have that  $r \geq \delta^*$  and thus  $B^* = [x]_{\delta^*} \subseteq [x]_r = B'$ .  $\square$

Now, we define a combinatorial tree  $T_X = (V_X, E_X)$  as follows: we let  $V_X := V(X)$ ; for any distinct  $B, B' \in V_X$ , we let  $(B, B') \in E_X$  iff either  $B = (B')^*$  or  $B' = B^*$ . We choose  $X \in V_X$  to be the root of  $T_X$ , then any  $B \neq X$  in  $V_X$  has a unique parent  $B^*$ . We define  $h_X : V_X \rightarrow [0, \infty)$  such that  $h_X(B) := \frac{\text{diam}(B)}{2}$  for any  $B \in V_X$ . Now,  $T_X$  endowed with the root  $X$  and the height function  $h_X$  is a synchronized rooted tree. It is easy to see that  $X$  can be isometrically identified with  $h_X^{-1}(0)$  of the so-called *metric completion* of  $T_X$  (see [50, Section 2.3] for details). With this construction Theorem 2.7 follows directly from [50, Lemma 3.1].

**A.3.  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  between compactly supported measures.** Next, we demonstrate that Theorem 2.9 extends naturally to the case of compactly supported probability measures in  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ . For this purpose, it is important to note that compact subsets of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$  have a very particular structure as shown by the subsequent lemma.

**Lemma A.2.** *Let  $X \subseteq (\mathbb{R}_{\geq 0}, \Lambda_\infty)$ .  $X$  is a compact subset if and only if  $X$  is either a finite set or a countable set with 0 being the unique cluster point (w.r.t. the usual Euclidean distance  $\Lambda_1$ ).*

*Proof.* If  $X$  is finite, then obviously  $X$  is compact. Assume that  $X$  is a countable set with 0 being the unique cluster point (w.r.t. the usual Euclidean distance  $\Lambda_1$ ). If  $\{x_n\}_{n \in \mathbb{N}} \subseteq X$  is a Cauchy sequence with respect to  $\Lambda_\infty$ , then either  $x_n$  is a constant when  $n$  is large or  $\lim_{n \rightarrow \infty} x_n = 0$ . In either case, the limit of  $\{x_n\}_{n \in \mathbb{N}}$  belongs to  $X$  and thus  $X$  is complete. Now for any  $\varepsilon > 0$ , by Lemma A.7,  $X_\varepsilon$  is a finite set. Denote  $X_\varepsilon = \{[x_1]_\varepsilon, \dots, [x_n]_\varepsilon\}$ . Then,  $\{x_1, \dots, x_n\}$  is a finite  $\varepsilon$ -net of  $X$ . Therefore,  $X$  is totally bounded and thus  $X$  is compact.

Now, assume that  $X$  is compact. Then, for any  $\varepsilon > 0$ ,  $X_\varepsilon$  is a finite set. Suppose  $X_\varepsilon = \{[x_1]_\varepsilon, \dots, [x_n]_\varepsilon\}$  where  $0 \leq x_1 < x_2 < \dots < x_n$ . Further, we have that  $\Lambda_\infty(x_i, x_j) = x_j$  whenever  $1 \leq i < j \leq n$ . This implies that

- (1)  $x_i > \varepsilon$  for all  $2 \leq i \leq n$ ;
- (2)  $[x_i]_\varepsilon = \{x_i\}$  for all  $2 \leq i \leq n$ .

Therefore,  $X \cap (\varepsilon, \infty) = \{x_2, \dots, x_n\}$  is a finite set. Since  $\varepsilon > 0$  is arbitrary,  $X$  is at most countable and has no cluster point (w.r.t. the usual Euclidean distance  $\Lambda_1$ ) other than 0. If  $X$  is countable, then 0 must be a cluster point and by compactness of  $X$ , we have that  $0 \in X$ .  $\square$

Based on the special structure of compact subsets of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ , we derive the following extension of Theorem 2.9.

**Theorem A.3** ( $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  between compactly supported measures). *Suppose  $\alpha, \beta$  are supported on a countable subset  $X := \{0\} \cup \{x_i \mid i \in \mathbb{N}\}$  of  $\mathbb{R}_{\geq 0}$  such that  $0 < \dots < x_n < x_{n-1} <$*

$\dots < x_1$  and 0 is the only cluster point with respect to the usual Euclidean distance. Let  $\alpha_i := \alpha(\{x_i\})$  for  $i \in \mathbb{N}$  and  $\alpha_0 := \alpha(\{0\})$ . Similarly, let  $\beta_i := \beta(\{x_i\})$  and  $\beta_0 := \beta(\{0\})$ . Then for  $p \in [1, \infty)$ ,

$$d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(\alpha, \beta) = 2^{-\frac{1}{p}} \left( \sum_{i=2}^{\infty} \left| \sum_{j=i}^{\infty} (\alpha_j - \beta_j) \right| \cdot |x_{i-1}^p - x_i^p| + \sum_{i=1}^{\infty} |\alpha_i - \beta_i| \cdot x_i^p \right)^{\frac{1}{p}}. \quad (29)$$

Let  $F_\alpha$  and  $F_\beta$  denote the cumulative distribution functions of  $\alpha$  and  $\beta$ , respectively. Then, we obtain

$$d_{\mathbb{W},\infty}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}(\alpha, \beta) = \max \left( \max_{2 \leq i < \infty, F_\alpha(x_i) \neq F_\beta(x_i)} x_{i-1}, \max_{1 \leq i < \infty, \alpha_i \neq \beta_i} x_i \right).$$

*Proof.* Note that  $V(X) = \{\{0\} \cup \{x_j | j \geq i\} | i \in \mathbb{N}\} \cup \{\{x_i\} | i \in \mathbb{N}\}$  (recall that each set corresponds to a closed ball). Thus, we conclude the proof by applying Lemma 2.7 and Lemma 2.8.  $\square$

A.3.1. *Closed-form solution for  $d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}$ .* In the following, we will derive the subsequent theorem.

**Theorem A.4.** *Given  $1 \leq p, q < \infty$  and two compactly supported probability measures  $\alpha$  and  $\beta$  on  $\mathbb{R}_{\geq 0}$ , we have that*

$$d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta) \leq \left( \int_0^1 \Lambda_q(F_\alpha^{-1}(t), F_\beta^{-1}(t))^p dt \right)^{\frac{1}{p}}.$$

When  $q \leq p$ , the equality holds whereas when  $q > p$ , the equality does not hold in general.

One important ingredient for the proof of Theorem A.4 is Lemma 3.2 of Chowdhury and Mémoli [23] which we restate here for convenience.

**Lemma A.5** (Chowdhury and Mémoli [23, Lemma 3.2]). *Let  $X, Y$  be two Polish metric spaces and let  $f : X \rightarrow \mathbb{R}$  and  $g : Y \rightarrow \mathbb{R}$  be measurable maps. Denote by  $f \times g : X \times Y \rightarrow \mathbb{R}^2$  the map  $(x, y) \mapsto (f(x), g(y))$ . Then, for any  $\mu_X \in \mathcal{P}(X)$  and  $\mu_Y \in \mathcal{P}(Y)$*

$$(f \times g)_\# \mathcal{C}(\mu_X, \mu_Y) = \mathcal{C}(f_\# \mu_X, g_\# \mu_Y).$$

Based on Lemma A.5, we can show the following auxiliary result.

**Lemma A.6.** *Let  $1 \leq q \leq p < \infty$ . Assume that  $\alpha$  and  $\beta$  are compactly supported probability measures on  $\mathbb{R}_{\geq 0}$ . Then,*

$$\left( d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta) \right)^p = \left( d_{\mathbb{W},\frac{p}{q}}^{(\mathbb{R}_{\geq 0}, \Lambda_1)}((S_q)_\# \alpha, (S_q)_\# \beta) \right)^{\frac{p}{q}},$$

where  $S_q : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{\geq 0}$  taking  $x$  to  $x^q$  is the  $q$ -snowflake transform defined in Section 3.3.

*Proof.*

$$\begin{aligned}
\left(d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta)\right)^p &= \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} (\Lambda_q(x, y))^p \mu(dx \times dy) \\
&= \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} |S_q(x) - S_q(y)|^{\frac{p}{q}} \mu(dx \times dy) \\
&= \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} |s - t|^{\frac{p}{q}} (S_q \times S_q)_{\#} \mu(ds \times dt) \\
&= \left(d_{\mathbb{W}, \frac{p}{q}}^{(\mathbb{R}_{\geq 0}, \Lambda_1)}((S_q)_{\#} \alpha, (S_q)_{\#} \beta)\right)^{\frac{p}{q}},
\end{aligned}$$

where we use  $\frac{p}{q} \geq 1$  and Lemma A.5 in the last equality.  $\square$

With Lemma A.6 at our disposal, we can demonstrate Theorem A.4.

*Proof of Theorem A.4.* We first note that  $d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta) = \inf_{(\xi, \eta)} (\mathbb{E}(\Lambda_q(\xi, \eta)^p))^{\frac{1}{p}}$ , where  $\xi$  and  $\eta$  are two random variables with marginal distributions  $\alpha$  and  $\beta$ , respectively. Moreover, let  $\zeta$  be the random variable uniformly distributed on  $[0, 1]$ , then  $F_{\alpha}^{-1}(\zeta)$  has distribution function  $F_{\alpha}$  and  $F_{\beta}^{-1}(\zeta)$  has distribution function  $F_{\beta}$  (see for example Vallender [97]). Let  $\xi = F_{\alpha}^{-1}(\zeta)$  and  $\eta = F_{\beta}^{-1}(\zeta)$ , then we have

$$d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta) \leq (\mathbb{E}(\Lambda_q(\xi, \eta)^p))^{\frac{1}{p}} = \left(\int_0^1 \Lambda_q(F_{\alpha}^{-1}(t), F_{\beta}^{-1}(t))^p dt\right)^{\frac{1}{p}}.$$

Next, we assume that  $q \leq p$ . By Lemma A.6, we have that

$$\left(d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta)\right)^p = \left(d_{\mathbb{W}, \frac{p}{q}}^{(\mathbb{R}_{\geq 0}, \Lambda_1)}((S_q)_{\#} \alpha, (S_q)_{\#} \beta)\right)^{\frac{p}{q}}.$$

Then,

$$\left(d_{\mathbb{W}, \frac{p}{q}}^{(\mathbb{R}_{\geq 0}, \Lambda_1)}((S_q)_{\#} \alpha, (S_q)_{\#} \beta)\right)^{\frac{p}{q}} = \int_0^1 |F_{\alpha,q}^{-1}(t) - F_{\beta,q}^{-1}(t)|^{\frac{p}{q}} dt,$$

where  $F_{\alpha,q}$  and  $F_{\beta,q}$  are distribution functions of  $(S_q)_{\#} \alpha$  and  $(S_q)_{\#} \beta$ , respectively. It is easy to verify that  $F_{\alpha,q}(t) = (F_{\alpha}^{-1}(t))^q$  and  $F_{\beta,q}(t) = (F_{\beta}^{-1}(t))^q$ . Therefore,

$$d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha, \beta) = \left(\int_0^1 \Lambda_q(F_{\alpha}^{-1}(t), F_{\beta}^{-1}(t))^p dt\right)^{\frac{1}{p}}$$

Finally, we demonstrate that for  $q > p$  the equality does not hold in general. We first consider the extreme case  $p = 1$  and  $q = \infty$  (though we require  $q < \infty$  in the assumptions of the theorem, we relax this for now). Let  $\alpha_0 = \frac{1}{2}\delta_1 + \frac{1}{2}\delta_2$  and  $\beta_0 = \frac{1}{2}\delta_2 + \frac{1}{2}\delta_3$  where  $\delta_x$  means the Dirac measure at point  $x \in \mathbb{R}_{\geq 0}$ . Then, we have that

$$d_{\mathbb{W},1}^{(\mathbb{R}_{\geq 0}, \Lambda_{\infty})}(\alpha_0, \beta_0) = \frac{3}{2} < \frac{5}{2} = \int_0^1 \Lambda_{\infty}(F_{\alpha}^{-1}(t), F_{\beta}^{-1}(t)) dt.$$

It is not hard to see that both  $d_{\mathbb{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha_0, \beta_0)$  and  $\left(\int_0^1 \Lambda_q(F_{\alpha}^{-1}(t), F_{\beta}^{-1}(t))^p dt\right)^{\frac{1}{p}}$  are continuous with respect to  $p \in [1, \infty)$  and  $q \in [1, \infty]$ . Then, for  $p$  close to 1 and  $q < \infty$  large enough,

and in particular,  $p < q$ , we have that

$$d_{\mathbf{W},p}^{(\mathbb{R}_{\geq 0}, \Lambda_q)}(\alpha_0, \beta_0) < \left( \int_0^1 \Lambda_q(F_\alpha^{-1}(t), F_\beta^{-1}(t))^p dt \right)^{\frac{1}{p}}.$$

□

**A.3.2. Miscellaneous.** In the remainder of this section, we collect several technical results that find implicit or explicit usage throughout Section 2.

**Lemma A.7.** *Let  $X$  be a complete ultrametric space. Then,  $X$  is compact ultrametric space if and only if for any  $t > 0$ ,  $X_t$  is a finite space.*

*Proof.* Wan [102, Lemma 2.3] proves that whenever  $X$  is compact,  $X_t$  is finite for any  $t > 0$ .

Conversely, we assume that  $X_t$  is finite for any  $t > 0$ . We only need to prove that  $X$  is totally bounded. For any  $\varepsilon > 0$ ,  $X_\varepsilon$  is a finite set and thus there exists  $x_1, \dots, x_n \in X$  such that  $X_\varepsilon = \{[x_1]_\varepsilon, \dots, [x_n]_\varepsilon\}$ . Now, for any  $x \in X$ , there exists  $x_i$  for some  $i = 1, \dots, n$  such that  $x \in [x_i]_\varepsilon$ . This implies that  $u_X(x, x_i) \leq \varepsilon$ . Therefore, the set  $\{x_1, \dots, x_n\} \subseteq X$  is an  $\varepsilon$ -net of  $X$ . Then,  $X$  is totally bounded and thus compact. □

**Lemma A.8.**  *$V(X)$  is the collection of all closed balls in  $X$  except for singletons  $\{x\}$  such that  $x$  is a cluster point in  $X$ . In particular,  $X \in V(X)$  and for any  $x \in X$ , if  $x$  is not a cluster point, then  $\{x\} \in V(X)$ .*

*Proof.* Given any  $t > 0$  and  $x \in X$ ,  $[x]_t = B_t(x) = \{x' \in X \mid u_X(x, x') \leq t\}$ . Therefore,  $V(X)$  is a collection of closed balls in  $X$ . On the contrary, any closed ball  $B_t(x)$  with positive radius  $t > 0$  coincides with  $[x]_t \in \theta_X(t)$  and thus belongs to  $V(X)$ . Now, for any singleton  $\{x\} = B_0(x)$ . If  $x$  is not a cluster point, then there exists  $t > 0$  such that  $B_t(x) = \{x\}$  which implies that  $\{x\} \in V(X)$ . If  $x$  is a cluster point, then for any  $t > 0$ ,  $\{x\} \subsetneq B_t(x) = [x]_t$ . In particular, this implies that  $\{x\} \neq [x]_t$  for all  $t > 0$  and thus  $\{x\} \notin V(X)$ . In conclusion,  $V(X)$  is the collection of all closed balls in  $X$  except for singletons  $\{x\}$  such that  $x$  is a cluster point in  $X$ .

If  $X$  is a one point space, then obviously  $X \in V(X) = \{X\}$ . Otherwise, let  $\delta := \text{diam}(X) > 0$ , then for any  $x \in X$  we have that  $X = [x]_\delta \in V(X)$ . As for singletons  $\{x\}$  where  $x \in X$  is not a cluster point, we have proved above that  $\{x\} \in V(X)$ . □

## APPENDIX B. MISSING DETAILS FROM SECTION 3

**B.1. Proofs from Section 3.1.** Next, we give the missing proofs of the results stated in Section 3.1.

**B.1.1. Proof of Proposition 3.3.**

- (1) This directly follows from the definitions of  $u_{\text{GW},p}^{\text{sturm}}$  and  $d_{\text{GW},p}^{\text{sturm}}$  (see Equation (10) and Equation (5)).
- (2) This simply follows from Jensen's inequality.

- (3) By (2), we know that  $\{u_{\text{GW},n}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})\}_{n \in \mathbb{N}}$  is an increasing sequence with a finite upper bound  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ . Therefore,  $L := \lim_{n \rightarrow \infty} u_{\text{GW},n}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  exists and  $L \leq u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .

Next, we come to the opposite inequality. By Proposition B.1, there exist  $u_n \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\mu_n \in \mathcal{C}(\mu_X, \mu_Y)$  such that

$$\left( \int_{X \times Y} (u_n(x, y))^n \mu_n(dx \times dy) \right)^{\frac{1}{n}} = u_{\text{GW},n}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}).$$

By Lemma B.19 and Lemma B.21, the sequence  $\{u_n\}_{n \in \mathbb{N}}$  uniformly converges to some  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\{\mu_n\}_{n \in \mathbb{N}}$  weakly converges to some  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  (after taking appropriate subsequences of both sequences). Let  $M := \sup_{(x,y) \in \text{supp}(\mu)} u(x, y)$ . Let  $\varepsilon > 0$  and let  $U = \{(x, y) \in X \times Y \mid u(x, y) > M - \varepsilon\}$ . Then,  $\mu(U) > 0$ . Since  $U$  is open, it follows that there exists a small  $\varepsilon_1 > 0$  such that  $\mu_n(U) > \mu(U) - \varepsilon_1 > 0$  for all  $n$  large enough (see e.g. Billingsley [7, Thm. 2.1]). Moreover, by uniform convergence of the sequence  $\{u_n\}_{n \in \mathbb{N}}$ , we have  $|u(x, y) - u_n(x, y)| \leq \varepsilon$  for any  $(x, y) \in X \times Y$  when  $n$  is large enough. Therefore, we obtain for  $n$  large enough

$$\left( \int_{X \times Y} (u_n(x, y))^n \mu_n(dx \times dy) \right)^{\frac{1}{n}} \geq (\mu_n(U))^{\frac{1}{n}} (M - 2\varepsilon) \geq (\mu(U) - \varepsilon_1)^{\frac{1}{n}} (M - 2\varepsilon).$$

Letting  $n \rightarrow \infty$ , we obtain  $L \geq M - 2\varepsilon$ . Since  $\varepsilon > 0$  is arbitrary, we obtain  $L \geq M \geq u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .

B.1.2. *Proof of Theorem 3.4.* In this section, we devote to prove Theorem 3.4. To this end, we will first verify the existence of optimal metrics and optimal couplings in Equation (18).

**Proposition B.1** (Existence of optimal couplings). *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be compact ultrametric measure spaces. Then, there always exist  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  such that for  $1 \leq p < \infty$*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \left( \int_{X \times Y} (u(x, y))^p \mu(dx \times dy) \right)^{\frac{1}{p}}$$

and such that

$$u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \sup_{(x,y) \in \text{supp}(\mu)} u(x, y).$$

*Proof.* The following proof is a suitable adaptation from proof of Lemma 3.3 in [92]. We will only prove the claim for the case  $p < \infty$  since the case  $p = \infty$  can be shown in a similar manner. Let  $u_n \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\mu_n \in \mathcal{C}(\mu_X, \mu_Y)$  be such that

$$\left( \int_{X \times Y} (u_n(x, y))^p \mu_n(dx \times dy) \right)^{\frac{1}{p}} \leq u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) + \frac{1}{n}.$$

By Lemma B.19,  $\{\mu_n\}_{n \in \mathbb{N}}$  weakly converges (after taking an appropriate subsequence) to some  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ . By Lemma B.21,  $\{u_n\}_{n \in \mathbb{N}}$  uniformly converges (after taking an appropriate subsequence) to some  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$ . Then, it is easy to verify that

$$\left( \int_{X \times Y} (u(x, y))^p \mu(dx \times dy) \right)^{\frac{1}{p}} \leq u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}).$$

□

As a direct consequence of the proposition, we get the subsequent result.

**Corollary B.2.** *Fix  $1 \leq p \leq \infty$ . Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be compact ultrametric measure spaces. Then, there exist a compact ultrametric space  $Z$  and isometric embeddings  $\phi : X \hookrightarrow Z$  and  $\psi : Y \hookrightarrow Z$  such that*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = d_{\text{W},p}^Z(\phi_{\#}\mu_X, \psi_{\#}\mu_Y).$$

Before we come to the proof of Theorem 3.4, it remains to establish another auxiliary result. We ensure that the Wasserstein pseudometric of order  $p$  on a compact pseudo-ultrametric space  $(X, u_X)$  is for  $p \in [1, \infty)$  a  $p$ -pseudometric and for  $p = \infty$  a pseudo-ultrametric, i.e., we prove for  $1 \leq p < \infty$  that for all  $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{P}(X)$

$$d_{\text{W},p}^{(X,u_X)}(\mu_1, \mu_3) \leq \left( \left( d_{\text{W},p}^{(X,u_X)}(\mu_1, \mu_2) \right)^p + \left( d_{\text{W},p}^{(X,u_X)}(\mu_2, \mu_3) \right)^p \right)^{1/p}$$

and for  $p = \infty$  that for all  $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{P}(X)$

$$d_{\text{W},p}^{(X,u_X)}(\mu_1, \mu_3) \leq \max \left( d_{\text{W},p}^{(X,u_X)}(\mu_1, \mu_2), d_{\text{W},p}^{(X,u_X)}(\mu_2, \mu_3) \right).$$

**Lemma B.3.** *Let  $(X, u_X)$  be a compact pseudo-ultrametric space. Then, for  $1 \leq p \leq \infty$  the  $p$ -Wasserstein metric  $d_{\text{W},p}^{(X,u_X)}$  is a  $p$ -pseudometric on  $\mathcal{P}(X)$ . In particular, when  $p = \infty$ , it is an pseudo-ultrametric on  $\mathcal{P}(X)$ .*

*Proof.* We prove the statement by adapting the proof of the triangle inequality for the  $p$ -Wasserstein distance (see e.g., [99, Theorem 7.3]). We only prove the case when  $p < \infty$  whereas the case  $p = \infty$  follows by analogous arguments.

Let  $\alpha_1, \alpha_2, \alpha_3 \in \mathcal{P}(X)$ , denote by  $\mu_{12}$  an optimal transport plan between  $\alpha_1$  and  $\alpha_2$  and by  $\mu_{23}$  an optimal transport plan between  $\alpha_2$  and  $\alpha_3$  (see [100, Theorem 4.1] for the existence of  $\mu_{12}$  and  $\mu_{23}$ ). Furthermore, let  $X_i$  be the support of  $\alpha_i$ ,  $1 \leq i \leq 3$ . Then, by the Gluing Lemma [99, Lemma 7.6] there exists a measure  $\mu \in \mathcal{P}(X_1 \times X_2 \times X_3)$  with marginals  $\mu_{12}$  on  $X_1 \times X_2$  and  $\mu_{23}$  on  $X_2 \times X_3$ . Clearly, we obtain

$$\begin{aligned} \left( d_{\text{W},p}^{(X,u_X)}(\alpha_1, \alpha_3) \right)^p &\leq \int_{X_1 \times X_2 \times X_3} u_X^p(x, z) \mu(dx \times dy \times dz) \\ &\leq \int_{X_1 \times X_2 \times X_3} (u_X^p(x, y) + u_X^p(y, z)) \mu(dx \times dy \times dz). \end{aligned}$$

Here, we used that  $u_X$  is an ultrametric, i.e., in particular a  $p$ -metric [70, Proposition 1.16]. With this we obtain that

$$\begin{aligned} \left( d_{\text{W},p}^{(X,u_X)}(\alpha_1, \alpha_2) \right)^p &\leq \int_{X_1 \times X_2} u_X^p(x, y) \mu_{12}(dx \times dy) + \int_{X_2 \times X_3} u_X^p(y, z) \mu_{23}(dy \times dz) \\ &= \left( d_{\text{W},p}^{(X,u_X)}(\alpha_1, \alpha_2) \right)^p + \left( d_{\text{W},p}^{(X,u_X)}(\alpha_2, \alpha_3) \right)^p. \end{aligned}$$

□

With Proposition B.1 and Lemma B.3 at our disposal we are now ready to prove Theorem 3.4 which states that  $u_{\text{GW},p}^{\text{sturm}}$  is indeed a  $p$ -metric on  $\mathcal{U}^w$ .

*Proof of Theorem 3.4.* It is clear that  $u_{\text{GW},p}^{\text{sturm}}$  is symmetric and that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = 0$  if  $\mathcal{X} \cong_w \mathcal{Y}$ . Furthermore, we remark that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \geq d_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  by Proposition 3.3. Since  $d_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = 0$  implies that  $\mathcal{X} \cong_w \mathcal{Y}$  ([93]), we have that  $u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = 0$  implies that  $\mathcal{X} \cong_w \mathcal{Y}$ . It remains to verify the  $p$ -triangle inequality. To this end, we only prove the case when  $p < \infty$  whereas the case  $p = \infty$  follows by analogous arguments.

Let  $\mathcal{X}, \mathcal{Y}, \mathcal{Z} \in \mathcal{U}^w$ . Suppose  $u_{XY} \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $u_{YZ} \in \mathcal{D}^{\text{ult}}(u_Y, u_Z)$  are optimal metric couplings such that

$$(u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}))^p = \left( d_{\text{W},p}^{(X \sqcup Y, u_{XY})}(\mu_X, \mu_Y) \right)^p \quad \text{and} \quad (u_{\text{GW},p}^{\text{sturm}}(\mathcal{Y}, \mathcal{Z}))^p = \left( d_{\text{W},p}^{(Y \sqcup Z, u_{YZ})}(\mu_Y, \mu_Z) \right)^p.$$

Further, define  $u_{XYZ}$  on  $X \sqcup Y \sqcup Z$  as

$$u_{XYZ}(x_1, x_2) = \begin{cases} u_{XY}(x_1, x_2) & x_1, x_2 \in X \sqcup Y \\ u_{YZ}(x_1, x_2) & x_1, x_2 \in Y \sqcup Z \\ \inf\{\max(u_{XY}(x_1, y), u_{YZ}(y, x_2)) \mid y \in Y\} & x_1 \in X, x_2 \in Z \\ \inf\{\max(u_{XY}(x_2, y), u_{YZ}(y, x_1)) \mid y \in Y\} & x_1 \in Z, x_2 \in X. \end{cases}$$

Then, by Lemma 1.1 of Zarichnyi [105]  $u_{XYZ}$  is a pseudo-ultrametric on  $X \sqcup Y \sqcup Z$  that coincides with  $u_{XY}$  on  $X \sqcup Y$  and with  $u_{YZ}$  on  $Y \sqcup Z$ . With this we obtain by Lemma B.3 that

$$\begin{aligned} (u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Z}))^p &\leq \left( d_{\text{W},p}^{(X \sqcup Y \sqcup Z, u_{XYZ})}(\mu_X, \mu_Z) \right)^p \\ &\leq \left( d_{\text{W},p}^{(X \sqcup Y \sqcup Z, u_{XYZ})}(\mu_X, \mu_Y) \right)^p + \left( d_{\text{W},p}^{(X \sqcup Y \sqcup Z, u_{XYZ})}(\mu_Y, \mu_Z) \right)^p \\ &= \left( d_{\text{W},p}^{(X \sqcup Y, u_{XY})}(\mu_X, \mu_Y) \right)^p + \left( d_{\text{W},p}^{(Y \sqcup Z, u_{YZ})}(\mu_Y, \mu_Z) \right)^p \\ &= (u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}))^p + (u_{\text{GW},p}^{\text{sturm}}(\mathcal{Y}, \mathcal{Z}))^p \end{aligned}$$

This gives the claim for  $p < \infty$ . □

**B.1.3. Proof of Theorem 3.7.** In order to proof Theorem 3.7, we will first establish the statement for *finite* ultrametric measure spaces. For this purpose, we need to introduce some notation. Given  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ , let  $\mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$  denote the collection of all admissible pseudo-ultrametrics on  $X \sqcup Y$ , where  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  is called *admissible*, if there exists no  $u^* \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  such that  $u^* \neq u$  and  $u^*(x, y) \leq u(x, y)$  for all  $x, y \in X \sqcup Y$ .

**Lemma B.4.** *For any  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ ,  $\mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y) \neq \emptyset$ . Moreover,*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)} d_{\text{W},p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y).$$

*Proof.* If  $\{u_n\}_{n \in \mathbb{N}} \subseteq \mathcal{D}^{\text{ult}}(u_X, u_Y)$  is a decreasing sequence (with respect to pointwise inequality), it is easy to verify that  $u := \inf_{n \in \mathbb{N}} u_n \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and thus  $u$  is a lower bound of  $\{u_n\}_{n \in \mathbb{N}}$ . Then, by Zorn's lemma  $\mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y) \neq \emptyset$ . Therefore, we obtain that

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)} d_{\text{W},p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y).$$

□

Combined with Example 3.6, the following result implies that each  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$  gives rise to an element in  $\mathcal{A}$ .

**Lemma B.5.** *Given finite spaces  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ , for each  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$ ,  $u^{-1}(0) \neq \emptyset$ .*

*Proof.* Assume otherwise that  $u^{-1}(0) = \emptyset$ . Then,  $u$  is a metric (instead of pseudo-metric). Let  $(x_0, y_0) \in X \times Y$  such that  $u(x_0, y_0) = \min_{x \in X, y \in Y} u(x, y)$ . The existence of  $(x_0, y_0)$  is guaranteed by the finiteness of  $X$  and  $Y$ . We define  $u_{(x_0, y_0)} : X \sqcup Y \times X \sqcup Y \rightarrow \mathbb{R}_{\geq 0}$  as follows:

- (1)  $u_{(x_0, y_0)}|_{X \times X} := u_X$  and  $u_{(x_0, y_0)}|_{Y \times Y} := u_Y$ ;
- (2) For  $(x, y) \in X \times Y$ ,  

$$u_{(x_0, y_0)}(x, y) := \min(u(x, y), \max(u_X(x, x_0), u_Y(y, y_0)))$$
;
- (3) For any  $(y, x) \in Y \times X$ ,  $u_{(x_0, y_0)}(y, x) := u_{(x_0, y_0)}(x, y)$ .

It is easy to verify that  $u_{(x_0, y_0)} \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$ . Further, it is obvious that  $u_{(x_0, y_0)}(x_0, y_0) = 0 < u(x_0, y_0)$  and that  $u_{(x_0, y_0)}(x, y) \leq u(x, y)$  for all  $x, y \in X \sqcup Y$  which contradicts with  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$ . Therefore,  $u^{-1}(0) \neq \emptyset$ .  $\square$

**Theorem B.6.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  be finite spaces. Then, we have for each  $p \in [1, \infty)$  that*

$$u_{\text{GW}, p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{(A, \varphi) \in \mathcal{A}} d_{\text{W}, p}^{Z_A} \left( (\phi_{(A, \varphi)}^X)_{\#} \mu_X, (\psi_{(A, \varphi)}^Y)_{\#} \mu_Y \right). \quad (30)$$

*Proof.* By Lemma B.4 it is sufficient to prove that each  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$  induces  $(A, \varphi) \in \mathcal{A}$  such that

$$d_{\text{W}, p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y) \geq d_{\text{W}, p}^{Z_A} \left( (\phi_{(A, \varphi)}^X)_{\#} \mu_X, (\psi_{(A, \varphi)}^Y)_{\#} \mu_Y \right).$$

Let  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$ . We define  $A_0 := \{x \in X \mid \exists y \in Y \text{ such that } u(x, y) = 0\}$  ( $A_0 \neq \emptyset$  by Lemma B.5). By Example 3.6, the map  $\varphi_0 : A_0 \rightarrow Y$  defined by taking  $x$  to  $y$  such that  $u(x, y) = 0$  is a well-defined isometric embedding. This means in particular that  $(A_0, \varphi_0) \in \mathcal{A}$ .

If  $u(x, y) \geq u_{Z_{A_0}} \left( \phi_{(A_0, \varphi_0)}^X(x), \psi_{(A_0, \varphi_0)}^Y(y) \right)$  holds for all  $(x, y) \in X \times Y$ , then we set  $A := A_0$  and  $\varphi := \varphi_0$ . This gives

$$d_{\text{W}, p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y) \geq d_{\text{W}, p}^{Z_A} \left( (\phi_{(A, \varphi)}^X)_{\#} \mu_X, (\psi_{(A, \varphi)}^Y)_{\#} \mu_Y \right).$$

Otherwise, there exists  $(x, y) \in X \setminus A_0 \times Y \setminus \varphi_0(A_0)$  such that

$$u(x, y) < u_{Z_{A_0}} \left( \phi_{(A_0, \varphi_0)}^X(x), \psi_{(A_0, \varphi_0)}^Y(y) \right)$$

(if  $x \in A_0$  or  $y \in \varphi_0(A_0)$ , then  $u(x, y) \geq u_{Z_{A_0}} \left( \phi_{(A_0, \varphi_0)}^X(x), \psi_{(A_0, \varphi_0)}^Y(y) \right)$  must hold). Let  $(x_1, y_1) \in X \setminus A_0 \times Y \setminus \varphi_0(A_0)$  be such that

$$u(x_1, y_1) = \min \left\{ u(x, y) \mid (x, y) \in X \setminus A_0 \times Y \setminus \varphi_0(A_0) \right. \\ \left. \text{and } u(x, y) < u_{Z_{A_0}} \left( \phi_{(A_0, \varphi_0)}^X(x), \psi_{(A_0, \varphi_0)}^Y(y) \right) \right\} > 0.$$

The existence of  $(x_1, y_1)$  follows from finiteness of  $X$  and  $Y$ . It is easy to check that  $\varphi_0$  extends to an isometry from  $A_0 \cup \{x_1\}$  to  $\varphi_0(A_0) \cup \{y_1\}$  by taking  $x_1$  to  $y_1$ . We denote the new isometry  $\varphi_1$  and set  $A_1 := A_0 \cup \{x_1\}$ . If for any  $(x, y) \in X \times Y$ , we have that  $u(x, y) \geq u_{Z_{A_1}} \left( \phi_{(A_1, \varphi_1)}^X(x), \psi_{(A_1, \varphi_1)}^Y(y) \right)$ , then we define  $A := A_1$  and  $\varphi := \varphi_1$ . Otherwise, we continue the process to obtain  $A_2, A_3, \dots$ . This process will eventually stop since we are considering finite spaces. Suppose the process stops at  $A_n$ , then  $A := A_n$  and  $\varphi := \varphi_n$  satisfy that  $u(x, y) \geq u_{Z_A} \left( \phi_{(A, \varphi)}^X(x), \psi_{(A, \varphi)}^Y(y) \right)$  for any  $(x, y) \in X \times Y$ . Therefore,

$$d_{W,p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y) \geq d_{W,p}^{Z_A} \left( (\phi_{(A, \varphi)}^X)_\# \mu_X, (\psi_{(A, \varphi)}^Y)_\# \mu_Y \right).$$

Since  $u \in \mathcal{D}_{\text{adm}}^{\text{ult}}(u_X, u_Y)$  is arbitrary, this gives the claim.  $\square$

As a direct consequence of Theorem B.6, we obtain that it is sufficient, as claimed in Remark 3.8, for finite spaces to infimize in Equation (30) over the collection of all maximal pairs  $\mathcal{A}^* \subseteq \mathcal{A}$ . Recall that a pair  $(A, \varphi_1) \in \mathcal{A}$  is denoted as *maximal*, if for all pairs  $(B, \varphi_2) \in \mathcal{A}$  with  $A \subseteq B$  and  $\varphi_2|_A = \varphi_1$  it holds  $A = B$ .

**Corollary B.7.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  be finite spaces. Then, we have for each  $p \in [1, \infty]$  that*

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \inf_{(A, \varphi) \in \mathcal{A}^*} d_{W,p}^{Z_A} \left( (\phi_{(A, \varphi)}^X)_\# \mu_X, (\psi_{(A, \varphi)}^Y)_\# \mu_Y \right). \quad (31)$$

By proving Theorem B.6, we have verified Theorem 3.7 for finite ultrametric measure spaces. In the following, we will use Theorem B.6 and weighted quotients to demonstrate Theorem 3.7. However, before we come to this, we need to establish the following two auxiliary results.

**Lemma B.8.** *Let  $X \in \mathcal{U}$  be a compact ultrametric space. Let  $t > 0$  and let  $p \in [1, \infty)$ . Then, for any  $\alpha, \beta \in \mathcal{P}(X)$ , we have that*

$$(d_{W,p}^{X_t}(\alpha_t, \beta_t))^p \geq (d_{W,p}^X(\alpha, \beta))^p - t^p,$$

where  $\alpha_t$  is the push forward of  $\alpha$  under the canonical quotient map  $Q_t : X \rightarrow X_t$  taking  $x \in X$  to  $[x]_t \in X_t$ .

*Proof.* For any  $\mu_t \in \mathcal{C}(\alpha_t, \beta_t)$ , it is easy to see that there exists  $\mu \in \mathcal{C}(\alpha, \beta)$  such that  $\mu_t = (Q_t \times Q_t)_\# \mu$  where  $Q_t \times Q_t : X \times X \rightarrow X_t \times X_t$  maps  $(x, x') \in X \times X$  to  $([x]_t, [x']_t)$ . For example, suppose  $X_t = \{[x_1]_t, \dots, [x_n]_t\}$ , then one can let

$$\mu := \sum_{i,j=1}^n \mu_t([x_i]_t, [x_j]_t) \frac{\alpha|_{[x_i]_t}}{\alpha([x_i]_t)} \otimes \frac{\beta|_{[x_j]_t}}{\beta([x_j]_t)},$$

where  $\alpha|_{[x_i]_t}$  is the restriction of  $\alpha$  on  $[x_i]_t$ .

For any  $x, x' \in X$ , we have that  $(u_X(x, x'))^p \leq (u_{X_t}([x]_t, [x']_t))^p + t^p$ . Then,

$$\begin{aligned} (d_{W,p}^X(\alpha, \beta))^p &\leq \int_{X \times X} (u_X(x, x'))^p \mu(dx \times dx') \\ &\leq \int_{X \times X} ((u_{X_t}([x]_t, [x']_t))^p + t^p) \mu(dx \times dx') \\ &= \int_{X \times X} (u_X(Q_t(x), Q_t(x')))^p \mu(dx \times dx') + t^p \\ &= \int_{X_t \times X_t} (u_{X_t}([x]_t, [x']_t))^p \mu_t(d[x]_t \times d[x']_t) + t^p \end{aligned}$$

Infimizing over all  $\mu_t \in \mathcal{C}(\alpha_t, \beta_t)$ , we obtain that

$$(d_{W,p}^{X_t}(\alpha_t, \beta_t))^p \geq (d_{W,p}^X(\alpha, \beta))^p - t^p.$$

□

**Lemma B.9.** *Let  $\mathcal{X} \in \mathcal{U}^w$  and let  $p \in [1, \infty]$ . Then, for any  $t > 0$ , we have that*

$$u_{GW,p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{X}) \leq t.$$

*In particular,  $\lim_{t \rightarrow 0} u_{GW,p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{X}) = 0$ .*

*Proof.* It is obvious that  $(\mathcal{X}_t)_t \cong_w \mathcal{X}_t$ . Hence, it holds by Theorem 3.13 that  $u_{GW,\infty}^{\text{sturm}}(\mathcal{X}_t, \mathcal{X}) \leq t$ . By Proposition 3.3 we have that for any  $p \in [1, \infty]$

$$u_{GW,p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{X}) \leq u_{GW,\infty}^{\text{sturm}}(\mathcal{X}_t, \mathcal{X}) \leq t.$$

□

With Lemma B.8 and Lemma B.9 available, we can come to the proof of Theorem 3.7.

*Proof of Theorem 3.7.* Clearly, it follows from the definition of  $u_{GW,p}^{\text{sturm}}$  (see Equation (10)) that

$$u_{GW,p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq \inf_{(A,\varphi) \in \mathcal{A}} d_{W,p}^{Z_A} \left( (\phi_{(A,\varphi)}^X)_\# \mu_X, (\psi_{(A,\varphi)}^Y)_\# \mu_Y \right)$$

Hence, we focus on proving the opposite inequality.

Given any  $t > 0$ , by Lemma A.7, both  $\mathcal{X}_t$  and  $\mathcal{Y}_t$  are finite spaces. By Theorem B.6 we have that

$$u_{GW,p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{Y}_t) = \inf_{(A_t, \varphi_t) \in \mathcal{A}_t} d_{W,p}^{Z_{A_t}} \left( \left( \phi_{(A_t, \varphi_t)}^{X_t} \right)_\# (\mu_X)_t, \left( \psi_{(A_t, \varphi_t)}^{Y_t} \right)_\# (\mu_Y)_t \right),$$

where

$$\mathcal{A}_t := \{(A_t, \varphi_t) \mid \emptyset \neq A_t \subseteq X_t \text{ is closed and } \varphi_t : A_t \hookrightarrow Y_t \text{ is an isometric embedding}\}.$$

For any  $(A_t, \varphi_t) \in \mathcal{A}_t$ , assume that  $A_t = \{[x_1]_t^X, \dots, [x_n]_t^X\}$  and that  $\varphi_t([x_i]_t) = [y_i]_t \in Y_t$  for all  $i = 1, \dots, n$ . Let  $A := \{x_1, \dots, x_n\}$ . Then, the map  $\varphi : A \rightarrow Y$  defined by  $x_i \mapsto y_i$  for  $i = 1, \dots, n$  is an isometric embedding. Therefore,  $(A, \varphi) \in \mathcal{A}$ .

*Claim 1:*  $((Z_A)_t, u_{(Z_A)_t}) \cong (Z_{A_t}, u_{Z_{A_t}})$ .

*Proof of the Claim.* We define a map  $\Psi : (Z_A)_t \rightarrow Z_{A_t}$  by  $[x]_t^{Z_A} \mapsto [x]_t^X$  for  $x \in X$  and  $[y]_t^{Z_A} \mapsto [y]_t^Y$  for  $y \in Y \setminus \varphi(A)$ . We first show that  $\Psi$  is well-defined. For any  $x' \in X$ , if  $u_{Z_A}(x, x') \leq t$ , then obviously we have that  $u_X(x, x') = u_{Z_A}(x, x') \leq t$  and thus  $[x]_t^X = [x']_t^X$ . Now, assume that there exists  $y \in Y \setminus \varphi(A)$  such that  $u_{Z_A}(x, y) \leq t$ , i.e.,  $[x]_t^{Z_A} = [y]_t^{Z_A}$ . Then, by finiteness of  $A$  and definition of  $Z_A$ , there exists  $x_i \in A$  such that  $u_{Z_A}(x, y) = \max(u_X(x, x_i), u_Y(\varphi(x_i), y)) \leq t$ . This gives that

$$u_{Z_{A_t}}([x]_t^X, [y]_t^Y) \leq \max(u_{X_t}([x]_t^X, [x_i]_t^X), u_{Y_t}([\varphi(x_i)]_t^Y, [y]_t^Y)) \leq t.$$

However, this happens only if  $u_{Z_{A_t}}([x]_t^X, [y]_t^Y) = 0$ , that is,  $[x]_t^X$  is identified with  $[y]_t^Y$  under the map  $\varphi_t$ . Therefore,  $\Psi$  is well-defined.

It is easy to see from the definition that  $\Psi$  is surjective. Thus, it suffices to show that  $\Psi$  is an isometric embedding to finish the proof. For any  $x, x' \in X$  such that  $u_X(x, x') > t$ , we have that

$$u_{(Z_A)_t}([x]_t^{Z_A}, [x']_t^{Z_A}) = u_{Z_A}(x, x') = u_X(x, x') = u_{X_t}([x]_t^X, [x']_t^X) = u_{Z_{A_t}}([x]_t^X, [x']_t^X).$$

Similarly, for any  $y, y' \in Y \setminus \varphi(A)$  such that  $u_Y(y, y') > t$ , we have that

$$u_{(Z_A)_t}([y]_t^{Z_A}, [y']_t^{Z_A}) = u_{Z_{A_t}}([y]_t^Y, [y']_t^Y).$$

Now, consider  $x \in X$  and  $y \in Y \setminus \varphi(A)$ . Assume that  $u_{Z_A}(x, y) > t$  (otherwise  $[x]_t^{Z_A} = [y]_t^{Z_A}$ ). Then, we have that

$$u_{Z_A}(x, y) = \min_{i=1, \dots, n} \max(u_X(x, x_i), u_Y(\varphi(x_i), y)) > t.$$

This implies that

$$\begin{aligned} u_{Z_{A_t}}([x]_t^X, [y]_t^Y) &= \min_{i=1, \dots, n} \max(u_{X_t}([x]_t^X, [x_i]_t^X), u_{Y_t}(\varphi_t([x_i]_t^X), [y]_t^Y)) \\ &= \min_{i=1, \dots, n} \max(u_X(x, x_i), u_Y(\varphi(x_i), y)) \\ &= u_{Z_A}(x, y) = u_{(Z_A)_t}([x]_t^{Z_A}, [y]_t^{Z_A}). \end{aligned}$$

Therefore,  $\Psi$  is an isometric embedding and thus we conclude the proof.  $\square$

By Lemma B.8 we have that

$$\begin{aligned} &\left( d_{W,p}^{Z_{A_t}} \left( \left( \phi_{(A_t, \varphi_t)}^{X_t} \right)_{\#} (\mu_X)_t, \left( \psi_{(A_t, \varphi_t)}^{Y_t} \right)_{\#} (\mu_Y)_t \right) \right)^p \\ &\geq \left( d_{W,p}^{Z_A} \left( \left( \phi_{(A, \varphi)}^X \right)_{\#} \mu_X, \left( \psi_{(A, \varphi)}^Y \right)_{\#} \mu_Y \right) \right)^p - t^p \end{aligned}$$

Therefore,

$$\begin{aligned} u_{GW,p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{Y}_t) &= \inf_{(A_t, \varphi_t) \in \mathcal{A}_t} d_{W,p}^{Z_{A_t}} \left( \left( \phi_{(A_t, \varphi_t)}^{X_t} \right)_{\#} (\mu_X)_t, \left( \psi_{(A_t, \varphi_t)}^{Y_t} \right)_{\#} (\mu_Y)_t \right) \\ &\geq \inf_{(A, \varphi) \in \mathcal{A}} \left( \left( d_{W,p}^{Z_A} \left( \left( \phi_{(A, \varphi)}^X \right)_{\#} \mu_X, \left( \psi_{(A, \varphi)}^Y \right)_{\#} \mu_Y \right) \right)^p - t^p \right)^{\frac{1}{p}}. \end{aligned}$$

Notice that the last inequality already holds when we only consider  $(A, \varphi)$  corresponding to  $(A_t, \varphi_t) \in \mathcal{A}_t$ .

By Lemma B.9, we have that

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \lim_{t \rightarrow 0} u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}_t, \mathcal{Y}_t) \geq \inf_{(A,\varphi) \in \mathcal{A}} d_{\text{W},p}^{Z_A} \left( (\phi_{(A,\varphi)}^X)_\# \mu_X, (\psi_{(A,\varphi)}^Y)_\# \mu_Y \right),$$

which concludes the proof.  $\square$

**B.2. Proofs from Section 3.2.** In the following, we give the complete proofs of the results stated in Section 3.2.

**B.2.1. Proof of Proposition 3.9.**

- (1) This follows directly from the definitions of  $u_{\text{GW},p}$  and  $d_{\text{GW},p}$  (see Equation (13) and Equation (7)).
- (2) By Jensen's inequality we have that  $\text{dis}_p^{\text{ult}}(\mu) \leq \text{dis}_q^{\text{ult}}(\mu)$  for any  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ . Therefore,  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \leq u_{\text{GW},q}(\mathcal{X}, \mathcal{Y})$ .
- (3) By (2), we know that  $\{u_{\text{GW},n}(\mathcal{X}, \mathcal{Y})\}_{n \in \mathbb{N}}$  is an increasing sequence with a finite upper bound  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ . Therefore,  $L := \lim_{n \rightarrow \infty} u_{\text{GW},n}(\mathcal{X}, \mathcal{Y})$  exists and it holds  $L \leq u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ .

To prove the opposite inequality, by Proposition B.10, there exists for each  $n \in \mathbb{N}$   $\mu_n \in \mathcal{C}(\mu_X, \mu_Y)$  such that

$$\left( \iint_{X \times Y \times X \times Y} \Lambda_\infty(u_X(x, x'), u_Y(y, y'))^n \mu_n(dx \times dy) \mu_n(dx' \times dy') \right)^{\frac{1}{n}} = u_{\text{GW},n}(\mathcal{X}, \mathcal{Y}).$$

By Lemma B.19,  $\{\mu_n\}_{n \in \mathbb{N}}$  weakly converges (after taking an appropriate subsequence) to some  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ . Let

$$M = \sup_{(x,y),(x',y') \in \text{supp}(\mu)} \Lambda_\infty(u_X(x, x'), u_Y(y, y'))$$

and for any given  $\varepsilon > 0$  let

$$U = \{((x, y), (x', y')) \in X \times Y \times X \times Y \mid \Lambda_\infty(u_X(x, x'), u_Y(y, y')) > M - \varepsilon\}.$$

Then, we have  $\mu \otimes \mu(U) > 0$ . As  $\mu_n$  weakly converges to  $\mu$ , we have that  $\mu_n \otimes \mu_n$  weakly converges to  $\mu \otimes \mu$ . Since  $U$  is open, there exists a small  $\varepsilon_1 > 0$  such that  $\mu_n \otimes \mu_n(U) > \mu \otimes \mu(U) - \varepsilon_1 > 0$  for  $n$  large enough (see e.g. Billingsley [7, Thm. 2.1]). Therefore,

$$\begin{aligned} & \left( \iint_{X \times Y \times X \times Y} \Lambda_\infty(u_X(x, x'), u_Y(y, y'))^n \mu_n(dx \times dy) \mu_n(dx' \times dy') \right)^{\frac{1}{n}} \\ & \geq (\mu_n \otimes \mu_n(U))^{\frac{1}{n}} (M - \varepsilon) \geq (\mu \otimes \mu(U) - \varepsilon_1)^{\frac{1}{n}} (M - \varepsilon). \end{aligned}$$

Letting  $n \rightarrow \infty$ , we obtain  $L \geq M - \varepsilon$ . Since  $\varepsilon > 0$  is arbitrary, we obtain  $L \geq M \geq u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ .

B.2.2. *Proof of Theorem 3.10.* One main step to verify Theorem 3.10 is to demonstrate the existence of optimal couplings.

**Proposition B.10.** *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be compact ultrametric measure spaces. Then, for any  $p \in [1, \infty]$ , there always exists an optimal coupling  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  such that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = \text{dis}_p^{\text{ult}}(\mu)$ .*

*Proof.* We will only prove the claim for the case  $p < \infty$  since the case  $p = \infty$  can be proven in a similar manner. Let  $\mu_n \in \mathcal{C}(\mu_X, \mu_Y)$  be such that

$$\left( \iint_{X \times Y \times X \times Y} \Lambda_\infty(u_X(x, x'), u_Y(y, y'))^p \mu_n(dx \times dy) \mu_n(dx' \times dy') \right)^{\frac{1}{p}} \leq u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) + \frac{1}{n}.$$

By Lemma B.19,  $\{\mu_n\}_{n \in \mathbb{N}}$  weakly converges to some  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  (after taking an appropriate subsequence). Then, by the boundedness and continuity of  $\Lambda_\infty(u_X, u_Y)$  on  $X \times Y \times X \times Y$  (cf. Lemma B.22) as well as the weak convergence of  $\mu_n \otimes \mu_n$ , we have that that

$$\text{dis}_p^{\text{ult}}(\mu) = \lim_{n \rightarrow \infty} \text{dis}_p^{\text{ult}}(\mu_n) \leq u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}).$$

Hence,  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = \text{dis}_p^{\text{ult}}(\mu)$ . □

Based on Proposition B.10, it is straightforward to prove Theorem 3.10.

*Proof of Theorem 3.10.* It is clear that  $u_{\text{GW},p}$  is symmetric and that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$  if  $\mathcal{X} \cong_w \mathcal{Y}$ . Furthermore, we remark that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \geq d_{\text{GW},p}(\mathcal{X}, \mathcal{Y})$  by Proposition 3.9. Since  $d_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$  implies that  $\mathcal{X} \cong_w \mathcal{Y}$  (see [67]), we have that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$  implies that  $\mathcal{X} \cong_w \mathcal{Y}$ . It remains to verify the  $p$ -triangle inequality. To this end, we only prove the case when  $p < \infty$  whereas the case  $p = \infty$  follows by analogous arguments.

Now let  $\mathcal{X}, \mathcal{Y}, \mathcal{Z}$  be three ultrametric measure spaces. Let  $\mu_{XY} \in \mathcal{C}(\mu_X, \mu_Y)$  and  $\mu_{YZ} \in \mathcal{C}(\mu_Y, \mu_Z)$  be optimal (cf. Proposition B.10). By the Gluing Lemma [99, Lemma 7.6], there exists a measure  $\mu_{XYZ} \in \mathcal{P}(X \times Y \times Z)$  with marginals  $\mu_{XY}$  on  $X \times Y$  and  $\mu_{YZ}$  on  $Y \times Z$ . Further, we define  $\mu_{XZ} = (\pi_{XZ})_\# \mu \in \mathcal{P}(X \times Z)$ , where  $\pi_{XZ}$  denotes the canonical projection

$X \times Y \times Z \rightarrow X \times Z$ . Then,

$$\begin{aligned}
(u_{\text{GW},p}(\mathcal{X}, \mathcal{Z}))^p &\leq \iint_{X \times Z \times X \times Z} (\Lambda_\infty(u_X(x, x'), u_Z(z, z')))^p \mu_{XZ}(dx \times dz) \mu_{XZ}(dx' \times dz') \\
&= \iint_{X \times Y \times Z \times X \times Y \times Z} (\Lambda_\infty(u_X(x, x'), u_Z(z, z')))^p \mu_{XYZ}(dx \times dy \times dz) \mu_{XYZ}(dx' \times dy' \times dz') \\
&\leq \iint_{X \times Y \times Z \times X \times Y \times Z} (\Lambda_\infty(u_X(x, x'), u_Y(y, y')))^p \mu_{XYZ}(dx \times dy \times dz) \mu_{XYZ}(dx' \times dy' \times dz') \\
&\quad + \iint_{X \times Y \times Z \times X \times Y \times Z} (\Lambda_\infty(u_Y(y, y'), u_Z(z, z')))^p \mu_{XYZ}(dx \times dy \times dz) \mu_{XYZ}(dx' \times dy' \times dz') \\
&= \iint_{X \times Y \times X \times Y} (\Lambda_\infty(u_X(x, x'), u_Y(y, y')))^p \mu_{XY}(dx \times dy) \mu_{XY}(dx' \times dy') \\
&\quad + \iint_{Y \times Z \times Y \times Z} (\Lambda_\infty(u_Y(y, y'), u_Z(z, z')))^p \mu_{YZ}(dy \times dz) \mu_{YZ}(dy' \times dz') \\
&= (u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}))^p + (u_{\text{GW},p}(\mathcal{Y}, \mathcal{Z}))^p,
\end{aligned}$$

where the second inequality follows from the fact that  $\Lambda_\infty$  is an ultrametric on  $\mathbb{R}_{\geq 0}$  (cf. [70, Remark 1.14]) and the observation that an ultrametric is automatically a  $p$ -metric for any  $p \in [1, \infty]$  [70, Proposition 1.16].  $\square$

**B.2.3. Proof of Theorem 3.13.** We first prove that

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\} \quad (32)$$

and then show that the infimum is attainable.

Since  $\mathcal{X}_0 \cong_w \mathcal{X}$  and  $\mathcal{Y}_0 \cong_w \mathcal{Y}$ , if  $\mathcal{X}_0 \cong_w \mathcal{Y}_0$ , then  $\mathcal{X} \cong_w \mathcal{Y}$  and thus by Theorem 3.10

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) = 0 = \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$$

Now, assume that for some  $t > 0$ ,  $\mathcal{X}_t \cong_w \mathcal{Y}_t$ . By Lemma A.7, for some  $n \in \mathbb{N}$  we can write  $X_t = \{[x_1]_t, \dots, [x_n]_t\}$  and  $Y_t = \{[y_1]_t, \dots, [y_n]_t\}$  such that  $u_{X_t}([x_i]_t, [x_j]_t) = u_{Y_t}([y_i]_t, [y_j]_t)$  and  $\mu_X([x_i]_t) = \mu_Y([y_i]_t)$ . Let  $\mu_X^i := \mu_X|_{[x_i]_t}$  and  $\mu_Y^i := \mu_Y|_{[y_i]_t}$  for all  $i = 1, \dots, n$ . Let  $\mu := \sum_{i=1}^n \mu_X^i \otimes \mu_Y^i$ . It is easy to check that  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  and  $\text{supp}(\mu) = \bigcup_{i=1}^n [x_i]_t \times [y_i]_t$ . Assume  $(x, y) \in [x_i]_t \times [y_i]_t$  and  $(x', y') \in [x_j]_t \times [y_j]_t$ . If  $i \neq j$ , then  $u_{X_t}([x_i]_t, [x_j]_t) = u_{Y_t}([y_i]_t, [y_j]_t)$  and thus

$$\Lambda_\infty(u_X(x, x'), u_Y(y, y')) = \Lambda_\infty(u_{X_t}([x_i]_t, [x_j]_t), u_{Y_t}([y_i]_t, [y_j]_t)) = 0.$$

If  $i = j$ , then  $u_X(x, x'), u_Y(y, y') \leq t$  and thus  $\Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq t$ . In either case, we have that

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) \leq \sup_{(x,y),(x',y') \in \text{supp}(\mu)} \Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq t.$$

Therefore,  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) \leq \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$ .

Conversely, suppose  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  and let  $t := \sup_{(x,y),(x',y') \in \text{supp}(\mu)} \Lambda_\infty(u_X(x, x'), u_Y(y, y'))$ . By Mémoli [67, Lemma 2.2], we know that  $\text{supp}(\mu)$  is a correspondence between  $X$  and  $Y$ .

We define a map  $f_t : X_t \rightarrow Y_t$  by taking  $[x]_t^X \in X_t$  to  $[y]_t^Y \in Y_t$  such that  $(x, y) \in \text{supp}(\mu)$ . It is easy to check that  $f_t$  is well-defined and moreover  $f_t$  is an isometry (see for example the proof of Mémoli et al. [70, Theorem 5.7]). Next, we prove that  $f_t$  is actually an isomorphism between  $\mathcal{X}_t$  and  $\mathcal{Y}_t$ . For any  $[x]_t^X \in X_t$ , let  $y \in Y$  be such that  $(x, y) \in \text{supp}(\mu)$  (in this case,  $[y]_t^Y = f_t([x]_t^X)$ ). If there exists  $(x', y') \in \text{supp}(\mu)$  such that  $x' \in [x]_t^X$  and  $y' \notin [y]_t^Y$ , then  $\Lambda_\infty(u_X(x, x'), u_Y(y, y')) = u_Y(y, y') > t$ , which is impossible. Consequently,  $\mu([x]_t^X \times (Y \setminus [y]_t^Y)) = 0$  and similarly,  $\mu((X \setminus [x]_t^X) \times [y]_t^Y) = 0$ . This yields that

$$\mu_X([x]_t^X) = \mu([x]_t^Y \times Y) = \mu([x]_t^X \times [y]_t^Y) = \mu(X \times [y]_t^Y) = \mu_Y([y]_t^Y).$$

Therefore,  $f_t$  is an isomorphism between  $\mathcal{X}_t$  and  $\mathcal{Y}_t$ . Hence, we have that  $u_{\text{GW}, \infty}(\mathcal{X}, \mathcal{Y}) \geq \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$  and hence  $u_{\text{GW}, \infty}(\mathcal{X}, \mathcal{Y}) = \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$ .

Finally, we show that the infimum of  $\inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$  is attainable. Let  $\delta := \inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$ . If  $\delta > 0$ , let  $\{t_n\}_{n \in \mathbb{N}}$  be a decreasing sequence converging to  $\delta$  such that  $\mathcal{X}_{t_n} \cong_w \mathcal{Y}_{t_n}$  for all  $t_n$ . Since  $\mathcal{X}_\delta$  and  $\mathcal{Y}_\delta$  are finite spaces, we actually have that  $\mathcal{X}_{t_n} = \mathcal{X}_\delta$  and  $\mathcal{Y}_{t_n} = \mathcal{Y}_\delta$  when  $n$  is large enough. This immediately implies that  $\mathcal{X}_\delta \cong_w \mathcal{Y}_\delta$ . Now, if  $\delta = 0$ , then by Equation (32) we have that  $u_{\text{GW}, \infty}(\mathcal{X}, \mathcal{Y}) = \delta = 0$ . By Theorem 3.10,  $\mathcal{X} \cong_w \mathcal{Y}$ . This is equivalent to  $\mathcal{X}_\delta \cong_w \mathcal{Y}_\delta$ . Therefore, the infimum of  $\inf \{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$  is always attainable.

**B.2.4. Proof of Theorem 3.17.** An important observation for the proof of Theorem 3.17 is that the snowflake transform relates the  $p$ -Wasserstein pseudometric on a pseudo-ultrametric space  $X$  with the 1-Wasserstein pseudometric on the space  $S_p(X)$ ,  $1 \leq p < \infty$ .

**Lemma B.11.** *Given a pseudo-ultrametric space  $(X, u_X)$  and  $p \geq 1$ , we have for any  $\alpha, \beta \in \mathcal{P}(X)$  that*

$$d_{\text{W}, p}^{(X, u_X)}(\alpha, \beta) = \left( d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta) \right)^{\frac{1}{p}}.$$

**Remark B.12.** Since  $S_p \circ u_X$  and  $u_X$  induce the same topology and thus the same Borel sets on  $X$ , we have that  $\mathcal{P}(X) = \mathcal{P}(S_p(X))$  and thus the expression  $d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta)$  in the lemma is well defined.

*Proof of Lemma B.11.* Suppose  $\mu_1, \mu_2 \in \mathcal{C}(\alpha, \beta)$  are optimal for  $d_{\text{W}, p}^X(\alpha, \beta)$  and  $d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta)$ , respectively (see Appendix B.5.1 for the existence of  $\mu_1$  and  $\mu_2$ ). Then,

$$\begin{aligned} \left( d_{\text{W}, p}^{(X, u_X)}(\alpha, \beta) \right)^p &= \int_{X \times X} (u_X(x, y))^p \mu_1(dx \times dy) \\ &= \int_{X \times X} S_p(u_X)(x, y) \mu_1(dx \times dy) \geq d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta), \end{aligned}$$

and

$$\begin{aligned} d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta) &= \int_{X \times X} S_p(u_X)(x, y) \mu_2(dx \times dy) \\ &= \int_{X \times X} (u_X(x, y))^p \mu_2(dx \times dy) \geq \left( d_{\text{W}, p}^{(X, u_X)}(\alpha, \beta) \right)^p. \end{aligned}$$

Therefore,  $d_{\text{W}, p}^{(X, u_X)}(\alpha, \beta) = \left( d_{\text{W}, 1}^{S_p(X)}(\alpha, \beta) \right)^{\frac{1}{p}}$ . □

With Lemma B.11 at our disposal we can prove Theorem 3.17.

*Proof of Theorem 3.17.* Let  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ . Then,

$$\begin{aligned} & \iint_{X \times Y \times X \times Y} (\Lambda_\infty(u_X(x, x'), u_Y(y, y')))^p \mu(dx \times dy) \mu(dx' \times dy') \\ &= \iint_{X \times Y \times X \times Y} \Lambda_\infty(u_X(x, x')^p, u_Y(y, y')^p) \mu(dx \times dy) \mu(dx' \times dy'). \end{aligned}$$

By infimizing over  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  on both sides, we obtain that

$$(u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}))^p = u_{\text{GW},1}(S_p(\mathcal{X}), S_p(\mathcal{Y})).$$

To prove the second part of the claim, let  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$ . By Lemma B.11 we have that

$$(d_{\text{W},p}^{(X \sqcup Y, u)}(\mu_X, \mu_Y))^p = d_{\text{W},1}^{(S_p(X) \sqcup S_p(Y), S_p(u))}(\mu_X, \mu_Y).$$

Finally, infimizing over  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  yields

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})^p = u_{\text{GW},1}^{\text{sturm}}(S_p(\mathcal{X}), S_p(\mathcal{Y})).$$

□

As a direct consequence of Theorem 3.17, we obtain the following relation between  $(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$  and  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  for  $p \in [1, \infty)$ .

**Corollary B.13.** *For each  $p \in [1, \infty)$ , the metric space  $(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$  is isometric to the snowflake transform of  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$ , i.e.,*

$$S_p(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}}) \cong (\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$$

*Proof.* Consider the snowflake transform map  $S_p : \mathcal{U}^w \rightarrow \mathcal{U}^w$  sending  $X \in \mathcal{U}^w$  to  $S_p(X) \in \mathcal{U}^w$ . It is obvious that  $S_p$  is bijective. By Theorem 3.17,  $S_p$  is an isometry from  $S_p(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  to  $(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$ . Therefore,  $S_p(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}}) \cong (\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$ . □

**B.3. Proofs from Section 3.3.** Throughout the following, we demonstrate the open claims from Section 3.3.

**B.3.1. Proof of Theorem 3.18.** First, we focus on the statement for  $p = 1$ , i.e., on showing

$$u_{\text{GW},1}(\mathcal{X}, \mathcal{Y}) \leq 2 u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}). \quad (33)$$

Let  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  be such that  $u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = \int u(x, y) \mu(dx \times dy)$ . The existence of  $u$  and  $\mu$  follows from Proposition B.1

*Claim 1:* For any  $(x, y), (x', y') \in X \times Y$ , we have

$$\Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq \max(u(x, y), u(x', y')) \leq u(x, y) + u(x', y').$$

*Proof.* We only need to show that

$$\Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq \max(u(x, y), u(x', y')).$$

If  $u_X(x, x') = u_Y(y, y')$ , then there is nothing to prove. Otherwise, we assume without loss of generality that  $u_X(x, x') < u_Y(y, y')$ . If  $\max(u(x, y), u(x', y')) < u_Y(y, y')$ , then by the strong triangle inequality we must have  $u(x, y') = u_Y(y, y') = u(x', y)$ . However,  $u(x', y) \leq \max(u_X(x, x'), u(x, y)) < u_Y(y, y')$ , which leads to a contradiction. Therefore,  $\Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq \max(u(x, y), u(x', y'))$ .  $\square$

By Claim 1, we have

$$\begin{aligned} & \iint_{X \times Y \times X \times Y} \Lambda_\infty(u_X(x, x'), u_Y(y, y')) \mu(dx \times dy) \mu(dx' \times dy') \\ & \leq \iint_{X \times Y \times X \times Y} u(x, y) \mu(dx \times dy) \mu(dx' \times dy') \\ & \quad + \iint_{X \times Y \times X \times Y} u(x', y') \mu(dx \times dy) \mu(dx' \times dy') \\ & = \int_{X \times Y} u(x, y) \mu(dx \times dy) + \int_{X \times Y} u(x', y') \mu(dx' \times dy') \leq 2u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}). \end{aligned}$$

Therefore,  $u_{\text{GW},1}(\mathcal{X}, \mathcal{Y}) \leq 2u_{\text{GW},1}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$ .

Applying Theorem 3.17 and Equation (33), yields that for any  $p \in [1, \infty)$

$$u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = (u_{\text{GW},1}(S_p(\mathcal{X}), S_p(\mathcal{Y})))^{\frac{1}{p}} \leq (2u_{\text{GW},1}^{\text{sturm}}(S_p(\mathcal{X}), S_p(\mathcal{Y})))^{\frac{1}{p}} = 2^{\frac{1}{p}} u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}).$$

B.3.2. *Proof of results in Example 3.20.* It follows from [67, Remark 5.17] that

$$d_{\text{GW},p}^{\text{sturm}}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \geq \frac{1}{4} \text{ and } d_{\text{GW},p}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \leq \frac{1}{2} \left( \frac{3}{2n} \right)^{\frac{1}{p}}.$$

Then, by Proposition 3.3, we have that

$$u_{\text{GW},p}^{\text{sturm}}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \geq d_{\text{GW},p}^{\text{sturm}}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \geq \frac{1}{4}.$$

Let  $\mu_n$  denote the uniform probability measure of  $\hat{\Delta}_n(1)$ . Since  $\hat{\Delta}_n(1)$  has the constant interpoint distance 1, it is obvious that for any coupling  $\mu \in \mathcal{C}(\mu_n, \mu_{2n})$ ,

$$\text{dis}_p(\mu) = \text{dis}_p^{\text{ult}}(\mu)$$

This implies that

$$u_{\text{GW},p}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) = 2d_{\text{GW},p}(\hat{\Delta}_n(1), \hat{\Delta}_{2n}(1)) \leq \left( \frac{3}{2n} \right)^{\frac{1}{p}}.$$

B.3.3. *Proof of Theorem 3.21.* First, we prove that  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \geq u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ . Indeed, for any  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ , we have that

$$\begin{aligned} \sup_{(x,y) \in \text{supp}(\mu)} u(x,y) &= \sup_{(x,y),(x',y') \in \text{supp}(\mu)} \max(u(x,y), u(x',y')) \\ &\geq \sup_{(x,y),(x',y') \in \text{supp}(\mu)} \Lambda_{\infty}(u_X(x, x'), u_Y(y, y')) \\ &\geq u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}), \end{aligned}$$

where the first inequality follows from Claim 1 in the proof of Theorem 3.18. Then, by a standard limit argument, we conclude that  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \geq u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ .

Next, we prove that  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq \min\{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$ . Let  $t > 0$  be such that  $\mathcal{X}_t \cong_w \mathcal{Y}_t$  and let  $\varphi : \mathcal{X}_t \rightarrow \mathcal{Y}_t$  denote such an isomorphism. Then, we define a function  $u : X \sqcup Y \times X \sqcup Y \rightarrow \mathbb{R}_{\geq 0}$  as follows:

$$(1) \quad u|_{X \times X} := u_X \text{ and } u|_{Y \times Y} := u_Y;$$

$$(2) \quad \text{for any } (x, y) \in X \times Y, \quad u(x, y) := \begin{cases} u_{Y_t}(\varphi([x]_t^X), [y]_t^Y), & \text{if } \varphi([x]_t^X) \neq [y]_t^Y \\ t, & \text{if } \varphi([x]_t^X) = [y]_t^Y. \end{cases}$$

$$(3) \quad \text{for any } (y, x) \in Y \times X, \quad u(y, x) := u(x, y).$$

Then, it is easy to verify that  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and that  $u$  is actually an ultrametric. Let  $Z := (X \sqcup Y, u)$ . By Lemma 2.8, we have

$$u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq d_{\text{W},\infty}^Z(\mu_X, \mu_Y) = \max_{B \in V(Z) \setminus \{Z\} \text{ and } \mu_X(B) \neq \mu_Y(B)} \text{diam}(B^*).$$

We verify that  $d_{\text{W},\infty}^Z(\mu_X, \mu_Y) \leq t$  in the following. It is obvious that  $Z_t \cong X_t \cong Y_t$ . Write  $X_t = \{[x_i]_t^X\}_{i=1}^n$  and  $Y_t = \{[y_i]_t^Y\}_{i=1}^n$  such that  $[y_i]_t^Y = \varphi([x_i]_t^X)$  for each  $i = 1, \dots, n$ . Then,  $[x_i]_t^Z = [y_i]_t^Z$  and  $Z_t = \{[x_i]_t^Z \mid i = 1, \dots, n\}$ . Since  $\varphi$  is an isomorphism, for any  $i = 1, \dots, n$  we have that  $\mu_X([x_i]_t^X) = \mu_Y([y_i]_t^Y)$  and thus  $\mu_X([x_i]_t^Z) = \mu_Y([y_i]_t^Z) = \mu_Y([x_i]_t^Z)$  when  $\mu_X$  and  $\mu_Y$  are regarded as pushforward measures under the inclusion map  $X \hookrightarrow Z$  and  $Y \hookrightarrow Z$ , respectively. Now for any  $B \in V(Z)$  (cf. Section 2.3), if  $\text{diam}(B) \geq t$ , then  $B$  is the union of certain  $[x_i]_t^Z$ 's in  $Z_t$  and thus  $\mu_X(B) = \mu_Y(B)$ . If  $\text{diam}(B) < t$  and  $\text{diam}(B^*) > t$ , then there exists some  $x_i$  such that  $B = [x_i]_s^Z$  and  $[x_i]_s^Z = [x_i]_t^Z$  where  $s := \text{diam}(B)$ . This implies that  $\mu_X(B) = \mu_Y(B)$ . In consequence, we have that  $d_{\text{W},\infty}^Z(\mu_X, \mu_Y) \leq t$  and thus  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq d_{\text{W},\infty}^{(X \sqcup Y, u)}(\mu_X, \mu_Y) \leq t$ . Therefore,  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq \inf\{t \geq 0 \mid \mathcal{X}_t \cong_w \mathcal{Y}_t\}$ .

Finally, by invoking Theorem 3.13, we conclude that  $u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ .

B.3.4. *Proof of Theorem 3.22.* We prove the result via an explicit construction. By Theorem 3.21, we have  $s = u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y})$ . By Theorem 3.13, there exists an isomorphism  $\varphi : \mathcal{X}_s \rightarrow \mathcal{Y}_s$ . Since  $s > 0$ , by Lemma A.7, both  $\mathcal{X}_s$  and  $\mathcal{Y}_s$  are finite spaces. Let  $X_s = \{[x_1]_s^X, \dots, [x_n]_s^X\}$ ,  $Y_s = \{[y_1]_s^Y, \dots, [y_n]_s^Y\}$  and assume  $[y_i]_s^Y = \varphi([x_i]_s^X)$  for each  $i = 1, \dots, n$ . Let  $A := \{x_1, \dots, x_n\}$  and define  $\phi : A \rightarrow Y$  by sending  $x_i$  to  $y_i$  for each  $i = 1, \dots, n$ . We prove that  $(A, \phi)$  satisfies the conditions in the statement.

Since  $\varphi$  is an isomorphism, for any  $1 \leq i < j \leq n$ ,

$$u_Y(y_i, y_j) = u_{Y_s}([y_i]_s^Y, [y_j]_s^Y) = u_{Y_s}(\varphi([x_i]_s^X), \varphi([x_j]_s^X)) = u_{X_s}([x_i]_s^X, [x_j]_s^X) = u_X(x_i, x_j).$$

This implies that  $\phi : A \rightarrow Y$  is an isometric embedding and thus  $(A, \phi) \in \mathcal{A}$ .

It is obvious that  $(Z_A)_s$  is isometric to both  $X_s$  and  $Y_s$ . In fact,  $[x_i]_s^{Z_A} = [y_i]_s^{Z_A}$  in  $Z_A$  for each  $i = 1, \dots, n$  and  $(Z_A)_s = \{[x_i]_s^{Z_A} \mid i = 1, \dots, n\}$ . Since  $\varphi$  is an isomorphism, for any  $i = 1, \dots, n$  we have that  $\mu_X([x_i]_s^X) = \mu_Y([y_i]_s^Y)$  and thus  $\mu_X([x_i]_s^{Z_A}) = \mu_Y([y_i]_s^{Z_A}) = \mu_Y([x_i]_s^{Z_A})$  when  $\mu_X$  and  $\mu_Y$  are regarded as pushforward measures under the inclusion maps  $X \rightarrow Z_A$  and  $Y \rightarrow Z_A$ , respectively. Now for any  $B \in V(Z_A)$  (cf. Section 2.3), if  $\text{diam}(B) \geq s$ , then  $B$  is the union of certain  $[x_i]_s^{Z_A}$ 's and thus  $\mu_X(B) = \mu_Y(B)$ . If otherwise  $\text{diam}(B) < s$  and  $\text{diam}(B^*) > s$ , then there exists  $x_i$  such that  $B = [x_i]_t^{Z_A}$  and  $[x_i]_t^{Z_A} = [x_i]_s^{Z_A}$  where  $t := \text{diam}(B)$ . This implies that  $\mu_X(B) = \mu_Y(B)$ . By Lemma 2.8, we have  $d_{W,\infty}^{Z_A}(\mu_X, \mu_Y) \leq s$  and thus  $d_{W,\infty}^{Z_A}(\mu_X, \mu_Y) = s$  since  $d_{W,\infty}^{Z_A}(\mu_X, \mu_Y)$  is an upper bound for  $s = u_{\text{GW},\infty}^{\text{sturm}}(\mathcal{X}, \mathcal{Y})$  due to Equation (10).

**B.3.5. Proof of Theorem 3.24.** In this section, we prove Theorem 3.24 by slightly modifying the proof of Proposition 5.3 in [67].

**Lemma B.14.** *Let  $(X, u_X)$  and  $(Y, u_Y)$  be compact ultrametric spaces and let  $S \subseteq X \times Y$  be non-empty. Assume that  $\sup_{(x,y),(x',y') \in S} \Lambda_\infty(u_X(x, x'), u_Y(y, y')) \leq \eta$ . Define  $u_S : X \sqcup Y \times X \sqcup Y \rightarrow \mathbb{R}_{\geq 0}$  as follows:*

- (1)  $u_S|_{X \times X} := u_X$  and  $u_S|_{Y \times Y} := u_Y$ ;
- (2) for any  $(x, y) \in X \times Y$ ,  $u_S(x, y) := \inf_{(x', y') \in S} \max(u_X(x, x'), u_Y(y, y'), \eta)$ .
- (3) for any  $(x, y) \in X \times Y$ ,  $u_S(y, x) := u_S(x, y)$ .

Then,  $u_S \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and  $u_S(x, y) \leq \eta$  for all  $(x, y) \in S$ .

*Proof.* That  $u_S \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  essentially follows by Zarichnyi [105, Lemma 1.1]. It remains to prove the second half of the statement. For  $(x, y) \in S$ , we set  $(x', y') := (x, y)$ . This yields

$$u_S(x, y) \leq \max(u_X(x, x'), u_Y(y, y'), \eta) = \max(0, 0, \eta) = \eta.$$

□

*Proof of Theorem 3.24.* Let  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  be a coupling such that  $\|\Gamma_{X,Y}^\infty\|_{L^p(\mu \otimes \mu)} < \delta^5$ . Set  $\varepsilon := 4v_\delta(X) \leq 4$ .

By Mémoli [67, Claim 10.1], there exist a positive integer  $N \leq [1/\delta]$  and points  $x_1, \dots, x_N$  in  $X$  such that  $\min_{i \neq j} u_X(x_i, x_j) \geq \frac{\varepsilon}{2}$ ,  $\min_i \mu_X(B_\varepsilon^X(x_i)) > \delta$  and  $\mu_X\left(\bigcup_{i=1}^N B_\varepsilon^X(x_i)\right) \geq 1 - \varepsilon$ .

*Claim 1:* For every  $i = 1, \dots, N$  there exists  $y_i \in Y$  such that

$$\mu(B_\varepsilon^X(x_i) \times B_{2(\varepsilon+\delta)}^Y(y_i)) \geq (1 - \delta^2)\mu_X(B_\varepsilon^X(x_i)).$$

*Proof.* Assume the claim is false for some  $i$  and let  $Q_i(y) = B_\varepsilon^X(x_i) \times (Y \setminus B_{2(\varepsilon+\delta)}^Y(y))$ . Then, as  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  it holds

$$\begin{aligned} \mu_X(B_\varepsilon^X(x_i)) &= \mu(B_\varepsilon^X(x_i) \times Y) \\ &= \mu(B_\varepsilon^X(x_i) \times B_{2(\varepsilon+\delta)}^Y(y)) + \mu(B_\varepsilon^X(x_i) \times (Y \setminus B_{2(\varepsilon+\delta)}^Y(y))). \end{aligned}$$

Consequently, we have that  $\mu(Q_i(y)) \geq \delta^2 \mu_X(B_\varepsilon^X(x_i))$ . Further, let

$$\mathcal{Q}_i := \{(x, y, x', y') \in X \times Y \times X \times Y \mid x, x' \in B_\varepsilon^X(x_i) \text{ and } u_Y(y, y') \geq 2(\varepsilon + \delta)\}.$$

Clearly, it holds for  $(x, y, x', y') \in \mathcal{Q}_i$  that

$$\Gamma_{X,Y}^\infty(x, y, x', y') = \Lambda_\infty(u_X(x, x'), u_Y(y, y')) = u_Y(y, y') \geq 2\delta.$$

Further, we have that  $\mu \otimes \mu(\mathcal{Q}_i) \geq \delta^4$ . Indeed, it holds

$$\begin{aligned} \mu \otimes \mu(\mathcal{Q}_i) &= \int_{B_\varepsilon^X(x_i) \times Y} \int_{Q_i(y)} 1 \mu(dx' \times dy') \mu(dx \times dy) \\ &= \int_{B_\varepsilon^X(x_i) \times Y} \mu(Q_i(y)) \mu(dx \times dy) \\ &= \mu_X(B_\varepsilon^X(x_i)) \int_Y \mu(Q_i(y)) \mu_Y(dy) \\ &\geq (\mu_X(B_\varepsilon^X(x_i)))^2 \delta^2 \\ &\geq \delta^4. \end{aligned}$$

However, this yields that

$$\|\Gamma_{X,Y}^\infty\|_{L^p(\mu \otimes \mu)} \geq \|\Gamma_{X,Y}^\infty\|_{L^1(\mu \otimes \mu)} \geq \|\Gamma_{X,Y}^\infty \mathbf{1}_{\mathcal{Q}_i}\|_{L^1(\mu \otimes \mu)} \geq 2\delta \cdot \mu \otimes \mu(\mathcal{Q}_i) \geq 2\delta^5,$$

which contradicts  $\|\Gamma_{X,Y}^\infty\|_{L^p(\mu \otimes \mu)} < \delta^5$ . □

Define for each  $i = 1, \dots, N$

$$S_i := B_\varepsilon^X(x_i) \times B_{2(\varepsilon+\delta)}^Y(y_i).$$

Then, by Claim 1,  $\mu(S_i) \geq \delta(1 - \delta^2)$ , for all  $i = 1, \dots, N$ .

*Claim 2:*  $\Gamma_{X,Y}^\infty(x_i, y_i, x_j, y_j) \leq 6(\varepsilon + \delta)$  for all  $i, j = 1, \dots, N$ .

*Proof.* Assume the claim fails for some  $(i_0, j_0)$ , i.e.,

$$\Lambda_\infty(u_X(x_{i_0}, x_{j_0}), u_Y(y_{i_0}, y_{j_0})) > 6(\varepsilon + \delta) > 0.$$

Then, we have  $\Lambda_\infty(u_X(x_{i_0}, x_{j_0}), u_Y(y_{i_0}, y_{j_0})) = \max(u_X(x_{i_0}, x_{j_0}), u_Y(y_{i_0}, y_{j_0}))$ . We assume without loss of generality that

$$u_X(x_{i_0}, x_{j_0}) = \Lambda_\infty(u_X(x_{i_0}, x_{j_0}), u_Y(y_{i_0}, y_{j_0})) > u_Y(y_{i_0}, y_{j_0}).$$

Consider any  $(x, y) \in S_{i_0}$  and  $(x', y') \in S_{j_0}$ . By the strong triangle inequality and the fact that  $u_X(x_{i_0}, x_{j_0}) > 6(\varepsilon + \delta) > \varepsilon$ , it is easy to verify that  $u_X(x, x') = u_X(x_{i_0}, x_{j_0})$ . Moreover,

$$\begin{aligned} u_Y(y, y') &\leq \max(u_Y(y, y_{i_0}), u_Y(y_{i_0}, y_{j_0}), u_Y(y_{j_0}, y')) \\ &< \max(2(\varepsilon + \delta), u_X(x_{i_0}, x_{j_0}), 2(\varepsilon + \delta)) = u_X(x_{i_0}, x_{j_0}) = u_X(x, x'). \end{aligned}$$

Therefore,

$$\Gamma_{X,Y}^\infty(x, y, x', y') = u_X(x, x') = u_X(x_{i_0}, x_{j_0}) = \Gamma_{X,Y}^\infty(x_{i_0}, y_{i_0}, x_{j_0}, y_{j_0}) > 6(\varepsilon + \delta) > 2\delta.$$

Consequently, we have that

$$\begin{aligned} \|\Gamma_{X,Y}^\infty\|_{L^p(\mu \otimes \mu)} &\geq \|\Gamma_{X,Y}^\infty\|_{L^1(\mu \otimes \mu)} \geq \|\Gamma_{X,Y}^\infty \mathbf{1}_{S_{i_0}} \mathbf{1}_{S_{j_0}}\|_{L^1(\mu \otimes \mu)} \geq 2\delta \mu(S_{i_0}) \mu(S_{j_0}) \\ &> 2\delta (\delta(1 - \delta^2))^2. \end{aligned}$$

However, for  $\delta \leq 1/2$ ,  $2\delta (\delta(1 - \delta^2))^2 \geq 2\delta^5$ . This leads to a contradiction.  $\square$

Consider  $S \subseteq X \times Y$  given by  $S := \{(x_i, y_i) \mid i = 1, \dots, N\}$ . Let  $u_S$  be the ultrametric on  $X \sqcup Y$  given by Lemma B.14. By Claim 2,  $\sup_{(x,y),(x',y') \in S} \Gamma_{X,Y}^\infty(x, y, x', y') \leq 6(\varepsilon + \delta)$ . Then, for all  $i = 1, \dots, N$  we have that  $u_S(x_i, y_i) \leq 6(\varepsilon + \delta)$  and for any  $(x, y) \in X \times Y$  we have that

$$u_S(x, y) \leq \max(\text{diam}(X), \text{diam}(Y), 6(\varepsilon + \delta)) \leq \max(\text{diam}(X), \text{diam}(Y), 27) =: M'.$$

Here in the second inequality we use the assumption that  $\delta < \frac{1}{2}$  and the fact that  $\varepsilon = 4v_\delta(X) \leq 4$ .

*Claim 3:* Fix  $i \in \{1, \dots, N\}$ . Then, for all  $(x, y) \in S_i$ , it holds  $u_S(x, y) \leq 6(\varepsilon + \delta)$ .

*Proof.* Let  $(x, y) \in S_i$ . Then,  $u_X(x, x_i) \leq \varepsilon$  and  $u_Y(y, y_i) \leq 2(\varepsilon + \delta)$ . Then, by the strong triangle inequality for  $u_S$  we obtain

$$\begin{aligned} u_S(x, y) &\leq \max\{u_X(x, x_i), u_Y(y, y_i), u_S(x_i, y_i)\} \\ &\leq \max\{\varepsilon, 2(\varepsilon + \delta), 6(\varepsilon + \delta)\} \leq 6(\varepsilon + \delta). \end{aligned}$$

$\square$

Let  $L := \bigcup_{i=1}^N S_i$ . The next step is to estimate the mass of  $\mu$  in the complement of  $L$ .

*Claim 4:*  $\mu(X \times Y \setminus L) \leq \varepsilon + \delta$ .

*Proof.* For each  $i = 1, \dots, N$ , let  $A_i := B_\varepsilon^X(x_i) \times (Y \setminus B_{2(\varepsilon+\delta)}^Y(y_i))$ . Then,

$$A_i = (B_\varepsilon^X(x_i) \times Y) \setminus (B_\varepsilon^X(x_i) \times B_{2(\varepsilon+\delta)}^Y(y_i)) = (B_\varepsilon^X(x_i) \times Y) \setminus S_i.$$

Hence,

$$\mu(A_i) = \mu(B_\varepsilon^X(x_i) \times Y) - \mu(S_i) = \mu_X(B_\varepsilon^X(x_i)) - \mu(S_i),$$

where the last equality follows from the fact that  $\mu \in \mathcal{M}(\mu_X, \mu_Y)$ . By Claim 1, we have that  $\mu(S_i) \geq \mu_X(B_\varepsilon^X(x_i)) (1 - \delta^2)$ . Consequently, we obtain

$$\mu(A_i) \leq \mu_X(B_\varepsilon^X(x_i)) \delta^2.$$

Notice that

$$X \times Y \setminus L \subseteq \left( X \setminus \bigcup_{i=1}^N B_\varepsilon^X(x_i) \right) \times Y \cup \left( \bigcup_{i=1}^N A_i \right).$$

Hence,

$$\begin{aligned}
\mu(X \times Y \setminus L) &\leq \mu_X \left( X \setminus \bigcup_{i=1}^N B_\varepsilon^X(x_i) \right) + \sum_{i=1}^N \mu(A_i) \\
&\leq 1 - \mu_X \left( \bigcup_{i=1}^N B_\varepsilon^X(x_i) \right) + \sum_{i=1}^N \delta^2 \mu_X(B_\varepsilon^X(x_i)) \\
&\leq \varepsilon + N \cdot \delta^2 \leq \varepsilon + \delta.
\end{aligned}$$

Here, the third inequality follows from the construction of  $x_i$ s in the beginning of this section and from the fact that  $N \leq [1/\delta]$ .  $\square$

Now,

$$\begin{aligned}
\int_{X \times Y} u_S^p(x, y) \mu(dx \times dy) &= \left( \int_L + \int_{X \times Y \setminus L} \right) u_S^p(x, y) \mu(dx \times dy) \\
&\leq (6(\varepsilon + \delta))^p + M'^p \cdot (\varepsilon + \delta).
\end{aligned}$$

Since we have for any  $a, b \geq 0$  and  $p \geq 1$  that  $a^{1/p} + b^{1/p} \geq (a + b)^{1/p}$ , we obtain

$$\begin{aligned}
u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) &\leq (\varepsilon + \delta)^{\frac{1}{p}} \left( 6(\varepsilon + \delta)^{1-\frac{1}{p}} + M' \right) \leq (\varepsilon + \delta)^{\frac{1}{p}} (27 + M') \\
&\leq (4v_\delta(\mathcal{X}) + \delta)^{\frac{1}{p}} \cdot M,
\end{aligned}$$

where we used  $\varepsilon = 4v_\delta(\mathcal{X})$  and  $M := 2 \max(\text{diam}(X), \text{diam}(Y)) + 54 \geq M' + 27$ . Since the roles of  $\mathcal{X}$  and  $\mathcal{Y}$  are symmetric, we have that

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) \leq (4 \min(v_\delta(\mathcal{X}), v_\delta(Y)) + \delta)^{\frac{1}{p}} \cdot M.$$

This concludes the proof.  $\square$

**B.4. Proofs from Section 3.4.** The subsequent section contains the full proofs of the statements in Section 3.4.

**B.4.1. Proof of Theorem 3.26.**

- (1) We first prove that  $(\mathcal{U}^w, u_{\text{GW},p})$  is non-separable for each  $p \in [1, \infty]$ . Recall notations in Example 3.5 and consider the family  $\{\hat{\Delta}_2(a)\}_{a \in [1,2]}$ .

*Claim 1:*  $\forall a \neq b \in [1, 2], u_{\text{GW},p}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) = 2^{-\frac{1}{p}} \Lambda_\infty(a, b) \geq 2^{-\frac{1}{p}}$ , where we let  $2^{-\frac{1}{\infty}} = 1$ .

*Proof of Claim 1.* First note by Theorem 4.1 that

$$u_{\text{GW},p}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) \geq \text{SLB}_p^{\text{ult}}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)).$$

It is easy to verify that  $\mathbf{SLB}_p^{\text{ult}}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) = 2^{-\frac{1}{p}} \Lambda_\infty(a, b)$ . On the other hand, consider the diagonal coupling between  $\mu_a$  and  $\mu_b$ , then for  $p \in [1, \infty)$

$$u_{\text{GW},p}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) \leq \left(2 \cdot \Lambda_\infty(a, b)^p \cdot \frac{1}{2} \cdot \frac{1}{2}\right)^{\frac{1}{p}} = 2^{-\frac{1}{p}} \Lambda_\infty(a, b),$$

and for  $p = \infty$

$$u_{\text{GW},\infty}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) \leq \Lambda_\infty(a, b).$$

Therefore,

$$u_{\text{GW},p}(\hat{\Delta}_2(a), \hat{\Delta}_2(b)) = 2^{-\frac{1}{p}} \Lambda_\infty(a, b).$$

□

By Claim 1, we have that  $\{\hat{\Delta}_2(a)\}_{a \in [1,2]}$  is an uncountable subset of  $\mathcal{U}^w$  with pairwise distance greater than  $2^{-\frac{1}{p}}$ , which implies that  $(\mathcal{U}^w, u_{\text{GW},p})$  is non-separable.

Now for  $p \in [1, \infty)$ , we show that  $u_{\text{GW},p}$  is not complete. Consider the family  $\{\Delta_{2^n}(1)\}_{n \in \mathbb{N}}$  of  $2^n$ -point spaces with unitary interpoint distances. Endow each space  $\Delta_{2^n}(1)$  with the uniform measure  $\mu_n$  and denote the corresponding ultrametric measure space by  $\hat{\Delta}_{2^n}(1)$ . It is proven in [93, Example 2.2] that  $\{\hat{\Delta}_{2^n}(1)\}_{n \in \mathbb{N}}$  is a Cauchy sequence with respect to  $d_{\text{GW},p}$  without a compact metric measure space as limit. It is not hard to check that

$$u_{\text{GW},p}(\hat{\Delta}_{2^m}(1), \hat{\Delta}_{2^n}(1)) = 2d_{\text{GW},p}(\hat{\Delta}_{2^m}(1), \hat{\Delta}_{2^n}(1)), \quad \forall n, m \in \mathbb{N}.$$

Therefore,  $\{\hat{\Delta}_{2^n}(1)\}_{n \in \mathbb{N}}$  is a Cauchy sequence with respect to  $u_{\text{GW},p}$  without limit in  $\mathcal{U}^w$ . This implies that  $(\mathcal{U}^w, u_{\text{GW},p})$  is not complete.

- (2) By Theorem 3.18 and (1), we have that  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is not separable. As for completeness, consider the subset  $X := \{1 - \frac{1}{n}\}_{n \in \mathbb{N}} \subseteq (\mathbb{R}_{\geq 0}, \Lambda_\infty)$ . By Lemma A.2,  $X$  is not a compact ultrametric space. Let  $\mu_0 \in \mathcal{P}(X)$  be a probability defined as follows:

$$\mu_0\left(\left\{1 - \frac{1}{n}\right\}\right) := 2^{-n}, \quad \forall n \in \mathbb{N}.$$

For each  $N \in \mathbb{N}$ , let  $X_N := \{1 - \frac{1}{n} \mid n = 1, \dots, N\}$ . Since each  $X_N$  is finite,  $(X_N, \Lambda_\infty)$  is a compact ultrametric space. Let  $\mu_N \in \mathcal{P}(X_N)$  be a probability defined as follows:

$$\mu_N\left(\left\{1 - \frac{1}{n}\right\}\right) := \begin{cases} 2^{-n}, & 1 \leq n < N \\ 2^{-N+1}, & n = N \end{cases}.$$

Then, it is easy to verify (e.g. via Theorem 3.7) that  $\{(X_N, \Lambda_\infty, \mu_N)\}_{N \in \mathbb{N}}$  is a  $u_{\text{GW},p}^{\text{sturm}}$  Cauchy sequence with  $(X, \Lambda_\infty, \mu_0)$  being the limit. Since the set  $X$  is not compact,  $(X, \Lambda_\infty, \mu_0) \notin \mathcal{U}^w$  and thus  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is not complete.

- (3) That  $(\mathcal{U}^w, u_{\text{GW},\infty})$  is non-separable is already proved in (1). Given a Cauchy sequence  $\{\mathcal{X}_n = (X_n, u_n, \mu_n)\}_{n \in \mathbb{N}}$  with respect to  $u_{\text{GW},\infty}$ , we have that the underlying ultrametric spaces  $\{X_n\}_{n \in \mathbb{N}}$  form a Cauchy sequence with respect to  $u_{\text{GH}}$  due to Corollary

3.15. Since  $(\mathcal{U}, u_{\text{GH}})$  is complete (see [105, Proposition 2.1]), there exists a compact ultrametric space  $(X, u_X)$  such that

$$\lim_{n \rightarrow \infty} u_{\text{GH}}(X_n, X) = 0.$$

For each  $n \in \mathbb{N}$ , let  $\delta_n := u_{\text{GH}}(X_n, X)$ . By Theorem 2.5, we have that  $(X_n)_{\delta_n} \cong X_{\delta_n}$ . Denote by  $\hat{\mu}_n \in \mathcal{P}(X_{\delta_n})$  the pushforward of  $(\mu_n)_{\delta_n}$  under the isometry. Furthermore, we have by Lemma A.7 that  $X_{\delta_n}$  is finite and we let  $X_{\delta_n} = \{[x_1]_{\delta_n}, \dots, [x_k]_{\delta_n}\}$  for  $x_1, \dots, x_k \in X$ . Based on this, we define

$$\nu_n := \sum_{i=1}^k \hat{\mu}_n([x_i]_{\delta_n}) \cdot \delta_{x_i} \in \mathcal{P}(X),$$

where  $\delta_{x_i}$  is the Dirac measure at  $x_i$ . Since  $X$  is compact,  $\mathcal{P}(X)$  is weakly compact. Therefore, the sequence  $\{\nu_n\}_{n \in \mathbb{N}}$  has a cluster point  $\nu \in \mathcal{P}(X)$ .

Now we show that  $\mathcal{X} := (X, u_X, \nu)$  is a  $u_{\text{GW}, \infty}$  cluster point of  $\{\mathcal{X}_n\}_{n \in \mathbb{N}}$  and thus the limit of  $\{\mathcal{X}_n\}_{n \in \mathbb{N}}$  since  $\{\mathcal{X}_n\}_{n \in \mathbb{N}}$  is a Cauchy sequence. Without loss of generality, we assume that  $\{\nu_n\}_{n \in \mathbb{N}}$  weakly converges to  $\nu$ . Fix any  $\varepsilon > 0$ , we need to show that  $u_{\text{GW}, \infty}(\mathcal{X}, \mathcal{X}_n) \leq \varepsilon$  when  $n$  is large enough. For any fixed  $x_* \in X$ ,  $[x_*]_\varepsilon$  is both an open and closed ball in  $X$ . Therefore,  $\nu([x_*]_\varepsilon) = \lim_{n \rightarrow \infty} \nu_n([x_*]_\varepsilon)$  (see e.g. Billingsley [7, Thm. 2.1]). Since  $\delta_n \rightarrow 0$  as  $n \rightarrow \infty$ , there exists  $N_1 > 0$  such that for any  $n > N_1$ ,  $\delta_n < \varepsilon$ . We specify an isometry  $\varphi_n : (X_n)_{\delta_n} \rightarrow X_{\delta_n}$  that gives rise to the construction of  $\nu_n$ . Then, we let  $\psi_n : (X_n)_\varepsilon \rightarrow X_\varepsilon$  be the isometry such that the following diagram commutes:

$$\begin{array}{ccc} (X_n)_{\delta_n} & \xrightarrow{\varphi_n} & X_{\delta_n} \\ \varepsilon\text{-quotient} \downarrow & & \downarrow \varepsilon\text{-quotient} \\ (X_n)_\varepsilon & \xrightarrow{\psi_n} & X_\varepsilon \end{array}$$

Assume that  $[x_*]_\varepsilon^X = \bigcup_{i=1}^l [x_i]_{\delta_n}^X$ . Let  $x_*^n \in X_n$  be such that  $\psi_n([x_*^n]_\varepsilon^{X_n}) = [x_*]_\varepsilon^X$  and let  $x_1^n, \dots, x_l^n \in X_n$  be such that  $\varphi_n([x_i^n]_{\delta_n}^{X_n}) = [x_i]_{\delta_n}^X$  for each  $i = 1, \dots, l$ . Then,  $[x_*]_\varepsilon^{X_n} = \bigcup_{i=1}^l [x_i^n]_{\delta_n}^{X_n}$ . Therefore,

$$\nu_n([x_*]_\varepsilon^X) = \sum_{i=1}^l \nu_n([x_i]_{\delta_n}^X) = \sum_{i=1}^l \hat{\mu}_n([x_i]_{\delta_n}^X) = \sum_{i=1}^l \mu_n([x_i^n]_{\delta_n}^{X_n}) = \mu_n([x_*^n]_\varepsilon^{X_n}).$$

Since  $\mathcal{X}_n$  is a Cauchy sequence, there exists  $N_2 > 0$  such that  $u_{\text{GW}, \infty}(\mathcal{X}_n, \mathcal{X}_m) < \varepsilon$  when  $n, m > N_2$ . Then, by Theorem 3.13,  $(\mathcal{X}_n)_\varepsilon \cong_w (\mathcal{X}_m)_\varepsilon$  for all  $n, m > N_2$ . By Lemma A.7,  $(X_n)_\varepsilon$  is finite, then  $(X_n)_\varepsilon$  has cardinality independent of  $n$  when  $n > N_2$ . For all  $n > N_2$ , we define the finite set  $A_n := \{\mu_n([x^n]_\varepsilon^{X_n}) \mid x^n \in X_n\}$ .  $A_n$  is independent of  $n$  since  $(\mathcal{X}_n)_\varepsilon \cong_w (\mathcal{X}_m)_\varepsilon$  for all  $n, m > N_2$ . This implies that  $\mu_n([x_*^n]_\varepsilon^{X_n})$  only takes value in a finite set  $A_n$ . Combining with the fact that  $\lim_{n \rightarrow \infty} \mu_n([x_*^n]_\varepsilon^{X_n}) = \lim_{n \rightarrow \infty} \nu_n([x_*]_\varepsilon^X) = \nu([x_*]_\varepsilon^X)$  exists, there exists  $N_3 > 0$  such that when  $n > N_3$ ,  $\mu_n([x_*^n]_\varepsilon^{X_n}) \equiv C$  for some constant  $C$ . This implies that

$$\nu([x_*]_\varepsilon^X) = \mu_n([x_*^n]_\varepsilon^{X_n}), \quad \text{when } n > \max(N_1, N_2, N_3).$$

Since  $X_\varepsilon$  is finite, there exists a common  $N > 0$  such that for all  $n > N$  and  $\forall [x_*]_\varepsilon \in X_\varepsilon$  we have

$$\nu([x_*]_\varepsilon^X) = \mu_n([x_*]_\varepsilon^{X_n}),$$

where  $[x_*]_\varepsilon^{X_n} = \psi_n^{-1}([x_*]_\varepsilon^X) \in (X_n)_\varepsilon$ . This indicates that  $\nu_\varepsilon = (\psi_n)_\#(\mu_n)_\varepsilon$  when  $n > N$ . Therefore,  $\mathcal{X}_\varepsilon \cong_w (\mathcal{X}_n)_\varepsilon$  and thus  $u_{\text{GW},\infty}(\mathcal{X}, \mathcal{X}_n) \leq \varepsilon$ .

**B.4.2. Proof of Theorem 3.27.** Next, we will demonstrate Theorem 3.27. However, before we come to this we recall some facts about  $p$ -metric and  $p$ -geodesic spaces.

**Lemma B.15** (Mémoli et al. [70, Proposition 7.10]). *Given  $p \in [1, \infty)$ , if  $X$  is a  $p$ -metric space, then  $X$  is not  $q$ -geodesic for all  $1 \leq q < p$ .*

**Lemma B.16** (Mémoli et al. [70, Theorem 7.7]). *Let  $X$  be a geodesic metric space. Then, for any  $p \geq 1$ ,  $S_{\frac{1}{p}}(X)$  is  $p$ -geodesic, where  $S_\alpha$  denotes the snowflake transform for  $\alpha > 0$  (cf. Section 3.3).*

For  $p = 1$ , the proof is based on the following property of the 1-Wasserstein space.

**Lemma B.17** (Bottou et al. [10, Theorem 5.1]). *Let  $X$  be a compact metric space. Then, the space  $W_1(X) := (\mathcal{P}(X), d_{W,1}^X)$  is a geodesic space.*

Based on the above results and Corollary B.2, the proof of Theorem 3.27 is straightforward.

*Proof of Theorem 3.27.* Let  $\mathcal{X}$  and  $\mathcal{Y}$  be two compact ultrametric measure spaces. First, we consider the case  $p = 1$ . By Corollary B.2, there exist a compact ultrametric space  $Z$  and isometric embeddings  $\phi : X \hookrightarrow Z$  and  $\psi : Y \hookrightarrow Z$  such that

$$u_{\text{GW},p}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}) = d_{W,p}^Z(\phi_\# \mu_X, \psi_\# \mu_Y).$$

The space  $W_1(Z)$  is geodesic (cf. Lemma B.17). Therefore, there exists a Wasserstein geodesic  $\tilde{\gamma} : [0, 1] \rightarrow W_1(Z)$  connecting  $\phi_\# \mu_X$  and  $\psi_\# \mu_Y$ . This induces a curve  $\gamma : [0, 1] \rightarrow \mathcal{U}^w$  where for each  $t \in [0, 1]$ ,  $\gamma(t) := (\text{supp}(\tilde{\gamma}(t)), u|_{\text{supp}(\tilde{\gamma}(t)) \times \text{supp}(\tilde{\gamma}(t))}, \tilde{\gamma}(t))$ . Note that  $\gamma(0) \cong_w \mathcal{X}$  and  $\gamma(1) \cong_w \mathcal{Y}$  and hence we simply replace  $\gamma(0)$  and  $\gamma(1)$  with  $\mathcal{X}$  and  $\mathcal{Y}$ , respectively. Now, for each  $s, t \in [0, 1]$ , we have that

$$d_{\text{GW},1}^{\text{sturm}}(\gamma(s), \gamma(t)) \leq d_{W,1}^Z(\tilde{\gamma}(s), \tilde{\gamma}(t)) = |s - t| d_{W,1}^Z(\tilde{\gamma}(0), \tilde{\gamma}(1)) = |s - t| d_{\text{GW},1}^{\text{sturm}}(\mathcal{X}, \mathcal{Y}).$$

Therefore,  $\gamma$  is a geodesic connecting  $\mathcal{X}$  and  $\mathcal{Y}$  and thus  $(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$  is geodesic.

Next, we come to the case  $p > 1$ . By Corollary B.13,  $S_p(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}}) \cong (\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}})$ . This implies that  $S_{\frac{1}{p}}(\mathcal{U}^w, u_{\text{GW},1}^{\text{sturm}}) \cong (\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$ . Hence, by Lemma B.16, we have that  $(\mathcal{U}^w, u_{\text{GW},p}^{\text{sturm}})$  is  $p$ -geodesic.  $\square$

**B.5. Technical issues from Section 3.** In the following, we address various technical issues from Section 3.

**B.5.1. The Wasserstein pseudometric.** Given a set  $X$ , a pseudometric is a symmetric function  $d_X : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying the triangle inequality and  $d_X(x, x) = 0$  for all  $x \in X$ . Note that if moreover  $d_X(x, y) = 0$  implies  $x = y$ , then  $d_X$  is a metric. There is a canonical identification on pseudometric spaces  $(X, d_X)$ :  $x \sim x'$  if  $d_X(x, x') = 0$ . Then,  $\sim$  is in fact an equivalence relation and we define the quotient space  $\tilde{X} = X / \sim$ . Define a function  $\tilde{d}_X : \tilde{X} \times \tilde{X} \rightarrow \mathbb{R}_{\geq 0}$  as follows:

$$\tilde{d}_X([x], [x']) := \begin{cases} d_X(x, x') & \text{if } d_X(x, x') \neq 0 \\ 0 & \text{otherwise} \end{cases}.$$

$\tilde{d}_X$  turns out to be a metric on  $\tilde{X}$ . In the following, the metric space  $(\tilde{X}, \tilde{d}_X)$  is referred to as the *metric space induced by the pseudometric space*  $(X, d_X)$ . Note that  $\tilde{d}_X$  preserves the induced topology (see e.g. [44]) and thus the quotient map  $\Psi : X \rightarrow \tilde{X}$  is continuous.

Analogously to the Wasserstein distance, which is defined for probability measures on metric spaces, we define the *Wasserstein pseudometric* for measures on compact pseudometric spaces as done in [94]. Let  $\alpha, \beta \in \mathcal{P}(X)$ . Then, we define for  $p \in [1, \infty)$  the Wasserstein pseudometric of order  $p$  as

$$d_{W,p}^{(X, d_X)}(\alpha, \beta) := \left( \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \int_{X \times X} d_X^p(x, y) \mu(dx \times dy) \right)^{\frac{1}{p}} \quad (34)$$

and for  $p = \infty$  as

$$d_{W,\infty}^{(X, d_X)}(\alpha, \beta) := \inf_{\mu \in \mathcal{C}(\alpha, \beta)} \sup_{(x, y) \in \text{supp}(\mu)} d_X(x, y). \quad (35)$$

It is easy to see that the Wasserstein pseudometric is closely related to the Wasserstein distance on the induced metric space. More precisely, one can show the following.

**Lemma B.18.** *Let  $(X, d_X)$  denote a compact pseudometric space, let  $\alpha, \beta \in \mathcal{P}(X)$ . Then, it follows for  $p \in [1, \infty]$  that*

$$d_{W,p}^{(X, d_X)}(\alpha, \beta) = d_{W,p}^{(\tilde{X}, \tilde{d}_X)}(\Psi_{\#}\alpha, \Psi_{\#}\beta) \quad (36)$$

*and in particular that the infimum in Equation (34) (resp. in Equation (35) if  $p = \infty$ ) is attained for some  $\mu \in \mathcal{C}(\alpha, \beta)$ .*

*Proof.* In the course of this proof we focus on the case  $p < \infty$  and remark that the case  $p = \infty$  follows by similar arguments. The quotient map allows us to define the map  $\theta : \mathcal{C}(\alpha, \beta) \rightarrow \mathcal{C}(\Psi_{\#}\alpha, \Psi_{\#}\beta)$  via  $\mu \mapsto (\Psi \times \Psi)_{\#}\mu$ . It is easy to see that  $\theta$  is well defined and surjective. Furthermore, it holds by construction that

$$\int_{X \times X} d_X^p(x, y) \mu(dx \times dy) = \int_{\tilde{X} \times \tilde{X}} \tilde{d}_X^p(x, y) \theta(\mu)(dx \times dy)$$

for all  $\mu \in \mathcal{C}(\alpha, \beta)$ . Hence, Equation (36) follows.

We come to the second part of the claim. By [100, Sec.4] there exists an optimal coupling  $\tilde{\mu}^* \in \mathcal{C}(\Psi_{\#}\alpha, \Psi_{\#}\beta)$  such that

$$d_{W,p}^{(\tilde{X}, \tilde{d}_X)}(\Psi_{\#}\alpha, \Psi_{\#}\beta) = \left( \int_{\tilde{X} \times \tilde{X}} \tilde{d}_X^p(x, y) \tilde{\mu}^*(dx \times dy) \right)^{\frac{1}{p}}.$$

In consequence, we find using our previous results that for any  $\mu^* \in \theta^{-1}(\tilde{\mu}^*)$  it holds

$$\begin{aligned} d_{W,p}^{(\tilde{X}, \tilde{d}_X)}(\Psi_{\#}\alpha, \Psi_{\#}\beta) &= \left( \int_{\tilde{X} \times \tilde{X}} \tilde{d}_X^p(x, y) \tilde{\mu}^*(dx \times dy) \right)^{\frac{1}{p}} \\ &= \left( \int_{X \times X} d_X^p(x, y) \mu^*(dx \times dy) \right)^{\frac{1}{p}} = d_{W,p}^{(X, d_X)}(\alpha, \beta). \end{aligned}$$

This yields the claim.  $\square$

**B.5.2. Regularity of the cost functionals of  $u_{GW,p}$  and  $u_{GW,p}^{\text{sturm}}$ .** In the remainder of this section, we collect various technical results required to demonstrate the existence of optimizers in the definitions of  $u_{GW,p}^{\text{sturm}}$  (see Equation (10)) and  $u_{GW,p}$  (see Equation (13)).

**Lemma B.19.** *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be compact ultrametric measure spaces. Then,  $\mu \in \mathcal{C}(\mu_X, \mu_Y) \subseteq \mathcal{P}(X \times Y, \max(u_X, u_Y))$  is compact with respect to weak convergence.*

*Proof.* The proof follows directly from Chowdhury and Mémoli [23, Lemma 2.2].  $\square$

**Lemma B.20.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$ . Let  $D_1 \subseteq \mathcal{D}^{\text{ult}}(u_X, u_Y)$  be a non-empty subset satisfying the following: there exist  $(x_0, y_0) \in X \times Y$  and  $C > 0$  such that  $u(x_0, y_0) \leq C$  for all  $u \in D_1$ . Then,  $D_1$  is pre-compact with respect to uniform convergence.*

*Proof.* Let  $\{u_n\}_{n \in \mathbb{N}} \subseteq D_1$  be a sequence. Note that  $X \times Y \subseteq X \sqcup Y \times X \sqcup Y$ . Let  $v_n := u_n|_{X \times Y}$ . For any  $n \in \mathbb{N}$  and any  $(x, y), (x', y') \in X \times Y$ , we have that

$$|u_n(x, y) - u_n(x', y')| \leq u_X(x, x') + u_Y(y, y') \leq 2 \max(u_X, u_Y)((x, y), (x', y')).$$

This means that  $\{v_n\}_{n \in \mathbb{N}}$  is equicontinuous with respect to the ultrametric  $\max\{u_X, u_Y\}$  on  $X \times Y$ . Now, since  $u_n(x_0, y_0) \leq C$ , we have that for any  $(x, y) \in X \times Y$ ,

$$u_n(x, y) \leq 2 \max(u_X, u_Y)((x, y), (x_0, y_0)) + u_n(x_0, y_0) \leq 2 \max(\text{diam}(X), \text{diam}(Y)) + C.$$

Consequently,  $\{v_n\}_{n \in \mathbb{N}}$  is uniformly bounded. By the Arzela-Ascoli theorem ([52, Theorem 7 on page 61]), we have that each subsequence of  $\{v_n\}_{n \in \mathbb{N}}$  has a uniformly convergent subsequence. Hence, we can assume without loss of generality that the sequence  $\{v_n\}_{n \in \mathbb{N}}$  converges to  $v : X \times Y \rightarrow \mathbb{R}_{\geq 0}$ .

Now, we define  $u : X \sqcup Y \times X \sqcup Y \rightarrow \mathbb{R}_{\geq 0}$  as follows:

- (1)  $u|_{X \times X} := u_X$  and  $u|_{Y \times Y} := u_Y$ ;
- (2)  $u|_{X \times Y} := v$ ;
- (3) for  $(y, x) \in Y \times X$ , we let  $u(y, x) := u(x, y)$ .

It is easy to verify that  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  and that  $u$  is a cluster point of the sequence  $\{u_n\}_{n \in \mathbb{N}}$ . Therefore,  $D_1$  is pre-compact.  $\square$

**Lemma B.21.** *Let  $\mathcal{X} = (X, u_X, \mu_X)$  and  $\mathcal{Y} = (Y, u_Y, \mu_Y)$  be compact ultrametric measure spaces. Let  $\{\mu_n\}_{n \in \mathbb{N}} \subseteq \mathcal{C}(\mu_X, \mu_Y)$  be a sequence weakly converging to  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$ . Let*

$\{u_n\}_{n \in \mathbb{N}} \subseteq \mathcal{D}^{\text{ult}}(u_X, u_Y)$ . Suppose that there exist a non-decreasing sequence  $\{p_n\}_{n \in \mathbb{N}} \subseteq [1, \infty)$  and  $C > 0$  such that

$$\left( \int_{X \times Y} (u_n(x, y))^{p_n} \mu_n(dx \times dy) \right)^{\frac{1}{p_n}} \leq C$$

for all  $n \in \mathbb{N}$ . Then,  $\{u_n\}_{n \in \mathbb{N}}$  uniformly converges to some  $u \in \mathcal{D}^{\text{ult}}(u_X, u_Y)$  (up to taking a subsequence).

*Proof.* The following argument adapts the proof of Lemma 3.3 in [92] to the current setting. For any  $(x_0, y_0) \in \text{supp}(\mu)$ , there exist  $\varepsilon, \delta > 0$  and  $N \in \mathbb{N}$  such that for all  $n \geq N$

$$\begin{aligned} C &\geq \left( \int_{X \times Y} (u_n(x, y))^{p_n} \mu_n(dx \times dy) \right)^{\frac{1}{p_n}} \geq \int_{X \times Y} u_n(x, y) \mu_n(dx \times dy) \\ &\geq \int_{B_\varepsilon^X(x_0) \times B_\varepsilon^Y(y_0)} u_n(x, y) \mu_n(dx \times dy) \geq \int_{B_\varepsilon^X(x_0) \times B_\varepsilon^Y(y_0)} (u_n(x_0, y_0) - 2\varepsilon) \mu_n(dx \times dy) \\ &\geq (u_n(x_0, y_0) - 2\varepsilon) (\mu(B_\varepsilon^X(x_0) \times B_\varepsilon^Y(y_0)) - \delta). \end{aligned}$$

Therefore,  $\{u_n(x_0, y_0)\}_{n \geq N}$  is uniformly bounded. By Lemma B.20, we have that  $\{u_n\}_{n \in \mathbb{N}}$  has a uniformly convergent subsequence.  $\square$

**Lemma B.22.** *Let  $X, Y$  be ultrametric spaces, then  $\Lambda_\infty(u_X, u_Y) : X \times Y \times X \times Y \rightarrow \mathbb{R}_{\geq 0}$  is continuous with respect to the product topology (induced by  $\max(u_X, u_Y, u_X, u_Y)$ ).*

*Proof.* Fix  $(x, y, x', y') \in X \times Y \times X \times Y$  and  $\varepsilon > 0$ . Choose  $0 < \delta < \varepsilon$  such that  $\delta < u_X(x, x')$  if  $x \neq x'$  and  $\delta < u_Y(y, y')$  if  $y \neq y'$ . Then, consider any point  $(x_1, y_1, x'_1, y'_1) \in X \times Y \times X \times Y$  such that  $u_X(x, x_1), u_Y(y, y_1), u_X(x', x'_1), u_Y(y', y'_1) \leq \delta$ . For  $u_X(x_1, x'_1)$ , we have the following two situations:

- (1)  $x = x'$ :  $u_X(x_1, x'_1) \leq \max(u_X(x_1, x), u_X(x, x'_1)) \leq \delta < \varepsilon$ ;
- (2)  $x \neq x'$ :  $u_X(x_1, x'_1) \leq \max(u_X(x_1, x), u_X(x, x'), u_X(x', x'_1)) = u_X(x, x')$ . Similarly,  $u_X(x, x') \leq u_X(x_1, x'_1)$  and thus  $u_X(x, x') = u_X(x_1, x'_1)$ .

Similar result holds for  $u_Y(y_1, y'_1)$ . This leads to four cases for  $\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1))$ :

- (1)  $x = x', y = y'$ : In this case we have  $u_X(x_1, x'_1), u_Y(y_1, y'_1) < \varepsilon$ . Then,
$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| = \Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) \leq \varepsilon;$$
- (2)  $x = x', y \neq y'$ : Now  $u_X(x_1, x'_1) < \varepsilon$  and  $u_Y(y_1, y'_1) = u_Y(y, y')$ . If  $u_Y(y, y') \geq \varepsilon > u_X(x_1, x'_1)$ , then
$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| = |u_Y(y, y') - u_Y(y, y')| = 0.$$
Otherwise  $u_Y(y, y') < \varepsilon$ , which implies that  $\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) \leq \varepsilon$  and  $\Lambda_\infty(u_X(x, x'), u_Y(y, y')) = u_Y(y, y') \leq \varepsilon$ . Therefore,
$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| \leq \varepsilon;$$
- (3)  $x \neq x', y = y'$ : Similar with (2) we have

$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| \leq \varepsilon;$$

(4)  $x \neq x', y \neq y'$ : Now  $u_X(x_1, x'_1) = u_X(x, x')$  and  $u_Y(y_1, y'_1) = u_Y(y, y')$ . Therefore,

$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| = 0.$$

In conclusion, whenever  $u_X(x, x_1), u_Y(y, y_1), u_X(x', x'_1), u_Y(y', y'_1) \leq \delta$  we have that

$$|\Lambda_\infty(u_X(x_1, x'_1), u_Y(y_1, y'_1)) - \Lambda_\infty(u_X(x, x'), u_Y(y, y'))| \leq \varepsilon.$$

Therefore,  $\Lambda_\infty(u_X, u_Y)$  is continuous with respect to the metric  $\max(u_X, u_Y, u_X, u_Y)$ .  $\square$

**B.5.3.  $u_{\text{GW},p}$  and the one point space.** It is possible to explicitly write down  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , in some simple settings. In the following, we derive an explicit formulation of  $u_{\text{GW},p}$ ,  $1 \leq p \leq \infty$ , between an arbitrary ultrametric measure space  $\mathcal{X}$  and the one point ultrametric measure space  $*$ . For this purpose, we need to introduce some notation. Let  $\mathcal{X} = (\mathcal{X}, d_X, \mu_X)$  be a ultrametric measure space. Let its  $p$ -diameter (see e.g., [67]) for  $1 \leq p < \infty$  be defined as

$$\text{diam}_p(\mathcal{X}) := \left( \iint_{X \times X} (d_X(x, x'))^p \mu_X(dx) \mu_X(dx') \right)^{1/p}$$

and for  $p = \infty$  as

$$\text{diam}_\infty(\mathcal{X}) := \sup_{(x, x') \in \text{supp}(\mu_X)} d_X(x, x').$$

Then, one can show the subsequent proposition.

**Proposition B.23.** *Let  $* \in \mathcal{U}^w$  be the one-point space. Then, it holds for any  $1 \leq p \leq \infty$  that*

$$u_{\text{GW},p}(\mathcal{X}, *) = \text{diam}_p(\mathcal{X}).$$

*Proof.* Denote by  $\mu$  the unique coupling  $\mu_X \otimes \delta_*$  between  $\mu_X$  and  $\delta_*$ . Then, for any  $p < \infty$  we have

$$\begin{aligned} u_{\text{GW},p}(\mathcal{X}, *) &= \left( \iint_{X \times * \times X \times *} (\Lambda_\infty(u_X(x, x'), u_*(y, y')))^p \mu(dx \times dy) \mu(dx' \times dy') \right)^{1/p} \\ &= \left( \iint_{X \times X} (u_X(x, x'))^p \mu_X(dx) \mu_X(dx') \right)^{1/p} = \text{diam}_p(\mathcal{X}). \end{aligned}$$

The case  $p = \infty$  follows by analogous arguments.  $\square$

## APPENDIX C. MISSING DETAILS FROM SECTION 4

**C.1. Proofs from Section 4.** In the following, we state the full proofs of the results from Section 4.

C.1.1. *Proof of Theorem 4.1.* We start by proving the first statement. To this end, we observe that for any point  $x$  in an ultrametric space  $X$ , there always exists a point  $x' \in X$  such that  $u_X(x, x') = \text{diam}(X)$  (see [30]). By assumption  $\mu_X$  is fully supported on  $X$ . Hence,  $s_{X,\infty} \equiv \text{diam}(X)$  is a constant function. Therefore,

$$\Lambda_\infty(s_{X,\infty}(x), s_{Y,\infty}(y)) \equiv \Lambda_\infty(\text{diam}(X), \text{diam}(Y)), \quad \forall x \in X, y \in Y.$$

This implies that  $\mathbf{FLB}_\infty^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \Lambda_\infty(\text{diam}(X), \text{diam}(Y))$ . By Corollary 5.8 of Mémoli et al. [70] and Corollary 3.15, we have that

$$u_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) \geq u_{\text{GH}}(X, Y) \geq \Lambda_\infty(\text{diam}(X), \text{diam}(Y)) = \mathbf{FLB}_\infty^{\text{ult}}(\mathcal{X}, \mathcal{Y}).$$

It remains to prove the second statement. The proof for  $d_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{TLB}_p(\mathcal{X}, \mathcal{Y})$  in [67, Sec. 6] can be used essentially without any change for showing  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y})$ . Hence, it only remains to show that  $\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y})$ , i.e., the claim follows once we have established Proposition C.1.

**Proposition C.1.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}^w$  and let  $p \in [1, \infty]$ . Then,*

$$\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) \geq \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}).$$

In order to prove Proposition C.1, we need the following technical lemma.

**Lemma C.2.** *Let  $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{U}^w$ . Then,  $\text{spec}(X) := \{u_X(x, x') \mid x, x' \in X\}$  is a compact subset of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ .*

*Proof.* By Lemma A.7, we have that for each  $t > 0$ ,  $X_t$  is a finite set. Let  $\{t_n\}_{n=1}^\infty$  be a positive sequence decreasing to 0. Then, it is easy to see that

$$\text{spec}(X) = \bigcup_{n=1}^{\infty} \text{spec}(X_{t_n}).$$

Since each  $\text{spec}(X_{t_n})$  is a finite set,  $\text{spec}(X)$  is a countable set.

Now, pick any  $0 \neq t \in \text{spec}(X)$ . Suppose  $t$  is a cluster point in  $\text{spec}(X)$ . Then, there exists infinitely many  $s \in \text{spec}(X)$  greater than  $\frac{t}{2}$ . However, this will result in  $X_{\frac{t}{2}}$  being an infinite set, which contradicts the fact that  $X_{\frac{t}{2}}$  is finite. Therefore, 0 is the only possible cluster point of  $\text{spec}(X)$ . By Lemma A.2, we have that  $\text{spec}(X)$  is compact.  $\square$

With the above auxiliary result available, we can demonstrate Proposition C.1 and hence finish the proof of Theorem 4.1.

*Proof of Proposition C.1.* We first prove the case when  $p < \infty$ . Let  $dh_{\mathcal{X}}(x) := u_X(x, \cdot)_{\#} \mu_X$  and let  $dh_{\mathcal{Y}}(y) := u_Y(y, \cdot)_{\#} \mu_Y$ . Further, define  $dH_{\mathcal{X}} := (u_X)_{\#}(\mu_X \otimes \mu_X)$  and  $dH_{\mathcal{Y}} := (u_Y)_{\#}(\mu_Y \otimes \mu_Y)$ . Lemma C.2 implies that the set  $S := \text{spec}(X) \cup \text{spec}(Y)$  is a compact subset of  $(\mathbb{R}_{\geq 0}, \Lambda_\infty)$ . It is easy to see that  $\text{supp}(dh_{\mathcal{X}}), \text{supp}(dh_{\mathcal{Y}}), \text{supp}(dH_{\mathcal{X}}), \text{supp}(dH_{\mathcal{Y}}) \subseteq S \subseteq \mathbb{R}_{\geq 0}$ . Now, recall that by Proposition 4.4

$$\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = d_{\text{W},p}^{(S, \Lambda_\infty)}(dH_{\mathcal{X}}, dH_{\mathcal{Y}})$$

and

$$\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \left( \inf_{\pi \in \mathcal{C}(\mu_X, \mu_Y)} \int_{X \times Y} \left( d_{W,p}^{(S, \Lambda_\infty)}(dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y)) \right)^p \mu(dx \times dy) \right)^{1/p}.$$

Further, we observe for any  $x \in X$  and  $y \in Y$  that

$$d_{W,p}^{(S, \Lambda_\infty)}(dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y)) = \inf_{\pi_{xy} \in \mathcal{C}(dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y))} \left( \int_{S \times S} \Lambda_\infty^p(s, t) \pi_{xy}(ds \times dt) \right)^{\frac{1}{p}}.$$

For the remainder of this proof, the metric on metric on  $S \subseteq \mathbb{R}_{\geq 0}$  is always given by  $\Lambda_\infty$ . Additionally,  $\mathcal{P}(S)$  denotes the set of probability measures on  $S$  and we equip  $\mathcal{P}(S)$  with the Borel  $\sigma$ -field with respect to the topology induced by weak convergence.

*Claim 1:* There is a measurable choice  $(x, y) \mapsto \pi_{xy}^*$  such that for each  $(x, y) \in X \times Y$ ,  $\pi_{x,y}^*$  is an optimal transport plan between  $dh_{\mathcal{X}}(x)$  and  $dh_{\mathcal{Y}}(y)$ .

*Proof of Claim 1.* It is easy to see that both  $\Lambda_1$  and  $\Lambda_\infty$  induce the same topology and thus Borel sets on  $S$ . This therefore implies that  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_1)}$  and  $d_{W,p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}$  metrize the same weak topology on  $\mathcal{P}(S)$ . By Mémoli and Needham [68, Remark 2.5], the following two maps are continuous with respect to the weak topology and thus measurable:

$$\Phi_1 : X \rightarrow \mathcal{P}(S), \quad x \mapsto dh_{\mathcal{X}}(x)$$

and

$$\Phi_2 : Y \rightarrow \mathcal{P}(S), \quad y \mapsto dh_{\mathcal{Y}}(y).$$

Since  $S$  is a compact space, the space  $(\mathcal{P}(S), d_{W,p}^{(S, \Lambda_\infty)})$  is separable [100, Theorem 6.18]. This yields that  $\mathcal{B}(\mathcal{P}(S) \times \mathcal{P}(S)) = \mathcal{B}(\mathcal{P}(S)) \otimes \mathcal{B}(\mathcal{P}(S))$  [36, Proposition 1.5]. Hence, the product  $\Phi$  of  $\Phi_1$  and  $\Phi_2$ , defined by

$$\Phi : X \times Y \rightarrow \mathcal{P}(S) \times \mathcal{P}(S), \quad (x, y) \mapsto (dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y))$$

is measurable [36, Proposition 2.4]. Since  $\Phi$  is measurable, a direct application of Villani [100, Corollary 5.22] gives the claim.  $\square$

Now, we have that for every  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  that

$$\begin{aligned} & \int_{X \times Y} \left( d_{W,p}^{(S, \Lambda_\infty)}(dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y)) \right)^p \mu(dx \times dy) \\ &= \int_{X \times Y} \int_{S \times S} \Lambda_\infty^p(s, t) \pi_{xy}^*(ds \times dt) \mu(dx \times dy) \\ &= \int_{S \times S} \Lambda_\infty^p(s, t) \bar{\mu}(ds \times dt), \end{aligned}$$

by Fubini's Theorem, where  $\bar{\mu} \in \mathcal{P}(S \times S)$  is defined as

$$\bar{\mu}(A) := \int_{X \times Y} \pi_{xy}^*(A) \mu(dx \times dy) \tag{37}$$

for measurable  $A \subseteq S \times S$ . We remark that by Claim 1 the measure  $\bar{\mu}$  in Equation (37) is well defined. Next, we verify that  $\bar{\mu} \in \mathcal{C}(dH_{\mathcal{X}}, dH_{\mathcal{Y}})$ . For any measurable  $A \subseteq (S, \Lambda_{\infty})$  we have

$$\begin{aligned} \bar{\mu}(A \times S) &= \int_{X \times Y} \pi_{x,y}^*(A \times S) \mu(dx \times dy) \\ &= \int_{X \times Y} dh_{\mathcal{X}}(x)(A) \mu(dx \times dy) \\ &= \int_X dh_{\mathcal{X}}(x)(A) \mu_X(dx) \\ &\stackrel{(i)}{=} \int_X \int_X \mathbb{1}_{\{d_{\mathcal{X}}(x,x') \in A\}} \mu_X(dx') \mu_X(dx) \\ &= dH_{\mathcal{X}}(A), \end{aligned}$$

where we have applied the marginal constraints for  $\pi_{xy}$  and  $\mu$ . Further, (i) follows by the change-of-variables formula. The analogous arguments give that

$$\bar{\mu}(S \times B) = dH_{\mathcal{Y}}(B),$$

for any measurable  $B \subseteq S$ . Thus, we conclude that for every  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$

$$\begin{aligned} \int_{X \times Y} \left( d_{W,p}^{(S, \Lambda_{\infty})}(dh_{\mathcal{X}}(x), dh_{\mathcal{Y}}(y)) \right)^p \mu(dx \times dy) &= \int_{S \times S} \Lambda_{\infty}^p(s, t) \bar{\mu}(ds \times dt) \\ &\geq \inf_{\pi \in \mathcal{C}(dH_{\mathcal{X}}, dH_{\mathcal{Y}})} \int_{S \times S} \Lambda_{\infty}(s, t) \pi(ds \times dt) \\ &= \left( d_{W,p}^{(S, \Lambda_{\infty})}(dH_{\mathcal{X}}, dH_{\mathcal{Y}}) \right)^p. \end{aligned}$$

This gives the claim for  $p < \infty$ .

Next, we prove the assertion for the case  $p = \infty$ . Note that for any  $p < \infty$

$$\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| d_{W,p}^{(S, \Lambda_{\infty})}(dh_{\mathcal{X}}(\cdot), dh_{\mathcal{Y}}(\cdot)) \right\|_{L^p(\mu)} \quad (38)$$

$$\leq \inf_{\mu \in \mathcal{C}(\mu_X, \mu_Y)} \left\| d_{W,\infty}^{(S, \Lambda_{\infty})}(dh_{\mathcal{X}}(\cdot), dh_{\mathcal{Y}}(\cdot)) \right\|_{L^{\infty}(\mu)} \quad (39)$$

$$= \mathbf{TLB}_{\infty}^{\text{ult}}(\mathcal{X}, \mathcal{Y}), \quad (40)$$

where the inequality holds since  $d_{W,p}^{(S, \Lambda_{\infty})} \leq d_{W,\infty}^{(S, \Lambda_{\infty})}$  and  $\|\cdot\|_{L^p(\mu)} \leq \|\cdot\|_{L^{\infty}(\mu)}$ .

By Givens and Shortt [38, Proposition 3] we have that

$$\mathbf{SLB}_{\infty}^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = d_{W,\infty}^{(S, \Lambda_{\infty})}(dH_{\mathcal{X}}, dH_{\mathcal{Y}}) = \lim_{p \rightarrow \infty} d_{W,p}^{(S, \Lambda_{\infty})}(dH_{\mathcal{X}}, dH_{\mathcal{Y}}) = \lim_{p \rightarrow \infty} \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}).$$

Therefore,

$$\mathbf{SLB}_{\infty}^{\text{ult}}(\mathcal{X}, \mathcal{Y}) = \lim_{p \rightarrow \infty} \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) \leq \limsup_{p \rightarrow \infty} \mathbf{TLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) \leq \mathbf{TLB}_{\infty}^{\text{ult}}(\mathcal{X}, \mathcal{Y}).$$

□

C.1.2. *Proof of Proposition 4.4.* We only prove the first statement for  $p \in [1, \infty)$ . The case  $p = \infty$  as well as the second statement can be proven in a similar manner.

By directly using the change-of-variables formula, we have the following:

$$\begin{aligned} \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) &= \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \int_{X \times X \times Y \times Y} (\Lambda_\infty(u_X(x, x'), u_Y(y, y')))^p \gamma(d(x, x') \times d(y, y')) \\ &= \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} (\Lambda_\infty(s, t))^p (u_X \times u_Y)_\# \gamma(ds \times dt), \end{aligned}$$

where  $u_X \times u_Y : X \times X \times Y \times Y \rightarrow \mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}$  maps  $(x, x', y, y')$  to  $(u_X(x, x'), u_Y(y, y'))$ . By Lemma A.5, we have that

$$(u_X \times u_Y)_\# \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y) = \mathcal{C}((u_X)_\#(\mu_X \otimes \mu_X), (u_Y)_\#(\mu_Y \otimes \mu_Y)).$$

Therefore,

$$\begin{aligned} \mathbf{SLB}_p^{\text{ult}}(\mathcal{X}, \mathcal{Y}) &= \inf_{\gamma \in \mathcal{C}(\mu_X \otimes \mu_X, \mu_Y \otimes \mu_Y)} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} (\Lambda_\infty(s, t))^p (u_X \times u_Y)_\# \gamma(ds \times dt) \\ &= \inf_{\tilde{\gamma} \in \mathcal{C}((u_X)_\#(\mu_X \otimes \mu_X), (u_Y)_\#(\mu_Y \otimes \mu_Y))} \int_{\mathbb{R}_{\geq 0} \times \mathbb{R}_{\geq 0}} (\Lambda_\infty(s, t))^p \tilde{\gamma}(ds \times dt) \\ &= d_{\mathbf{W}, p}^{(\mathbb{R}_{\geq 0}, \Lambda_\infty)}((u_X)_\#(\mu_X \otimes \mu_X), (u_Y)_\#(\mu_Y \otimes \mu_Y)). \end{aligned}$$

C.1.3. *The relation between  $\mathbf{SLB}^{\text{ult}}$  and  $\mathbf{TLB}^{\text{ult}}$ .* Next, we will demonstrate that there are ultrametric measure spaces  $\mathcal{X}_1$  and  $\mathcal{X}_2$  such that  $\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}_1, \mathcal{X}_2) = 0$ , while it holds  $\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}_1, \mathcal{X}_2) > 0$ . To this end, consider the three point space  $\Delta_3(1) = (\{x_1, x_2, x_3\}, u)$  where  $u(x_i, x_j) = 1$  whenever  $i \neq j$ . Let  $\mu_1 := \frac{2}{3}\delta_{x_1} + \frac{1}{6}\delta_{x_2} + \frac{1}{6}\delta_{x_3}$  and let  $\mu_2 := \frac{1}{3}\delta_{x_1} + \left(\frac{1}{3} - \frac{1}{2\sqrt{3}}\right)\delta_{x_2} + \left(\frac{1}{3} + \frac{1}{2\sqrt{3}}\right)\delta_{x_3}$ . Both  $\mu_1$  and  $\mu_2$  are probability measures on  $\Delta_3(1)$ . We then let  $\mathcal{X}_1 := (\Delta_3(1), \mu_1)$  and  $\mathcal{X}_2 := (\Delta_3(1), \mu_2)$ . It is easy to check that

$$u_\#(\mu_1 \otimes \mu_1) = u_\#(\mu_2 \otimes \mu_2) = \frac{1}{2}\delta_0 + \frac{1}{2}\delta_1.$$

Then, by Proposition 4.4 we immediately have that  $\mathbf{SLB}_p^{\text{ult}}(\mathcal{X}_1, \mathcal{X}_2) = 0$  for any  $p \in [1, \infty]$ . Now, note that

$$u(x_1, \cdot)_\# \mu_1 = \frac{2}{3}\delta_0 + \frac{1}{3}\delta_1,$$

which is obviously different from all  $u(x_i, \cdot)_\# \mu_2$  for  $i = 1, 2, 3$ . This implies (by Proposition 4.4) that we have  $\mathbf{TLB}_p^{\text{ult}}(\mathcal{X}_1, \mathcal{X}_2) > 0$  for any  $p \in [1, \infty]$ .

In fact, this example works as well for showing that  $\mathbf{TLB}_p(\mathcal{X}_1, \mathcal{X}_2) > \mathbf{SLB}_p(\mathcal{X}_1, \mathcal{X}_2) = 0$ .

## APPENDIX D. MISSING DETAILS FROM SECTION 5

D.1. **Proofs from Section 5.** Next, we give the complete proofs of the results stated in Section 5.

D.1.1. *Proof of Theorem 5.6.* The first step to prove this is to verify the existence of an optimal coupling. To this end, we make the following obvious observation.

**Lemma D.1.** *Let  $X, Y$  be finite ultra-dissimilarity spaces, then  $\Lambda_\infty(u_X, u_Y) : X \times Y \times X \times Y \rightarrow \mathbb{R}_{\geq 0}$  is continuous with respect to the discrete topology.*

This allows us to verify the subsequent analogue to Proposition B.10.

**Proposition D.2.** *Let  $\mathcal{X}, \mathcal{Y} \in \mathcal{U}_{\text{dis}}^w$ . Then, for any  $p \in [1, \infty]$ , there always exists an optimal coupling  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  such that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = \text{dis}_p^{\text{ult}}(\mu)$ .*

*Proof.* The proof is essentially the same as the one for Proposition B.10. We only replace Lemma B.22 with Lemma D.1. The details are left to the reader.  $\square$

With Proposition D.2 available and Theorem 3.10 already proven, it is immediately clear how to verify the symmetry and the  $p$ -triangle inequality for  $u_{\text{GW},p}$  on  $\mathcal{U}_{\text{dis}}^w$ . Hence it only remains to demonstrate identity of indiscernibles.

*Proof of Theorem 5.6.* Due to the similarity between Theorem 5.6 and Theorem 3.10, we only verify that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$  if and only if  $\mathcal{X} \cong_w \mathcal{Y}$ . If  $\mathcal{X} \cong_w \mathcal{Y}$ , then obviously  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$ .

Next, we assume that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = 0$ . By Proposition D.2 there exists  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  such that  $u_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) = \text{dis}_p^{\text{ult}}(\mu) = 0$ . Now, we define a map  $\varphi : X \rightarrow Y$  as follows: For any  $x \in X$  we have  $\mu_X(\{x\}) > 0$ , since  $\mu_X$  has full support and  $X$  is finite. As a result,  $\mu(\{(x, y)\}) > 0$  for some  $y \in Y$ , then we let  $\varphi(x) \mapsto y$ . This map is well-defined. Indeed, if there are  $x \in X$  and  $y, y' \in Y$  such that  $\mu(\{(x, y)\}), \mu(\{(x, y')\}) > 0$ , then by  $\text{dis}_p^{\text{ult}}(\mu) = 0$  we must have that

$$\Delta_\infty(u_X(x, x), u_Y(y, y')) = \Delta_\infty(u_X(x, x), u_Y(y, y)) = \Delta_\infty(u_X(x, x), u_Y(y', y')) = 0.$$

This implies that  $u_Y(y, y') = u_Y(y, y) = u_Y(y', y') = u_X(x, x)$ . Since  $u_Y$  is an ultra-dissimilarity, we have that  $y = y'$  (cf. condition (3) in Definition 5.1). Essentially the same argument gives that  $\varphi : X \rightarrow Y$  is an injective map. As  $\mu \in \mathcal{C}(\mu_X, \mu_Y)$  and  $\varphi$  is injective, it follows  $\mu_X(\{x\}) = \mu(\{(x, \varphi(x))\}) \leq \mu_Y(\{\varphi(x)\})$  for any  $x \in X$ . Since

$$1 = \sum_{x \in X} \mu_X(\{x\}) \leq \sum_{x \in X} \mu_Y(\{\varphi(x)\}) \leq 1,$$

we have that  $\mu_X(\{x\}) = \mu_Y(\{\varphi(x)\})$  for all  $x \in X$ . Since  $\mu_Y$  is fully supported, this implies that  $\varphi$  is a bijective measure preserving map. Now, for any  $x, x' \in X$ ,  $\text{dis}_p^{\text{ult}}(\mu) = 0$  implies that  $\Delta_\infty(u_X(x, x'), u_Y(\varphi(x), \varphi(x'))) = 0$  and thus  $u_X(x, x') = u_Y(\varphi(x), \varphi(x'))$ . Therefore,  $\varphi$  is also an isometry and thus an isomorphism. In consequence,  $\mathcal{X} \cong_w \mathcal{Y}$ .  $\square$

## APPENDIX E. MISSING DETAILS FROM SECTION 6

E.1. **Missing details from Section 6.2.** Here, we list the precise results for the comparisons of the spaces  $\mathcal{X}_i$ ,  $1 \leq i \leq 4$ , illustrated in Figure 6. They are gathered in Table 2 and Table 3.

	$u_{\text{GW},1}$				$u_{\text{GW},\infty}$			
	$\mathcal{X}_1$	$\mathcal{X}_2$	$\mathcal{X}_3$	$\mathcal{X}_4$	$\mathcal{X}_1$	$\mathcal{X}_2$	$\mathcal{X}_3$	$\mathcal{X}_4$
$\mathcal{X}_1$	0.0000	0.9333	0.2444	0.7071	0.0000	2.1000	1.1000	2.000
$\mathcal{X}_2$	0.9333	0.0000	1.1778	1.5107	2.1000	0.0000	2.1000	2.1000
$\mathcal{X}_3$	0.2444	1.1778	0.0000	0.4493	1.1000	2.1000	0.0000	2.0000
$\mathcal{X}_4$	0.7071	1.5107	0.4493	0.0000	2.0000	2.1000	2.0000	0.0000

TABLE 2. **Comparison of different ultrametric measure spaces I:** The values of  $u_{\text{GW},1}(\mathcal{X}_i, \mathcal{X}_j)$  (approximated by Algorithm 1) and  $u_{\text{GW},\infty}(\mathcal{X}_i, \mathcal{X}_j)$ ,  $1 \leq i \leq j \leq 4$ , where  $\mathcal{X}_i$ ,  $1 \leq i \leq 4$ , denote the ultrametric measure spaces displayed in Figure 6.

	$\text{SLB}_1^{\text{ult}}$			
	$\mathcal{X}_1$	$\mathcal{X}_2$	$\mathcal{X}_3$	$\mathcal{X}_4$
$\mathcal{X}_1$	0.0000	0.9333	0.2444	0.0778
$\mathcal{X}_2$	0.9333	0.0000	1.1778	1.4522
$\mathcal{X}_3$	0.2444	1.1778	0.0000	0.2764
$\mathcal{X}_4$	0.0778	1.5107	0.2764	0.0000

TABLE 3. **Comparison of different ultrametric measure spaces II:** The values of  $\text{SLB}_1^{\text{ult}}(\mathcal{X}_i, \mathcal{X}_j)$ ,  $1 \leq i \leq j \leq 4$ , where  $\mathcal{X}_i$ ,  $1 \leq i \leq 4$ , denote the ultrametric measure spaces displayed in Figure 6.

E.2. **Missing details from Section 6.3.** Here, we state more results for the comparison of the ultrametric measure spaces illustrated in Figure 6 and give the precise construction of the ultrametric spaces  $Z_{k,t}^i$ ,  $2 \leq k \leq 5$ ,  $t = 0, 0.2, 0.4, 0.4$ ,  $1 \leq i \leq 15$ .

**The ultrametric measure spaces from Figure 6.** First, we give the precise results for comparing the ultrametric dissimilarity spaces in Figure 6 based on  $d_{\text{GW},1}$  and  $\text{SLB}_1$ . They are gathered in Table 4.

	$d_{\text{GW},1}$				$\text{SLB}_1$			
	$\mathcal{X}_1$	$\mathcal{X}_2$	$\mathcal{X}_3$	$\mathcal{X}_4$	$\mathcal{X}_1$	$\mathcal{X}_2$	$\mathcal{X}_3$	$\mathcal{X}_4$
$\mathcal{X}_1$	0.0000	0.0444	0.0222	0.2111	0.0000	0.0444	0.0222	0.0422
$\mathcal{X}_2$	0.0444	0.0000	0.0667	0.2556	0.0444	0.0000	0.0667	0.0867
$\mathcal{X}_3$	0.0222	0.0667	0.0000	0.2253	0.0222	0.0667	0.0000	0.0573
$\mathcal{X}_4$	0.2111	0.2556	0.2253	0.0000	0.0422	0.0867	0.0573	0.0000

TABLE 4. **Comparison of different ultrametric measure spaces III:** The values of  $d_{\text{GW},1}(\mathcal{X}_i, \mathcal{X}_j)$  (approximated by Algorithm 1) and  $\text{SLB}_1(\mathcal{X}_i, \mathcal{X}_j)$ ,  $1 \leq i \leq j \leq 4$ , where  $(X_i, d_{X_i}, \mu_{X_i})$ ,  $1 \leq i \leq 4$ , denote the ultrametric measure spaces displayed in Figure 6.

**Perturbations at level  $t$ .** Next, we give the precise construction of the ultrametric measure spaces  $Z_{k,t}^i$ ,  $2 \leq k \leq 5$ ,  $t = 0, 0.2, 0.4, 0.6$ ,  $1 \leq i \leq 15$ . For each  $k = 2, 3, 4, 5$  we first draw a sample with  $100 \times k$  points from the mixture distribution

$$\sum_{i=0}^k \frac{1}{k} U[1.5(k-1), 1.5(k-1)+1],$$

where  $U[a, b]$  denotes the uniform distribution on  $[a, b]$ . For each sample, we employ the single linkage algorithm to create a dendrogram, which then induces an ultrametric on the given sample. We further draw a 30-point subspace from each ultrametric space and denote it by  $Z_k$ . These four spaces have similar diameter values between 0.5 and 0.6. Each space  $Z_k$  is equipped with the uniform probability measure and the resulting ultrametric measure spaces are denoted by  $\mathcal{Z}_k = (Z_k, u_{Z_k}, \mu_{Z_k})$ ,  $k = 2, 3, 4, 5$ . We remark that  $k$  can be regarded as the number of blocks in the dendrogram representation of the obtained ultrametric measure spaces (see the top row of Figure 7 for a visualization of three 3-block spaces).

Finally, we introduce our method for perturbing ultrametric spaces. Given a perturbation level  $t \geq 0$  and an ultrametric space  $X$ , we consider the quotient space  $X_t$ . Each equivalence class  $[x]_t \subseteq X$  is an ultrametric subspace of  $X$ . If  $|[x]_t| > 1$ , we let  $m := |\text{spec}([x]_t)| - 1$  and write  $\text{spec}([x]_t) = \{0 < s_1 < \dots < s_m\}$ . Let  $\delta := \text{diam}([x]_t)$ . We generate  $m$  uniformly distributed numbers from  $[0, t - \delta]$  and sort them according to ascending order to obtain  $a_1 \leq \dots \leq a_m$ . We then perturb  $u_X|_{[x]_t \times [x]_t}$  by replacing  $s_i$  with  $s_i + a_i$  for each  $i = 1, \dots, m$ . We do the same for all equivalence classes  $[x]_t$  and thus obtain a new ultrametric on  $X$ .

DEPARTMENT OF MATHEMATICS AND DEPARTMENT OF COMPUTER SCIENCE AND ENGINEERING, THE OHIO STATE UNIVERSITY

*Email address:* `memoli@math.osu.edu`

INSTITUTE FOR MATHEMATICAL STOCHASTICS, UNIVERSITY OF GÖTTINGEN

*Email address:* `munk@math.uni-goettingen.de`

DEPARTMENT OF MATHEMATICS, THE OHIO STATE UNIVERSITY

*Email address:* `wan.252@osu.edu`

INSTITUTE FOR MATHEMATICAL STOCHASTICS, UNIVERSITY OF GÖTTINGEN

*Email address:* `cweitka@mathematik.uni-goettingen.de`