Soliton resolution for the Hirota equation with weighted Sobolev initial data *

Jin-Jie Yang, Shou-Fu Tian* and Zhi-Qiang Li

School of Mathematics, China University of Mining and Technology, Xuzhou 221116, People's Republic of China

Abstract

In this work, the $\overline{\partial}$ steepest descent method is employed to investigate the soliton resolution for the Hirota equation with the initial value belong to weighted Sobolev space $H^{1,1}(\mathbb{R}) = \{f \in L^2(\mathbb{R}) : f', xf \in L^2(\mathbb{R})\}$. The long-time asymptotic behavior of the solution q(x, t) is derived in any fixed space-time cone $C(x_1, x_2, v_1, v_2) = \{(x, t) \in \mathbb{R} \times \mathbb{R} : x = x_0 + vt \text{ with } x_0 \in [x_1, x_2]\}$. We show that solution resolution conjecture of the Hirota equation is characterized by the leading order term $O(t^{-1/2})$ in the continuous spectrum, $\mathcal{N}(\mathcal{I})$ soliton solutions in the discrete spectrum and error order $O(t^{-3/4})$ from the $\overline{\partial}$ equation.

Keywords: The Hirota equation, $\bar{\partial}$ -steepest descent method, Matrix Riemann-Hilbert problem, Soliton resolution.

Contents

1	Introduction	2
2	The spectral analysis	7
3	Conjugation	10
4	A mixed $\bar\partial$ -problem	13
5	The decomposition of the mixed $\overline{\partial}$ -RH probelm	18
6	Outer model RH problem	22

^{*}Corresponding author.

E-mail addresses: sftian@cumt.edu.cn, shoufu2006@126.com (S. F. Tian)

	6.1	The existence of soliton solution	22
	6.2	The long-time behavior of soliton solution	25
7	A lo	cal RH problem near phase points	29
8	The small norm RH problem for the error function		
9	Pure	$b\bar{\partial}$ -Problem	34
10	Solit	on resolution for the Hirota equation	36
11	App	endix A: The parabolic cylinder model problem	37
12	App	endix B: Detailed calculations for the pure $ar\partial$ -Problem	39

1. Introduction

In 1974, employing inverse scattering transformation to study the long-time asymptotic solutions of nonlinear equations was first proposed by Manakov [1]. Subsequently, Zakharov and Manakov obtained the long-time asymptotic solution of the nonlinear Schrödinger (NLS) equation with decaying initial value for the first time [2]. This method is used to study the long-time asymptotic solutions of nonlinear integrable systems, including the Landau-Lifshitz equation [3] and the Maxwell-Bloch equation [4], etc. Until 1993, Deift and Zhou proposed the nonlinear steepest method rigorously obtained the long-time asymptotic behavior of the matrix modified Korteweg-de Vries (mKdV) with decaying initial value [5], which is called Deift-Zhou (DZ) method. Since then, this method has been gradually applied to other integrable models with many excellent results including focusing NLS [6], short pulse equation [7, 8], Fokas-Lenells equation [9], Camassa-Holm equation [10], derivative nonlinear Schrödinger (DNLS) equation [11], and other equations [12]-[14].

Recently, McLaughlin and Miller [15, 16] proposed a new method to analyze the oscillatory Riemann-Hilbert (RH) problem based on the DZ method and the $\overline{\partial}$ dressing method [17]-[22], namely the $\overline{\partial}$ steepest descent method. This method is different from the DZ method in the following aspects: (i) For DZ method, it is necessary to extend the jump matrix analytically, while for the $\overline{\partial}$ descent method, it is only necessary to extend continuously. (ii) The $\overline{\partial}$ method avoids the delicate estimation involving the L^p estimation of Cauchy projection operator. (iii) The $\overline{\partial}$ method improves the error estimation without adding additional conditions to the initial data.

For finite mass initial data [23], this method was successfully applied to the defocusing NLS equation. In addition, this method was used by Jenkins et.al to study the soliton decomposition conjecture of the focusing NLS equation with the initial value belongs to a weighted Sobolev space [24], which decomposes the solution into the sum of a finite number of separated solitons and a radiative parts as $t \rightarrow \infty$. Furthermore, Cuccagna and Jenkins proved the asymptotic stability of *N*-soliton solutions of the de-focusing NLS equation by using the $\overline{\partial}$ steepest decent method [25]. Of course, in recent years, some excellent results have been obtained by using this method, which not only further improves the estimation accuracy of the DZ method, but also verifies the soliton decomposition conjecture, including DNLS [24, 26], short pulse equation [27], Fokas-Lenells equation [28], Kundu-Eckhaus equation [29], mKdV equation [30] etc.

In this work, we apply the $\overline{\partial}$ steepest decent method to investigate the Hirota equation [31] with the initial value $q_0(x)$ belongs to the weighted Sobolev space $H^{1,1}(\mathbb{R})$

$$iq_t + \alpha \left(q_{xx} + 2|q|^2 q \right) + i\beta \left(q_{xxx} + 6|q|^2 q_x \right) = 0, \quad \alpha, \beta \in \mathbb{R},$$

$$(1.1)$$

$$q(x,0) = q_0(x) \in H^{1,1}(\mathbb{R}), \tag{1.2}$$

where α , β denote the second-order and third-order dispersions, and $H^{1,1}(\mathbb{R})$ is the weighted Sobolev space, i.e.,

$$H^{1,1}(\mathbb{R}) = \left\{ f \in L^2(\mathbb{R}) : f', xf \in L^2(\mathbb{R}) \right\}.$$
 (1.3)

The Hirota equation can be regarded as modified Schrödinger equation with high-order dispersion and time-delay corrections to the cubic nonlinearity. It is a more accurate approximation than the NLS equation in describing wave propagation in the ocean and optical fiber [31]. Many properties and different kinds of exact solutions of the Hirota equation have been obtained, including rouge wave via Darboux transformation [32], solutions under non-zero boundary conditions via RH probelm [33], other aspects of research [34, 35]. In addition, the conservation laws have been investigated in detail in [36]. From Lax pair representation, the spectrum problem is transformed into Riccati equation

$$W_x = q_x \frac{W}{q} - |q|^2 + 2izW - W^2,$$

by introducing a transformation $W = q\Gamma$ with $\Gamma = \psi_2 \psi_1^{-1}$, where ψ_j (j = 1, 2) are the column of ψ . The following formula can be obtained from the compatibility condition

$$(-iz+W)_t = (A+B\Gamma)_x, \tag{1.4}$$

where the *A* and *B* denote the elements V_{11} and V_{12} of matrix *V* defined by (2.2), respectively, and *W* such that $W = \sum_{n=1}^{\infty} \varpi_n z^{-n}$. After calculation, it is known that ϖ_n satisfies the following recurrence relation

$$-2i\varpi_{j+1} + \sum_{k=1}^{j} \varpi_k \varpi_{j-k} - \frac{\varpi_j q_x}{q} + \varpi_{j,x} = 0, \quad j = 2, 3 \cdots,$$

and then the first three conserved quantities can be expressed as

$$I_{1} = -\frac{1}{2} \int_{-\infty}^{+\infty} iq(x,t)q^{*}(x,t)dx,$$

$$I_{2} = -\frac{1}{4} \int_{-\infty}^{+\infty} q(x,t)q^{*}_{x}(x,t)dx,$$

$$I_{3} = \frac{1}{8} \int_{-\infty}^{+\infty} iq(x,t) \left[q(x,t)(q^{*}(x,t))^{2} + q^{*}_{xx}(x,t)\right]dx, \dots$$

Beside that, Hamiltonian function can also be calculated by Lax representation [37]

$$\frac{d\psi_1}{dx} = \frac{\partial H}{\partial \psi_2}, \quad \frac{d\psi_1}{dx} = -\frac{\partial H}{\partial \psi_1}, \tag{1.5}$$

with the real valued Hamiltonian function

$$H = z\psi_1\psi_2 + \overline{z}\overline{\psi}_1\overline{\psi}_2 + \frac{1}{2}(\psi_1^2 + \overline{\psi}_2^2)(\psi_2^2 + \overline{\psi}_1^2), \qquad (1.6)$$

where $\psi = (\psi_1 \ \psi_2)$ is a nonzero solution of the Lax pair, and the $\overline{\psi}$ denotes complex conjugation of ψ .

It is worth noting that the DZ method was used to study the long-time asymptotic solution of Hirota equation with the error estimate $\epsilon(x, t) = O(t^{-1} \log t)$ in [38]. The initial boundary value problem of (1.1) was studied in [39]. In addition, the Hirota equation can be regarded as a mixture of nonlinear Schrödinger equation and complex mKdV equation, because when the parameters are $\alpha = 1$, $\beta = 0$ and $\alpha = 0$, $\beta = 1$ respectively, the equation (1.1) degenerates to nonlinear Schrödinger equation and complex mKdV equation, namely

• Taking $\alpha = 1$, $\beta = 0$, equation (1.1) reduces to the NLS equation

$$iq_t + q_{xx} + 2|q|^2 q = 0, (1.7)$$

• Taking $\alpha = 0$, $\beta = 1$, equation (1.1) reduces to the mKdV equation

$$iq_t + q_{xxx} + 6|q|^2 q_x = 0. (1.8)$$

For NLS equation (1.7), the long-time asymptotic solution was investigated by Deift and Zhou via DZ method in [6, 40], which show that the error term $\varepsilon(x, t) = O(t^{-1} \log t)$ as the initial value $q_0(x)$ has a high degree of smoothness and decay. In [41], Borghese and Jenkins et.al studied the initial value in the weighted Sobolev space by using the $\overline{\partial}$ steepest decent method, the error estimate is $\varepsilon(x, t) = O(t^{-3/4})$. In addition, the soliton resolution conjecture was proved. On the other hand, Tian employed Fokas method to study the initial boundary value problem of coupled NLS equations [42, 43]. For mKdV equation, similar results were obtained in [30]. The soliton resolution conjecture about NLS equation and mKdV equation has been proved. Therefore in this work, we will prove that the soliton resolution conjecture about Hirota equation is also true based on the relationship between the two equations and Hirota equation.

A brief description of the scattering data for the Hirota equation (1.1) is given as follow: generally, the points where the diagonal element $s_{11}(z)$ of the scattering matrix are zero are called spectral points. When these spectral points satisfy $s_{11}(z_n) =$ 0, $s'_{11}(z_n) \neq 0$, which are simple spectral points. It is worth noting that these spectral points are distributed in the whole complex *z*-plane. When $s_{11}(z) = 0$, $z \in R$ holds, these points are called spectral singularities. If there are spectral singularities, the spectral points may be infinite. When there are no spectral singularities, the spectral points are finite, which consist of the finite discrete spectrum set. c_k is called a norming constant if it is a nonzero constant associated with a simple discrete spectrum. For the initial data $q_0(x)$ with only simple zeros and no spectral singularities, the minimum scattering data set of Hirota equation is $D = \{\gamma(z), \{(z_k, c_k)\}_{k=1}^N\}$. We will not discuss the case of higher order zeros and spectral singularities, which is described in [44].

In this work, we consider the long-time asymptotic solution of Hirota equation by using the $\overline{\partial}$ steepest decent method for initial value $q_0(x)$ with simple zeros and spectral singularities. However, it should be noted that the steady-state phase points z_0 and z_1 are asymmetric. We use the similar method in [30], that is, adding a jump contour, and we can prove that the new jump contour has no effect on the solution.

The main result and remark is as follows.

We will give the main theorem of this work, the soliton resolution conjecture, that is, the solution is decomposed into the sum of a finite number of separated solitons and a radiative parts as $t \to \infty$.

Theorem 1.1. Let q(x, t) be the solution of equation (1.1) whose initial value $q(x, 0) = q_0(x)$ belongs to the weighted Sobolev space $H^{1,1}(\mathbb{R})$ and satisfies Assumption 2.2.

Denoting $D = \{\gamma(z), \{(z_k, c_k)\}_{k=1}^N\}$ is the scattering data corresponding to the initial value without spectral singularities. Fixed $x_1, x_2, v_1, v_2 \in \mathbb{R}$ with $x_1 < x_2, v_1 < v_2$,

$$I = \{z : f(v_2) < |z| < f(v_1)\},$$
(1.9)

$$\mathcal{K}(I) = \{ z_j \in \mathcal{K} : z_j \in I \}, \quad \mathcal{N}(I) = |\mathcal{K}(I)|, \tag{1.10}$$

and a cone shown in Fig.5. Let $m_{sol}(x, t|D(I))$ be the N(I) soliton solution corresponding to the scattering data $\{z_j, c_j(I)\}_{j=1}^N$ defined by (6.20), it follows from (2.18), (6.16), (6.36), (8.10) and (9.17) that

$$m(x,t) = m_{sol}(x,t|D(I)) + t^{-1/2}f(x,t) + O(t^{-3/4}),$$
(1.11)

which implies that the long-time asymptotic behaviour can be expressed by

$$q(x,t) = m_{sol}(x,t|D(I)) + t^{-1/2}f(x,t) + O(t^{-3/4}),$$
(1.12)

where f(x, t) is defined by (8.14).

Remark 1.2. Here we point out that the result of **Theorem** 1.1 degenerates to the work of Borghese and Jenkins [41] when the parameters $\alpha = 1$, $\beta = 0$, but it is worth noting that when the parameters $\alpha = 0$, $\beta = 1$, it is different from the work of Chen and Liu, whose work is real mKdV in [30]. In addition, dealing with the real mKdV, they gave different solutions according to different regions.

Organization of the work

In Section 2, we construct the analytic, asymptotic and symmetry of the eigenfunctions and scattering matrix to establish a suitable RH problem for M(z), from which we establish the relationship between the solution of Hirota equation and the solution of RH problem. In Section 3, a new matrix valued function $M^{(1)}(z)$ is defined by introducing T(z), which has regular discrete spectrum and two triangular decompositions at steady-state phase points. In Section 4, by introducing $R^{(2)}(z)$, we get a mixed $\overline{\partial}$ -RH problem about $M^{(2)}(z)$, whose jump path changes from the real axis \mathbb{R} to the the descent line. In Section 5, we decompose the mixed RH problem $M^{(2)}(z)$ into a pure RH problem and a pure $\overline{\partial}$ problem corresponding to $M^{(2)}_{RHP}(z)$ and $M^{(3)}(z)$, respectively. In section 6, the $M^{(2)}_{RHP}(z)$ can be solved by an out model $M^{(out)}(z)$ for the soliton components under the reflection-less condition. In section 7, the internal models $M^{(z_0)}(z)$ and $M^{(z_1)}(z)$ near the steady-state phase points z_0 and z_1 can be approximated by the known solvable parabolic cylinder model. In Section 8, the error function E(z) is considered by using the small norm RH problem. In Section 9, with the help of known results, the pure $\overline{\partial}$ problem $M^{(3)}(z)$ is considered. Finally, the soliton resolution and long-time asymptotic solution of Hirota equation are obtained via the inverse transformation

$$M(z) \leftrightarrows M^{(1)}(z) \leftrightarrows M^{(2)}(z) \leftrightarrows M^{(3)}(z) \leftrightarrows E(z).$$

2. The spectral analysis

In this section, we mainly establish the eigenfunctions and scattering matirx related to the initial value $q(x, 0) \in H^{1,1}$, and further obtain the corresponding analysis, asymmetry and symmetry. Note that the Hirota equation admits the Lax pair

$$\psi_x(x,t;z) + iz[\sigma_3,\psi(x,t;z)] = Q\psi(x,t;z),$$
(2.1)

$$\psi_t(x,t;z) + \left(4i\beta z^3 + 2i\alpha z^2\right)[\sigma_3,\psi(x,t;z)] = V\psi(x,t;z),$$
(2.2)

where

$$Q = -i[\sigma_3, \langle \psi R \rangle] = \begin{pmatrix} 0 & q \\ -q^* & 0 \end{pmatrix},$$
$$V = \alpha \left[2zQ + i\sigma_3(Q_x - Q^2) \right] + \beta \left[4z^2Q + 2iz\sigma_3(Q_x - Q^2) + [Q_x, Q] - Q_{xx} + 2Q^3 \right].$$

Considering the asymptotic spectrum problem as $x \to \infty$, we introduce the two eigenfunctions

$$\phi_{\pm} = \psi_{\pm} e^{i \left[zx + (4\beta z^3 + 2\alpha z^2)t \right] \sigma_3} \to \mathbb{I}, \text{ as } x \to \infty.$$
(2.3)

The linear spectrum problem can be written as

$$\phi_x + iz[\sigma_3, \phi] = Q\phi,$$

$$\phi_t + (4i\beta z^3 + 2i\alpha z^2)[\sigma_3, \phi] = V\phi,$$
(2.4)

which can be written as a fully differential form

$$d\left(e^{i\left[zx+(4\beta z^3+2\alpha z^2)t\right]\widehat{\sigma}_3}\phi\right) = e^{i\left[zx+(4\beta z^3+2\alpha z^2)t\right]\widehat{\sigma}_3}\left(Qdx+Vdt\right)\phi.$$
(2.5)

From Eq.(2.5), we know that the integral has nothing to do with the path, so we choose two special paths $(x_1, t_1) = (-\infty, t)$ and $(x_2, t_2) = (+\infty, t)$ shown in Fig. 1. Then ϕ_{\pm} are determined by the following integral equations

$$\phi_{-}(x;z) = \mathbb{I} + \int_{-\infty}^{x} e^{-iz(x-y)\widehat{\sigma}_{3}} Q(y;z)\phi_{-}(y;z)dy,$$

$$\phi_{+}(x;z) = \mathbb{I} - \int_{x}^{+\infty} e^{-iz(x-y)\widehat{\sigma}_{3}} Q(y;z)\phi_{+}(y;z)dy.$$
(2.6)




Figure 1. The contour of integral (2.6).

From the integral equations (2.6), the following results can be proved similar to those in [45, 46].

Proposition 2.1. Assume that the initial value $q(x) - q_0 \in H^{1,1}$, the eigenfunctions ϕ_{\pm} admit

• $\phi_{-,1}$ and $\phi_{+,2}$ are analytic in $\mathbb{C}^+ = \{z | Imz > 0\}$, and continuous to the real axis \mathbb{R} ,

• $\phi_{-,2}$ and $\phi_{+,1}$ are analytic in $\mathbb{C}^- = \{z | Imz < 0\}$, and continuous to the real axis \mathbb{R} , where $\phi_{\pm,j}$ (j = 1, 2) denote the j – th column of ϕ_{\pm} .

For the first-order linear spectral problem, since both ϕ_- and ϕ_+ are solutions, there exists a matrix $S(z) = (s_{ij})_{2\times 2}$ which is independent of time and space such that

$$\phi_{-}(x;z) = \phi_{+}(x;z)S(z), \quad x, t, z \in R.$$
(2.7)

Note that trQ = 0, it follows from the Liouville's formula [47] that $(\det \phi)_x = (\det \phi)_t = 0$, furthermore, $\det \psi_{\pm} = \det \phi_{\pm} = 1$ can be derived by Eq.(2.3). The elements of the scattering matrix S(z) can be expressed as eigenfunctions by direct calculation as follows

$$s_{11}(z) = Wr(\phi_{-,1}, \phi_{+,2}), \quad s_{22}(z) = Wr(\phi_{+,1}, \phi_{-,2}), \quad (2.8)$$

where Wr represents the Wronskians determinant. Combining with the Proposition (2.1) and Eq.(2.8), we have that the scattering data s_{11} is analytic in \mathbb{C}^+ , and s_{22} is analytic in \mathbb{C}^- .

Next, we consider the symmetry of the eigenfunctions and the scattering matrix, which is related to the distribution of the zeros of the scattering data. It plays a decisive role in the analysis of the corresponding RH problem and the establishment of the residue condition. Assuming that $\phi(x, t; z)$ is the solution of the first formula for Eq.(2.4), so is $\sigma_2 \phi^*(x, t; z^*) \sigma_2$, where $\sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}$. It follows from asymptotic behavior (2.3) that

$$\phi(x,t;z) = \sigma_2 \phi^*(x,t;z)\sigma_2, \quad z \in \mathbb{R},$$
(2.9)

from which together with (2.7), we obtain the symmetry of the scattering matrix

$$S(z) = \sigma_2 S^*(z^*) \sigma_2,$$
 (2.10)

which leads to

$$s_{11}(z) = s_{22}^*(z^*), \quad s_{12}(z) = -s_{21}^*(z^*).$$
 (2.11)

According to the symmetry of scattering data, we further introduce the reflection coefficient and its symmetry

$$\gamma(z) = \frac{s_{21}(z)}{s_{11}(z)}, \quad \frac{s_{12}(z)}{s_{22}(z)} = -\gamma^*(z), \ z \in \mathbb{R}.$$
(2.12)

In order to ensure that the discrete spectrum is finite, and to avoid the need to deal with many pathologies possible when discussing general initial value problems, we will make the following assumptions

Assumption 2.2. The initial value q(x, 0) for the Hirota equation satisfies the following assumptions:

(*I*) The scattering data $s_{11}(z) \neq 0$ for all $z \in \mathbb{R}$.

(*II*) The scattering data $s_{11}(z)$ only is of finite number of simple zeros, namely, assume that the set $\mathcal{Z} = \{z_j\}_{j=1}^N$ is composed of *N* zeros of scattering data $s_{11}(z)$. From the symmetry Eq.(2.11), the set $\overline{\mathcal{Z}} = \{z_j^*\}_{j=1}^N$ is composed of *N* zeros of scattering data $s_{22}(z)$.

To get a suitable RH problem, we introduce piecewise meromorphic function M(z) = M(x, t; z)

$$M(x,t;z) = \begin{cases} \left(\frac{\phi_{-,1}(x,t;z)}{s_{11}(z)}, \phi_{+,2}(x,t;z)\right), & z \in \mathbb{C}^+, \\ \left(\phi_{+,1}(x,t;z), \frac{\phi_{-,2}(x,t;z)}{s_{22}(z)}\right), & z \in \mathbb{C}^-, \end{cases}$$
(2.13)

which satisfies the following RH problem.

RH Problem 2.3. Find an analytic function M(z) = M(x, t; z) that satisfies the following properties:

- (1) M(z) is analytic in $C \setminus (R \cup Z \cup \overline{Z})$.
- (II) The jump condition:

$$M^{+}(z) = M^{-}(z)V(z), \quad z \in \mathbb{R} \triangleq \Sigma^{(1)},$$
 (2.14)

where $M^{\pm}(x, t; z) = \lim_{\varepsilon \to 0^+} M(x, t; z \pm i\varepsilon), \ \varepsilon \in \mathbb{R}$, and

$$V(z) = \begin{pmatrix} 1 + |r(z)|^2 & -r^*(z)e^{-2it\theta(z)} \\ r(z)e^{2it\theta(z)} & 1 \end{pmatrix},$$
(2.15)

where $\theta(z) = z\frac{x}{t} + 2\alpha z^2 + 4\beta z^3$.

(III) Asymptotic behavior: $M(z) = \mathbb{I} + O(z^{-1})$ as $z \to \infty$.

(*IV*) For any $z_j \in \mathbb{Z}$ and $z_j \in \overline{\mathbb{Z}}$, the residue conditions for M(z) are determined by

$$\operatorname{Res}_{z=z_{j}} M = \lim_{z \to z_{j}} M \begin{pmatrix} 0 & 0 \\ c_{j} e^{2it\theta(z)} & 0 \end{pmatrix}, \quad \operatorname{Res}_{z=z_{j}^{*}} M = \lim_{z \to z_{j}^{*}} M \begin{pmatrix} 0 & -c_{j}^{*} e^{-2it\theta(z)} \\ 0 & 0 \end{pmatrix}.$$
(2.16)

Remark 2.4. The Zhou's vanishing lemma [44] can guarantee the existence of the solution of the **RH Problem** 2.3 by transforming the poles into jumps on a sufficiently small circular contours. The uniqueness of the solution is a natural result of Liouville's theorem when the solution exists.

Expanding M(z) as follows

$$M(z) = \mathbb{I} + \frac{M_1(z)}{z} + O(z^{-1}), \quad z \to \infty,$$
(2.17)

and substituting it into Eq.(2.4), we have the following reconstruction formula of q(x, t)

$$q(x,t) = 2i \lim_{z \to \infty} (zM(z))_{12},$$
(2.18)

by comparing the coefficients of z^i (i = 0, 1, 2, 3), where A_{ij} denote the i - th row and j - th column of the matrix A.

3. Conjugation

In order to study the long-time asymptotic behavior of solutions, we first deal with the oscillatory term in the jump matrix defined in Eq.(2.15), that is, we decompose the jump matrix according to the sign change graph of $Re(it\theta)$ a to ensure that any jump matrix is bounded in a given region. In this work, we only consider the case $\beta > 0$ for our analysis convenient.

As shown in [5], the stationary points of the function $\theta(z)$ should be considered

$$\theta(z) = z\frac{x}{t} + 2\alpha z^2 + 4\beta z^3.$$
(3.1)

Letting

$$\frac{d(\theta(z))}{dz} = 0$$

which leads to

$$z_0 = \frac{-\alpha - \sqrt{\alpha^2 - 3\beta x/t}}{6\beta}, \quad z_1 = \frac{-\alpha + \sqrt{\alpha^2 - 3\beta x/t}}{6\beta}.$$
 (3.2)

The sign change diagram of the function $Re(it\theta(z))$ is shown in Fig. 2.

$$Re(it\theta(z)) > 0$$


$$Re(it\theta(z)) < 0$$

$$Re(it\theta(z)) < 0$$

$$Re(it\theta(z)) > 0$$

$$Re(it\theta(z)) > 0$$

$$Re(it\theta(z)) < 0$$



For $z \in (z_0, z_1)$, the upper and lower triangular decomposition of jump matrix V(z) will produce a diagonal matrix, therefore similar to [38] we first introduce a scalar RH problem about $\delta(z)$ to deal with this case

$$\begin{cases} \delta_{+}(z) = \delta_{-}(z)(1 + |\gamma(z)|^{2}), & z \in (z_{0}, z_{1}), \\ \delta_{+}(z) = \delta_{-}(z) = \delta(z), & \mathbb{R} \setminus (z_{0}, z_{1}), \\ \delta(z) \to \mathbb{I}, & z \to \infty. \end{cases}$$
(3.3)

Then the solution of Eq.(3.3) can be written as

$$\delta(z) = exp\left[\frac{1}{2\pi i} \int_{z_0}^{z_1} \frac{\log(1+|\gamma|^2)}{s-z} ds\right] \triangleq exp\left(i \int_{z_0}^{z_1} \frac{\nu(s)}{s-z} ds\right),$$
(3.4)

where $v(z) = -\frac{1}{2\pi} \log(1 + |\gamma(z)|^2)$.

For the convenience of the following analysis, we introduce the following notation

$$\Delta_{z_1}^- = \{k \in \{1, 2, \cdots, N\}, \ z_0 < z_k < z_1\},\tag{3.5}$$

$$\Delta_{z_1}^+ = \{k \in \{1, 2, \cdots, N\}, \ z_0 < z_k, z_k > z_1\},\tag{3.6}$$

$$T(z) = T(z, z_1) = \prod_{z \in \Delta_{z_1}^-} \frac{z - z_k^*}{z - z_k} \cdot \delta(z),$$
(3.7)

$$T_0(z_1) = T(z_1, z_1) = \prod_{z \in \Delta_{z_1}^-} \frac{z_1 - z_k^*}{z_1 - z_k} \cdot e^{i\beta^+(z_1, z_1)},$$
(3.8)

$$T_0(z_0) = T(z_0, z_1) = \prod_{z \in \Delta_{z_1}^-} \frac{z_0 - z_k^*}{z_0 - z_k} \cdot e^{i\beta^-(z_0, z_1)},$$
(3.9)

$$\beta^{+}(z,z_{1}) = -\nu(z_{1})\ln(z-z_{1}+1) + \int_{z_{0}}^{z_{1}} \frac{\nu(s) - \chi_{1}(s)\nu(z_{1})}{s-z} ds, \qquad (3.10)$$

$$\beta^{-}(z, z_0) = -\nu(z_0)\ln(z - z_0 + 1) + \int_{z_0}^{z_1} \frac{\nu(s) - \chi_0(s)\nu(z_0)}{s - z} ds,$$
(3.11)

$$\chi_1(s) = \begin{cases} 1, & z_1 < s < z_1 + 1, \\ 0, & \text{elsewhere,} \end{cases}, \quad \chi_0(s) = \begin{cases} 1, & z_0 < s < z_0 + 1, \\ 0, & \text{elsewhere.} \end{cases}.$$
(3.12)

Proposition 3.1. *The function* T(z) *defined by* Eq.(3.7) *such that*

(I) T(z) is meromorphic function in $\mathbb{C} \setminus (z_0, z_1)$, and has a simple pole at z_n and a simple zero at z_n^* for $n \in \Delta_{z_1}^-$.

(II) For $z \in \mathbb{C} \setminus (z_0, z_1)$, $T^*(z^*)T(z) = 1$. (III) For $z \in \mathbb{C} \setminus (z_0, z_1)$, T(z) is of the jump condition

$$T_+(z) = T_-(z)(1 + |\gamma(z)|^2), \ z \in (z_0, z_1),$$

where $T_{\pm}(z) = \lim_{\varepsilon \to 0^+} T(z \pm i\varepsilon)$. (IV) As $|z| \to \infty$ with $|arg(z)| \le c < \pi$,

$$T(z) = 1 + \frac{i}{z} \left[2 \sum_{k \in \Delta_{z_0}^-} Im(z_k) - \int_{z_0}^{z_1} v(s) ds \right] + O(z^{-2}).$$
(3.13)

(V) As $z \to z_0$ along any ray $z_0 + e^{i\phi}R_+$ and $z \to z_1$ along any ray $z_1 + e^{i\phi}R_+$ with $|\phi| \le c < \pi$

$$|T(z-z_1) - T_0(z_1)(z-z_1)^{i\nu(z_1)}| \le C ||r||_{H^1(R)} |z-z_1|^{\frac{1}{2}},$$
(3.14)

$$|T(z-z_0) - T_0(z_0)(z-z_0)^{i\nu(z_0)}| \le C ||r||_{H^1(R)} |z-z_0|^{\frac{1}{2}},$$
(3.15)

In order to redistribute the poles and eliminate the diagonal matrix generated by the matrix triangular decomposition of the jumping matrix, we employ T(z) to define a new matrix function $M^{(1)}(z)$ such that

$$M^{(1)}(z) = M(z)T(z)^{-\sigma_3},$$
(3.16)

which admits the RH problem

RH Problem 3.2. The new matrix $M^{(1)}(z)$ such that:

(*I*) Analyticity: $M^{(1)}(z)$ is analytic in $z \in \mathbb{C} \setminus \{\mathbb{R} \cup \mathbb{Z} \cup \mathbb{Z}^*\}$.

(*II*) Jump condition: as z approaches the real axis from above and below, $M^{(1)}(z)$ satisfies

$$M_{+}^{(1)}(z) = M_{-}^{(1)}(z)V^{(1)}(z), \quad z \in \mathbb{R},$$
(3.17)

where

$$V^{(1)} = \begin{cases} \begin{pmatrix} 1 & \gamma^*(z)T(z)^2 e^{-2it\theta} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ \gamma(z)T(z)^{-2}e^{2it\theta} & 1 \end{pmatrix}, & z \in C \setminus \Gamma, \\ \begin{pmatrix} 1 & 0 \\ \frac{\gamma(z)T_-^{-2}(z)}{1+|\gamma(z)|^2}e^{2it\theta} & 1 \end{pmatrix} \begin{pmatrix} 1 & \frac{\gamma^*(z)T_+^2(z)}{1+|\gamma(z)|^2}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Gamma. \end{cases}$$
(3.18)

(III) Asymptotic behaviours:

$$M^{(1)}(z) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$$
 (3.19)

(IV) The residue conditions

$$\begin{split} Res_{z=z_{k}}^{es} M^{(1)} &= \begin{cases} \lim_{z \to z_{k}} M^{(1)} \begin{pmatrix} 0 & c_{k}^{-1} \left((\frac{1}{T})'(z_{k}) \right)^{-2} e^{-2it\theta} \\ 0 & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}} M^{(1)} \begin{pmatrix} 0 & 0 \\ c_{k}^{-1} T^{-2}(z_{k}) e^{2it\theta} & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{+}, \end{cases} \end{split}$$

$$\begin{aligned} Res_{z=z_{k}^{*}}^{es} M^{(1)} &= \begin{cases} \lim_{z \to z_{k}^{*}} M^{(1)} \begin{pmatrix} 0 & 0 \\ -\bar{c}_{k}^{-1} (T'(z_{k}^{*}))^{-2} e^{2it\theta^{*}(z)} & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}^{*}} M^{(1)} \begin{pmatrix} 0 & -c_{k}^{*} (T(z_{k}^{*}))^{2} e^{-2it\theta^{*}(z)} \\ 0 & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{+}. \end{cases} \end{aligned}$$

$$(3.20)$$

4. A mixed $\bar{\partial}$ -problem

We now will deform the jump contour of **RH Problem** 3.2 so that the oscillation term in the jump matrix is bounded on the transformed contour. It follows that from

Eq.(3.2), the phase function Eq.(3.1) has two phase points at z_0 and z_1 . As usual, the new contour is chosen as follows

$$\Sigma^{(2)} = \Sigma_1 \cup \Sigma_2 \cup \Sigma_3 \cup \Sigma_4 \cup \Sigma_5 \cup \Sigma_6 \cup \Sigma_7 \cup \Sigma_8, \tag{4.1}$$

and consists of the rays $z_0 + e^{i\phi}R^+$ and $z_1 + e^{i\phi}R^+$ with $\phi = \frac{\pi}{4}, \frac{3\pi}{4}, \frac{5\pi}{4}, \frac{7\pi}{4}$. Then the contour $\Sigma^{(2)}$ and the real axis *R* separate the complex into ten sectors denoted by Ω_j , $(j = 1, 2, \dots, 6, 7^{\pm}, 8^{\pm})$ shown in Fig. 3.




Figure 3. The contour of deformation from \mathbb{R} to $\Sigma^{(2)}$.

For technical reasons, we define

$$\Xi_{\mathcal{Z}}(z) = \begin{cases} 1, & dist(z, \mathcal{Z} \cup \mathcal{Z}^*) < \rho/3, \\ 0, & dist(z, \mathcal{Z} \cup \mathcal{Z}^*) > 2\rho/3. \end{cases}$$
(4.2)

where

$$\rho = \frac{1}{2} \min_{\lambda, \zeta \in \mathcal{Z} \cup \mathcal{Z}^* \lambda \neq \mu} |\lambda - \zeta|.$$
(4.3)

Note that the discrete spectrum appear as conjugate pairs from Eq.(2.11), moreover one has $dist(\mathcal{Z} \cup \mathcal{Z}^*, R) > \rho$ from the Assumption (2.2).

Now we introduce a new matrix $M^{(2)}$ such that the jump contour of the **RH Prob**lem 3.2 is transformed from $\Sigma^{(1)}$ to $\Sigma^{(2)}$


$$M^{(2)}(z) = M^{(1)}(z)R^{(2)}(z), (4.4)$$


Remark 4.1. The selection of matrix $R^{(2)}(z)$ shown in Fig.4 and Fig.5 needs to meet the following conditions:

► To remove the jump on the real axis and the new analytic jump matrix has the expected exponential decay along the contour $\Sigma^{(2)}$.

► The norm of $R^{(2)}(z)$ should be controlled to ensure that the long-time asymptotic behavior of $\overline{\partial}$ -contribution to the solution q(x, t) is negligible.

► The residues are not affected by this transformation.





Proposition 4.2. The matrices $R_j : \overline{\Omega} \to C$ have the following boundary values:

$$R_{1}(z) = \begin{cases} \gamma(z)T^{-2}(z), & z \in (z_{1}, \infty), \\ \gamma(z_{1})T_{0}^{-2}(z_{1})(z-z_{1})^{-2i\nu(z_{1})}(1-\Xi_{Z_{1}}), & z \in \Sigma_{1}, \end{cases}$$

$$(4.5)$$

$$R_{6}(z) = \begin{cases} \gamma(z)T(z), & z \in (z_{1}, \omega), \\ \gamma^{*}(z_{1})T_{0}^{2}(z_{1})(z - z_{1})^{2i\nu(z_{1})}(1 - \Xi_{Z_{1}}), & z \in \Sigma_{4}, \end{cases}$$
(4.6)

$$R_{3}(z) = \begin{cases} \gamma(z)T^{-2}(z), & z \in (-\infty, z_{0}), \\ \gamma(z_{0})T_{0}^{-2}(z_{0})(z - z_{0})^{-2i\nu(z_{0})}(1 - \Xi_{\mathcal{I}_{0}}), & z \in \Sigma_{2}, \end{cases}$$
(4.7)

$$R_4(z) = \begin{cases} \gamma^*(z)T^2(z), & z \in (-\infty, z_0), \\ \gamma^*(z_0)T_0^2(z_0)(z - z_0)^{2i\nu(z_0)}(1 - \Xi_{\mathcal{Z}_0}), & z \in \Sigma_3, \end{cases}$$
(4.8)

$$R_{7^{+}}(z) = \begin{cases} \frac{\gamma^{*}(z)T_{+}^{-2}(z)}{1+|\gamma(z)|^{2}}, & z \in (z_{0}, z_{1}), \\ \frac{\gamma^{*}(z_{1})T_{0}^{-2}(z_{1})}{1+|\gamma(z_{1})|^{2}}(z-z_{1})^{-2i\nu(z_{1})}(1-\Xi_{Z_{1}}), & z \in \Sigma_{6}, \end{cases}$$
(4.9)

$$R_{8^{+}}(z) = \begin{cases} \frac{\gamma(z)T_{-}^{-2}(z)}{1+|\gamma(z)|^{2}}, & z \in (z_{0}, z_{1}), \\ \frac{\gamma(z_{1})T_{0}^{-2}(z_{1})}{1+|\gamma(z_{1})|^{2}}(z-z_{1})^{-2i\nu(z_{1})}(1-\Xi_{Z_{1}}), & z \in \Sigma_{8}, \\ \left(\gamma^{*}(z)T_{+}^{2}(z)\right) & z \in (z_{0}, z_{1}) \end{cases}$$
(4.10)

$$R_{7^{-}}(z) = \begin{cases} \frac{1}{1+|\gamma(z)|^2}, & z \in (z_0, z_1), \\ \frac{\gamma^*(z_0)T_0^2(z_0)}{1+|\gamma(z_0)|^2} (z-z_0)^{2i\nu(z_0)} (1-\Xi_{Z_0}), & z \in \Sigma_5, \end{cases}$$
(4.11)
$$\left(\frac{\gamma(z)T_-^{-2}(z)}{1+|\gamma(z_0)|^2} (z-z_0)^{2i\nu(z_0)} (1-\Xi_{Z_0}), & z \in \Sigma_5, \end{cases}$$

$$R_{8^{-}}(z) = \begin{cases} \frac{\gamma(z_{0}-z_{0})}{1+|\gamma(z_{0})|^{2}}, & z \in (z_{0}, z_{1}), \\ \frac{\gamma(z_{0})T_{0}^{-2}(z_{0})}{1+|\gamma(z_{0})|^{2}}(z-z_{0})^{-2i\nu(z_{0})}(1-\Xi_{Z_{0}}), & z \in \Sigma_{7}, \end{cases}$$
(4.12)

These matrices R_j are estimated as follows

$$|R_j| \lesssim \sin^2(\arg(z-z_1)) + \langle Re(z) \rangle^{-\frac{1}{2}}, \qquad (4.13)$$

$$|\bar{\partial}R_j(z)| \leq |\bar{\partial}\Xi_1(z)| + |p'_j(Re(z))| + |z - z_1|^{-1/2},$$
(4.14)

for $j = 1, 6, 7^+, 8^+$, similarly for $j = 3, 4, 7^-, 8^-$, we have

$$|R_j| \lesssim \sin^2(\arg(z-z_0)) + \langle Re(z) \rangle^{-\frac{1}{2}}, \tag{4.15}$$

$$|\bar{\partial}R_j(z)| \le \left|\bar{\partial}\Xi_0(z)\right| + \left|p'_j(Re(z))\right| + |z - z_0|^{-1/2},\tag{4.16}$$

where

$$p_1(z) = p_3(z) = \gamma(z), \quad p_{7^+}(z) = p_{7^-}(z) = \frac{\gamma^*(z)}{1 + |\gamma(z)|^2},$$
 (4.17)

$$p_4(z) = p_6(z) = \gamma^*(z), \quad p_{8^+}(z) = p_{8^-}(z) = \frac{\gamma(z)}{1 + |\gamma(z)|^2},$$
 (4.18)

$$\langle (\cdot) \rangle = \sqrt{1 + (\cdot)^2}. \tag{4.19}$$

Employing $M^{(2)}(z)$ defined by $R^{(2)}(z)$, we can get a new RH problem.

RH Problem 4.3. The matrix-valued function $M^{(2)}(z)$ such that:

(*I*) Analyticity: $M^{(2)}(z)$ is continuous in complex \mathbb{C} , sectionally continuous first partial derivatives in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathbb{Z} \cup \overline{\mathbb{Z}})$.

(II) Jump condition:

$$M_{+}^{(2)}(z) = M_{-}^{(2)}(z)V^{(2)}(z)$$
(4.20)

where the jump matrix

$$V^{(2)}(z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ R_1 e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_1, \\ \begin{pmatrix} 1 & R_{7^+} e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_6, \\ \begin{pmatrix} 1 & 0 \\ R_{8^+} e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_8, \\ \begin{pmatrix} 1 & R_6 e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_4, \\ \begin{pmatrix} 1 & R_{7^-} e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_5, \\ \begin{pmatrix} 1 & R_3 e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_2, \\ \begin{pmatrix} 1 & R_4 e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_3, \\ \begin{pmatrix} 1 & 0 \\ R_8 - e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_7. \end{cases}$$

(III) Analyticity behavior:

$$M^{(2)}(z) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$$
 (4.21)

(*IV*) The $\bar{\partial}$ -derivative: For $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathcal{Z} \cup \overline{\mathcal{Z}})$

$$\bar{\partial}M^{(2)}(z) = M^{(1)}\bar{\partial}R^{(2)}(z),$$
 (4.22)

where

$$\bar{\partial}R^{(2)}(z) = \begin{cases} \begin{pmatrix} 1 & (-1)^m \bar{\partial}R_j(z)e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & j = 7^{\pm}, 4, 6, \\ \begin{pmatrix} 1 & 0 \\ (-1)^m \bar{\partial}R_j(z)e^{2it\theta} & 1 \end{pmatrix}, & j = 8^{\pm}, 1, 3, \\ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, & z \in \Omega_2 \cup \Omega_5, \end{cases}$$
(4.23)

and $m = \begin{cases} 1, j = 7^{\pm}, 1, 3, \\ 0, j = 8^{\pm}, 4, 6, \end{cases}$

(V) The residue conditions: $M^{(2)}$ has simple poles at the set $\mathcal{Z} \cup \overline{\mathcal{Z}}$

$$\begin{split} R_{z=z_{k}}^{es} M^{(2)} &= \begin{cases} \lim_{z \to z_{k}} M^{(2)} \begin{pmatrix} 0 & c_{k}^{-1} \left((\frac{1}{T})'(z_{k}) \right)^{-2} e^{-2it\theta} \\ 0 & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}} M^{(2)} \begin{pmatrix} 0 & 0 \\ c_{k}^{-1} T^{-2}(z_{k}) e^{2it\theta} & 0 \end{pmatrix}, \quad k \in \Delta_{z_{0}}^{+}, \end{cases} \end{split}$$

$$\begin{aligned} R_{es}^{es} M^{(2)} &= \begin{cases} \lim_{z \to z_{k}^{*}} M^{(2)} \begin{pmatrix} 0 & 0 \\ -\overline{c}_{k}^{-1} (T'(z_{k}^{*}))^{-2} e^{2it\theta^{*}(z)} & 0 \end{pmatrix}, \quad k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}^{*}} M^{(2)} \begin{pmatrix} 0 & -c_{k}^{*} (T(z_{k}^{*}))^{2} e^{-2it\theta^{*}(z)} \\ 0 & 0 \end{pmatrix}, \quad k \in \Delta_{z_{0}}^{+}. \end{cases} \end{split}$$

$$\end{split}$$

5. The decomposition of the mixed $\overline{\partial}$ -RH probelm

In this section, the pure RH problem in the mixed **RH problem** 4.3 is separated from setting $\overline{\partial}R^{(2)}(z) = 0$, that is, the soliton solution corresponding to the discrete spectrum is denoted as $M_{RHP}^{(2)}(z)$. We then establish the following RH problem

RH Problem 5.1. Find a matrix-valued function $M_{RHP}^{(2)}(z)$ admits the following properties:

(*I*) Analyticity: $M_{RHP}^{(2)}(z)$ is analytic in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathbb{Z} \cup \overline{\mathbb{Z}})$. (*II*) Analytic behavior:

$$M_{RHP}^{(2)}(z) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$$
(5.1)

(III) The jump condition:

$$M_{RHP,+}^{(2)}(z) = M_{RHP,-}^{(2)}(z)V^{(2)}(z), \quad z \in \mathbb{R}.$$
(5.2)

(*IV*) The $\bar{\partial}$ -derivative: $\overline{\partial}R^{(2)}(z) = 0$ for $z \in \mathbb{C}$.

(V) The residue conditions: $M_{RHP}^{(2)}(z)$ has simple poles at the set $\mathcal{Z} \cup \overline{\mathcal{Z}}$.

$$\begin{split} Res_{z=z_{k}}^{es} M_{RHP}^{(2)}(z) &= \begin{cases} \lim_{z \to z_{k}} M_{RHP}^{(2)}(z) \begin{pmatrix} 0 & c_{k}^{-1} \left(\left(\frac{1}{T}\right)'(z_{k}) \right)^{-2} e^{-2it\theta} \\ 0 & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}} M_{RHP}^{(2)}(z) \begin{pmatrix} 0 & 0 \\ c_{k}^{-1} T^{-2}(z_{k}) e^{2it\theta} & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{+}, \end{cases} \end{split}$$

$$\begin{aligned} Res_{z=z_{k}^{*}} M_{RHP}^{(2)}(z) &= \begin{cases} \lim_{z \to z_{k}^{*}} M_{RHP}^{(2)}(z) \begin{pmatrix} 0 & 0 \\ -\bar{c}_{k}^{-1} (T'(z_{k}^{*}))^{-2} e^{2it\theta^{*}(z)} & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{-}, \\ \lim_{z \to z_{k}^{*}} M_{RHP}^{(2)}(z) \begin{pmatrix} 0 & -c_{k}^{*} (T(z_{k}^{*}))^{2} e^{-2it\theta^{*}(z)} \\ 0 & 0 \end{pmatrix}, \ k \in \Delta_{z_{0}}^{+}. \end{cases} \end{aligned}$$

$$\begin{aligned} (5.3)$$

The existence and asymptotic of $M_{RHP}^{(2)}(z)$ will shown in next section. In virtue of $M_{RHP}^{(2)}(z)$, a new matrix function $M^{(3)}(z)$ is constructed

$$M^{(3)}(z) = M^{(2)}(z)M^{(2)}_{RHP}(z)^{-1},$$
(5.4)

which satisfies a pure $\overline{\partial}$ -problem.

RH Problem 5.2. The matrix $M^{(3)}(z)$ defined in (5.4) such that: (*I*) $M^{(3)}(z)$ is continuous with sectionally continuous first partial derivatives in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathbb{Z} \cup \mathbb{Z}^*)$. (*II*) For $z \in \mathbb{C}$, one has

$$\bar{\partial}M^{(3)}(z) = M^{(3)}(z)M^{(2)}_{RHP}(z)\bar{\partial}R^{(2)}M^{(2)}_{RHP}(z)^{-1} \triangleq M^{(3)}(z)W^{(3)}(z),$$
(5.5)

where $\bar{\partial}R^{(2)}(z)$ is defined by (4.23). (*III*) Asymptotic behavior:

$$M^{(3)}(z) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$$
 (5.6)

Proof. The analytic and asymptotic properties of $M^{(2)}(z)$ and $M^{(2)}_{RHP}(z)$ are exactly given in **RH Problem** 4.3 and **RH Problem** 5.1. According to Eq.(5.3), we can derive these properties of $M^{(3)}(z)$. Next, we will show that the matrix $M^{(3)}(z)$ has no jumps

through the real axis and has no poles. It follows $M^{(2)}(z)$ and $M^{(2)}_{RHP}(z)$ admit the same jump condition that

$$\begin{aligned} M_{-}^{(3)}(z)^{-1}M_{+}^{(3)}(z) &= M_{RHP,-}(z)M_{-}^{(2)}(z)^{-1}M_{+}^{(2)}(z)M_{RHP,+}(z)^{-1} \\ &= M_{RHP,-}(z)V^{(2)}(z)\left(M_{RHP,-}(z)V^{(2)}(z)\right)^{-1} = \mathbb{I} \end{aligned}$$

Note that for each $z_k \in \mathbb{Z} \cup \overline{\mathbb{Z}}$, $M^{(2)}(z)$ and $M^{(2)}_{RHP}(z)$ admit the same residue conditions, which appear in the left side of (5.3) denoted by a nilpotent matrix Q_k . Taking the Laurent expansions yields

$$M^{(2)}(z) = C(z_k) \left[\frac{Q_k}{z - z_k} + \mathbb{I} \right] + O(z - z_k),$$

$$M_{RHP}(z) = \widehat{C}(z_k) \left[\frac{Q_k}{z - z_k} + \mathbb{I} \right] + O(z - z_k),$$

where $C(z_k)$ and $\hat{C}(z_k)$ are constant terms, we then can derive that

$$M^{(2)}(z)M^{(2)}_{RHP}(z)^{-1} = O(1)$$

which means has removable singularities at z_k . From the definition of $M^{(3)}(z)$, it can be obtained by direct calculation

$$\begin{split} \bar{\partial}M^{(3)}(z) &= \bar{\partial}(M^{(2)}(z)M^{(2)}_{RHP}(z)^{-1}) = \bar{\partial}M^{(2)}(z)M^{(2)}_{RHP}(z)^{-1} = M^{(2)}(z)\bar{\partial}R^{(2)}(z)M^{(2)}_{RHP}(z)^{-1} \\ &= M^{(2)}(z)M^{(2)}_{RHP}(z)^{-1}(M^{(2)}_{RHP}(z)\bar{\partial}R^{(2)}(z)M^{(2)}_{RHP}(z)^{-1}) = M^{(3)}(z)W^{(3)}(z). \end{split}$$

Remark 5.3. We transform jump contour $\Sigma^{(2)}$ into contour $\Sigma^{(3)}$ shown in Fig. 6. By comparing the contours $\Sigma^{(2)}$ and $\Sigma^{(3)}$, we find that a new curve is introduced on contour $\Sigma^{(3)}$ with jump matrix defined by

$$v_{9} = \begin{cases} \mathbb{I}, & z \in \left(-i\frac{z_{1}-z_{0}}{2}\tan(\pi/12), i\frac{z_{1}-z_{0}}{2}\tan(\pi/12)\right), \\ \left(1 - (R_{7^{-}} - R_{7^{+}})e^{-2i\theta}\right), & z \in \left(i\frac{z_{1}-z_{0}}{2}\tan(\pi/12), iz_{1}\right), \\ \left(1 - (R_{7^{-}} - R_{7^{+}})e^{2i\theta}-1\right), & z \in \left(-iz_{1}, -i\frac{z_{1}-z_{0}}{2}\tan(\pi/12)\right). \end{cases}$$

Similar to [30], we can derive that

 $|v_9-\mathbb{I}| \lesssim e^{-t},$

which implies that the contour does not contribute to the long-time asymptotic solution as $t \to \infty$.




Figure 6. The contour of $\Sigma^{(3)}$.

To construct the solution of RH Problem 5.1, we define

$$M_{RHP}^{(2)}(z) = \begin{cases} E(z)M^{(out)}(z), & z \notin \{\mathcal{U}_0 \cup \mathcal{U}_1\}, \\ E(z)M^{(z_0)}(z), & z \in \mathcal{U}_0, \\ E(z)M^{(z_1)}(z), & z \in \mathcal{U}_1, \end{cases}$$
(5.7)

where \mathcal{U}_0 and \mathcal{U}_1 denote the neighborhoods of z_0 and z_1 , respectively,

$$\mathcal{U}_0 = \left\{ z : |z - z_0| \le \min\left\{\frac{z_0}{2}, \rho/3\right\} \triangleq \varepsilon \right\},$$

$$\mathcal{U}_1 = \left\{ z : |z - z_1| \le \min\left\{\frac{z_1}{2}, \rho/3\right\} \triangleq \varepsilon \right\},$$

which imply that $M_{RHP}^{(2)}(z)$, $M^{(z_0)}(z)$ and $M^{(z_1)}(z)$ have no poles in \mathcal{U}_0 and \mathcal{U}_1 from dist $(\mathcal{Z} \cup \overline{\mathcal{Z}}, R) > \rho$. The matrix $M_{RHP}^{(2)}(z)$ is divided into two parts by the decomposition: one can be called the external model RH problem denoted by $M^{(out)}(z)$, which can be solved directly by considering the standard RH problem under the condition of reflection-less potential. The other is in the neighborhood of the phase points $M^{(z_0)}(z)$ and $M^{(z_1)}(z)$, which can be matched to the known model, namely the parabolic cylinder model in \mathcal{U}_0 and \mathcal{U}_1 , to solve in section 8. In addition, the matrix E(z) is a error function, which can be solved by a small-norm RH problem in section 9.

Proposition 5.4. *The jump matrices defined by RH**Problem 5.1 satisfy the following estimates:*

For fixed $\epsilon > 0$, we define

$$\mathcal{L}_{\epsilon} = \left\{ z : z = z_1 + uz_1 e^{3i\pi/4}, \epsilon \le u \le \sqrt{2} \right\},$$

$$\cup \left\{ z : z = z_1 + uz_1 e^{i\pi/4}, \epsilon \le u \le \infty \right\},$$

$$\cup \left\{ z : z = z_0 + uz_0 e^{i\pi/4}, \epsilon \le u \le \sqrt{2} \right\},$$

$$\cup \left\{ z : z = z_0 + uz_0 e^{3i\pi/4}, \epsilon \le u \le \infty \right\},$$

then the estimates can be obtained

$$\left\| V^{(2)} - \mathbb{I} \right\|_{L^{\infty}(\Sigma_{+}^{(2)} \cap \mathcal{U}_{1})} = O\left(t^{-5/6} |z - z_{1}|^{-5/6} \right),$$
(5.8)

$$\left\|V^{(2)} - \mathbb{I}\right\|_{L^{\infty}(\Sigma_{-}^{(2)} \cap \mathcal{U}_{0})} = O\left(t^{-5/6} |z - z_{0}|^{-5/6}\right),\tag{5.9}$$

$$\left\|V^{(2)} - \mathbb{I}\right\|_{L^{\infty}(\Sigma^{(2)} \setminus (\mathcal{U}_1 \cup \mathcal{U}_0))} = O\left(e^{-2t\varepsilon}\right),\tag{5.10}$$

where the contours are defined by

$$\Sigma_{+}^{(2)} = \Sigma_1 \cup \Sigma_6 \cup \Sigma_8 \cup \Sigma_4, \quad \Sigma_{-}^{(2)} = \Sigma_5 \cup \Sigma_2 \cup \Sigma_3 \cup \Sigma_7.$$

The proposition implies that the jump matrix $V^{(2)}(z)$ uniformly goes to I on both $\Sigma^{(2)} \setminus (\mathcal{U}_1 \cup \mathcal{U}_0)$, in addition, outside the $\mathcal{U}_0 \cup \mathcal{U}_1$ there is only exponentially small error (in t) by completely ignoring the jump condition of $M^{(2)}_{RHP}(z)$.

6. Outer model RH problem

In this section, an external RH problem will be established and its solution can be approximated by a finite number of soliton solutions.

6.1. The existence of soliton solution

RH Problem 6.1. The matrix value function $M^{(out)}(x, t; z)$, satisfing (*I*) $M^{(out)}(x, t; z)$ is analytical in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathbb{Z} \cup \mathbb{Z}^*)$. (*II*) As $z \to \infty$,

$$M^{(out)}(x,t;z) = \mathbb{I} + O(z^{-1}).$$
(6.1)

(*III*) $M^{(out)}(x, t; z)$ has simple poles at each point in $\mathbb{Z} \cup \mathbb{Z}^*$ admitting the same residue condition in **RH Problem** 4.3 with $M^{(out)}(x, t; z)$ replacing $M^{(2)}(x, t; z)$.

Before proving the existence and uniqueness of the solution of the **RH Problem** 6.1, we first consider the case of the RH problem without reflection, where **RH Problem** 2.3 is reduced to the following RH problem

RH Problem 6.2. For the discrete date $\sigma_d = \{(z_k, c_k)\}_{k=1}^N$, and the set $\mathcal{Z} = \{z_k\}_{k=1}^N$, a new matrix valued function $m(x, t; z | \sigma_d)$ such that: (*I*) $m(x, t; z | \sigma_d)$ is analytic in $\mathbb{C} \setminus (\Sigma^{(2)} \cup \mathcal{Z} \cup \mathcal{Z}^*)$. (*II*) The asymptotic behaviour:

$$m(x,t;z|\sigma_d) \to \mathbb{I} + O(z^{-1}), \quad as \ z \to \infty.$$
 (6.2)

(III) Symmetry:

$$\overline{m(x,t;\overline{z}|\sigma_d)} = \sigma_2 m(x,t;z|\sigma_d)\sigma_2.$$
(6.3)

(III) The residue conditions: $m(x, t; z | \sigma_d)$ has simple poles at each point in $\mathbb{Z} \cup \mathbb{Z}^*$ satisfying

$$\begin{aligned} & \operatorname{Res}_{z=z_k} m(x,t;z|\sigma_d) = \lim_{z \to z_k} m(x,t;z|\sigma_d) Q_k, \\ & \operatorname{Res}_{z=z_k^*} m(x,t;z|\sigma_d) = \lim_{z \to z_k^*} m(x,t;z|\sigma_d) \sigma_2 Q_k^* \sigma_2, \end{aligned}$$
(6.4)

where Q_k is a nilpotent matrix

$$Q_k = \begin{pmatrix} 0 & 0 \\ \gamma_k(x,t) & 0 \end{pmatrix}, \quad \gamma_k(x,t) = c_k e^{2it\theta(z_k)},$$

$$\theta(z) = z \frac{x}{t} + 2\alpha z^2 + 4\beta z^3.$$
(6.5)

The uniqueness of **RH Problem** 6.2 is the direct result of Liouville theorem. From the symmetry (6.3), taking the following expansion

$$m(x,t;z|\sigma_d) = \mathbb{I} + \sum_{k=1}^{N} \left[\frac{1}{z - z_k} \begin{pmatrix} \zeta_k(x,t) & 0\\ \eta_k(x,t) & 0 \end{pmatrix} + \frac{1}{z - z_k^*} \begin{pmatrix} 0 & -\eta_k^*(x,t)\\ 0 & \zeta_k^*(x,t) \end{pmatrix} \right], \quad (6.6)$$

where $\zeta_k(x, t)$ and $\eta_k(x, t)$ are unknown coefficients to be determined. Similar to [41] we can prove the existence of the solution for the **RH Problem** 6.2.

Note that the trace formula can be written as under the reflection-less case

$$s_{11}(z) = \prod_{k=1}^{N} \frac{z - z_k}{z - z_k^*}.$$
(6.7)

Let $\triangle \subseteq \{1, 2, \dots, N\}$ and define

$$s_{11,\Delta}(z) = \prod_{k \in \Delta} \frac{z - z_k}{z - z_k^*}.$$
(6.8)

Taking the following transformation, we obtain a new matrix valued function $m^{\Delta}(z|D)$ in which the poles in the column are divided according to the choice of Δ

$$m^{\Delta}(z|D) = m(z|\sigma_d)(z)s_{11,\Delta}(z)^{\sigma_3},$$
 (6.9)

where the scattering data

$$D = \{(z_k, c'_k)\}_{k=1}^N, \quad c'_k = c_k s_{11,\Delta}(z)^2.$$
(6.10)

RH Problem 6.3. For the scattering data defined by (6.10), the matrix $m^{\Delta}(z|D)$ satisfies (*I*) Analyticity: $m^{\Delta}(x, t; z|D)$ is analytic in $\mathbb{C} \setminus (\mathcal{Z} \bigcup \mathcal{Z}^*)$. (*II*) Symmetry:

$$\overline{m^{\Delta}(x,t;z|D)} = \sigma_2 m(x,t;z|\sigma_d)\sigma_2.$$
(6.11)

(*III*) Asymptotic behavior: $m^{\Delta}(x, t; z|D) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$

(*IV*) The residue conditions: $m^{\Delta}(x, t; z|D)$ has simple poles at the set $\mathcal{Z} \cup \mathcal{Z}^*$

$$\begin{aligned} & \operatorname{Res}_{z=z_k} m^{\Delta}(x,t;z|D) = \lim_{z \to z_k} m(x,t;z|\sigma_d) Q_k^{\Delta}, \\ & \operatorname{Res}_{z=z_k^*} m^{\Delta}(x,t;z|D) = \lim_{z \to z_k^*} m(x,t;z|\sigma_d) \sigma_2 Q_k^{\Delta*} \sigma_2, \end{aligned}$$
(6.12)

where $Q_k^{\scriptscriptstyle \Delta}$ is a nilpotent matrix

$$\mathbf{Q}_{k}^{\Delta} = \begin{cases} \begin{pmatrix} 0 & \gamma_{k}^{\Delta} \\ 0 & 0 \end{pmatrix}, & k \in \Delta, \\ \begin{pmatrix} 0 & 0 \\ \gamma_{k}^{\Delta} & 0 \end{pmatrix}, & k \notin \Delta, \end{cases} \qquad \gamma_{k}^{\Delta} = \begin{cases} c_{k}^{-1}(s_{11,\Delta}^{'}(z_{k}))^{-2}e^{-2it\theta(z_{k})}, & k \in \Delta, \\ c_{k}(s_{11,\Delta}(z_{k}))^{2}e^{2it\theta(z_{k})}, & k \notin \Delta, \end{cases}$$

$$\theta(z) = z\frac{x}{t} + 2\alpha z^{2} + 4\beta z^{3}.$$
(6.13)

Remark 6.4. It is noticed that the solution of **RH Problem** 6.2 is existent and unique, and it follows from the transformation (6.9) that the solution of **RH Problem** 6.3 is also existent and unique.

In **RH Problem** 6.3, taking $\triangle = \triangle_{z_1}^-$ and replacing the scattering data *D* with

$$\widetilde{D} = \{(z_k, \widetilde{c}_k)\}_{k=1}^N, \quad \widetilde{c}_k = c_k \delta(z_k)^2, \tag{6.14}$$

we then obtain

Corollary 6.5. There exists a unique solution to the RH Problem 6.3 satisfying

$$M^{(out)}(z)(x,t;z) = m^{\Delta_{z_1}^-}(x,t;z|\tilde{D}),$$
(6.15)

where the scattering data \widetilde{D} is defined by (6.6).

Corollary 6.6. The soliton solution of the Hirota equation (1.1) is determined by

$$q_{sol}(x,t) = 2i \lim_{z \to \infty} (zM(z))_{12} = 2i \lim_{z \to \infty} (zm(x,t;z|\sigma_d))_{12}$$

= $2i \lim_{z \to \infty} \left(zm^{\Delta_{z_1}^-} (z|\sigma_d^{\Delta_{z_1}^-}) s_{11,\Delta_{z_1}^-} (z)^{-\sigma_3} \right)_{12}$
= $2i \lim_{z \to \infty} \left((zm^{\Delta_{z_1}^-} z|\sigma_d^{\Delta_{z_1}^-}) \right)_{12} = 2i \lim_{z \to \infty} (zM^{(out)}(z))_{12}.$ (6.16)

6.2. The long-time behavior of soliton solution

Taking N = 1, $\sigma_d = \{z_1 = \xi + i\eta, c_1\}$, and employing the residue conditions (6.12) we have the 1-soliton solution

$$q_{sol}(x,t) = 2\eta \operatorname{sech} \left(2\eta (x + 4\alpha\xi t + 4\beta(3\xi\eta^2 - \eta^2)t + x_0) \right) h(x,t).$$
(6.17)

where $x_0 = -\frac{1}{2\eta} \log\left(\frac{|c_1|}{2\eta}\right)$, and $h(x,t) = e^{-2i\left[x\xi + 2\alpha(\eta^2 - \eta^2)t + 4\beta(\xi^3 - 3\xi\eta^2)\right] - i(\frac{\pi}{2} + arg(c_1))}$, from which the speed of the soliton solution is

$$v_1 = -4 \left[\alpha Rez_1 + \beta Imz_1^2 (3Rez_1 - 1) \right].$$
(6.18)

For given $x_1 \leq x_2$, and velocities $v_1 \leq v_2$ with $x_1, x_2, v_1, v_2 \in \mathbb{R}$, we define a cone and obtain the discrete spectral distribution shown in Fig. 7 and Fig. 8.

$$C(x_1, x_2, v_1, v_2) = \{(x, t), x = x_0 + vt \text{ with } x_0 \in [x_1, x_2], v \in [v_1, v_2]\}.$$
(6.19)




Figure 7. Space-time $C(x_1, x_2, v_1, v_2)$.




Figure 8. For fixed $v_1 < v_2$, $I = [f(v_2), f(v_1)]$.

In additional, we denote

$$I = \{z : f(v_2) < |z| < f(v_1)\}, \quad f(v_m) = -\left(\frac{v_m}{4\alpha} + \frac{\beta}{\alpha}Imz_m^2(3Rez_m - 1)\right),$$

$$\mathcal{K}(I) = \{z_j \in \mathcal{K} : z_j \in I\}, \quad N(I) = |\mathcal{K}(I)|,$$

$$\mathcal{K}_+(I) = \{z_j \in \mathcal{K} : |z_j| > f(v_1)\},$$

$$\mathcal{K}_-(I) = \{z_j \in \mathcal{K} : |z_j| < f(v_2)\},$$

$$c_j^{\pm}(I) = c_j \prod_{Rez_n \in I_{\pm} \setminus I} \left(\frac{z_j - z_n}{z_j - z_n^*}\right)^2 \exp\left[\pm \frac{1}{\pi i} \int_{I_{\pm}} \frac{\log(1 + |\gamma(\zeta)|^2)}{\zeta - z_j} d\zeta\right].$$

(6.20)

Proposition 6.7. The choice of $\triangle = \triangle_{z_1}^{\mp}$ in **RH Problem** 6.3 ensures the following estimation with $(x, t) \in C(x_1, x_2, v_1, v_2)$ as $t \to \infty$

$$\left\| \boldsymbol{Q}_{k}^{\boldsymbol{\Delta}_{z_{1}}^{\pi}} \right\| = \begin{cases} O(1), & z_{k} \in \mathcal{K}(I), \\ O(e^{-8\mu t}), & z_{k} \in \mathcal{K} \setminus \mathcal{K}(I). \end{cases}$$
(6.21)

where $\mu(\mathcal{I}) = \min_{z_k \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})} \{Im(z_k) \cdot dist(Rez_k, I)\}.$

Proof. Taking $\triangle = \triangle_{z_1}^-$ in **RH Problem** 6.3, for $z_j \in \mathcal{K}_-(\mathcal{I})$ and $(x, t) \in C(x_1, x_2, v_1, v_2)$ we have

$$\left\| \boldsymbol{Q}^{\boldsymbol{\Delta}_{\boldsymbol{z}_{1}}^{-}} \right\| \lesssim |\boldsymbol{e}^{-2it\theta(\boldsymbol{z}_{j})}|, \tag{6.22}$$

with

$$-2it\theta(z_j) = -2i(x_0 z_j + v t z_j + 2\alpha z_j^2 + 4\beta z_j^3),$$
(6.23)

which implies that

$$Re(-2it\theta(z_j)) = 2Imz_j t \left[x_0 + vt + 4\alpha Rez_j + 4\beta Imz_j (3Rez_j - 1) \right].$$
(6.24)

Then we can obtain

$$\left\| Q^{\Delta_{z_1}^-} \right\| = O(e^{-8\mu t}). \tag{6.25}$$

Based on the above estimates, the following results are derived.

Proposition 6.8. For the reflection-less data $D = \{(z_j, c_j)\}_{j=1}^N$, $D^{\pm}(I) = \{(z_j, c_j^{\pm}(I)) | z_j \in \mathcal{K}(I)\}$, we have the following relationship with $(x, t) \in C(x_1, x_2, v_1, v_2)$ as $t \to \infty$

$$m^{\Delta_{z_1}^{\mp}}(x,t;z|D) = \left(\mathbb{I} + O(e^{-8\mu t})\right) m^{\Delta_{z_1}^{\mp}}(x,t;z|D^{\pm}(I)),$$
(6.26)

with

$$c_{j}^{\pm}(I) = c_{j} \prod_{Rez_{n} \in I_{\pm \setminus I}} \left(\frac{z_{j} - z_{n}}{z_{j} - z_{n}^{*}} \right)^{2}.$$
 (6.27)

Proof. For each $z_j \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})$, suppose S_j is a circle centered on z_j and the radius is smaller than $\mu(\mathcal{I})$. ∂S is the boundary of S. We introduce a transformation to transform the residue conditions at each $z_j \in \mathcal{K} \setminus \mathcal{K}(\mathcal{I})$ into a jump condition along the circle S_j .

$$\widetilde{m}^{\Delta_{z_1}^-}(z|D) = \begin{cases} m^{\Delta_{z_1}^-}(z|D) \left(\mathbb{I} - \frac{Q_j}{z - z_j} \right), & z \in S_j, \\ m^{\Delta_{z_1}^-}(z|D) \left(\mathbb{I} - \frac{\sigma_2 Q_j \sigma_2}{z - z_j^*} \right), & z \in S_j, \\ m^{\Delta_{z_1}^-}(z|D), & \text{elsewhere.} \end{cases}$$
(6.28)

The new matrix $\widetilde{m}^{\Delta_{z_1}^-}(z|D)$ has new jump condition in each ∂S_i denoted by $\widetilde{V}(z)$

$$\widetilde{m}_{+}^{\Delta_{z_{1}}^{-}}(z|D) = \widetilde{m}_{-}^{\Delta_{z_{1}}^{-}}(z|D)\widetilde{V}(z), \qquad z \in \widetilde{\Sigma},$$
(6.29)

where

$$\widetilde{\Sigma} = \bigcup_{z_j \in \mathcal{K} \setminus \mathcal{K}(I)} (\partial S_j \cup \partial S_j^*).$$

Using (6.28) and Proposition 6.8 leads to

$$\left\|\widetilde{V}(z) - \mathbb{I}\right\|_{L^{\infty}(\widetilde{\Sigma})} = O(e^{-8\mu t}).$$
(6.30)

It follows that from $\widetilde{m}_{+}^{\Delta_{z_1}^-}(z|D)$ and $m_{+}^{\Delta_{z_1}^-}(z|D)$ has the same poles and residue conditions

$$\epsilon(z) = \widetilde{m}^{\tilde{\Delta}_{z_1}}(z|D)m^{\tilde{\Delta}_{z_1}(I)}(z|D^{\pm}(I))^{-1}$$
(6.31)

has no poles, but the jump condition for $\epsilon(z)$ can be written as as $z \in \widetilde{\Sigma}$

$$\epsilon_{+}(z) = \epsilon_{-}(z)V_{\epsilon}(z), \tag{6.32}$$

where the jump matrix

$$V_{\epsilon}(z) = m(z|D^{\pm}(\mathcal{I}))\widetilde{V}(z)m(z|D^{\pm}(\mathcal{I}))^{-1}$$

from which together with Eq.(6.30) yields

$$\left\| V_{\epsilon}(z) - \mathbb{I} \right\|_{L^{\infty}(\widetilde{\Sigma})} = O(e^{-8\mu t}), \quad t \to \infty.$$
(6.33)

Corollary 6.9. Suppose that $m_{sol}(x, t; D)$ and $m_{sol}(x, t; D(I))$ represent the N-soliton solutions of the Hirota equation (1.1) corresponding to scattering data D and D(I), respectively. For given data I, D(I) and $C(x_1, x_2, v_1, v_2)$, the relationship can be derived

$$2i \lim_{z \to \infty} z(m(x,t;z|\sigma_d))_{12} = m_{sol}(x,t;D) = m_{sol}(x,t;D(I)) + O(e^{-8\mu t}).$$
(6.34)

Furthermore the outer model can be expressed as

Corollary 6.10. The **RH Problem** 6.1 exists an unique solution $M^{(out)}(z)$ satisfying

$$M^{(out)}(z) = m^{\Delta_{z_1}^-}(z|D^{(out)})$$

= $m^{\Delta_{z_1}^-}(x, t, z|D(I)) \prod_{Rez_n \in I_{\pm} \setminus I} \left(\frac{z_j - z_n}{z_j - z_n^*}\right)^{-\sigma_3} \delta^{-\sigma_3} + O(e^{-8\mu t}),$ (6.35)

where $D^{(out)} = \left\{z_j, c_j(z_1)\right\}_{j=1}^N$ with

$$c_j(z_1) = c_j exp\left[-\frac{1}{\pi i} \int_{I_+} \frac{\log(1+|\gamma(\varsigma)|^2)}{\varsigma-z} d\varsigma\right].$$

We further obtain the reconstruction formula

$$2i \lim_{z \to \infty} z(M^{(out)})_{12} = m(x, t; D^{(out)}) = m(x, t; D^{(out)}(I)) + O(e^{-8\mu t}),$$
(6.36)

with

$$m_{sol}(x,t;D^{(out)}) = m_{sol}(x,t;D(I)) + O(e^{-8\mu t}), \quad t \to \infty.$$
 (6.37)

7. A local RH problem near phase points

From the **Proposition** 5.4, we know that $V^{(2)} - \mathbb{I}$ doesn't have a uniformly small jump in the neighborhood \mathcal{U}_{z_0} and \mathcal{U}_{z_1} , therefore we establish a local model for the error function E(z) with a uniformly small jump. In addition, in this section, we separate the pure RH problem from the mixed RH problem and obtain the pure $\overline{\partial}$ problem.

RH Problem 7.1. Find a matrix-valued function $M^{(HR)}(x, t; z)$ satisfying:

- (*I*) Analyticity: $M^{(HR)}(x, t; z)$ is analytic in $\mathbb{C} \setminus \Sigma^{(2)}$.
- (II) Asymptotic behavior:

$$M^{(HR)}(x,t;z) = \mathbb{I} + O(z^{-1}), \quad z \to \infty.$$

$$(7.1)$$

(III) Jump condition: $M^{(HR)}(x, t; z)$ has continuous boundary values on $\Sigma^{(2)}$ and

$$M_{+}^{(HR)}(x,t;z) = M_{-}^{(HR)}(x,t;z)V^{(HR)}(z), \quad z \to \infty.$$
(7.2)

where the jump matrix is determined by

$$V^{(HR)}(z) = \begin{cases} \begin{pmatrix} 1 & 0 \\ \gamma(z_1)\delta^{-2}(z_1)(z-z_1)^{-2i\nu(z_1)}e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_1, \\ \begin{pmatrix} 1 & \frac{\gamma^*(z_1)\delta^{-2}(z_1)}{1+|\gamma(z_1)|^2}(z-z_1)^{2i\nu(z_1)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_6, \\ \begin{pmatrix} 1 & 0 \\ \frac{\gamma(z_1)\delta^{-2}(z_1)}{1+|\gamma(z_1)|^2}(z-z_1)^{-2i\nu(z_1)}e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_8, \\ \begin{pmatrix} 1 & \gamma^*(z_1)\delta^2(z_1)(z-z_1)^{2i\nu(z_1)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_4, \\ \begin{pmatrix} 1 & \frac{\gamma^*(z_0)\delta^2(z_0)}{1+|\gamma(z_0)|^2}(z-z_0)^{2i\nu(z_0)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_5, \\ \begin{pmatrix} 1 & 0 \\ \gamma(z_0)\delta^{-2}(z_0)(z-z_0)^{-2i\nu(z_0)}e^{2it\theta} & 1 \end{pmatrix}, & z \in \Sigma_2, \\ \begin{pmatrix} 1 & \gamma^*(z_0)\delta^2(z_0)(z-z_0)^{2i\nu(z_0)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_3, \\ \begin{pmatrix} 1 & \gamma^*(z_0)\delta^{-2}(z_0)(z-z_0)^{-2i\nu(z_0)}e^{-2it\theta} \\ 0 & 1 \end{pmatrix}, & z \in \Sigma_7. \end{cases}$$

To solve the RH problem, we need to consider the parabolic cylinder (PC) model with two stationary-phase points shown in Fig. 9.




Figure 9. The jump contour for the local RH problem near z_0 and z_1 .

Taking z_1 as an example, the other stationary-phase point can be derived similarly. The jump contour of the standard parabolic cylindrical function is shown in **Appendix A** in detail.

Expanding the function $\theta(z)$, one has

$$\theta(z) = 4\beta(z-z_1)^3 + (12\beta z_1 + 2\alpha)(z-z_1)^2 - 8\beta z_1^3 - 2\alpha z_1^2.$$
(7.3)

To match the standard PC-model, we first do the following scaling transformation

$$N_A: f(z) \to (N_A f)(z) = f\left(\frac{\lambda}{\sqrt{8(6\beta z_1 + \alpha)t}} + z_1\right).$$
(7.4)

Setting

$$\gamma_0 = \gamma(z_1) T_0^{-2}(z_1) e^{2i\nu \log \sqrt{8(\alpha + 6\beta z_1)}} e^{2it \left[4\beta(z - z_1)^3 - 2z_1^2(\alpha + 4\beta z_1)\right]},$$
(7.5)

similar to [41], we have

$$M_{z_1}^{PC}(\lambda) = \mathbb{I} + \frac{M_1^{PC}(z_1)}{i\lambda} + O(\lambda^{-2}),$$
(7.6)

where

$$M_1^{PC}(z_1) = \begin{pmatrix} 0 & \beta_{12}(\gamma_{z_1}) \\ \beta_{21}(\gamma_{z_1}) & 0 \end{pmatrix},$$
(7.7)

with

$$\beta_{12}(\gamma_{z_1}) = \frac{\sqrt{2\pi}e^{i\pi/4}e^{-\pi\nu(z_1)/2}}{\gamma_{z_1}\Gamma(-i\nu(z_1))}, \quad \beta_{21}(\gamma_{z_1}) = \frac{-\sqrt{2\pi}e^{-i\pi/4}e^{-\pi\nu(z_1)/2}}{r_{z_1}^*\Gamma(i\nu(z_1))} = \frac{\nu}{\beta_{12}(\gamma_{z_1})}.$$

Remark 7.2. For the other stationary-phase point z_0 , in the same way, we take the following scaling transformation

$$N_B: f(z) \to (N_B f)(z) = f\left(\frac{\lambda}{\sqrt{8t(6\beta z_1 + \alpha)}} + z_0\right).$$
(7.8)

Taking

$$M_{z_0}^{PC}(\lambda) = \mathbb{I} + \frac{M_1^{PC}(z_0)}{i\lambda} + O(\lambda^{-2}),$$
(7.9)

we have

$$M_1^{PC}(z_0) = \begin{pmatrix} 0 & \beta_{12}(\gamma_{z_0}) \\ \beta_{21}(\gamma_{z_0}) & 0 \end{pmatrix},$$
(7.10)

with

$$\beta_{12}(\gamma_{z_0}) = \frac{\sqrt{2\pi}e^{i\pi/4}e^{-\pi\nu(z_0)/2}}{\gamma_{z_0}\Gamma(-i\nu(z_0))}, \quad \beta_{21}(\gamma_{z_0}) = \frac{-\sqrt{2\pi}e^{-i\pi/4}e^{-\pi\nu(z_0)/2}}{\gamma_{z_0}^*\Gamma(i\nu(z_0))} = \frac{\nu}{\beta_{12}(\gamma_{z_0})}.$$

The matrix-valued $M^{(HR)}(x, t; z)$ satisfies the asymptotic behavior

$$M^{(HR)}(x,t;z) = \mathbb{I} + \frac{1}{\lambda} \left(M_A^{(PC)} + M_B^{(PC)} \right) + O(\lambda^{-2}),$$
(7.11)

substituting (7.4) into (7.11) yields

$$M^{(HR)}(x,t;z) = \mathbb{I} + \frac{1}{\sqrt{8t(6\beta z_1 + \alpha)}} \frac{M_{1A}^{(PC)}(z_1)}{z - z_1} + \frac{1}{\sqrt{8t(6\beta z_0 + \alpha)}} \frac{M_{1B}^{(PC)}(z_0)}{z - z_0}.$$
 (7.12)

For the local circular domain of z_n (n = 0, 1), there exists a constant c, which is independent of z such that

$$\left|\frac{1}{z-z_n}\right| < c,\tag{7.13}$$

We can obtain a consistent estimate

$$\left| M^{(HT)} - \mathbb{I} \right| \lesssim O(t^{-1/2}), \tag{7.14}$$

which implies that

$$\|M^{(HT)}\|_{\infty} \lesssim 1.$$
 (7.15)

By using $M^{(HT)}$ to define the local model in the circles $z \in \mathcal{U}_{z_0}$ and $z \in \mathcal{U}_{z_1}$

$$M^{(z_0,z_1)}(z) = M^{(out)}M^{(HT)},$$
(7.16)

which is a bounded function in the \mathcal{U}_{z_0} and \mathcal{U}_{z_1} , and has the same jump matrix as $M_{RHP}^{(2)}$.

8. The small norm RH problem for the error function

From the definition (5.7) and (7.16), we find a error function E(z) satisfying

RH Problem 8.1. The matrix-valued function E(z) such that:

(*I*) Analyticity: E(z) is analytic in $\mathbb{C} \setminus \Sigma^{(E)}$, where

$$\Sigma^{(E)} = \partial \mathcal{U}_{z_0} \cup \partial \mathcal{U}_{z_1} \cup (\Sigma^{(2)} \setminus (\mathcal{U}_{z_0} \cup \mathcal{U}_{z_1}))$$

(II) Symmetry: $\overline{E(\overline{z})} = \sigma_2 E(z)\sigma_2$.

(III) Jump condition: E(z) has the continuous boundary value on $\Sigma^{(E)}$

$$E_{+}(z) = E_{-}(z)V^{(E)}(z), \qquad (8.1)$$

where the jump matrix $V^{(E)}(z)$

$$V^{(E)}(z) = \begin{cases} M^{(out)}(z)V^{(2)}(z)M^{(out)}(z)^{-1}, & z \in \Sigma^{(2)} \setminus (\mathcal{U}_{z_0} \cup \mathcal{U}_{z_1}), \\ M^{(out)}(z)M^{(HT)}(z)M^{(out)}(z)^{-1}, & z \in \partial \mathcal{U}_{z_0} \cup \partial \mathcal{U}_{z_1}. \end{cases}$$
(8.2)

(IV) Asymptotic behavior:

$$E(z) = \mathbb{I} + \frac{E_1(z)}{z} + O(^{-2}), \tag{8.3}$$

Note that from the **Proposition 5.4**, we have the estimate as follow

$$\left\|V^{(E)} - \mathbb{I}\right\|_{L^{\infty}(\Sigma^{(2)} \setminus (\mathcal{U}_1 \cup \mathcal{U}_0))} \lesssim O\left(e^{-2t\varepsilon}\right).$$
(8.4)

Next we will show that the error function E(z) can solve the small norm RH problem for a sufficiently large time.

Proposition 8.2. For $z \in \partial \mathcal{U}_{z_0} \cup \partial \mathcal{U}_{z_1}$, $M^{(out)}$ is bounded, the following estimate can be obtained

$$\left| V^{(E)} - \mathbb{I} \right| = \left| M^{(out)}(z)^{-1} (M^{(HT)} - \mathbb{I}) M^{(out)}(z) \right| = O(t^{-1/2}).$$
(8.5)

The **RH Problem** 8.1 then exists a unique solution which can be expressed by a small norm RH problem

$$E(z) = \mathbb{I} + \frac{1}{2\pi i} \int_{\Sigma^{(E)}} \frac{(\mathbb{I} + \rho(s))(V^{(E)} - \mathbb{I})}{s - z} ds,$$
(8.6)

where $\rho \in L^2(\Sigma^{(E)})$ is the unique solution of the following formula

$$(\mathbb{I} - C_E)\rho = C_E(\mathbb{I}),\tag{8.7}$$

where C_E is a integral operator defined by

$$C_E f(z) = C_-(f(V^{(E)} - \mathbb{I})),$$
 (8.8)

with the Cauchy projection operator on $\Sigma^{(E)}$

$$C_{-}(f(s)) = \lim_{z \to \Sigma_{-}^{(E)}} \frac{1}{2\pi i} \int_{\Sigma^{(E)}} \frac{f(s)}{s - z} ds.$$

Combining with (8.5), we have $||C_E|| < 1$ for the large *t*, further $1 - C_E$ is invertible, and ρ exists and is unique. In additional

$$\|\rho\|_{L^{2}(\Sigma^{(E)})} \lesssim \frac{\|C_{E}\|}{1 - \|C_{E}\|} \lesssim |t|^{-1/2}.$$
(8.9)

Then the existence and boundedness of function E(z) can be obtained. To construct the solution q(x, t), the asymptotic behavior of function E(z) needs to be considered.

Proposition 8.3. As $z \to \infty$, the asymptotic behavior of function E(z) can be expressed by

$$E(z) = \mathbb{I} + \frac{E_1}{z} + O(z^{-2}), \qquad (8.10)$$

where

$$E_{1} = \frac{1}{\sqrt{8(6\beta z_{1} + \alpha)}} \left\{ M^{(out)}(z_{1}) \begin{pmatrix} 0 & \beta_{12}(\gamma_{z_{1}}) \\ -\beta_{21}(\gamma_{z_{1}}) & 0 \end{pmatrix} M^{(out)}(z_{1})^{-1} \right\} + \frac{1}{\sqrt{8(6\beta z_{0} + \alpha)}} \left\{ M^{(out)}(z_{0}) \begin{pmatrix} 0 & \beta_{12}(\gamma_{z_{0}}) \\ -\beta_{21}(\gamma_{z_{0}}) & 0 \end{pmatrix} M^{(out)}(z_{0})^{-1} \right\}.$$
(8.11)

Proof. Employing the relationships (8.7), (8.9) and the estimate (8.4), we have

$$E_{1} = -\frac{1}{2\pi i} \oint_{\mathcal{U}_{z_{0}} \cup \mathcal{U}_{z_{1}}} (V^{(E)}(s) - I)ds + O(t^{-1}),$$

$$= \frac{1}{i\sqrt{8(6\beta z_{0} + \alpha)}} M^{(out)}(z_{0}) M_{1}^{(PC)}(z_{0}) M^{(out)}(z_{0})^{-1}$$

$$+ \frac{1}{i\sqrt{8(6\beta z_{1} + \alpha)}} M^{(out)}(z_{1}) M_{1}^{(PC)}(z_{1}) M^{(out)}(z_{1})^{-1} + O(t^{-1}).$$
(8.12)

Taking the notation

$$2i(E_1)_{12} = t^{-1/2} f(x, t) + O(t^{-1}),$$
(8.13)

where

$$f(x,t) = \frac{1}{\sqrt{2(6\beta z_1 + \alpha)}} \left(\beta_{12}(\gamma_{z_1}) M_{11}^{(out)}(z_1)^2 + \beta_{21}(\gamma_{z_1}) M_{12}^{(out)}(z_1)^2 \right) + \frac{1}{\sqrt{2(6\beta z_0 + \alpha)}} \left(\beta_{12}(\gamma_{z_0}) M_{11}^{(out)}(z_0)^2 + \beta_{21}(\gamma_{z_0}) M_{12}^{(out)}(z_0)^2 \right).$$

$$(8.14)$$

9. Pure $\bar{\partial}$ -Problem

In this section we will consider the long-time asymptotic behavior of $M^{(3)}(z)$, and the solution of the RH Problem 5.2 can be expressed by the following integral equation

$$M^{(3)}(z) = \mathbb{I} - \frac{1}{\pi} \int_{\mathbb{C}} \frac{\partial M^{(3)}(s)}{z - s} dA(s) = \mathbb{I} - \frac{1}{\pi} \int_{\mathbb{C}} \frac{M^{(3)}(s) W^{(3)}(s)}{z - s} dA(s),$$
(9.1)

where dA(s) is Lebesgue measure on the contour \mathbb{C} . Denoting C_z is the left Cauchy-Green integral equation, i.e.,

$$fC_z(z) = -\frac{1}{\pi} \int_{\mathbb{C}} \frac{f(s)W^{(3)}(s)}{z-s} dA(s).$$
(9.2)

we transform equation (9.1) into a compact form

$$(\mathbb{I} - C_z)M^{(3)}(z) = \mathbb{I},$$
 (9.3)

which is equivalent to

$$M^{(3)}(z) = \mathbb{I} \cdot (\mathbb{I} - C_z)^{-1}.$$
(9.4)

Regarding the existence of operator $(\mathbb{I} - C_z)^{-1}$, we give the following proposition

Proposition 9.1. The norm of the Cauchy-Green integral operator C_z decays to zero as $t \to \infty$

$$\|C_z\|_{L^{\infty} \to L^{\infty}} \lesssim t^{-1/4}, \tag{9.5}$$

which indicates that $(\mathbb{I} - C_z)^{-1}$ exists.

Proof. Taking the region Ω_1 (s = u + iv, z = x + iy) with $|u| \ge |v|$ as an example, the other regions are similarly to discuss. For any $f \in L^{\infty}$, we have

$$\begin{split} \|fC_{z}(z)\|_{L^{\infty}} &\leq \|f\|_{L^{\infty}} \frac{1}{\pi} \int_{\Omega_{1}} \frac{|W^{(3)}(s)|}{|z-s|} dA(s) \\ &\leq \|f\|_{L^{\infty}} \frac{1}{\pi} \int_{\Omega_{1}} \frac{|\overline{\partial}R^{(2)}(s)|}{|z-s|} dA(s), \end{split}$$
(9.6)

by using the fact

$$\left| W^{(3)}(s) \right| \le \left\| M^{(2)}_{RHP} \right\|_{L^{\infty}} \left| \overline{\partial} R^{(2)}(s) \right| \left\| M^{(2)}_{RHP} \right\|_{L^{\infty}}^{-1} \le \left| \overline{\partial} R^{(2)}(s) \right|.$$

$$(9.7)$$

Obviously we only consider the integral equation

$$\frac{1}{\pi} \int_{\Omega_1} \frac{|\overline{\partial}R^{(2)}(s)|}{|z-s|} dA(s), \tag{9.8}$$

from which together with (4.14) and

$$\begin{aligned} Re(2it\theta) &= 8it[4\beta(3iu^2v - iv^3) + 2i(12\beta z_1 + 2\alpha)uv] \\ &\lesssim 8t[4\beta(-3u^2v + u^2v) - 2(12\beta z_1 + 2\alpha)uv] \\ &\lesssim 8t[-8\beta u^2v - 24\beta z_1uv] \\ &\lesssim -8tz_1uv, \end{aligned}$$

we have

$$\int_{\Omega_1} \frac{|\overline{\partial}R^{(2)}(s)|}{|z-s|} dA(s) \lesssim I_1 + I_2 + I_3, \tag{9.9}$$

where

$$I_1 = \int_0^\infty \int_v^\infty \frac{1}{|s-z|} |\overline{\partial}\Xi_1| e^{-tz_1 u v} du dv, \qquad (9.10)$$

$$I_2 = \int_0^\infty \int_v^\infty \frac{1}{|s-z|} |p_1'(Rez)| e^{-tz_1 u v} du dv,$$
(9.11)

$$I_3 = \int_0^\infty \int_v^\infty \frac{1}{|s-z|} |s-z_1|^{-1/2} e^{-tz_1 u v} du dv.$$
(9.12)

Through direct calculation we can get the following estimate

$$\left\|\frac{1}{|s-z|}\right\|_{L^{\infty}} = \left(\int_{v}^{\infty} \frac{1}{(u-x)^{2} + (v-y)^{2}} du\right)^{1/2} \le \frac{\pi}{|v-y|}.$$
(9.13)

It follows from the **Appendix B** that

$$|I_1|, |I_2|, |I_3| \leq t^{-1/4}.$$
 (9.14)

It follows that

$$\int_{\Omega_1} \frac{|\overline{\partial}R^{(2)}(s)|}{|z-s|} dA(s) \lesssim t^{-1/4},$$
(9.15)

which implies that the inequality (9.5) holds.

In what follows, we consider the asymptotic behavior of $M^{(3)}(z)$

$$M^{(3)}(z) = \mathbb{I} + \frac{M_1^{(3)}(x,t)}{z} + O(z^{-2}),$$
(9.16)

where

$$M_1^{(3)}(x,t) = \frac{1}{\pi} \int_C M^{(3)}(s) W^{(3)}(s) dA(s).$$
(9.17)

In addition, $M_1^{(3)}(x, t)$ satisfies the estimate

Proposition 9.2. For $t \neq 0$, the matrix $M_1^{(3)}(x, t)$ such that

$$\left|M_{1}^{(3)}(x,t)\right| \lesssim t^{-3/4},$$
(9.18)

Proof. Similar to the proof of **Proposition** 9.1, we only consider the region $\Omega_1 = \{(u + z_1, v) : v \ge 0, v \le u < \infty\}$, the other case can be proved in the same way. Note that $M^{(3)}$ and $M^{(2)}_{RHP}$ are bounded, and employing formulas (9.17) and (9.18), we obtain

$$\left\|M_{1}^{(3)}(x,t)\right\| \leq \frac{1}{\pi} \iint_{\Omega_{1}} \left|M^{(3)}(s)M_{RHP}^{(2)}(s)\overline{\partial}R^{(2)}(s)M_{RHP}^{(2)}(s)^{-1}\right| dA(s),$$
(9.19)

$$\leq I_4 + I_5 + I_6, \tag{9.20}$$

where

$$I_4 = \int_0^\infty \int_v^\infty |\overline{\partial}\Xi_1| e^{-tz_1 u v} du dv, \qquad (9.21)$$

$$I_{5} = \int_{0}^{\infty} \int_{v}^{\infty} |p_{1}'(Rez)| e^{-tz_{1}uv} du dv, \qquad (9.22)$$

$$I_6 = \int_0^\infty \int_v^\infty |s - z_1|^{-1/2} e^{-tz_1 u v} du dv.$$
(9.23)

It follows from similar to the prove of I_i (i = 1, 2, 3) that

$$|I_4|, |I_5|, |I_6| \leq t^{-3/4},$$
 (9.24)

which imply the proof of the proposition is completed.

10. Soliton resolution for the Hirota equation

In this section we will construct the long-time asymptotic solution of the Hirota equation. According to the inverse transformation of (3.16), (4.4), (5.4) and (5.7), one has

$$M(z) = M^{(3)}(z)E(z)M^{(out)}(z)R^{(2)}(z)^{-1}T^{\sigma_3}(z), \ z \in \mathbb{C} \setminus (\mathcal{U}_{z_0} \cup \mathcal{U}_{z_1}),$$
(10.1)

note that $T^{\sigma_3}(z)$ is a diagonal matrix.

To construct the solution q(x, t), we take $z \to \infty$ along the imaginary axis. The advantage of this is that $R^{(2)}(z)$ is the identity matrix. By using these relationships (3.13), (6.37), (8.3) and (8.14), we have

$$M = \left(\mathbb{I} + \frac{M_1^{(3)}}{z} + \cdots\right) \left(\mathbb{I} + \frac{E_1}{z} + \cdots\right) \left(\mathbb{I} + \frac{M_1^{(out)}}{z} + \cdots\right) \left(\mathbb{I} + \frac{T_1^{\sigma_3}}{z} + \cdots\right), \quad (10.2)$$

from which, comparing the coefficients of z^{-1} to get

$$M_1 = M_1^{(out)} + E_1 + M_1^{(3)} + T_1^{\sigma_3}.$$
 (10.3)

Acknowledgements

This work was supported by the National Natural Science Foundation of China under Grant No. 11975306, the Natural Science Foundation of Jiangsu Province under Grant No. BK20181351, the Six Talent Peaks Project in Jiangsu Province under Grant No. JY-059, and the Fundamental Research Fund for the Central Universities under the Grant Nos. 2019ZDPY07 and 2019QNA35.

11. Appendix A: The parabolic cylinder model problem

Here we describe in detail the construction of parabolic cylindrical function solution [48, 49], which is frequently used in the literature of long-time asymptotic solution. Define the contour $\Sigma^{(pc)} = \bigcup_{i=1}^{4} \Sigma_{i}^{(pc)}$ shown in Fig. 10 where

$$\Sigma_{j}^{(pc)} = \left\{ \lambda \in \mathbb{C} | \arg \lambda = \frac{2j-1}{4} \pi \right\}.$$
 (A.1)




Figure 10. The contour of $\Sigma_j^{(pc)}$ which is the jump condition of the following RH problem.

For $r_0 \in \mathbb{C}$, let $v(r) = -\frac{1}{2\pi} \log(1 + |\gamma_0|^2)$, we then consider the following parabolic cylinder model Riemann-Hilbert problem.

RH Problem 11.1. Find a matrix-valued function $M^{(pc)}(\lambda)$ satisfying

• $M^{(pc)}(\lambda)$ is analytic in $\mathbb{C} \setminus \Sigma^{(pc)}$, (A.2)

•
$$M^{(pc)}_+(\lambda) = M^{(pc)}_-(\lambda)V^{(pc)}(\lambda), \quad \lambda \in \Sigma^{pc},$$
 (A.3)

•
$$M^{(pc)}(\lambda) = \mathbb{I} + \frac{M_1}{\lambda} + O(\lambda^2), \quad \lambda \to \infty.$$
 (A.4)

where

$$V^{(pc)}(\lambda) = \begin{cases} \lambda^{i\nu\hat{\sigma}_3} e^{-\frac{i\lambda^2}{4}\hat{\sigma}_3} \begin{pmatrix} 1 & 0 \\ r_0 & 1 \end{pmatrix}, & \lambda \in \Sigma_1^{(pc)}, \\ \lambda^{i\nu\hat{\sigma}_3} e^{-\frac{i\lambda^2}{4}\hat{\sigma}_3} \begin{pmatrix} 1 & \frac{r_0^*}{1+|r_0|^2} \\ 0 & 1 \end{pmatrix}, & \lambda \in \Sigma_2^{(pc)}, \\ \lambda^{i\nu\hat{\sigma}_3} e^{-\frac{i\lambda^2}{4}\hat{\sigma}_3} \begin{pmatrix} 1 & 0 \\ \frac{r_0}{1+|r_0|^2} & 1 \end{pmatrix}, & \lambda \in \Sigma_3^{(pc)}, \\ \lambda^{i\nu\hat{\sigma}_3} e^{-\frac{i\lambda^2}{4}\hat{\sigma}_3} \begin{pmatrix} 1 & r_0^* \\ 0 & 1 \end{pmatrix}, & \lambda \in \Sigma_4^{(pc)}, \end{cases}$$
(A.5)

As shown in [50], the parabolic cylinder equation reads

$$\left(\frac{\partial^2}{\partial z^2} + \left(\frac{1}{2} - \frac{z^2}{2} + a\right)\right) D_a = 0.$$

The explicit solution $M^{(pc)}(\lambda, r_0)$ of the **RH Problem** 11.1 can be expressed by in the literatures [5, 51]

$$M^{(pc)}(\lambda, r_0) = \Phi(\lambda, r_0) \mathcal{P}(\lambda, r_0) e^{\frac{i}{4}\lambda^2 \sigma_3} \lambda^{-i\nu\sigma_3},$$

where

$$\mathcal{P}(\lambda, r_0) = \begin{cases} \begin{pmatrix} 1 & 0 \\ -r_0 & 1 \end{pmatrix}, & \lambda \in \Omega_1, \\ \begin{pmatrix} 1 & -\frac{r_0^*}{1+|r_0|^2} \\ 0 & 1 \end{pmatrix}, & \lambda \in \Omega_3, \\ \begin{pmatrix} 1 & 0 \\ \frac{r_0}{1+|r_0|^2} & 1 \end{pmatrix}, & \lambda \in \Omega_4, \\ \begin{pmatrix} 1 & r_0^* \\ 0 & 1 \end{pmatrix}, & \lambda \in \Omega_6, \\ \mathbb{I}, & \lambda \in \Omega_2 \cup \Omega_5, \end{cases}$$

and

$$\Phi(\lambda, r_0) = \begin{cases} \begin{pmatrix} e^{-\frac{3\pi \nu}{4}} D_{i\nu} \left(e^{-\frac{3i\pi}{4}} \lambda \right) & -i\beta_{12} e^{-\frac{\pi}{4}(\nu-i)} D_{-i\nu-1} \left(e^{-\frac{i\pi}{4}} \lambda \right) \\ i\beta_{21} e^{-\frac{3\pi (\nu+i)}{4}} D_{i\nu-1} \left(e^{-\frac{3i\pi}{4}} \lambda \right) & e^{\frac{\pi \nu}{4}} D_{-i\nu} \left(e^{-\frac{i\pi}{4}} \lambda \right) \\ \begin{pmatrix} e^{\frac{\pi \nu}{4}} D_{i\nu} \left(e^{\frac{i\pi}{4}} \lambda \right) & -i\beta_{12} e^{-\frac{3\pi (\nu-i)}{4}} D_{-i\nu-1} \left(e^{\frac{3i\pi}{4}} \lambda \right) \\ i\beta_{21} e^{\frac{\pi}{4}(\nu+i)} D_{i\nu-1} \left(e^{\frac{i\pi}{4}} \lambda \right) & e^{-\frac{3\pi \nu}{4}} D_{-i\nu} \left(e^{\frac{3\pi \nu}{4}} \lambda \right) \end{pmatrix}, \quad \lambda \in \mathbb{C}^-, \end{cases}$$

with

$$\beta_{12} = \frac{\sqrt{2\pi}e^{i\pi/4}e^{-\pi\nu/2}}{r_0\Gamma(-i\nu)}, \quad \beta_{21} = \frac{-\sqrt{2\pi}e^{-i\pi/4}e^{-\pi\nu/2}}{r_0^*\Gamma(i\nu)} = \frac{\nu}{\beta_{12}}.$$

It follows from the well known asymptotic behavior of $D_a(z)$ that the asymptotic behavior of the solution can be determined by

$$M^{(pc)}(r_0, \lambda) = \mathbb{I} + \frac{M_1^{(pc)}}{i\lambda} + O(\lambda^{-2}),$$
(A.6)

where

$$M_1^{(pc)} = \begin{pmatrix} 0 & \beta_{12} \\ -\beta_{21} & 0 \end{pmatrix}.$$

12. Appendix B: Detailed calculations for the pure $\bar\partial$ -Problem

For $z \in \Omega_1$, let s = u + iv, z = x + iy, then we have

$$\begin{aligned} |I_{1}| &= \int_{0}^{\infty} \int_{v}^{\infty} \frac{1}{|s-z|} |\overline{\partial}\Xi_{1}| e^{-tz_{1}uv} du dv \\ &\lesssim \int_{0}^{\infty} e^{-tz_{1}uv} \left\| \overline{\partial}\Xi_{1} \right\|_{L^{2}} \left\| \frac{1}{|s-z|} \right\|_{L^{2}} dv \\ &\lesssim \int_{0}^{\infty} \frac{1}{\sqrt{|v-\eta|}} e^{-tz_{1}v^{2}} dv \\ &= \int_{0}^{\eta} \frac{1}{\sqrt{|v-\eta|}} e^{-tz_{1}v^{2}} dv + \int_{\eta}^{\infty} \frac{1}{\sqrt{|v-\eta|}} e^{-tz_{1}v^{2}} dv \\ &\triangleq F_{1} + F_{2} \end{aligned}$$
(B.1)

Note that taking $w = \frac{v}{\eta}$ yields

$$F_{1} = \int_{0}^{\eta} \frac{1}{\sqrt{\eta} \sqrt{1 - \frac{v}{\eta}}} e^{-tz_{1}v^{2}} dv = \int_{0}^{\eta} \frac{\sqrt{\eta} e^{-tz_{1}w^{2}\eta^{2}}}{\sqrt{1 - w}} dw$$
$$\lesssim t^{-1/4} \int_{0}^{\eta} \frac{1}{\sqrt{w(1 - w)}} dw \lesssim t^{-1/4}.$$
(B.2)

Similarly taking $v = w + \eta$ yields

$$F_{2} = \int_{\eta}^{\infty} \frac{1}{\sqrt{v - \eta}} e^{-tz_{1}v^{2}} dv = \int_{\eta}^{\infty} \frac{1}{\sqrt{w}} e^{-tz_{1}(w + \eta)^{2}} dv$$
$$\lesssim \int_{\eta}^{\infty} \frac{1}{\sqrt{w}} e^{-tz_{1}w^{2}} dv \lesssim t^{-1/4}.$$
(B.3)

For I_2 , we can obtain the estimate in the same way

$$I_2 \lesssim t^{-1/4}.\tag{B.4}$$

It follows from the Hölder inequality (taking q > 2) that

$$\left\|\frac{1}{s-z}\right\|_{L^{q}} \lesssim |v-\eta|^{\frac{1}{q}-1}.$$
(B.5)

Then

$$I_{3} \lesssim \int_{0}^{\infty} e^{-tz_{1}v^{2}} \left\| \frac{1}{\sqrt{s-z_{1}}} \right\|_{L^{p}} \left\| \frac{1}{s-z} \right\|_{L^{q}} dv$$

$$\lesssim \int_{0}^{\infty} e^{-tz_{1}v^{2}} v^{\frac{1}{p}-\frac{1}{2}} |v-\eta|^{\frac{1}{q}-1} dv$$

$$\lesssim t^{-1/4}.$$
(B.6)

Subsequently combining with (B.2), (B.3), (B.4) and (B.6), we can obtain (9.14).

References

- S.V. Manakov, Nonlinear Fraunhofer diffraction, Sov. Phys.-JETP 38 (1974) 693-696.
- [2] V. E. Zakharov, S. V. Manakov, Asymptotic behavior of nonlinear wave systems integrated by the inverse scattering method, Soviet Physics JETP. 44 (1976) 106-112.
- [3] R. F. Bikbaev, Asymptotic-behavior as *t*-infinity of the solution to the cauchyproblem for the landau-lifshitz equation, Theor. Math. Phys. 77 (1988) 1117-1123.
- [4] R. F. Bikbaev, Soliton generation for initial-boundary-value problems, Phys. Rev. Lett. 68 (1992) 3117-3120.
- [5] P. Deift, X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation, Ann. Math. 137(2) (1993) 295-368.
- [6] X. Zhou, P. Deift, Long-time behavior of the non-focusing nonlinear Schrödinger equation-a case study, Lectures in Mathematical Sciences, Graduate School of Mathematical Sciences, University of Tokyo, 1994.
- [7] J. Xu, Long-time asymptotics for the short pulse equation, J. Differential Equations. 265 (2018) 3494-3532.
- [8] J. Xu, E.G. Fan, Long-time asymptotic behavior for the complex short pulse equation, J. Differential Equations. 269 (2020) 10322-10349.

- [9] J. Xu, E.G. Fan, Long-time asymptotics for the Fokas-Lenells equation with decaying initial value problem: Without solitons, J. Differential Equations. 259 (2015) 1098-1148.
- [10] A.B. de Monvel, A. Kostenko, D. Shepelsky, G. Teschl, Long-time asymptotics for the Camassa-Holm equation, SIAM J. Math. Anal. 41 (2009) 1559-1588.
- [11] S.F. Tian, T.T. Zhang, Long-time asymptotic behavior for the Gerdjikov-Ivanov type of derivative nonlinear Schrödinger equation with time-periodic boundary condition, Proc. Amer. Math. Soc. 146 (2018) 1713-1729.
- [12] H. Liu, X.G. Geng, B. Xue, The Deift-Zhou steepest descent method to long-time asymptotics for the Sasa-Satsuma equation, J. Differential Equations 265 (2018) 5984-6008.
- [13] D.S. Wang, B.L. Guo, X.L. Wang, Long-time asymptotics of the focusing Kundu-Eckhaus equation with nonzero boundary conditions, J. Differential Equations 266(9) (2019) 5209-5253.
- [14] W.X. Ma, Long-time asymptotics of a three-component coupled nonlinear Schrödinger system, J. Geom. Phys. 153 (2020) 103669.
- [15] K.T.R. McLaughlin, P.D. Miller, The $\bar{\partial}$ steepest descent method and the asymptotic behavior of polynomials orthogonal on the unit circle with fixed and exponentially varying non-analytic weights, Int. Math. Res. Not. (2006), Art. ID 48673.
- [16] K.T.R. McLaughlin, P.D. Miller, The $\bar{\partial}$ steepest descent method for orthogonal polynomials on the real line with varying weights, Int. Math. Res. Not. IMRN (2008), Art. ID 075.
- [17] V.E. Zakharov and A.B. Shabat, A scheme for integrating the nonlinear equations of mathematical physics by the method of the inverse scattering problem, I Funct. Anal. Appl. 8 (1974) 226-235.
- [18] V.E. Zakharov and A.B. Shabat, Integration of the nonlinear equations of mathematical physics by the method of the inverse scattering transform, II Funct. Anal. Appl. 13 (1979) 166-174.
- [19] M.J. Ablowitz, D. Bar Yaacov and A.S. Fokas, On the inverse scattering transform for the Kadomtsev-Petviashvili equation Stud. Appl. Math. 69 (1983) 135-143.
- [20] R. Beals, R. Coifman, Linear spectral problems, non-linear equations and the $\bar{\partial}$ -method, Inverse Probl. 5 (1989) 87-130.
- [21] J.Y. Zhu and X.G. Geng, A hierarchy of coupled evolution equations with selfconsistent sources and the dressing method, J. Phys. A: Math. Theor. 46 (2013) 035204.

- [22] J.H. Luo, E.G. Fan, ∂-dressing method for the coupled Gerdjikov-Ivanov equation, Appl. Math. Lett. 110 (2020) 106589.
- [23] M. Dieng, K. McLaughlin, Long-time asymptotics for the NLS equation via dbar methods, arXiv:0805.2807.
- [24] R. Jenkins, J. Liu, P. Perry, C. Sulem, Soliton resolution for the derivative nonlinear Schrödinger equation, Commun. Math. Phys. 363 (2018) 1003-1049.
- [25] S. Cuccagna, R. Jenkins, On the asymptotic stability of *N*-soliton solutions of the defocusing nonlinear Schrödinger equation, Commun. Math. Phys. 343(3) (2016) 921-969.
- [26] R. Jenkins, J. Liu, P. Perry, C. Sulem, Global well-posedness for the derivative nonlinear Schrödinger equation, Commun. Part. Diff. Equ. 43(8) (2018) 1151-1195.
- [27] Y. L. Yang, E.G. Fan, Soliton Resolution for the Short-pluse Equation, arXiv:2005.12208.
- [28] Q. Y. Cheng, E.G. Fan, Soliton resolution for the focusing Fokas-Lenells equation with weighted Sobolev initial data, arXiv:2010.08714.
- [29] R. H. Ma, E.G. Fan, Long time asymptotic behavior of the focusing nonlinear Kundu-Eckhaus equation, arXiv:1912.01425.
- [30] G. Chen and J.Q. Liu, Soliton resolution for the modified KdV equation, arXiv:1907.07115.
- [31] R. Hirota, Exact envelope-soliton solutions of a nonlinear wave equation, J. Math. Phys. 14 (1973) 805-809.
- [32] Y. Tao, J. He, Multisolitons, breathers, and rogue waves for the Hirota equation generated by the Darboux transformation, Phys. Rev. E 85 (2012) 026601.
- [33] G. Zhang, S. Chen, Z. Yan, Focusing and defocusing Hirota equations with nonzero boundary conditions: Inverse scattering transforms and soliton solutions, Commun. Nonlinear Sci. Numer. Simul. 80 (2020) 104927.
- [34] M. Lakshmanan, S. Ganesan, Equivalent forms of a generalized Hirota's equation with linear inhomogeneities, J. Phys. Soc. Japan. 52 (1983) 4031-4034.
- [35] G.L. Lamb, Elements of Soliton Theory, Wiley, New York, 1980.
- [36] H.Q. Hao, J.W. Zhang, Conservation Laws for the Continuous and Discrete Hirota Equation, J. TaiYuan University of Technology, 048(006) (2017) 1025-1028.
- [37] W.Q. Peng, S.F. Tian, X.B. Wang, T.T. Zhang, Characteristics of rogue waves on a periodic background for the Hirota equation, Wave Motion. 93 (2020) 102454.
- [38] L. Huang, J. Xu, E.G. Fan, Long-time asymptotic for the Hirota equation via nonlinear steepest descent method, Nonlinear Anal. RWA 26 (2015) 229-262.

- [39] B.L. Guo, N. Liu, Y.F. Wang, Long-time asymptotics for the Hirota equation on the half-line, Nonlinear Anal. RWA 174 (2018) 118-140.
- [40] P. Deift, X. Zhou, Long-time asymptotics for integrable systems, Higher order theory, Comment. Math. Phys. 165(1) (1994) 175-191,
- [41] M. Borghese, R. Jenkins, K. T. R. McLaughlin, Long-time asymptotic behavior of the focusing nonlinear Schrödinger equation, Ann. I. H. Poincaré Anal, 35 (2018) 887-920.
- [42] S.F. Tian, T.T. Zhang, Long-time asymptotic behavior for the Gerdjikov-Ivanov type of derivative nonlinear Schrödinger equation with time-periodic boundary condition, Proc. Am. Math. Soc. 146 (2018) 1713-1729.
- [43] S.F. Tian, Initial-boundary value problems for the general coupled nonlinear Schrödinger equation on the interval via the Fokas method, J. Differential Equations 262 (2017) 506-558.
- [44] X. Zhou, Direct and inverse scattering transforms with arbitrary spectral singularities, Commun. Pure Appl. Math. 42(7) (1989) 895-938.
- [45] R. Beals, R. Coifman, Scattering and inverse scattering for first order systems, Commun. Pure Appl. Math. 37(1) (1984) 39-90.
- [46] R. Beals, P. Deift, C. Tomei, Direct and Inverse Scattering on the Line, Math. Surv. Monogr., vol. 28, American Mathematical Society, Providence, RI, ISBN 0-8218-1530-X, 1988.
- [47] Liouville's formula, wikipedia, https://en.wikipedia.org/wiki/Liouville%27s_formula.
- [48] A. Its, Asymptotic behavior of the solutions to the nonlinear Schrödinger equation, and isomonodromic deformations of systems of linear differential equations, Dokl. Akad. Nauk SSSR. 261(1) (1981) 14-18.
- [49] J. Liu, P. Perry, C. Sulem, Long-time behavior of solutions to the derivative nonlinear Schrödinger equation for soliton-free initial data, Ann. I. H. Poincaré, Anal. Non Linéaire, 35 (2018) 217-265.
- [50] F.W.J. Olver, A.B. Olde Daalhuis, D.W. Lozier, B.I. Schneider, R.F. Boisvert, C.W. Clark, B.R. Miller, B.V. Saunders, NIST Digital Library of Mathematical Functions, 2016.
- [51] R. Jenkins, K. McLaughlin, Semiclassical limit of focusing NLS for a family of square barrier initial data, Commun. Pure Appl. Math. 67(2) (2014) 246-320.