

Dynamical complexity of the Teleparallel gravity cosmology

Geovanny A. Rave-Franco

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Circuito Exterior C.U., A.P. 70-543, México D.F. 04510, México.

E-mail: geovanny.rave@ciencias.unam.mx

Celia Escamilla-Rivera

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Circuito Exterior C.U., A.P. 70-543, México D.F. 04510, México.

E-mail: celia.escamilla@nucleares.unam.mx

Jackson Levi Said

Institute of Space Sciences and Astronomy, University of Malta, Msida, MSD 2080, Malta.
Department of Physics, University of Malta, Msida, MSD 2080, Malta.

E-mail: jackson.said@um.edu.mt

Abstract. The exploration of teleparallel gravity has been done from a dynamical systems point of view in order to be tested against the cosmological evolution currently observed. So far, the proposed autonomous systems have been restrictive over a constant dynamical variable, which contains information related to the dynamics on the H_0 value. It is therefore that in this paper we consider a generalisation of the dynamical system by imposing a non-constant degree of freedom over it which allows us to rewrite a generic autonomous dynamical analysis. We will describe the treatment of our non-linear autonomous system by study the hyperbolic critical points and discuss an interesting phenomenological feature in regards to H_0 : the possibility to obtain a best fit value for this parameter in a cosmological viable $f(T, B)$ model, a mixed power law. This result allows to present a generic scenario in which it is possible to fix constraints to alleviate H_0 tension at present time.

1. Introduction

Λ CDM cosmology offers a very successful model structure in which to study galactic dynamics through the prism of cold dark matter [1], whereas cosmological scale physics is dominated by a cosmological constant [2]. Together with an inflationary epoch [3], Λ CDM cosmology largely reproduces the correct observations measured at various scales of the Universe. The necessity of adding both dark matter and dark energy stems from the lack of suitable predictions from general relativity (GR) which is the gravitational foundations of Λ CDM. Despite the success of Λ CDM as a cosmological model, some crucial observations have started to show important tensions between the model and observational evidence, which are apart from the well-known theoretical problems internal to the theory [4, 5]. The prominent contention in recent years has been the H_0 tension in which local and early Universe observations predict different values of the H_0 expansion parameter [6]. Another growing tension in recent cosmological data is related to the growth of large scale structure in the $f\sigma_8$ parameter [7]. Also, a more novel tension has been suggested wherein cosmic birefringence would be present in the Planck Collaboration data 2018 release [8] which would pose an even more serious problem for Λ CDM.

The growing observational tensions motivates us to explore the possible landscape of gravitational theories beyond GR which can each be distinguished through Lovelock's theorem [9] which provides a clear set of criteria through which to introduce different forms of gravity. One of these potential theories of gravity is that of teleparallel gravity (TG) where the curvature produced by the Levi-Civita connection is replaced by the torsion associated with the teleparallel connection [10, 11]. This novel foundation of gravity does not depend on GR but can produce a teleparallel equivalent to general relativity (TEGR) which is dynamically equivalent to GR in terms of its field equations [12, 13] but that originates from a distinct action in which the Ricci scalar Lagrangian is replaced by a torsion scalar [14, 15]. The core difference between these two Lagrangians is that the Ricci scalar also contains total divergence terms which has the impact that the teleparallel analogue of the Lovelock theorem will produce a much larger class of gravitational models which retain generically second order derivatives in their field equations [16, 17]. TG has also been shown to have a number of other attractive features such as its likeness to a Yang-mills theory [11] giving it a strong similarity to a particle physics theory, its potential definition of a gravitational energy-momentum tensor [18, 19], as well as it not requiring a Gibbons-Hawking-York boundary term giving TG a well defined Hamiltonian description, among other properties. Observationally, TG also has a number of other attractive features [20, 21, 22, 23].

In the same vein as $f(\overset{\circ}{R})$ gravity [24, 25] (over-circles denote quantities determined using the Levi-Civita connection), TEGR can be directly generalised to $f(T)$ gravity [26, 27, 28, 29, 30, 31, 32], which is a second order gravitational theory that has shown several potentially positive observational predictions [13, 33, 14, 22, 15, 34, 35, 36, 37, 21, 38]. However, to fully embrace the generalisations offered by TG and relate this

generalisation to $f(\overset{\circ}{R})$ gravity, we must also consider the inclusion of the boundary (or total divergence) term, B , through $f(T, B)$ gravity [39, 40, 41, 42, 43, 44, 44, 45]. This relation becomes an equivalence when the arguments are combined in a particular prescribed way, namely when the limit $f(T, B) = f(-T + B) = f(\overset{\circ}{R})$ is selected. Similar to $f(T)$ gravity, there have been a number of studies in this extension to TEGR [46, 47, 43, 39, 42, 41, 44, 39, 48, 49] which have shown a number of promising results for the theory.

TG also has a number of other extensions to TEGR such as $f(T, T_G)$ gravity where the TG analogue of the Gauss-Bonnet scalar is incorporated into the gravitational model [50, 51, 52, 53]. Another recent proposal of TG was proposed in Ref.[54] where a new term in addition to the standard Horndeski terms was introduced which was later found to allow for a number of Horndeski models to be revived despite the recent speed of light constraint on the propagation of gravitational waves [55]. This new realisation of Horndeski gravity has also been explored through the parameterized post-Newtonian (PPN) formalism [56] where it was found that most model largely survive current PPN constraints.

Dynamical systems offers a vital approach in which the background equations of cosmological models can be probed against cosmological observations [57]. In this work, we probe $f(T, B)$ gravity in the context of a homogeneous and isotropic Universe using the Friedmann–Lemaître–Robertson–Walker (FLRW) metric. This approach has been used to explore higher order modified TG models in Ref.[58] where a number of important reconstructions were investigated together with their dynamical systems evolution, and where they also compare their results with Supernova type 1a data. In Ref.[41], $f(T, B)$ gravity is directly studied in the context of Noether’s theorem where several reconstruction approaches are explored along with some stability conditions. Another important work where Noether’s theorem was used to determine new solutions is Ref.[59], some of which have shown promise in terms of producing cosmologically viable models [41]. $f(T, B)$ gravity has also been investigated in terms of its thermodynamics in Ref.[44], where the matter perturbation equation is derived. Other reconstructions [60, 61] have been developed in this class of theories of gravity where energy condition information was also provided.

In Ref.[48] a number of literature $f(T, B)$ gravity models were fitted using cosmic chronometers, supernova and baryonic acoustic oscillation data which resulted in model parameter constraints. In Ref.[49], the viable models were explored in terms of their dynamical systems. One of these realisations was the mixed power law model where the torsion scalar and boundary term are coupled with independent indices. This is interesting because it is one of the core studied models in literature that is novel from $f(\overset{\circ}{R})$ gravity, i.e. that cannot be produced by any choice of this free function. However, due to the complexity of the ensuing dynamical system, one of the dynamical variables had been set to be a constant which overly constrained the ensuing system. In this work, we have found an approach in which this assumption does not need to be taken and the fully autonomous dynamical system can be explored for this model. The paper

is structured as follows: in section 2, we develop the technical details of $f(T, B)$ gravity and provide the background cosmology equations. The dynamical system is then defined in section 3 for the mixed power law model of $f(T, B)$ gravity. It is this section that we generalise the dynamical variable from being a constant. In section 4, we describe the dynamical systems treatment in further detail and expose the hyperbolic critical points of the system. The possibility of reducing the H_0 tension is discussed in section 5 where we obtain a best fit value for this observational parameter. Finally, in section 6, we summarise our core results and provide a discussion of these results.

2. $f(T, B)$ cosmology

TG is built on the exchange of the curvature associated with the Levi-Civita connection $\overset{\circ}{\Gamma}_{\mu\nu}^{\sigma}$ (we use over-circles to denote quantities calculated with the Levi-Civita connection throughout) with the torsion linked to the teleparallel connection $\Gamma_{\mu\nu}^{\sigma}$ [13, 12]. GR and its modifications are principally built from scalars based on the Riemann tensor [5]. However, the Riemann tensor identically vanishes when the teleparallel connection is used since it is curvatureless and satisfies metricity.

The construction of the teleparallel connection is formulated by building the metric tensor $g_{\mu\nu}$ from tetrad fields e^A_{μ} (and its inverses E_A^{μ}) [11]. In this setting, Latin indices refer to local Minkowski space coordinates while Greek indices refer to the general manifold, where the tetrad components solder both spaces together and provide a direct approach to transform indices between these manifolds. These concepts come together in the relations that produce the metric tensor from the tetrad fields which takes the form

$$g_{\mu\nu} = e^A_{\mu} e^B_{\nu} \eta_{AB}, \eta_{AB} = E_A^{\mu} E_B^{\nu} g_{\mu\nu}, \quad (1)$$

where the tetrads satisfy orthogonality conditions

$$e^A_{\mu} E_B^{\mu} = \delta_A^B, e^A_{\mu} E_A^{\nu} = \delta_{\mu}^{\nu}, \quad (2)$$

for internal consistency. Then, the teleparallel connection can be defined as [10]

$$\Gamma_{\nu\mu}^{\sigma} := E_A^{\sigma} \partial_{\mu} e^A_{\nu} + E_A^{\sigma} \omega^A_{B\mu} e^B_{\nu}, \quad (3)$$

where $\omega^A_{B\mu}$ represents the spin connection. The spin connection appears in the connection to sustain the general covariance of the ensuing theories [62]. The components of the spin connection are totally inertial and represent the local Lorentz transformations (LLTs) [11, 63]. Naturally, one may choose a Lorentz frame in which the spin connection components vanish which is called the Weitzenböck gauge. However, one must be careful when applying this gauge in order to not lose diffeomorphism invariance of the equations of motion.

Taking the same rationale as GR where scalars are built from the Riemann tensor, TG theories are constructed using contractions of the torsion tensor defined as [12, 13]

$$T^{\sigma}_{\mu\nu} := -2\Gamma^{\sigma}_{[\mu\nu]}, \quad (4)$$

where square brackets denote the usual antisymmetric operator. The torsion tensor represents the field strength of TG, and can produce a number of other measures of torsion [11]. There exists a particular choice of quadratic contractions of the torsion tensor called the torsion scalar defined as

$$T := \frac{1}{4}T^\sigma{}_{\mu\nu}T^\mu{}_{\sigma\nu} + \frac{1}{2}T^\sigma{}_{\mu\nu}T^{\nu\mu}{}_\sigma - T^\sigma{}_{\mu\sigma}T^{\nu\mu}{}_\nu, \quad (5)$$

which is the TG analogue of the standard gravity Ricci scalar \mathring{R} . The introduction of torsion scalar is necessary because the Ricci scalar identically vanishes when the teleparallel connection is assumed, i.e. $\mathring{R} = 0$.

Calculating the Ricci scalar in both connections leads to a revealing relation from which the torsion scalar emerges, namely [64, 65]

$$R = \mathring{R} + T - 2B = 0, \quad (6)$$

where $B := \frac{1}{e}\partial_\mu(eT^\mu) = \mathring{\nabla}_\mu(T^\mu)$ is a boundary term, and $e = \det(e^A{}_\mu) = \sqrt{-g}$ is the tetrad determinant. Hence, a teleparallel equivalent of general relativity (TEGR) action can be written as

$$\mathcal{S}_{\text{TEGR}} = -\frac{1}{2\kappa^2} \int d^4x eT + \int d^4x e\mathcal{L}_m, \quad (7)$$

where $\kappa^2 = 8\pi G$ and \mathcal{L}_m is the regular matter Lagrangian, which are guaranteed to reproduce the Einstein equations.

The boundary term plays no role in the ensuing TEGR field equations as one would expect. However, in standard gravity GR the boundary term is the reason why many modified theories of gravity turn out to be fourth order in their field equations, such as in $f(\mathring{R})$ gravity [66, 25]. Thus, in TG, we decouple the second and fourth order contributions to the field equations. In the context of the central role that the torsion scalar and the boundary term play in TG and its relation to GR, the generalization of TEGR to $f(T, B)$ gravity has gained significant momentum in recent years [67, 39, 40, 41, 42, 43, 44, 44, 45]. The theory has an associated action

$$\mathcal{S}_{f(T,B)} = \frac{1}{2\kappa^2} \int d^4x e f(T, B) + \int d^4x e\mathcal{L}_m. \quad (8)$$

where it is important to note that $f(T, B) = f(-T + B) = f(\mathring{R})$, but more generally $f(T, B) \neq f(\mathring{R})$.

In the present work, we work in a flat Friedmann-Lemaître-Robertson-Walker (FLRW) Universe where the tetrad field

$$e^A{}_\mu = \text{diag}(1, a(t), a(t), a(t)), \quad (9)$$

tuns out to be compatible with the Weitzenböck gauge in $f(T, B)$ gravity [44, 68, 69], and where $a(t)$ is the scale factor. Through Eq.(1), this leads directly to the regular FLRW metric

$$ds^2 = dt^2 - a(t)^2(dx^2 + dy^2 + dz^2). \quad (10)$$

Through the definitions of the torsion scalar and boundary term, it follows that for a flat FLRW scenario

$$T = 6H^2, \quad B = 6(3H^2 + \dot{H}), \quad (11)$$

which together reproduce the Ricci scalar, i.e. $\mathring{R} = -T + B = 6(\dot{H} + 2H^2)$. These scalars contribute to produce the $f(T, B)$ gravity Friedmann equations

$$-3H^2(3f_B + 2f_T) + 3H\dot{f}_B - 3\dot{H}f_B + \frac{1}{2}f = \kappa^2\rho, \quad (12)$$

$$-\left(3H^2 + \dot{H}\right)(3f_B + 2f_T) - 2H\dot{f}_T + \ddot{f}_B + \frac{1}{2}\ddot{f} = -\kappa^2P, \quad (13)$$

where overdots refer to derivatives with respect to cosmic time t , and where ρ and P respectively represent the energy density and pressure of matter.

In order to better explore the contributions from the modified Lagrangian, we consider $f(T, B)$ gravity as an effective fluid that appears in addition toTEGR through $f(T, B) \rightarrow -T + \tilde{f}(T, B)$. Using this setting, the $f(T, B)$ Friedmann equations can be written as

$$3H^2 = \kappa^2(\rho + \rho_{\text{eff}}), \quad (14)$$

$$3H^2 + 2\dot{H} = -\kappa^2(P + P_{\text{eff}}), \quad (15)$$

where the effective energy density and pressure terms take the form

$$\kappa^2\rho_{\text{eff}} := 3H^2(3\tilde{f}_B + 2\tilde{f}_T) - 3H\dot{\tilde{f}}_B + 3\dot{H}\tilde{f}_B + \frac{1}{2}\tilde{f}, \quad (16)$$

$$\kappa^2P_{\text{eff}} := -\frac{1}{2}\tilde{f} - \left(3H^2 + \dot{H}\right)(3\tilde{f}_B + 2\tilde{f}_T) - 2H\dot{\tilde{f}}_T + \ddot{\tilde{f}}_B. \quad (17)$$

An interesting property of this effective fluid is that it continues to observe continuity equation

$$\dot{\rho}_{\text{eff}} + 3H(\rho_{\text{eff}} + P_{\text{eff}}) = 0, \quad (18)$$

and also defines an effective equation of state (EoS) parameter

$$\omega_{\text{eff}} := \frac{P_{\text{eff}}}{\rho_{\text{eff}}} \quad (19)$$

$$= -1 + \frac{\ddot{\tilde{f}}_B - 3H\dot{\tilde{f}}_B - 2\dot{H}\tilde{f}_T - 2H\dot{\tilde{f}}_T}{3H^2(3\tilde{f}_B + 2\tilde{f}_T) - 3H\dot{\tilde{f}}_B + 3\dot{H}\tilde{f}_B - \frac{1}{2}\tilde{f}}. \quad (20)$$

As expected, this limits to the Λ CDM model when $f(T, B)$ is a constant.

3. Generic $f(T, B)$ cosmological dynamical system

In our previous work [67] it was noted that the mixed power law model

$$f(T, B) = f_0 B^k T^m, \quad (21)$$

leads to a degeneration in the dynamical variables, which allow us to study the general case of $\lambda = \frac{\ddot{H}}{H^3}$ and still working with an autonomous dynamical system. In fact, since

$$X = f_B = k f_0 B^{k-1} T^m = \frac{k}{B} f = \frac{k}{6(3H^2 + \dot{H})} f = \frac{f}{6H^2} \frac{k}{3 + \frac{\dot{H}}{H^2}} = -\frac{Wk}{3 + Z}, \quad (22)$$

with $X = f_B$, $W = -\frac{f}{6H^2}$ and $Z = \frac{H'}{H}$, where the prime denotes the derivative with respect to $N = \ln a$. It can be noted that X is not an independent variable, and it depends on W and Z , with the restriction $Z \neq -3$. This corresponds to a non-vanishing boundary term $B \neq 0$, since $B = 6H^2(3 + Z)$.

Under the same reasoning, it is easy to show that $Y = f'_B$ is not an independent variable. Then, using the second Friedmann equation it is possible obtain the derivative $\frac{d}{dN}\lambda$ in terms of Z, W, λ and close the dynamical system. Nevertheless, in doing so, it is possible but cumbersome to actually solve the system but this would require a different approach. From Eq.(22) it is required that $Z \neq -3$, where it is then possible to isolate W

$$W = -\frac{X}{k}(Z + 3), \quad (23)$$

then, instead of considering Z, W and λ as dependent variables, we consider X, Y and Z as independent variables and W, λ as dependent ones. This approach is equivalent to the other one but easier to work with in practical terms. In fact, a direct calculation using Eq.(23) and its derivative, shows that $\lambda = \frac{\ddot{H}}{H^3}$ can be rewritten as

$$\lambda = \frac{1}{1-k} \left(6kZ + 2(m-1)(Z+3)Z - \frac{Y(Z+3)}{X} + 2Z^2 \right). \quad (24)$$

Therefore, if we consider a Universe filled with two fluids $\rho = \rho_\omega + \rho_r$ where ρ_ω is a perfect fluid in which the EoS ω does not necessarily vanish, and ρ_r is the EoS for radiation, then the independent dynamical variables are

$$X \equiv f_B, \quad Y \equiv f'_B, \quad Z = \frac{H'}{H} = \frac{\dot{H}}{H^2}, \quad V \equiv \Omega_r \equiv \frac{\kappa\rho_r}{3H^2}. \quad (25)$$

In this context, the Friedmann equations can be recast in terms of a dynamical system given through the set of couple differential equations

$$Z' = \lambda - 2Z^2, \quad (26)$$

$$X' = Y, \quad (27)$$

$$Y' = -3\omega\Omega_\omega - V + 3W + (9 + 3Z)X + f_T(6 + 2Z) + 2f'_T - ZY - 3 - 2Z, \quad (28)$$

$$V' = -4V - 2ZV, \quad (29)$$

where $f_T = -mW$, W and λ given by Eqs.(23) and Eq.(24) respectively, and

$$\Omega_\omega = 1 - V - \Omega_{eff}, \quad (30)$$

where

$$\Omega_{eff} = (3 + Z)X + 2f_T - Y + W. \quad (31)$$

The critical point of the dynamical system is

$$\mathbf{x}_* = (Z, X, Y, V) = \left(0, \frac{k}{3(k + 2m - 1)}, 0, 0 \right), \quad (32)$$

and the stability matrix evaluated at the critical point is

$$\mathcal{M}|_{\mathbf{x}_*} = \begin{pmatrix} -\frac{6m}{k-1} - 6 & 0 & \frac{9(k+2m-1)}{(k-1)k} & 0 \\ 0 & 0 & 1 & 0 \\ -\frac{2m}{k-1} + \omega - 1 & \frac{9(\omega+1)(k+2m-1)}{k} & \frac{6m}{k-1} - 3\omega & 3\omega - 1 \\ 0 & 0 & 0 & -4 \end{pmatrix}. \quad (33)$$

The eigenvalues of Eq.(33) are given by

$$\mu_1 = -4, \quad (34)$$

$$\mu_2 = -\frac{3}{2} \left(1 + \frac{\sqrt{\alpha}}{k(k-1)} \right), \quad (35)$$

$$\mu_3 = -\frac{3}{2} \left(1 - \frac{\sqrt{\alpha}}{k(k-1)} \right), \quad (36)$$

$$\mu_4 = -3(1 + \omega), \quad (37)$$

where $\alpha = (k-1)k(9k^2 + 24(k-1)m - 17k + 16m^2 + 8)$. In order to obtain a stable critical point using linear theory [70, 71], we have to determine what values of ω , k and m make the real part of the eigenvalues vanish, i.e. make the critical point non-hyperbolic. From μ_4 , we observe that the value $\omega = -1$ lead us to a non-hyperbolic critical point, then the condition $\omega \neq -1$ is required.

On the other hand, observe that if $\alpha < 0$ then all eigenvalues have non-vanishing real parts, therefore lets focus on the case $\alpha > 0$ that makes all eigenvalues real. Considering each eigenvalue in turn

- $\mu_2 = 0$:

$$(i) \quad 0 < m < \frac{1}{2} \wedge (k = 1 - 2m \vee k = 1 - m). \quad (38)$$

$$(ii) \quad \frac{1}{2} \leq m < 1 \wedge k = 1 - m. \quad (39)$$

- $\mu_3 = 0$:

$$(i) \quad m < 0 \wedge (k = 1 - m \vee k = 1 - 2m). \quad (40)$$

$$(ii) \quad \frac{1}{2} < m \leq 1 \wedge k = 1 - 2m. \quad (41)$$

$$(iii) \quad m > 1 \wedge (k = 1 - 2m \vee k = 1 - m). \quad (42)$$

Now, let us examine the conditions over ω , k and m which provide a stable critical point. The region of any kind of instability (saddle-like or repulsor-like) are not going to be stated explicitly since, as we shall see later, the critical point is a de Sitter acceleration phase, then any kind of instability of the critical point is not supported by observations.

As stated earlier, the condition that $\alpha \leq 0$ gives, without any further condition, a stable critical point and, based on the eigenvalue μ_4 , in order to have a stable hyperbolic critical point, the inequality $\omega > -1$ is a necessary condition. Hence any kind of ghost-like fluid generates an inevitable instability. The conditions over k and m such that $\alpha \leq 0$ are

$$(i) \quad m \leq -\frac{1}{48} \wedge 0 < k < 1. \quad (43)$$

(ii)

$$\begin{aligned}
 & -\frac{1}{48} < m < 0 \wedge \\
 & \left(0 < k \leq \frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \vee \frac{1}{18}(17 - 24m) \right. \\
 & \left. + \frac{1}{18}\sqrt{48m + 1} \leq k < 1 \right).
 \end{aligned} \tag{44}$$

$$\text{(iii)} \quad m = 0 \wedge 0 < k \leq \frac{8}{9}. \tag{45}$$

(iv)

$$\begin{aligned}
 & 0 < m < \frac{1}{2} \wedge \\
 & \left(0 \leq k \leq \frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \vee \frac{1}{18}(17 - 24m) \right. \\
 & \left. + \frac{1}{18}\sqrt{48m + 1} \leq k \leq 1 \right).
 \end{aligned} \tag{46}$$

$$\text{(v)} \quad m = \frac{1}{2} \wedge \frac{5}{9} \leq k < 1. \tag{47}$$

(vi)

$$\begin{aligned}
 & \frac{1}{2} < m < 1 \wedge \\
 & \left(\frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \leq k < 0 \vee \frac{1}{18}(17 - 24m) \right. \\
 & \left. + \frac{1}{18}\sqrt{48m + 1} \leq k < 1 \right).
 \end{aligned} \tag{48}$$

$$\text{(vii)} \quad m = 1 \wedge -\frac{7}{9} \leq k < 1. \tag{49}$$

(viii)

$$\begin{aligned}
 & m > 1 \wedge \\
 & \left(\frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \leq k \leq \frac{1}{18}(17 - 24m) \right. \\
 & \left. + \frac{1}{18}\sqrt{48m + 1} \vee 0 < k < 1 \right).
 \end{aligned} \tag{50}$$

Now, the condition $\alpha > 0$ leads to a set of real eigenvalues, then to determine the stability of the critical point, we have to analyse the regions such that $\mu_2 < 0$ and $\mu_3 < 0$. These regions are

$$\text{(i)} \quad m \leq -\frac{1}{48} \wedge 1 - m < k < 1 - 2m. \tag{51}$$

(ii)

$$\begin{aligned}
 & -\frac{1}{48} < m \leq 0 \wedge \\
 & \left(\frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} < k < \frac{1}{18}(17 - 24m) + \frac{1}{18}\sqrt{48m + 1} \vee \right. \\
 & \left. 1 - m < k < 1 - 2m \right), \tag{52}
 \end{aligned}$$

(iii)

$$\begin{aligned}
 & 0 < m \leq \frac{1}{2} \wedge \\
 & \left(\frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} < k < 1 - 2m \vee \right. \\
 & \left. 1 - m < k < \frac{1}{18}(17 - 24m) + \frac{1}{18}\sqrt{48m + 1} \right), \tag{53}
 \end{aligned}$$

(iv)

$$\begin{aligned}
 & \frac{1}{2} < m \leq 1 \wedge \\
 & \left(1 - 2m < k < \frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \vee \right. \\
 & \left. 1 - m < k < \frac{1}{18}(17 - 24m) + \frac{1}{18}\sqrt{48m + 1} \right), \tag{54}
 \end{aligned}$$

(v)

$$\begin{aligned}
 & m > 1 \wedge \\
 & \left(1 - 2m < k < \frac{1}{18}(17 - 24m) - \frac{1}{18}\sqrt{48m + 1} \vee \right. \\
 & \left. \frac{1}{18}(17 - 24m) + \frac{1}{18}\sqrt{48m + 1} < k < 1 - m \right). \tag{55}
 \end{aligned}$$

The regions different to the aforementioned values lead to either saddle-like instability or repulsor-like instability.

While evaluating the density parameter of the effective fluid in Eq.(31) at the critical point, it is straightforward to find that

$$\Omega_{eff} = 1, \tag{56}$$

and $\Omega_r = 0$, $\Omega_\omega = 0$. Therefore, the critical point corresponds to an effective fluid dominated era, and since $Z = 0$ at the critical point and $Z = \frac{\dot{H}}{H^2}$, then $H = H_0$, which corresponds to a de Sitter Universe. Actually, when evaluating the deceleration parameter $q = -\ddot{a}/\dot{a}^2$ and the jerk parameter $j = \ddot{\ddot{a}}/aH^3$ at the critical point, we obtain

$$q = -1, \quad j = 1, \tag{57}$$




Figure 1. Two dimensional phase portrait of the dynamical system Eq.(29) for $m = -2/5$, $k = 1/2$ and $\omega = 0$ with $X = \frac{k}{3(-1+k+2m)}$, $V = 0$. Here we notice the attractor nature of the critical point of the 4D dynamical system.

respectively. Here, it is important to point out that this is radically different to our previous result in Ref.[49]. It was shown in Ref.[67] that the mixed power law model, when considering λ as a constant, the critical points corresponded to a matter-dominated era, therefore, when turning on λ to change dynamically, the matter-dominated critical points are shifted to a single de Sitter point. Therefore, the mixed power law model is able to recover a late-time accelerated Universe as an attractor point. In Fig.(1) a two dimensional phase portrait of the dynamical system from Eq.(29) is shown.

4. Linearised $f(T, B)$ solutions near the critical point

Consider an autonomous dynamical system of the form

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}), \quad (58)$$

where $\mathbf{x} \in \mathbb{R}^n$, \mathbf{x}_0 is a critical point of the system and $\mathbf{f} \in C^r(U_{\mathbf{x}_0})$, $r \geq 1$ with $U_{\mathbf{x}_0}$ a neighbourhood of \mathbf{x}_0 . Since $\mathbf{f} \in C^r(U_{\mathbf{x}_0})$ we could approximate \mathbf{f} in the neighbourhood $U_{\mathbf{x}_0}$ by a Taylor polynomial, then we have that the autonomous dynamical system is

$$\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}) \approx \mathbf{f}(\mathbf{x}_0) + \mathcal{M}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)^T + \mathcal{O}(\|\mathbf{x} - \mathbf{x}_0\|^2), \quad (59)$$

where $\mathcal{M}(\mathbf{x}_0) = D\mathbf{f}(\mathbf{x}_0)$ is the stability matrix or Jacobian matrix evaluated at the critical point $\mathbf{x} = \mathbf{x}_0$. Since \mathbf{x}_0 is a critical point $\mathbf{f}(\mathbf{x}_0) = 0$, then, near the critical point, the generally non-linear autonomous dynamical system in Eq.(58) can be approximated by the linear system

$$\dot{\mathbf{x}} \approx \mathcal{M}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)^T. \quad (60)$$

This result was actually proven by Hartmann and Grobmann separately and it is known as the *Hartmann-Grobmann theorem* [70, 72] or *the linearisation theorem*. This states that if \mathbf{x}_0 is a hyperbolic critical point of the non-linear dynamical system Eq.(58) then

the dynamical system and its linearisation Eq.(60) are topologically equivalent, i.e. there exists an homeomorphism $H : U_{\mathbf{x}_0} \rightarrow V$, where V is an open set containing the origin, mapping trajectories of Eq.(58) in $U_{\mathbf{x}_0}$ onto trajectories of Eq.(60) in V and preserves orientation in the sense that if a trajectory is directed from p_1 to p_2 in $U_{\mathbf{x}_0}$, then its image is directed from $H(p_1)$ to $H(p_2)$ in V . An even more powerful theorem was proven only by Hartmann and states that if $\mathbf{f} \in C^2(U_{\mathbf{x}_0})$ and the critical point is hyperbolic, then it is possible to find a C^1 -diffeomorphism between the non-linear dynamical system and its linearisation, thus the differential structure near the critical point is preserved. The latter theorem allows us to study the linearised solution of the dynamical system Eq.(29) when the Universe is near to an accelerated de Sitter expansion. Moreover, this solution is not a condition for a stable critical point, wherever the case, this should be confronted with observational data. A similar approach was made in Ref.[73] in the context of Brans-Dicke cosmology. Notice that the vector function defined by the right hand side part of the dynamical system in Eq.(29) is of class C^∞ over a neighbourhood of the critical point in Eq.(32) where $k \neq 0$, then it is possible to use the linearisation theorems. The linearisation of the dynamical system Eq.(29) in a neighbourhood of the critical point Eq.(32), defining $\tilde{X} \equiv X - \frac{k}{3(k+2m-1)}$, is

$$\begin{pmatrix} Z' \\ \tilde{X}' \\ Y' \\ V' \end{pmatrix} = \begin{pmatrix} -\frac{6m}{k-1} - 6 & 0 & \frac{9(k+2m-1)}{(k-1)k} & 0 \\ 0 & 0 & 1 & 0 \\ -\frac{2m}{k-1} + \omega - 1 & \frac{9(\omega+1)(k+2m-1)}{k} & \frac{6m}{k-1} - 3\omega & 3\omega - 1 \\ 0 & 0 & 0 & -4 \end{pmatrix} \begin{pmatrix} Z \\ \tilde{X} \\ Y \\ V \end{pmatrix}. \quad (61)$$

To obtain non-complex solutions for the Hubble parameter, in the next subsections we analyse the following two cases: $\alpha \geq 0$ and $\alpha < 0$.

4.1. Case $\alpha \geq 0$

If $\alpha \geq 0$ the solutions of the linearisation Eq.(61) are given by

$$\begin{aligned} Z(N) \approx & be^{-3N(\omega+1)} + Fe^{-4N} + c \exp \left[-\frac{3}{2} \left(1 + \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right] \\ & + d \exp \left[-\frac{3}{2} \left(1 - \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right], \end{aligned} \quad (62)$$

$$\begin{aligned} X(N) \approx & \frac{k}{3(k+2m-1)} \\ & - \frac{1}{18(k+2m-1)} \left(-\frac{2bke^{-3N(\omega+1)}(k\omega - k - 2m - \omega + 1)}{\omega + 1} + \right. \\ & \left. \frac{ck(\sqrt{\alpha} + k(-3k - 4m + 5) + 4m - 2) \exp \left[-\frac{3}{2} \left(1 + \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right]}{k + 2m - 1} \right. \\ & \left. - \frac{dk(\sqrt{\alpha} + k(3k + 4m - 5) - 4m + 2) \exp \left[-\frac{3}{2} \left(1 - \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right]}{k + 2m - 1} \right) \\ & + Fke^{-4N}(k + 3m - 1), \end{aligned} \quad (63)$$

$$\begin{aligned}
 Y(N) \approx & \frac{(k-1)k}{9(k+2m-1)} \left(\frac{3be^{-3N(\omega+1)}(-k\omega+k+2m+\omega-1)}{k-1} \right. \\
 & - \frac{3c(\sqrt{\alpha}+k(-3k-4m+3)) \exp\left[-\frac{3}{2}\left(1+\frac{\sqrt{\alpha}}{k(k-1)}\right)N\right]}{2(k-1)k} + \\
 & \left. \frac{3d(\sqrt{\alpha}+k(3k+4m-3)) \exp\left[-\frac{3}{2}\left(1-\frac{\sqrt{\alpha}}{k(k-1)}\right)N\right]}{2(k-1)k} + \right. \\
 & \left. Fe^{-4N}\left(\frac{6m}{k-1}+2\right)\right), \tag{64}
 \end{aligned}$$

$$V(N) \equiv \Omega_r(N) \approx \Omega_{0,r} \exp(-4N), \tag{65}$$

with b, c, d integration constants and

$$F = \frac{18\Omega_{0,r}(k+2m-1)}{7k^2+27km-16k+18m^2-27m+9}. \tag{66}$$

Notice that these solutions are only valid in a neighbourhood of the critical point, i.e. when $\|\mathbf{x} - \mathbf{x}_0\| < \varepsilon$ with $\mathbf{x} = (Z, X, Y, V)$. Based on these solutions, it is possible to compute some important cosmological quantities such as functions of $N = \ln a$.

We can see directly from the solution Eq.(65) that the evolution of the radiation density near the critical point is

$$\Omega_r(N) = \Omega_{0,r} \exp(-4N), \tag{67}$$

with $\Omega_{0,r}$ the radiation density today. When substituting the variable $N = \ln a$ this expression can be rewritten as

$$\Omega_r(a) = \frac{\Omega_{0,r}}{a^4}, \tag{68}$$

which is the standard evolution of the radiation density parameter in a de Sitter Universe with respect to the scale factor. Since the solutions are in a neighbourhood of the critical point, every non-linear term in the cosmological parameters vanishes and then only linear solutions remain.

The density parameter of the effective fluid is

$$\Omega_{eff} = 3f_B + 2f_T - f'_B - \frac{f}{6H^2} + \left(\frac{H'}{H}\right) f_B, \tag{69}$$

which can be written in terms of the dynamical variables as

$$\Omega_{eff} = \frac{(3+Z)}{k} \tilde{X}(k+2m-1) + \frac{3+Z}{3} - Y, \tag{70}$$

so that its linearised solution is

$$\Omega_{eff} \approx 1 + \frac{1}{3}be^{-3N(\omega+1)} \left(\frac{(k-1)k\omega}{k+2m-1} - \frac{2(k+m-1)}{\omega+1} \right) - \Omega_{0,r}e^{-4N}. \tag{71}$$

Analogously, since the first Friedmann equation reads as $\Omega_\omega + \Omega_r + \Omega_{eff} = 1$, we have

$$\Omega_\omega \approx -\frac{1}{3}be^{-3N(\omega+1)} \left(\frac{(k-1)k\omega}{k+2m-1} - \frac{2(k+m-1)}{\omega+1} \right) \equiv \Omega_{0,\omega}e^{-3N(\omega+1)}. \tag{72}$$

On the other hand, we have that

$$\begin{aligned} \frac{H'}{H} = Z(N) \approx & be^{-3N(\omega+1)} + c \exp \left[-\frac{3}{2} \left(1 + \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right] + \\ & d \exp \left[-\frac{3}{2} \left(1 - \frac{\sqrt{\alpha}}{k(k-1)} \right) N \right] + Fe^{-4N}. \end{aligned} \quad (73)$$

Integrating both sides of the later equation and applying the exponential function, we find that the linearised solution of the Hubble parameter is

$$\frac{H(N)}{H_i} \approx 1 - \frac{be^{-3N(\omega+1)}}{3(\omega+1)} - \frac{2ce^{-\frac{3}{2}N\left(\frac{\sqrt{\alpha}}{(k-1)k}+1\right)}}{3\left(\frac{\sqrt{\alpha}}{(k-1)k}+1\right)} - \frac{2de^{-\frac{3}{2}N\left(\frac{\sqrt{\alpha}}{k-k^2}+1\right)}}{3\left(\frac{\sqrt{\alpha}}{k-k^2}+1\right)} - \frac{1}{4}Fe^{-4N}, \quad (74)$$

where H_i is an integration constant.

Finally, since the EoS of the effective fluid is

$$\omega_{eff} = -1 + \frac{\ddot{f}_B - 3H\dot{f}_B - 2\dot{H}f_T - 2H\dot{f}_T}{3H^2(3f_B + 2f_T) - 3H\dot{f}_B + 3\dot{H}f_B - \frac{1}{2}f}, \quad (75)$$

this EoS can be written in terms of the dynamical variables as

$$\omega_{eff} = -1 + \frac{Y' + Y(Z-3) - \frac{2mXZ}{k}(3+Z) - \frac{2m}{k}[Y(3+Z) + X(\lambda - 2Z^2)]}{3\frac{x}{k}(3+Z)(k+2m-1) - 3Y}, \quad (76)$$

and then, its linearised solution will be given by

$$\begin{aligned} \omega_{eff} \approx & -1 + \\ & \left(-\frac{4e^{-4N}(18(k+1)m^2 + 3(k(2k+7) - 3)m + 7(k-1)k)\Omega_{0,r}}{7k^2 + 27(k-1)m - 16k + 18m^2 + 9} \right. \\ & + \frac{be^{-3N(\omega+1)}((k-1)k(2m+1)\omega - 2(k+2m-1)(km+k+m) + (k-1)k\omega^2)}{k+2m-1} \\ & + \frac{c(-3k^2m + k((3-4m)m-2) + (\sqrt{\alpha}-4)m+2) \exp\left[-\frac{3}{2}\left(1 + \frac{\sqrt{\alpha}}{k(k-1)}\right)N\right]}{k+2m-1} \\ & \left. - \frac{d(3k^2m + k(m(4m-3)+2) + (\sqrt{\alpha}+4)m-2) \exp\left[-\frac{3}{2}\left(1 - \frac{\sqrt{\alpha}}{k(k-1)}\right)N\right]}{k+2m-1} \right) \div \\ & \left(\frac{be^{-3N(\omega+1)}(-2(k+m-1)(k+2m-1) + (k-1)k\omega^2 + (k-1)k\omega)}{(\omega+1)(k+2m-1)} \right. \\ & \left. - 3\Omega_{0,r}e^{-4N} + 3 \right). \end{aligned} \quad (77)$$

4.2. Case $\alpha < 0$

In this case, the solutions of the linearisation in Eq.(61) give real solutions on the cosmological parameters, which are

$$\begin{aligned} Z(N) & \approx be^{-3N(\omega+1)} + 2ce^{-3N/2} \cos(N\epsilon) + Fe^{-4N}, \\ X(N) & \approx \frac{bke^{-3N(\omega+1)}(k(\omega-1) - 2m - \omega + 1)}{9(\omega+1)(k+2m-1)} - \frac{Fke^{-4N}(k+3m-1)}{18(k+2m-1)} \end{aligned} \quad (78)$$

$$\begin{aligned}
 & + e^{-3N/2} \left(\frac{c(k-1)k(3k+4m-2)\cos(\epsilon N)}{9(k+2m-1)^2} - \frac{\sqrt{\alpha}ck\sin(\epsilon N)}{9(k+2m-1)^2} \right), \quad (79) \\
 Y(N) \approx & \frac{bke^{-3N(\omega+1)}(-k\omega+k+2m+\omega-1)}{3(k+2m-1)} + \frac{2Fke^{-4N}(k+3m-1)}{9(k+2m-1)} \\
 & + e^{-3N/2} \left(\frac{ck(3k+4m-3)\cos(\epsilon N)}{3(k+2m-1)} - \frac{\sqrt{\alpha}c\sin(\epsilon N)}{3(k+2m-1)} \right), \quad (80)
 \end{aligned}$$

$$V(N) \equiv \Omega_r(N) \approx \Omega_{0,r}e^{-4N}, \quad (81)$$

where

$$\epsilon = \frac{3\sqrt{|\alpha|}}{2(k-1)k}, \quad (82)$$

and the same restriction as Eq.(66) on F .

In this case, the density parameters near the critical point are

$$\begin{aligned}
 \Omega_{\text{eff}} \approx & 1 + \frac{be^{-3N(\omega+1)}(-2(k+m-1)(k+2m-1) + (k-1)k\omega^2 + (k-1)k\omega)}{3(\omega+1)(k+2m-1)} \quad (83) \\
 & - \Omega_{0,r}e^{-4N},
 \end{aligned}$$

$$\Omega_\omega \approx -\frac{be^{-3N(\omega+1)}(-2(k+m-1)(k+2m-1) + (k-1)k\omega^2 + (k-1)k\omega)}{3(\omega+1)(k+2m-1)}, \quad (84)$$

$$\Omega_r \approx \Omega_{0,r}e^{-4N}, \quad (85)$$

where the Hubble parameter is then given by

$$\frac{H(N)}{H_i} \approx 1 - \frac{be^{-3N(\omega+1)}}{3(\omega+1)} + \frac{4ce^{-3N/2}(2\epsilon\sin(N\epsilon) - 3\cos(N\epsilon))}{4\epsilon^2 + 9} - \frac{1}{4}Fe^{-4N}, \quad (86)$$

and the EoS of the effective fluid is

$$\begin{aligned}
 \omega_{\text{eff}} \approx & -1 + \left(-\frac{4e^{-4N}(18(k+1)m^2 + 3(k(2k+7) - 3)m + 7(k-1)k)\Omega_{0,r}}{7k^2 + 27(k-1)m - 16k + 18m^2 + 9} \right. \\
 & + \frac{be^{-3N(\omega+1)}((k-1)k\omega(k-2m-1) - 2((k-1)k-m)(k+2m-1) + (k-1)^2k\omega^2)}{(k-1)(k+2m-1)} \\
 & \left. + e^{-3N/2} \left(\frac{2\sqrt{\alpha}cm\sin(N\epsilon)}{k+2m-1} - \frac{2c(k(m(3k+4m-3) + 2) + 4m-2)\cos(N\epsilon)}{k+2m-1} \right) \right) \div \\
 & \left(\frac{be^{-3N(\omega+1)}(-2(k+m-1)(k+2m-1) + (k-1)k\omega^2 + (k-1)k\omega)}{(\omega+1)(k+2m-1)} \right. \\
 & \left. - 3\Omega_{0,r}e^{-4N} + 3 \right). \quad (87)
 \end{aligned}$$

We have to keep in mind that the solutions described for both branches $\alpha \geq 0$ and $\alpha < 0$ are only valid in a neighbourhood of the critical point, i.e., they describe the dynamics of a Universe close to be a perfect de Sitter Universe, which in terms of the parameter $N = \ln a$, corresponds to $N \rightarrow \infty$, and in the terms of the redshift, corresponds to $z \rightarrow -1$. The theory of dynamical systems cannot guarantee these solutions to work well for a wide range of redshift but a narrow one, as we shall see, when comparing to observational data, we will notice that some models of $f(T, B)$ can describe the dynamics of the Universe today and may give a glimpse into the solution to the H_0 tension.

5. Phenomenological hints on H_0 tension from the generic $f(T, B)$ dynamical system

As we discussed, the theory of dynamical systems guarantees that the equations obtained above for the cosmological parameters are a solution of the Friedmann equations when $N \rightarrow \infty$, or $z \rightarrow -1$ in terms of the redshift. Therefore, it is viable to study them in phenomenological scenarios where should be possible to test with current astrophysical observations.

In this section, we will use our results derived from our non-linear dynamical system and link them with current $H(z)$ observational data to investigate in which redshift range this system of dynamical equations still holds as a valid solution for the Friedmann equations.

As a first step in that direction, we consider in this proposal Hubble parameter measurements $H(z)$ [74], in which the current sample consists of 51 measurements in the redshift range $0.07 < z < 2.0$ of which 31 points correspond to Cosmic Chronometers and the rest are from BAO estimates from the late time. Afterwards, we will constrain the values for $\Omega_{0,r}$, k , m , ω , and the integration constants, and also perform this for the $\alpha < 0$ case to assure that the obtained values of k and m provide a stable critical point.




Figure 2. Reconstruction of the evolution of $H(z)$ using the best fit obtained with the model Eq.(86) (blue color solid line) in comparison to the observational data described [74, 75] (red color dots).

To obtain a convenient set of free parameters appearing in Eq.(86) we must perform the change in variable $N = \ln \frac{1}{1+z}$, in order to obtain an expression in terms of the redshift and then fit this equation to the available data of the $H(z)$ [74] with respect to the redshift. If we consider the free parameters without any constrictions, a nonphysical set of parameters may be obtained from the fitting, like a negative value for the mean value of the radiation density $\Omega_{0,r}$ or an EoS for matter less than -1 which leads to an instability, to name a few, then a more mindful approach of fitting has to be performed. To obtain a convenient set of parameters with physical relevance, we have to restrict

or associate certain values to these parameters. More to the point, it has been shown that the radiation density today has a small value of $\Omega_{0,r} \approx 10^{-5}$ [76], then in order to simplify the model, we considered $\Omega_{0,r} = 0$. We restricted, based upon observations in Refs.[77, 78, 79], H_0 to vary in the range of $60 \leq H_0 \leq 90$. Finally, we considered the case $\alpha < 0$ for this analysis since any values of k and m in this case provide a stable critical point. Under these considerations, it is possible to avoid fitting k and m directly since it may lead to problems in the numerical code, due to the square root in Eq.(86), but it is possible and feasible to fit ϵ and then, based on Eq.(82), obtain values of k and m in the level curve of the obtained value for ϵ . Based on these considerations, the best fit parameters are

$$\begin{aligned} \omega &= 0.000 \pm 2.951, & \epsilon &= 1.749 \pm 5.671, & b &= -0.200 \pm 2.735, \\ c &= -0.213 \pm 2.221, & H_i &= 60.000 \pm 10.973, \end{aligned} \quad (88)$$

and the possible set of values of k and m in the level curve $\epsilon(k, m) = 1.749$ are

$$k \approx -0.294, \quad m = 0.75. \quad (89)$$

If we evaluate the best fit at $z = 0$ we obtain that $H_0 \approx 71.236 \pm 144.229$. In Fig.(2) the evolution of the approximated solution for the Hubble parameter Eq.(86) with the fitted values Eq.(88) and the observational data from [74] is shown.

The method described above leads to high error regions (C.L) in the fitted parameters and a high error to the H_0 value due to the large number of free parameters and the small amount of observational data. A different way to link the observational data with the approximated solutions obtained for the dynamical system can be performed. By taking the stability analysis performed in section 3, we can choose some values for k and m from the analysis and reduce the number of free parameters in the fitting procedure, but we still obtain a high error C.L are obtained for the remaining free parameters. To reduce further this error propagation we have to set some free parameters to a specific value, e.g. setting $\omega = 0$ which corresponds to a dust model, reduce the error C.L to the remaining parameters and the H_0 value.

In Fig.(3) we show a plot with different values of H_0 obtained under different conditions for some values of k and m obtained from the dynamical analysis. Describing the regions denote by color, we have that: The cyan region shows different values of H_0 for a dust $\omega = 0$ and zero radiation $\Omega_{0,r} = 0$ model in which the fitted parameters satisfy $-0.580 \leq c \leq -0.006$, $-0.449 \leq b \leq -0.017$ and $42.21 \leq H_i \leq 75.670$.

The pink region shows different values for H_0 for a dust model with radiation, in this case the fitted parameters satisfy $-0.078 \leq c \leq 0.292$, $-0.944 \leq b \leq -0.024$, $51.789 \leq H_i \leq 85.471$ and $0 \leq \Omega_{0,r} \leq 0.029$.

In the gray region only one value of H_0 is shown, since no values are defined a priori, the error C.L becomes large, and only one value is plotted to show explicitly the large error in this case. The obtained value for H_0 corresponds to the fitted parameters $b = -1.229 \pm 3.464$, $c = 0.203 \pm 1.916$, $H_i = 46.178 \pm 15.736$, $\Omega_{0,r} = 0.013 \pm 0.023$ and $\omega = -0.297 \pm 1.961$.




Figure 3. Values obtained for H_0 with specific choices of k and m denoted in the inner box label. *Top:* H_0 best fit value for a dynamical system with all the variables settled free. *Middle:* H_0 best fit values for a dust model $\omega = 0$. *Bottom:* H_0 best fit values for a dust and zero radiation model.

With the results obtained it is also possible to plot, e.g. the density parameters, the EoS of the effective fluid and deceleration parameter with respect to the redshift, making

the change of variable $N = \ln \frac{1}{1+z}$, e.g. with the fitted values (88) and null radiation, the graph of these cosmological parameters are shown in Fig.(4) for the density parameters (Left) and for the EoS of the effective fluid (Right).




Figure 4. *Left:* Evolution of the density parameters given by Eqs. (83) and (84) in terms of the redshift z . *Right:* Evolution of $\omega_{\text{eff}}(z)$ given by Eq.(87). The inflexion point occurs at $z \approx 1.397$ for this model.

6. Conclusions

As we have explored, a number of $f(T, B)$ models have been fitted using cosmic chronometers, supernova and baryonic acoustic oscillation data which resulted in model parameter constraints, that in some cases can alleviate the H_0 tension which has become very topical in recent years. Due to the complexity of the evolution cosmological system, some viable $f(T, B)$ models can be analysed via their dynamical systems. One of these results was the mixed power law model, where the torsion scalar and boundary term are coupled with independent indices. This aspect is remarkable since it includes one of the core studied models in literature that is novel from $f(\overset{\circ}{R})$ gravity. Moreover, due to the complexity of the ensuing dynamical system, one of the dynamical variables had been set to be a constant which overly constrained the system. With the aim to find a generic formulation of this dynamical system, in this work we have explored an approach in which this assumption does not need to be taken and the fully autonomous dynamical system can be explored using their hyperbolic critical points. We used the Hartmann-Grobmann theorem in order to study our non-linear dynamical system Eq.(29) whose solution is near to an accelerated de Sitter expansion. One of the most interesting features presented in this work is the possibility to describe the dynamics of the Universe today using a narrow branch of our dynamical system, and in this case to explore the nature of the H_0 tension from an autonomous system point of view.

To achieve the last goal, we employed in this work a $H(z)$ observational sample from Cosmic Chronometers plus BAO estimates. Our purpose is to constrain the characteristic values which define the mixed power law only. According to this phenomenological analysis, it was possible to find an optimal set of cases where the H_0 tension can be alleviated using specific values for the mixed $f(T, B)$ power law as we can see from Fig.(3). In these analyses, we considered three scenarios: with all

the variables settled free, for a dust model and for a dust and zero radiation model. According to this, cases where $k < 0.9$ seems to be in agreement with Planck 2018 H_0 constraint at $1-\sigma$, while for the specific case $k = 7457/8192$ and $1/4$, the H_0 is quite high with a considerable error propagation. In conclusion, the mixed power law model analysed here is novel in that it is not reproduced by an $f(\overset{\circ}{R})$ gravity Lagrangian, and has shown promise in reproducing a cosmology that resonates with current observations, giving motivation to further studies in this direction.

7. Acknowledgments

CE-R is supported by the Royal Astronomical Society as FRAS 10147. CE-R and GAR-F are supported by PAPIIT Project IA100220. This article is based upon work from CANTATA COST (European Cooperation in Science and Technology) action CA15117, EU Framework Programme Horizon 2020. The authors would like to acknowledge networking support by the COST Action CA18108 and funding support from Cosmology@MALTA which is supported by the University of Malta.

8. References

- [1] L. Baudis, *Dark matter detection*, *J. Phys.* **G43** (2016) 044001.
- [2] S. Weinberg, *Cosmology*, Cosmology. OUP Oxford, 2008.
- [3] L. Perenon, F. Piazza, C. Marinoni and L. Hui, *Phenomenology of dark energy: general features of large-scale perturbations*, *JCAP* **1511** (2015) 029 [1506.03047].
- [4] S. Weinberg, *The Cosmological Constant Problem*, *Rev. Mod. Phys.* **61** (1989) 1.
- [5] T. Clifton, P. G. Ferreira, A. Padilla and C. Skordis, *Modified Gravity and Cosmology*, *Phys. Rept.* **513** (2012) 1 [1106.2476].
- [6] E. Di Valentino et al., *Cosmology Intertwined II: The Hubble Constant Tension*, 2008.11284.
- [7] E. Di Valentino et al., *Cosmology Intertwined III: $f\sigma_8$ and S_8* , 2008.11285.
- [8] Y. Minami and E. Komatsu, *New Extraction of the Cosmic Birefringence from the Planck 2018 Polarization Data*, *Phys. Rev. Lett.* **125** (2020) 221301 [2011.11254].
- [9] D. Lovelock, *The Einstein tensor and its generalizations*, *J. Math. Phys.* **12** (1971) 498.
- [10] R. Weitzenböck, *‘Invariantentheorie’*. Noordhoff, Gronningen, 1923.
- [11] R. Aldrovandi and J. G. Pereira, *Teleparallel Gravity*, vol. 173. Springer, Dordrecht, 2013, 10.1007/978-94-007-5143-9.
- [12] M. Krssak, R. van den Hoogen, J. Pereira, C. Böhrmer and A. Coley, *Teleparallel theories of gravity: illuminating a fully invariant approach*, *Class. Quant. Grav.* **36** (2019) 183001 [1810.12932].
- [13] Y.-F. Cai, S. Capozziello, M. De Laurentis and E. N. Saridakis, *$f(T)$ teleparallel gravity and cosmology*, *Rept. Prog. Phys.* **79** (2016) 106901 [1511.07586].
- [14] G. Farrugia and J. L. Said, *Stability of the flat FLRW metric in $f(T)$ gravity*, *Phys. Rev.* **D94** (2016) 124054 [1701.00134].
- [15] G. Farrugia, J. L. Said and M. L. Ruggiero, *Solar System tests in $f(T)$ gravity*, *Phys. Rev.* **D93** (2016) 104034 [1605.07614].
- [16] P. Gonzalez and Y. Vasquez, *Teleparallel Equivalent of Lovelock Gravity*, *Phys. Rev. D* **92** (2015) 124023 [1508.01174].
- [17] P. González, S. Reyes and Y. Vásquez, *Teleparallel Equivalent of Lovelock Gravity, Generalizations and Cosmological Applications*, *JCAP* **07** (2019) 040 [1905.07633].
- [18] D. Blixt, M. Hohmann and C. Pfeifer, *Hamiltonian and primary constraints of new general relativity*, *Phys. Rev. D* **99** (2019) 084025 [1811.11137].
- [19] D. Blixt, M. Hohmann and C. Pfeifer, *On the gauge fixing in the Hamiltonian analysis of general teleparallel theories*, *Universe* **5** (2019) 143 [1905.01048].
- [20] R. Briffa, S. Capozziello, J. Levi Said, J. Mifsud and E. N. Saridakis, *Constraining Teleparallel Gravity through Gaussian Processes*, 2009.14582.
- [21] J. Levi Said, J. Mifsud, D. Parkinson, E. N. Saridakis, J. Sultana and K. Z. Adami, *Testing the violation of the equivalence principle in the electromagnetic sector and its consequences in $f(T)$ gravity*, *JCAP* **11** (2020) 047 [2005.05368].
- [22] A. Finch and J. L. Said, *Galactic Rotation Dynamics in $f(T)$ gravity*, *Eur. Phys. J. C* **78** (2018) 560 [1806.09677].
- [23] S. Bahamonde, J. Levi Said and M. Zubair, *Solar system tests in modified teleparallel gravity*, *JCAP* **10** (2020) 024 [2006.06750].
- [24] A. De Felice and S. Tsujikawa, *$f(R)$ theories*, *Living Rev. Rel.* **13** (2010) 3 [1002.4928].
- [25] S. Capozziello and M. De Laurentis, *Extended Theories of Gravity*, *Phys. Rept.* **509** (2011) 167 [1108.6266].
- [26] R. Ferraro and F. Fiorini, *Modified teleparallel gravity: Inflation without inflaton*, *Phys. Rev.* **D75** (2007) 084031 [gr-qc/0610067].
- [27] R. Ferraro and F. Fiorini, *On Born-Infeld Gravity in Weitzenböck spacetime*, *Phys. Rev.* **D78** (2008) 124019 [0812.1981].
- [28] G. R. Bengochea and R. Ferraro, *Dark torsion as the cosmic speed-up*, *Phys. Rev.* **D79** (2009)

- 124019 [0812.1205].
- [29] E. V. Linder, *Einstein's Other Gravity and the Acceleration of the Universe*, *Phys. Rev.* **D81** (2010) 127301 [1005.3039].
- [30] S.-H. Chen, J. B. Dent, S. Dutta and E. N. Saridakis, *Cosmological perturbations in $f(T)$ gravity*, *Phys. Rev.* **D83** (2011) 023508 [1008.1250].
- [31] S. Bahamonde, K. Flathmann and C. Pfeifer, *Photon sphere and perihelion shift in weak $f(T)$ gravity*, *Phys. Rev. D* **100** (2019) 084064 [1907.10858].
- [32] U. Ualikhanova and M. Hohmann, *Parametrized post-Newtonian limit of general teleparallel gravity theories*, *Phys. Rev. D* **100** (2019) 104011 [1907.08178].
- [33] S. Nesseris, S. Basilakos, E. N. Saridakis and L. Perivolaropoulos, *Viable $f(T)$ models are practically indistinguishable from Λ CDM*, *Phys. Rev.* **D88** (2013) 103010 [1308.6142].
- [34] L. Iorio and E. N. Saridakis, *Solar system constraints on $f(T)$ gravity*, *Mon. Not. Roy. Astron. Soc.* **427** (2012) 1555 [1203.5781].
- [35] M. L. Ruggiero and N. Radicella, *Weak-Field Spherically Symmetric Solutions in $f(T)$ gravity*, *Phys. Rev.* **D91** (2015) 104014 [1501.02198].
- [36] X.-M. Deng, *Probing $f(T)$ gravity with gravitational time advancement*, *Class. Quant. Grav.* **35** (2018) 175013.
- [37] S.-F. Yan, P. Zhang, J.-W. Chen, X.-Z. Zhang, Y.-F. Cai and E. N. Saridakis, *Interpreting cosmological tensions from the effective field theory of torsional gravity*, *Phys. Rev.* **D101** (2020) 121301 [1909.06388].
- [38] A. Paliathanasis, J. L. Said and J. D. Barrow, *Stability of the Kasner Universe in $f(T)$ Gravity*, *Phys. Rev.* **D97** (2018) 044008 [1709.03432].
- [39] S. Bahamonde, C. G. Böhmmer and M. Wright, *Modified teleparallel theories of gravity*, *Phys. Rev.* **D92** (2015) 104042 [1508.05120].
- [40] S. Capozziello, M. Capriolo and M. Transirico, *The gravitational energy-momentum pseudotensor: the cases of $f(R)$ and $f(T)$ gravity*, *Int. J. Geom. Meth. Mod. Phys.* **15** (2018) 1850164 [1804.08530].
- [41] S. Bahamonde and S. Capozziello, *Noether Symmetry Approach in $f(T, B)$ teleparallel cosmology*, *Eur. Phys. J.* **C77** (2017) 107 [1612.01299].
- [42] A. Paliathanasis, *de Sitter and Scaling solutions in a higher-order modified teleparallel theory*, *JCAP* **1708** (2017) 027 [1706.02662].
- [43] G. Farrugia, J. Levi Said, V. Gakis and E. N. Saridakis, *Gravitational Waves in Modified Teleparallel Theories*, *Phys. Rev. D* **97** (2018) 124064 [1804.07365].
- [44] S. Bahamonde, M. Zubair and G. Abbas, *Thermodynamics and cosmological reconstruction in $f(T, B)$ gravity*, *Phys. Dark Univ.* **19** (2018) 78 [1609.08373].
- [45] M. Wright, *Conformal transformations in modified teleparallel theories of gravity revisited*, *Phys. Rev.* **D93** (2016) 103002 [1602.05764].
- [46] G. Farrugia, J. Levi Said and A. Finch, *Gravitoelectromagnetism, Solar System Test and Weak-Field Solutions in $f(T, B)$ Gravity with Observational Constraints*, *Universe* **6** (2020) 34 [2002.08183].
- [47] S. Capozziello, M. Capriolo and L. Caso, *Weak Field Limit and Gravitational Waves in $f(T, B)$ Teleparallel Gravity*, *Eur. Phys. J. C* **80** (2020) 156 [1912.12469].
- [48] C. Escamilla-Rivera and J. Levi Said, *Cosmological viable models in $f(T, B)$ theory as solutions to the H_0 tension*, *Class. Quant. Grav.* **37** (2020) 165002 [1909.10328].
- [49] G. A. R. Franco, C. Escamilla-Rivera and J. Levi Said, *Stability analysis for cosmological models in $f(T, B)$ gravity*, *Eur. Phys. J. C* **80** (2020) 677 [2005.14191].
- [50] G. Kofinas and E. N. Saridakis, *Cosmological applications of $F(T, T_G)$ gravity*, *Phys. Rev.* **D90** (2014) 084045 [1408.0107].
- [51] S. Capozziello, M. De Laurentis and K. F. Dialektopoulos, *Noether symmetries in Gauss–Bonnet-teleparallel cosmology*, *Eur. Phys. J.* **C76** (2016) 629 [1609.09289].
- [52] A. de la Cruz-Dombriz, G. Farrugia, J. L. Said and D. Sáez-Chillón Gómez, *Cosmological*

- bouncing solutions in extended teleparallel gravity theories, *Phys. Rev. D* **97** (2018) 104040 [1801.10085].
- [53] A. de la Cruz-Dombriz, G. Farrugia, J. L. Said and D. Saez-Gomez, *Cosmological reconstructed solutions in extended teleparallel gravity theories with a teleparallel Gauss–Bonnet term*, *Class. Quant. Grav.* **34** (2017) 235011 [1705.03867].
- [54] S. Bahamonde, K. F. Dialektopoulos and J. Levi Said, *Can Horndeski Theory be recast using Teleparallel Gravity?*, *Phys. Rev. D* **100** (2019) 064018 [1904.10791].
- [55] S. Bahamonde, K. F. Dialektopoulos, V. Gakis and J. Levi Said, *Reviving Horndeski theory using teleparallel gravity after GW170817*, *Phys. Rev. D* **101** (2020) 084060 [1907.10057].
- [56] S. Bahamonde, K. F. Dialektopoulos, M. Hohmann and J. Levi Said, *Post-Newtonian limit of Teleparallel Horndeski gravity*, 2003.11554.
- [57] S. Bahamonde, C. G. Böhmer, S. Carloni, E. J. Copeland, W. Fang and N. Tamanini, *Dynamical systems applied to cosmology: dark energy and modified gravity*, *Phys. Rept.* **775–777** (2018) 1 [1712.03107].
- [58] L. Karpathopoulos, S. Basilakos, G. Leon, A. Paliathanasis and M. Tsamparlis, *Cartan symmetries and global dynamical systems analysis in a higher-order modified teleparallel theory*, *Gen. Rel. Grav.* **50** (2018) 79 [1709.02197].
- [59] S. Capozziello, M. De Laurentis and R. Myrzakulov, *Noether Symmetry Approach for teleparallel-curvature cosmology*, *Int. J. Geom. Meth. Mod. Phys.* **12** (2015) 1550095 [1412.1471].
- [60] A. Pourbagher and A. Amani, *Thermodynamics and stability of $f(T, B)$ gravity with viscous fluid by observational constraints*, *Astrophys. Space Sci.* **364** (2019) 140 [1908.11595].
- [61] M. Zubair, S. Waheed, M. Atif Fayyaz and I. Ahmad, *Energy constraints and the phenomenon of cosmic evolution in the $f(T, B)$ framework*, *Eur. Phys. J. Plus* **133** (2018) 452 [1807.07399].
- [62] M. Krššák and E. N. Saridakis, *The covariant formulation of $f(T)$ gravity*, *Class. Quant. Grav.* **33** (2016) 115009 [1510.08432].
- [63] F. W. Hehl, J. D. McCrea, E. W. Mielke and Y. Ne’eman, *Metric affine gauge theory of gravity: Field equations, Noether identities, world spinors, and breaking of dilation invariance*, *Phys. Rept.* **258** (1995) 1 [gr-qc/9402012].
- [64] K. Hayashi and T. Shirafuji, *New General Relativity*, *Phys. Rev. D* **19** (1979) 3524.
- [65] F. W. Hehl, P. Von Der Heyde, G. D. Kerlick and J. M. Nester, *General Relativity with Spin and Torsion: Foundations and Prospects*, *Rev. Mod. Phys.* **48** (1976) 393.
- [66] T. P. Sotiriou and V. Faraoni, *$f(R)$ Theories Of Gravity*, *Rev. Mod. Phys.* **82** (2010) 451 [0805.1726].
- [67] G. A. R. Franco, C. Escamilla-Rivera and J. L. Said, *Stability analysis for cosmological models in $f(T, B)$ gravity*, *The European Physical Journal C* **80** (2020) .
- [68] N. Tamanini and C. G. Boehmer, *Good and bad tetrads in $f(T)$ gravity*, *Phys. Rev.* **D86** (2012) 044009 [1204.4593].
- [69] M. Caruana, G. Farrugia and J. Levi Said, *Cosmological bouncing solutions in $f(T, B)$ gravity*, *Eur. Phys. J. C* **80** (2020) 640 [2007.09925].
- [70] L. Perko, *Differential Equations and Dynamical Systems*. Springer-Verlag GmbH, 2008.
- [71] S. Bahamonde, C. G. Böhmer, S. Carloni, E. J. Copeland, W. Fang and N. Tamanini, *Dynamical systems applied to cosmology: Dark energy and modified gravity*, *Physics Reports* **775–777** (2018) 1.
- [72] D. Arrowsmith and C. Place, *Dynamical Systems: Differential Equations, Maps, and Chaotic Behaviour*, Chapman Hall/CRC Mathematics Series. Taylor & Francis, 1992.
- [73] O. Hrycyna and M. Szydłowski, *Dynamical complexity of the brans-dicke cosmology*, *Journal of Cosmology and Astroparticle Physics* **2013** (2013) 016.
- [74] J. Magaña, M. H. Amante, M. A. Garcia-Aspeitia and V. Motta, *The cardassian expansion revisited: constraints from updated hubble parameter measurements and type ia supernova data*, *Monthly Notices of the Royal Astronomical Society* **476** (2018) 1036.

- [75] C. Escamilla-Rivera, A. Hernández-Almada, M. A. García-Aspeitia and V. Motta, *Linear perturbations spectra for dynamical dark energy inspired by modified gravity*, 2005.13957.
- [76] O. Lahav and A. R. Liddle, *The Cosmological Parameters (2019)*, 1912.03687.
- [77] N. Aghanim, Y. Akrami, M. Ashdown, J. Aumont, C. Baccigalupi, M. Ballardini et al., *Planck 2018 results, Astronomy & Astrophysics* **641** (2020) A6.
- [78] R. Abbott, T. D. Abbott, S. Abraham, F. Acernese, K. Ackley, C. Adams et al., *GW190814: Gravitational waves from the coalescence of a 23 solar mass black hole with a 2.6 solar mass compact object*, *The Astrophysical Journal* **896** (2020) L44.
- [79] S. H. Suyu, V. Bonvin, F. Courbin, C. D. Fassnacht, C. E. Rusu, D. Sluse et al., *H0LiCOW I. H₀ lenses in COSMOGRAIL's wellspring: program overview*, *Monthly Notices of the Royal Astronomical Society* **468** (2017) 2590.