

KNUTSON IDEALS OF GENERIC MATRICES

LISA SECCIA

ABSTRACT. In this paper we show that determinantal ideals of generic matrices are Knutson ideals. This fact leads to a useful result about Gröbner bases of certain sums of determinantal ideals. More specifically, given $I = I_1 + \dots + I_k$ a sum of ideals of minors on adjacent columns or rows, we prove that the union of the Gröbner bases of the I_j 's is a Gröbner basis of I .

1. INTRODUCTION

Let \mathbb{K} be a field of any characteristic. Fix $f \in S = \mathbb{K}[x_1, \dots, x_n]$ a polynomial such that its leading term $\text{in}_{\prec}(f)$ is a squarefree monomial for some term order \prec . We can define many more ideals starting from the principal ideal (f) and taking associated primes, intersections and sums. Thereby, if \mathbb{K} has characteristic p , we obtain a family of ideals which are compatibly split with respect to $\text{Tr}(f^{p-1}\bullet)$ (see [Kn] for more details).

Geometrically this means that we start from the hypersurface defined by f and we construct a family of new subvarieties $\{Y_i\}_i$ by taking irreducible components, intersections and unions.

Definition 1 (Knutson ideals). Let $f \in S = \mathbb{K}[x_1, \dots, x_n]$ be a polynomial such that its leading term $\text{in}_{\prec}(f)$ is a squarefree monomial for some term order \prec . Define \mathcal{C}_f to be the smallest set of ideals satisfying the following conditions:

1. $(f) \in \mathcal{C}_f$;
2. If $I \in \mathcal{C}_f$ then $I : J \in \mathcal{C}_f$ for every ideal $J \subseteq S$;
3. If I and J are in \mathcal{C}_f then also $I + J$ and $I \cap J$ must be in \mathcal{C}_f .

This class of ideals has some interesting properties which were first proved by Knutson in the case $\mathbb{K} = \mathbb{Z}/p\mathbb{Z}$ and then generalized to fields of any characteristic in [Se]:

- i) Every $I \in \mathcal{C}_f$ has a squarefree initial ideal, so every Knutson ideal is radical.
- ii) If two Knutson ideals are different their initial ideals are different. So \mathcal{C}_f is finite.
- iii) The union of the Gröbner bases of Knutson ideals associated to f is a Gröbner basis of their sum.

Remark 1. Actually, assuming that every ideal of \mathcal{C}_f is radical, the second condition in Definition 1 can be replaced by the following:

- 2'. If $I \in \mathcal{C}_f$ then $\mathcal{P} \in \mathcal{C}_f$ for every $\mathcal{P} \in \text{Min}(I)$.

In this paper we will continue the study undertaken in [Se] about Knutson ideals.

So far, it has been proved that determinantal ideals of Hankel matrices are Knutson ideals for a suitable choice of f ([Se, Theorems 3.1,3.2]). As a consequence of these results, one can derive an alternative proof (see [Se, Corollary 3.3]) of the F -purity of

Hankel determinantal rings, a result recently proved in [CMSV].

In this paper we are going to show that also determinantal ideals of generic matrices are Knutson ideals (see Theorem 2.1). In particular, they define F -split rings. This was already known since the 1990's from a result by Hochster and Huneke ([HH]).

As a corollary we obtain an interesting result about Gröbner bases of certain sums of determinantal ideals. More specifically, given $I = I_1 + \dots + I_k$ a sum of ideals of minors on adjacent columns or rows, we will prove that the union of the Gröbner bases of the I_j 's is a Gröbner basis of I (see Corollary 2.4).

Example 1. Let $X = (x_{ij})$ be the generic square matrix of size 6 and consider the ideal

$$J = I_3(X_{[1,3]}) + I_3(X^{[1,3]})$$

in the polynomial ring $S = \mathbb{K}[X]$. Then J is the ideal generated by the 3-minors of the following highlighted ladder

$$X = \begin{bmatrix} x_{11} & x_{12} & x_{13} & x_{14} & x_{15} & x_{16} \\ x_{21} & x_{22} & x_{23} & x_{24} & x_{25} & x_{26} \\ x_{31} & x_{32} & x_{33} & x_{34} & x_{35} & x_{36} \\ x_{41} & x_{42} & x_{43} & x_{44} & x_{45} & x_{46} \\ x_{51} & x_{52} & x_{53} & x_{54} & x_{55} & x_{56} \\ x_{61} & x_{62} & x_{63} & x_{64} & x_{65} & x_{66} \end{bmatrix}.$$

From Corollary 2.4, we get that set of 3-minors that generate J is a Gröbner basis of J with respect to any diagonal term order. Actually, this result was already known for ladder determinantal ideals (see [Na, Corollary 3.4]). Nonetheless, Corollary 2.4 can be applied to more general sums of ideals. Consider for instance the ideal

$$J = I_2(X_{[12]}) + I_2(X^{[12]}) + I_2(X_{[56]}) + I_2(X^{[56]})$$

that is the ideal generated by the 2-minors inside the below coloured region of X

$$X = \begin{bmatrix} x_{11} & x_{12} & x_{13} & x_{14} & x_{15} & x_{16} \\ x_{21} & x_{22} & x_{23} & x_{24} & x_{25} & x_{26} \\ x_{31} & x_{32} & x_{33} & x_{34} & x_{35} & x_{36} \\ x_{41} & x_{42} & x_{43} & x_{44} & x_{45} & x_{46} \\ x_{51} & x_{52} & x_{53} & x_{54} & x_{55} & x_{56} \\ x_{61} & x_{62} & x_{63} & x_{64} & x_{65} & x_{66} \end{bmatrix}.$$

In this case, J is not a ladder determinantal ideal but we can use Corollary 2.4 to prove that the 2-minors that generate J form a Gröbner basis for the ideal J with respect to any diagonal term order. In fact, J is a sum of ideals of the form $I_t(X_{[a,b]})$ or $I_t(X^{[c,d]})$ which are Knutson ideals from Theorem 2.1. Then a Gröbner basis for J is given by the union of their Gröbner bases.

Furthermore, we can also consider sums of ideals of minors of different sizes, such as

$$J = I_2(X_{[2,4]}) + I_3(X^{[2,5]}).$$

In this case, J is generated by the 2-minors of the blue rectangular submatrix and the 3-minors of the red rectangular submatrix illustrated below

$$X = \begin{bmatrix} x_{11} & x_{12} & x_{13} & x_{14} & x_{15} & x_{16} \\ x_{21} & x_{22} & x_{23} & x_{24} & x_{25} & x_{26} \\ x_{31} & x_{32} & x_{33} & x_{34} & x_{35} & x_{36} \\ x_{41} & x_{42} & x_{43} & x_{44} & x_{45} & x_{46} \\ x_{51} & x_{52} & x_{53} & x_{54} & x_{55} & x_{56} \\ x_{61} & x_{62} & x_{63} & x_{64} & x_{65} & x_{66} \end{bmatrix}.$$

Again from Corollary 2.4, being $I_2(X_{[2,4]})$ and $I_3(X^{[2,5]})$ Knutson ideals, the union of their Gröbner bases is a Gröbner basis for J . So, a Gröbner basis of J is given by the 2-minors of $X_{[2,4]}$ and the 3-minors of $X^{[2,5]}$.

Unlike in the case of Hankel matrices, a characterization of all the ideals belonging to the family C_f has not been found yet. A first step towards this result would be to understand the primary decomposition of certain sums belonging to the family. Some known results (see [HS], [MR]) suggest us what these primary decompositions might be and computer experiments seem to confirm this guess. Finding this characterization could lead to interesting properties on the Gröbner bases of determinantal-like ideals and it would also answer to the question asked by F.Mohammadi and J. Rhau in [MR].

Acknowledgments. The author is grateful to her advisor Matteo Varbaro for his valuable comments and precious advice.

2. KNUTSON IDEALS AND DETERMINANTAL IDEALS OF GENERIC MATRICES

Let m, n be two positive integers with $m < n$, we will denote by X_{mn} the generic matrix of size $m \times n$ with entries x_{ij} , that is

$$X_{mn} = \begin{bmatrix} x_{11} & x_{12} & x_{13} & \dots & x_{1n} \\ x_{21} & x_{22} & x_{23} & \dots & x_{2n} \\ x_{31} & x_{32} & x_{33} & \dots & x_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ x_{m1} & x_{m2} & x_{m3} & \dots & x_{mn} \end{bmatrix}.$$

Moreover, for any $1 \leq i < j \leq n$ and $1 \leq k < l \leq m$, we denote by $X_{[i,j]}^{[k,l]}$ the submatrix of X_{mn} with column indices $i, i+1, \dots, j$ and row indices $k, k+1, \dots, l$. In the case $[k, l] = [1, m]$, we omit the superscript and we simply write $X_{[i,j]}$.

Given a generic matrix X_{mn} and an integer $t \leq \min(m, n)$, we will denote by $I_t(X)$ the determinantal ideal in $S = \mathbb{K}[X] = \mathbb{K}[x_{i,j} \mid 1 \leq i \leq m, 1 \leq j \leq n]$ generated by all the t -minors of X .

We are going to prove that determinantal ideals of a generic matrix are Knutson ideals for a suitable choice of the polynomial f .

Theorem 2.1. Let $X = X_{mn}$ be the generic matrix of size $m \times n$ with entries x_{ij} and $m < n$. Consider the polynomial

$$f = \prod_{k=0}^{m-2} \left(\det X_{[1,k+1]}^{[m-k,m]} \cdot \det X_{[n-k,n]}^{[1,k+1]} \right) \cdot \prod_{k=1}^{n-m+1} \left(\det X_{[k,m+k-1]} \right)$$

in $S = \mathbb{K}[x_{ij} \mid 1 \leq i \leq m, 1 \leq j \leq n]$. Then $I_t(X) \in \mathcal{C}_f$ for $t = 1, \dots, m$.

Furthermore, $I_t(X_{[a,b]})$ (respectively, $I_t(X^{[a,b]})$) are Knutson ideals associated to f for $t = 1, \dots, m$ and $0 \leq a < b \leq n$ (respectively, $0 \leq a < b \leq m$) with $b - a + 1 \geq t$.

A first step towards the proof of Theorem 2.1 is showing that all the ideals generated by the t -minors on t adjacent columns are in \mathcal{C}_f . This fact is formally stated in the lemma below.

Lemma 2.2. Let $X = X_{m \times n}$ be the generic square matrix of size $m \times n$ with entries x_{ij} and let f to be as in Theorem 2.1. If we fix $t \leq m$, then:

$$I_t(X_{[i,t+i-1]}) \in \mathcal{C}_f \quad \forall i = 1, \dots, n - t + 1.$$

Proof. It is known (e.g. see [Co] and [BC]) that every determinantal ideal of a generic matrix $X = X_{m \times n}$ is prime and its height is given by the following formula:

$$(1) \quad \text{ht}(I_t(X)) = (n - t + 1)(m - t + 1).$$

Therefore

$$\text{ht}(I_t(X_{[i,t+i-1]})) = (t + i - 1 - i + 1 - t + 1)(m - t + 1) = m - t + 1.$$

We have three possibilities for i .

1st case: $m - t + 1 \leq i \leq n - m + 1$. Then

$$I_t(X_{[i,t+i-1]}) \supseteq \left(\det X_{[i,i+m-1]}, \det X_{[i-1,i+m-2]}, \dots, \det X_{[i-m+t,i+t-1]} \right).$$

2nd case: $i \leq m - t$

$$I_t(X_{[i,t+i-1]}) \supseteq \left(\det X_{[1,m]}, \det X_{[2,m+1]}, \dots, \det X_{[i,i+m-1]}, \det X_{[1,t+i+1]}^{[m-t-i+2,m]}, \right. \\ \left. \det X_{[1,t+i]}^{[m-t-i+1,m]}, \dots, \det X_{[1,m-1]}^{[2,m]} \right).$$

3rd case: $i \geq n - m + 2$

$$I_t(X_{[i,t+i-1]}) \supseteq \left(\det X_{[n-m+1,n]}, \det X_{[n-m,n-1]}, \dots, \det X_{[t+i-m,t+i-1]}, \det X_{[i,n]}^{[1,n-i+1]}, \right. \\ \left. \det X_{[i-1,n]}^{[1,n-i+2]}, \dots, \det X_{[n-m+2,n]}^{[1,m-1]} \right).$$

Define H to be the right hand side ideal for each of the previous cases. Note that the initial ideal of H is given by some of the diagonals of the matrix X . Since these monomials are coprime, this ideal is a complete intersection and

$$\text{ht}(H) = m - t + 1$$

in each of the above mentioned cases. So $I_t(X_{[i,t+i-1]})$ is minimal over H .

By Definition 1, $(f) : J \in \mathcal{C}_f$ for every ideal $J \subseteq S$. Taking J to be the principal ideal generated by the product of some of the factors of f , we have that all the principal ideals

generated by one of the factors of f are Knutson ideal associated to f . Being H a sum of these ideals, $H \in \mathcal{C}_f$.

In conclusion, we get that $I_t(X_{[i,t+i-1]})$ is a minimal prime over an ideal of \mathcal{C}_f . So it is in \mathcal{C}_f . \square

Using Lemma 2.2, we can then prove Theorem 2.1.

Proof. Fix $t \in \{1, \dots, m\}$. We want to prove that $I_t(X) \in \mathcal{C}_f$. By lemma 2.2, we know that $I_t(X_{[1,t]}), I_t(X_{[2,t+1]}) \in \mathcal{C}_f$ and so their sum.

We claim that the minimal prime decomposition of the sum is given by

$$I_t(X_{[1,t]}) + I_t(X_{[2,t+1]}) = I_t(X_{[1,t+1]}) \cap I_{t-1}(X_{[2,t]}).$$

To simplify the notation, we set $I_1 := I_t(X_{[1,t]}), I_2 := I_t(X_{[2,t+1]}), P_1 = I_t(X_{[1,t+1]})$ and $P_2 = I_{t-1}(X_{[2,t]})$. We want to prove that the minimal prime decomposition is given by:

$$I_1 + I_2 = P_1 \cap P_2.$$

We already know that $I_1 + I_2 \subseteq P_1 \cap P_2$. Passing to the correspondent algebraic varieties, we get the reverse inclusion

$$\mathcal{V}(I_1 + I_2) \supseteq \mathcal{V}(P_1 \cap P_2)$$

If we prove that $\mathcal{V}(I_1 + I_2) \subseteq \mathcal{V}(P_1 \cap P_2)$, then

$$\mathcal{V}(I_1 + I_2) = \mathcal{V}(P_1 \cap P_2)$$

and this is equivalent to say that $\sqrt{I_1 + I_2} = \sqrt{P_1 \cap P_2}$. Since $I_1 + I_2 \in \mathcal{C}_f$, it is radical and $P_1 \cap P_2$ is radical because P_1 and P_2 are both radical ideals, then

$$I_1 + I_2 = P_1 \cap P_2$$

and we are done.

For this aim, let $\mathbf{X} \in \mathcal{V}(I_1 + I_2) = \mathcal{V}(I_1) \cap \mathcal{V}(I_2)$. This means that $\mathbf{X}_{[1,t]}$ and $\mathbf{X}_{[2,t+1]}$ have rank less or equal than $t - 1$. Now we consider two cases:

Case 1. Suppose that $\mathbf{X}_{[2,t]}$ has rank less or equal than $t - 2$. This implies that all the $(t - 1) \times (t - 1)$ -minors corresponding to this interval vanish on \mathbf{X} . So $\mathbf{X} \in \mathcal{V}(P_2)$.

Case 2. Suppose that $\mathbf{X}_{[2,t]}$ has full rank, namely $t - 1$. Then it generates a vector space V of dimension $t - 1$. But by assumption, $\mathbf{X}_{[1,t]}$ and $\mathbf{X}_{[2,t+1]}$ have rank less or equal than $t - 1$, so they also generate the vector space V . Consequently, $\mathbf{X}_{[1,t+1]}$ generates the vector space V and this means that all the $t \times t$ -minors of our matrix X vanish on \mathbf{X} . Therefore we have proved that $\mathbf{X} \in \mathcal{V}(P_1)$.

This proves the claim and shows that $I_t(X_{[1,t+1]}) \in \mathcal{C}_f$, being a minimal prime over a Knutson ideal.

In the same way, simply shifting the submatrices, we get that $I_t(X_{[k,t+k]}) \in \mathcal{C}_f$ for every $k = 1, \dots, n - t$.

In particular $I_t(X_{[2,t+2]}) \in \mathcal{C}_f$; therefore the sum $I_t(X_{[1,t+1]}) + I_t(X_{[2,t+2]})$ belongs to \mathcal{C}_f .

Using a similar argument to that used to prove the claim, it can be shown that the primary decomposition of the latter sum is given by

$$I_t(X_{[1,t+1]}) + I_t(X_{[2,t+2]}) = I_t(X_{[1,t+2]}) \cap I_{t-1}(X_{[2,t+1]}).$$

Therefore $I_t(X_{[1,t+2]})$ is a Knutson ideal associated to f .

Again, shifting the submatrices, the same argument shows that $I_t(X_{[k,t+k+1]}) \in \mathcal{C}_f$ for every $k = 1, \dots, n - t - 1$.

Iterating this procedure we get that $I_t(X_{[a,b]}) \in \mathcal{C}_f$ for every $1 \leq a < b \leq n$ such that $b - a + 1 \geq t$. In particular, $I_t(X_{[1,n-1]}), I_t(X_{[2,n]}) \in \mathcal{C}_f$. Hence their sum belongs to \mathcal{C}_f .

Again, one can show that the primary decomposition of the sum is given by

$$I_t(X_{[1,n-1]}) + I_t(X_{[2,n]}) = I_t(X_{[1,n]}) \cap I_{t-1}(X_{[2,n-1]}).$$

This shows that $I_t(X_{[1,n]}) \in \mathcal{C}_f$ and we are done.

Notice that an identical proof shows that $I_t(X^{[a,b]}) \in \mathcal{C}_f$ for every $0 \leq a < b \leq m$ with $b - a + 1 \geq t$. □

As an immediate consequence of the previous theorem, we get an alternative proof of F -purity of determinantal ideals of generic matrices.

Corollary 2.3. *Assume that \mathbb{K} is a field of characteristic p and let X be a generic matrix of size $m \times n$. Then $S/I_t(X)$ is F -pure.*

Proof. We may assume that \mathbb{K} is a perfect field of positive characteristic. In fact, we can always reduce to this case by tensoring with the algebraic closure of \mathbb{K} and the F -purity property descends to the non-perfect case. Using Lemma 4 in [Kn], we know that the ideal (f) is compatibly split with respect to the Frobenius splitting defined by $\text{Tr}(f^{p-1}\bullet)$ (where f is taken to be as in the previous theorems). Thus all the ideals belonging to \mathcal{C}_f are compatibly split with respect to the same splitting, in particular $I_t(X)$. This implies that such Frobenius splitting of S provides a Frobenius splitting of $S/I_t(X)$. Being $S/I_t(X)$ F -split, it must be also F -pure. □

Furthermore, we obtain an interesting and useful result about Gröbner bases of certain sums of determinantal ideals.

Corollary 2.4. *Let X be a generic matrix of size $m \times n$ and let I be a sum of ideals, say $I = I_1 + I_2 + \dots + I_k$, where each I_i is of the form either $I_{t_i}(X_{[a_i,b_i]})$ or $I_{t_i}(X^{[a_i,b_i]})$. Then*

$$\mathcal{G}_I = \mathcal{G}_{I_1} \cup \mathcal{G}_{I_2} \cup \dots \cup \mathcal{G}_{I_k}$$

where \mathcal{G}_J denotes a Gröbner basis of the ideal J .

Furthermore, if \mathbb{K} has positive characteristic, I is also F -pure.

Proof. By Theorem 2.1, we know that $I_{t_i}(X_{[a_i,b_i]})$ and $I_{t_i}(X^{[a_i,b_i]})$ are Knutson ideals. From property (iii) of Knutson ideals, we get the thesis. □

REFERENCES

- [BC] W.Bruns, A.Conca, *Algebra of minors*, Journal of Algebra 246, 311–330, 2001
- [BH] W.Bruns, J. Herzog, *Cohen-Macaulay rings*, Cambridge University Press, 1993.
- [CMSV] A.Conca, M. Mostafazadehfard, A. K. Singh, M.Varbaro, *Hankel determinantal rings have rational singularities*, Adv. Math. 335 (2018), 111–129.
- [Co] A. Conca, *Straightening law and powers of determinantal ideals of Hankel matrices*, Adv. Math. 138 (1998), no. 2, 263–292.
- [CV] A.Conca, M.Varbaro, *Squarefree Grobner degeneration*, arXiv:1805.11923, 2018.
- [HH] M.Hochster, C.Huneke, *Tight closure of parameter ideals and splitting in module-finite extensions*, Journal of Algebraic Geometry 3 (4), p. 599–670, 1994.

- [HS] S.Hosten, S.Sullivant *Ideals of adjacent minors*, J.Algebra vol.277,no.2, 2004.
- [Kn] A.Knutson, *Frobenius splitting, point-counting, and degeneration*, arXiv:0911.4941, 2009.
- [MR] F.Mohammadi, J.Rauh, *Prime splitting of determinantal ideals*, Communications in Algebra, 46:5, p. 2278-2296, 2018.
- [Na] H.Narasimhan, *The irreducibility of ladder determinantal varieties*, J. Algebra 102, p. 162-185, (1986).
- [Se] L.Seccia, *Knutson ideals and determinantal ideals of Hankel matrices*, arXiv:2003.14232 , 2020.

UNIVERSITÀ DI GENOVA, DIPARTIMENTO DI MATEMATICA. VIA DODECANESO 35, 16146 GENOVA, ITALY

Email address: `seccia@dima.unige.it`