

On Graded 1-Absorbing Prime Submodules

Ahmad KA'ABNEH and Rashid ABU-DAWWAS

ABSTRACT. Let G be a group with identity e , R be a commutative G -graded ring with unity 1 and M be a G -graded unital R -module. In this article, we introduce the concept of graded 1-absorbing prime submodule. A proper graded R -submodule N of M is said to be a graded 1-absorbing prime R -submodule of M if for all non-unit homogeneous elements x, y of R and homogeneous element m of M with $xym \in N$, either $xy \in (N :_R M)$ or $m \in N$. We show that the new concept is a generalization of graded prime submodules at the same time it is a special graded 2-absorbing submodule. Several properties of a graded 1-absorbing prime submodule have been obtained. We investigate graded 1-absorbing prime submodules when the components $\{M_g : g \in G\}$ are multiplication R_e -modules.

1. Introduction

Throughout this article, G will be a group with identity e and R a commutative ring with nonzero unity 1. Then R is said to be G -graded if $R = \bigoplus_{g \in G} R_g$ with $R_g R_h \subseteq R_{gh}$ for all $g, h \in G$ where R_g is an additive subgroup of R for all $g \in G$. The elements of R_g are called homogeneous of degree g . If $x \in R$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in R_g . Moreover, R_e is a subring of R and $1 \in R_e$ and $h(R) = \bigcup_{g \in G} R_g$. Let I be an ideal of a graded ring R . Then I is said to be graded ideal if $I = \bigoplus_{g \in G} (I \cap R_g)$, i.e., for $x \in I$, $x = \sum_{g \in G} x_g$ where $x_g \in I$ for all $g \in G$. An ideal of a graded ring need not be graded.

Assume that M is a left unital R -module. Then M is said to be G -graded if $M = \bigoplus_{g \in G} M_g$ with $R_g M_h \subseteq M_{gh}$ for all $g, h \in G$ where M_g is an additive subgroup of M for all $g \in G$. The elements of M_g are called homogeneous of degree g . It is clear that M_g is an R_e -submodule of M for all $g \in G$. If $x \in M$, then x can be written uniquely as $\sum_{g \in G} x_g$, where x_g is the component of x in M_g . Moreover, $h(M) = \bigcup_{g \in G} M_g$. Let N be an R -submodule of a graded R -module M . Then N is

2020 *Mathematics Subject Classification.* Primary 16W50; Secondary 13A02.

Key words and phrases. Graded 1-absorbing prime ideal; graded 1-absorbing prime submodule; graded prime submodule.

said to be graded R -submodule if $N = \bigoplus_{g \in G} (N \cap M_g)$, i.e., for $x \in N$, $x = \sum_{g \in G} x_g$ where $x_g \in N$ for all $g \in G$. An R -submodule of a graded R -module need not be graded.

For more details and terminology, see [12] and [14].

LEMMA 1.1. ([10], Lemma 2.1) *Let R be a G -graded ring and M be a G -graded R -module.*

- (1) *If I and J are graded ideals of R , then $I + J$ and $I \cap J$ are graded ideals of R .*
- (2) *If N and K are graded R -submodules of M , then $N + K$ and $N \cap K$ are graded R -submodules of M .*
- (3) *If N is a graded R -submodule of M , $r \in h(R)$, $x \in h(M)$ and I is a graded ideal of R , then Rx , IN and rN are graded R -submodules of M . Moreover, $(N :_R M) = \{r \in R : rM \subseteq N\}$ is a graded ideal of R .*

Similarly, if M is a graded R -module, N a graded R -submodule of M and $m \in h(M)$, then $(N :_R m)$ is a graded ideal of R . Also, in particular, $\text{Ann}_R(M) = (0_M :_R M)$ is a graded ideal of R .

The concept of graded prime ideals and its generalizations have a significant place in graded commutative algebra since they are used in understanding the structure of graded rings. Recall that a proper graded ideal P of R is said to be graded prime ideal if $x, y \in h(R)$ such that $xy \in P$ implies $x \in P$ or $y \in P$ ([16]). Graded prime ideals have been extended to graded modules in [6]. A proper graded R -submodule N of M is said to be a graded prime submodule if whenever $x \in h(R)$ and $m \in h(M)$ with $xm \in N$, then $m \in N$ or $x \in (N :_R M)$. The notion of graded 2-absorbing ideal, which is a generalization of graded prime ideal, was introduced in [4] as the following: a proper graded ideal P of R is said to be graded 2-absorbing if whenever $x, y, z \in h(R)$ such that $xyz \in P$, then either $xy \in P$ or $xz \in P$ or $yz \in P$. Also, Graded 2-absorbing ideals have been deeply studied in [13]. Graded 2-absorbing submodules have been introduced and studied in [3]. A proper graded R -submodule N of a graded R -module M is said to be graded 2-absorbing if whenever $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in N$, then either $xm \in N$ or $ym \in N$ or $xy \in (N :_R M)$. Graded 2-absorbing submodules have been generalized into graded n -absorbing submodules in [11]. Actually, the concept of graded 2-absorbing submodule is a generalization of graded prime submodules.

Recently, in [2], a new class of graded ideals has been introduced, which is an intermediate class of graded ideals between graded prime ideals and graded 2-absorbing ideals. A proper graded ideal P of R is said to be a graded 1-absorbing prime ideal if for each non-units $x, y, z \in h(R)$ with $xyz \in P$, then either $xy \in P$ or $z \in P$. Clearly, every graded prime ideal of R is graded 1-absorbing prime and every graded 1-absorbing prime ideal of R is graded 2-absorbing. The next example shows that not every graded 2-absorbing ideal of R is graded 1-absorbing prime.

EXAMPLE 1.2. [2] *Consider $R = \mathbb{Z} + 3x\mathbb{Z}[x]$ and $G = \mathbb{Z}$. Then R is G -graded by $R_0 = \mathbb{Z}$, $R_j = 3\mathbb{Z}x^j$ for $j \geq 1$ and $R_j = \{0\}$ otherwise. Now, $P = 3x\mathbb{Z}[x]$ is a graded ideal of R , so by Lemma 1.1, P^2 is a graded ideal of R . By ([18], Example 2.2), P^2 is a 2-absorbing ideal of R , and then it is graded 2-absorbing. On the other hand, P^2 is not graded 1-absorbing prime as $3, 9, 3x^2 \in h(R)$ such that $(3)(9)(3x^2) \in P^2$ while $(3)(9) \notin P^2$ and $3x^2 \notin P^2$.*

Thus we have the following chain:

graded prime ideals \Rightarrow *graded 1-absorbing prime ideals* \Rightarrow *graded 2-absorbing ideals*.

On the other hand, we have another chain:

graded prime submodules \Rightarrow *graded 2-absorbing submodules*.

Thus we realize that there is a missing part in the second chain, which is between graded prime submodules and graded 2-absorbing submodules. Then we define the missing part of the chain as graded 1-absorbing prime submodules. In this article, we are motivated by [17] to introduce and study the concept of graded 1-absorbing prime submodule. A proper graded R -submodule N of M is said to be a graded 1-absorbing prime R -submodule of M if for all non-unit elements $x, y \in h(R)$ and $m \in h(M)$ with $xym \in N$, either $xy \in (N :_R M)$ or $m \in N$. We show that the new concept is a generalization of graded prime submodules at the same time it is a special graded 2-absorbing submodule. Several properties of a graded 1-absorbing prime submodule have been obtained. We investigate graded 1-absorbing prime submodules when the components $\{M_g : g \in G\}$ are multiplication R_e -modules.

2. Graded 1-Absorbing Prime Submodules

In this section, we introduce and study the concept of graded 1-absorbing prime submodules.

DEFINITION 2.1. *Let M be a graded R -module and N be a proper graded R -submodule of M . Then N is said to be a graded 1-absorbing prime R -submodule of M if for all non-unit elements $x, y \in h(R)$ and $m \in h(M)$ with $xym \in N$, either $xy \in (N :_R M)$ or $m \in N$.*

EXAMPLE 2.2. *Let R be a graded local ring with graded maximal ideal X satisfies $X^2 = \{0\}$. If M is a graded R -module, then every proper graded R -submodule of M is graded 1-absorbing prime. To prove that, let N be a proper graded R -submodule of M , $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in N$. Since $xy \in X^2 = \{0\}$, $xy \in (N :_R M)$, and hence N is a graded 1-absorbing prime R -submodule of M .*

PROPOSITION 2.3. *Let M be a graded R -module and N be a graded R -submodule of M .*

- (1) *If N is a graded prime R -submodule of M , then N is a graded 1-absorbing prime R -submodule of M .*
- (2) *If N is a graded 1-absorbing prime R -submodule of M , then N is a graded 2-absorbing R -submodule of M .*

PROOF. (1) Let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in N$. Then $z = xy \in h(R)$ with $zm \in N$. Since N is graded prime, either $z = xy \in (N :_R M)$ or $m \in N$. Hence, N is a graded 1-absorbing prime R -submodule of M .

- (2) Let $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in N$. If x is unit, then $ym \in N$. If y is unit, then $xm \in N$. Suppose that x and y are non-units. Since N is graded 1-absorbing prime, either $xy \in (N :_R M)$ or $m \in N$, as needed. Hence, N is a graded 2-absorbing R -submodule of M . □

The next example shows that the converse of Proposition 2.3 (1) is not true in general.

EXAMPLE 2.4. Consider $R = \mathbb{Z}_4$, $M = \mathbb{Z}_4[x]$ and $G = \mathbb{Z}$. Then R is G -graded by $R_0 = \mathbb{Z}_4$ and $R_j = \{0\}$ otherwise. Also, M is G -graded by $M_j = \mathbb{Z}_4x^j$ for $j \geq 0$ and $M_j = \{0\}$ otherwise. As $x \in h(M)$, $N = \langle x \rangle$ is a graded R -submodule of M . By Example 2.2, N is a graded 1-absorbing prime R -submodule of M . On the other hand, N is not graded prime R -submodule of M .

The next example shows that the converse of Proposition 2.3 (2) is not true in general.

EXAMPLE 2.5. Consider $R = \mathbb{Z}$, $M = \mathbb{Z}_{30}[i]$ and $G = \mathbb{Z}_2$. Then R is G -graded by $R_0 = \mathbb{Z}$ and $R_1 = \{0\}$. Also, M is G graded by $M_0 = \mathbb{Z}_{30}$ and $M_1 = i\mathbb{Z}_{30}$. Then $N = \langle \bar{6} \rangle$ is a graded 2-absorbing R -submodule of M . On the other hand, $2 \in h(R)$ is non-unit and $\bar{3} \in h(M)$ with $(2)(2)(\bar{3}) \in N$, but $(2)(2) \notin (N :_R M)$ and $\bar{3} \notin N$. Hence, N is not graded 1-absorbing prime R -submodule of M .

So, we have the following chain:

$$\text{graded prime submodules} \Rightarrow \text{graded 1-absorbing prime submodules} \Rightarrow \text{graded 2-absorbing submodules.}$$

And the converse of each implication is not true in general.

PROPOSITION 2.6. Let M be a graded R -module and N be a graded R -submodule of M . If N is a graded 1-absorbing prime R -submodule of M , then $(N :_R M)$ is a graded 1-absorbing prime ideal of R .

PROOF. Let $x, y, z \in h(R)$ be non-units such that $xyz \in (N :_R M)$. Assume that $m \in M$. Then $xyzm_g \in N$ for all $g \in G$. Since N is graded 1-absorbing prime, either $xy \in (N :_R M)$ or $zm_g \in N$ for all $g \in G$. If $zm_g \in N$ for all $g \in G$, then $zm = z \left(\sum_{g \in G} m_g \right) = \sum_{g \in G} zm_g \in N$, which implies that $z \in (N :_R M)$. Hence, $(N :_R M)$ is a graded 1-absorbing prime ideal of R . \square

Similarly, one can prove the following:

PROPOSITION 2.7. Let M be a graded R -module and N be a graded R -submodule of M . If N is a graded 1-absorbing prime R -submodule of M , then $(N :_R m)$ is a graded 1-absorbing prime ideal of R for all $m \in h(M) - N$.

The next example shows that the converse of Proposition 2.6 is not true in general.

EXAMPLE 2.8. Consider $R = \mathbb{Z}$, $M = \mathbb{Z} \times \mathbb{Z}$ and $G = \mathbb{Z}_2$. Then R is G -graded by $R_0 = \mathbb{Z}$ and $R_1 = \{0\}$. Also, M is G -graded by $M_0 = \mathbb{Z} \times \{0\}$ and $M_1 = \{0\} \times \mathbb{Z}$. As $(3, 0) \in h(M)$, $N = \langle (3, 0) \rangle$ is a graded R -submodule of M such that $(N :_R M) = \{0\}$ is a graded prime ideal of R , so $(N :_R M)$ is a graded 1-absorbing prime ideal of R . On the other hand, $3, 2 \in h(R)$ are non-units and $(1, 0) \in h(M)$ with $(3)(2)(1, 0) \in N$, but $(3)(2) \notin (N :_R M)$ and $(1, 0) \notin N$. Hence, N is not graded 1-absorbing prime R -submodule of M .

LEMMA 2.9. Let M be a graded R -module and N be a graded 1-absorbing prime R -submodule of M . For a graded R -submodule L of M and for non-unit elements $x, y \in h(R)$, if $xyL \subseteq N$, then either $xy \in (N :_R M)$ or $L \subseteq N$.

PROOF. Assume that $L \not\subseteq N$. Then there is an element $0 \neq m \in L - N$, and then there exists $g \in G$ such that $m_g \notin N$. Note that, $m_g \in L$ as L is a graded

R -submodule. By assumption, we have $xym_g \in N$. Since N is a graded 1-absorbing prime R -submodule, either $xy \in (N :_R M)$ or $m_g \in N$. The second choice implies a contradiction, we conclude that $xy \in (N :_R M)$, as desired. \square

THEOREM 2.10. *Let M be a graded R -module and N be a proper graded R -submodule of M . Then N is a graded 1-absorbing prime R -submodule of M if and only if whenever I, J are proper graded ideals of R and L is a graded R -submodule of M such that $IJL \subseteq N$, then either $IJ \subseteq (N :_R M)$ or $L \subseteq N$.*

PROOF. Suppose that N is a graded 1-absorbing prime R -submodule of M . Assume that I, J are proper graded ideals of R and L is a graded R -submodule of M such that $IJL \subseteq N$. Suppose that $IJ \not\subseteq (N :_R M)$. Then there exist $x \in I$ and $y \in J$ such that $xy \notin (N :_R M)$, and then there exist $g, h \in G$ such that $x_g y_h \notin (N :_R M)$. Note that, $x_g \in I$ and $y_h \in J$ as I and J are graded ideals. Since I and J are proper, x_g and y_h are non-units. Now, $x_g y_h L \subseteq N$, so by Lemma 2.9, $L \subseteq N$, as needed. Conversely, let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in N$. Then $I = \langle x \rangle$, $J = \langle y \rangle$ are proper graded ideals of R and $L = \langle m \rangle$ is a graded R -submodule of M such that $IJL \subseteq N$. By assumption, either $IJ \subseteq (N :_R M)$ or $L \subseteq N$. If $IJ \subseteq (N :_R M)$, then $xy \in (N :_R M)$. Hence, N is a graded 1-absorbing prime R -submodule of M . \square

Let M be a G -graded R -module, N be a graded R -submodule of M and $g \in G$ such that $N_g \neq M_g$. In [6], N is said to be a g -prime R -submodule of M if whenever $r \in R_e$ and $m \in M_g$ such that $rm \in N$, then either $r \in (N :_R M)$ or $m \in N$. Also, in [2], A graded ideal P of R is said to be a g -1-absorbing prime ideal of R if $P_g \neq R_g$ and whenever non-unit elements x, y, z in R_g such that $xyz \in P$, then $xy \in P$ or $z \in P$. We introduce the following definition:

DEFINITION 2.11. *Let M be a G -graded R -module, N be a graded R -submodule of M and $g \in G$ such that $N_g \neq M_g$. Then N is said to be a g -1-absorbing prime R -submodule of M if for all non-unit elements $x, y \in R_e$ and $m \in M_g$ with $xym \in N$, either $xy \in (N :_R M)$ or $m \in N$.*

REMARK 2.12. *Clearly, every g -prime R -submodule is g -1-absorbing prime. However, the converse is not true in general; $N = \langle x \rangle$ in Example 2.4 is a graded 1-absorbing prime R -submodule, so it is a g -1-absorbing prime R -submodule for all $g \in G = \mathbb{Z}$, but if we choose $g = 1$, then N will be not graded g -prime R -submodule. Now, we are going to prove that if R_e is not local ring, then a graded R -submodule will be an e -1-absorbing prime if and only if it is an e -prime R -submodule.*

LEMMA 2.13. *Let M be a graded R -module. If M has an e -1-absorbing prime R -submodule that is not an e -prime R -submodule, then the sum of every non-unit element of R_e and every unit element of R_e is a unit element of R_e .*

PROOF. Let N be an e -1-absorbing prime R -submodule of M that is not an e -prime R -submodule. Then there exist a non-unit $r \in R_e$ and $m \in M_g$ such that $rm \in N$ but $r \notin (N :_R M)$ and $m \notin N$. Choose a non-unit element $a \in R_e$. Then we have that $ram \in N$ and $m \notin N$. Since N is e -1-absorbing prime, $ra \in (N :_R M)$. Let $u \in R_e$ be a unit element. Assume that $a+u$ is non-unit. Then $r(a+u)m \in N$. As N is e -1-absorbing prime, $r(a+u) \in (N :_R M)$. This means that $ru \in (N :_R M)$, i.e., $r \in (N :_R M)$, which is a contradiction. Thus, we have $a+u$ is a unit element. \square

THEOREM 2.14. *Let M be a graded R -module. If M has an e -1-absorbing prime R -submodule that is not an e -prime R -submodule, then R_e is a local ring.*

PROOF. By Lemma 2.13, the sum of every non-unit element of R_e and every unit element of R_e is a unit element of R_e , and then by ([8], Lemma 1), R_e is a local ring. \square

COROLLARY 2.15. *Let R be a graded ring such that R_e is not local ring. Suppose that M is a graded R -module. Then a graded R -submodule N of M is an e -1-absorbing prime R -submodule if and only if N is an e -prime R -submodule of M .*

PROPOSITION 2.16. *Let $\{N_k\}_{k \in \Delta}$ be a chain of graded 1-absorbing prime R -submodules of M . Then $\bigcap_{k \in \Delta} N_k$ is a graded 1-absorbing prime R -submodule of M .*

PROOF. Let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in \bigcap_{k \in \Delta} N_k$. Suppose that $m \notin \bigcap_{k \in \Delta} N_k$. Then $m \notin N_i$ for some $i \in \Delta$. Since $xym \in N_i$ and N_i is graded 1-absorbing prime, $xy \in (N_i :_R M)$. For any $k \in \Delta$, if $N_i \subseteq N_k$, then $xy \in (N_k :_R M)$, and then $xy \in \left(\bigcap_{k \in \Delta} N_k :_R M \right)$. If $N_k \subseteq N_i$, then $m \notin N_k$. Since $xym \in N_k$ and N_k is graded 1-absorbing prime, $xy \in (N_k :_R M)$, and then $xy \in \left(\bigcap_{k \in \Delta} N_k :_R M \right)$. Hence, $\bigcap_{k \in \Delta} N_k$ is a graded 1-absorbing prime R -submodule of M . \square

LEMMA 2.17. *Let M be a graded R -module. If M is finitely generated, then the union of every chain of proper graded R -submodules of M is a proper graded R -submodule of M .*

PROOF. Suppose that $M = \langle x_1, x_2, \dots, x_n \rangle$ for some $x_1, \dots, x_n \in M$. Let $\{N_k\}_{k \in \Delta}$ be a chain of proper graded R -submodules of M . If $\bigcup_{k \in \Delta} N_k = M$, then $x_1, x_2 \in \bigcup_{k \in \Delta} N_k$, and then $x_1 \in N_j$ and $x_2 \in N_r$ for some $j, r \in \Delta$. If $N_j \subseteq N_r$, then $x_1, x_2 \in N_r$. If $N_r \subseteq N_j$, then $x_1, x_2 \in N_j$. In fact, this works for all x_i 's. So, there exists $i \in \Delta$ such that N_i contains all the generators x_1, \dots, x_n of M , and then $N_i = M$, which is a contradiction. \square

PROPOSITION 2.18. *Let $\{N_k\}_{k \in \Delta}$ be a chain of graded 1-absorbing prime R -submodules of M . If M is finitely generated, then $\bigcup_{k \in \Delta} N_k$ is a graded 1-absorbing prime R -submodule of M .*

PROOF. By Lemma 2.17, $\bigcup_{k \in \Delta} N_k$ is a proper graded R -submodule of M . Let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in \bigcup_{k \in \Delta} N_k$. Suppose that $m \notin \bigcup_{k \in \Delta} N_k$. Then $m \notin N_k$ for all $k \in \Delta$. Now, since $xym \in \bigcup_{k \in \Delta} N_k$, $xym \in N_i$ for some $i \in \Delta$, and since N_i is graded 1-absorbing prime, $xy \in (N_i :_R M) \subseteq$

$\left(\bigcup_{k \in \Delta} N_k :_R M\right)$. Hence, $\bigcup_{k \in \Delta} N_k$ is a graded 1-absorbing prime R -submodule of M . \square

Let M and S be two G -graded R -modules. An R -homomorphism $f : M \rightarrow S$ is said to be a graded R -homomorphism if $f(M_g) \subseteq S_g$ for all $g \in G$ ([14]).

LEMMA 2.19. ([1], Lemma 2.16) Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules. If K is a graded R -submodule of S , then $f^{-1}(K)$ is a graded R -submodule of M .

LEMMA 2.20. ([7], Lemma 4.8) Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules. If L is a graded R -submodule of M , then $f(L)$ is a graded R -submodule of $f(M)$.

PROPOSITION 2.21. Suppose that $f : M \rightarrow S$ is a graded R -homomorphism of graded R -modules.

- (1) If K is a graded 1-absorbing prime R -submodule of S and $f^{-1}(K) \neq M$, then $f^{-1}(K)$ is a graded 1-absorbing prime R -submodule of M .
- (2) Assume that f is a graded epimorphism. If L is a graded 1-absorbing prime R -submodule of M containing $\text{Ker}(f)$, then $f(L)$ is a graded 1-absorbing prime R -submodule of S .

PROOF. (1) By Lemma 2.19, $f^{-1}(K)$ is a graded R -submodule of M . Let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in f^{-1}(K)$. Then $f(m) \in h(S)$ such that $xyf(m) = f(xym) \in K$. Since K is graded 1-absorbing prime, either $xy \in (K :_R S)$ or $f(m) \in K$, and then either $xy \in (f^{-1}(K) :_R M)$ or $m \in f^{-1}(K)$. Hence, $f^{-1}(K)$ is a graded 1-absorbing prime R -submodule of M .

- (2) By Lemma 2.20, $f(L)$ is a graded R -submodule of S . Let $x, y \in h(R)$ be non-units and $s \in h(S)$ such that $xys \in f(L)$. Since f is graded epimorphism, there exists $m \in h(M)$ such that $s = f(m)$, and then $f(xym) = xyf(m) = xys \in f(L)$, which implies that $f(xym) = f(t)$ for some $t \in L$, and then $xym - t \in \text{Ker}(f) \subseteq L$, which gives that $xym \in L$. Since L is graded 1-absorbing prime, either $xy \in (L :_R M)$ or $m \in L$, and then either $xy \in (f(L) :_R S)$ or $s = f(m) \in f(L)$. Hence, $f(L)$ is a graded 1-absorbing prime R -submodule of S . \square

If M is a G -graded R -module and L is a graded R -submodule of M , then M/L is a G -graded R -module by $(M/L)_g = (M_g + L)/L$ for all $g \in G$ ([14]).

COROLLARY 2.22. Let M be a graded R -module and $L \subseteq N$ be graded R -submodules of M . If N is a graded 1-absorbing prime R -submodule of M , then N/L is a graded 1-absorbing prime R -submodule of M/L .

PROOF. Define $f : M \rightarrow M/L$ by $f(x) = x + L$. Then f is a graded epimorphism with $\text{Ker}(f) = L \subseteq N$. So, by Proposition 2.21, $f(N) = N/L$ is a graded 1-absorbing prime R -submodule of M/L . \square

DEFINITION 2.23. Let M be a graded R -module, L be a proper graded R -submodule of M and N be a graded 1-absorbing prime R -submodule of M with $L \subseteq N$. Then N is said to be a minimal graded 1-absorbing prime R -submodule with respect to L if there is no a graded 1-absorbing prime R -submodule P of M such that $L \subseteq P \subset N$.

THEOREM 2.24. *Let M be a graded R -module and L be a proper graded R -submodule of M . If N is a graded 1-absorbing prime R -submodule of M such that $L \subseteq N$, then there exists a minimal graded 1-absorbing prime R -submodule with respect to L that it is contained in N .*

PROOF. Let X be the set of all graded 1-absorbing prime R -submodules N_i of M such that $L \subseteq N_i \subseteq N$. Then X is non-empty as it contains N . Consider (X, \supseteq) . Let $\{N_k\}_{k \in \Delta}$ be a chain in X . Then by Proposition 2.16, $\bigcap_{k \in \Delta} N_k$ is a graded 1-absorbing prime R -submodule of M , and then by Zorn's Lemma, X has a maximal element K . So, K is a graded 1-absorbing prime R -submodule of M such that $L \subseteq K \subseteq N$. If K is not minimal graded 1-absorbing prime R -submodule with respect to L , then there exists a graded 1-absorbing prime R -submodule P of M such that $L \subseteq P \subseteq K$, and then $P \in X$, which implies that $K \subseteq P$, and hence $P = K$. Thus, K is a minimal graded 1-absorbing prime R -submodule with respect to L . \square

Let I be a graded ideal of R . Then $\text{Grad}(I)$ is the intersection of all graded prime ideals of R containing I ([5]). Similarly, if N is a graded R -submodule of M , then the graded radical of N is $\text{Grad}(N)$ which is the intersection of all graded prime R -submodules of M containing N . Motivated by this, we have the following definition.

DEFINITION 2.25. (1) *Let I be a graded ideal of R . Then the 1-graded radical of I is the intersection of all graded 1-absorbing prime ideals of R containing I , and is denoted by $\text{Grad}_1(I)$. If $I = R$ or R has no graded 1-absorbing prime ideals, we define $\text{Grad}_1(I) = R$.*
 (2) *Let N be a graded R -submodule of M . Then the 1-graded radical of N is the intersection of all graded 1-absorbing prime R -submodules of M containing N , and is denoted by $\text{Grad}_1(N)$. If $N = M$ or M has no graded 1-absorbing prime R -submodules, we define $\text{Grad}_1(N) = M$.*

REMARK 2.26. *Since every graded prime ideal is graded 1-absorbing prime, then $\text{Grad}_1(I) \subseteq \text{Grad}(I)$ for all graded ideal I of R . Similarly, $\text{Grad}_1(N) \subseteq \text{Grad}(N)$ for all graded R -submodule N of M .*

Motivated by ([15], Proposition 1.2), we state the following proposition, and the proof is elementary.

PROPOSITION 2.27. *Let M be a graded R -module and N, K be two graded R -submodules of M . Then the following statements hold:*

- (1) $N \subseteq \text{Grad}_1(N)$.
- (2) $\text{Grad}_1(\text{Grad}_1(N)) \subseteq \text{Grad}_1(N)$.
- (3) $\text{Grad}_1(N \cap K) \subseteq \text{Grad}_1(N) \cap \text{Grad}_1(K)$.

PROPOSITION 2.28. *Let M be a graded R -module and N be a graded R -submodule of M . Then $\text{Grad}_1((N :_R M)) \subseteq (\text{Grad}_1(N) :_R M)$.*

PROOF. If $\text{Grad}_1(N) = M$, then it is done. Suppose that $\text{Grad}_1(N) \neq M$. Then there exists a graded 1-absorbing prime R -submodule L of M such that $N \subseteq L$, and then by Proposition 2.6, $(L :_R M)$ is a graded 1-absorbing prime ideal of R such that $(N :_R M) \subseteq (L :_R M)$, which implies that $\text{Grad}_1((N :_R M)) \subseteq (L :_R M)$, and hence $\text{Grad}_1((N :_R M))M \subseteq L$. In fact, in similar way, $\text{Grad}_1((N :_R M))M \subseteq L_i$ for every

graded 1-absorbing prime R -submodule L_i of M such that $N \subseteq L_i$. So, $\text{Grad}_1((N :_R M))M \subseteq \text{Grad}_1(N)$, which means that $\text{Grad}_1((N :_R M)) \subseteq (\text{Grad}_1(N) :_R M)$. \square

PROPOSITION 2.29. *Let R be a graded ring and I be a graded ideal of R . If I is a graded 1-absorbing prime ideal of R , then I_e is a 1-absorbing prime ideal of R_e .*

PROOF. Let $x, y, z \in R_e$ be non-units such that $xyz \in I_e$. Then $x, y, z \in h(R)$ such that $xyz \in I$. Since I is graded 1-absorbing prime, either $xy \in I$ or $z \in I$. If $xy \in I$, then $xy \in R_e R_e \cap I \subseteq R_e \cap I = I_e$. If $z \in I$, then $z \in R_e \cap I = I_e$. Hence, I_e is a 1-absorbing prime ideal of R_e . \square

PROPOSITION 2.30. *Let M be a G -graded R -module and N be a graded R -submodule of M . If N is a graded 1-absorbing prime R -submodule of M , then N_g is a 1-absorbing prime R_e -submodule of M_g for all $g \in G$.*

PROOF. Let $g \in G$. Suppose that $x, y \in R_e$ be non-units and $m \in M_g$ such that $xym \in N_g$. Then $x, y \in h(R)$ and $m \in h(M)$ such that $xym \in N$. Since N is graded 1-absorbing prime, either $xy \in (N :_R M)$ or $m \in N$. If $xy \in (N :_R M)$, then $xy \in R_e$ such that $xyM_g \subseteq xyM \subseteq N$, also, $xyM_g \subseteq R_e M_g \subseteq M_g$, and hence $xyM_g \subseteq N \cap M_g = N_g$, which implies that $xy \in (N_g :_{R_e} M_g)$. If $m \in N$, then $m \in N \cap M_g = N_g$. Hence, N_g is a 1-absorbing prime R_e -submodule of M_g . \square

Let M be a G -graded R -module and $g \in G$. In the sense of [9], M_g is said to be a multiplication R_e -module if whenever N is an R_e -submodule of M_g , then $N = IM_g$ for some ideal I of R_e . Note that, since $I \subseteq (N :_{R_e} M_g)$, $N = IM_g \subseteq (N :_{R_e} M_g)M_g \subseteq N$. So, if M_g is a multiplication R_e -module, then $N = (N :_{R_e} M_g)M_g$ for every R_e -submodule N of M_g .

PROPOSITION 2.31. *Let M be a G -graded R -module and $g \in G$ such that M_g is a faithful multiplication R_e -module. Suppose that I is a 1-absorbing prime ideal of R_e . Then whenever $x, y \in R_e$ are non-units and $m \in M_g$ such $xym \in IM_g$, then either $xy \in I$ or $m \in IM_g$.*

PROOF. Apply ([17], Theorem 5) on the R_e -module M_g . \square

THEOREM 2.32. *Let M be a G -graded R -module and $g \in G$ such that M_g is a faithful multiplication R_e -module. If I is a graded 1-absorbing prime ideal of R and $(IM)_g \neq M_g$, then IM is a g -1-absorbing prime R -submodule of M .*

PROOF. By Lemma 1.1 (3), IM is a graded R -submodule of M . Since I is a graded 1-absorbing prime ideal of R , by Proposition 2.29, I_e is a 1-absorbing prime ideal of R_e . Let $x, y \in R_e$ be non-units and $m \in M_g$ such that $xym \in IM$. Then $xym = x_e y_e m_g = (xym)_g \in (IM)_g = I_e M_g$. By Proposition 2.31, either $xy \in I_e$ or $m \in I_e M_g \subseteq IM$. If $xy \in I_e$, then $xy \in I$ and then $xyM \subseteq IM$, which implies that $xy \in (IM :_R M)$. Hence, IM is a g -1-absorbing prime R -submodule of M . \square

PROPOSITION 2.33. *Let M be a graded R -module and N be a graded 1-absorbing prime $R/\text{Ann}_R(M)$ -submodule of M as M is a graded $R/\text{Ann}_R(M)$ -module. If $U(R/\text{Ann}_R(M)) = \{r + \text{Ann}_R(M) : r \in U(R)\}$, then N is a graded 1-absorbing prime R -submodule of M as M is a graded R -module.*

PROOF. Let $x, y \in h(R)$ be non-units and $m \in h(M)$ such that $xym \in N$. Then $x + \text{Ann}_R(M), y + \text{Ann}_R(M) \in h(R/\text{Ann}_R(M))$ are non-units such that $(x + \text{Ann}_R(M))(y + \text{Ann}_R(M))m = xym + \text{Ann}_R(M)m \in N$. Since N is a graded 1-absorbing prime $R/\text{Ann}_R(M)$ -submodule of M , either $xy + \text{Ann}_R(M) \in$

$(N :_{R/Ann_R(M)} M)$ or $m \in N$. If $xy + Ann_R(M) \in (N :_{R/Ann_R(M)} M)$, then $xyM \subseteq N$, which implies that $xy \in (N :_R M)$. Hence, N is a graded 1-absorbing prime R -submodule of M as M is a graded R -module. \square

THEOREM 2.34. *Let M be a G -graded R -module and $g \in G$ such that M_g is a multiplication R_e -module and $U(R_e/Ann_{R_e}(M_g)) = \{r + Ann_{R_e}(M_g) : r \in U(R_e)\}$. Assume that I is a graded 1-absorbing prime ideal of R containing $Ann_R(M)$. Then $(I_e/Ann_{R_e}(M_g))M_g$ is a g -1-absorbing prime R -submodule of M .*

PROOF. Since M_g is a multiplication R_e -module, M_g is a faithful multiplication $R_e/Ann_{R_e}(M_g)$ -module (see [9], page 759). Also, since I is a graded 1-absorbing prime ideal of R , I_e is a 1-absorbing prime ideal of R_e by Proposition 2.29, and then $I_e/Ann_{R_e}(M_g)$ is a 1-absorbing prime ideal of $R_e/Ann_{R_e}(M_g)$ by ([17], Proposition 7). So, $(I_e/Ann_{R_e}(M_g))M_g$ is a g -1-absorbing prime $R_e/Ann_{R_e}(M_g)$ -submodule of M by Theorem 2.32, which implies that $(I_e/Ann_{R_e}(M_g))M_g$ is a g -1-absorbing prime R -submodule of M by ([17], Proposition 8). \square

References

- [1] R. Abu-Dawwas, M. Bataineh and H. Shashan, Graded generalized 2-absorbing submodules, *Beiträge zur Algebra und Geometrie / Contributions to Algebra and Geometry*, (2020), <https://doi.org/10.1007/s13366-020-00544-1>.
- [2] R. Abu-Dawwas, Ü. Tekir, S. Koç and E. Yıldız, Graded 1-absorbing prime ideals, accepted in the *Sao Paulo Journal of Mathematical Sciences*, in press.
- [3] K. Al-Zoubi and R. Abu-Dawwas, On graded 2-absorbing and weakly graded 2-absorbing submodules, *Journal of Mathematical Sciences: Advances and Applications*, 28 (2014), 45-60.
- [4] K. Al-Zoubi, R. Abu-Dawwas and S. Çeken, On graded 2-absorbing and graded weakly 2-absorbing ideals, *Hacettepe Journal of Mathematics and Statistics*, 48 (3) (2019), 724-731.
- [5] K. Al-Zoubi and F. Qarqaz, An intersection condition for graded prime ideals, *Bollettino dell'Unione Matematica Italiana*, 11 (2018), 483-488.
- [6] S. E. Atani, On graded prime submodules, *Chiang Mai Journal of Science*, 33 (1) (2006), 3-7.
- [7] S. E. Atani and F. E. K. Saraei, Graded modules which satisfy the gr-radical formula, *Thai Journal of Mathematics*, 8 (1) (2010), 161-170.
- [8] A. Badawi and E. Y. Celikel, On 1-absorbing primary ideals of commutative rings, *Journal of Algebra and Its Applications*, 19 (6), 2050111 (2020).
- [9] Z. A. El-Bast and P. F. Smith, Multiplication modules, *Communications in Algebra*, 16 (1988), 755-779.
- [10] F. Farzalipour, P. Ghiasvand, On the union of graded prime submodules, *Thai Journal of Mathematics*, 9 (1) (2011), 49-55.
- [11] M. Hamoda and A. E. Ashour, On graded n -ansorbing submodules, *Le Matematiche*, (2015), 243-254, doi:10.4418/2015.70.2.16.
- [12] R. Hazrat, *Graded rings and graded Grothendieck groups*, Cambridge University press, 2016.
- [13] S. R. Naghani and H. F. Moghimi, On graded 2-absorbing and graded weakly 2-absorbing ideals of a commutative ring, *Çankaya University Journal of Science and Engineering*, 13 (2) (2016), 11-17.
- [14] C. Nastasescu and F. Oystaeyen, *Methods of graded rings*, Lecture Notes in Mathematics, 1836, Springer-Verlag, Berlin, 2004.
- [15] M. Refai and K. Al-Zoubi, On graded primary ideals, *Turkish Journal of Mathematics*, 28 (2004), 217-229.
- [16] M. Refai, M. Hailat and S. Obiedat, Graded radicals and graded prime spectra, *Far East Journal of Mathematical Sciences*, (2000), 59-73.
- [17] E. A. Ugurlu, On 1-absorbing prime submodules, (2020), arXiv:2007.01103v1.
- [18] A. Yassine, M. J. Nikmehr, R. Nikandish, On 1-absorbing prime ideals of commutative rings, *Journal of Algebra and its Applications*, (2020), doi:10.1142/S0219498821501759.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF JORDAN, AMMAN, JORDAN
Email address: `ahmadkabeya@yahoo.com`

DEPARTMENT OF MATHEMATICS, YARMOUK UNIVERSITY, IRBID, JORDAN
Email address: `rrashid@yu.edu.jo`