# Least resolved trees for two-colored best match graphs

David Schaller<sup>1,2</sup>, Manuela Geiß<sup>3</sup>, Marc Hellmuth<sup>4</sup>, and Peter F. Stadler<sup>1,2,5,6,7</sup>

<sup>1</sup>Max Planck Institute for Mathematics in the Sciences, Inselstraße 22, D-04103 Leipzig, Germany

sdavid@bioinf.uni-leipzig.de

<sup>2</sup>Bioinformatics Group, Department of Computer Science & Interdisciplinary Center for Bioinformatics, Universität Leipzig,

Härtelstraße 16–18, D-04107 Leipzig, Germany

studla@bioinf.uni-leipzig.de

 $^3{\rm Software}$  Competence Center Hagenberg GmbH, Software<br/>park 21, A-4232 Hagenberg, Austria

manuela.geiss@scch.at

<sup>4</sup>Department of Mathematics, Faculty of Science, Stockholm University, SE - 106 91 Stockholm, Sweden marc.hellmuth@math.su.se

<sup>5</sup>Institute for Theoretical Chemistry, University of Vienna, Währingerstraße 17, A-1090 Wien, Austria

<sup>6</sup>Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia

<sup>7</sup>The Santa Fe Institute, 1399 Hyde Park Rd., Santa Fe, NM 87501, USA

#### Abstract

2-colored best match graphs (2-BMGs) form a subclass of sink-free bi-transitive graphs that appears in phylogenetic combinatorics. There, 2-BMGs describe evolutionarily most closely related genes between a pair of species. They are explained by a unique least resolved tree (LRT). Introducing the concept of support vertices we derive an  $O(|V| + |E|\log^2 |V|)$ -time algorithm to recognize 2-BMGs and to construct its LRT. The approach can be extended to also recognize binary-explainable 2-BMGs with the same complexity. An empirical comparison emphasizes the efficiency of the new algorithm.

### 1 Introduction

Best match graphs recently have been introduced in phylogenetic combinatorics to formalize the notion of a gene y in species 2 being an evolutionary closest relative of a gene x in species 1, i.e., y is a best match for x [5]. The best matches between genes of two species form a bipartite directed graph, the 2-colored best match graph or 2-BMG, that is determined by the phylogenetic tree describing the evolution of the genes. 2-BMGs are characterized by four local properties [5, 8] that relate them to previously studied classes of digraphs:

**Definition 1.** A bipartite digraph  $\vec{G} = (L, E)$  is a 2-BMG if it satisfies

- (N0) Every vertex has at least one out-neighbor, i.e.,  $\vec{G}$  is sink-free.
- (N1) If u and v are two independent vertices, then there exist no vertices w and t such that  $(u,t), (v,w), (t,w) \in E$ .
- (N2) For any four vertices  $u_1, u_2, v_1, v_2$  with  $(u_1, v_1), (v_1, u_2), (u_2, v_2) \in E$  we have  $(u_1, v_2) \in E$ , *i.e.*,  $\vec{G}$  is bi-transitive.
- (N3) For any two vertices u and v with a common out-neighbor, if there exists no vertex w such that either  $(u, w), (w, v) \in E$ , or  $(v, w), (w, u) \in E$ , then u and v have the same in-neighbors and either all out-neighbors of u are also out-neighbors of v or all out-neighbors of v are also out-neighbors of u.

Sink-free graphs have appeared in particular in the context of graph semigroups [1] and graph orientation problems [3]. Bi-transitive graphs were introduced in [4] in the context of oriented bipartite graphs and investigated in more detail in [7, 8]. The class of graphs satisfying (N1), (N2), and (N3) are characterized by a system of forbidden induced subgraphs [12], see Thm. 2 below.

In general, best match graphs (BMGs) are defined as vertex-colored digraphs  $(\bar{G}, \sigma)$ , where the vertex coloring  $\sigma$  assigns to each gene x the species  $\sigma(x)$  in which it resides. The subgraphs of a



Figure 1: Example for a 2-BMG  $(\vec{G}, \sigma)$  and its explaining least resolved tree  $(T^*, \sigma)$ .

BMG induced by vertices of two distinct colors form a 2-BMG. Note that in this context the vertex coloring is assigned a priori, while Def. 1 induces a coloring that is unique only up to relabeling of the colors independently on each (weakly) connected component of  $\vec{G}$ . For each BMG  $(\vec{G}, \sigma)$ , there is a unique least resolved leaf-colored tree  $(T^*, \sigma)$  with leaves corresponding to the vertices of  $(\vec{G}, \sigma)$  such that the arcs in  $(\vec{G}, \sigma)$  are the best matches w.r.t.  $(T^*, \sigma)$  (cf. Def. 2 below). Fig. 1 shows an example for a 2-BMG together with its least resolved tree. Using certain sets of rooted triples that can be inferred from the 2-colored induced subgraphs of  $(\vec{G}, \sigma)$  with three vertices, it is possible to determine whether  $(G, \sigma)$  is a BMG in polynomial time and, if so, to construct the least resolved tree  $(T^*, \sigma)$  [5, 9]. This work also describes  $O(|V|^3)$ -time algorithms for the recognition of 2-BMGs and the construction of the LRT for a given 2-BMG.

In this contribution, we derive an alternative characterization of 2-BMGs that avoids the use of rooted triples. This will give rise to an alternative, efficient algorithm for the recognition of 2-BMGs and the construction of the least resolved tree. The contribution is organized as follows: In Sec. 2, we introduce the necessary notation and review some results from the published literature that are needed later on. Sec. 3 is concerned with a more detailed analysis of the least resolved trees (LRTs) of BMGs with an arbitrary number of colors. We then turn to the peculiar properties of the LRTs of 2-BMGs in Sec. 4. To this end, we introduce the concept of "support leaves" that uniquely determine the LRT. The main result of this section is Thm. 4, which shows that the support leaves of the root can be identified directly in the 2-BMG. In Sec. 5, we then turn Thm. 4 into an efficient algorithm for recognizing 2-BMGs and constructing their LRTs. Computational experiments demonstrate the performance gain in practise. In Sec. 6 we extend the algorithmic approach to binary-explainable 2-BMGs, a subclass that features an additional forbidden induced subgraph.

### 2 Preliminaries

Let T = (V, E) be a tree with root  $\rho$  and leaf set  $L := L(T) \subset V$ . The set of inner vertices of T is  $V^0(T) := V \setminus L$ , in particular  $\rho$  is an inner vertex. An edge  $e = uv \in E(T)$  is called an *inner* edge of T if u and v are both inner vertices. Otherwise it is called an *outer* edge. We consider leaf-colored trees  $(T, \sigma)$  and write  $\sigma(L') := \{\sigma(v) \mid v \in L'\}$  for subsets  $L' \subseteq L$ . A vertex  $u \in V$  is an *ancestor* of  $v \in V$  in T, in symbols  $v \preceq_T u$ , if u lies on the path from  $\rho$  to v. For the edges  $uv \in E(T)$  we use the convention that  $uv \in E$ ,  $v \prec_T u$ , v is a child of u. We write  $\operatorname{child}_T(u)$  for the set of children of u in T and T(u) for the subtree of T rooted in u. The least common ancestor  $\operatorname{lca}_T(A)$  is the unique  $\preceq_T$ -smallest vertex that is an ancestor of all genes in A. Writing  $\operatorname{lca}_T(x, y) := \operatorname{lca}_T(\{x, y\})$ , we have

**Definition 2.** Let  $(T, \sigma)$  be a leaf-colored tree. A leaf  $y \in L(T)$  is a best match of the leaf  $x \in L(T)$  if  $\sigma(x) \neq \sigma(y)$  and  $\operatorname{lca}(x, y) \preceq_T \operatorname{lca}(x, y')$  holds for all leaves y' of color  $\sigma(y') = \sigma(y)$ .

Given  $(T, \sigma)$ , the graph  $\vec{G}(T, \sigma) = (V, E)$  with vertex set V = L(T), vertex coloring  $\sigma$ , and with arcs  $(x, y) \in E$  if and only if y is a best match of x w.r.t.  $(T, \sigma)$  is called the *best match graph* (BMG) of  $(T, \sigma)$  [5].

**Definition 3.** An arbitrary vertex-colored graph  $(\vec{G}, \sigma)$  is a best match graph (BMG) if there exists a leaf-colored tree  $(T, \sigma)$  such that  $(\vec{G}, \sigma) = \vec{G}(T, \sigma)$ . In this case, we say that  $(T, \sigma)$  explains  $(\vec{G}, \sigma)$ .

**Theorem 1** ([5], Thm. 9). If  $(\vec{G}, \sigma)$  is a BMG, then there is a unique least-resolved tree  $(T, \sigma)$  that explains  $(\vec{G}, \sigma)$ .

We say that  $(\vec{G}, \sigma)$  is an  $\ell$ -BMG if  $\sigma : V(\vec{G}) \to S$  is surjective and  $|S| = \ell$ . Given a directed graph  $\vec{G} = (V, E)$  we denote the set of out-neighbors of a vertex  $x \in V$  by  $N(x) := \{y \in V | (x, y) \in E(\vec{G})\}$  and the out-degree |N(x)| of x by outdeg(x). Similarly,  $N^{-}(x) := \{y \in V | (y, x) \in E(\vec{G})\}$  denotes the set of in-neighbors. By construction, the coloring  $\sigma$  of a BMGs  $(\vec{G}, \sigma)$  is proper, i.e.,  $x \in N(y)$  implies  $\sigma(x) \neq \sigma(y)$ , and there is at least one best match of x for every color  $s \in \sigma(V) \setminus \{\sigma(x)\}$ . In particular, therefore, we have  $N(x) \neq \emptyset$  for every 2-BMG, i.e., every 2-BMG is sink-free. Note that BMGs will in general have sources, i.e.,  $N^{-}(x)$  may be empty. We write  $\vec{G}[W]$  for the subgraph of  $\vec{G} = (V, E)$  induced by  $W \subseteq V$  and  $\vec{G} - W$  for  $\vec{G}[V \setminus W]$ . A directed graph is (weakly) connected if its underling undirected graph is connected. A connected component is a maximal connected subgraph of  $\vec{G}$ .

Following [13] we say that T' is displayed by T, in symbols  $T' \leq T$ , if the tree T' can be obtained from a subtree of T by contraction of edges. For leaf-colored trees we say that  $(T, \sigma)$  displays or is a refinement of  $(T', \sigma')$ , whenever  $T' \leq T$  and  $\sigma(v) = \sigma'(v)$  for all  $v \in L(T')$ .

**Definition 4.** An edge  $e \in E(T)$  is redundant with respect to  $\vec{G}(T,\sigma)$  if the tree  $T_e$  obtained by contracting the edge e satisfies  $\vec{G}(T_e,\sigma) = \vec{G}(T,\sigma)$ .

We will need the following characterization of redundant edges:

**Lemma 1** ([11], Lemma 2.10). Let  $(\vec{G}, \sigma)$  be a BMG explained by a tree  $(T, \sigma)$ . The edge e = uvin  $(T, \sigma)$  is redundant w.r.t.  $(\vec{G}, \sigma)$  if and only if (i) e is an inner edge of T and (ii) there is no arc  $(a, b) \in E(\vec{G})$  such that  $lca_T(a, b) = v$  and  $\sigma(b) \in \sigma(L(T(u)) \setminus L(T(v)))$ .

In the following we will frequently need the restriction of the coloring  $\sigma$  on  $\vec{G}$  or L(T) to a subset of vertices or leaves. Since in situations like  $(G_i, \sigma_{|V(G_i)})$  the set to which  $\sigma$  is restricted is clear, we will write  $\sigma_{|}$  to keep the notation less cluttered.

BMGs can also be understood in terms of their connected components:

**Proposition 1** ([5], Prop. 1). A digraph  $(\vec{G}, \sigma)$  is an  $\ell$ -BMG if and only if all its connected components are  $\ell$ -BMGs.

As a simple consequence of Prop. 1 and by definition of  $\ell$ -BMGs, all connected components  $(G_i, \sigma_{|.})$  and  $(G_j, \sigma_{|.})$  of an  $\ell$ -BMG satisfy  $\sigma(V(G_i)) = \sigma(V(G_j))$  and  $|\sigma(V(G_j))| = \ell$ . For our purposes it will also be important to relate the structure of a tree  $(T, \sigma)$  to the connectedness of the BMG  $\vec{G}(T, \sigma)$  that it explains.

**Proposition 2** ([5], Thm. 1). Let  $(T, \sigma)$  be a leaf-labeled tree and  $\vec{G}(T, \sigma)$  its BMG. Then  $\vec{G}(T, \sigma)$  is connected if and only if there is a child v of the root  $\rho$  such that  $\sigma(L(T(v))) \neq \sigma(L(T))$ . Furthermore, if  $\vec{G}(T, \sigma)$  is not connected, then for every connected component  $\vec{G}_i$  of  $\vec{G}(T, \sigma)$  there is a child v of the root  $\rho$  such that  $V(G_i) \subseteq L(T(v))$ .

Moreover, 2-BMGs can be characterized by three types of forbidden subgraphs [12]. To this end we will need the following classes of small bipartite graphs:

Definition 5 (F1-, F2-, and F3-graphs).

- (F1) A properly 2-colored graph on four distinct vertices  $V = \{x_1, x_2, y_1, y_2\}$  with coloring  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y_1) = \sigma(y_2)$  is an F1-graph if  $(x_1, y_1), (y_2, x_2), (y_1, x_2) \in E$  and  $(x_1, y_2), (y_2, x_1) \notin E$ .
- (F2) A properly 2-colored graph on four distinct vertices  $V = \{x_1, x_2, y_1, y_2\}$  with coloring  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y_1) = \sigma(y_2)$  is an F2-graph if  $(x_1, y_1), (y_1, x_2), (x_2, y_2) \in E$  and  $(x_1, y_2) \notin E$ .
- (F3) A properly 2-colored graph on five distinct vertices  $V = \{x_1, x_2, y_1, y_2, y_3\}$  with coloring  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y_1) = \sigma(y_2) = \sigma(y_3)$  is an F3-graph if  $(x_1, y_1), (x_2, y_2), (x_1, y_3), (x_2, y_3) \in E$  and  $(x_1, y_2), (x_2, y_1) \notin E$ .

**Theorem 2** ([12], Thm. 3.4). A properly 2-colored graph is a 2-BMG if and only if it is sink-free and does not contain an induced F1-, F2-, or F3-graph.

As noted in [12], the forbidden induced F1-, F2-, and F3-subgraphs characterize exactly the class of bipartite directed graphs satisfying the Axioms (N1), (N2), and (N3) mentioned in the introduction.

Although we aim at avoiding the use of triples in the final results, we will need them during our discussion. A triple ab|c is a rooted tree t on three pairwise distinct vertices  $\{a, b, c\}$  such that  $lca_t(a, b) \prec_t lca_t(a, c) = lca_t(b, c) = \rho$ , where  $\rho$  denotes the root of t. A set  $\mathcal{R}$  of triples is consistent

if there is a tree T that displays all triples in  $\mathcal{R}$ . Given a vertex-colored graph  $(\vec{G}, \sigma)$ , we define its set of *informative triples* [5, 11] as

$$\mathcal{R}(\vec{G},\sigma) \coloneqq \left\{ ab|b': \sigma(a) \neq \sigma(b) = \sigma(b'), \ (a,b) \in E(\vec{G}); \ (a,b') \notin E(\vec{G}) \right\}.$$
(1)

**Lemma 2** ([11], Lemma 2.8 and 2.9). If  $(\vec{G}, \sigma)$  is a BMG, then every tree  $(T, \sigma)$  that explains  $(\vec{G}, \sigma)$  displays all triples  $t \in \Re(\vec{G}, \sigma)$ .

Moreover, if the triples ab|b' and cb'|b are informative for  $(\vec{G}, \sigma)$ , then every tree  $(T, \sigma)$  that explains  $(\vec{G}, \sigma)$  contains two distinct children  $v_1, v_2 \in \text{child}_T(\text{lca}_T(a, c))$  such that  $a, b \prec_T v_1$  and  $b', c \prec_T v_2$ .

**Observation 3.** Let  $(T, \sigma)$  be a tree explaining the BMG  $(\vec{G}, \sigma)$ , and  $v \in V(T)$  a vertex such that  $\sigma(L(T(v))) = \sigma(L(T))$ . Then  $(a, b) \in E(\vec{G})$  and  $a \in L(T(v))$  implies  $b \in L(T(v))$ .

Finally, there is a close connection between subtrees of T and subgraphs of  $\vec{G}(T, \sigma)$ . We have

**Lemma 3** ([9], Lemma 22 and 23). Let  $(T, \sigma)$  be a tree explaining an BMG  $(\vec{G}, \sigma)$ . Then  $\vec{G}(T(u), \sigma_{|.}) = (\vec{G}[L(T(u))], \sigma_{|.})$  holds for every  $u \in V(T)$ . Moreover, if  $(T, \sigma)$  is least resolved for  $(\vec{G}, \sigma)$ , then the subtree T(u) is least resolved for  $\vec{G}(T(u), \sigma_{|.})$ .

### **3** Properties of Least Resolved Trees

In this short section we derive some helpful properties of LRTs which we will use repeatedly throughout this work.

**Lemma 4.** Let  $(\vec{G}, \sigma)$  be a BMG and  $(T, \sigma)$  its least resolved tree. Then the BMG  $\vec{G}(T(v), \sigma_{|.})$  is connected for every  $v \in V(T)$  with  $v \prec_T \rho_T$ .

Proof. By Lemma 3,  $\vec{G}(T(v), \sigma_{|.})$  is a BMG. First observe that the BMG  $\vec{G}(T(v), \sigma_{|.})$  is trivially connected if v is a leaf. Now let  $v \prec_T \rho_T$  be an arbitrary inner vertex of T. Thus, there exists a vertex  $u \succ_T v$  such that uv is an inner edge. Since  $(T, \sigma)$  is least resolved, it does not contain any redundant edges. Hence, by contraposition of Lemma 1, there is an arc  $(a,b) \in E(\vec{G})$  such that  $\operatorname{lca}_T(a,b) = v$  and  $\sigma(b) \in \sigma(L(T(u)) \setminus L(T(v)))$ . Since  $a, b \in L(T(v))$ , Lemma 3 implies that (a,b) is also an arc in  $\vec{G}(T(v), \sigma_{|.})$ . Moreover,  $\operatorname{lca}_{T(v)}(a, b) = v$  clearly also holds in the subtree rooted at v. Now consider the child  $w \in \operatorname{child}_{T(v)}(v)$  such that  $a \preceq_{T(v)} w$ . There cannot be a leaf  $b' \in L(T(w))$  with  $\sigma(b') = \sigma(b)$  since otherwise  $\operatorname{lca}_{T(v)}(a, b') \preceq_{T(v)} w \prec_{T(v)} v$  would contradict that (a, b) is an arc in  $\vec{G}(T(v), \sigma_{|.})$ . Thus  $\sigma(b) \notin \sigma(L(T(w)))$ . Since  $\sigma(b) \in \sigma(L(T(v)))$ , we thus conclude  $\sigma(L(T(w))) \neq \sigma(L(T(v)))$ . The latter together with Prop. 2 implies that  $\vec{G}(T(v), \sigma_{|.})$  is connected.

The converse of Lemma 4, however, is not true, i.e., a tree  $(T, \sigma)$  for which  $\vec{G}(T(v), \sigma_{|.})$  is connected for every  $v \in V(T)$  with  $v \prec_T \rho_T$  is not necessarily least resolved. To see this, consider the caterpillar tree  $(T, \sigma)$  given by (x'', (x', (x, y))) with  $\sigma(x) = \sigma(x') = \sigma(x'') \neq \sigma(y)$  and  $u = lca_T(x, x')$ . It is an easy task to verify that the BMG of each subtree of T is connected. However, the edge  $\rho_T u$  is redundant.

**Lemma 5.** Let  $(T, \sigma)$  be the least resolved tree of some BMG  $(\vec{G}, \sigma)$ . Then every vertex  $v \prec_T \rho_T$  with  $|\sigma(L(T(v)))| = 1$  is a leaf.

*Proof.* Let  $v \prec_T \rho_T$  with  $|\sigma(L(T(v)))| = 1$  and assume, for contradiction, that v is not a leaf. Hence, |L(T(v))| > 1. By Lemma 3  $\vec{G}(T(v), \sigma_{|.})$  is a BMG and, therefore, properly colored. But then  $\vec{G}(T(v), \sigma_{|.})$  is disconnected; a contradiction to Lemma 4.

As a consequence we find

**Corollary 1.** Let  $(T, \sigma)$  be the least resolved tree of some BMG  $(\tilde{G}, \sigma)$ . Then any vertex  $v \in V(T)$  with  $v \prec_T \rho_T$  is an inner vertex if and only if  $|\sigma(L(T(v)))| > 1$ .

*Proof.* If  $|\sigma(L(T(v)))| = 1$ , Lemma 5 implies that v is a leaf. Otherwise, if  $|\sigma(L(T(v)))| > 1$ , T(v) clearly must contain at least two leaves and thus v cannot be a leaf.

### 4 Support Leaves

In this section we introduce "support leaves" as a means to recursively construct the LRT of a 2-BMG. The main result of this section shows that these leaves can be inferred directly from the BMG without any further knowledge of the corresponding LRT. We start with a technical result similar to Cor. 3 in [5]; here we use a much simpler, more convenient notation.

**Lemma 6.** Let  $(T, \sigma)$  be the least resolved tree of a 2-colored BMG  $(\vec{G}, \sigma)$ . Then, for every vertex  $u \in V^0(T) \setminus \{\rho_T\}$ , it holds  $\operatorname{child}_T(u) \cap L(T) \neq \emptyset$ . If  $(\vec{G}, \sigma)$  is connected, then  $\operatorname{child}_T(u) \cap L(T) \neq \emptyset$  holds for every  $u \in V^0(T)$ .

Proof. Suppose first that  $(\vec{G}, \sigma)$  is disconnected and let  $u \in V^0(T) \setminus \{\rho_T\}$ . Since  $(T, \sigma)$  is least resolved, Lemma 4 implies that  $\vec{G}(T(u), \sigma_{|.})$  is connected for every  $u \in V(T)$  with  $u \prec_T \rho_T$ . Hence, we can apply Prop. 2 to  $\vec{G}(T(u), \sigma_{|.})$  and conclude that there is a child  $v \in \text{child}_{T(u)}(u)$  such that  $\sigma(L(T(v))) \neq \sigma(L(T(u)))$ , hence in particular  $\sigma(L(T(v))) \subsetneq \sigma(L(T(u)))$ . Since  $(T, \sigma)$  is 2colored, the latter immediately implies  $|\sigma(L(T(v)))| = 1$  and, by Cor. 1, v is a leaf. Thus every  $u \in V^0(T) \setminus \{\rho_T\}$  has a leaf v among its children, i.e.  $\text{child}_T(u) \cap L(T) \neq \emptyset$ . If in addition  $(\vec{G}, \sigma)$  is connected, we can apply the same argumentation to  $u = \rho_T$  and conclude that a leaf v is attached to  $\rho_T$ .

Lemma 6 states that, in the least resolved tree of a connected 2-colored BMG, every inner vertex u is adjacent to at least one leaf, and thus in a way "supported" by it.

**Definition 6** (Support Leaves). For a given tree T, the set  $S_u := \text{child}_T(u) \cap L(T)$  is the set of all support leafs of vertex  $u \in V(T)$ .

Note that Lemma 6 is in general not true for  $\ell$ -BMGs with  $\ell \geq 3$ , as exemplified by the (least-resolved) tree ((a, b), (c, a')) with three distinct leaf colors  $\sigma(a) = \sigma(a') \neq \sigma(b) \neq \sigma(c)$ .

As a simple consequence of Prop. 2 and Cor. 1, we find

**Corollary 2.** Let  $(T, \sigma)$  be the least resolved tree (with root  $\rho$ ) of some 2-colored BMG  $\vec{G}(T, \sigma)$ . Then,  $\vec{G}(T, \sigma)$  is connected if and only if  $S_{\rho} \neq \emptyset$ .

*Proof.* By Prop. 2,  $\vec{G}(T, \sigma)$  is connected if and only if there exists a child v of the root  $\rho$  of T,  $v \in \text{child}_T(\rho)$ , such that T(v) does not contain all colors. Thus  $|\sigma(L(T(v)))| = 1$ . By Cor. 1, we have  $|\sigma(L(T(v)))| = 1$  if and only if v is a leaf, i.e.  $v \in S_{\rho}$ . Hence,  $\vec{G}(T, \sigma)$  is connected if and only if  $S_{\rho} \neq \emptyset$ .

**Lemma 7.** Let  $(T, \sigma)$  be the least resolved tree of a 2-BMG  $(\vec{G}, \sigma)$ , and  $S_{\rho}$  the set of support leaves of the root  $\rho$ . Then the connected components of  $(\vec{G} - S_{\rho}, \sigma_{|.})$  are exactly the BMGs  $\vec{G}(T(v), \sigma_{|.})$ with  $v \in \text{child}(\rho) \setminus S_{\rho}$ .

Proof. Let  $v \in \operatorname{child}_T(\rho) \cap V^0(T) = \operatorname{child}_T(\rho) \setminus S_\rho$  and consider the BMG  $\vec{G}(T(v), \sigma_{|.})$ . By Lemma 4 and Lemma 3,  $\vec{G}(T(v), \sigma_{|.})$  is connected and we have  $\vec{G}(T(v), \sigma_{|.}) = (\vec{G}[L(T(v))], \sigma_{|.})$ . Moreover, it holds  $((\vec{G} - S_\rho)[L(T(v))], \sigma_{|.}) = (\vec{G}[L(T(v))], \sigma_{|.})$  since  $L(T(v)) = V(\vec{G}[L(T(v))]) = V(H[L(T(v))])$  for  $H \coloneqq \vec{G} - S_\rho = \vec{G}[V(\vec{G}) \setminus S_\rho]$ .

If  $\operatorname{child}_T(\rho) \setminus S_\rho = \{v\}$ , then the statement is trivially satisfied. Therefore, suppose that  $|\operatorname{child}_T(\rho) \setminus S_\rho| > 1$ . Hence, it remains to show that there are no arcs between  $\vec{G}(T(v), \sigma_{|})$  and  $\vec{G}(T(w), \sigma_{|})$  for any  $w \in \operatorname{child}_T(\rho) \setminus S_\rho$ ,  $w \neq v$ . Cor. 1 and  $v \prec_T \rho$  imply that T(v) contains both colors. Thus, by Obs. 3, there are no out-arcs to any vertex in  $L(T) \setminus L(T(v))$ , hence in particular there are no out-arcs (x, y) with  $x \preceq_T v$ ,  $y \preceq_T w$ . By symmetry, the same holds for w, thus we can conclude that there are no arcs (y, x). From the observation that each  $x \in L(T) \setminus S_\rho$  must be located below some  $v \in \operatorname{child}_T(\rho) \cap V^0(T)$ , it now immediately follows that  $(\vec{G} - S_\rho, \sigma_{|})$  consists exactly of these connected components as stated.  $\Box$ 

As a consequence, we have

**Corollary 3.** Let  $(T, \sigma)$  with root  $\rho$  be the LRT of a 2-BMG  $(\vec{G}, \sigma)$ . Then each child of  $\rho$  is either one of the support leaves  $S_{\rho}$  of  $\rho$  or the root of the LRT for a connected component of  $(\vec{G} - S_{\rho}, \sigma_{\parallel})$ .

*Proof.* Let  $(T, \sigma)$  with root  $\rho$  be the least resolved tree for  $(\vec{G}, \sigma)$ . The support leaves  $S_{\rho}$  are children of  $\rho$  by definition. By Lemma 7, the connected components of  $(\vec{G} - S_{\rho}, \sigma_{|.})$  are exactly the BMGs  $\vec{G}(T(v), \sigma_{|.})$  with  $v \in \text{child}_T(\rho) \setminus S_{\rho}$ . Moreover, by Lemma 3, the subtrees T(v) with  $v \in \text{child}_T(\rho) \setminus S_{\rho}$  are exactly the unique LRTs for these BMGs.

In order to use this property as a means of constructing the LRT in a recursive manner, we need to identify the support leaves of the root  $S_{\rho}$  directly from the 2-BMG  $(\vec{G}, \sigma)$  without constructing the LRT first. To this end, we consider the set of *umbrella vertices*  $U(\vec{G}, \sigma)$  comprising all vertices x for which N(x) consists of all vertices of  $V(\vec{G})$  that have the color distinct from  $\sigma(x)$ .

**Definition 7** (Umbrella Vertices). For an arbitrary 2-colored graph  $(\vec{G}, \sigma)$ , the set

$$U(\vec{G},\sigma) \coloneqq \left\{ x \in V(\vec{G}) \mid y \in N(x) \text{ if } \sigma(y) \neq \sigma(x) \text{ and } y \in V(\vec{G}) \right\}$$

is the set umbrella vertices of  $(\vec{G}, \sigma)$ .

The intuition behind this definition is that every support leaf of the root of the LRT of a 2-BMG must have all differently colored vertices as out-neighbors, i.e., they are umbrella vertices. We now define "support sets" of graphs as particular subsets of umbrella vertices. As we shall see later, support sets are closely related to support vertices in  $S_{\rho}$ .

**Definition 8** (Support Set of  $(\vec{G}, \sigma)$ ). Let  $(\vec{G}, \sigma)$  be a 2-colored graph. A support set  $S := S(\vec{G}, \sigma)$ of  $(\vec{G}, \sigma)$  is a maximal subset  $S \subseteq U(\vec{G}, \sigma)$  of umbrella vertices such that  $x \in S$  implies  $N^{-}(x) \subseteq S$ .

**Lemma 8.** Every 2-colored graph  $(\vec{G}, \sigma)$  has a unique support set  $S(\vec{G}, \sigma)$ .

*Proof.* Assume, for contradiction, that  $(\vec{G}, \sigma)$  has (at least) two distinct support sets  $S, S' \subseteq U(\vec{G}, \sigma)$ . Clearly neither of them can be a subset of the other, since supports sets are maximal. We have  $N^-(x) \subseteq S$  for all  $x \in S$  and and  $N^-(x') \subseteq S'$  for all  $x' \in S'$ , which implies that  $N^-(z) \subseteq S \cup S'$  for all  $z \in S \cup S'$ . Together with the fact that S, S', and thus  $S \cup S'$ , are all subsets of  $U(\vec{G}, \sigma)$ , this contradicts the maximality of both S and S'.

For the construction of the support set  $S := S(\vec{G}, \sigma)$ , we consider the following sequence of sets, defined recursively by

$$S^{(k)} := \{ x \in S^{(k-1)} \mid N^{-}(x) \subseteq S^{(k-1)} \} \text{ for } k \ge 1 \text{ and } S^{(0)} = U(\vec{G}, \sigma).$$

$$(2)$$

By construction  $S^{(k+1)} \subseteq S^{(k)}$ . Furthermore, there is a  $k < |V(\vec{G})|$  such that  $S^{(k+1)} = S^{(k)}$ . Next we show that in a 2-BMG, S is obtained in a single iteration.

**Lemma 9.** If  $(\vec{G}, \sigma)$  is a 2-BMG, then  $S = S^{(1)}$ .

Proof. Let  $(\vec{G} = (V, E), \sigma)$  be a 2-BMG and  $U = U(\vec{G}, \sigma)$ . Assume for contradiction that  $S \neq S^{(1)}$ , and thus  $S^{(2)} \subsetneq S^{(1)}$ . We will show that this implies the existence of a forbidden F2-graph. By assumption, there is a vertex  $x_2 \in S^{(1)} \setminus S^{(2)}$ . Thus, there must be a vertex  $y_1 \in N^-(x_2)$  (and thus  $(y_1, x_2) \in E$ ) with  $\sigma(y_1) \neq \sigma(x_2)$  such that  $y_1 \notin S^{(1)}$ . However, by definition,  $y_1 \in N^-(x_2)$  and  $x_2 \in S^{(1)}$  implies  $y_1 \in U$ . Now, it follows from  $y_1 \in U \setminus S^{(1)}$  that there is a vertex  $x_1 \in N^-(y_1)$  with  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y_1)$  such that  $x_1 \notin U$ . The latter together with  $x_2 \in S^{(1)} \subseteq U$  implies  $x_1 \neq x_2$ . In particular, since  $x_1 \notin U$ , the vertex  $x_1$  does not have an out-arc to every differently colored vertex, thus there must be a vertex  $y_2$  with  $\sigma(y_2) = \sigma(y_1)$  such that  $(x_1, y_2) \notin E$ . Since  $x_1 \in N^-(y_1)$ , we have  $(x_1, y_1) \in E$  and  $y_1 \neq y_2$ . Finally,  $x_2 \in U$  and  $\sigma(y_2) = \sigma(y_1) \neq \sigma(x_2)$  implies that  $(x_2, y_2) \in E$ . In summary, we have four distinct vertices  $x_1, x_2, y_1, y_2$  with  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y_1) = \sigma(y_2)$  and (non-)arcs  $(x_1, y_1), (y_1, x_2), (x_2, y_2) \in E$  and  $(x_1, y_2) \notin E$ , and hence an induced F2-graph in  $(\vec{G}, \sigma)$ . By Thm. 2, we can conclude that  $(\vec{G}, \sigma)$  is not a BMG; a contradiction.

In general,  $S = S^{(0)} = U(\vec{G}, \sigma)$  is not satisfied. To see this consider the BMG  $(\vec{G}, \sigma)$  that is explained by the triple  $x_1y|x_2$  with  $\sigma(x_1) = \sigma(x_2) \neq \sigma(y)$ . One easily verifies that  $U(\vec{G}, \sigma) = \{x_1, x_2\}$  but  $S = \{x_2\}$ .

**Theorem 4.** Let  $(T, \sigma)$  be the least resolved tree of a 2-BMG  $(\vec{G}, \sigma)$ . Then, the set of support leaves  $S_{\rho}$  of the root  $\rho$  equals the support set S of  $(\vec{G}, \sigma)$ . In particular  $S \neq \emptyset$  if and only if  $(\vec{G}, \sigma)$ is connected. *Proof.* Let  $(T, \sigma)$  be the LRT of a 2-BMG  $(\vec{G} = (V, E), \sigma)$ . We set  $U := U(\vec{G}, \sigma)$  and note first that  $S = S^{(1)}$  by Lemma 9.

First, suppose that  $(\vec{G}, \sigma)$  is not connected. Then it immediately follows from Prop. 2 that  $\sigma(L(T(v))) = \sigma(L(T))$  and thus  $|\sigma(L(T(v)))| > 1$  for any  $v \in \text{child}_T(\rho)$ . The latter together with Cor. 1 implies that any child of  $\rho$  must be an inner vertex in T. Hence,  $S_{\rho} = \emptyset$ . On the other hand, since  $(\vec{G}, \sigma)$  is not connected, each of its connected components is a 2-BMG (cf. Prop. 1), and thus, contains both colors. Therefore, for each vertex x in  $\vec{G}$ , we can find a vertex y with  $\sigma(x) \neq \sigma(y)$  such that  $(x, y), (y, x) \notin E$ , and thus  $x \notin S$ . Since this is true for any vertex in  $\vec{G}$ , we can conclude  $S = \emptyset = S_{\rho}$ .

Now, suppose that  $(\vec{G}, \sigma)$  is connected. By Cor. 2, we have  $S_{\rho} \neq \emptyset$ . We first show  $S_{\rho} \subseteq S$ . Let  $x \in S_{\rho}$ . By definition, x satisfies  $\operatorname{lca}_{T}(x, y) = \rho$  and therefore  $(x, y) \in E$  for all  $y \in L(T)$  with  $\sigma(y) \neq \sigma(x)$ , i.e., x has an out-arc to every differently colored vertex in  $\vec{G}$ . By definition, we thus have  $x \in U$ . Now assume for contradiction that  $x \notin S = S^{(1)} = \{z \in U \mid N^{-}(z) \subseteq U\}$ . The latter implies that there exists a vertex  $y \in N^{-}(x)$  such that  $y \notin U$ . In particular,  $(y, x) \in E$ . Since  $y \notin U$ , there is some vertex x' with  $\sigma(x') = \sigma(x)$  such that  $(y, x') \notin E$ . Together this implies that xy|x' is an informative triple. By Lemma 2, we obtain  $\operatorname{lca}_{T}(x, y) \prec_{T} \operatorname{lca}_{T}(x, x') = \operatorname{lca}_{T}(x', y) \preceq_{T} \rho$ ; a contradiction to the assumption that x is a support leaf of  $\rho$ . Thus  $x \in S$ .

Next, we show by contraposition that  $S \subseteq S_{\rho}$ . To this end, suppose that x is not a support leaf of  $\rho$ , i.e.  $x \notin S_{\rho}$ . Hence, there is an inner vertex  $v \in \operatorname{child}_{T}(\rho) \cap V^{0}(T)$  such that  $x \prec_{T} v$ . By Cor. 1, we conclude that  $|\sigma(L(T(v)))| = 2$ , i.e., the subtree T(v) contains both colors. We now distinguish two cases: (i) there is a leaf  $y' \in L(T) \setminus L(T(v))$  with  $\sigma(y') \neq \sigma(x)$ , and (ii) there is no leaf  $y' \in L(T) \setminus L(T(v))$  with  $\sigma(y') \neq \sigma(x)$ .

Case(i): Since T(v) contains both colors, there is a leaf  $y \in L(T(v))$ , with  $y \neq y'$  and  $\sigma(y) = \sigma(y') \neq \sigma(x)$ . Since, by construction, we have  $lca_T(x,y) \preceq_T v \prec_T \rho = lca_T(x,y')$ , it follows  $(x,y') \notin E$ . Together with  $\sigma(x) \neq \sigma(y')$ , this immediately implies  $x \notin U$ . From  $S^{(2)} \subseteq S^{(1)} \subseteq U$ , we conclude  $x \notin S^{(1)} = S$ .

 $Case(ii): \text{ Suppose that there is no leaf } y' \in L(T) \setminus L(T(v)) \text{ with } \sigma(y') \neq \sigma(x). \text{ We will continue by showing that there is a support leaf y of vertex v with } \sigma(y) \neq \sigma(x). \text{ Assume, for contradiction, that the latter is not the case. Since <math>(T, \sigma)$  is least resolved, the inner edge  $\rho v$  is not redundant. Hence, by Lemma 1, there must be an arc  $(a,b) \in E$  such that  $\operatorname{lca}_T(a,b) = v$  and  $\sigma(b) \in \sigma(L(T) \setminus L(T(v)))$ . Since there is no leaf  $y' \in L(T) \setminus L(T(v))$  with  $\sigma(y') \neq \sigma(x)$ , we conclude that  $\sigma(b) = \sigma(x)$  and  $\sigma(a) \neq \sigma(x)$ . Clearly, it holds  $a, b \in L(T(v))$ . Now consider an arbitrary  $a' \in L(T(v))$  with  $\sigma(a') \neq \sigma(x)$ . Since, by assumption, every such a' is not a support leaf of v, there must be an inner vertex  $w \in \operatorname{child}_{T(v)}(v)$  with  $a' \prec_T w$ . By Cor. 1 and since  $w \prec_T v \prec_T \rho$ , we conclude that  $|\sigma(L(T(w)))| = 2$ , i.e., the subtree T(w) contains both colors. Thus there is some b' with  $\sigma(b') = \sigma(x)$  and  $\operatorname{lca}_T(a', b') \preceq_T w \prec_T v$ . Since a' was chosen arbitrarily, we conclude that there is a support leaf y of vertex v with  $\sigma(y) \neq \sigma(x)$ . Hence,  $\operatorname{lca}_T(x, y) = v \preceq_T \operatorname{lca}_T(x'', y)$  for all  $x'' \in L(T)$  with  $\sigma(x'') = \sigma(x)$ , and thus  $(y, x) \in E$  and  $y \in N^-(x)$ . Since  $S_{\rho} \neq \emptyset$  and  $\sigma(y) \notin \sigma(L(T) \setminus L(T(v)))$ , there must be a leaf  $x' \in S_{\rho}$  with  $\sigma(x') = \sigma(x)$ . The fact that  $\operatorname{lca}_T(x, y) = v \prec_T \rho = \operatorname{lca}_T(x', y)$  implies  $(y, x') \notin E$ . Therefore and since  $\sigma(x') \neq \sigma(y)$ , it follows  $y \notin U$ . Together with  $y \in N^-(x)$ , we conclude that  $x \notin S^{(1)} = S$ .

In summary, we have shown  $S = S_{\rho}$  for any BMG  $(\vec{G}, \sigma)$ . Finally,  $S = S_{\rho}$  together with Cor. 2 implies that  $S \neq \emptyset$  if and only if  $(\vec{G}, \sigma)$  is connected, which completes the proof.

#### 5 Algorithmic Considerations

Thm. 4 provides not only a convenient necessary condition for connected 2-BMGs but also a fast way of determining the support set  $S = S_{\rho}$  and thus also a fast recursive approach to construct the LRT for a 2-BMG. It is formalized in Alg. 1 and illustrated in Fig. 2.

**Lemma 10.** Let  $(\vec{G}, \sigma)$  be a connected 2-BMG. Then Alg. 1 returns the least resolved tree for  $(\vec{G}, \sigma)$ .

Proof. Let  $(T, \sigma)$  be the (unique) least resolved tree of  $(\vec{G}, \sigma)$  with root  $\rho$ . The latter is supplied to Alg. 1 to initialize the tree. By Thm. 4, Lemma 9 and since  $(\vec{G}, \sigma)$  is connected, the set of support leaves  $S_{\rho} = S^{(2)} = S^{(1)} \neq \emptyset$  for the root  $\rho$  is correctly identified in the top-level recursion of Alg. 1 (Line 2-4) and attached to the root  $\rho$  (Line 8-9). According to Cor. 3, one can now proceed to Algorithm 1: LRT for connected 2-colored BMGs  $(\vec{G}, \sigma)$ .

**Input:** Connected properly 2-colored digraph  $(\vec{G} = (L, E), \sigma)$ , vertex  $\rho$ **Output:** LRT of  $(\vec{G}, \sigma)$  if  $(\vec{G}, \sigma)$  is a BMG 1 Function Build2ColLRT( $\vec{G}, \sigma, \rho$ )  $U \leftarrow \{x \in L \mid \text{outdeg}(x) = |L| - |L[\sigma(x)]|\}$ // umbrella vertices  $\mathbf{2}$  $S^{(1)} \leftarrow \{ x \in U \mid N^-(x) \subseteq U \}$ // all in-neighbors in U3  $S^{(2)} \leftarrow \{x \in S^{(1)} \mid N^-(x) \subseteq S^{(1)}\}$ // all in-neighbors in  $S^{(1)}$ 4 if  $S^{(1)} = \emptyset$  or  $S^{(2)} \neq S^{(1)}$  then  $\mathbf{5}$ exit false 6 7 else foreach  $x \in S^{(2)}$  do 8 add x as a child of  $\rho$ 9 foreach connected component  $\vec{G}_v$  of  $\vec{G} - S^{(2)}$  do 10 if  $|V(\vec{G}_v)| = 1$  then 11 exit false 12 create vertex v13  $T_v \leftarrow \texttt{Build2ColLRT}(\vec{G}_v, \sigma_{|.}, v)$  $\mathbf{14}$ connect the root v of  $T_v$  as a child to  $\rho$  $\mathbf{15}$ 



**Figure 2:** Illustration of Alg. 1 with input  $(\vec{G}, \sigma)$  (uppermost box). The boxes indicate the five recursion steps that are necessary to decompose  $(\vec{G}, \sigma)$ , and correspond to the five inner vertices of the LRT shown on the right. Note that, in the recursion step on  $(\vec{G}[\{a_3, a_4, b_5\}], \sigma_{|})$ , we have  $U \neq S^{(2)}$ .

recursively construct the LRTs for the connected components of  $(\vec{G} - S_{\rho}, \sigma_{|.})$ , which is done in Line 10-15. By Lemma 7, these connected components  $(\vec{G}_v, \sigma_{|.})$  are exactly the BMGs  $\vec{G}(T(v), \sigma_{|.})$ with  $v \in \text{child}_T(\rho) \setminus \{S_{\rho}\}$  (Line 14). In particular, therefore, we have  $V(\vec{G}_v) = L(T(v))$ . Since  $v \notin S_{\rho}$ , i.e., v is an inner vertex, Cor. 1 and  $v \prec_T \rho$  imply  $|\sigma(L(T(v)))| > 1$ . Hence, in particular, the condition  $|V(\vec{G}_v)| > 1$  (cf. Line 11) to proceed recursively is satisfied for each connected component.

**Theorem 5.** Given a connected properly 2-colored digraph  $(\vec{G}, \sigma)$  as input, Alg. 1 returns a tree

T if and only if  $(\vec{G}, \sigma)$  is a 2-colored BMG. In particular, T is the unique least resolved tree for  $(\vec{G}, \sigma)$ .

*Proof.* By Lemma 10, Alg. 1 returns the unique least resolved tree T if  $(\vec{G}, \sigma)$  is a connected 2colored BMG. To prove the converse, suppose that Alg. 1 returns a tree T given the connected properly 2-colored digraph  $(\vec{G}, \sigma)$  as input. We will show that  $(\vec{G}, \sigma) = \vec{G}(T, \sigma)$ , and thus  $(\vec{G}, \sigma)$  is a BMG.

It is easy to see that  $L(T) = V(\vec{G})$  must hold since, in each step of Alg. 1 every vertex is either attached to some inner vertex or passed down to a deeper-level recursion as part of some connected component. Therefore, every vertex of  $\vec{G}$  eventually appears in the output. Thus  $\sigma(L(T)) = \sigma(V(\vec{G}))$  and  $|\sigma(L(T))| = |\sigma(V(\vec{G}))| = 2$ . It remains to show  $E(\vec{G}) = E(\vec{G}(T, \sigma))$ .

Note first that neither  $(\vec{G}, \sigma)$  nor  $\vec{G}(T, \sigma)$  contain arcs between vertices of the same color. Moreover, since Alg. 1 eventually returns a tree, we have  $S^{(1)} = S^{(2)} \neq \emptyset$  in every recursion step. Throughout the remainder of the proof, we will write  $S_i^{(1)}$  and  $S_i^{(2)}$  for the sets  $S^{(1)}$  and  $S^{(2)}$  of the  $i^{th}$  recursion step. Likewise, in every step, each connected component  $(\vec{G}_v, \sigma_{|.})$  computed in Line 10 must contain at least two vertices (cf. Line 11), and thus  $|\sigma(V(\vec{G}_v))| = 2$  because  $(\vec{G}, \sigma)$  is properly 2-colored.

First, let S be the support set of  $\vec{G}(T,\sigma)$  and  $x \in S$  be arbitrary. Note that the support set is computed in the first iteration step of the algorithm as  $S = S_1^{(2)}$ , hence  $S = S_1^{(2)} \neq \emptyset$ . By construction of T, x is attached as a leaf to  $\rho$ , i.e.  $lca_T(x,y) = \rho$ . Consequently, (x,y) is an arc in  $\vec{G}(T,\sigma)$  for all  $y \in V(\vec{G})$  with  $\sigma(y) \neq \sigma(x)$ . By construction of S in Alg. 1, we have  $x \in S \subseteq U$ , i.e. x is an umbrella vertex in  $(\vec{G}, \sigma)$  and has out-arcs to every vertex  $y \in V(\vec{G})$  with  $\sigma(y) \neq \sigma(x)$ . Hence, all arcs of the form (x, y) with  $x \in S$  and  $\sigma(x) \neq \sigma(y)$  exist both in  $(\vec{G}, \sigma)$ and in  $\vec{G}(T,\sigma)$ . The latter property is in particular satisfied for all vertices in S and hence, all arcs between differently colored elements in S exist both in  $(\vec{G}, \sigma)$  and in  $\vec{G}(T, \sigma)$ . Now consider an arbitrary vertex  $y \in V(\vec{G}) \setminus S$ . Clearly, all in-neighbors in  $(\vec{G}, \sigma)$  of the elements in  $S = S_1^{(2)}$  must be contained in S, as a consequence of the condition  $S_1^{(1)} = S_1^{(2)}$  (cf. Line 5) and the construction of  $S_1^{(1)}$  and  $S_1^{(2)}$ . Hence,  $y \notin S$  and  $x \in S$  implies that (y, x) is not an arc in  $(\vec{G}, \sigma)$ . Moreover,  $y \notin S$ also implies that y is part of some connected component  $(\vec{G}_v, \sigma_{\perp})$  of  $(\vec{G} - S, \sigma_{\perp})$ . Therefore, and because Alg. 1 returns T, we must have  $y \in V(\vec{G}_v) = L(T(v))$  for some inner vertex  $v \in \text{child}_T(\rho)$ . As argued above,  $(\vec{G}_v, \sigma_{|.})$  and thus also the subtree T(v) contain both colors. Together with Obs. 3 and  $x \notin L(T(v))$ , this implies that  $\vec{G}(T,\sigma)$  does not contain the arc (y,x). By the same arguments, there is no arc (y, x') in  $\vec{G}(T, \sigma)$  such that the vertex x' is contained in a different connected component  $(\vec{G}_{v'}, \sigma_{|.}) \neq (\vec{G}_{v}, \sigma_{|.})$  of  $(\vec{G} - S, \sigma_{|.})$  than y. Since  $x \in S$  and  $y \notin S$  were chosen arbitrarily, we conclude that (i) any arc incident to some vertex in S exists in  $(\vec{G}, \sigma)$  if and only if it exists in  $\vec{G}(T,\sigma)$ , and (ii)  $\vec{G}(T,\sigma)$  contains no arcs between distinct connected components of  $(\hat{G} - S, \sigma_{\rm L})$ . Hence, it remains to consider the arcs within a connected component  $(\hat{G}_v, \sigma_{\rm L})$  of  $(\vec{G}-S,\sigma_{|.}).$ 

Alg. 1 recurses on each such connected component  $(\vec{G}_v, \sigma_{|.})$  using a newly created vertex  $v \in \text{child}_T(\rho)$  to initialize the tree T(v). By Lemma 3, it clearly holds that, for any  $x, y \in L(T(v)) = V(\vec{G}_v)$ , (x, y) is an arc in  $\vec{G}(T, \sigma)$  if and only it is an arc in  $\vec{G}(T(v), \sigma)$ . Thus, it suffices to consider only the subtree T(v). Now, we can apply the same arguments as in the previous recursion step to conclude that all arcs incident to the support set  $S_2^{(2)}$  constructed in the current recursion step are the same in  $(\vec{G}, \sigma)$  and  $\vec{G}(T, \sigma)$  and that neither  $(\vec{G}, \sigma)$  nor  $\vec{G}(T, \sigma)$  contain arcs between distinct connected components of  $(\vec{G}_v - S_2^{(2)}, \sigma_{|.})$ . Hence, it suffices to consider the connected components of  $(\vec{G}_v - S_2^{(2)}, \sigma_{|.})$ . Repeated application of this argumentation results in a chain of connected components that are contained in each other. Since Alg. 1 finally returns a tree, this chain is finite, say with a last element  $(\vec{G}_w - S_k^{(2)}, \sigma_{|.})$ , and thus  $S_k^{(2)} = V(\vec{G}_w)$ . In particular, therefore, every vertex in  $V(\vec{G})$  is contained in the support set of some recursion step.

In summary, we have shown that  $\vec{G}(T,\sigma) = (\vec{G},\sigma)$ . Hence,  $(\vec{G},\sigma)$  is a connected 2-BMG and, by Lemma 10, T is the unique least resolved tree of  $(\vec{G},\sigma)$ .

The construction in Lines 2-4 in Alg. 1 naturally produces two cases,  $U = S^{(1)} = S^{(2)}$  and  $S^{(2)} \subseteq S^{(1)} \subsetneq U$ . The following result shows that the latter case implies that the corresponding interior node in the LRT has only a single non-leaf descendant:

**Lemma 11.** Let  $(\vec{G}, \sigma)$  be a 2-BMG and  $S_{\rho}$  the support leaves of the root  $\rho$  of its LRT  $(T, \sigma)$ . If  $W := U(\vec{G}, \sigma) \setminus S_{\rho} \neq \emptyset$ , then the following statements are true:

- 1.  $S_{\rho} \neq \emptyset$ ,  $\vec{G}$  is connected, and  $\vec{G} S_{\rho}$  is connected.
- 2. All vertices in  $U(\vec{G}, \sigma) = S_{\rho} \cup W$  have the same color,
- 3. The set of support leaves  $S_v$  of the unique inner vertex child v of  $\rho$  contains vertices of both colors, and
- 4.  $W \subsetneq S_v$ .

Proof. First recall that, by Thm. 4 and the definition of the support set S of  $(\vec{G}, \sigma)$ , we have  $S_{\rho} = S \subseteq U(\vec{G}, \sigma)$ , and thus  $U(\vec{G}, \sigma) = S_{\rho} \cup W$ . Moreover, by Lemma 7, the connected components of  $(\vec{G}-S_{\rho}, \sigma_{|.})$  are exactly the BMGs  $\vec{G}(T(v), \sigma_{|.})$  with  $v \in \text{child}(\rho) \setminus S_{\rho}$ . The vertices  $v \in \text{child}(\rho) \setminus S_{\rho}$  are all inner vertices of T since, by definition, the support leaves  $S_{\rho}$  are exactly the children of  $\rho$  that are leaves. Together with the contraposition of Lemma 5 this implies that T(v) contains both colors.

Statement 1: Let  $x \in W$ , which exists due to the assumption  $W := U(\vec{G}, \sigma) \setminus S_{\rho} \neq \emptyset$ . Since  $x \notin S_{\rho}$ , it must be part of some connected component of  $(\vec{G} - S_{\rho}, \sigma_{|})$ , say  $\vec{G}(T(v), \sigma_{|})$  for some  $v \in \operatorname{child}_{T}(\rho) \setminus S_{\rho}$ . Now assume, for contradiction, that  $\vec{G} - S_{\rho}$  consists of more than one connected component. By Lemmas 7 and 5, there is a vertex  $v' \in \operatorname{child}_{T}(\rho) \setminus S_{\rho}$  such that  $v \neq v'$  and both subtrees T(v) and T(v') contain both colors. Hence, there are distinct  $y \in L(T(v))$  and  $y' \in L(T(v'))$  with  $\sigma(y) = \sigma(y') \neq \sigma(x)$ . Together with  $x \in L(T(v))$ , we therefore have  $\operatorname{lca}_{T}(x, y) \preceq_{T} v \prec_{T} \rho = \operatorname{lca}_{T}(x, y')$ , which implies  $(x, y') \notin E(\vec{G})$ . However,  $x \in W \subseteq U(\vec{G}, \sigma)$  and  $\sigma(y') \neq \sigma(x)$  imply  $(x, y') \in E(\vec{G})$ ; a contradiction. Hence, we conclude that  $\vec{G} - S_{\rho}$  has exactly one connected component, and thus  $\rho$  has a single inner vertex child v. Since T is phylogenetic, the latter implies that  $\rho$  must be incident to at least one leaf, i.e.  $S_{\rho} \neq \emptyset$ . Together with Thm. 4 this in turn implies that  $\vec{G}$  is connected. In summary, Statement 1 is true.

Statement 2: Let  $x \in W$  as in the proof of Statement 1. By arguments analogous to those used for Statement 1, we conclude that  $\sigma(x) = \sigma(y)$  for every  $y \in S_{\rho}$ , since otherwise we would obtain  $(x, y) \notin E(\vec{G})$ , and thus a contradiction to  $x \in U(\vec{G}, \sigma)$ . Since  $x \in W$  was chosen arbitrarily and  $S_{\rho}$  is non-empty, we immediately obtain that all vertices in  $U(\vec{G}, \sigma) = S_{\rho} \cup W$  have the same color, i.e., Statement 2 is true.

Statement 3: Now consider the single inner vertex child v of  $\rho$ , and its set of support leaves  $S_v$ , which must be non-empty by Lemma 6. Note that W must be entirely contained in L(T(v))and recall that all vertices in  $S_{\rho} \cup W$  are of the same color (cf. Statement 2). First suppose, for contradiction, that  $S_v$  only contains vertices of the *opposite* color as the vertices in  $S_{\rho} \cup W$ . This immediately implies  $S_v \cap W = \emptyset$ , thus every vertex  $x \in W$  must be located in a subtree T(w) of some inner vertex child w of v. Again by contraposition of Lemma 5, every such T(w) contains both colors. However, this contradicts  $(x,y) \in E(\vec{G})$  for every  $y \in S_v$ , which must hold as a consequence of  $x \in W \subset U(\vec{G}, \sigma)$  and  $\sigma(y) \neq \sigma(x)$ . Next suppose, for contradiction, that  $S_v$  only contains vertices of the same color as the vertices in  $S_{\rho} \cup W$ . In this case, we obtain that the edge  $\rho v$  is redundant w.r.t.  $(\vec{G}, \sigma)$ . To see this, consider an arc  $(x, y) \in E(\vec{G})$  such that  $lca_T(x, y) = v$ . Clearly, x must be directly incident to v, since otherwise the subtree below v to which x belongs would contain both colors, and thus contradict  $(x, y) \in E(\vec{G})$ . In other words, every such vertex x is a support leaf of v, thus  $\sigma(x) = \sigma(S_v) = \sigma(S_\rho)$  and  $\sigma(y) \neq \sigma(S_\rho)$ . In particular, there exists no arc  $(x,y) \in E(\vec{G})$  such that  $\operatorname{lca}_T(x,y) = v$  and  $\sigma(y) \in \sigma(L(T) \setminus L(T(v))) = \sigma(S_{\rho})$  and therefore, by Lemma 1, the inner edge  $\rho v$  is redundant. However, this contradicts the fact that T is least resolved. In summary, only the case in which  $S_v \neq \emptyset$  contains vertices of both colors is possible, and thus Statement 3 is true.

Statement 4: First, recall from the proof of Statement 3 that  $W \subseteq L(T(v))$  for the single inner vertex child v of  $\rho$ . In order to see that  $W \subseteq S_v$ , assume for contradiction that this is not the case. By similar arguments as used for showing Statement 3, this implies that some  $x \in W$  lies in a 2-colored subtree T(w) for some  $w \in \text{child}_T(v) \setminus S_v$ . Together with the above established fact that  $S_v$  contains both colors, this contradicts  $x \in U(\vec{G}, \sigma)$ . Finally,  $W \neq S_v$  is a consequence of the fact that  $S_v$  contains both colors (Statement 3) but  $W \subseteq S_\rho \cup W$  contains only one color (Statement 2).

We now use this result to consider the performance of Alg. 1.



**Figure 3:** Illustration of Lemma 11. (A) The (local) situation if  $W = U \setminus S_{\rho} \neq \emptyset$  as implied by Lemma 11. In particular,  $\rho$  only as a single inner vertex child v, all vertices in  $U = S_{\rho} \cup W$  have the same color,  $S_v$  contains vertices of both colors, and  $W \subsetneq S_v$ . (B) There cannot be a second inner vertex child v', since then none of the vertices except those in  $S_{\rho}$  can be umbrella vertices, e.g. (a, b) is not an arc in the graph explained by the tree in (B). Hence, this situation is not possible for  $W \neq \emptyset$ . (C) If  $S_v$  does not contain vertices of both colors, then the edge  $\rho v$  is redundant in the tree, contradicting that  $(T, \sigma)$  in Lemma 11 is the LRT.

**Lemma 12.** Alg. 1 can be implemented to run in  $O(|E|\log^2 |V|)$  time for a connected input graph.

Proof. Since  $\vec{G}$  is connected by assumption, we have  $|V| \in O(|E|)$ . Starting from  $(\vec{G}, \sigma)$ , the list of out-degrees can be constructed in O(|E|). The initial umbrella set U is then obtained by listing the vertices with maximal out-degree in the color class. The initial set  $S^{(1)}$  is constructed by checking, for each  $u \in U$ , the in-neighbors of u for membership in U in O(|V| + |E|) operations. Then  $S^{(2)}$  is obtained in the same manner from  $S^{(1)}$ , requiring O(|V| + |E|) operations. The initial umbrella set U and the sets  $S^{(1)}$  and  $S^{(2)}$  thus can be constructed in linear time. In each recursive call of Build2ColLRT, at least one leaf is split off, hence the recursion depth is |V| - 1 in the worst case. Since the support vertices removed in each step have all of their in-neighbors in U, their removal does not affect the out-neighborhood for any  $x \in V(\vec{G} - U) \subseteq V(\vec{G} - S^{(2)})$ , and hence, outdeg(x) does not require updates. The in-neighborhoods  $N^-(x)$  can be updated by removing the arcs between  $\vec{G} - S^{(2)}$  and  $S^{(2)}$  as a consequence of Lemma 7 and Thm. 4. Since every arc appears exactly once in the removal, the total effort for these updates is O(|E|).

We continue by showing that every vertex needs to be considered as an umbrella vertex at most twice, and that the total effort of constructing all sets  $S^{(1)}$  and  $S^{(2)}$  is O(|E|), given that the umbrella vertices U can be obtained efficiently, which we discuss afterwards. To this end, we distinguish, for each of the single recursion steps, two cases:  $S^{(1)} = U$  and  $S^{(1)} \subseteq U$ . First if  $S^{(1)} = U$ , and thus also  $S^{(2)} = S^{(1)} = U$ , we consider each in-arc of  $x \in U$ . Since these vertices and their corresponding arcs are removed when constructing  $\vec{G} - S^{(2)}$ , they are not considered again in a deeper recursion step. In the second case, we have  $S^{(1)} \subseteq U$ , which together with  $S^{(2)} = S^{(1)}$ implies  $W \coloneqq U \setminus S^{(2)} \neq \emptyset$ , and only the vertices in  $U \setminus W$  are removed. However, Lemma 11 guarantees that, for a 2-BMG as input graph, the vertices in W appear as support leaves in the next step and thus appear in the update of  $U, S^{(1)}$ , and  $S^{(2)}$  no more than a second time. In order to use the properties in Lemma 11 for the general case (i.e.  $(\vec{G}, \sigma)$  is not necessarily a BMG), we can, whenever  $W \neq \emptyset$ , (i) check that  $\vec{G} - S^{(2)}$  only has a single connected component  $\vec{G}_v$ , and (ii) pass down the set W to the recursion step on  $\vec{G}_v$  in which the condition  $W \subseteq S^{(2)}$  is checked. If any of these checks fails, we can exit false. This way, we ensure that every vertex appears at most two times as an umbrella vertex in the general case. To construct  $S^{(1)}$  from U, we have to scan the in-neighborhood  $N^{-}(x)$  of each vertex  $x \in U$  and check whether  $N^{-}(x) \subset U$ . We repeat this step to construct  $S^{(2)}$  from  $S^{(1)}$ . Membership in U and  $S^{(1)}$ , resp., can be checked in constant time (e.g. by marking the vertices in the current set U). Since we have to consider each vertex, and hence, each in-neighborhood at most twice, all sets  $S^{(1)}$  and  $S^{(2)}$  can be obtained with a total effort of O(|E|).

It remains to show that the input graph can be decomposed efficiently in such a way that the connectivity information is maintained and the candidates for umbrella vertices in each component are updated. The connected components  $\vec{G}_v$  can be obtained by using the dynamic data structure described in [6], often called HDT data structure. It maintains a maximal spanning forest representing the underlying undirected graph with edge set  $\tilde{E} = \{xy \mid (x, y) \in E \text{ or } (y, x) \in E\}$ , and allows deletion of all  $|\tilde{E}| \in O(|E|)$  edges with amortized cost  $O(\log^2 |V|)$  per edge deletion.

The explicit traversal of the connected components to compute U can be avoided as follows: Since outdeg(x) does not require updates, we can maintain a doubly-linked list of vertices x for each color  $i \in \{1, 2\}$ , and each value of outdeg(x) where  $\sigma(x) = i$ . In order to be able to access the highest value of the out-degrees, we maintain these values together with pointer to the respective doubly-linked list in balanced binary search trees (BST), one for each color and each connected component. The BSTs for the two colors are computed first for  $(\vec{G}, \sigma)$  in  $O(|V|\log(|V|))$  time and afterwards updated to fit with the out-degree of the currently considered component  $\vec{G}_v$ . To update these lists and BSTs for  $\vec{G}_v$ , observe first that  $\vec{G}_v$  can be obtained from G by stepwise deletion of single arcs, i.e. edges in the HDT data structure representing the underlying undirected versions. We update, resp., construct the pair of BSTs (one for each color) for each connected component as follows: Since a single arc deletion splits a connected component  $\vec{G}'$  into at most two connected components  $\vec{G}_1$ , and  $\vec{G}_2$ , we can apply the well-known technique of traversing the smaller component [14]. The size of each connected component can be queried in O(1) time in the HDT data structure. Suppose w.l.o.g. that  $|V(\vec{G}_1)| \leq |V(\vec{G}_2)|$ . We construct a new pair of BSTs for  $\vec{G}_1$ , and delete the vertices  $V(\vec{G}_1)$  and the respective degrees from the two original BSTs for  $\vec{G}$ , which then become the BSTs for  $\vec{G}_2$ . More precisely, we delete each vertex  $x \in V(\vec{G}_1)$  in the respective list corresponding to outdeg(x), and if the length of this list drops to zero, we also remove the corresponding out-degree in the BST. Likewise, we insert the out-degree of x and an empty doubly-linked list into the newly-created BST for  $\vec{G}_1$ , if it is not yet present, and append x to this list. Note that the number of out-degree deletions and insertions does not exceed  $|V(\vec{G}_1)|$ . Due to the technique of traversing the smaller component, every vertex is deleted and inserted at most  $\log |V||$  times. Therefore, we obtain an overall complexity of  $O(|V| \log^2 |V|)$  for the maintenance of the BSTs where the additional log-factor originates from rebalancing the BSTs whenever necessary.

In each recursion step, the set U can now be obtained by listing (at most) the vertices with the maximal out-degree for each of the two colors. Finding the two out-degrees and corresponding lists in the BSTs requires  $O(\log |V|)$  in each step, and thus  $O(|V|\log |V|)$  in total. In order to determine whether these candidates x are actually umbrella vertices, we have to check whether outdeg $(x) = |V(G_v)| - |V(G_v)[\sigma(x)]|$ . The HDT data structure allows constant-time query of the size of a given connected component, since this information gets updated during the maintenance of the spanning forest. By the same means, we can keep track of the number of vertices of a specific color in each connected components. Note that we only need to do this for one color rsince  $|V(G_v)[s]| = |V(G_v)| - |V(G_v)[r]|$ . This does not increase the overall effort for maintaining the data structure since it happens alongside the update of  $|V(G_v)|$ .

In summary, the total effort is dominated by maintaining the connectedness information while deleting O(|E|) arcs, i.e.,  $O(|E|\log^2 |V|)$  time.

As a direct consequence of Thm. 4 the LRT of a disconnected graph  $\vec{G}$  is obtained by connecting the roots of the LRTs  $T_v$  of the connected components  $G_v$  to an additional root vertex, see also [5, Cor. 4]. Lemma 12 thus implies

**Theorem 6.** The LRT of a 2-BMG can be computed in  $O(|V| + |E| \log^2 |V|)$ .

*Proof.* The connected components  $G_i = (V_i, E_i)$  of G = (V, E) can be enumerated in O(|V| + |E|) operations, e.g. using a breadth-first search on the underlying undirected graph. By Lemma 12,  $O(|E_i|\log^2 |V_i|) \leq O(|E_i|\log^2 |V|)$  operations are required for each  $G_i$ . Hence, the total effort is  $O(|V| + |E| + \log^2 |V| \sum_i |E_i|) = O(|V| + |E| \log^2 |V|)$ .

In order to illustrate the improved complexity for the construction of LRTs of 2-BMGs, we implemented both the well-known triple-based approach, i.e., the application of BUILD [2] with the informative triples defined in Eq. (1) as input, and the new approach of Alg. 1. As input, we used 2-BMGs that where randomly generated as follows: First, we simulate random trees T recursively, starting from a single vertex, by attaching to a randomly chosen vertex v either a single leaf if vis an inner vertex of T or a pair of leaves if v was a leaf. The construction stops when the desired number of leaves is reached. Note that the resulting tree is phylogenetic by construction. Each leaf is then colored by selecting at random one of the two colors. Finally, we compute the 2-BMG  $\vec{G}(T, \sigma)$  from each of the simulated leaf-colored trees  $(T, \sigma)$ .

Both methods for the LRT computation were implemented in Python. Moreover, we note that we did not implement the sophisticated dynamic data structures used in the proof of Lemma 12, but a rather naïve implementation of Alg. 1. Nevertheless, Fig. 4 shows a remarkable improvement



Figure 4: Running time comparison of the general approach for constructing an LRT using BUILD (blue) vs. Alg. 1 (green). For each number of leaves, 200 2-BMGs where generated as described in the text. In the left panel, the median values are shown with logarithmic axes. The additional dotted line indicates the median values of the size of the simulated BMGs, i.e. the number of arcs, scaled by a factor  $10^{-3}$ . We did not compute the LRTs with the first method for instances with more than 1000 leaves because of the excessive computational cost.



Figure 5: The tree on the r.h.s. explains the hourglass graph on the l.h.s.

of the running time when compared to the general  $O(|V| |E| \log^2 |V|)$  approach for  $\ell$ -BMGs detailed in [5]. Empirically, we observe that the running time of Alg. 1 indeed scales nearly linearly with the number of edges.

### 6 Binary-explainable 2-BMGs

Binary phylogenetic trees are of particular interest in practical applications. Not every 2-BMG can be explained by a binary tree. The subclass of *binary-explainable*  $(\ell)BMG$  are characterized among all BMGs by the absence of single forbidden subgraph called *hourglass* [10, 11], illustrated in Fig. 5. In this section we briefly describe a modification of Alg. 1 that allows the efficient recognition of binary-explainable 2-BMGs.

**Definition 9.** An hourglass in a properly vertex-colored graph  $(\vec{G}, \sigma)$ , denoted by  $[xy \Join x'y']$ , is a subgraph  $(\vec{G}[Q], \sigma_{|Q})$  induced by a set of four pairwise distinct vertices  $Q = \{x, x', y, y'\} \subseteq V(\vec{G})$ such that (i)  $\sigma(x) = \sigma(x') \neq \sigma(y) = \sigma(y')$ , (ii) (x, y), (y, x) and (x'y'), (y', x') are bidirectional arcs in  $\vec{G}$ , (iii)  $(x, y'), (y, x') \in E(\vec{G})$ , and (iv)  $(y', x), (x', y) \notin E(\vec{G})$ .

A graph  $(\vec{G}, \sigma)$  is called *hourglass-free* if it does not contain an hourglass as an induced subgraph. We summarize Lemma 31 and Prop. 8 in [11] as

**Proposition 3.** For every BMG  $(\vec{G}, \sigma)$ , the following three statements are equivalent:

- 1.  $(\vec{G}, \sigma)$  is binary-explainable.
- 2.  $(\vec{G}, \sigma)$  is hourglass-free.
- 3. If  $(T, \sigma)$  is a tree explaining  $(\vec{G}, \sigma)$ , then there is no vertex  $u \in V^0(T)$  with three distinct children  $v_1, v_2$ , and  $v_3$  and two distinct colors r and s satisfying
  - (a)  $r \in \sigma(L(T(v_1))), r, s \in \sigma(L(T(v_2))), and s \in \sigma(L(T(v_3))), and$ (b)  $s \notin \sigma(L(T(v_1))), and r \notin \sigma(L(T(v_3))).$

The following Lemma shows that the third condition in Prop. 3 can be translated to a much simpler statement in terms of the support leaves of its LRT.

**Lemma 13.** A 2-BMG  $(\vec{G}, \sigma)$  contains an induced hourglass if and only if its LRT  $(T, \sigma)$  contains an inner vertex u such that  $S_u$  contains support vertices of both colors and  $V(\vec{G}(T(u)) - S_u) \neq \emptyset$ .

Proof. By Thm. 5, Alg. 1 returns the LRT  $(T, \sigma)$  for  $(\vec{G}, \sigma)$  if and only if  $(\vec{G}, \sigma)$  is a 2-BMG. Hence, we assume in the following that the latter is satisfied. As a consequence of Prop. 3 and the fact that  $(T, \sigma)$  explains  $(\vec{G}, \sigma)$ , we know that  $(\vec{G}, \sigma)$  is binary-explainable if and only if there is no vertex  $u \in V^0(T)$  with three distinct children  $v_1, v_2$ , and  $v_3$  and two distinct colors r and ssatisfying (a)  $r \in \sigma(L(T(v_1))), r, s \in \sigma(L(T(v_2)))$ , and  $s \in \sigma(L(T(v_3)))$ , and (b)  $s \notin \sigma(L(T(v_1)))$ , and  $r \notin \sigma(L(T(v_3)))$ .

First, suppose that  $(\vec{G}, \sigma)$  contains an hourglass, i.e., by Prop. 3 there is a vertex  $u \in V^0(T)$ with distinct children  $v_1, v_2$ , and  $v_3$  and two distinct colors r and s satisfying (a) and (b). Since  $(\vec{G}, \sigma)$  is 2-colored and  $(T, \sigma)$  its LRT, Lemma 5 together with  $s \notin \sigma(L(T(v_1)))$  and  $r \notin \sigma(L(T(v_3)))$ implies that  $v_1$  of color r and  $v_2$  of color s, respectively, are both leaves. In particular, therefore, we know that  $v_1, v_2 \in S_u$  are support leaves. By Lemma 7 and since  $\vec{G}(T(u), \sigma_{|})$  is also a BMG, the connected components of  $(\vec{G}(T(u)) - S_u, \sigma_{|}) = (\vec{G}[L(T(u))] - S_u, \sigma_{|})$  (cf. Lemma 3) are exactly the BMGs  $\vec{G}(T(v), \sigma_{|})$  with  $v \in \text{child}(u) \setminus S_u$ . Together with the fact that  $v_2 \in V^0(T)$  as a consequence of  $L(T(v_2))$  containing both colors r and s, this implies that  $(\vec{G}(T(u)) - S_u, \sigma_{|})$  is not the empty graph.

Conversely, suppose there a vertex  $u \in V^0(T)$  such that  $S_u$  contains support vertices  $v_1$  and  $v_3$ with distinct colors  $\sigma(v_1) \neq \sigma(v_3)$  and  $V(\vec{G}(T(u)) - S_u) \neq \emptyset$ , i.e., u has a child  $v_2 \in \text{child}(u) \setminus S_u$ that is not a support leaf and hence satisfies  $v_2 \in V^0(T)$ . Lemma 5 implies that  $L(T(v_2))$  contains both colors since  $v_2 \in V^0(T)$ . Hence, the three children  $v_1$ ,  $v_2$ , and  $v_3$  of u satisfy conditions (a) and (b) of Prop. 3(3), and thus  $(\vec{G}, \sigma)$  contains an induced hourglass.

**Corollary 4.** It can be checked in  $O(|V| + |E| \log^2 |V|)$  whether or not a properly 2-colored graph  $(\vec{G}, \sigma)$  is a binary-explainable BMG.

Proof. Recall that there is a one-to-one correspondence between the recursion step in Alg. 1 and the inner vertices  $u \in V^0(T)$ . As argued in the proof of Lemma 12, every vertex appears at most twice in an umbrella set U. Therefore, it can be checked in O(|V|) total time whether  $S = S^{(2)}$  contains vertices of both colors. Since the vertex set of  $\vec{G}_u - S_u$  is maintained in the dynamic graph HDT data structure, it can be checked in constant time for each u whether  $\vec{G}_u - S_u$  is non-empty. The additional effort to check the condition of Lemma 13 is therefore only O(|V|). Hence, we still require a total effort of  $O(|V| + |E| \log^2 |V|)$  (cf. Thm. 6).

Cor. 4 improves the complexity for the decision whether a 2-BMG is binary-explainable as compared to the  $O(|V|^3 \log^2 |V|)$ -time algorithm for (general) BMGs presented in [10].

## 7 Concluding Remarks

We have shown here that 2-BMGs have a recursive structure that is reflected in certain induced subgraphs that correspond to subtrees of the LRT. The leaves connected directly to the root of a given subtree play a special role as support vertices in the corresponding subgraph of the 2-BMG. Since the support vertices of the root can be identified efficiently in a given input graph, there is a recursive decomposition of  $(\vec{G}, \sigma)$  that directly yields the LRT. With the help of a dynamic data structure to maintain connectedness information [6], this provides an  $O(|V|+|E|\log^2 |V|)$  algorithm to recognize both 2-BMGs and binary explainable 2-BMGs and to construct the corresponding LRT. This provides a considerable speed-up compared to the previously known  $O(|V||E|\log^2 |V|)$  and  $O(|V|^3)$  algorithms. Empirically, we observe a substantial speed-up even if simpler data structures are used to implement Alg. 1.

Both the theoretical insights and Alg. 1 itself have potential applications to the analysis of gene families in computational biology. Real-life data necessarily contain noise, and thus likely will deviate from perfect BMGs, naturally leading to graph editing problems for BMGs. Like many combinatorial problems in phylogenetics, these are NP-complete [12] and hence require approximation algorithms and heuristics. The support leaves introduced here provide an avenue to a new class of heuristics, conceptually distinct from approaches that attempt to extract consistent subsets of triples from  $\Re(\vec{G}, \sigma)$ .

## Acknowledgments

This work was supported in part by the Austrian Federal Ministries BMK and BMDW and the Province of Upper Austria in the frame of the COMET Programme managed by FFG, and the *Deutsche Forschungsgemeinschaft*.

### References

- G. Abrams and J. K. Sklar. The graph menagerie: Abstract algebra and the mad veterinarian. Math. Mag., 83:168–179, 2010. doi:10.4169/002557010X494814.
- [2] A. Aho, Y. Sagiv, T. Szymanski, and J. Ullman. Inferring a tree from lowest common ancestors with an application to the optimization of relational expressions. SIAM J Comput, 10:405–421, 1981. doi:10.1137/0210030.
- [3] H. Cohn, R. Pemantle, and J. G. Propp. Generating a random sink-free orientation in quadratic time. *Electr. J. Comb.*, 9:R10, 2002. doi:10.37236/1627.
- [4] S. Das, P. Ghosh, S. Ghosh, and S. Sen. Oriented bipartite graphs and the Goldbach graph. Technical Report math.CO/1611.10259v6, arXiv, 2020.
- [5] M. Geiß, E. Chávez, M. González Laffitte, A. López Sánchez, B. M. R. Stadler, D. I. Valdivia, M. Hellmuth, M. Hernández Rosales, and P. F. Stadler. Best match graphs. J. Math. Biol., 78:2015–2057, 2019. doi:10.1007/s00285-019-01332-9.
- [6] J. Holm, K. de Lichtenberg, and M. Thorup. Poly-logarithmic deterministic fully-dynamic algorithms for connectivity, minimum spanning tree, 2-edge, and biconnectivity. J. ACM, 48:723-760, 2001. doi:10.1145/502090.502095.
- [7] A. Korchmaros. Circles and paths in 2-colored best match graphs. Technical Report math.CO/2006.04100v1, arXiv, 2020.
- [8] A. Korchmaros. The structure of 2-colored best match graphs. Technical Report math.CO/2009.00447v2, arXiv, 2020.
- [9] D. Schaller, M. Geiß, E. Chávez, M. González Laffitte, A. López Sánchez, B. M. R. Stadler, D. I. Valdivia, M. Hellmuth, M. Hernández Rosales, and P. F. Stadler. Best match graphs (corrigendum). arxiv.org/1803.10989v4, 2020.
- [10] D. Schaller, M. Geiß, M. Hellmuth, and P. F. Stadler. Best match graphs with binary trees. 2020. submitted; arXiv 2011.00511.
- [11] D. Schaller, M. Geiß, P. F. Stadler, and M. Hellmuth. Complete characterization of incorrect orthology assignments in best match graphs. J. Math. Biol., 2021. accepted; arXiv: 2006.02249.
- [12] D. Schaller, P. F. Stadler, and M. Hellmuth. Complexity of modification problems for best match graphs. *Theor. Comp. Sci.*, 2021. in revision; arxiv: 2006.16909.
- [13] C. Semple and M. Steel. *Phylogenetics*. Oxford University Press, Oxford UK, 2003.
- [14] Y. Shiloach and S. Even. An on-line edge-deletion problem. J. ACM, 28:1–4, 1981. doi: 10.1145/322234.322235.