

Energetic Stability of the Solutions of the Einstein Field Equations for Spherically Symmetric Liquid Shells

Jorge L. deLyra*

Universidade de São Paulo
Instituto de Física
Rua do Matão, 1371,
05508-090 São Paulo, SP, Brazil

March 18, 2021

Abstract

We interpret the exact solutions previously obtained for spherically symmetric shells of liquid fluid in General Relativity in terms of the energies involved. In order to do this we make a change of variables in the field equations in order to introduce some integral expressions that are related to various parts of the energy. We then use these integrals in order to show that a certain parameter with dimensions of length, that was necessarily introduced into the solutions by the interface boundary conditions, is related to the binding energies of the gravitational systems.

In sequence, we use this representation of the gravitational binding energy in order to discuss the energetic stability of the new solutions found. We include in the stability discussion the well-known interior Schwarzschild solution for a liquid sphere, which can be obtained as a specific limit of the solutions that were previously obtained for the liquid shells. We show that this particular family of solutions turns out to have zero binding energy and therefore to be a maximally unstable one, from the energetic point of view discussed here.

We also perform a numerical exploration of the energetic stability criterion of the liquid shell solutions, all of which have strictly positive binding energies, and show that indeed there is a particular subset of the solutions which are energetically stable. All these solutions have the form of shells with non-vanishing internal radii. This reduces the original three-parameter family of liquid shell solutions to a two-parameter family of energetically stable solutions.

*Email: delyra@lmail.if.usp.br

1 Introduction

The issue of the energy in General Relativity is a difficult one, and its discussion in specific examples quite often becomes involved and obscure. The difficulties start at the very foundations of the theory, with the impossibility of defining an energy-momentum tensor density for the gravitational field itself, a problem which apparently is related to the impossibility of localizing the energy of the gravitational field in the general case [1].

However, a recently discovered new class of static and time-independent exact solutions [2] provides us with an opportunity to discuss the subject in a clear, precise and complete manner. It leads to a simple and clear characterization of all the energies involved in this class of solutions, as well as a characterization of the relations among them, which establishes an important connection with the fundamental concept of the conservation of energy.

It is noteworthy that results similar to the ones we presented in [2] were obtained for the case of neutron stars, with the Chandrasekhar equation of state [3], by Ni [4] and Neslušan [5]. Just as in [2], the analysis of that case also led to an inner vacuum region containing a singularity at the origin and a gravitational field which is repulsive with respect to that origin. This tends to indicate that these results are general at least to some extent. It is to be expected that the ideas regarding the energy that we present here will be useful in that case as well.

This paper is organized as follows: in the remainder of this introduction we quickly review the new class of static and time-independent exact solutions for liquid shells, as well as the interior Schwarzschild solution, which can be obtained from the new shell solutions in a certain limit; in Section 2 we establish certain general integral formulas for all the energies involved; in Section 3 we establish the general physical interpretation of the energies involved, including for both the shell solutions and the interior Schwarzschild solution; in Section 4 we perform a small numerical exploration of the energetic stability of the shell solutions, and in Section 5 we state our conclusions.

1.1 The Liquid Shell Solutions

In a previous paper [2] we established the solution of the Einstein field equations for the case of a spherically symmetric shell of liquid fluid located between the radial positions r_1 and r_2 of the Schwarzschild system of coordinates. This is a three-parameter family of solutions, which can be taken as any three of the four parameters r_1 , r_2 , M and ρ_0 . The matter distribution is characterized by the radii r_1 and r_2 , by its total asymptotic gravitational mass M , associated to the Schwarzschild radius r_M , and by a matter energy density ρ_0 which is constant with the radial Schwarzschild coordinate r within (r_1, r_2) , and zero outside that interval. In this work we will use the time-like signature $(+, -, -, -)$, following [1]. In terms of the coefficients of the metric, for an invariant interval given in terms of the Schwarzschild coordinates (t, r, θ, ϕ) by

$$ds^2 = e^{2\nu(r)} c^2 dt^2 - e^{2\lambda(r)} dr^2 - r^2 [d\theta^2 + \sin^2(\theta) d\phi^2], \quad (1)$$

where $\exp[\nu(r)]$ and $\exp[\lambda(r)]$ are two positive functions of only r , as was explained in [2] the Einstein field equations reduce to the set of three first-order differential equations

$$\left\{ 1 - 2 [r\lambda'(r)] \right\} e^{-2\lambda(r)} = 1 - \kappa r^2 \rho(r), \quad (2)$$

$$\left\{ 1 + 2 [r\nu'(r)] \right\} e^{-2\lambda(r)} = 1 + \kappa r^2 P(r), \quad (3)$$

$$[\rho(r) + P(r)] \nu'(r) = -P'(r), \quad (4)$$

where $\rho(r)$ is the energy density of the matter, $P(r)$ is the isotropic pressure, $\kappa = 8\pi G/c^4$, G is the universal gravitational constant and c is the speed of light. In these equations the primes indicate differentiation with respect to r . Given these equations, as presented in [2] the complete solution for $\lambda(r)$ is given by

$$\lambda(r) = \begin{cases} -\frac{1}{2} \ln\left(\frac{r+r_\mu}{r}\right) & \text{for } 0 \leq r \leq r_1, \\ -\frac{1}{2} \ln\left[\frac{\kappa\rho_0(r_2^3 - r^3) + 3(r - r_M)}{3r}\right] & \text{for } r_1 \leq r \leq r_2, \\ -\frac{1}{2} \ln\left(\frac{r - r_M}{r}\right) & \text{for } r_2 \leq r < \infty, \end{cases} \quad (5)$$

where $r_M = 2GM/c^2$, while for $\nu(r)$ we have

$$\nu(r) = \begin{cases} \frac{1}{2} \ln\left(\frac{1 - r_M/r_2}{1 + r_\mu/r_1}\right) + \frac{1}{2} \ln\left(\frac{r+r_\mu}{r}\right) & \text{for } 0 \leq r \leq r_1, \\ \frac{1}{2} \ln\left(\frac{r_2 - r_M}{r_2}\right) + \ln[z(r)] & \text{for } r_1 \leq r \leq r_2, \\ \frac{1}{2} \ln\left(\frac{r - r_M}{r}\right) & \text{for } r_2 \leq r < \infty, \end{cases} \quad (6)$$

and finally the pressure within the shell, that is, for $r_1 \leq r \leq r_2$, is given by

$$P(r) = \rho_0 \frac{1 - z(r)}{z(r)}. \quad (7)$$

This solution is valid under the condition that $r_2 > r_M$. In all these expressions we have that r_μ is given in terms of the parameters characterizing the system by

$$r_\mu = \frac{\kappa\rho_0}{3} (r_2^3 - r_1^3) - r_M, \quad (8)$$

we have that ρ_0 is determined algebraically in terms of r_1 , r_2 and r_M as the solution of the transcendental algebraic equation

$$\begin{aligned} \sqrt{\frac{r_2}{3(r_2 - r_M)}} &= \sqrt{\frac{r_1}{\kappa\rho_0(r_2^3 - r_1^3) + 3(r_1 - r_M)}} + \\ &+ \frac{3}{2} \int_{r_1}^{r_2} dr \frac{\kappa\rho_0 r^{5/2}}{[\kappa\rho_0(r_2^3 - r^3) + 3(r - r_M)]^{3/2}}, \end{aligned} \quad (9)$$

and we have that the real function $z(r)$ is determined in terms of a non-trivial elliptic real integral by the relation

$$\begin{aligned} z(r) &= \sqrt{\frac{\kappa\rho_0(r_2^3 - r^3) + 3(r - r_M)}{r}} \times \\ &\times \left\{ \sqrt{\frac{r_2}{3(r_2 - r_M)}} + \frac{3}{2} \int_{r_2}^r ds \frac{\kappa\rho_0 s^{5/2}}{[\kappa\rho_0(r_2^3 - s^3) + 3(s - r_M)]^{3/2}} \right\}. \end{aligned} \quad (10)$$

The relation shown in Equation (8) is a direct consequence of the field equations and of the interface boundary conditions associated to them. In [2] we proved that, so long as

the pressure of the liquid is positive, we must have $r_\mu > 0$. In fact, the hypotheses of that proof can be weakened to require only that the pressure be strictly positive at a single point. This strictly positive value of r_μ implies that the solution has a singularity at the origin. However, that singularity is not associated to an infinite concentration of matter, but rather, as explained in [2], to zero energy density at that point. Also, the solution introduces into the system the new physical parameter r_μ with dimensions of length, which can be associated to a mass parameter μ in the same way that M is associated to r_M .

1.2 The Interior Schwarzschild Solution

It is an interesting and somewhat remarkable fact that the well-known interior Schwarzschild solution [6, 7] can be obtained from our solution for a shell, even though the interior Schwarzschild solution has no singularity at the origin, while our solution always has that singularity. Curiously enough, we must start by assuming that $r_\mu = 0$, even though we proved in [2] that one must have $r_\mu > 0$ in the shell solutions. The subtle point here is that the proof given in [2] relies on the existence of a shell with $r_1 > 0$, while in the case of the interior Schwarzschild solution we will have to use $r_1 = 0$, so that the shell becomes a filled sphere. If we start by first putting $r_\mu = 0$ and then making $r_1 \rightarrow 0$ in Equation (8), we are led to the relation

$$\kappa\rho_0 = \frac{3r_M}{r_2^3}, \quad (11)$$

so that we may substitute $\kappa\rho_0$ in terms of r_M and the radius r_2 of the resulting sphere. Following the usual notation for the interior Schwarzschild solution, we now define a parameter R , with dimensions of length, such that $R^2 = r_2^3/r_M$, in terms of which we have

$$\kappa\rho_0 = \frac{3}{R^2}. \quad (12)$$

Note that the required condition that $r_2 > r_M$ is translated here as the condition that $R > r_2$. Making this substitution we have for $\lambda(r)$ inside the resulting sphere, directly from the line in Equation (5) for the case of the matter region, in the case in which $r_\mu = 0$ and $r_1 \rightarrow 0$,

$$\lambda_i(r) = -\frac{1}{2} \ln \left[1 - \left(\frac{r}{R} \right)^2 \right], \quad (13)$$

which implies that for the radial metric coefficient we have

$$e^{-\lambda_i(r)} = \sqrt{1 - \left(\frac{r}{R} \right)^2}. \quad (14)$$

In order to obtain $\nu(r)$ inside the sphere we must first work out the function $z(r)$. Making the substitution of $\kappa\rho_0$ in terms of R in the result for $z(r)$ given in Equation (10) we get

$$z(r) = \sqrt{1 - \left(\frac{r}{R} \right)^2} \left[\sqrt{\frac{r_2}{r_2 - r_M}} + \frac{3}{2} \int_{r_2}^r ds \frac{s/R^2}{(1 - s^2/R^2)^{3/2}} \right]. \quad (15)$$

It is now easy to see that in this case the remaining integral can be done, and we get

$$z(r) = \frac{3}{2} - \frac{1}{2} \sqrt{\frac{r_2}{r_2 - r_M}} \sqrt{1 - \left(\frac{r}{R} \right)^2}. \quad (16)$$

Using again the definition of R , which implies that we have $r_M/r_2 = (r_2/R)^2$, we may write this as

$$z(r) = \frac{3}{2} - \frac{1}{2} \sqrt{\frac{1 - (r/R)^2}{1 - (r_2/R)^2}}. \quad (17)$$

Note that we have $z(r_2) = 1$, which corresponds to $P(r_2) = 0$, so that the boundary conditions for $z(r)$ and $P(r)$ at r_2 are still satisfied. From this we may now obtain all the remaining results for the interior Schwarzschild solution. From the line in Equation (6) for the case of the matter region, in the case in which $r_\mu = 0$ and $r_1 \rightarrow 0$, we get for $\nu(r)$ in the interior of the sphere

$$\nu_i(r) = \frac{1}{2} \ln \left[1 - \left(\frac{r_2}{R} \right)^2 \right] + \ln \left[\frac{3}{2} - \frac{1}{2} \sqrt{\frac{1 - (r/R)^2}{1 - (r_2/R)^2}} \right], \quad (18)$$

which implies that for the temporal metric coefficient we have

$$e^{\nu_i(r)} = \frac{3}{2} \sqrt{1 - \left(\frac{r_2}{R} \right)^2} - \frac{1}{2} \sqrt{1 - \left(\frac{r}{R} \right)^2}. \quad (19)$$

Finally, from Equation (7), in the case in which $r_\mu = 0$ and $r_1 \rightarrow 0$, we get for the pressure $P(r)$ within the sphere

$$P(r) = \rho_0 \frac{\sqrt{1 - (r/R)^2} - \sqrt{1 - (r_2/R)^2}}{3\sqrt{1 - (r_2/R)^2} - \sqrt{1 - (r/R)^2}}. \quad (20)$$

These are indeed the correct results for the case of the interior Schwarzschild solution. Note that all the arguments of the logarithms and of the square roots are positive due to the conditions that $R > r_2 > r$. Note also that in the $r_1 \rightarrow 0$ limit the lines in Equations (5) and (6) for the case of the inner vacuum region become irrelevant, since this region reduces to a single point. On the other hand, the lines for the case of the outer vacuum region do not change at all.

It is therefore apparent that the $r_1 \rightarrow 0$ limit of our shell solutions does reproduce the interior Schwarzschild solution, so long as we adopt the value zero for r_μ . Our interpretation of these facts is that the $r_1 \rightarrow 0$ limit to the interior Schwarzschild solution is a *non-uniform* one, in which we have to leave out one point, the origin. In the $r_1 \rightarrow 0$ limit the singularity of the shell solutions becomes a strictly point-like one, and therefore a removable one, by a simple continuity criterion. This is certainly the case for the energy density $\rho(r)$, which in the limit is non-zero everywhere around the origin but at a single point, the origin itself. The same is true for the pressure $P(r)$, which in the limit is also non-zero around the origin but at the origin itself. Similar situations hold for $\lambda(r)$ and $\nu(r)$, as is not difficult to see numerically. It seems that all these functions converge in the $r_1 \rightarrow 0$ limit to functions with a point-like removable discontinuity at the origin.

2 Integral Expressions for the Energies

It is possible to express the masses M and μ , as well as the corresponding energies Mc^2 and μc^2 , which are associated to the parameters with dimensions of length $r_M = 2MG/c^2$ and $r_\mu = 2\mu G/c^2$ that appear in the exact solutions described in Section 1, as integrals of the matter energy density $\rho(r)$ over coordinate volumes, in a way similar to what is usually

done for M in the literature [3, 8], but leading to very different results in the case of the shell solutions. In order to do this in a simple and organized way, we first change variables in the field equations from $\lambda(r)$ to $\beta(r)$, which is defined to be such that

$$e^{2\lambda(r)} = \frac{r}{r - r_M \beta(r)}, \quad (21)$$

which then implies that we have for the corresponding derivatives

$$2r\lambda'(r) = -r_M \frac{\beta(r) - r\beta'(r)}{r - r_M \beta(r)}. \quad (22)$$

Note that $\beta(r) = 0$ corresponds to $\lambda(r) = 0$ and therefore to $\exp[2\lambda(r)] = 1$ for the radial coefficient of the metric. In such cases the variations of the radial coordinate are equal to the variations of the corresponding proper lengths. Substituting these expressions in the component field equation shown in Equation (2) a very simple relation giving the derivative of $\beta(r)$ in terms of $\rho(r)$ results,

$$\beta'(r) = \frac{\kappa r^2 \rho(r)}{r_M}. \quad (23)$$

Therefore, wherever $\rho(r) = 0$, we have that $\beta(r)$ is a constant. Note that these facts are completely general for the spherically symmetric static case, in the sense that they are not limited to the case in which $\rho(r)$ is constant within the matter region. It then follows from Equation (5) that we have that $\beta(r) = 1 > 0$ in the outer vacuum region, and in particular at r_2 , and that we have that $\beta(r) = -r_\mu/r_M < 0$ in the inner vacuum region, and in particular at r_1 . Since $\beta(r)$ is a continuous function that goes from negative values at r_1 to positive values at r_2 , it follows that there is a radial position r_z within the matter region where $\beta(r_z) = 0$, regardless of whether or not $\rho(r)$ is constant within the shell. At this particular radial position we also have that $\lambda(r_z) = 0$.

Let us now consider the integral of the energy density over a coordinate volume within the matter region, where $\rho(r) \neq 0$, say from an arbitrary point r_a to another point $r_b > r_a$,

$$\int_{r_a}^{r_b} dr \int_0^\pi d\theta \int_0^{2\pi} d\phi r^2 \sin(\theta) \rho(r) = 4\pi \int_{r_a}^{r_b} dr r^2 \rho(r), \quad (24)$$

where we integrated over the angles. Note that this is not an integral over the proper volume, but just an integral over the coordinate volume, since we are missing here the remaining factor $\exp[\lambda(r) + \nu(r)]$ of the Jacobian $\sqrt{-g}$. Since we have the three special points r_1 , r_z and r_2 where the values of $\beta(r)$ are known, let us consider now the integral of the energy density over the coordinate volume from r_z to r_2 . Using Equation (23) we get

$$4\pi \int_{r_z}^{r_2} dr r^2 \rho(r) = 4\pi \frac{r_M}{\kappa} \int_{r_z}^{r_2} dr \beta'(r). \quad (25)$$

One can now see that the integral is trivial, and since we have that $\beta(r_z) = 0$ and that $\beta(r_2) = 1$, we get

$$Mc^2 = 4\pi \int_{r_z}^{r_2} dr r^2 \rho(r), \quad (26)$$

where we have replaced κ and r_M by their values in terms of M and c . We have therefore an expression for the energy Mc^2 in terms of a coordinate volume integral of the energy density. Note however that the integral does not run over the whole matter region, since it

starts at r_z rather than at r_1 . In a similar way, if we consider the integral from r_1 to r_z , we get

$$4\pi \int_{r_1}^{r_z} dr r^2 \rho(r) = 4\pi \frac{r_M}{\kappa} \int_{r_1}^{r_z} dr \beta'(r). \quad (27)$$

Once again one can see that the integral is trivial, and since we have that $\beta(r_z) = 0$ and that $\beta(r_1) = -r_\mu/r_M$, we now get

$$\mu c^2 = 4\pi \int_{r_1}^{r_z} dr r^2 \rho(r), \quad (28)$$

where we have replaced κ and r_μ by their values in terms of μ and c . We have therefore an expression for the energy μc^2 in terms of a coordinate volume integral of the energy density.

If we now consider the integral over the whole matter region, due to the additive property of the integrals over the union of disjoint domains, using Equations (26) and (28) we obtain the result that

$$4\pi \int_{r_1}^{r_2} dr r^2 \rho(r) = \mu c^2 + M c^2. \quad (29)$$

This is a sum of energies, and is therefore also an energy, to which we will associate a mass parameter M_u , such that this energy is given by $M_u c^2$, so that we have the relation

$$M_u c^2 = \mu c^2 + M c^2. \quad (30)$$

We see therefore that the point r_z where $\beta(r_z) = 0$ and therefore $\lambda(r_z) = 0$ plays a particular role when it comes to the determination of the energies involved.

Note that all this is true for any function $\rho(r)$ within the matter region. For our specific case here, with a constant ρ_0 , we find from Equation (5) that we have within the matter region

$$\beta(r) = 1 - \frac{\kappa \rho_0}{3r_M} (r_2^3 - r^3), \quad (31)$$

so that in this case we have for the zero r_z of $\beta(r)$

$$r_z = \left(r_2^3 - \frac{3r_M}{\kappa \rho_0} \right)^{1/3}. \quad (32)$$

Note that, although all these integrals are written in terms of the energy density $\rho(r)$ of the matter, none of them represents just the energy of only the matter itself. In fact we must now interpret the meaning of each one of these expressions, which is what we will do in the next section.

3 Physical Interpretation of the Energies

Of the three energies at play here, namely $M_u c^2$, μc^2 and $M c^2$, only the last one has a well established meaning at this point. Since M is the asymptotic gravitational mass of the system, that is, the gravitational mass seen as the source of the gravitational field at large radial distances, the standard interpretation in General Relativity is that the energy $M c^2$ is the total energy of this gravitational system, bound into the shell by the gravitational interactions, and which from now on we will simply call the *bound system*. It includes both the energy of the matter in the bound state and the energy stored in the gravitational field

itself, also in this bound state. The energy density $\rho(r)$ is the amount of energy of the matter, per unit volume, as seen by a stationary local observer at the radial position r .

Our first task here is to establish the physical interpretation of the energy $M_u c^2$. In order to do this, the first thing to be done is to define an *unbound system* related to our bound system as defined above. This unbound system is what we get when we scatter all the elements of the shell to very large distances from each other, in order to eliminate all the gravitational interactions, but without making any changes in the energy content of the matter. We will show here that the energy $M_u c^2$ is the total energy of this unbound system. We will do this by performing a mathematical transformation on the integral in Equation (29), which with the use of Equation (30) leads to the following expression in terms of a volume integral

$$M_u c^2 = \int_{r_1}^{r_2} dr \int_0^\pi d\theta \int_0^{2\pi} d\phi r^2 \sin(\theta) \rho(r). \quad (33)$$

The transformation, applied to the right-hand side of this equation, will allow us to interpret the meaning of the left-hand side. This will be done in a general way, for any function $\rho(r)$ within the matter region. This transformation will consist in fact of the construction of a second integral, based on the concept of the Riemann sums of the volume integral shown in Equation (33).

Let us consider therefore an arbitrary Riemann partition of the integral in Equation (33), consisting of a finite number of cells δV_n with coordinate volume and linear coordinate dimensions below certain maximum values, where $n \in \{1, \dots, N\}$. By definition of a partition the sum of all these volume elements is equal to the coordinate volume V of the shell,

$$V = \sum_{n=1}^N \delta V_n, \quad (34)$$

where we will assume that each volume element is at the spatial position \vec{r}_n , as illustrated in Figure 1. The energy $M_u c^2$ can therefore be written as the integration limit of the Riemann sum over this partition,

$$M_u c^2 = \lim_{N \rightarrow \infty} \sum_{n=1}^N \rho(r_n) \delta V_n, \quad (35)$$

where $r_n = |\vec{r}_n|$. We now consider the mathematical transformation in which we map each volume element δV_n at \vec{r}_n onto an identical volume element $\delta V'_n$ at the coordinate position $\vec{r}'_n = \alpha \vec{r}_n$, for some large positive real number α , without changing the coordinate volume of the volume elements. The result is a new set of volume elements, all at large distances from each other, whose sum is still equal to the coordinate volume of the shell,

$$V = \sum_{n=1}^N \delta V'_n, \quad (36)$$

The geometrical transformation leading to the construction of the new integral is illustrated in Figure 1. Note that no physical transport of the matter or of the energy within the volume elements δV_n of the shell is meant here, so that there are no actual physical transformations involved.

After defining the volume elements $\delta V'_n$ at large distances in this fashion, we now put within each one of these new volume elements exactly the same amount of mass and energy




Figure 1: Illustration of the geometrical transformation of the integral over the shell.

that we have in the corresponding coordinate volume elements δV_n of the shell. This means putting into each volume element $\delta V'_n$ at infinity the same numbers of the same types of particles, as well as the same amount of thermal energy and pressure, as seen by a stationary local observer at the position \vec{r}'_n , that a stationary local observer at \vec{r}_n sees within δV_n . In other words, we associate to each volume element at infinity the same value of the energy density $\rho(r'_n) = \rho(r_n)$ that we had for the corresponding volume element of the shell, where $r'_n = |\vec{r}'_n|$ and $r_n = |\vec{r}_n|$.

For large values of α these elements of mass and energy within $\delta V'_n$ are all at large distances from each other, so as to render the gravitational interactions among them negligible. In the $\alpha \rightarrow \infty$ limit all the gravitational interactions among the volume elements $\delta V'_n$ go to zero. Besides, in the integration limit each element of mass and energy so constructed tends to zero, so that the gravitational self-interactions within each volume element also become negligible. However, independently of either limit, by construction the total coordinate volume of the elements of volume at infinity remains equal to the coordinate volume of the shell. Therefore, by construction the corresponding sum of all the energy elements of energy at infinity is the same as the Riemann sum that appears in Equation (35),

$$\sum_{n=1}^N \rho(r'_n) \delta V'_n = \sum_{n=1}^N \rho(r_n) \delta V_n. \quad (37)$$

Now, at radial infinity spacetime is flat, so that the coordinate volume of each volume element $\delta V'_n$ coincides with its proper volume, and hence the energy element $\rho(r'_n) \delta V'_n$ is the total energy of that element of matter, so that the sum of all these energy elements is the total energy of the matter at infinity. In other words, once we take the integration limit the integral given in Equation (33) gives us the total energy of the system at infinity, which is free from all gravitational bindings. Hence we will name the quantity $M_u c^2$ the *total energy of the unbound system*. This is the total energy of the system when all gravitational inter-

actions have been eliminated by increasing without limit the distances among its elements. This is in both analogy and contrast with the quantity Mc^2 , which is the *total energy of the bound system*, after all its parts have been brought together to form the shell.

Note that this whole argument is general, in the sense that it is not limited to the case in which $\rho(r) = \rho_0$ is a constant. In our case here, since $\rho(r) = \rho_0$ is a constant, the total energy of the unbound system is just the product of ρ_0 by the coordinate volume V of the shell,

$$M_u c^2 = \rho_0 V. \quad (38)$$

Our next task here is to establish the physical interpretation of the energy μc^2 . From Equation (30) we have that the energy parameter μc^2 is the difference between the total energy of the unbound system and the total energy of the bound system,

$$\mu c^2 = M_u c^2 - M c^2, \quad (39)$$

and therefore we conclude that it is the *binding energy* of the system. It is the amount of energy that must be given to the system in order to disperse its elements to infinity, thus eliminating all the gravitational bindings between those elements. It is also the amount of energy that must be dissipated by the system during the process of its assembly into the bound system, starting from the unbound system at infinity. The theorem we proved in [2], in the $\rho(r) = \rho_0$ case that we have here, namely that we must have $r_\mu > 0$, is equivalent to the statement that the bound system must have a finite, positive and non-zero binding energy. This is, of course, closely related to the attractive nature of the gravitational interaction between particles.

Note that, although all these integrals are written in terms of the energy density $\rho(r)$ of the matter, the energy Mc^2 is *not* the energy $M_m c^2$ of just the matter within the bound system. That would be given by the integral with the full Jacobian factor $\sqrt{-g}$, where g is the determinant of $g_{\mu\nu}$, which in our case here results in

$$M_m c^2 = 4\pi \int_{r_1}^{r_2} dr r^2 e^{\lambda(r)+\nu(r)} \rho(r). \quad (40)$$

As a partial consistency check, it is not difficult to verify that this energy is always smaller than $M_u c^2$, due to the fact that the exponent $\lambda(r) + \nu(r)$ is always negative within the matter region. In order to show this we just take the difference between the component field equations shown in Equations (3) and (2), thus obtaining

$$[\lambda(r) + \nu(r)]' = \frac{\kappa}{2} e^{2\lambda(r)} r [\rho(r) + P(r)]. \quad (41)$$

Since all quantities appearing on the right-hand side are positive or zero, we may conclude that the derivative of the exponent is non-negative. However, we have that $\lambda(r_2) + \nu(r_2) = 0$, since this exponent is identically zero within the outer vacuum region. It follows that

$$\lambda(r) + \nu(r) < 0, \quad (42)$$

and therefore that

$$e^{\lambda(r)+\nu(r)} < 1, \quad (43)$$

throughout the whole matter region, with the exception of the single point r_2 where the exponential is equal to one. Therefore, it follows for the two integrals that

$$4\pi \int_{r_1}^{r_2} dr r^2 e^{\lambda(r)+\nu(r)} \rho(r) < 4\pi \int_{r_1}^{r_2} dr r^2 \rho(r), \quad (44)$$

and therefore that $M_m c^2 < M_u c^2$. The difference $M c^2 - M_m c^2$ is the part of the energy of the bound system which is not the energy of the matter itself, but rather the energy stored in the gravitational field. In general, in order to determine this difference, $M_m c^2$ has to be calculated numerically.

3.1 Energetic Stability

This interpretation of the parameters involved leads right away to the idea that we may define a notion of *energetic stability* of the solutions obtained, in the general spirit of the principle of virtual work. Given certain constraints regarding some of the parameters of the solutions, we may obtain the parameter r_μ as a function of the remaining parameters of the system. Within this class of solutions, if there are two with different values of r_μ , which is proportional to the binding energy μc^2 , then in principle the constrained system will tend to go from the one with the smaller value of r_μ to the one with the larger value, given the existence of a permissible path between the two solutions. This type of analysis allows us to acquire some information about the dynamical behavior of the system, without having to find explicitly the corresponding time-dependent solutions.

Let us exemplify this with our current system, in a way that is physically illustrative. Our system contains four parameters, namely r_1 , r_2 , r_M and ρ_0 , of which only three are independent. As was explained in [2], these four parameters are related by the condition in Equation (9). Given any three of the parameters, that equation can be used to determine the fourth in terms of those three. Let us assume that we are given fixed values of both M and ρ_0 , thus determining the local properties of the matter and the total amount of energy of the bound system. This is equivalent to fixing r_M and ρ_0 , and therefore the result of solving Equation (9) is to establish r_1 as a function of r_2 . We therefore are left with a collection of solutions parametrized by a single real parameter, the external radius r_2 . We may then determine $r_\mu(r_2)$ and verify whether this function has a single local maximum at a certain value of r_2 . This then identifies that particular solution which is stable, or that has the largest binding energy, among all others, given the constraints described.

Another approach, slightly more indirect, but perhaps simpler and more physically compelling, would be to keep constant the local parameter ρ_0 and the energy $M_u c^2$ of the unbound system. This fixes the local properties of the matter and the total energy of the unbound system that we start with, and we may then ask which is the solution that corresponds to the most tightly bound system that can be assembled from that unbound system. Since the energy of the unbound system is the product of ρ_0 by the coordinate volume V of the shell, as can be seen in Equation (38), keeping fixed both ρ_0 and M_u corresponds to keeping fixed at a value V_0 that coordinate volume, which is given by

$$V_0 = \frac{4\pi}{3} (r_2^3 - r_1^3). \quad (45)$$

This immediately determines r_2 as a simple function $r_2(r_1)$ of r_1 . Then solving Equation (9) results in r_M being given as a function $r_M(r_1)$ of r_1 for the fixed value of ρ_0 and the fixed coordinate volume V_0 . This corresponds to the energy of the bound system with internal radius r_1 , for the given fixed values of ρ_0 and V_0 . The minimum of this function gives us the value of r_1 that corresponds to the most tightly bound system that can be assembled from a given unbound system. Other solutions in the same family, with other values of r_1 ,

will tend to decay into this one, given a permissible decay path between the two solutions involved. We will execute this program numerically in Section 4.

We saw that in the case of the interior Schwarzschild solution we have the value zero for r_μ . This implies that the resulting solution has zero gravitational binding energy, and that its energy is the same as the energy of the corresponding unbound system, which is a very strange and even bizarre situation indeed. This means that the resulting solution is not only energetically unstable, but that it is in fact *maximally* energetically unstable, since the bound system cannot possibly have more energy than the unbound system. Given a permissible path, in principle one would be able to disassemble the matter distribution of the interior Schwarzschild solution, taking every element of matter to infinity, without giving any energy at all to the system. This is quite unrealistic, and may be the reason why this solution has never proved to be a very useful one.

4 Numerical Exploration of the Binding Energy

Here we will explore numerically the issues of the binding energy and of the energetic stability of the shell solutions. In this exploration we will keep fixed the local energy density parameter ρ_0 , as well as the total energy $M_u c^2$ of the unbound system. Our objective will be then to determine the existence and the parameters of the maximally bound shell solution. We will do this by calculating the energy Mc^2 of the bound system and showing that it has a point of minimum as a function of r_1 . Since we keep fixed the parameter ρ_0 , and since the energy of the unbound system is given by $M_u c^2 = \rho_0 V_0$, this implies that we also keep fixed the coordinate volume V_0 of the shell, given in Equation (45), which immediately establishes r_2 as a given function of r_1 ,

$$r_2(r_1) = \left(r_1^3 + \frac{3V_0}{4\pi} \right)^{1/3}. \quad (46)$$

Therefore, of the three free parameters of our solutions, which can be taken to be r_1 , r_2 and ρ_0 , one is being kept fixed and another is a given function, so that we are left with only one free parameters, which we will take to be r_1 . Under these circumstances we have that r_M , and therefore both the mass M and the energy Mc^2 of the bound system, are functions of r_1 , with values that are left to be determined numerically.

In order to perform the numerical work it is convenient to first rescale the variables, creating a set of equivalent dimensionless variables. Since under these conditions $\kappa\rho_0$ is a constant which has dimensions of inverse square length, we will define a constant r_0 with dimensions of length by

$$r_0 = \frac{1}{\sqrt{\kappa\rho_0}}. \quad (47)$$

Having now the known constant r_0 , we use it in order to define the set of dimensionless parameters given by

$$\begin{aligned} \xi_1 &= \frac{r_1}{r_0}, \\ \xi_2 &= \frac{r_2}{r_0}, \\ \xi_M &= \frac{r_M}{r_0}, \\ \vartheta_0 &= \frac{3V_0}{4\pi r_0^3}, \end{aligned} \quad (48)$$




Figure 2: Graph of the energy of the bound system as a function of ξ_1 , for a fixed energy of the unbound system, given by $\vartheta_0 = 2$, and with ξ_1 in $[1, 5]$.

where ϑ_0 is the ratio between the coordinate volume V_0 of the shell and the volume of an Euclidean sphere of radius r_0 . The expression in Equation (46) giving r_2 as a function of r_1 is now translated as

$$\xi_2(\xi_1) = (\vartheta_0 + \xi_1^3)^{1/3}. \quad (49)$$

Note, for subsequent use, that this can also be written as $\xi_2^3 - \xi_1^3 = \vartheta_0$. The relation which we must now use in order to determine ξ_M is that given in Equation (9), which upon rescalings by r_0 can be written as

$$\sqrt{\frac{\xi_2}{3(\xi_2 - \xi_M)}} = \sqrt{\frac{\xi_1}{\xi_2^3 - \xi_1^3 + 3(\xi_1 - \xi_M)}} + \frac{3}{2} \int_{\xi_1}^{\xi_2} d\xi \frac{\xi^{5/2}}{[\xi_2^3 - \xi^3 + 3(\xi - \xi_M)]^{3/2}}, \quad (50)$$

where we changed variables in the integral from r to $\xi = r/r_0$. Substituting for ϑ_0 where possible we have the following non-trivial algebraic equation that determines ξ_M and therefore r_M ,

$$\sqrt{\frac{\xi_1}{\vartheta_0 + 3(\xi_1 - \xi_M)}} - \sqrt{\frac{\xi_2}{3(\xi_2 - \xi_M)}} + \frac{3}{2} \int_{\xi_1}^{\xi_2} d\xi \frac{\xi^{5/2}}{[\xi_2^3 - \xi^3 + 3(\xi - \xi_M)]^{3/2}} = 0. \quad (51)$$

Our objective here is to solve this equation in order to get $\xi_M(\xi_1)$, given a fixed value of ϑ_0 and with ξ_2 given by Equation (49). Note that, due to the homogeneous scalings leading




Figure 3: Graph of the energy of the bound system as a function of ξ_1 , for a fixed energy of the unbound system, given by $\vartheta_0 = 5$, and with ξ_1 in $[1, 5]$.

from the dimensionfull quantities to the dimensionless ones, shown in Equation (48), each solution of this equation is valid for any value of ρ_0 , which no longer appears explicitly. The same is true of the graphs to be generated using this equation. Given a value of ϑ_0 , the corresponding graph represents the results for all the possible strictly positive values of the energy density ρ_0 .

There are two main numerical tasks here, the calculation of the integral and the resolution of this algebraic equation for ξ_M . The integral can be readily and efficiently calculated by a cubic interpolation method, using the values of the integrand and of its derivative at the two ends of each integration interval. So long as we can return the value of the integral without too much trouble, Equation (51) can be readily and efficiently solved by an exponential sandwich (or bisection) method [9]. There are two readily available and robust initial upper and lower bounds for the value of ξ_M , the minimum possible lower bound being zero, and the maximum possible upper bound being the energy of the unbound system, since we must have that $Mc^2 < M_uc^2$, which in terms of the dimensionless parameters translates as $\xi_M < \vartheta_0/3$. We may therefore start the process with a lower bound $\xi_{M\ominus} = 0$ and an upper bound $\xi_{M\oplus} = \vartheta_0/3$ for ξ_M . In practice, the efficiency of this algorithm may be highly dependent on the use of a tighter pair of bounds.

A few examples of the functions obtained in this way can be seen in Figures 2 through 5, which show ξ_M as a function of ξ_1 , for fixed values of the energy of the unbound system, that is, for fixed values of ϑ_0 . Each graph consists of 81 data points. In order to ensure good numerical precision we used 10^6 integration intervals in the domain $[\xi_1, \xi_2]$. The exponential sandwich was iterated until a relative precision of the order of 10^{-12} was reached. The four graphs shown were generated on a high-end PC in approximately 25 hours, 15 hours,




Figure 4: Graph of the energy of the bound system as a function of ξ_1 , for a fixed energy of the unbound system, given by $\vartheta_0 = 10$, and with ξ_1 in $[1, 5]$.

62 hours and 154 hours, respectively, without too much preoccupation with efficiency. As one can see, the graphs clearly display minima of ξ_M , which are located at certain values of ξ_1 . At these minima the pairs of values (ξ_1, ξ_2) are given approximately, in each case, by $(2.79, 2.87)$, $(2.72, 2.93)$, $(2.60, 3.02)$ and $(2.35, 3.21)$, respectively. There is freely available an open-source program [10] that can be used to perform these calculations for any set of input parameters.

The minima of these functions give us the value of ξ_1 that corresponds to the most tightly bound system that can be assembled from the given unbound system in each case. With the given values of ρ_0 and $M_u c^2$, in each case this establishes the value of r_1 for the most tightly bound and therefore energetically stable solution, and hence determines the values of r_2 , r_M and of all the functions describing both the spacetime geometry and the matter for that stable solution. The limiting value of ξ_M when $\xi_1 \rightarrow 0$, not shown in these graphs, corresponds to the interior Schwarzschild solution and thus to the energy of the unbound system in each case, which in terms of the variables shown in the graphs is given by $\vartheta_0/3$. The $\xi_1 \rightarrow \infty$ limit to the other side rises fairly slowly and does not seem to approach this same value asymptotically, a situation that is probably due to the fact that an infinitesimally thin shell at infinity still has some binding energy, as compared to the corresponding set of isolated infinitesimal point masses.




Figure 5: Graph of the energy of the bound system as a function of ξ_1 , for a fixed energy of the unbound system, given by $\vartheta_0 = 20$, and with ξ_1 in $[1, 5]$.

5 Conclusions

In this paper we have established the energetic interpretation of the exact solutions obtained in a previous paper for spherically symmetric shells of liquid fluid [2]. All the energies involved were precisely characterized, including the total energies of the unbound systems, the total energies of the bound systems, the gravitational binding energies, and the energies stored in the gravitational field. This led to a characterization of the stability of the bound systems in terms of their binding energies. We have identified a two-parameter family of energetically stable solutions, within the original three-parameter family of solutions. In a few cases the stable solutions were identified numerically. It is to be expected that the interpretations of the energies that were introduced here will be useful in other cases, such as those involving polytropes, white dwarfs and neutron stars.

In order to accomplish this, integral expressions for all the energies involved were presented, as integrals of the matter energy density over various coordinate volumes. All these expressions hold more generally than just in the case of constant energy density $\rho(r) = \rho_0$ that we are directly dealing with here. A particular radial position r_z within the matter region, at which we have $\lambda(r_z) = 0$ and therefore $\exp[\lambda(r_z)] = 1$ for the radial coefficient of the metric, was identified as playing a special role in relation to the integral expressions for the various energies. This is the single finite radial position where the three-dimensional space is neither stretched nor contracted, as compared to the behavior of the radial coordinate r .

The energetic interpretation was extended to the case of the two-parameter family of interior Schwarzschild solutions for filled spheres [6, 7], which can be obtained as a particular limit of the shell solutions, and which turn out to be maximally unstable ones. This means

that there is a strong tendency of the solution for a filled sphere to spontaneously generate an internal vacuum region and thus become a shell solution. This is clearly connected to the repulsive character of the gravitational field around the origin, in the case of the shell solutions, pushing matter and energy away from that origin, as was discussed and characterized in the previous paper [2]. Any small perturbation of the interior Schwarzschild solution will put this mechanism in action, thus leading to an energetic decay from that filled sphere solution to a shell solution.

The crucial development leading to all this was the introduction of the parameter r_μ in the previous paper, which was shown there to be necessarily strictly positive in that case, for the correct resolution of the differential equations and the corresponding interface boundary conditions, as implied by the Einstein field equations. The apparently traditional routine of choosing $r_\mu = 0$ in order to eliminate the singularity at the origin not only is often incompatible with the correct resolution of the differential system but, when it is not thus incompatible, it is tantamount to selecting a solution which has no binding energy at all and is therefore maximally unstable from the energetic point of view. Both from the purely mathematical point of view and from the physical point of view, this is more often than not the incorrect choice, which we are simply not at liberty to make.

Acknowledgments

The author would like to thank his friends Prof. C. E. I. Carneiro and Mr. Rodrigo de A. Orselli for their helpful criticism and careful reading of the manuscript.

References

- [1] P. A. M. Dirac, *General Theory of Relativity*. John Wiley & Sons, Inc., 1975. ISBN 0-471-21575-9.
- [2] J. L. deLyra, R. de A. Orselli, and C. E. I. Carneiro, “Exact solution of the einstein field equations for a spherical shell of fluid matter,” *arXiv*, vol. gr-qc/2101.02012, 2021. Submitted to Physical Review D.
- [3] S. Weinberg, *Gravitation and Cosmology*. New York: John Wiley and Sons, 1972.
- [4] J. Ni, “Solutions without a maximum mass limit of the general relativistic field equations for neutron stars,” *Science China*, vol. 54, no. 7, pp. 1304–1308, 2011.
- [5] L. Neslušan, “Solutions without a maximum mass limit of the general relativistic field equations for neutron stars,” *Journal of Modern Physics*, vol. 6, pp. 2164–2183, 2015.
- [6] K. Schwarzschild, “Über das gravitationsfeld einer kugel aus inkompressibler flüssigkeit nach der einsteinschen theorie (on the gravitational field of a ball of incompressible fluid following einstein’s theory),” *Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften*, vol. 7, pp. 424–434, 1916.
- [7] R. Wald, *General Relativity*. University of Chicago Press, 2010.
- [8] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. San Francisco: W.H. Freeman and Co., 1973.

- [9] W. Press, B. Flannery, S. Teukolsky, and W. Vetterling, *Numerical Recipes in FORTRAN 77: Volume 1, Volume 1 of Fortran Numerical Recipes: The Art of Scientific Computing*. Cambridge University Press, 1992.
- [10] “Energetic stability program for liquid shells.” Direct web access for dowload.
<http://sft.if.usp.br/scientific/>.