

Companion varieties for root systems and Fermat arrangements

Roberta Di Gennaro, Giovanna Ilardi, Rosa Maria Miró-Roig
Tomasz Szemberg, Justyna Szpond

February 20, 2021

Abstract

Unexpected hypersurfaces are a brand name for some special linear systems. They were introduced around 2017 and are a field of intensive study since then. They attracted a lot of attention because of their close ties to various other areas of mathematics including vector bundles, arrangements of hyperplanes, geometry of projective varieties. Our research is motivated by the what is now known as the BMSS duality, which is a new way of deriving projective varieties out of already constructed. The last author coined the concept of companion surfaces in the setting of unexpected curves admitted by the B_3 root system. Here we extend this construction in various directions. We revisit the configurations of points associated to either root systems or to Fermat arrangements and we study the geometry of the associated varieties and their companions.

1 Introduction

In the present note we study companion varieties of unexpected hypersurfaces associated to some symmetric configurations of points in projective spaces. Companion varieties were introduced by the last author in [21] in the course of her study of Togliatti-type surfaces.

It is a classical problem in algebraic geometry to determine the dimension of linear systems. A lot of attention was given to linear systems with imposed base loci, i.e., of the form

$$H^0(X; L \otimes I(Z)), \quad (1)$$

where L is a positive (e.g. ample or very ample) line bundle on a smooth variety X and Z is a subscheme of X . Even in the very simple situation when X is the projective plane, L is the line bundle $\mathcal{O}_{\mathbb{P}^2}(d)$ for $d > 0$ and Z is a zero-dimensional subscheme of \mathbb{P}^2 the problem of computing the dimension of the vector space in (1) is far from being completely understood. Indeed, the two open conjectures: one due to Nagata (1959) [19] and the other the SHGH-Conjecture package due to Segre (1969), Harbourne (1986), Gimigliano (1987) and Hirschowitz (1989) (see [8] for introduction and recent progress) provide sufficient evidence for this claim. In any case, it is well-understood and clear that a single general point, or a fat point scheme concentrated in a single general point impose independent conditions on homogeneous polynomials of any fixed degree in a projective space of arbitrary dimension. With this in mind, it came as a surprise when Cook II, Harbourne, Migliore and Nagel [3], inspired by earlier results due to Di Gennaro, Ilardi and Vallès [5], announced that a single general fat point might impose less conditions than

Key words and phrases. unexpected hypersurfaces, arithmetically Cohen-Macaulay varieties, root systems, Fermat-type arrangements, Veronese varieties.

2010 *Mathematics subject classification.* 13A15, 13C70, 14C20, 14E05, 14J70, 14N20

expected on the linear system of homogeneous polynomials with assigned base loci. The elements arising this way are called *unexpected hypersurfaces*, see Section 2 for precise definitions.

The unexpected hypersurfaces are interesting not only because they violate the naive conditions count when determining the dimension of a linear system. In a private communication to the last author Igor Dolgachev suggested that, under additional positivity conditions, there should be a relation between the existence of unexpected hypersurfaces and varieties with defective osculating spaces. This has been indeed established in [22] for a configuration of points determined by the B_3 root system. The linear system of quadrics vanishing at configuration points determines, after passing to the blow up of \mathbb{P}^2 in these points, a morphism to \mathbb{P}^5 whose image is a surface S such that at every point P of S , there is a hyperplane in \mathbb{P}^5 tangent to P to order 2 (in other words: cutting out on S a curve which passes through P with multiplicity at least 3). It has been observed additionally, that there is another surface S' , which we call a companion surface of S , which exhibits also interesting geometrical properties. This example motivates our present work.

The general yoga is as follows. Assume that there is a set of points Z in \mathbb{P}^N which admits a unique unexpected hypersurface $H_{Z,P}$ of degree d and multiplicity m at a general point $P = (a_0 : \dots : a_N) \in \mathbb{P}^N$. Let

$$F_Z((x_0 : \dots : x_N), (a_0 : \dots : a_N)) = 0$$

be a homogeneous polynomial equation of $H_{Z,P}$. Let g_0, \dots, g_M be a basis of the vector space $[I(Z)]_d$ of homogeneous polynomials of degree d vanishing at all points of Z . Under some mild hypothesis the unexpected hypersurface $H_{Z,P}$ comes from a bi-homogeneous polynomial $F_Z((x_0 : \dots : x_N), (a_0 : \dots : a_N))$ of bidegree (m, d) [17]. Indeed, F_Z can be written in a unique way as a combination

$$F_Z = h_0(a_0 : \dots : a_N)g_0(x_0 : \dots : x_N) + \dots + h_M(a_0 : \dots : a_N)g_M(x_0 : \dots : x_N) \quad (2)$$

where $g_0(x_0 : \dots : x_N), \dots, g_M(x_0 : \dots : x_N)$ are homogeneous polynomials of degree d and $h_0(a_0 : \dots : a_N), \dots, h_M(a_0 : \dots : a_N)$ are homogeneous polynomials of degree m . Therefore, there are two rational maps naturally associated to equation (2)

$$\varphi : \mathbb{P}^N \ni (x_0 : \dots : x_N) \dashrightarrow (g_0(x_0 : \dots : x_N) : \dots : g_M(x_0 : \dots : x_N)) \in \mathbb{P}^M$$

and

$$\psi : \mathbb{P}^N \ni (a_0 : \dots : a_N) \dashrightarrow (h_0(a_0 : \dots : a_N) : \dots : h_M(a_0 : \dots : a_N)) \in \mathbb{P}^M.$$

The images of these maps are the *companion varieties*. The purpose of this note is to study their properties and relations.

We focus here on three natural generalizations of the aforementioned B_3 configuration. In what follows H is always the pull-back of the hyperplane bundle under the appropriate blow up.

First, it was established already in [17] that all B_n configurations give rise to unexpected hypersurfaces in \mathbb{P}^{n-1} . In case of $n = 4$ there is also a unique unexpected hypersurface and we determine its explicit equation in (4) and in slightly different generators in (6). It turns out that the unexpected surface for B_4 is also unexpected for F_4 , a non-crystallographic root system. The other difference when compared to the B_3 root system is that now the degree of the unexpected surface equals its multiplicity in the general point, i.e., the unexpected surfaces are cones.

Theorem B_4/F_4 . *Let Y be the blow up of \mathbb{P}^3 at points in the F_4 root system with exceptional divisors E_1, \dots, E_{24} . Then the linear system $4H - \sum E_i$ embeds Y into \mathbb{P}^{11} as a smooth threefold X of degree 40. The companion threefold X' is in this case isomorphic to X .*

This is proved in Proposition 4.3 and Remark 4.4.

Secondly, we study the H_3 root system. Similarly as B_3 for $m = 3$, it admits an unexpected curve of degree $m + 1$ with multiplicity m at the general point, this time with $m = 5$.

Theorem H_3 . *Let Y be the blow up of P^2 at points in the H_3 root system with exceptional divisors E_1, \dots, E_{15} . Then the linear system $6H - \sum E_i$ embeds Y as a smooth aCM surface X of degree 21. The companion surface X' is not linearly normal embedded into \mathbb{P}^{11}*

This result is proved in Proposition 4.5 and Remarks 4.6 and 4.7. A very surprising feature of this example is that the companion surface X'' of X' is not Y again but it is rather its projection.

Finally, we study companion surfaces related to Fermat arrangements. The B_3 configuration belongs also to this family. Our main results in this direction are Theorem 5.6, Proposition 5.7 and Remark 5.8. The precise statements are slightly too technical to quote them here.

We work over the field of complex numbers \mathbb{C} .

2 Unexpected hypersurfaces

In the ground-breaking work [3] by Cook II, Harbourne, Migliore and Nagel introduced the concept of unexpected curves. This notion was generalized to arbitrary hypersurfaces in the subsequent article [17] by Harbourne, Migliore, Nagel and Teitler.

Definition 2.1. We say that a reduced set of points $Z \subset \mathbb{P}^N$ admits an *unexpected hypersurface* of degree d if there exists a sequence of non-negative integers m_1, \dots, m_s such that for general points P_1, \dots, P_s the zero-dimensional subscheme $P = m_1P_1 + \dots + m_sP_s$ fails to impose independent conditions on forms of degree d vanishing along Z and the set of such forms is non-empty. In other words, we have

$$h^0(\mathbb{P}^N; \mathcal{O}_{\mathbb{P}^N}(d) \otimes I(Z) \otimes I(P)) > \max \left\{ 0, h^0(\mathbb{P}^N; \mathcal{O}_{\mathbb{P}^N}(d) \otimes I(Z)) - \sum_{i=1}^s \binom{N + m_i - 1}{N} \right\}.$$

Let us illustrate the above definition with an example.

Example 2.2. We consider the extended Fermat arrangement of planes in \mathbb{P}^3 :

$$F = xyz t(x^4 - y^4)(x^4 - z^4)(x^4 - t^4)(y^4 - z^4)(y^4 - t^4)(z^4 - t^4)$$

and we denote by $Z \subset \mathbb{P}^3$ the set of 28 points dual to the linear factors of F . We have:

$$I(Z) = (xyz, xyt, xzt, yzt, \\ xy(x^4 - y^4), xz(x^4 - z^4), xt(x^4 - t^4), yz(y^4 - z^4), yt(y^4 - t^4), zt(z^4 - t^4)).$$

Using Macaulay2 [12], we easily check that there is one and only one unexpected surface of degree 6 containing Z and three general points P_1, P_2, P_3 of multiplicities $m_1 = m_2 = m_3 = 4$.

Following [2, Definition 2.5] we introduce also the following notion.

Definition 2.3 (Unexpected cone property). Let Z be a finite set of points in \mathbb{P}^N and let d be a positive integer. We say that Z has the *unexpected cone property* $\mathcal{C}(d)$, if for a general point $P \in \mathbb{P}^N$, there exists an unexpected (in the sense of Definition 2.1) hypersurface S_P of degree d and multiplicity d at P passing through all points in Z .

Example 2.4. We consider the set $Z \subset \mathbb{P}^3$ of 28 points introduced in Example 2.2. Using again Macaulay2 [12], we check that Z has the unexpected cone property $\mathcal{C}(6)$ since for a general point $P \in \mathbb{P}^3$, there exists an unexpected surface $S_P \subset \mathbb{P}^3$ of degree 6 and multiplicity 6 at P passing through all 28 points in Z .

3 BMSS duality

For seek of completeness let us explain what we understand for BMSS duality a notion which comes from [1] and [17].

We consider integers $0 < m \leq d \in \mathbb{Z}$, a set $Z \subset \mathbb{P}^n$ of points and a general point $P = (a_0 : a_1 : \dots : a_n) \in \mathbb{P}^n$. We assume that there is a unique hypersurface $H_Z \subset \mathbb{P}^n$ of degree d containing Z and having multiplicity m at P . The hypersurface H_Z is defined by a bihomogeneous form

$$F_Z(\underline{y}, \underline{x}) := F_Z((y_0 : y_1 : \dots : y_n), (x_0 : x_1 : \dots : x_n)) \in \mathbb{C}[\underline{y}, \underline{x}].$$

of bidegree (t, d) with $t \geq m$ (see [17, Lemma 3.1(d)]).

Note that the form must be bihomogeneous since if it were not, changing the projective coordinates of each of the two set of variables, the form would vary. Moreover, under mild hypothesis on Z the bidegree of $F_Z(\underline{y}, \underline{x})$ is (m, d) i.e. $t = m$ (See, for instance, [17] and [7]). It is important to point out that the bidegree of $F_Z(\underline{y}, \underline{x})$ is not always (m, d) . In fact, in [1, Theorem 6] the authors considered a set $W \subset \mathbb{P}^3$ of 31 points with an expected quartic surface $S_W \subset \mathbb{P}^3$ having a general point $Q \in \mathbb{P}^3$ of multiplicity 3 and defined by a bihomogeneous polynomial $F_W(\underline{y}, \underline{x})$ of bidegree $(5, 4)$.

The BMSS duality allows us to view the bihomogeneous polynomial $F_Z(\underline{y}, \underline{x}) \in \mathbb{C}[\underline{y}][\underline{x}]$ of bidegree (t, m) as a polynomial of degree d in the indeterminates \underline{x} with coefficients the point $P \in \mathbb{P}^n$ of multiplicity m , i.e. a family of hypersurfaces in the variables \underline{x} parameterized by P (the unexpected hypersurface):

$$F_{Z,P}(\underline{x}) := F_Z((a_0 : \dots : a_n), \underline{x}) \in \mathbb{C}[a_0, \dots, a_n][\underline{x}]$$

or, as a homogeneous polynomial of degree $t \geq m$ in the variables \underline{y} with coefficients a point $Q = (b_0 : \dots : b_n) \in \mathbb{P}^n$, i.e. a family of hypersurfaces in the variables \underline{y} parameterized by Q :

$$F_{Z,Q}(\underline{y}) := F_Z(\underline{y}, (b_0 : \dots : b_n)) \in \mathbb{C}[b_0, \dots, b_n][\underline{y}].$$

The BMSS duality establishes when the following assertions are true:

- The tangent cone of $F_{Z,P}(\underline{x})$ at Q coincide with the tangent cone of $F_{Z,Q}(\underline{y})$ at P .
- $F_{Z,P}(\underline{x})$ has multiplicity m at a general point Q and $F_{Z,Q}(\underline{y})$ has also multiplicity m at a general point P .

In next section we discuss in detail an example which comes from the free arrangement of planes in \mathbb{P}^3 determined by linear factors of:

$$xyz(x+y)(x-y)(x+z)(x-z)(x+t)(x-t)(y+z)(y-z)(y+t)(y-t)(z+t)(z-t)$$

as well as an example which comes from the root system H_3 .

4 Companion varieties for root systems

In [17] the authors study unexpected hypersurfaces related to root systems. A vector v in the root system in the vector space V determines a point in the projective space $\mathbb{P}(V)$, which corresponds to the line spanned by v . As the set Z one takes all points in $\mathbb{P}(V)$ determined in this way by all vectors in the root system.

The study in [17] was motivated by [5], where the first example of an unexpected curve coming from the B_3 root system was found. Computer experiments carried out in [17] provide the following list of root systems admitting a unique unexpected hypersurface. In the table, N stands for the dimension of the ambient projective space \mathbb{P}^N , d for the degree of an unexpected hypersurface and m for the multiplicity in the general point.

root system	N	d	m
B_3	2	4	3
B_4	3	4	4
D_4	3	3	3
F_4	3	4	4
H_3	2	6	5
H_4	3	6	6

Table 1: Root systems determining a unique unexpected hypersurface

As [17] and [1] study the B_3 case in detail, we turn our attention to the B_4 root system and to the H_3 root system. They are the only root systems determining a set of points Z with an unexpected hypersurface of degree d and multiplicity $m < d$ at a general point.

4.1 The B_4 root system

We will encounter a couple of new phenomena in the course. The set $Z(B_4)$ consists of 16 points, which can be assigned the following coordinates:

$$\begin{aligned}
 P_1 &= [1 : 0 : 0 : 0] & P_2 &= [0 : 1 : 0 : 0] & P_3 &= [0 : 0 : 1 : 0] & P_4 &= [0 : 0 : 0 : 1] \\
 P_5 &= [1 : 1 : 0 : 0] & P_6 &= [1 : 0 : 1 : 0] & P_7 &= [1 : 0 : 0 : 1] \\
 P_8 &= [0 : 1 : 1 : 0] & P_9 &= [0 : 1 : 0 : 1] & P_{10} &= [0 : 0 : 1 : 1] \\
 P_{11} &= [1 : -1 : 0 : 0] & P_{12} &= [1 : 0 : -1 : 0] & P_{13} &= [1 : 0 : 0 : -1] \\
 P_{14} &= [0 : 1 : -1 : 0] & P_{15} &= [0 : 1 : 0 : -1] & P_{16} &= [0 : 0 : 1 : -1]
 \end{aligned}$$

The saturated ideal $I(B_4)$ is generated by

$$xyz, xyw, xzw, yzw,$$

and

$$xy(x^2 - y^2), xz(x^2 - z^2), xw(x^2 - w^2), yz(y^2 - z^2), yw(y^2 - w^2), zw(z^2 - w^2). \quad (3)$$

In degree 4, the ideal $I(B_4)$ has, additionally to generators in (3), the following generators:

$$xyzw, x^2yz, xy^2z, xyz^2, x^2yw, xy^2w, xyw^2, x^2zw, xz^2w, xzw^2, y^2zw, yz^2w, yzw^2.$$

The unexpected cone, written down with the aid of above generators has then the following equation

$$\begin{aligned}
F = & 3ad(c^2 - b^2)x^2yz + 3bd(a^2 - c^2)xy^2z + 3cd(b^2 - a^2)xyz^2 + \\
& 3ac(b^2 - d^2)x^2yw + 3bc(d^2 - a^2)xy^2w + 3dc(a^2 - b^2)xyw^2 + \\
& 3ab(d^2 - c^2)x^2zw + 3cb(a^2 - d^2)xz^2w + 3db(c^2 - a^2)xzw^2 + \\
& 3ba(c^2 - d^2)y^2zw + 3ca(d^2 - b^2)yz^2w + 3da(b^2 - c^2)yzw^2 + \\
& cd(d^2 - c^2)xy(x^2 - y^2) + bd(b^2 - d^2)xz(x^2 - z^2) + bc(c^2 - b^2)xw(x^2 - w^2) + \\
& ad(d^2 - a^2)yz(y^2 - z^2) + ac(a^2 - c^2)yw(y^2 - w^2) + ab(b^2 - a^2)zw(z^2 - w^2).
\end{aligned} \tag{4}$$

It is immediately clear, that the generator $xyzw$ is not involved in the equation. We do not have an explanation for this phenomenon. The second property we observe is also quite surprising.

Proposition 4.1. *All unexpected cones for the B_4 root system, have in their base locus 8 additional points:*

$$\begin{aligned}
Q_1 &= [1 : 1 : 1 : 1] & Q_2 &= [1 : 1 : 1 : -1] & Q_3 &= [1 : 1 : -1 : 1] \\
Q_4 &= [1 : 1 : -1 : -1] & Q_5 &= [1 : -1 : 1 : 1] & Q_6 &= [1 : -1 : 1 : -1] \\
Q_7 &= [1 : -1 : -1 : 1] & Q_8 &= [1 : -1 : -1 : -1].
\end{aligned}$$

Proof. The vanishing of F at the given points is easy to check. That the base locus is not larger can be verified by computing equations of unexpected cones for some specific values of a, b, c, d . \square

The points in B_4 together with the points Q_1, \dots, Q_8 give the root system F_4 .

Corollary 4.2. *The unexpected cones for B_4 and F_4 are the same.*

This observation has interesting implications for the companion threefolds. First, let us note, that the ideal $I(F_4)$ is generated in degree 4 by the following forms:

$$\begin{aligned}
m_0 &= xy(x^2 - y^2), & m_1 &= xy(z^2 - w^2), & m_2 &= xz(x^2 - z^2), \\
m_3 &= xz(y^2 - w^2), & m_4 &= xw(x^2 - w^2), & m_5 &= xw(z^2 - y^2), \\
m_6 &= yz(y^2 - z^2), & m_7 &= yz(x^2 - w^2), & m_8 &= yw(y^2 - w^2), \\
m_9 &= yw(x^2 - z^2), & m_{10} &= zw(z^2 - w^2), & m_{11} &= zw(x^2 - y^2).
\end{aligned} \tag{5}$$

Using these generators the equation (4) becomes

$$\begin{aligned}
F = & cd(d^2 - c^2)xy(x^2 - y^2) + 3cd(b^2 - a^2)xy(z^2 - w^2) + bd(b^2 - d^2)xz(x^2 - z^2) + \\
& 3bd(a^2 - c^2)xz(y^2 - w^2) + bc(c^2 - b^2)xw(x^2 - w^2) + 3bc(a^2 - d^2)xw(z^2 - y^2) + \\
& ad(d^2 - a^2)yz(y^2 - z^2) + 3ad(c^2 - b^2)yz(x^2 - w^2) + ac(a^2 - c^2)yw(y^2 - w^2) + \\
& 3ac(b^2 - d^2)yw(x^2 - z^2) + ab(b^2 - a^2)zw(z^2 - w^2) + 3ab(d^2 - c^2)zw(x^2 - y^2).
\end{aligned} \tag{6}$$

Let $\varphi : \mathbb{P}^3 \dashrightarrow \mathbb{P}^{11}$ be the rational map defined by the generators in (5).

Proposition 4.3. *The image of φ is a smooth threefold of degree 40.*

Proof. Let $\sigma : X \rightarrow \mathbb{P}^3$ be the simultaneous blow up of each of 24 points in F_4 with the exceptional divisor \mathbb{E} (which splits into 24 projective planes, one over each of the points blown up) and as usual let $H = \sigma^*\mathcal{O}_{\mathbb{P}^3}(1)$. Since $\{p \in \mathbb{P}^3 \mid m_i(p) = 0 \text{ for } i = 0, \dots, 11\} = \{P_1, \dots, P_{16}, Q_1, \dots, Q_8\}$, the linear system $L = 4H - \mathbb{E}$ is base point free and it defines a morphism onto its image $\varphi_L : X \rightarrow \mathbb{P}^{11}$ which lifts the map

$$\varphi : \mathbb{P}^3 \dashrightarrow \mathbb{P}^{11} \quad (x : y : z : t) \mapsto (m_0 : m_1 : \dots : m_{11}).$$

Therefore, we have the commutative diagram

$$\begin{array}{ccc} Y & & \\ \downarrow \sigma & \searrow \varphi_L & \\ \mathbb{P}^3 & \dashrightarrow \varphi & \mathbb{P}^{11}. \end{array}$$

Let us call X the image of φ . To check that the map $\varphi : \mathbb{P}^3 \dashrightarrow Y$ has degree 1 and even more it is a birational map onto its image, we use the packages "Cremona" and "RationalMaps" of Macaulay2 [12]. In fact, we have the following code block:

```
loadPackage "Cremona"
loadPackage "RationalMaps"
R = QQ[x,y,z,w]
I = ideal(x*y*(x^2-y^2),x*y*(z^2-w^2),x*z*(x^2-z^2),x*z*(y^2-w^2),
x*w*(x^2-w^2),x*w*(y^2-z^2),y*z*(y^2-z^2),y*z*(x^2-w^2),
y*w*(y^2-w^2),y*w*(x^2-z^2),z*w*(z^2-w^2),z*w*(x^2-y^2))
degree rationalMap gens I
(Using the package "Cremona" we get that the degree is 1)
S=QQ[a_0..a_((numgens I)-1)]
L = apply(numgens I, j -> (gens I)_(0,j))
varphi = map(R, S, L)
isBirationalOntoImage(varphi)
(Using the package "RationalMaps" we get that the map is birational onto its image)
```

Once we know that φ is a birational map onto its image we compute the degree of Y : $\text{degree}(X) = L^3$. The smoothness of X follows from the Jacobian criterion and a straightforward computation which shows that outside $\{P_1, \dots, P_{16}, Q_1, \dots, Q_8\}$ the rank of $\left(\frac{\partial m_i}{\partial x} \quad \frac{\partial m_i}{\partial y} \quad \frac{\partial m_i}{\partial z} \quad \frac{\partial m_i}{\partial w}\right)_{i=0, \dots, 11}$ is 4.

It is worthwhile to point out that the Jacobian dual criterion as stated in [6, Theorem 2.18 and 3.2] or [20, Theorem 1.4] gives us a computer free proof of the birationality of φ . In fact, the ideal $I = (m_0, m_1, \dots, m_{11}) \subset R := \mathbb{C}[x, y, z, w]$ has a linear presentation:

$$\dots \longrightarrow R(-5)^{16} \xrightarrow{\psi} R(-4)^{12} \longrightarrow R \longrightarrow R/I \longrightarrow 0$$

with

$$\psi = \begin{pmatrix} -z & 0 & 0 & 0 & 0 & 0 & -w & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ y & -y & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -w & 0 & 0 & 0 & 0 & 0 \\ 0 & x & -y & 0 & -z & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -w & 0 & 0 & 0 \\ -y & 0 & x & -y & 0 & -z & 0 & 0 & 0 & 0 & 0 & w & 0 & 0 & -w & 0 \\ 0 & 0 & 0 & x & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & w & 0 & -w \\ z & -z & 0 & z & x & y & -w & w & 0 & -w & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & y & -y & 0 & 0 & z & -z & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & w & 0 & 0 & x & -y & 0 & 0 & 0 & z & -z & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & w & -y & 0 & x & -y & 0 & 0 & 0 & 0 & z & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & x & 0 & 0 & 0 & 0 & 0 & z \\ 0 & 0 & t & 0 & 0 & 0 & 0 & 0 & z & 0 & 0 & x & 0 & y & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & -x & 0 & y & 0 & x & -y \end{pmatrix}$$

Since I has a linear presentation and $\text{rank}(\psi)$ attains its maximal possible value, we apply the Jacobian dual criterion and we conclude that φ is a birational map onto its image. \square

Remark 4.4. It follows from the equation (6) of the unexpected cone that $Y = im(\varphi) \subset \mathbb{P}^{11}$ is selfdual under BMSS duality. This turns out to be true in a much more general set up. Indeed, let $Z \subset \mathbb{P}^n$ be a finite set of points and let $S_P \subset \mathbb{P}^n$ be an unexpected hypersurface of degree d and multiplicity d at a general point $P = (a_0 : \dots : a_n) \in \mathbb{P}^n$ passing through all points in Z . Consider

$$0 = F_Z((a_0 : \dots : a_n), (x_0 : \dots : x_n)) = \sum_{i=0}^N q_i(a_0 : \dots : a_n) f_i(x_0 : \dots : x_n)$$

the equation of the unexpected cone S_P . Set $X = im(\varphi)$ being $\varphi : \mathbb{P}^n \dashrightarrow \mathbb{P}^N$ the rational map defined by f_0, \dots, f_N . By [7], X is selfdual under the BMSS duality.

4.2 The H_3 root system

Associated to the root system H_3 we have a set $Z(H_3)$ of 15 points, which can be assigned the following coordinates:

$$P_1 = [1 : 0 : 0] \quad P_2 = [0 : 1 : 0] \quad P_3 = [0 : 0 : 1]$$

$$P_4 = [1 : u : u^2] \quad P_5 = [-1 : u : u^2] \quad P_6 = [1 : -u : u^2] \quad P_7 = [1 : u : -u^2]$$

$$P_8 = [u : -u^2 : 1] \quad P_9 = [-u : u^2 : 1] \quad P_{10} = [u : u^2 : -1] \quad P_{11} = [u : u^2 : 1]$$

$$P_{12} = [u^2 : 1 : -u] \quad P_{13} = [u^2 : -1 : u] \quad P_{14} = [-u^2 : 1 : u] \quad P_{15} = [u^2 : 1 : u]$$

where $u^2 - u - 1 = 0$, so that u is the golden ratio or its Galois conjugate $\bar{u} = 1 - u$. The configuration together with 6 lines, each of them containing 5 configuration points (2 of them are at the infinity) is visualised at Figure 1. The 10 thin lines contain 3 configuration points each. In order to increase the readability of the figure, only numbers of points are indicated.




Figure 1: The H_3 configuration of points with P_1 and P_2 at the infinity

The lines with 5 configuration points have rather simple equations:

$$L_1 : y - (u - 1)z = 0, \quad L_2 : y + (u - 1)z = 0, \quad L_3 : x - uz = 0, \quad L_4 : x + uz = 0,$$

$$L_5 : x - (u - 1)y = 0, \quad L_6 : x + (u - 1)y = 0.$$

The saturated ideal $I(H_3)$ is generated by six quintics. One can choose as the generators the products of all but one line:

$$\begin{aligned} f_1 &= L_2L_3L_4L_5L_6, & f_2 &= L_1L_3L_4L_5L_6, & f_3 &= L_1L_2L_4L_5L_6, \\ f_4 &= L_1L_2L_3L_5L_6, & f_5 &= L_1L_2L_3L_4L_6, & f_6 &= L_1L_2L_3L_4L_5. \end{aligned} \quad (7)$$

This follows from the fact that the lines L_1, \dots, L_6 form the so called star-configuration, see [11, Proposition 2.9]. Below we present equivalent argument based on the Hilbert-Burch theorem.

In degree 6, $I(H_3)$ has 13 generators. Indeed, $I(H_3)$ is generated by the 6 maximal minors of the 6×5 matrix U with linear entries:

$$U = \begin{pmatrix} L_1 & L_2 & 0 & 0 & 0 & 0 \\ 0 & L_2 & L_3 & 0 & 0 & 0 \\ 0 & 0 & L_3 & L_4 & 0 & 0 \\ 0 & 0 & 0 & L_4 & L_5 & 0 \\ 0 & 0 & 0 & 0 & L_5 & L_6 \end{pmatrix}.$$

Therefore, $I(H_3)$ has a minimal free resolution:

$$0 \longrightarrow S(-6)^5 \xrightarrow{U} S(-5)^6 \longrightarrow I(H_3) \longrightarrow 0 \quad (8)$$

with $S = \mathbb{C}[x, y, z]$ and we conclude that $\dim I(H_3)_6 = 13$. It is convenient to write down explicitly generators of $I(H_3)_6$:

$$g_i = L_1 f_i, \text{ for } i = 1, \dots, 6,$$

$$g_7 = L_2 f_1, \quad g_8 = L_2 f_3, \quad g_9 = L_2 f_4, \quad g_{10} = L_2 f_5, \quad g_{11} = L_2 f_6, \quad g_{12} = L_3 f_1, \quad g_{13} = L_3 f_2.$$

The set of points $Z(H_3)$ admits a unique unexpected curve $C \subset \mathbb{P}^2$ of degree 6 and multiplicity 5 at a general point $P = (a : b : c) \in \mathbb{P}^2$ defined by the following equation:

$$F_{Z(H_3)}((a : b : c), (x : y : z)) = \sum_{i=1}^{13} h_i(a : b : c) g_i(x : y : z),$$

where

$$h_1 = \left(\frac{7}{2}u + 2 \right) \cdot b \cdot (b^4 + (-44u + 72)ab^2c + (-6u + 8)b^2c^2 + (-4u + 12)ac^3 + (u - 3)c^4),$$

$$h_2 = \frac{-10u - 5}{4} \left(a + \frac{u + 2}{5}b + \frac{-3u - 1}{5}c \right) (b + (u - 1)c)^4,$$

$$h_3 = \frac{2u - 1}{4} (a + (u + 2)b + (u - 1)c) (a - uc)^4,$$

$$h_4 = \frac{-2u + 1}{4} (a - (u + 2)b - (u - 1)c) (a + uc)^4,$$

$$h_5 = \frac{4u + 3}{4} (a - (u - 1)b)^4 (a + ub - (u - 3)c),$$

$$h_6 = \frac{-4u - 3}{4} (a + (u - 1)b)^4 (a - ub + (u - 3)c),$$

$$h_7 = \frac{10u + 5}{4} \left(a + \frac{-u - 2}{5}b + \frac{-3u - 1}{5}c \right) (b - (u - 1)c)^4,$$

$$\begin{aligned}
h_8 &= \frac{2u-1}{4} (a - (u+2)b + (u-1)c) (a-uc)^4, \\
h_9 &= \frac{-2u+1}{4} (a + (u+2)b - (u-1)c) (a+uc)^4, \\
h_{10} &= \frac{-4u-3}{4} (a - (u-1)b)^4 (a+ub + (u-3)c), \\
h_{11} &= \frac{4u+3}{4} (a + (u-1)b)^4 (a-ub - (u-3)c), \\
h_{12} &= \frac{-3u-1}{2} (b+uc) (b - (u-1)c)^4, \\
h_{13} &= \frac{3u+1}{2} (b-uc) (b + (u-1)c)^4.
\end{aligned}$$




Figure 2: Unexpected sextic with a point of multiplicity 5

Figure 2 indicates the unexpected sextic curve with multiplicity 5 at a general point. The configuration points of the H_3 root system are arranged as in Figure 1. The lines through only 3 configuration points and the names of the points are omitted for transparency.

Let $\varphi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^{12}$ be the rational map defined by the 13 generators g_1, \dots, g_{13} of $I(Z(H_3))$.

Proposition 4.5. *The image of φ is an arithmetically Cohen-Macaulay smooth surface $X \subset \mathbb{P}^{12}$ of degree 21 whose homogeneous ideal $I(X)$ is generated by quadrics.*

Proof. We consider $\pi : Y \rightarrow \mathbb{P}^2$ the blow up of \mathbb{P}^2 at the 15 points of $Z(H_3)$. Set $H = \pi^* \mathcal{O}_{\mathbb{P}^2}(1)$ and denote by E_i , $i = 1, \dots, 15$, the exceptional divisors.

Claim 1: The linear system $L = 6H - \sum_{i=1}^{15} E_i$ is base point free and very ample.

Proof of Claim 1. We sheafify the exact sequence (8) and we get the exact sequence:

$$0 \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-6)^5 \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-5)^6 \longrightarrow \mathcal{J}_{Z(H_3)} \longrightarrow 0.$$

We compute the cohomology and we get $H^1(\mathcal{J}_{Z(H_3)}(5)) = 0$. Therefore, the linear system $L = 6H - \sum_{i=1}^{15} E_i$ is base point free. Moreover, since no line in \mathbb{P}^2 contains 6 points of $Z(H_3)$, we can apply [4, Theorem 3.1] and we conclude that L is very ample.

Since $L = 6H - \sum_{i=1}^{15} E_i$ is base point free and very ample, it defines an embedding, i.e. an isomorphism onto its image $\varphi_L : Y \rightarrow \mathbb{P}^{12}$ which lifts the rational map $\varphi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^{12}$

Therefore, we have the commutative diagram

$$\begin{array}{ccc} Y & & \\ \downarrow \pi & \searrow \varphi_L & \\ \mathbb{P}^2 & \dashrightarrow \varphi & \mathbb{P}^{12}. \end{array}$$

and the image of φ is a smooth surface $X \subset \mathbb{P}^{12}$ of degree $L^2 = 21$.

Let us now prove that X is an arithmetically Cohen-Macaulay surface. First we compute the h -vector (h_i) of the set of points $Z(H_3) \subset \mathbb{P}^2$ where

$$\begin{aligned} h_i &= \Delta H(Z(H_3), i) = H(Z(H_3), i) - H(Z(H_3), i-1) \\ &= \dim \left(\frac{k[x, y, z]}{I(Z(H_3))} \right)_i - \dim \left(\frac{k[x, y, z]}{I(Z(H_3))} \right)_{i-1}. \end{aligned}$$

The h -vector of $Z(H_3)$ is $(1, 2, 3, 4, 5)$. Therefore, $h_i = 0$ for all $i \geq 5$ and $\sigma = 5$, where σ is the least integer t such that $\Delta H(Z(H_3), t) = 0$. By [10, Theorem B], X is an arithmetically Cohen-Macaulay surface and by [10, Theorem C], $I(X)$ is generated by quadrics which proves what we want. \square

Turning to the companion surface we take a closer look at the polynomials h_1, h_2, \dots, h_{13} . First of all, it can be checked directly that h_1 is a linear combination of the remaining 12 polynomials. These polynomials, in turn, can be paired according to their fourth power factors. Removing the constant terms and taking linear combinations, we obtain the following 12 quintics in variables $(a : b : c)$:

$$\begin{aligned} q_1 &= a(b - (u-1)c)^4, & q_2 &= (b + uc)(b - (u-1)c)^4, \\ q_3 &= a(b + (u-1)c)^4, & q_4 &= (b - uc)(b + (u-1)c)^4, \\ q_5 &= b(a - uc)^4, & q_6 &= (a + (u-1)c)(a - uc)^4, \\ q_7 &= b(a + uc)^4, & q_8 &= (a - (u-1)c)(a + uc)^4, \\ q_9 &= c(a - (u-1)b)^4, & q_{10} &= (a + ub)(a - (u-1)b)^4, \\ q_{11} &= c(a + (u-1)b)^4, & q_{12} &= (a - ub)(a + (u-1)b)^4, \end{aligned}$$

Interestingly, the unexpected curve written down in these generators has the following form

$$\begin{aligned} F_{Z(H_3)} &= \frac{1}{2} \left[(3u+1) [M_1 f_1 q_1 - x f_1 q_2 - M_2 f_2 q_3 + x f_2 q_4] \right. \\ &\quad + (1-2u) [M_3 f_3 q_5 - y f_3 q_6 - M_4 f_4 q_7 + y f_4 q_8] \\ &\quad \left. + (3u+1) [M_5 f_5 q_9 - z f_5 q_{10} - M_6 f_6 q_{11} + z f_6 q_{12}] \right], \end{aligned}$$

where M_1, \dots, M_6 are Galois conjugate of lines L_1, \dots, L_6 and f_1, \dots, f_6 are defined in (7):

$$\begin{aligned} M_1 : y + uz &= 0, & M_2 : y - uz &= 0, \\ M_3 : x - \bar{u}z &= 0, & M_4 : x + \bar{u}z &= 0, \\ M_5 : x + uy &= 0, & M_6 : x - uy &= 0. \end{aligned}$$

These lines are indicated as dashed lines in Figure 3, the L -lines are indicated there as the thin lines. The dashed lines intersect in the Galois conjugate H_3 configuration of points. In particular the origin and the two points at the infinity belong to both configurations.




Figure 3: The H_3 configuration of points and the Galois conjugate lines

Remark 4.6. We describe the companion surface X' of X .

Let $\psi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^{11}$ be the rational map defined by the 12 generators q_1, \dots, q_{12} and we call X' the image of ψ . To check that the map $\psi : \mathbb{P}^2 \dashrightarrow \mathbb{P}^{11}$ has degree one and even more it is a birational map onto its image we use the packages "Cremona and birational maps" of Macaulay2 as before.

```

loadPackage "Cremona"
loadPackage "RationalMaps"
kk = ZZ/22621
R = kk[a,b,c]
u = 1873 -- golden ratio
I = ideal(a(b-(u-1)c)^4, (b+uc)(b-(u-1)c)^4, a(b+(u-1)c)^4, (b-uc)(b+(u-1)c)^4,
b(a-uc)^4, (a+(u-1)c)(a-uc)^4, b(a+uc)^4, (a-(u-1)c)(a+uc)^4, c(a-(u-1)b)^4,
(a+ub)(a-(u-1)b)^4, c(a+(u-1)b)^4, (a-ub)(a+(u-1)b)^4)
degree rationalMap gens I
(Using the package "Cremona" we get that the degree is 1)
S=kk[q_1..q_(numgens I)]
varphi = map(R, S, gens I)
isBirationalOntoImage(varphi) -- true
(Using the package "RationalMaps" we get that the map is birational onto its image)

```

Once we know that ψ is a birational map onto its image we compute the degree of X' , it is 25. The smoothness of X' follows from the Jacobian criterion and a computation which shows that the rank of $(\frac{\partial q_i}{\partial a} \quad \frac{\partial q_i}{\partial b} \quad \frac{\partial q_i}{\partial c})_{i=1, \dots, 12}$ is 3. It is a projection of the Veronese $V_{2,5}$ by a subspace of dimension eight that does not intersect the Secant variety. Moreover, the ideal of X' is generated by quadrics and cubics.

Remark 4.7. We consider the rational map $\bar{\varphi} : \mathbb{P}^2 \dashrightarrow \mathbb{P}^{11}$ where

$$\bar{\varphi} = (M_1 f_1, x f_1, M_2 f_2, x f_2, M_3 f_3, y f_3, M_4 f_4, y f_4, M_5 f_5, z f_5, M_6 f_6, z f_6)$$

and we call \bar{X} its image. Using Macaulay2 we get that $\bar{\varphi}$ is a birational map onto \bar{X} , $\text{degree}(\bar{X}) = 21$ and $I(\bar{X})$ is generated by 32 quadrics. The surface $\bar{X} \subset \mathbb{P}^{11}$ can be seen

as a projection of the surface $X \subset \mathbb{P}^{12}$ described in Proposition 4.5. However, \bar{X} is not aCM, as its h -vector is $h : (1, 9, 13, -3, 1)$, so it is not positive.

5 Companion surfaces for Fermat arrangements of lines

Recall that in suitable coordinates the points in the root system B_3 are dual to linear factors of the Fermat-type polynomial

$$F_{2,3}(x : y : z) = xyz(x^2 - y^2)(y^2 - z^2)(z^2 - x^2).$$

Thus it is natural to generalize B_3 from the perspective of Fermat-type arrangements. We follow this path of thoughts in this section. Our notation is consistent with the notation introduced in [22].

Let

$$F_{m,0}(x : y : z) = (x^m - y^m)(y^m - z^m)(z^m - x^m)$$

and let $Z(F_{m,0})$ be the set of points dual to the linear factors of $F_{m,0}$. Thus $Z(F_{m,0})$ is a set of $3m$ points distributed evenly on the coordinate lines. This is visualised in Figure 4a for $m = 5$.




Figure 4: Fermat configurations of points for $m = 5$

We recall a result [22, Theorem 4.6] which motivated our present research.

Theorem 5.1 (Szpond). *Let $m \geq 5$ be an integer and let $R = (a : b : c)$ be a general point in \mathbb{P}^2 . The set $Z(F_{m,0})$ admits a unique irreducible unexpected curve $C_{R,m}$ of degree $m + 2$ and multiplicity $m + 1$. Moreover, all these curves $C_{R,m}$ pass through the coordinate points.*

It turns out that for the augmented configuration

$$Z(F_{m,3}) = Z(F_{m,0}) \cup (1 : 0 : 0) \cup (0 : 1 : 0) \cup (0 : 0 : 1)$$

the curve $C_{R,m}$ is unexpected as soon as $m \geq 2$. For $m = 2$ we get exactly the unexpected curve associated to the B_3 root system.

Proposition 5.2. *The ideal $I(Z(F_{m,3})) \subset S = \mathbb{C}[x, y, z]$ is generated by*

$$xyz, G_x = yz(y^m - (-z)^m), G_y = zx(z^m - (-x)^m), G_z = xy(x^m - (-y)^m).$$

Even more, $I(Z(F_{m,3}))$ is defined by the maximal minors of the matrix

$$M = \begin{pmatrix} y^m - (-z)^m & z^m - (-x)^m & x^m - (-y)^m \\ x & 0 & 0 \\ 0 & y & 0 \\ 0 & 0 & z \end{pmatrix}$$

and, hence, it has a minimal free resolution of the following type:

$$0 \longrightarrow S(-m-3)^3 \xrightarrow{M} S(-3) \oplus S(-m-2)^3 \longrightarrow I(Z(F_{m,3})) \longrightarrow 0.$$

Proof. First we note that the reason for the sign in the three binomial generators depending on the parity of m is due to the convention that a point $(p : q : r)$ is dual to the line $px + qy + rz = 0$. Thus, for example, if $m = 3$ and ε is a primitive root of the unity of order 3, then

$$(x - \varepsilon y)(x - \varepsilon^2 y)(x - y) = x^3 - y^3,$$

and the dual points are

$$(1 : -\varepsilon : 0), (1 : -\varepsilon^2 : 0), (1 : -1 : 0).$$

Their ideal is generated by $(z, x^3 + y^3)$.

It is now easy to see that for arbitrary m the trace of $Z(F_{m,0})$ on the coordinate line, say $x = 0$, is a set of zeroes of a complete intersection ideal generated by

$$x \text{ and } yz(y^m + (-1)^{m+1}z^m)$$

and similarly on the other lines. Thus

$$\begin{aligned} I(Z(F_{m,3})) &= (xy, yz, zx) \cap (x, yz(y^m + (-1)^{m+1}z^m)) \\ &\cap (y, xz(z^m + (-1)^{m+1}x^m)) \cap (z, xy(x^m + (-1)^{m+1}y^m)). \end{aligned}$$

and the claim follows readily. \square

We are interested in homogeneous elements in $I(Z(F_{m,3}))$ of degree $(m+2)$. They are G_x, G_y, G_z and monomials of the form

$$xyz \cdot \text{monomial of degree } m-1.$$

Hence

$$\dim[I(Z(F_{m,3}))]_{m+2} = \frac{1}{2}m^2 + \frac{1}{2}m + 3 \quad (9)$$

Geometrically the elements in $[I(Z(F_{m,3}))]_{m+2}$ correspond to elements of the linear system of forms of degree $m+2$ vanishing at all points of the set $Z(F_{m,3})$.

5.1 Positivity on anticanonical surfaces

Before we continue the study of the linear system determined by $[I(Z(F_{m,3}))]_{m+2}$, we need to recall some results on anticanonical systems on surfaces. Results presented here were obtained by Harbourne in [16] and [15]. We begin by the following useful Lemma, which is a combination of Lemma II.2 and Corollary II.3 in [15] and Lemma 2.2 and Corollary 2.3 in [16].

Lemma 5.3 (Harbourne). *Let Y be a smooth rational surface and let D be a nef class on Y . Then*

$$h^2(Y, D) = 0 \quad \text{and} \quad D^2 \geq 0.$$

Moreover, if $-K_Y$ is effective, then so is D .

Assuming the effectivity of the anticanonical class, one can say in fact much more. The following statement is extracted from a much more precise (and complicated) result of Harbourne [15, Theorem III.1] and [16, Theorem 2.11].

Proposition 5.4 (Harbourne). *Let Y be a smooth anticanonical surface and let D be a nef class on Y such that $-K_Y.D \geq 2$. Then D is non-special, i.e., $h^1(Y, D) = 0$ and the linear system $|D|$ is base point free.*

Thus the anticanonical degree at least 2 implies that $|D|$ defines a morphism. If this degree is at least 3, then the positivity of $|D|$ increases, see [16, Proposition 3.2].

Proposition 5.5 (Harbourne). *Let Y be a smooth anticanonical surface and let D be a nef class on Y such that $-K_Y.D \geq 3$ and $D^2 > 0$. Then the morphism φ_D defined by the elements of $|D|$ is birational and its image Y' is a normal surface obtained by contracting all curves perpendicular to $|D|$. Moreover a general member of $|D|$ is smooth and irreducible.*

5.2 On the positivity of maps determined Fermat arrangements

Let m be fixed and let ε be a primitive root of the unity of order m . It is convenient to introduce the following notation:

$$P_{x,\alpha} = (0 : 1 : \varepsilon^\alpha), \quad P_{y,\alpha} = (1 : 0 : \varepsilon^\alpha), \quad P_{z,\alpha} = (1 : \varepsilon^\alpha : 0)$$

for $\alpha = 0, \dots, m-1$ and

$$P_{x,y} = (0 : 0 : 1), \quad P_{x,z} = (0 : 1 : 0), \quad P_{y,z} = (1 : 0 : 0).$$

Our next result concerns the positivity of the linear system corresponding to $[I(Z(F_{m,3}))]_{m+2}$.

Theorem 5.6. *Let $f : Y \rightarrow \mathbb{P}^2$ be the blow up at the points of the set $Z(F_{m,3})$ with the exceptional divisors $E_{a,\alpha}$, $E_{a,b}$ over the points $P_{a,\alpha}$ and $P_{a,b}$ respectively, where $a, b \in \{x, y, z\}$ and $\alpha \in \{0, \dots, m-1\}$. Let $H = f^*\mathcal{O}_{\mathbb{P}^2}(1)$ be the pull-back of the hyperplane bundle and let \mathbb{E} be the union of all exceptional divisors of f . Then the linear system*

$$M = (m+2)H - \mathbb{E}$$

is base point free and defines a morphism φ_M birational onto its image

$$\varphi_M : Y \rightarrow X \subset \mathbb{P}^N, \quad (N = \frac{1}{2}(m^2 + m) + 2),$$

which is an isomorphism away of the proper transforms of the coordinate lines.

Proof. Let

$$\begin{aligned} N_x &= H - E_{x,y} - E_{x,z} - E_{x,0} - \dots - E_{x,m-1} \\ N_y &= H - E_{x,y} - E_{y,z} - E_{y,0} - \dots - E_{y,m-1} \\ N_z &= H - E_{x,z} - E_{y,z} - E_{z,0} - \dots - E_{z,m-1} \end{aligned}$$

be the proper transformations of the coordinate lines.

Our first claim is that M is nef and the only curves having the intersection number with M equal zero are N_x , N_y and N_z .

It is obvious that $M.N_a = 0$ for all $a \in \{x, y, z\}$, so it is enough to check that M has positive intersection with all other irreducible curves on Y . Certainly it is $M.E = 1$ for all exceptional divisors of f . Let C be an irreducible plane curve of degree d , different from

the coordinate lines, passing through the points $P_{a,\alpha}$ and $P_{a,b}$ with multiplicities $m_{a,\alpha}$ and $m_{a,b}$ respectively. Let

$$\tilde{C} = dH - \sum m_{a,\alpha} E_{a,\alpha} - \sum m_{a,b} E_{a,b}$$

be the proper transform of C . Then

$$0 \leq \tilde{C}.N_x = d - m_{x,y} - m_{x,z} - m_{x,0} - \dots - m_{x,m-1}, \quad (10)$$

$$0 \leq \tilde{C}.N_y = d - m_{x,y} - m_{y,z} - m_{y,0} - \dots - m_{y,m-1}, \quad (11)$$

$$0 \leq \tilde{C}.N_z = d - m_{x,z} - m_{y,z} - m_{z,0} - \dots - m_{z,m-1}. \quad (12)$$

Adding the inequalities (10), (11) and (12) and rearranging terms we get

$$3d - 2(m_{x,y} + m_{x,z} + m_{y,z}) - \sum m_{a,\alpha} \geq 0.$$

Since

$$M.\tilde{C} = 4d - m_{x,y} - m_{x,z} - m_{y,z} - \sum m_{a,\alpha}$$

we conclude that $M.\tilde{C} > 0$ as asserted.

For the rest of the proof, the key observation is that the surface Y is an anticanonical surface, i.e., $-K_Y$ is an effective divisor. Indeed

$$-K_Y = N_x + N_y + N_z + E_{x,y} + E_{x,z} + E_{y,z}.$$

Since

$$-K_Y.M = 3$$

Proposition 5.4 implies that $|M|$ is non-special and base point free. Revoking Lemma 5.3 we have thus the vanishing

$$h^1(Y, M) = h^2(Y, D) = 0,$$

which together with the Riemann-Roch formula on Y reconfirms the computation in (9). Since

$$M^2 = m^2 + m + 1 > 0,$$

it follows from Proposition 5.5 that the morphism defined by $|M|$ is birational and its image is a normal surface of degree $m^2 + m + 1$ in \mathbb{P}^N with $N = \frac{1}{2}(m^2 + m) + 2$ singular in exactly 3 points, which are images of the proper transforms of the coordinate lines in \mathbb{P}^2 . \square

We now turn to the equations defining X . To this end let us be a little bit more specific about the coordinates of the rational map φ determined by the commutative diagram

$$\begin{array}{ccc} Y & & \\ \downarrow f & \searrow \varphi_M & \\ \mathbb{P}^2 & \dashrightarrow \varphi & X \subset \mathbb{P}^N, \end{array}$$

which is not just incidentally similar to that in Proposition 4.3. We can assume that

$$\varphi = (xyz\mu_d : G_x : G_y : G_z),$$

where $d = m - 1$ and

$$\mu_d : \mathbb{P}^2 \hookrightarrow \mathbb{P}^{N-3}$$

denotes the Veronese embedding of degree d and the rational map $xyz\mu_d$ is given by coordinates of μ_d multiplied by the monomial xyz .

It is well known, see for example [23, Corollary 7.2.3], that ideal of the image V_d of μ_d is generated by $\frac{(d-1)d(d+1)(d+6)}{8}$ quadratic binomials.

Consider the projection $\pi : \mathbb{P}^N \dashrightarrow \mathbb{P}^{N-3}$ determined by the plane Π spanned by the last three coordinate points in \mathbb{P}^N . Then $\pi(X) = V_d$ and the mapping φ (or equivalently φ_M) can be considered as the ‘‘unprojection’’ of the Veronese surface in \mathbb{P}^{N-3} . We use the word unprojection to indicate a morphism inverse to a projection, not quite in the sense as introduced to the birational geometry by Miles Reid.

Proposition 5.7. *Let $m \geq 4$ be a fixed integer and let $I = I(X)$ be the ideal of $X \subset \mathbb{P}^N$. It holds:*

- (1) $X \subset \mathbb{P}^N$ is an aCM surface.
- (2) The ideal I is generated by forms of degree ≤ 3 . More precisely, I is generated by $3\binom{m+2}{4}$ quadrics and one cubic C_m , where for m odd

$$C_m : 2H_xH_yH_z + G_xH_yH_z + H_xG_yH_z + H_xH_yG_z - G_xG_yG_z,$$

and for m even

$$C_m : G_xH_yH_z + H_xG_yH_z + H_xH_yG_z - G_xG_yG_z,$$

where $H_x = x^m y z$, $H_y = x y^m z$ and $H_z = x y z^m$.

Proof. (1) Let $L \cong \mathbb{P}^{N-1} \subset \mathbb{P}^N$ be a general hyperplane. Since $X \subset \mathbb{P}^N$ is an aCM surface if and only if a general hyperplane section $C = X \cap L$ of X is an aCM curve, it is enough to check that indeed $C \subset L = \mathbb{P}^{N-1}$ is an aCM curve. We observe that

$$\deg(C) = \deg(X) = m^2 + m + 1.$$

By the adjunction formula we have:

$$2g(C) - 2 = C(C + K_X) = C^2 - 3$$

(because $CK_X = MK_X = ((m+2)H - \sum_{i=1}^{3m+3} E_i)(-3H + \sum_{i=1}^{3m+3} E_i) = -3$). Therefore, $C^2 = 2g(C) + 1$. Thus, $\mathcal{O}_C(C)$ is a very ample line bundle on C and by the Corollary to [18, Theorem 6], $\mathcal{O}_C(C)$ is projectively normal. So, $\mathcal{O}_C(C)$ embeds C in a projective space as a projectively normal curve. Since X is rational, it is regular (i.e. $H^1(X, \mathcal{O}_X) = 0$) and from the exact cohomology sequence associated to the exact sequence

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_X(C) \longrightarrow \mathcal{O}_C(C) \longrightarrow 0$$

we deduce that the projective space into which $\mathcal{O}_C(C)$ embeds C is just L and we are done.

(2) Let us first compute the dimension of $I(X)_2$. By [23, Corollary 7.2.3] there are $\frac{(m-2)(m-1)m(m+5)}{8}$ quadrics involving the monomials $xyz(x, y, z)^{m-1}$. A straightforward computation shows that there are $\binom{m}{2}$ quadrics involving G_x (resp., G_y and G_z) and the monomials in $xyz(x, y, z)^{m-1}$. Therefore $\dim I(X)_2 = 3\binom{m+2}{4}$. To prove that $I(X)$ is generated by quadrics and cubics, we consider the h -vector of X . Recall that the h -vector of X is a sequence $(h_i)_{i \geq 0}$ of non-negative integers satisfying:

- (i) For all $i \geq 0$, $h_i = \Delta^3 H(X, i)$ where $H(X, i) = \dim(\mathbb{C}[y_0, \dots, y_N]/I(X))_i$.
- (ii) $\deg(X) = \sum_{i \geq 0} h_i$.

(iii) If σ is the least integer such that $h_i = 0$, then $I(X)$ is generated by forms of degree $\leq \sigma$.

Since X is non-degenerate and $\dim I(X)_2 = 3\binom{m+2}{4}$, we have $h_0 = 1$, $h_1 = h_2 = N - 2$. From the equality

$$\deg(X) = m^2 + m + 1 = \sum_{i \geq 0} h_i = 2N - 3 + \sum_{i \geq 3} h_i,$$

we conclude that $h_i = 0$ for $i \geq 3$, $\sigma = 3$ and $I(X)$ is generated by forms of degree ≤ 3 . Let us now prove that $I(X)$ is not generated by quadrics i.e. at least one cubic is required. Since X is an aCM surface, we know that the graded Betti numbers of X and of a general hyperplane section $C = X \cap \mathbb{P}^{N-1}$ are equal; in particular, the degrees of a minimal set of generators of C and X coincide. So, it will be enough to check that a minimal set of generators of $I(C)$ contains a cubic. In [13, Theorem 2], Green and Lazarsfeld proved that $I(C)$ fails to be generated by quadrics if and only if C is hyperelliptic or $\mathcal{O}_C(C)$ embeds C with a trisecant line, i.e. $H^0(C; C - K_C) \neq 0$ where K_C denotes the canonical divisor of C . We consider the exact sequence

$$0 \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_X(C) \longrightarrow \mathcal{O}_C(C) \longrightarrow 0$$

and its associated exact cohomology sequence:

$$0 \longrightarrow H^0(X; -C - K_X) \longrightarrow H^0(X; -K_X) \longrightarrow H^0(C; \mathcal{O}_C(-K_X)) \longrightarrow H^1(X; -C - K_X) \cdots$$

From the adjunction formula $K_C \cong \mathcal{O}_C(C + K_X)$, we deduce that $H^0(C; \mathcal{O}_C(-K_X)) \cong H^0(C; \mathcal{O}_C(C - K_C))$. On the other hand, $H^0(X; -C - K_X) \cong H^0(X; -M - K_X) \cong H^0(X; -(m-1)H) = 0$ and $H^1(X; -C - K_X) \cong H^1(X; -M - K_X) \cong H^1(X; M) = 0$ (since $|M|$ is non-special). Therefore, we have

$$H^0(C; \mathcal{O}_C(C - K_C)) \cong H^0(X; -K_X) \cong H^0\left(X; 3H - \sum_{i=1}^{3m+3} E_i\right) \cong \mathbb{C} \quad (13)$$

and we are done.

Let us prove that C_m is part of a minimal system of generators of $I(X)$. To this end, we first describe $I(X)_2$. Recall that X is the image of the rational map

$$\begin{aligned} \varphi : \mathbb{P}^2 &\dashrightarrow \mathbb{P}^N \\ (x : y : z) &\mapsto (m_0 : m_1 : \dots : m_{N-3} : G_x : G_y : G_z), \end{aligned}$$

where $m_i = x^{i_0} y^{i_1} z^{i_2}$ with $i_0 + i_1 + i_2 = m + 2$ and $i_0, i_1, i_2 \geq 1$. Fix homogeneous coordinates $\omega_0, \dots, \omega_N$ in \mathbb{P}^N . $I(X)_2$ is generated by

- All $\frac{(m-2)(m-1)m(m+5)}{8}$ quadrics $\omega_i \omega_j - \omega_k \omega_l$, $0 \leq i, j, k, l \leq N - 3$ satisfying $i_u + j_u = k_u + l_u$ for $u = 0, 1, 2$.
- All $\binom{m}{2}$ quadrics $\omega_i H_y - (-1)^m \omega_k H_z - \omega_s \omega_{N-2}$ (Analogous quadrics with ω_{N-1} and ω_N) where $\omega_s = x^{s_0} y^{s_1} z^{s_2}$, $\omega_i = x^{s_0-1} y^{s_1+1} z^{s_2}$ and $\omega_k = x^{s_0-1} y^{s_1} z^{s_2+1}$ with $s_0 + s_1 + s_2 = m + 2$, $2 \leq s_0$ and $1 \leq s_1, s_2$.

Note that all quadrics in $I(X)_2$ vanish on the plane Π : $\omega_0 = \dots = \omega_{N-3} = 0$ while C_m does not vanish on Π . Therefore C_m is not a linear combination of the quadrics in $I(X)_2$.

The existence of a trisecant line ℓ for C is now clear. If $C = X \cap \mathbb{P}^{N-1}$ then the trisecant line to C is the line $\mathbb{P}^{N-1} \cap \Pi$.

It remains to see that there is only one cubic in a minimal system of generators of $I(C)$ but this follows from the fact that $H^0(C; \mathcal{O}_C(C - K_C)) = \mathbb{C}$ i.e. the span of the divisor $\mathcal{O}_C(C - K_C)$ is the unique trisecant line. Indeed, assume that C has a unique trisecant line ℓ and call C_1 the union of C and ℓ . Then C_1 is a canonical singular curve and its ideal is generated by quadrics (Petri's theorem) which are the quadrics in the ideal of $I(C)$ (In fact, $h^0(C; \mathcal{O}_C(2)) = h^0(C_1; \mathcal{O}_{C_1}(2))$ by Riemann-Roch theorem for singular curves (see [9, Example 18.3.4]). Hence, to cut out C from C_1 , it suffices to consider one cubic in the ideal of $I(C)$ which is not a linear combination of quadrics. \square

Remark 5.8. (1) We note that the unexpected curve $C_{R,m}$ in [22, Theorem 4.6] can be written

$$C_{R,m} = xyz \left[(m+1) \left(a(b^m + (-1)^{m+1}c^m)x^{m-1} \right. \right. \\ \left. \left. + b(a^m + (-1)^{m+1}c^m)y^{m-1} + c(a^m + (-1)^{m+1}b^m)z^{m-1} \right) \right. \\ \left. + \sum_{i=1}^{m-2} (-1)^{i+1} \binom{m+1}{i+1} \left(a^{m-i}b^{i+1}x^i y^{m-i-1} + a^{m-i}c^{i+1}x^i z^{m-i-1} + b^{m-i}c^{i+1}y^i z^{m-i-1} \right) \right] \\ + a^{m+1}G_x + b^{m+1}G_y + c^{m+1}G_z$$

and this form does not involve monomials that are product of $(xyz)^2$ and a monomial of degree $m-4$. So, let J be the ideal generated by all coordinates ω_i corresponding to the monomials in $(xyz)^2\mu_{m-4}$, we can consider the projection π_J and consider the map $\bar{\varphi}_m = \pi_J \circ \varphi : \mathbb{P}^2 \rightarrow \mathbb{P}^{3m-1}$. In this way the ideal of the image $\bar{X} = \pi_J(X)$ of this map is the elimination ideal of $I(X)$ and so also $I(\bar{X})$ is generated by quadrics and a unique cubic which turns out to be the same as in Proposition 5.7. With a simple calculation we get that the number of quadrics decreases to $\frac{1}{2}(5m^2 - m - 12)$.

Moreover, \bar{X} is the result of subsequent projections from external points. So, the degree is preserved and $\deg(\bar{X}) = \deg(X) = m^2 + m + 1$.

Now we recall that $\deg(\bar{X}) = \sum_i h_i$ and, as we know the Hilbert function in degree 1 and 2, we can compute h_1 and h_2 so we get that $\sum_{i \geq 3} h_i = -(m-2)(m-3) < 0$ for any $m \geq 4$. So we deduce that the h -vector is not positive and \bar{X} is not aCM. Actually, computations with Macaulay2 supports the following conjecture:

$$h : (1, 3(m-1), 2m^2 - 7m + 9, -\frac{3}{2}(m-2)(m-3), \frac{1}{2}(m-2)(m-3)).$$

(2) In Proposition 5.7 we have seen that for all $m \geq 4$ the image of the map $\varphi : \mathbb{P}^2 \rightarrow \mathbb{P}^N$ where

$$\varphi = (xyz\mu_d : G_x : G_y : G_z)$$

is a smooth surface X of degree $m^2 + m + 1$, its ideal $I(X)$ is generated by quadrics and only one cubic and it is aCM. Using Macaulay2 we have checked what happens with their companion surfaces $X' \subset \mathbb{P}^{3m-1}$. Recall that X' is the image of $\psi : \mathbb{P}^2 \rightarrow \mathbb{P}^{3m-1}$ where

$$\psi = (a^{m+1}, b^{m+1}, c^{m+1}, a^{m-1}b^2, \dots, a^2b^{m-1}, a^{m-1}c^2, \dots, a^2c^{m-1}, b^{m-1}c^2, \dots, b^2c^{m-1}, \\ a(b^m - (-c)^m), b(a^m - (-c)^m), c(b^m - (-a)^m).$$

Our computation, for $m \leq 8$, strongly supports the following facts: ψ is a birational map onto its image, the ideal of X' is generated by $\frac{1}{2}(5m^2 - 5m - 12)$ quadrics and a unique cubic (unless for $m = 4$ where 10 independent cubics are necessary). Moreover, the companion

has degree $(m+1)^2$ and with the same arguments as for \bar{X} we conjecture that the h -vector is

$$h : (1, 3(m-1), 2m^2 - 5m + 9, -\frac{3}{2}(m-2)(m-3), \frac{1}{2}(m-1)(m-6)).$$

Therefore, X' is not an aCM surface. Indeed, the projective dimension of the coordinate ring of X' is equal to $\text{codim}(X') + 1$.

Summarizing, for $m \leq 8$, we have that $X \subset \mathbb{P}^N$ is an aCM projection of the Veronese surface $V_{2,m+2}$ in $\mathbb{P}^{\binom{m+4}{2}-1}$ while its companion surface X' is a non-aCM projection of the Veronese surface $V_{2,m+2}$ in \mathbb{P}^{3m-1} . This example makes more intriguing Gröbner's problem: To determine when a projection of a Veronese variety $V_{n,d} \subset \mathbb{P}^{\binom{n+d}{d}-1}$ is aCM and when it is not aCM (see [14]).

Remark 5.9. If we consider the ideal generated only by the quadrics in $I(\bar{X})$ the degree increases by 1. Instead if we consider the ideal generated only by the quadrics in $I(X')$, the degree is still the $(m+1)^2$.

Finally, we remark explicitly, that the embedding of \bar{X} is never too positive.

Corollary 5.10. *The surface $X \subset \mathbb{P}^N$ does not satisfy the N_2 condition of Green.*

Acknowledgement. Our work began during Oberwolfach workshop on Lefschetz Properties in Algebra, Geometry and Combinatorics held in September/October 2020. We thank the MFO for providing, during the COVID-19 crisis, excellent working conditions for collaboration between on site and remote participants of the workshop.

The first two authors have been partially supported by GNSAGA-INDAM. The third author has been partially supported by MTM201678623-P. The last two authors were partially supported by National Science Centre, Poland, Opus Grant 2019/35/B/ST1/00723. Our collaboration was partially supported by National Science Centre, Poland, Harmonia Grant 2018/30/M/ST1/00148.

References

- [1] T. Bauer, G. Malara, T. Szemberg, and J. Szpond. Quartic unexpected curves and surfaces. *Manuscripta Math.*, 161(3-4):283–292, 2020.
- [2] L. Chiantini and J. Migliore. Sets of points which project to complete intersections. *Trans. AMS*, to appear. With an appendix by the authors and A. Bernardi, G. Denham, G. Favacchio, B. Harbourne, T. Szemberg and J. Szpond.
- [3] D. Cook II, B. Harbourne, J. Migliore, and U. Nagel. Line arrangements and configurations of points with an unexpected geometric property. *Compos. Math.*, 154(10):2150–2194, 2018.
- [4] E. Davis and A. Geramita. Birational morphisms to \mathbb{P}^2 : An ideal-theoretic perspective. *Math. Ann.*, 279:435–448, 1988.
- [5] R. Di Gennaro, G. Ilardi, and J. Vallès. Singular hypersurfaces characterizing the Lefschetz properties. *J. Lond. Math. Soc. (2)*, 89(1):194–212, 2014.
- [6] A. V. Doria, S. H. Hassanzadeh, and A. Simis. A characteristic-free criterion of birationality. *Adv. Math.*, 230(1):390–413, 2012.

- [7] M. Dumnicki, L. Farnik, B. Harbourne, G. Malara, J. Szpond, and H. Tutaj-Gasińska. A matrixwise approach to unexpected hypersurfaces. *Linear Algebra Appl.*, 592:113–133, 2020.
- [8] M. Dumnicki, A. Küronya, C. Maclean, and T. Szemberg. Rationality of Seshadri constants and the Segre-Harbourne-Gimigliano-Hirschowitz conjecture. *Adv. Math.*, 303:1162–1170, 2016.
- [9] W. Fulton. *Intersection theory*, volume 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, second edition, 1998.
- [10] A. Geramita and A. Gimigliano. Generators for the defining ideal of certain rational surfaces. *Duke Math. J.*, 62:61–83, 1991.
- [11] A. V. Geramita, B. Harbourne, and J. Migliore. Star configurations in \mathbb{P}^n . *J. Algebra*, 376:279–299, 2013.
- [12] D. Grayson and M. Stillman. Macaulay2, a software system for research in algebraic geometry. <http://www.math.uiuc.edu/Macaulay2/>, 2020.
- [13] M. Green and R. Lazarsfeld. Some results on the syzygies of finite sets and algebraic curves. *Compositio Math.*, 67(3):301–314, 1988.
- [14] W. Gröbner. Über Veronesesche Varietäten und deren Projektionen. *Arch. Math. (Basel)*, 16:257–264, 1965.
- [15] B. Harbourne. Anticanonical rational surfaces. *Trans. Amer. Math. Soc.*, 349(3):1191–1208, 1997.
- [16] B. Harbourne. Birational morphisms of rational surfaces. *J. Algebra*, 190(1):145–162, 1997.
- [17] B. Harbourne, J. Migliore, U. Nagel, and Z. Teitler. Unexpected hypersurfaces and where to find them. *Michigan Math. J.*, to appear.
- [18] D. Mumford. Varieties defined by quadratic equations. In *Questions on Algebraic Varieties (C.I.M.E., III Ciclo, Varenna, 1969)*, pages 29–100. Edizioni Cremonese, Rome, 1970.
- [19] M. Nagata. On the 14-th problem of Hilbert. *Amer. J. Math.*, 81:766–772, 1959.
- [20] F. Russo and A. Simis. On birational maps and jacobian matrices. *Compositio Mathematica*, 126:335–358, 2001.
- [21] J. Szpond. Unexpected curves and Togliatti-type surfaces. *Math. Nachr.*, 293:158–168, 2020.
- [22] J. Szpond. Fermat-type arrangements, In: Stamate D., Szemberg T. (eds) *Combinatorial Structures in Algebra and Geometry*. NSA 2018. Springer Proceedings in Mathematics & Statistics, vol 331. p. 161 – 182, Springer, Cham, 2020.
- [23] J. Weyman. *Cohomology of vector bundles and syzygies*, volume 149 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2003.

Roberta Di Gennaro,
Dipartimento di Matematica e Applicazioni “Renato Caccioppoli”
Università degli Studi di Napoli Federico II
Via Cintia, Complesso Universitario Monte Snt’Angelo
80126 Napoli, Italy.
E-mail address: digennar@unina.it

Giovanna Ilardi,
Dipartimento di Matematica e Applicazioni “Renato Caccioppoli”
Università degli Studi di Napoli Federico II
Via Cintia, Complesso Universitario Monte Snt’Angelo
80126 Napoli, Italy.
E-mail address: giovanna.ilardi@unina.it

Rosa Maria Miró Roig,
Facultat de Matemàtiques i Informàtica, Universitat de Barcelona, Gran Via des les Corts
Catalanes 585, 08007 Barcelona, Spain.
E-mail address: miro@ub.edu

Tomasz Szemberg,
Department of Mathematics, Pedagogical University of Cracow, Podchorążych 2, PL-30-
084 Kraków, Poland.
E-mail address: tomasz.szemberg@gmail.com

Justyna Szpond,
Institute of Mathematics, Polish Academy of Sciences, Śniadeckich 8, PL-00-656 Warszawa,
Poland.
E-mail address: szpond@gmail.com