

SOME RESULTS ON HIGHER ORDER ISOSYMMETRIES IN SEMI-HILBERTIAN SPACES

RCHID RABAOUI

ABSTRACT. In this paper, we introduce the class of $(A, (m, n))$ -isosymmetric operators and we study some of their properties, for a positive semi-definite operator A and $m, n \in \mathbb{N}$, which extend, by changing the initial inner product with the semi-inner product induced by A , the well-known class of (m, n) -isosymmetric operators introduced by Mark Stankus ([18], [19]). In particular, we characterize a family of A -isosymmetric (2×2) upper triangular operator matrices. Moreover, we show that if T is $(A, (m, n))$ -isosymmetric and if Q is a nilpotent operator of order r doubly commuting with T , then T^p is $(A, (m, n))$ -isosymmetric for any $p \in \mathbb{N}$ and $(T + Q)$ is $(A, (m + 2r - 2, n + 2r - 1))$ -isosymmetric. Some properties of the spectrum are also investigated.

1. INTRODUCTION AND PRELIMINARIES

Let $\mathcal{B}(H)$ be the algebra of all bounded linear operators on a separable complex Hilbert space H with the inner product $\langle \cdot | \cdot \rangle$. For $T \in \mathcal{B}(H)$, we write $\mathcal{R}(T)$, $\mathcal{N}(T)$, $\sigma(T)$, $\sigma_p(T)$ and $\sigma_{ap}(T)$ for the range, the null space, the spectrum, the point spectrum and the approximate point spectrum of T , respectively. A nonzero positif operator A defines a semi-norm $\|u\|_A := \langle Au | u \rangle^{\frac{1}{2}}$. Observe that $\|\cdot\|_A$ is a norm if and only if A is injective. For $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$, we consider the following symbols (for more details see again [8] and [15])

$$\begin{aligned} (\{(yx - 1)^m\}_{\mathbf{a}})(T, A) &:= \left(\sum_{k=0}^m (-1)^{m-k} \binom{m}{k} y^k \mathbf{a} x^k \right) (T, A), \\ (\{(x - y)^n\}_{\mathbf{a}})(T, A) &:= \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} y^k \mathbf{a} x^{n-k} \right) (T, A), \\ (\{(x + y)^n\}_{\mathbf{a}})(T, A) &= i^n (\{(x - y)^n\}_{\mathbf{a}})(iT, A). \end{aligned}$$

In 1995, J. Agler and M. Stankus ([1], [2], [3]) introduced the class of m -isometric operators and showed many properties of such a family. In fact, $T \in \mathcal{B}(H)$ is said to be m -isometric if

$$\mathcal{I}_T^m(T) := (\{(yx - 1)^m\}_{\mathbf{a}})(T, I) = 0, \quad \mathcal{I}_T^0(T) = I.$$

Sid Ahmed et al. ([20]) generalized the concept of those operators on a Hilbert space. They introduced the (A, m) -isometric operators in a semi-hilbertian space. For $m \geq 1$, an operator $T \in \mathcal{B}(H)$ is an (A, m) -isometry if

$$(1.1) \quad \mathcal{I}_A^m(T) := (\{(yx - 1)^m\}_{\mathbf{a}})(T, A) = 0, \quad \mathcal{I}_A^0(T) = A.$$

They extend well known properties and showed important results related to such a family ([8], [12]). The dynamic of the orbits of (A, m) -isometries was studied in [4] and [17]. We have shown (see [13]) that most of those properties hold true for (A, n) -symmetries (resp. skew (A, n) -symmetries). An operator T is (A, n) -symmetric (resp. skew (A, n) -symmetric) if

$$(1.2) \quad \begin{aligned} \mathcal{S}_A^n(T) &:= (\{(y - x)^n\}_{\mathbf{a}})(T, A) = 0, \quad \mathcal{S}_A^{(0)}(T) = A \\ (\text{resp. } \zeta_A^n(T) &:= (\{(y + x)^n\}_{\mathbf{a}})(T, A) = 0, \quad \zeta_A^0(T) = A). \end{aligned}$$

2010 *Mathematics Subject Classification.* 47A55, 47B25, 47B37, 47B65, 46C05.

Key words and phrases. : Semi-Hilbert space, Isosymmetric operators, $(A; (m, n))$ -isosymmetric operators, (A, m) -isometric operators, (A, m) -symmetric operators, spectrum, nilpotent perturbation.

Note that (A, n) -symmetries seems to be a natural generalization of n -symmetries, which satisfy an identity of the form $\mathcal{S}_I^n(T) = 0$, equivalently, $T \in \text{Helton}_m(T^*)$. The study of such a family was initiated by Helton in 1970 (see [11]).

In [18], M. Stankus introduced and intensively studied isosymmetries, namely bounded linear operators satisfying

$$(yx - 1)(y - x)(T) = T^{*2}T - T^*T^2 - T^* + T = 0.$$

In quest of generality, the author in [19] extends this definition to higher order, and introduces a class of operators termed (m, n) -isosymmetries. Such a family of bounded operators satisfies an identity of the form

$$(yx - 1)^m(y - x)^n(T) = 0.$$

Note that this class includes n -symmetries (i.e. T such that $(y - x)^n(T) = 0$), m -isometries (i.e. T such that $(yx - 1)^m(T) = 0$) and some classes of non-normal operators (see [18]).

Recently, M. Chō et al. ([6], [7]) introduced (m, n, C) -isosymmetries: an operator $T \in \mathcal{B}(H)$ is said to be (m, n, C) -isosymmetric if there exists a conjugation C on H such that $\gamma_{m,n}(T, C) = 0$, where

$$\gamma_{m,n}(T, C) := \begin{cases} \sum_{j=0}^m (-1)^j \binom{m}{j} T^{*m-j} \alpha_n(T, C) C T^{m-j} C, \\ \sum_{k=0}^n (-1)^k \binom{n}{k} T^{*n-k} \beta_m(T, C) C T^k C. \end{cases}$$

Recall that a conjugation on H is an antilinear operator $C : H \rightarrow H$ which satisfies $\langle Cx | Cy \rangle = \langle y | x \rangle$ for all $x, y \in H$ and $C^2 = I$.

In the light of (m, n) -isosymmetric and (m, n, C) -isosymmetric operators, using the identities (1.1) and (1.2), we aim in this paper to introduce the higher order isosymmetries in semi-Hilbertian spaces and to initiate this study by giving some of their basic properties and results. For $A \in \mathcal{B}(H)^+$, $T \in \mathcal{B}(H)$ and $m, n \in \mathbb{N}$, set

$$\Omega_A^{m,n}(T) := \begin{cases} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^n(T) T^j, \\ \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} T^{*k} \mathcal{I}_A^m(T) T^{n-k}, \end{cases}, \Lambda_A^{m,n}(T) := \begin{cases} \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} T^{*k} \zeta_A^n(T) T^k, \\ \sum_{k=0}^n \binom{n}{k} T^{*k} \mathcal{I}_A^m(T) T^{n-k}. \end{cases}$$

We define $(A; (m, n))$ -isosymmetric and skew $(A; (m, n))$ -isosymmetric operators as follows:

- (1) T is said to be $(A; (m, n))$ -isosymmetric if $\Omega_A^{m,n}(T) = 0$. Let $(A; (m, n))$ denote the set of $(A; (m, n))$ -isosymmetries. Then $(A; (m, 0))$ is the set of (A, m) -isometries and $(A; (0, n))$ is the set of (A, n) -symmetries.
- (2) T is said to be skew $(A; (m, n))$ -isosymmetric if $\Lambda_A^{m,n}(T) = 0$.

Remark 1.1.

- (1) Recall by definition

$$T \in (A; (m, n)) \iff \mathcal{S}_A^n(T) \in (I; (m, 0)) \iff \mathcal{I}_A^m(T) \in (I; (0, n)).$$

Note that:

- (a) If $\mathcal{S}_A^n(T) \geq 0$, then $T \in (A; (m, n))$ if and only if $T \in (\mathcal{S}_A^n(T); (m, 0))$.
- (b) If $\mathcal{I}_A^m(T) \geq 0$, then $T \in (A; (m, n))$ if and only if $T \in (\mathcal{I}_A^m(T); (0, n))$.
- (2) An $(A; (1, 1))$ -isosymmetry is an A -isosymmetry, and if $A = I$ then an $(A; (m, n))$ -isosymmetry is an (m, n) -isosymmetry.
- (3) If $T \in (A; (m, 0))$, then $T \in (A; (m, n'))$ for all integer $n' \geq 0$.
- (4) If $T \in (A; (0, n))$, then $T \in (A; (m', n))$ for all integer $m' \geq 0$.

- (5) If $S \in \mathcal{B}(H)$ is unitarily equivalent to T , then $S \in (A; (m, n))$ if and only if $T \in (VAV^*; (m, n))$, with V is unitary. Indeed, since T and S are unitary equivalent, there exists $V \in \mathcal{B}(H)$ unitary with $\mathcal{S}_A^n(S) = V^* \mathcal{S}_{VAV^*}^n(T)V$. It holds

$$\Omega_A^{m,n}(S) = V^*(\Omega_{VAV^*}^{m,n}(T))V.$$

- (6) From the following identity

$$(\{(x+y)^n\}_{\mathbf{a}})(T, A) = i^n(\{(x-y)^n\}_{\mathbf{a}})(iT, A)$$

it can be seen that T is skew $(A; (m, n))$ -isosymmetric if and only if $iT \in (A; (m, n))$. So, several parallel results on skew $(A; (m, n))$ -isosymmetries follow from the results of $(A; (m, n))$ -isosymmetries.

- (7) The class of $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric) operators is closed in norm, that is: let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$, if $(T_p)_p$ is a sequence of $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric) operators such that $\lim_{p \rightarrow +\infty} \|T_p - T\| = 0$, then T is also $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric). Indeed, assume $T_p \rightarrow T$ in norm as $p \rightarrow \infty$, then $0 = \Omega_A^{m,n}(T_p) \rightarrow \Omega_A^{m,n}(T)$. Thus $\Omega_A^{m,n}(T) = 0$ and $T \in (A; (m, n))$.

Let

$$\mathcal{B}_A(H) := \{T \in \mathcal{B}(H) : \mathcal{R}(T^*A) \subset \mathcal{R}(A)\}.$$

Recall that, if $T \in \mathcal{B}_A(H)$, then T admits an A -adjoint operator $T^\sharp$. This operator verifies

$$AT^\sharp = T^*A, \mathcal{R}(T^\sharp) \subseteq \overline{\mathcal{R}(A)} \text{ and } \mathcal{N}(T^\sharp) = \mathcal{N}(T^*A).$$

- (1) Let $T \in \mathcal{B}_A(H)$. Then, T is $(A; (m, n))$ -isosymmetric if and only if the following operator identity holds

$$\sum_{j=0}^m \sum_{k=0}^n (-1)^{m+n-(j+k)} \binom{m}{j} \binom{n}{k} T^{\sharp k+j} T^{n+j-k} = 0.$$

- (2) Under some conditions, the classes of (m, n) -isosymmetries and $(A, (m, n))$ -isosymmetries coincide. Let $T \in \mathcal{B}(H)$ and A be injective. Then, the following claims hold.
- (a) If A and T commute, then T is $(A, (m, n))$ -isosymmetric if and only if T is (m, n) -isosymmetric.
 - (b) If T is A -normal (i.e $T^\sharp T = TT^\sharp$), then T is $(A, (m, n))$ -isosymmetric if and only if $T^\sharp$ is $(A, (m, n))$ -isosymmetric.

Example 1.2.

- (1) The identity operator and the orthogonal projection on $\overline{\mathcal{R}(A)}$ are in $(A; (m, n))$, for every $m, n \in \mathbb{N}$.
- (2) Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. If T is nilpotent of order $r > 2$, then $T \in (A; (2r-2, 2r-1))$. Indeed, since T is nilpotent of order r , it gives that $T^*j = T^j = 0$ for all $j \geq r$. Then, since $\max\{k+j, (2r-1)-j+k\} \geq r$ for any $j \in \{0, 1, \dots, 2r-1\}$ and $k \in \{0, 1, \dots, 2r-2\}$, we get

$$\Omega_A^{2r-2, 2r-1}(T) = \sum_{k=0}^{2r-2} \sum_{j=0}^{2r-1} (-1)^{4r-3-k-j} \binom{2r-2}{k} \binom{2r-1}{j} T^{*k+j} AT^{2r-1-j+k} = 0.$$

- (3) Generally, an $(A; (m, n))$ -isosymmetry is not necessarily an (m, n) -isosymmetry. Let $H = \mathbb{C}^2$ with norm $\|(x, y)\|^2 = |x|^2 + |y|^2$. Let $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ and $T = \begin{pmatrix} 0 & -\frac{1}{2} \\ 0 & \frac{1}{2} \end{pmatrix}$. Since $T \in (A; (0, 1))$, it holds that $T \in (A; (1, 1))$. Moreover, T is not isosymmetric.
- (4) An operator $T \in \mathcal{B}(H)$ can be in $(A; (m, n))$, without being in $(A; (m, 0))$ or in $(A; (0, n))$. In fact, Let $H = \mathbb{C}^3$ with norm $\|(x, y, z)\|^2 = |x|^2 + |y|^2 + |z|^2$, $A =$

$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{pmatrix}$ and $T = \begin{pmatrix} 0 & 0 & -1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$. Then, a simple calculation gives $T \in (A; (1, 1))$ but neither in $(A; (1, 0))$, nor in $(A; (0, 1))$.

We now outline the plan of this paper. In Section 2, we restrict our selves to investigate general properties of $(A, (m, n))$ -isosymmetric operators. In particular, we show that every integer power of an $(A, (m, n))$ -isosymmetry is also $(A, (m, n))$ -isosymmetric. Some relations between $(A, (m, n))$ -isosymmetries (resp. skew $(A, (m, n))$ -isosymmetries) and their exponential operators are established. At the end of the section, we study some spectral properties. In Section 3, we investigate the stability of an $(A, (m, n))$ -isosymmetric (resp. skew $(A, (m, n))$ -isosymmetric) operator T under some perturbation by a nilpotent operator. More precisely, we show that if T is $(A, (m, n))$ -isosymmetric and if Q is a nilpotent operator of order r doubly commuting with T , then T^p is $(A, (m, n))$ -isosymmetric symmetric for any $p \in \mathbb{N}$ and $(T + Q)$ is $(A, (m + 2r - 2, n + 2r - 1))$ -isosymmetric. In the closing section, we characterize a family of A -isosymmetric upper triangular operator matrices. We first show that, under suitable conditions on $N, E, X \in \mathcal{B}(H)$, if N is A -isometric, then $\mathbf{T} = \begin{pmatrix} N & E \\ 0 & X \end{pmatrix}$ on $\mathbf{H} = H \oplus H$ is A -isosymmetric if and only if X is A -isosymmetric. Moreover, if N is A -symmetric and $NE = EX$, then, $\mathbf{T}$ is A -isosymmetric if and only if X is A -isosymmetric and $AE = ANEX$. We then prove that if $\mathcal{M} = \ker(\mathcal{I}_A^1(\mathbf{T}))$ is invariant for A and for $\mathbf{T}$, then $\mathbf{T} = \begin{pmatrix} N & E \\ 0 & X \end{pmatrix}$ on $\mathcal{M} \oplus \mathcal{M}^\perp$, with N is an $A|_{\mathcal{M}}$ -isometry and $E^*A|_{\mathcal{M}}N = 0$.

2. BASIC PROPERTIES OF $(A; (m, n))$ -ISOSYMMETRIES

The main purpose of this section is to develop some properties for the classes of $(A; (m, n))$ -isosymmetries and skew $(A; (m, n))$ -isosymmetries on Hilbert spaces.

Proposition 2.1. *Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Then, the following assertions hold.*

(1) *T is $(A; (m, n))$ -isosymmetric if and only if*

$$\begin{aligned} \langle \Omega_A^{m,n}(T)x | x \rangle &= \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \langle \mathcal{S}_A^n(T)T^k x | T^k x \rangle \\ (2.1) \quad &= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \langle \mathcal{I}_A^m(T)T^{n-k} x | T^k x \rangle = 0, \quad \forall x \in H. \end{aligned}$$

(2) *If T is $(A; (m, n))$ -isosymmetric, then T is $(A; (m', n'))$ -isosymmetric ($m' \geq m, n' \geq n$).*

(3) *If T is (A, n) -symmetric (resp. skew A -symmetric), then $(T - s)$ is $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric) for $s \in \mathbb{R}$.*

Proof. (1) A simple computation shows that

$$(\Omega_A^{m,n}(T))^* = \begin{cases} \Omega_A^{m,n}(T), & \text{if } n \text{ is even;} \\ -\Omega_A^{m,n}(T), & \text{if } n \text{ is odd.} \end{cases}$$

If n is odd then $(i\Omega_A^{m,n}(T))^* = i\Omega_A^{m,n}(T)$ and, consequently, $\mathcal{K} := i\Omega_A^{m,n}(T)$ is a symmetric operator, which allows to conclude.

(2) Since the following identities hold

$$\mathcal{I}_A^{m+1}(T) = (y\{(yx-1)^m\}_{\mathbf{a}}x - \{(yx-1)^m\}_{\mathbf{a}})(T, A) = T^*\mathcal{I}_A^m(T) - \mathcal{I}_A^m(T)T,$$

$$\mathcal{S}_A^{n+1}(T) = (y\{(y-x)^n\}_{\mathbf{a}} - \{(y-x)^n\}_{\mathbf{a}}x)(T, A) = T^*\mathcal{S}_A^n(T)T - \mathcal{S}_A^n(T),$$

we obtain immediately

$$\Omega_A^{m+1,n}(T) = T^*\Omega_A^{m,n}(T) - \Omega_A^{m,n}(T)T,$$

$$\Omega_A^{m,n+1}(T) = T^*\Omega_A^{m,n}(T)T - \Omega_A^{m,n}(T).$$

The above two equalities completes the proof.

(3) The following identity follows from the multinomial theorem

$$\begin{aligned}
& ((y-s)(x-s)-1)^m \{((y-s)-(x-s))^n\}_{\mathbf{a}} \\
&= ((yx-1) + (y-s)(-s) + (-sx))^m \{(y-x)^n\}_{\mathbf{a}} \\
&= \left(\sum_{\alpha_1+\alpha_2+\alpha_3=m} \binom{m}{\alpha_1, \alpha_2, \alpha_3} (y-s)^{\alpha_2} (-s)^{\alpha_3+\alpha_2} (yx-1)^{\alpha_1} x^{\alpha_3} \right) \{(y-x)^n\}_{\mathbf{a}} \\
&= \sum_{\alpha_1+\alpha_2+\alpha_3=m} \binom{m}{\alpha_1, \alpha_2, \alpha_3} (y-s)^{\alpha_2} (-s)^{\alpha_3+\alpha_2} (yx-1)^{\alpha_1} \{(y-x)^n\}_{\mathbf{a}} x^{\alpha_3} \\
&= \sum_{k=0}^m \sum_{j=0}^{m-k} \binom{m}{k} \binom{m-k}{j} (y-s)^k (-s)^{k+j} (yx-1)^{m-k-j} \{(y-x)^n\}_{\mathbf{a}} x^j.
\end{aligned}$$

Moreover, we have

$$\begin{aligned}
& ((y-s)(x-s)-1)^m \{((y-s)+(x-s))^n\}_{\mathbf{a}} \\
&= \sum_{k=0}^m \sum_{j=0}^{m-k} \sum_{i=0}^{n-i} \binom{m}{k} \binom{m-k}{j} \binom{n}{i} (y-s)^k (-s)^{k+j} (yx-1)^{m-k-j} \{(y+x)^i\}_{\mathbf{a}} (-2s)^{n-i} x^j.
\end{aligned}$$

By using these identities with x replaced by T , y replaced by T^* and a replaced by A , we obtain

$$(2.2) \quad \Omega_A^{m,n}(T-s) = \sum_{k=0}^m \sum_{j=0}^{m-k} \binom{m}{k} \binom{m-k}{j} (T^*-s)^k (-s)^{k+j} \Omega_A^{m-k-j,n}(T) T^j,$$

$$(2.3) \quad \Lambda_A^{m,n}(T-s) = \sum_{k=0}^m \sum_{j=0}^{m-k} \sum_{i=0}^{n-i} \binom{m}{k} \binom{m-k}{j} \binom{n}{i} (T^*-s)^k (-s)^{k+j} \Lambda_A^{m-k-j,i}(T) (-2s)^{n-i} T^j.$$

This completes the proof. $\square$

Proposition 2.2. *Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Then, the following statements hold.*

- (1) *If T is (A, m) -isometric and $\Omega_A^{m-1,n}(T) = 0$, then T is $\left(\frac{1}{(m-1)!} \mathcal{T}_A^{(m-1)}(T), n\right)$ -symmetric.*
- (2) *If T is (A, n) -symmetric and $\Omega_A^{m,n-1}(T) = 0$, then T is $\left(\frac{(-i)^{n-1}}{(n-1)!} \mathcal{S}_A^{(n-1)}(T), m\right)$ -isometric.*

Proof. (1) Let consider the following identity

$$\Omega_A^{m,n}(T) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} T^{*k} \mathcal{T}_A^m(T) T^{n-k}.$$

Since T is (A, m) -isometric, $\frac{1}{(m-1)!} \mathcal{T}_A^{m-1}(T)$ is positive and T is $(A; (m, n))$ -isosymmetric. Moreover, we have

$$\frac{1}{(m-1)!} \Omega_A^{m-1,n}(T) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} T^{*k} \left\{ \frac{1}{(m-1)!} \mathcal{T}_A^{m-1}(T) \right\} T^{n-k} = 0.$$

Then, $\frac{(-i)^{n-1}}{(m-1)!(n-1)!} \Omega_A^{m-1,n-1}(T)$ is positive and T is $\left(\frac{1}{(m-1)!} \mathcal{T}_A^{(m-1)}(T), n\right)$ -symmetric.

(2) Let consider the following identity

$$\Omega_A^{m,n}(T) = \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} T^{*k} \mathcal{S}_A^n(T) T^k.$$

Since T is (A, n) -symmetric, $\frac{(-i)^{n-1}}{(n-1)!} \mathcal{S}_A^{(n-1)}(T)$ is a positive operator and T is $(A; (m, n))$ -isosymmetric. On the other hand, we have

$$\frac{(-i)^{n-1}}{(n-1)!} \Omega_A^{m, n-1}(T) = \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} T^{*k} \left\{ \frac{(-i)^{n-1}}{(n-1)!} \mathcal{S}_A^{(n-1)}(T) \right\} T^k = 0.$$

Hence, $\frac{(-i)^{n-1}}{(m-1)!(n-1)!} \Omega_A^{m-1, n-1}(T)$ is positive and T is $\left(\frac{(-i)^{n-1}}{(n-1)!} \mathcal{S}_A^{(n-1)}(T), m \right)$ -isometric. $\square$

Corollary 2.3. *Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Assume that T is (A, n) -symmetric and $\Omega_A^{m, n-1}(T) = 0$. Then, the following properties hold.*

- (1) *If $n = 2p + 1$, then $\Omega_A^{m-1, 2p}(T) \geq 0$ if p is even.*
- (2) *If n is even, then T is $(A; (m-1, n-1))$ -isosymmetric.*

Proof. (1) Assume $n = 2p + 1$. Since T is (A, n) -symmetric and $\Omega_A^{m, n-1}(T) = 0$, we have

$$0 \leq \frac{(-i)^{n-1}}{(m-1)!(n-1)!} \Omega_A^{m-1, n-1}(T) = \frac{(-1)^p}{(m-1)!(2p)!} \Omega_A^{m, 2p}(T)$$

which allows to conclude.

- (2) Assume that n is even. Since T is (A, n) -symmetric, by Theorem 2.5-[13] we have $\mathcal{S}_A^{n-1}(T) = 0$. Hence, $\Omega_A^{m-1, n-1}(T) = 0$, that is T is $(A, (m-1, n-1))$ -isosymmetric. $\square$

Theorem 2.4. *Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Then, the following assertions hold.*

- (1) *If T is invertible, then T is $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric) if and only if T^{-1} is $(A; (m, n))$ -isosymmetric (resp. skew $(A; (m, n))$ -isosymmetric).*
- (2) *If T is $(A; (m, n))$ -isosymmetric, then T^k is $(A; (m, n))$ -isosymmetric ($\forall k \in \mathbb{N}$).*
- (3) *If T is skew $(A; (m, n))$ -isosymmetric, then T^k is skew $(A; (m, n))$ -isosymmetric (k odd).*

Proof. (1) Assume that T is $(A; (m, n))$ -isosymmetric and invertible. Since if m is even, then $(-1)^p = (-1)^{m-p}$ and if m is odd, then $(-1)^p = (-1)^{m+1-p}$, we obtain

$$\begin{aligned} 0 &= (T^{-1})^{*m+n} \Omega_A^{m, n}(T) (T^{-1})^{m+n} \\ &= \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*-m-n} T^{*j} \mathcal{S}_A^n(T) T^j T^{-m-n} \\ &= \sum_{j=0}^m (-1)^{m-j} \binom{m}{m-j} T^{*-(m-j)} (T^{*-n} \mathcal{S}_A^n(T) T^{-n}) T^{-(m-j)} \\ &= \sum_{p=0}^m (-1)^p \binom{m}{p} (T^{-1})^{*p} ((T^{-1})^{*n} \mathcal{S}_A^n(T) (T^{-1})^n) (T^{-1})^{-p} \\ &= \begin{cases} \Omega_{T^{-1}}^{m, n}(A), & \text{if } m \text{ is even;} \\ -\Omega_{T^{-1}}^{m, n}(A), & \text{if } m \text{ is odd.} \end{cases} \end{aligned}$$

By applying a similar argument, we can show the corresponding property for skew $(A; (m, n))$ -isosymmetric operators.

(2) Note that, for $k \in \mathbb{N}$ and for some constants α_l and β_j , we have

$$\begin{aligned}
& (y^k x^k - 1)^m \{(y^k - x^k)^n\}_{\mathbf{a}} \\
&= ((yx - 1)(y^{k-1} x^{k-1} + y^{k-2} x^{k-2} + \cdots + 1))^m \{((y - x)(y^{k-1} + y^{k-2} x + \cdots + x^{k-1}))^n\}_{\mathbf{a}} \\
&= \left(\sum_{l=0}^{m(k-1)} \alpha_l y^{m(k-1)-l} (yx - 1)^m x^{m(k-1)-l} \right) \left\{ \sum_{j=0}^{n(k-1)} \beta_j y^{n(k-1)-j} (y - x)^n x^j \right\}_{\mathbf{a}} \\
&= \left(\sum_{l=0}^{m(k-1)} \alpha_l y^{m(k-1)-l} (yx - 1)^m x^{m(k-1)-l} \right) \left(\sum_{j=0}^{n(k-1)} \beta_j y^{n(k-1)-j} \{(y - x)^n\}_{\mathbf{a}} x^j \right) \\
&= \sum_{l=0}^{m(k-1)} \sum_{j=0}^{n(k-1)} \alpha_l \beta_j y^{m(k-1)-l} y^{n(k-1)-j} (yx - 1)^m \{(y - x)^n\}_{\mathbf{a}} x^j x^{m(k-1)-l}.
\end{aligned}$$

The above identity gives

$$\begin{aligned}
\Omega_A^{m,n}(T^k) &= ((y^k x^k - 1)^m \{(y^k - x^k)^n\}_{\mathbf{a}})(T^*, T, A) \\
&= \sum_{l=0}^{m(k-1)} \sum_{j=0}^{n(k-1)} \alpha_l \beta_j T^{*m(k-1)-l+n(k-1)-j} \Omega_A^{m,n}(T) T^{j+m(k-1)-l},
\end{aligned}$$

and the desired claim follows from that.

(3) For an odd integer k and for some constants α_l and ξ_j , the following formula holds true

$$\begin{aligned}
& (y^k x^k - 1)^m \{(y^k + x^k)^n\}_{\mathbf{a}} \\
&= ((yx - 1)(y^{k-1} x^{k-1} + y^{k-2} x^{k-2} + \cdots + 1))^m \left\{ ((y + x)(y^{k-1} - y^{k-2} x + \cdots - y x^{k-2} + x^{k-1}))^n \right\}_{\mathbf{a}} \\
&= \left(\sum_{l=0}^{m(k-1)} \eta_l y^{m(k-1)-l} (yx - 1)^m x^{m(k-1)-l} \right) \left\{ \sum_{j=0}^{n(k-1)} \xi_j y^{n(k-1)-j} (y + x)^n x^j \right\}_{\mathbf{a}} \\
&= \left(\sum_{l=0}^{m(k-1)} \eta_l y^{m(k-1)-l} (yx - 1)^m x^{m(k-1)-l} \right) \left(\sum_{j=0}^{n(k-1)} \xi_j y^{n(k-1)-j} \{(y + x)^n\}_{\mathbf{a}} x^j \right) \\
&= \sum_{l=0}^{m(k-1)} \sum_{j=0}^{n(k-1)} \eta_l \xi_j y^{m(k-1)-l} y^{n(k-1)-j} (yx - 1)^m \{(y + x)^n\}_{\mathbf{a}} x^j x^{m(k-1)-l}.
\end{aligned}$$

It holds from the above identity

$$\begin{aligned}
\Lambda_A^{m,n}(T^k) &= ((y^k x^k - 1)^m \{(y^k + x^k)^n\}_{\mathbf{a}})(T^*, T, A) \\
&= \sum_{l=0}^{m(k-1)} \sum_{j=0}^{n(k-1)} \eta_l \xi_j T^{*m(k-1)-l+n(k-1)-j} \Lambda_A^{m,n}(T) T^{j+m(k-1)-l}.
\end{aligned}$$

If T is skew $(A; (m, n))$ -isosymmetric, that is $\Lambda_A^{m,n}(T) = 0$, then $\Lambda_A^{m,n}(T^k) = 0$. Hence, T^k is skew $(A; (m, n))$ -isosymmetric. □

In the following corollary, we state some immediate consequences of Theorem 2.4.

Corollary 2.5. *Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Then, the following statements hold.*

- (1) *If T is (A, m) -isometric, then T^k is (A, m) -isometric for any $k \in \mathbb{N}$.*
- (2) *If T is (A, n) -symmetric, then T^k is (A, n) -symmetric for any $k \in \mathbb{N}$.*
- (3) *If T is skew (A, n) -symmetric, then T^k is skew (A, n) -symmetric for k odd.*

Proposition 2.6. *Let T be $(A, (m, n))$ -isosymmetric. Then, the following statements hold.*

(1) If $p \geq m$, then

$$(2.4) \quad \sum_{k=0}^p (-1)^{p-k} \binom{p}{k} k^i T^{*k} \mathcal{S}_A^n(T) T^k = 0, \quad i = 0, 1, \dots, p-m.$$

(2) If $p \geq n$, then

$$(2.5) \quad \sum_{k=0}^p (-1)^{p-k} \binom{p}{k} k^i T^{*k} \mathcal{I}_A^m(T) T^{p-k} = 0, \quad i = 0, 1, \dots, p-n.$$

Proof. (1) We aim to prove (2.4) by induction on p . Assume that T is $(A, (m, n))$ -isosymmetric, then (2)-Proposition (2.1) yields that T is $(A, (m', n))$ -isosymmetric for each $m' \geq m$. Thus, for $i = 0$ the proof of (2.4) is immediate. Moreover, the result is true for $p = m$. Assume now that (2.4) is true for $i \in \{1, 2, \dots, p-m\}$ and prove it for $i \in \{1, 2, \dots, p+1-m\}$. By the induction hypothesis, we obtain

$$\begin{aligned} & \sum_{k=0}^{p+1} (-1)^{p+1-k} \binom{p+1}{k} k^i T^{*k} \mathcal{S}_A^n(T) T^k \\ &= \sum_{k=1}^{p+1} (-1)^{p+1-k} \binom{p+1}{k} k^i T^{*k} \mathcal{S}_A^n(T) T^k \\ &= \sum_{k=0}^p (-1)^{p-k} \binom{p+1}{k+1} (k+1)^i T^{*k+1} \mathcal{S}_A^n(T) T^{k+1} \\ &= (p+1) T^* \left\{ \sum_{k=0}^p (-1)^{p-k} \binom{p}{k} (k+1)^{i-1} T^{*k} \mathcal{S}_A^n(T) T^k \right\} T \\ &= (p+1) T^* \left\{ \sum_{k=0}^p (-1)^{p-k} \binom{p}{k} \left(\sum_{j=0}^{i-1} \binom{i-1}{j} k^j \right) T^{*k} \mathcal{S}_A^n(T) T^k \right\} T \\ &= (p+1) T^* \left\{ \sum_{j=0}^{i-1} \binom{i-1}{j} \underbrace{\left(\sum_{k=0}^p (-1)^{p-k} \binom{p}{k} k^j T^{*k} \mathcal{S}_A^n(T) T^k \right)}_{=0} \right\} T = 0. \end{aligned}$$

(2) The remaining identity (2.5) in the statement (2) also holds by a similar method. $\square$

Now, we devote most of our interest to distinguish some connection between $(A, (m, n))$ -isosymmetries, skew $(A, (m, n))$ -isosymmetries and their associated exponential operator. We begin our study by introducing the notion of left A -invertible operators which will play a central role in characterizing skew $(A; (m, n))$ -isosymmetries and giving some their interesting properties.

Definition 2.7. Let $A \in \mathcal{B}(H)^+$ and $T, S \in \mathcal{B}(H)$. We say that the operator T is left (resp., right) (A, m) -invertible by S , for some integer $m \geq 1$, if $\{(yx - 1)^m\}_{\mathbf{a}}(S, T, A) = 0$ (resp. $\{(yx - 1)^m\}_{\mathbf{a}}(T, S, A) = 0$).

Proposition 2.8. Let $A \in \mathcal{B}(H)^+$ and $R_1, R_2, S_1, S_2 \in \mathcal{B}(H)$ such that $R_1 R_2 = R_2 R_1$ and $S_1 S_2 = S_2 S_1$. Assume that R_1 is a left (A, m) -inverse of S_1 and R_2 is a left (A, m) -inverse of S_2 , then $R_1 R_2$ is a left $(A, m+n-1)$ -inverse of $S_1 S_2$.

Proof. Fix $x, y \in H$ and let $a_{i,j} = \langle R_1^i R_2^j A S_1^i S_2^j x \mid y \rangle$. Then, for all non-negative integers k and l , we have

$$a_{k+i, l} = \langle R_1^i A S_1^i (S_1^k S_2^l x) \mid (R_1^{*k} R_2^{*l} y) \rangle, \quad a_{k, l+j} = \langle R_2^j A S_2^j (S_1^k S_2^l x) \mid (R_1^{*k} R_2^{*l} y) \rangle.$$

The left (A, m) -invertibility of S_1 by R_1 and the left (A, n) -invertibility of S_2 by R_2 gives

$$\sum_{i=0}^m (-1)^{m-i} a_{k+i, l} = 0, \quad \sum_{j=0}^n (-1)^{n-j} a_{k, l+j} = 0.$$

Applying [9, Corollary 2.5] we deduce that

$$0 = \sum_{s=0}^{m+n-1} (-1)^{m+n-1-s} a_{s, s} = \sum_{s=0}^{m+n-1} (-1)^{m+n-1-s} \binom{m+n-1}{s} \langle (R_1 R_2)^s A (S_1 S_2)^s x | y \rangle$$

Since x and y are arbitrary in H , it holds

$$\sum_{s=0}^{m+n-1} (-1)^{m+n-1-s} \binom{m+n-1}{s} (R_1 R_2)^s A (S_1 S_2)^s = 0.$$

Hence, $R_1 R_2$ is a left $(A, m+n-1)$ -inverse of $S_1 S_2$. □

Proposition 2.9. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Then, the following statements hold.*

(1) *T is $(A, (m, n))$ -isosymmetric if and only if, for each positive integer k , it holds*

$$(2.6) \quad (e^{isT})^{*k} \mathcal{I}_A^m(T) (e^{isT})^k = \sum_{h=0}^{n-1} \frac{(-isk)^h}{h!} \Omega_A^{m, h}(T).$$

(2) *T is skew $(A, (m, n))$ -isosymmetric if and only if, for each positive integer k , it holds*

$$(2.7) \quad (e^{isT^*})^k \mathcal{I}_A^m(T) (e^{isT})^k = \sum_{h=0}^{n-1} \frac{(isk)^h}{h!} \Lambda_A^{m, h}(T).$$

Proof. (1) We have shown, in [13], that

$$(2.8) \quad (e^{isT})^* A (e^{isT}) = \mathcal{S}_A^0(T) + (-is) \mathcal{S}_A^1(T) + \frac{(-is)^2}{2!} \mathcal{S}_A^2(T) + \frac{(-is)^3}{3!} \mathcal{S}_A^3(T) + \dots$$

It holds from (2.8) that

$$\begin{aligned} & (e^{isT})^{*k} \mathcal{I}_A^m(T) (e^{isT})^k = \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} (e^{isT})^{*k} A (e^{isT})^k T^j \\ &= \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^0(T) T^j + (-is) k \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^1(T) T^j \\ &+ \frac{(-is)^2 k^2}{2!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^2(T) T^j + \frac{(-is)^3 k^3}{3!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^3(T) T^j + \dots + \\ &+ \dots + \frac{(-is)^{n-1} k^{n-1}}{(n-1)!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^{n-1}(T) T^j \\ &+ \frac{(-is)^n k^n}{n!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \mathcal{S}_A^n(T) T^j + \dots \\ &= \Omega_A^{m, 0}(T) + (-is) k \Omega_A^{m, 1}(T) + \frac{(-is)^2 k^2}{2!} \Omega_A^{m, 2}(T) + \frac{(-is)^3 k^3}{3!} \Omega_A^{m, 3}(T) + \dots \\ &+ \dots + \frac{(-is)^{n-1} k^{n-1}}{(n-1)!} \Omega_A^{m, n-1}(T) + \frac{(-is)^n k^n}{n!} \Omega_A^{m, n}(T) + \dots \end{aligned}$$

Since T is $(A, (m, n))$ -isosymmetric, $\Omega_A^{m, n'}(T) = 0$ for $n' \geq n$. This completes the proof.

(2) A simple calculation shows that

$$(2.9) \quad e^{isT^*} A e^{isT} = \zeta_T^0(A) + (is)\zeta_T^1(A) + \frac{(is)^2}{2!}\zeta_T^2(A) + \frac{(is)^3}{3!}\zeta_T^3(A) + \dots$$

It holds from (2.9) that

$$\begin{aligned} & \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \left(e^{isT^*}\right)^k A \left(e^{isT}\right)^k T^j \\ &= \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} A T^j + (is)k \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \zeta_A^1(T) T^j \\ & \quad + \frac{(is)^2 k^2}{2!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \zeta_A^2(T) T^j + \frac{(is)^3 k^3}{3!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \zeta_A^3(T) T^j + \dots + \\ & \quad + \dots + \frac{(is)^{n-1} k^{n-1}}{(n-1)!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \zeta_A^{n-1}(T) T^j \\ & \quad + \frac{(is)^n k^n}{n!} \sum_{j=0}^m (-1)^{m-j} \binom{m}{j} T^{*j} \zeta_A^n(T) T^j + \dots \\ &= \Lambda_A^{m,0}(T) + (is)k \Lambda_A^{m,1}(T) + \frac{(is)^2 k^2}{2!} \Lambda_A^{m,2}(T) + \frac{(is)^3 k^3}{3!} \Lambda_A^{m,3}(T) + \dots \\ & \quad + \dots + \frac{(is)^{n-1} k^{n-1}}{(n-1)!} \Lambda_A^{m,n-1}(T) + \frac{(is)^n k^n}{n!} \Lambda_A^{m,n}(T) + \dots \end{aligned}$$

Since T is skew $(A, (m, n))$ -isosymmetric, $\Lambda_A^{m,n}(T) = 0$ for $n' \geq n$, and this allows to conclude. $\square$

Theorem 2.10. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Then the following statements hold.*

(1) *If T is $(A, (m, n))$ -isosymmetric, then for every $s \in \mathbb{R}$, it holds*

$$(2.10) \quad \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT}\right)^{*k} \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = 0.$$

(2) *If T is skew $(A, (m, n))$ -isosymmetric, then for every $s \in \mathbb{R}$, it holds*

$$\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = 0.$$

Proof. (1) By using (2.7) and taking into account the identity

$$(2.11) \quad \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^j = \begin{cases} 0, & \text{if } 0 \leq j \leq n-1; \\ n!, & \text{if } j = n, \end{cases}$$

it follows

$$\begin{aligned}
& \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT} \right)^{*k} \mathcal{I}_A^m(T) \left(e^{isT} \right)^k \\
&= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left\{ \Omega_A^{m,0}(T) + (-is)k \Omega_A^{m,1}(T) + \frac{(-is)^2 k^2}{2!} \Omega_A^{m,2}(T) + \frac{(-is)^3 k^3}{3!} \Omega_A^{m,3}(T) + \dots \right. \\
&\quad \left. + \dots + \frac{(-is)^{n-1} k^{n-1}}{(n-1)!} \Omega_A^{m,n-1}(T) + \frac{(-is)^n k^n}{n!} \Omega_A^{m,n}(T) + \dots \right\} \\
&= \Omega_A^{m,0}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \right) + (-is) \Omega_A^{m,1}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k \right) \\
&\quad + \frac{(-is)^2}{2!} \Omega_A^{m,2}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^2 \right) + \dots + \\
&\quad + \dots + \frac{(-is)^{n-1}}{(n-1)!} \Omega_A^{m,n-1}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^{n-1} \right) = 0.
\end{aligned}$$

(2) By using (2.7) and (2.11), it follows

$$\begin{aligned}
& \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT^*} \right)^k \mathcal{I}_A^m(T) \left(e^{isT} \right)^k \\
&= \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left\{ \Lambda_A^{m,0}(T) + (is)k \Lambda_A^{m,1}(T) + \frac{(is)^2 k^2}{2!} \Lambda_A^{m,2}(T) + \frac{(is)^3 k^3}{3!} \Lambda_A^{m,3}(T) + \dots \right. \\
&\quad \left. + \dots + \frac{(is)^{n-1} k^{n-1}}{(n-1)!} \Lambda_A^{m,n-1}(T) + \frac{(is)^n k^n}{n!} \Lambda_A^{m,n}(T) + \dots \right\} \\
&= \Lambda_A^{m,0}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \right) + (is) \Lambda_A^{m,1}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k \right) \\
&\quad + \frac{(is)^2}{2!} \Lambda_A^{m,2}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^2 \right) + \dots + \\
&\quad + \dots + \frac{(is)^{n-1}}{(n-1)!} \Lambda_A^{m,n-1}(T) \left(\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} k^{n-1} \right) = 0.
\end{aligned}$$

□

The following corollary is an immediate consequence of Theorem 2.10.

Corollary 2.11. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Assume that $\mathcal{I}_A^n(T) \geq 0$. Then, we have:*

- (1) *If T is $(A, (m, n))$ -isosymmetric, then for every $s \in \mathbb{R}$, the operator e^{isT} is $(\mathcal{I}_A^m(T), n)$ -isometric. In particular, if T is (A, n) -symmetric, then e^{isT} is (A, n) -isometric.*
- (2) *If T is skew $(A, (m, n))$ -isosymmetric, then for every $s \in \mathbb{R}$, the operator e^{isT} is left $(\mathcal{I}_A^m(T), n)$ -invertible. In particular, if T is skew (A, n) -symmetric, then T is left (A, n) -invertible with a left inverse e^{isT^*} .*

Theorem 2.12. *Let $A \in \mathcal{B}(H)^+$ and $R, S \in \mathcal{B}(H)$ such that $RS = SR$. If R is skew (A, m) -symmetric and S is skew (A, n) -symmetric, then $e^{is(R+S)}$ is left $(A, m+n-1)$ -invertible with left inverse $e^{is(R+S)^*}$.*

Proof. Since R is skew (A, m) -symmetric and S is skew (A, n) -symmetric, it holds from Corollary 2.11 that $S_1 = e^{isR}$ is left (A, m) -invertible with left inverse $R_1 = e^{isR^*}$ and $S_2 = e^{isS}$ is left (A, n) -invertible with left inverse $R_2 = e^{isS^*}$, for every $s \in \mathbb{R}$. It is clear that $R_1 R_2 = R_2 R_1$ and $S_1 S_2 = S_2 S_1$. Applying Proposition 2.8, we deduce that $S_1 S_2 = e^{is(S+R)}$ is left $(A, m+n-1)$ -invertible with left inverse $R_1 R_2 = e^{is(R+S)^*}$.

□

In [13], we have introduced the following algebraic condition on $T \in \mathcal{B}(H)$

$$(2.12) \quad (\mathbf{POL}_n(\mathbf{A})) : \quad (e^{isT})^* A (e^{isT}) = A + \sum_{l=1}^n s^l B_l, \quad B_l := B_l(A, T, T^*) \in \mathcal{B}(H).$$

Note that an induction argument gives

$$\frac{d^{n+1}}{ds^{n+1}} (e^{isT})^* A (e^{isT}) = (-i)^{n+1} (e^{isT})^* \mathcal{S}_A^{n+1}(T) (e^{isT}).$$

The $(\mathbf{POL}_n(\mathbf{A}))$ condition is equivalent to $\frac{d^{n+1}}{ds^{n+1}} (e^{isT})^* A (e^{isT}) = 0$ and consequently to $\mathcal{S}_A^{n+1}(T) = 0$. Hence, $T \in \mathcal{B}(\mathbb{H})$ is (A, n) -symmetric if it satisfies $(\mathbf{POL}_{n-1}(\mathbf{A}))$.

Inspired by (2.12), we introduce

$$(2.13) \quad (\mathbf{IPOL}_n(\mathbf{A})) : \quad (e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) = \Omega_A^{m,0}(T) + \sum_{l=1}^n s^l C_l, \quad C_l := C_l(A, T, T^*) \in \mathcal{B}(H).$$

For $m \in \mathbb{N}$, we obtain

$$\frac{d^{n+1}}{ds^{n+1}} (e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) = (-i)^{n+1} (e^{isT})^* \Omega_A^{m,n+1}(T) (e^{isT}).$$

Moreover,

$$\begin{aligned} (e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) &= \sum_{h \geq 0} \frac{s^h}{h!} \left(\frac{d^h}{ds^h} (e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) \Big|_{s=0} \right) \\ &= \sum_{h \geq 0} \frac{(-is)^h}{h!} \Omega_A^{m,h}(T). \end{aligned}$$

Referring to Proposition 2.9 and applying the identity (2.6) for $k = 1$, we deduce that $(\mathbf{IPOL}_n(\mathbf{A}))$ is equivalent to $\frac{d^{n+1}}{ds^{n+1}} (e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) = 0$ and consequently to $\Omega_A^{m,n+1}(T) = 0$.

Remark 2.13. For $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$, T is $(A, (m, n))$ -isosymmetric if and only if T satisfies $(\mathbf{IPOL}_{n-1}(\mathbf{A}))$.

Theorem 2.14. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Then the following properties hold.*

- (1) *If m is even, then every invertible $(A; (m, n))$ -isosymmetric operator T satisfying $\mathcal{S}_A^n(T) \geq 0$, is $(A; (m-1, n))$ -isosymmetric.*
- (2) *If n is even, then every $(A, (m, n))$ -isosymmetric operator T satisfying $\mathcal{I}_A^m(T) \geq 0$ is $(A, (m, n-1))$ -isosymmetric.*

Proof. (1) If $\mathcal{S}_A^n(T) \geq 0$ for some positive integer n , then T is $(A; (m, n))$ -isosymmetric if and only if T is $(\mathcal{S}_A^n(T), m)$ -isometric. Note that if T is invertible, then $\mathcal{S}_A^n(T^{-1}) \geq 0$ and, consequently, T^{-1} is $(A; (m, n))$ -isosymmetric if and only if T^{-1} is $(\mathcal{S}_A^n(T^{-1}), m)$ -isometric. It follows from that $\Omega_A^{m-1,n}(T) \geq 0$ and $\Omega_A^{m-1,n}(T^{-1}) \geq 0$. Hence, the following holds

$$(2.14) \quad 0 \leq (T^{-1})^{*m-1+n} \Omega_A^{m-1,n}(T) (T^{-1})^{m-1+n} = \begin{cases} \Omega_{T^{-1}}^{m-1,n}(A), & \text{if } m-1 \text{ is even;} \\ -\Omega_{T^{-1}}^{m-1,n}(A), & \text{if } m-1 \text{ is odd.} \end{cases}$$

Since m is an even number, $m-1$ is odd and (2.14) gives $\Omega_A^{m-1,n}(T) \leq 0$. Therefore, we have $\Omega_A^{m-1,n}(T) = 0$. Hence, T is $(A; (m-1, n))$ -isosymmetric.

- (2) Since T is $(A, (m, n))$ -isosymmetric, it holds for n even (that is $n = 2k$),

$$(e^{isT})^* \mathcal{I}_A^m(T) (e^{isT}) = \sum_{h=0}^{2k-1} \frac{(-is)^h}{h!} \Omega_A^{m,h}(T).$$

The positivity of $\mathcal{I}_A^m(T)$ implies

$$\begin{aligned} 0 &\leq \frac{(e^{isT})^* \mathcal{I}_A^m(T) (e^{isT})}{s^{2k-1}} = \frac{1}{s^{2k-1}} \Omega_A^{m,0}(T) + \sum_{h=0}^{2k-2} \frac{(-is)^h}{h!} \Omega_A^{m,h}(T) \\ &\quad + \frac{(-i)^{2k-1}}{(2k-1)!} \Omega_A^{m,2k-1}(T) \longrightarrow \frac{(-i)^{2k-1}}{(2k-1)!} \Omega_A^{m,2k-1}(T) \quad (s \longrightarrow +\infty). \end{aligned}$$

On the other hand, we have

$$\begin{aligned} 0 &\leq \frac{-(e^{isT})^* \mathcal{I}_A^m(T) (e^{isT})}{s^{2k-1}} = \frac{1}{s^{2k-1}} (-\Omega_A^{m,0}(T)) + \sum_{h=0}^{2k-2} \frac{(-is)^h}{h!} (-\Omega_A^{m,h}(T)) \\ &\quad + \frac{(-i)^{2k-1}}{(2k-1)!} (-\Omega_A^{m,2k-1}(T)) \longrightarrow \frac{(-i)^{2k-1}}{(2k-1)!} (-\Omega_A^{m,2k-1}(T)) \quad (s \longrightarrow -\infty). \end{aligned}$$

Hence, $\Omega_A^{m,2k-1}(T) = 0$, which implies that T is $(A, (m, 2k-1))$ -isosymmetric. $\square$

The following corollary follows immediately from Theorem 2.14 and summarizes few known facts.

Corollary 2.15. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Then, the following properties hold.*

- (1) *If m is even, then every invertible (A, m) -isometric operator is $(A, m-1)$ -isometric.*
- (2) *If n is even, then every (A, n) -symmetric operator is $(A, n-1)$ -symmetric.*

Let consider, now,

$$(2.15) \quad (\widetilde{\mathbf{POL}_n(\mathbf{A})}) : (e^{isT^*})A(e^{isT}) = A + \sum_{l=1}^n s^l D_l, \quad D_l := D_l(A, T, T^*) \in \mathcal{B}(H).$$

Note that

$$\frac{d^n}{ds^n} (e^{isT^*})A(e^{isT}) = i^n e^{isT^*} \zeta_T^n(A) e^{isT}, \quad n \in \mathbb{N}.$$

The $(\widetilde{\mathbf{POL}_n(\mathbf{A})})$ condition is equivalent to $\frac{d^{n+1}}{ds^{n+1}} (e^{isT^*} A e^{isT}) = 0$ and consequently to $\zeta_T^{n+1}(A) = 0$. Hence, $T \in \mathcal{B}(H)$ is skew (A, n) -symmetric if it satisfies $(\widetilde{\mathbf{POL}_{n-1}(\mathbf{A})})$.

Inspired by (2.15), we introduce

$$(2.16) \quad (\widetilde{\mathbf{IPOL}_n(\mathbf{A})}) : (e^{isT^*})\mathcal{I}_A^m(T)(e^{isT}) = \Lambda_A^{m,0}(T) + \sum_{l=1}^n s^l E_l, \quad E_l := E_l(A, T, T^*) \in \mathcal{B}(H).$$

Referring to Proposition 2.9, T is skew $(A, (m, n))$ -isosymmetric if and only if

$$\begin{aligned} (e^{isT^*})\mathcal{I}_A^m(T)(e^{isT}) &= \sum_{h=0}^{n-1} \frac{s^h}{h!} \left(\frac{d^h}{ds^h} (e^{isT^*})\mathcal{I}_A^m(T)(e^{isT}) \Big|_{s=0} \right) \\ &= \sum_{h=0}^{n-1} \frac{(is)^h}{h!} \Lambda_A^{m,h}(T), \end{aligned}$$

that is T satisfies $(\widetilde{\mathbf{IPOL}_{n-1}(\mathbf{A})})$.

In Corollary 2.15 ([8, Theorem 2.5.]), we have shown that if T is (A, n) -symmetric and n is even, then T is always $(A, n-1)$ -symmetric. Under suitable conditions, we extend this property to skew (A, m) -symmetric operators, and more generally to skew $(A, (m, n))$ -isosymmetries. For $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$, let consider the following hypothesis:

$$\begin{aligned}
(1) \quad (H_1) : & \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k \geq 0. \\
(2) \quad (H_2) : & \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{isT^*}\right)^{n-1-k} \mathcal{I}_A^m(T) \left(e^{isT}\right)^{n-1-k} \geq 0.
\end{aligned}$$

Proposition 2.16. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Assume (H_1) and (H_2) are fulfilled and*

$$\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = 0.$$

If n is even, then it holds

$$\sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = 0.$$

Proof. Since n is even, we have $(-1)^k = -(-1)^{n-1-k}$. Then, we obtain

$$\begin{aligned}
& \left(e^{isT^*}\right)^{n-1} \left\{ \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{-isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{-isT}\right)^k \right\} \left(e^{isT}\right)^{n-1} \\
&= \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{isT^*}\right)^{n-1-k} \mathcal{I}_A^m(T) \left(e^{isT}\right)^{n-1-k} \\
&= - \sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} \left(e^{isT^*}\right)^{n-1-k} \mathcal{I}_A^m(T) \left(e^{isT}\right)^{n-1-k} \geq 0.
\end{aligned}$$

This completes the proof. $\square$

Theorem 2.17. *Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$. Assume (H_1) and (H_2) are fulfilled. If T is skew $(A, (m, n))$ -isosymmetric where n is even, then T is skew $(A, (m, n-1))$ -isosymmetric.*

Proof. Since T is skew $(A, (m, n))$ -isosymmetric, $\Lambda_A^{m,p}(T) = 0$ for all $p \geq n$ and, by Theorem 2.10, the operator e^{isT} satisfies

$$\sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k = 0.$$

On the other hand, since n is even, it follows from Proposition 2.16

$$\begin{aligned}
0 &= \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left(e^{isT^*}\right)^k \mathcal{I}_A^m(T) \left(e^{isT}\right)^k \\
&= \sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \left\{ \Lambda_A^{m,0}(T) + (is)(n-1-k) \Lambda_A^{m,1}(T) + \frac{(is)^2(n-1-k)^2}{2!} \Lambda_A^{m,2}(T) \right. \\
&\quad \left. + \frac{(is)^3(n-1-k)^3}{3!} \Lambda_A^{m,3}(T) + \dots + \frac{(is)^{n-1}(n-1-k)^{n-1}}{(n-1)!} \Lambda_A^{m,n-1}(T) \right\} \\
&= \Lambda_A^{m,0}(T) \left(\sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} \right) + (is) \Lambda_A^{m,1}(T) \left(\sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} (n-1-k) \right) \\
&\quad + \frac{(is)^2}{2!} \Lambda_A^{m,2}(T) \left(\sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} (n-1-k)^2 \right) + \dots + \\
&\quad + \dots + \frac{(is)^{n-1}}{(n-1)!} \Lambda_A^{m,n-1}(T) \left(\sum_{k=0}^{n-1} (-1)^{n-1-k} \binom{n-1}{k} (n-1-k)^{n-1} \right).
\end{aligned}$$

Using (2.11), we obtain $\Lambda_A^{m,n-1}(T) = 0$. Hence T is skew $(A, (m, n-1))$ -isosymmetric. $\square$

In the rest of this section, we investigate some spectral properties of $(A, (m, n))$ -isosymmetric operators. The showed results extends those of [8], [12], [13], [19] and [20].

Theorem 2.18. *Let $A \in \mathcal{B}(H)^+$, $T \in \mathcal{B}(H)$. If T is an $(A, (m, n))$ -isosymmetric operator and $0 \notin \sigma_{ap}(A)$, then the following statements hold.*

- (1) $\partial\sigma(T) \subset \partial\mathbf{D} \cup \mathbb{R}$.
- (2) Let x, y be unit vectors and x_k, y_k be sequences of unit vectors of H . Then, we have
 - (a) If $Tx = \lambda x$, $Ty = \mu y$, $\lambda \neq \mu$ and $\lambda \neq \bar{\mu}$, then $\langle Ax | y \rangle = 0$.
 - (b) If $(T - \lambda)x_k \rightarrow 0$, $(T - \mu)y_k \rightarrow 0$ ($k \rightarrow \infty$, $\lambda \neq \mu$, $\lambda \neq \bar{\mu}$), then $\lim_{k \rightarrow \infty} \langle Ax_k | y_k \rangle = 0$.

Proof. (1) If $\lambda \in \mathbb{C}$ is in the approximate point spectrum of T , then there exists a sequence $(x_j) \subset H$ such that for all j , $\|x_j\| = 1$, and $(T - \lambda)x_j \rightarrow 0$ as $j \rightarrow \infty$. Thus, for each integer i , $\lim_{j \rightarrow \infty} (T^i - \lambda^i)x_j \rightarrow 0$. Moreover,

$$\begin{aligned} 0 &= \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \langle \mathcal{S}_A^n(T) T^k x | T^k y \rangle \\ &= (\lambda \bar{\lambda} - 1)^m ((\lambda - \bar{\lambda})^n) \lim_{j \rightarrow \infty} \langle Ax_j | x_j \rangle \\ &= (|\lambda|^2 - 1)^m (2\Im m\lambda)^n \lim_{j \rightarrow \infty} \|x_j\|_A^2, \end{aligned}$$

which implies that either $|\lambda| = 1$ or $\Im m\lambda = 0$ and so, either $\lambda \in \mathbf{D}$ or $\lambda \in \mathbb{R}$. Hence, $\sigma_{ap}(T) \subset \partial\mathbf{D} \cup \mathbb{R}$. Moreover, $\partial\sigma(T) \subset \sigma_{ap}(T)$. So, $\partial\sigma(T) \subset \partial\mathbf{D} \cup \mathbb{R}$.

- (2) (a) Let λ and μ be two distinct eigenvalues of T and suppose that $Tx = \lambda x$ and $Ty = \mu y$. Then, it holds

$$\begin{aligned} 0 &= \langle \Omega_A^{m,n}(T)x | y \rangle \\ &= \left(\sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \lambda^k \bar{\mu}^k \right) \left(\sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \lambda^{n-i} \bar{\mu}^i \right) \langle Ax | y \rangle \\ &= (\lambda \bar{\mu} - 1)^m (\lambda - \bar{\mu})^n \langle Ax | y \rangle. \end{aligned}$$

Since $\lambda \neq \mu$ and $\lambda \neq \bar{\mu}$, we obtain $\langle Ax | y \rangle = 0$.

- (b) By similar arguments as in the proof of (3), it holds

$$0 = (\lambda \bar{\mu} - 1)^m (\lambda - \bar{\mu})^n \lim_{k \rightarrow \infty} \langle Ax_k | y_k \rangle.$$

Since $\lambda \neq \mu$ and $\lambda \neq \bar{\mu}$, it follows $\lim_{k \rightarrow \infty} \langle Ax_k | y_k \rangle = 0$. $\square$

Corollary 2.19. *Let m, n be positive integers, $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$ with $0 \notin \sigma_{ap}(A)$. Then the following statements holds.*

- (1) If T is an (A, m) -isometry, then $\partial\sigma(T) \subset \partial\mathbf{D}$ and either $\sigma(T) \subset \partial\mathbf{D}$ or $\sigma(T) = \mathbf{D}$.
- (2) If T is an (A, n) -symmetry, then $\sigma(T) \subset \mathbb{R}$.
- (3) If T is (A, m) -isometric or (A, n) -symmetric, then
 - (a) $\sigma_p(T) \subset \sigma_p(T^*)$.
 - (b) $\sigma_{ap}(T) \subset \sigma_{ap}(T^*)$.
- (4) If T is an $(A, (m, n))$ -isosymmetry and either $\sigma(T) \subset \{e^{it} \in \partial\mathbf{D} : 0 < t < \pi\}$ or $\sigma(T) \subset \{e^{it} \in \partial\mathbf{D} : \pi < t < 2\pi\}$, then T is an (A, m) -isometry.
- (5) If T is an $(A, (m, n))$ -isosymmetry and either $\sigma(T) \subset \{t \in \mathbb{R} : |t| > 1\}$ or $\sigma(T) \subset \{t \in \mathbb{R} : |t| < 1\}$, then T is an (A, n) -symmetry.

An operator $T \in \mathcal{B}(H)$ is said to have the single valued extension property at λ (abbreviated SVEP at λ) if for every open set D containing λ the only analytic function $f : D \rightarrow H$ which

satisfies the equation

$$(2.17) \quad (T - \lambda)f(\lambda) = 0 \quad \text{on } D$$

is the constant function $f \equiv 0$. T has the SVEP if T has the SVEP at every $\lambda \in \mathbb{C}$ ([14]).

Theorem 2.20. *Let m, n be positive integers and let $A \in \mathcal{B}(H)^+$ with $0 \notin \sigma_p(A)$. Then, every $(A, (m, n))$ -isosymmetric operator has the SVEP.*

Proof. Assume that $f : D \rightarrow H$ is any analytic function on D such that (2.17) holds true. From (2.17), it follows that $(T^k - \lambda^k)f(\lambda) = 0$ on D for all positive integer k . This implies that

$$\begin{aligned} 0 = \langle \Omega_A^{m,n}(T)f(\lambda) | f(\lambda) \rangle &= \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} \langle \mathcal{S}_A^n(T)T^k f(\lambda) | T^k f(\lambda) \rangle \\ &= \sum_{k=0}^m (-1)^{m-k} \binom{m}{k} |\lambda|^{2k} \langle \mathcal{S}_A^n(T)f(\lambda) | f(\lambda) \rangle \\ &= (1 - |\lambda|^2)^m \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} \langle A^{\frac{1}{2}}T^{n-j}f(\lambda) | A^{\frac{1}{2}}T^j f(\lambda) \rangle \\ &= (1 - |\lambda|^2)^m \sum_{j=0}^n (-1)^{n-j} \binom{n}{j} \lambda^{n-j} \bar{\lambda}^j \langle A^{\frac{1}{2}}f(\lambda) | A^{\frac{1}{2}}f(\lambda) \rangle \\ &= (1 - |\lambda|^2)^m (2\Im m \lambda)^n \|A^{\frac{1}{2}}f(\lambda)\|^2. \end{aligned}$$

This completes the proof. □

3. SUM WITH A NILPOTENT OPERATOR

The paper [7] proves that, for positive integers m, n , $(T + Q)$ is $(m + 2r - 2, n + 2r - 1)$ -isosymmetric if T is an (m, n) -isosymmetry, Q is a nilpotent operator with order r , and Q doubly commutes with T (i.e $TQ = QT$ and $TQ^* = Q^*T$). The authors, in [7], shows this property for (m, n, C) -isosymmetric operators, with C a conjugation on a Hilbert space and T and S are C -doubly commuting (i.e $TQ = QT$ and $CTCQ^* = Q^*CTC$). Theorem 3.2 extend this result to the sum of an $(A, (m, n))$ -isosymmetry and a nilpotent operator.

Lemma 3.1. *Let $T, S \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. If T and S are doubly commuting, then the following two operator identities hold*

$$(3.1) \quad \Omega_A^{m,n}(T+S) = \sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i+l+h=m} (-1)^k \binom{n}{k} \binom{n-k}{j} \binom{m}{i, l, h} (T^* + S^*)^i S^{*l+j} \Omega_A^{h, n-k-j}(T) T^l S^{i+k}.$$

$$(3.2) \quad \Lambda_A^{m,n}(T+S) = \sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i+l+h=m} \binom{n}{k} \binom{n-k}{j} \binom{m}{i, l, h} (T^* + S^*)^i S^{*l+j} \Lambda_A^{h, n-k-j}(T) T^l S^{i+k}.$$

Proof. The multinomial theorem gives

$$\begin{aligned}
& ((y_1 + y_2)(x_1 + x_2) - 1)^m \{((y_1 + y_2) - (x_1 + x_2))^n\}_{\mathbf{a}} \\
&= ((y_1 x_1 - 1) + (y_1 + y_2)x_2 + y_2 x_1)^m \{((y_1 - x_1) + (y_2 - x_2))^n\}_{\mathbf{a}} \\
&= \left(\sum_{i+l+h=m} \binom{m}{i, l, h} (y_1 + y_2)^i y_2^l (y_1 x_1 - 1)^h x_1^l x_2^i \right) \\
&\quad \cdot \left\{ \sum_{k=0}^n \sum_{j=0}^{n-k} (-1)^k \binom{n}{k} \binom{n-k}{j} y_2^j (y_1 - x_1)^{(n-k-j)} x_2^k \right\}_{\mathbf{a}} \\
&= \left(\sum_{i+l+h=m} (y_1 + y_2)^i y_2^l (y_1 x_1 - 1)^h x_1^l x_2^i \right) \\
&\quad \cdot \left(\sum_{k=0}^n \sum_{j=0}^{n-k} (-1)^k \binom{n}{k} \binom{n-k}{j} y_2^j \left\{ (y_1 - x_1)^{(n-k-j)} \right\}_{\mathbf{a}} x_2^k \right) \\
&= \sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i+l+h=m} (-1)^k \binom{n}{k} \binom{n-k}{j} \binom{m}{i, l, h} (y_1 + y_2)^i y_2^{l+j} (y_1 x_1 - 1)^h \left\{ (y_1 - x_1)^{(n-k-j)} \right\}_{\mathbf{a}} x_1^l x_2^{i+k}.
\end{aligned}$$

On the other hand, we have

$$\begin{aligned}
& ((y_1 + y_2)(x_1 + x_2) - 1)^m \{((y_1 + y_2) - (x_1 + x_2))^n\}_{\mathbf{a}} \\
&= ((y_1 x_1 - 1) + (y_1 + y_2)x_2 + y_2 x_1)^m \{((y_1 + x_1) + y_2 + x_2)^n\}_{\mathbf{a}} \\
&= \sum_{k=0}^n \sum_{j=0}^{n-k} \sum_{i+l+h=m} \binom{n}{k} \binom{n-k}{j} \binom{m}{i, l, h} (y_1 + y_2)^i y_2^{l+j} (y_1 x_1 - 1)^h \left\{ (y_1 + x_1)^{(n-k-j)} \right\}_{\mathbf{a}} x_1^l x_2^{i+k}.
\end{aligned}$$

If T and S are doubly commuting, then replacing x_1 by T , x_2 by S , y_1 by T^* , y_2 by S^* and $\mathbf{a}$ by A , we obtain (3.1) and (3.2). $\square$

Theorem 3.2. *Let $A \in \mathcal{B}(H)^+$, $T \in \mathcal{B}(H)$ and Q is a nilpotent operator of order r . If T and Q are doubly commuting, then the following statements hold.*

- (1) *If T is $(A, (m, n))$ -isosymmetric, then $(T+Q)$ is $(A, (m+2r-2, n+2r-1))$ -isosymmetric.*
- (2) *If T is skew $(A, (m, n))$ -isosymmetric, then $(T+Q)$ is skew $(A, (m+2r-2, n+2r-1))$ -isosymmetric.*

Proof. (1) By (3.1),

$$\begin{aligned}
\Omega_A^{m+2r-2, n+2r-1}(T+Q) &= \sum_{k=0}^{n+2r-1} \sum_{j=0}^{n+2r-1-k} \sum_{i+l+h=m+2r-2} (-1)^k \binom{n+2r-1}{k} \binom{n+2r-1-k}{j} \\
&\quad \binom{m+2r-2}{i, l, h} (T^* + Q^*)^i Q^{*l+j} \Omega_A^{h, n+2r-1-k-j}(T) T^l Q^{i+k},
\end{aligned}$$

Note that if $(k+j) \geq 2r$ or $i \geq r$ or $l \geq r$, then $(T^* + Q^*)^i Q^{*l+j} \Omega_A^{h, n+2r-1-k-j}(T) T^l Q^{i+k} = 0$ since either $Q^{*l+j} = 0$ or $Q^{i+k} = 0$. On the other hand, if $(k+j) \leq 2r-1$, then

$$n + 2r - 1 - (k + j) \geq n + 2r - 1 - (2r - 1) = n,$$

and, if $i \leq r-1$ and $l \leq r-1$, then

$$h = m + 2r - 2 - i - l \geq m + 2r - 2 - (r-1) - (r-1) \geq m.$$

Hence, T is an $(A, (h, n+2r-1-k-j))$ -isosymmetry and $\Omega_A^{h, n+2r-1-k-j}(T) = 0$. Therefore, $\Omega_A^{m+2r-2, n+2r-1}(T+Q) = 0$ and $(T+Q)$ is $(A, (m+2r-2, n+2r-1))$ -isosymmetric.

(2) By (3.2),

$$\begin{aligned} \Lambda_A^{m+2r-2, n+2r-1}(T+Q) &= \sum_{k=0}^{n+2r-1} \sum_{j=0}^{n+2r-1-k} \sum_{i+l+h=m+2r-2} \binom{n+2r-1}{k} \binom{n+2r-1-k}{j} \\ &\quad \binom{m+2r-2}{i, l, h} (T^* + Q^*)^i Q^{*l+j} \Lambda_A^{h, n+2r-1-k-j}(T) T^l Q^{i+k}. \end{aligned}$$

By using this identity and arguing in the same way as in the first statement we complete the proof. $\square$

Theorem 3.2 enjoys a number of interesting consequences that we now state. Some of these results have appeared in the literature with different proofs.

Corollary 3.3. *Let $T, Q \in \mathcal{B}(H)$ satisfying $TQ = QT$ and Q is r -nilpotent and fix an integer $l \geq 1$. Then, it holds:*

- (1) *If T is (A, m) -isometric, then $(T+Q)^l$ is $(A, m+2r-2)$ -isometric.*
- (2) *If T is (A, n) -symmetric, then $(T+Q)^l$ is $(A, n+2r-1)$ -symmetric.*
- (3) *If T is skew (A, n) -symmetric, then $(T+Q)^l$ is skew $(A, n+2r-1)$ -symmetric (l odd).*

Corollary 3.4. *Let $A \in \mathcal{B}(H)^+$ and $R, S \in \mathcal{B}(H)$. Assume that R and S are doubly commuting. If R is $(A, (m, n))$ -isosymmetric, then the operator $\mathbf{K} = \begin{pmatrix} R & S \\ 0 & R \end{pmatrix}$ on $\mathbf{H} = H \oplus H$ is $(A \oplus A, (m+2, n+3))$ -isosymmetric.*

Proof. Put $\mathbf{T} = \begin{pmatrix} R & 0 \\ 0 & R \end{pmatrix}$ and $\mathbf{N} = \begin{pmatrix} 0 & S \\ 0 & 0 \end{pmatrix}$. It follows that $\mathbf{A}$ is positive on $\mathbf{H}$, $\mathbf{K} = \mathbf{T} + \mathbf{N}$, $\mathbf{N}$ is 2-nilpotent and $\mathbf{T}$ and $\mathbf{N}$ are doubly commuting. Applying Theorem 3.2, we deduce that $\mathbf{K}$ is $(\mathbf{A}, (m+2, n+3))$ -isosymmetric. $\square$

4. A CHARACTERIZATION OF (2×2) UPPER TRIANGULAR OPERATOR MATRICES IN $(A; (1, 1))$

For $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$, let consider the following identities:

$$(4.1) \quad \Omega_A^{1,1}(T) = T^* \mathcal{I}_A^1(T) - \mathcal{I}_A^1(T) T = T^{*2} A T - T^* A T^2 - T^* A + A T,$$

$$(4.2) \quad \Lambda_A^{1,1}(T) = \mathcal{I}_A^1(T) T + T^* \mathcal{I}_A^1(T) T = T^{*2} A T + T^* A T^2 - T^* A - A T.$$

It holds, from (4.1) and (4.2), that

$$(4.3) \quad \Omega_A^{1,1}(T) + \Lambda_A^{1,1}(T) = 2(T^{*2} A T - T^* A) = 2T^* \mathcal{I}_A^1(T),$$

Note that $\Omega_A^{1,1}(T)$ is skew symmetric and $\Lambda_A^{1,1}(T)$ is symmetric. An operator T is said to be A -isosymmetric (resp. skew A -isosymmetric) if $\Omega_A^{1,1}(T) = 0$ (resp. $\Lambda_A^{1,1}(T) = 0$).

Example 4.1.

(1) Let $H = \mathbb{C}^2$, $T = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ ($a, b, c, d \in \mathbb{R}$) and $A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$. It holds that,

$$\begin{aligned} \Omega_T^{1,1}(T) &= \begin{pmatrix} 0 & (b-c)(cb-ad+1) \\ (b-c)(ad-bc-1) & 0 \end{pmatrix}, \\ \Omega_A^{1,1}(T) &= \begin{pmatrix} 0 & -c(bc-ad+1) \\ -c(ad-bc-1) & 0 \end{pmatrix}. \end{aligned}$$

- (a) T is I -isosymmetric if and only if $(b = c \text{ or } ad - bc = 1)$.
- (b) T is A -isosymmetric if and only if $(c = 0 \text{ or } ad - bc = 1)$.

(2) Let $H = \mathbb{C}^2$, $T = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$ ($a, b, d \in \mathbb{R}$) and $A = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$. It holds from (4.3) that,

$$\begin{aligned}\Lambda_I^{1,1}(T) &= \begin{pmatrix} 2a(a^2 - 1) & b(2a^2 - 1 + ad) \\ b(2a^2 + ad - 1) & 2(b^2(a + d) + d^3 - d) \end{pmatrix}, \\ \Lambda_A^{1,1}(T) &= \begin{pmatrix} 0 & 0 \\ 0 & 2d(1 - d^2) \end{pmatrix}.\end{aligned}$$

- (a) T is skew I -isosymmetric if and only if one of the following situations holds true
- (i) $a = 0$, $b = 0$ and $d = \pm 1$.
 - (ii) $a = 1$, $b = 0$ and ($d = 0$ or $d = \pm 1$).
 - (iii) $a = 1$ and $d = -1$.
 - (iv) $a = -1$, $b = 0$ and ($d = 0$ or $d = \pm 1$).
 - (v) $a = -1$ and $d = 1$.
- (b) T is skew A -isosymmetric if and only if ($d = 0$ or $d = \pm 1$).

Remark 4.2. Let $A \in \mathcal{B}(H)^+$ and $T \in \mathcal{B}(H)$.

- (1) T is an A -isosymmetry (resp. a skew A -isosymmetry) if and only if $-T$ is an A -isosymmetry (resp. a skew A -isosymmetry).
- (2) If T is A -expansive, then: T is A -isosymmetric (resp. skew A -isosymmetric) if and only if T is $\mathcal{I}_A^1(T)$ -symmetric (resp. skew $\mathcal{I}_A^1(T)$ -symmetric).
- (3) The following statements are equivalent
 - (a) T is A -isosymmetric.
 - (b) $T^* \mathcal{S}_A^1(T) T = \mathcal{S}_A^1(T)$.
 - (c) $T^* \mathcal{I}_A^1(T) = \mathcal{I}_A^1(T) T$.
- (4) For an invertible operator A , let

$$R := \frac{1}{2}(T + A^{-1}T^*A), \quad S := \frac{1}{2}(T - A^{-1}T^*A).$$

Then, R is A -isosymmetric, S is skew A -isosymmetric and $T = R + S$.

In the following results, we consider a class of (2×2) upper triangular operator matrices. We aim to extend [18, Lemma 4.10, Lemma 4.27], [6, Theorem 2.10, Corollary 2.11, Theorem 2.12] established for isosymmetries and complex isosymmetries, respectively, to our developed approach about A -isosymmetries.

Theorem 4.3. Let $A \in \mathcal{B}(H)^+$, $N, E, X \in \mathcal{B}(H)$ and $\mathbf{T} = \begin{pmatrix} N & E \\ 0 & X \end{pmatrix}$ on $\mathbf{H} = H \oplus H$. Then, the following statements hold.

- (1) Assume that $E = NEX$, $N^*AE = AEX$ and N is A -isometric. Then, $\mathbf{T}$ is A -isosymmetric if and only if X is A -isosymmetric.
- (2) Assume that N is A -symmetric and $NE = EX$. Then, $\mathbf{T}$ is A -isosymmetric if and only if X is A -isosymmetric and $AE = ANEX$ holds.

Proof. (1) Since $A \in \mathcal{B}(H)^+$, we have $\mathbf{A} = A \oplus A \in \mathcal{B}(H \oplus H)^+$. On the other hand,

$$\mathcal{I}_A^1(\mathbf{T}) = \begin{pmatrix} \mathcal{I}_A^1(N) & N^*AE \\ E^*AN & E^*AE + \mathcal{I}_A^1(X) \end{pmatrix}.$$

It holds from that

$$(4.4) \quad \mathbf{T}^* \mathcal{I}_A^1(\mathbf{T}) = \begin{pmatrix} N^* \mathcal{I}_A^1(N) & N^{*2}AE \\ E^* \mathcal{I}_A^1(N) + X^* E^* AN & E^* N^* AE + X^* E^* AE + X^* \mathcal{I}_A^1(X) \end{pmatrix},$$

$$(4.5) \quad \mathcal{I}_A^1(\mathbf{T}) \mathbf{T} = \begin{pmatrix} \mathcal{I}_A^1(N)N & \mathcal{I}_A^1(N)E + N^*AEX \\ E^*AN^2 & E^*ANE + E^*AEX + \mathcal{I}_A^1(X)X \end{pmatrix}.$$

By equations (4.4) and (4.5), $\mathbf{T}$ is A -isosymmetric if and only if

$$(4.6) \quad \begin{cases} N^* \mathcal{I}_A^1(N) = \mathcal{I}_A^1(N)N, \\ E^* \mathcal{I}_A^1(N) + X^* E^* AN = E^* AN^2, \\ N^{*2} AE = \mathcal{I}_A^1(N)E + N^* AEX, \\ (E^* N^* AE + X^* E^* AE) + X^* \mathcal{I}_A^1(X) = (E^* ANE + E^* AEX) + \mathcal{I}_A^1(X)X. \end{cases}$$

Assume that N is A -isometric, that is $\mathcal{I}_A^1(N) = N^* AN - A = 0$. It holds from that

$$(4.7) \quad \begin{cases} X^* E^* AN = E^* AN^2, \\ N^{*2} AE = N^* AEX, \\ (E^* N^* AE + X^* E^* AE) + X^* \mathcal{I}_A^1(X) = (E^* ANE + E^* AEX) + \mathcal{I}_A^1(X)X. \end{cases}$$

If $E = NEX$ and $N^* AE = AEX$, then the first and the second equation of (4.7) hold. Moreover, we have

$$\begin{aligned} E^* ANE + E^* AEX &= X^* E^* \underbrace{N^* AN}_E E + E^* N^* AE \\ &= X^* E^* AE + E^* N^* AE. \end{aligned}$$

The above equation gives

$$X^* \mathcal{I}_A^1(X) = \mathcal{I}_A^1(X)X.$$

Hence, X is A -isosymmetric. Arguing as above, we can show the converse of the claim.

- (2) Assume that N is A -symmetric, that is $\mathcal{S}_A^1(N) = N^* A - AN = 0$ and $\mathbf{T}$ is A -isosymmetric. From the assumptions, it holds $X^* E^* A = E^* N^* A = E^* AN$, and so $N^* AE = AEX$. Hence, it follows

$$(4.8) \quad \begin{cases} X^* E^* AE = E^* ANE, \\ E^* N^* AE = E^* AEX. \end{cases}$$

From (4.6) and (4.8), we obtain

$$E^* N^* AE + X^* E^* AE = E^* ANE + E^* AEX.$$

So, $X^* \mathcal{I}_A^1(X) = \mathcal{I}_A^1(X)X$. Hence, X is A -isosymmetric. On the other hand, (4.6) becomes

$$\begin{cases} E^*(N^{*2} - I)A + X^* E^* N^* A = E^* N^{*2} A, \\ AN^2 E = A(N^2 - I)E + ANEX. \end{cases}$$

This gives that

$$\begin{cases} E^* A = X^* E^* N^* A, \\ AE = ANEX. \end{cases}$$

By similar arguments, we can show the other implication. □

Corollary 4.4. *Let $A \in \mathcal{B}(H)^+$, $N, E, X \in \mathcal{B}(H)$ and $\mathbf{T} = \begin{pmatrix} N & E \\ 0 & X \end{pmatrix}$ on $\mathbf{H} = H \oplus H$ with N an A -isometry. If $N^* AE = 0$, then $\mathbf{T}$ is A -isosymmetric.*

Proof. Since N is an A -isometry and $N^* AE = 0$, it holds from (4.7) that

$$X^*(E^* AE + X^* AX - A) = (E^* AE + X^* AX - A)X.$$

On the other hand,

$$(T \text{ is } A\text{-isosymmetric}) \iff (X^*(E^* AE + X^* AX - A) = (E^* AE + X^* AX - A)X).$$

This completes the proof. □

In the following theorem, we describe the block operator form of an A -isosymmetry such that $\mathcal{M} = \ker(\mathcal{I}_A^1(\mathbf{T})) \neq \{0\}$.

Theorem 4.5. Let $T \in \mathcal{B}(H)$ and $A \in \mathcal{B}(H)^+$. Assume that $\mathcal{M}$ is invariant for A and for T . Then T has the form

$$(4.9) \quad T = \begin{pmatrix} N & E \\ 0 & X \end{pmatrix} \text{ on } \mathcal{M} \oplus \mathcal{M}^\perp, \quad N \in \mathcal{B}(\mathcal{M}), E \in \mathcal{B}(\mathcal{M}^\perp, \mathcal{M}), X \in \mathcal{B}(\mathcal{M}^\perp),$$

such that N is an $A|_{\mathcal{M}}$ -isometry and $E^*A|_{\mathcal{M}}N = 0$.

Proof. Since $\mathcal{M}$ is invariant for A , it follows that $\mathcal{M}^\perp$ is invariant for A . Moreover, since $A \in \mathcal{B}(H)^+$, we have $A_1 = A|_{\mathcal{M}} \geq 0$, $A_2 = A|_{\mathcal{M}^\perp} \geq 0$ and $A = A_1 \oplus A_2$. If $\mathcal{M}$ is invariant for T , then T has the block operator form (4.9). Hence it follows that

$$\mathcal{I}_A^1(T) = \begin{pmatrix} N^*A_1N - A_1 & N^*A_1E \\ E^*A_1N & E^*A_1E + X^*A_2X - A_2 \end{pmatrix} \text{ on } \mathcal{M} \oplus \mathcal{M}^\perp.$$

Since $\mathcal{M} = \ker(\mathcal{I}_A^1(T))$, it holds $\mathcal{I}_A^1(T)(x \oplus 0) = 0$ for $x \in \mathcal{M}$. Moreover, we have $N^*A_1N - A_1 = 0$ and $E^*A_1N = 0$ on $\mathcal{M}$. Hence, N is an A_1 -isometry. $\square$

REFERENCES

- [1] J. Agler, M. Stankus, *m-Isometric Transformations of Hilbert Spaces I*. Integral Equations and Operator Theory **21** (1995), no. 4, 383-429.
- [2] J. Agler, M. Stankus, *m-isometric transformations of Hilbert space II*. Integral Equations and Operator Theory **23** (1995), no. 1, 1-48.
- [3] J. Agler, M. Stankus, *m-isometric transformations of Hilbert space III*. Integral Equations and Operator Theory **24** (1996), no. 4, 379-421.
- [4] M. F. Ahmadi, *Powers of A-m-Isometric Operators and Their Supercyclicity*, Bull. Malays. Math. Sci. Soc. (2016) 39: 901911, DOI 10.1007/s40840-015-0201-6
- [5] M. Chō, E. Ko and J. E. Lee, *Skew m-Complex Symmetric Operators*, Filomat 33:10 (2019), 29752983.
- [6] M. Chō, J. E. Lee, T. Prasad, and K. Tanahashi, *Complex isosymmetric operators*, Adv. Oper. Theory. Volume 3, Number 3 (2018), 620-631.
- [7] M. Chō, J. E. Lee, K. Tanahashi and J. Tomiyama, *On $[m, C]$ -symmetric Operators*, KYUNGPOOK Math. J. 58(2018), 637-650.
- [8] M. Chō and S. A. O. Mahmoud, *(A, m)-Symmetric commuting tuples of operators on a Hilbert Space*, MIA **22** (2019), no. 3, 931-947.
- [9] B. P. Duggal and V. Müller, *Tensor product of left n-invertible operators*, STUD. MATH. 215, (2), (2013).
- [10] J. W. Helton, *Jordan operators in infinite dimensions and Surm Liouville conjugate point theory*, Bul. Amer. Math. Soc **78** (1972), no 1, 57-61.
- [11] J. W. Helton, *Operators with a representation as multiplication by x on a Sobolev space*, Colloquia Math. Soc. Janos Bolyai **5**, Hilbert Space Operators, Tihany, Hungary (1970), 279-287.
- [12] S. Jung, Y. Kim, E. Ko and J. E. Lee, *On (A, m)-expansive operators*, Stud. Math 213 (1) (2012).
- [13] N. Jeridi and R. Rabaoui, *On (A, m)-Symmetric Operators in a Hilbert Space*, Results Math (2019) 74: 124. <https://doi.org/10.1007/s00025-019-1049-0>.
- [14] Y. Kim, E. Ko, and J. Lee, *Operators with the single valued extension property*, Bull. Korean Math. Soc. **43** (2006), no. 3, 509-517.
- [15] T. Le, *Decomposing algebraic m-isometric tuples*, Journal of Functional Analysis. Volume 278, Issue 8, 1 May 2020, 108424.
- [16] Scott A. McCullough and Leiba Rodman, *Hereditary classes of operators and matrices*, Amer. Math. Monthly **104** (1997), no. 5, 415-430.
- [17] R. Rabaoui and A. Saddi, *On the Orbit of an A-m-isometry*, Annales Mathematicae Silesianae, 26 (2012), 75-91.
- [18] M. Stankus, *Isosymmetric linear transformations on complex Hilbert Space*, Thesis (Ph.D.) University of California, San Diego. Proquest LLC, Ann Arbor, MI, 1993. 80 pp.
- [19] M. Stankus, *m-Isometries, n-Symmetries and Other Linear Transformations Which are Hereditary Roots*, Integral Equations and Operator Theory 75(3) (2013), DOI: 10.1007/s00020-012-2026-0.
- [20] O. A. M. Sid Ahmed, A. Saddi, *A-m-Isometric operators in semi-Hilbertian spaces*, Linear Algebra and its Applications, Vol 436, Issue 10 (2012), 3930-3942.
- [21] A. Tokmakoff, *Exponential operators, Time-Dependent Quantum Mechanics and Spectroscopy*, Reference material, Univ. of Chicago, <https://tdqms.uchicago.edu/>

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCES. GABES 6072. TUNISIA, UNIVERSITY OF TUNIS EL MANAR, FACULTY OF SCIENCES OF TUNIS, MATHEMATICAL ANALYSIS AND APPLICATIONS LABORATORY LR11ES11, 2092, EL MANAR I, TUNIS, TUNISIA.

Email address: `rchid.rabaoui@fsg.rnu.tn`