

Very stable Higgs bundles, equivariant multiplicity and mirror symmetry

Tamás Hausel
IST Austria

Nigel Hitchin
University of Oxford

tamas.hausel@ist.ac.at hitchin@maths.ox.ac.uk

February 15, 2021

Abstract

We define and study the existence of very stable Higgs bundles on Riemann surfaces, how it implies a precise formula for the multiplicity of the very stable components of the global nilpotent cone and its relationship to mirror symmetry. The main ingredients are the Bialynicki-Birula theory of \mathbb{C}^* -actions on semiprojective varieties, \mathbb{C}^* characters of indices of \mathbb{C}^* -equivariant coherent sheaves, Hecke transformation for Higgs bundles, relative Fourier-Mukai transform along the Hitchin fibration, hyperholomorphic structures on universal bundles and cominuscule Higgs bundles.

Contents

| | | |
|----------|--|-----------|
| 1 | Introduction | 2 |
| 2 | Bialynicki-Birula partition and decomposition | 9 |
| 2.1 | Bialynicki-Birula theory for semi-projective varieties | 9 |
| 2.2 | Very stable upward flows | 17 |
| 3 | Bialynicki-Birula theory on the moduli space of Higgs bundles | 18 |
| 3.1 | Fixed points of the \mathbb{T} -action $\mathcal{M}^{s\mathbb{T}}$ | 19 |
| 3.2 | Upward flows on \mathcal{M} | 21 |
| 3.3 | Downward flows on \mathcal{M} | 25 |

| | | |
|----------|---|-----------|
| 4 | Very stable Higgs bundles | 26 |
| 4.1 | Definition and basic properties | 26 |
| 4.2 | Hecke transformations | 27 |
| 4.2.1 | The Hecke transform of a full compatible filtration | 29 |
| 4.2.2 | Very stable Higgs bundles via Hecke transformations | 31 |
| 4.2.3 | Example in rank 2 | 34 |
| 4.2.4 | Hecke transforms of Lagrangians | 35 |
| 5 | Multiplicities in the nilpotent cone | 39 |
| 5.1 | Multiplicities of very stable components | 39 |
| 5.2 | Intersection of upward flows with generic fibres of h | 48 |
| 6 | Mirror of upward flows | 53 |
| 6.1 | Jacobians, duality, Fourier-Mukai transform | 53 |
| 6.2 | Fourier-Mukai transform of $\mathcal{O}_{W_\varepsilon^+}$ | 56 |
| 6.3 | An agreement for \mathbb{T} -equivariant Euler forms | 62 |
| 6.3.1 | \mathbb{T} -equivariant Euler characteristics and Euler forms | 63 |
| 6.3.2 | \mathbb{T} -equivariant universal bundles | 63 |
| 7 | A hyperholomorphic structure on \mathbb{E}_c | 69 |
| 7.1 | Hyperholomorphic connections | 69 |
| 7.2 | The connection on a universal bundle | 71 |
| 7.3 | The universal Higgs field | 73 |
| 8 | Further issues | 74 |
| 8.1 | Simple Lie groups | 74 |
| 8.2 | Multiple zeros | 78 |
| 8.3 | The cotangent fibre | 81 |

1 Introduction

Drinfeld [Dr1] and Laumon [La1] call a vector bundle on a smooth complex projective curve C of genus g a *very stable bundle* if there is no non-zero nilpotent Higgs field $\Phi \in H^0(C; \text{End}(E) \otimes K)$ on E . Laumon [La1, Proposition 3.5] proved that a very stable bundle is semi-stable for $g > 0$ and stable for $g > 1$ in the sense of Mumford and that very stable bundles form a non-empty open subset in the moduli space of stable bundles on C .

We will assume $g > 1$, $d \in \mathbb{Z}$ and $n \in \mathbb{Z}_{>1}$. Let $\mathcal{M} := \mathcal{M}_{n,d}$ denote the moduli space of semi-stable rank n degree d Higgs bundles on C i.e. pairs (E, Φ) with $\Phi \in$

$H^0(C; \text{End}(E) \otimes K)$. It was constructed in Hitchin [Hi5] by gauge theoretic and in [Ni, Si2] by algebraic geometric techniques. Pauly and Peon-Nieto [PPN] proved that E is very stable if and only if

$$U_E = \{(E, \Phi) \mid \Phi \in H^0(C; \text{End}(E) \otimes K)\} \subset \mathcal{M} \quad (1.1)$$

is closed in \mathcal{M} .

There is a generalization of $U_E \subset \mathcal{M}$ for more general Higgs bundles. Recall that there is a \mathbb{T} -action on \mathcal{M} by $(E, \Phi) \mapsto (E, \lambda\Phi)$ for $\lambda \in \mathbb{T}$. For a stable Higgs bundle $\mathcal{E} = (E, \Phi) \in \mathcal{M}^{s\mathbb{T}}$ fixed by the \mathbb{T} -action we define its *upward flow* as

$$W_{\mathcal{E}}^+ := \{(E', \Phi') \in \mathcal{M} \mid \lim_{\lambda \rightarrow 0} (E', \lambda\Phi') = (E, \Phi)\}. \quad (1.2)$$

In particular, for a rank n degree d vector bundle E we have $W_{(E,0)}^+ = U_E$ from (1.1).

We define a \mathbb{T} -fixed stable Higgs bundle $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ to be a *very stable Higgs bundle* if the only Higgs bundle $(E', \Phi') \in W_{\mathcal{E}}^+$ with nilpotent Higgs field is \mathcal{E} . Our first result is a generalization of [PPN, Theorem 1.1]

Theorem 1.1. $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable if and only if $W_{\mathcal{E}}^+ \subset \mathcal{M}$ is closed.

From Laumon's result we know that a generic rank n vector bundle is very stable, thus there always exists a very stable Higgs bundle of the form $(E, 0)$ with E very stable, that is, very stable Higgs bundles in the subvariety $\mathcal{N} \in \pi_0(\mathcal{M}^{\mathbb{T}})$ parametrizing semi-stable vector bundles on C .

There are however many other fixed point components. Namely, when $E = E_1 \oplus \dots \oplus E_j$ is a direct sum of vector bundles and $\Phi|_{E_i} \rightarrow E_{i+1}K \subset EK$. We call

$$(\text{rank}(E_1), \text{rank}(E_2), \dots, \text{rank}(E_j))$$

– an ordered partition of n – the *type* of the \mathbb{T} -invariant semistable Higgs bundle (E, Φ) . For example, for a rank n semistable bundle E the semistable Higgs bundle $(E, 0) \in \mathcal{M}^{\mathbb{T}}$ is of type (n) .

In this paper we classify all type $(1, \dots, 1)$ very stable Higgs bundles:

Theorem 1.2. *Let*

$$E = L_0 \oplus \dots \oplus L_{n-1}$$

be a direct sum of line bundles L_i on C . Furthermore, let the Higgs field be given by

$$\Phi|_{L_{i-1}} = b_i : L_{i-1} \rightarrow L_i K \subset EK$$

where

$$b_i \in \text{Hom}(L_{i-1}, L_i K) = H^0(L_{i-1}^* L_i K).$$

Assume that (E, Φ) is stable. Then (E, Φ) is very stable if and only if

$$b = b_{n-1} \circ \cdots \circ b_1 \in H^0(L_0^* L_{n-1} K)$$

has no repeated zero.

The proof of this proceeds by first verifying it in the case when b has no zeroes. The corresponding Higgs bundle is called a uniformising Higgs bundle, the upward flow from which is nothing but a Hitchin section [Hi3], which was already observed to be closed in [Hi3]. In particular when $L_0 \cong \mathcal{O}_C$ then we get the *canonical uniformising Higgs bundle*

$$\mathcal{E}_0 := (\mathcal{O}_C \oplus K^{-1} \oplus \cdots \oplus K^{-n+1}, \Phi|_{K^{-i+1}} = 1 : K^{-i+1} \rightarrow K^{-i} \otimes K). \quad (1.3)$$

Then we apply Hecke transformations at points of the curve to these Hitchin sections, and observe that we can reach all type $(1, \dots, 1)$ upward flows with subsequent Hecke transformations, and can infer Theorem 1.2 in turn. This argument is detailed in subsection 4.2.

From Theorem 1.2 we deduce that in every type $(1, \dots, 1)$ component $F \in \pi_0(\mathcal{M}^\mathbb{T})$ there exists a very stable Higgs bundle. Thus when $n = 2$ every component, being either of type (2) or type (1, 1), contains very stable Higgs bundles. However, already for rank $n = 3$ there are type (1, 2) components $F \in \pi_0(\mathcal{M}^\mathbb{T})$ which contain no very stable Higgs bundles. This can be seen by computing the multiplicity of a very stable component of the nilpotent cone, which we now explain.

We recall the Hitchin map [Hi2]

$$\begin{aligned} h : \mathcal{M} &\rightarrow \mathcal{A} = \bigoplus_{i=1}^n H^0(C; K^i) \\ (E, \Phi) &\mapsto \det(\Phi - x) \end{aligned}$$

given by the characteristic polynomial. It is a proper completely integrable system [Ni, Hi2] with respect to a natural symplectic structure on \mathcal{M} . In particular, $\dim(\mathcal{A}) = \dim(\mathcal{M})/2$ and the fibres of h are Lagrangian subvarieties. The generic fibre is isomorphic to an Abelian variety – the Jacobian of a spectral curve – while the most singular fibre $N := h^{-1}(0)$, the so-called *nilpotent cone*, is non-reduced as a subscheme. Its components can be understood from the *core* of \mathcal{M} . Namely, for $F \in \pi_0(\mathcal{M}^\mathbb{T})$ one defines the *downward flow*

$$W_F^- := \{(E', \Phi') \in \mathcal{M} \mid \lim_{\lambda \rightarrow \infty} (E', \lambda \Phi') \in F\}.$$

Note that the Hitchin map is proper and \mathbb{T} -equivariant with an appropriate weighted \mathbb{T} -action on \mathcal{A} . It follows that the *core*

$$\mathcal{C} := \bigcup_{F \in \pi_0(\mathcal{M}^\mathbb{T})} W_F^- \cong N_{red}$$

is a proper subvariety isomorphic to N_{red} the reduction of the nilpotent cone. Thus each component N_F of the nilpotent cone N is labelled by $F \in \pi_0(\mathcal{M}^{\mathbb{T}})$ and its reduction is isomorphic to $N_{F_{red}} \cong \overline{W_F^-}$. We denote by $m_F := \ell(\mathcal{O}_{N, N_F})$ the multiplicity of the component $N_F \subset N$. We have the following

Theorem 1.3. *Let $\mathcal{E} \in \mathcal{M}^{\mathbb{T}}$ be a very stable Higgs bundle. Denote by $T_{\mathcal{E}}^+ \subset T_{\mathcal{E}}\mathcal{M}$ the positive part of the \mathbb{T} -module $T_{\mathcal{E}}\mathcal{M}$ and by $F_{\mathcal{E}}$ the \mathbb{T} -fixed point component containing \mathcal{E} . We have*

1. *the multiplicity of the corresponding component $N_{F_{\mathcal{E}}}$ of the nilpotent cone equals*

$$m_{F_{\mathcal{E}}} = \frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))} \Big|_{t=1}, \quad (1.4)$$

2. *for generic $a \in \mathcal{A}$ the intersection $W_{\mathcal{E}}^+ \cap h^{-1}(a)$ is transversal and has cardinality $m_{F_{\mathcal{E}}}$.*

Here for a positive \mathbb{T} -module V with weight space decomposition $V = \bigoplus_{\lambda > 0} V_{\lambda}$ we have denoted the \mathbb{T} -character of its symmetric algebra by

$$\chi_{\mathbb{T}}(\mathrm{Sym}(V^*)) = \sum_{\lambda \leq 0} \dim(\mathrm{Sym}(V^*)_{\lambda}) t^{-\lambda} = \prod_{\lambda > 0} \frac{1}{(1 - t^{\lambda})^{\dim(V_{\lambda})}} \in \mathbb{Z}[[t]], \quad (1.5)$$

where t denotes the \mathbb{T} -character of weight -1 . The computation of (1.4) is straightforward and when it is non-integer valued we deduce the non-existence of very stable Higgs bundles in a given component. This is the case for certain types of (1, 2) components in rank 3, see Remark 5.12.

It is straightforward to compute (1.4) for a very stable Higgs bundle. Consequently, we deduce the multiplicity of type (n) and type $(1, \dots, 1)$ components of the nilpotent cone.

Corollary 1.4. *The multiplicity of \mathcal{N} , the moduli space of semi-stable rank n degree d bundles, in the nilpotent cone is*

$$m_{\mathcal{N}} = 2^{3g-3} 3^{5g-5} \dots n^{(2n-1)(g-1)}.$$

Moreover let $N_{F_{\mathcal{E}}}$ be the component of the nilpotent cone for a type $(1, \dots, 1)$ Higgs bundle $\mathcal{E} \in \mathcal{M}^{\mathbb{T}}$ as in Theorem 1.2. Let $m_i = \deg(b_i)$. Then its multiplicity in the nilpotent cone is

$$m_{F_{\mathcal{E}}} = \prod_{i=1}^{n-1} \binom{n}{n-i}^{m_i}.$$

The main motivation of this paper was to try to understand the mirror of upward flows (1.2) in the framework of Kapustin and Witten [KW]. Fixing $\mathcal{M} := \mathcal{M}_{n, -n(n-1)(g-1)}$ we expect that the mirror of the structure sheaf $\mathcal{O}_{W_{\mathcal{E}}^+}$ of the Lagrangian upward flow

$W_{\mathcal{E}}^+ \subset \mathcal{M}$ will be a hyperholomorphic vector bundle on \mathcal{M} if and only if \mathcal{E} is very stable, in which case we expect the mirror to be of rank $m_{F_{\mathcal{E}}}$.

More precisely, for any $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ we introduce the *virtual multiplicity*

$$m_{\mathcal{E}}(t) := \frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))} \quad (1.6)$$

which is a priori a rational function in t . In the course of proving Theorem 1.3 we find that for a very stable \mathcal{E} the pushforward $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})$ is a \mathbb{T} -equivariant vector bundle on \mathcal{A} , with \mathbb{T} -equivariant fibre at 0 satisfying $\chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0) = m_{\mathcal{E}}(t)$. In particular, for a very stable \mathcal{E} the quantity $m_{\mathcal{E}}(t)$ is a polynomial in t , which we call the *equivariant multiplicity*.

Its significance for mirror symmetry is that we expect (see Remark 6.12) that the mirror, denoted $\Lambda_{\mathcal{E}}$, of a very stable upward flow $\mathcal{O}_{W_{\mathcal{E}}^+}$ should be a \mathbb{T} -equivariant vector bundle on \mathcal{M} whose restriction to the canonical Hitchin section is isomorphic to $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})$ and therefore its fibre at the canonical uniformising Higgs bundle \mathcal{E}_0 should have \mathbb{T} -character

$$\chi_{\mathbb{T}}(\Lambda_{\mathcal{E}}|_{\mathcal{E}_0}) = m_{\mathcal{E}}(t). \quad (1.7)$$

In particular, its rank should be

$$\mathrm{rank}(\Lambda_{\mathcal{E}}) = \dim(\Lambda_{\mathcal{E}}|_{\mathcal{E}_0}) = m_{\mathcal{E}}(1) = m_{F_{\mathcal{E}}}$$

the multiplicity of $N_{F_{\mathcal{E}}}$ in the nilpotent cone N .

In the case of type $(1, \dots, 1)$ fixed points, where we have a complete classification of very stable Higgs bundles in Theorem 1.2, we find a candidate for the mirror, which is supported by the following results sketched below. For details see Section 6.

Let $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1})$ with $\delta_0 := c_{01} + \dots + c_{0m_0}$ a divisor of degree ℓ , where we assume that either $c_{0j} \in C$ or $-c_{0j} \in C$. For $i > 0$ we have the effective divisor $\delta_i = c_{i1} + \dots + c_{im_i} \in C^{[m_i]}$ of degree $m_i \geq 0$, where $c_{ij} \in C$ are points. This data gives rise to a type $(1, \dots, 1)$ Higgs bundle as in Theorem 1.2 by setting

$$L_i := (\delta_0 + \delta_1 + \dots + \delta_i)K^{-i}$$

and

$$b_i := s_{\delta_i} \in H^0(C; L_{i-1}^* L_i K) \cong H^0(C; \mathcal{O}_C(\delta_i))$$

the defining section of $\mathcal{O}_C(\delta_i)$. We denote this Higgs bundle by \mathcal{E}_{δ} . Let us assume that $\mathcal{E}_{\delta} \in \mathcal{M}^{s\mathbb{T}}$ is very stable, i.e. it is stable and the divisor $\delta_1 + \dots + \delta_{n-1}$ is reduced. Then $W_{\mathcal{E}_{\delta}}^+ \subset \mathcal{M}$ is a closed Lagrangian, and we denote its structure sheaf

$$\mathcal{L}_{\delta} := \mathcal{O}_{W_{\mathcal{E}_{\delta}}^+}.$$

We construct

$$\Lambda_\delta := \bigotimes_{i=0}^{n-1} \bigotimes_{j=0}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}), \quad (1.8)$$

using a certain (twisted by a gerbe and appropriately normalised) universal Higgs bundle (\mathbb{E}, Φ) on $C \times \mathcal{M}^s$ and denote $\mathbb{E}_{c_{ij}} = \mathbb{E}|_{\mathcal{M}^s \times \{c_{ij}\}}$ when $c_{ij} \in C$ and $\mathbb{E}_{c_{0j}} = \mathbb{E}^*|_{\mathcal{M}^s \times \{-c_{0j}\}}$, when $-c_{0j} \in C$. When n divides $\sum_{i=1}^{n-1} (n-i)m_i$ (1.8) is a well-defined vector bundle of rank $m_{\mathcal{E}_\delta}$.

For simplicity we assume that

$$d = n\ell + \sum_{i=1}^{n-1} (n-i)m_i - n(n-1)(g-1) = -n(n-1)(g-1).$$

In this case for a generic $a \in \mathcal{A}^\# \subset \mathcal{A}$ we can identify $h^{-1}(a) = J(C_a)$ with the Jacobian of the smooth spectral curve $C_a \subset T^*C$. Define

$$\iota : \begin{array}{ccc} \mathcal{M} & \rightarrow & \mathcal{M} \\ (E, \Phi) & \mapsto & (E^*K^{1-n}, \Phi^T) \end{array},$$

which on $h^{-1}(a)$ will induce the inverse map. In particular, we will have $\iota(\mathcal{E}_\delta) = \mathcal{E}_{\delta^\iota}$ where $\delta^\iota = (\delta_0^\iota, \delta_1^\iota, \dots, \delta_{n-1}^\iota) = (-\delta_{n-1} - \dots - \delta_0, \delta_{n-1}, \dots, \delta_1)$.

For $\mathcal{M}^\# := h^{-1}(\mathcal{A}^\#)$ we construct a relative Poincaré bundle $\tilde{\mathbb{P}}$ on the fibre product $\mathcal{M}^\# \times_{\mathcal{A}^\#} \mathcal{M}^\#$, which has fibre $J(C_a) \times J(C_a)$ over $a \in \mathcal{A}^\#$. We then have the following

Theorem 1.5. *Let $\mathcal{E}_\delta \in \mathcal{M}^{s\mathbb{T}}$ be a very stable Higgs bundle of degree $-n(n-1)(g-1)$. The relative Fourier-Mukai transform over $\mathcal{A}^\#$ satisfies*

$$S(\mathcal{L}_\delta|_{\mathcal{M}^\#}) = (\pi_2)_*(\pi_1^*(\mathcal{L}_\delta|_{\mathcal{M}^\#}) \otimes \tilde{\mathbb{P}}) = \Lambda_\delta|_{\mathcal{M}^\#}$$

and

$$S(\Lambda_\delta|_{\mathcal{M}^\#}) = \mathcal{L}_{\delta^\iota}[-n^2(g-1) - 1]|_{\mathcal{M}^\#}$$

Moreover, Λ_δ extends to a vector bundle with a hyperholomorphic connection on \mathcal{M}^s .

This theorem gives the mirror of \mathcal{L}_δ as a hyperholomorphic vector bundle of rank $m_{F_{\mathcal{E}_\delta}}$. The following will give evidence of the mirror relationship globally, in particular over the nilpotent cone. To motivate it recall that mirror symmetry should be an equivalence of categories. Not just objects should match but morphisms between them. If two objects have a \mathbb{T} -equivariant structure so will the vector space of morphisms between them. Thus its \mathbb{T} -character should agree with the \mathbb{T} -character of the vector space of morphisms between the mirror objects.

To formalise this for two \mathbb{T} -equivariant coherent sheaves \mathcal{F}_1 and \mathcal{F}_2 on \mathcal{M} we denote the *equivariant Euler pairing* as

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{F}_1, \mathcal{F}_2) = \sum_{k,l} \dim(H^k(R\mathcal{H}om(\mathcal{F}_1, \mathcal{F}_2))_l) (-1)^k t^{-l},$$

which on a semi-projective variety like \mathcal{M} we expect to be a Laurent series in $\mathbb{Z}((t))$.

Theorem 1.6. *Let $\mathcal{E}_\delta, \mathcal{E}_{\delta'} \in \mathcal{M}^{s\mathbb{T}}$ be two type $(1, \dots, 1)$ very stable Higgs bundles of degree $-n(n-1)(g-1)$. Then there exist \mathbb{T} -equivariant structures on our coherent sheaves such that*

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'}, \Lambda_\delta) = \chi_{\mathbb{T}}(\det(\mathbb{C}_1 \otimes \mathcal{A}^*)) \chi_{\mathbb{T}}(\mathcal{M}; \Lambda_{\delta'}, \mathcal{L}_{\delta'}[-n^2(g-1) - 1]).$$

Thus up to the q -power $\chi_{\mathbb{T}}(\det(\mathbb{C}_1 \otimes \mathcal{A}^*))$, where \mathbb{C}_1 denotes the weight 1 \mathbb{T} -character, we find that the \mathbb{T} -character of morphisms between our Lagrangian and hyperholomorphic branes agree with that of the conjectured mirror branes.

The content of the paper is as follows. In Section 2.1 we work out from scratch the Bialynicki-Birula theory for a semi-projective variety, to conclude that the upward flows and the core of a semiprojective variety with a homogeneity 1 symplectic form are Lagrangian. Here we introduce very stable upward flows and prove some properties including the general form of Theorem 1.1. In Section 3 we determine which Higgs bundles belong to which upward and downward flows in the moduli space of Higgs bundles. In Section 4 we examine in detail the Hecke transformation of a type $(1, \dots, 1)$ \mathbb{T} -fixed Higgs bundle and apply this to classify all very stable Higgs bundles of this type proving Theorem 1.2. Moreover we show that the Hecke transform takes a Lagrangian subvariety to another in general. In Section 5 we investigate the multiplicity of a very stable component of the nilpotent cone, by deriving an explicit linear algebra formula for it, and study the implications. Here we prove Theorem 1.3 and Corollary 1.4. In Section 6 we study the relative Fourier-Mukai transform of our type $(1, \dots, 1)$ very stable upward flows proving the first part of Theorem 1.5, and prove an agreement of \mathbb{T} -equivariant Euler forms for pairs of mirror branes settling Theorem 1.6. In Section 7 using the twistor space construction of the moduli space of Higgs bundles we endow the universal bundle at a point in C with a hyperholomorphic connection and from the expression (1.8) prove the second part of Theorem 1.5.

In the final Section 8 we discuss three issues relevant to further progress. First in Subsection 8.1 we compute the virtual multiplicity (1.6) for the analogue of the type $(1, \dots, 1)$ fixed points for a simple group G instead of GL_n . This is the case where the \mathbb{T} -action at the fixed point set has the same form as the Hitchin section for the group G as in [Hi3], which is of course a closed upward flow. The b_i which we studied for the general linear group are now replaced by sections b_1, \dots, b_ℓ of line bundles associated to the simple roots of G . We find that unlike the GL_n case considered in the body of this

paper, the virtual multiplicity is rarely a polynomial. We prove in Proposition 8.1 that for *cominuscule* Higgs bundles where b_i are nonvanishing sections except at cominuscule roots it is in fact a polynomial. In this case we offer Conjecture 8.2 that under precise conditions these are very stable and find a potential mirror in terms of the minuscule representations of the Langlands dual group, satisfying the expectation of (1.7). Secondly in Subsection 8.2 we study the first non-trivial wobbly type $(1, 1)$ SL_2 Higgs bundle where b has a single double 0. We find the closure of its upward flow and formulate a conjecture about the mirror of the universal SO_3 bundle in the adjoint representation. Finally in Subsection 8.3 we discuss the problem of the mirror of a very stable cotangent fibre to the moduli of semi-stable vector bundles. We relate this to considerations of Donagi–Pantev and to Drinfeld–Laumon in the geometric Langland correspondence.

Acknowledgements. We would like to thank Brian Collier, Davide Gaiotto, Peter Gothen, Jochen Heinloth, Daniel Huybrechts, Quoc Ho, Joel Kamnitzer, Gérard Laumon, Luca Migliorini, Alexander Minets, Brent Pym, Peng Shan, Carlos Simpson, András Szenes, Fernando R. Villegas, Richard Wentworth, Edward Witten and Kōta Yoshioka for interesting comments and discussions. We would also like to thank the organizers of the Summer School on Higgs bundles in Hamburg in September 2018, where the authors and Richard Wentworth were giving lectures and where the work in this paper started by considering the mirror of the Lagrangian upward flows $W_{\mathcal{E}}^+$ investigated in [CW]. The second author wishes to thank EPSRC and ICMAT for support.

2 Bialynicki-Birula partition and decomposition

2.1 Bialynicki-Birula theory for semi-projective varieties

Let M be a normal, complex, semi-projective algebraic variety. Semi-projective means that M is quasi-projective and we have a $\mathbb{T} := \mathbb{C}^\times$ action on M such that the fixed point locus $M^{\mathbb{T}}$ is projective and for every $x \in M$ there is a $p \in M^{\mathbb{T}}$ such that $\lim_{\lambda \rightarrow 0} \lambda x = p$. By the latter we mean that there exists a \mathbb{T} -equivariant morphism $g : \mathbb{A}^1 \rightarrow M$ with $g(1) = x$ and $g(0) = p$, where \mathbb{T} acts on \mathbb{A}^1 in the standard way.

Examples of semi-projective varieties include cotangent bundles of smooth projective varieties, moduli spaces of Higgs bundles and Nakajima quiver varieties.

For every $\alpha \in M^{\mathbb{T}}$ denote by

$$W_{\alpha}^+ := \{x \in M \mid \lim_{\lambda \rightarrow 0} \lambda x = \alpha\}$$

the *upward flow* from α and

$$W_{\alpha}^- := \{x \in M \mid \lim_{\lambda \rightarrow \infty} \lambda x = \alpha\}$$

the *downward flow* from α . We also define for connected components $F \in \pi_0(M^\mathbb{T})$ of the fixed point locus

$$W_F^+ := \cup_{\alpha \in F} W_\alpha^+$$

the *attractor* of F and

$$W_F^- := \cup_{\alpha \in F} W_\alpha^-.$$

the *repeller* of F .

Let $M^s \subset M$ denote the smooth locus of M . Let $\alpha \in M^{s\mathbb{T}}$ be a smooth fixed point. Then \mathbb{T} acts linearly on the tangent space $T_\alpha M$. We can decompose this representation into weight spaces $T_\alpha M \cong \oplus_{k \in \mathbb{Z}} (T_\alpha M)_k$, where $(T_\alpha M)_k \subset T_\alpha M$ denotes the isotypical component, where $\lambda \in \mathbb{T}$ acts as multiplication by λ^k . We will denote by $T_\alpha^+ M = \oplus_{k > 0} (T_\alpha M)_k$ the positive and by $T_\alpha^- M = \oplus_{k < 0} (T_\alpha M)_k$ the negative part of this decomposition. Let $T_\alpha^0 M = (T_\alpha M)_0$ denote the zero weight component then we have the decomposition

$$T_\alpha M \cong T_\alpha^+ M \oplus T_\alpha^0 M \oplus T_\alpha^- M.$$

Proposition 2.1. *For a smooth fixed point $\alpha \in M^{s\mathbb{T}}$ the upward and downward flows W_α^+ and W_α^- are \mathbb{T} -invariant locally closed subvarieties in M . Furthermore we have $W_\alpha^+ \cong T_\alpha^+ M$ and $W_\alpha^- \cong T_\alpha^- M$ as varieties with \mathbb{T} -action.*

Proof. This is proved in [B-B] for a smooth complete M . The statement above could be reduced to that case by equivariantly compactifying M (see below or [Su, Theorem 3]) and equivariantly resolving the singularities of the compactification ([Vi, Corollary 7.6.3]).

For completeness, and to prepare the ground for the proof of Proposition 2.10, we present a proof inspired by the approach of [BDMN, Lemma 2.2] see also [Dr3, §1]. Let $\alpha \in M^{s\mathbb{T}}$. As the singular locus $\text{Sing}(M) \subset M$ is \mathbb{T} -invariant and closed we have $W_\alpha^+ \subset M^s$. Take $\alpha \in U_0 \subset M^s$ a \mathbb{T} -invariant open affine neighbourhood (using [Su, Corollary 2]). In particular, U_0 is smooth. The \mathbb{T} -action on U_0 induces a \mathbb{Z} -grading on $\mathbb{C}[U_0] \cong H^0(U_0, \mathcal{O}_{U_0})$, defined by setting $f \in \mathbb{C}[U_0]_i$ if and only if

$$f(\lambda p) = \lambda^i f(p) \tag{2.1}$$

for all $\lambda \in \mathbb{T}$ and $p \in U_0$. In this case we will say that f is homogeneous of degree $\text{deg}(f) = i$.

Let $\mathfrak{m}_\alpha \triangleleft \mathbb{C}[U_0]$ be the maximal ideal of functions vanishing at α . As α is \mathbb{T} -fixed \mathfrak{m}_α is a homogeneous ideal. We can find homogeneous elements $x_1, \dots, x_n \in \mathfrak{m}_\alpha \subset \mathbb{C}[U_0]$, where $n = \dim(M)$ such that their image in $\mathfrak{m}_\alpha / \mathfrak{m}_\alpha^2 \cong T_\alpha^* M$ form a basis. The one-forms $dx_1, \dots, dx_n \subset H^0(U_0, \Omega_{U_0})$ then form a basis at α and we let U be the complement of the divisor of zeroes of the section $dx_1 \wedge \dots \wedge dx_n \in H^0(U_0, \Omega^n(U_0))$. It is a \mathbb{T} -invariant affine open subset $U \subset U_0$, where dx_1, \dots, dx_n remains a basis at every point. In particular, U

is non-singular. As U is invariant under the \mathbb{T} -action and the closure of any \mathbb{T} -orbit in $\mathcal{M} \setminus U$ is still contained in $\mathcal{M} \setminus U$, as it is closed, we have

$$W_\alpha^\pm \subset U.$$

Therefore we can prove Proposition 2.1 only working in U .

Let $A := \mathbb{C}[U]$ and $A = \bigoplus_{n \in \mathbb{Z}} A_n$ be the grading induced by the \mathbb{T} -action. Let $A_+ = \bigoplus_{n \in \mathbb{Z}_{>0}} A_n$ and $A_- = \bigoplus_{n \in \mathbb{Z}_{<0}} A_n$. Thus in particular $A = A_- \oplus A_0 \oplus A_+$. First we observe that

Lemma 2.2. $U^\mathbb{T} = V(f | f \in A_+ + A_-) \subset U$ i.e. the subvariety of zeroes of homogenous f of non-zero degree. Thus $\mathbb{C}[U^\mathbb{T}] = A/(A_+, A_-)$.

Proof. Clearly if $p \in U^\mathbb{T}$, $f \in A_i$ with $i \in \mathbb{Z} \setminus \{0\}$ and $\lambda \in \mathbb{T}$ such that $\lambda^i \neq 1$ then

$$f(p) = f(\lambda p) = \lambda^i f(p) = 0.$$

On the other hand if $p \in U$ is such that $f(p) = 0$ for all $f \in A_i$ with $i \in \mathbb{Z} \setminus \{0\}$ then the maximal ideal at p satisfies $(A_+ + A_-) \subset \mathfrak{m}_p$ and so \mathfrak{m}_p is a homogenous ideal, thus $p \in U^\mathbb{T}$. \square

Let $F := U^\mathbb{T}$ and as above, $W_F^+ = \prod_{\beta \in F} W_\beta^+ = \{p \in U | \lim_{\lambda \rightarrow 0} \lambda p \in F\}$ the attractor of F in U . Then we have

Lemma 2.3. $W_F^+ = V(f | f \in A_-)$ i.e. the subvariety of zeroes of homogenous f of negative degree. Thus $\mathbb{C}[W_F^+] = A/(A_-)$.

Proof. Again it is clear that if $p \in W_F^+$ then $f(p) = 0$ for $f \in A_-$. Namely if $f \in A_i$ for $i < 0$ then

$$f(\lim_{\lambda \rightarrow 0} \lambda p) = \lim_{\lambda \rightarrow 0} f(\lambda p) = \lim_{\lambda \rightarrow 0} \lambda^i f(p)$$

implies that $f(p) = 0$.

On the other hand assume that for some $p \in U$ we have $f(p) = 0$ for all $f \in A_-$. Then let $\pi_p : A \rightarrow A/\mathfrak{m}_p \cong \mathbb{C}$ and define

$$g : \begin{array}{ccc} A & \rightarrow & \mathbb{C}[x] \\ \sum_i a_i & \mapsto & \sum_i \pi_p(a_i) x^i \end{array},$$

where we write every element in A in the form of a finite sum of homogeneous elements $\sum_i a_i$ where $a_i \in A_i$. Because $f(a_i) = 0$ for $i < 0$, the definition makes sense. We see that g is a graded ring homomorphism. Observe also that for $z \in \mathbb{A}^1 = \text{Spec}(\mathbb{C}[x])$ the composition

$$(\pi_z \circ g) \left(\sum_i a_i \right) = \sum_i \pi_p(a_i) z^i.$$

When $z = 1$ we see that $\ker(\pi_1 \circ g) = \mathfrak{m}_p$ and that $\ker(\pi_0 \circ g) = (\ker(\pi_p|_{A_0}), A_-, A_+)$ is a maximal ideal containing A_- and A_+ thus corresponds to a fixed point of the \mathbb{T} -action. Indeed the ring homomorphism g gives a \mathbb{T} -equivariant map $\mathbb{A}^1 \rightarrow U$, such that $g(1) = p$ and $g(0) = \lim_{\lambda \rightarrow 0} \lambda p \in F$ by definition. Thus $p \in W_F^+$. The result follows. \square

Remark 2.4. We can build a map $\pi : W_F^+ \rightarrow F$ by sending $p \in W_F^+$ to $\lim_{\lambda \rightarrow 0} \lambda p \in F$. By the construction above we see that the maximal ideal of $\pi(p)$ is $(\ker(\pi_p|_{A_0}), A_-, A_+) \triangleleft A$. Thus we see that π is induced by the map

$$\begin{aligned} \pi^* : \quad A/(A_+, A_-) &\rightarrow A/(A_-) \\ a \bmod (A_+, A_-) &\mapsto a_0 \bmod (A_-) \end{aligned} \quad (2.2)$$

where $a = \sum_{i \in \mathbb{Z}} a_i$ is the sum of homogeneous elements $a_i \in A_i$. It is straightforward to check that π^* is a well-defined graded ring homomorphism and that $(\pi^*)^{-1}(\ker(\pi_p)) = (\ker(\pi_p|_{A_0}), A_-, A_+)$.

Next we determine the tangent spaces of F , W_F^+ , W^-F , W_α^+ and W_α^- . To do this we will denote by $\partial_i \in H^0(U, TU)$ the dual basis to dx_i . Recall that $\lambda \in \mathbb{T}$ acts on the function $x_i \in \mathbb{C}[U]$ via the formula

$$\lambda \cdot x_i(p) = x_i(\lambda^{-1}p) = \lambda^{-\deg(x_i)} x_i(p),$$

i.e. $\lambda \cdot x_i = \lambda^{-\deg(x_i)} x_i$ which should not be confused with the transformation rule in (2.1). In particular, the induced action on one-forms gives

$$\lambda \cdot dx_i = (m_\lambda)_*(dx_i) = \lambda^{-\deg(x_i)} dx_i, \quad (2.3)$$

where m_λ denotes the action isomorphisms:

$$\begin{aligned} m_\lambda : \quad U &\rightarrow U \\ p &\mapsto \lambda p \end{aligned}$$

for $\lambda \in \mathbb{T}$. One can compare¹ (2.3) to

$$m_\lambda^*(dx_i) = d(m_\lambda^*(x_i)) = \lambda^{\deg(x_i)} dx_i.$$

This way we see that ∂_i will have weight $\deg(x_i)$ in the sense that

$$(m_\lambda)_*(\partial_i) = Tm_\lambda(\partial_i) = \lambda^{\deg(x_i)} (\partial_i). \quad (2.4)$$

¹There seems to be a related confusion in [BDMN, Definition 1.1]. Their definition of elliptic \mathbb{T} -action should either require the existence of limit points in the $t \rightarrow 0$ limit as in this paper, or replace the definition of positive weight using $(m_t)_*$ instead of m_t^* . In fact, when they use the definition of positive weight in the displayed line after (2-1) on page 2613 of [BDMN], they use the correct $(m_t)_*$ instead of m_t^* as in [BDMN, Definition 1.1].

As a reality check for $\lambda \in \mathbb{T}$ we compute, using (2.1)

$$dx_j(Tm_\lambda(\delta_i)) = m_\lambda^*(dx_j)(\delta_i) = d(x_j \circ m_\lambda)(\delta_i) = \lambda^{\deg x_j} dx_j(\delta_i) = \lambda^{\deg x_j} \delta_{i,j},$$

showing (2.4).

Lemma 2.5. *For $\beta \in F$ we have*

$$T_\beta F = (T_\beta U)^\mathbb{T} \cong \text{span}(\partial_i|_\beta | \deg(x_i) = 0) \subset T_\beta U.$$

In particular, $F \subset U$ is a non-singular subvariety.

Proof. Dually, we will prove that the kernel of the surjective map $g : T_\beta^* U \rightarrow T_\beta^* F$ is

$$\text{span}(dx_i|_\beta | \deg(x_i) \neq 0) < T_\beta^* U.$$

It is clear that $dx_i|_\beta \in \ker(g)$ when $\deg(x_i) \neq 0$, because g is a map of \mathbb{T} -modules when we endow $T_\beta^* F$ with the trivial \mathbb{T} -module structure. Let $\eta \in (T_\beta^* U)^\mathbb{T} \cap \ker(g)$. We can represent $\eta = dy_0$ with the homogeneous degree 0 element $y_0 \in \mathfrak{m}_{\beta,U} \triangleleft A$. We will denote by

$$\bar{y}_0 \in \mathfrak{m}_{\beta,F} \triangleleft A/(A_+, A_-) \cong \mathbb{C}[F]$$

the image of y_0 in the quotient $A/(A_+, A_-)$. By assumption $0 = g(\eta) = d\bar{y}_0$ thus $\bar{y}_0 \in \mathfrak{m}_{\beta,F}^2$. Thus there exists some $\bar{c}_i, \bar{d}_i \in \mathfrak{m}_{\beta,F}$ such that $\bar{y}_0 = \sum_i \bar{c}_i \bar{d}_i$. Denoting by c_i and d_i some lift of \bar{c}_i and \bar{d}_i in $\mathfrak{m}_{\beta,U}$, we get that $y_0 - \sum_i c_i d_i \in (A_+, A_-)$. We can assume that c_i and d_i are homogeneous and in turn that $\deg(c_i) = \deg(d_i) = 0$. We have some $a_j \in A_+, b_j \in A_-$ such that $y_0 - \sum_i c_i d_i = \sum_j a_j b_j$. We can assume that a_j and b_j are homogeneous of non-zero degree and in turn that $\deg(a_j) + \deg(b_j) = 0$. Denoting $\pi_\beta(a_j) = \lambda_j \in A/\mathfrak{m}_{\beta,U} \cong \mathbb{C}$ and $\pi_\beta(b_j) = \mu_j \in A/\mathfrak{m}_{\beta,U} \cong \mathbb{C}$ we get that $a_j - \lambda_j$ and $b_j - \mu_j$ are in $\mathfrak{m}_{\beta,U}$. Thus we conclude that

$$y_0 - \left(\sum_j a_j \mu_j + \lambda_j b_j \right) \in \mathfrak{m}_{\beta,U}^2.$$

As $\mathfrak{m}_{\beta,U}^2$ is a homogeneous ideal we deduce that $y_0 \in \mathfrak{m}_{\beta,U}^2$ proving the first claim.

Finally, we deduce that $\dim(\mathfrak{m}_{\beta,U}/\mathfrak{m}_{\beta,U}^2) = \#\{dx_i|_\beta | \deg(x_i) = 0\}$ is independent of $\beta \in F$, thus indeed F is non-singular. \square

Lemma 2.6. *For $p \in W_F^+$ we have $T_p W_F^+ = \text{span}(\partial_i|_p | \deg(x_i) \geq 0) < T_p U$. Similarly for $p \in W_F^-$ we have $T_p W_F^- = \text{span}(\partial_i|_p | \deg(x_i) \leq 0) < T_p U$.*

Proof. Again, we will prove dually that the kernel of the surjective map $h : T_p^* U \rightarrow T_p^* W_F^+$ is

$$\text{span}(dx_i|_p | \deg(x_i) < 0) < T_p^* U.$$

By Lemma 2.3 for $\deg(x_i) < 0$ the function x_i vanishes on W_F^+ and thus $dx_i|_p$ is in the kernel of h . Assume now that $\sum_{\deg x_i \geq 0} \alpha_i dx_i|_p \in \ker(h)$ for some $\alpha_i \in \mathbb{C}$. This means that

$$\sum_{\deg x_i \geq 0} \alpha_i (x_i - \pi_p(x_i)) \in \mathfrak{m}_{p, W_F^+}^2.$$

It follows that there are $c_j, d_j \in \mathfrak{m}_{p, U}$ such that

$$\sum_{\deg x_i \geq 0} \alpha_i (x_i - \pi_p(x_i)) - \sum_j c_j d_j \in (A_-).$$

So we have homogeneous $a_k \in A_-$ and $r_k \in A$ such that

$$\sum_{\deg x_i \geq 0} \alpha_i (x_i - \pi_p(x_i)) - \sum_k r_k a_k \in \mathfrak{m}_{p, W_F^+}^2.$$

By writing $r_k = r_k - \pi_p(r_k) + \pi_p(r_k)$ and noting that $\pi_p(a_k) = 0$ as $p \in W_F^+$ and $a_k \in A_-$ we get

$$\sum_{\deg x_i \geq 0} \alpha_i (x_i - \pi_p(x_i)) - \sum_k \beta_k a_k \in \mathfrak{m}_{p, W_F^+}^2,$$

where $\beta_k = \pi_p(r_k) \in \mathbb{C}$. In other words

$$0 = \sum_{\deg x_i \geq 0} \alpha_i dx_i|_p - \sum_k \beta_k da_k|_p \in T_p^*(U). \quad (2.5)$$

Our final observation is that for $\deg(x_i) \geq 0$ the function $\partial_i(a_k)$ has degree $-\deg(x_i) + \deg(a_k) < 0$. This is because $da_k = \sum_j \partial_j(a_k) dx_j$ therefore

$$da_k \wedge dx_1 \wedge \cdots \wedge dx_{i-1} \wedge dx_{i+1} \wedge \cdots \wedge dx_n = (-1)^{i-1} \partial_i(a_k) dx_1 \wedge \cdots \wedge dx_n.$$

Thus $\partial_i(a_k)$ vanishes on W_F^+ . This means that $da_k|_p$ is a linear combination of the dx_i with $\deg(x_i) < 0$ and so $\sum_{\deg x_i \geq 0} \alpha_i dx_i|_p$ and $\sum_k \beta_k da_k|_p$ are linearly independent. Thus (2.5) implies that $\sum_{\deg x_i \geq 0} \alpha_i dx_i|_p = 0$. The first statement follows.

The proof of the second statement on the downward flows is the same as above for the inverse action of \mathbb{T} on U . \square

Lemma 2.7. For $\alpha \in F$, and $p \in W_\alpha^+$ we have $T_p W_\alpha^+ = \text{span}(\partial_i|_p | \deg(x_i) > 0) < T_p U$. Similarly, for $p \in W_\alpha^-$ we have $T_p W_\alpha^- = \text{span}(\partial_i|_p | \deg(x_i) < 0) < T_p U$.

Proof. First we note that the locus of zeroes of (x_1, \dots, x_n) in U is zero dimensional, because dx_1, \dots, dx_n is a basis at every point. Also this locus is invariant by \mathbb{T} , they are thus fixed points in F . All non-negative degree elements vanish on F therefore the zeroes of the 0 degree elements

$$\mathbf{x}_0 := \{x_i | \deg(x_i) = 0\}$$

on F are precisely the zeroes of (x_1, \dots, x_n) on U .

By construction one such common zero is α and so W_α^+ is a connected component of the variety of zeroes of \mathbf{x}_0 on W_F^+ . In particular, $W_\alpha^+ \subset U$ is closed and thus it is locally closed in M .

Thus we have that W_α^+ is a connected component of $\text{Spec}(A/(A_-, \mathbf{x}_0))$. For $p \in W_\alpha^+$ we determine the kernel of the surjective map $f : T_p^*U \rightarrow T_p^*W_\alpha^+$. All functions in A_- and \mathbf{x}_0 restrict to W_α^+ as zero thus $dx_i|_p$ is in the kernel when $\deg(x_i) \leq 0$. Let now $\sum_{\deg x_i > 0} \alpha_i dx_i|_p \in \ker(f)$. This means that

$$\sum_{\deg x_i > 0} \alpha_i (x_i - \pi_p(x_i)) \in \mathfrak{m}_{p, W_\alpha^+}^2,$$

hence there are $c_j, d_j \in \mathfrak{m}_{p, U}$ such that

$$\sum_{\deg x_i > 0} \alpha_i (x_i - \pi_p(x_i)) - \sum_j c_j d_j \in (A_-, \mathbf{x}_0).$$

In turn, this means that there are homogeneous elements $a_k \in A_-$ and scalars $\beta_j \in \mathbb{C}$ such that

$$0 = \sum_{\deg x_i > 0} \alpha_i dx_i|_p - \sum_k da_k|_p - \sum_{\deg x_j = 0} \beta_j dx_j|_p \in T_p^*(U). \quad (2.6)$$

Just as in the proof of Lemma 2.6 above we can argue that as $a(p) = 0$ for all $a \in A_-$, $da_k|_p \in \text{span}(dx_i|_p | \deg x_i < 0)$, thus the linear independence of $\{dx_1|_p, \dots, dx_n|_p\}$ and (2.6) implies that $\sum_{\deg x_i > 0} \alpha_i dx_i|_p = 0$ proving the first claim.

The second claim again follows from the first by inverting the \mathbb{T} -action on U . \square

To finish the proof of Proposition 2.1 we define the map

$$f_\alpha : W_\alpha^+ \rightarrow T_\alpha^+ \quad (2.7)$$

sending

$$p \mapsto \sum_{\deg(x_i) > 0} x_i(p) \partial_i$$

where $\partial_i \in T_\alpha M$ is the dual basis of dx_1, \dots, dx_n at α . By Lemma 2.7 the derivative of f_α is an isomorphism, thus it is étale and so open. Furthermore f_α is \mathbb{T} -equivariant and its image is an open \mathbb{T} -invariant subvariety in the positive \mathbb{T} -module $T_\alpha^+ M$, containing the origin, thus f_α is surjective. As $T_\alpha^+ M$ is simply connected, it follows that f_α is a trivial covering, and W_α^+ being connected implies that f_α is an isomorphism. \square

Remark 2.8. If we put the maps f_α in a family over F we get a \mathbb{T} -equivariant isomorphism $f : W_F^+ \cong T^+ F$ showing that $U_{F_\alpha}^s \subset M$ is a locally trivial affine fibration, another main result of [B-B].

Definition 2.9. We call $M = \coprod_{\alpha} W_{\alpha}^{+}$ the *Bialynicki-Birula partition* and $M = \coprod_F W_F^{+}$ the *Bialynicki-Birula decomposition*. Define also

$$\mathcal{C} := \coprod_{\alpha \in M^{\mathbb{T}}} W_{\alpha}^{-} = \coprod_{F \in \pi_0(M^{\mathbb{T}})} W_F^{-} \quad (2.8)$$

to be the *core* of M .

Additionally, in all our examples (M^s, ω) with $\omega \in \Omega^2(M)$ will be symplectic s.t.

$$\lambda^*(\omega) = \lambda\omega. \quad (2.9)$$

Then we have

Proposition 2.10. *Let M be a normal, semi-projective complex variety with (M^s, ω) symplectic with $\omega \in \Omega^2(M^s)$. Assume (2.9) that ω is homogeneous of weight 1. For a smooth point $\alpha \in M^{s\mathbb{T}}$ the subspaces $T_{\alpha}^{+}M, T_{\alpha}^{\leq 0}M := T_{\alpha}^0M \oplus T_{\alpha}^{+}M < T_{\alpha}M$ and subvarieties $W_{\alpha}^{+}, W_{F_{\alpha}}^{-} \subset M^s$ are Lagrangian.*

Proof. As $\alpha \in M^{s\mathbb{T}}$ the tangent space $T_{\alpha}M$ is a \mathbb{T} -module. Let $X, Y \in T_{\alpha}M$ be homogeneous with weight ν_1 and ν_2 respectively. That means that $\lambda \cdot X = \lambda^{\nu_1}X$ and $\lambda \cdot Y = \lambda^{\nu_2}Y$. Then from (2.9)

$$\lambda\omega(X, Y) = \lambda^*(\omega)(X, Y) = \omega(\lambda \cdot X, \lambda \cdot Y) = \lambda^{\nu_1 + \nu_2}\omega(X, Y).$$

Consequently $\omega(X, Y) = 0$ unless $\nu_1 + \nu_2 = 1$. Thus ω is trivial on T_{α}^{+} and $T_{\alpha}^{\leq 0}$. As $T_{\alpha}M = T_{\alpha}^{+} \oplus T_{\alpha}^{\leq 0}$ both subspaces are Lagrangian.

Recall the construction of $U \subset M^s$, the \mathbb{T} -invariant affine neighbourhood of $\alpha \in U$ from the proof of Proposition 2.1 above. Let $\omega = \sum_{i < j} f_{i,j} dx_i \wedge dx_j$ for some unique $f_{i,j} \in \mathbb{C}[U]$. As ω, x_i and x_j are homogeneous so is $f_{i,j}$ and $\deg f_{i,j} = 1 - \deg(x_i) - \deg(x_j)$. Let now $p \in W_{\alpha}^{+}$. By Lemma 2.7 the tangent vectors $\partial_k|_p \in T_pU$ with positive weights will be a basis for $T_pW_{\alpha}^{+}$. Take two such $\partial_k|_p, \partial_l|_p \in T_pW_{\alpha}^{+}$ and compute

$$\omega(\partial_k|_p, \partial_l|_p) = \sum_{i < j} f_{i,j} dx_i \wedge dx_j (\partial_k|_p, \partial_l|_p) = f_{k,l}(p).$$

The degree of x_l and x_k are positive, so the degree of $f_{k,l}$ is negative in particular $f_{k,l}$ vanishes on W_{α}^{+} thus $f_{k,l}(p) = 0$, and $T_pW_{\alpha}^{+} \subset T_pU$ is isotropic, and half-dimensional as $T_{\alpha}W_{\alpha}^{+} = T_{\alpha}^{+} < T_{\alpha}M$ is Lagrangian, thus indeed W_{α}^{+} is Lagrangian in $U \subset M^s$. Similarly we can prove that W_F^{-} is Lagrangian in M^s . The result follows. \square

Remark 2.11. Our approach in the proof of Proposition 2.10 above was inspired by [BDMN, §2.2]. Similar results were mentioned in [Na, Proposition 7.1].

2.2 Very stable upward flows

Definition 2.12. Let $\alpha \in M^{s\mathbb{T}}$. We say that α is *very stable* if the only point in its upward flow which is also in the core \mathcal{C} of (2.8) is α :

$$W_\alpha^+ \cap \mathcal{C} = \{\alpha\}.$$

Remark 2.13. The origin of the term *very stable* is in the work of Drinfeld [Dr1] and Laumon [La1], which was applied to vector bundles on curves (see Section 4). However there is an accidental coincidence in terminology. Namely, Bialynicki-Birula [B-B2] using terminology of Smale [Sm, II.2] calls our upward flow W_α^+ the *stable* subscheme of α while our downward flow W_α^- is called the *unstable* subscheme of α . Our definition above can be formulated to say that the stable subscheme W_α^+ is *very stable* if and only if it is disjoint from the unstable subschemes W_β^- for $\beta \neq \alpha$.

Proposition 2.14. $\alpha \in M^{s\mathbb{T}}$ is *very stable* if and only if $W_\alpha^+ \subset M$ is closed.

Proof. If α is not very stable then there exists $\beta \neq \alpha \in W_\alpha^+ \cap \mathcal{C}$ in the upward flow of α . Then $\lim_{\lambda \rightarrow \infty} (\lambda \cdot \beta) \in M^{\mathbb{T}}$ is fixed by the \mathbb{T} -action. Moreover a linearized \mathbb{T} -action on a very ample line bundle on M (which exists, because M is normal and quasi-projective and [Do, Corollary 7.2]) embeds it equivariantly in some projective space $M \subset \mathbb{P}^N$ with linear \mathbb{T} -action, where the closure of every non-trivial \mathbb{T} -orbit is linear, thus has two distinct \mathbb{T} -fixed points. Consequently,

$$\lim_{\lambda \rightarrow \infty} (\lambda \cdot \beta) \neq \lim_{\lambda \rightarrow 0} (\lambda \cdot \beta) = \alpha$$

and thus W_α^+ is not closed.

Assume now that $\alpha \in M^{s\mathbb{T}}$ is very stable i.e.

$$W_\alpha^+ \cap \mathcal{C} = \{\alpha\}$$

as sets. Then we will determine the image of $W_\alpha^+ \setminus \mathcal{C}$ in the geometric quotient

$$Z = (M \setminus \mathcal{C})/\mathbb{T}.$$

By construction it is

$$\mathbb{P}(W_\alpha^+) = W_\alpha^+ \setminus \{\alpha\}/\mathbb{T} \subset Z$$

isomorphic to the weighted projective space $\mathbb{P}(T_\alpha^+) = (T_\alpha^+ \setminus \{0\})/\mathbb{T}$. By the construction of [HV, (1.2.3)] (c.f. also [Si3, §11] and [Ha1] in the Higgs moduli space case) we have the compactification

$$\overline{M} = (M \times \mathbb{C})//\mathbb{T} = (M \times \mathbb{C} \setminus \mathcal{C} \times \{0\})/\mathbb{T} \cong M \sqcup Z.$$

We see that the geometric quotient

$$\mathbb{P}(W_\alpha^+ \times \mathbb{C}) = ((W_\alpha^+ \times \mathbb{C}) \setminus (\alpha, 0)) / \mathbb{T} = W_\alpha^+ \sqcup \mathbb{P}(W_\alpha^+) \subset M \sqcup Z = \overline{M}$$

is also a weighted projective space, thus projective, thus closed in \overline{M} . It follows that all boundary points of the closure \overline{U}_α in \overline{M} lie on the divisor at infinity $Z \subset \overline{M} \setminus M$, thus W_α^+ is closed in M . \square

3 Bialynicki-Birula theory on the moduli space of Higgs bundles

Fix C a smooth complex projective curve of genus $g > 1$. Recall that a vector bundle E on C is called *stable* (resp. *semi-stable*) if for all proper subbundles $F \subset E$

$$\frac{\deg(F)}{\text{rank}(F)} < \frac{\deg(E)}{\text{rank}(E)} \quad (3.1)$$

(resp. $\deg(F)/\text{rank}(F) \leq \deg(E)/\text{rank}(E)$).

Similarly, a Higgs bundle (E, Φ) where $\Phi \in H^0(\text{End}(E) \otimes K)$, is stable (resp. semi-stable) if (4.14) holds for Φ -invariant proper subbundles $F \subset E$.

We fix integers $n \in \mathbb{Z} > 0$ and $d \in \mathbb{Z}$. The moduli space of semistable Higgs bundles $\mathcal{M} := \mathcal{M}_{n,d}$ of rank n and degree d was constructed by [Hi1] by gauge-theoretic and by [Ni] and [Si1] by algebraic geometric methods. At its smooth points it carries a hyperkähler metric, in particular an algebraic symplectic structure. It is a normal [Si1, Corollary 11.7] quasi-projective variety. The open subset $\mathcal{M}^s \subset \mathcal{M}$ of stable points is precisely the smooth locus. It is a coarse moduli space, but carries a universal Higgs bundle étale locally on \mathcal{M}^s [Si1, Theorem 4.5.(4)].

For a Higgs bundle (E, Φ) we can compute the characteristic polynomial of the Higgs field as $\det(t - \Phi) = t^n + a_1 t^{n-1} + \dots + a_n$, where $a_i \in H^0(C; K^i)$. This leads to the map

$$h : \mathcal{M} \rightarrow \bigoplus_{i=1}^n H^0(C; K^i),$$

a proper, completely integrable Hamiltonian system [Hi2, Si2, Ni] in particular, the fibres are Lagrangian at their smooth points.

\mathcal{M} carries a \mathbb{C}^\times -action defined by $(E, \Phi) \mapsto (E, \lambda\Phi)$ for $\lambda \in \mathbb{C}^\times$. If we let \mathbb{C}^\times act with weight i on $H^0(C; K^i)$ then h becomes \mathbb{C}^\times -equivariant. As h is proper and all the weights on $\bigoplus_{i=1}^n H^0(C; K^i)$ are positive, we deduce that \mathcal{M} is semi-projective with respect to this \mathbb{T} -action.

Finally, we recall that there is a canonical symplectic structure on \mathcal{M}^s originally arising as part of its hyperkähler structure. The tangent space at $(E, \Phi) \in \mathcal{M}^s$ can be identified (see e.g. [BR])

$$T_{(E,\Phi)}\mathcal{M}^s \cong \mathbb{H}^1 \left(C; \text{End}(E) \xrightarrow{ad(\Phi)} \text{End}(E) \otimes K \right). \quad (3.2)$$

Serre duality will give an isomorphism between this hypercohomology and its dual. This defines a symplectic form on $T_{(E,\Phi)}$. To see that it gives rise to a closed form ω , we recall [BR, Theorem 4.3] that $\omega = d\theta$ where

$$\theta : \mathbb{H}^1 \left(C; \text{End}(E) \xrightarrow{ad(\Phi)} \text{End}(E) \otimes K \right) \rightarrow \mathbb{C}$$

is given by first mapping to $H^1(C; \text{End}(E))$ and pairing this element with $\Phi \in H^0(C; \text{End}(E) \otimes K)$ by Serre duality. In particular ω is closed. With respect to the \mathbb{T} -action we clearly have that $\lambda^*(\theta) = \lambda\theta$ and thus $\lambda^*(\omega) = \lambda\omega$ is also of weight 1.

In this paper we will study the upward and downward flows of the Bialynicki-Birula partition on \mathcal{M} .

3.1 Fixed points of the \mathbb{T} -action $\mathcal{M}^{s\mathbb{T}}$

First we recall a parametrisation of $\mathcal{M}^{\mathbb{T}}$. If $(E, \Phi) \in \mathcal{M}^{s\mathbb{T}}$ is a stable Higgs bundle fixed by the \mathbb{T} -action then we have for every $\lambda \in \mathbb{T}$

$$(E, \lambda\Phi) \cong (E, \Phi).$$

The isomorphism above gives a vector bundle automorphism $f_\lambda : E \rightarrow E$ making the diagram

$$\begin{array}{ccc} E & \xrightarrow{\lambda\Phi} & EK \\ f_\lambda \downarrow & & \downarrow f_\lambda \\ E & \xrightarrow{\Phi} & EK \end{array} \quad (3.3)$$

commutative. These define a fibrewise \mathbb{T} -action on E . Let $E = L_0 \oplus \dots \oplus L_k$ be the weight space decomposition of this \mathbb{T} -action, for subbundles $L_i < E$ and $1 \leq k \leq n-1$. Let \mathbb{T} act on L_i by weight $w_i \in \mathbb{Z}$, in other words f_λ acts on L_i as multiplication by λ^{w_i} . From the diagram (3.3) we see that Φ maps the weight w_i space to the $w_i - 1$ weight space. Using an overall scaling we can assume that $w_i = -i$. Then $\Phi(L_i) \subset L_{i+1}K$. We call the ordered partition $(\text{rank}(L_0), \dots, \text{rank}(L_k))$ of n the *type* of the fixed point $(E, \Phi) \in \mathcal{M}^{s\mathbb{T}}$. Similarly we can talk about the type of a component of the fixed point set $F \in \pi_0(\mathcal{M}^{s\mathbb{T}})$ as the type is locally constant on $\mathcal{M}^{s\mathbb{T}}$. The latter can be seen by noting that étale locally we have a \mathbb{T} -equivariant universal bundle on \mathcal{M}^s and on the connected component $F \in \pi_0(\mathcal{M}^{s\mathbb{T}})$ the weight space for a given weight of the \mathbb{T} -action forms a vector bundle.

Example 3.1. For example when $E = L_0$ the type is (n) and $\Phi = 0$, thus E is a semistable bundle. We will denote by \mathcal{N} the moduli space of rank n degree d semistable bundles which is embedded $\mathcal{N} \subset \mathcal{M}^{\mathbb{T}}$, the component of $\mathcal{M}^{\mathbb{T}}$ of type (n) .

Example 3.2. At the other extreme are the fixed points of type $(1, \dots, 1)$. Let $(E, \Phi) \in \mathcal{M}^{\mathbb{T}}$ be a type $(1, \dots, 1)$ fixed point of the \mathbb{T} -action. In other words $E = L_0 \oplus \dots \oplus L_{n-1}$ is a direct sum of line bundles and the Higgs field Φ satisfies $\Phi(L_{i-1}) \subset L_i K$. Such a Higgs bundle is determined by L_0 and the choice of

$$b_i := \Phi|_{L_{i-1}} \in H^0(C; L_{i-1}^* L_i K).$$

Note that the isomorphism class of such a Higgs bundle only depends on the effective divisors $\delta_i := \text{div}(b_i) \in C^{[m_i]}$, where $m_i = \deg(L_{i-1}^* L_i K)$, and L_0 . To simplify notation we will choose a divisor δ_0 so that $L_0 = \mathcal{O}(\delta_0)$. We will write $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1})$. We also denote by \mathcal{E}_δ the type $(1, \dots, 1)$ Higgs bundle for which

$$L_i := (\delta_0 + \delta_1 + \dots + \delta_{i-1}) K^{-i+1}$$

and

$$b_i := s_{\delta_i} \in H^0(C; \mathcal{O}(\delta_i)) \cong H^0(C; L_{i-1}^* L_i K)$$

the defining section of $\mathcal{O}(\delta_i)$.

Let $\ell := \deg(L) = \deg(\delta_0)$ and $m := (m_1, \dots, m_{n-1})$ where $m_i = \deg(b_i) = \deg(\delta_i)$. Then the ambient component $\mathcal{E}_\delta \in F_{\ell, m} \in \pi_0(\mathcal{M}^{\mathbb{T}})$ of the fixed point set $\mathcal{M}^{\mathbb{T}}$ is isomorphic to

$$F_{\ell, m} \cong J_\ell(C) \times C^{[m_1]} \times \dots \times C^{[m_{n-1}]}.$$
 (3.4)

Here the Jacobian $J_\ell(C)$ means the moduli space of degree ℓ divisors on C modulo principal ones, when we represent its elements by degree ℓ divisors δ_0 , or equivalently $J_\ell(C)$ can mean the moduli space of degree ℓ line bundles when we represent its elements by the line bundle L_0 .

Remark 3.3. For rank $n = 2$ we have just two different types. The type (2) fixed point component is $\mathcal{N} \subset \mathcal{M}$ the moduli space of semistable bundles with zero Higgs field. The type (1, 1) components $F_{\ell, m_1} \cong J_\ell(C) \times C^{[m_1]}$ parametrise Higgs bundles of the form $E = L_0 \oplus L_1$ and

$$\Phi = \begin{pmatrix} 0 & 0 \\ b & 0 \end{pmatrix},$$

where L_0 and L_1 are line bundles of degrees $\ell_0 := \ell$ and ℓ_1 respectively, $b \in H^0(C; L_0^* L_1 K)$ and $m_1 = \deg(b) = \ell_1 - \ell_0 + 2g - 2$. For stability we need $m_1 < 2g - 2$. When $d = \ell_0 + \ell_1$ is fixed then $2\ell_0 = d - m_1 + (2g - 2)$ and we will abbreviate $F_{m_1} := F_{\ell, m_1}$. We then have $g - 1$ type (1, 1) fixed point components $F_1, F_3, \dots, F_{2g-3}$ when d is odd and $g - 1$ of them F_0, \dots, F_{2g-4} when d is even.

3.2 Upward flows on \mathcal{M}

We now describe the upward flows in the Bialynicki-Birula partition of \mathcal{M} . We know from Proposition 2.1 that the upward flow $W_{\mathcal{E}}^+$ for $\mathcal{E} \in \mathcal{M}^{\mathbb{T}}$ is isomorphic to a vector space modelled on $T_{\mathcal{E}}^+$. The following proposition describes which Higgs bundles belong to $W_{\mathcal{E}}^+$.

Proposition 3.4. *Let $\mathcal{E} = (E, \Phi) \in \mathcal{M}$ then*

$$\lim_{\lambda \rightarrow 0} (E, \lambda\Phi) = (E', \Phi') \quad (3.5)$$

for some $(E', \Phi') \in \mathcal{M}^{s\mathbb{T}}$ if and only if there exists a filtration

$$0 = E_0 \subset E_1 \subset \cdots \subset E_{k-1} \subset E_k = E \quad (3.6)$$

by subbundles such that for all i

$$\Phi(E_i) \subset E_{i+1}K. \quad (3.7)$$

This induces maps

$$gr_0(\Phi) : E_i/E_{i+1} \rightarrow E_{i+1}/E_{i+2}K$$

and the corresponding graded Higgs bundle agrees with the limiting Higgs bundle

$$(E_1/E_0 \oplus E_2/E_1 \oplus \cdots \oplus E_k/E_{k-1}, gr(\Phi)) \cong (gr(E), gr_0(\Phi)) \cong (E', \Phi'). \quad (3.8)$$

Furthermore the filtration (3.6) with these properties is unique.

Proof. The proof is by a Higgs bundle analogue of the Rees construction. The vector bundle version is discussed in [As, He, Hal].

First let us assume that we have a filtration (3.6) which is compatible with the Higgs field i.e. (3.7) holds for all i . We denote by $E' := gr(E)$ and $\Phi' = gr_0(\Phi)$ and assume that $(E', \Phi') \in \mathcal{M}^{s\mathbb{T}}$ is a stable Higgs bundle.

We can define a vector bundle \tilde{E} over $\mathbb{C} \times C = \text{Spec}(\mathbb{C}[x]) \times C$ given by the \mathbb{Z} -graded $\mathcal{O}_C[x]$ -module

$$\tilde{E} := \bigoplus_{i \in \mathbb{Z}} x^{-i} E_{-i} \quad (3.9)$$

on C , where x acts via the embedding $x^{-i} E_{-i} \subset x^{-i+1} E_{-i+1}$. Here $E_k = E$ for $k \geq n$ and $E_k = 0$ for $k \leq 0$. Note, that we only have non-positive weights in the \mathbb{Z} -graded module (3.9).

We also get the vector bundle homomorphism $\tilde{\Phi} : \tilde{E} \rightarrow \tilde{E} \otimes K$ by $\tilde{\Phi} : x^i E_i \rightarrow x^{i+1} E_{i+1} \otimes K$ given by $\Phi|_{E_i}$. Furthermore the \mathbb{Z} -grading (3.9) defines a \mathbb{T} -action on \tilde{E}

covering the standard action of \mathbb{T} on $\mathbb{C} \cong \text{Spec}(\bigoplus_{i \in \mathbb{Z}_{\leq 0}} x^{-i})$. Under this \mathbb{T} -action $\lambda \in \mathbb{T}$ sends $\tilde{\Phi}$ to $\lambda^{-1}\tilde{\Phi}$. This gives us a family of Higgs bundles over C parametrized by \mathbb{C} . Over $\{1\} \times C$ we get the vector bundle

$$\tilde{E}/(x-1)\tilde{E} \cong E$$

back on C , while $\tilde{\Phi}$ will induce precisely Φ , yielding our original Higgs bundle

$$(\tilde{E}, \tilde{\Phi})_{\{1\} \times C} \cong (E, \Phi)$$

The \mathbb{T} -equivariant structure on $(\tilde{E}, \tilde{\Phi})$ shows that over $\{\lambda\} \times C$ for $0 \neq \lambda \in \mathbb{C}$ we get

$$(\tilde{E}, \tilde{\Phi})_{\{\lambda\} \times C} \cong (E, \lambda\Phi) \quad (3.10)$$

Over $\{0\} \times C \in \mathbb{C}$ the vector bundle \tilde{E} restricts as

$$\tilde{E}/(x)\tilde{E} \cong \text{gr}(E) = E'$$

while $\tilde{\Phi}$ restricts as $\text{gr}(\Phi) = \Phi'$, giving the Higgs bundle

$$(\tilde{E}, \tilde{\Phi})_{\{0\} \times C} \cong (E', \Phi'). \quad (3.11)$$

As \mathcal{M} is a coarse moduli space we have a morphism $f : \mathbb{C} \rightarrow \mathcal{M}$ such that

$$f(\lambda) = \begin{cases} (E, \lambda\Phi) & \text{when } \lambda \neq 0 \\ (E', \Phi') & \text{when } \lambda = 0 \end{cases} \quad (3.12)$$

which exactly means (3.5).

For the other direction we assume (3.5) and note that as (E', Φ') is a stable Higgs bundle $(E, \lambda\Phi)$ is stable too, because stability is an open condition (c.f. [Si1, Lemma 3.7] or [Ni, Proposition 3.1]). Let us assume the existence of a map like f . Then, as the obstruction in $H^2(\mathcal{M}^s, \mathbb{T})$ vanishes on $f(\mathbb{C})$, we get a family of stable Higgs bundles $(\tilde{E}, \tilde{\Phi})$ over $\mathbb{C} \times C$ with the properties (3.10) and (3.11). As in §6.3.2 and [HT, §4] we can construct a \mathbb{T} -equivariant structure on \tilde{E} where $\lambda \in \mathbb{T}$ acts on $\tilde{\Phi}$ as $\lambda^{-1}\tilde{\Phi}$. In other words if $v \in \tilde{E}$ then

$$\tilde{\Phi}(\lambda(v)) = \lambda(\tilde{\Phi}(v)),$$

if the \mathbb{T} -action on $\tilde{E}K$ is induced from the \mathbb{T} -action on \tilde{E} and the weight one action on K , i.e. for $\omega \in K$ $\lambda(\omega) := \lambda\omega$; compare it with (6.23). Denote by

$$E_i = \{v \in E \cong \tilde{E}|_{\{1\}} \mid \lim_{\lambda \rightarrow 0} \lambda^i \lambda(v) \text{ exists}\}. \quad (3.13)$$

Note that

$$\lambda^i \lambda(\Phi(v)) = \lambda^i \lambda(\tilde{\Phi}|_{\{1\}}(v)) = \lambda^i \tilde{\Phi}(\lambda(v)) = \tilde{\Phi}(\lambda^i \lambda(v)). \quad (3.14)$$

So if $v \in E_i$ then $\lim_{\lambda \rightarrow 0} \lambda^i \lambda(v)$ exists and so does $\lim_{\lambda \rightarrow 0} \lambda^i \lambda(\Phi(v))$. Recalling that λ acts on K with weight one, we see that $\Phi(v) \in E_{i+1}K$. As the construction is locally trivial over C , we get an increasing filtration $E_i \subset E_{i+1} \subset E$ by subbundles of E . Furthermore $\Phi(E_i) \subset E_{i+1}K$ thus Φ is compatible with the filtration. For $\mu \in \mathbb{T}$ we compute

$$\mu(\lim_{\lambda \rightarrow 0} \lambda^i \lambda(v)) = \lim_{\lambda \rightarrow 0} \lambda^i \mu(\lambda(v)) = \mu^{-i} \lim_{\lambda \rightarrow 0} (\mu \lambda)^i (\mu \lambda)(v) = \mu^{-i} \lim_{\lambda \rightarrow 0} \lambda^i \lambda(v).$$

This way we get a map

$$f_i : \begin{array}{ccc} E_i & \rightarrow & (\tilde{E}|_{\{0\}})_{-i} \\ v & \mapsto & \lim_{\lambda \rightarrow 0} \lambda^i \lambda(v) \end{array}$$

from E_i to the weight $-i$ \mathbb{T} -isotypical component of the vector bundle with \mathbb{T} -action $\tilde{E}|_{\{0\}}$ on C . The kernel of this map consists of those $v \in E_i$ for which

$$\lim_{\lambda \rightarrow 0} \lambda^i \lambda(v) = 0,$$

i.e there exists a section $s \in \Gamma(\mathbb{C}, \tilde{E}|_{\mathbb{C} \times \{\pi(v)\}})$, where $\pi : E \rightarrow C$ is the projection of the vector bundle, $s(0) = 0$ and $s(\lambda) = \lambda^i \lambda(v)$. Then we can write $s = \lambda s'$ for another $s' \in \Gamma(\mathbb{C}, \tilde{E}|_{\mathbb{C} \times \{\pi(v)\}})$ and so $\lim_{\lambda \rightarrow 0} \lambda^{i-1} \lambda(v) = s'(0)$ exists showing $\ker(f_i) = E_{i-1}$. Thus f_i induces

$$f_i : E_i/E_{i-1} \hookrightarrow (\tilde{E}|_{\{0\}})_i \subset \tilde{E}|_{\{0\}},$$

and if we put them together we get $f : gr(E) \hookrightarrow \tilde{E}|_{\{0\}}$ an injective bundle map between vector bundles of the same rank, which thus must be an isomorphism.

Finally, we note that (3.14) implies that for $v \in E_i$ we have

$$f_{i+1}(\Phi(v)) = \tilde{\Phi}|_0(f_i(v))$$

thus

$$(gr(E), gr(\Phi)) \cong (\tilde{E}|_{\{0\}}, \tilde{\Phi}|_{\{0\}})$$

completing the proof of existence of the required compatible filtration.

In order to prove the uniqueness of the filtration we start with a compatible filtration (4.3) as in the first part of the proof. Then we construct the \mathbb{T} -equivariant family of stable Higgs bundles \tilde{E} over $\mathbb{C} \times C$ as in (3.9). Then we will show that the original filtration agrees with the one constructed in the second part of the proof with the definition (3.13). First we reformulate (3.13) by saying that $E'_i \subset E$ is the subsheaf whose sections $s \in H^0(U, E'_i) \subset H^0(U, E)$ on open subsets $U \subset C$ satisfy the condition that the section $\lambda^{-i} \lambda(s) \in H^0(\mathbb{C}^* \times U, \tilde{E})$ extends to a section in $H^0(\mathbb{C} \times U, \tilde{E})$. We note that the $\mathbb{C}[x, x^{-1}]$ -module $H^0(\mathbb{C}^* \times U, \tilde{E})$ is the localization of the $\mathbb{C}[x]$ -module $\bigoplus_{i \in \mathbb{Z}} x^{-i} E_{-i}$ at x , thus

$$H^0(\mathbb{C}^* \times U, \tilde{E}) \cong \bigoplus_{i \in \mathbb{Z}} x^{-i} \left(\bigoplus_{j \in \mathbb{Z}_{\geq 0}} H^0(U; E_{-i+j}) \right).$$

We see that the \mathbb{T} -invariant section $\lambda(s) \in H^0(\mathbb{C}^* \times U; \tilde{E})^\mathbb{T}$ agrees with $s = x^0 s$ in this description and so the section $\lambda^{-i} \lambda(s) \in H^0(\mathbb{C}^* \times U; \tilde{E})$ agrees with $x^{-i} s$ in this description also. Finally, we see that $x^{-i} s$ extends to a section over $\mathbb{C} \times U$ if and only if it is in the image of the restriction map $H^0(\mathbb{C} \times U, \tilde{E}) \rightarrow H^0(\mathbb{C}^* \times U, \tilde{E})$. This happens exactly when $s \in H^0(U; E_{-i})$. Thus $E'_{-i} = E_{-i}$. Therefore the filtration in (4.3) satisfying (3.8) is unique.

The proof of the Proposition is complete. \square

Remark 3.5. When $n = 2$ and $(E, \Phi) \in \mathcal{M}$ then we can find the filtration in Proposition 3.4, giving the limiting Higgs bundle $\lim_{\lambda \rightarrow 0} (E, \lambda \Phi) \in \mathcal{M}^\mathbb{T}$ as follows. If E is semistable then the limiting Higgs bundle is simply $(E, 0)$. When E is not stable then it has a unique (maximal) destabilizing subbundle $L \subset E$. The compatibility condition (3.7) in this case is vacuous, and thus the limiting Higgs bundle is $\mathcal{E}_{\delta_0, \delta_1}$ where $L = \mathcal{O}(\delta_0)$, $\delta_1 = \text{div}(b)$ with $b := \Phi|_L : L \rightarrow (E/L) \otimes K$.

Remark 3.6. If we projectivise the total space of the \mathbb{T} -equivariant vector bundle \tilde{E} as $\mathbb{P}(\tilde{E})$, then the map $\mathbb{P}(\tilde{E}) \rightarrow \mathbb{C}$ will be proper, and \mathbb{T} -equivariant with respect to the induced \mathbb{T} -action on $\mathbb{P}(\tilde{E})$. It follows that this \mathbb{T} -action on $\mathbb{P}(\tilde{E})$ is semi-projective. The fixed point set of the action is

$$(\mathbb{P}(\tilde{E}))^\mathbb{T} = \cup_i \mathbb{P}((\tilde{E}|_{\{0\}})_i).$$

The Bialynicki-Birula decomposition $\mathbb{P}(\tilde{E}) = \cup_i W_i^+$ gives

$$\mathbb{P}(E_i) \setminus \mathbb{P}(E_{i-1}) = W_{-i}^+ \cap \mathbb{P}(\tilde{E})|_{\{1\}},$$

which follows from the proof of Proposition 3.4. Thus, amusingly, the filtration $E_i \subset E_{i+1} \subset E$ in Proposition 3.4 describing the Bialynicki-Birula upward flows on \mathcal{M} , can be recovered from the Bialynicki-Birula decomposition on $\mathbb{P}(\tilde{E})$.

In fact, one can construct the Bialynicki-Birula partition on the total space of the projectivised equivariant universal bundle (which always exist over the stable locus, and extends over the whole of \mathcal{M} using Simpson's framed moduli Higgs moduli space [Si1, Theorem 4.10], see (4.18) for $k = 1$) on $\mathcal{M} \times C$ which will contain the information on the filtration in Proposition 3.4 which determines the limiting Higgs bundle.

Remark 3.7. In a differential geometric language Proposition 3.4 appeared in [CW, Proposition 4.2]. Also a closely related partition of \mathcal{M}_{DR} appeared in [Si2]. In fact the algorithm in [Si2] can be used verbatim to find the filtration in Proposition 3.4 producing the limiting Higgs bundle.

Remark 3.8. Let us recall the construction of a section of the Hitchin map from [Hi3]. Let $a = (a_1, \dots, a_n) \in H^0(C; K) \oplus \dots \oplus H^0(C; K^n) = \mathcal{A}$ be a point in the Hitchin base. For a line bundle L on C let

$$E_L := L \oplus LK^{-1} \oplus \dots \oplus LK^{1-n}.$$

Take the Higgs field $\Phi_a : E_L \rightarrow E_L K$ given by the companion matrix:

$$\Phi_a := \begin{pmatrix} 0 & 0 & \dots & 0 & -a_n \\ 1 & 0 & \dots & 0 & -a_{n-1} \\ 0 & 1 & \dots & 0 & -a_{n-2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & -a_1 \end{pmatrix}. \quad (3.15)$$

Denote the Higgs bundle $\mathcal{E}_{L,a} := (E_L, \Phi_a)$. Define $E_i := L \oplus LK^{-1} \oplus \dots \oplus LK^{-i+1}$. The filtration $0 \subset E_1 \cdots \subset E_n = E_L$ satisfies $\Phi_a(E_i) \subset E_{i+1}K$ and the associated graded is $(gr(E_L), gr_0(\Phi_a)) = (E_L, \Phi_0) = \mathcal{E}_{L,0}$. It is straightforward to check this is stable. We call $\mathcal{E}_{L,0}$ a *uniformising Higgs bundle*². Now Proposition 3.4 applies and shows that $\mathcal{E}_{L,a}$ is stable too and $\lim_{\lambda \rightarrow 0} (E_L, \lambda \Phi_a) = (E_L, \Phi_0) = \mathcal{E}_{L,0}$. Thus we have a map $s : \mathcal{A} \rightarrow \mathcal{M}$ given by $a \mapsto \mathcal{E}_a$. By construction $h(\mathcal{E}_{L,a}) = a$ thus s is a section of the Hitchin map $h : \mathcal{M} \rightarrow \mathcal{A}$. Thus we see that $\mathcal{A} \cong s(\mathcal{A}) \subset W_{\mathcal{E}_{L,0}}^+$ is a subvariety of an affine variety which is isomorphic to a vector space of the same dimension, therefore $s(\mathcal{A}) = W_{\mathcal{E}_{L,0}}^+$, and so the upward flow $W_{\mathcal{E}_{L,0}}^+$ is just a Hitchin section.

In particular, this implies that if $\mathcal{E} \in W_{\mathcal{E}_{L,0}}^+$, i.e. it has a full filtration as in Proposition 3.4 such that the associated graded is isomorphic to $\mathcal{E}_{L,0}$, then it has to be isomorphic to a Higgs bundle of the form (E_L, Φ_a) . For example, the underlying bundle must split as a direct of line bundles.

If we choose $L = \mathcal{O}_C$ then we have

$$\mathcal{E}_0 := \mathcal{E}_{\mathcal{O}_C,0} = E_{\mathcal{O}_C} \xrightarrow{\Phi_0} E_{\mathcal{O}_C} \otimes K \quad (3.16)$$

the *canonical uniformising Higgs bundle*. This will give a *canonical section* $W_{\mathcal{E}_0}^+ \subset \mathcal{M}$ of the Hitchin map, corresponding to the structure sheaves of the spectral curves under the BNR correspondence, see (5.18).

3.3 Downward flows on \mathcal{M}

Recall that the core of \mathcal{M} is defined as $\mathcal{C} := \coprod_{\alpha} W_{\alpha}^- = \coprod_F W_F^-$. From Proposition 2.10 we know that $W_F^- \subset M^s$ is a Lagrangian subvariety. Thus the core \mathcal{C} is Lagrangian at its smooth points.

By recalling the \mathbb{T} -equivariant map $h : \mathcal{M} \rightarrow \mathcal{A}$ where \mathbb{T} acts on \mathcal{A} with positive weights we see that the core of \mathcal{A} is the origin $0 \in \mathcal{A}$, and so $\mathcal{C} \subset h^{-1}(0)$. On the other hand h is proper, therefore $h^{-1}(0)$ is projective, thus $h^{-1}(0) \subset \mathcal{C}$. The zero fibre of the

²The terminology is motivated by the $n = 2$ case when the PGL_2 Higgs bundle associated to our uniformising Higgs bundle corresponds by non-abelian Hodge theory to the representation $\pi_1(C) \rightarrow \mathrm{PGL}_2(\mathbb{R}) \subset \mathrm{PGL}_2(\mathbb{C})$ giving the uniformising hyperbolic metric on C .

Hitchin map $h^{-1}(0)$ is called the *nilpotent cone*, because $h(E, \Phi) = 0$ if and only if Φ is nilpotent. We will think of the nilpotent cone $N := h^{-1}(0)$ as a subscheme of \mathcal{M} . In particular, we will see later that most of its components are non-reduced. The argument above then implies

Proposition 3.9. *The reduced scheme of the nilpotent cone coincides with the core: $N_{red} = \mathcal{C}$.*

Remark 3.10. The core is a Lagrangian subvariety. Thus Proposition 3.9 shows that the nilpotent cone is Lagrangian, which is a result of Laumon [La1, Theorem 3].

We finish with the analogue of Proposition 3.4 for the downward flow.

Proposition 3.11. *Let $\mathcal{E} = (E, \Phi) \in \mathcal{M}$ then*

$$\lim_{\lambda \rightarrow \infty} (E, \lambda\Phi) = (E', \Phi')$$

for some $(E', \Phi') \in \mathcal{M}^{s\mathbb{T}}$ if and only if there exists a filtration

$$0 = E_0 \subset E_1 \subset \cdots \subset E_{k-1} \subset E_k = E \tag{3.17}$$

by subbundles such that for all i

$$\Phi(E_{i+1}) \subset E_i K.$$

Thus it induces maps $gr_\infty(\Phi) : E_i/E_{i-1} \rightarrow E_{i-1}/E_{i-2}K$ and the corresponding graded Higgs bundle agrees with the limiting Higgs bundle

$$(E_1/E_0 \oplus \cdots \oplus E_{k-1}/E_{k-2} \oplus E_k/E_{k-1}, gr_\infty(\Phi)) \cong (gr(E), gr_\infty(\Phi)) \cong (E', \Phi').$$

Proof. The proof is an appropriate modification of the proof of Proposition 3.4. Just as in Remark 3.6 we can think of the filtration (3.17) on the Higgs bundle induced from the downward flows on the projectivised universal bundle. \square

4 Very stable Higgs bundles

4.1 Definition and basic properties

Drinfeld and Laumon in [La1, Definition 3.4] call a vector bundle E on C *very stable*, if the only nilpotent Higgs field $\Phi \in H^0(\text{End}(E) \otimes K)$ is the trivial one. [La1, Proposition 3.5] proves that a rank n very stable bundle is stable, and that very stable bundles form an open dense subset of the moduli space of stable bundles.

Definition 4.1. Let $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ be a stable \mathbb{T} -invariant Higgs bundle. We call \mathcal{E} a *very stable Higgs bundle* if and only if the only nilpotent Higgs bundle in $W_{\mathcal{E}}^+$ is \mathcal{E} . On the other hand we call \mathcal{E} *wobbly*, when it is not very stable.

In this language [La1, Proposition 3.5] implies

Theorem 4.2. *There exist very stable type (n) Higgs bundles $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$. In fact, they form an open dense subset of $\mathcal{N} \in \pi_0(\mathcal{M}^{s\mathbb{T}})$.*

The following was proved for very stable bundles E in [PPN, Theorem 1.1].

Lemma 4.3. *The stable Higgs bundle $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable if and only if the upward flow $W_{\mathcal{E}}^+ \subset \mathcal{M}$ is closed.*

Proof. This is Proposition 2.14 applied to the semi-projective variety \mathcal{M} . \square

Remark 4.4. Recall from Remark 3.8 that the upward flow $W_{\mathcal{E}_{L,0}}^+ \subset \mathcal{M}$ of the uniformising Higgs bundle $\mathcal{E}_{L,0}$ is a Hitchin section $h : W_{\mathcal{E}_{L,0}}^+ \cong s(\mathcal{A})$. Thus $W_{\mathcal{E}_{L,0}}^+ \cap h^{-1}(0) = \{\mathcal{E}_{L,0}\}$. Therefore the Higgs bundle $\mathcal{E}_{L,0}$ is very stable. In particular, by the Lemma above, $W_{\mathcal{E}_{L,0}}^+ \subset \mathcal{M}$ is closed. This also follows from the fact that it is a section of the Hitchin map, which was already pointed out in [Hi3, p. 454].

Lemma 4.5. *When $\mathcal{E} \in \mathcal{M}^{\mathbb{T}}$ is very stable then $h : W_{\mathcal{E}}^+ \rightarrow \mathcal{A}$ is finite, flat, surjective and generically étale. In particular, when $\mathcal{E} \in \mathcal{M}^{\mathbb{T}}$ is very stable then $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})$ is locally free.*

Proof. As $W_{\mathcal{E}}^+ \subset \mathcal{M}$ is closed by Lemma 4.3 the restriction of the Hitchin map $h : W_{\mathcal{E}}^+ \rightarrow \mathcal{A}$ is proper. Furthermore since $h : W_{\mathcal{E}}^+ \rightarrow \mathcal{A}$ is a map between affine varieties (cf. Proposition 2.1), it is quasi-finite. As a proper and quasi-affine map it is finite (c.f [Gr, Theorem 8.11.1]). Finally, it is a map between smooth equidimensional varieties and so by miracle flatness (c.f. [Ma, Corollary 23.1]) h is flat. Thus h is finite and flat thus $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})$ is locally free.

Finally, as $h : W_{\mathcal{E}}^+ \rightarrow \mathcal{A}$ is finite and between equidimensional varieties, it is surjective, and thus generically smooth, and so generically étale. \square

4.2 Hecke transformations

For a point $c \in C$ recall the exact sequence of sheaves

$$0 \rightarrow \mathcal{O}_C(-c) \xrightarrow{s_c} \mathcal{O}_C \rightarrow \mathcal{O}_c \rightarrow 0. \quad (4.1)$$

Let $E \rightarrow C$ be a vector bundle. By abuse of notation we will also denote by E its locally-free sheaf of sections. Tensoring (4.1) with the flat E gives

$$0 \rightarrow E(-c) \rightarrow E \xrightarrow{f} (E_c)_c \rightarrow 0.$$

Here E_c denotes the fibre of the vector bundle E at c and for any vector space U we denote $U_c := U \otimes \mathcal{O}_c$. In particular, $(E_c)_c = E \otimes \mathcal{O}_c$. If now $V \subset E_c$ then $V_c \subset (E_c)_c$ is a subsheaf and if $W := E_c/V$ then W_c is a quotient sheaf of $(E_c)_c$. Denoting by $\pi_V : (E_c)_c \rightarrow W_c$ the quotient map and $f_V = f \circ \pi_V$ we make the following

Definition 4.6. Let E be a vector bundle on the curve C . Let $V < E_c$ be a subspace at a point $c \in C$. The *Hecke transform* of E at V is defined to be $E_V := \ker f_V$.

Equivalently, the following short exact sequence of sheaves defines the Hecke transformed vector bundle:

$$0 \rightarrow E_V \rightarrow E \xrightarrow{f_V} W_c \rightarrow 0.$$

As a subsheaf of a torsion free sheaf E_V is torsion free and so a locally free sheaf, i.e. the sheaf of sections of a vector bundle which we also denote by E_V or sometimes $E' = E_V$.

Remark 4.7. By construction $E(-c) \subset E_V$ is a subsheaf and fits into a short exact sequence

$$0 \rightarrow E(-c) \rightarrow E' \rightarrow V_c \rightarrow 0.$$

This way we get a surjective map $E'_c \rightarrow V$. Denote by $V' < E'_c$ its kernel. Then we see that $E'_{V'} = E(-c)$.

Let now (E, Φ) be a Higgs bundle. let $V < E_c$ be a Φ_c -invariant subspace of dimension $n-k$. Then $W := E_c/V$ is of dimension k . Denote by $\bar{\Phi}_c$ the morphism $W_c \rightarrow W_c \otimes K$ induced by Φ . This data induces a unique Higgs bundle (E', Φ') making the following diagram commutative.

$$\begin{array}{ccccccc} 0 & \rightarrow & E' & \rightarrow & E & \xrightarrow{f_V} & W_c & \rightarrow & 0 \\ & & \Phi' \downarrow & & \Phi \downarrow & & \bar{\Phi}_c \downarrow & & \\ 0 & \rightarrow & E'K & \rightarrow & EK & \rightarrow & W_cK & \rightarrow & 0 \end{array} \quad (4.2)$$

Definition 4.8. Let (E, Φ) be a Higgs bundle on the curve C and $V < E_c$ a subspace at $c \in C$. The *Hecke transform* of (E, Φ) at $V < E_c$ is the unique Higgs bundle (E', Φ') making the diagram (4.2) commutative. Sometimes we will denote it $\mathcal{H}_V(E, \Phi) = (E', V')$ or by $(E_V, \Phi_V) = (E', \Phi')$.

Remark 4.9. As in Remark 4.7, the Hecke transformation will produce an $n-k$ dimensional subspace $V' < E'_c$ such that the Hecke transform of E' at V' is $E(-c)$. We note that the Hecke transformed Higgs field $\Phi'' : E(-c) \rightarrow E(-c)K$ as a section of $\text{End}(E(-c))K = \text{End}(E)K$ agrees with $\Phi \in \text{End}(E)K$ away from c , and so agrees everywhere. Thus the Hecke transform of (E', Φ') at V' is the Higgs bundle $(E(-c), \Phi)$.

Remark 4.10. We also note that as Φ' and Φ agree away from c so their characteristic polynomial will also agree away from c , therefore $h(\Phi') = h(\Phi) \in \oplus_{i=1}^n H^0(C; K^i)$. Therefore the Hecke transformation is along the Hitchin fibration. In the case of a point $a \in \mathcal{A}$ for which the spectral curve C_a is smooth, the fibre is isomorphic to the Jacobian of C_a and at a generic point $c \in C$, the subspace $V < E_c$ is a sum of $(n-k)$ eigenspaces of Φ_c . Since C_a is the curve of eigenvalues this is a degree $(n-k)$ effective divisor D and the Hecke transform takes a line bundle U on the spectral curve to $U(-D)$.

4.2.1 The Hecke transform of a full compatible filtration

Here we investigate what happens to a compatible full filtration on a Higgs bundle, when we perform a Hecke transformation at a point $c \in C$ where $b(c) \neq 0$.

Let a rank n Higgs bundle (E, Φ) carry a full filtration

$$E_0 = 0 \subset E_1 \subset \cdots \subset E_n = E \quad (4.3)$$

by $\text{rank}(E_i) = i$ subbundles, such that

$$\Phi(E_i) \subset E_{i+1} \otimes K. \quad (4.4)$$

Then we have the induced maps

$$b_i =: E_i/E_{i-1} \rightarrow E_{i+1}/E_i \otimes K \quad (4.5)$$

between line bundles, and the composition

$$b = b_{n-1} \circ \cdots \circ b_1 : E_1 \rightarrow E/E_{n-1} \otimes K. \quad (4.6)$$

Suppose that b does not vanish at $c \in C$. This means that for all i the map $\Phi_c : E_i|_c \rightarrow E_{i+1}|_c \otimes K|_c$ does not preserve $E_i|_c$. To perform the Hecke transformation we choose a Φ_c -invariant subspace $V < E_c$ of $\dim(V) = k < n$. We need the following linear algebra

Lemma 4.11. *Let $U_0 = 0 < U_1 < \dots < U_n < U$ be a full flag of subspaces of an n -dimensional complex vector space U . Furthermore let A be a linear transformation $A : U \rightarrow U$ such that $A(U_i) \subset U_{i+1}$ but U_i is not A -invariant. If $V < U$ of $\dim(V) = k < n$ is A -invariant, then $\dim(V \cap U_i) = \max(0, k + i - n)$.*

Proof. By assumption $U_i + A(U_i) = U_{i+1}$. Thus $V + U_i = V + U_{i+1}$ if and only if $V + U_i$ is invariant under A . On the other hand $V + U_i = V + U_{i+1}$ in turn implies $V + U_i = V + U_n = U$. Thus we see that the chain of subspaces

$$V = V + U_0 \leq V + U_1 \leq \cdots \leq V + U_{n-1} \leq V + U_n = U$$

must be strictly increasing until it reaches the whole ambient space U . Therefore $\dim(V + U_i) = \min(k + i, n)$, which implies the result. \square

The Φ_c -invariant subspace $V < E_c$ induces a map of sheaves $f_V : E \rightarrow W_c$, where $W := E_c/V$. This induces the Hecke transformed Higgs bundle (E', Φ') :

$$\begin{array}{ccccccc} 0 & \rightarrow & E' & \rightarrow & E & \xrightarrow{f_V} & W_c & \rightarrow 0 \\ & & \Phi' \downarrow & & \Phi \downarrow & & \overline{\Phi}_c \downarrow & \\ 0 & \rightarrow & E'K & \rightarrow & EK & \rightarrow & W_cK & \rightarrow 0 \end{array} .$$

Each subbundle E_i will also induce a Hecke transformed line bundle with respect to the subspace $V_i := E_i|_c \cap V < E_i|_c$. Denoting $W^i := E_i|_c/V_i$ we get a new vector bundle E'_i from the kernel of the map $f_{V_i} : E_i \rightarrow W^i_c$. In other words we have the top row of the following commutative diagram

$$\begin{array}{ccccccccc}
0 & \rightarrow & E'_i & \rightarrow & E_i & \xrightarrow{f_{V_i}} & W^i_c & \rightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \rightarrow & E'_{i+1} & \rightarrow & E_{i+1} & \xrightarrow{f_{V_{i+1}}} & W^{i+1}_c & \rightarrow & 0 \\
& & \downarrow & & \downarrow & & \downarrow & & \\
0 & \rightarrow & E'_{i+1}/E'_i & \rightarrow & E_{i+1}/E_i & \xrightarrow{\bar{f}_{V_{i+1}}} & (W^{i+1}/W^i)_c & \rightarrow & 0
\end{array} \tag{4.7}$$

Here the embedding $E'_i \hookrightarrow E'_{i+1}$ is uniquely induced from the rest of the first two top rows of the diagram. The quotient maps connecting the second and third rows of the diagram are the ones making the columns into short exact sequences. The last row, and in particular the map \bar{f}_{V_i} , is then uniquely induced from the rest of the diagram to make it commutative.

From Lemma 4.11 we have

$$\dim(W^{i+1}/W^i) = \begin{cases} 1 & \text{for } i < n - k \\ 0 & \text{for } i \geq n - k \end{cases} \tag{4.8}$$

Thus from the bottom row of (4.7) we obtain

$$E'_{i+1}/E'_i \cong \begin{cases} (E_{i+1}/E_i)(-c) & \text{for } i < n - k \\ (E_{i+1}/E_i) & \text{for } i \geq n - k \end{cases} \tag{4.9}$$

In particular, E'_{i+1}/E'_i is a line bundle, and so $E'_i \subset E'_{i+1}$ is a subbundle. Thus we get a full filtration of E' by subbundles:

$$E'_0 = 0 \subset E'_1 \subset \dots \subset E'_n = E \tag{4.10}$$

By construction it is compatible with Φ' i.e.

$$\Phi'(E'_i) \subset E'_{i+1}K. \tag{4.11}$$

Finally we will determine the maps $b'_i : E'_i/E'_{i-1} \rightarrow E'_{i+1}/E'_i \otimes K$ induced from Φ' . We have the commutative diagram

$$\begin{array}{ccccccccc}
0 & \rightarrow & E'_i/E'_{i-1} & \rightarrow & E_i/E_{i-1} & \xrightarrow{\bar{f}_{V_i}} & (W^i/W^{i-1})_c & \rightarrow & 0 \\
& & b'_i \downarrow & & b_i \downarrow & & \bar{b}_i \downarrow & & \\
0 & \rightarrow & (E'_{i+1}/E'_i) \otimes K & \rightarrow & (E_{i+1}/E_i) \otimes K & \xrightarrow{\bar{f}_{V_{i+1}}} & (W^{i+1}/W^i)_c \otimes K & \rightarrow & 0
\end{array}$$

Here \bar{b}_i is induced from Φ_c . When $i > n - k$ then from (4.8) we see that \bar{b}_i maps between zero-dimensional spaces, which shows that

$$b'_i = b_i \in H^0(C; (E'_i/E'_{i-1})^*(E'_{i+1}/E'_i)K) = H^0(C; (E_i/E_{i-1})^*(E_{i+1}/E_i)K). \quad (4.12)$$

When $i < n - k$ then \bar{b}_i is induced from an isomorphism of 1-dimensional vector spaces, and we get again the agreement (4.12).

Furthermore, when $i = n - k$ then W^i/W^{i-1} is one-dimensional and $W^{i+1}/W^i = 0$. This implies that $b'_i = b_i s_c$ where $s_c \in H^0(C; \mathcal{O}(c)) \cong H^0(C; \text{Hom}(\mathcal{O}(-c), \mathcal{O}))$ is the defining (4.1) section vanishing at c .

Finally by tracing the kernel of the induced maps on the fibres at c in the diagram (4.7) and using (4.8) we see that $V' = (E'_{n-k})_c < E'_c$. Thus $\mathcal{H}_{V'}(E', \Phi') = (E(-c), \Phi)$.

We proved the following

Proposition 4.12. *Let a Higgs bundle (E, Φ) carry a full (4.3) and compatible (4.4) filtration, such that b (4.6) does not vanish at c . Let $V < E_c$ be Φ_c -invariant. Then the Hecke transformed Higgs bundle $\mathcal{H}_V(E, \Phi) = (E', \Phi')$ carries an induced full filtration (4.10) which is compatible with Φ' (4.11). Furthermore we have (4.9) and $b'_i = b_i$ unless $i = n - k$ when $b'_{n-k} = b_{n-k} s_c$ acquires a simple zero at c . Finally, the subspace of Remark 4.9 is $V' = (E'_{n-k})_c < E'_c$. Thus $\mathcal{H}_{V'}(E', \Phi') = (E(-c), \Phi)$.*

4.2.2 Very stable Higgs bundles via Hecke transformations

For $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1}) \in J_\ell(C) \times C^{[m_1]} \times \dots \times C^{[m_{n-1}]}$ let $\mathcal{E}_\delta \in \mathcal{M}^{s\mathbb{T}}$ be a type $(1, \dots, 1)$ fixed point of the \mathbb{T} -action constructed in Example 3.2.

Theorem 4.13. *Let $\mathcal{E}_\delta \in \mathcal{M}^{s\mathbb{T}}$. Then \mathcal{E}_δ is very stable if and only if $b = b_1 \circ \dots \circ b_{n-1} \in H^0(C; L_0^* L_{n-1} K)$ has no repeated zero, i.e. if and only if the effective divisor $\text{div}(b) = \delta_1 + \dots + \delta_{n-1}$ is reduced.*

Proof. First we prove the if part. The proof is by induction on $\text{deg}(b)$. When b has no zeroes, then $\mathcal{E}_\delta \cong \mathcal{E}_{L_0, 0}$ are the uniformising Higgs bundles from Remark 3.8. Thus $W_{\mathcal{E}_\delta}^+$ is a Hitchin section. In particular \mathcal{E}_δ is its intersection with the nilpotent cone $h^{-1}(0)$ (c.f. Remark 4.4). Thus \mathcal{E}_δ is very stable when $\text{deg}(b) = 0$.

Let $\mathcal{E}_\delta \in \mathcal{M}^{s\mathbb{T}}$ be such that b has no repeated zero, but at least one zero say $c \in C$ with $b(c) = 0$. Assume that it is $b_k(c) = 0$. Then $V_{1k} := (L_0 \oplus \dots \oplus L_{k-1})_c < E_c$ is Φ_c invariant. Thus $\mathcal{H}_{V_{1k}}(\mathcal{E}_\delta) = \mathcal{E}_{\delta'}$ where $\delta'_i = \delta_i$ i.e. $b'_i = b_i$ except if $i = k$ when $b'_k = b_k/s_c$ i.e. $\delta'_k = \delta_k - c$. In particular, b' does not vanish at c . We also have

Lemma 4.14. *$\mathcal{E}_{\delta'}$ is a stable Higgs bundle.*

Proof. As by assumption \mathcal{E}_δ is stable all proper Φ -invariant subbundles have slope less than the slope of E . As the only Φ -invariant subbundles are $L_i \oplus \cdots \oplus L_{n-1} \subset E$, we have for every $0 \leq i \leq n-1$ the following inequality

$$\frac{\ell_i + \cdots + \ell_{n-1}}{n-i} < \frac{\ell_0 + \cdots + \ell_{n-1}}{n}.$$

To check that $\mathcal{E}_{\delta'}$ is stable we check that

$$\begin{aligned} \frac{\ell'_i + \cdots + \ell'_{n-1}}{n-i} &= \frac{\ell_i + \cdots + \ell_{n-1} - \min(n-i, n-k)}{n-i} < \frac{\ell_0 + \cdots + \ell_{n-1}}{n} - \frac{\min(n-i, n-k)}{n-i} \\ &< \frac{\ell'_0 + \cdots + \ell'_{n-1} + n-k}{n} - \frac{\min(n-i, n-k)}{n-i} < \frac{\ell'_0 + \cdots + \ell'_{n-1}}{n} \end{aligned}$$

because

$$\frac{n-k}{n} \leq \frac{\min(n-i, n-k)}{n-i}.$$

□

Assume now that we have a nilpotent Higgs bundle $\mathcal{E} = (E, \Phi) \in W_{\mathcal{E}_\delta}^+$. From Proposition 3.4 let $E_1 \subset \cdots \subset E_n = E$ be the Φ -invariant filtration inducing $\lim_{\lambda \rightarrow 0} \lambda \cdot \mathcal{E} = \mathcal{E}_\delta$. Let $V := (E_k)_c < E_c$, which by the assumption of $b_k(c) = 0$ is Φ_c -invariant. As we saw in the previous subsection $\mathcal{H}_V(E, \Phi) = (E', \Phi')$ is nilpotent in the upward flow of $\mathcal{H}_{V_{1k}}(\mathcal{E}_\delta) = \mathcal{E}_{\delta'}$, because $\mathcal{E}_{\delta'}$ is stable by Lemma 4.14. But $\deg(b') = \deg(b) - 1$ and so by induction we can assume that $\mathcal{E}_{\delta'}$ is very stable and hence the nilpotent (E', Φ') in its upward flow should be isomorphic to it: $(E', \Phi') \cong \mathcal{E}_{\delta'}$. Now $b'(c) \neq 0$ therefore Φ'_c is regular nilpotent. Hence there is a unique Φ'_c -invariant $(n-k)$ -dimensional subspace of E' , namely $V'_{k, n-1} = (L_k)_c \oplus \cdots \oplus (L_{n-1})_c$. The Hecke transform of $(E', \Phi') \cong \mathcal{E}_{\delta'}$ at this subspace gives $(E(-c), \Phi) \cong \mathcal{E}_{\delta-c}$, where

$$\delta - c := (\delta_0 - c, \delta_1, \dots, \delta_{n-1}).$$

This shows that \mathcal{E}_δ is also very stable. This proves the “if” part of the theorem.

For the only if part, when b has repeated zeroes, the strategy will be to produce a Hecke curve [NR] - a one-parameter family of Higgs bundles produced as Hecke transforms of a fixed Higgs bundle - in the intersection of the nilpotent cone and the upward flow $W_{\mathcal{E}_\delta}^+$ originating at \mathcal{E}_δ .

Take a multiple zero c of b . Assume that $b_k(c) = 0$. Then $V := (L_0 \oplus \cdots \oplus L_{k-1})_c < E_c$ is Φ_c -invariant. The Hecke transformed Higgs bundle $\mathcal{H}_V(\mathcal{E}_\delta) = \mathcal{E}_{\delta'}$ has $b'(c) = 0$ otherwise $\mathcal{H}_{V'}(\mathcal{E}_{\delta'})$ would have $b(c) = 0$ a simple zero from Proposition 4.12.

Thus $b'(c) = 0$ and as before in Lemma 4.14 $\mathcal{E}_{\delta'}$ is still stable. Now however the nilpotent Φ'_c is no longer regular due to $b'(c) = 0$. This implies that, besides $V' = (L'_k \oplus \cdots \oplus L'_{n-1})_c$, there are more than one Φ'_c -invariant $n-k$ -dimensional subspaces

in E'_c . By [Sh, Theorem 6] the subvariety $S_{n-k}(\Phi'_c) \subset Gr_{n-k}(E'_c)$ of Φ'_c -invariant $(n-k)$ -dimensional subspaces in the Grassmannian of $(n-k)$ -planes in E'_c is connected.

We will find a curve in $S_{n-k}(\Phi'_c)$, leading to a Hecke curve in \mathcal{M} by studying a natural \mathbb{T} -action on the projective variety $S_{n-k}(\Phi'_c)$. We note that $\mathcal{E}_{\delta'} = (E', \Phi')$ is \mathbb{T} -invariant, meaning that

$$(E', \Phi') \cong (E', \lambda\Phi') \quad (4.13)$$

for all $\lambda \in \mathbb{C}$. The isomorphism in (4.13) is induced by an automorphism of (E', Φ') . Let \mathbb{T} act on $E' = L'_0 \oplus \cdots \oplus L'_{n-1}$ with weight $-i$ on L'_i . This \mathbb{T} -action induces the weight -1 action of \mathbb{T} on Φ' , showing (4.13). Restricting this \mathbb{T} -action to E'_c we get an action of \mathbb{T} on $Gr_{n-k}(E'_c)$. We note that $\Phi'_c(\lambda \cdot v) = \lambda \cdot \Phi'_c(v)$ and thus if $V \subset E'_c$ is a Φ'_c -invariant subspace then $\lambda \cdot V$ is also Φ'_c -invariant. Therefore this \mathbb{T} -action leaves $S_{n-k}(\Phi'_c)$ invariant. We have $S_{n-k}(\Phi'_c)^\mathbb{T} \subset Gr_{n-k}(E'_c)^\mathbb{T}$. Fixed points of the \mathbb{T} -action on $Gr_{n-k}(E'_c)$ are of the form $(L_{i_1} \oplus \cdots \oplus L_{i_{n-k}})_c$ for any $(n-k)$ -element subset $\{i_1, \dots, i_{n-k}\} \subset \{0, \dots, n-1\}$. We have

$$V' = (L'_{k}, \dots, L'_{n-1})_c \in S_{n-k}(\Phi'_c)^\mathbb{T} \subset Gr_{n-k}(E'_c).$$

As $S_{n-k}(\Phi'_c)$ is projective, connected and contains more than one point, there is $V' \neq V \in S_{n-k}(\Phi'_c)$ such that $\lim_{\lambda \rightarrow 0} \lambda V = V'$. Let

$$V_\infty := \lim_{\lambda \rightarrow \infty} \lambda \cdot V \in S_{n-k}(\Phi'_c)^\mathbb{T}.$$

As the \mathbb{T} -action on $S_{n-k}(\Phi'_c)$ is linear we see that $V' \neq V_\infty$. Let $V_\infty = (L_{i_1} \oplus \cdots \oplus L_{i_{n-k}})_c$ for some

$$\{k, \dots, n-1\} \neq \{i_1, \dots, i_{n-k}\} \subset \{0, \dots, n-1\}. \quad (4.14)$$

We thus have a map $\mathbb{P}^1 \rightarrow S_{n-k}(\Phi'_c)$ sending 0 to V' , ∞ to V_∞ and $\lambda \in \mathbb{T}$ to $\lambda \cdot V$. We will denote the corresponding subspaces V_t for $t \in \mathbb{C} \cup \infty$.

First we need the following

Lemma 4.15. $\mathcal{H}_{V_\infty}(\mathcal{E}_{\delta'})$ is a \mathbb{T} -fixed stable Higgs bundle.

Proof. Let $\mathcal{H}_{V_\infty}(\mathcal{E}_{\delta'}) = (E'', \Phi'')$. Then $E'' \cong L''_0 \oplus \cdots \oplus L''_{n-1}$ where $L''_i \cong L'_i$ when $i \in \{i_1, \dots, i_{n-k}\}$ and $L''_i \cong L'_i(-c)$ when $i \notin \{i_1, \dots, i_{n-k}\}$. Thus

$$\ell''_i := \deg(L''_i) = \begin{cases} \ell'_i & \text{when } i \in \{i_1, \dots, i_{n-k}\} \\ \ell'_i - 1 & \text{when } i \notin \{i_1, \dots, i_{n-k}\} \end{cases} \quad (4.15)$$

To check stability we compute

$$\begin{aligned} \frac{\ell''_i + \cdots + \ell''_{n-1}}{n-i} &\leq \frac{\ell'_i + \cdots + \ell'_{n-1} - \max(0, k-i)}{n-i} = \frac{\ell_i + \cdots + \ell_{n-1} - (n-i)}{n-i} \\ &< \frac{\ell_0 + \cdots + \ell_{n-1} - n}{n} = \frac{\ell''_0 + \cdots + \ell''_{n-1}}{n} \end{aligned}$$

□

Let $V \in S_{n-k}(E_c)$ and denote $V_\lambda := \lambda \cdot V$. Denote by $f_\lambda : E' \rightarrow E'$ the automorphism induced by our \mathbb{T} -action on E' as in (3.3). Finally identify $(K)_c \cong \mathbb{C}$. We now have the commutative diagram:

$$\begin{array}{ccccccc}
 & & E'_V K & \longrightarrow & E' K & \longrightarrow & (E_c/V)_c \\
 & \nearrow \lambda \Phi'_V & \downarrow & & \nearrow \lambda \Phi' & \downarrow & \nearrow \lambda \bar{\Phi}'_c \\
 E'_V & \longrightarrow & & \longrightarrow & E' & \longrightarrow & (E_c/V)_c \\
 & \downarrow f''_\lambda & & & \downarrow f'_\lambda & & \downarrow \bar{f}'_{\lambda c} \\
 & & E'_{V_\lambda} K & \longrightarrow & E' K & \longrightarrow & (E_c/V_\lambda)_c \\
 & \nearrow \Phi'_{V_\lambda} & \downarrow f'_\lambda & & \nearrow \Phi' & \downarrow \bar{f}'_{\lambda c} & \nearrow \bar{\Phi}'_c \\
 E'_{V_\lambda} & \longrightarrow & & \longrightarrow & E' & \longrightarrow & (E_c/V_\lambda)_c
 \end{array}$$

The map $f''_\lambda : E'_V \rightarrow E'_{V_\lambda}$ is defined uniquely so as to make the diagram commutative. The diagram shows that f''_λ induces $(E'_{V_\lambda}, \Phi'_{V_\lambda}) \cong (E'_V, \lambda \cdot \Phi'_V)$.

Therefore $\mathcal{H}_{V_i}(\mathcal{E}_\delta)$ is stable for all t since stability is an open condition for Higgs bundles (c.f. [Si1, Lemma 3.7] or [Ni, Proposition 3.1]). Finally, we note that $\mathcal{H}_{V_0}(\mathcal{E}_\delta) \cong \mathcal{E}_{\delta-c} \not\cong \mathcal{H}_{V_\infty}(\mathcal{E}_\delta)$ because there exists an i such that $\deg(L''_i) \neq \deg(L_i(-c))$ because of (4.14) and (4.15). Thus we found a \mathbb{T} -invariant family of nilpotent stable Higgs bundles in \mathcal{M}^s parametrized by \mathbb{P}^1 connecting $\mathcal{E}_{\delta-c}$ and $\mathcal{H}_{V_\infty}(\mathcal{E}_\delta)$. Thus $W_{\mathcal{E}_{\delta-c}}^+ \cap h^{-1}(0)$ is not trivial. Therefore $\mathcal{E}_{\delta-c}$ and so \mathcal{E}_δ is not very stable. The theorem follows. \square

Corollary 4.16. *There is a very stable Higgs bundle in every type $(1, \dots, 1)$ component $F_{\ell, m} \in \pi_0(\mathcal{M}^{s\mathbb{T}})$ from (3.4). In fact, they form a dense open subset.*

Proof. From Theorem 4.13 the very stable locus in $F_{\ell, m}$ are Higgs bundles \mathcal{E}_δ where b has no repeated zero. Thus they form the complement of some divisor. The result follows. \square

4.2.3 Example in rank 2

Let L_0 and L_1 be line bundles on C and $E = L_0 \oplus L_1$ together with a lower triangular Higgs field: $\Phi = \begin{pmatrix} 0 & 0 \\ b & 0 \end{pmatrix}$ with $b = b_1 \in H^0(C; L_0^* L_1 K)$. Assume that $\deg(b) < 2g - 2$ so that (E, Φ) is stable. Suppose further that $b = s_c^2 b''$ has a double zero at $c \in C$ where $b'' \in H^0(C; L_0^* L_1 K(-2c))$. Let $V := (L_0)_c < E_c$. It is Φ_c -invariant as $\Phi_c = 0$. We have

$$E' = L'_0 \oplus L'_1 = L_0 \oplus L_1(-c)$$

and $V' = V_0 = (L'_1)_c$. As $b' = s_c b''$ still vanishes at c , $\Phi'_c = 0$. Thus $S_1(\Phi'_c) = \mathbb{P}(E_c)$ the whole \mathbb{P}^1 . Let $V_\infty = (L_0)_c < E'_c$ which is Φ'_c -invariant. Fix basis vectors $\langle v_1 \rangle = (L'_1)_c$ and $\langle v_2 \rangle = (L'_2)_c$ then let $V_t := \langle tv_1 + v_2 \rangle$. We obtain $\lim_{t \rightarrow 0} V_t = V'$ and $\lim_{t \rightarrow \infty} V_t = V_\infty$. Then we see that $\mathcal{H}_{V'}(E', \Phi') = (E(-c), \Phi)$ and $\mathcal{H}_{V_\infty}(E', \Phi') = (E'', \Phi'')$. Here

$$E'' = L''_0 \oplus L''_1 = L_0 \oplus L_1(-2c)$$

and $\Phi'' = \begin{pmatrix} 0 & 0 \\ b'' & 0 \end{pmatrix}$. For $t \neq 0, \infty$ the Hecke transforms E'_{V_t} are no longer direct sums but extensions

$$0 \rightarrow L_0(-c) \rightarrow E'_{V_t} \rightarrow L_1(-c) \rightarrow 0, \quad (4.16)$$

or

$$0 \rightarrow L_1(-2c) \rightarrow E'_{V_t} \rightarrow L_0 \rightarrow 0. \quad (4.17)$$

The first one corresponds to the modification of $E'_1 = L'_0 = L_0$ in E'_{V_t} . In other words (4.16) is the modified filtration of Proposition 4.12. This shows that

$$\lim_{\lambda \rightarrow 0} \lambda \cdot \mathcal{H}_{V_t}(E', \Phi') = (E(-c), \Phi).$$

In particular, as the latter is stable so is $\mathcal{H}_{V_t}(E', \Phi')$.

The second extension (4.17) is induced from $L_1(-2c) \subset E'_{V_t}$ the Hecke modification of the subbundle $L'_1 \cong L_1(-c) \subset E'$. As Φ' was trivial on L'_1 the modification Φ'_{V_t} will be trivial on $L_1(-2c)$. Thus Φ'_{V_t} will be given by projection to L_0 in (4.17) followed by $b'' : L_0 \rightarrow L_1(-2c)K$. As in Proposition 3.11 this shows that

$$\lim_{\lambda \rightarrow \infty} \lambda \cdot \mathcal{H}_{V_t}(E', \Phi') = (E'', \Phi'').$$

Thus $\mathcal{H}_{V_t}(E', \Phi')$ is a \mathbb{T} equivariant family of stable nilpotent Higgs bundles parametrized by $t \in \mathbb{P}^1 = \mathbb{C} \cup \infty$, connecting $\mathcal{H}_{V_0}(E', \Phi') = (E(-c), \Phi)$ and $\mathcal{H}_{V_\infty}(E', \Phi') = (E'', \Phi'')$.

This shows that $(E(-c), \Phi)$ and thus (E, Φ) are not very stable.

4.2.4 Hecke transforms of Lagrangians

We have seen above how the different upward flows of type $(1, \dots, 1)$ very stable Higgs bundles are related by Hecke transforms. We also showed they were Lagrangian in Proposition 2.10. It is a general fact that the Hecke transform takes Lagrangians to Lagrangians, as we show next.

Simpson constructed [Si1, Theorem 4.10] the fine moduli space \mathcal{R}^s of stable rank n degree d Higgs bundles, framed at the point $c \in C$. The group GL_n acts on the framing

with quotient \mathcal{M}^s so that $\mathcal{R}^s \rightarrow \mathcal{M}^s$ is a PGL_n -principal bundle, associated to the projective universal bundle $\mathbb{P}(\mathbb{E}_c)$ restricted to c . Let $H_k \subset \mathrm{GL}_n$ be the stabilizer of $\mathbb{C}^k < \mathbb{C}^n$ - a maximal parabolic subgroup of GL_n . The quotient GL_n/H_k is isomorphic to the Grassmannian of k -planes in \mathbb{C}^n and we can construct the Grassmannian bundle

$$\mathrm{Gr}_k(\mathbb{E}_c) := \mathcal{R}^s/H_k \quad (4.18)$$

over \mathcal{M}^s .

By identifying $K_c \cong \mathbb{C}$, étale locally the universal Higgs field Φ_c restricted to c , gives an endomorphism $\Phi_c \in H^0(\mathcal{M}^s; \mathrm{End}(\mathbb{E}_c))$. We will denote the set of Φ_c -invariant k -dimensional subspaces of E_c by

$$\mathcal{H}^s := \{V < \mathbb{E}_{((E, \Phi), c)} \mid \text{s.t. } (E, \Phi) \in \mathcal{M}^s, \dim(V) = k \text{ and } \Phi_c(V) \subset V\} \subset \mathrm{Gr}_k(\mathbb{E}_c).$$

The subset \mathcal{H}^s has the structure of a variety, as étale-locally in \mathcal{M}^s we can assume that some $\lambda \in \mathbb{C}$ is never an eigenvalue of Φ_c in other words $\det(\Phi_c - \lambda) \neq 0$. That means that $\Phi_c - \lambda$ is invertible and thus acts on $\mathrm{Gr}_k(\mathbb{E}_c)$ and \mathcal{H}^s in this étale neighbourhood is the fixed point variety of a morphism. By construction

$$\pi : \mathcal{H}^s \rightarrow \mathcal{M}^s \quad (4.19)$$

is a proper map.

A point of \mathcal{H}^s is represented by a pair $((E, \Phi), V) \in \mathcal{H}^s$ of a Higgs bundle (E, Φ) and k -subspace $V \in \mathrm{Gr}_k(E_c)$ such that $\Phi_c(V) \subset V$. We also consider ${}^s\mathcal{H}^s \subset \mathcal{H}^s$ the subset of points $((E, \Phi), V) \in \mathcal{H}^s$ where $V \in \mathrm{Gr}_k(E_c)$ such that $\mathcal{H}_V(E, \Phi)$ is stable. As stability is an open condition for Higgs bundles ${}^s\mathcal{H}^s \subset \mathcal{H}^s$ is an open subset. Further we define $\mathcal{M}^\# \subset \mathcal{M}$ the open subset where the spectral curve is smooth and the Higgs field Φ_c at the point $c \in C$ has distinct eigenvalues. Then we set $\mathcal{H}^\# := \pi^{-1}(\mathcal{M}^\#) \subset \mathcal{H}^s$ an open subset. The Hecke transform does not change the spectral curve, and when it is smooth the Higgs bundle is automatically stable. Thus $\mathcal{H}^\# \subset {}^s\mathcal{H}^s$.

By abuse of notation we will still denote $\pi : \mathcal{H}^\# \rightarrow \mathcal{M}^\#$ the restriction of (4.19). If $\mathcal{M} = \mathcal{M}_{n,d}$ then we denote $\mathcal{M}' := \mathcal{M}_{n,d-k}$ so that $\mathcal{H}_V(E, \Phi) \in \mathcal{M}'^s$ when $((E, \Phi), V) \in {}^s\mathcal{H}^s$. This defines $\pi' : \mathcal{H}^\# \rightarrow \mathcal{M}'^\#$. We have the following

Proposition 4.17. *The map $(\pi', \pi) : \mathcal{H}^\# \rightarrow \mathcal{M}'^\# \times \mathcal{M}^\#$ is injective.*

Proof. When the Higgs field Φ_c has n distinct eigenvalues λ_i , a Hecke transform depends on the choice of a k -element subset or equivalently a subspace $\mathbb{C}^k < \mathbb{C}^n$ spanned by the corresponding eigenspaces. If the equation of the spectral curve is given by $\det(x - \Phi) = 0$ and (E, Φ) is the direct image of the line bundle U on C_a then the Hecke transform is the direct image of $U(-D)$ where D is the effective divisor $(\lambda_{i_1}, c) + (\lambda_{i_2}, c) + \dots + (\lambda_{i_k}, c)$.

The map will be injective if for two choices of subspace the divisor classes D_1, D_2 are distinct. This will follow if $\dim H^0(C_a, \mathcal{O}(D)) = 1$. By Riemann-Roch and Serre duality this condition is equivalent to $\dim H^0(C_a, K_{C_a}(-D)) = \tilde{g} - k$ where \tilde{g} is the genus of C_a , or alternatively if the restriction $H^0(C_a, K_{C_a}) \rightarrow H^0(D, K_{C_a}) \cong \mathbb{C}^k$ is surjective.

By adjunction on the total space of K , which has trivial canonical bundle, we have $K_{C_a} \cong \pi^* K^n$ and hence sections of $\pi^* K^n$ of the form $a_0 + a_1x + \cdots + a_{n-1}x^{n-1}$ where a_i is the pullback of a section of $H^0(C, K^{n-i})$. This space has dimension $n^2(g-1) + 1 = \tilde{g}$ and so represents uniquely every section of K_{C_a} . Restricting to D gives (v_1, \dots, v_k) where

$$v_m = a_0(c) + a_1(c)\lambda_{i_m} + \cdots + a_{n-1}(c)\lambda_{i_m}^{n-1}.$$

Since the λ_i are distinct, Lagrange interpolation provides a polynomial of degree $(n-1)$ which agrees with any (v_1, \dots, v_k) and because K^i on C has no base points, there exist sections a_i agreeing at c with the coefficients of the polynomial. Hence restriction is surjective. \square

We can define a correspondence on $\mathcal{M}'^s \times \mathcal{M}^s$ by saying that two points (m, m') are equivalent if they are represented by Higgs bundles $(E, \Phi), (E', \Phi')$ where (E', Φ') is related to (E, Φ) by a k -plane Hecke transformation at c . Then, from Proposition 4.20, at a generic point this is locally given by the graph of a map and hence has a dense open set which is a manifold of half the dimension.

Recall that a *Lagrangian correspondence* between symplectic manifolds M_1, M_2 is a Lagrangian submanifold $M_1 \times M_2$ with respect to the symplectic form $p_1^*\omega_1 - p_2^*\omega_2$, for example the graph of a symplectic transformation. Then points in M_2 corresponding to those in a Lagrangian in M_1 will, under transversality conditions, form a Lagrangian submanifold. The following result was already discussed in [KW, pp.185-186].

Theorem 4.18. *The Hecke correspondence on $\mathcal{M}'^s \times \mathcal{M}^s$ is Lagrangian at a generic point.*

Proof. We use a differential-geometric approach and assume that Φ_c has distinct eigenvalues. As defined above, when we have a rank n vector bundle E on the curve C and a point $c \in C$ the Hecke transform E' is the vector bundle defined by the kernel subsheaf of a homomorphism $h : \mathcal{O}(E) \rightarrow \mathcal{O}_c^{n-k}$ with the kernel at c the vector space V .

Let v_1, \dots, v_n be a local basis of sections of E with v_1, \dots, v_k spanning V at c . Using a local coordinate where c is $z = 0$, a local basis for E' is given by u_1, \dots, u_n where $v_i = u_i$ for $i > k$ and $v_i = zu_i$ for $1 \leq i \leq k$. We shall use these two local bases to make calculations.

Varying $V \subset E_c$ gives a variation of the holomorphic structure on E' as in [NR]. A first order deformation of V in a parameter t is described by local holomorphic sections of the form

$$v_i + t\dot{v}_i = v_i + t \sum_{j=k+1}^n a_{ij}v_j$$

for $1 \leq i \leq k$. The variation of a local basis of the subsheaf is then

$$\dot{u}_i = \frac{1}{z} \sum_{j=k+1}^n a_{ij} u_j.$$

The matrix $a = a_{ij}$ in the basis u_1, \dots, u_n for E' gives the section

$$\begin{pmatrix} 0 & 0 \\ a/z & 0 \end{pmatrix}$$

over a punctured disc.

For a fixed holomorphic structure on E , this is a Čech cocycle for a class in $H^1(C, \text{End}(E'))$ defining the variation of the holomorphic structure on the Hecke transform corresponding to $a(c) \in \text{Hom}(V, E_c/V)$, a tangent vector to the Grassmannian. Here however we need to vary E also and it is convenient to use a Dolbeault representative $\dot{A} \in \Omega^{0,1}(C, \text{End}(E))$. We may make the assumption that the deformation of holomorphic structure on E is trivial in a neighbourhood of c so that we can take $\dot{A} = 0$ in this neighbourhood, extend a to a C^∞ section supported in a neighbourhood of c and holomorphic in a smaller one and take $\dot{A} + \bar{\partial}\dot{a}/z \in \Omega^{0,1}(C, \text{End}(E'))$ giving the combined deformation of holomorphic structure on E' .

Let $(\dot{A}, \dot{\Phi})$ be a first order deformation of a Higgs bundle preserving the deformation of the subsheaf. A local section of $\text{End}(E)$ can be written in block matrix form with respect to the subspaces spanned by v_1, \dots, v_k and v_{k+1}, \dots, v_n respectively as

$$\begin{pmatrix} P & Q \\ R & S \end{pmatrix}$$

and in the basis u_1, \dots, u_n it becomes

$$\begin{pmatrix} P & zQ \\ z^{-1}R & S \end{pmatrix} \tag{4.20}$$

From this we see explicitly that a Higgs field $\Phi \in H^0(C, \text{End}(E) \otimes K)$ extends to $\text{End}(E') \otimes K$ if and only if the entry R is divisible by z , i.e. Φ preserves V at c , which is the definition of \mathcal{H}^s .

To preserve the subsheaf to first order we need $(\Phi + t\dot{\Phi})(v + ta(v)) = w + ta(w)$ modulo t^2 for v and w linear combinations of v_1, \dots, v_k . It follows that $\dot{\Phi} + [\dot{\Phi}, a]$ preserves the subspace V at c .

Write

$$\Phi = \begin{pmatrix} P & Q \\ R & S \end{pmatrix} \quad \dot{\Phi} = \begin{pmatrix} \dot{P} & \dot{Q} \\ \dot{R} & \dot{S} \end{pmatrix}$$

Since $\dot{A} = 0$ in a neighbourhood the entries are holomorphic. Then

$$\dot{\Phi} + [\Phi, a] = \begin{pmatrix} \dot{P} - Qa & \dot{Q} \\ \dot{R} + Sa - aP & \dot{S} - aQ \end{pmatrix}.$$

This preserves V , so $\dot{R} + Sa - aP$ must be divisible by z . Since Φ_c has distinct eigenvalues λ_i we can take the basis vectors v_i to be eigenvectors of Φ in a neighbourhood and then the condition is that $\dot{R}_{ij} + (\lambda_i - \lambda_j)a_{ij}$ be divisible by z and accordingly $a_{ij}(0)$ is uniquely determined by $\dot{\Phi}$.

In the basis for E' we have

$$\dot{\Phi} + [\Phi, a] = \begin{pmatrix} \dot{P} - Qa & \dot{Q}z \\ (\dot{R} + Sa - aP)z^{-1} & \dot{S} - aQ \end{pmatrix}.$$

which is now holomorphic in a neighbourhood of c .

Let $(\dot{A}, \dot{\Phi}) \in \Omega^{0,1}(C, \text{End}(E)) \oplus \Omega^{1,0}(C, \text{End}(E))$ be a Dolbeault representative of a tangent vector to \mathcal{M}^s . The holomorphic symplectic form ω is defined by

$$\int_C \text{tr}(\dot{A}_1 \dot{\Phi}_2) - \text{tr}(\dot{A}_2 \dot{\Phi}_1).$$

With $\dot{A}' = \dot{A} + \bar{\partial}a/z$ as above we obtain

$$\int_C \text{tr}(\dot{A}'_1 \dot{\Phi}'_2) = \int_C \text{tr}(\dot{A}_1 \dot{\Phi}_2) + \text{tr}(\bar{\partial}a_1 Q_2/z) - \text{tr} \bar{\partial}a_1 [\Phi, a_2]/z$$

But Q_2 is holomorphic on the support of a_1 and divisible by z so by Stokes' theorem the second term vanishes. By the Jacobi identity $\text{tr}(a_1 [\Phi, a_2]) - \text{tr}(a_2 [\Phi, a_1]) = \text{tr}(\Phi [a_1, a_2])$ but the off-diagonal matrices a_1, a_2 commute so $\text{tr} \bar{\partial}a_1 [\Phi, a_2]/z = \text{tr} \bar{\partial}a_2 [\Phi, a_1]/z$. Then

$$\int_C \text{tr}(\dot{A}'_1 \dot{\Phi}'_2) = \int_C \text{tr}(\dot{A}_1 \dot{\Phi}_2)$$

and the two symplectic structures coincide. \square

5 Multiplicities in the nilpotent cone

5.1 Multiplicities of very stable components

Let $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}} \subset h^{-1}(0)$ be a very stable Higgs bundle. Denote by $F_{\mathcal{E}} \in \pi_0(\mathcal{M}^{\mathbb{T}})$ the component of the \mathbb{T} -fixed point set containing \mathcal{E} . Let $N := h^{-1}(0) \subset \mathcal{M}$ denote the subscheme of nilpotent Higgs bundles, the so called (global) nilpotent cone. Then the corresponding subvariety $N_{red} \cong \mathcal{C} \subset \mathcal{M}$ is isomorphic to the core of \mathcal{M} . Denote by $\mathcal{C}_{F_{\mathcal{E}}} := \overline{W_{F_{\mathcal{E}}^-}}$ the closure of the downward flow from $F_{\mathcal{E}}$, which is an irreducible component of \mathcal{C} . Denote by $N_{F_{\mathcal{E}}} \subset N$ the corresponding irreducible component of the subscheme N .

Definition 5.1. The *multiplicity* of the component N_{F_ε} in the subscheme N is defined to be the length of the local ring $\mathcal{O}_{N, N_{F_\varepsilon}}$ at a generic point and is denoted by $m_{F_\varepsilon} := \ell(\mathcal{O}_{N, N_{F_\varepsilon}})$.

Further we let $T_\varepsilon^+ \mathcal{M} < T_\varepsilon \mathcal{M}$ denote the part of the \mathbb{T} -module $T_\varepsilon \mathcal{M}$ with positive weights. Then $T_\varepsilon W_\varepsilon^+ \cong T_\varepsilon^+ \mathcal{M}$ which we will abbreviate as T_ε^+ . Finally, recall that for a finite dimensional positive \mathbb{T} -module $V = \bigoplus_{\lambda > 0} V_\lambda$ where V_λ is the weight space corresponding to $\lambda \in \text{Hom}(\mathbb{T}, \mathbb{T}) \cong \mathbb{Z}$, the character of the symmetric algebra $\text{Sym}(V^*) = \bigoplus_{\lambda \leq 0} \text{Sym}(V^*)_\lambda$ of its dual satisfies

$$\chi_{\mathbb{T}}(\text{Sym}(V^*)) = \sum_{\lambda \leq 0} \dim(\text{Sym}(V^*)_\lambda) t^{-\lambda} = \prod_{\lambda > 0} \frac{1}{(1 - t^\lambda)^{\dim(V_\lambda)}} \in \mathbb{Z}[[t]]. \quad (5.1)$$

In particular, the equivariant Euler characteristic (see §6.3.1 for more details) of the structure sheaf \mathcal{O}_V on V can be computed as

$$\chi_{\mathbb{T}}(V; \mathcal{O}_V) = \sum_{i, k} (-1)^i t^{-k} \dim H^i(\mathcal{O}_V)_k = \sum_k t^{-k} \dim H^0(\mathcal{O}_V)_k = \chi_{\mathbb{T}}(\text{Sym}(V^*)), \quad (5.2)$$

where we recall that \mathbb{T} acts on $s \in H^0(\mathcal{O}_V)$ with non-positive weights by the formula $(\lambda \cdot s)(v) = s(\lambda^{-1} \cdot v)$.

Theorem 5.2. *When $\mathcal{E} \in \mathcal{M}^{\text{st}\mathbb{T}} \subset N$ is a very stable Higgs bundle then the multiplicity of the component N_{F_ε} in the nilpotent cone N satisfies*

$$m_{F_\varepsilon} = \left. \frac{\chi_{\mathbb{T}}(\text{Sym}(T_\varepsilon^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))} \right|_{t=1}.$$

Moreover, for a generic $a \in \mathcal{A}$ the intersection $W_\varepsilon^+ \cap h^{-1}(a)$ is transversal and has cardinality m_{F_ε} .

Proof. We shall use basic properties of the K-theory of coherent sheaves with supports as developed in e.g. [BF, GS, PW]. For $Z \subset X$ a closed algebraic subset of a complex algebraic variety we will denote by $K_Z^\circ(X)$ (resp. by $K_\circ^Z(X)$) the Grothendieck group of bounded complexes of locally free sheaves (resp. coherent sheaves) with homology supported in Z .

We shall compute the intersection product

$$[\mathcal{O}_N] \cap [\mathcal{O}_{W_\varepsilon^+}] \in K_\circ(\{\mathcal{E}\}) \cong \mathbb{Z} \quad (5.3)$$

in two different ways. Here $[\mathcal{O}_N] = h^*([\mathcal{O}_0]) \in K_\circ^\circ(\mathcal{M})$, where $[\mathcal{O}_0] \in K_{\{0\}}^\circ(\mathcal{A})$ and $[\mathcal{O}_{W_\varepsilon^+}] \in K_\circ^{W_\varepsilon^+}(\mathcal{M})$, where $W_\varepsilon^+ \subset \mathcal{M}$ is closed by Lemma 4.3. Thus the cap-product

$$[\mathcal{O}_N] \cap [\mathcal{O}_{W_\varepsilon^+}] = [\mathcal{O}_N \otimes^L \mathcal{O}_{W_\varepsilon^+}] \in K_\circ^{c \cap U_\varepsilon}(\mathcal{M}) \cong K_\circ^{\{\mathcal{E}\}}(\mathcal{M}) \cong K_\circ(\{\mathcal{E}\}) \cong \mathbb{Z}$$

is defined. The last isomorphism is given by the Euler characteristic $\chi : K_\circ(\{\mathcal{E}\}) \rightarrow \mathbb{Z}$.

First we note that $W_\mathcal{E}^+$ intersects $\mathcal{C} = N_{red}$ transversally at the single point \mathcal{E} , since both $W_\mathcal{E}^+$ and \mathcal{C} are smooth at \mathcal{E} and the tangent spaces $T_\mathcal{E}^+$ and $T_\mathcal{E}^{\leq 0}$ are complementary. This also follows from Bialynicki-Birula's local description of the upward flow (see Remark 2.8). Then [Sta, Lemma 42.14.3] implies that the intersection multiplicity

$$i(\mathcal{M}, W_\mathcal{E}^+ \cdot \mathcal{C}, \{\mathcal{E}\}) = i(\mathcal{M}, W_\mathcal{E}^+ \cdot \mathcal{C}_{F_\mathcal{E}}, \{\mathcal{E}\}) = 1. \quad (5.4)$$

We can also find an affine open neighbourhood $\text{Spec}(R)$ of $\mathcal{E} \in \mathcal{M}$ so that $\mathcal{O}_{W_\mathcal{E}^+}|_{\text{Spec}(R)}$ is represented by the R -algebra A and $\mathcal{O}_{N_{F_\mathcal{E}}}$ by the R -algebra B . As $N_{F_\mathcal{E}}$ is irreducible B has a unique minimal prime ideal $I_{min} \triangleleft B$ which corresponds to the generic point of $N_{F_\mathcal{E}}$ and coincides with the nilradical of B . In particular, the reduced ring $B_{red} = B/I_{min}$ and

$$\text{Spec} B_{red} \cong \mathcal{C}_{F_\mathcal{E}} \cap \text{Spec}(R). \quad (5.5)$$

By [Bou, Theorem 2 of IV.1.4] we know that B as a B -module has a composition series

$$0 = B_k \subset B_{k-1} \subset \cdots \subset B_0 = B$$

with factors $B_j/B_{j+1} \cong B/P_j$, where $P_j \triangleleft B$ are prime ideals. Moreover we know from *loc. cit.* that the factor B/I_{min} appears $\ell(B_{I_{min}})$ times in this composition series, which further agrees with

$$\ell(B_{I_{min}}) = \ell(\mathcal{O}_{N, N_{F_\mathcal{E}}}) = m_{F_\mathcal{E}}.$$

Now $\mathcal{O}_{W_\mathcal{E}^+} \otimes^L \mathcal{O}_{N_{F_\mathcal{E}}}$ has support $\{\mathcal{E}\}$. The Euler characteristic of its stalk at $\{\mathcal{E}\}$ is given by Serre's Tor formula

$$\chi(A_{I_\mathcal{E}}, B_{I_\mathcal{E}}) = \sum (-1)^i \ell(\text{Tor}_i(A_{I_\mathcal{E}}, B_{I_\mathcal{E}})),$$

where $I_\mathcal{E} \triangleleft R$ is the maximal ideal corresponding to the point $\mathcal{E} \in \text{Spec}(R) \subset \mathcal{M}$. Computing this further

$$\sum_i (-1)^i \ell(\text{Tor}_i(A_{I_\mathcal{E}}, B_{I_\mathcal{E}})) = \sum_i \sum_j (-1)^i \ell(\text{Tor}_i(A_{I_\mathcal{E}}, (B/P_j)_{I_\mathcal{E}})) = \sum_j \chi(A_{I_\mathcal{E}}, (B/P_j)_{I_\mathcal{E}}).$$

We know that $P_j \supset I_{min}$. When $P_j \neq I_{min}$ then $\dim(B/P_j) < \dim(B/I_{min})$ and so Serre's [Se, Theorem C.1.1.a] shows that $\chi(A_{I_\mathcal{E}}, (B/P_j)_{I_\mathcal{E}}) = 0$ in this case. Thus the Euler characteristic of the stalk of $\mathcal{O}_{W_\mathcal{E}^+} \otimes^L \mathcal{O}_{N_{F_\mathcal{E}}}$ at $\{\mathcal{E}\}$ equals

$$\chi(A_{I_\mathcal{E}}, B_{I_\mathcal{E}}) = m_{F_\mathcal{E}} \chi(A_{I_\mathcal{E}}, (B/I_{min})_{I_\mathcal{E}}) = m_{F_\mathcal{E}},$$

as

$$\chi(A_{I_\mathcal{E}}, (B/I_{min})_{I_\mathcal{E}}) = \chi(A_{I_\mathcal{E}}, (B_{red})_{I_\mathcal{E}}) = i(\mathcal{M}, W_\mathcal{E}^+ \cdot \mathcal{C}_{F_\mathcal{E}}, \{\mathcal{E}\}) = 1$$

from (5.4),(5.5) and [Se, Theorem C.1.1.b]. This implies that

$$\chi([\mathcal{O}_{W_\varepsilon^+}] \cap [\mathcal{O}_N]) = m_{F_\varepsilon}. \quad (5.6)$$

On the other hand, by the projection formula in K -theory [PW, (1)]

$$h_*([\mathcal{O}_N] \cap [\mathcal{O}_{W_\varepsilon^+}]) = [\mathcal{O}_0] \cap h_*([\mathcal{O}_{W_\varepsilon^+}]) \in K_o^{\{0\}}(\mathcal{A}). \quad (5.7)$$

By Lemma 4.5 $h_*([\mathcal{O}_{W_\varepsilon^+}])$ is locally free and we compute

$$\chi([\mathcal{O}_N] \cap [\mathcal{O}_{W_\varepsilon^+}]) = \chi(h_*([\mathcal{O}_N] \cap [\mathcal{O}_{W_\varepsilon^+}])) = \chi([\mathcal{O}_0] \cap h_*([\mathcal{O}_{W_\varepsilon^+}])) = \chi([h_*([\mathcal{O}_{W_\varepsilon^+}])_0]) = \text{rank}(h_*(\mathcal{O}_{W_\varepsilon^+})).$$

Thus the multiplicity m_{F_ε} in (5.6) agrees with $\text{rank}(h_*(\mathcal{O}_{W_\varepsilon^+}))$ the rank of the locally free sheaf $h_*(\mathcal{O}_{W_\varepsilon^+})$.

To determine this rank, we notice that W_ε^+ is \mathbb{T} -invariant. We compute the equivariant Euler characteristic of $\mathcal{O}_{W_\varepsilon^+}$ as

$$\chi_{\mathbb{T}}(\mathcal{O}_{W_\varepsilon^+}) = \chi_{\mathbb{T}}(\mathcal{O}_{T_\varepsilon^+}) = \chi_{\mathbb{T}}(\text{Sym}(T_\varepsilon^{+*})) \in \mathbb{Z}((t)) \quad (5.8)$$

since the \mathbb{T} -action on W_ε^+ is modelled on the \mathbb{T} -module T_ε^+ i.e. $W_\varepsilon^+ \cong T_\varepsilon^+$ are \mathbb{T} -equivariantly isomorphic (c.f. (2.7)).

On the other hand because $h : W_\varepsilon^+ \rightarrow \mathcal{A}$ is finite the \mathbb{T} -equivariant sheaf $h_*(\mathcal{O}_{W_\varepsilon^+})$ is locally free. By the equivariant Serre conjecture proved for \mathbb{T} in [MMP] the \mathbb{T} -equivariant locally free sheaf $h_*(\mathcal{O}_{W_\varepsilon^+})$ on the affine space \mathcal{A} is trivial. In other words

$$h_*(\mathcal{O}_{W_\varepsilon^+}) = h_*(\mathcal{O}_{W_\varepsilon^+})_0 \times \mathcal{A}, \quad (5.9)$$

where the \mathbb{T} -action on the 0 fibre $h_*(\mathcal{O}_{W_\varepsilon^+})_0$ is inherited from the \mathbb{T} -equivariant structure on $h_*(\mathcal{O}_{W_\varepsilon^+})$ and the \mathbb{T} action on \mathcal{A} is the standard one. Thus we compute

$$\chi_{\mathbb{T}}(\mathcal{O}_{W_\varepsilon^+}) = \chi_{\mathbb{T}}(Rh_*(\mathcal{O}_{W_\varepsilon^+})) = \chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_\varepsilon^+})) = \chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_\varepsilon^+})_0) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)) \in \mathbb{Z}((t)) \quad (5.10)$$

Here $\chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_\varepsilon^+})_0)$ is thought of as the \mathbb{T} -equivariant Euler characteristic over the point $0 \in \mathcal{A}$, or more generally if W is a finite dimensional vector space with a \mathbb{T} -representation and $W = \bigoplus_{\lambda \in \mathbb{Z}} W_\lambda$ is the weight space decomposition then

$$\chi_{\mathbb{T}}(W) = \sum_{\lambda \in \mathbb{Z}} t^{-\dim(W_\lambda)} \in \mathbb{Z}[t, t^{-1}].$$

Comparing (5.8) and (5.10) we see that the equivariant Euler characteristic of the \mathbb{T} -module $h_*(\mathcal{O}_{W_\varepsilon^+})_0$ can be expressed as

$$\chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_\varepsilon^+})_0) = \frac{\chi_{\mathbb{T}}(\text{Sym}(T_\varepsilon^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))} \in \mathbb{Z}((t)). \quad (5.11)$$

In particular, the dimension of $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0$ is

$$\text{rank}(h_*(\mathcal{O}_{W_{\mathcal{E}}^+})) = \dim(h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0) = \frac{\chi_{\mathbb{T}}(\text{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))} \Big|_{t=1}.$$

This proves the first statement. For the second statement note that when $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable then $W_{\mathcal{E}}^+ \subset \mathcal{M}$ is closed and for $a \in \mathcal{A}$

$$h_*([\mathcal{O}_{W_{\mathcal{E}}^+}] \cap [h^{-1}(a)]) = [h_*(\mathcal{O}_{W_{\mathcal{E}}^+}) \cap \mathcal{O}_a] = [h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_a] = m_{F_{\mathcal{E}}} \in K(\{a\}) \cong \mathbb{Z}. \quad (5.12)$$

When $a \in \mathcal{A}$ generic then by Lemma 4.5 the intersection of $W_{\mathcal{E}}^+$ and $h^{-1}(a)$ is transversal and thus has cardinality $m_{F_{\mathcal{E}}}$ from (5.12). □

Definition 5.3. For $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ define the rational function

$$m_{\mathcal{E}}(t) := \frac{\chi_{\mathbb{T}}(\text{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))} \in \mathbb{Z}((t)).$$

We call it the *virtual equivariant multiplicity* or *virtual multiplicity* for short.

Corollary 5.4. When $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable $m_{\mathcal{E}}(t)$ is a palindromic and monic polynomial with non-negative integer coefficients, such that $m_{\mathcal{E}}(1) = m_{F_{\mathcal{E}}}$ is the multiplicity of the component $N_{F_{\mathcal{E}}} \subset N$ in the nilpotent cone.

Proof. When \mathcal{E} is very stable by (5.11) $m_{\mathcal{E}}(t)$ is the Laurent polynomial $\chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0) \in \mathbb{Z}[t, t^{-1}]$, the character of the finite dimensional \mathbb{T} -module $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0$. In particular, $m_{\mathcal{E}}(t)$ has non-negative coefficients.

When all the weights of \mathbb{T} on a finite dimensional \mathbb{T} -module V are positive, then $1/\chi_{\mathbb{T}}(\text{Sym}(V^*))$ is a monic polynomial i.e.

$$\left(\frac{1}{\chi_{\mathbb{T}}(\text{Sym}(V^*))} \right)_{t=0} = 1$$

from (5.1). It follows that $m_{\mathcal{E}}(t) \in \mathbb{Z}[t]$ is a polynomial and $m_{\mathcal{E}}(0) = 1$.

As

$$\prod_{\lambda > 0} (1 - t^{\lambda})^{\dim(V_{\lambda})} = (-1)^{\dim V} t^{\sum \lambda \dim(V_{\lambda})} \prod_{\lambda > 0} (1 - t^{-\lambda})^{\dim(V_{\lambda})}$$

we get

$$m_{\mathcal{E}}(t) = m_{\mathcal{E}}(t^{-1}) t^{\sum \lambda (\dim(\mathcal{A}_{\lambda}) - \dim((T_{\mathcal{E}}^+)_{\lambda}))} = m_{\mathcal{E}}(t^{-1}) t^{\deg(m_{\mathcal{E}}(t))} \quad (5.13)$$

where

$$\deg(m_{\mathcal{E}}(t)) = \sum \lambda (\dim(\mathcal{A}_{\lambda}) - \dim((T_{\mathcal{E}}^+)_{\lambda}))$$

is the degree of the polynomial $m_{\mathcal{E}}(t)$. Thus $m_{\mathcal{E}}(t)$ is palindromic. As $m_{\mathcal{E}}(0) = 1$ it is monic too. The result follows. □

Remark 5.5. In Section 7 we shall see that, in the cases considered here, $m_{\mathcal{E}}(t^2)$ is a product of Poincaré polynomials of compact homogeneous spaces and hence monic and palindromic by Poincaré duality.

Remark 5.6. For a very stable $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ we have $m_{\mathcal{E}}(t) = \chi_{\mathbb{T}}(h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0)$ and so \mathbb{T} acts on the vector space $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0$ with non-positive weights and the multiplicity of the trivial character of \mathbb{T} is 1 in $h_*(\mathcal{O}_{W_{\mathcal{E}}^+})_0$.

Definition 5.7. When $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable we call $m_{\mathcal{E}}(t)$ the *equivariant multiplicity* of $N_{F_{\mathcal{E}}} \subset N$.

Remark 5.8. Note that by the \mathbb{T} -equivariant analogue of the projection formula (5.7) we see that the equivariant multiplicity at a very stable Higgs bundle satisfies

$$m_{\mathcal{E}}(t) = [\mathcal{O}_N]^{\mathbb{T}} \cap [\mathcal{O}_{W_{\mathcal{E}}^+}]^{\mathbb{T}} \in K_{\circ}^{\mathbb{T}}(\{\mathcal{E}\}) \cong \mathbb{Z}[t, t^{-1}].$$

Hence the name.

Remark 5.9. We know from [La1, Proposition 3.5] that $E \in \mathcal{N}_n$ very stable rank n vector bundles exist. Then $\mathcal{E} = (E, 0)$ is a very stable Higgs bundle, and $T_{\mathcal{E}}^+ \cong T_E^* \mathcal{N}$ and \mathbb{T} acts with weight one. Thus

$$\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*})) = \frac{1}{(1-t)^{n^2(g-1)+1}} \quad (5.14)$$

while

$$\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*)) = \frac{1}{(1-t)^g (1-t^2)^{3g-3} (1-t^3)^{5g-5} \dots (1-t^n)^{(2n-1)(g-1)}}.$$

Thus the multiplicity in this case is

$$m_{\mathcal{E}}(t) = \frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))} = [2]_t^{3g-3} [3]_t^{5g-5} \dots [n]_t^{(2n-1)(g-1)}, \quad (5.15)$$

where

$$[n]_t := \frac{1-t^n}{1-t} = 1 + t + \dots + t^{n-1}$$

are the quantum integers. So by Theorem 5.2 the multiplicity of the component of the nilpotent cone N whose reduced subvariety is isomorphic with \mathcal{N} , the moduli space of rank n stable bundles, is

$$m_{\mathcal{N}} = 2^{3g-3} 3^{5g-5} \dots n^{(2n-1)(g-1)}.$$

This number was also computed in [BNR] as the degree of the dominant map $h^{-1}(a) \rightarrow \mathcal{N}$. We will see in Subsection 8.3 below that this is not a coincidence.

Finally, we note that for any other $\mathcal{F} \in \mathcal{M}^{s\mathbb{T}}$

$$\frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{F}}^{+*}))} \in \mathbb{Z}[t]$$

is a polynomial from the definition (5.1), thus we can deduce the following.

Corollary 5.10. *For $\mathcal{F} \in \mathcal{M}^{s\mathbb{T}}$ very stable, the equivariant multiplicity divides (5.15):*

$$m_{\mathcal{F}}(t) \mid [2]_t^{3g-3} [3]_t^{5g-5} \dots [n]_t^{(2n-1)(g-1)}.$$

In particular, it is a product of cyclotomic polynomials.

Remark 5.11. Let $n = 2$. By Corollary 4.16 and Theorem 4.2 we know that there exists a very stable Higgs bundle \mathcal{E} in every component \mathcal{N} , $F_i \in \pi_0(\mathcal{M}_2)$ of the fixed point set of the \mathbb{T} -action on \mathcal{M} (see Remark 3.3). We see that $T_{\mathcal{E}}^+ = V_2 \oplus V_1$ has only weights 1 and 2 and $V_1 \cong T^*F_i$, thus

$$\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*})) = \frac{1}{(1-t)^{\dim F_i} (1-t^2)^{\dim \mathcal{M}/2 - \dim F_i}} = \frac{1}{(1-t)^{i+g} (1-t^2)^{3g-i-3}}.$$

On the other hand

$$\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*)) = \frac{1}{(1-t)^g (1-t^2)^{3g-3}}.$$

Consequently

$$m_{\mathcal{E}}(t) = \frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))} = (1+t)^i = [2]_t^i.$$

Thus from Theorem 5.2 we get that

$$m_{F_i} = 2^i.$$

This result was proved in [Hi5, Proposition 6] for SL_2 Higgs bundles (i even) and for twisted SL_2 Higgs bundles (i odd) in [HT].

Remark 5.12. One can compute the rational function $\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*})) / \chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))$ for every $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$, which by Corollary 6.29 equals the polynomial $m_{\mathcal{E}}(t)$, when \mathcal{E} is very stable. This quantity depends only on the ambient $F_{\mathcal{E}} \in \pi_0(\mathcal{M}^{\mathbb{T}})$ and when it is not a polynomial, we deduce that there exists no very stable Higgs bundle in $F_{\mathcal{E}}$.

As an example recall [Go] that a type $(1, 2)$ fixed point \mathcal{E} of the \mathbb{T} -action has underlying rank 3 vector bundle of the form $E = L \oplus V$, where L is a rank 1 and V is a rank 2 vector bundle on C and the Higgs field only non-trivial in $H^0(C; L^*VK)$. If $\ell = \deg(L)$ and $v = \deg(V)$ then $d = \deg(E) = \ell + v$. It is shown in [Go, Proposition 2.5] that there is a stable Higgs bundle of this form if $1/3d < \ell < 1/3d + g - 1$ or equivalently $0 < 2\ell - v < 3g - 3$.

We can easily compute the weight spaces of the \mathbb{T} -module $T_{\mathcal{E}}^+$. It has two weights 1 and 2, and we can compute the 2-weight space isomorphic to $H^0(C; V^*LK)$ whose dimension is $2\ell - v + 2g - 2$ from Hirzebruch-Riemann-Roch and the fact that $H^1(C; V^*LK) \cong H^0(C; VL^*)^* \cong 0$ from [Go, proof of Proposition 4.2]. Because of the homogeneity 1 symplectic form, the weight 2 space is isomorphic with the dual of the weight -1 space,

thus $2\ell - v + 2g - 2$ also agrees with half of the Morse index of the ambient fixed point component, matching [Go, Proposition 4.2.i]. It follows that

$$\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*})) = \frac{1}{(1-t)^{9g-8-(2g-2+2\ell-v)}(1-t^2)^{2\ell-v+2g-2}}.$$

This implies

$$\frac{\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))} = (1+t)^{g-1-2\ell+v}(1+t+t^2)^{5g-5}.$$

When $2\ell - v > g - 1$ this expression is not a polynomial in t . This shows that the type $(1, 2)$ components of $\mathcal{M}_3^{\mathbb{T}}$ where $g - 1 < 2\ell - v < 3g - 3$ do not contain very stable Higgs bundles.

In fact, we can see this fact directly as follows. Our fixed point is $L \oplus V$ with Higgs field $\phi : L \rightarrow VK$. Assume for simplicity that the rank 3 bundle has degree 0. The upward flow consists of extensions $L \rightarrow E \rightarrow V$ where Φ acting on L and projecting to V is ϕ .

Let $U \subset V$ be the line bundle generated by ϕ , or the image of $\phi : LK^* \rightarrow V$. Then

$$\deg U \geq \deg L - \deg K = \ell - (2g - 2).$$

Its annihilator is $(V/U)^* \subset V^*$ which has degree $\deg U - \deg V = \deg U + \ell$ so the degree of $L(V/U)^*K$ is

$$\deg U + 2\ell + (2g - 2) \geq 3\ell.$$

If $\ell > (g-1)/3$ then $L(V/U)^*K$ always has a non-trivial section ψ which we can think of as lying in $\mathrm{Hom}(V, LK)$. Then $(\phi, \psi) : L \oplus V \rightarrow (V \oplus L) \otimes K$ is a Higgs field on $L \oplus V$ and $\langle \phi, \psi \rangle = 0$ means it is nilpotent.

Remark 5.13. For rank 2 we know that every component of $\mathcal{M}^{\mathbb{T}}$ contains a very stable Higgs bundle, thus every component of the nilpotent cone is very stable (see Remark 5.11). When a bundle is not very stable, the current term for it is *wobbly*. So For rank 3 above we found wobbly components of the nilpotent cone by computing $m_F(t)$ and finding that it is not a polynomial. In fact, for rank 3 we conjecture that all other components of the nilpotent cone are very stable.

However, adapting the argument for the type $(1, 2)$ wobbly fixed point components above to the rank 4 case we find that all stable type $(1, 3)$ \mathbb{T} -fixed Higgs bundle are wobbly. However the corresponding $m_{\mathcal{E}}(t)$ is a polynomial, using the description in [GHS, Example 6.6]. Thus the obstruction of integrality of $m_{\mathcal{E}}(t)$ for very stable Higgs bundles discussed in Remark 5.12 is not sufficient. In particular, already for rank 4 we do not have a conjectured list of very stable components of the nilpotent cone.

Remark 5.14. We know from Theorem 4.13 that all type $(1, \dots, 1)$ components of the nilpotent cone are very stable and we can explicitly compute $\chi_{\mathbb{T}}(\mathrm{Sym}(T_{\mathcal{E}}^{+*}))/\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*))$

for a fixed point $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ of type $(1, \dots, 1)$. In this case $E = L_0 \oplus \dots \oplus L_{n-1}$ and $\Phi(L_{i-1}) \subset L_i K$ is determined by $b_i := \Phi|_{L_{i-1}} : L_{i-1} \rightarrow L_i K$ non-zero. We denote $\ell_i = \deg(L_i)$ and $m_i = 2g - 2 - \ell_i + \ell_{i+1}$. As b_i is non-zero $m_i \geq 0$.

By (3.2) the tangent space $T_{\mathcal{E}}$ to \mathcal{M}^s at this point is the hypercohomology group

$$\begin{aligned} T_{\mathcal{E}} &\cong \mathbb{H}^1 \left(\text{End}(E) \xrightarrow{ad(\Phi)} \text{End}(E) \otimes K \right) \cong \\ &\cong \bigoplus_{k=-n+1}^{k=n} \mathbb{H}^1 \left(\bigoplus_{i-j=k} \text{Hom}(L_i, L_j) \xrightarrow{ad(\Phi)} \bigoplus_{i-j=k-1} \text{Hom}(L_i, L_j) \otimes K \right) \end{aligned}$$

indexed according to the weights of the \mathbb{T} action on $T_{\mathcal{E}}$. Thus we see that

$$T_{\mathcal{E}}^+ = \bigoplus_{k=1}^{k=n} (T_{\mathcal{E}})_k = \bigoplus_{k=1}^{k=n} \mathbb{H}^1 \left(\bigoplus_{i-j=k} \text{Hom}(L_i, L_j) \xrightarrow{ad(\Phi)} \bigoplus_{i-j=k-1} \text{Hom}(L_i, L_j) \otimes K \right).$$

Because of the stability of the Higgs bundle \mathcal{E} [Ha2, Theorem 4.3] we get that the zeroth and second hypercohomology of all the complexes vanish, thus

$$\begin{aligned} \dim T_{\mathcal{E}}^k &= - \sum_{i-j=k} \chi(\text{Hom}(L_i, L_j)) + \sum_{i-j=k-1} \chi(\text{Hom}(L_i, L_j) \otimes K) \\ &= - \sum_{i-j=k} (-\ell_i + \ell_j + 1 - g) + \sum_{i-j=k-1} (-\ell_i + \ell_j + g - 1) \\ &= (2n - 2k + 1)(g - 1) - \ell_k + \ell_{n-k+1} \\ &= (2k - 1)(g - 1) + m_{k, n-k+1}, \end{aligned}$$

where

$$m_{a,b} = \begin{cases} \sum_{a \leq j < b} m_j & \text{when } a < b \\ 0 & \text{when } a = b \\ - \sum_{b \leq j < a} m_j & \text{when } a > b \end{cases}$$

Then a straightforward computation gives

$$m_{\mathcal{E}}(t) = \frac{\chi_{\mathbb{T}}(\text{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))} = \frac{\prod_{j=\lfloor \frac{n}{2} \rfloor}^{n-1} (1 - t^{j+1})^{m_{n-j, j+1}}}{\prod_{j=1}^{\lfloor \frac{n}{2} \rfloor} (1 - t^j)^{m_{j, n-j+1}}} = \prod_{i=1}^{n-1} \left[\begin{matrix} n \\ i \end{matrix} \right]_t^{m_i}. \quad (5.16)$$

Here

$$\left[\begin{matrix} n \\ k \end{matrix} \right]_t = \prod_{j=1}^k \frac{1 - t^{n-j+1}}{1 - t^j}$$

is the quantum binomial coefficient, in particular a polynomial. Incidentally, this is the Poincaré polynomial of the Grassmanian $\text{Gr}_k(\mathbb{C}^n)$, which coincidence will be further discussed in Subsection 8.1.

We thus see that the rational function in (5.16) is always a polynomial. The polynomiality also follows from the existence of very stable Higgs bundles in each type $(1, 1, \dots, 1)$ component of $\mathcal{M}^{\mathbb{T}}$, as proved in Corollary 4.16.

We obtain immediately

Corollary 5.15. *For a fixed point component $F_{\mathcal{E}} \in \pi_0(\mathcal{M}^{\mathbb{T}})$ of type $(1, 1, \dots, 1)$ as in Remark 5.14 the multiplicity of $N_{F_{\mathcal{E}}}$ in the nilpotent cone N is*

$$m_{F_{\mathcal{E}}} = m_{\mathcal{E}}(1) = \prod_{i=1}^{n-1} \binom{n}{i}^{m_i}.$$

Another consequence is the following

Corollary 5.16. *For $\mathcal{E} \in \mathcal{M}_n^{s\mathbb{T}}$ very stable the following are equivalent*

1. $m_{F_{\mathcal{E}}} = 1$
2. $m_{F_{\mathcal{E}}}(t) = 1$
3. \mathcal{E} is a uniformising Higgs bundle of Remark 3.8
4. $W_{\mathcal{E}}^+$ is a section of the Hitchin map h

Proof. Corollary 6.29 implies that $m_{\mathcal{E}}(t) \in \mathbb{Z}_{\geq 0}[t]$ is palindromic with non-negative coefficients showing 1. \Rightarrow 2.. By Definition 5.3 $m_{\mathcal{E}}(t) = 1$ implies that there are n distinct weights in the \mathbb{T} -module $T_{\mathcal{E}}^+$, to match those in \mathcal{A} . This implies in turn that \mathcal{E} is of type $(1, \dots, 1)$. For a type $(1, \dots, 1)$ \mathcal{E} we will have $m_{\mathcal{E}}(t) = 1$ if all $m_i = 0$ from (5.16). This only happens for a uniformising Higgs bundle. This shows 2. \Rightarrow 3.. From Remark 3.8 for a uniformising Higgs bundle $W_{\mathcal{E}}^+$ is a section of the Hitchin map. This shows 3. \Rightarrow 4.. Finally the second part of Theorem 5.2 implies 4. \Rightarrow 1.. \square

5.2 Intersection of upward flows with generic fibres of h

Let $a = (a_1, \dots, a_n) \in H^0(C; K) \oplus \dots \oplus H^0(C; K^n) = \mathcal{A}$ be coefficients that define a smooth spectral curve $C_a \subset T^*C$. Assume that $\text{div}(b)$ for \mathcal{E} of type $(1, \dots, 1)$ avoids the ramification divisor of $\pi_a : C_a \rightarrow C$. In other words over any zero $c \in C$ of b we have $|\pi_a^{-1}(c)|$ has n distinct preimages of c in C_a .

We recall that the spectral curve is a subscheme of T^*C defined as

$$C_a = \det(\pi^*\Phi - x)^{-1}(0),$$

where $\pi : T^*C \rightarrow C$ is the projection and $x \in H^0(T^*C, \pi_a^*(K))$ is the tautological section. For $(E, \Phi) \in h^{-1}(a)$ denote by

$$U := \text{coker}(\pi_a^*(E \otimes K^{-1}) \xrightarrow{\pi_a^*\Phi - x} \pi_a^*(E)). \quad (5.17)$$

Then by [Hi2, §5.1] and [BNR, §3.7] we have the following proposition, which we prove for completeness as the proof is not readily available in the literature.

Proposition 5.17. 1. *The direct image of the trivial bundle*

$$(\pi_a)_*(\mathcal{O}_{C_a} \xrightarrow{x} \pi_a^*(K)) \cong \mathcal{E}_{\mathcal{O}_C, a} = (E_{\mathcal{O}_C}, \Phi_a) \in W_{\mathcal{E}_0}^+ \quad (5.18)$$

defines the canonical section $W_{\mathcal{E}_0}^+$ of the Hitchin map (see (3.16)).

2. *If a Higgs bundle (E, Φ) is the direct image of a line bundle U :*

$$(\pi_a)_*(U \xrightarrow{x} U \otimes \pi_a^*(K)) = (E, \Phi). \quad (5.19)$$

then on C_a we have the exact sequence

$$0 \rightarrow U \otimes \pi_a^*(K^{-n}) \rightarrow \pi_a^*(E \otimes K^{-1}) \xrightarrow{\pi_a^*\Phi - x} \pi_a^*(E) \rightarrow U \rightarrow 0. \quad (5.20)$$

Proof. To prove (5.18) we note that the pushforward $(\pi_a)_*(\mathcal{O}_{C_a})$ is the \mathcal{O}_C -algebra $\text{Sym}(K^{-1})/\mathcal{I}$ where the ideal sheaf \mathcal{I} is generated by the image of $u : K^{-n} \rightarrow \text{Sym}(K^{-1})$ the sum of $a_i : K^{-n} \rightarrow K^{-n+i}$. Thus we see that as a sheaf $(\pi_a)_*(\mathcal{O}_C) \cong \mathcal{E}_{\mathcal{O}_C}$ and the algebra structure on it induces the Higgs field Φ_a (3.15) the companion matrix of a .

For the proof of the second statement we also denote by

$$U' := \ker(\pi_a^*(E \otimes K^{-1}) \xrightarrow{\pi_a^*\Phi - x} \pi_a^*(E))$$

the kernel. Then the pushforward of the middle square of the diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & U' & \rightarrow & \pi_a^*(E \otimes K^{-1}) & \xrightarrow{\pi_a^*\Phi - x} & \pi_a^*(E) & \rightarrow & U & \rightarrow & 0 \\ & & x \downarrow & & x \downarrow & & x \downarrow & & x \downarrow & & \\ 0 & \rightarrow & U' \otimes \pi_a^*(K) & \rightarrow & \pi_a^*(E) & \xrightarrow{\pi_a^*\Phi - x} & \pi_a^*(E \otimes K) & \rightarrow & U \otimes \pi_a^*(K) & \rightarrow & 0 \end{array}$$

is the middle square of the following diagram

$$\begin{array}{ccccccccc} 0 & \rightarrow & EK^{-n} & \xrightarrow{\Psi} & E_{\mathcal{O}_C} \otimes EK^{-1} & \xrightarrow{Id_{E_{\mathcal{O}_C}} \otimes \Phi - \Phi_a \otimes Id_{EK^{-1}}} & E_{\mathcal{O}_C} \otimes E & \xrightarrow{\Xi} & E & \rightarrow & 0 \\ & & \Phi \downarrow & & \Phi_a \otimes Id_E \downarrow & & \Phi_a \otimes Id_E \downarrow & & \Phi \downarrow & & \\ 0 & \rightarrow & EK^{-n+1} & \xrightarrow{\Psi} & E_{\mathcal{O}_C} \otimes E & \xrightarrow{Id_{E_{\mathcal{O}_C}} \otimes \Phi - \Phi_a \otimes Id_{EK^{-1}}} & E_{\mathcal{O}_C} \otimes EK & \xrightarrow{\Xi} & EK & \rightarrow & 0 \end{array} .$$

The rest of the diagram is given by the maps

$$\Psi = \begin{pmatrix} \Psi_1 \\ \Psi_2 \\ \vdots \\ \Psi_n \end{pmatrix} : EK^{-n} \rightarrow EK^{-1} \oplus EK^{-2} \oplus \dots \oplus EK^{-n}$$

with

$$\Psi_i = \Phi^{n-i} + a_1 \Phi^{n-i-1} + \cdots + a_{n-i} \text{Id}_{EK^{-n}} : EK^{-n} \rightarrow EK^{-i}$$

and $\Xi = (\Xi_1, \Xi_2, \dots, \Xi_n) : E \oplus EK^{-1} \oplus \cdots \oplus EK^{-n+1} \rightarrow E$ is the map given by $\Xi_i = \Phi^{i-1} : EK^{i-1} \rightarrow E$. It is straightforward to check that the second diagram commutes due to the Cayley-Hamilton identity $0 = \Phi^n + a_1 \Phi^{n-1} + \cdots + a_n \text{Id}_E$. Therefore the second diagram is the pushforward of the first. This implies (5.19) and that $U' \cong U \otimes \pi_a^*(K^{-n})$. The result follows. \square

Now let \mathcal{E} be, as in Remark 5.14, a very stable Higgs bundle of type $(1, 1, \dots, 1)$. We know from Proposition 3.4 that $(E, \Phi) \in W_{\mathcal{E}}^+$ if and only if there exists a filtration with a full flag of subbundles as in (4.3) which is compatible with the Higgs field (4.4). Recall the maps b_i and b from (4.5) and (4.6) respectively.

Assume that $(E, \Phi) \in W_{\mathcal{E}}^+ \cap h^{-1}(a)$. Denote by $f_i : \pi_a^*(E_i) \rightarrow U$ the composition of the embedding $\pi_a^*(E_i) \subset \pi_a^*(E)$ and projection $\pi_a^*(E) \rightarrow U$ arising from the definition (5.17) of U . Furthermore, denote by $U_i = \text{im}(f_i)$ the image sheaf. As a non-trivial subsheaf of the invertible sheaf U it is itself an invertible sheaf. We have thus constructed a filtration

$$U_1 \subset U_2 \subset \cdots \subset U_{n-1} \subset U_n = U$$

by invertible subsheaves. Denote the induced map $\tilde{f}_i : U_i \rightarrow U$ and by $\Delta_i := \text{div}(\tilde{f}_i)$ the divisor of zeroes of $\tilde{f}_i \in H^0(C; U_i^*U)$. Note that the divisor Δ_1 determines

$$U = \pi_a^*(E_1)(\Delta_1),$$

and so in turn by (5.19) it also determines our Higgs bundle $(E, \Phi) \in W_{\mathcal{E}}^+ \cap h^{-1}(a)$. We get the chain of divisors on C

$$\Delta_1 \geq \Delta_2 \geq \cdots \geq \Delta_{n-1} \geq \Delta_n = 0.$$

Proposition 5.18. *Let $\mathcal{E} \in \mathcal{M}^{\text{st}}$ be a type $(1, \dots, 1)$ very stable Higgs bundle. Let $a \in \mathcal{A}$ be such that the spectral curve $C_a \subset K$ is smooth and $\text{div}(b)$ avoids the ramification divisor of π_a .*

1. *If $(E, \Phi) \in W_{\mathcal{E}}^+ \cap h^{-1}(a)$ then the effective divisor $\Delta_i - \Delta_{i+1} \geq 0$ is reduced and supported on $(n-i)$ preimages in C_a of π_a over each zero of b_i in C . More precisely if $c \in C$ is a zero of b_i then Φ_c leaves $(E_i)_c$ invariant and $\tilde{c} \in \pi_a^{-1}(c)$ is in the support of $\Delta_i - \Delta_{i+1}$ if and only if $\tilde{c} \in K_c$ is not an eigenvalue for $\Phi_c|_{(E_i)_c} : (E_i)_c \rightarrow (E_i)_c K_c$.*
2. *On the other hand for any choice of an $(n-i)$ element subset of $\pi_a^{-1}(c)$ for all zeroes c of all the b_i , there is a unique $(E, \Phi) \in W_{\mathcal{E}}^+ \cap h^{-1}(a)$ such that the corresponding invertible sheaf on the spectral curve C_a has the form $U = \pi_a^*(E_1)(\Delta_1)$ for Δ_1 effective and reduced and supported at our chosen points in C_a .*

3. Finally, any two distinct choices of such $(n - i)$ element subsets of $\pi_a^{-1}(c)$ over all zeroes of b_i for all i gives non-isomorphic Higgs bundles. In particular,

$$|W_{\mathcal{E}}^+ \cap h^{-1}(a)| = \prod_{i=1}^{n-1} \binom{n}{i}^{m_i},$$

where $\deg(b_i) = m_i$.

Proof. First we prove that $\Delta_i - \Delta_{i+1} \geq 0$ is supported at precisely $(n - i)$ preimages in C_a over each zero of b_i in C with no multiplicity, i.e. $\Delta_i - \Delta_{i+1}$ is reduced.

Note that $\Phi(E_i) \subset E_{i+1}K$ thus we have the map

$$p_i := \pi_a^*(\Phi) - x : \pi_a^*(E_i K^*) \rightarrow \pi_a^*(E_{i+1}).$$

Let $\overline{\text{im}(p_i)} \leq \pi_a^*(E_{i+1})$ denote the subbundle which is the saturation of the subsheaf $\text{im}(p_i) \subset \pi_a^*(E_{i+1})$. Let $V_{i+1} := \pi_a^*(E_{i+1})/\overline{\text{im}(p_i)}$ be the quotient bundle and

$$g_{i+1} : \pi_a^*(E_{i+1}) \rightarrow V_{i+1}$$

the quotient map. Another way to think about V_{i+1} is that it is the quotient of the sheaf $\text{coker}(p_i)$ by its torsion:

$$V_{i+1} = \text{coker}(p_i)/T(\text{coker}(p_i)).$$

As p_i is generically an embedding V_{i+1} is a line bundle. We have a map

$$h_i : V_i \rightarrow V_{i+1}.$$

This can be seen by noting that the embedding $\pi_a^*(E_i) \subset \pi_a^*(E_{i+1})$ induces a map $\text{coker}(p_i) \rightarrow \text{coker}(p_{i+1})$ which induces a map on the quotients by torsion. By chasing around definitions we see that

$$\begin{array}{ccc} \pi_a^*(E_{i+1}) & \xrightarrow{g_{i+1}} & V_{i+1} \\ \uparrow & & \uparrow h_i \\ \pi_a^*(E_i) & \xrightarrow{g_i} & V_i \end{array} \quad (5.21)$$

is commutative. We define $V_1 := \pi_a^*(E_1)$ and $h_1 : V_1 \rightarrow V_2$ as $g_2|_{\pi_a^*(E_1)}$ so that the diagram (5.21) is still commutative for $i = 1$. Finally, we have a map $h_n : V_n \rightarrow U$ induced by the map

$$\text{coker}(p_{n-1}) \rightarrow \text{coker}(\pi_a^*(\Phi) - x) = U.$$

This induces the commutative diagram

$$\begin{array}{ccc} & & V_n \\ & \nearrow g_n & \searrow h_n \\ \pi_a^*(E) & \xrightarrow{f_n} & U \end{array} \quad (5.22)$$

If we denote by

$$h_{\geq i} := h_n \circ \cdots \circ h_i : V_i \rightarrow U,$$

then we see from the commutativity of the diagrams (5.21) and (5.22) that $f_i = h_{\geq i} \circ g_i$ and because g_i is surjective we can deduce $\text{im}(h_{\geq i}) = \text{im}(f_i) = U_i$. In particular, $h_{\geq i} : V_i \rightarrow U_i$ is surjective and thus an isomorphism. Thus $\Delta_i - \Delta_{i+1} = \text{div}(h_i)$.

The map $g_{i+1}|_{\pi_a^*(E_i)} : \pi_a^*(E_i) \rightarrow V_{i+1}$ vanishes at $\tilde{c} = (c, \lambda) \in C_a$ provided that

$$\pi_a^*(E_i)_{\tilde{c}} \subset p_i(\pi_a^*(E_i K^*)_{\tilde{c}}).$$

This happens when

$$(E_i)_c \subset (\Phi - \lambda \text{Id}_E)((E_i K^*)_c).$$

As $(E_i)_c$ and $(E_i K^*)_c$ are of the same dimension this is equivalent to

$$(E_i)_c = (\Phi - \lambda \text{Id}_E)((E_i K^*)_c). \quad (5.23)$$

This implies $\Phi((E_i)_c) \subset (E_i K^*)_c$, i.e. that $(E_i)_c$ is Φ -invariant, which implies that $(b_i)_c = 0$. As by assumption c avoids the ramification divisor of π_a we see that Φ_c has distinct eigenvalues, i.e. it is regular semisimple. Finally, when λ is not an eigenvalue of $\Phi|_{(E_i)_c}$ then $(\Phi - \lambda \text{Id}_E)$ is injective so gives an isomorphism between $(E_i K^*)_c$ and $(E_i)_c$ and thus (5.23) holds.

Thus we see that $g_{i+1}|_{\pi_a^*(E_i)}$ vanishes at $n - i$ preimages of each zero of b_i , the ones corresponding to the eigenvectors of Φ_c not in $(E_i)_c$. From the commutativity of (5.21) and surjectivity of g_i we see that the same follows for h_i .

We compute

$$\deg(\Delta_1) = \sum_{i=1}^{n-1} \deg(\Delta_i - \Delta_{i+1}) = \sum_{i=1}^{n-1} \deg(h_i) \geq \sum_{i=1}^{n-1} m_i(n - i) = \deg(\Delta_1),$$

as by assumption $\text{div}(b_i)$ is reduced, $\deg(\text{div}(b_i)) = m_i$, so $\deg(h_i) \geq m_i(n - i)$ and we have

$$\begin{aligned} \sum_{i=1}^{n-1} m_i(n - i) &= \sum_{i=1}^{n-1} (-\ell_i + \ell_{i+1} + 2g - 2)(n - i) = -n\ell_1 + \sum_{i=1}^{n-1} \ell_i + n(n - 1)(g - 1) \\ &= -n \deg(E_1) + \deg(E) + \deg(K^{\binom{n}{2}}) \\ &= -\deg \pi_a^*(E_1) + \deg(U) = \deg(\Delta_1). \end{aligned}$$

Here we used $\deg(E) = \deg(U) - \deg(K^{\binom{n}{2}})$ for which see [BNR, §4]. It follows that $\Delta_i - \Delta_{i-1} \geq 0$ is reduced. Over a zero of b_i the morphism $h_i : V_i \rightarrow V_{i+1}$ is zero at precisely those points which correspond to eigenvectors of Φ_c not in $(E_i)_c$. This proves the first part of the statement.

For the second part of Proposition 5.18, we will construct a Higgs bundle such that $\Delta_1 \subset C_a$ is the union of the prescribed choices of an $(n - i)$ tuple in $\pi_a^{-1}(c)$ over every zero c of every b_i . The construction is by induction on $\deg(b_i)$. When $\deg(b_i) = 0$ then $U = \pi_a^*(E_1)$ is unique and given by the Higgs bundle $\mathcal{E}_{E_1, a}$ defined in Remark 3.8.

Recall the notation \mathcal{E}_δ for a type $(1, \dots, 1)$ Higgs bundle from Example 3.2. Suppose that we have constructed a Higgs bundle in $(E, \Phi) \in W_{\mathcal{E}_\delta}^+ \cap h^{-1}(a)$ for any choice of an $(n - i)$ tuple in $\pi_a^{-1}(c)$ over every zero c of every b_i . Let $c \in C$ be a point which avoids the ramification divisor of π_a and $b(c) \neq 0$. Let $b'_j := b_j$ i.e. $\delta'_j = b_j$ except when $j = i$, when we let $b'_i := b_i s_c$ or equivalently $\delta'_i := \delta + c$. Assume that $\mathcal{E}_{\delta'}$ is still stable. Then Φ_c is regular semisimple and we can take an i -dimensional Φ_c -invariant subspace $V_i \subset E_c$ corresponding to any i element subset of $\pi_a^{-1}(c)$. Then by Proposition 4.12 the Hecke transform $\mathcal{H}_{V_i}(E, \Phi) = (E', \Phi')$ will be contained in $W_{\mathcal{E}_{\delta'}}^+ \cap h^{-1}(a)$, and $b_i(c) = 0$ so that Φ'_c leaves $(E_i)_c$ invariant. By induction the second part of the proposition follows.

For the last part we note that there are $\binom{n}{n-i}^{m_i} = \binom{n}{i}^{m_i}$ choices for $\Delta_i - \Delta_{i-1}$. This gives us $\prod_{i=1}^{n-1} \binom{n}{i}^{m_i}$ possibilities for Δ_1 . In theory, two distinct divisors could represent the same line bundle, but in our case, this cannot happen. Namely, a Higgs bundle $(E, \Phi) \in W_{\mathcal{E}_\delta}^+ \cap h^{-1}(a)$ will have a unique filtration yielding the limiting Higgs bundle \mathcal{E}_δ from Proposition 3.4. Then in turn the filtration defines a unique subset $\Delta_1 \subset C_a$ as above in the proof of part one of the proposition. The result follows. \square

Remark 5.19. When $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is very stable then $W_{\mathcal{E}}^+ \subset \mathcal{M}$ is closed and we have (5.12). When \mathcal{E} is of type $(1, \dots, 1)$ from Corollary 5.15 and Proposition 5.18 we get that $|W_{\mathcal{E}}^+ \cap h^{-1}(a)| = m_{\mathcal{E}}$, provided that C_a is smooth and $\text{div}(b)$ avoids the ramification divisor of π_a . Consequently, in this case $W_{\mathcal{E}}^+$ and $h^{-1}(a)$ intersect transversally.

Similarly as above in Proposition 5.18 one can prove that $|W_{\mathcal{E}}^+ \cap h^{-1}(a)| < m_{\mathcal{E}}$ when b has repeated 0's. The second part of Theorem 5.2 implies that such type $(1, \dots, 1)$ Higgs bundles must be wobbly. This gives an alternative proof of the "only if" part of Theorem 4.13.

6 Mirror of upward flows

6.1 Jacobians, duality, Fourier-Mukai transform

The Jacobian $J := J_0(C)$ of a smooth complex projective curve C is the moduli space of degree 0 line bundles on C . It is an Abelian variety with multiplication of line bundles. Its dual Abelian variety is defined as $\text{Pic}^0(J)$ the moduli space of translation invariant line bundles on J . One can identify $\text{Pic}^0(J)$ with J in two different ways.

First, we fix a point $c_0 \in C$ and define the *Abel-Jacobi map* as $\alpha_{c_0} : C \rightarrow J$ by

$$\alpha_{c_0}(c) := \mathcal{O}(c - c_0).$$

Then it turns out [BL, Lemma 11.3.1] that the restriction map

$$\alpha_{c_0}^* : \text{Pic}^0(J) \rightarrow J \tag{6.1}$$

is an isomorphism.

Another isomorphism between $\text{Pic}^0(J)$ and J is given by \mathcal{L} the line bundle of the Theta divisor Θ . One defines

$$\phi_L : J \rightarrow \text{Pic}^0(J) \tag{6.2}$$

by $L \mapsto t_L^*(\mathcal{L}) \otimes \mathcal{L}^*$. The map (6.2) is an isomorphism as \mathcal{L} defines a principal polarization on J . It follows from [BL, Proposition 11.3.5] that $-\phi_L = (\alpha_{c_0}^*)^{-1}$. Thus $-\phi_L$ and $\alpha_{c_0}^*$ defines the same isomorphism between J and $\text{Pic}^0(J)$. We will identify J with $\text{Pic}^0(J)$ using this isomorphism below.

We have the *Poincaré bundle* \mathbb{P} on $J \times \text{Pic}^0(J)$ unique with the properties that

$$\mathbb{P}|_{J \times \{L\}} \cong L \tag{6.3}$$

for every $L \in \text{Pic}^0(J)$ and

$$\mathbb{P}|_{\{\mathcal{O}_C\} \times \text{Pic}^0(J)} \cong \mathcal{O}_{\text{Pic}^0(J)}.$$

For $L \in J$ we will denote the line bundle

$$\mathbb{P}_L := \mathbb{P}|_{\{L\} \times \text{Pic}^0(J)}$$

on $\text{Pic}^0(J) \cong J$, where we identified $\text{Pic}^0(J)$ and J with $\alpha_{c_0}^*$. This translation invariant line bundle on $\text{Pic}^0(J)$ is characterised by the property that

$$\alpha_{c_0}^*(\mathbb{P}_L) \cong L. \tag{6.4}$$

In particular, for $L, L_2 \in J$ we have

$$\mathbb{P}_{LL_2} \cong \mathbb{P}_L \otimes \mathbb{P}_{L_2}, \tag{6.5}$$

and

$$\mathbb{P}_{\mathcal{O}_C} = \mathcal{O}_J.$$

Using the Poincaré bundle \mathbb{P} on $J \times \text{Pic}^0(J) \cong J \times J$ we can define the Fourier-Mukai transform [Mu]. For a bounded complex $\mathcal{F}^\bullet \in D_{coh}^b(J)$ of coherent sheaves on J we denote by

$$S(\mathcal{F}^\bullet) := (\pi_2)_*(\pi_1^*(\mathcal{F}^\bullet) \otimes^L \mathbb{P}) \in D_{coh}^b(J)$$

the Fourier-Mukai transform of \mathcal{F}^\bullet . For example we have

$$S(\mathcal{O}_L) = \mathbb{P}_L.$$

We recall three properties of the Fourier-Mukai transform. First by [Mu, Theorem 3.13.1] for any $\mathcal{F}^\bullet \in D_{coh}^b(J)$

$$S(S(\mathcal{F}^\bullet)) = (-1)^*(\mathcal{F}^\bullet)[-g], \quad (6.6)$$

where $-1 : J \rightarrow J$ denotes the inverse map on the abelian variety J and $g = \dim(J)$ is the genus of C . The second [Mu, Theorem 3.13.4]

$$S(\mathcal{F}^{\bullet \vee}) \cong (-1)^*S(\mathcal{F}^\bullet)^\vee[g]. \quad (6.7)$$

Here $(\mathcal{F}^\bullet)^\vee := R\mathcal{H}om(\mathcal{F}^\bullet, \mathcal{O}_J)$ is the derived dual complex. Finally, by proper base change and (6.3) we see that the restriction of the Fourier-Mukai transform to the identity satisfies

$$S(\mathcal{F}^\bullet)|_{\{\mathcal{O}_C\}} \cong R\pi_*(\mathcal{F}^\bullet), \quad (6.8)$$

where $\pi : J \rightarrow \{\mathcal{O}_C\}$ is the projection to the point.

We can pull back

$$\mathbb{L} := (\alpha_{c_0} \times \alpha_{c_0}^{*-1})^*(\mathbb{P}) \quad (6.9)$$

the Poincaré bundle on $J \times \text{Pic}_0(J)$ to $C \times J$ via the map

$$\alpha_{c_0} \times \alpha_{c_0}^{*-1} : C \times J \rightarrow J \times \text{Pic}_0(J)$$

to get a universal line bundle on $C \times J$ satisfying

$$\mathbb{L}|_{C \times \{L\}} \cong L \quad (6.10)$$

for all $L \in J$ and normalized by an isomorphism

$$\beta : \mathbb{L}|_{\{c_0\} \times J} \xrightarrow{\cong} \mathcal{O}_J. \quad (6.11)$$

Remark 6.1. As a universal line bundle on $C \times J$ the line bundle \mathbb{L} is not unique as we can tensor it with the pull back of any line bundle on J , and the universal property still holds. However (\mathbb{L}, β) is the unique universal object for the functor assigning to a scheme T families of framed line bundles (L, β) , where L is a line bundle on $C \times T$ and $\beta : L|_{\{c_0\} \times T} \cong \mathcal{O}_T$.

Finally, we denote by $\pi : \mathbf{C} \rightarrow C$ a finite morphism of smooth projective curves, by \tilde{J} the Jacobian of \mathbf{C} and $\tilde{\mathbb{P}}$ denotes the Poincaré bundle on $\tilde{J} \times \tilde{J}$. Then, in this setting, we have the norm map $Nm_\pi : \tilde{J} \rightarrow J$ by

$$L \mapsto \det(\pi_*(M)) \otimes \det(\pi_*(\mathcal{O}_{\mathbf{C}}))^{-1}, \quad (6.12)$$

which is a group homomorphism.

Lemma 6.2. *For $M \in J$ we have $\pi^*(M) \in \tilde{J}$ and*

$$\tilde{\mathbb{P}}_{\pi^*(M)} \cong Nm_\pi^*(\mathbb{P}_M).$$

Proof. Fix $\tilde{c}_0 \in \mathbf{C}$ a lift such that $c_0 = \pi(\tilde{c}_0)$. Then by [BL, p. 331] the diagram

$$\begin{array}{ccc} \text{Pic}_0(\tilde{J}) & \xrightarrow{\alpha_{\tilde{c}_0}^*} & \tilde{J} \\ Nm_\pi^* \uparrow & & \uparrow \pi^* \\ \text{Pic}_0(J) & \xrightarrow{\alpha_{c_0}^*} & J \end{array}$$

commutes. Thus

$$\alpha_{\tilde{c}_0}^*(Nm_\pi^*(\mathbb{P}_M)) = \pi^*(\alpha_{c_0}^*(\mathbb{P}_M)) = \pi^*(M).$$

By (6.4) this implies the statement. \square

6.2 Fourier-Mukai transform of $\mathcal{O}_{W_{\mathcal{E}}^+}$

In this section we will fix $d = -n(n-1)(g-1)$ the degree of our Higgs bundles. This will ensure that the line bundle U defined in (5.17) corresponding to our Higgs bundles on a spectral curve will have degree 0. In particular, the canonical uniformising Higgs bundle $\mathcal{E}_0 \in \mathcal{M}$ and the canonical Hitchin section $W_{\mathcal{E}_0}^+ \subset \mathcal{M}$ will be contained in our Higgs moduli space \mathcal{M} .

Consider now a very stable type $(1, \dots, 1)$ Higgs bundle $\mathcal{E} \in \mathcal{M}^{\text{st}}$. We shall compute the relative Fourier-Mukai transform of $\mathcal{O}_{W_{\mathcal{E}}^+}$ over a certain open subset of \mathcal{A} . Denote by $\mathbf{C} \subset T^*C \times \mathcal{A}$ the universal spectral curve. We denote by $\mathcal{A}^\# \subset \mathcal{A}$ the open subset of characteristic equations $a \in \mathcal{A}^\#$ for which the following properties hold

1. C_a is smooth
2. $\text{div}(b)$ avoids the ramification divisor of the spectral cover $\pi_a : C_a \rightarrow C$.

We let

$$\mathbf{C}^\# \subset T^*C \times \mathcal{A}^\#$$

denote the family of spectral curves over $\mathcal{A}^\#$. By the BNR correspondence we can identify $\mathcal{M}^\# := h^{-1}(\mathcal{A}^\#)$ with the relative Jacobian of the family $\mathbf{C}^\# \rightarrow \mathcal{A}^\#$:

$$\mathcal{M}^\# \cong J(\mathbf{C}^\#/\mathcal{A}^\#).$$

We shall denote the relative Poincaré bundle on

$$J(\mathbf{C}^\#/\mathcal{A}^\#) \times_{\mathcal{A}^\#} J(\mathbf{C}^\#/\mathcal{A}^\#)$$

by $\tilde{\mathbb{P}}$. In this section we will determine the relative Fourier-Mukai transform

$$S(\mathcal{O}_{W_\mathcal{E}^+}|_{\mathcal{M}^\#}) = (\pi_2)_*(\pi_1^*(\mathcal{O}_{W_\mathcal{E}^+}|_{\mathcal{M}^\#}) \otimes^L \tilde{\mathbb{P}})$$

over $\mathcal{A}^\#$.

Étale locally $U_i \rightarrow \mathcal{M}^s$ we can define [Si1, Theorem 4.7.(4)] universal Higgs bundles $(\mathbb{E}_{U_i}, \Phi_{U_i})$ on $U_i \times C$. More precisely, choosing a point $c_0 \in C$, we use the framed moduli space \mathcal{R}^s [Si1, Theorem 4.10]. Then $C \times \mathcal{R}^s$ carries a unique universal framed Higgs bundle $(\mathbb{E}_{\mathcal{R}^s}, \Phi_{\mathcal{R}^s}, \beta_{\mathcal{R}^s})$, where the map $\mathbb{E}_{\mathcal{R}^s}$ is a rank n vector bundle on $C \times \mathcal{R}^s$, the universal Higgs field $\Phi_{\mathcal{R}^s} \in H^0(C \times \mathcal{R}^s; \text{End}(\mathbb{E}_{\mathcal{R}^s} \otimes K))$ and the universal framing $\beta_{\mathcal{R}^s} : \mathbb{E}_{\mathcal{R}^s}|_{\{c_0\} \times \mathcal{R}^s} \xrightarrow{\cong} \mathcal{O}_{\mathcal{R}^s}^n$. We can take slices $U_i \subset \mathcal{R}^s$ of the PGL_n -principal bundle $\mathcal{R}^s \rightarrow \mathcal{M}^s$ by Luna's slice theorem [Lu, Theorem III.1] such that $U_i \rightarrow \mathcal{M}^s$ is étale and $\coprod U_i \rightarrow \mathcal{M}^s$ covers \mathcal{M}^s . Then $(\mathbb{E}_{U_i}, \Phi_{U_i})$ is just the Higgs bundle part of the family of framed Higgs bundles $(\mathbb{E}_{U_i}, \Phi_{U_i}, \beta_{U_i})$ which is obtained as the pull back of the universal framed Higgs bundle $(\mathbb{E}_{\mathcal{R}^s}, \Phi_{\mathcal{R}^s}, \beta_{\mathcal{R}^s})$ from $C \times \mathcal{R}^s$. However denoting the *normalized determinant*

$$\text{Norm}(E) := \det(E) \otimes K^{\binom{n}{2}}$$

for a vector bundle (or a family of vector bundles) on C this gives us the additional isomorphism

$$\phi_i : \text{Norm}(\mathbb{E}_{U_i}) \xrightarrow{\cong} \text{Norm}^*(\mathbb{L}), \quad (6.13)$$

where $\text{Norm} : \mathcal{M}^s \rightarrow J$ is given by

$$\text{Norm}(E, \Phi) := \text{Norm}(E). \quad (6.14)$$

This is because $(\text{Norm}(\mathbb{E}_{U_i}), \text{Norm}(\beta_{U_i}))$ is a U_i -family of framed degree 0 line bundles on C , which is unique, see Remark 6.1.

On the overlap $U_i \times_{\mathcal{M}^s} U_j$ of two of our étale covers [HT, (4.2)] implies that we get a line bundle $L_{ij} \in \text{Pic}(U_i \times_{\mathcal{M}^s} U_j)$ which satisfies

$$(\mathbb{E}_{U_i}, \Phi_{U_i}) \cong (L_{ij}\mathbb{E}_{U_j}, \Phi_{U_j}).$$

Passing to a refinement of the étale covering if necessary we can assume that all L_{ij} are trivial. Namely, we can trivialise L_{ij} on $U_i \times_{\mathcal{M}^s} U_j$ even on a Zariski covering, and then we can use [Ar, Theorem 4.1] (see also [Mi, Lemma III.2.19]) to refine the original coverings so that the connecting line bundles will trivialise on the overlaps. It is obvious how to do this in the analytic topology, the étale version which we need is more complicated.

This way we can assume that

$$(\mathbb{E}_{U_i}, \Phi_{U_i}) \cong (\mathbb{E}_{U_j}, \Phi_{U_j})$$

on $U_i \times_{\mathcal{M}^s} U_j$ for all ij . Taking such an isomorphism

$$\phi_{ij} : \mathbb{E}_{U_i} \xrightarrow{\cong} \mathbb{E}_{U_j}$$

and recalling the fixed isomorphisms (6.13) we consider

$$\phi_j \circ \text{Norm}(\phi_{ij}) \circ \phi_i^{-1} : \text{Norm}^*(\mathbb{L}) \rightarrow \text{Norm}^*(\mathbb{L}).$$

By passing to an étale cover if needed we can assume that this function has an n th root (for this we use again [Ar, Theorem 4.1]), and thus we can rescale ϕ_{ij} so that we can assume that

$$\phi_j \circ \text{Norm}(\phi_{ij}) \circ \phi_i^{-1} = \text{Id}_{\text{Norm}^*(\mathbb{L})}.$$

The isomorphisms ϕ_{ij} with this property are unique only up to multiplying with an n th root of unity, but we can make sure that $\phi_{ji} = \phi_{ij}^{-1}$. Thus $\phi_{jk} \circ \phi_{ij}$ agrees with ϕ_{ik} up to an n th root of unity, which we denote by $\theta_{ijk} \in \mu_n \subset \mathbb{C}^\times$ so that

$$\phi_{jk} \circ \phi_{ij} = \theta_{ijk} \phi_{ik}.$$

Computing

$$\theta_{ijk} \theta_{kli} \text{Id}_{\mathbb{E}_{U_k}} = \theta_{ijk} \phi_{ik} \circ \theta_{kli} \phi_{ki} = \phi_{jk} \circ \phi_{ij} \circ \phi_{li} \circ \phi_{kl} \in H^0(U_i \times_{\mathcal{M}^s} U_j \times_{\mathcal{M}^s} U_k \times_{\mathcal{M}^s} U_l; \text{End}(\mathbb{E}_{U_k}))$$

and similarly

$$\theta_{jkl} \theta_{lij} \text{Id}_{\mathbb{E}_{U_l}} = \phi_{kl} \circ \phi_{jk} \circ \phi_{ij} \circ \phi_{li} \in H^0(U_i \times_{\mathcal{M}^s} U_j \times_{\mathcal{M}^s} U_k \times_{\mathcal{M}^s} U_l; \text{End}(\mathbb{E}_{U_l})).$$

These two scalar endomorphisms are conjugate by ϕ_{kl} and therefore the scalars agree:

$$\theta_{ijk} \theta_{kli} = \theta_{jkl} \theta_{lij}.$$

It follows that the n th roots of unity θ_{ijk} form a Čech 2-cocycle θ , thus define a cohomology class $[\theta] \in H^2(\mathcal{M}^s; \mu_n)$, that is a μ_n -gerbe. By definition [Ca, §1.2] the data of the vector bundles \mathbb{E}_{U_i} and ϕ_{ij} constitute a θ -twisted vector bundle \mathbb{E} on $\mathcal{M}^s \times C$, when θ is

considered a Čech 2-cocycle for the sheaf $\mathcal{O}_{\mathcal{M}^s}^*$ via the embedding $\mathbb{P}_n \subset \mathbb{C}^\times$. We can multiply ϕ_{ij} by $\alpha_{ij} \in \mathbb{P}_n$ for a 1-cocycle representing $[\alpha] \in H^1(\mathcal{M}^s; \mathbb{P}_n)$, i.e. an order n line bundle, then all of the above will hold. Thus the twisted bundle \mathbb{E} constructed above is well-defined up to tensoring with an order n line bundle on \mathcal{M}^s but normalized such that

$$\text{Norm}(\mathbb{E}) \cong \text{Norm}^*(\mathbb{L}). \quad (6.15)$$

We also note that the 2-cocycle θ does not change if we rescale with the 1-cocycle α as above.

As for a rank n vector bundle E and line bundle M we have $\Lambda^i(ME) = M^i \Lambda^i(E)$ and it follows that $\Lambda^i(\mathbb{E})$ is naturally a θ^i -twisted sheaf. For $(\ell, m_1, \dots, m_{n-1}) \in \mathbb{Z} \times \mathbb{Z}_{\geq 0}$ and $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1}) \in J_\ell(C) \times C^{[m_1]} \times \dots \times C^{[m_{n-1}]}$ with $\delta_0 = c_{01} + \dots + c_{0m_0}$ where $c_{0j} \in C$ or $-c_{0j} \in C$ is a point in C and $\delta_i = c_{i1} + \dots + c_{im_i} \in C^{[m_i]}$ for points $c_{ij} \in C$ we consider

$$\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}), \quad (6.16)$$

where $\mathbb{E}_{c_{ij}} = \mathbb{E}|_{\mathcal{M}^s \times \{c_{ij}\}}$ when $c_{ij} \in C$ and $\mathbb{E}_{c_{0j}} = \mathbb{E}^*|_{\mathcal{M}^s \times \{-c_{0j}\}}$, when $-c_{0j} \in C$. Then for a line bundle M on \mathcal{M}^s we have

$$\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(M\mathbb{E}_{c_{ij}}) = M^D \bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}),$$

where

$$D = n\ell + \sum_{i=1}^n (n-i)m_i. \quad (6.17)$$

Thus (6.16) is a θ^D -twisted sheaf on \mathcal{M}^s . In particular, when $n|D$ the 2-cocycle $\theta^D = 1$ is trivial, thus in this case (6.16) descends as a vector bundle to $C \times \mathcal{M}^s$. Additionally, when $n|D$ the ambiguity of tensoring \mathbb{E} with an order n line bundle disappears, and so (6.16) is a well-defined vector bundle on \mathcal{M}^s , which is normalised such that

$$\det \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}) \right) \cong \text{Norm}^* \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \mathbb{L}_{c_{ij}}^{\binom{n-1}{n-i-1}} \right).$$

Let $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1}) \in J_\ell(C) \times C^{[m_1]} \times \dots \times C^{[m_{n-1}]} \subset \mathcal{M}$ such that \mathcal{E}_δ is very stable. In other words $\delta_1 + \dots + \delta_{n-1}$ is reduced, and we denote $\delta_i = c_{i1} + \dots + c_{im_i}$, for points $c_{ij} \in C$ or $-c_{0j} \in C$. We also introduce the notation

$$\mathcal{L}_\delta := \mathcal{O}_{W_{\mathcal{E}_\delta}^+}$$

for the structure sheaf of the Lagrangian upward flow of \mathcal{E}_δ . We recall that in this section we set $d = -n(n-1)(g-1)$ for our Higgs bundles and clearly $d = D - n(n-1)(g-1)$ of (6.17) thus $D = 0$ and

$$\Lambda_\delta := \bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})$$

is a well-defined vector bundle on \mathcal{M}^s . With this notation we have

Theorem 6.3. *The relative Fourier-Mukai transform over $\mathcal{A}^\#$ satisfies*

$$S(\mathcal{L}_\delta|_{\mathcal{M}^\#}) = \Lambda_\delta|_{\mathcal{M}^\#}. \quad (6.18)$$

Proof. First we check the theorem fibrewise along h . We let $a \in \mathcal{A}^\#$ so that C_a is smooth, $\text{div}(b)$ avoids the ramification divisor of the spectral cover $\pi_a : C_a \rightarrow C$ and where $W_\mathcal{E}^+ \cap h^{-1}(a)$ is transversal.

As \mathcal{E} is a very stable type $(1, \dots, 1)$ Higgs bundle, we have from Theorem 4.13 that $b = b_{n-1} \circ \dots \circ b_1$ has no repeated 0. We will denote by c_{i1}, \dots, c_{im_i} the m_i zeroes of b_i . For $1 \leq i \leq n-1$ and $1 \leq j \leq m_i$ denote by $\tilde{c}_{ij}^1, \dots, \tilde{c}_{ij}^n \subset C_a$ the n points in the preimage $\pi_a^{-1}(c_{ij})$ of the zero c_{ij} of b_i . From Proposition 5.18 we see that $L \in W_\mathcal{E}^+ \cap h^{-1}(a) \cong W_\mathcal{E}^+ \cap J(C_a)$ is a degree 0 line bundle on C_a of the form $L \cong \pi^*(E_1)(\Delta_1)$ where $\Delta_1 = \sum_{i=1}^{n-1} \Delta_i - \Delta_{i+1}$. We know all possibilities for the divisor $\Delta_i - \Delta_{i+1}$ on C_a ; namely it is effective and reduced and it has the form

$$\Delta_i - \Delta_{i+1} = \sum_{j=1}^{m_i} \tilde{c}_{ij}^{k_{ij}^1} + \dots + \tilde{c}_{ij}^{k_{ij}^{n-i}}$$

where $1 \leq k_{ij}^1 < \dots < k_{ij}^{n-i} \leq n$. Thus we see that

$$\mathcal{O}_{W_\mathcal{E}^+ \cap h^{-1}(a)} = \bigoplus_{\{1 \leq k_{ij}^1 < \dots < k_{ij}^{n-i} \leq n\}_{1 \leq j \leq m_i, 1 \leq i \leq n}} \mathcal{O}_{\pi^*(E_1)} \left(\sum_{i=1}^{n-1} \sum_{j=1}^{m_i} \tilde{c}_{ij}^{k_{ij}^1} + \dots + \tilde{c}_{ij}^{k_{ij}^{n-i}} \right) \quad (6.19)$$

is a direct sum of skyscraper sheaves. Fixing a lift $\tilde{c}_0 \in C_a$ of $c_0 \in C$ we can compute the

Fourier-Mukai transform of (6.19) as follows:

$$\begin{aligned}
S(\mathcal{O}_{W_{\mathcal{E}}^+ \cap h^{-1}(a)}) &= \bigoplus_{\{1 \leq k_{ij}^1 < \dots < k_{ij}^{n-i} \leq n\}_{1 \leq j \leq m_i, 1 \leq i \leq n}} \bigoplus S \left(\mathcal{O}_{\pi^*(E_1)} \left(\sum_{i=1}^{n-1} \sum_{j=1}^{m_i} \tilde{c}_{ij}^{k_{ij}^1} + \dots + \tilde{c}_{ij}^{k_{ij}^{n-i}} \right) \right) \\
&= \bigoplus_{\{1 \leq k_{ij}^1 < \dots < k_{ij}^{n-i} \leq n\}_{1 \leq j \leq m_i, 1 \leq i \leq n}} \bigoplus S \left(\mathcal{O}_{\pi^*(E_1)} \left((\sum_{i=1}^n (n-i)m_i) \tilde{c}_0 + \sum_{i=1}^{n-1} \sum_{j=1}^{m_i} \tilde{c}_{ij}^{k_{ij}^1} - \tilde{c}_0 + \dots + \tilde{c}_{ij}^{k_{ij}^{n-i}} - \tilde{c}_0 \right) \right) \\
&= \bigoplus_{\{1 \leq k_{ij}^1 < \dots < k_{ij}^{n-i} \leq n\}_{1 \leq j \leq m_i, 1 \leq i \leq n}} \tilde{\mathbb{P}}_{\pi^*(E_1)(-n\ell\tilde{c}_0)} \otimes \bigotimes_{i=1}^{n-1} \bigotimes_{j=1}^{m_i} \tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^1} - \tilde{c}_0)} \otimes \dots \otimes \tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^{n-i}} - \tilde{c}_0)} \\
&= \tilde{\mathbb{P}}_{\pi^*(E_1)(-n\ell\tilde{c}_0)} \otimes \bigotimes_{i=1}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i} \left(\tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^1} - \tilde{c}_0)} \oplus \dots \oplus \tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^{n-i}} - \tilde{c}_0)} \right)
\end{aligned}$$

We see that

$$\tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^1} - \tilde{c}_0)} = \tilde{\mathbb{L}}|_{J(C_a) \times \{\tilde{c}_{ij}^{k_{ij}^1}\}},$$

where $\tilde{\mathbb{L}}$ is a universal Poincaré bundle on $J(C_a) \times C_a$ normalized at \tilde{c}_0 as defined in (6.9). Furthermore,

$$\tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^1} - \tilde{c}_0)} \oplus \dots \oplus \tilde{\mathbb{P}}_{(\tilde{c}_{ij}^{k_{ij}^{n-i}} - \tilde{c}_0)} = \pi_*(\tilde{\mathbb{L}})|_{h^{-1}(a) \times \{c_{ij}\}}, \quad (6.20)$$

and $(\pi_*(\tilde{\mathbb{L}}), \Phi_{h^{-1}(a)})$ with $\Phi_{h^{-1}(a)} := \pi_*(x : \tilde{\mathbb{L}} \rightarrow \tilde{\mathbb{L}} \otimes \pi^*(K))$ is a universal Higgs bundle on $h^{-1}(a) \times C$.

Lemma 6.4. For $L \in J(C_a)$ let $Nm : J(C_a) \rightarrow J(C)$ given by $Nm(L) = \det(\pi_*(L)) \otimes K^{\binom{n}{2}}$ the usual norm map. Then on $J(C_a) \times C$ we have

$$\text{Norm}((\pi_a)_*(\tilde{\mathbb{L}})) = Nm^*(\mathbb{L}),$$

where $\tilde{\mathbb{L}}$ is a universal line bundle on $J(C_a) \times \tilde{C}_a$ normalized at $\tilde{c}_0 \in \tilde{C}_a$ and \mathbb{L} is a universal line bundle on $J(C) \times C$ normalized at $c_0 = \pi_a(\tilde{c}_0)$.

Proof. For $L \in J(C_a)$ we have

$$Nm^*(\mathbb{L})|_{\{L\} \times C} = Nm(L) = \text{Norm}(\pi_*(\tilde{\mathbb{L}}))|_{\{L\} \times C}.$$

On the other hand for $c \in C$ we have

$$Nm^*(\mathbb{L})|_{J(C_a) \times \{c\}} = Nm^*(\mathbb{L}|_{J(C) \times \{c\}}) = Nm^*(\mathbb{P}_{(c-c_0)}).$$

While for the other side

$$\text{Norm}(\pi_*(\tilde{\mathbb{L}}))|_{J(C_a) \times \{c\}} = \det(\pi_*(\tilde{\mathbb{L}})|_{J(C_a) \times \{c\}}) = \det \left(\tilde{\mathbb{P}}_{(\tilde{c}_1 - \tilde{c}_0)} + \dots + \tilde{\mathbb{P}}_{(\tilde{c}_n - \tilde{c}_0)} \right) = \tilde{\mathbb{P}}_{\pi^*(c-c_0)}$$

where $\pi^{-1}(c) = \{\tilde{c}_1, \dots, \tilde{c}_n\}$.

Now Lemma 6.2 implies that

$$\det(\pi_*(\tilde{\mathbb{L}}))|_{J(C_a) \times \{c\}} = \det^*(\mathbb{L})|_{J(C_a) \times \{c\}}.$$

The result follows. \square

The norm map $Nm : J(C_a) \rightarrow J(C)$ in Lemma 6.4 agrees with the one restricted from the normalised determinant map in (6.14) to $h^{-1}(a) \cong J(C_a)$. Thus $\text{Norm}(\pi_*(\tilde{\mathbb{L}})) = \text{Norm}^*(\mathbb{L})$ and therefore $\pi_*(\tilde{\mathbb{L}}) \cong \mathbb{E}$ on $h^{-1}(a)$. Furthermore $\tilde{\mathbb{P}}_{\pi^*(E_1)(-nl\tilde{c}_0)} \cong \text{Norm}^*(\mathbb{P}_{E_1(-lc_0)})$ from Lemma 6.2. By the normalisation (6.15), the definition (6.9) and $E_1 \cong \mathcal{O}(\delta_0) = \mathcal{O}(c_{01} + \dots + c_{0m_0})$ this agrees with

$$\text{Norm}^*(\mathbb{P}_{E_1(-lc_0)}) \cong \bigotimes_{j=1}^{m_0} \Lambda^n(\mathbb{E}_{c_{0j}}).$$

Thus (6.20) and Lemma 6.4 verify Theorem 6.20 on $h^{-1}(a)$.

By noting that étale-locally on $\mathcal{A}^\#$ we have sections of the universal spectral curve corresponding to the preimages of zeroes of b and c_0 , we see that the same argument can be repeated relative to étale neighbourhoods $U_i \rightarrow \mathcal{A}^\#$, proving Theorem 6.3 étale-locally, which as argued above glue together to prove Theorem 6.3. \square

Remark 6.5. Vector bundle duality defines a natural involution

$$\begin{aligned} \iota : \mathcal{M} &\rightarrow \mathcal{M} \\ (E, \Phi) &\mapsto (E^*K^{1-n}, \Phi^T) \end{aligned} \quad (6.21)$$

where Φ^T is the induced action of Φ on E^* . This is well-defined as $\deg(E^*K^{1-n}) = -\deg(E) - n(n-1)2(g-1) = -n(n-1)(g-1)$. From the exact sequence (5.20) we see that $(\pi_a)_*(U^*) = \iota(E, \Phi)$ and therefore ι restricts to the abelian variety $h^{-1}(a) \cong J(C_a)$ as the inverse map -1 . If $\mathcal{E} \in \mathcal{M}$ we will denote $\mathcal{E}^\iota := \iota(\mathcal{E})$, then $\iota(W_{\mathcal{E}}^+) = W_{\mathcal{E}^\iota}^+$.

For example for a type $(1, \dots, 1)$ very stable Higgs bundle \mathcal{E}_δ we can determine $\mathcal{E}_\delta^\iota = \mathcal{E}_{\delta^\iota}$, where $\delta^\iota = (\delta_0^\iota, \delta_1^\iota, \dots, \delta_{n-1}^\iota) = (-\delta_0 - \delta_1 - \dots - \delta_{n-1}, \delta_{n-1}, \dots, \delta_1)$. We thus see that the skew involutive property (6.6) of the Fourier-Mukai transform yields

$$S(\Lambda_\delta|_{\mathcal{M}^\#}) = \mathcal{L}_{\delta^\iota}[-n^2(g-1) - 1]|_{\mathcal{M}^\#}. \quad (6.22)$$

6.3 An agreement for \mathbb{T} -equivariant Euler forms

As a motivation recall that mirror symmetry should be an equivalence of categories. Not only objects should match but morphisms between them. If two objects have \mathbb{T} -equivariant structure so will the vector space of morphisms between them. Thus its \mathbb{T} -character should agree with the \mathbb{T} -character of the vector space of morphisms between

the mirror objects. In this subsection we find the agreement of these \mathbb{T} -characters (aka Euler pairing) for our mirror pairs of branes. Incidentally, it is easier to pair a Lagrangian brane with a hyperholomorphic one because the computation reduces to a computation on the Lagrangian support which in our case is just a vector space.

6.3.1 \mathbb{T} -equivariant Euler characteristics and Euler forms

Recall that for a \mathbb{T} -equivariant coherent sheaf \mathcal{F} on a semiprojective variety M the *equivariant Euler characteristic* is defined by

$$\chi_{\mathbb{T}}(M; \mathcal{F}) := \sum_{ij} (-1)^i \dim(H^i(M; \mathcal{F})^j) t^{-j},$$

when the weight spaces $H^i(M; \mathcal{F})^j$, where \mathbb{T} acts with weight j , are finite dimensional. The finiteness condition holds for smooth semiprojective varieties and then $\chi_{\mathbb{T}}(M; \mathcal{F}) \in \mathbb{Z}((t))$, i.e. $H^i(M; \mathcal{F})^j = 0$ for i sufficiently large. More generally for a \mathbb{T} -equivariant bounded complex of coherent sheaves \mathcal{F}^\bullet we define $\chi_{\mathbb{T}}(M; \mathcal{F}^\bullet) = (-1)^k \sum_k \chi_{\mathbb{T}}(M; \mathcal{F}^k)$.

Furthermore, given two \mathbb{T} -equivariant coherent sheaves \mathcal{F}_1 and \mathcal{F}_2 on M we define the *equivariant Euler pairing* as

$$\begin{aligned} \chi_{\mathbb{T}}(M; \mathcal{F}_1, \mathcal{F}_2) &= \sum_{k,l} \dim(H^k(R\mathcal{H}om(\mathcal{F}_1, \mathcal{F}_2))^l) (-1)^k t^{-l} = \\ &= \sum_{k,l} \dim(\mathrm{Hom}_{\mathbf{D}_{coh}(M)}(\mathcal{F}_1, \mathcal{F}_2[k])^l) (-1)^k t^{-l} = \sum_{k,l} \dim(\mathrm{Ext}^k(M; \mathcal{F}_1, \mathcal{F}_2)^l) (-1)^k t^{-l}. \end{aligned}$$

As $\chi_{\mathbb{T}}(M; \mathcal{F}_1, \mathcal{F}_2) = \chi_{\mathbb{T}}(M; \mathcal{F}_1^\vee \otimes^L \mathcal{F}_2)$, where \mathcal{F}_1^\vee is the derived dual and \otimes^L is the derived tensor product, we expect that $\chi_{\mathbb{T}}(M; \mathcal{F}_1, \mathcal{F}_2) \in \mathbb{Z}((t))$ on a semiprojective variety M .

We note that the upward flow $W_{\mathcal{E}}^+$ for $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ is invariant under the \mathbb{T} -action and we shall endow $\mathcal{O}_{W_{\mathcal{E}}^+}$ with the trivial \mathbb{T} -equivariant structure.

6.3.2 \mathbb{T} -equivariant universal bundles

On the other hand we would like to endow the sheaf (6.16), when $n|D$, on $\mathcal{M}^s \times C$ with a \mathbb{T} -equivariant structure. For this we need

Lemma 6.6. *We can choose the étale cover $\coprod_i U_i \rightarrow \mathcal{M}^s$ so that the U_i are \mathbb{T} -equivariant and the local universal Higgs bundle (E_{U_i}, Φ_{U_i}) on $U_i \times C$ carries a \mathbb{T} -equivariant structure with \mathbb{T} acting on Φ_{U_i} with weight -1 . Furthermore the isomorphism (6.13) is \mathbb{T} -equivariant with some \mathbb{T} -equivariant structure on \mathbb{L} .*

Proof. First we note that over $\mathcal{R} \times C$ there exists, from [Si1, Theorem 4.10], the universal framed Higgs bundle $(\mathbb{E}_{\mathcal{R}}, \Phi_{\mathcal{R}}, \beta_{\mathcal{R}})$. We have the \mathbb{T} -action on \mathcal{R} multiplying the Higgs field and we denote the morphism $m_{\lambda} : \mathcal{R} \rightarrow \mathcal{R}$ given by scaling the Higgs field with $\lambda \in \mathbb{T}$. Then

$$(m_{\lambda}^*(\mathbb{E}_{\mathcal{R}}), \lambda^{-1}m_{\lambda}^*(\Phi_{\mathcal{R}}), m_{\lambda}^*(\beta_{\mathcal{R}}))$$

is also a universal framed Higgs bundle and as \mathcal{R} is a fine moduli space with a unique universal object we get that $\mathbb{E}_{\mathcal{R}} \cong m_{\lambda}^*(\mathbb{E}_{\mathcal{R}})$ canonically, inducing $f_{\lambda} : \mathbb{E}_{\mathcal{R}} \rightarrow \mathbb{E}_{\mathcal{R}}$ covering m_{λ} and making the diagram

$$\begin{array}{ccc} \mathbb{E}_{\mathcal{R}} & \xrightarrow{\Phi_{\mathcal{R}}} & \mathbb{E}_{\mathcal{R}} \otimes K \\ f_{\lambda} \downarrow & & \downarrow f_{\lambda} \\ \mathbb{E}_{\mathcal{R}} & \xrightarrow{\lambda^{-1}\Phi_{\mathcal{R}}} & \mathbb{E}_{\mathcal{R}} \otimes K \end{array} \quad (6.23)$$

commute. This will endow $\mathbb{E}_{\mathcal{R}}$ with a canonical \mathbb{T} -equivariant structure which acts on $\Phi_{\mathcal{R}}$ with weight -1 .

Now we take a point $\mathcal{E} \in \mathcal{M}^s$ such that the orbit $\mathbb{T}\mathcal{E} \subset \mathcal{M}^s$ is closed. This means that either $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ or that neither $\lim_{\lambda \rightarrow 0} \lambda\mathcal{E}$ nor (if it exists) $\lim_{\lambda \rightarrow \infty} \lambda\mathcal{E}$ is stable. We take a lift $\tilde{\mathcal{E}} \in \mathcal{R}^s$. Recall that PGL_n acts freely on \mathcal{R}^s by changing the framing and \mathbb{T} acts by multiplying the Higgs field. The PGL_n quotient $p : \mathcal{R}^s \rightarrow \mathcal{M}^s$ is \mathbb{T} -equivariant. It follows that for $G := \mathbb{T} \times \mathrm{PGL}_n$ the orbit $G\tilde{\mathcal{E}} \subset \mathcal{R}^s$ is closed. We let $G_{\tilde{\mathcal{E}}} \subset G$ denote the stabilizer which projects injectively into \mathbb{T} as PGL_n acts freely. We apply Luna's slice theorem [Lu, Theorem III.1] for the action of G on \mathcal{R}^s around the point $\tilde{\mathcal{E}}$. This way we get a locally closed $G_{\tilde{\mathcal{E}}}$ -invariant affine slice $\tilde{\mathcal{E}} \in V \subset \mathcal{R}^s$, such that the map $G \times_{G_{\tilde{\mathcal{E}}}} V \rightarrow \mathcal{R}^s$ is étale and whose image $U \subset \mathcal{R}^s$ is a saturated affine open G -invariant subset, the morphism

$$V/G_{\tilde{\mathcal{E}}} \rightarrow U/G \quad (6.24)$$

is also étale and the natural morphism induces

$$G \times_{G_{\tilde{\mathcal{E}}}} V \cong U \times_{U/G} V/G_{\tilde{\mathcal{E}}} \quad (6.25)$$

a G -isomorphism. We let $V' := \mathbb{T} \times_{G_{\tilde{\mathcal{E}}}} V = (G \times_{G_{\tilde{\mathcal{E}}}} V)/\mathrm{PGL}_n$ then $V'/\mathbb{T} = V/G_{\tilde{\mathcal{E}}}$ and dividing (6.25) by PGL_n gives

$$V' = \mathbb{T} \times_{G_{\tilde{\mathcal{E}}}} V \cong U/\mathrm{PGL}_n \times_{U/G} V/G_{\tilde{\mathcal{E}}}.$$

By base change of the étale morphism (6.24) by $U/\mathrm{PGL}_n \rightarrow U/G = (U/\mathrm{PGL}_n)/\mathbb{T}$

$$V' \rightarrow U/\mathrm{PGL}_n \subset \mathcal{M}^s \quad (6.26)$$

is also étale and \mathbb{T} -equivariant. Finally, the \mathbb{T} -equivariant universal Higgs bundle $(\mathbb{E}_{\mathcal{R}}, \Phi_{\mathcal{R}})$ restricts to the \mathbb{T} -equivariant $V' \rightarrow \mathcal{R}$ giving \mathbb{T} -equivariant universal bundles $(\mathbb{E}_{V'}, \Phi_{V'})$ étale locally (6.26) around $\mathcal{E} \in U/\mathrm{PGL}_n$.

Finally, we note that a \mathbb{T} -invariant open covering of the closed \mathbb{T} -orbits in \mathcal{M}^s covers the whole \mathcal{M}^s and so the result follows. \square

Thus we have local \mathbb{T} -equivariant universal Higgs bundles on $U_k \rightarrow \mathcal{M}^s$. On overlaps $C \times U_k \times_{\mathcal{M}^s} U_l$ they differ by a \mathbb{T} -equivariant line bundle by [HT, (4.2)]. Thus the \mathbb{T} -equivariant projective bundles $\mathbb{P}\mathbb{E}_{U_i}$ and $\mathbb{P}\mathbb{E}_{U_j}$ are canonically isomorphic over the overlap. Therefore they descend as a \mathbb{T} -equivariant projective bundle $\mathbb{P}\mathbb{E}$ over $C \times \mathcal{M}^s$.

Lemma 6.7. *In the notation of (6.16) when $n|D = n\ell + \sum_{i=1}^{n-1} (n-i)m_i$ there is a \mathbb{T} -equivariant structure on the vector bundle $\Lambda_\delta = \bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})$ normalised such that*

$$\det \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}) \right) \cong \text{Norm}^* \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \mathbb{L}_{c_{ij}}^{\binom{n-1}{n-i-1}} \right) \quad (6.27)$$

as equivariant line bundles, where the line bundle $\mathbb{L}_{c_{ij}}^{\binom{n-1}{n-i-1}}$ on J is endowed with a weight $\binom{n-i}{2} - \binom{n}{2} - \binom{n-1}{n-i-1}$ \mathbb{T} -action.

Proof. Similarly to $\mathbb{P}\mathbb{E}$ above, we can argue that $\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{U_k}|_{c_{ij} \times U_k})$ on $U_k \rightarrow \mathcal{M}^s$ carries a \mathbb{T} -equivariant structure, and they differ by a \mathbb{T} -equivariant line bundle on the overlaps. Thus the total space of the projective bundle $\mathbb{P} \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}) \right)$ descends to \mathcal{M}^s with a \mathbb{T} -equivariant structure. When $n|D = \sum_{i=0}^{n-1} (n-i)m_i$ then we saw before Theorem 6.3 that even the vector bundle $\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})$ descends to \mathcal{M}^s . Thus the projective bundle $\mathbb{P} \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}) \right)$ has a sheaf $\mathcal{O}(1)$ on it, such that

$$\pi_*(\mathcal{O}(1))^* \cong \bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}). \quad (6.28)$$

By [Br, Theorem 4.2.2] the \mathbb{T} -action on $\mathbb{P} \left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}}) \right)$ lifts to $\mathcal{O}(1)$ because $\text{Pic}(\mathbb{T})$ is trivial and thus the last map in [Br, (12)] is an isomorphism. Thus we have a \mathbb{T} -equivariant structure on $\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})$ from (6.28).

We can modify this \mathbb{T} -equivariant structure by tensoring with a \mathbb{T} -equivariant structure on $\mathcal{O}_{\mathcal{M}^s}$. We can choose it such that at the canonical uniformising Higgs bundle $\mathcal{E}_0 = \mathcal{E}_{\mathcal{O}_C, 0}$ we get that $\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})|_{\mathcal{E}_0}$ has only non-positive weights and non-trivial weight 0 space. We can endow $E_{\mathcal{O}_C} = \mathcal{O} \oplus K^{-1} \oplus \dots \oplus K^{-n+1}$ with a \mathbb{T} -action which acts by weight $-i$ on the i th summand K^{-i} . This will act on Φ_0 with weight -1 . Therefore any \mathbb{T} -equivariant family of stable Higgs bundles containing \mathcal{E}_0 will restrict to \mathcal{E}_0 as this \mathbb{T} -equivariant structure, up to a rescaling of the \mathbb{T} -action by an overall 1-dimensional

\mathbb{T} -module. Therefore our normalisation of the \mathbb{T} -action on $\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})$ will satisfy

$$\chi_{\mathbb{T}}\left(\bigotimes_{i=0}^{n-1} \bigotimes_{j=1}^{m_i} \Lambda^{n-i}(\mathbb{E}_{c_{ij}})\right)|_{\mathcal{E}_0} = \prod_{i=0}^{n-1} \begin{bmatrix} n \\ n-i \end{bmatrix}_t^{m_i} \quad (6.29)$$

which incidently also agrees with the equivariant multiplicity $m_{\mathcal{E}_\delta}(t)$ of (5.16). This will be related to mirror symmetry in (6.33).

A straightforward computation checks that this normalisation of the \mathbb{T} -action will induce the one on the determinant as claimed in the Lemma. \square

The following is an indication that the mirror relationships (6) and (6.22) over $\mathcal{A}^\#$ hold over the whole \mathcal{A} .

Theorem 6.8. *For $\delta = (\delta_0, \delta_1, \dots, \delta_{n-1}) \in J_\ell(C) \times C^{[m_1]} \times \dots \times C^{[m_{n-1}]}$ and $\delta' = (\delta'_0, \delta'_1, \dots, \delta'_{n-1}) \in J_{\ell'}(C) \times C^{[m'_1]} \times \dots \times C^{[m'_{n-1}]}$ let $\mathcal{E}_\delta, \mathcal{E}_{\delta'} \in \mathcal{M}^{\text{st}}$ be two very stable type $(1, \dots, 1)$ Higgs bundles. Then*

$$\begin{aligned} \chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'}, \Lambda_\delta) &= \chi_{\mathbb{T}}(\det(\mathbb{C}_1 \otimes \mathcal{A}^*)) \chi_{\mathbb{T}}(\mathcal{M}; \Lambda_{\delta'}, \mathcal{L}_{\delta'}[-n^2(g-1) - 1]) \\ &= (-1)^{n^2(g-1)+1} t^{(4n+1)(n-1)n(g-1)/6} \chi_{\mathbb{T}}(\mathcal{M}; \Lambda_{\delta'}, \mathcal{L}_{\delta'}). \end{aligned}$$

Proof. We compute

$$\begin{aligned} \chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'}, \Lambda_\delta) &= \chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'}^\vee \otimes^L \Lambda_\delta) = \\ &= \chi_{\mathbb{T}}\left(W_{\delta'}^+; \det(N_{W_{\delta'}^+})[-n^2(g-1) - 1] \otimes \Lambda_\delta|_{W_{\delta'}^+}\right), \end{aligned}$$

where in the last equation we used [Hu, Corollary 3.40] and the simplified notation $W_{\delta'}^+ := W_{\mathcal{E}_{\delta'}}^+$.

As $\det(N_{W_{\delta'}^+}) \otimes \Lambda_\delta|_{W_{\delta'}^+}$ is a \mathbb{T} -equivariant vector bundle on the affine space $W_{\delta'}^+$ its \mathbb{T} -equivariant index can be computed as

$$\chi_{\mathbb{T}}\left(W_{\delta'}^+; \det(N_{W_{\delta'}^+}) \otimes \Lambda_\delta|_{W_{\delta'}^+}\right) = \chi_{\mathbb{T}}\left((\det(N_{W_{\delta'}^+}) \otimes \Lambda_\delta)|_{\mathcal{E}_{\delta'}}\right) \chi_{\mathbb{T}}(W_{\delta'}^+; \mathcal{O}_{W_{\delta'}^+}).$$

The same argument as above for (6.29) gives

$$\chi_{\mathbb{T}}(\Lambda_\delta|_{\mathcal{E}_{\delta'}}) = m_{\mathcal{E}_\delta}(t).$$

From Definition 5.3 we have

$$\chi_{\mathbb{T}}(W_{\delta'}^+; \mathcal{O}_{W_{\delta'}^+}) = m_{\mathcal{E}_{\delta'}}(t) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)).$$

Finally $N_{W_{\delta'}^+}|_{\mathcal{E}_{\delta'}} \cong T_{\mathcal{E}_{\delta'}}^{\leq 0}$ consequently

$$\chi_{\mathbb{T}}(\det(N_{W_{\delta'}^+}|_{\mathcal{E}_{\delta'}})) = \chi_{\mathbb{T}}(\det(T_{\mathcal{E}_{\delta'}}^{\leq 0}))$$

as $\dim \mathcal{A} = \dim T_{\mathcal{E}_{\delta'}}^+ = \dim \mathcal{M}/2$. Thus for the left hand side we have

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'}, \Lambda_{\delta}) = (-1)^{\dim \mathcal{A}} m_{\mathcal{E}_{\delta'}}(t) m_{\mathcal{E}_{\delta'}}(t) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)) \chi_{\mathbb{T}}(\det(T_{\mathcal{E}_{\delta'}}^{\leq 0})). \quad (6.30)$$

For the other side we compute similarly

$$\begin{aligned} \chi_{\mathbb{T}}(\mathcal{M}; \Lambda_{\delta'}, \mathcal{L}_{\delta'}) &= \chi_{\mathbb{T}}(\mathcal{M}; \Lambda_{\delta'}^{\vee} \otimes^L \mathcal{L}_{\delta'}) \\ &= \chi_{\mathbb{T}}(W_{\delta'}^+; \Lambda_{\delta'}^{\vee}|_{W_{\delta'}^+}) \\ &= \chi_{\mathbb{T}}(\Lambda_{\delta'}^{\vee}|_{\mathcal{E}_{\delta'}}) \chi_{\mathbb{T}}(W_{\delta'}^+; \mathcal{O}_{W_{\delta'}^+}) \\ &= m_{\mathcal{E}_{\delta'}}(t^{-1}) m_{\mathcal{E}_{\delta'}}(t) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)) \\ &= \frac{\chi_{\mathbb{T}}(\det(\mathcal{A}))}{\chi_{\mathbb{T}}(\det(T_{\mathcal{E}_{\delta'}}^+))} m_{\mathcal{E}_{\delta'}}(t) m_{\mathcal{E}_{\delta'}}(t) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)) \\ &= \frac{\chi_{\mathbb{T}}(\det(T_{\mathcal{E}_{\delta'}}^{\leq 0}))}{\chi_{\mathbb{T}}(\det(\mathbb{C}_1 \otimes \mathcal{A}^*))} m_{\mathcal{E}_{\delta'}}(t) m_{\mathcal{E}_{\delta'}}(t) \chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*)), \end{aligned} \quad (6.31)$$

where in the fourth equation we used (5.13). For the last equation we can argue as follows. As ι of (6.21) is a \mathbb{T} -equivariant isomorphism, it follows that $T_{\mathcal{E}_{\delta'}}^+ \cong T_{\mathcal{E}_{\delta}}^+$ and thus $m_{\mathcal{E}_{\delta'}}(t) = m_{\mathcal{E}_{\delta}}(t)$. Using the homogeneity 1 symplectic form on $T_{\mathcal{E}_{\delta'}}$ we see that $\chi_{\mathbb{T}}(\mathbb{C}_{-1} \otimes \det(T_{\mathcal{E}_{\delta'}}^+)) \chi_{\mathbb{T}}(\det(T_{\mathcal{E}_{\delta'}}^{\leq 0})) = 1$. This implies the last equation. Comparing (6.30) and (6.31) implies the first equation in the Theorem.

For the second equation we note that $\dim \mathcal{A} = \sum_{i=1}^n \dim H^0(C; K^i) = n^2(g-1) + 1$ and that a straightforward computation gives

$$\chi_{\mathbb{T}}(\mathbb{C}_1 \otimes \mathcal{A}^*) = t^{\sum_i (i-1) \dim \mathcal{A}_i} = t^{\sum_{i=1}^n (i-1) \dim H^0(C; K^i)} = t^{\sum_{i=1}^n (i-1)(2i-1)(g-1)} = t^{(4n+1)(n-1)n(g-1)/6}.$$

The result follows. □

Remark 6.9. We proved in Theorem 6.3 that the relative Fourier-Mukai transform $S(\mathcal{L}_{\delta}) = \Lambda_{\delta}$ holds over $\mathcal{A}^{\#}$ and in Remark 6.5 that $S(\Lambda_{\delta}) = \mathcal{L}_{\delta'}[-n^2(g-1) - 1]$ also holds over $\mathcal{A}^{\#}$. The above theorem would be a consequence of a \mathbb{T} -equivariant extension of the mirror symmetry conjecture in the "classical limit" as in [DP1, Conjecture 2.5].

Remark 6.10. Similar agreement of \mathbb{T} -equivariant Euler pairings between certain conjectured mirror pairs of Lagrangian and hyperholomorphic branes of [Hi4] were observed in [HMP, §7], when $n = 2$. When $n = 2$ we have $\chi_{\mathbb{T}}(\mathbb{C}_1 \otimes \mathcal{A}^*) = t^{3g-3}$ in Theorem 6.8 which we cannot get rid of by just renormalising the \mathbb{T} -action on our branes when $3g-3$ is odd. The same phenomenon was observed in [HMP, Remark 7.7].

In fact, one can also pair certain $n = 2$ analogues of the mirror pairs of this paper with the ones in [HMP] and find the expected agreement.

Remark 6.11. Just as in [HMP, Corollary 4.11] we get a clearly symmetric expression in δ and δ' if we compute, equivalently as in (6.30),

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\delta'} \otimes \Lambda_{\delta}) = m_{\mathcal{E}_{\delta'}}(t)m_{\mathcal{E}_{\delta}}(t)\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*)). \quad (6.32)$$

When deriving the agreement of the corresponding Euler forms [HMP, Corollary 7.1] we basically argued that it follows from the fact that both of the branes \mathcal{L}_i and Λ_j were self-dual. In our present situation $\mathcal{L}_{\delta'}$ is self-dual (up to a shift) for the trivial reason that it is supported on an affine space. On the other hand Λ_{δ} is typically not self-dual, rather it satisfies $\Lambda_{\delta}^{\vee} = \iota^*(\Lambda_{\delta}) = \Lambda_{\delta'}$ using the normalised inverse map ι of Remark 6.5. However in [HMP] both types of our branes \mathcal{L}_i and Λ_j were secretly invariant under ι as well. In general, what (6.6) tells us is that self-dual branes should be mirror to branes invariant under $\vee \circ \iota$. The former is satisfied by $\mathcal{L}_{\delta'}$ and the latter by Λ_{δ} .

Remark 6.12. It is instructive to take $\delta' = 0 = (0, 0, \dots, 0)$ in (6.32). Then $\mathcal{E}_0 = \mathcal{E}_{\mathcal{O}_C, 0}$ is the canonical uniformising Higgs bundle of (3.16). It has $m_{\mathcal{E}_0}(t) = 1$ and $\mathcal{L}_0 \cong \mathcal{O}_{W_0^+}$ is the structure sheaf of the canonical Hitchin section, while $\Lambda_0 \cong \mathcal{O}_{\mathcal{M}^s}$. We get

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{O}_{W_0^+} \otimes \Lambda_0) = m_{\mathcal{E}_0}(t)\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*)).$$

On the other hand

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{O}_{W_0^+} \otimes \Lambda_{\delta}) = \chi_{\mathbb{T}}(\mathcal{O}_{W_0^+}; \Lambda_{\delta}|_{W_0^+}) = \chi_{\mathbb{T}}(\Lambda_{\delta}|_{\mathcal{E}_0})\chi_{\mathbb{T}}(\mathrm{Sym}(\mathcal{A}^*)).$$

Thus $\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{O}_{W_0^+} \otimes \Lambda_0) = \chi_{\mathbb{T}}(\mathcal{M}; \mathcal{O}_{W_0^+} \otimes \Lambda_{\delta})$ (or the equivalent Theorem 6.8) implies that

$$\chi_{\mathbb{T}}(\Lambda_{\delta}|_{\mathcal{E}_0}) = m_{\mathcal{E}_0}(t). \quad (6.33)$$

This was observed already in (6.29), but here we derived it as a consequence of mirror symmetry. Indeed such a relationship is expected from the mirror in general. Because of the restriction property (6.8) of the Fourier-Mukai transform to the identity we expect from the fibrewise Fourier-Mukai transform that the mirror of a bounded complex of coherent sheaves \mathcal{F}^{\bullet} on \mathcal{M} should restrict to the canonical Hitchin section as $h^*(h_*(\mathcal{F}^{\bullet}))$. The latter was computed for the upward flow of a very stable Higgs bundle in (5.9) and (5.11). Thus (6.33) verifies this expectation from the mirror of \mathcal{L}_{δ} .

Remark 6.13. We can also generalise these ideas further for very stable upward flows of type not $(1, \dots, 1)$. For $\mathcal{E} \in \mathcal{M}^{s\mathbb{T}}$ a very stable Higgs bundle let $\mathcal{L}_{\mathcal{E}} := \mathcal{O}_{W_{\mathcal{E}}^+}$ denote the structure sheaf of its upward flow and $\Lambda_{\mathcal{E}}$ denote its mirror, which we expect to be a \mathbb{T} -equivariant vector bundle on \mathcal{M}^s of rank $m_{\mathcal{E}}$. If $\mathcal{F} \in \mathcal{M}^{s\mathbb{T}}$ is another very stable Higgs bundle then denote by

$$m_{\mathcal{E}, \mathcal{F}}(t) := \chi_{\mathbb{T}}(\Lambda_{\mathcal{E}}|_{\mathcal{F}}) \in \mathbb{Z}[t]$$

the \mathbb{T} -character of the fibre $\Lambda_{\mathcal{E}}|_{\mathcal{F}}$, which we expect to contain only non-positive weights. In particular, by (6.33) we have $m_{\mathcal{E},\mathcal{E}_0}(t) = m_{\mathcal{E}}(t)$ the equivariant multiplicity. As above from mirror symmetry we expect

$$\chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\mathcal{E}} \otimes \Lambda_{\mathcal{F}}) = \chi_{\mathbb{T}}(\mathcal{M}; \mathcal{L}_{\mathcal{F}} \otimes \Lambda_{\mathcal{E}})$$

which in turn leads to the identity

$$m_{\mathcal{E}}(t)m_{\mathcal{F},\mathcal{E}}(t) = m_{\mathcal{F}}(t)m_{\mathcal{E},\mathcal{F}}(t). \quad (6.34)$$

When both \mathcal{E} and \mathcal{F} are of type $(1, \dots, 1)$ then $m_{\mathcal{E},\mathcal{F}}(t) = m_{\mathcal{E}}(t)$ is just the equivariant multiplicity as $\Lambda_{\mathcal{E}}|_{\mathcal{F}} \cong \Lambda_{\mathcal{E}}|_{\mathcal{E}_0}$ and so (6.34) holds. If only \mathcal{F} is of type $(1, \dots, 1)$ we know what $\Lambda_{\mathcal{F}}$ should be and how it restricts to any \mathcal{E} giving a formula for $m_{\mathcal{F},\mathcal{E}}(t)$ only depending on the type of \mathcal{E} . From (6.34) we can deduce

$$m_{\mathcal{E},\mathcal{F}}(t) = \frac{m_{\mathcal{E}}(t)m_{\mathcal{F},\mathcal{E}}(t)}{m_{\mathcal{F}}(t)}.$$

As it is again expected to be a polynomial in t we get more obstructions on \mathcal{E} being very stable.

To illustrate the above take the example of $n = 2$ and $d = -2(g - 1)$ and let \mathcal{E} be a very stable Higgs bundle of type (2) , i.e. a very stable bundle E together with the zero Higgs field. For some $0 \leq i < g$ let $\mathcal{F} \in F_{l,2i}$ be a very stable type $(1, 1)$ Higgs bundle. Then $\Lambda_{\mathcal{F}}$ is a tensor product of universal bundles restricted to points of C . By carefully following the normalisation of the \mathbb{T} -action on the tensor product of these restricted universal bundles given by Lemma 6.27 we find that

$$m_{\mathcal{F},\mathcal{E}}(t) = \chi_{\mathbb{T}}(\Lambda_{\mathcal{F}}|_{\mathcal{E}}) = 2^{2i}t^i.$$

By (6.34) we get that

$$m_{\mathcal{E},\mathcal{F}}(t) = \frac{m_{\mathcal{E}}(t)m_{\mathcal{F},\mathcal{E}}(t)}{m_{\mathcal{F}}(t)} = \frac{(1+t)^{3g-3}2^{2i}t^i}{(1+t)^{2i}} = 2^{2i}t^i(1+t)^{3g-3-2i}. \quad (6.35)$$

Thus even though we do not know (see 8.3 for more on this) what vector bundle $\Lambda_{\mathcal{E}}$ the mirror of the cotangent fibre at E should be we can infer its \mathbb{T} -character at type $(1, 1)$ fixed points from (6.35).

7 A hyperholomorphic structure on \mathbb{F}_C

7.1 Hyperholomorphic connections

Thus far we have considered \mathcal{M} as a variety with a holomorphic symplectic structure in which the upward flows are complex Lagrangian subvarieties. At this point we bring

into play the hyperkähler metric on \mathcal{M} . This is a metric which is Kähler with respect to complex structures I, J, K which obey the quaternionic relations $I^2 = J^2 = K^2 = IJK = -1$. It exists as a consequence of the theorem [Hi1],[Si1],[Si1] that a stable Higgs bundle admits a Hermitian metric, unique up to gauge equivalence, satisfying the equation $F_A + [\Phi, \Phi^*] = 0$.

Within this context, mirror symmetry predicts that a BAA-brane should correspond to a BBB-brane on the mirror of \mathcal{M} , which is to be understood as the statement that to a flat vector bundle on a complex Lagrangian on \mathcal{M} there should exist a hyperholomorphic connection on a hyperkähler submanifold of the mirror. A hyperholomorphic connection is a unitary connection whose curvature is of type $(1, 1)$ with respect to each complex structure in the hyperkähler family of complex structures $x_1I + x_2J + x_3K$ with $(x_1, x_2, x_3) \in S^2 \subset \mathbb{R}^3$. At this point in time, we have few constructions of hyperholomorphic connections on \mathcal{M} so verifying the predictions is an important issue.

Since the fibration structure of \mathcal{M} gives a natural setting for the Strominger-Yau-Zaslow approach to mirror symmetry, it is the Fourier-Mukai transform approach which can be applied. In the original SYZ context of a special Lagrangian fibration Arinkin and Polishchuk [1] showed how a real Lagrangian can describe a holomorphic bundle on the mirror. This could be applied to the current situation but would give at most a hyperholomorphic connection relative to the semi-flat hyperkähler metric which is defined over the regular locus in \mathcal{A} and is only an approximation to the actual metric. We have described the Fourier-Mukai transform of the trivial bundle over an upward flow Lagrangian in terms of universal bundles over the mirror moduli space, so to verify the prediction we need to construct a hyperholomorphic connection, using the genuine hyperkähler metric, on a universal bundle. Twistor theory allows us to express this by holomorphic means using the direct analogy of the Atiyah-Ward description of instantons [AHS], [MS] – a hyperholomorphic connection on \mathbb{R}^4 is precisely an anti-self-dual connection.

Recall that the twistor space Z of a hyperkähler manifold M is the manifold $M \times S^2$ endowed with a complex structure such that the fibre over $(x_1, x_2, x_3) \in S^2$ is M with complex structure $x_1I + x_2J + x_3K$. The projection onto the second factor is a holomorphic fibration $p : Z \rightarrow \mathbb{P}^1$ with a real structure σ covering the antipodal map on S^2 and each 2-sphere (m, S^2) is a real holomorphic section, called a twistor line. Then a unitary hyperholomorphic connection on a vector bundle is equivalent to a holomorphic bundle on Z which is trivial on each twistor line and real with respect to σ .

A description of the twistor space for Higgs bundles due to Deligne and Simpson [Si3] is phrased in terms of λ -connections. A holomorphic λ -connection on a vector bundle E is a differential operator $D : E \rightarrow E \otimes K$ satisfying $D(fs) = fDs + \lambda s \otimes df$ for a local holomorphic section s and function f . If $\lambda \neq 0$ then $\lambda^{-1}D$ is a holomorphic, and therefore flat, connection and for $\lambda = 0$ this is a Higgs field. There is an algebraic

construction of a moduli space $\mathcal{M}_{\text{Hod}}^s$ of stable λ -connections and the parameter λ gives a projection to \mathbb{C} .

A solution to the Higgs bundle equations gives rise to a $\bar{\partial}$ -operator $\bar{\partial}_A + \zeta\Phi^*$ parametrized by $\zeta \in \mathbb{C}$ which commutes with $\zeta\partial_A + \Phi$, so defining a λ -connection with $\lambda = \zeta$ for the holomorphic structure on E defined by the $\bar{\partial}$ -operator. This is a holomorphic section of $\mathcal{M}_{\text{Hod}} \rightarrow \mathbb{C}$ and the twistor space is constructed so as to extend this to $\zeta = \infty \in \mathbb{P}^1$ and make this a twistor line.

When $\zeta \neq 0$ we have the flat connection

$$\nabla_A + \zeta^{-1}\Phi + \zeta\Phi^*. \quad (7.1)$$

Now, as ζ tends to ∞ , $\partial_A + \zeta^{-1}\Phi$ defines a holomorphic structure with respect to the conjugate complex structure on C and $\zeta^{-1}\bar{\partial}_A + \Phi^*$ a λ -connection for $\lambda = \zeta^{-1}$. The twistor space \mathcal{Z} is defined by taking the moduli space of λ -connections and the corresponding one for the conjugate complex structure on C , and identifying with the moduli space of flat connections over $\zeta \in \mathbb{C}^* \subset \mathbb{P}^1$, identifying $\lambda = \zeta$ on one side with $\lambda = \zeta^{-1}$ on the other. Then (7.1) defines a section of the fibration $p : \mathcal{Z} \rightarrow \mathbb{P}^1$.

We may also reinterpret the flat connection as an operator

$$\zeta\nabla_A + \Phi + \zeta^2\Phi^*$$

with coefficients in holomorphic sections of $\mathcal{O}(2)$ and observe from this that the antilinear involution $A \mapsto -A^*$ on $\text{End}(E)$ coupled with the antipodal map $\zeta \mapsto -1/\bar{\zeta}$ acting on $H^0(\mathbb{P}^1, \mathcal{O}(2))$ transforms this expression to itself, so a solution to the Higgs bundle equations defines a real section, and this corresponds to a point of \mathcal{M} .

7.2 The connection on a universal bundle

As described above, the Fourier-Mukai transform of the upward flow of a very stable fixed point of type $(1, 1, \dots, 1)$ can be expressed as a tensor product of exterior powers of the universal bundle corresponding to points $c \in C$. A hyperholomorphic connection on the universal bundle will then induce one on the associated tensor or exterior powers.

In the complex structure $\zeta \neq 0, \infty$ there is a straightforward description of the universal bundle: given a framing of a bundle E at c , the holonomy of a flat connection on E around closed curves with base point c defines a homomorphism from $\pi_1(C, c)$ to $\text{GL}_n(\mathbb{C})$. We denote $\mathcal{R}_B = \text{Hom}(\pi_1(C, c) \rightarrow \text{GL}_n(\mathbb{C}))$, the open locus of irreducible representations \mathcal{R}_B^s . We have the action of $\text{GL}_n(\mathbb{C})$ on \mathcal{R}_B by conjugation, and denote the affine quotient by \mathcal{M}_B , the character variety. Over the open subset $\mathcal{M}_B^s = \mathcal{R}_B^s/\text{GL}_n(\mathbb{C}) \subset \mathcal{M}_B$ the variety \mathcal{R}_B^s is a principal $\text{PGL}_n(\mathbb{C})$ -bundle, representing irreducible local systems framed at the point $c \in C$.

There is an obstruction in $H^2(\mathcal{M}_B^s, \mathbb{Z}_n)$ to lifting to a principal $\mathrm{GL}_n(\mathbb{C})$ -bundle. Given local liftings over open sets U_i , the obstruction class is represented by a flat line bundle L_{ij} over $U_i \cap U_j$ with L_{ij}^n trivial – the data for a gerbe α . If the universal vector bundle \mathbb{E} does not exist globally, it does over the neighbourhoods U_i , and a connection is a locally defined notion. Moreover tensoring with a flat line bundle L_{ij} acts trivially on the curvature and so takes a local hyperholomorphic connection to another one, so the notion of hyperholomorphic passes over to the situation where we need to twist by a flat gerbe.

For the complex structure $\zeta = 0$ or ∞ , the moduli space of framed Higgs bundles was constructed by Simpson in [Si1, Theorem 4.10], and in [Si3, Proposition 4.1] he outlined, by the same approach, a construction for λ -connections. We denote by $\mathcal{R}_{\mathrm{Hod}}^s$ the framed moduli space of stable λ -connections. This carries a $\mathrm{GL}_n(\mathbb{C})$ -action on the framing, so that it becomes a principal $\mathrm{PGL}_n(\mathbb{C})$ -bundle over $\mathcal{M}_{\mathrm{Hod}}^s$. Over $\lambda = 0$ this principal bundle $\mathcal{R}_{\mathrm{Hod}}^s \rightarrow \mathcal{M}_{\mathrm{Hod}}^s$ restricts to $\mathcal{R}_{\mathrm{Dol}}^s \rightarrow \mathcal{M}_{\mathrm{Dol}}^s = \mathcal{M}^s$ the framed moduli space of stable rank n degree 0 Higgs bundles, which is a $\mathrm{PGL}_n(\mathbb{C})$ -principal bundle over \mathcal{M}^s .

When $\lambda \neq 0$ we have the complex analytically equivalent alternative description $\mathcal{R}_B^s \rightarrow \mathcal{M}_B^s$ of the framed moduli space in terms of the character variety, using the holonomy of the flat connection. This is clearly compatible with the identification in the construction of the twistor space \mathcal{Z} , so we have a holomorphic $\mathrm{PGL}_n(\mathbb{C})$ -bundle over \mathcal{Z} . The final point is to show this is trivial on a real twistor line.

A real twistor line is defined by a solution to the Higgs bundle equations which, as ζ varies, is a pair $\bar{\partial}_A + \zeta\Phi^*$, $\zeta\partial_A + \Phi$ on the same C^∞ vector bundle E . A basis e_1, \dots, e_n of the fibre E_c then defines a framing for all holomorphic structures and we need this to be holomorphic in ζ . But from the integrability theorem of the holomorphic structure $\bar{\partial}_A + \zeta\Phi^*$ we can find a basis of local sections s_1, \dots, s_n in a neighbourhood of c with $(\bar{\partial}_A + \zeta\Phi^*)s_i = 0$ and these vary holomorphically with ζ . Then evaluation at c gives a frame which is holomorphic in ζ and similarly using the conjugate complex structure for $\zeta \neq 0$.

Remark 7.1. A formal explanation for the existence of the hyperholomorphic connection is provided by the infinite-dimensional hyperkähler quotient construction of \mathcal{M} itself [Hi1]. This considers the action of the group \mathcal{G} of unitary gauge transformations on the infinite-dimensional flat hyperkähler manifold which is the cotangent bundle of the space \mathfrak{A} of $\bar{\partial}$ -operators $\bar{\partial}_A$ on the underlying C^∞ vector bundle E . The zero set of the hyperkähler moment map for this action consists of pairs (A, Φ) satisfying the Higgs bundle equations. This space is formally a principal $\mathcal{G}/U(1)$ bundle over the quotient \mathcal{M} and the restriction of the flat L^2 metric on $T^*\mathfrak{A}$ defines a distribution orthogonal to the orbit directions, and this is a connection on the principal bundle. In any hyperkähler quotient this is hyperholomorphic. Now given a framing at c , the evaluation map gives

a homomorphism from $\mathcal{G}/U(1)$ to $PU(n)$ and an associated hyperholomorphic connection on the universal bundle.

Thus we have proved the following.

Theorem 7.2. *There is a hyperholomorphic connection on the $PGL_n(\mathbb{C})$ bundle $\mathcal{R}_{\text{Dol}}^s$ on \mathcal{M}^s , which is associated to the projective bundle $\mathbb{P}(\mathbb{E}_c)$. Moreover, there is a hyperholomorphic structure on the B -twisted vector bundle \mathbb{E}_c on \mathcal{M}^s .*

7.3 The universal Higgs field

For each point $c \in C$ we obtain the universal bundle \mathbb{E}_c with a hyperholomorphic connection on \mathcal{M} , and so C itself is the parameter space for a family of such bundles with connection. Equivalently, these correspond to holomorphic bundles on the twistor space \mathcal{Z} . In particular, considering first order deformations, there is a natural map from the tangent space of C at c to $H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c))$. This space can be interpreted in terms of variations of hyperholomorphic connections by using the Penrose transform in the current quaternionic context [MS]. It is isomorphic to the first cohomology group of an elliptic complex generalizing the deformation complex for anti-self-dual connections in four dimensions in [AHS].

Since \mathcal{M} is the moduli space of pairs (E, Φ) the universal bundle should have a universal Higgs field to accompany it and indeed a global construction of the universal pair can be found in [Ha2] for the $(d, n) = 1$ case and Theorem 4.5.(4) for a local construction in general. Given the role of the universal bundle, the reader may wonder how the universal Higgs field at c enters the picture. We shall see here that it defines the infinitesimal variation of the hyperholomorphic connection on the universal bundle.

The starting point is the observation that the space $\text{Hom}(\pi_1(C, c), PGL_n(\mathbb{C}))$ is independent of c . More accurately, choosing a path from c to y , there is a biholomorphism between the two spaces which only depends on the homotopy class of the path. In our situation it is the holonomy of the flat connection along that path which identifies the two. We have a flat connection when $\zeta \neq 0$ or ∞ and this suggests that the deformation class in $H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c))$ is supported on the fibres $Z_0, Z_\infty \subset \mathcal{Z}$ over these two points in \mathbb{P}^1 .

Given a tangent vector $X \in T_c(C)$, we can contract with the Higgs field $\Phi_c \in (\text{End}(E) \otimes K)_c$ to obtain $\Phi(X)$ which is a section of the universal bundle $\text{End}(\mathbb{E}_c)$ over the moduli space of Higgs bundles \mathcal{M} which we now think of as the fibre Z_0 over $\zeta = 0$ in the twistor space. Similarly $-\Phi^*$ defines a holomorphic section $-\Phi^*(X)$ over Z_∞ , the moduli space for the conjugate complex structure. Let s be the pullback to $\mathcal{Z} \rightarrow \mathbb{P}^1$ of the holomorphic section $\zeta d/d\zeta$ of $T\mathbb{P}^1 \cong \mathcal{O}(2)$ on \mathbb{P}^1 , then we have an exact sequence of

sheaves on \mathcal{Z} :

$$0 \rightarrow \mathcal{O}(\text{End}(\mathbb{E}_c)) \xrightarrow{s} \mathcal{O}(\text{End}(\mathbb{E}_c)(2)) \rightarrow \mathcal{O}_{Z_0+Z_\infty}(\text{End}(\mathbb{E}_c)(2)) \rightarrow 0.$$

The vector field $d/d\zeta$ trivializes $\mathcal{O}(2)$ at $\zeta = 0$ and similarly at infinity so the pair $(\Phi(X), -\Phi^*(X))$ may be regarded as a section of $\text{End}(\mathbb{E}_c)(2)$ over $Z_0 + Z_\infty$. Then we have:

Proposition 7.3. *The map $TC_c \rightarrow H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c))$ defining the first order deformation of the universal bundle at c is $\delta(\Phi(X), -\Phi^*(X))$ where Φ is the universal Higgs field at c and $\delta : H^0(Z_0 + Z_\infty, \text{End}(\mathbb{E}_c)(2)) \rightarrow H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c))$ is the connecting homomorphism in the cohomology of the above exact sequence.*

Proof. For $y \in C$ in a neighbourhood of c , parallel translation of a flat connection takes the principal bundle $\text{Hom}(\pi_1(C, c), \text{PGL}_n(\mathbb{C}))$ to $\text{Hom}(\pi_1(C, y), \text{PGL}_n(\mathbb{C}))$. This is an isomorphism of the frame bundle of the universal bundle at c to the universal bundle at y , and to first order at c it is the horizontal lift of a tangent vector X to the universal bundle over $\mathcal{M} \times C$, the lift defined by the connection. This is a trivial deformation of holomorphic principal bundles where \mathcal{M} has the complex structure of the moduli space of flat connections, so since $\zeta \neq 0$ describes the points of \mathcal{Z} via flat connections, it gives a trivialization of the deformation of the universal bundle on this part of \mathcal{Z} . The flat connection is $\partial_A + \Phi/\zeta$ and so the trivialization does not extend to $\zeta = 0$.

We may consider the first order deformation of the universal bundle as the Kodaira-Spencer class for the deformation of complex structure of the principal bundle $\mathcal{P}_c \rightarrow \mathcal{Z}$ as $c \in C$ varies. In Čech cohomology, the deformation class in $H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c)) \subset H^1(\mathcal{P}_c, T)$ is given using an open covering $\{U_i\}$, taking local lifts X_i of the tangent vector X and defining the cocycle $X_j - X_i$, a section of $\text{End}(\mathbb{E}_c)$ over $U_i \cap U_j$. In our case, for $\zeta \neq 0$, there is a global lift given by the horizontal space of the connection $\partial_A + \Phi/\zeta$. If we multiply by ζ then this is global for all ζ . Repeating this process near $\zeta = \infty$ shows that the class maps to zero in $H^1(\mathcal{Z}, (\text{End}(\mathbb{E}_c)) \xrightarrow{s} H^1(\mathcal{Z}, \text{End}(\mathbb{E}_c)(2))$ and consequently is in the image of the coboundary map $\delta : H^0(Z_0 + Z_\infty, (\text{End}(\mathbb{E}_c)(2)) \rightarrow H^1(\mathcal{Z}, (\text{End}(\mathbb{E}_c))$.

But the polar part of $\partial_A + \Phi/\zeta$ at $\zeta = 0$ is $\Phi(X)$ and we see that the class is given by extending $\Phi(X)$ and $-\Phi^*(X)$ as sections of $\text{End}(\mathbb{E}_c)(2)$ and dividing by s , which is the coboundary map in Čech cohomology. \square

8 Further issues

8.1 Simple Lie groups

The simplest example of a very stable upward flow of type $(1, 1, \dots, 1)$ is, as we have seen, the canonical section of the Hitchin fibration. In fact the moduli space of Higgs bundles

for any simple Lie group G admits a corresponding section, constructed in [Hi3]. For the adjoint group this is unique. This points towards a generalization of the $(1, 1, \dots, 1)$ type in this context, where the mirror should be a hyperholomorphic bundle on the moduli space for the Langlands dual group G^\vee .

The key point in [Hi3] is to use the principal three-dimensional subgroup of G [Ko]. The fixed point \mathcal{E} is then obtained from the canonical uniformising SL_2 Higgs bundle via the homomorphism $SL_2 \rightarrow G$. At a fixed point the scalar action on the Higgs field arises from the action of a subgroup of the adjoint group of G and in this case it is the action of the image of $\mathbb{C}^\times \subset SL_2$.

The characteristic feature of the subgroup is that it breaks up the Lie algebra \mathfrak{g} into ℓ irreducible components where ℓ is the rank of G and the dimensions are $2d_i - 1$ where the d_i are the degrees of a basis of invariant polynomials. These are odd-dimensional and hence representations of SO_3 . Therefore $t \in \mathbb{T}$ acts through $(t^{-1}, 1, t) \subset SO_3$.

The Higgs field has homogeneity one with respect to the \mathbb{T} . Each irreducible representation of SO_3 has a single weight one subspace so the Higgs field must take values in an ℓ -dimensional subspace of \mathfrak{g} . From [Ko, Lemma 5.2], the action has weight 1 on the root spaces of the ℓ simple roots $\alpha_1, \dots, \alpha_\ell$ which therefore form a basis for this subspace. In the case of the general linear group treated in this paper the simple roots are $x_{i+1} - x_i$ which correspond to line bundles $L_{i+1}L_i^*$ and we dealt with sections b_i of $L_{i+1}L_i^*K$. The direct analogue for the group G of the $(1, 1, \dots, 1)$ type fixed point is then a line bundle L_{α_i} for each simple root and a section b_i of $L_{\alpha_i}K$. Writing any root α in terms of simple roots then defines by tensor product a line bundle L_α and a Lie algebra bundle

$$H \oplus \bigoplus_{\alpha \in \Delta} L_\alpha$$

where H is a trivial rank ℓ bundle associated to the Cartan subalgebra. For the canonical fixed point \mathcal{E}_0 in the moduli space of G -bundles each b_i is nonvanishing and the upward flow is a section of the Hitchin fibration, and in particular is closed. We shall discuss next the case of a Higgs bundle \mathcal{E} when b_i is of degree $m_i \geq 0$ by studying the virtual multiplicity

$$m_{\mathcal{E}}(t) = \frac{\chi_{\mathbb{T}}(\text{Sym}(T_{\mathcal{E}}^{+*}))}{\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))}.$$

Here $T_{\mathcal{E}}^+$ is the sum of root space line bundles for roots of positive weight under the \mathbb{C}^* -action. Since the weight is 1 on the simple roots, these are the positive roots. If $\alpha = c_1\alpha_1 + \dots + c_\ell\alpha_\ell$ in terms of the simple roots then $w(\alpha) = c_1 + \dots + c_\ell$. Since $m_i = \deg L_{\alpha_i}K$ we have

$$\deg L_\alpha = \sum_{i=1}^{\ell} c_i(\alpha)(m_i - (2g - 2)).$$

Following the procedure of Remark 5.12 we obtain

$$\chi_{\mathbb{T}}(T_{\mathcal{E}}^+) = \sum_{\alpha > 0} t^{-w(\alpha)} \left(\sum_{w(\alpha)=k} -(\deg L_{\alpha} + (1 - g)) + \sum_{w(\alpha)=k-1} \deg(L_{\alpha} + (g - 1)) \right)$$

or

$$\chi_{\mathbb{T}}(T_{\mathcal{E}}^+) = \sum_{\alpha > 0} t^{-w(\alpha)} \left(\sum_{w(\alpha)=k} \sum_{i=1}^{\ell} -m_i c_i(\alpha) + \sum_{w(\alpha)=k-1} \sum_{i=1}^{\ell} m_i c_i(\alpha) \right) + (g - 1)N(t).$$

When all the $m_i = 0$, the multiplicity is 1 because then we have an equivariant section of $\mathcal{A} \rightarrow \mathcal{M}$ (one may also easily do the calculation – see below the generating function for N_j) so the $(g - 1)N(t)$ term above cancels the denominator $\chi_{\mathbb{T}}(\text{Sym}(\mathcal{A}^*))$ and the multiplicity reduces to

$$m_{\mathcal{E}}(t) = \prod_{i=1}^{\ell} \prod_{\alpha > 0} \left(\frac{1 - t^{w(\alpha)+1}}{1 - t^{w(\alpha)}} \right)^{m_i c_i(\alpha)}. \quad (8.1)$$

By experimentation one finds that this is rarely a polynomial. In the case of G_2 this happens independently of the m_i , supposing they are not identically zero, for other groups there are constraints on the m_i . So there is a distinct difference from the case of GL_n or SL_n . One case however always gives a polynomial as we see next.

A simple root α_i is called *cominuscule* if the coefficient $c_i(\alpha) = 0, \pm 1$ for every root α . Every simple root of SL_n has this property but for types B_n, C_n, D_n, E_6, E_7 they are special. For other groups, like G_2 , there are none. The coroot α_i^{\vee} in the dual root system, the root system of the Langlands dual group, is called *minuscule* (see for example [BS]).

Suppose then that α_i is cominuscule and the Higgs field has $m_j = 0$ for $j \neq i$ and $m_i = 1$. The formula (8.1) is multiplicative in m_i so this case is sufficient. Since $c_i(\alpha) = 0, 1$ for a positive root, the multiplicity is

$$\prod_{\alpha > 0, c_i(\alpha) \neq 0} \left(\frac{1 - t^{w(\alpha)+1}}{1 - t^{w(\alpha)}} \right).$$

Consider this product over all the positive roots of G :

$$\prod_{\alpha > 0} \left(\frac{1 - t^{w(\alpha)+1}}{1 - t^{w(\alpha)}} \right).$$

Let N_j be the number of root spaces of weight j , then this expression is

$$q_G(t) = \prod_{j \geq 1} \left(\frac{1 - t^{j+1}}{1 - t^j} \right)^{N_j} = \prod_{j \geq 1} (1 - t^j)^{N_{j-1} - N_j}.$$

The Lie algebra \mathfrak{g} splits under the SO_3 -action into ℓ irreducibles of dimension $2d_i - 1$ where d_i is the degree of a basis element of the ring of invariant polynomials. In each irreducible of dimension $2d_i - 1$, there is a one-dimensional space of weight $1, 2, \dots, d_i - 1$. Then

$$N_1x + N_2x^2 + \dots = \sum_{i=1}^{\ell} (x + x^2 + \dots + x^{d_i-1}) = \frac{x}{(1-x)} \sum_{i=1}^{\ell} (1 - x^{d_i-1}).$$

So the generating function for $N_{j-1} - N_j$ is $-\ell x + \sum_1^{\ell} x^{d_i}$, hence

$$q_G(t) = (1-t)^{-\ell} \prod_{j=1}^{\ell} (1-t^{d_j}). \quad (8.2)$$

Associated to each cominuscle root is a maximal parabolic subgroup P whose Levi subgroup L is obtained by deleting α_i from the Dynkin diagram of G to produce a semisimple group and has an additional \mathbb{C}^\times factor corresponding to the coroot α_i^\vee (see [BS] for example). All simple roots of type A_ℓ are minuscule and we have dealt with those in the main body of the paper. For the other groups the deleted Dynkin diagram is connected and so we get a simple group. The roots for which $c_i(\alpha) = 0$ are the roots of L . The \mathbb{T} -action has weight one on the simple root spaces of G and hence also on those of L . This means that \mathbb{T} acts with weight one on a linear combination $\sum_{j \neq i} \lambda_j e_{\alpha_j}$ of root vectors. From [Ko] if $\lambda_j \neq 0$, this is a principal nilpotent in the Levi subalgebra and so \mathbb{T} acts as a semisimple element in a principal three-dimensional subgroup of the Levi group L .

Let n_j be the degrees of generators of the invariant polynomials on L . The rank of L is ℓ and the simple component $\ell - 1$. Then

$$q_L(t) = (1-t)^{-\ell} \prod_{j=1}^{\ell-1} (1-t^{n_j})$$

and

$$m_{\mathcal{E}}(t) = \prod_{\alpha > 0, c_i(\alpha) \neq 0} \left(\frac{1-t^{w(\alpha)+1}}{1-t^{w(\alpha)}} \right) = \frac{q_G(t)}{q_L(t)} = \prod_{j=1}^{\ell} \frac{(1-t^{d_j})}{(1-t^{n_j})}. \quad (8.3)$$

Proposition 8.1. *The virtual multiplicity $m_{\mathcal{E}}(t)$ in equation (8.3) is a polynomial.*

Proof. From [Bor, Theorem 26.1] if K, U are compact Lie groups of the same rank then the Poincaré polynomial of K/U is

$$P(K/U, t) = \frac{(1-t^{s_1})(1-t^{s_2}) \dots (1-t^{s_\ell})}{(1-t^{r_1})(1-t^{r_2}) \dots (1-t^{r_\ell})}.$$

where $s_i - 1, r_i - 1$ are the degrees of generators of the cohomology of K, U . Taking K and U to be the maximal compact subgroups of P, L then $s_j = 2d_j, r_j = 2n_j$ and $P(K/U, t)$ is a polynomial $Q(t^2)$. Then $m_{\mathcal{E}}(t) = Q(t)$. \square

A theorem (which also follows from the geometric Satake theorem of [MV], c.f. [Gi, Theorem 1.6.3]) of B.Gross [Gro, Corollary 6.4] states that the Lefschetz action of SL_2 on the cohomology of the flag variety G/P is isomorphic to the representation of the principal SL_2 of the Langlands dual G^\vee on the representation given by the minuscule weight dual to α_i . In the light of our results on exterior powers for the linear group this suggests the following.

Conjecture 8.2. *Let $\alpha_{i_1}, \dots, \alpha_{i_k}$ be distinct cominuscule roots and suppose that the corresponding sections b_{i_j} of $L_{\alpha_{i_j}} K$ have m distinct zeros x_1, \dots, x_m , and for other simple roots b_i has no zero. Then the fixed point is very stable and the mirror of its upward flow is the tensor product of the vector bundles associated to the universal principal bundle for the Langlands dual group at x_j in the corresponding minuscule representation.*

8.2 Multiple zeros

We showed in Theorem 1.2 that a necessary and sufficient condition for a fixed point of type $(1, 1, \dots, 1)$ to be very stable is that b should have no multiple zeros. It is natural to ask what happens when we do have a multiple zero. What is the closure of the upward flow, and is there a corresponding hyperholomorphic bundle on the mirror moduli space?

We already know from Remark 5.15 that the intersection of the upward flow with a generic fibre has fewer points than the situation where b has simple zeros. We also see in the proof of Lemma 4.13 that a multiple zero leads to a Hecke curve in the closure – the closure of the flow tangent to a nilpotent Higgs field.

We illustrate this here with the simplest example of this situation in rank 2, for the group SL_2 . Let \mathcal{E} be a fixed point of the \mathbb{T} -action defined by the vector bundle $L \oplus L^*$ and the off-diagonal Higgs field $b : L \rightarrow L^* K$, defining a section of $L^{-2} K$ which we assume has a single double zero at $c \in C$, and hence $L \cong K^{1/2}(-c)$. The upward flow is given by Higgs bundles (E, Φ) where E is an extension $L \rightarrow E \xrightarrow{p} L^*$, $\text{tr} \Phi = 0$ and $b = p\Phi : L \rightarrow L^* K$.

The spectral curve C_a is defined by $\det(x - \Phi) = x^2 + \det \Phi = 0$ and $\sigma(x) = -x$ is the covering involution. Then if C_a is smooth, as we have seen, $E = \pi_* U$ where the line bundle U is the cokernel of $x - \Phi : \pi^*(E \otimes K^*) \rightarrow \pi^* E$. The inclusion $L \subset E \in H^0(C, \text{Hom}(L, E))$ is canonically defined by the direct image construction as a section of the line bundle $\text{Hom}(\pi^* L, U)$ on C_a .

Now b vanishes at $c \in C$ and suppose further that $\det \Phi(c) \neq 0$. Since L is preserved by Φ at c , it maps to zero in the cokernel of $\Phi - x$ for one of the two eigenvalues $\pm x$. Then $p = (x, c) \in C_a$ or $\sigma(p) = (-x, c)$ are zeros of the holomorphic section of $(\pi^*L)^*U$. Since $\pi : C_a \rightarrow C$ is unramified at these points, these zeros have multiplicity 2 since b has a double zero at c .

The bundle $\Lambda^2 E$ is trivial which implies $U \cong M\pi^*K^{1/2}$ where $\sigma^*M \cong M^*$ (i.e. M lies in the Prym variety) and so since $L \cong K^{1/2}(-c)$, $\pi^*L^*U \cong M(c)$ which has degree 2. But we have a section which vanishes with multiplicity 2 hence $M(c) \sim 2p$ or $2\sigma(p)$ as divisors, or, since $c \sim p + \sigma(p)$, $M \sim \pm(p - \sigma(p))$. Thus the intersection with the generic fibre consists of two points in the Prym variety whereas if b had two distinct zeros the intersection would be four points.

Note that if $q(c) = 0$ and C_a is smooth, there is a single point p with $\pi(p) = c$ and π locally has the form $z \mapsto z^2$. Since b has a double zero at c , the section of π^*L^*U has a simple zero which is a contradiction since it has degree 2. Thus the upward flow does not intersect these fibres, but as we shall see next, the closure does.

Suppose (E, Φ) is a limit point of the upward flow. There is still a non-zero homomorphism $K^{1/2}(-c) \rightarrow E$ but its image may not be an embedding. It will generate a subbundle L_0 of larger degree. Since it has positive degree by stability it cannot be invariant by the Higgs field Φ_0 so there is a nonvanishing section of $L_0^{-2}K$. But $\deg L_0 > \deg K^{1/2}(-c) = g - 2$ hence $\deg L_0^{-2}K \leq 0$. The existence of a section means that we have equality and $L_0 = K^{1/2}$, so the bundle is an extension $K^{1/2} \rightarrow E \rightarrow K^{-1/2}$ and b is an isomorphism. But from [Hi1, Proposition 3.3] the extension is trivial and the limit lies on the canonical section. Since $\det \Phi \in H^0(C, K^2)$ parametrizes the section the limit lies on the $(3g - 4)$ -dimensional linear subspace consisting of sections which vanish at c .

Consider now the uniformising Higgs bundle $K^{1/2} \oplus K^{-1/2}$ with Higgs field $1 : K^{1/2} \rightarrow K^{-1/2}K$. The *downward* flow from this point consists of extensions $K^{-1/2} \rightarrow E \rightarrow K^{1/2}$ with the same nilpotent Higgs field mapping the quotient to the subbundle. This is a Lagrangian submanifold and the extensions are parametrized by a class $e \in H^1(C, K^*)$ (of dimension $(3g - 3) = \dim \mathcal{M}/2$). The section s of $\mathcal{O}(c)$ vanishing at c defines a homomorphism $s : K^{1/2}(-c) \rightarrow K^{1/2}$ which lifts to E if $se = 0 \in H^1(C, K^*)$. Equivalently $e = \delta(u)$ for $u \in H^0(c, \mathcal{O}_c(K^*)) \cong \mathbb{C}$ where δ is the connecting homomorphism in the cohomology exact sequence.

In Dolbeault terms with a C^∞ splitting $K^{1/2} \oplus K^{-1/2}$ we have

$$\bar{\partial} + \begin{pmatrix} 0 & \bar{\partial}v/s \\ 0 & 0 \end{pmatrix} \quad \Phi = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$$

where (as in 4.1.4) v is a local extension of u , zero outside a neighbourhood and holomorphic in a smaller one. Then the embedding of $K^{1/2}(-c) \subset E$ is $(v, -s)$ so that

$\Phi(-v, s) = (s, 0)$ and the skew form on E gives $\langle (v, -s), (s, 0) \rangle = s^2$. This is equivalent to projecting onto the quotient and clearly has a double zero at c . Then each extension E lies on the upward flow from \mathcal{E} and has as limit the canonical fixed point. This is a projective line in the closure, a Hecke curve.

We see therefore that the union of the upward flow W_c^+ from c and W_0^+ from the canonical point is a closed subspace, and the symplectic form vanishes at smooth points.

Conjecture 8.3. *The mirror of the closed Lagrangian cycle $W_0^+ \cup W_c^+$ is the universal adjoint bundle for c in the SO_3 moduli space.*

Here is some motivation. The subvariety $W_0^+ \cup W_c^+$ intersects a generic fibre in three points and so the Fourier-Mukai transform is generically a rank 3 vector bundle. The virtual multiplicity $m_{\mathcal{E}}(t)$ for a double zero of b is the same as for two distinct zeros and, from the point of view of the previous subsection this is $(1+t)^2$, from the Poincaré polynomial of $\mathbb{P}^1 \times \mathbb{P}^1$. Moreover the Lefschetz action of SL_2 on the cohomology is the action on $V \otimes V$ where V is the defining 2-dimensional representation. This is the direct sum of the adjoint and the trivial one-dimensional representation so on the adjoint component the action is consistent with the conjecture.

Now the adjoint bundle for SO_3 is a rank 3 bundle V and $\Phi : K^* \rightarrow V$ is injective when Φ is regular at each point of C , i.e. when the spectral curve is smooth. Trivializing K at c , the universal bundle admits a homomorphism from the trivial bundle. It follows that the canonical section, which corresponds to the trivial line bundle on the spectral curve, forms a component of the support of the inverse transform. The other component clearly relates to the quotient sheaf which is generically a rank 2 bundle.

One point to note is that the hyperholomorphic connection on the universal bundle is not reducible (its Pontryagin class is a generator of $H^4(\mathcal{M}, \mathbb{R})$) and this casts a question mark over the issue of whether the mirror of the closure of W_c^+ actually has a hyperholomorphic connection.

More generally, there is a Hecke correspondence for every irreducible representation of the Langlands dual group, which has a natural compactification (see [KW, §9]). In our case, the image of the Hitchin section in the SL_2 Higgs moduli space under this compactified Hecke correspondence, attached to the adjoint representation of SO_3 , is precisely our Lagrangian cycle $W_0^+ \cup W_c^+$. Its mirror then should be the universal adjoint bundle at c . The \mathbb{T} -equivariant index of its structure sheaf then should be $1+t+t^2$, which is also the \mathbb{T} -character of the mirror adjoint bundle at the canonical uniformising Higgs bundle. Incidentally, it also matches the intersection Poincaré polynomial of a certain weighted projective space, which is the closure of the space of Hecke modifications at a point in our case (c.f. [KW, 9.2]). In fact, from the geometric Satake correspondence [MV, Gi] this generalises to the general case of simple (even reductive) groups, and we expect the analogous results to hold there too.

8.3 The cotangent fibre

The upward flow from a very stable bundle $(E, 0)$ is an important example. We have calculated the rank of its transform in Remark 5.9 using the multiplicity formula. The alternative method, as in [BNR] is to argue that if there are no nilpotent Higgs fields then locally the moduli space is given by the vanishing of invariant polynomials. For a simple Lie group this is therefore the vanishing of $(2d_i - 1)(g - 1)$ functions of degree d_i for $i = 1, \dots, \ell$ and so the rank of this bundle is

$$\prod_1^\ell d_i^{(2d_i - 1)(g - 1)},$$

which Laumon in [La2, Remark 6.2.3] called the global analogue of the order of the Weyl group. The structure of this bundle even for the 2^{3g-3} case of SL_2 is largely unknown, and less so a way to construct a hyperholomorphic connection on it. In (6.35) we offered evidence from mirror symmetry for a conjecture of the \mathbb{T} -character of the fibre of this bundle with a \mathbb{T} -equivariant structure at the fixed points of type $(1, 1)$.

It is an important example as it relates to the work of Donagi–Pantev [DP2] on the geometric Langlands programme, where they proposed a geometric approach of constructing automorphic sheaves on the moduli stack of bundles, which was, in fact, also the motivation of Drinfeld [Dr1] and Laumon [La1] for the introduction of very stable bundles, see also [La2, Conjecture 6.2.1]. This programme aims to associate to a point $m \in \mathcal{M}^\vee$, the mirror of the Higgs bundle moduli space \mathcal{M} , a flat connection on \mathcal{N} , the moduli space of stable bundles, with singularities on the “wobbly locus”. A generic point in the mirror consists of a spectral curve C_a and (according to the SYZ procedure) a line bundle L over its Jacobian. In \mathcal{M} , the spectral curve defines a fibre of the Hitchin fibration. This fibre over $a \in \mathcal{A}$ intersects $T^*\mathcal{N} \subset \mathcal{M}$ in the complement of a subvariety and the projection to \mathcal{N} extends to a rational map as in [BNR]. The idea is that the vector bundle V on \mathcal{N} is associated to a direct image of L , and the canonical 1-form on the cotangent bundle $T^*\mathcal{N}$ defines a Higgs field on V , a section of $\text{End}(V) \otimes T^*$. Then, using the higher dimensional nonlinear Hodge theory due to Simpson and Mochizuki [Mo], one hopes to obtain a flat connection.

What is the link with this paper? The complement of the wobbly locus in \mathcal{N} is the set of very stable vector bundles E . The fibre of V at a generic $[E] \in \mathcal{N}$ is the direct sum of the fibres of L over the intersection of the upward flow of $(E, 0)$ with the Hitchin fibre. Put another way, if $\mathcal{N}^{vs} \subset \mathcal{N}$ denotes the moduli space of very stable bundles, then the Fourier-Mukai transform of the upward flows over \mathcal{N}^{vs} defines a bundle on $\mathcal{N}^{vs} \times \mathcal{M}^\vee$. In this paper we have been concerned with the restriction to $[E] \times \mathcal{M}^\vee$, for Donagi and Pantev it is the restriction to $\mathcal{N}^{vs} \times \{m\}$.

In fact what we have encountered is a miniature version of this in the case of very

stable Higgs bundles of type $(1, 1, \dots, 1)$. Instead of the \mathbb{T} -fixed point set \mathcal{N} we have another component: a subvariety defined by the zero set of (b_1, \dots, b_{n-1}) in the product of symmetric products $C^{[m_1]} \times C^{[m_2]} \times \dots \times C^{[m_{n-1}]}$. Moreover we have explicitly identified the wobbly locus, namely where the product $b_1 \dots b_{n-1}$ has multiple zeroes. This includes the various diagonals. The hyperholomorphic bundle we identified as a tensor product of exterior powers of universal bundles corresponding to the zeros of the b_i . By nonlinear Hodge theory on C , a point $m \in \mathcal{M}^\vee$ defines a flat connection on the curve C and hence also on the product $C^{m_1} \times C^{m_2} \times \dots \times C^{m_{n-1}}$, invariant by the action of the product of symmetric groups, hence descending to a flat connection, but with singularities over the diagonals.

References

- [1] ARINKIN, D. and POLISHCHUK, A.: Fukaya category and Fourier transform, (in *Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds* (Cambridge, MA, 1999)), AMS/IP Stud. Adv. Math., **23**, AMS, (2001), 261–274.
- [Ar] ARTIN, M.: On the joins of Hensel rings, *Advances in Math.* **7** (1971), 282–296.
- [As] ASOK, A.: Equivariant vector bundles on certain affine G-varieties, *Pure Appl. Math. Q.* **2** (2006), no. 4, Special Issue: In honor of Robert D. MacPherson. Part 2, 1085–1102.
- [AHS] ATIYAH, M.F., HITCHIN, N.J. and SINGER, I.M.: Self-duality in four-dimensional Riemannian geometry, *Proc. Roy. Soc. London Ser A* **362** (1978), 425–461.
- [BF] BAUM, P. , FULTON, W. and MACPHERSON, R.: Riemann-Roch and topological K-theory for singular varieties, *Acta Math.* Volume **143** (1979), 155–192.
- [BNR] BEAUVILLE, A., NARASIMHAN, M.S., and RAMANAN, S.: Spectral curves and the generalised theta divisor, *J. Reine Angew. Math.* **398** (1989), 169–179.
- [BDMN] BELLAMY, G, DODD, C., MCGERTY, K. and NEVINS, T.: Categorical cell decomposition of quantized symplectic algebraic varieties, *Geom. Topol.* **21** (2017), no. 5, 2601–2681
- [B-B] BIALYNICKI-BIRULA, A.: Some theorems on actions of algebraic groups, *Ann. of Math.* (2) **98** (1973), 480–497.
- [B-B2] BIALYNICKI-BIRULA, A: Some properties of the decompositions of algebraic varieties determined by actions of a torus, *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **24** (1976), no. 9, 667–674.

- [BR] BISWAS, I.; RAMANAN, S.: An infinitesimal study of the moduli of Hitchin pairs, *J. London Math. Soc. (2)* **49** (1994), no. 2, 219–231.
- [BL] BIRKENHAKE, CH. and LANGE, L.: *Complex Abelian Varieties* 2nd ed., Grundlehren Math. Wiss. **302**, Springer (2004)
- [Bor] BOREL, A., Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts, *Ann. of Math.* **57** (1953) 115–207.
- [Bou] BOURBAKI, N.: *Commutative algebra*, Chapters 1–7. Elements of Mathematics. Springer-Verlag, Berlin, 1989
- [Br] BRION, M.: Linearization of algebraic group actions, *Handbook of group actions*. Vol. IV, 291–340, Adv. Lect. Math. (ALM), **41**, Int. Press, Somerville, MA, 2018.
- [BS] BUCH, A. & SAMUEL, M.: K-theory of minuscule varieties, *J. Reine Angew. Math.* **719** (2016) 133–171.
- [Ca] CALDARARU, A.: Derived categories of twisted sheaves on Calabi-Yau manifolds, Thesis (Ph.D.) Cornell University 2000 196 pp.
- [CW] COLLIER, B. and WENTWORTH, R. : Conformal limits and the Bialynick-Birula stratification of the space of lambda-connections, *Adv. Math.* **350** (2019), 1193–1225. [arXiv:1808.01622](https://arxiv.org/abs/1808.01622)
- [Do] DOLGACHEV, I.: *Lectures on invariant theory*, Springer 2004
- [DP1] DONAGI, R. and PANTEV, T.: Langlands duality for Hitchin systems, *Invent. Math.* **189** 3, 653–735. [arXiv:math/0604617](https://arxiv.org/abs/math/0604617)
- [DP2] DONAGI, R. and PANTEV, T.: Geometric Langlands and non-abelian Hodge theory, 85–116, *Surv. Differ. Geom.*, **13**, Int. Press, Somerville, MA, 2009.
- [Dr1] DRINFELD, V.G.: Letter to P. Deligne June 1981
- [Dr3] DRINFELD, V.: On algebraic spaces with an action of \mathbb{G}_m , preprint [arXiv:1308.2604](https://arxiv.org/abs/1308.2604)
- [GHS] GARCÍA-PRADA, O.; HEINLOTH, J. and SCHMITT, A.: On the motives of moduli of chains and Higgs bundles, *J. Eur. Math. Soc.* **16** (2014), no. 12, 2617–2668.
- [GS] GILLET, H. and SOULÉ, C.: Intersection theory using Adams operations, *Invent. math.* **90** (1987) 243–277
- [Gi] GINZBURG, V.: Perverse sheaves on a loop group and Langlands’ duality. preprint [arXiv:alg-geom/9511007](https://arxiv.org/abs/alg-geom/9511007)

- [Go] GOTHEN, P.B., The Betti numbers of the moduli space of rank 3 Higgs bundles, *Internat. J. Math.* **5** (1994) 861-875
- [Gro] GROSS B.: On minuscule representations and the principal SL_2 , *Representation Theory* **4** (2000) 225–244.
- [Gr] GROTHENDIECK, A.: Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné) : IV. Étude locale des schémas et des morphismes de schémas, Troisième partie . *Publications Mathématiques de l’IHÉS*, **28** (1966), 5-255
- [Hal] HALPERN-LEISTNER, D.: On the structure of instability in moduli theory, [arXiv:1411.0627](https://arxiv.org/abs/1411.0627)
- [Ha1] HAUSEL, T.: Compactification of moduli of Higgs bundles, *J. Reine Angew. Math.*, **503** (1998) 169-192.
- [Ha2] HAUSEL, T. Vanishing of intersection numbers on the moduli space of Higgs bundles. *Adv. Theor. Math. Phys.*, 2(5):1011–1040, 1998.
- [HMP] HAUSEL, T; MELLIT, A. and PEI, D.: Mirror symmetry with branes by equivariant Verlinde formulas. *Geometry and physics*, J.E.Andersen et al (eds) Vol. I, 189–218, Oxford Univ. Press, Oxford, 2018.
- [HT] HAUSEL, T. and THADDEUS, M.: Generators for the cohomology ring of the moduli space of rank 2 Higgs bundles, *Proc. London Math. Soc.* **88** (2004) 632-658.
- [HV] HAUSEL, T. and R. VILLEGAS, F. .: Cohomology of large semiprojective hyperkähler varieties, *Astérisque* No. **370** (2015), 113–156,
- [He] HEINLOTH, J.: Hilbert-Mumford stability on algebraic stacks and applications to G -bundles on curves, *Épjournal Geom. Algébrique* 1 (2017), Art. 11, 37 pp.
- [Hi1] HITCHIN, N.: The self-duality equations on a Riemann surface, *Proc. London Math. Soc.* (3) **55** (1987) 59–126.
- [Hi2] HITCHIN, N.: Stable bundles and integrable systems. *Duke Math. J.*, 54(1):91–114, 1987.
- [Hi3] HITCHIN, N. : Lie groups and Teichmüller space. *Topology* **31** (1992), no. 3, 449–473
- [Hi4] HITCHIN, N.: Higgs bundles and characteristic classes, in *Arbeitstagung Bonn 2013; In Memory of Friedrich Hirzebruch*, Progress in Mathematics, Birkhäuser, 2016, 247–264. [arXiv:1308.4603](https://arxiv.org/abs/1308.4603)

- [Hi5] HITCHIN, N.: Critical loci for Higgs bundles. *Comm. Math. Phys.* **366** (2019), no. 2, 841–864.
- [Hu] HUYBRECHTS, D.: *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2006.
- [KW] KAPUSTIN, K. and WITTEN, E.: Electric-magnetic duality and the geometric Langlands program, *Commun. Number Theory Phys.* **1** (2007), no. 1, 1–236.
- [Ko] KOSTANT, B.: The principal three-dimensional subgroup and the Betti numbers of a complex simple Lie group, *Amer.J.Math.* **81** (1959) 973–1032.
- [Mu] MUKAI, S.: Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves, *Nagoya Math. J.* **81** (1981), 153–175.
- [La1] LAUMON G.: Un analogue du cone nilpotent, *Duke Math. J.* (2) **57** (1988) 647-671
- [La2] LAUMON, G.: Faisceaux automorphes liés aux séries d’Eisenstein, in *Automorphic forms, Shimura varieties, and L-functions*, Vol. I (Ann Arbor, MI, 1988), 227–281, *Perspect. Math.*, 10, Academic Press, Boston, MA, 1990.
- [Lu] LUNA, D.: Slices étales, *Mémoires de la S. M. F.*, tome **33** (1973), 81–105
- [MMP] MASUDA, M.; MOSER-JAUSLIN, L. and PETRIE, T. The equivariant Serre problem for abelian groups, *Topology* **35** (1996), no. 2, 329–334.
- [MS] MAMONE-CAPRIA, M. and SALAMON, S.M.: Yang-Mills fields on quaternionic spaces, *Nonlinearity* **1** (1988) 517–530.
- [Ma] MATSUMURA, H.: *Commutative ring theory*, Cambridge Studies in Advanced Mathematics, 8 (2nd ed.), Cambridge University Press, (1989)
- [Mi] MILNE, J.: Étale cohomology. *Princeton Mathematical Series*, **33**. Princeton University Press, Princeton, N.J., 1980.
- [MV] MIRKOVIĆ, I. and Vilonen, K.: Geometric Langlands duality and representations of algebraic groups over commutative rings, *Annals of Mathematics*, Second Series, **166** (1): 95–143
- [Mo] MOCHIZUKI, T.: Kobayashi-Hitchin correspondence for tame harmonic bundles and an application, *Astérisque*, **309**: viii+117, 2006.
- [Na] NAKAJIMA, H.: *Lectures on Hilbert schemes of points on surfaces*, University Lecture Series, 18. American Mathematical Society, Providence, RI, 1999.

- [NR] NARASIMHAN, M.S. and RAMANAN S.: Geometry of Hecke cycles I, in “C.P.Ramanujam – a tribute” *Tata Inst. Fund. Res. Studies in Math.* **8**, Springer-Verlag, Berlin (1978) 291–345.
- [Ni] NITSURE, N., Moduli space of semistable pairs on a curve, *Proc. London Math. Soc.* (3) **62** (1991) 275–300.
- [PPN] PAULY, C. and PEÓN-NIETO A.: Very stable bundles and properness of the Hitchin map, [arXiv:1710.10152](https://arxiv.org/abs/1710.10152) *Geometriae Dedicata* **198**, (2019) (143-145)
- [PW] PIEPMAYER, G. and WALKER, M.E.: A new proof of the New Intersection Theorem, *Journal of Algebra* **322** (2009) 3366–3372
- [Se] SERRE, J-P. : *Local algebra*, Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2000.
- [Sh] SHAYMAN, M. A. : On the variety of invariant subspaces of a finite-dimensional linear operator. *Trans. Amer. Math. Soc.* **274** (1982), no. 2, 721–747
- [Si1] SIMPSON, C.: Moduli of representations of the fundamental group of a smooth projective variety. I, *Inst. Hautes Études Sci. Publ. Math.* No. **79** (1994), 47–129.
- [Si1] SIMPSON, C.: Moduli of representations of the fundamental group of a smooth projective variety. II, *Inst. Hautes Études Sci. Publ. Math.* No. **80** (1994), 5–79.
- [Si2] SIMPSON, C.: Iterated destabilizing modifications for vector bundles with connection, in *Vector bundles and complex geometry*, 183–206, *Contemp. Math.*, **522**, Amer. Math. Soc., Providence, RI, 2010.
- [Si3] SIMPSON C.: The Hodge filtration on nonabelian cohomology. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, 217–281. Amer. Math. Soc., Providence, RI, 1997.
- [Sm] SMALE, S.: Differentiable dynamical systems, *Bull. Amer. Math. Soc.* **73** (1967), 747–817.
- [Sta] THE STACKS PROJECT AUTHORS: *Stacks Project*, <https://stacks.math.columbia.edu>
- [Su] SUMIHIRO, H.: Equivariant completion, *J. Math. Kyoto Univ.* **14** (1974), no. 1, 1–28.
- [Vi] VILLAMAYOR, O. : Patching local uniformizations, *Annales scientifiques de l'É.N.S. 4e série*, **25**, no 6 (1992), 629–677