

A NOTE ON HESSENBERG VARIETIES

KARI VILONEN AND TING XUE

We give a short proof based on Lusztig’s generalized Springer correspondence of some results of [BrCh, BaCr, P].

Let \mathfrak{g} denote a complex semisimple Lie algebra and let us write $G = \text{Aut}(\mathfrak{g})^0$ for the identity component of its group of automorphisms. Note that the group G is adjoint. Let us write \mathcal{B} for the flag variety of \mathfrak{g} and let us fix a Borel subalgebra $\mathfrak{b} \subset \mathfrak{g}$. Let $B \subset G$ be the corresponding Borel subgroup. Consider a Hessenberg subspace $V \subset \mathfrak{g}$, i.e., a B -invariant subspace of \mathfrak{g} . It gives rise to a G -equivariant bundle

$$(1) \quad \mathcal{H} := G \times^B V \rightarrow \mathcal{B}.$$

We have a G -equivariant projective morphism

$$(2) \quad \pi : \mathcal{H} \rightarrow \mathfrak{g}, \quad (g, x) \mapsto \text{Ad}(g)x$$

the “universal” family of Hessenberg varieties. We can also consider a perpendicular Hessenberg subspace V^\perp and we similarly get

$$(3) \quad \check{\pi} : \check{\mathcal{H}} := G \times^B V^\perp \rightarrow \mathfrak{g}.$$

We now assume that $\mathfrak{b} \subset V$ so that $V^\perp \subset \mathfrak{n} = [\mathfrak{b}, \mathfrak{b}]$. By the decomposition theorem [BBD] we have

$$(4) \quad R\pi_* \mathbb{C}_{\mathcal{H}} = \bigoplus \text{IC}(\mathfrak{g}^{rs}, \mathcal{L}_i)[-] \oplus \text{terms with smaller support};$$

here the \mathcal{L}_i are irreducible G -equivariant local systems on the regular semisimple locus \mathfrak{g}^{rs} and $[-]$ denotes cohomological shifts. We also have

$$(5) \quad R\check{\pi}_* \mathbb{C}_{\check{\mathcal{H}}} = \bigoplus \text{IC}(\mathcal{O}, \mathcal{E})[-];$$

here the \mathcal{O} are nilpotent orbits of G and the \mathcal{E} are G -equivariant irreducible local systems on the \mathcal{O} . By considering the Fourier transform we conclude that all the IC’s appearing in (4) are Fourier transforms of the IC’s appearing in (5).

We have the following

Conjecture 1 (Patrick Brosnan [X]). *All the $\text{IC}(\mathcal{O}, \mathcal{E})$ in (5) appear in the Springer correspondence. In particular, all the terms in (4) have full support.*

This conjecture has been proved by Martha Precup and Eric Sommers (for type G_2 see also [X]). It is not difficult to prove the following weaker statement which suffices for some applications.

KV was supported in part by the ARC grants DP150103525 and DP180101445 and the Academy of Finland.

TX was supported in part by the ARC grant DP150103525.

Proposition 2. *We have*

$$(R\pi_*\mathbb{C}_{\mathcal{H}})|_{\mathfrak{g}^{reg}} = \bigoplus \mathrm{IC}(\mathfrak{g}^{rs}, \mathcal{L}_i)|_{\mathfrak{g}^{reg}}[-].$$

Proof. The terms in (4) which do not appear in the Springer correspondence come by induction from the (nontrivial) cuspidals in Lusztig's generalized Springer correspondence [L]. Recall that we are in the context of the adjoint group. One can now verify, case-by-case using [L], that the supports of such terms do not meet the regular locus \mathfrak{g}^{reg} . \square

The \mathcal{L}_i come from representations of the Weyl group W and we will now denote them as such by ρ_i . Let us consider $\mathrm{IC}(\mathfrak{g}^{rs}, \rho_i)$ restricted to the regular locus. It is a direct summand in $\tilde{p}_*\mathbb{C}_{\tilde{\mathfrak{g}}^{reg}}$ where $\tilde{p} : \tilde{\mathfrak{g}}^{reg} \rightarrow \mathfrak{g}^{reg}$ is the Grothendieck simultaneous resolution restricted to the regular locus. Let \mathfrak{t} be a Cartan subspace of \mathfrak{g} . Then we have the following Cartesian square

$$(6) \quad \begin{array}{ccc} \tilde{\mathfrak{g}}^{reg} & \xrightarrow{\tilde{p}} & \mathfrak{g}^{reg} \\ \downarrow & & \downarrow f \\ \mathfrak{t} & \xrightarrow[p]{} & \mathfrak{t}/W \end{array}$$

and so we have $\tilde{p}_*\mathbb{C}_{\tilde{\mathfrak{g}}^{reg}} = f^*p_*\mathbb{C}_{\mathfrak{t}}$. Thus, as f is smooth, it suffices to analyze $p_*\mathbb{C}_{\mathfrak{t}}$.

Let us now consider a stratum \mathcal{O}_s in \mathfrak{t}/W associated to a semisimple element $s \in \mathfrak{t}$. Note that the fundamental group of \mathcal{O}_s is the braid group of ${}^sW = N_G(L)/L$ where $L = Z_G(s)$. We also write $W_s = \mathrm{Stab}_W(s)$. By observing that $N_G(L)/L = (N_G(L) \cap N_G(T))/N_L(T)$ we have an exact sequence

$$(7) \quad 1 \rightarrow W_s = N_L(T)/T \rightarrow (N_G(L) \cap N_G(T))/T \rightarrow {}^sW = N_G(L)/L \rightarrow 1.$$

The fiber of p over this stratum is W/W_s . This implies:

$$(8) \quad \mathrm{IC}(\mathfrak{t}/W, \rho_i)|_{\mathcal{O}_s} = \rho_i^{W_s}.$$

The righthand side can be viewed as a representation of sW by (7) and hence as a local system on \mathcal{O}_s . Note that, of course, all the IC's are just sheaves so in the whole analysis talking about IC's is not really necessary.

The considerations above allow us to conclude that

$$(9) \quad H^*(\mathcal{H}_x) = H^*(\mathcal{H}_y)^{W_s}$$

where x is a regular element with semisimple part s and y is a regular semisimple element. In particular, this covers some of the results of [BrCh, BaCr] as well as the palindromicity of the cohomology of regular Hessenberg varieties [P].

REFERENCES

- [BaCr] Ana Bălibanu and Peter Crooks, Perverse sheaves and the cohomology of regular Hessenberg varieties, arXiv:2004.07970.
- [BBD] A. A. Beilinson, J. Bernstein and P. Deligne, Faisceaux pervers, in *Analysis and topology on singular spaces, I (Luminy, 1981)*, 5–171, Astérisque, 100, Soc. Math. France, Paris.
- [BrCh] Patrick Brosnan and Timothy Chow, Unit interval orders and the dot action on the cohomology of regular semisimple Hessenberg varieties, *Adv. Math.*, 329:955–1001 (2018).

- [L] George Lusztig, Intersection cohomology complexes on a reductive group. *Invent. Math.* 75 (1984), no.2, 205–272.
- [P] Martha Precup, The Betti numbers of regular Hessenberg varieties are palindromic, *Transf. Groups*, 23(2):491–499 (2017).
- [X] Ke Xue, Affine pavings of Hessenberg ideal fibers, arXiv: 2007.08712.

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA,
ALSO DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI, HELSINKI, FINLAND

Email address: kari.vilonen@unimelb.edu.au, kari.vilonen@helsinki.fi

SCHOOL OF MATHEMATICS AND STATISTICS, UNIVERSITY OF MELBOURNE, VIC 3010, AUSTRALIA,
ALSO DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF HELSINKI, HELSINKI, FINLAND

Email address: ting.xue@unimelb.edu.au