COUNTING CENTRALIZERS OF A FINITE GROUP WITH AN APPLICATION IN CONSTRUCTING THE COMMUTING CONJUGACY CLASS GRAPH

A. R. ASHRAFI AND M. A. SALAHSHOUR*

ABSTRACT. The set of all centralizers of elements in a finite group G is denoted by Cent(G) and G is called n-centralizer if |Cent(G)| = n. In this paper, the structure of centralizers in a non-abelian finite group G with this property that $\frac{G}{Z(G)} \cong Z_{p^2} \rtimes Z_{p^2}$ is obtained. As a consequence, it is proved that such a group has exactly $[(p+1)^2 + 1]$ element centralizers and the structure of the commuting conjugacy class graph of G is completely determined.

Keywords: Commuting graph, commuting conjugacy class graph, element centralizer.

2010 Mathematics Subject Classification: Primary: 20C15; Secondary: 20D15.

1. INTRODUCTION

Throughout this paper all groups are assumed to be finite and $C_G(x)$ denotes the centralizer of an element x in G. If G is a group containing two subgroups H and N such that $N \triangleleft G$, $H \cap N = 1$ and G = HN then G is said to be the semi-direct product of H by N and we write $G = N \rtimes H$. The group G is called capable if there exists another group H such that $G \cong \frac{H}{Z(H)}$. Our other notations are standard and taken mainly from [9] and our calculations are done with the aid of GAP [12].

The center of G is denoted by Z(G) and $Cent(G) = \{C_G(x) \mid x \in G\}$. The group G is called *n*-centralizer if n = |Cent(G)| [1]. The study of finite groups with respect to the number of distinct element centralizers was started by Belcastro and Sherman [3]. It is clear that |Cent(G)| = 1 if and only if G is abelian and there is no finite group with exactly two or three element centralizers. They proved that if $\frac{G}{Z(G)} \cong \mathbb{Z}_p \times \mathbb{Z}_p$ then |Cent(G)| = p + 2 [3, Theorem 5].

Let H be a graph with vertex set $\{1, 2, \dots, k\}$ and G_i 's, $1 \leq i \leq k$, are disjoint graphs of order n_i . Following Sabidussi [10], the graph $H[G_1, G_2, \dots, G_k]$ is formed by taking the graphs G_1, G_2, \dots, G_k and connect a vertex of G_i to another vertex in G_j whenever

^{*}Corresponding author (Email: salahshour@iausk.ac.ir).

i is adjacent to *j* in *H*. The graph $H[G_1, G_2, \dots, G_k]$ is called the *H*-join of the graphs G_1, \dots, G_k .

The notion of **commuting conjugacy class graph** of a non-abelian group G, $\Gamma(G)$, was introduced by Mohammadian et al. [8]. This is a simple graph with non-central conjugacy classes of G as its vertex set and two distinct vertices A and B are adjacent if and only if there are $x \in A$ and $y \in B$ such that xy = yx. In such a case, we also say that two conjugacy classes commute to each other. The authors of the mentioned paper obtained some interesting properties of this graph among them a classification of triangle-free commuting conjugacy class graph of a finite groups is given.

A **CA-group** is a group in which every noncentral element has an abelian centralizer. The present authors [11], obtained the structure of the commuting conjugacy class graph of finite CA-groups. They proved that this graph is a union of some complete graphs. As a consequence of this work, the commuting conjugacy class graph of dihedral, semi-dihedral, dicyclic and three other families of meta-cyclic groups were constructed.

The following two theorems are the main results of this paper:

Theorem 1.1. Suppose p is a prime number and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$. Then G has exactly $[(p+1)^2 + 1]$ element centralizers.

Theorem 1.2. Suppose p is a prime number and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$. The commuting conjugacy class graph of G has one of the following types:

- (1) $\frac{G}{Z}$ is abelian. In this case, $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$ and the commuting conjugacy class graph of G is isomorphic to $\mathcal{M}_1[K_{n(p-1)}, \cdots, K_{n(p-1)}, K_{m(p^2-p)}, \cdots, K_{m(p^2-p)}]$, where there are p+1 copies of $K_{n(p-1)}$, $p^2 + p$ copies of $K_{m(p^2-p)}$ and \mathcal{M}_1 is the graph depicted in Figure 1a. Here, $m = \frac{|Z|}{p^2}$ and $n = \frac{|Z|}{p}$.
- (2) $\frac{G}{Z}$ is not abelian. In this case, the commuting conjugacy class graph of G is isomorphic to $\mathcal{M}_2[K_{n(p-1)}, \cdots, K_{n(p-1)}, K_{n(p^2-p)}]$, where there are $p^2 + p + 1$ copies of $K_{n(p-1)}$, a copy of $K_{n(p^2-p)}$ and \mathcal{M}_2 is depicted in Figure 1b. Here, p is an odd prime and $n = \frac{|Z|}{p}$.

2. Preliminary Results

The aim of this section is to prove some preliminary results which are crucial in next sections. The greatest common divisor of positive integers r and s is denoted by (r, s). We start this section by the following simple lemma:



FIGURE 1. The Graphs of \mathcal{M}_1 and \mathcal{M}_2 .

Lemma 2.1. Let p be a prime number and G be a group with center Z such that $x^{p^2}, y^p \in Z$. Then, for each positive integer m with this condition that $m \not\equiv 0 \pmod{p}$, we have $C_G(x^m) = C_G(x)$ and $C_G(y^m) = C_G(y)$.

Proof. By our assumption, $(m, p^2) = 1$ and there are n and k such that $mn + kp^2 = 1$. Since $x^{p^2} \in Z$, $C_G(x^m) \subseteq C_G(x^{mn}) = C_G(x^{1-kp^2}) = C_G(xz) = C_G(x)$, as desired. The proof of second one is similar and so it is omitted.

Lemma 2.2. Let G be a finite group with center Z, $p \mid |G|$ and $a, b \in G \setminus Z$ such that $o(aZ) = o(bZ) = p^2$ and $G = \{a^i b^j z \mid z \in Z\}$. Then the following hold:

(a) Suppose $1 \le i, j \le p^2 - 1$ and at least one of i, j is not divisible by p. Then $a^i b^j \ne b^j a^i$. (b) $C_G(a^s b) = C_G(a^t b), \ 1 \le s, t \le p^2 - 1$, if and only if s = t.

Proof. Our argument consists in showing:

- (a) Suppose $1 \leq i, j \leq p^2 1$, $a^i b^j = b^j a^i$ and at least one of i, j is not divisible by p. Without loss of generality we can assume that $j \not\equiv 0 \pmod{p}$. Then $(j, p^2) = 1$ and so there are integers m and n such that $mj + np^2 = 1$. Since $b^{p^2} \in Z$, $a^i b = ba^i$ which means that $a^i \in Z$. This implies that $p^2 = o(aZ) \mid i$, contradict by our condition that $1 \leq i \leq p^2 - 1$.
- (b) If $s \neq t$ and $C_G(a^s b) = C_G(a^t b)$ then $a^s b \in C_G(a^t b)$ and so $a^{s-t} b = ba^{s-t}$. Again $p^2 = o(aZ) \mid s t$ which is impossible.

Hence the result.

The following result is a particular case of [7, Theorem 8.1(c)].

Theorem 2.3. The non-abelian split extension $\mathbb{Z}_4 \rtimes \mathbb{Z}_4$ is not capable.

Suppose p is a prime number. Then it is easy to see that there are only two groups of order p^4 that can be written as a semidirect product of two cyclic groups of order p^2 . One of these groups is the abelian group $L_1 = \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$ and another one is a group presented as $L_2 = \langle x, y | x^{p^2} = y^{p^2} = 1, yx = x^{p+1}y \rangle$, see [5, pp. 145,146] for details. For simplicity of our argument, it is useful to write the presentations of L_1 and L_2 in the form $\langle x, y | x^{p^2} = y^{p^2} = 1, yx = x^{rp+1}y \rangle$, where r = 0, 1.

Lemma 2.4. Suppose $G = L_1$ or L_2 . Then for every $i, j, 0 \le i, j \le p^2 - 1$, which are not simultaneously zero, we have:

$$o(x^{i}y^{j}) = \begin{cases} p & p \mid (i,j) \\ p^{2} & Otherwise \end{cases}.$$

Proof. The result is trivial for L_1 . So, it is enough to assume that r = 1. By definition, for every integers i and j, $0 \le i, j \le p^2 - 1$, such that i, j are not simultaneously zero, we have

$$yx^{2} = yxx$$

= $(x^{p+1}y)x = x^{p+1}(yx)$
= $x^{p+1}(x^{p+1}y)$
= $x^{2(p+1)}y.$

By the same way for each $i, yx^i = x^{i(p+1)}y$ and so

$$y^{2}x^{i} = y(yx^{i})$$

= $y(x^{i(p+1)}y) = (yx^{i(p+1)})y$
= $(x^{i(p+1)^{2}}y)y = x^{i(p+1)^{2}}y^{2}$.

On the other hand, $y^j x^i = x^{i(p+1)^j} y^j$. Since $x^{p^2} = 1$ and $(p+1)^j = p^j + jp^{j-1} + \cdots + \frac{j(j-1)}{2}p^2 + jp + 1$. So,

This shows that

$$\begin{array}{rcl} (x^{i}y^{j})^{2} & = & (x^{i}y^{j})(x^{i}y^{j}) = x^{i}(y^{j}x^{i})y^{j} \\ \\ & = & x^{i}(x^{ijp+i}y^{j})y^{j} \\ \\ & = & x^{ijp}x^{2i}y^{2j}. \end{array}$$

An inductive argument now proves that, for each k,

(2.2)
$$(x^i y^j)^k = x^{\frac{k(k-1)}{2}ijp} x^{ki} y^{kj}.$$

Therefore, for every i, j with $0 \le i, j \le p^2 - 1$ and i, j are not simultaneously zero, we have:

$$o(x^{i}y^{j}) = \begin{cases} p & p \mid (i,j) \\ p^{2} & \text{otherwise} \end{cases}$$

This completes the proof.

3. Proof of Theorem 1.1

In this section, the proof of our first main result will be presented. To do this, we need some information about a group G with this property that $\left|\frac{G}{Z(G)}\right| = p^4$, p is prime, and $\frac{G}{Z(G)}$ can be generated by two elements of order p^2 .

Lemma 3.1. Let G be a finite group with center Z, p be prime and $a, b \in G \setminus Z$. Moreover, we assume that $a^{p^2}, b^{p^2} \in Z$ and $(a^i b^j)^k = a^{\frac{k(k-1)}{2}ijrp}a^{ki}b^{kj}z$ where $z \in Z$, r = 0, 1 and i, j, k are positive integers. The following statements hold:

- i) If $1 \leq i, j \leq p-1$, then there are n_j and s such that $a^{ip}b^{jp} = (a^{sp}b^p)^{n_j}z$, where $n_j \neq 0 \pmod{p}, 1 \leq s \leq p-1$ and $z \in Z$.
- ii) If $1 \leq i \leq p-1$, $1 \leq j \leq p^2-1$ and $j \not\equiv 0 \pmod{p}$, then there are n_j and s such that $a^{ip}b^j = (a^{sp}b)^{n_j}z$, where $n_j \not\equiv 0 \pmod{p}$, $1 \leq s \leq p-1$ and $z \in Z$.
- iii) If $1 \leq j \leq p-1$, $1 \leq i \leq p^2-1$ and $i \not\equiv 0 \pmod{p}$, then there are n_i and s such that $a^i b^{jp} = (ab^{sp})^{n_i} z$, where $n_i \not\equiv 0 \pmod{p}$, $1 \leq s \leq p-1$ and $z \in Z$.
- iv) If $1 \leq i, j \leq p^2 1$, $i \not\equiv 0 \pmod{p}$ and $j \not\equiv 0 \pmod{p}$, then for there are n_j and s such that $a^i b^j = (a^s b)^{n_j} z$, where $n_j \not\equiv 0 \pmod{p}$, $s \not\equiv 0 \pmod{p}$, $1 \leq s \leq p^2 1$ and $z \in \mathbb{Z}$.
- v) If $1 \leq i, j \leq p^2 1$, $i \not\equiv 0 \pmod{p}$ and $j \not\equiv 0 \pmod{p}$, then there are n_i and s such that $a^i b^j = (ab^s)^{n_i} z$, where $n_i \not\equiv 0 \pmod{p}$, $s \not\equiv 0 \pmod{p}$, $1 \leq s \leq p^2 1$ and $z \in Z$.

Proof. By assumption, $(a^i b^j)^k = a^{\frac{k(k-1)}{2}ijrp}a^{ki}b^{kj}z$. In this equality, we assume that i = k = p and j = 1. Since $a^{p^2} \in Z$, $(a^p b)^p = b^p z'$ in which $z' \in Z$. Also $(a^p b)^p (a^p b) = (a^p b)(a^p b)^p$ then $(b^p z')(a^p b) = (a^p b)(b^p z')$ so $a^p b^p = b^p a^p$. Since $a^{p^2}, a^{p^2} \in Z$,

$$(a^{ip}b^{jp})^p = a^{\frac{p(p-1)}{2}ijrp^3}a^{ip^2}b^{jp^2}z \in Z$$

then $(a^{ip}b^{jp})^p \in \mathbb{Z}$. Also we can show that $(a^{ip}b^j)^{p^2}$, $(a^ib^j)^{p^2} \in \mathbb{Z}$. Now, each part of the lemma will be proved separately.

i) In this case (j, p) = 1 and so there are integers m_j and k such that $m_j j + kp = 1$. By our assumption, $a^p b^p = b^p a^p$ and $b^{p^2} \in \mathbb{Z}$ and hence,

$$(a^{ip}b^{jp})^{m_j} = a^{im_jp}b^{jm_jp}$$
$$= a^{im_jp}b^{(1-kp)p}$$
$$= a^{im_jp}b^p z_1.$$

Since $(a^{ip}b^{jp})^p \in Z$, $a^{ip}b^{jp} = (a^{im_jp}b^p)^{n_j}z_3$. It is clear that $p \nmid im_j$ and so there are integers s and k such that $im_j = kp + s$. It is now easy to see that for a fixed positive integer j, $i_1 = i_2$ if and only if $s_1 = s_2$. Since $1 \le i \le p - 1$, $1 \le s \le p - 1$. So, by above discussion and this fact that $a^{p^2} \in Z$,

$$a^{ip}b^{jp} = (a^{im_jp}b^p)^{n_j}z_3 = (a^{kp^2+sp}b^p)^{n_j}z_3 = (a^{sp}b^p)^{n_j}z_3$$

as desired.

ii) Since $p \nmid j$, $(j, p^2) = 1$. So, there are m_j and k such that $m_j j + kp^2 = 1$ in which $(m_j, p) = 1$. By our assumption $a^{p^2}, b^{p^2} \in \mathbb{Z}$ and $(a^i b^j)^k = a^{\frac{k(k-1)}{2}ijtp}a^{ki}b^{kj}$. Thus,

$$(a^{ip}b^{j})^{m_{j}} = a^{\frac{m_{j}(m_{j}-1)}{2}ijtp^{2}}a^{im_{j}p}b^{jm_{j}}z_{1}$$
$$= a^{im_{j}p}b^{1-kp^{2}}z_{2}$$
$$= a^{im_{j}p}bz_{3}.$$

Again since $(m_j, p) = 1$, $(m_j, p^2) = 1$ and so there are n_j and t such that $n_j m_j + tp^2 = 1$ in which $p \nmid n_j$. We now apply above discussion and this fact that $(a^{ip}b^j)^{p^2} \in Z$ to derive the equality $a^{ip}b^j = (a^{im_jp}b)^{n_j}z_5$. A similar argument shows that there exists an integer s such that $1 \leq s \leq p-1$ and $a^{ip}b^j = (a^{sp}b)^{n_j}z$ which completes our argument.

iii) The proof is similar to (ii) and so it is omitted.

iv) Since $p \nmid j$, $(j, p^2) = 1$. So there are m_j and k such that $m_j j + kp^2 = 1$ in which $(m_j, p) = 1$. By our assumption $b^{p^2} \in Z$ and $(a^i b^j)^k = a^{\frac{k(k-1)}{2}ijtp}a^{ki}b^{kj}$. Therefore,

$$(a^{i}b^{j})^{m_{j}} = a^{\frac{m_{j}(m_{j}-1)}{2}ijtp}a^{im_{j}}b^{jm_{j}}z_{1}$$
$$= a^{\frac{m_{j}(m_{j}-1)}{2}ijtp+im_{j}}b^{1-kp^{2}}z_{1}$$
$$= a^{u}bz_{2},$$

where

(3.1)
$$u = \frac{m_j(m_j - 1)}{2}ijtp + im_j = i\left(\frac{m_j(m_j - 1)}{2}jtp + m_j\right).$$

Since $(m_j, p) = 1$, $(m_j, p^2) = 1$ and again there are n_j and t such that $m_j n_j + tp^2 = 1$ in which $p \nmid n_j$. By above discussion and this fact that $(a^i b^j)^{p^2} \in Z$, $a^i b^j = (a^u b)^{n_j} z_4$. It is now clear that $p \nmid u$ and so $p^2 \nmid u$. Choose integers s and k such that $u = kp^2 + s$ and $p \nmid s$. By Equation 3.1, j is fixed, one can see that $i_1 = i_2$ if and only if $u_1 = u_2$ if and only if $s_1 = s_2$. Since $1 \leq i \leq p^2 - 1$, $1 \leq s \leq p^2 - 1$ and by above discussion and the fact that $a^{p^2} \in Z$, $a^i b^j = (a^u b)^{n_j} z_4 = (a^{kp^2 + s} b)^{n_j} z_4 = (a^s b)^{n_j} z$ which completes the proof of this part.

v) The proof is similar to iv and so it is omitted.

Hence the result.

Suppose p is a prime number. By a result of Baer [2], the abelian group $\mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$ is capable. If p is an odd prime, then the non-abelian group $\mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$ is also capable [13]. This means that there exists a group G such that $\frac{G}{Z(G)} \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$ or $\mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$. In the next theorem, we obtain the number and structure of the centralizers of G. We also proved in Theorem 2.3 that the non-abelian group $\mathbb{Z}_4 \rtimes \mathbb{Z}_4$ is not capable. In the next theorem, if p = 2, then we will assume that $\frac{G}{Z} \cong \mathbb{Z}_4 \times \mathbb{Z}_4$.

Theorem 3.2. Suppose p is a prime number and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$. Then the number of centralizers of G is $(p+1)^2 + 1$.

Proof. By our assumption, it is enough to assume that G is non-abelian and $Z \neq 1$. By our discussion before Lemma 2.4, we can see that

$$\frac{G}{Z} \cong Z_{p^2} \rtimes Z_{p^2} = \langle x, y \mid x^{p^2} = y^{p^2} = 1 , yx = x^{rp+1}y \rangle.$$

Thus, there are $a, b \in G \setminus Z$ such that

$$\frac{G}{Z} = \langle aZ, bZ \mid (aZ)^{p^2} = (bZ)^{p^2} = 1 , (bZ)(aZ) = (aZ)^{rp+1}(bZ) \rangle$$
7

and so

$$G = \{a^{i}b^{j}z \mid 0 \le i, j \le p^{2} - 1, ba = a^{rp+1}bz', z, z', a^{p^{2}}, b^{p^{2}} \in Z\}.$$

We now apply Equation 2.2 to prove that

(3.2)
$$(a^i b^j)^k = a^{\frac{k(k-1)}{2}ijrp} a^{ki} b^{kj} z,$$

where $z \in Z$. In Equation 3.2, set i = k = p and j = 1. Since $a^{p^2} \in Z$, there exists $z \in Z$ such that $(a^p b)^p = b^p z$. Therefore,

$$(3.3) a^p b^p = b^p a^p.$$

So, for every $0 \le i, j \le p - 1$,

(3.4)
$$\{a^{sp}b^{tp}z \mid 0 \le s, t \le p-1, \ z \in Z\} \subseteq C_G(a^{ip}b^{jp}z).$$

Note that $|G| = p^4 |Z|$ and for each $x \in G \setminus Z$, $Z \nleq C_G(x) \gneqq G$. So,

(3.5)
$$|C_G(x)| = p|Z|$$
 or $p^2|Z|$ or $p^3|Z|$

Suppose i and j are fixed and $0 \le i, j \le p^2 - 1$. By Equation 3.2, $(a^i b^j)^{p^2} \in \mathbb{Z}$. Thus,

(3.6)
$$\{(a^i b^j)^k z \mid 0 \le k \le p^2 - 1, \ z \in Z\} \subseteq C_G(a^i b^j).$$

Next we partition the elements of G into the following eight parts:

$$V_{1} = \{a^{i}z \mid p \mid i \& z \in Z\},\$$

$$V_{2} = \{a^{i}z \mid p \nmid i \& z \in Z\},\$$

$$V_{3} = \{b^{j}z \mid p \mid j \& z \in Z\},\$$

$$V_{4} = \{b^{j}z \mid p \nmid j \& z \in Z\},\$$

$$V_{5} = \{a^{i}b^{j}z \mid p \mid (i,j) \& z \in Z\},\$$

$$V_{6} = \{a^{i}b^{j}z \mid p \mid i, p \nmid j \& z \in Z\},\$$

$$V_{7} = \{a^{i}b^{j}z \mid p \nmid i, p \mid j \& z \in Z\},\$$

$$V_{8} = \{a^{i}b^{j}z \mid p \nmid i, p \nmid j \& z \in Z\}.$$

To compute the centralizer of non-central elements of G, the following cases will be considered:

a)
$$i \neq 0$$
 and $j = 0$. Then for every $1 \le i \le p^2 - 1$,
(3.7) $\{a^k z \mid 0 \le k \le p^2 - 1, z \in Z\} \subseteq C(a^i).$

1) $p \mid i$. Suppose i = sp such that $1 \leq s \leq p-1$. By Equation 3.4,

$$A = \{a^k b^{tp} z \mid 0 \le k \le p^2 - 1, \ 0 \le t \le p - 1, \ z \in Z\} \subseteq C_G(a^i) = C_G(a^{sp}).$$

Since $|A| = p^3 |Z|$, by Equation 3.5 and Lemma 2.1,

$$C_G(a^p) = C_G(a^{sp}) = \{a^k b^{tp} z \mid 0 \le k \le p^2 - 1, \ 0 \le t \le p - 1, \ z \in Z\}.$$

2) $p \nmid i$. By Lemma 2.2(a), the element a commutes only with elements of $V_1 \cup V_2$ and by Equations 3.7, 3.5 and Lemma 2.1,

$$C_G(a) = C_G(a^i) = \{a^k z \mid 0 \le k \le p^2 - 1, \ z \in Z\} \le C(a^p).$$

- b) $j \neq 0$ and i = 0. Apply a similar argument as the case of (1). To do this, we consider two cases that $p \mid j$ or $p \nmid j$.
 - 3) $p \mid j$. Suppose $j = tp, 1 \le t \le p 1$. Then,

$$C_G(b^p) = C_G(b^{tp}) = \{ a^{sp} b^k z \mid 0 \le k \le p^2 - 1, \ 0 \le s \le p - 1, \ z \in Z \}.$$

4) $p \nmid j$. In this case, we can easily see that,

$$C_G(b) = C_G(b^j) = \{b^k z \mid 0 \le k \le p^2 - 1, \ z \in Z\} \le C(b^p).$$

c) $i \neq 0$ and $j \neq 0$. The following four subcases are considered into account.

5) $p \mid i \text{ and } p \mid j$. Suppose $i = i_1 p$ and $j = j_1 p$ in which $1 \leq i_1, j_1 \leq p - 1$. By Lemma 3.1(I), there are n_{j_1} and s such that $a^{i_1 p} b^{j_1 p} = (a^{sp} b^p)^{n_{j_1}} z$, where $p \nmid n_{j_1}$, $1 \leq s \leq p - 1$ and $z \in Z$. By Equation 3.2, $(a^{sp} b^p)^p \in Z$ and by Lemma (2.1),

$$C_G(a^{i_1p}b^{j_1p}) = C_G((a^{sp}b^p)^{n_{j_1}}z) = C_G((a^{sp}b^p)^{n_{j_1}}) = C_G(a^{sp}b^p).$$

Since $1 \le s \le p-1$, the number of distinct proper centralizers constructed from the elements of V_5 is at most p-1. We now obtain the structures of these centralizers. In Equation 3.2, we put i = s, j = 1 and k = p. Then $(a^s b)^p = a^{sp} b^p z$ in which $z \in Z$. Thus, $(a^{sp} b^p)(a^s b) = (a^s b)(a^{sp} b^p)$ and by Equation 3.4,

$$A = \{a^{ip}(a^{s}b)^{k}b^{jp}z \mid 0 \le i, j, k \le p - 1, \ z \in Z\} \subseteq C_{G}(a^{sp}b^{p}).$$

It can easily see that $|A| = p^3 |Z|$ and by applying Equations 3.2 and 3.5,

(3.8)
$$C_G(a^{sp}b^p) = \{a^{ip}a^{sk}b^kb^{jp}z \mid 0 \le i, j, k \le p-1, z \in Z\}.$$

Suppose $1 \le t \le p-1$. By Equation 3.8,

(3.9)
$$a^t b \in C_G(a^{sp}b^p)$$
 if and only if $t = s$.

Therefore, $C_G(a^{tp}b^p) = C_G(a^{sp}b^p)$, $1 \le s, t \le p-1$, if and only if t = s. This shows that the number of distinct proper centralizers is exactly p-1.

6) $p \mid i \text{ and } p \nmid j$. Suppose $i = i_1 p$, where $1 \leq i_1 \leq p-1$. By Lemma 3.1(II), there are n_j and s such that $a^{i_1 p} b^j = (a^{sp} b)^{n_j} z$ in which $p \nmid n_j$, $1 \leq s \leq p-1$ and $z \in Z$. Now by Equation 3.2, $(a^{sp} b)^{p^2} \in Z$. Hence by Lemma 2.1, $C_G(a^{i_1 p} b^j) = C_G((a^{sp} b)^{n_j} z)$ $= C_G((a^{sp} b)^{n_j}) = C_G(a^{sp} b)$. Since $1 \leq s \leq p-1$, by Lemma 2.2(b), the number of distinct proper centralizers is exactly p-1. To obtain the structures of these centralizers, it is enough to apply Equation 3.6. This implies that

$$A = \{ (a^{sp}b)^k z \mid 0 \le k \le p^2 - 1 , z \in Z \} \subseteq C(a^{sp}b).$$

On the other hand, by Equation 3.2, $(a^{sp}b)^k = a^{spk}b^kz'$ in which $z' \in Z$. We assume that $k = tp + k_1$ in which $0 \le t, k_1 \le p - 1$. Thus, $(a^{sp}b)^k = a^{spk}b^kz' = a^{spk_1}b^{k_1}b^{tp}z''$, where $z'' \in Z$. So,

$$A = \{ a^{spk_1} b^{k_1} b^{t_p} z \mid 0 \le k_1, t \le p - 1 , z \in Z \} \subseteq C(a^{sp}b).$$

By Lemma 2.2(a), $a^{sp}b$ cannot commute with an element of $V_1 \cup V_2 \cup V_4 \cup V_5 \cup V_7$. Furthermore, it is easy to see that $V_3 \subseteq A$. We claim that there are no elements $x \in V_6$ and $y \in V_8$ such that xy = yx. To see this, we assume that $x = a^{i_1 p} b^{j_1} \in V_6$ and $y = a^{i_2} b^{j_2} \in V_8$ are arbitrary. By Lemma 3.1(*iii*,*iv*), $a^{i_1p}b^{j_1} = (a^{sp}b)^{n_{j_1}}z_1$ and $a^{i_2}b^{j_2} = (a^tb)^{n_{j_2}}z_2$, where $p \nmid n_{j_1}, p \nmid n_{j_2}, p \nmid s$ and $p \nmid t$. By Equation 3.2, $(a^{sp}b)^{p^2}$, $(a^tb)^{p^2} \in Z$ and by Lemma 2.1, $C_G(a^{i_1p}b^{j_1}) = C_G(a^{sp}b)$ and $C_G(a^{i_2}b^{j_2}) = C_G(a^t b)$. Thus, $(a^{i_1p}b^{j_1})(a^{i_2}b^{j_2}) = (a^{i_2}b^{j_2})(a^{i_1p}b^{j_1})$ if and only if $(a^{sp}b)(a^tb) = (a^tb)(a^{sp}b)$ if and only if $a^{sp}ba^t = a^tba^{sp}$ if and only if $a^{sp-t}b = ba^{sp-t}$. We now apply Lemma 2.2(a) to deduce that $p \mid t$ which is impossible. Next we prove that $C(a^{sp}b) = A$. Choose $x \in C(a^{sp}b) \setminus A$. By above discussion, $x \in V_6$ and so we can write $x = a^{ip}b^{j}$, where $1 \le i \le p-1$. By Lemma 3.1(*ii*), there are n_{i} and m such that $a^{ip}b^j = (a^{mp}b)^{n_j}z$ with this condition that $p \nmid n_j, 1 \leq m \leq p-1$, and $z \in Z$. On the other hand, By Equation 3.2, $(a^{mp}b)^{p^2} \in Z$. So, by Lemma 2.1, $C(a^{ip}b^j) = C(a^{mp}b)$. Since $x \in C(a^{sp}b)$, $a^{sp}b \in C_G(x) = C(a^{mp}b)$. Hence, $a^{(s-m)p}b$ $= ba^{(s-m)p}$. Again by Lemma 2.2(a), $p \mid s-m$ and since $1 \leq s, m \leq p-1, s=m$. This proves that $C(a^{sp}b) = A$. Therefore,

$$C(a^{sp}b) = \{a^{spk}b^k b^{tp}z \mid 0 \le k, t \le p-1, z \in Z\}.$$

7) $p \mid j \text{ and } p \nmid i$. A similar argument as Case (6) shows that the number of distinct proper centralizers of G is equal to p-1 and

$$C_G(ab^{tp}) = \{ a^{sp} a^k b^{tpk} z \mid 1 \le k, s \le p - 1, \ z \in Z \}.$$

8) $p \nmid i$ and $p \nmid j$. By Lemma 3.1(*iv*), there are n_j and s such that $a^i b^j = (a^s b)^{n_j} z$ in which $p \nmid n_j$, $p \nmid s$, $1 \leq s \leq p^2 - 1$ and $z \in Z$. By Equation 3.2, $(a^s b)^{p^2} \in Z$. Thus, by Lemma 2.1, $C_G(a^i b^j) = C_G((a^s b)^{n_j} z) = C_G((a^s b)^{n_j}) = C_G(a^s b)$. Since $p \nmid s, 1 \leq s \leq p^2 - 1$ and Lemma 2.2(b), the number of distinct proper centralizers contained in V_8 is equal to $p^2 - p$. We now obtain the structure of centralizers. By Equations 3.6 and (3.2),

(3.10)
$$A = \{ (a^{s}b)^{k}z \mid 0 \le k \le p^{2} - 1 , z \in Z \} \subseteq C(a^{s}b) \\ = \{ a^{\frac{k(k-1)}{2}sp+sk}b^{k}z \mid 0 \le k \le p^{2} - 1 , z \in Z \}.$$

By Lemma 2.2(a) and definition of V_7 , one can see that $a^s b$ cannot be commuted with an elements in $V_1 \cup V_2 \cup V_3 \cup V_4 \cup V_6 \cup V_7$. We claim that $C(a^{sp}b) = A$. To prove, we assume that $x \in C_G(a^s b) \setminus A$. By above discussion, $x \in V_5 \cup V_8$. If $x \in V_8$, then $x = a^{i'}b^{j'}$ and by Lemma 3.1(*iv*), there are $n_{j'}$ and t such that $a^{i'}b^{j'} = (a^t b)^{n_{j'}}z$ in which $p \nmid n_{j'}, p \nmid t$ and $z \in Z$. Hence by Equation 3.2, $(a^t b)^{p^2} \in Z$ and by Lemma 2.1, we have $C_G(a^{i'}b^{j'}) = C_G(a^t b)$. Since $x \in C_G(a^s b)$, $a^s b \in C_G(x) = C_G(a^t b)$ and so $a^{s-t}b = ba^{s-t}$. Apply Lemma 2.2(a) to deduce that $p^2 \mid s - t$. Since $1 \leq s, t \leq p^2 - 1$, s = t and $x = a^{i'}b^{j'} = (a^s b)^{n_{j'}}z \in A$. If $x \in V_5$, then $x = a^{i'p}b^{j'p}$ and by Lemma 3.1(*i*), there are $n_{j'}$ and t such that $a^{i'p}b^{j'p} = (a^{tp}b^p)^{n_{j'}}z$ in which $p \nmid n_{j'}, p \nmid t$ and $z \in Z$. Next by Equation 3.2, $(a^{tp}b^p)^p \in Z$ and Lemma 2.1 implies that $C_G(a^{i'p}b^{j'p}) = C_G(a^{tp}b^p)$. Since $x \in C_G(a^s b), a^s b \in C_G(x) = C_G(a^{tp}b^p)$ and by Equation 3.9, one can see that t = s. Hence Equation 3.2 shows that $x = a^{i'p}b^{j'p} = (a^{sp}b^p)^{n_{j'}}z = (a^s b)^{n_{j'}p}z \in A$. Therefore, in both cases $x \in A$ which means that

$$C_G(a^s b) = A = \{ (a^s b)^k z \mid 0 \le k \le p^2 - 1, z \in Z \}.$$

In addition, by Equations 3.8 and 3.10, we can see that $C_G(a^{sb}) \leq C_G(a^{sp}b^p)$.

Therefore, the structure of non-trivial proper centralizers of G is as follows:

$$\begin{array}{lll} C_{G}(a^{p}) &=& \{a^{k}b^{jp}z \mid 0 \leq j \leq p-1, \ 0 \leq k \leq p^{2}-1, \ z \in Z\}, \\ C_{G}(a) &=& \{a^{k}z \mid 0 \leq k \leq p^{2}-1\} \leq C_{G}(a^{p}), \\ C_{G}(b^{p}) &=& \{a^{ip}b^{k}z \mid 0 \leq i \leq p-1, \ 0 \leq k \leq p^{2}-1, \ z \in Z\}, \\ C_{G}(b) &=& \{b^{k}z \mid 0 \leq k \leq p^{2}-1\} \leq C_{G}(b^{p}), \\ C_{G}(a^{sp}b^{p}) &=& \{a^{ip}a^{sk}b^{k}b^{jp}z \mid 0 \leq i, j, k \leq p-1, \ z \in Z\}, 1 \leq s \leq p-1 \\ C_{G}(a^{sp}b) &=& \{a^{ip}a^{k}b^{spk}z \mid 0 \leq i, k \leq p-1, \ z \in Z\} \leq C_{G}(a^{p}), 1 \leq s \leq p-1 \\ C_{G}(a^{sp}b) &=& \{a^{spk}b^{k}b^{jp}z \mid 0 \leq j, k \leq p-1, \ z \in Z\} \leq C_{G}(b^{p}), 1 \leq s \leq p-1 \\ C_{G}(a^{sp}b) &=& \{(a^{s}b)^{k}a^{isp}b^{ip}z \mid 0 \leq i, k \leq p-1, \ z \in Z\} \leq C_{G}(a^{sp}b^{p}), 1 \leq s \leq p^{2}-1, p \nmid s \\ \end{array}$$

So, the number of distinct proper centralizers of G is $(p^2-p)+3(p-1)+4=p^2+2p+1=(p+1)^2$ and therefore $|Cent(G)|=(p+1)^2+1$. This completes the proof.

We end this section by the following conjecture:

Conjecture 3.3. Suppose p is a prime number, n is a positive integer and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^n} \rtimes \mathbb{Z}_{p^n}$. Then $|Cent(G)| = (p+1)^n + 1$.

4. Proof of Theorem 1.2

In this section, we apply the results of Section 3 to obtain the structure of the commuting conjugacy class graph of G when $\frac{G}{Z(G)} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$.

Theorem 4.1. Suppose p is a prime and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$. Then the commuting conjugacy classes graph of G has the graph structure

$$\mathcal{M}_1[\overbrace{K_{n(p-1)},\cdots,K_{n(p-1)}}^{p+1},\overbrace{K_{m(p^2-p)},\cdots,K_{m(p^2-p)}}^{p^2+p}]$$

where \mathcal{M}_1 is the graph depicted in Figure 2. Here, $m = \frac{|Z|}{p^2}$ and $n = \frac{|Z|}{p}$.



FIGURE 2. The Graph \mathcal{M}_1 .

Proof. Since $\frac{G}{Z}$ is abelian, xgZ = xZgZ = gZxZ = gxZ, where $x, g \in G \setminus Z$ are arbitrary. Hence there exists $z \in Z$ such xg = gxz and so for every $x \in G \setminus Z$,

(4.1)
$$x^{G} = \{g^{-1}xg \mid g \in G\} = \{g^{-1}gxz \mid z \in Z\} = \{xz \mid z \in Z\}.$$

We now apply Theorem 3.2 to prove that the group G has the following structure:

$$G = \{a^{i}b^{j}z \mid 0 \le i, j \le p^{2} - 1, \ ba = abz_{t}, \ z, z_{t} \in Z\}$$

Hence for each $x \in G \setminus Z$, $x = a^i b^j z$, where $0 \le i, j \le p^2 - 1$ and i, j are not simultaneously zero. To obtain the non-central conjugacy classes of G, the following have to be investigated:

- a) $i \neq 0$ and j = 0. We will consider two cases that $p \mid i$ and $p \nmid i$.
 - 1) $p \mid i$. In this case, i = sp, where $1 \le s \le p-1$. By Theorem 3.2, $|C_G(a^{sp})| = p^3|Z|$ and so $|(a^{sp})^G| = p$. By Equation 4.1, $(a^{sp})^G = \{a^{sp}z_1, a^{sp}z_2, \cdots, a^{sp}z_p\} = a^{sp}H$, where $H = \{z_1, z_2, \cdots, z_p\} \subseteq Z$. Suppose |Z| > p and choose $z_{r_1} \in Z \setminus H$. It is easy to see that $a^{sp}z_{r_1} \notin (a^{sp})^G$ and so $(a^{sp}z_{r_1})^G \neq (a^{sp})^G$. Since $C_G(a^{sp}z_{r_1}) = C_G(a^{sp})$, $|(a^{sp}z_{r_1})^G| = |(a^{sp})^G| = p$. Also, $(a^{sp}z_{r_1})^G = (a^{sp})^G z_{r_1} = a^{sp}Hz_{r_1}$. Since $(a^{sp})^G \cap$ $(a^{sp}z_{r_1})^G = \emptyset$, $H \cap Hz_{r_1} = \emptyset$ and $H \cup Hz_{r_1} \subseteq Z$. We now choose an element $z_{r_2} \in$ $Z \setminus (H \cup Hz_{r_1})$ and by above method we will see that $p \mid |Z|$. Suppose $n = \frac{|Z|}{p}$. Thus, there are n distinct conjugacy classes of the form $(a^{sp}z_{r_1})^G, (a^{sp}z_{r_2})^G, \dots, (a^{sp}z_{r_n})^G$. Since $1 \le s \le p-1$, there are n(p-1) distinct conjugacy classes with $n = \frac{|Z|}{p}$ and each of which has p elements.
 - 2) $p \nmid i$. Then by Theorem 3.2, $|C_G(a^i)| = p^2 |Z|$ and so $|(a^i)^G| = p^2$. Also, by Equation 4.1, $(a^i)^G = \{a^i z_1, a^i z_2, \cdots, a^i z_{p^2}\} = a^i H$, where $H = \{z_1, z_2, \cdots, z_{p^2}\} \subseteq Z$. Suppose that $|Z| > p^2$ and choose $z_{r_1} \in Z \setminus H$. It is easy to see that $a^i z_{r_1} \notin (a^i)^G$ and hence $(a^i z_{r_1})^G \neq (a^i)^G$. Since $C_G(a^i z_{r_1}) = C_G(a^i)$, $|(a^i z_{r_1})^G| = |(a^i)^G| = p^2$. Also, $(a^i z_{r_1})^G = (a^i)^G z_{r_1} = a^i H z_{r_1}$. Note that $(a^i)^G \cap (a^i z_{r_1})^G = \emptyset$. Thus, $H \cap H z_{r_1} = \emptyset$ and $H \cup H z_{r_1} \subseteq Z$. Choose the element $z_{r_2} \in Z \setminus (H \cup H z_{r_1})$. By repeated applications of above method we can see that $p^2 \mid |Z|$. Define $m = \frac{|Z|}{p^2}$. Then for constant i, there are m distinct conjugacy classes in the form of $(a^i z_{r_1})^G, (a^i z_{r_2})^G, \dots, (a^i z_{r_m})^G$. Since $1 \leq i \leq p^2 - 1$ and $p \nmid i$, there are $m(p^2 - p)$ distinct conjugacy classes, where $m = \frac{|Z|}{p^2}$ and each of which has p^2 elements.
- b) i = 0 and $j \neq 0$.
 - 3) If $p \mid j$, then j = tp, where $1 \le t \le p-1$. By Theorem 3.2, $|C_G(b^{tp})| = p^3 |Z|$ and so $|(b^{tp})^G| = p$. Then by Equation 4.1 and a similar argument as (1), for each t there are n distinct conjugacy classes of the form $(b^{tp}z_{r_1})^G, (b^{tp}z_{r_2})^G, \ldots, (b^{tp}z_{r_n})^G$, where $n = \frac{|Z|}{p}$. Since $1 \le t \le p-1$, there are n(p-1) distinct conjugacy classes with $n = \frac{|Z|}{p}$ and each class has p elements.
 - 4) If $p \nmid j$, then by Theorem 3.2, $|C_G(b^j)| = p^2 |Z|$ and so $|(b^j)^G| = p^2$. Apply again Equation 4.1 and a similar argument as (2) to result that for each constant j, there are m distinct conjugacy classes of the form $(b^j z_{r_1})^G, (b^j z_{r_2})^G, \ldots, (b^j z_{r_m})^G$, where $m = \frac{|Z|}{p^2}$. Since $1 \leq j \leq p^2 - 1$ and $p \nmid j$, there are $m(p^2 - p)$ distinct conjugacy classes with $m = \frac{|Z|}{p^2}$ and each class has p^2 elements.

- c) $i \neq 0$ and $j \neq 0$. This case can be separated into the following four cases:
 - 5) $p \mid i \text{ and } p \mid j$. In this case, i = sp and j = tp in which $1 \leq s, t \leq p 1$. By Theorem 3.2, $|C_G(a^{sp}b^{tp})| = p^3|Z|$ and so $|(a^{sp}b^{tp})^G| = p$. We now apply Equation 4.1 and a similar argument as (1) for constants s and t to deduce that there are n distinct conjugacy classes of the form $(a^{sp}b^{tp}z_{r_1})^G, (a^{sp}b^{tp}z_{r_2})^G, \ldots, (a^{sp}b^{tp}z_{r_n})^G,$ where $n = \frac{|Z|}{p}$. Since $1 \leq s, t \leq p - 1$, there are $n(p-1)^2$ distinct conjugacy classes with $n = \frac{|Z|}{p}$ and each class has p members.
 - 6) $p \nmid i \text{ and } p \mid j$. Clearly $j = tp, 1 \leq t \leq p-1$. By Theorem 3.2, $|C_G(a^i b^{tp})| = p^2 |Z|$ and hence $|(a^i b^{tp})^G| = p^2$. On the other hand, Equation 4.1 and a similar argument as (2) for constants i and t, show that there are m distinct conjugacy classes of the form $(a^i b^{tp} z_{r_1})^G, (a^i b^{tp} z_{r_2})^G, \ldots, (a^i b^{tp} z_{r_m})^G$, where $m = \frac{|Z|}{p^2}$. Since $1 \leq t \leq p-1$, $1 \leq i \leq p^2 - 1$ and $p \nmid i$, there are $mp(p-1)^2$ distinct conjugacy classes with $m = \frac{|Z|}{p^2}$ and each class has p^2 members.
 - 7) $p \mid i \text{ and } p \nmid j$. In this case i = sp, for some integer s such that $1 \leq s \leq p-1$. By Theorem 3.2, $|C_G(a^{sp}b^j)| = p^2|Z|$ and so $|(a^{sp}b^j)^G| = p^2$. Then by Equation 4.1 and a similar argument as (2) for constants s and j, one can prove that there are m distinct conjugacy classes of the form $(a^{sp}b^jz_{r_1})^G, (a^{sp}b^jz_{r_2})^G, \ldots, (a^{sp}b^jz_{r_m})^G,$ where $m = \frac{|Z|}{p^2}$. Since $1 \leq s \leq p-1, 1 \leq j \leq p^2-1$ and $p \nmid j$, there are $mp(p-1)^2$ distinct conjugacy classes with $m = \frac{|Z|}{p^2}$ and each class has p^2 members.
 - 8) $p \nmid i$ and $p \nmid j$. By Theorem 3.2, $|C_G(a^i b^j)| = p^2 |Z|$ and so $|(a^i b^j)^G| = p^2$. On the other hand, by Equation 4.1 and applying a similar argument as (2) for constants *i* and *j*, one can see that there are *m* distinct conjugacy classes of the form $(a^i b^j z_{r_1})^G, (a^i b^j z_{r_2})^G, \ldots, (a^i b^j z_{r_m})^G$, where $m = \frac{|Z|}{p^2}$. Since $1 \leq i, j \leq p^2 - 1$ and $p \nmid i, j$, there are $mp^2(p-1)^2$ distinct conjugacy classes with $m = \frac{|Z|}{p^2}$ and each class has length p^2 .

The above discussion are summarized in Table 1.

We now obtain the commuting conjugacy class graph of G. To do this, we consider the following cases:

(1) Suppose a^{s_1p} and a^{s_2p} are representatives of two conjugacy classes of Type 1 in Table 1. It is clear that $a^{s_1p}a^{s_2p} = a^{s_2p}a^{s_1p}$ and so all such classes are commuting together. Hence, the commuting conjugacy class graph has a subgraph isomorphic to the complete graph $K_{n(p-1)}$. We now assume that a^{sp} is a representative of a conjugacy class of Type 1 and $a^u b^v z$ is an arbitrary element of Gsuch that $(a^{sp})(a^u b^v z) = (a^u b^v z)(a^{sp})$. Thus $a^{sp}b^v = b^v a^{sp}$ and by Lemma 2.2(a),

Type	The R	tepresentatives of Conjugacy Classes	$ x^G $	# Conjugacy Classes
1	a^{sp}	$1 \le s \le p - 1$	p	n(p-1)
2	a^i	$1 \le i \le p^2 - 1, i \not\equiv 0$	p^2	mp(p-1)
3	b^{tp}	$1 \le t \le p - 1$	p	n(p-1)
4	b^j	$1 \leq j \leq p^2 - 1, j \stackrel{p}{\not\equiv} 0$	p^2	mp(p-1)
5	$a^{sp}b^{tp}$	$1 \leq s,t \leq p-1$	p	$n(p-1)^2$
6	$a^i b^{tp}$	$1 \le t \le p-1, 1 \le i \le p^2 - 1, i \not\equiv 0$	p^2	$mp(p-1)^2$
7	$a^{sp}b^j$	$1 \le s \le p-1, 1 \le j \le p^2 - 1, j \not\equiv 0$	p^2	$mp(p-1)^2$
8	$a^i b^j$	$1 \leq i,j \leq p^2 - 1, i \neq 0, j \neq 0$	p^2	$mp^2(p-1)^2$

TABLE 1. Conjugacy classes of G with $n = \frac{|Z|}{p}$ and $m = \frac{|Z|}{p^2}$.

 $v \equiv 0 \pmod{p}$ or v = tp in which $0 \le t \le p - 1$. Hence a^{sp} commutes with all elements in the form $a^u b^{tp} z$. Now the following cases can be occurred:

- (a) If t = 0, then the conjugacy classes of Types 1 and 2 in Table 1 are commuting to each other.
- (b) If $t \neq 0$ and u = 0, then the conjugacy classes of Types 1 and 3 in Table 1 are commuting to each other.
- (c) If $u, t \neq 0$ and $p \mid u$, then the conjugacy classes of Types 1 and 5 in Table 1 can be commuted to each other.
- (d) If $t \neq 0$ and $p \nmid u$, then the conjugacy classes of Types 1 and 6 in Table 1 are commuting together.
- (2) It is clear that two conjugacy classes of Type 2 are commuted to each other and so the commuting conjugacy class graph has a subgraph isomorphic to K_{mp(p-1)}. We now determine the relationship between conjugacy classes of this and other types. To do this, we assume that aⁱ is a representatives of a conjugacy class of Type 2 and a^ub^vz is an arbitrary element of G such that (aⁱ)(a^ub^vz) = (a^ub^vz)(aⁱ). So aⁱb^v = b^vaⁱ and since p ∤ i, by Lemma 2.2(a) the latter is true if and only if v = 0. Therefore, aⁱ commutes with all elements in the form a^uz. This shows that the conjugacy classes of Type 2 are commuted only with the conjugacy classes of Type 1.
- (3) Suppose b^{t_1p} and b^{t_2p} are the representative of two conjugacy classes of Type 3. It is clear that $b^{t_1p}b^{t_2p} = b^{t_2p}b^{t_1p}$ and so all conjugacy classes of Type 3 are commuting together in the graph. Hence by Table 1, the graph has a subgraph isomorphic

to $K_{n(p-1)}$. To determine the relationship between conjugacy classes of this and other types, we choose b^{tp} to be a representative of a conjugacy class of Type 3 and $a^u b^v z$ as an arbitrary element of G such that $(b^{tp})(a^u b^v z) = (a^u b^v z)(b^{tp})$. Thus, $b^{tp}a^u = a^u b^{tp}$ and by Lemma 2.2(a), the later is true if and only if $u \equiv 0 \pmod{p}$ or u = sp in which $0 \le s \le p - 1$. Therefore, b^{tp} can be commuted with all elements in the form $a^{sp}b^v z$. Hence, the following are satisfied:

- (a) If s = 0, then the conjugacy classes of Types 3 and 4 are commuting together.
- (b) If $s \neq 0$ and v = 0, then the conjugacy classes of Types 3 and 1 are commuting to each other.
- (c) If $v, s \neq 0$ and $p \mid v$, then the conjugacy classes of Types 3 and 5 are commuting together.
- (d) If $s \neq 0$ and $p \nmid v$, then the conjugacy classes of Types 3 and 7 are commuting to each other.
- (4) Suppose b^{j_1} and b^{j_2} are the representatives of two conjugacy classes of Type 4. It is obvious that $b^{j_1}b^{j_2} = b^{j_2}b^{j_1}$ which shows that all conjugacy classes of Type 4 are commuted to each other. So, the commuting conjugacy class graph has a subgraph isomorphic to $K_{mp(p-1)}$. To determine the relationship between this and other types in Table 1, we assume that b^j is a representative of a conjugacy class of type 4 and $a^u b^v z$ is an arbitrary element of G such that $(b^j)(a^u b^v z) = (a^u b^v z)(b^j)$. Thus, $b^j a^u = a^u b^j$ and since $p \nmid j$, by Lemma 2.2(a), the latter is true if and only if u = 0. Therefore, b^j commutes with all elements in the form $b^v z$ and hence the conjugacy classes of Type 4 can be commuted only with the conjugacy classes of Type 3.
- (5) Suppose $a^{s_1p}b^{t_1p}$ and $a^{s_2p}b^{t_2p}$ are the representatives of two conjugacy classes of Type 5. By Equation 3.3, $(a^{s_1p}b^{t_1p})(a^{s_2p}b^{t_1p}) = (a^{s_2p}b^{t_2p})(a^{s_1p}b^{t_2p})$ and so all conjugacy classes of Type 5 are commuted to each other. This gives us the complete graph $K_{n(p-1)^2}$ as a subgraph of the commuting conjugacy class graph. Hence, it is enough to determine the relationship between this and conjugacy classes of Types 6, 7 and 8. Suppose $a^{sp}b^{tp}$ and a^ub^{vp} are the representatives of the conjugacy classes of Types 5 and 6, respectively, such that they are commuting together. By Lemma 3.1, $a^{sp}b^{tp} = (a^{ip}b^p)^{n_t}z_1$ and $a^ub^{vp} = (ab^{jp})^{n_u}z_2$ and so by Lemma 2.1, $C_G(a^{sp}b^{tp}) = C_G(a^{ip}b^p)$ and $C_G(a^ub^{vp}) = C_G(ab^{jp})$. It is now easy to prove that $(a^{sp}b^{tp})(a^ub^{vp}) = (a^ub^{vp})(a^{sp}b^{tp})$ if and only if $(a^{ip}b^p)(a^{bjp}) = (ab^{jp})(a^{ip}b^p)$ if and only if $b^pa = ab^p$ which is a contradiction with Lemma 2.2(a). Therefore, a conjugacy class of Type 5 is not adjacent with a conjugacy class of Type 6. Similarly,

a conjugacy class of Type 5 is not adjacent to another one of Type 7. We now assume that $a^{sp}isb^{tp}$ and a^ub^v are representatives of the conjugacy classes of Types 5 and 8, respectively. By Lemma 3.1, $a^{sp}b^{tp} = (a^{ip}b^p)^{n_t}z_1$ and $a^ub^v = (a^jb)^{n_v}z_2$. On the other hand, By Lemma 2.1, $C_G(a^{sp}b^{tp}) = C_G(a^{ip}b^p)$ and $C_G(a^ub^v) = C_G(a^jb)$ and by Equation 3.9, we can see $(a^{sp}b^{tp})(a^ub^v) = (a^ub^v)(a^{sp}b^{tp})$ if and only if $(a^{ip}b^p)(a^jb) = (a^jb)(a^{ip}b^p)$ if and only if $j \equiv i(mod \ p)$. Since $1 \le i \le p - 1$, the conjugacy classes of Types 5 and 8 can be divided into p - 1 parts. Moreover, this process constructs a complete subgraph of size n(p - 1) in the commuting conjugacy class graph.

(6) Suppose $a^{i_1}b^{t_1p}$ and $a^{i_2}b^{t_2p}$ are the representative of two conjugacy classes of Types 6 such that they are commute to each other. By Lemma 3.1, $a^{i_1}b^{t_1p} = (ab^{s_1p})^{n_{i_1}}z_1$ and $a^{i_2}b^{t_2p} = (ab^{s_2p})^{n_{i_2}}z_2$. On the other hand, by Lemma (2.1), $C_G(a^{i_1}b^{t_1p}) =$ $C_G(ab^{s_1p})$ and $C_G(a^{i_2}b^{t_2p}) = C_G(ab^{s_2p})$. Therefore, $(a^{i_1}b^{t_1p})(a^{i_2}b^{t_2p}) = (a^{i_2}b^{t_2p})(a^{i_1}b^{t_1p})$ if and only if $(ab^{s_1p})(ab^{s_2p}) = (ab^{s_2p})(ab^{s_1p})$ if and only if $ab^{(s_2-s_1)p} = b^{(s_2-s_1)p}a$. By Lemma 2.2(a), the last equality is satisfied if and only if $s_1 = s_2$. This proves that in the commuting conjugacy class graph, two conjugacy classes of Type 6 are adjacent, when the centralizers of their representatives is equal. By Theorem 3.2, the number of centralizer of Types 6 is p-1 and so the conjugacy classes of this type can be divided into p-1 parts. On the other hand, by Table 1, the number of conjugacy classes of Type 6 is $mp(p-1)^2$. Hence, each part of Type 6 is a clique of size mp(p-1). Now it is enough to determine the relationship between conjugacy classes of Type 6 with other conjugacy classes of Types 7 and 8. To do this, we assume that $a^i b^{tp}$ and $a^{sp} b^j$ are the representatives of conjugacy classes of Types 6 and 7, respectively, such that they are commute to each other. By Equation 3.3, $(a^i b^{tp})(a^{sp} b^j) = (a^{sp} b^j)(a^i b^{tp})$ if and only if $a^i b^j = b^j a^i$. Since $p \nmid i, j$, Lemma 2.2(a) implies that the last equality cannot be occurred. This means that a conjugacy class of Type 6 is not adjacent with another one of Type 7 in the commuting conjugacy class graph. Next, we assume that $a^i b^{tp}$ and $a^u b^v$ are the representatives of two conjugacy classes of Types 6 and 8, respectively and they are commute to each other. By Lemma 3.1, $a^i b^{tp} = (ab^{sp})^{n_i} z_1$ and $a^u b^v = (ab^j)^{n_u} z_2$ and by Lemma 2.1, $C_G(a^i b^{tp}) = C_G(a b^{sp})$ and $C_G(a^u b^v) = C_G(a b^j)$. Therefore, $(a^i b^{tp})(a^u b^v) = (a^u b^v)(a^i b^{tp})$ if an only if $(ab^{sp})(ab^j) = (ab^j)(ab^{sp})$ if and only if $b^{sp-j}a = ab^{sp-j}$. We now apply Lemma 2.2(a) to prove that the last equality is satisfied if and only if $p^2 \mid (sp - j)$ or $p \mid j$ which is a contradiction. Since $p \nmid j$,

the conjugacy classes of Type 6 is not adjacent with another one of Type 8 in the commuting conjugacy class graph.

- (7) Suppose that $a^{s_1p}b^{j_1}$ and $a^{s_2p}b^{j_2}$ are representatives of two conjugacy classes of Type 7 such that they are commuting to each other. By Lemma 3.1, $a^{s_1p}b^{j_1} =$ $(a^{t_1p}b)^{n_{j_1}}z_1$ and $a^{s_2p}b^{j_2} = (a^{t_2p}b)^{n_{j_2}}z_2$ and by Lemma 2.1, $C_G(a^{s_1p}b^{j_1}) = C_G(a^{t_1p}b)$ and $C_G(a^{s_2p}b^{j_2}) = C_G(a^{t_2p}b)$. Therefore, $(a^{s_1p}b^{j_1})(a^{s_2p}b^{j_2}) = (a^{s_2p}b^{j_2})(a^{s_1p}b^{j_1})$ if and only if $(a^{t_1p}b)(a^{t_2p}b) = (a^{t_2p}b)(a^{t_1p}b)$ if and only if $ba^{(t_2-t_1)p} = a^{(t_2-t_1)p}b$. By Lemma 2.2(a), the last equality is satisfied if and only if $t_1 = t_2$. This means that two conjugacy classes of Types 7 are adjacent in the commuting conjugacy class graph, when their centralizers is equal. Also, by Theorem 3.2, the number of centralizers of Type 7 is p-1. So, the conjugacy classes of this type can be divided into p-1 parts. Now by Table 1, the number of conjugacy classes of Type 7 is equal to $mp(p-1)^2$ and the graph structure of each part of Type 7 gives a complete subgraph of size mp(p-1). It is enough to find the relationship between this and other conjugacy classes of Type 8. To do this, we assume that $a^{sp}b^{j}$ and $a^{u}b^{v}$ are representatives of two conjugacy classes of Types 7 and 8, respectively such that they are commute to each other. By Lemma 3.1, $a^{sp}b^j = (a^{tp}b)^{n_j}z_1$ and $a^u b^v = (a^i b)^{n_v} z_2$. On the other hand, by Lemma 2.1, $C_G(a^{sp} b^j) = C_G(a^{tp} b)$ and $C_G(a^u b^v) = C_G(a^i b)$. Therefore, $(a^{sp} b^j)(a^u b^v) = (a^u b^v)(a^{sp} b^j)$ if and only if $(a^{tp}b)(a^{i}b) = (a^{i}b)(a^{tp}b)$ if and only if $a^{tp-i}b = ba^{tp-i}$. By Lemma 2.2(a), the last equality is satisfied if and only if $p^2 \mid (tp - i)$ or $p \mid i$ which is a contradiction. Thus, the conjugacy classes of Type 7 is not adjacent with a conjugacy class of Type 8 in the commuting conjugacy class graph.
- (8) Suppose $a^{i_1}b^{j_1}$ and $a^{i_2}b^{j_2}$ are the representatives of two conjugacy classes of Type 8 such that they are commuting together. By Lemma 3.1, $a^{i_1}b^{j_1} = (a^sb)^{n_{j_1}}z_1$ and $a^{i_2}b^{j_2} = (a^tb)^{n_{j_2}}z_2$ and by Lemma 2.1, $C_G(a^{i_1}b^{j_1}) = C_G(a^sb)$ and $C_G(a^{i_2}b^{j_2}) = C_G(a^tb)$. Therefore, $(a^{i_1}b^{j_1})(a^{i_2}b^{j_2}) = (a^{i_2}b^{j_2})(a^{i_1}b^{j_1})$ if and only if $(a^sb)(a^tb) = (a^tb)(a^sb)$ if and only if $a^{s-t}b = ba^{s-t}$. By Lemma 2.2(a), the last equality is satisfied if and only if s = t. This means that two conjugacy classes of Type 8 are adjacent in the commuting conjugacy class graph, when their centralizers is equal. Also, by Theorem 3.2, the number of centralizers of Type 8 is p(p-1) and so the conjugacy classes of this type can be divided into p(p-1) parts. On the other hand, by Table 1, the number of conjugacy classes of Type 8 is $mp^2(p-1)^2$ which proves that each part of Type 8 gives a clique of size mp(p-1) in the commuting conjugacy class of Type 8 signal.

8 are adjacent only with conjugacy classes of Type 5. Since the conjugacy classes of Types 5 and 8 can be divided into p-1 and p(p-1) parts, every part of Type 5 is adjacent with p parts of Type 8.

By above discussion in eight parts, the commuting conjugacy class graph of G is a connected graph with $n(p^2 - 1) + m(p^4 - p^2)$ vertices. Suppose \mathcal{M}_1 is a graph depicted in Figure 2. Therefore, the commuting conjugacy class graph of G is a \mathcal{M}_1 -join of graphs as follows:

$$\Gamma(G) = \mathcal{M}_1[\overbrace{K_{n(p-1)}, \cdots, K_{n(p-1)}}^{p+1}, \overbrace{K_{m(p^2-p)}, \cdots, K_{m(p^2-p)}}^{p^2+p}],$$

where $n = \frac{|Z|}{n}$ and $m = \frac{|Z|}{n^2}$. The graph $\Gamma(G)$ is depicted in Figure 3.



FIGURE 3. The Commuting Conjugacy Class Graph of G with $\frac{G}{Z(G)} \cong \mathbb{Z}_{p^2} \times \mathbb{Z}_{p^2}$.

In the following theorem, we study the commuting conjugacy class graph of G with this condition that $\frac{G}{Z(G)} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$, when the semidirect product is non-abelian. We recall that by Theorem 2.3, p is an odd prime number.

Theorem 4.2. Suppose p is an odd prime number and G is a group with center Z such that $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$ and $\frac{G}{Z}$ is non-abelian. Then the commuting conjugacy classes of G is as follows:

$$\mathcal{M}_2[\overbrace{K_{n(p-1)},\cdots,K_{n(p-1)}}^{p^2+p+1},K_{np(p-1)}],$$

where \mathcal{M}_2 is depicted in Figure 4. Here, $n = \frac{|Z|}{p}$.



FIGURE 4. The Graph \mathcal{M}_2 .

Proof. We first apply Equation 2.1 to deduce that

$$(4.2) b^j a^i Z = a^{ijp+i} b^j Z$$

which implies that

$$(4.3) b^j Z a^i Z = (a^p Z)^{ij} a^i Z b^j Z,$$

(4.4)
$$a^i Z b^j Z = (a^p Z)^{-ij} b^j Z a^i Z.$$

Choose elements x and g in $G \setminus Z$. By the proof of Theorem 3.2, $x = a^i b^j z_1$ and $g = a^u b^v z_2$, where $z_1, z_2 \in Z$, $0 \le i, j, u, v \le p^2 - 1$ and i, j, as well as u, v, are not simultaneously zero. In Equation 4.2, we put i = p and j = 1. Since $a^{p^2} \in Z$, $bZa^pZ = a^pZbZ$. Thus $a^pZ \in Z\left(\frac{G}{Z}\right)$ and by Equations 4.3 and 4.4,

So, there exists $z \in Z$ such that

g

(4.5)
$$(a^{u}b^{v})^{-1}(a^{i}b^{j})(a^{u}b^{v}) = a^{(uj-vi)p}a^{i}b^{j}z$$

Suppose that $x = a^i b^j z$ is an arbitrary element of G such that $0 \le i, j \le p^2 - 1$ and i, j are not simultaneously zero. To compute the number of non-central conjugacy classes of G and obtain their structure, we will consider three separate cases that all together can be divided into eight subcases:

a) $i \neq 0$ and j = 0. This case is separated into two subcases that $p \mid i$ and $p \nmid i$ as follows: 1) $p \mid i$. Suppose i = sp, where $1 \leq s \leq p - 1$. By Equation 4.5 and this fact that $a^{p^2} \in Z$, $(a^u b^v)^{-1}(a^{sp})(a^u b^v) = a^{-vsp^2}a^{sp}z_1 = a^{sp}z_2$ and so $(a^{sp})^G = \{a^{sp}z \mid z \in Z\}$. On the other hand, by Theorem 3.2, $|C_G(a^{sp})| = p^3|Z|$ and $|(a^{sp})^G| = p$. Thus, $(a^{sp})^G = \{a^{sp}z_1, \cdots, a^{sp}z_p\} = a^{sp}H$ in which $H = \{z_1, \cdots, z_p\} \subseteq Z$. Suppose |Z| > p and choose the elements $z_r \in Z \setminus H$ and an element in the form $a^{sp}z_r$. It is easy to see that $a^{sp}z_r \notin (a^{sp})^G$ and hence $(a^{sp}z_r)^G \neq (a^{sp})^G$. Furthermore, $C(a^{sp}z_r) = C(a^{sp})$ and so $|(a^{sp}z_r)^G| = |(a^{sp})^G| = p$. Also, $(a^{sp}z_{r_1})^G = (a^{sp})^G z_{r_1} = a^{sp}Hz_{r_1}$. Since $(a^{sp})^G \cap (a^{sp}z_{r_1})^G = \emptyset$, $H \cap Hz_{r_1} = \emptyset$ and $H \cup Hz_{r_1} \subseteq Z$. We now choose the element $z_{r_2} \in Z \setminus (H \cup Hz_{r_1})$ and continue this process to prove that $p \mid |Z|$. If $n = \frac{|Z|}{p}$ then there are n distinct conjugacy classes in the form of $(a^{sp}z_{r_1})^G, (a^{sp}z_{r_2})^G \cdots (a^{sp}z_{r_n})^G$. Since $1 \leq s \leq p - 1$, there are n(p-1) distinct conjugacy classes with $n = \frac{|Z|}{p}$ and each conjugacy class has p elements. 2) $p \nmid i$. By Equation 4.5,

(4.6)
$$(a^u b^v)^{-1} (a^i) (a^u b^v) = a^{-vip} a^i z$$

By Theorem 3.2, $|C_G(a^i)| = p^2 |Z|$ and so $|(a^i)^G| = p^2$. Note that $0 \le v \le p^2 - 1$ and so there are v' and k such that -vi = v'p + k and $0 \le k \le p - 1$. By Equation 4.6 and the fact that $a^{p^2} \in Z$,

(4.7)
$$(a^i)^G = \{ a^{kp} a^i z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z \}.$$

Set $A_i = \{a^{kp}a^iz \mid 0 \leq k \leq p-1, z \in Z\}$ and $H = \{z_{(0,l)} \mid 0 \leq l \leq p-1\}$. Obviously, $|A_i| = p|Z|$, $(a^i)^G \subseteq A_i$ and $H \subseteq Z$. If |Z| > p, then we choose an element $z_r \in Z \setminus H$ and an element $a^i z_r \in A_i$. It can be easily seen that $a^i z_r \notin (a^i)^G$ and $(a^i z_r)^G \neq (a^i)^G$. Since $C(a^i z_r) = C(a^i)$, $|(a^i z_r)^G| = |(a^i)^G| = p^2$ and $(a^i z_{r_1})^G = (a^i)^G z_{r_1}$. On the other hand, since $(a^i)^G \cap (a^i z_{r_1})^G = \emptyset$, $H \cap H z_{r_1} = \emptyset$ and $H \cup H z_{r_1} \subseteq Z$. We now choose an element $z_{r_2} \in Z \setminus (H \cup H z_{r_1})$ and continue this process to prove that $(a^i z_{r_2})^G \subseteq A_i$. This shows that $p^2 \mid |A_i|$. Put $n = \frac{|A_i|}{p^2} = \frac{|Z|}{p}$. Then, for each *i* there are *n* distinct conjugacy classes as $(a^i z_{r_1})^G, \cdots, (a^i z_{r_n})^G$. Again since $1 \leq i \leq p^2 - 1$ and $p \nmid i$, there are non-negative integers k' and i' such that i = k'p + i' and $1 \leq i' \leq p - 1$. By Equation 4.7,

$$(a^{i})^{G} = (a^{k'p+i'})^{G}$$

= $\{a^{kp}a^{k'p+i'}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}$
= $\{a^{(k+k')p}a^{i'}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}.$

Therefore, there exists $z_{i'} \in Z$ such that $(a^i)^G = (a^{i'}z_{i'})^G$. This proves that the number of distinct conjugacy classes is equal to n(p-1) in which $n = \frac{|Z|}{p}$ and each class has p^2 elements.

- b) i = 0 and $j \neq 0$. Again there are two cases that $p \mid j$ and $p \nmid j$.
 - 3) $p \mid j$. Suppose j = tp, where $1 \leq t \leq p-1$. By Equation 4.5 and the fact that a^{p^2} is an element of Z, we can conclude that $(a^u b^v)^{-1}(b^{tp})(a^u b^v) = a^{utp^2}b^{tp}z_1 = b^{tp}z_2$ and so $(b^{tp})^G = \{b^{tp}z \mid z \in Z\}$. By Theorem 3.2, $|C_G(b^{tp})| = p^3|Z|$ and hence $|(b^{tp})^G| = p$. Therefore, similar to the Case 1, there are n distinct conjugacy classes $(b^{tp}z_{r_1})^G, \cdots, (b^{tp}z_{r_n})^G$, where $n = \frac{|Z|}{p}$. So, the number of distinct conjugacy classes is equal to n(p-1) and each class has p elements.
 - 4) $p \nmid j$. Then by Equation 4.5, $(a^u b^v)^{-1}(b^j)(a^u b^v) = a^{ujp}b^j z$. By Theorem 3.2, $|C_G(b^j)| = p^2 |Z|$ and so $|(b^j)^G| = p^2$. A similar argument as in the Case 2 shows that $(b^j)^G = \{a^{kp}b^j z_{(k,l)} \mid 0 \leq k, l \leq p-1, z_{(k,l)} \in Z\}$. Thus, there are *n* distinct conjugacy classes $(b^j z_{r_1})^G, \dots, (b^j z_{r_n})^G$, when *j* is a fixed number and $n = \frac{|Z|}{p}$.

Since $1 \le j \le p^2 - 1$ and $p \nmid j$, the number of distinct conjugacy classes is equal to $n(p^2 - p)$ and each class has p^2 elements.

- c) $i \neq 0$ and $j \neq 0$. We have four subcases as follows:
 - 5) $p \mid i \text{ and } p \mid j$. Suppose i = sp and j = tp such that $1 \leq s, t \leq p-1$. By Equation 4.5 and the fact that $a^{p^2} \in Z$, $(a^u b^v)^{-1}(a^{sp}b^{tp})(a^u b^v) = a^{(utp-vsp)p}a^{sp}b^{tp}z_1$ $= a^{(ut-vs)p^2}a^{sp}b^{tp}z_1 = a^{sp}b^{tp}z_2$. Thus, $(a^{sp}b^{tp})^G = \{a^{sp}b^{tp}z \mid z \in Z\}$. Apply Theorem 3.2 to deduce that $|C_G(a^{sp}b^{tp})| = p^3|Z|$ which implies that $|(a^{sp}b^{tp})^G| = p$. Similar to the Case 1, for fixed non-negative integers s and t, there are n distinct conjugacy classes as $(a^{sp}b^{tp}z_{r_1})^G$, ..., $(a^{sp}b^{tp}z_{r_n})^G$ in which $n = \frac{|Z|}{p}$. By above discussion, we result that the number of the distinct conjugacy classes is equal to $n(p-1)^2$ and each class has p elements.
 - 6) $p \nmid i \text{ and } p \mid j$. Suppose j = tp, where $1 \leq t \leq p-1$. By Equation 4.5 and the fact that $a^{p^2} \in Z$, $(a^u b^v)^{-1}(a^i b^{tp})(a^u b^v) = a^{(utp-vi)p}a^i b^{tp} z_1 = a^{-vip}a^i b^{tp} z_2$. By Theorem 3.2, $|C_G(a^i b^{tp})| = p^2 |Z|$ and so $|(a^i b^{tp})^G| = p^2$. Now a similar argument as the Case 2,

(4.8)
$$(a^{i}b^{tp})^{G} = \{a^{kp}a^{i}b^{tp}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}.$$

Hence, for fixed positive integers i and t, there are n distinct conjugacy classes as $(a^i b^{tp} z_{r_1})^G$, ..., $(a^i b^{tp} z_{r_n})^G$ in which $n = \frac{|Z|}{p}$. Since $1 \le i \le p^2 - 1$ and $p \nmid i$, there are integers k' and i' such that i = k'p + i' and $1 \le i' \le p - 1$. By Equation 4.8,

$$(a^{i}b^{tp})^{G} = (a^{k'p+i'}b^{tp})^{G}$$

= $\{a^{kp}a^{k'p+i'}b^{tp}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}$
= $\{a^{(k+k')p}a^{i'}b^{tp}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}.$

This shows that there exists $z_{(i',t)} \in Z$ such that $(a^i b^{tp})^G = (a^{i'} b^{tp} z_{(i',t)})^G$. Since $1 \leq i', t \leq p-1$, the number of the distinct conjugacy classes is equal to $n(p-1)^2$ and each class has p^2 elements.

- 7) $p \mid i \text{ and } p \nmid j$. Suppose i = sp such that $1 \leq s \leq p-1$. By Equation 4.5 and the fact that $a^{p^2} \in Z$, we have $(a^u b^v)^{-1} (a^{sp} b^j) (a^u b^v) = a^{(uj-vsp)p} a^{sp} b^j z_1 = a^{ujp} a^{sp} b^j z_2$ = $a^{(uj+s)p} b^j z_2$. In this case, there is no new conjugacy class and the argument is similar to the Case 4.
- 8) $p \nmid i$ and $p \nmid j$. By Equation 4.5, $(a^u b^v)^{-1} (a^i b^j) (a^u b^v) = a^{(uj-vi)p} a^i b^j z$. By Theorem 3.2, $|C_G(a^i b^j)| = p^2 |Z|$ and so $|(a^i b^j)^G| = p^2$. Now by a similar argument as in the Case 2, we can see that

(4.9)
$$(a^{i}b^{j})^{G} = \{a^{kp}a^{i}b^{j}z_{(k,l)} \mid 0 \leq k, l \leq p-1, z_{(k,l)} \in Z\}.$$

Type	The I	Representative of Conjugacy Classes	$ x^G $	# Conjugacy Classes
1	a^{sp}	$1 \le s \le p-1$	p	n(p-1)
2	a^i	$1 \le i \le p-1$	p^2	n(p-1)
3	b^{tp}	$1 \le t \le p-1$	p	n(p-1)
4	b^j	$1 \leq j \leq p^2 - 1, j \not \equiv 0$	p^2	np(p-1)
5	$a^{sp}b^{tp}$	$1 \le s,t \le p-1$	p	$n(p-1)^2$
6	$a^i b^{tp}$	$1 \le i,t \le p-1$	p^2	$n(p-1)^2$
7	$a^{sp}b^j$	$1 \le s \le p-1, 1 \le j \le p^2 - 1, j \not\equiv 0$	Г	The same of Type 4
8	$a^i b^j$	$1 \le i \le p - 1, 1 \le j \le p^2 - 1, j \not\equiv 0$	p^2	$np(p-1)^{2}$

TABLE 2. Conjugacy Classes of G such that $n = \frac{|Z|}{p}$.

Hence, for fixed positive integers i and j, there are n distinct conjugacy classes $(a^i b^j z_{r_1})^G, \dots, (a^i b^j z_{r_n})^G$, where $n = \frac{|Z|}{p}$. Since $1 \le i \le p^2 - 1$ and $p \nmid i$, there are integers k' and i' such that i = k'p + i' and $1 \le i' \le p - 1$. Therefore, by Equation 4.9,

$$(a^{i}b^{j})^{G} = (a^{k'p+i'}b^{j})^{G}$$

= $\{a^{kp}a^{k'p+i'}b^{j}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}$
= $\{a^{(k+k')p}a^{i'}b^{j}z_{(k,l)} \mid 0 \le k, l \le p-1, z_{(k,l)} \in Z\}.$

Thus, there exists $z_{(i',j)} \in Z$ such that $(a^i b^j)^G = (a^{i'} b^j z_{(i',j)})^G$. Since $1 \leq i' \leq p-1$, $1 \leq j \leq p^2 - 1$ and $p \nmid j$, the number of distinct conjugacy classes is equal to $np(p-1)^2$ and each class has p^2 elements.

We now investigate the commuting conjugacy class graph of G. To do this, it is enough to determine the relationship between conjugacy classes in Table 2. We consider the following cases:

(1) Suppose a^{s_1p} and a^{s_2p} are the representatives of two classes of Type 1 in Table 2. It is clear that $a^{s_1p}a^{s_2p} = a^{s_2p}a^{s_1p}$ and so all conjugacy classes of Type 1 are commuting together. Hence, the commuting conjugacy class graph has a subgraph isomorphic to $K_{n(p-1)}$. We now determine the relationship between conjugacy classes of this and another types. For this purpose, we suppose a^{sp} is a representative of a conjugacy class of Type 1 and $a^u b^v z$ is an arbitrary element of G such that they are commuted to each other. This means that $(a^{sp})(a^u b^v z) =$

 $(a^u b^v z)(a^{sp})$ and hence $a^{sp}b^v = b^v a^{sp}$. By Lemma 2.2(a), the last equality is true if and only if $p \mid v$ or v = tp in which $0 \le t \le p - 1$. Then a^{sp} commutes with all elements of the form $a^u b^{tp} z$. Therefore, the following cases can be occurred:

- (a) t = 0. Then the conjugacy classes of Types 1 and 2 in Table 2 are commuting together.
- (b) $t \neq 0$ and u = 0. In this case, the conjugacy classes of Types 1 and 3 in Table 2 are commuting to each other.
- (c) $u, t \neq 0$ and $p \mid u$. Then the conjugacy classes of Types 1 and 5 in Table 2 are commuting together.
- (d) $t \neq 0$ and $p \nmid u$. By this condition, the conjugacy classes of Types 1 and 6 are commuting together.
- (2) Suppose a^{i_1} and a^{i_2} are the representatives of two conjugacy classes of Type 2. Obviously, $a^{i_1}a^{i_2} = a^{i_2}a^{i_1}$ and so all conjugacy classes of Type 2 are commuted to each other in the group. Hence, the commuting conjugacy class graph has a clique of size n(p-1). We now determine the relation between this and another types. To do this, we suppose that a^i is a representative of a conjugacy class of Type 2 and $a^u b^v z$ is an arbitrary element of G such that $(a^i)(a^u b^v z) = (a^u b^v z)(a^i)$. Thus, $a^i b^v = b^v a^i$. Since $p \nmid i$, then By Lemma (2.2)(a) the last equality is satisfied if and only if v = 0. Then a^i commutes with all elements of the form $a^u z$ which shows that the conjugacy classes of Type 2 are commuting only with conjugacy classes of Type 1.
- (3) Suppose b^{t_1p} and b^{t_2p} are representatives of two conjugacy classes of Type 3. It is clear that $b^{t_1p}b^{t_2p} = b^{t_2p}b^{t_1p}$. This shows that all conjugacy classes of Type 3 are commuting together and hence the commuting conjugacy class graph has a subgraph isomorphic to $K_{n(p-1)}$. We now determine the relation between this and another types. Suppose b^{tp} is a representatives of a conjugacy class of Type 3 and $a^u b^v z$ is an arbitrary element of G such that $(b^{tp})(a^u b^v z) = (a^u b^v z)(b^{tp})$. Thus, $b^{tp}a^u = a^u b^{tp}$. By Lemma 2.2(a), the last equality is satisfied if and only if $p \mid u$ or u = sp in which $0 \leq s \leq p - 1$. Then b^{tp} commutes with all elements in the form of $a^{sp}b^v z$. Again, the following cases can be occurred:
 - (a) s = 0. Then the conjugacy classes of Types 3 and 4 in Table 2 are commuting together.
 - (b) $s \neq 0$ and v = 0. Under this condition, the conjugacy classes of Types 3 and 1 in Table 2 are commuting together.

- (c) $v, s \neq 0$ and $p \mid v$. In this case, the conjugacy classes of Types 3 and 5 in Table 2 are commuting to each other.
- (d) $s \neq 0$ and $p \nmid v$. The classes classes of Types 3 and 7 in Table 2 are commuting together.
- (4) Suppose b^{j_1} and b^{j_2} are representatives of two conjugacy classes of Type 4. Since $b^{j_1}b^{j_2} = b^{j_2}b^{j_1}$, all conjugacy classes of Type 4 are commuting together and hence the commuting conjugacy class graph has a clique of size np(p-1). We now determine the relationship between this and another types. To do this, we assume that b^j is a representative of a conjugacy class of Type 4 and $a^u b^v z$ is an arbitrary element of G such that $(b^j)(a^u b^v z) = (a^u b^v z)(b^j)$. Thus, $b^j a^u = a^u b^j$. Since $p \nmid j$, by Lemma 2.2(a) the last equality is satisfied if and only if u = 0. Then b^j commutes with all elements in the form of $b^v z$. So, the conjugacy classes of Type 4 are commuting only with the conjugacy classes of Type 3.
- (5) Suppose $a^{s_1p}b^{t_1p}$ and $a^{s_2p}b^{t_2p}$ are representatives of two conjugacy classes of Type 5. By Equation 3.3, $(a^{s_1p}b^{t_1p})(a^{s_2p}b^{t_1p}) = (a^{s_2p}b^{t_2p})(a^{s_1p}b^{t_2p})$. This proves that all conjugacy classes of Type 5 are commuting to each other and hence the commuting conjugacy class graph has a clique of size $n(p-1)^2$. It is now enough to determine the relationship between this and conjugacy classes of types 6, 7 and 8. Suppose $a^{sp}b^{tp}$ and $a^{u}b^{vp}$ are representatives of conjugacy classes of Types 5 and 6, respectively, which are commuting together. By Lemma 3.1, $a^{sp}b^{tp} = (a^{ip}b^p)^{n_t}z_1$ and $a^u b^{vp} = (ab^{jp})^{n_u} z_2$. On the other hand, By Lemma 2.1, $C_G(a^{sp}b^{tp}) = C_G(a^{ip}b^p)$ and $C_G(a^u b^{vp}) = C_G(a b^{jp})$. We note that $(a^{sp} b^{tp})(a^u b^{vp}) = (a^u b^{vp})(a^{sp} b^{tp})$ if and only if $(a^{ip}b^p)(ab^{jp}) = (ab^{jp})(a^{ip}b^p)$ if and only if $b^p a = ab^p$, contradicts by Lemma 2.2(a). Therefore, a conjugacy class of Type 5 is not adjacent with a conjugacy class of Type 6. We now assume that $a^{sp}b^{tp}$ and $a^{u}b^{v}$ are representatives of the conjugacy classes of Types 5 and 8, respectively, which are commute to each other. By Lemma 3.1, $a^{sp}b^{tp} = (a^{ip}b^p)^{n_t}z_1$ and $a^ub^v = (a^jb)^{n_v}z_2$. On the other hand, By Lemma 2.1, $C_G(a^{sp}b^{tp}) = C_G(a^{ip}b^p)$ and $C_G(a^ub^v) = C_G(a^jb)$. By Equation 3.9, $(a^{sp}b^{tp})(a^ub^v) = (a^ub^v)(a^{sp}b^{tp})$ if and only if $(a^{ip}b^p)(a^jb) = (a^jb)(a^{ip}b^p)$ if and only if $j \equiv i \pmod{p}$. Since $1 \leq i \leq p-1$, the conjugacy classes of Types 5 and 8 can be divided into p-1 parts and hence each part of Type 5 gives a complete subgraph of size n(p-1).
- (6) Suppose $a^{i_1}b^{t_1p}$ and $a^{i_2}b^{t_2p}$ are representatives of two commuting conjugacy classes of Type 6. By Lemma 3.1, $a^{i_1}b^{t_1p} = (ab^{s_1p})^{n_{i_1}}z_1$ and $a^{i_2}b^{t_2p} = (ab^{s_2p})^{n_{i_2}}z_2$. On the other hand, by Lemma 2.1, $C_G(a^{i_1}b^{t_1p}) = C_G(ab^{s_1p})$ and $C_G(a^{i_2}b^{t_2p}) = C_G(ab^{s_2p})$.

This proves that $(a^{i_1}b^{t_1p})(a^{i_2}b^{t_2p}) = (a^{i_2}b^{t_2p})(a^{i_1}b^{t_1p})$ if and only if $(ab^{s_1p})(ab^{s_2p}) =$ $(ab^{s_2p})(ab^{s_1p})$ if and only if $ab^{(s_2-s_1)p} = b^{(s_2-s_1)p}a$. By Lemma 2.2(a), the last equality is satisfied if and only if $s_1 = s_2$ and so in the commuting conjugacy class graph two conjugacy classes of Type 6 are adjacent, when their centralizers is equal. Also, by Theorem 3.2, the number of centralizers of Type 6 is p-1. So, the conjugacy classes of this type can be divided into p-1 parts. On the other hand, by Table 2, the number of conjugacy classes of Type 6 are $n(p-1)^2$. Hence, each part of Type 6 gives a complete subgraph isomorphic to $K_{n(p-1)}$. It is now enough to find the relationship between conjugacy classes of Types 6 and 8. To see this, we assume that $a^i b^{tp}$ and $a^u b^v$ are representatives of conjugacy classes of Types 6 and 8, respectively, such that they are commuting together. By Lemma 3.1, $a^i b^{tp} = (ab^{sp})^{n_i} z_1$ and $a^u b^v = (ab^j)^{n_u} z_2$. On the other hand, by Lemma 2.1, $C_G(a^i b^{tp}) = C_G(a b^{sp})$ and $C_G(a^u b^v) = C_G(a b^j)$. Therefore, $(a^i b^{tp})(a^u b^v) =$ $(a^u b^v)(a^i b^{tp})$ if and only if $(ab^{sp})(ab^j) = (ab^j)(ab^{sp})$ if and only if $b^{sp-j}a = ab^{sp-j}$. By Lemma 2.2(a), the last equality is satisfied if and only if $p^2 \mid (sp - j)$ or $p \mid j$ which is impossible. This shows that in the commuting conjugacy class graph, the conjugacy classes of Type 6 is not adjacent with any conjugacy classes of Type 8. (7) Suppose $a^{i_1}b^{j_1}$ and $a^{i_2}b^{j_2}$ are representatives of two commuting conjugacy classes of

Type 8 such that they are commuting together. By Lemma 3.1, $a^{i_1}b^{j_1} = (a^sb)^{n_{j_1}}z_1$ and $a^{i_2}b^{j_2} = (a^tb)^{n_{j_2}}z_2$. On the other hand, by Lemma 2.1, $C_G(a^{i_1}b^{j_1}) = C_G(a^sb)$ and $C_G(a^{i_2}b^{j_2}) = C_G(a^tb)$. Therefore, $(a^{i_1}b^{j_1})(a^{i_2}b^{j_2}) = (a^{i_2}b^{j_2})(a^{i_1}b^{j_1})$ if and only if $(a^sb)(a^tb) = (a^tb)(a^sb)$ if and only if $a^{s-t}b = ba^{s-t}$. By Lemma 2.2(a), the last equality is satisfied if and only if s = t and so in the commuting conjugacy class graph two conjugacy classes of Type 8 are adjacent, when their centralizers is equal. Also, by Theorem 3.2, the number of centralizers of Type 8 is equal to p(p-1) and so the conjugacy classes of this type can be divided into p(p-1)parts. By Table 1, the number of conjugacy classes of Type 8 is $np(p-1)^2$ and hence each part of Type 8 gives a complete subgraph of size n(p-1). Therefore, we can see that in commuting conjugacy classes of Type 5. Finally, we know that the conjugacy classes of Types 5 and 8 can be divided into p-1 and p(p-1)parts, respectively. Therefore, every conjugacy class of Type 5 is adjacent with pconjugacy classes of Type 8.

By above discussion, the commuting conjugacy class graph of G is a connected graph with $n(p-1)(p+1)^2$ vertices. Suppose \mathcal{M}_2 is the graph depicted in Figure 3. Then the commuting conjugacy class graph of G can be written as a \mathcal{M}_2 -join, i.e.

$$\Gamma(G) = \mathcal{M}_2[K_{n(p-1)}, \cdots, K_{n(p-1)}, K_{np(p-1)}].$$

The graph $\Gamma(G)$ is depicted in Figure 5.



FIGURE 5. The Graph $\Gamma(G)$, when $\frac{G}{Z(G)}$ is non-abelian and $\frac{G}{Z} \cong \mathbb{Z}_{p^2} \rtimes \mathbb{Z}_{p^2}$

Acknowledgement. The research of the first author is partially supported by the University of Kashan under grant no. 364988/64.

References

- [1] A. R. Ashrafi, On finite groups with a given number of centralizers, Algebra Colloq. 7 (2000) 139–146.
- [2] R. Baer, Groups with preassigned central and central quotient groups, Trans. Amer. Math. Soc. 44 (1938) 387–412.
- [3] S. M. Belcastro and G. J. Sherman, Counting centralizers in finite groups, Math. Mag. 5 (1994) 111–114.
- [4] R. Brauer and K. A. Fowler. On groups of even order. Ann. of Math. 62(3)(1955) 565-583.
- [5] W. Burnside, Theory of Groups of Finite Order, Cambridge University Press, Cambridge, 1897.
- [6] M. Herzog, P. Longobardi and M. Maj, On a commuting graph on conjugacy classes of groups, Comm. Algebra 37(10)(2009) 3369–3387.
- [7] A. Magidin, Capable 2-generator 2-groups of class 2, Comm. Algebra 34 (2006) 2183–2193.

- [8] A. Mohammadian, A. Erfanian, M. Farrokhi D. G. and B. Wilkens, Triangle-free commuting conjugacy class graphs, J. Group Theory 19 (3) (2016) 1049–1061.
- [9] D. J. Robinson, A Course in the Theory of Groups, Springer-Verlag, New York, 1996.
- [10] G. Sabidussi, Graph derivatives, Math. Z. 76 (1961) 385–401.
- [11] M. A. Salahshour and A. R. Ashrafi, Commuting conjugacy class graph of finite CA-groups, *Khayyam J. Math.* 6 (1) (2020) 108–118.
- [12] The GAP Team, Group, GAP Groups, Algorithms, and Programming, Version 4.5.5, 2012, http://www.gap-system.org.
- [13] R. Zainal, N. M. Mohd Ali, N. H. Sarmin and S. Rashid, On the capability of nonabelian groups of order p⁴, Proceedings of the 21st National Symposium on Mathematical Sciences (SKSM21), AIP Conf. Proc. 1605 (2014) 575–579.

Ali Reza Ashrafi: DEPARTMENT OF PURE MATHEMATICS, FACULTY OF MATHEMATICAL SCI-ENCES, UNIVERSITY OF KASHAN, KASHAN 87317–53153, I. R. IRAN

Mohammad Ali Salahshour: DEPARTMENT OF MATHEMATICS, SAVADKOOH BRANCH, ISLAMIC Azad University, Savadkooh, I. R. Iran