

MEMBRANES WITH THIN AND HEAVY INCLUSIONS: ASYMPTOTICS OF SPECTRA

YURIY GOLOVATY

ABSTRACT. We study the asymptotic behaviour of eigenvalues of 2D vibrating systems with mass density perturbed in a vicinity of closed curves. The threshold case in which the resonance frequencies of the membrane and the frequencies of thin inclusion coincide is investigated. The perturbed eigenvalue problem can be realized as a family of self-adjoint operators acting on varying Hilbert spaces. However the so-called limit operator is non-self-adjoint and possesses the Jordan chains of length 2. Apart from the lack of self-adjointness, the operator has non-compact resolvent. As a consequence, its spectrum has a complicated structure, for instance, the spectrum contains a countable set of eigenvalues with infinite multiplicity. The complete asymptotic analysis of eigenvalues has been carried out.

1. INTRODUCTION AND STATEMENT OF PROBLEM

The mechanical systems with strongly inhomogeneous mass distributions have become the subject of intensive experimental and theoretical studies since the time of Poisson and Bessel [1, Ch.2], and a lot of research has been devoted to the analysis of vibrating systems with so-called added masses. Historically, the first relevant mathematical models in classical mechanics go back to the first half of the 20th century (see e.g. [2] and the references given there). Many authors have investigated properties of strings and rods with the mass densities perturbed by finite or infinite sums $\sum M_k \delta(x - x_k)$, where δ is Dirac's delta-function and M_k is an added mass at the point x_k . Recently, such models in dimensions two and three with heavy inclusions of a different geometry are widely used not only in mechanics, but in various fields of science and technology such as physics of liquid crystals, physical chemistry of polymers, micelles and microemulsions, molecular theory, cell membrane theory [3–5]. For instance, cell membranes are known to contain embedded proteins and various colloidal particles [6].

In higher dimensions, the perturbation of mass densities by the δ -functions often leads to incorrect mathematical models, because the formal differential equations which appear have no mathematical meaning. As an example of such ill-posed problem we can consider the eigenvalue problem for the Laplace operator

$$-\Delta u = \lambda(1 + M\delta(x))u \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega,$$

where Ω is a bounded domain in \mathbb{R}^3 containing the origin. The equation has no non-trivial solution, because any such solution u has a singularity at $x = 0$ and therefore the product $\delta(x)u(x) = u(0)\delta(x)$ is not defined. The new and at the

2020 *Mathematics Subject Classification.* 35B25, 35P05, 74K15.

Key words and phrases. Asymptotics of eigenvalues, eigenvalue of infinite multiplicity, quasi-mode, non-self-adjoint operator, concentrated mass, singular perturbation.

same time obvious idea was instead to replace the δ -function with its regularization $\varepsilon^{-3}q(x/\varepsilon)$, where q is a function of compact support, and study the asymptotic behaviour of eigenvalues and eigenfunctions as $\varepsilon \rightarrow 0$. The problem was first investigated by E. Sánchez-Palencia [7, 8], who proved the existence of the so-called local eigenvibrations: the eigenfunctions are significant in a small neighborhood of the origin only.

The model in which the density is perturbed by $\sum M_k \delta(x - x_k)$ is not adequate even in the one-dimensional case, when dealing with the large masses M_k . The very heavy inclusions cause a strong local reaction of vibrating system, but this phenomenon can not be described on the discrete set which is a support of the sum of Dirac's functions. The geometry of small domains where the large masses are loaded should also have an effect on the form of eigenvibrations. In [9], asymptotic analysis was applied to a spectral problem for the Sturm-Liouville operator with weight function of the form $\rho_\varepsilon(x) = \rho(x) + \varepsilon^{-m}q(x/\varepsilon)$, where q is a function of compact support and $m \in \mathbb{R}$. For the case $m = 1$ the perturbation is a δ -like sequence, but the most interesting cases of the limit behaviour of eigenfunctions as $\varepsilon \rightarrow 0$ are those when the power m is greater than 1.

These advanced models have attracted considerable attention in the mathematical literature over the last three decades (see e.g. [10] for a review). The spectral properties of differential operators with weight functions having the form

$$\rho_\varepsilon(x) = \rho(x) + \sum \varepsilon^{-m_k} q_{k,\varepsilon}(x),$$

where $q_{k,\varepsilon}$ are compactly supported in vicinity of different sets, have been investigated in numerous articles. We mention here [11] for the Laplace operator in dimension 3, [12, 13] for ordinary differential operators of the fourth order and the biharmonic operator, [14–16] for boundary value problems on junctions of a very complicated geometry, [17, 18] for the Sturm-Liouville operators on metric graphs. The spectral properties of strings with rapidly oscillating and periodic densities have been treated in [19–21] in the framework of homogenization theory. Another model in which the heavy inclusions were regarded as rigid ones has been studied in [22]. Since the 90s of the last century, a series of papers was published concerning 2D and 3D elastic systems with many concentrated masses near the boundary [23–30]. New asymptotic results for the spectral problems in domains surrounded by thin stiff and heavy bands, when the mass density and stiffness are simultaneously perturbed in a neighbourhood of the boundary, were obtained in [31–34]. The Neumann eigenvalue problem for a membrane, almost the entire mass of which is concentrated around the boundary, was studied in [35].

In this paper we study the eigenvibration characteristics of a membrane with heavy and thin inclusion inside. Let Ω be a smooth bounded domain of \mathbb{R}^2 , and let $\gamma \subset \Omega$ be a smooth closed curve. We will denote by ω_ε the ε -neighborhood of γ (i.e., the union of all open balls of radius ε around a point on γ). Consider $\varepsilon > 0$ sufficiently small such that $\omega_\varepsilon \subset \Omega$ and the boundary $\partial\omega_\varepsilon$ is smooth. Assume ρ is a smooth uniformly positive functions in Ω , and

$$\rho_\varepsilon(x) = \begin{cases} \rho(x), & \text{if } x \in \Omega \setminus \omega_\varepsilon, \\ \varepsilon^{-2}q_\varepsilon(x), & \text{if } x \in \omega_\varepsilon. \end{cases} \quad (1.1)$$

To specify explicit dependence of q_ε on ε we introduce the Fermi normal coordinates in ω_ε (see Fig. 1). Let $\alpha: [0, |\gamma|) \rightarrow \mathbb{R}^2$ be the unit-speed smooth parametrization




FIGURE 1. Membrane with heavy and thin inclusion

of γ with the natural parameter s , and $|\gamma|$ is the length of γ . Then the vector $\nu = (-\dot{\alpha}_2, \dot{\alpha}_1)$ is a unit normal on γ . Set $x = \alpha(s) + r\nu(s)$ for $(s, r) \in [0, |\gamma|] \times (-\varepsilon, \varepsilon)$, where r is the signed distance from x to γ . Suppose that

$$q_\varepsilon(x) = q\left(\frac{r}{\varepsilon}\right),$$

where q is a smooth positive function in $[-1, 1]$. We see that the perturbation of mass density varies on the normal direction to γ only. Let us consider the spectral problem

$$-\Delta u_\varepsilon + au_\varepsilon = \lambda^\varepsilon \rho_\varepsilon u_\varepsilon \quad \text{in } \Omega, \quad \ell u_\varepsilon = 0 \quad \text{on } \partial\Omega, \quad (1.2)$$

where $a \in C^\infty(\bar{\Omega})$ and $\ell v = 0$ denotes the Dirichlet, Neumann or Robin boundary conditions on $\partial\Omega$. Our goal is to describe the asymptotic behaviour of the eigenvalues λ^ε of (1.2) as $\varepsilon \rightarrow 0$.

This paper is a continuation of [36, 37], where the problem with the Dirichlet boundary condition and more general perturbation $\varepsilon^{-m}q_\varepsilon$ of mass density has been treated using the variational approach and the case $m = 3$ has been completely investigated by asymptotic methods. Also, it has been shown that there exist five different limit behaviours for the spectrum and eigenspaces depending on m : $m < 1$, $m = 1$, $1 < m < 2$, $m = 2$ and $m > 2$. These cases differ by the form of eigenvibrations and place on the membrane where the main part of their “energy” is concentrated in the limit. From the mathematical viewpoint, the difference between the cases is that the spectral parameter in the limit eigenvalue problems appears alternately both in a differential equation on Ω and in an equation on the strip $\gamma \times (-1, 1)$, which is a dilatation of ω_ε , and even in coupling conditions on γ .

The threshold case $m = 2$ is the most difficult and interesting one to analyse, because the spectral parameter is included simultaneously in two differential equations, and the limit spectral problem is associated with a non-self-adjoint operator. The main insight of the present work is to exhibit this non-self-adjoint operator, its spectrum and generalized eigenspaces.

To better understand what happens in dimension 2, we briefly discuss the similar model for a string. Recently, we have revised some results from [9] concerning the case $m = 2$. In [38], we have studied the limiting behavior, as $\varepsilon \rightarrow 0$, of eigenvalues

and eigenfunctions of the problem

$$-y''_\varepsilon + Qy_\varepsilon = \lambda^\varepsilon \rho_\varepsilon y_\varepsilon, \quad x \in (a, b), \quad (1.3)$$

$$y_\varepsilon(a) \cos \alpha + y'_\varepsilon(a) \sin \alpha = 0, \quad y_\varepsilon(b) \cos \beta + y'_\varepsilon(b) \sin \beta = 0 \quad (1.4)$$

with the weight function ρ_ε given by (1.1), where $\Omega = (a, b)$ is an interval containing the origin, and $\omega_\varepsilon = (-\varepsilon, \varepsilon)$. For each real α and β the problem can be associated with a family of self-adjoint operators S_ε in the weighted space $L_2(\rho_\varepsilon, \Omega)$. The operators S_ε are defined by $S_\varepsilon \phi = \rho_\varepsilon^{-1}(-\phi'' + Q\phi)$ on functions $\phi \in W_2^2(\Omega)$ obeying boundary conditions (1.4). The spectra of S_ε are real, discrete and simple. It is worth noting that the study of operator families acting on varying spaces entails some mathematical difficulties. First of all, the question arises how to understand the convergence of such families. Next, if these operators do converge in some sense, does this convergence implies the convergence of their spectra (see, e.g., [19, III.1], [39, 40]). By abandoning the self-adjointness, we have realized (1.3), (1.4) as a family of non-self-adjoint matrix operators \mathcal{A}_ε acting on the fixed Hilbert space $L_2(\rho, (a, b)) \times L_2(q, (-1, 1))$ (see details in [38]). The operators are certainly similar to self-adjoint ones and their resolvents are compact. Moreover the spectra of \mathcal{A}_ε and S_ε coincide and the corresponding eigenspaces are isomorphic. It has been proved that \mathcal{A}_ε converge in the norm resolvent sense (the resolvents of \mathcal{A}_ε converge in the uniform norm as $\varepsilon \rightarrow 0$) to the matrix operator \mathcal{A} associated with the eigenvalue problem

$$\begin{aligned} -v'' + Qv &= \lambda \rho v \text{ in } (a, 0) \cup (0, b), & -w'' &= \lambda q w \text{ in } (-1, 1), \\ w'(-1) &= 0, & w'(1) &= 0, & v(-0) &= w(-1), & v(+0) &= w(1), \\ v(a) \cos \alpha + v'(a) \sin \alpha &= 0, & v(b) \cos \beta + v'(b) \sin \beta &= 0. \end{aligned}$$

Surprisingly enough, \mathcal{A} is not similar to a self-adjoint one, because it possesses multiple eigenvalues with the Jordan chains of length 2. In view of [41, Ch.1], [42], the norm resolvent convergence $\mathcal{A}_\varepsilon \rightarrow \mathcal{A}$ implies the “number-by-number” convergence of the corresponding eigenvalues and some results on the convergence of eigenspaces.

Problem (1.2) can also be associated with a family of matrix operators acting on the same Hilbert space. However, in two dimensions there is no uniform convergence of resolvents, because the limit operator \mathcal{P} which can be obtained in the strong resolvent topology has non-compact resolvent [43, Theorem IV.2.26]. It is known that the strong resolvent convergence of operators does not imply the convergence of their spectra. For this reason, we study the convergence of eigenvalues of (1.2) without touching on the convergence of corresponding operators. The operator \mathcal{P} is non-self-adjoint and possesses nontrivial Jordan chains. In Sec. 3, we describe the spectrum of \mathcal{P} and the generalized eigenspaces. The spectrum is discrete and real; it consists of eigenvalues of three types: (a) eigenvalues of finite multiplicity with generalized eigenspaces generated by eigenvectors only; (b) infinite-fold eigenvalues with generalized eigenspaces generated by eigenvectors only; (c) infinite-fold eigenvalues with generalized eigenspaces containing both eigenvectors and generalized eigenvectors. In fact, this spectrum is a union of spectra of two operators that correspond to two different parts of the vibrating system, namely, the membrane clamped along the curve γ and the thin heavy inclusion. Using the method of quasimodes we describe the asymptotic behaviour of eigenvalues of (1.2) as perturbation of $\sigma(\mathcal{P})$. In Sec. 4 and 5, we prove the existence of a countable number

of eigenvalues with the asymptotics $\lambda_{\varepsilon,k} = \lambda + \varepsilon\lambda_1^{(k)} + O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$, $k \in \mathbb{N}$, where λ is an eigenvalue of type (b) or (c). The set of correctors $\{\lambda_1^{(k)}\}_{k=1}^\infty$ is a spectrum of some pseudodifferential operator N_λ acting in $L_2(\gamma)$. In addition, if λ is an eigenvalue of type (c), there also exists $2d$ eigenvalues with the half-integer power asymptotics $\mu_{\varepsilon,j}^\pm = \lambda \pm \varepsilon^{1/2}\lambda_{1/2,j} + \varepsilon\lambda_{1,j}^\pm + O(\varepsilon^{3/2})$, as $\varepsilon \rightarrow 0$, where d is a number of Jordan chains of length 2 corresponding to λ . In Sec. 6 we prove that there exist K eigenvalues of (1.2) with the asymptotics $\lambda_{\varepsilon,k} = \lambda + \varepsilon\lambda_1^{(k)} + O(\varepsilon^2)$, where λ is an eigenvalue of type (a) with multiplicity K .

2. FORMAL CONSTRUCTION OF THE LIMIT OPERATOR

It will be convenient to parameterize the curve γ by points of a circle. It will allow us not to indicate every time that functions on γ are periodic on s . Let S be the circle of the length $|\gamma|$. Then ω_ε is diffeomorphic to the cylinder $S \times (-\varepsilon, \varepsilon)$. We also set $\omega = S \times (-1, 1)$. Here and subsequently, writing “in ω ” after an equation, we mean that the equation is considered in the rectangle $(0, |\gamma|) \times (-1, 1)$ and the corresponding solution is a $|\gamma|$ -periodic function on s .

Let us denote by γ_t the curve that is obtained from γ by flowing for “time” t along the normal vector field, i.e., $\gamma_t = \{x \in \mathbb{R}^2: x = \alpha(s) + t\nu(s), s \in S\}$. Then the boundary of ω_ε consists of two curves $\gamma_{-\varepsilon}$ and γ_ε . The domain Ω is divided by γ into subdomains Ω_- and Ω_+ . Suppose that $\partial\Omega_- = \partial\Omega \cup \gamma_-$ and $\partial\Omega_+ = \gamma_+$, where γ_- and γ_+ are two edges of the cut γ . In the sequel, the coordinate r increases in the direction from Ω^- to Ω^+ .

For any $\varepsilon > 0$, problem (1.2) admits a self-adjoint operator realization in the weighted Lebesgue space $L^2(\rho_\varepsilon, \Omega)$. We introduce operator T_ε defined by

$$T_\varepsilon\phi = \rho_\varepsilon^{-1}(-\Delta\phi + a\phi)$$

on functions $\phi \in W_2^2(\Omega)$ obeying the boundary condition $\ell\phi = 0$ on $\partial\Omega$. Obviously, the spectrum of T_ε is real and discrete.

We look for the approximation to the eigenvalue λ^ε and the corresponding eigenfunction u_ε of (1.2) in the form

$$\lambda^\varepsilon \sim \lambda + \dots, \quad u_\varepsilon(x) \sim \begin{cases} v(x) + \dots & \text{if } x \in \Omega \setminus \omega_\varepsilon, \\ w(s, \frac{r}{\varepsilon}) + \dots & \text{if } x \in \omega_\varepsilon. \end{cases} \quad (2.1)$$

Since u_ε solves (1.2) and the domain ω_ε shrinks to γ as $\varepsilon \rightarrow 0$, the function v must be a solution of the equation $-\Delta v + av = \lambda\rho v$ in $\Omega \setminus \gamma$ that satisfies the boundary condition $\ell v = 0$ on $\partial\Omega$. Of course, v must also satisfy appropriate transmission conditions on γ . To find these conditions, we must examine more closely the equation in (1.2) in a vicinity of γ .

Returning to the local coordinates (s, r) , we see that the vectors $\alpha = (\dot{\alpha}_1, \dot{\alpha}_2)$, $\nu = (-\dot{\alpha}_2, \dot{\alpha}_1)$ give the Frenet frame for γ . The Jacobian of transformation

$$x_1 = \dot{\alpha}_1(s) - r\dot{\alpha}_2(s), \quad x_2 = \dot{\alpha}_2(s) + r\dot{\alpha}_1(s)$$

has the form $J(s, r) = 1 - r\kappa(s)$, where $\kappa = \det(\dot{\alpha}, \ddot{\alpha})$ is the signed curvature of γ . We see that J is positive for sufficiently small r , because the curvature κ is bounded on γ . In addition, the Laplace-Beltrami operator becomes

$$\Delta\phi = J^{-1}(\partial_s(J^{-1}\partial_s\phi) + \partial_r(J\partial_r\phi)).$$

In the local coordinates (s, n) , where $n = r/\varepsilon$, the Laplacian can be written as

$$\Delta = (1 - \varepsilon n \varkappa)^{-1} (\varepsilon^{-2} \partial_n (1 - \varepsilon n \varkappa) \partial_n + \partial_s ((1 - \varepsilon n \varkappa)^{-1} \partial_s)).$$

From this we readily deduce the representation

$$\Delta = \varepsilon^{-2} \partial_n^2 - \varepsilon^{-1} \varkappa \partial_n - n \varkappa^2 \partial_n + \partial_s^2 + \varepsilon (2n \varkappa \partial_s^2 + n \varkappa' \partial_s - n^2 \varkappa^3 \partial_n) + \varepsilon^2 P_\varepsilon, \quad (2.2)$$

where P_ε is a PDE of the second order on s and the first one on n whose coefficients are uniformly bounded in ω with respect to ε . Then using this representation and (2.1) we obtain the equation $-\partial_n^2 w = \lambda q w$ in ω . Next, the eigenfunction u_ε as an element of $\text{dom } T_\varepsilon$ satisfies the conditions

$$[u_\varepsilon]_{-\varepsilon} = 0, \quad [u_\varepsilon]_\varepsilon = 0, \quad [\partial_r u_\varepsilon]_{-\varepsilon} = 0, \quad [\partial_r u_\varepsilon]_\varepsilon = 0, \quad (2.3)$$

where $[\cdot]_t$ stands for the jump of a function across γ_t . These conditions imply $v^\pm = w(\cdot, \pm 1)$ and $\partial_n w(\cdot, \pm 1) = 0$, where v^\pm denote the one-side traces of v on γ , i.e., $v^\pm = v|_{\gamma_\pm}$. Combining the latter equalities, we can formally deduce that the pair (v, w) must be an eigenvector of the spectral problem

$$-\Delta v + av = \lambda \rho v \quad \text{in } \Omega \setminus \gamma, \quad \ell v = 0 \quad \text{on } \partial\Omega, \quad (2.4)$$

$$-\partial_n^2 w = \lambda q w \quad \text{in } \omega, \quad \partial_n w(s, -1) = 0, \quad \partial_n w(s, 1) = 0, \quad (2.5)$$

$$v^- = w(s, -1), \quad v^+ = w(s, 1), \quad s \in S \quad (2.6)$$

with the spectral parameter λ ; (2.4)-(2.6) will be regarded as the *limit problem*.

We will show that the problem can be associated with a non-self-adjoint matrix operator. Moreover this operator is not similar to a self-adjoint one, because it possesses generalized eigenvectors. It means that the limit problem admits no self-adjoint realization. It is worth mentioning that the similar non-self-adjoint problem appeared in [31], where the spectral problem for a membrane surrounded by a thin stiff band was studied. Actually perturbed spectral problem (1.2) and the problem in [31] give us nontrivial examples of self-adjoint operators acting on varying Hilbert spaces that converge in some sense to a non-self-adjoint operator.

3. PROPERTIES OF THE LIMIT OPERATOR

We use the following notation. The spectrum and resolvent set of a linear operator T are denoted by $\sigma(T)$ and $\varrho(T)$, respectively. Let T^* denote the adjoint operator of T . For any $z \in \varrho(T)$, $R_z(T) = (T - zI)^{-1}$ is the resolvent operator. Here and subsequently, I is an identity operator.

3.1. Spectrum of the limit operator. We introduce the operators

$$\mathring{A} = \rho^{-1}(-\Delta + a) \quad \text{in } L_2(\rho, \Omega), \quad \text{dom } \mathring{A} = \{f \in W_2^2(\Omega \setminus \gamma) : \ell f = 0 \text{ on } \partial\Omega\},$$

$$B = -q^{-1} \partial_n^2 \quad \text{in } L_2(q, \omega), \quad \text{dom } B = \left\{ g \in W_2^{2,0}(\omega) : \partial_n g(\cdot, -1) = \partial_n g(\cdot, 1) = 0 \right\},$$

where $W_2^{2,0}(\omega)$ is the anisotropic Sobolev space

$$W_2^{2,0}(\omega) = \{g \in L_2(\omega) : \partial_n^k g \in L_2(\omega) \text{ for } k = 1, 2\}.$$

In the space $\mathcal{L} = L_2(\rho, \Omega) \times L_2(q, \omega)$ we consider the matrix operator

$$\mathcal{P} = \begin{pmatrix} \mathring{A} & 0 \\ 0 & B \end{pmatrix}, \quad \text{dom } \mathcal{P} = \{(f, g) \in \text{dom } \mathring{A} \times \text{dom } B :$$

$$f^- = g(\cdot, -1), \quad f^+ = g(\cdot, 1)\}.$$

Now (2.4)–(2.6) can be written as $\mathcal{P}u = \lambda u$ with the notation $u = (v, w)^\top$. For simplicity of notation we often write (v, w) instead of $(v, w)^\top$. Note that the operator \mathcal{P} is non-self-adjoint. Direct computations show that

$$\mathcal{P}^* = \begin{pmatrix} A & 0 \\ 0 & \mathring{B} \end{pmatrix}, \quad \text{dom } \mathcal{P}^* = \{(f, g) \in \text{dom } A \times W_2^{2,0}(\omega) : \\ \partial_r f^- = -\partial_n g(\cdot, -1), \quad \partial_r f^+ = \partial_n g(\cdot, 1)\},$$

where A is the restrictions of \mathring{A} to $\text{dom } A = \{f \in \text{dom } \mathring{A} : f = 0 \text{ on } \gamma\}$ and \mathring{B} is the extension of B to the whole space $W_2^{2,0}(\omega)$. Here $\partial_r f^\pm$ stand for the one-side traces of the normal derivative of f on γ .

Lemma 3.1. *The spectrum of B consists of a countable set of real eigenvalues of infinite multiplicity. Moreover, λ belongs to $\sigma(B)$ if and only if λ is an eigenvalue of the Sturm-Liouville problem*

$$y'' + \lambda q(n)y = 0, \quad n \in (-1, 1), \quad y'(-1) = 0, \quad y'(1) = 0. \quad (3.1)$$

Proof. Obviously, the operator B is self-adjoint. For given $\lambda \in \mathbb{C}$ and $g \in L_2(q, \omega)$, the equation $(B - \lambda I)\phi = g$ can be treated as the boundary value problem for the ordinary differential equation

$$-(\partial_n^2 + \lambda q(n))\phi = q(n)g(s, n) \text{ in } \omega, \quad \partial_n \phi(s, -1) = 0, \quad \partial_n \phi(s, 1) = 0$$

with the parameter $s \in S$. If λ is not an eigenvalue of (3.1), then the problem has a unique solution $\phi(s, \cdot)$ for each $g(s, \cdot) \in L_2(-1, 1)$ and almost all $s \in S$. Moreover ϕ belongs to $L_2(q, \omega)$ in view of its integral representation via the Green function. Otherwise, if λ is an eigenvalue of (3.1) with eigenfunction y , the problem is unsolvable for some right-hand sides g , because the corresponding homogeneous problem has infinitely many linearly independent solutions of the form $b(s)y(n)$, where $b \in L_2(\gamma)$. Therefore the spectrum of B consists of all eigenvalues of (3.1), and the corresponding eigenspaces are infinite-dimensional. \square

Theorem 3.2. *The operator \mathcal{P} has real discrete spectrum. Moreover*

$$\sigma(\mathcal{P}) = \sigma(A) \cup \sigma(B).$$

Proof. Let us construct the resolvent of \mathcal{P} in an explicit form. Given $\mu \in \mathbb{C}$, $f \in L_2(\rho, \Omega)$ and $g \in L_2(q, \omega)$, we write $(\mathring{A} - \mu I)v = f$, $(B - \mu I)w = g$. The second equation admits a unique solution $w = R_\mu(B)g$ if $\mu \in \varrho(B)$, and then v is a solution of the problem

$$\begin{aligned} -\Delta v + av - \mu \rho v &= \rho f \quad \text{in } \Omega \setminus \gamma, \quad \ell v = 0 \quad \text{on } \partial\Omega, \\ v^- &= w(\cdot, -1), \quad v^+ = w(\cdot, 1). \end{aligned} \quad (3.2)$$

Suppose the operator $T(\mu) : W_2^2(\omega) \rightarrow L_2(\rho, \Omega)$ solves the problem

$$\begin{aligned} -\Delta \phi + a\phi - \mu \rho \phi &= 0 \quad \text{in } \Omega \setminus \gamma, \quad \ell \phi = 0 \text{ on } \partial\Omega, \\ \phi^- &= \psi(\cdot, -1), \quad \phi^+ = \psi(\cdot, 1) \end{aligned} \quad (3.3)$$

for a given function $\psi \in W_2^2(\omega)$. If $\mu \in \varrho(A)$, then the problem has a unique solution $\phi = T(\mu)\psi$ and hence $T(\mu)$ is bounded. Next, v can be represented as the sum of a solution of (3.2) subject to the homogeneous boundary conditions on γ_\pm and a solution of (3.3) with $\psi = w$. Therefore

$$v = R_\mu(A)f + T(\mu)w = R_\mu(A)f + T(\mu)R_\mu(B)g,$$

provided $\mu \in \varrho(A) \cap \varrho(B)$. Then the resolvent of \mathcal{P} can be written in the form

$$R_\mu(\mathcal{P}) = \begin{pmatrix} R_\mu(A) & T(\mu)R_\mu(B) \\ 0 & R_\mu(B) \end{pmatrix}.$$

The equality $\sigma(\mathcal{P}) = \sigma(A) \cup \sigma(B)$ follows directly from this representation and the fact that $T(\mu)$ is bounded for $\mu \in \varrho(A)$. Obviously, $\sigma(A) \cup \sigma(B) \subset \sigma(\mathcal{P})$. Suppose, contrary to our claim, that $\sigma(\mathcal{P})$ is not contained in $\sigma(A) \cup \sigma(B)$. Then there exists $\mu \in \sigma(\mathcal{P})$ such that $\mu \in \rho(A) \cap \rho(B)$. Hence, the operators $R_\mu(A)$, $R_\mu(B)$ and $T(\mu)$ are bounded, and therefore, the operator $R_\mu(\mathcal{P})$ is also bounded, a contradiction. Clearly, $\sigma(\mathcal{P})$ is real and discrete, because A and B are self-adjoint operators with discrete spectra. \square

Corollary 3.3. *The operator \mathcal{P} has non-compact resolvent.*

Proof. The entry $R_\mu(B)$ of the matrix $R_\mu(\mathcal{P})$ is a non-compact operator, which is clear from Lemma 3.1. The main reason for which $R_\mu(B)$ is not compact is that the second derivative ∂_n^2 is a non-elliptic PDE in the two-dimensional domain ω . \square

Remark 3.4. The operator A can be represent as the direct sum $A_- \oplus A_+$, where

$$\begin{aligned} A_- &= \rho^{-1}(-\Delta + a), \quad \text{dom } A_- = \{f \in W_2^2(\Omega_-) : \ell f = 0 \text{ on } \partial\Omega, f = 0 \text{ on } \gamma\}, \\ A_+ &= \rho^{-1}(-\Delta + a), \quad \text{dom } A_+ = \{f \in W_2^2(\Omega_+) : f = 0 \text{ on } \gamma\}. \end{aligned}$$

Hence $\sigma(\mathcal{P}) = \sigma(A_-) \cup \sigma(A_+) \cup \sigma(B)$.

3.2. Structure of generalized eigenspaces. Let X_μ be the generalized eigenspace corresponding to $\mu \in \sigma(\mathcal{P})$, i.e., $X_\mu = \{h \in \mathcal{L} : h \in \ker(\mathcal{P} - \mu I)^k \text{ for some } k \in \mathbb{N}\}$. If a vector h belongs to $\ker(\mathcal{P} - \mu I)^j$ and $(\mathcal{P} - \mu I)^{j-1}h \neq 0$, one says that h is a generalized eigenvector of rank j . The eigenspace $X_\mu^0 = \ker(\mathcal{P} - \mu I)$ is a subspace of X_μ ; eigenvectors are precisely the generalized eigenvectors of rank 1.

3.2.1. Case $\lambda \in \sigma(A) \setminus \sigma(B)$. We look first for non-trivial solutions $u = (v, w)$ of $\mathcal{P}u = \lambda u$. Since λ does not belong to $\sigma(B)$, we have $w = 0$. This in turn implies $v|_\gamma = 0$, by (2.6). Therefore v must be a eigenvector of A corresponding to λ . All eigenvalues of A have finite multiplicities. Assume that λ is an eigenvalue of multiplicity K and V_1, \dots, V_K are the eigenfunctions of A such that

$$\int_\Omega \rho V_i V_j dx = \delta_{i,j}, \quad i, j = 1, \dots, K. \quad (3.4)$$

Here $\delta_{i,j}$ is the Kronecker symbol. Then X_λ^0 is spanned by

$$u_1 = (V_1, 0), \quad u_2 = (V_2, 0), \quad \dots, \quad u_K = (V_K, 0). \quad (3.5)$$

The generalized eigenvectors of rank 2 satisfy the equation $(\mathcal{P} - \lambda I)u_* = u$, where u is an eigenvector of \mathcal{P} . There is no such vectors $u_* = (v_*, w_*)$ in this case, because the second components of all the eigenvectors u_k are zero. In fact, $Bw_* = \lambda w_*$ implies $w_* = 0$. Therefore $(A - \lambda I)v_* = v$, where v is a linear combination of V_j . The last equation is unsolvable, since A is self-adjoint.

3.2.2. *Case $\lambda \in \sigma(B) \setminus \sigma(A)$.* Suppose that y is an eigenfunction of (3.1) corresponding to λ . Then B has a countable collection of linearly independent eigenfunctions $\{b_j(s)y(n)\}_{j=1}^\infty$, where $\{b_j\}_{j=1}^\infty$ is a basis in $L_2(\gamma)$ consisting of smooth functions. On the other hand, the problem

$$-\Delta v + av = \lambda \rho v \text{ in } \Omega \setminus \gamma, \quad \ell v = 0 \text{ on } \partial\Omega, \quad v^- = y(-1)b, \quad v^+ = y(1)b \quad (3.6)$$

is uniquely solvable for any $b \in C^\infty(\gamma)$, since λ does not belong to $\sigma(A)$. Therefore \mathcal{P} possesses the countable set of linearly independent eigenvectors

$$u_1 = (v_1, b_1 y), \quad u_2 = (v_2, b_2 y), \quad \dots, \quad u_j = (v_j, b_j y), \quad \dots, \quad (3.7)$$

where v_j is a solution of (3.6) with b_j in place of b . Note that the values $y(\pm 1)$ are different from zero and hence all v_j are non-trivial solutions. In this case, finding generalized eigenvectors of rank 2 leads to the equation $(B - \lambda I)w_* = by$, which is unsolvable for $b \neq 0$. Hence $X_\lambda = X_\lambda^0$ and $\dim X_\lambda = \infty$.

3.2.3. *Case $\lambda \in \sigma(A) \cap \sigma(B)$.* Since λ belongs to $\sigma(B)$, any eigenvector of \mathcal{P} has the form $(v(x), b(s)y(n))$, where v is a solution of (3.6). But now we can not solve problem (3.6) for any function b , since λ is a point of $\sigma(A)$.

Proposition 3.5. *Let $f \in L_2(\Omega)$ and $b_\pm \in W_2^{3/2}(\gamma)$. Assume that λ is an eigenvalue of A of multiplicity K and the corresponding eigenfunctions V_1, \dots, V_K satisfy (3.4). Then the problem*

$$-\Delta v + av = \lambda \rho v + f \text{ in } \Omega \setminus \gamma, \quad \ell v = 0 \text{ on } \partial\Omega, \quad v^- = b_-, \quad v^+ = b_+ \quad (3.8)$$

admits a solution $v \in W_2^2(\Omega \setminus \gamma)$ if and only if

$$\int_\gamma (b_+ \partial_r V_k^+ - b_- \partial_r V_k^-) d\gamma + \int_\Omega f V_k dx = 0 \quad (3.9)$$

for all $k = 1, \dots, K$.

Proof. This statement is a simple consequence of the Fredholm alternative for the self-adjoint operator A with compact resolvent. Conditions (3.9) can be obtained by multiplying the equation in (3.8) by V_1, \dots, V_K in turn and then integrating by parts twice in view of the boundary conditions. \square

In view of Proposition 3.5, problem (3.6) is solvable if and only if $(b, \Psi_k)_{L_2(\gamma)} = 0$ for all $k = 1, \dots, K$, where

$$\Psi_k = y(1) \partial_r V_k^+ - y(-1) \partial_r V_k^-. \quad (3.10)$$

Let H_λ be the subspace in $L_2(\gamma)$ spanned by Ψ_1, \dots, Ψ_K . Hence the solvability of (3.6) is equivalent to the orthogonality of b to H_λ .

Proposition 3.6. *Assume λ is an eigenvalue of $A = A_- \oplus A_+$ and k_\pm is the multiplicity of λ in the spectrum of A_\pm . Then $\dim H_\lambda \geq \max(k_+, k_-)$.*

Proof. Since the operator A is a direct sum of A_+ and A_- , we can choose a basis in the eigenspace of A such that V_1, \dots, V_{k_+} are identically equal to zero in Ω_- and the rest of k_- eigenfunctions V_{k_++1}, \dots, V_K are identically equal to zero in Ω_+ . Note that $k_+ + k_- = K$. Then H_λ is the linear span of

$$\partial_r V_1^+, \dots, \partial_r V_{k_+}^+, \partial_r V_{k_++1}^-, \dots, \partial_r V_K^-. \quad (3.11)$$

In general, these normal derivatives are linearly dependent in $L_2(\gamma)$, but the first of k_+ derivatives are always linearly independent as well as the last of k_- ones.

Suppose, contrary to our claim, that the functions $\partial_r V_1^+, \dots, \partial_r V_{k_+}^+$ are linearly dependent in $L_2(\gamma)$. Then there exists an eigenfunction v of the Dirichlet type problem $-\Delta v + av = \lambda \rho v$ in Ω_+ , $v = 0$ on $\partial\Omega_+$ such that $\partial_r v = 0$ on the whole boundary of Ω_+ , but this is impossible. The same conclusion can be drawn for the second part of the normal derivatives. \square

Assume $\dim H_\lambda = d$ and choose a basis $\{b_j\}_{j=1}^\infty$ in $L_2(\gamma)$ such that b_1, \dots, b_d belong to H_λ , while b_k for $j > d$ are elements of H_λ^\perp , and $b_j \in C^\infty(\gamma)$. Then

$$(V_1, 0), \dots, (V_K, 0), (v_{d+1}, b_{d+1}y), (v_{d+2}, b_{d+2}y), \dots \quad (3.12)$$

is a countable set of linearly independent eigenvectors of \mathcal{P} . Here v_j is a solution of (3.6) for $b = b_j$ which is orthogonal to the span of V_1, \dots, V_K in $L_2(\rho, \Omega)$. So X_λ^0 is infinite-dimensional.

In this case, we can also find the generalized eigenvectors $u_* = (v_*, w_*)$ of rank 2. They satisfy the equation $(\mathcal{P} - \lambda)u_* = u$, where $u = (v, w)$ is an eigenvector of \mathcal{P} . Hence v_* and w_* solve the problem

$$-\Delta v_* + av_* = \lambda \rho v_* + \rho v \quad \text{in } \Omega \setminus \gamma, \quad \ell v_* = 0 \quad \text{on } \partial\Omega, \quad (3.13)$$

$$-\partial_n^2 w_* = \lambda q w_* + q w \quad \text{in } \omega, \quad \partial_n w_*(\cdot, -1) = 0, \quad \partial_n w_*(\cdot, 1) = 0, \quad (3.14)$$

$$v_*^- = w_*(\cdot, -1), \quad v_*^+ = w_*(\cdot, 1). \quad (3.15)$$

It is easily seen that (3.13)–(3.15) is unsolvable if $w \neq 0$. Therefore the generalized eigenvectors can be associated with some non-trivial linear combinations of the eigenvectors $(V_1, 0), \dots, (V_K, 0)$ only. We set $w = 0$ and $v = c_1 V_1 + \dots + c_K V_K$. Then $w_*(s, n) = b_*(s)y(n)$ and

$$-\Delta v_* + av_* = \lambda \rho v_* + \rho \sum_{j=1}^K c_j V_j \quad \text{in } \Omega \setminus \gamma, \quad \ell v_* = 0 \quad \text{on } \partial\Omega, \quad (3.16)$$

$$v_*^- = y(-1)b_*, \quad v_*^+ = y(1)b_*. \quad (3.17)$$

By Proposition 3.5 and (3.4), the problem admits a solution if and only if

$$c_1 = -(b_*, \Psi_1)_{L_2(\gamma)}, \dots, c_K = -(b_*, \Psi_K)_{L_2(\gamma)}.$$

Since v is non-zero, b_* must have a non-trivial projection onto the subspace H_λ . But b_1, \dots, b_d have this property. Putting $b_* = b_j$ for $j = 1, \dots, d$ into (3.16), (3.17), we obtain the Jordan chains of length 2

$$(V^{(j)}, 0) \mapsto (v_*^{(j)}, b_j y), \quad j = 1, \dots, d,$$

where $V^{(j)} = c_1^{(j)} V_1 + \dots + c_K^{(j)} V_j$ with $c_i^{(j)} = -(b_j, \Psi_i)_{L_2(\gamma)}$.

There are no generalized eigenvectors of rank 3, because all the eigenvectors of rank 2 have nonzero second components and the equation $(B - \lambda I)y_* = b_j y$ is unsolvable for $b_j \neq 0$. Hence the space X_λ has the basis consisting of:

- d Jordan chains of length 2

$$(V^{(1)}, 0) \mapsto (v_*^{(1)}, b_1 y), \dots, (V^{(d)}, 0) \mapsto (v_*^{(d)}, b_d y), \quad (3.18)$$

- the series of eigenvectors of the form

$$(v_{d+1}, b_{d+1}y), (v_{d+2}, b_{d+2}y), \dots, (v_j, b_j y), \dots, \quad (3.19)$$

- $K - d$ eigenvectors of the form

$$(V^{(d+1)}, 0), \dots, (V^{(K)}, 0). \quad (3.20)$$

Here $V^{(1)}, \dots, V^{(K)}$ are linearly independent eigenfunctions of A corresponding to eigenvalue λ .

We summarize the information about the spectrum and generalized eigenspaces of the limit operator \mathcal{P} that we have obtained.

Theorem 3.7. *Let X_λ be the generalized eigenspace corresponding to $\lambda \in \sigma(\mathcal{P})$ and X_λ^0 be the eigenspace.*

(i) *If $\lambda \in \sigma(A) \setminus \sigma(B)$, then X_λ is finite dimensional, $X_\lambda = X_\lambda^0$, and the basis in X_λ is given by (3.5).*

(ii) *The part $\sigma(B) \setminus \sigma(A)$ of $\sigma(\mathcal{P})$ consists of eigenvalues λ of infinite multiplicity with eigenspaces $X_\lambda = X_\lambda^0$ generated by vectors (3.7).*

(iii) *The part $\sigma(A) \cap \sigma(B)$ is also consists of eigenvalues λ of infinite multiplicity, but $X_\lambda \neq X_\lambda^0$. Apart from the eigenvectors, there exist the generalized eigenvectors of rank 2, and the basis in X_λ is given by (3.18)–(3.20). The dimension of factor space X_λ/X_λ^0 does not exceed the multiplicity of λ in the spectrum of A , namely*

$$\max(k_-, k_+) \leq \dim X_\lambda/X_\lambda^0 \leq k_- + k_+,$$

where k_- and k_+ are the multiplicity of λ in the spectra of A_- and A_+ respectively.

Remark 3.8. The operator \mathcal{P} has always the eigenvalue $\lambda = 0$ of infinite multiplicity, since $0 \in \sigma(B)$. The zero eigenvalue is the smallest infinite-fold one, because B is non-negative. All negative eigenvalues, if they exist, have finite multiplicities.

4. ASYMPTOTICS OF EIGENVALUES IN THE CASE $\lambda_0 \in \sigma(B) \setminus \sigma(A)$

We will first focus our attention on perturbations of infinite-fold eigenvalues. In this section, we construct the asymptotics of countable set of eigenvalues λ_ε of (1.2) that converge to an infinite-fold eigenvalue λ_0 of \mathcal{P} when $\lambda_0 \notin \sigma(A)$. We look for the asymptotics in the form

$$\lambda^\varepsilon \sim \lambda_0 + \lambda_1 \varepsilon + \dots, \quad u_\varepsilon(x) \sim \begin{cases} v_0(x) + \varepsilon v_1(x) + \dots & \text{if } x \in \Omega \setminus \omega_\varepsilon, \\ w_0(s, \frac{r}{\varepsilon}) + \varepsilon w_1(s, \frac{r}{\varepsilon}) + \dots & \text{if } x \in \omega_\varepsilon, \end{cases} \quad (4.1)$$

where (v_0, w_0) is a non-zero element of X_{λ_0} . To match the expansions on $\gamma_{\pm\varepsilon}$, we write v_k in the local coordinates (s, n) . Then (2.3) becomes

$$\begin{aligned} w_0(s, \pm 1) + \varepsilon w_1(s, \pm 1) - v_0(s, \pm \varepsilon) - \varepsilon v_1(s, \pm \varepsilon) + \dots &\sim 0, \\ \varepsilon^{-1} \partial_n w_0(s, \pm 1) + \partial_n w_1(s, \pm 1) - \partial_r v_0(s, \pm \varepsilon) - \varepsilon \partial_r v_1(s, \pm \varepsilon) + \dots &\sim 0. \end{aligned}$$

After expanding $v_k(s, \pm \varepsilon)$ and their derivatives into the Taylor series about $n = \pm 0$ for fixed s , we in particular derive $w_1(\cdot, \pm 1) = v_1^\pm \pm \partial_r v_0^\pm$ and $\partial_n w_1(\cdot, \pm 1) = \partial_r v_0^\pm$. Of course, $w_0(\cdot, \pm 1) = v_0^\pm$ and $\partial_n w_0(\cdot, \pm 1) = 0$, since $(v_0, w_0) \in \text{dom } \mathcal{P}$.

Substituting (4.1) into (1.2), and taking into account representation (2.2), we obtain that the pair (v_1, w_1) solves the problem

$$-\Delta v_1 + a v_1 = \lambda_0 \rho v_1 + \lambda_1 \rho v_0 \quad \text{in } \Omega \setminus \gamma, \quad \ell v_1 = 0 \quad \text{on } \partial\Omega, \quad (4.2)$$

$$-\partial_n^2 w_1 = \lambda_0 q w_1 - \varkappa \partial_n w_0 + \lambda_1 q w_0 \quad \text{in } \omega, \quad (4.3)$$

$$\partial_n w_1(\cdot, -1) = \partial_r v_0^-, \quad \partial_n w_1(\cdot, 1) = \partial_r v_0^+, \quad (4.4)$$

$$v_1^- = w_1(\cdot, -1) + \partial_r v_0^-, \quad v_1^+ = w_1(\cdot, 1) - \partial_r v_0^+. \quad (4.5)$$

The compatibility condition for (4.3), (4.4) reads

$$y(-1) \partial_r v_0^- - y(1) \partial_r v_0^+ + \varkappa \int_{-1}^1 \partial_n w_0 y \, dn = \lambda_1 \int_{-1}^1 q w_0 y \, dn \quad \text{on } \gamma, \quad (4.6)$$

where y is an eigenfunction of (3.1) corresponding to λ_0 . We have applied the Fredholm alternative for the operator with compact resolvent associated with the Sturm-Liouville problem (3.1). The relation (4.6) can be treated as a spectral equation on λ_1 for a pseudodifferential operator on γ . We introduce the Dirichlet-to-Neumann maps N_λ^\pm in $L_2(\gamma)$ as follows. Let z be the solution of problem

$$-\Delta z + az = \lambda \rho z \quad \text{in } \Omega_-, \quad z = \varphi \quad \text{on } \gamma, \quad \ell z = 0 \quad \text{on } \partial\Omega \quad (4.7)$$

for given φ . Set $N_\lambda^- \varphi = \partial_r z|_\gamma$. We follow [44] in assuming that

$$\text{dom } N_\lambda^- = \{\varphi \in L_2(\gamma) : z \in W_2^1(\Omega_-) \text{ and } \partial_r z|_\gamma \in L_2(\gamma)\}.$$

Likewise, we set $N_\lambda^+ \varphi = -\partial_r z|_\gamma$, where z is a solution of the problem

$$-\Delta z + az = \lambda \rho z \quad \text{in } \Omega_+, \quad z = \varphi \quad \text{on } \gamma, \quad (4.8)$$

and $\text{dom } N_\lambda^+ = \{\varphi \in L_2(\gamma) : z \in W_2^1(\Omega_+) \text{ and } \partial_r z|_\gamma \in L_2(\gamma)\}$. The operators N_λ^\pm transform the Dirichlet data on γ for solutions of the corresponding boundary value problems into the Neumann ones. Both operators are well-defined if $\lambda \notin \sigma(A)$. The minus sing in definition of N_λ^+ indicates that the direction of axis r coincides with the inward normal on $\gamma = \partial\Omega_+$.

It follows from (3.7) that $w_0(s, n) = b_0(s)y(n)$ and v_0 is a solution of (3.6) for $b = b_0$. Then $\partial_r v_0^- = y(-1) N_{\lambda_0}^- b_0$ and $\partial_r v_0^+ = -y(1) N_{\lambda_0}^+ b_0$. Consequently, condition (4.6) reads

$$y^2(-1) N_{\lambda_0}^- b_0 + y^2(1) N_{\lambda_0}^+ b_0 + \varkappa b_0 \int_{-1}^1 y y' \, dn = \lambda_1 b_0 \int_{-1}^1 q y^2 \, dn.$$

Suppose $\|y\|_{L_2(q, (-1, 1))} = 1$ and write $\theta_{\lambda_0}^\pm = y^2(\pm 1)$. Since

$$\int_{-1}^1 y y' \, dn = \frac{1}{2} \int_{-1}^1 (y^2)' \, dn = \frac{1}{2} (\theta_{\lambda_0}^+ - \theta_{\lambda_0}^-),$$

we can finally rewrite (4.6) in the form $N_{\lambda_0} b_0 = \lambda_1 b_0$, where

$$N_\lambda = \theta_\lambda^- N_\lambda^- + \theta_\lambda^+ N_\lambda^+ + \frac{1}{2} (\theta_\lambda^+ - \theta_\lambda^-) \varkappa. \quad (4.9)$$

Note that the values $\theta_{\lambda_0}^-$ and $\theta_{\lambda_0}^+$ do not depend on our choice of y , because all eigenvalues of the Sturm-Liouville problem are simple.

Proposition 4.1. *The operator N_λ is self-adjoint, bounded below and has compact resolvent for all $\lambda \notin \sigma(A)$.*

Proof. The operators N_λ^- and N_λ^+ are self-adjoint in $L_2(\gamma)$, bounded below and have compact resolvents [44, Th.3.1]. The linear combination $\theta_\lambda^- N_\lambda^- + \theta_\lambda^+ N_\lambda^+$ has the same properties, since θ_λ^- and θ_λ^+ are positive. Finally, N_λ is a perturbation of this linear combination by the operator of multiplication by the bounded function $\frac{1}{2} (\theta_\lambda^+ - \theta_\lambda^-) \varkappa(s)$, which completes the proof. \square

Denote by $\{\lambda_1^{(k)}\}_{k=1}^\infty$ the eigenvalues of N_{λ_0} . So we have calculated the countable set of correctors $\lambda_1^{(k)}$ in asymptotics (4.1). To keep the mathematics rather simple we suppose that the spectrum of N_{λ_0} is simple. It means that the infinite-fold

eigenvalue λ_0 of \mathcal{P} asymptotically splits into an infinite number of simple eigenvalues of T_ε under perturbation. Let $\{b^{(k)}\}_{k=1}^\infty$ be the collection of orthonormal eigenfunctions of N_{λ_0} .

Let us fix the corrector $\lambda_1^{(k)}$ and set $w_0 = w^{(k)} = b^{(k)}y$. Then $v_0 = v^{(k)}$ is a unique solution of (3.6) with λ_0 and $b^{(k)}$ in place of λ and b respectively. Such a choice of $(v^{(k)}, w^{(k)})$ ensures that problem (4.3), (4.4) is solvable, since the compatibility condition $N_{\lambda_0}b^{(k)} = \lambda_1^{(k)}b^{(k)}$ holds. The problem admits the family of solutions

$$w_1(s, n) = \hat{w}_1^{(k)}(s, n) + b_1(s)y(n),$$

where b_1 is an arbitrary $L_2(\gamma)$ -function and $\hat{w}_1^{(k)}$ is the partial solution subject to the condition

$$\int_{-1}^1 qy\hat{w}_1^{(k)} dn = 0 \quad \text{for all } s \in S. \quad (4.10)$$

According to (4.2) and (4.5), with this choice of w_1 the function v_1 can be written as $v_1 = \hat{v}_1^{(k)} + \tilde{v}_1$, where $\hat{v}_1^{(k)}$ (resp. \tilde{v}_1) is a solution of the problem

$$-\Delta\phi + a\phi = \lambda_0\rho\phi + \lambda_1^{(k)}\rho v^{(k)} \text{ in } \Omega \setminus \gamma, \quad \ell\phi = 0 \text{ on } \partial\Omega, \quad \phi^- = g_-, \quad \phi^+ = g_+ \quad (4.11)$$

with the Dirichlet data $g_\pm = \hat{w}_1^{(k)}(\cdot, \pm 1) \mp \partial_n v^{(k)}|_{\gamma_\pm}$ (resp. $g_\pm = y(\pm 1)b_1$). The term $\hat{v}_1^{(k)}$ is uniquely defined, while \tilde{v}_1 along with b_1 will be fixed below.

Set $a_0(s) = a(s, 0)$. The pair (v_2, w_2) in the asymptotics of u_ε solves the problem

$$-\Delta v_2 + av_2 = \lambda_0\rho v_2 + \lambda_1\rho v_1 + \lambda_2\rho v_0 \text{ in } \Omega \setminus \gamma, \quad \ell v_2 = 0 \text{ on } \partial\Omega, \quad (4.12)$$

$$-\partial_n^2 w_2 = \lambda_0 q w_2 - (\varkappa \partial_n - \lambda_1 q)w_1 - (\varkappa^2 n \partial_n - \partial_s^2 + a_0 - \lambda_2 q)w_0 \text{ in } \omega, \quad (4.13)$$

$$\partial_n w_2(\cdot, -1) = \partial_r v_1^- - \partial_r^2 v_0^-, \quad \partial_n w_2(\cdot, 1) = \partial_r v_1^+ + \partial_r^2 v_0^+, \quad (4.14)$$

$$v_2^- = w_2(\cdot, -1) + \partial_r v_1^- - \partial_r^2 v_0^-, \quad v_2^+ = w_2(\cdot, 1) - \partial_r v_1^+ - \partial_r^2 v_0^+. \quad (4.15)$$

By reasoning as above, we obtain that the solution w_2 exists if and only if

$$(N_{\lambda_0} - \lambda_1^{(k)}I)b_1 = \lambda_2 b^{(k)} + h_1^{(k)}, \quad (4.16)$$

where

$$\begin{aligned} h_1^{(k)} &= y(1)(\partial_r \hat{v}_1^{(k)}|_{\gamma_+} + \partial_r^2 v^{(k)}|_{\gamma_+}) - y(-1)(\partial_r \hat{v}_1^{(k)}|_{\gamma_-} - \partial_r^2 v^{(k)}|_{\gamma_-}) \\ &- \int_{-1}^1 (\varkappa \partial_n - \lambda_1^{(k)}q)y\hat{w}_1^{(k)} dn - \int_{-1}^1 (\varkappa^2 n y' b^{(k)} - y^2(\partial_s^2 b^{(k)} - a_0 b^{(k)})) dn. \end{aligned} \quad (4.17)$$

Since $\lambda_1^{(k)}$ is a simple eigenvalue of N_{λ_0} , the second corrector $\lambda_2 = \lambda_2^{(k)}$ can be uniquely calculated from the solvability condition for equation (4.16)

$$\lambda_2^{(k)} = -(h_1^{(k)}, b^{(k)})_{L_2(\gamma)}.$$

Moreover, there exists a unique solution $b_1^{(k)}$ of (4.16) satisfying the condition $(b_1^{(k)}, b^{(k)})_{L_2(\gamma)} = 0$. Hence $v_1^{(k)}$ is now uniquely defined by choosing \tilde{v}_1 to be a solution of (4.11) with the Dirichlet data $g_\pm = y(\pm 1)b_1^{(k)}$. The compatibility condition (4.16) allows us to solve problems (4.13), (4.14) and (4.12), (4.15) one after another and to find solutions $w_2^{(k)}$ and $v_2^{(k)}$.

The process used to find the leading terms of asymptotic expansions (4.1) can be continued to systematically construct all other terms. For fixed $\lambda_0 \in \sigma(B) \setminus \sigma(A)$

we consider the countable collection of formal approximations to eigenvalues and eigenfunctions of the perturbed problem

$$\Lambda_\varepsilon^{(k)} = \lambda_0 + \lambda_1^{(k)}\varepsilon + \lambda_2^{(k)}\varepsilon^2 + \lambda_3^{(k)}\varepsilon^3, \quad (4.18)$$

$$\hat{u}_\varepsilon^{(k)}(x) = \begin{cases} v_0^{(k)}(x) + \varepsilon v_1^{(k)}(x) + \varepsilon^2 v_2^{(k)}(x) + \varepsilon^3 v_3^{(k)}(x) & \text{if } x \in \Omega \setminus \omega_\varepsilon, \\ w_0^{(k)}(s, \frac{r}{\varepsilon}) + \varepsilon w_1^{(k)}(s, \frac{r}{\varepsilon}) + \varepsilon^2 w_2^{(k)}(s, \frac{r}{\varepsilon}) + \varepsilon^3 w_3^{(k)}(s, \frac{r}{\varepsilon}) & \text{if } x \in \omega_\varepsilon. \end{cases} \quad (4.19)$$

Let H be a Hilbert space with norm $\|\cdot\|$ and let T be a self-adjoint operator in H with a domain $\text{dom } T$. We say a pair $(\mu, u) \in \mathbb{R} \times \text{dom } T$ is a *quasimode* of T with the accuracy δ , if $\|u\| = 1$ and $\|(T - \mu I)u\| \leq \delta$. Of course, if $\delta = 0$, then μ is an eigenvalue of T with the eigenvector u .

Proposition 4.2 ([45, p.139]). *Assume (μ, u) is a quasimode of T with accuracy $\delta > 0$ and the spectrum of T is discrete in the interval $[\mu - \delta, \mu + \delta]$. Then there exists an eigenvalue μ_* of T such that $|\mu_* - \mu| \leq \delta$.*

In order to construct the quasimodes of T_ε , we must modify the approximations of eigenfunctions, because they do not belong to $\text{dom } T_\varepsilon$. By construction, the functions $v_j^{(k)}$ and $w_j^{(k)}$ are smooth, since the coefficients a , ρ and q are smooth. But, in general, $\hat{u}_\varepsilon^{(k)}$ have jump discontinuities on $\partial\omega_\varepsilon$.




FIGURE 2. Plot of the function ζ .

Let us define the function ζ plotted in Fig. 2. This function is smooth outside the origin, $\zeta(r) = 1$ for $r \in [0, \beta/2]$ and $\zeta(r) = 0$ in the set $\mathbb{R} \setminus [0, \beta)$. We can choose β small enough such that the local coordinates (s, r) are well defined in $\omega_{2\beta}$. Set

$$\eta_\varepsilon^{(k)} = ([\hat{u}_\varepsilon^{(k)}]_\varepsilon + [\partial_r \hat{u}_\varepsilon^{(k)}]_\varepsilon (r - \varepsilon)) \zeta(r - \varepsilon) + ([\hat{u}_\varepsilon^{(k)}]_{-\varepsilon} + [\partial_r \hat{u}_\varepsilon^{(k)}]_{-\varepsilon} (r + \varepsilon)) \zeta(-r - \varepsilon). \quad (4.20)$$

The function is different from zero in the set $\omega_{\beta+\varepsilon} \setminus \omega_\varepsilon$ only. And it is easy to check that $\eta_\varepsilon^{(k)}$ and $\partial_r \eta_\varepsilon^{(k)}$ have the same jumps across the boundary of ω_ε as $\hat{u}_\varepsilon^{(k)}$ and $\partial_r \hat{u}_\varepsilon^{(k)}$ respectively. Therefore the function $U_\varepsilon^{(k)} = \hat{u}_\varepsilon^{(k)} - \eta_\varepsilon^{(k)}$ belongs to the domain of T_ε . Moreover we have not changed $\hat{u}_\varepsilon^{(k)}$ too much, since

$$\sup_{x \in \Omega \setminus \bar{\omega}_\varepsilon} (|\eta_\varepsilon^{(k)}(x)| + |\Delta \eta_\varepsilon^{(k)}(x)|) \leq c\varepsilon^3. \quad (4.21)$$

It follows from the explicit formula for $\eta_\varepsilon^{(k)}$ and smallness of jumps of $\hat{u}_\varepsilon^{(k)}$, $\partial_r \hat{u}_\varepsilon^{(k)}$ across $\partial\omega_\varepsilon$. All the jumps are of order $O(\varepsilon^3)$, as $\varepsilon \rightarrow 0$, by construction.

We will henceforth write $n_\varepsilon^{(k)} = \|U_\varepsilon^{(k)}\|_{L^2(\rho_\varepsilon, \Omega)}^{-1}$.

Lemma 4.3. *For each $k \in \mathbb{N}$, the pair $(\Lambda_\varepsilon^{(k)}, n_\varepsilon^{(k)} U_\varepsilon^{(k)})$ constructed above is a quasimode of the operator T_ε with the accuracy $O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$.*

Proof. For simplicity we shall drop the index k in the sequel and write $(\Lambda_\varepsilon, U_\varepsilon)$, v_j , w_j and λ_j instead of $(\Lambda_\varepsilon^{(k)}, U_\varepsilon^{(k)})$, $v_j^{(k)}$, $w_j^{(k)}$ and $\lambda_j^{(k)}$. Set $F_\varepsilon = (T_\varepsilon - \Lambda_\varepsilon)U_\varepsilon$. Then we have

$$F_\varepsilon = (-\Delta + a - \Lambda_\varepsilon \rho)U_\varepsilon = \sum_{j=0}^3 \varepsilon^j (-\Delta v_j + a v_j - \rho \sum_{i=0}^j \lambda_i v_{j-i}) + \Delta \eta_\varepsilon - a \eta_\varepsilon + \Lambda_\varepsilon \rho \eta_\varepsilon$$

outside ω_ε . From our choice of v_j , we derive that the first sum in the right-hand side vanishes. Therefore $\sup_{x \in \Omega \setminus \omega_\varepsilon} |F_\varepsilon(x)| \leq c_1 \varepsilon^3$, because of (4.21). Applying representation (2.2) of the Laplace operator in ω , we have

$$\Delta - a + \varepsilon^{-2} \Lambda_\varepsilon q = \sum_{j=0}^3 \varepsilon^{j-2} p_j + \varepsilon^2 P_\varepsilon - a_\varepsilon,$$

where $p_0 = \partial_n^2 + \lambda_0 q$, $p_1 = -\kappa \partial_n + \lambda_1 q$, $p_2 = -n \kappa^2 \partial_n + \partial_s^2 - a(s, 0) + \lambda_2 q$,

$$p_3 = 2n \kappa \partial_s^2 + n \kappa' \partial_s - n^2 \kappa^3 \partial_n - n a(s, 0) \partial_n + \lambda_3 q,$$

and $a_\varepsilon(s, n) = a(s, \varepsilon n) - a(s, 0) - \varepsilon n \partial_n a(s, 0)$. Then, in the domain ω_ε , we obtain

$$F_\varepsilon = (-\Delta + a - \varepsilon^{-2} \Lambda_\varepsilon q)U_\varepsilon = \sum_{j=0}^3 \varepsilon^{j-2} \sum_{i=0}^{j-i} p_i w_{j-i} - \varepsilon^2 P_\varepsilon U_\varepsilon + a_\varepsilon = -\varepsilon^2 P_\varepsilon U_\varepsilon + a_\varepsilon,$$

since the functions w_0, \dots, w_3 solve the equations $\sum_{i=0}^{j-i} p_i w_{j-i} = 0$, by construction. Consequently $\sup_{x \in \omega_\varepsilon} |F_\varepsilon(x)| = \varepsilon^2 \sup_{x \in \omega_\varepsilon} |P_\varepsilon U_\varepsilon(x)| + \sup_{x \in \omega_\varepsilon} |a_\varepsilon(x)| \leq c_2 \varepsilon^2$. Hence

$$\|F_\varepsilon\|_{L^2(\rho_\varepsilon, \Omega)}^2 \leq \int_{\Omega \setminus \omega_\varepsilon} \rho |F_\varepsilon|^2 dx + \varepsilon^{-2} \int_{\omega_\varepsilon} q |F_\varepsilon|^2 dx \leq c_3 \varepsilon^6 + c_4 |\omega_\varepsilon| \varepsilon^2 \leq c_5 \varepsilon^3.$$

The main contribution to the $L^2(\rho_\varepsilon, \Omega)$ -norm of U_ε is given by the integral $\varepsilon^{-2} \int_{\omega_\varepsilon} q |w_0|^2 dx$ which is of order $O(\varepsilon^{-1})$, as $\varepsilon \rightarrow 0$. Hence $\|U_\varepsilon\|_{L^2(\rho_\varepsilon, \Omega)} \geq c_6 \varepsilon^{-1/2}$ for ε small enough, i.e., $n_\varepsilon \leq c_7 \varepsilon^{1/2}$. This bound along with $\|F_\varepsilon\|_{L^2(\rho_\varepsilon, \Omega)} \leq c_8 \varepsilon^{3/2}$ yields

$$\|(T_\varepsilon - \Lambda_\varepsilon I)(n_\varepsilon U_\varepsilon)\|_{L^2(\rho_\varepsilon, \Omega)} = n_\varepsilon \|F_\varepsilon\|_{L^2(\rho_\varepsilon, \Omega)} \leq c_9 \varepsilon^2, \quad (4.22)$$

and this is precisely the assertion of the lemma. \square


Theorem 4.4. *Suppose that λ_0 is an eigenvalue of the limit operator \mathcal{P} such that $\lambda_0 \in \sigma(B) \setminus \sigma(A)$. Assume the spectrum $\{\lambda_1^{(k)}\}_{k \in \mathbb{N}}$ of the operator N_{λ_0} is simple. Then there exists a countable set of eigenvalues $\lambda_{\varepsilon, k}$, $k \in \mathbb{N}$, in the spectrum of T_ε that converge to λ_0 and admit the asymptotics*

$$\lambda_{\varepsilon, k} = \lambda_0 + \varepsilon \lambda_1^{(k)} + O(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0. \quad (4.23)$$

Proof. Fix $I \in \mathbb{N}$. In view of Proposition 4.2 and Lemma 4.3, there exist eigenvalues $\lambda_{\varepsilon, 1}, \dots, \lambda_{\varepsilon, I}$ of T_ε and a constant c_I such that

$$|\lambda_{\varepsilon, k} - \lambda_0 - \varepsilon \lambda_1^{(k)}| \leq c_I \varepsilon^2 \quad (4.24)$$

for $k = 1, 2, \dots, I$ and ε small enough. Moreover these eigenvalues are pairwise different. Suppose, contrary to our claim, that some eigenvalue λ_ε of T_ε simultaneously satisfies two estimates (4.24), for example when $k = 1$ and $k = 2$. Then $\lambda_\varepsilon - \lambda_0 - \varepsilon \lambda_1^{(1)} \leq c_I \varepsilon^2$ and $\lambda_0 + \varepsilon \lambda_1^{(2)} - \lambda_\varepsilon \leq c_I \varepsilon^2$. Adding these inequalities

FIGURE 3. Bifurcation of the eigenvalue $\lambda_0 \in \sigma(B) \setminus \sigma(A)$.

yields $\lambda_1^{(2)} - \lambda_1^{(1)} \leq 2c_I \varepsilon$ for all ε small enough. But this is impossible, because the spectrum of N_{λ_0} is simple and therefore $\lambda_1^{(1)} < \lambda_1^{(2)}$. Write

$$\delta_I^\varepsilon = \varepsilon \max_{k=1, \dots, I} |\lambda_1^{(k)}| + c_I \varepsilon^2.$$

Hence the interval $[\lambda_0 - \delta_I^\varepsilon, \lambda_0 + \delta_I^\varepsilon]$ contains at least I eigenvalues of T_ε that possess the asymptotics (4.23). \square

5. ASYMPTOTICS OF EIGENVALUES IN THE CASE $\lambda_0 \in \sigma(A) \cap \sigma(B)$

In view of Theorem 3.7, if the set $\sigma(A) \cap \sigma(B)$ is non-empty, the operator \mathcal{P} possesses generalized eigenvectors of rank 2. This requires changing the structure of asymptotics

$$\lambda^\varepsilon \sim \lambda_0 + \lambda_{1/2} \varepsilon^{1/2} + \lambda_1 \varepsilon + \dots, \quad (5.1)$$

$$u_\varepsilon(x) \sim \begin{cases} v_0(x) + \varepsilon^{1/2} v_{1/2}(x) + \varepsilon v_1(x) + \dots & \text{for } x \in \Omega \setminus \omega_\varepsilon, \\ w_0(s, \frac{r}{\varepsilon}) + \varepsilon^{1/2} w_{1/2}(s, \frac{r}{\varepsilon}) + \varepsilon w_1(s, \frac{r}{\varepsilon}) + \dots & \text{for } x \in \omega_\varepsilon. \end{cases} \quad (5.2)$$

Here (v_0, w_0) is a non-zero element of the eigenspace spanned by vectors (3.12). If y is a normalized eigenfunction of (3.1) corresponding to λ_0 , then $w_0(s, n) = b_0(s)y(n)$ for some $b_0 \in L_2(\gamma)$. As above, substituting the series in (1.2) in particular yields

$$-\Delta v_{1/2} + a v_{1/2} = \lambda_0 \rho v_{1/2} + \lambda_{1/2} \rho v_0 \text{ in } \Omega \setminus \gamma, \quad \ell v_{1/2} = 0 \text{ on } \partial\Omega, \quad (5.3)$$

$$-\partial_s^2 w_{1/2} = \lambda_0 q w_{1/2} + \lambda_{1/2} q w_0 \text{ in } \omega, \quad (5.4)$$

$$\partial_n w_{1/2}(\cdot, -1) = 0, \quad \partial_n w_{1/2}(\cdot, 1) = 0, \quad (5.5)$$

$$v_{1/2}^- = w_{1/2}(\cdot, -1), \quad v_{1/2}^+ = w_{1/2}(\cdot, 1). \quad (5.6)$$

Since (5.4), (5.5) can be written as $(B - \lambda_0)w_{1/2} = \lambda_{1/2}b_0y$ and B is self-adjoint, a solution $w_{1/2}$ exists if only if

$$\lambda_{1/2}b_0 = 0. \quad (5.7)$$

This condition is a branching point of our algorithm.

Integer power asymptotics: case $w_0 \neq 0$. If b_0 is a non-zero function, then $\lambda_{1/2} = 0$ by necessity. Then problem (5.3)–(5.6) turns into the limit problem (2.4)–(2.6) and $(v_{1/2}, w_{1/2}) \in X_{\lambda_0}^0$. Without loss of generality we assume that $(v_{1/2}, w_{1/2}) = 0$, i.e., this vector is absorbed by the leading term of the asymptotics. Moreover, a trivial verification shows that all terms $\lambda_{j/2}$, $(v_{j/2}, w_{j/2})$ with half-integer indexes in (5.1), (5.2) can be treated as equal to zero. In this case, we come back to the integer power asymptotics (4.1), but the construction of quasimodes needs a slight modification. The next terms λ_1 , (v_1, w_1) solve problem (4.2)–(4.5), and therefore condition (4.6) must hold. But now we cannot rewrite (4.6) in the form of the spectral equation for N_{λ_0} , because this operator is not defined for $\lambda_0 \in \sigma(A)$.

We will “extend” N_λ to $\sigma(A)$ by means of a restriction of the space in which it acts. A slight change in the proof of Proposition 3.5 actually shows that both problems (4.7) and (4.8) are solvable for $\lambda \in \sigma(A)$ if the function φ in the boundary condition on γ is orthogonal to the subspace $H_\lambda \subset L_2(\gamma)$ spanned by functions (3.10). Although solutions of the problems, in this case, are ambiguously determined, we can subject them to some additional condition. Namely, there exists a unique solution z of (4.7) (or (4.8)) satisfying the condition $\partial_r z|_\gamma \perp H_\lambda$. So we can define the Dirichlet-to-Neumann map $M_\lambda^- \varphi = \partial_r z|_\gamma$ on H_λ^\perp , where z is a solution of (4.7) belonging to H_λ^\perp . Similarly, we define $M_\lambda^+ \varphi = -\partial_r z|_\gamma$, where $z \in H_\lambda^\perp$ is a solution of (4.8). Both operators are well defined for $\lambda \in \mathbb{C}$. In fact, we have

$$M_\lambda^\pm = (I - P_\lambda)N_\lambda^\pm(I - P_\lambda),$$

where P_λ is the orthogonal projector onto the subspace H_λ . Moreover, $M_\lambda^\pm = N_\lambda^\pm$ for $\lambda \in \mathbb{C} \setminus \sigma(A)$, since the subspace H_λ is trivial in this case.

Now solvability condition (4.6) becomes $M_{\lambda_0} b_0 = \lambda_1 b_0$, where

$$M_\lambda = \theta_\lambda^- M_\lambda^- + \theta_\lambda^+ M_\lambda^+ + \frac{1}{2}(\theta_\lambda^+ - \theta_\lambda^-) \mathcal{K}.$$

The pseudodifferential operator M_λ has the same properties as N_λ . Suppose that the spectrum of M_{λ_0} is simple and $\lambda_1^{(k)}$ is the k -th eigenvalue of M_λ with the normalized eigenfunction $b^{(k)}$. We set $w_0 = b^{(k)} y$.

Assume λ_0 is an K -fold eigenvalue of A with the eigenspace \mathcal{V}_{λ_0} . In order to shorten notation, we introduce the vector $\mathbf{V} = (V_1, \dots, V_K)$, where the eigenfunctions V_k are subject to condition (3.4). Then the leading term $v_0 = v_0^{(k)}$ in (5.2) solves (3.6) for $\lambda = \lambda_0$ and $b = b^{(k)}$ and has the form

$$v_0^{(k)}(x) = \hat{v}_0^{(k)}(x) + \alpha_0 \cdot \mathbf{V}(x),$$

where α_0 is an arbitrary vector in \mathbb{R}^K and $\hat{v}_0^{(k)}$ is a partial solution of (3.6) such that $\hat{v}_0^{(k)} \perp \mathcal{V}_{\lambda_0}$. The dot denotes the scalar product in \mathbb{R}^K . To determine $v_0^{(k)}$ uniquely, we should calculate α_0 . Next, we have

$$w_1(s, n) = \hat{w}_1^{(k)}(s, n) + b_1(s)y(n), \quad (5.8)$$

where $\hat{w}_1^{(k)}$ is a partial solution of (4.3), (4.4) subject to condition (4.10), and b_1 is an arbitrary $L_2(\gamma)$ -function. Assume that $b_1 = g_0 + g$, where $g_0 \in H_{\lambda_0}$ and $g \in H_{\lambda_0}^\perp$. Then problem (4.2), (4.5) for v_1 admits solutions

$$v_1 = \phi + \psi + \alpha_1 \cdot \mathbf{V}, \quad \alpha_1 \in \mathbb{R}^K, \quad (5.9)$$

where ϕ and ψ solve the problems

$$\begin{aligned} -\Delta\phi + a\phi &= \lambda_0\rho\phi \quad \text{in } \Omega \setminus \gamma, \quad \ell\phi = 0 \quad \text{on } \partial\Omega, \quad \phi = y(\pm 1)g \quad \text{on } \gamma_{\pm}; \\ -\Delta\psi + a\psi &= \lambda_0\rho\psi + \lambda_1^{(k)}\rho\left(\hat{v}_0^{(k)} + \alpha_0 \cdot \mathbf{V}\right) \quad \text{in } \Omega \setminus \gamma, \quad \ell\psi = 0 \quad \text{on } \partial\Omega, \\ \psi^{\pm} &= \hat{w}_1^{(k)}(\cdot, \pm 1) \mp \partial_r \hat{v}_0^{(k)} + y(\pm 1)g_0 \mp \alpha_0 \cdot \partial_r \mathbf{V} \quad \text{on } \gamma \end{aligned}$$

respectively. Since g is orthogonal to H_{λ_0} , the first problem is solvable and admits a solution belonging to $\mathcal{V}_{\lambda_0}^{\perp}$. As for the second one, its solvability conditions can be written in the form $(C_{\lambda_0} - \lambda_1^{(k)}I)\alpha_0 = f$, where C_{λ_0} is a matrix with the entries

$$c_{ij} = \int_{\gamma_+} \partial_r V_i \partial_r V_j d\gamma + \int_{\gamma_-} \partial_r V_i \partial_r V_j d\gamma, \quad i, j = 1, \dots, K,$$

and f is a vector with the components

$$\begin{aligned} f_i &= \lambda_1^{(k)} \int_{\Omega} \rho \hat{v}_0^{(k)} V_i dx + \int_{\gamma_+} (\hat{w}_1^{(k)} - \partial_r \hat{v}_0^{(k)} + y(1)g_0) \partial_r V_i d\gamma \\ &\quad - \int_{\gamma_-} (\hat{w}_1^{(k)} + \partial_r \hat{v}_0^{(k)} + y(-1)g_0) \partial_r V_i d\gamma, \quad i = 1, \dots, K. \end{aligned}$$

If we suppose that $\lambda_1^{(k)}$ is not an eigenvalue of C_{λ_0} , then the solvability conditions for ψ can be fulfilled for any $g \in H_{\lambda_0}$. It is enough to set $\alpha_0 = (C_{\lambda_0} - \lambda_1^{(k)}I)^{-1}f$. Then the problem has a solution $\phi \in \mathcal{V}_{\lambda_0}^{\perp}$.

Note that the vector α_0 has not yet been defined, because f depends on the unknown function g_0 . Using representations (5.8) and (5.9) along with the fact that $g \in H_{\lambda_0}^{\perp}$, we can write the solvability condition for (4.13), (4.14) in the form

$$(M_{\lambda_0} - \lambda_1^{(k)}I)g = \lambda_2^{(k)}b^{(k)} + \lambda_1^{(k)}g_0 + h_1^{(k)}, \quad (5.10)$$

where $h_1^{(k)}$ is given by (4.17). For the equation (5.10) to be meaningful, we need to ensure that the right hand side is orthogonal to H_{λ_0} . Obviously, $b^{(k)}$ belongs to $H_{\lambda_0}^{\perp}$. Next, if $\lambda_1^{(k)}$ is different from zero, there exists a unique vector $g_0 \in H_{\lambda_0}$ such that $\lambda_1^{(k)}g_0 + h_1^{(k)} \in H_{\lambda_0}^{\perp}$. With g_0 in hand, we can uniquely defined f , α_0 , and finally the leading term $(v_0^{(k)}, w_0^{(k)})$. The solvability condition for (5.10) has the form $\lambda_2^{(k)} = -(h_1^{(k)} + \lambda_1^{(k)}g_0, b^{(k)})_{L_2(\gamma)}$. Then there exists a unique solution g satisfying the condition $(g, b^{(k)})_{L_2(\gamma)} = 0$. And finally, we can uniquely define $w_1^{(k)}$ and $v_1^{(k)}$ (up to the vector α_1). This process can be continued to systematically construct all other terms $\lambda_j^{(k)}$, $v_j^{(k)}$ and $w_j^{(k)}$ in the approximation $\Lambda_{\varepsilon}^{(k)}$, $\hat{u}_{\varepsilon}^{(k)}$ of the form (4.18), (4.19). As in the previous section, $\hat{u}_{\varepsilon}^{(k)}$ can be improved to the element $U_{\varepsilon}^{(k)} = \hat{u}_{\varepsilon}^{(k)} - \eta_{\varepsilon}^{(k)}$ from the domain of operator T_{ε} , where $\eta_{\varepsilon}^{(k)}$ is given by (4.21).

Summarizing results of the above calculations, we obtain the following statement.

Lemma 5.1. *The pairs $(\Lambda_{\varepsilon}^{(k)}, \|U_{\varepsilon}^{(k)}\|_{L^2(\rho_{\varepsilon}, \Omega)}^{-1}U_{\varepsilon}^{(k)})$, $k \in \mathbb{N}$, are quasimodes of T_{ε} with the accuracy $O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$ that approximate the part of spectrum lying in a vicinity of $\lambda_0 \in \sigma(A) \cap \sigma(B)$.*

The lemma can be proved similarly to Lemma 4.3.

Integer power asymptotics: case $w_0 = 0$. Suppose now that $\lambda_{1/2}$ and b_0 are equal to zero simultaneously. In this case, we also construct the integer power asymptotics (4.1). Since $w_0 = 0$, problem (4.2)–(4.5) becomes

$$-\Delta v_1 + av_1 = \lambda_0 \rho v_1 + \lambda_1 \rho v_0 \quad \text{in } \Omega \setminus \gamma, \quad \ell v_1 = 0 \quad \text{on } \partial\Omega, \quad (5.11)$$

$$-\partial_n^2 w_1 = \lambda_0 q w_1 \quad \text{in } \omega, \quad \partial_n w_1(\cdot, -1) = \partial_r v_0^-, \quad \partial_n w_1(\cdot, 1) = \partial_r v_0^+, \quad (5.12)$$

$$v_1^- = w_1(\cdot, -1) + \partial_r v_0^-, \quad v_1^+ = w_1(\cdot, 1) - \partial_r v_0^+. \quad (5.13)$$

Compatibility condition (4.6) for problem (5.12) takes the form

$$y(-1) \partial_r v_0^- - y(1) \partial_r v_0^+ = 0. \quad (5.14)$$

If we set $v_0 = \alpha \cdot \mathbf{V}$, then the last condition can be written as $\alpha \cdot \Psi = 0$, where $\alpha \in \mathbb{R}^K \setminus \{0\}$, $\Psi = (\Psi_1, \dots, \Psi_K)$ and Ψ_k are given by (3.10). Let y_* be the solution of the Cauchy problem $-y_*'' = \lambda_0 q y_*$ in $(-1, 1)$, $y_*(-1) = 0$, $y_*'(1) = 1$. Problem (5.12) admits the solution $w_1(s, n) = y_*(n) \alpha \cdot \partial_r \mathbf{V}^-(s) + y(n) b_1(s)$, where b_1 is an arbitrary $L_2(\gamma)$ -function. Obviously, $\partial_n w_1(\cdot, -1) = \partial_r v_0^-$. The Lagrange identity yields $(yy_*' - y'y_*)|_{-1}^1 = 0$, and hence, $y_*'(1) = y(-1)/y(1)$. Then condition (5.14) becomes $\alpha \cdot \partial_r \mathbf{V}^+ = y_*'(1) \alpha \cdot \partial_r \mathbf{V}^-$, and we have

$$\partial_n w_1(\cdot, 1) = y_*'(1) \alpha \cdot \partial_r \mathbf{V}^- = \alpha \cdot \partial_r \mathbf{V}^+ = \partial_r v_0^+.$$

Next, we can rewrite (5.11) and (5.13) in the form

$$-\Delta v_1 + av_1 = \lambda_0 \rho v_1 + \lambda_1 \rho \alpha \cdot \mathbf{V} \quad \text{in } \Omega \setminus \gamma, \quad \ell v_1 = 0 \quad \text{on } \partial\Omega, \quad (5.15)$$

$$v_1^- = \alpha \cdot \partial_r \mathbf{V}^- + y(-1) b_1, \quad v_1^+ = (y_*(1) - y_*'(1)) \alpha \cdot \partial_r \mathbf{V}^+ + y(1) b_1. \quad (5.16)$$

To achieve solvability of the problem one needs to choose proper vectors α along with the parameter λ_1 . Multiplying the equation in (5.15) by V_1, \dots, V_K in turn and integrating by parts twice in view of the boundary conditions (5.16) yield the equation $L\alpha = \lambda_1 \alpha$, where the symmetric matrix $L = (l_{ij})_{i,j=1}^K$ has the entries

$$l_{ij} = \int_{\gamma} ((y_*(1) - y_*'(1)) \partial_r V_i^+ \partial_r V_j^+ + \partial_r V_i^- \partial_r V_j^-) d\gamma. \quad (5.17)$$

Note that L does not depend on b_1 , because of (5.14).

Hence, the number λ_1 and non-zero vector α satisfy two conditions $L\alpha = \lambda_1 \alpha$ and $\alpha \cdot \Psi = 0$. We must find the eigenvectors of L which are “orthogonal” to Ψ . The dimension of the space $h_{\lambda_0} = \{\alpha \in \mathbb{C}^K : \alpha \cdot \Psi = 0\}$ is $K - d$, since $\dim H_{\lambda_0} = d$. Assume that L has the simple eigenvalues ν_1, \dots, ν_p with the eigenvectors $\alpha_1, \dots, \alpha_p$ belonging to h_{λ_0} . Obviously, $p \leq K - d$. For any pair (ν_k, α_k) , we can solve (5.15), (5.16) and find

$$v_1^{(k)} = \hat{v}_1^{(k)} + \alpha_1^{(k)} \cdot \mathbf{V}$$

up to the vector $\alpha_1^{(k)} \in \mathbb{R}^K$. Also, it follows from the first condition in (5.16) that

$$b_1^{(k)}(s) = y(-1)^{-1} \hat{v}_1^{(k)}(s, -1) - \alpha_k \cdot \partial_r \mathbf{V}(s, -1),$$

and now the function $w_1^{(k)} = y_* \alpha_k \cdot \partial_r \mathbf{V}^- + y b_1^{(k)}$ is uniquely defined. We continue in this fashion obtaining p quasimodes of T_ε of the form

$$\Lambda_\varepsilon^{(k)} = \lambda_0 + \nu_k \varepsilon + \lambda_{2,k} \varepsilon^2 + \lambda_{3,k} \varepsilon^3, \quad (5.18)$$

$$W_\varepsilon^{(k)}(x) = \begin{cases} \alpha_k \cdot \mathbf{V}(x) + \varepsilon v_1^{(k)}(x) + \varepsilon^2 v_2^{(k)}(x) + \varepsilon^3 v_3^{(k)}(x) - \eta_\varepsilon^{(k)}(x) & \text{if } x \in \Omega \setminus \omega_\varepsilon, \\ \varepsilon w_1^{(k)}(s, \frac{r}{\varepsilon}) + \varepsilon^2 w_2^{(k)}(s, \frac{r}{\varepsilon}) + \varepsilon^3 w_3^{(k)}(s, \frac{r}{\varepsilon}) & \text{if } x \in \omega_\varepsilon. \end{cases} \quad (5.19)$$

Now we summarize the results of our calculations. The next lemma can be proved similarly to Lemma 4.3.

Lemma 5.2. *The pairs $(\Lambda_\varepsilon^{(k)}, \|W_\varepsilon^{(k)}\|_{L^2(\rho_\varepsilon, \Omega)}^{-1} W_\varepsilon^{(k)})$, $k = 1, \dots, p$, are quasimodes of T_ε with the accuracy $O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$ that approximate a part of spectrum lying in a vicinity of $\lambda_0 \in \sigma(A) \cap \sigma(B)$.*

Half-integer power asymptotics. The set of quasimodes in Lemma 5.1 does not approximate all eigenvalues of T_ε that converge to λ_0 . We assume that $\lambda_{1/2}$ in (5.2) is different from zero, and then $b_0 = 0$, by (5.7). Recalling now (3.12), we have $v_0 = \beta_0 \cdot \mathbf{V}$, where β_0 is an arbitrary vector in \mathbb{R}^K such that $\|\beta_0\| = 1$. In this case, we will use some finite-dimensional operator instead of M_{λ_0} to split the limit multiple eigenvalue λ_0 . Reasoning as above we deduce that the problem (5.3)-(5.6) has a solution of the form $w_{1/2}(s, n) = b_{1/2}(s)y(n)$, $v_{1/2} = \hat{v}_{1/2} + \beta_{1/2} \cdot \mathbf{V}$ where $\beta_{1/2} \in \mathbb{R}^K$ and $\hat{v}_{1/2}$ is a partial solution of the problem

$$-\Delta v_{1/2} + a v_{1/2} = \lambda_0 \rho v_{1/2} + \lambda_{1/2} \rho v_0 \quad \text{in } \Omega \setminus \gamma, \quad \ell v_{1/2} = 0 \quad \text{on } \partial\Omega, \quad (5.20)$$

$$v_{1/2}^- = y(-1) b_{1/2}, \quad v_{1/2}^+ = y(1) b_{1/2} \quad (5.21)$$

that is orthogonal to \mathcal{V}_{λ_0} in $L_2(\rho, \Omega)$. By Proposition 3.5, the solvability conditions for (5.20), (5.21) can be written in the vector form

$$\int_\gamma b_{1/2} \Psi d\gamma + \lambda_{1/2} \beta_0 = 0. \quad (5.22)$$

For the next terms we have

$$-\Delta v_1 + a v_1 = \rho(\lambda_0 v_1 + \lambda_{1/2} v_{1/2} + \lambda_1 v_0) \quad \text{in } \Omega \setminus \gamma, \quad \ell v_1 = 0 \quad \text{on } \partial\Omega, \quad (5.23)$$

$$-\partial_n^2 w_1 = \lambda_0 q w_1 + \lambda_{1/2} q w_{1/2} \quad \text{in } \omega, \quad (5.24)$$

$$\partial_n w_1(\cdot, -1) = \partial_r v_0^-, \quad \partial_n w_1(\cdot, 1) = \partial_r v_0^+, \quad (5.25)$$

$$v_1^- = w_1(\cdot, -1) + \partial_r v_0^-, \quad v_1^+ = w_1(\cdot, 1) - \partial_r v_0^+. \quad (5.26)$$

Problem (5.24), (5.25) is solvable if $y(1) \partial_r v_0^+ - y(-1) \partial_r v_0^- + \lambda_{1/2} b_{1/2} = 0$. Since $\partial_r v_0^\pm = \beta_0 \cdot \partial_r \mathbf{V}^\pm$, it can be written in the form

$$\beta_0 \cdot (y(1) \partial_r \mathbf{V}^+ - y(-1) \partial_r \mathbf{V}^-) + \lambda_{1/2} b_{1/2} = 0.$$

Using notation (3.10), we have

$$\beta_0 \cdot \Psi + \lambda_{1/2} b_{1/2} = 0. \quad (5.27)$$

Multiplying this equality by Ψ^\top , integrating over γ and recalling (5.22), we finally discover $(G_{\lambda_0} - \lambda_{1/2}^2) \beta_0 = 0$, where G_{λ_0} is the Gram matrix of Ψ_1, \dots, Ψ_K . This matrix is semi-positive and its rank is equal to the dimension of H_{λ_0} .

Suppose ω^2 is a positive simple eigenvalue of G_{λ_0} with the eigenvector β_0 . So there exist two different correctors $\lambda_{1/2} = \omega$ and $\lambda_{1/2} = -\omega$ in asymptotics (5.1)

with the same leading term $v_0 = \beta_0 \cdot \mathbf{V}$ in approximation (5.2). First assume that $\lambda_{1/2} = \omega$. From (5.27), we have $b_{1/2} = -\omega^{-1}\beta_0 \cdot \Psi$. Up to a function b_1 , we can find $w_1(s, n) = \hat{w}_1(s, n) + b_1(s)y(n)$, where \hat{w}_1 is a partial solution of (5.24), (5.25) subject to the condition $(\hat{w}_1, y)_{L_2(q, (-1, 1))} = 0$. Next, problem (5.23), (5.26) is solvable if and only if $\int_{\gamma} b_1 \Psi d\gamma + \omega \beta_{1/2} + \lambda_1 \beta_0 = h$, where

$$h = \int_{\gamma_+} (\hat{w}_1 - \partial_r v_0) \partial_r \mathbf{V}_k d\gamma - \int_{\gamma_-} (\hat{w}_1 + \partial_r v_0) \partial_r \mathbf{V}_k d\gamma - \omega \int_{\Omega} \rho \hat{v}_{1/2} \mathbf{V}_k dx.$$

Also, a solution of the problem

$$\begin{aligned} -\partial_n^2 w_{3/2} &= q(\lambda_0 w_{3/2} + \omega w_1 + \lambda_1 w_{1/2}) \quad \text{in } \omega, \\ \partial_n w_{3/2}(\cdot, -1) &= \partial_r v_{1/2}^-, \quad \partial_n w_{3/2}(\cdot, 1) = \partial_r v_{1/2}^+. \end{aligned}$$

exists if and only if

$$y(-1) \partial_r v_{1/2}^- - y(1) \partial_r v_{1/2}^+ = \omega b_1 + \lambda_1 b_{1/2}. \quad (5.28)$$

Reasoning as in the previous step, we can rewrite this condition in the form

$$(G_{\lambda_0} - \omega^2 I) \beta_{1/2} = 2\omega \lambda_1 \beta_0 - \omega h + f,$$

where

$$f = y(-1) \int_{\gamma_-} \partial_r \hat{v}_{1/2} \Psi d\gamma - y(1) \int_{\gamma_+} \partial_r \hat{v}_{1/2} \Psi d\gamma.$$

Since ω^2 is an eigenvalue of G_{λ_0} , the system admits a solution $\beta_{1/2}$ if and only if

$$\lambda_1 = \frac{1}{2\omega} \beta_0 \cdot (\omega h - f). \quad (5.29)$$

Although the unit vector β_0 is defined up to the change of sign, λ_1 is uniquely defined by (5.29), because the transformation $\beta_0 \mapsto -\beta_0$ implies that the vectors h and f also change their signs. We fix this solution such that $\beta_{1/2} \cdot \beta_0 = 0$. Then

$$b_1 = \frac{1}{\omega} \left(y(-1) \partial_r \hat{v}_{1/2}^- - y(1) \partial_r \hat{v}_{1/2}^+ - \beta_{1/2} \cdot \Psi - \lambda_1 b_{1/2} \right),$$

by (5.28). Assuming that $\lambda_{1/2}^+ = \omega$ we have calculated λ_1^+ , $w_{1/2}^{(+)}$, $w_1^{(+)}$ and $v_{1/2}^{(+)}$ in (5.1), (5.2). We can continue in this fashion obtaining the next terms of the asymptotics. Taking $\lambda_{1/2}^- = -\omega$ we can compute analogously the terms λ_k^- , $w_k^{(-)}$ and $v_k^{(-)}$ in the asymptotics of other eigenvalue and eigenfunction. A simple analysis of the foregoing formulas shows that $w_{1/2}^{(-)} = -w_{1/2}^{(+)}$ and $v_{1/2}^{(-)} = -v_{1/2}^{(+)}$. Moreover, by construction, we have

$$w_{1/2}^{(\pm)}(s, n) = \mp \omega^{-1} y(n) \beta_0 \cdot \Psi(s), \quad v_{1/2}^{(\pm)}(x) = \pm v_*(x),$$

where v_* is a solution of the problem

$$-\Delta v + av = \lambda_0 \rho v + \omega \rho \beta_0 \cdot \mathbf{V} \text{ in } \Omega \setminus \gamma, \quad \ell v = 0 \text{ on } \partial\Omega, \quad v = -\omega^{-1} y(\pm 1) \beta_0 \cdot \Psi \text{ on } \gamma_{\pm}$$

that is orthogonal to \mathcal{V}_{λ_0} .

Let us summarize the above considerations in the following lemma.

Lemma 5.3. *Let d be the dimension of H_{λ_0} , where $\lambda_0 \in \sigma(A) \cap \sigma(B)$. Suppose that all non-zero eigenvalues $\omega_1^2, \dots, \omega_d^2$ of the matrix G_{λ_0} are simple and $\beta_{0,1}, \dots, \beta_{0,d}$ are the corresponding normalized eigenvectors. Then the operator T_{ε} possesses d*

pairs of the quasimodes $(\hat{\mu}_{\varepsilon,j}^{\pm}, n_{\varepsilon,j}^{\pm} V_{\varepsilon,j}^{(\pm)})$, $j = 1, \dots, d$, with the accuracy $O(\varepsilon^{3/2})$ as $\varepsilon \rightarrow 0$, where $n_{\varepsilon,j}^{\pm}$ is a normalizing factor and

$$\begin{aligned}\hat{\mu}_{\varepsilon,j}^{\pm} &= \lambda_0 \pm \varepsilon^{1/2} \omega_j + \sum_{k=2}^6 \varepsilon^{k/2} \lambda_{k/2,j}^{\pm}, \\ V_{\varepsilon,j}^{(\pm)} &= \beta_{0,j} \cdot \mathbf{V} \pm \varepsilon^{1/2} v_{*,j} + \sum_{k=2}^6 \varepsilon^{k/2} v_{k/2,j}^{(\pm)} - \eta_{\varepsilon,j}^{(\pm)} \quad \text{in } \Omega \setminus \omega_{\varepsilon}, \\ V_{\varepsilon,j}^{(\pm)} &= \mp \varepsilon^{1/2} \omega_j^{-1} y\left(\frac{r}{\varepsilon}\right) \beta_{0,j} \cdot \Psi(s) + \sum_{k=2}^6 \varepsilon^{k/2} w_{k/2,j}^{(\pm)}\left(s, \frac{r}{\varepsilon}\right) \quad \text{in } \omega_{\varepsilon}.\end{aligned}$$

The small correctors $\eta_{\varepsilon,j}^{(\pm)}$ are defined as in (4.20) with $V_{\varepsilon,j}^{(\pm)}$ in place of $\hat{u}_{\varepsilon}^{(k)}$.

Proof. The proof differs from the proof of Lemma 4.3 only by estimate (4.22). The approximations to eigenvalues and eigenfunctions have been constructed up to order $O(\varepsilon^3)$ and therefore the remainder term F_{ε} (with the notation of Lemma 4.3) can be also estimated by $\|F_{\varepsilon}\|_{L^2(\rho_{\varepsilon}, \Omega)} \leq c_1 \varepsilon^{3/2}$. But the leading term w_0 in this asymptotics is equal to zero and hence the $L^2(\rho_{\varepsilon}, \Omega)$ -norm of $V_{\varepsilon,j}^{(\pm)}$ is bounded uniformly with respect to ε . Then the normalizing factor $n_{\varepsilon,j}^{\pm}$ tends to some positive number as $\varepsilon \rightarrow 0$ and

$$\|(T_{\varepsilon} - \hat{\mu}_{\varepsilon,j}^{\pm} I)(n_{\varepsilon,j}^{\pm} V_{\varepsilon,j}^{(\pm)})\|_{L^2(\rho_{\varepsilon}, \Omega)} = n_{\varepsilon,j}^{\pm} \|F_{\varepsilon}\|_{L^2(\rho_{\varepsilon}, \Omega)} \leq c_2 \varepsilon^{3/2},$$

which is the desired conclusion. \square

In view of Lemmas 5.1–5.3, we can prove the following result in the same way as Theorem 4.4.

Theorem 5.4. *Suppose that the set $\sigma(A) \cap \sigma(B)$ is non empty and λ_0 is an eigenvalue of the limit operator \mathcal{P} belonging to this intersection.*

- (i) *Assume the spectrum $\{\lambda_1^{(k)}\}_{k \in \mathbb{N}}$ of the operator M_{λ_0} is simple. Then there exists a countable set of eigenvalues $\lambda_{\varepsilon,k}$, $k \in \mathbb{N}$, in the spectrum of T_{ε} that admit the asymptotics*

$$\lambda_{\varepsilon,k} = \lambda_0 + \varepsilon \lambda_1^{(k)} + O(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0.$$

- (ii) *Assume $\dim H_{\lambda_0} = d$ and all positive eigenvalues ω_j^2 of the matrix G_{λ_0} are simple. Then operator T_{ε} possesses $2d$ eigenvalues with the asymptotics*

$$\begin{aligned}\mu_{\varepsilon,j}^{-} &= \lambda_0 - \varepsilon^{1/2} \omega_j + \varepsilon \lambda_{1,j}^{-} + O(\varepsilon^{3/2}), \\ \mu_{\varepsilon,j}^{+} &= \lambda_0 + \varepsilon^{1/2} \omega_j + \varepsilon \lambda_{1,j}^{+} + O(\varepsilon^{3/2}), \quad j = 1, \dots, d,\end{aligned}$$


as $\varepsilon \rightarrow 0$, where $\lambda_{1,j}^{\pm}$ can be calculated from (5.29).

- (iii) *Also, the operator T_{ε} can have no more than $K - d$ eigenvalues of possessing the asymptotics*

$$\lambda_i^{\varepsilon} = \lambda_0 + \nu_i \varepsilon + O(\varepsilon^2), \quad \text{as } \varepsilon \rightarrow 0,$$

where ν_i are simple non-zero eigenvalues of the matrix in (5.17).

In this case, the generalized eigenspace X_{λ_0} contains the Jordan chains. Note that problem (5.3)–(5.6) for $(v_{1/2}, w_{1/2})$ coincides up to the multiplier $\lambda_{1/2}$ in the

FIGURE 4. Bifurcation of the eigenvalue $\lambda_0 \in \sigma(A) \cap \sigma(B)$.

right-hand side with problem (3.13)–(3.15) for generalized eigenvectors. Inspecting the structure of $V_{\varepsilon,j}^{(\pm)}$ more closely we see that

$$V_{\varepsilon,j}^{(\pm)} = U_j \pm \varepsilon^{1/2} U_j^* + O(\varepsilon), \text{ as } \varepsilon \rightarrow 0,$$

where the vectors $U_j = (\beta_{0,j} \cdot \mathbf{V}(x), 0)$, $U_j^* = (v_{*,j}(x), -\omega_j^{-1} y(n) \beta_{0,j} \cdot \Psi(s))$ form a Jordan chain of \mathcal{P} corresponding to λ_0 . This observation has the following geometric interpretation. The operator T_ε possesses pairs of eigenvalues with the asymptotics $\mu_{\varepsilon,j}^\pm = \lambda_0 \pm \omega_j \varepsilon^{1/2} + O(\varepsilon)$ for which the corresponding normalized eigenfunctions $u_{\varepsilon,j}^{(-)}$ and $u_{\varepsilon,j}^{(+)}$ converge in $L_2(\Omega)$ to the same function $v_{0,j} = \beta_{0,j} \cdot \mathbf{V}$, as $\varepsilon \rightarrow 0$. Although these eigenfunctions remain orthogonal for all ε in the weighted space $L_2(\rho_\varepsilon, \Omega)$, they make an infinitely small angle between them in $L_2(\Omega)$ with the standard norm, and stick together at the limit. In particular, it leads to the loss of completeness in $L_2(\Omega)$ for the limit eigenfunction collection. Interestingly enough, however, the plane $\pi_{\varepsilon,j}$ that is the span of $u_{\varepsilon,j}^{(-)}$ and $u_{\varepsilon,j}^{(+)}$ has regular behaviour as $\varepsilon \rightarrow 0$. The limit position of $\pi_{\varepsilon,j}$ is the 2-dimensional space π_j spanned by the functions $\beta_{0,j} \cdot \mathbf{V}$ and $v_{*,j}$.

We actually have an example of singular perturbations in which the completeness property of perturbed eigenfunction collection passes in some sense into the completeness of generalized eigenfunctions of the limit non-self-adjoint operator. Although we didn't justify the asymptotics of eigenfunctions of (1.2), we can formally state that the non-self-adjoint operator \mathcal{P} contains all the information about the asymptotic behaviour of eigenvalues of the perturbed problem:

- the spectrum of \mathcal{P} is a limit set for the spectra $\sigma(T_\varepsilon)$ as $\varepsilon \rightarrow 0$;
- knowing the multiplicity of eigenvalues of \mathcal{P} , we can divide $\sigma(T_\varepsilon)$ into finite or infinite subsets of eigenvalues with the same limits as $\varepsilon \rightarrow 0$;
- in the case $\lambda \in \sigma(A) \cap \sigma(B)$, the dimension of space H_λ indicates how many eigenvalues of T_ε possess the half-integer power asymptotics;
- the Jordan chains of \mathcal{P} are involved in the quasimodes of T_ε (in the formal asymptotics of eigenfunctions of T_ε).

As we pointed out in the introduction, the problem

$$-\Delta u_\varepsilon = \lambda^\varepsilon \rho_\varepsilon u_\varepsilon \quad \text{in } \Omega, \quad u_\varepsilon = 0 \quad \text{on } \partial\Omega \quad (5.30)$$

with the perturbed density $\rho_\varepsilon(x; m) = \rho(x)$ for $x \in \Omega \setminus \omega_\varepsilon$ and $\rho_\varepsilon(x; m) = \varepsilon^{-m} q_\varepsilon(x)$ for $x \in \omega_\varepsilon$, the Dirichlet boundary condition and without the potential a in the differential equation has been considered in [37]. The variational methods have been applied to study the asymptotic behaviour of eigenvalues and eigenfunctions. In this case the problem can be realized as the family of bounded self-adjoint operators A_ε in $\dot{W}_2^1(\Omega) = \{v \in W_2^1(\Omega) : u_\varepsilon = 0 \text{ on } \partial\Omega\}$. Due to the second representation theorem [43, Theorem VI.2.23], A_ε is defined by the identity $\mathfrak{a}(A_\varepsilon u, \phi) = (\rho_\varepsilon u, \phi)_{L_2(\Omega)}$ for all $\phi \in \dot{W}_2^1(\Omega)$, because the Dirichlet form $\mathfrak{a}(u, \phi) = \int_\Omega \nabla u \cdot \nabla \phi \, dx$ is a scalar product in $\dot{W}_2^1(\Omega)$. Problem (5.30) can be written in the form $A_\varepsilon u_\varepsilon = \lambda_\varepsilon^{-1} u_\varepsilon$. For $m > 1$ the operators A_ε diverge as $\varepsilon \rightarrow 0$, and it has been shown that $\|A_\varepsilon\| \leq c\varepsilon^{1-m}$ with a constant c being independent of ε .

The complete asymptotic analysis of eigenvalues λ_ε of (5.30) has been carried out for $m = 3$. We have proved that the spectrum consists of a countable number of infinite series of eigenvalues with the asymptotics

$$\lambda_{kj}^\varepsilon = \mu_k \varepsilon + \nu_{kj} \varepsilon^2 + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0, \quad k, j \in \mathbb{N}, \quad (5.31)$$

where μ_k is an infinite-fold eigenvalue of B and $\{\nu_{kj}\}_{j=1}^\infty$ is a spectrum of pseudo-differential operator N_{μ_k} acting in $L_2(\gamma)$ (see formula (4.9)). The operator N_{μ_k} is a self-adjoint realization of a boundary value problem for the Laplace operator in Ω with coupling conditions on γ .

Since the first eigenvalue μ_1 of B is equal to zero, the low-lying eigenvalues admit the asymptotics $\lambda_{1j}^\varepsilon = \varepsilon^2(\nu_{1j} + o(1))$, $\varepsilon \rightarrow 0$. In this case, $\{\nu_{1j}\}_{j=1}^\infty$ are eigenvalues of the spectral problem $-\Delta u = \nu q_0 \delta_\gamma u$ in Ω , $u = 0$ on $\partial\Omega$, where $q_0 = \int_{-1}^1 q(r) \, dr$ and δ_γ is a distribution in $\mathcal{D}'(\Omega)$ acting as $\langle \delta_\gamma, \phi \rangle = \int_\gamma \phi \, d\gamma$ for any $\phi \in C^\infty(\Omega)$. The problem describes the eigenvibrations of a membrane with the mass density $\rho_0(x) = q_0 \delta_\gamma(x)$, i.e., the whole mass of the membrane is supported on γ and the rest part of vibrating system is massless. Such series of infinitesimal eigenvalues with the asymptotics $\lambda_{1j}^\varepsilon = \varepsilon^{m-1}(\nu_{1j} + o(1))$, $\varepsilon \rightarrow 0$, exists for any $m > 1$. Hence, the estimate $\|A_\varepsilon\| \leq c\varepsilon^{1-m}$ is sharp. The similar result is obtained in Theorems 4.4 and 5.4 for problem (1.2) when $m = 2$ (see Corollary 5.5).

In [37] some results on the convergence of eigenfunctions have been also obtained. Interestingly enough, the complete asymptotic description of λ^ε and u_ε in (5.30) includes not only the operator B but also the operator A which is associated with the problem $-\Delta u = \lambda u$ in $\Omega \setminus \gamma$, $u = 0$ on $\partial\Omega$ and γ in this case. In view of (5.31), any point of the positive spectral half-line is an accumulation point for eigenvalues λ^ε , i.e., for each $\lambda > 0$ there exists a sequence $\{\lambda^{\varepsilon_i}\}_{i=1}^\infty$ of eigenvalues such that $\lambda^{\varepsilon_i} \rightarrow \lambda$ as $\varepsilon_i \rightarrow 0$. We have proved that only the points of $\sigma(A)$ can be approximated by eigenvalues λ^{ε_i} so that the corresponding eigenfunctions u_{ε_i} converge to nontrivial limits in $L^2(\Omega)$. These limits are eigenfunctions of A corresponding to λ .

Corollary 5.5. *Problem (1.2) has a series of small eigenvalues that admit the asymptotics $\lambda_{\varepsilon,k} = \varepsilon \nu_k + O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$, where ν_k are eigenvalues of the problem*

$$-\Delta v + av = 0 \text{ in } \Omega \setminus \gamma, \quad \ell v = 0 \text{ on } \partial\Omega, \quad [v]_\gamma = 0, \quad [\partial_r v]_\gamma + \nu q_0 v = 0 \text{ on } \gamma \quad (5.32)$$

with the spectral parameter ν in the coupling conditions. In addition, if $0 \in \sigma(A)$, then (1.2) has also a finite number of small eigenvalues with asymptotics

$$\mu_{\varepsilon,j}^\pm = \pm \omega_j \sqrt{\varepsilon} + O(\varepsilon) \text{ as } \varepsilon \rightarrow 0, \quad j = 1, \dots, d.$$

Proof. The existence of such eigenvalues follows from Theorems 4.4 and 5.4 along with the observation that $\lambda_0 = 0$ belongs to $\sigma(B)$ (see Remark 3.8). It remains to derive the eigenvalue problem for ν_k . Let us look at (2.4)–(2.6) and put $\lambda = 0$:

$$\begin{aligned} -\Delta v + av &= 0 \text{ in } \Omega \setminus \gamma, \quad \ell v = 0 \text{ on } \partial\Omega, \quad v^- = w(\cdot, -1), \quad v^+ = w(\cdot, 1), \text{ on } \gamma, \\ \partial_n^2 w &= 0 \text{ in } \omega, \quad \partial_n w(\cdot, -1) = 0, \quad \partial_n w(\cdot, 1) = 0, \end{aligned}$$

We see that $w = b(s)$ and hence $v^- = v^+$, i.e., v is continuous on γ . Since $y = 1$ in this case, condition (4.6) for $v_0 = v$ reads as $\partial_r v^+ - \partial_r v^- + \lambda_1 q_0 b = 0$ on γ . To complete the proof we replace λ_1 with ν and recall that $b = v|_\gamma$. Note that problem (5.32) can be written in the form

$$-\Delta v + av = \nu q_0 \delta_\gamma v \text{ in } \Omega, \quad \ell v = 0 \text{ on } \partial\Omega.$$

Hence, ν_k are eigenvalues of a membrane with the mass density $\rho_0(x) = q_0 \delta_\gamma(x)$ supported by the curve γ . \square

6. ASYMPTOTICS OF EIGENVALUES IN THE CASE $\lambda_0 \in \sigma(A) \setminus \sigma(B)$

For the sake of completeness, we briefly discuss the perturbation of eigenvalue λ_0 of \mathcal{P} which does not belong to $\sigma(B)$. In view of part (i) of Theorem 3.7, λ_0 is an eigenvalue of finite multiplicity. Also, we have $v_0 = \alpha_0 \cdot \mathbf{V}$ and $w_0 = 0$ in asymptotics (4.1), where $\alpha_0 \in \mathbb{R}^K \setminus \{0\}$. Then (4.3) and (4.4) imply

$$-\partial_n^2 w_1 = \lambda_0 q w_1 \text{ in } \omega, \quad \partial_n w_1(\cdot, -1) = \alpha_0 \cdot \partial_r \mathbf{V}^-, \quad \partial_n w_1(\cdot, 1) = \alpha_0 \cdot \partial_r \mathbf{V}^+.$$

Since $\lambda_0 \notin \sigma(B)$, there exists a unique solution of the problem. Suppose the functions W_k , $k = 1, \dots, K$, solve the problems

$$-\partial_n^2 w = \lambda_0 q w \text{ in } \omega, \quad \partial_n w(\cdot, -1) = \partial_r V_k^-, \quad \partial_n w(\cdot, 1) = \partial_r V_k^+,$$

and $\mathbf{W} = (W_1, \dots, W_K)$. Thus we have $w_1 = \alpha_0 \cdot \mathbf{W}$. Next, we can rewrite (4.2) and (4.5) in the form

$$-\Delta v_1 + av_1 = \lambda_0 \rho v_1 + \lambda_1 \rho \alpha_0 \cdot \mathbf{V} \text{ in } \Omega \setminus \gamma, \quad \ell v_1 = 0 \text{ on } \partial\Omega, \quad (6.1)$$

$$v_1^- = \alpha_0 \cdot (\mathbf{W}(\cdot, -1) + \partial_r \mathbf{V}^-), \quad v_1^+ = \alpha_0 \cdot (\mathbf{W}(\cdot, 1) - \partial_r \mathbf{V}^+). \quad (6.2)$$

In general, the problem is unsolvable, because λ_0 belongs to the spectrum of A which is the direct sum of A_- and A_+ . To achieve the solvability one needs to choose proper vectors α_0 along with the parameter λ_1 . Multiplying the equation in (6.1) by V_1, \dots, V_K in turn and integrating by parts twice in view of the boundary conditions (6.2) yield the spectral matrix equation $R\alpha_0 = \lambda_1 \alpha_0$, where the matrix $R = (r_{ij})_{i,j=1}^K$ has the entries

$$r_{ij} = \int_\gamma ((W_i(\cdot, 1) - \partial_r V_i^+) \partial_r V_j^+ + (W_i(\cdot, -1) + \partial_r V_i^-) \partial_r V_j^-) d\gamma.$$

The matrix R is symmetric, because it is easy to check that

$$\int_\gamma W_i(\cdot, \pm 1) \partial_r V_j^\pm d\gamma = \int_\gamma W_j(\cdot, \pm 1) \partial_r V_i^\pm d\gamma.$$

Assume that R has K simple eigenvalues $\lambda_1^{(1)}, \dots, \lambda_1^{(K)}$ with the corresponding eigenvalues $\alpha_0^{(1)}, \dots, \alpha_0^{(K)}$. For any pair $(\lambda_1^{(k)}, \alpha_0^{(k)})$, we can solve (6.1), (6.2) and find $v_1^{(k)} = \hat{v}_1^{(k)} + \alpha_1^{(k)} \cdot \mathbf{V}$ up to the vector $\alpha_1^{(k)} \in \mathbb{R}^K$. We continue in this fashion obtaining K different quasimodes with high enough accuracy of the operator T_ε that approximate the part of spectrum lying in a vicinity of λ_0 .

Theorem 6.1. *Suppose that λ_0 is an eigenvalue of the limit operator \mathcal{P} such that $\lambda_0 \in \sigma(A) \setminus \sigma(B)$. Assume λ_0 has multiplicity K and the matrix R possesses the simple eigenvalues $\lambda_1^{(1)}, \dots, \lambda_1^{(K)}$. Then there exist K eigenvalues in the spectrum of T_ε that converge to λ_0 and admit the asymptotics $\lambda_{\varepsilon,k} = \lambda_0 + \varepsilon \lambda_1^{(k)} + O(\varepsilon^2)$ as $\varepsilon \rightarrow 0$, $k = 1, \dots, K$.*

REFERENCES

- [1] T. Sarpkaya (2010). Wave forces on offshore structures. Cambridge University Press.
- [2] F. R. Gantmakher, M. G. Krein (1950). Oscillation matrices and kernels and small oscillations of mechanical systems. Gostekhizdat, Moscow; English transl., American Mathematical Soc., 2002.
- [3] Bates, F. S., Fredrickson, G. H. (1990). Block copolymer thermodynamics: theory and experiment. Annual review of physical chemistry, 41(1), 525-557.
- [4] Sens, P., Turner, M. S. (1997). Inclusions in Thin Smectic Films. Journal de Physique II, 7(12), 1855-1870.
- [5] Pratibha, R., Park, W., Smalyukh, I. I. (2010). Colloidal gold nanosphere dispersions in smectic liquid crystals and thin nanoparticle-decorated smectic films. Journal of Applied Physics, 107(6), 063511.
- [6] Ladbroke, B. D., Chapman, D. (1969). Thermal analysis of lipids, proteins and biological membranes a review and summary of some recent studies. Chemistry and Physics of Lipids, 3(4), 304-356.
- [7] E. Sánchez-Palencia(1984). Perturbation of eigenvalues in thermoelasticity and vibration of systems with concentrated masses. In Trends and applications of pure mathematics to mechanics (pp. 346-368). Springer, Berlin, Heidelberg.
- [8] J. Sanchez Hubert and E. Sánchez-Palencia. Vibration and coupling of continuous systems. Springer-Verlag, Berlin, 1989.
- [9] Yu. D. Golovaty, S. A. Nazarov, O. A. Oleinik, and T. S. Soboleva. Natural oscillations of a string with an additional mass. *Sibirsk. Mat. Zh.*, 29(5):71–91, 237, 1988.
- [10] M. Lobo, E. Pérez (2003). Local problems for vibrating systems with concentrated masses: a review. Comptes Rendus Mécanique, V. 331(4), pp. 303-317.
- [11] Yu. D. Golovaty, S. A. Nazarov, O. A. Oleinik (1990). Asymptotic expansions of eigenvalues and eigenfunctions in problems on oscillations of a medium with concentrated perturbations. Trudy Matematicheskogo Instituta imeni VA Steklova, 192, 42-60.
- [12] Yu. D. Golovaty. Spectral properties of oscillatory systems with added masses. Trudy Moskov. Mat. Obshch., 54:29–72, 278, 1992.
- [13] Yu. D. Golovaty, A. S. Lavrenyuk (2000). Asymptotic expansions of local eigen vibrations for plate with density perturbed in neighbourhood of one-dimensional manifold. Matematychni Studii, 13(1), 51-62.
- [14] T. A. Mel'nik, S. A. Nazarov, Asymptotic analysis of the Neumann problem on the junction of a body and thin heavy rods, Algebra i Analiz, 12:2 (2000), 188–238; St. Petersburg Math. J., 12:2 (2001), 317–351
- [15] T. A. Mel'nik Vibrations of a thick periodic junction with concentrated masses. Mathematical Models and Methods in Applied Sciences, (2001) 11:06, 1001-1027.
- [16] G. A. Chechkin, T. A. Mel'nyk (2012) Asymptotics of eigenelements to spectral problem in thick cascade junction with concentrated masses, Applicable Analysis, 91:6, 1055-1095.
- [17] Yu. Golovaty, H. Hrabchak (2007). Asymptotics of the spectrum of a Sturm-Liouville operator on networks with perturbed density. *Visnyk Lviv. Univ., Ser. Mech. Math.* Vol. 67, 66-83.
- [18] Yu. Golovaty, H. Hrabchak (2010). On Sturm-Liouville problem on starlike graphs with “heavy” nodes. *Visnyk Lviv. Univ., Ser. Mech. Math.* Vol. 72, 63-78.
- [19] O. A. Oleinik, A. S. Shamaev, G. A. Yosifian, Mathematical Problems in Elasticity and Homogenization, North-Holland, Amsterdam, 1992.
- [20] Castro, C., Zuazua, E. (2000). Low frequency asymptotic analysis of a string with rapidly oscillating density. SIAM Journal on Applied Mathematics, 60(4), 1205-1233.
- [21] C. Castro, E. Zuazua (2000). High frequency asymptotic analysis of a string with rapidly oscillating density. European Journal of Applied Mathematics, 11(6), 595-622.

- [22] V. Rybalko (2002). Vibrations of elastic systems with a large number of tiny heavy inclusions. *Asymptotic Analysis*, 32(1), 27-62.
- [23] M. Lobo and E. Perez, On vibrations of a body with many concentrated masses near the boundary, *Math. Models Methods Appl. Sci.* 3 (1993) 249-273.
- [24] M. Lobo and E. Perez, Vibrations of a body with many concentrated masses near the boundary: High frequency vibrations, *Spectral Analysis of Complex Structures* (Hermann, 1995) pp. 85-101.
- [25] M. Lobo and E. Perez (1995). Vibrations of a membrane with many concentrated masses near the boundary. *Mathematical Models and Methods in Applied Sciences*, 5(5), 565-585.
- [26] Chechkin, G. A. (2004). Estimation of solutions of boundary-value problems in domains with concentrated masses located periodically along the boundary: Case of light masses. *Mathematical Notes*, 76(5-6), 865-879.
- [27] G. A. Chechkin, (2005). Asymptotic expansions of eigenvalues and eigenfunctions of an elliptic operator in a domain with many “light” concentrated masses situated on the boundary. Two-dimensional case. *Izvestiya: Mathematics*, 69(4), 805.
- [28] S. A. Nazarov, E. Perez, New asymptotic effects for the spectrum of problems on concentrated masses near the boundary. *Comptes Rendus Mécanique*, V. 337, Is. 8, 2009, pp 585-590.
- [29] G. A. Chechkin, M. E. Pérez, E. I. Yablokova (2005). Non-periodic boundary homogenization and “light” concentrated masses. *Indiana University Mathematics Journal*, 321-348.
- [30] Yablokova, E. I. (2005). On vibrations of bodies with concentrated masses near boundaries. *Journal of Mathematical Sciences*, 129(1), 3688-3695.
- [31] Gómez, D., Lobo, M., Nazarov, S. A., Pérez, E. (2006). Asymptotics for the spectrum of the Wentzell problem with a small parameter and other related stiff problems. *Journal de mathématiques pures et appliquées*, 86(5), 369-402.
- [32] Gómez, D., Lobo, M., Nazarov, S. A., Pérez, E. (2006). Spectral stiff problems in domains surrounded by thin bands: Asymptotic and uniform estimates for eigenvalues. *Journal de mathématiques pures et appliquées*, 85(4), 598-632.
- [33] Nazarov, S. A., Pérez, M. E. (2018). On multi-scale asymptotic structure of eigenfunctions in a boundary value problem with concentrated masses near the boundary. *Revista Matemática Complutense*, 31(1), 1-62.
- [34] Gómez, D., Nazarov, S. A., Pérez-Martínez, M. E. (2021). Localization effects for Dirichlet problems in domains surrounded by thin stiff and heavy bands. *Journal of Differential Equations*, 270, 1160-1195.
- [35] Riva, M. D., Provenzano, L. (2018). On vibrating thin membranes with mass concentrated near the boundary: an asymptotic analysis. *SIAM Journal on Mathematical Analysis*, 50(3), 2928-2967.
- [36] Yu. Golovaty, D. Gómez, M. Lobo, and E. Pérez. Asymptotics for the eigenelements of vibrating membranes with very heavy thin inclusions. *C. R., Méc., Acad. Sci. Paris*, 330(11):777-782, 2002.
- [37] Yu. D. Golovaty, D. Gómez, M. Lobo, and E. Pérez. On vibrating membranes with very heavy thin inclusions. *Math. Models Methods Appl. Sci.*, 14(7):987-1034, 2004.
- [38] Yu. Golovaty (2020). On spectrum of strings with δ' -like perturbations of mass density. *Visnyk Lviv. Univ., Ser. Mech. Math.*, Vol. 89, 60-79.
- [39] T. A. Mel'nyk (2001). Hausdorff convergence and asymptotic estimates of the spectrum of a perturbed operator. *Zeitschrift für Analysis und ihre Anwendungen*, 20(4), 941-957.
- [40] D. Mugnolo, R. Nittka, O. Post (2010). Convergence of sectorial operators on varying Hilbert space. *Operators and Matrices*, Vol.7(4), 955-995.
- [41] I. Gohberg, M. G. Kreĭn. Introduction to the theory of linear nonselfadjoint operators (Vol. 18). American Mathematical Soc., 1978.
- [42] J. H. Bramble, J. E. Osborn (1973). Rate of convergence estimates for nonselfadjoint eigenvalue approximations. *Mathematics of Computation*, 27(123), 525-549.
- [43] T. Kato, *Perturbation Theory for Linear Operators*, 2nd edition, Grundlehren der Mathematischen Wissenschaften, Band 132, Springer-Verlag, Berlin-New York, 1976.
- [44] W. Arendt and R. Mazzeo. Spectral properties of the Dirichlet-to-Neumann operator on Lipschitz domains. *Ulmer Seminare Heft*, 12:23-38, 2007.
- [45] Fedoruk M.V., Babich V.M., Lazutkin, V.F., ... & Vainberg, B. R. (1999). *Partial Differential Equations V: Asymptotic Methods for Partial Differential Equations* (Vol. 5). Springer Science & Business Media.

DEPT. OF MECHANICS AND MATHEMATICS, IVAN FRANKO NATIONAL UNIVERSITY OF LVIV,
UNIVERSYTETSKA STR.1, LVIV, 79000, UKRAINE