

2-GENERATED AXIAL ALGEBRAS OF MONSTER TYPE

CLARA FRANCHI, MARIO MAINARDIS, SERGEY SHPECTOROV

ABSTRACT. In this paper we provide the basic setup for a project, initiated by Felix Rehren in [16], aiming at classifying all 2-generated primitive axial agebras of Monster type (α, β) . We first revise Rehren's construction of an initial object in the category of primitive n -generated axial algebras giving a formal one, filling some gaps and, though correcting some inaccuracies, confirm Rehren's results. Then we focus on 2-generated algebras which naturally part into three cases: the two *critical* cases $\alpha = 2\beta$ and $\alpha = 4\beta$, and the *generic* case (i.e. all the rest). About these cases, which will be dealt in detail in subsequent papers, we give bounds on the dimensions (the generic case already treated by Rehren) and classify all 2-generated primitive axial algebras of Monster type (α, β) over $\mathbb{Q}(\alpha, \beta)$ for α and β algebraically independent indeterminates over $\mathbb{Q}$. Finally we restrict to the 2-generated *Majorana algebras* (i.e. when $\alpha = \frac{1}{4}$ and $\beta = \frac{1}{32}$), showing that these fall precisely into the nine isomorphism types of the Norton-Sakuma algebras.

1. INTRODUCTION

Axial algebras constitute a class of commutative non-associative algebras generated by idempotent elements called *axes* such that their adjoint action is semisimple and the relative eigenvectors satisfy a prescribed fusion law. Let R be a ring, $\{\alpha, \beta\} \subseteq R \setminus \{0, 1\}$ and $\alpha \neq \beta$. An axial algebra over R is called of *Monster type* (α, β) if it satisfies the fusion law $\mathcal{M}(\alpha, \beta)$ given in Table 1. This means that the adjoint action of every axis has spectrum $\{1, 0, \alpha, \beta\}$ and, for any two eigenvectors v_γ, v_δ with relative eigenvalues $\gamma, \delta \in \{1, 0, \alpha, \beta\}$, the product $v_\gamma \cdot v_\delta$ is a sum of eigenvectors relative to eigenvalues contained in $\gamma \star \delta$. This class was introduced by J. Hall, F. Rehren and S. Shpectorov [7] in order to axiomatise some key features of important classes of algebras, including the weight-2 components of OZ-type vertex operator algebras, Jordan algebras and Matsuo algebras (see the introductions of [7], [16] and [5]). They are also of particular interest for finite group theorists as most of the finite simple groups, or their automorphism groups, can be faithfully and effectively represented as automorphism groups of these algebras.

$\star$	1	0	α	β
1	1	$\emptyset$	α	β
0	$\emptyset$	0	α	β
α	α	α	$1, 0$	β
β	β	β	β	$1, 0, \alpha$

TABLE 1. Fusion table $\mathcal{M}(\alpha, \beta)$

The motivating example is the Griess algebra $V_2^\sharp$ which is a real axial algebra of Monster type $(\frac{1}{4}, \frac{1}{32})$ and coincides with the weight-2 component of the Monster vertex operator algebra $V^\sharp$. Here axes are associated to the involutions of type $2A$ in the Monster (i.e. those having the double cover of the Baby Monster as centraliser). The subalgebras of the Griess algebra which are generated by two axes were first classified by S. Norton in [14] who showed that there are nine isomorphism classes of such algebras, corresponding to the 9 conjugacy classes in the Monster group M of the dihedral subgroups generated by the pairs of $2A$ involutions associated to the two generating axes. These algebras are labelled $1A$, $2A$, $3A$, $4A$, $5A$, $6A$, $4B$, $2B$.

On the basis of an earlier work by M. Miyamoto [12], who observed that Ising vectors in a vertex operator algebra of CFT-type satisfy the Monster fusion law, in [17] S. Sakuma classified all OZ -type vertex operator algebras generated by a pair of two Ising conformal vectors showing that, up to rescaling, the isomorphism types of their weight-2 subspaces match precisely the 9 classes of Norton, now often called Norton-Sakuma algebras.

By extracting the relevant properties of these weight-2 subspaces, A. A. Ivanov introduced in 2009 the concept of Majorana algebras [10], which are real axial algebras of Monster type $(\frac{1}{4}, \frac{1}{32})$ satisfying some additional properties, in particular they are endowed with a positive definite associative bilinear form. In 2010 A. Ivanov, D. Pasechnik, A. Seress and S. Shpectorov obtained Norton's classification within the axiomatic context of Majorana algebras (see [11]). A further development was achieved by J. Hall, F. Rehren and S. Shpectorov [7] who constructed a universal object for primitive Frobenius axial algebras with a prescribed fusion law and extended Norton-Sakuma theorem to 2-generated primitive Frobenius axial algebras of Monster type $(\frac{1}{4}, \frac{1}{32})$. Subsequently F. Rehren [16, 15] addressed the general case dropping the assumption on the existence of the bilinear form (which characterises Frobenius axial algebras) and described a universal object for primitive axial algebras with a prescribed fusion law. In the particular case of 2-generated algebras of Monster type (α, β) with $\alpha \notin \{2\beta, 4\beta\}$ and in characteristic other than 2, he produced a spanning set of 8 elements and computed the structure constants with respect to these elements. Finally he produced new examples of 2-generated primitive Frobenius axial algebras of Monster type. This paper is part of a project of the authors aiming to classifying 2-generated primitive axial algebras of Monster type over fields. We start by giving, for any positive integer n , a formal construction of a universal n -generated primitive axial algebra mapping epimorphically onto every n -generated axial algebra with a prescribed fusion law. We then focus on 2-generated primitive axial algebras of Monster type. We say that such an algebra is *symmetric* if the map that swaps the two generating axes extends to an automorphism of the entire algebra. All the algebras considered by Rehren in [15, 16] are symmetric. In Section 4, we re-prove Rehren's result on the number of generators and get the following bound for symmetric algebras in the case $\alpha = 4\beta$.

Theorem 1.1. *Every 2-generated primitive axial algebra of Monster type (α, β) over a field $\mathbb{F}$ of characteristic other than 2 has dimension at most 8, provided $\alpha \notin \{2\beta, 4\beta\}$.*

Theorem 1.2. *Every 2-generated symmetric primitive axial algebra of Monster type $(4\beta, \beta)$ over a field $\mathbb{F}$ of characteristic other than 2 has dimension at most 8, except possibly when $(\alpha, \beta) = (2, \frac{1}{2})$.*

The case $(\alpha, \beta) = (2, \frac{1}{2})$ is truly exceptional, as the infinite dimensional 2-generated symmetric primitive axial algebra of Monster type $(2, \frac{1}{2})$ constructed in [4] shows. On the other hand, the same bound 8 holds for 2-generated primitive axial algebras of Monster type $(2\beta, \beta)$ over any ring in which 2 and β are invertible and it is the best possible (see [3]).

In Section 5 we consider in more details the case when $\alpha - 2\beta$ and $\alpha - 4\beta$ are invertible in the field $\mathbb{F}$, which we call the *generic case*. Denote by $\mathbb{F}_0$ the prime subfield of $\mathbb{F}$ and let $\mathbb{F}_0(\alpha, \beta)[x, y, z, t]$ be the polynomial ring in 4 variables over $\mathbb{F}_0(\alpha, \beta)$. We prove the following result.

Theorem 1.3. *There exists a subset $T \subseteq \mathbb{F}_0(\alpha, \beta)[x, y, z, t]$ of size 4, depending only on $\mathbb{F}_0$, α , and β , such that every 2-generated primitive axial algebra of Monster type (α, β) over a field $\mathbb{F}$ of characteristic other than 2, with $\alpha, \beta \in \mathbb{F}$ and $\alpha \notin \{2\beta, 4\beta\}$ is completely determined, up to homomorphic images, by a quadruple $(x_0, y_0, z_0, t_0) \in \mathbb{F}^4$ which is a common zero of all the elements of T .*

Using Theorem 1.3, we classify the algebras defined over the field $\mathbb{Q}(\alpha, \beta)$ with α and β independent indeterminates over $\mathbb{Q}$. We refer to [8] for the definition of the algebras of type 1A, 2B, 3C(η), $\eta \in \mathbb{F}$. We denote by 3A(α, β) the algebra of dimension 4 defined in [16, Table 9] for $\alpha \neq \frac{1}{2}$.

Theorem 1.4. *Let V be a 2-generated primitive axial algebra of Monster type (α, β) over the field $\mathbb{Q}(\alpha, \beta)$, with α and β algebraically independent indeterminates over $\mathbb{Q}$. Then we have one of the following:*

- (1) V is the trivial algebra $\mathbb{Q}(\alpha, \beta)$ of type 1A;
- (2) V is an algebra of type 2B;
- (3) V is an algebra of Jordan type α of type 3C(α);
- (4) V is an algebra of Jordan type β of type 3C(β);
- (5) V is an algebra of dimension 4 of type 3A(α, β).

In particular, V is symmetric.

Recall that the fusion law $\mathcal{M}(\alpha, \beta)$ admits a $\mathbb{Z}_2$ -grading $\mathcal{M}(\alpha, \beta)_+ = \{1, 0, \alpha\}$ and $\mathcal{M}(\alpha, \beta)_- = \{\beta\}$ and this implies that every axis induces an automorphism of the algebra called Miyamoto involution. The Miyamoto group is the group generated by all Miyamoto involutions (see [9]).

Corollary 1.5. *Let V be a primitive finitely generated axial algebra of Monster type (α, β) over the field $\mathbb{Q}(\alpha, \beta)$, with α and β independent indeterminates over $\mathbb{Q}$. Then the Miyamoto group of V is a group of 3-transpositions.*

Finally, as a consequence of Theorem 1.3, we get that the Norton-Sakuma Theorem holds in general for 2-generated primitive axial algebra of Monster type $(\frac{1}{4}, \frac{1}{32})$ over a field of characteristic zero, without any assumption on the existence of a Frobenius form. This fact has been also checked computationally in [19].

Theorem 1.6. *Every 2-generated primitive axial algebra of Monster type $(\frac{1}{4}, \frac{1}{32})$ over a field of characteristic zero is a Norton-Sakuma algebra.*

Throughout this paper R is a commutative associative ring with 1.

While this paper was in preparation, Yabe posted a preprint in arXiv giving an almost complete classification of the primitive symmetric 2-generated axial algebras of Monster type (α, β) [18].

2. PRIMITIVE AXIAL ALGEBRAS

We begin with some basic results about endomorphisms of R -modules, which are well known for vector spaces. The main difference is that, when considering R -modules instead of vector spaces, it is no longer true in general that eigenvectors relative to different eigenvalues are linearly independent.

Let V be an R -module. For $\xi \in \text{End}(V)$, $\lambda \in R$, and $\Gamma \subseteq R$, define

$$V_\lambda^\xi := \{v \in V \mid \xi(v) = \lambda v\} \text{ and } V_\Gamma^\xi := \sum_{\lambda \in \Gamma} V_\lambda^\xi.$$

If V is also an R -algebra and $a \in V$, denote by ad_a the endomorphism of V induced by multiplication by a :

$$\begin{aligned} \text{ad}_a : V &\rightarrow V \\ x &\mapsto ax \end{aligned}$$

In this case, we'll write simply V_λ^a and V_Γ^a instead of $V_\lambda^{\text{ad}_a}$ and $V_\Gamma^{\text{ad}_a}$, respectively.

Two elements α and β of R are called *distinguishable* if $\alpha - \beta$ is a unit in R . In the remainder of this section we assume that Γ is a finite set of pairwise distinguishable elements of R . Note that every nontrivial ring homomorphism maps sets of pairwise distinguishable elements into sets of pairwise distinguishable elements. Let $R[x]$ be the ring of polynomials over R with indeterminate x .

Lemma 2.1. *If $g \in R[x]$ satisfies $g(\lambda) = 0$ for every $\lambda \in \Gamma$ and $\deg g < |\Gamma|$, then $g = 0$.*

Proof. We proceed by induction on $|\Gamma|$. If $|\Gamma| = 1$, then g is a constant and since the value of g is zero in at least one element (the element of Γ), it must be $g = 0$. Suppose $|\Gamma| > 1$. Assume by contradiction that g is not the zero polynomial and set $k := \deg g$. Then $k < |\Gamma|$. Let $\lambda \in \Gamma$, then, since $x - \lambda$ is monic, we have $g = q \cdot (x - \lambda) + r$, with $r = g(\lambda) = 0$, so $g = q \cdot (x - \lambda)$. Then clearly $\deg q = k - 1 < |\Gamma'|$, where $\Gamma' = \Gamma \setminus \{\lambda\}$. Also for $\mu \in \Gamma'$, $0 = g(\mu) = q(\mu) \cdot (\mu - \lambda)$. Since μ and λ are distinguishable, $(\mu - \lambda)$ is a unit and so $q(\mu) = 0$. By induction this means that $q = 0$, whence also $g = 0$, a contradiction. $\square$

For $\mu \in \Gamma$, define

$$(1) \quad f_\mu := \prod_{\lambda \in \Gamma \setminus \{\mu\}} (x - \lambda),$$

and

$$(2) \quad f := \prod_{\lambda \in \Gamma} (x - \lambda),$$

clearly

$$f = (x - \mu)f_\mu$$

for every $\mu \in \Gamma$. Note that, since elements of Γ are pairwise distinguishable, $f_\mu(\mu)$ is a unit in R .

Corollary 2.2.

$$\sum_{\mu \in \Gamma} \frac{1}{f_\mu(\mu)} f_\mu = 1.$$

Proof. Define

$$g := \sum_{\mu \in \Gamma} \frac{1}{f_\mu(\mu)} f_\mu - 1,$$

then $\deg g < |\Gamma|$ and clearly $g(\lambda) = 0$ for every $\lambda \in \Gamma$. Hence, by Lemma 2.1, $g = 0$. $\square$

Lemma 2.3. *For every $\xi \in \text{End}(V)$, the following statements are equivalent:*

- (1) $V = \bigoplus_{\lambda \in \Gamma} V_\lambda^\xi$;
- (2) $V = V_\Gamma^\xi$;
- (3) $f(\xi)V = 0$.

Proof. Clearly (1) implies (2) and (2) implies (3). Suppose (3) is satisfied. Then, for every $\mu \in \Gamma$, and $v \in V$,

$$\begin{aligned} 0 &= (f(\xi))(v) = \left(\prod_{\lambda \in \Gamma} (\xi - \lambda) \right) (v) \\ &= (\xi - \mu) \left(\prod_{\lambda \in \Gamma \setminus \{\mu\}} (\xi - \lambda) \right) (v) \\ &= (\xi - \mu)((f_\mu(\xi))(v)), \end{aligned}$$

whence $(f_\mu(\xi))(v) \in V_\mu^\xi$. Set

$$v_\mu := \frac{1}{f_\mu(\mu)} (f_\mu(\xi))(v).$$

By Corollary 2.2,

$$id_V = \sum_{\mu \in \Gamma} \frac{1}{f_\mu(\mu)} f_\mu(\xi)$$

and so

$$v = \sum_{\mu \in \Gamma} \frac{1}{f_\mu(\mu)} (f_\mu(\xi))(v) = \sum_{\mu \in \Gamma} v_\mu.$$

Now, assume $v = \sum_{\mu \in \Gamma} w_\mu$ for some $w_\mu \in V_\mu^\xi$. Since

$$\frac{1}{f_\mu(\mu)} (f_\mu(\xi))(w_\lambda) = \delta_{\lambda\mu} w_\mu$$

(where $\delta_{\lambda\mu}$ is the Kronecker delta), we get

$$v_\mu = \frac{1}{f_\mu(\mu)} (f_\mu(\xi))(v) = \sum_{\mu \in \Gamma} \frac{1}{f_\mu(\mu)} (f_\mu(\xi))(w_\lambda) = w_\mu,$$

giving (1). $\square$

Recall [2] that a *fusion law* is a pair $(\mathcal{S}, *)$ such that $\mathcal{S}$ is a set and $*$ is a map from the cartesian product $\mathcal{S} \times \mathcal{S}$ to the power set $2^{\mathcal{S}}$. A *morphism* between two fusion laws $(\mathcal{S}_1, *_1)$ and $(\mathcal{S}_2, *_2)$ is a map

$$\phi: \mathcal{S}_1 \rightarrow \mathcal{S}_2$$

such that, for $\alpha, \beta \in \mathcal{S}_1$,

$$\phi(\alpha *_1 \beta) \subseteq \phi(\alpha) *_2 \phi(\beta).$$

An *isomorphism* of fusion laws is a bijective morphism such that its inverse is also a morphism. A fusion law $(\mathcal{S}, *)$ is said to be *finite* if $\mathcal{S}$ is a finite set. In this paper we deal with fusion laws $(\mathcal{S}, *)$ where $\mathcal{S}$ is a finite set containing the spectrum of the adjoint action of an idempotent element in an R -algebra. Therefore, we assume $1_R \in \mathcal{S} \subseteq R$. Without loss of generality, we may also assume that $0_R \in \mathcal{S}$. Further, for every morphism ϕ of fusion laws, we'll assume that $1^\phi = 1$ and $0^\phi = 0$. More generally, when considering morphisms between fusion laws, one may want to preserve some possible algebraic relations between the elements of the set $\mathcal{S}$. To this aim, we call a morphism ϕ of fusion laws an *algebraic morphism* if it is a $\mathbb{Z}$ -linear map. An *axial algebra over R with generating set $\mathcal{A}$ and fusion law $(\mathcal{S}, \star)$* is a quadruple

$$(R, V, \mathcal{A}, (\mathcal{S}, \star))$$

such that

- (1) R is an associative commutative ring with identity 1;
- (2) $\mathcal{S}$ is a subset of R containing 1 and 0;
- (3) V is a commutative non associative R -algebra;
- (4) $\mathcal{A}$ is a set of idempotent elements (called *axes*) of V that generate V as an R -algebra and such that

$$\text{Ax1 } V = V_{\mathcal{S}}^a \text{ for every } a \in \mathcal{A};$$

$$\text{Ax2 } V_{\lambda}^a V_{\mu}^a \subseteq V_{\lambda \star \mu}^a \text{ for every } \lambda, \mu \in \mathcal{S} \text{ and } a \in \mathcal{A}.$$

Further, V is called *primitive* if,

$$\text{Ax3 } V_1^a = Ra \text{ for every } a \in \mathcal{A}.$$

A *Frobenius* axial algebra is an axial algebra $(R, V, \mathcal{A}, (\mathcal{S}, \star))$ endowed with an associative bilinear form $\kappa : V \times V \rightarrow R$ such that the map $a \mapsto \kappa(a, a)$ is constant on the set of axes.

Let $(R, V, \mathcal{A}, (\mathcal{S}, \star))$ be an axial algebra and assume the elements of $\mathcal{S}$ are pairwise distinguishable. As in the proof of Lemma 2.3, for every $v \in V$, denote by v_1 the projection of v into V_1^a with respect to the decomposition of V into ad_a -eigenspaces. If V is primitive, we have

$$v_1 = \lambda_a(v)a$$

for some $\lambda_a(v) \in R$ which is generally not unique. On the other hand, if the annihilator ideal

$$\text{Ann}_R(a) := \{r \in R \mid ra = 0\}$$

of a in R is trivial, then $\lambda_a(v)$ is unique, and we say that a is a *free* axis. Clearly this condition is satisfied when R is a field. As an immediate consequence we have the main result of this section.

Proposition 2.4. *Let $(R, V, \mathcal{A}, (\mathcal{S}, \star))$ be a primitive axial algebra and assume that the elements of $\mathcal{S}$ are pairwise distinguishable and the axes in $\mathcal{A}$ are free. Then, for every $a \in \mathcal{A}$, there is a well defined R -linear map*

$$(3) \quad \begin{array}{ccc} \lambda_a & : & V \rightarrow R \\ & & v \mapsto \lambda_a(v) \end{array}$$

such that every $v \in V$ decomposes uniquely as

$$(4) \quad v = \lambda_a(v)a + \sum_{\mu \in \mathcal{S} \setminus \{1\}} v_\mu,$$

with $v_\mu \in V_\mu^a$.

We say that V is *weak primitive* if, for every $a \in \mathcal{A}$ and every element $v \in V$, v can be decomposed as

$$(5) \quad v = l_a(v)a + \sum_{\mu \in \mathcal{S} \setminus \{1\}} v_\mu$$

where $l_a(v) \in R$ depends on v and a , and $v_\mu \in V_\mu^a$. Note that in general the decomposition in (5) and $l_a(v)$ are not uniquely determined by v .

Lemma 2.5. *If the elements of $\mathcal{S}$ are pairwise distinguishable, in particular, if R is a field, then weak primitivity is equivalent to primitivity.*

Proof. Trivially, primitivity implies weak primitivity. Conversely, suppose that V is weak primitive, fix $a \in \mathcal{A}$ and let $v_1 \in V_1^a$. Then, by weak primitivity, there exist $l \in R$, $v_\mu \in V_\mu^a$, such that

$$v_1 = la + \sum_{\mu \in \mathcal{S} \setminus \{1\}} v_\mu.$$

Hence

$$\sum_{\mu \in \mathcal{S} \setminus \{1\}} v_\mu = v_1 - la \in \left(\sum_{\mu \in \mathcal{S} \setminus \{1\}} V_\mu^a \right) \cap V_1^a,$$

and the last intersection is trivial by Lemma 2.3. Thus $v_1 = la \in Ra$. $\square$

We conclude this section with the following straightforward observation.

Lemma 2.6. *Let $(R, V, \mathcal{A}, (\mathcal{S}, \star))$ be a primitive axial algebra and assume that the elements of $\mathcal{S}$ are pairwise distinguishable, and the axes in $\mathcal{A}$ are free. Then, for every $a \in \mathcal{A}$, $\gamma, \delta \in \mathcal{S}$, and $v, w \in V$, we have*

$$(f_1(\text{ad}_a))(w - \lambda_a(w)a) = 0$$

and

$$\left(\prod_{\eta \in \gamma \star \delta} (\text{ad}_c - \eta) \right) ((f_\gamma(\text{ad}_c))(v) \cdot (f_\delta(\text{ad}_c))(w)) = 0.$$

3. UNIVERSAL PRIMITIVE AXIAL ALGEBRAS

In this section we fix a positive integer k , a fusion law $(\mathcal{S}_0, *_{\mathcal{S}_0})$, and denote by $\mathcal{O}$ the class whose objects are the primitive axial algebras

$$(R, V, \mathcal{A}, (\mathcal{S}, *))$$

such that

- O1 $\mathcal{A}$ has size at most k and its elements are free axes,
- O2 $(\mathcal{S}, *)$ is isomorphic to $(\mathcal{S}_0, *_{\mathcal{S}_0})$, and
- O3 the elements of $\mathcal{S}$ are pairwise distinguishable in R .

For two elements

$$(6) \quad \mathcal{V}_1 := (R_1, V_1, \mathcal{A}_1, (\mathcal{S}_1, *_1)) \text{ and } \mathcal{V}_2 := (R_2, V_2, \mathcal{A}_2, (\mathcal{S}_2, *_2))$$

in $\mathcal{O}$, let $\mathcal{H}om(\mathcal{V}_1, \mathcal{V}_2)$ be the set of maps

$$\phi: R_1 \cup V_1 \rightarrow R_2 \cup V_2$$

satisfying the following conditions:

- H1 $\phi|_{R_1}$ is an homomorphism of rings with identity between R_1 and R_2 that induces by restriction an isomorphism of fusion laws between $(\mathcal{S}_1, *_1)$ and $(\mathcal{S}_2, *_2)$;
- H2 $\phi|_{V_1}$ is a (non-associative) ring homomorphism between V_1 and V_2 such that $\phi(\mathcal{A}_1) \subseteq \mathcal{A}_2$;
- H3 $(\gamma v)^\phi = \gamma^\phi v^\phi$, for every $\gamma \in R_1$ and $v \in V_1$.

Note that, since $\phi|_{R_1}$ is a ring homomorphism, the induced isomorphism of fusion law in (H1) is in fact an algebraic isomorphism. Denote by $\mathcal{H}$ the class of all $\phi \in \mathcal{H}om(\mathcal{V}_1, \mathcal{V}_2)$ where $\mathcal{V}_1$ and $\mathcal{V}_2$ range in $\mathcal{O}$. Then clearly $(\mathcal{O}, \mathcal{H})$ is a category.

It will turn convenient to define some subcategories of $(\mathcal{O}, \mathcal{H})$ in the following way. Let

- $n := |\mathcal{S} \setminus \{0, 1\}|$;
- $x_i, y_i, w_i, z_{i,j}, t_h$ be algebraically independent indeterminates over $\mathbb{Z}$, for $i, j \in \{1, \dots, n\}$, with $i < j$, $h \in \mathbb{N}$;
- D be the polynomial ring

$$\mathbb{Z}[x_i, y_i, w_i, z_{i,j}, t_h \mid i, j \in \{1, \dots, n\}, i < j, h \in \mathbb{N}];$$

- L be a proper ideal of D containing the set

$$\Sigma := \{x_i y_i - 1, (1 - x_i)w_i - 1, \text{ and } (x_i - x_j)z_{i,j} - 1 \text{ for all } 1 \leq i < j \leq n\};$$

- $\hat{D} := D/L$. For $d \in D$, we denote the element $L + d$ by $\hat{d}$.

The extra indeterminates t_h in the definition of D have been introduced here in order to guarantee, when necessary, the invertibility of certain elements in the ring $\hat{D}$.

Since L is proper and, for $1 \leq i < j \leq n$, the elements $\hat{x}_i - \hat{x}_j$ are invertible in $\hat{D}$, the elements $\hat{x}_1, \dots, \hat{x}_n$ are still pairwise distinguishable in $\hat{D}$. Define $(\mathcal{O}_L, \mathcal{H}_L)$ as the full subcategory of $(\mathcal{O}, \mathcal{H})$ whose objects are the primitive axial algebras $(R, V, \mathcal{A}, (\mathcal{S}, *)) \in \mathcal{O}$ that satisfy the further condition

- O4 R is a ring containing a subring isomorphic to a factor of $\hat{D}$.

Clearly, if (Σ) is the ideal of D generated by the set Σ , then $\mathcal{O}_{(\Sigma)} = \mathcal{O}$.

Lemma 3.1. *Let $\mathcal{V}_1, \mathcal{V}_2 \in \mathcal{O}_L$ be as in Equation (6) and let $\phi \in \mathcal{H}om(\mathcal{V}_1, \mathcal{V}_2)$. Then, for every $a \in \mathcal{A}_1, v \in V_1$, we have*

$$\lambda_{a^\phi}(v^\phi) = (\lambda_a(v))^\phi.$$

Proof. Since $\mathcal{V}_1, \mathcal{V}_2$ are primitive axial algebras, the result follows applying ϕ to the decomposition of v in Equation (4). $\square$

We construct a universal object for each category $(\mathcal{O}_L, \mathcal{H}_L)$ as follows. Let

- $\mathcal{A}$ be a set of size k ;
- W be the free commutative magma generated by the elements of $\mathcal{A}$ subject to the condition that every element of $\mathcal{A}$ is idempotent;

- $\hat{R} := \hat{D}[\Lambda]$ be the ring of polynomials with coefficients in $\hat{D}$ and indeterminates set $\Lambda := \{\lambda_{c,w} \mid c \in \mathcal{A}, w \in W, c \neq w\}$ where $\lambda_{c,w} = \lambda_{c',w'}$ if and only if $c = c'$ and $w = w'$.
- $\hat{V} := \hat{R}[W]$ be the set of all formal linear combinations $\sum_{w \in W} \gamma_w w$ of the elements of W with coefficients in $\hat{R}$, with only finitely many coefficients different from zero. Endow $\hat{V}$ with the usual structure of a commutative non associative $\hat{R}$ -algebra;
- $\hat{\mathcal{S}}$ be the set $\{1, 0, \hat{x}_1, \dots, \hat{x}_n\}$.

Let $\star: \hat{\mathcal{S}} \times \hat{\mathcal{S}} \rightarrow 2^{\hat{\mathcal{S}}}$ be a map such that $(\hat{\mathcal{S}}, \star)$ is a fusion law isomorphic to $(\mathcal{S}, *)$. Since, obviously, a fusion law is isomorphic to $(\mathcal{S}, *)$ if and only if it is isomorphic to $(\hat{\mathcal{S}}, \star)$, we may assume $(\mathcal{S}, *) = (\hat{\mathcal{S}}, \star)$. For $\mu \in \hat{\mathcal{S}}$, let f_μ be the polynomial defined in Equation (1), for every $c \in \mathcal{A}$, let $\lambda_{c,c} := 1$, and let J be the ideal of $\hat{V}$ generated by all the elements

$$(7) \quad (f_1(\text{ad}_c))(w - \lambda_{c,w}c) \quad \text{for all } c \in \mathcal{A} \text{ and } w \in W$$

and

$$(8) \quad \left(\prod_{\eta \in \gamma \star \delta} (\text{ad}_c - \eta \text{id}_{\hat{V}}) \right) ((f_\gamma(\text{ad}_c))(v) \cdot (f_\delta(\text{ad}_c))(w))$$

for all $v, w \in \hat{V}$, $\gamma, \delta \in \hat{\mathcal{S}}$, $c \in \mathcal{A}$. Let I_0 be the ideal of $\hat{R}$ generated by all the elements

$$(9) \quad \sum_{w \in W} \gamma_w \lambda_{c,w} \quad \text{for all } c \in \mathcal{A}, w \in W, \gamma_w \in \hat{R} \text{ such that } \sum_{w \in W} \gamma_w w \in J.$$

Finally, set

- $\mathcal{A}/J := \{c + J \mid c \in \mathcal{A}\}$
- $J_0 := J + I_0 \hat{V}$,
- $\overline{R} := \hat{R}/I_0$,
- $\overline{V} := \hat{V}/J_0$,
- $\alpha_i := x_i + I_0$, for $i \in \{1, \dots, n\}$
- $\bar{c} := c + J_0$, for $c \in \mathcal{A}$,
- $\overline{\mathcal{A}} := \{\bar{c} \mid c \in \mathcal{A}\}$,
- $\overline{\mathcal{S}} := \{1 + I_0, 0 + I_0, \alpha_i \mid i \in \{1, \dots, n\}\}$.

Remark 3.2. Since $\hat{D} \cap I_0 = \{0\}$ and $\overline{\mathcal{S}} \leq \hat{D}I_0/I_0$, $(\overline{\mathcal{S}}, \overline{\star})$ is a fusion law isomorphic to $(\mathcal{S}, *)$ and the elements of $\overline{\mathcal{S}}$ are pairwise distinguishable.

Lemma 3.3. $(\hat{R}, \hat{V}/J, \mathcal{A}/J, (\hat{\mathcal{S}}, \star))$ and $(\hat{R}, \overline{V}, \overline{\mathcal{A}}, (\hat{\mathcal{S}}, \star))$ are primitive axial algebras such that, for $J_* \in \{J, J_0\}$,

$$\lambda_{c+J_*}(w + J_*) = \lambda_{c,w}$$

for every $c \in \mathcal{A}$ and $w \in W$.

Proof. Clearly $\hat{V}/J$ is generated by $\mathcal{A}/J$ and $\overline{V}$ is generated by $\overline{\mathcal{A}}$. Let $J_* \in \{J, J_0\}$ and let f be as in Equation (2). By the definition of J , for every $a \in \mathcal{A}$, $f(\text{ad}_a)\hat{V} \subseteq J$, whence, by Lemma 2.3, $\hat{V}/J_*$ satisfies condition Ax1. By Equation (8), $\hat{V}/J_*$

also satisfies the fusion law $(\hat{\mathcal{S}}, \star)$. Furthermore, for every $c \in \mathcal{A}$ and $w \in W$, since $f_1(\text{ad}_c)(w - \lambda_{c,w}c) \in J_*$, we get

$$w + J_* = (\lambda_{c,w}c + \sum_{\mu \in \hat{\mathcal{S}} \setminus \{1\}} w_\mu) + J_*,$$

where $w_\mu + J_*$ is a μ -eigenvector for ad_{c+J_*} , so $\hat{V}/J_*$ is also weak primitive. By Remark 3.2 and Lemma 2.5, $\hat{V}/J_*$ is also primitive. $\square$

Lemma 3.4. *Let $\mathcal{V} := (R, V, \mathcal{A}, (\mathcal{S}, *))$ be an element of $\mathcal{O}_L$, let $\phi: \hat{R} \cup \hat{V} \rightarrow R \cup V$ be a map that satisfies conditions (H1) (with respect to $(\hat{\mathcal{S}}, \star)$ and $(\mathcal{S}, *)$), (H2), and (H3). Then $I_0 \subseteq \ker \phi|_{\hat{R}}$, $J_0 \subseteq \ker \phi|_{\hat{V}}$.*

Proof. By Lemma 2.6, $J \subseteq \ker \phi|_{\hat{V}}$, so ϕ induces an $\hat{R}$ -algebra homomorphism

$$\begin{aligned} \phi_{\hat{V}/J}: \hat{V}/J &\rightarrow V, \\ v + J &\mapsto v^\phi. \end{aligned}$$

Since, by Lemma 3.3, $\hat{V}/J$ is a primitive axial algebra over the ring $\hat{R}$, for every $c \in \mathcal{A}$ and $w \in W$, we can write

$$w + J = (\lambda_{c,w}c + \sum_{\mu \in \hat{\mathcal{S}} \setminus \{1\}} w_\mu) + J,$$

where, for every $\mu \in \hat{\mathcal{S}} \setminus \{1\}$, $w_\mu + J$ is a μ -eigenvector for ad_{c+J} . Condition (H3) implies that, for every $\mu \in \mathcal{S}$, $a \in \mathcal{A}$, $\phi_{\hat{V}/J}$ maps μ -eigenvectors for ad_{a+J} to μ^ϕ -eigenvectors for ad_{a^ϕ} . Thus, if $v = \sum_{w \in W} \gamma_w w \in J$, then $v^\phi = 0$, whence

$$\begin{aligned} 0 &= \lambda_{a^\phi}(v^\phi) \\ &= \lambda_{a^\phi} \left(\sum_{w \in W} \gamma_w^\phi w^\phi \right) \\ &= \sum_{w \in W} \gamma_w^\phi \lambda_{a^\phi}(w^\phi) \\ &= \sum_{w \in W} \gamma_w^\phi \lambda_{a^\phi}((\lambda_{a,w}a + \sum_{\mu \in \hat{\mathcal{S}} \setminus \{1\}} w_\mu)^\phi) \\ &= \sum_{w \in W} \gamma_w^\phi \lambda_{a^\phi}((\lambda_{a,w})^\phi a^\phi) + \sum_{w \in W} \gamma_w^\phi \lambda_{a^\phi} \left(\sum_{\mu \in \hat{\mathcal{S}} \setminus \{1\}} w_\mu^\phi \right) \\ &= \sum_{w \in W} \gamma_w^\phi (\lambda_{a,w})^\phi = \left(\sum_{w \in W} \gamma_w \lambda_{a,w} \right)^\phi. \end{aligned}$$

Thus $I_0 \subseteq \ker \phi|_{\hat{R}}$. Finally, by condition (H3), $(I_0 \hat{V})^\phi = I_0^\phi V^\phi = 0V_1 = 0$, whence $J_0 \subseteq \ker \phi|_{\hat{V}}$. $\square$

Lemma 3.5. *We have $J_0 \neq \hat{V}$, in particular $|\overline{\mathcal{A}}| = k$.*

Proof. Let $\overline{R}^k$ be the direct sum of k copies of $\overline{R}$. Set $\mathcal{B} := \{e_1, \dots, e_k\}$, where $(e_1, \dots, e_k)$ is the canonical basis of $\overline{R}^k$. Then, for every $i \in \{1, \dots, k\}$, e_i is an idempotent and

$$\overline{R}^k = \overline{R}e_i \oplus \ker \text{ad}_{e_i}.$$

Therefore, for every fusion law $(\mathcal{S}_{\overline{R}}, *_{\overline{R}})$, the 4-tuple

$$(\overline{R}, \overline{R}^k, \mathcal{B}, (\mathcal{S}_{\overline{R}}, *_{\overline{R}}))$$

is obviously a primitive (associative) axial algebra. By the construction of $\hat{V}$, any bijection from $\mathcal{A}$ to $\mathcal{B}$ extends uniquely to a map $\phi_{\hat{V}}: \hat{V} \rightarrow \overline{R}^k$. Let

$$\phi: \hat{R} \cup \hat{V} \rightarrow \overline{R} \cup \overline{R}^k$$

be the map whose restrictions to $\hat{R}$ and $\hat{V}$ are the canonical projection on $\overline{R}$ and $\phi_{\hat{V}}$, respectively. Then ϕ satisfies the conditions (H1), (H2), and (H3). Therefore, by Lemma 3.4, $J_0 \subseteq \ker \phi|_{\hat{V}} \neq \hat{V}$. Since $k = |\mathcal{A}| \geq |\overline{\mathcal{A}}| \geq |\mathcal{B}| = k$, we get $|\overline{\mathcal{A}}| = k$. $\square$

Theorem 3.6. $\overline{\mathcal{V}} := (\overline{R}, \overline{V}, \overline{\mathcal{A}}, (\overline{\mathcal{S}}, \overline{\star}))$ is a universal object in the category $(\mathcal{O}_L, \mathcal{H}_L)$.

Proof. Clearly $\overline{V}$ is an algebra over $\overline{R}$ generated by the set of idempotents $\overline{\mathcal{A}}$. Since by Lemma 3.3, $(\hat{R}, \overline{V}, \overline{\mathcal{A}}, (\hat{\mathcal{S}}, \overline{\star}))$ is a primitive axial algebra, and $I_0 \subseteq \text{Ann}_{\hat{R}}(\overline{V})$, we get that $(\overline{R}, \overline{V}, \overline{\mathcal{A}}, (\overline{\mathcal{S}}, \star))$ is a primitive axial algebra. Finally, the axes in $\overline{\mathcal{A}}$ are free, for, if $c \in \mathcal{A}$ and $r \in \hat{R}$ are such that $rc \in J_0$, then there exist $j \in J, i_0 \in I_0$ and $\sum_{w \in W} \gamma_w w \in \hat{V}$ such that

$$rc - i_0 \left(\sum_{w \in W} \gamma_w w \right) = j,$$

whence, by the definition of I_0 ,

$$r \in i_0 \left(\sum_{w \in W} \gamma_w \lambda_{c,w} \right) + I_0 = I_0.$$

By Lemma 3.5, the elements of $\hat{\mathcal{S}}$ are pairwise distinguishable in $\overline{R}$, whence $\overline{\mathcal{V}} \in \mathcal{O}_L$.

Now assume $\mathcal{V}_1 := (R_1, V_1, \mathcal{A}_1, (\mathcal{S}_1, *_1))$ is an object in $\mathcal{O}_L$ and let $\bar{t}: \overline{\mathcal{A}} \rightarrow \mathcal{A}_1$ a map of sets. Since every non-empty subset of $\mathcal{A}_1$ generates a primitive axial subalgebra of V_1 with fusion law $(\mathcal{S}_1, *_1)$ and free axes, without loss of generality we may assume that $\bar{t}$ is surjective. Let t be the composition of $\bar{t}$ with the (bijective) projection of $\mathcal{A}$ to $\overline{\mathcal{A}}$. Since W is the free commutative magma generated by the set of idempotents $\mathcal{A}$ there is a unique magma homomorphism

$$\chi: W \rightarrow V_1,$$

inducing the map $t: \mathcal{A} \rightarrow \mathcal{A}_1$. Since the elements of Λ are algebraically independent over $\hat{D}$, there is a unique homomorphism of $\hat{D}$ -algebras

$$\hat{\psi}: \hat{R} \rightarrow R_1$$

such that, for $c \in \mathcal{A}$ and $w \in W \setminus \{c\}$,

$$(10) \quad \lambda_{c,w}^{\hat{\psi}} = \lambda_{c^t}(w^\chi),$$

where λ_{c^t} is the function defined in Proposition 2.4. Define

$$\begin{aligned} \hat{\chi} : \quad \hat{V} &\rightarrow V_1 \\ \sum_{w \in W} \gamma_w w &\mapsto \sum_{w \in W} \gamma_w^{\hat{\psi}} w^\chi. \end{aligned}$$

Then $\hat{\chi}$ is a ring homomorphism extending t and such that $(\gamma v)^{\hat{\chi}} = \gamma^{\hat{\psi}} v^{\hat{\chi}}$ for every $\gamma \in \hat{R}$ and $v \in \hat{V}$. Since, for every $c \in \mathcal{A}$, $v \in \hat{V}$, and $\gamma \in \mathcal{S}$,

$$[(\text{ad}_c - \gamma \text{id}_{\hat{V}})(v)]^{\hat{\chi}} = (\text{ad}_{c^t} - \gamma^{\hat{\psi}} \text{id}_{V_1})(v^{\hat{\chi}}),$$

by Lemma 2.6, it follows that J_0 is contained in $\ker \hat{\chi}$ and so $\hat{\chi}$ induces a ring homomorphism

$$\phi_V : \overline{V} \rightarrow V_1$$

extending $\bar{t}$ and such that $(\gamma \bar{w})^{\phi_V} = \gamma^{\hat{\psi}} \bar{w}^{\phi_V}$ for every $\gamma \in \hat{R}$ and $\bar{w} \in \overline{V}$. As in the proof of Lemma 3.4 we get that $I_0 \subseteq \ker \hat{\psi}$. Let $\phi_R : \overline{R} \rightarrow R_1$ be the homomorphism of $\hat{D}$ -algebras induced by $\hat{\psi}$. Then $(\phi_R, \phi_V) \in \mathcal{H}om(\overline{\mathcal{V}}, \mathcal{V}_1)$.

Since $\hat{R} = \hat{D}[\Lambda]$, ϕ_R is completely determined by its values on the elements $\lambda_{c,w} + I_0$. Further, by Equation (10), for every $c \in \mathcal{A}$, $w \in W \setminus \{c\}$,

$$(\lambda_{c,w} + I_0)^{\phi_R} = \lambda_{c^t}((w + J_0)^{\phi_V}),$$

whence ϕ_R is completely determined by the images $(w + J_0)^{\phi_V}$, with $w \in W$. Since ϕ_V is a ring homomorphism extending t , such images are uniquely determined, whence the uniqueness of (ϕ_R, ϕ_V) . $\square$

Corollary 3.7. *Every permutation σ of the set $\overline{\mathcal{A}}$ extends to a unique automorphism $f_\sigma \in \mathcal{H}om(\overline{\mathcal{V}}, \overline{\mathcal{V}})$.*

Note that, for a generic object $\mathcal{V}$, the above assertion is false (see e.g. the algebra $Q_2(\eta)$ constructed in [5, Section 5.3]). We say that $\mathcal{V} = (R, V, \mathcal{A}, (\mathcal{S}, *)) \in \mathcal{O}_L$ is *symmetric* if every permutation σ of the set $\mathcal{A}$ extends to a unique automorphism $f_\sigma \in \mathcal{H}om(\mathcal{V}, \mathcal{V})$.

Corollary 3.8. *Let $\mathcal{V} := (R, V, \mathcal{A}, (\mathcal{S}, *)) \in \mathcal{O}_L$, then*

- (1) $\overline{\mathcal{V}} \otimes R := (\overline{R} \otimes_{\hat{D}} R, \overline{V} \otimes_{\hat{D}} R, \overline{\mathcal{A}} \otimes_{\hat{D}} 1, (\overline{\mathcal{S}} \otimes_{\hat{D}} 1, \overline{*})) \in \mathcal{O}_L$,
- (2) R is isomorphic to a factor of $\overline{R} \otimes_{\hat{D}} R$,
- (3) V is isomorphic to a factor of $\overline{V} \otimes_{\hat{D}} R$.

Remark 3.9. *Note that, with the notation of Corollaries 3.7 and 3.8, $f_\sigma \otimes \text{id}_R$ is an automorphism of $\overline{\mathcal{V}} \otimes R$.*

Questions

- (1) *Can we define a variety of axial algebras corresponding to the fusion law $(\mathcal{S}, *)$?*
- (2) *Is it true that any ideal I of $\hat{R}$ containing I_0 defines an axial algebra?*

4. 2-GENERATED PRIMITIVE AXIAL ALGEBRAS OF MONSTER TYPE (α, β)

In this section we keep the notation of Section 3, with $k = n = 2$, $(\mathcal{S}, *)$ equal to the Monster fusion law $\mathcal{M}(\alpha, \beta)$, and L an ideal of D containing $2t_1 - 1$, so that (the class of) 2 is invertible in $\hat{D}$. In order to simplify notation we'll also identify α_1 with α and α_2 with β .

Let $\mathcal{V} = (V, R, \mathcal{A}, (\mathcal{S}, *)) \in \mathcal{O}_L$ and $a \in \mathcal{A}$. Let $\mathcal{S}^+ := \{1, 0, \alpha\}$ and $\mathcal{S}^- := \{\beta\}$. The partition $\{\mathcal{S}^+, \mathcal{S}^-\}$ of $\mathcal{S}$ induces a $\mathbb{Z}_2$ -grading on $\mathcal{S}$ which, on turn, induces, a $\mathbb{Z}_2$ -grading $\{V_+^a, V_-^a\}$ on V where $V_+^a := V_1^a + V_0^a + V_\alpha^a$ and $V_-^a = V_\beta^a$. It follows

that, if τ_a is the map from $R \cup V$ to $R \cup V$ such that $\tau_{a|_V}$ inverts V_β^a and leaves invariant the elements of V_+^a and $\tau_{a|R}$ is the identity, then τ_a is an involutory automorphism of $\mathcal{V}$ (see [7, Proposition 3.4]). The map τ_a is called the *Miyamoto involution* associated to the axis a . By definition of τ_a , the element $av - \beta v$ of V is τ_a -invariant and, since a lies in $V_+^a \leq C_V(\tau_a)$, also $av - \beta(a + v)$ is τ_a -invariant. In particular, by symmetry,

Lemma 4.1. *Let a and b be axes of V . Then $ab - \beta(a + b)$ is fixed by the 2-generated group $\langle \tau_a, \tau_b \rangle$.*

Let $\mathcal{A} := \{a_0, a_1\}$ and, for $i \in \{1, 2\}$, let τ_i be the Miyamoto involutions associated to a_i . Set $\rho := \tau_0\tau_1$, and for $i \in \mathbb{Z}$, $a_{2i} := a_0^{\rho^i}$ and $a_{2i+1} := a_1^{\rho^i}$. Since ρ is an automorphism of V , for every $j \in \mathbb{Z}$, a_j is an axis. Denote by $\tau_j := \tau_{a_j}$ the corresponding Miyamoto involution.

Lemma 4.2. *For every $n \in \mathbb{N}$, and $i, j \in \mathbb{Z}$ such that $i \equiv j \pmod n$ we have*

$$a_i a_{i+n} - \beta(a_i + a_{i+n}) = a_j a_{j+n} - \beta(a_j + a_{j+n}),$$

Proof. This follows immediately from Lemma 4.1. $\square$

For $n \in \mathbb{N}$ and $r \in \{0, \dots, n-1\}$ set

$$(11) \quad s_{r,n} := a_r a_{r+n} - \beta(a_r + a_{r+n}).$$

For every $a \in \mathcal{A}$, let λ_a be as in Proposition 2.4.

Lemma 4.3. *For $i \in \{1, 2, 3\}$ we have*

$$a_0 s_{0,i} = (\alpha - \beta) s_{0,i} + [(1 - \alpha) \lambda_{a_0}(a_i) + \beta(\alpha - \beta - 1)] a_0 + \frac{1}{2} \beta(\alpha - \beta)(a_i + a_{-i}).$$

Proof. This is [16, Lemma 3.1]. $\square$

For $i \in \{1, 2, 3\}$, let

$$(12) \quad a_i = \lambda_{a_0}(a_i) a_0 + u_i + v_i + w_i$$

be the decomposition of a_i into ad_{a_0} -eigenvectors, where u_i is a 0-eigenvector, v_i is an α -eigenvector and w_i is a β -eigenvector.

Lemma 4.4. *With the above notation,*

- (1) $u_i = \frac{1}{\alpha}((\lambda_{a_0}(a_i) - \beta - \alpha \lambda_{a_0}(a_i)) a_0 + \frac{1}{2}(\alpha - \beta)(a_i + a_{-i}) - s_{0,i});$
- (2) $v_i = \frac{1}{\alpha}((\beta - \lambda_{a_0}(a_i)) a_0 + \frac{\beta}{2}(a_i + a_{-i}) + s_{0,i});$
- (3) $w_i = \frac{1}{2}(a_i - a_{-i}).$

Proof. (3) follows from the definitions of τ_0 and a_i , (2) is just a rearranging of $a_0 a_i = \lambda_{a_0}(a_i) a_0 + \alpha v_i + \beta w_i$ using Equation (11), and (1) follows rearranging Equation (12). $\square$

For $i, j \in \{1, 2, 3\}$, set

$$P_{ij} := u_i u_j + u_i v_j \quad \text{and} \quad Q_{ij} := u_i v_j - \frac{1}{\alpha^2} s_{0,i} s_{0,j}.$$

Lemma 4.5. *For $i, j \in \{1, 2, 3\}$ we have*

$$(13) \quad s_{0,i} \cdot s_{0,j} = \alpha(a_0 P_{ij} - \alpha Q_{ij}).$$

Proof. Since u_i and v_j are a 0-eigenvector and an α -eigenvector for ad_{a_0} , respectively, by the fusion rule, we have $a_0 P_{ij} = \alpha(u_i \cdot v_j)$ and the result follows. $\square$

From now on we assume $\mathcal{V} = \overline{\mathcal{V}}$. Let f be the automorphism of $\overline{\mathcal{V}}$ induced by the permutation that swaps a_0 with a_1 as defined in Corollary 3.7. For $i \in \mathbb{N}$ define

$$(14) \quad \lambda_i := \lambda_{a_0}(a_i).$$

Note that, by Lemma 3.1, for every $i \in \mathbb{N}$, we have

$$\lambda_{a_0}(a_{-i}) = \lambda_i, \quad \lambda_{a_1}(a_0) = \lambda_1^f, \quad \text{and} \quad \lambda_{a_1}(a_{-1}) = \lambda_2^f.$$

Set $T_0 := \langle \tau_0, \tau_1 \rangle$ and $T := \langle \tau_0, f \rangle$.

Lemma 4.6. *The groups T_0 and T are dihedral groups, T_0 is a normal subgroup of T such that $|T : T_0| \leq 2$. For every $n \in \mathbb{N}$, the set $\{s_{0,n}, \dots, s_{n-1,n}\}$ is invariant under the action of T . In particular, if K_n is the kernel of this action, we have*

- (1) $K_1 = T$;
- (2) $K_2 = T_0$, in particular $s_{0,2}^f = s_{1,2}$;
- (3) T/K_3 induces the full permutation group on the set $\{s_{0,3}, s_{1,3}, s_{2,3}\}$ with point stabilisers generated by $\tau_0 K_3$, $\tau_1 K_3$ and $f K_3$, respectively. In particular $s_{0,3}^f = s_{1,3}$ and $s_{0,3}^{\tau_1} = s_{2,3}$.

Proof. This follows immediately from the definitions. $\square$

Lemma 4.7. *In the algebra $\overline{\mathcal{V}}$, the following equalities hold:*

$$\begin{aligned} & (\alpha - 2\beta)a_0s_{1,2} = \\ & \quad \beta^2(\alpha - \beta)(a_{-2} + a_2) \\ & \quad + \left[-2\alpha\beta\lambda_1 + 2\beta(1 - \alpha)\lambda_1^f + \frac{\beta}{2}(4\alpha^2 - 2\alpha\beta - \alpha + 4\beta^2 - 2\beta) \right] (a_1 + a_{-1}) \\ & \quad + \frac{1}{(\alpha - \beta)} \left[(6\alpha^2 - 8\alpha\beta - 2\alpha + 4\beta)\lambda_1^2 + (2\alpha^2 - 2\alpha)\lambda_1\lambda_1^f \right. \\ & \quad + 2(-2\alpha^2 - 2\alpha\beta + \alpha)(\alpha - \beta)\lambda_1 - 4\beta(\alpha - 1)(\alpha - \beta)\lambda_1^f - \alpha\beta(\alpha - \beta)\lambda_2 \\ & \quad \left. + (4\alpha^2\beta - 2\alpha\beta + 2\beta^3)(\alpha - \beta) \right] a_0 \\ & \quad + \left[-4\alpha\lambda_1 - 4(\alpha - 1)\lambda_1^f + (4\alpha^2 - 2\alpha\beta - \alpha + 4\beta^2 - 2\beta) \right] s_{0,1} \\ & \quad + 2\beta(\alpha - \beta)s_{0,2} \end{aligned}$$

and

$$\begin{aligned} & 4(\alpha - 2\beta)s_{0,1} \cdot s_{0,1} = \\ & \quad \beta(\alpha - \beta)^2(\alpha - 4\beta)(a_{-2} + a_2) \\ & \quad + \left[4\alpha\beta(\alpha - \beta)\lambda_1 + 2(-\alpha^3 + 5\alpha^2\beta + \alpha^2 - 4\alpha\beta^2 - 5\alpha\beta + 4\beta^2)\lambda_1^f \right. \\ & \quad \left. + \beta(-10\alpha^2\beta - \alpha^2 + 14\alpha\beta^2 + 7\alpha\beta - 4\beta^3 - 6\beta^2) \right] (a_{-1} + a_1) \\ & \quad + 2 \left[2(-3\alpha^2 + 4\alpha\beta + \alpha - 2\beta)\lambda_1^2 + 2\alpha(1 - \alpha)\lambda_1\lambda_1^f \right. \\ & \quad + 2(\alpha^3 + 4\alpha^2\beta - 6\alpha\beta^2 - 3\alpha\beta + 4\beta^2)\lambda_1 + 2\alpha\beta(\alpha - 1)\lambda_1^f + \alpha\beta(\alpha - \beta)\lambda_2 \\ & \quad \left. + \beta(-\alpha^3 - 8\alpha^2\beta + 13\alpha\beta^2 + 4\alpha\beta - 4\beta^3 - 4\beta^2) \right] a_0 \\ & \quad + 4 \left[2\alpha(\alpha - \beta)\lambda_1 + \alpha(\alpha - 1)\lambda_1^f + (-6\alpha^2\beta + 10\alpha\beta^2 + \alpha\beta - 4\beta^3) \right] s_{0,1} \\ & \quad - 2\alpha\beta(\alpha - \beta)s_{0,2} + 2\beta(\alpha - \beta)(\alpha - 2\beta)s_{1,2}. \end{aligned}$$

Proof. By the fusion law,

$$(15) \quad a_0(u_1 \cdot u_1 - v_1 \cdot v_1 + \lambda_{a_0}(v_1 \cdot v_1)a_0) = 0.$$

Substituting in the left side of (15) the values for u_1 and v_1 given in Lemma 4.4 we get the first equality. The expression for $(\alpha - 2\beta)a_{s_{1,2}}$ allows us to write explicitly as a linear combination of $a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1}, s_{0,2}, s_{1,2}$ the vector

$$(\alpha - 2\beta)(a_0P_{11} - \alpha Q_{11}).$$

Thus, the second equality then follows from Equation (13) in Lemma 4.5. $\square$

Lemma 4.8. *In the algebra $\overline{\mathcal{V}}$, the following equalities hold:*

- (1) $\beta(\alpha - \beta)^2(\alpha - 4\beta)(a_3 - a_{-2}) = c,$
- (2) $\beta(\alpha - \beta)^2(\alpha - 4\beta)(a_4 - a_{-1}) = -c^{\tau_1},$

where

$c =$

$$\begin{aligned} & (\alpha - \beta) \left[4\alpha\beta\lambda_1 - 2(\alpha - 1)(\alpha - 4\beta)\lambda_1^f + \beta(-\alpha^2 - 5\alpha\beta - \alpha + 6\beta) \right] a_{-1} \\ & + \left[4(-3\alpha^2 + 4\alpha\beta + \alpha - 2\beta)\lambda_1^2 - 4\alpha(\alpha - 1)\lambda_1\lambda_1^f \right. \\ & + (6\alpha^3 + 6\alpha^2\beta - 2\alpha^2 - 16\alpha\beta^2 - 2\alpha\beta + 8\beta^2)\lambda_1 + 4\alpha\beta(\alpha - 1)\lambda_1^f + 2\alpha\beta(\alpha - \beta)\lambda_2 \\ & + \left. \beta(\alpha - \beta)(-2\alpha^2 - 8\alpha\beta + \alpha + 4\beta^2 + 2\beta) \right] a_0 \\ & + \left[4\alpha(\alpha - 1)\lambda_1\lambda_1^f + 4(13\alpha^2 - 4\alpha\beta - \alpha + 2\beta)\lambda_1^{f^2} - 4\alpha\beta(\beta - 1)\lambda_1 \right. \\ & + (-6\alpha^3 - 6\alpha^2\beta + 2\alpha^2 + 16\alpha\beta^2 + 2\alpha\beta - 8\beta^2)\lambda_1^f - 2\alpha\beta(\alpha - \beta)\lambda_2^f \\ & + \left. \beta(\alpha - \beta)(2\alpha^2 + 8\alpha\beta - \alpha - 4\beta^2 - 2\beta) \right] a_1 \\ & + (\alpha - \beta) \left[2(\alpha - 1)(\alpha - 4\beta)\lambda_1 - 4\alpha\beta\lambda_1^f + \beta(\alpha^2 + 5\alpha\beta + \alpha - 6\beta) \right] a_2 \\ & + (\alpha - \beta) \left[4\alpha(\alpha - 2\beta + 1)(\lambda_1 - \lambda_1^f) \right] s_{0,1} \\ & - 4\beta(\alpha - \beta)^2(s_{0,2} - s_{1,2}). \end{aligned}$$

Proof. Since $s_{0,1}s_{0,1}$ is invariant under f , we have $4(\alpha - 2\beta)[s_{0,1}s_{0,1} - (s_{0,1}s_{0,1})^f] = 0$. Then, equality (1) follows from the expression for $4(\alpha - 2\beta)s_{0,1}s_{0,1}$ given in Lemma 4.7. By applying τ_1 to equation in (1) we get (2). $\square$

From the formulae in Lemmas 4.7 and 4.8, it is clear that we have different pictures according whether $\alpha - 2\beta$ and $\alpha - 4\beta$ are invertible in $\overline{\mathcal{R}}$ or not. Since we are most concerned with algebras over a field, later we will also assume that $\alpha - 2\beta$ and $\alpha - 4\beta$ are either invertible or zero. Thus we deal separately with the following three cases:

- (1) The generic case: the ideal L contains the elements t_h for $h \in \mathbb{N}$, and $h \geq 3$, $2t_1 - 1$, $(x_1 - 2x_2)t_2 - 1$, and $(x_1 - 4x_2)t_3 - 1$. In this case we set $\mathcal{O}_g := \mathcal{O}_L$.
- (2) The $\alpha = 2\beta$ case: the ideal L contains the elements t_h for $h \in \mathbb{N}$, and $h \geq 1$, $2t_1 - 1$, and $x_1 - 2x_2$. In this case we set $\mathcal{O}_{2\beta} := \mathcal{O}_L$.
- (3) The $\alpha = 4\beta$ case: the ideal L contains the elements t_h for $h \in \mathbb{N}$, and $h \geq 1$, $2t_1 - 1$, and $x_1 - 4x_2$. In this case we set $\mathcal{O}_{4\beta} := \mathcal{O}_L$.

In [3] the case $\alpha = 2\beta$ is considered in details: in particular it is shown that every 2-generated primitive axial algebra in $\mathcal{O}_{2\beta}$ is at most 8 dimensional and this

bound is attained. The following result, which can be compared to Theorem 3.7 in [16], is a consequence of the *resurrection principle* [11, Lemma 1.7].

Proposition 4.9. *Let $\overline{\mathcal{V}} = (\overline{R}, \overline{V}, \overline{A}, (\overline{\mathcal{S}}, \overline{\kappa}))$ be the universal object in the category $\mathcal{O}_g$. Then $\overline{V}$ is linearly spanned by the set $\{a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1}, s_{0,2}, s_{1,2}\}$.*

Proof. Let U be the linear span in $\overline{V}$ of the set

$$B := \{a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1}, s_{0,2}, s_{1,2}\}$$

with coefficients in $\overline{R}$. From Lemmas 4.7 and 4.8, since $\alpha - 2\beta$ and $\alpha - 4\beta$ are invertible in $\overline{R}$, we get that

$$a_0 \cdot s_{1,2}, a_3 \in U.$$

The set B is clearly invariant under the action of τ_0 and since $a_{-2}^f = a_3$, U is also invariant under f . By Equation (11), $a_0 a_1$ and $a_0 a_2$ are contained in U ; by applying alternatively τ_0 and f we get that U contains all the products $a_i a_j$ for $i, j \in \mathbb{Z}$. Similarly, since by Lemma 4.3, for $i \in \{1, 2\}$, $a_0 s_{0,i} \in U$, and by Lemma 4.7, $a_0 s_{1,2} \in U$, we get that U contains all the products $a_j s_{0,i}$ and $a_j s_{1,2}$ for $j \in \mathbb{Z}, i \in \{1, 2\}$.

It follows that, for $i, j \in \{1, 2\}$, the expression on the righthand side of Equation (13) is contained in U , whence $s_{0,i} \cdot s_{0,j}$ is contained in U . As U is invariant under f , we have also $s_{1,2} \cdot s_{1,2} \in U$. Finally, with a similar argument, using the identity

$$a_0 \cdot (u_i \cdot u_{-1} + u_i \cdot v_{-1}) = \alpha(u_i \cdot v_{-1}),$$

we can express $s_{0,i} \cdot s_{1,2}$ as linear combination of elements of B . Hence U is a subalgebra of $\overline{V}$, and since $\overline{V}$ is generated by a_0 and a_1 , we get that $U = \overline{V}$. $\square$

Remark 4.10. *Note that the above proof gives an explicit way to compute the structure constants of the algebra $\overline{V}$ relative to the generating set B . This has been done with the use of GAP [6]. The explicit expressions however are far too long to be written explicitly here.*

Corollary 4.11. *Let $\overline{\mathcal{V}} = (\overline{R}, \overline{V}, \overline{A}, (\overline{\mathcal{S}}, \overline{\kappa}))$ be the universal object in the category $\mathcal{O}_g$. Then, $\overline{R}$ is generated as a $\hat{D}$ -algebra by λ_1 , λ_2 , λ_1^f , and λ_2^f .*

Proof. Since, for every $v \in \overline{V}$, $\lambda_{a_1}(v) = (\lambda_{a_0}(v^f))^f$, λ_{a_0} is a linear function, and $\overline{R} = \overline{R}^f$, by Proposition 4.9, we just need to show that, for every $v \in \{a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1}, s_{0,2}, s_{1,2}\}$, $\lambda_{a_0}(v)$ can be written as a linear combination, with coefficients in $\hat{D}$, of products of λ_1 , λ_2 , λ_1^f , and λ_2^f . By definition we have

$$\lambda_{a_0}(a_0) = 1, \quad \lambda_{a_0}(a_1) = \lambda_1, \quad \text{and} \quad \lambda_{a_0}(a_2) = \lambda_2.$$

Since τ_0 is an $\overline{R}$ -automorphism of $\overline{V}$ fixing a_0 , we get

$$\lambda_{a_0}(a_{-1}) = \lambda_{a_0}((a_1)^{\tau_0}) = \lambda_1,$$

$$\lambda_{a_0}(a_{-2}) = \lambda_{a_0}((a_2)^{\tau_0}) = \lambda_2,$$

and

$$\lambda_{a_0}(s_{0,1}) = \lambda_{a_0}(a_0 a_1 - \beta a_0 - \beta a_1) = \lambda_1 - \beta - \beta \lambda_1,$$

$$\lambda_{a_0}(s_{0,2}) = \lambda_{a_0}(a a_2 - \beta a - \beta a_2) = \lambda_2 - \beta - \beta \lambda_2.$$

Finally, by the fusion law, $u_1u_1 + u_1v_1$ is a sum of a 0 and an α -eigenvector for ad_{a_0} , whence $\lambda_{a_0}(u_1u_1 + u_1v_1) = 0$. By Lemma 4.3, we can compute $u_1u_1 + u_1v_1$ and find

$$u_1u_1 + u_1v_1 = w + \frac{(\alpha - \beta)}{2\alpha} s_{1,2},$$

with $w \in \langle a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1} \rangle$. So, we can express $\lambda_{a_0}(s_{1,2})$ and we obtain

$$\lambda_{a_0}(s_{1,2}) = \frac{2(\alpha - 1)}{\alpha - \beta} \lambda_1^2 - \frac{2(\alpha - 1)}{\alpha - \beta} \lambda_1 \lambda_1^f + (1 - 2\beta) \lambda_1 + \beta \lambda_2 - \beta.$$

□

We conclude this section with a similar result for symmetric algebras over a field in the case $\alpha = 4\beta$. Note that, in this case, since we are assuming $\alpha \neq \beta$, we also assume that the field is of characteristic other than 3.

Proposition 4.12. *Let V be a primitive symmetric axial algebra of Monster type $(4\beta, \beta)$ over a field $\mathbb{F}$ of characteristic greater than 3, generated by two axes $\bar{a}_0$ and $\bar{a}_1$. Then V has dimension at most 8, unless $2\beta - 1 = 0$ and*

$$\lambda_{\bar{a}_0}(\bar{a}_1) = \lambda_{\bar{a}_1}(\bar{a}_0) = \lambda_{\bar{a}_0}(\bar{a}_0^{\tau_{\bar{a}_1}}) = \lambda_{\bar{a}_1}(\bar{a}_1^{\tau_{\bar{a}_0}}) = 1.$$

Proof. Let $\bar{\mathcal{V}} = (\bar{R}, \bar{V}, \bar{\mathcal{A}}, (\bar{\mathcal{S}}, \bar{\tau}))$ be the universal object in the category $\mathcal{O}_{4\beta}$. Since $\alpha - 2\beta = 2\beta$ is invertible in $\bar{R}$, Lemma 4.7 yields that, $a_0 \cdot s_{1,2}$ is contained in $\langle a_{-2}, a_{-1}, a_0, a_1, a_2, s_{0,1}, s_{0,2}, s_{1,2} \rangle$. Since $\alpha = 4\beta$, Equation (1) in Lemma 4.8 becomes

$$\begin{aligned} 0 &= [(48\beta^3)\lambda_1 - (108\beta^4 - 6\beta^3)] a_{-1} \\ &\quad + [(-128\beta^2 + 8\beta)\lambda_1^2 + (-64\beta^2 + 16\beta)\lambda_1\lambda_1^f + (416\beta^3 - 32\beta^2)\lambda_1 \\ &\quad + (16\beta^3 - 16\beta^2)\lambda_1^f + (24\beta^3)\lambda_2 + (-180\beta^4 + 18\beta^3)] a_0 \\ (16) \quad &\quad + [(64\beta^2 - 16\beta)\lambda_1\lambda_1^f + (128\beta^2 - 8\beta)\lambda_1^f{}^2 + (-16\beta^3 + 16\beta^2)\lambda_1 \\ &\quad + (-416\beta^3 + 32\beta^2)\lambda_1^f + (-24\beta^3)\lambda_2^f + (180\beta^4 - 18\beta^3)] a_1 \\ &\quad + [(-48\beta^3)\lambda_1^f + (108\beta^4 - 6\beta^3)] a_2 \\ &\quad + 48\beta^2(2\beta + 1)(\lambda_1 - \lambda_1^f) s_{0,1} \\ &\quad - 36\beta^3(s_{0,2} - s_{1,2}). \end{aligned}$$

By Corollary 3.8, V is a homomorphic image of $\bar{V} \otimes_{\bar{D}} \mathbb{F}$, via a homomorphism ϕ_V mapping a_i to $\bar{a}_i$, for $i \in \{0, 1\}$ and $\mathbb{F}$ is a homomorphic image of $\bar{R} \otimes_{\bar{D}} \mathbb{F}$ via ϕ_R . We use the bar notation to denote the images of the elements of $\bar{V} \otimes_{\bar{D}} \mathbb{F}$ via ϕ_V , while we identify the image under ϕ_R of an element of $\bar{R} \otimes_{\bar{D}} \mathbb{F}$ with the element itself. When we apply ϕ_V to the relation (16) we get a similar relation in V . If the coefficient of a_2 is not zero in $\mathbb{F}$, then we get $\bar{a}_2 \in U_0 := \langle \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{s}_{0,1}, \bar{s}_{0,2}, \bar{s}_{1,2} \rangle$. Since V is symmetric, f induces an automorphism $\bar{f}$ of V and U_0 is $\bar{f}$ invariant. Since U_0 is also $\tau_{\bar{a}_0}$ -invariant, we get also $\bar{a}_{-2} \in U_0$. More generally, by applying alternatively $\tau_{\bar{a}_0}$ and $\bar{f}$ we get that $\bar{a}_i \in U_0$ for every $i \in \mathbb{Z}$. The argument used in the proof of Proposition 4.9 yields $V = U_0$. If the coefficient of $\bar{a}_2$ in the Equation (16) is zero in $\mathbb{F}$, then we may consider the coefficient of $\bar{a}_{-1}$ and if it is not zero we deduce as above that $\bar{a}_{-1} \in \langle \bar{a}_0, \bar{a}_1, \bar{s}_{0,1}, \bar{s}_{0,2}, \bar{s}_{1,2} \rangle$. By proceeding as in the previous case, we

get $V = \langle \bar{a}_0, \bar{a}_1, \bar{s}_{0,1}, \bar{s}_{0,2}, \bar{s}_{1,2} \rangle$. If the coefficients of $\bar{a}_2$ and $\bar{a}_{-1}$ are both zero, then we get

$$\lambda_{\bar{a}_0}(\bar{a}_1) = \lambda_{\bar{a}_1}(\bar{a}_0) = \frac{18\beta - 1}{8}.$$

As above, if the coefficient of $\bar{a}_0$ (or the coefficient of $\bar{a}_1$) is not zero, we can express $\bar{a}_0$ (or $\bar{a}_1$ respectively) as a linear combination of $\bar{a}_{-1}, \bar{s}_{0,1}, \bar{s}_{0,2} - \bar{s}_{1,2}$ ($\bar{a}_0, \bar{s}_{0,1}, \bar{s}_{0,2} - \bar{s}_{1,2}$ respectively). In both cases, it follows that $V = \langle \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{s}_{0,1}, \bar{s}_{0,2}, \bar{s}_{1,2} \rangle$. If also the coefficients of $\bar{a}_0$ and $\bar{a}_1$ are both zero, then we get

$$\lambda_{\bar{a}_0}(\bar{a}_2) = \lambda_{\bar{a}_1}(\bar{a}_{-1}) = \frac{480\beta^3 - 228\beta^2 + 28\beta - 1}{64\beta^2}$$

and Equation (16) becomes

$$0 = 36\beta^2(\bar{s}_{0,2} - \bar{s}_{1,2}).$$

Hence, since $\mathbb{F}$ has characteristic greater than 3, $\bar{s}_{0,2} = \bar{s}_{1,2}$ and the identity $\lambda_{\bar{a}_0}(\bar{s}_{0,2}) = \lambda_{\bar{a}_0}(\bar{s}_{1,2})$ gives that β satisfies the relation

$$(17) \quad (2\beta - 1)^2(12\beta - 1)(14\beta - 1) = 0.$$

From now on assume $\beta \in \{\frac{1}{12}, \frac{1}{14}\} \setminus \{\frac{1}{2}\}$, in particular $\mathbb{F}$ has characteristic other than 5. Set $U_1 := \langle \bar{a}_{-3}, \bar{a}_{-2}, \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{s}_{0,1}, \bar{s}_{0,2}, \bar{s}_{0,3} \rangle$. From the identity $\bar{a}_0(\bar{u}_1\bar{u}_2 - \bar{v}_1\bar{v}_2 + \lambda_{\bar{a}_0}(\bar{v}_1\bar{v}_2)\bar{a}_0) = 0$ we can express $\bar{a}_0(\bar{s}_{1,3} + \bar{s}_{3,2})$ as a linear combination of $\bar{a}_{-3}, \bar{a}_{-2}, \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{s}_{0,1}$, and $\bar{s}_{0,2}$ and then, by Lemma 4.5, we get that $\bar{s}_{0,1}\bar{s}_{0,2} \in U_1$. Then, from the identity $\bar{s}_{0,1}\bar{s}_{0,2} - (\bar{s}_{0,1}\bar{s}_{0,2})^f = 0$, we derive that $\bar{s}_{1,3} \in U_1$, whence also $\bar{s}_{2,3} = (\bar{s}_{1,3})^{\tau_{\bar{a}_0}} \in U_1$. From the identity $\bar{s}_{2,3} - (\bar{s}_{2,3})^f = 0$ we then get $\bar{a}_4 \in U_1$. It follows that U_1 is invariant under f and $\tau_{\bar{a}_0}$, hence $\bar{a}_{\pm i} \in U_1$ for $i \geq 4$. Since U_1 is also $\text{ad}_{\bar{a}_0}$ -invariant, it follows that U_1 contains $\bar{s}_{r,n}$ for every $n \geq 1$, $r \in \{0, \dots, n-1\}$. Thus U_1 is a subalgebra of V , whence $V = U_1$. From the identity $\bar{a}_0(\bar{v}_1\bar{v}_2 - \lambda_{\bar{a}_0}(\bar{v}_1\bar{v}_2)\bar{a}_0) = 0$ we get $\bar{s}_{0,3} \in \langle \bar{a}_{-3}, \bar{a}_{-2}, \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{s}_{0,1}, \bar{s}_{0,2} \rangle$. Finally, from the identity $\bar{a}_5^2 - \bar{a}_5 = 0$ we get $\bar{a}_{-3} \in \langle \bar{a}_{-2}, \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{a}_3, \bar{s}_{0,1}, \bar{s}_{0,2} \rangle$, that is V has dimension at most 8. $\square$

5. THE GENERIC CASE

Let $\bar{\mathcal{V}} = (\bar{R}, \bar{V}, \bar{\mathcal{A}}, (\bar{\mathcal{S}}, \bar{\pi}))$ be the universal object in the category $\mathcal{O}_g$. Note that in this case we have

$$\hat{D} = \mathbb{Z}[1/2, x_1, x_2, x_1^{-1}, x_2^{-1}, (x_1 - x_2)^{-1}, (x_1 - 2x_2)^{-1}, (x_1 - 4x_2)^{-1}].$$

By Corollary 4.11, $\bar{R} = \hat{D}[\lambda_1, \lambda_1^f, \lambda_2, \lambda_2^f]$. The elements $\lambda_1, \lambda_1^f, \lambda_2, \lambda_2^f$ are not necessarily indeterminates on $\hat{D}$, as they have to satisfy various relations imposed by the definition of $\bar{R}$. In particular, since by Lemma 4.6, $s_{2,3} - s_{2,3}^f = 0$, in the ring $\bar{R}$ the following relations hold

- (1) $\lambda_{a_0}(s_{2,3} - (s_{2,3})^f) = 0$,
- (2) $\lambda_{a_0}((s_{2,3} - (s_{2,3})^f)^{\tau_1}) = 0$,
- (3) $\lambda_{a_0}(a_3a_3 - a_3) = 0$,
- (4) $\lambda_{a_1}(s_{2,3} - (s_{2,3})^f) = 0$.

By Remark 4.10, the four expressions on the left hand side of the above identities can be computed explicitly and produce respectively four polynomials $p_i(x, y, z, t)$ for $i \in \{1, \dots, 4\}$ in $\hat{D}[x, y, z, t]$ (with x, y, z, t indeterminates on $\hat{D}$), that simultaneously annihilate on the quadruple $(\lambda_1, \lambda_1^f, \lambda_2, \lambda_2^f)$. Define also, for $i \in \{1, 2\}$,

$q_i(x, z) := p_i(x, x, z, z)$. The polynomials p_i 's and q_i 's are too long to be displayed here but can be computed using [1] or [6].

Suppose V is a primitive axial algebra of Monster type (α, β) over a field $\mathbb{F}$ of odd characteristic, with $\alpha, \beta \in \mathbb{F}$ and $\alpha \notin \{2\beta, 4\beta\}$, generated by two axes $\bar{a}_0$ and $\bar{a}_1$. Then, by Corollary 3.8, V is a homomorphic image of $\bar{V} \otimes_{\bar{D}} \mathbb{F}$ and $\mathbb{F}$ is a homomorphic image of $\bar{R} \otimes_{\bar{D}} \mathbb{F}$. We denote the images of an element δ of $\bar{R} \otimes_{\bar{D}} \mathbb{F}$ in $\mathbb{F}$ by $\bar{\delta}$ and by $\bar{p}_i$ and $\bar{q}_i$ the polynomials in $\mathbb{F}[x, y, z, t]$ and $\mathbb{F}[x, z]$ corresponding to p_i and q_i , respectively. Set

$$T := \{\bar{p}_1, \bar{p}_2, \bar{p}_3, \bar{p}_4\}$$

and

$$T_s := \{\bar{p}_1(x, z), \bar{p}_2(x, z)\}.$$

Moreover, for $P \in \{T, T_s\}$, denote by $Z(P)$ the set of common zeroes of all the elements of P in $\mathbb{F}^4$ and $\mathbb{F}^2$ respectively. It is clear from the definition that the $\bar{p}_i$'s have the coefficients in the field $\mathbb{F}_0(\alpha, \beta)$. By Proposition 4.9 and Corollary 4.11, the algebra V is completely determined, up to homomorphic images, by the quadruple

$$(\lambda_{\bar{a}_0}(\bar{a}_1), \lambda_{\bar{a}_1}(\bar{a}_0), \lambda_{\bar{a}_0}(\bar{a}_2), \lambda_{\bar{a}_1}(\bar{a}_{-1})).$$

Furthermore, this quadruple is the homomorphic image in $\mathbb{F}^4$ of the quadruple $(\lambda_1, \lambda_1^f, \lambda_2, \lambda_2^f)$ and so it is a common zero of the elements of T .

If, in addition, the algebra V is symmetric, then

$$\lambda_{\bar{a}_0}(\bar{a}_1) = \lambda_{\bar{a}_1}(\bar{a}_0) \text{ and } \lambda_{\bar{a}_0}(\bar{a}_2) = \lambda_{\bar{a}_1}(\bar{a}_{-1})$$

and the pair $(\lambda_{\bar{a}_0}(\bar{a}_1), \lambda_{\bar{a}_0}(\bar{a}_2))$ is a common zero of the elements of the set T_s .

We thus have proved Theorem 1.3.

Computing the resultant of the polynomials $\bar{p}_1(x, z)$ and $\bar{p}_2(x, z)$ with respect to z one obtains a polynomial in x of degree 10, which is the product of the five linear factors

$$x, \quad x - 1, \quad 2x - \alpha, \quad 2x - \beta, \quad 4(2\alpha - 1)x - (3\alpha^2 + 3\alpha\beta - \alpha - 2\beta)$$

and a factor of degree at most 5. This last factor has degree 5 and is irreducible in $\mathbb{Q}(\alpha, \beta)$, if α and β are indeterminates over $\mathbb{Q}$. On the other hand, for certain values of α and β , this factor can be reducible: for example, it even completely splits in $\mathbb{Q}(\alpha, \beta)[x]$ when $\alpha = 2\beta$ (see [3]), or in the Norton-Sakuma case, when $(\alpha, \beta) = (1/4, 1/32)$ (see the proof of Theorem 1.6 below).

Fixed a field $\mathbb{F}$, in order to classify primitive generic axial algebras of Monster type (α, β) over $\mathbb{F}$ generated by two axes $\bar{a}_0$ and $\bar{a}_1$ we can proceed as follows. We first find all the zeroes of the set T_s and classify all symmetric algebras. Then we observe that, the even subalgebra $\langle\langle \bar{a}_0, \bar{a}_2 \rangle\rangle$ and the odd subalgebra $\langle\langle \bar{a}_{-1}, \bar{a}_1 \rangle\rangle$ are symmetric, since the automorphisms $\tau_{\bar{a}_0}$ and $\tau_{\bar{a}_1}$ respectively, swap the generating axes. Hence, from the classification of the symmetric case, we know all possible values for the pairs

$$(\lambda_{\bar{a}_0}(\bar{a}_2), \lambda_{\bar{a}_1}(\bar{a}_{-1}))$$

and we can look for common zeros (x_0, y_0, z_0, t_0) of the set T with those prescribed values for (x_0, z_0) .

Using this method, we now classify 2-generated primitive axial algebras of Monster type (α, β) over the field $\mathbb{Q}(\alpha, \beta)$, with α and β independent indeterminates over $\mathbb{Q}$.

Lemma 5.1. *If $\mathbb{F} = \mathbb{Q}(\alpha, \beta)$, with α and β independent indeterminates over $\mathbb{Q}$, the set $Z(T_s)$ consists exactly of the 5 points*

$$(1, 1), (0, 1), \left(\frac{\beta}{2}, \frac{\beta}{2}\right), \left(\frac{\alpha}{2}, 1\right),$$

and

$$(q(\alpha, \beta), q(\alpha, \beta)), \text{ with } q(\alpha, \beta) = \frac{(3\alpha^2 + 3\alpha\beta - \alpha - 2\beta)}{4(2\alpha - 1)}.$$

Proof. The system can be solved in $\mathbb{Q}(\alpha, \beta)$ using [1] giving the five solutions of the statement. $\square$

Lemma 5.2. *Let $\mathbb{F} = \mathbb{Q}(\alpha, \beta)$, with α and β independent indeterminates over $\mathbb{Q}$ and let $(x_0, z_0) \in Z(T_s)$. Then $(x_0, y_0, z_0, t_0) \in Z(T)$ if and only if*

$$(y_0, t_0) = (x_0, z_0).$$

Proof. This have been checked using [1]. $\square$

Lemma 5.3. *The algebras $3C(\alpha)$, $3C(\beta)$, and $3A(\alpha, \beta)$ over the field $\mathbb{Q}(\alpha, \beta)$ are simple.*

Proof. The claim follows from [9, Theorem 4.11 and Corollary 4.6]. For algebras $3C(\alpha)$ and $3C(\beta)$ it is proved in [8, Example 3.4]. The algebra $3A(\alpha, \beta)$ is the same as the algebra $3A'_{\alpha, \beta}$ defined by Reheren in [16]. By [16, Lemma 8.2], it admits a Frobenius form wich is non degenerate over the field $\mathbb{Q}(\alpha, \beta)$ and such that all the generating axes are non-singular with respect to this form. Hence, by Theorem 4.11 in [9], every non trivial ideal contains at least one of the generating axes. Then, Corollary 4.6 in [9] yields that the algebra is simple. $\square$

Proof of Theorem 1.4. It is straightforward to check that the algebras $1A$, $2B$, $3C(\alpha)$, $3C(\beta)$, and $3A(\alpha, \beta)$ are 2-generated symmetric axial algebras of Monster type (α, β) over the field $\mathbb{Q}(\alpha, \beta)$ and their corresponding values of $(\lambda_{\bar{a}_0}(\bar{a}_1), \lambda_{\bar{a}_0}(\bar{a}_2))$ are respectively

$$(1, 1), (0, 1), \left(\frac{\beta}{2}, \frac{\beta}{2}\right), \left(\frac{\alpha}{2}, 1\right), \text{ and } (q(\alpha, \beta), q(\alpha, \beta)),$$

where $q(\alpha, \beta)$ is defined in Lemma 5.1. Let V be an axial algebra of Monster type (α, β) over the field $\mathbb{Q}(\alpha, \beta)$ generated by the two axes $\bar{a}_0$ and $\bar{a}_1$. Set

$$\bar{\lambda}_1 := \lambda_{\bar{a}_0}(\bar{a}_1), \bar{\lambda}'_1 := \lambda_{\bar{a}_1}(\bar{a}_0), \bar{\lambda}_2 := \lambda_{\bar{a}_0}(\bar{a}_2), \text{ and } \bar{\lambda}'_2 := \lambda_{\bar{a}_1}(\bar{a}_{-1}).$$

By Theorem 1.3, V is determined, up to homomorphic images, by the quadruple $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2)$, which must be in $Z(T)$. By Lemma 5.2, we get the five quadruples

$$(1, 1, 1, 1), (0, 1, 0, 1), \left(\frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2}\right), \left(\frac{\alpha}{2}, 1, \frac{\alpha}{2}, 1\right),$$

and

$$(q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta)).$$

By Corollary 3.8 and Proposition 4.9, V is linearly generated on $\mathbb{Q}(\alpha, \beta)$ by the set $\bar{a}_{-2}, \bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{s}_{0,1}, \bar{s}_{0,2}, \text{ and } \bar{s}_{1,2}$. Define

$$\bar{d}_0 := \bar{s}_{2,3} - \bar{s}_{1,3}^{\tau_0}, \bar{d}_1 := \bar{d}_0^f, \bar{d}_2 := \bar{d}_0^{\tau_1},$$

and, for $i \in \{0, 1, 2\}$,

$$\bar{D}_i := \bar{d}_i^{\tau_0} - \bar{d}_i.$$

By Lemma 4.6, all vectors $\bar{d}_i, \bar{D}_i$ for $i \in \{0, 1, 2\}$ are zero. For all the admissible values of $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2)$, the coefficient of $\bar{a}_{-2}$ in D_0 is non zero, hence we can express $\bar{a}_{-2}$ as a linear combination of $\bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{s}_{0,1}, \bar{s}_{0,2}$, and $\bar{s}_{1,2}$. Similarly, from identity $\bar{d}_0 = 0$ we can express $\bar{s}_{1,2}$ as a linear combination of $\bar{a}_{-1}, \bar{a}_0, \bar{a}_1, \bar{a}_2, \bar{s}_{0,1}$, and $\bar{s}_{0,2}$.

For $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2)$ in

$$\left\{ \left(\frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2} \right), (q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta)) \right\},$$

from identity $\bar{d}_2 = 0$ we get $\bar{a}_{-1} = \bar{a}_2$ and consequently $\bar{s}_{0,2} = \bar{s}_{0,1}$. Thus in this two cases the dimension of V is at most 4. Moreover, if $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2) = (\frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2})$, then from the identity $\bar{s}_{0,2}\bar{s}_{0,1} - \bar{s}_{0,1}\bar{s}_{0,1} = 0$ we get $\bar{s}_{0,1} = -\frac{\beta}{2}(\bar{a}_0 + \bar{a}_1 + \bar{a}_2)$ and hence the dimension is at most 3.

For $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2)$ in

$$\left\{ (1, 1, 1, 1), (0, 1, 0, 1), \left(\frac{\alpha}{2}, 1, \frac{\alpha}{2}, 1 \right) \right\},$$

from the identity $\bar{D}_2 = 0$ we get $\bar{a}_{-1} = \bar{a}_1$. Then, from the identity $\bar{d}_2 = 0$ we deduce $\bar{a}_2 = \bar{a}_0$ and hence $\bar{s}_{0,2} = (1 - 2\beta)\bar{a}_0$. Hence in this cases V has dimension at most 3. Suppose $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2) = (0, 1, 0, 1)$. Then, from the identity $\bar{s}_{0,1}\bar{s}_{0,2} + (2\beta - 1)\bar{a}_0$ we get $\bar{s}_{0,1} = -\beta(\bar{a}_0 + \bar{a}_1)$ and so $\bar{a}_0\bar{a}_1 = 0$. Hence in this case V is isomorphic to the algebra $2B$. Finally, suppose $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2) = (1, 1, 1, 1)$. From the identity $\bar{s}_{0,2} - \bar{s}_{1,2}^f = 0$ we get $\bar{a}_0 = \bar{a}_1$, that is V is the algebra $1A$.

Thus, for each $(\bar{\lambda}_1, \bar{\lambda}'_1, \bar{\lambda}_2, \bar{\lambda}'_2)$ in the set

$$\left\{ \left(\frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2}, \frac{\beta}{2} \right), (q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta), q(\alpha, \beta)), \left(\frac{\alpha}{2}, 1, \frac{\alpha}{2}, 1 \right) \right\},$$

we get that V satisfies the same multiplication table as the algebras $3C(\beta)$, $3C(\alpha)$ and $3A(\alpha, \beta)$ respectively and has at most the same dimension. Therefore, to conclude the proof, we need only to show that the algebras $3C(\alpha)$, $3C(\beta)$ and $3A(\alpha, \beta)$ are simple. This follows from Lemma 5.3. $\square$

As a corollary of Theorem 1.3, we can prove now Theorem 1.6.

Proof of Theorem 1.6. Let $\mathbb{F}$ be a field of characteristic zero. Then $\mathbb{F}$ contains $\mathbb{Q}$. The resultant with respect to z of the polynomials in T_s has degree 9 and splits in $\mathbb{Q}[x]$ as the product of a constant and the linear factors

$$x, x - 1, x - \frac{1}{8}, \left(x - \frac{1}{64} \right)^2, x - \frac{13}{2^8}, x - \frac{1}{32}, x - \frac{3}{2^7}, x - \frac{5}{2^8}.$$

In $\mathbb{Q}^4$, the set $Z(T)$ consists of the 9 points

$$\begin{aligned} & (1, 1, 1, 1), (0, 0, 1, 1), \left(\frac{1}{8}, \frac{1}{8}, 1, 1 \right), \left(\frac{1}{64}, \frac{1}{64}, \frac{1}{64}, \frac{1}{64} \right), \left(\frac{13}{2^8}, \frac{13}{2^8}, \frac{13}{2^8}, \frac{13}{2^8} \right), \\ & \left(\frac{1}{32}, \frac{1}{32}, 0, 0 \right), \left(\frac{1}{64}, \frac{1}{64}, \frac{1}{8}, \frac{1}{8} \right), \left(\frac{3}{2^7}, \frac{3}{2^7}, \frac{3}{2^7}, \frac{3}{2^7} \right), \left(\frac{5}{2^8}, \frac{5}{2^8}, \frac{13}{2^8}, \frac{13}{2^8} \right). \end{aligned}$$

By [11, 7], each quadruple of the above list corresponds to a Norton-Sakuma algebra. By Corollary 4.13 in [9], every Norton-Sakuma algebra is simple, provided it is not of type $2B$. Hence, the thesis follows from Theorem 1.3, once we prove that in each case the dimension of V is at most equal to the dimension of the corresponding Norton-Sakuma algebra. This can be done, by Remark 4.10, using [6]. $\square$

REFERENCES

- [1] Decker, W.; Greuel, G.-M.; Pfister, G.; Schönemann, H.: SINGULAR 4-1-1 — A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2018).
- [2] De Medts, T., Peacock S.F., Shpectorov, S. and van Couwenberghe M.: Decomposition algebras and axial algebras, arXiv:1905.03481v1
- [3] Franchi, C., Mainardis M., Shpectorov, S.: 2-generated axial algebras of Monster type $(2\beta, \beta)$, in preparation.
- [4] Franchi, C., Mainardis M., Shpectorov, S.: An infinite dimensional 2-generated axial algebras of Monster type, arXiv:2007.02430
- [5] Galt, A., Joshi, V., Mamontov, A., Shpectorov, S., Staroletov, A.: Double axes and subalgebras of Monster type in Matsuo algebras, arXiv:2004.11180
- [6] GAP – Groups, Algorithms, and Programming, Version 4.8.6, The GAP group, 2016.
- [7] Hall, J., Rehren, F., Shpectorov, S.: Universal Axial Algebras and a Theorem of Sakuma, *J. Algebra* **421** (2015), 394-424.
- [8] Hall, J., Rehren, F., Shpectorov, S.: Primitive axial algebras of Jordan type, *J. Algebra* **437** (2015), 79-115.
- [9] Khasraw, S.M.S., McInroy, J., Shpectorov, S.: On the structure of axial algebras, *Trans. Amer. Math. Soc.*, **373** (2020), 2135-2156
- [10] Ivanov, A. A.: The Monster group and Majorana involutions. Cambridge Tracts in Mathematics 176, Cambridge Univ. Press, Cambridge (2009)
- [11] Ivanov, A. A., Pasechnik, D. V., Seress, Á., Shpectorov, S.: Majorana representations of the symmetric group of degree 4, *J. Algebra* **324** (2010), 2432-2463
- [12] Miyamoto, M.: Griess algebras and conformal vectors in vertex operator algebras, *J. Algebra* **179** (1996), 523-548
- [13] Norton, S. P.: The uniqueness of the Fischer-Griess Monster. In: McKay, J. (ed.) Finite groups-coming of age (Montreal, Que.,1982). Contemp. Math. 45, pp. 271–285. AMS, Providence, RI (1985)
- [14] Norton, S. P.: The Monster algebra: some new formulae. In Moonshine, the Monster and related topics (South Hadley, Ma., 1994), Contemp. Math. 193, pp. 297-306. AMS, Providence, RI (1996)
- [15] Rehren, F.: Axial algebras, PhD thesis, University of Birmingham, 2015.
- [16] Rehren, F.: Generalized dihedral subalgebras from the Monster, *Trans. Amer. Math. Soc.* **369** (2017), 6953-6986.
- [17] Sakuma, S.: 6-transposition property of τ -involutions of vertex operator algebras. Int. Math. Res. Not. (2007). doi:10.1093/imrn/rmn030
- [18] Yabe, T.: On the classification of 2-generated axial algebras of Majorana type, arXiv:2008.01871
- [19] Whybrow, M.: Graded 2-generated axial algebras, arXiv:2005.03577