

T-Quadratic Forms and Spectral Analysis of T-Symmetric Tensors

Liqun Qi* and Xinzhen Zhang†

November 13, 2021

Abstract

An $n \times n \times p$ tensor is called a T-square tensor. It arises from many applications, such as the image feature extraction problem and the multi-view clustering problem. We may symmetrize a T-square tensor to a T-symmetric tensor. For each T-square tensor, we define a T-quadratic form, whose variable is an $n \times p$ matrix, and whose value is a p -dimensional vector. We define eigentuples and eigenmatrices for T-square tensors. We show that a T-symmetric tensor has unique largest and smallest eigentuples, and a T-quadratic form is positive semi-definite (definite) if and only if its smallest eigentuple is nonnegative (positive). The relation between the eigen-decomposition of T-symmetric tensors, and the TSVD of general third order tensors are also studied.

Key words. T-square tensors, T-symmetric tensors, T-quadratic forms, eigentuples.

AMS subject classifications. 15A69, 15A18

1 Introduction

We call a third order tensor $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ a T-square tensor. It was called an f-square tensor in [6]. The representation tensor $\mathcal{Z} \in \mathbb{R}^{n \times n \times p}$ arising in the multi-view clustering problem [2] and the multi-view image feature extraction problem [9, 10] is a T-square tensor. Here n is the number of the samples in the database, p is the number of the views.

*Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong, China; (liqun.qi@polyu.edu.hk).

†School of Mathematics, Tianjin University, Tianjin 300354 China; (xzzhang@tju.edu.cn). This author's work was supported by NSFC (Grant No. 11871369).

Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is a T-square tensor. Let $X \in \mathbb{R}^{n \times p}$. We may regard X as a tensor $\mathcal{X} \in \mathbb{R}^{n \times 1 \times p}$. Define

$$F_{\mathcal{A}}(X) := \mathcal{X}^{\top} * \mathcal{A} * \mathcal{X}, \quad (1.1)$$

where $*$ is the T-product operation introduced in [1, 3, 4], and \top is the transpose operation in the T-product sense. In the next section, we will review the definition of T-product and its transpose concept. We call $F_{\mathcal{A}}$ the T-quadratic form defined by \mathcal{A} . Then for any $X \in \mathbb{R}^{n \times p}$, $F_{\mathcal{A}}(X) \in \mathbb{R}^p$. If $F_{\mathcal{A}}(X) \geq \mathbf{0}$ for any $X \in \mathbb{R}^{n \times p}$, then we say that the T-quadratic form $F_{\mathcal{A}}$ is T-positive semi-definite. If $F_{\mathcal{A}}(X) > \mathbf{0}$ for any $X \in \mathbb{R}^{n \times p}$, then we say that the T-quadratic form $F_{\mathcal{A}}$ is T-positive definite.

The T-positive semidefiniteness (definiteness) concept here is different from the T-positive semidefiniteness (definiteness) concept discussed in [12]. The T-positive semidefiniteness (definiteness) concept in [12] is in the sense of nonnegative (positive) scalars. Here, the concept is in the sense of nonnegative (positive) vectors. Thus, the T-positive semidefiniteness (definiteness) concept here is stronger and may reflect more correlative properties of T-square tensors.

The T-product operation, TSVD decomposition and tubal ranks were introduced by Kilmer and her collaborators in [1, 3, 4]. It is now widely used in engineering [2, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16]. In [1], Bradman defined real eigentuples and eigenmatrices for third order tensors in $\mathbb{R}^{n \times n \times n}$. Viewing the wide applications of T-product, TSVD decomposition and tubal ranks, the theory of eigentuples and eigenmatrices deserves to be further studied.

In this paper, we extend the concepts of eigentuples and eigenmatrices to T-square tensors and allow complex eigentuples and eigenmatrices. We show that an $n \times n \times p$ T-symmetric tensor has unique largest eigentuple $\mathbf{s}_1 \in \mathbb{R}^p$ and unique smallest eigentuple $\mathbf{s}_n \in \mathbb{R}^p$ such that any real eigentuple \mathbf{s} of \mathcal{A} satisfies $\mathbf{s}_1 \geq \mathbf{s} \geq \mathbf{s}_n$. We further show that a T-quadratic form is positive semidefinite (definite) if and only if the smallest eigentuple of the corresponding T-symmetric tensor is nonnegative (positive).

The T-quadratic function $F_{\mathcal{A}}$ maps $\mathbb{R}^{n \times p}$ to \mathbb{R}^p . Its positive semidefiniteness (definiteness) requires p quadratic polynomials of np variables to be nonnegative (positive) simultaneously. We present its spectral conditions. This theory is novel.

We then further study the relation between the eigen-decomposition of T-symmetric tensors, and the TSVD of general third order tensors.

The rest of this paper is distributed as follows. We deliver some preliminary knowledge of T-product operations in the next section. In Section 3, we define eigentuples and eigenmatrices for a T-square tensor, and show the existence of the largest and the smallest eigentuples of a T-symmetric tensor. In Section 4, we prove that a T-symmetric tensor is positive semidefinite (definite) if and only if its smallest eigentuple is nonnegative (positive). We study the relation between the eigen-decomposition of

T-symmetric tensors, and the TSVD of general third order tensors in Section 5.

2 Preliminaries

Let $\mathbf{a} = (a_1, a_2, \dots, a_p)^\top \in \mathbb{C}^p$. Then

$$\text{circ}(\mathbf{a}) := \begin{pmatrix} a_1 & a_p & a_{p-1} & \cdots & a_2 \\ a_2 & a_1 & a_p & \cdots & a_3 \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ \cdot & \cdot & \cdot & \cdots & \cdot \\ a_p a_{p-1} & a_{p-2} & \cdots & a_1 \end{pmatrix},$$

and $\text{circ}^{-1}(\text{circ}(\mathbf{a})) := \mathbf{a}$.

Suppose that $\mathbf{a}, \mathbf{b} \in \mathbb{C}^p$. Define

$$\mathbf{a} \odot \mathbf{b} = \text{circ}(\mathbf{a})\mathbf{b}.$$

In [3], $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p$ are called tubal scalars. Here, we extend them to \mathbb{C}^p .

In general, $\mathbf{a} \odot \mathbf{b} \neq \mathbf{b} \odot \mathbf{a}$. We denote

$$\mathbf{a}^{\odot 2} := \mathbf{a} \odot \mathbf{a}.$$

If $\mathbf{a} \in \mathbb{R}^p$ is nonnegative, then $\mathbf{a}^{\odot 2}$ is also nonnegative. However, if $\mathbf{b} \in \mathbb{R}^p$ is nonnegative, there may be no $\mathbf{a} \in \mathbb{R}^p$ such that $\mathbf{a}^{\odot 2} = \mathbf{b}$. For example, let $p = 2$, $\mathbf{a} = (a_1, a_2)^\top$, $\mathbf{b} = (b_1, b_2)^\top$ and $\mathbf{a}^{\odot 2} = \mathbf{b}$. Then we have $b_1 = a_1^2 + a_2^2$ and $b_2 = 2a_1a_2$. To satisfy these two equations, we must have $b_1 \geq b_2$. We say that $\mathbf{b} \in \mathbb{R}^p$ is a square tubal scalar if it is nonnegative and there is an $\mathbf{a} \in \mathbb{R}^p$, such that \mathbf{a} is nonnegative and $\mathbf{a}^{\odot 2} = \mathbf{b}$.

For $\mathbf{a} = (a_1, \dots, a_p)^\top \in \mathbb{R}^p$, denote $|\mathbf{a}| := (|a_1|, \dots, |a_p|)^\top$.

Question 1 Suppose that $\mathbf{b} \in \mathbb{R}^p$ is a square tubal scalar. Is there a unique $\mathbf{a} \in \mathbb{R}^p$, such that \mathbf{a} is nonnegative and $\mathbf{a}^{\odot 2} = \mathbf{b}$?

Proposition 2.1 $(\mathbb{C}^p, +, \odot)$ is a commutative ring with unity $\mathbf{e} = (1, 0, \dots, 0)^\top \in \mathbb{C}^p$, where $+$ is the vector addition.

Proposition 2.1 extends Theorem 3.2 of [1] from \mathbb{R}^p to \mathbb{C}^p , as we need to consider complex eigentuples for third order real tensors. The proof is almost the same. Hence, we omit the proof.

Note that the operation \odot is different from vector convolution. For $\mathbf{a}, \mathbf{b} \in \mathbb{C}^p$, the vector convolution of \mathbf{a} and \mathbf{b} is in \mathbb{C}^{2p-1} .

For $X \in \mathbb{C}^{n \times p}$ and $\mathbf{a} \in \mathbb{C}^p$, define

$$\mathbf{a} \circ X = X \text{circ}(\mathbf{a}).$$

Proposition 2.2 *Let $\mathbf{a}, \mathbf{b} \in \mathbb{C}^p$, and $X, Y \in \mathbb{C}^{n \times p}$. Then*

1. $\mathbf{a} \circ (X + Y) = \mathbf{a} \circ X + \mathbf{a} \circ Y$;
2. $(\mathbf{a} + \mathbf{b}) \circ X = \mathbf{a} \circ X + \mathbf{b} \circ X$;
3. $\mathbf{a} \circ (\mathbf{b} \circ X) = (\mathbf{a} \odot \mathbf{b}) \circ X$;
4. *Let $\mathbf{e} = (1, 0, \dots, 0)^\top \in \mathbb{C}^p$ as in Proposition 2.1. Then $\mathbf{e} \circ X = X$ for all $X \in \mathbb{C}^{n \times p}$. Furthermore, \mathbf{e} is the unique element in \mathbb{C}^p with this property.*

Proof This proposition extends Theorem 3.5 of [1] from $\mathbb{C}^{p \times p}$ to $\mathbb{C}^{n \times p}$, except the second half of item 4 is additional. The proof of the other part except the second half of item 4 is almost the same as the proof of Theorem 3.5 of [1]. We omit this part and now prove the second half of item 4. Suppose $\mathbf{a} \circ X = X$ for all $X \in \mathbb{C}^{n \times p}$. Then $X \text{circ}(\mathbf{a}) = X$ for all $X \in \mathbb{C}^{n \times p}$. This implies $\text{circ}(\mathbf{a}) = I_p$, the identity matrix of $\mathbb{R}^{p \times p}$. Thus, $\mathbf{a} = \text{circ}^{-1}(I_p) = \mathbf{e}$. \square

For a third order tensor $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$, its frontal slices are denoted as $A^{(1)}, \dots, A^{(p)} \in \mathbb{R}^{m \times n}$. As in [1, 3, 4], define

$$\text{bcirc}(\mathcal{A}) := \begin{pmatrix} A^{(1)} & A^{(p)} & & A^{(p-1)} \dots & A^{(2)} \\ A^{(2)} & A^{(1)} & & A^{(p)} \dots & A^{(3)} \\ \cdot & \cdot & & \cdot \dots & \cdot \\ \cdot & \cdot & & \cdot \dots & \cdot \\ A^{(p)} & A^{(p-1)} & & A^{(p-2)} \dots & A^{(1)} \end{pmatrix},$$

and $\text{bcirc}^{-1}(\text{bcirc}(\mathcal{A})) := \mathcal{A}$.

Various T-product structured properties of third order tensors are based upon their block circulant matrix versions. For a third order tensor $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$, its transpose can be defined as

$$\mathcal{A}^\top = \text{bcirc}^{-1}[(\text{birc}(\mathcal{A}))^\top].$$

This will be the same as the definition in [1, 3, 4]. The identity tensor \mathcal{I}_{nnp} may also be defined as

$$\mathcal{I}_{nnp} = \text{bcirc}^{-1}(I_{np}),$$

where I_{np} is the identity matrix in $\mathbb{R}^{np \times np}$.

However, a third order tensor \mathcal{S} in $\mathbb{R}^{m \times n \times p}$ is f-diagonal in the sense of [1, 3, 4] if all of its frontal slices $S^{(1)}, \dots, S^{(p)}$ are diagonal. In this case, $\text{bcirc}(\mathcal{S})$ may not be diagonal.

For a third order tensor $\mathcal{A} \in \mathfrak{R}^{m \times n \times p}$, it is defined [1, 4] that

$$\text{unfold}(\mathcal{A}) := \begin{pmatrix} A^{(1)} \\ A^{(2)} \\ \cdot \\ \cdot \\ A^{(p)} \end{pmatrix} \in \mathfrak{R}^{mp \times n},$$

and $\text{fold}(\text{unfold}(\mathcal{A})) := \mathcal{A}$. For $\mathcal{A} \in \mathfrak{R}^{m \times s \times p}$ and $\mathcal{B} \in \mathfrak{R}^{s \times n \times p}$, the T-product of \mathcal{A} and \mathcal{B} is defined as $\mathcal{A} * \mathcal{B} := \text{fold}(\text{bcirc}(\mathcal{A})\text{unfold}(\mathcal{B})) \in \mathfrak{R}^{m \times n \times p}$. Then, we see that

$$\mathcal{A} * \mathcal{B} = \text{bcirc}^{-1}(\text{bcirc}(\mathcal{A})\text{bcirc}(\mathcal{B})).$$

Thus, the bcirc and bcirc^{-1} operations not only form a one-to-one relationship between third order tensors and block circulant matrices, but their product operation is reserved.

The Standard Form of a Real f-Diagonal Tensor. Let $\mathcal{S} = (s_{ijk}) \in \mathfrak{R}^{m \times n \times p}$ be a f-diagonal tensor. Let $\mathbf{s}_j = (s_{jj1}, s_{jj2}, \dots, s_{jjp})^\top$ be the jj th tube of \mathcal{S} for $j = 1, \dots, \min\{m, n\}$. We say that \mathcal{S} is in its standard form if $\mathbf{s}_1 \geq \mathbf{s}_2 \geq \dots \geq \mathbf{s}_{\min\{m, n\}}$.

3 Eigentuples and Eigenmatrices of T-Square Tensors

For a matrix $X \in \mathbb{C}^{n \times p}$, let its column vectors be $\mathbf{x}^{(1)}, \dots, \mathbf{x}^{(p)}$. Define

$$\text{unfold}(X) := \begin{pmatrix} \mathbf{x}^{(1)} \\ \mathbf{x}^{(2)} \\ \cdot \\ \cdot \\ \mathbf{x}^{(p)} \end{pmatrix} \in \mathbb{C}^{np},$$

and $\text{fold}(\text{unfold}(X)) := X$. Then we define the T-product of \mathcal{A} and X as

$$\mathcal{A} * X = \text{fold}(\text{bcirc}(\mathcal{A})\text{unfold}(X)).$$

Thus, $\mathcal{A} * X \in \mathbb{C}^{m \times p}$.

We now define eigentuples and eigenmatrices of T-square tensors. Suppose that $\mathcal{A} \in \mathfrak{R}^{n \times n \times p}$ is a T-square tensor, $X \in \mathbb{C}^{n \times p}$ and $X \neq O$, $\mathbf{d} \in \mathbb{C}^p$, and

$$\mathcal{A} * X = \mathbf{d} \circ X. \tag{3.2}$$

Then we call \mathbf{d} an eigentuple of \mathcal{A} , and X an eigenmatrix of \mathcal{A} , corresponding to the eigentuple \mathbf{d} .

The eigentuple and eigenmatrix concepts extend the eigentuple and eigenmatrix concepts of [1] from $\mathbb{R}^{p \times p \times p}$ to $\mathbb{R}^{n \times n \times p}$ and allow complex eigentuples and eigenmatrices.

We aim to study T-positive semi-definiteness and T-positive definiteness of the T-quadratic form $F_{\mathcal{A}}$, defined in (1.1). This would not be easy by using the eigentuples of \mathcal{A} , as even for real square matrices, their eigenvalues may not be real. Thus, as in the matrix case, we symmetrize the T-square tensor \mathcal{A} .

Let $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ be a T-square tensor. We say that \mathcal{A} is T-symmetric if $\mathcal{A} = \mathcal{A}^\top$. T-symmetric tensors have been studied in [12]. We have the following propositions.

Proposition 3.1 *Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$. Then $\mathcal{A} + \mathcal{A}^\top$ is a T-symmetric tensor. Then \mathcal{A} is positive semidefinite (definite) if and only if the T-symmetric tensor $\mathcal{A} + \mathcal{A}^\top$ is positive semidefinite (definite).*

Proof Since

$$(\mathcal{A} + \mathcal{A}^\top)^\top = \mathcal{A}^\top + (\mathcal{A}^\top)^\top = \mathcal{A} + \mathcal{A}^\top,$$

$\mathcal{A} + \mathcal{A}^\top$ is T-symmetric.

For $X \in \mathbb{R}^{n \times p}$, regard it as a tensor $\mathcal{X} \in \mathbb{R}^{n \times 1 \times p}$. We have

$$F_{\mathcal{A}}(X) = \mathcal{X}^\top * \mathcal{A} * \mathcal{X} = (\mathcal{X}^\top * \mathcal{A} * \mathcal{X})^\top = \mathcal{X}^\top * \mathcal{A}^\top * \mathcal{X} = \frac{1}{2} \mathcal{X}^\top * (\mathcal{A} + \mathcal{A}^\top) * \mathcal{X}.$$

Thus, \mathcal{A} is positive semidefinite (definite) if and only if the T-symmetric tensor $\mathcal{A} + \mathcal{A}^\top$ is positive semidefinite (definite). \square

We thus study the eigentuples of T-symmetric tensors, and use them to analyze positive semidefiniteness (definiteness) of these tensors.

The following proposition holds obviously.

Proposition 3.2 *A T-square tensor $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is T-symmetric if and only if $\text{bcirc}(\mathcal{A})$ is symmetric. A T-square tensor $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is invertible if and only if $\text{bcirc}(\mathcal{A})$ is invertible. In this case, we have*

$$\mathcal{A}^{-1} = \text{bcirc}^{-1}(\text{bcirc}(\mathcal{A}^{-1})).$$

Furthermore, \mathcal{A} is orthogonal in the sense of [1, 3, 4] if and only if $\text{bcirc}(\mathcal{A})$ is orthogonal.

We have the following theorem.

Theorem 3.3 Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is a T -symmetric tensor. Then there are orthogonal tensor $\mathcal{U} \in \mathbb{R}^{n \times n \times p}$ and T -symmetric f -diagonal tensor $\mathcal{D} \in \mathbb{R}^{n \times n \times p}$ such that

$$\mathcal{A} = \mathcal{U} * \mathcal{D} * \mathcal{U}^\top. \quad (3.3)$$

Let the frontal slices of \mathcal{D} be $D^{(1)}, \dots, D^{(p)}$. If $\hat{\mathcal{D}} \in \mathbb{R}^{n \times n \times p}$ is another T -symmetric f -diagonal tensor, whose frontal slices $\hat{D}^{(1)}, \dots, \hat{D}^{(p)}$ are resulted from switching some diagonal elements of $D^{(1)}, \dots, D^{(p)}$, then there is an orthogonal tensor $\hat{\mathcal{U}} \in \mathbb{R}^{n \times n \times p}$, such that

$$\mathcal{A} = \hat{\mathcal{U}} * \hat{\mathcal{D}} * \hat{\mathcal{U}}^\top. \quad (3.4)$$

Proof Block circulant matrices can be block diagonalized with normalized discrete Fourier transformation (DFT) matrix, which is unitary. Then, as in (3.1) of [4], we have

$$(F_p \otimes I_n) \cdot \text{bcirc}(\mathcal{A}) \cdot (F_p^* \otimes I_n) = \text{diag}(D_1, \dots, D_p), \quad (3.5)$$

where F_p is the $p \times p$ DFT matrix, F_p^* is its conjugate transpose, \cdot is the standard matrix multiplication, \otimes denotes the Kronecker product. Since $\text{bcirc}(\mathcal{A})$ is symmetric, by taking conjugate transpose of (3.5), we see that D_1, \dots, D_p in (3.1) of [4] are all hermite. Applying the eigen-decomposition of $D_i = U_i \Sigma_i U_i^\top$ for $i = 1, \dots, p$, we have

$$\text{diag}(D_1, \dots, D_p) = \text{diag}(U_1, \dots, U_p) \text{diag}(\Sigma_1, \dots, \Sigma_p) \text{diag}(U_1^\top, \dots, U_p^\top). \quad (3.6)$$

Apply $(F_p^* \otimes I_n)$ to the left and $(F_p \otimes I_n)$ to the right of each of the block diagonal matrices in (3.6). In each of the three cases, the resulting triple product results in a block circulant matrix. We have

$$\text{bcirc}(\mathcal{A}) = \text{bcirc}(\mathcal{U}) \text{bcirc}(\mathcal{D}) \text{bcirc}(\mathcal{U}^\top).$$

This implies (3.3). Then we have

$$\mathcal{D} = \mathcal{U}^\top * \mathcal{A} * \mathcal{U},$$

and

$$\mathcal{D}^\top = (\mathcal{U}^\top * \mathcal{A} * \mathcal{U})^\top = \mathcal{U}^\top * \mathcal{A}^\top * \mathcal{U} = \mathcal{U}^\top * \mathcal{A} * \mathcal{U} = \mathcal{D},$$

as \mathcal{A} is T -symmetric. Thus, \mathcal{D} is also T -symmetric.

Switching the order of eigenvalues in the eigen-decomposition $D_i = U_i \Sigma_i U_i^\top$ for $i = 1, \dots, p$, we have (3.4). \square

We call (3.3) a T -eigen-decomposition (TED) of \mathcal{A} .

Corollary 3.4 Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is a T -symmetric tensor and (3.3) holds. Denote $\mathcal{A}^{*2} = \mathcal{A} * \mathcal{A}$ and $\mathcal{A}^{*k} = \mathcal{A}^{*(k-1)} * \mathcal{A}$ for any integer $k \geq 3$. Then for any positive integer k , \mathcal{A}^{*k} is still T -symmetric, and we have

$$\mathcal{A}^{*k} = \mathcal{U} * \mathcal{D}^{*k} * \mathcal{U}^\top.$$

Corollary 3.5 Suppose that $\mathcal{A} \in \mathfrak{R}^{n \times n \times p}$ is a T -symmetric tensor and (3.3) holds. Then \mathcal{A}^{-1} exists if and only if \mathcal{D}^{-1} exists. If they exist, then they are T -symmetric and

$$\mathcal{A}^{-1} = \mathcal{U} * \mathcal{D}^{-1} * \mathcal{U}^\top.$$

We may rewrite (3.3) as

$$\mathcal{A} * \mathcal{U} = \mathcal{U} * \mathcal{D}, \quad (3.7)$$

or

$$\text{bcirc}(\mathcal{A})\text{bcirc}(\mathcal{U}) = \text{bcirc}(\mathcal{U})\text{bcirc}(\mathcal{D}), \quad (3.8)$$

Denote the j th lateral slice of \mathcal{U} by $U_j \in \mathfrak{R}^{n \times p}$ for $j = 1, \dots, n$. Consider the j th column of (3.8) for $j = 1, \dots, n$. Let $\mathcal{D} = (d_{ijk})$. Then $d_{ijk} = 0$ if $i \neq j$. Let $d_{11k} \geq d_{22k} \geq \dots \geq d_{nnk}$ for $k = 1, \dots, p$. We have

$$\mathcal{A} * U_j = \mathbf{d}_j \circ U_j, \quad (3.9)$$

where $\mathbf{d}_j = (d_{jj1}, d_{jjp}, d_{jj(p-1)}, \dots, d_{jj2})^\top$. Since \mathcal{U} is orthogonal, $U_j \neq O$. Thus, \mathbf{d}_j is an eigentuple of \mathcal{A} with an eigenmatrix U_j .

For a matrix $U \in \mathfrak{R}^{n \times p}$, let its column vectors are $\mathbf{u}^{(1)}, \dots, \mathbf{u}^{(p)}$. Then

$$U = (\mathbf{u}^{(1)}, \mathbf{u}^{(2)}, \dots, \mathbf{u}^{(p)}).$$

Denote $U^{[0]} = U$,

$$\begin{aligned} U^{[1]} &= (\mathbf{u}^{(p)}, \mathbf{u}^{(1)}, \dots, \mathbf{u}^{(p-1)}), \\ U^{[2]} &= (\mathbf{u}^{(p-1)}, \mathbf{u}^{(2)}, \dots, \mathbf{u}^{(p-1)}), \\ &\dots, \\ U^{[p-1]} &= (\mathbf{u}^{(2)}, \mathbf{u}^{(3)}, \dots, \mathbf{u}^{(1)}). \end{aligned}$$

Consider the $(n + j)$ th column of (3.8). We have

$$\mathcal{A} * U_j^{[1]} = \mathbf{d}_j \circ U_j^{[1]}. \quad (3.10)$$

Thus, $U_j^{[1]}$ is also an eigenmatrix of \mathcal{A} , associated with the eigentuple \mathbf{d}_j . Similarly, $U_j^{[2]}, \dots, U_j^{[p-1]}$ are also eigenmatrices of \mathcal{A} , associated with the eigentuple \mathbf{d}_j .

Consider the set of eigenmatrices

$$T = \left\{ U_j^{[k]} : j = 1, \dots, n, k = 0, \dots, p-1 \right\}.$$

Then T forms an orthonormal basis of $\mathfrak{R}^{n \times p}$. For any two distinct members $W, V \in T$, let \mathcal{W} and \mathcal{V} be the corresponding $n \times 1 \times p$ tensors. Then we have

$$\mathcal{W}^\top * \mathcal{W} = \mathcal{I}_{11p}, \quad (3.11)$$

and

$$\mathcal{W}^\top * \mathcal{V} = \mathcal{O}_{11p}. \quad (3.12)$$

Viewing (3.4), we may switch the order in $\{d_{11k}, d_{22k}, \dots, d_{nnk}\}$ for any $k = 1, \dots, p$. The resulted $\hat{\mathbf{d}}_j$, $j = 1, \dots, n$ are still eigentuples of \mathcal{A} . Hence, the number of eigentuples of \mathcal{A} is large. But we may always take $\mathbf{d}_1, \dots, \mathbf{d}_n$ in its standard form.

Combining the orthogonality of \mathcal{U} , we have the following theorem.

Theorem 3.6 *Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is a T-symmetric tensor. Then \mathcal{A} has real eigentuples $\mathbf{d}_1, \dots, \mathbf{d}_n$, such that $\mathbf{d}_1 \geq \mathbf{d}_2 \geq \dots \geq \mathbf{d}_n$. For each j , $j = 1, \dots, n$, there are real eigenmatrices $U_j^{[0]}, \dots, U_j^{[p-1]}$, of \mathcal{A} , associated with the eigentuple \mathbf{d}_j . These np eigenmatrices form an orthonormal basis of $\mathbb{R}^{n \times p}$.*

We call the eigentuples $\{\mathbf{d}_1, \dots, \mathbf{d}_n\}$, satisfying $\mathbf{d}_1 \geq \mathbf{d}_2 \geq \dots \geq \mathbf{d}_n$, in Theorem 3.6 the set of the principal eigentuples of \mathcal{A} .

If $\mathcal{A} = \mathcal{I}_{nnp}$, then $\mathcal{U} = \mathcal{D} = \mathcal{I}_{nnp}$. Therefore, $\mathbf{d}_1 = \dots = \mathbf{d}_n = (1, 0, \dots, 0)^\top$. If \mathcal{A} has a set of principal eigentuples $\mathbf{d}_j = (d_{j1}, \dots, d_{jp})^\top$ for $j = 1, \dots, n$, then $\mathcal{A} + \lambda \mathcal{I}_{nnp}$ has a set of principal eigentuples $\mathbf{d}_j = (d_{j1} + \lambda, \dots, d_{jp} + \lambda)^\top$ for $j = 1, \dots, n$.

We are not sure whether all eigentuples of a T-symmetric tensor are real and two eigenmatrices associated with two distinct eigentuples of a T-symmetric tensor are orthogonal to each other. However, but we can prove the following theorem.

Theorem 3.7 *Suppose that $\mathcal{A} \in \mathbb{R}^{n \times n \times p}$ is a T-symmetric tensor, and $\{\mathbf{d}_1, \dots, \mathbf{d}_n\}$ is a set of principal eigentuples of \mathcal{A} such that $\mathbf{d}_1 \geq \dots \geq \mathbf{d}_n$. Then for any real eigentuple \mathbf{d}_0 of \mathcal{A} , we have*

$$\mathbf{d}_1 \geq \mathbf{d}_0 \geq \mathbf{d}_n. \quad (3.13)$$

Proof Assume that there is an eigenmatrix $V \in \mathbb{C}^{n \times p}$ such that

$$\mathcal{A} * V = \mathbf{d}_0 \circ V.$$

Taking conjugate, we have

$$\mathcal{A} * \bar{V} = \mathbf{d}_0 \circ \bar{V}.$$

Let $W = V + \bar{V}$. Then W is real and

$$\mathcal{A} * W = \mathbf{d}_0 \circ W.$$

If W is nonzero, then W is a real eigenmatrix of \mathcal{A} , associated with \mathbf{d}_0 . Otherwise, V is pure imaginary. Letting $\hat{W} = \sqrt{-1}V$, we still have a real eigenmatrix of \mathcal{A} , associated with \mathbf{d}_0 . Without loss of generality, assume W is such a real eigenmatrix.

Let $U_j^{[0]}, \dots, U_j^{[p-1]}$ be the eigenmatrices of \mathcal{A} in Theorem 3.6. Then we have real coefficients $\alpha_j^{[0]}, \dots, \alpha_j^{[p-1]}$, for $j = 1, \dots, n$, such that

$$W = \sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} U_j^{[k-1]}.$$

Let $\mathcal{U}_j^{[k]}$ be the $n \times 1 \times p$ tensors corresponding to $U_j^{[k]}$ for $j = 1, \dots, n$ and $k = 0, \dots, p-1$. Let \mathcal{W} be the $n \times 1 \times p$ tensor corresponding to W . Let \mathcal{D}_j be the $1 \times 1 \times p$ tensors corresponding to \mathbf{d}_j for $j = 0, \dots, n$. Then

$$\begin{aligned} & \mathcal{W}^\top * \mathcal{A} * \mathcal{W} \\ &= \mathcal{W}^\top * \mathcal{W} * \mathcal{D}_0 \\ &= \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} \right)^\top * \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} \right) * \mathcal{D}_0 \\ &= \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{I}_{11p} * \mathcal{D}_0 \\ &= \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{D}_0. \end{aligned}$$

On the other hand,

$$\begin{aligned} & \mathcal{W}^\top * \mathcal{A} * \mathcal{W} \\ &= \mathcal{W}^\top * \mathcal{A} * \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} \right) \\ &= \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} \right)^\top * \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{A} * \mathcal{U}_j^{[k-1]} \right) \\ &= \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} \right)^\top * \left(\sum_{j=1}^n \sum_{k=1}^p \alpha_j^{[k-1]} \mathcal{U}_j^{[k-1]} * \mathcal{D}_j \right) \\ &= \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{I}_{11p} * \mathcal{D}_j \\ &= \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{D}_j. \end{aligned}$$

From this, we have

$$\sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{D}_0 = \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathcal{D}_j,$$

i.e.,

$$\sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathbf{d}_0 = \sum_{j=1}^n \left(\sum_{k=1}^p \alpha_j^{[k-1]} \right)^2 \mathbf{d}_j.$$

The inequality (3.13) is obtained. \square

Corollary 3.8 *The eigentuples \mathbf{d}_1 and \mathbf{d}_n are unique to \mathcal{A} .*

We call \mathbf{d}_1 the largest eigentuple of \mathcal{A} , and \mathbf{d}_n the smallest eigentuple of \mathcal{A} .

4 T-Symmetric Positive Semidefinite Tensors

Suppose that $\mathcal{A} \in \Re^{n \times n \times p}$ is a T-square tensor. Then by Proposition ??, the T-quadratic form is positive semidefinite (definite) if and only if the T-symmetric tensor $\mathcal{A} + \mathcal{A}^\top$ is positive semidefinite (definite). This stimulates us to study positive semidefiniteness (definiteness) of a T-symmetric tensor.

Theorem 4.1 *Suppose that $\mathcal{A} \in \Re^{n \times n \times p}$ is a T-symmetric tensor and it has a set of principal eigentuples $\mathbf{d}_1, \dots, \mathbf{d}_n$, such that $\mathbf{d}_1 \geq \mathbf{d}_2 \cdots \geq \mathbf{d}_n$. Then \mathcal{A} is positive semidefinite (definite) if and only if the smallest eigentuple $\mathbf{d}_n \geq (>)\mathbf{0}$.*

Proof By Theorem 3.6, $\mathbf{d}_1 \geq \mathbf{d}_2 \cdots \geq \mathbf{d}_n$, and for each j , $j = 1, \dots, n$, there are real eigenmatrices $U_j^{[0]}, \dots, U_j^{[p-1]}$, of \mathcal{A} , associated with the eigentuple \mathbf{d}_j , such that these np eigenmatrices form an orthonormal basis of $\Re^{n \times p}$.

If \mathbf{d}_n is not nonnegative, let $U = U_n^{[0]}$ and $\mathcal{U}_n^{[0]}$ be the corresponding $n \times 1 \times p$ tensor. Let \mathbf{L} be the $1 \times 1 \times p$ tensor corresponding to \mathbf{d}_n . Then

$$F_{\mathcal{A}}(U) = (\mathcal{U}_n^{[0]})^\top * \mathcal{A} * \mathcal{U}_n^{[0]} = (\mathcal{U}_n^{[0]})^\top * \mathcal{U}_n^{[0]} * \mathbf{L} = \mathcal{I}_{nnp} * \mathbf{L} \not\geq \mathbf{0},$$

which implies that $F_{\mathcal{A}}$ is not positive semi-definite. Similarly, if \mathbf{d}_n is not positive, then $F_{\mathcal{A}}$ is not positive definite.

On the other hand, suppose that $\mathbf{d}_n \geq \mathbf{0}$. Let $\mathcal{U}_j^{[k]}$ be the $n \times 1 \times p$ tensor corresponding to $U_j^{[k]}$ for $j = 1, \dots, n$ and $k = 0, \dots, p-1$. Let $X \in \Re^{n \times p}$. Then there are real coefficients $\alpha_j^{[k]}$ for $j = 1, \dots, n$ and $k = 0, \dots, p-1$, such that

$$X = \sum_{j=1}^n \sum_{k=0}^{p-1} \alpha_j^{[k]} U_j^{[k]}.$$

Let \mathbf{L}_j be the $1 \times 1 \times p$ tensor corresponding to \mathbf{s}_j for $j = 1, \dots, n$. We have

$$\begin{aligned}
F_{\mathcal{A}}(X) &= \left(\sum_{i=1}^n \sum_{l=0}^{p-1} \alpha_i^{[l]} \mathcal{U}_i^{[l]} \right)^{\top} * \mathcal{A} * \left(\sum_{j=1}^n \sum_{k=0}^{p-1} \alpha_j^{[k]} \mathcal{U}_j^{[k]} \right) \\
&= \sum_{i=1}^n \sum_{j=1}^n \sum_{l=0}^{p-1} \sum_{k=0}^{p-1} \alpha_i^{[l]} \alpha_j^{[k]} (\mathcal{U}_i^{[l]})^{\top} * \mathcal{A} * \mathcal{U}_j^{[k]} \\
&= \sum_{i=1}^n \sum_{j=1}^n \sum_{l=0}^{p-1} \sum_{k=0}^{p-1} \alpha_i^{[l]} \alpha_j^{[k]} (\mathcal{U}_i^{[l]})^{\top} * \mathcal{U}_j^{[k]} * \mathbf{L}_j \\
&= \sum_{j=1}^n \sum_{k=0}^{p-1} \left(\alpha_j^{[k]} \right)^2 \mathbf{d}_j \\
&\geq \mathbf{0}.
\end{aligned}$$

Thus, $F_{\mathcal{A}}$ is positive semidefinite. Similarly, if $\mathbf{d}_n \geq \mathbf{0}$, then $\mathcal{F}_{\mathcal{A}}$ is positive definite. \square

5 Relation with TSVD of General Third Order Tensors

Suppose that $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$. By [4], \mathcal{A} has a T-singular value decomposition (TSVD):

$$\mathcal{A} = \mathcal{U} * \mathcal{S} * \mathcal{V}^{\top}, \quad (5.14)$$

where $\mathcal{U} \in \mathbb{R}^{m \times m \times p}$ and $\mathcal{V} \in \mathbb{R}^{n \times n \times p}$ are orthogonal tensors, $\mathcal{S} \in \mathbb{R}^{m \times n \times p}$ is a f-diagonal tensor.

Theorem 5.1 *Suppose that $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$ with TSVD (5.14). Then $\mathcal{A} * \mathcal{A}^{\top} \in \mathbb{R}^{m \times m \times p}$ and $\mathcal{A}^{\top} * \mathcal{A} \in \mathbb{R}^{n \times n \times p}$ are T-symmetric positive semi-definite tensors with TED*

$$\mathcal{A} * \mathcal{A}^{\top} = \mathcal{U} * (\mathcal{S} * \mathcal{S}^{\top}) * \mathcal{U}^{\top},$$

and

$$\mathcal{A}^{\top} * \mathcal{A} = \mathcal{V} * (\mathcal{S}^{\top} * \mathcal{S}) * \mathcal{V}^{\top},$$

respectively.

We now define singular tuples and singular matrices of general third order tensors. Suppose that $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$ is a third order tensor, $X \in \mathbb{R}^{n \times p}$, $X \neq O$, $Y \in \mathbb{R}^{m \times p}$, $Y \neq O$, $\mathbf{s} \in \mathbb{R}^p$, and

$$\mathcal{A} * X = \mathbf{s} \circ Y \quad (5.15)$$

and

$$\mathcal{A}^\top * Y = \mathbf{s} \circ X. \quad (5.16)$$

Then we call \mathbf{s} a singular tuple of \mathcal{A} , X a right singular matrix of \mathcal{A} , Y a right singular matrix of \mathcal{A} , corresponding to the singular tuple \mathbf{s} .

Theorem 5.2 *Suppose that $\mathcal{A} \in \mathbb{R}^{m \times n \times p}$ is a third order tensor. Without loss of generality, assume that $n \leq m$. Then \mathcal{A} has singular tuples $\mathbf{s}_1 \geq \mathbf{s}_2 \geq \dots \geq \mathbf{s}_n \geq \mathbf{0}$. For each j , $j = 1, \dots, n$, there are right singular matrices $U_j^{[0]}, \dots, U_j^{[p-1]}$, and left singular matrices $V_j^{[0]}, \dots, V_j^{[p-1]}$, of \mathcal{A} , associated with the singular tuple \mathbf{s}_j . The np singular matrices $U_j^{[0]}, \dots, U_j^{[p-1]}$ form an orthonormal basis of $\mathbb{R}^{n \times p}$, and the np singular matrices $V_j^{[0]}, \dots, V_j^{[p-1]}$ form a part of an orthonormal basis of $\mathbb{R}^{m \times p}$, respectively.*

*Furthermore, $\mathcal{A}^\top * \mathcal{A} \in \mathbb{R}^{n \times n \times p}$ and $\mathcal{A} * \mathcal{A}^\top \in \mathbb{R}^{m \times m \times p}$ are two T -symmetric positive semidefinite tensors. The tensor $\mathcal{A}^\top * \mathcal{A}$ has n nonnegative eigentuples $\mathbf{s}_1^{\odot 2}, \dots, \mathbf{s}_n^{\odot 2}$. For each j , $j = 1, \dots, n$, there are real eigenmatrices $U_j^{[0]}, \dots, U_j^{[p-1]}$, of \mathcal{A} , associated with the eigentuple $\mathbf{s}_j^{\odot 2}$. The tensor $\mathcal{A} * \mathcal{A}^\top$ has n nonnegative eigentuples $\mathbf{s}_1^{\odot 2}, \dots, \mathbf{s}_n^{\odot 2}$. For each j , $j = 1, \dots, n$, there are real eigenmatrices $V_j^{[0]}, \dots, V_j^{[p-1]}$, of \mathcal{A} , associated with the eigentuples $\mathbf{s}_j^{\odot 2}$. If $n < m$, for each j , $j = n + 1, \dots, m$, there are real eigenmatrices $V_j^{[0]}, \dots, V_j^{[p-1]}$, of \mathcal{A} , associated with the zero eigentuple $\mathbf{0} \in \mathbb{R}^p$. The mp singular matrices $V_j^{[0]}, \dots, V_j^{[p-1]}$ form an orthonormal basis of $\mathbb{R}^{m \times p}$.*

Acknowledgment We are thankful to Prof. Yicong Zhou and Dr. Dongdong Chen for the discussion on the multi-view clustering problem and the image feature extraction problem.

References

- [1] K. Bradman, “Third-Order tensors as linear operators on a space of matrices”, *Linear Algebra and Its Applications* **433** (2010) 1241-1253.
- [2] Y. Chen, X. Xiao and Y. Zhou, “Multi-view subspace clustering via simultaneously learning the representation tensor and affinity matrix”, *Pattern Recognition* **106** (2020) 107441.
- [3] M. Kilmer, K. Braman, N. Hao and R. Hoover, “Third-order tensors as operators on matrices: A theoretical and computational framework with applications in imaging”, *SIAM Journal on Matrix Analysis and Applications* **34** (2013) 148-172.
- [4] M. Kilmer and C.D. Martin, “Factorization strategies for third-order tensors”, *Linear Algebra and Its Applications* **435** (2011) 641-658.

- [5] Y. Miao, L. Qi and Y. Wei, “Generalized tensor function via the tensor singular value decomposition based on the T-product”, *Linear Algebra and Its Applications* **590** (2020) 258-303.
- [6] Y. Miao, L. Qi and Y. Wei, “T-Jordan canonical form and T-Drazin inverse based on the T-product”, *Communications on Applied Mathematics and Computation* **3** (2021) doi.org/10.1007/s42967-019-00055-4.
- [7] O. Semerci, N. Hao, M.E. Kilmer and E.L. Miller, “Tensorbased formulation and nuclear norm regularization for multienergy computed tomography”, *IEEE Transactions on Image Processing* **23** (2014) 1678–1693.
- [8] G. Song, M.K. Ng and X. Zhang, “Robust Tensor Completion Using Transformed Tensor SVD”, *Numerical Linear Algebra with Applications* doi.org/10.1002/nla.2299.
- [9] X. Xiao, Y. Chen, Y.J. Gong and Y. Zhou, “Low-rank reserving t-linear projection for robust image feature extraction”, *IEEE Transactions on image processing* **30**, (2021) 108-120.
- [10] X. Xiao, Y. Chen, Y.J. Gong and Y. Zhou, “Prior knowledge regularized multiview self-representation and its applications”, *IEEE Transactions on neural networks and learning systems*, in press.
- [11] L. Yang, Z.H. Huang, S. Hu and J. Han, “An iterative algorithm for third-order tensor multi-rank minimization”, *Computational Optimization and Applications* **63** (2016) 169-202.
- [12] M. Zheng, Z. Huang and Y. Wang, “T-positive semidefiniteness of third-order symmetric tensors and T-semidefinite programming”, *Computational Optimization and Applications* doi.org/10.1007/s10589-020-00231-w.
- [13] J. Zhang, A.K. Saibaba, M.E. Kilmer and S. Aeron, “A randomized tensor singular value decomposition based on the t-product”, *Numerical Linear Algebra with Applications* **25** (2018) e2179.
- [14] Z. Zhang and S. Aeron, “Exact tensor completion using t-SVD”, *IEEE Transactions on Signal Processing* **65** (2017) 1511-1526.
- [15] Z. Zhang, G. Ely, S. Aeron, N. Hao and M. Kilmer, “Novel methods for multilinear data completion and de-noising based on tensor-SVD”, *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition, ser. CVPR ’14* (2014) 3842-3849.

- [16] P. Zhou, C. Lu, Z. Lin and C. Zhang, “Tensor factorization for low-rank tensor completion”, *IEEE Transactions on Image Processing* **27** (2018) 1152-1163.