A Positive Energy Theorem for Fourth-Order Gravity

R. Avalos, P. Laurain, J. H. Lira*

December 1, 2021

Abstract

In this paper we prove a positive energy theorem related to fourth-order gravitational theories, which is a higher-order analogue of the classical ADM positive energy theorem of general relativity. We will also show that, in parallel to the corresponding situation in general relativity, this result intersects several important problems in geometric analysis. For instance, it underlies positive mass theorems associated to the Paneitz operator, playing a similar role in the positive Q-curvature conformal prescription problem as the Schoen-Yau positive energy theorem does for the Yamabe problem. Several other links to Q-curvature analysis and rigidity phenomena are established.

1 Introduction

In this paper we will analyse the properties of a recently proposed energy associated to higher-order gravitational theories in the stationary limit, see [3]. Specifically, in parallel work, we have analysed gravitational theories described on globally hyperbolic space-times $(V \doteq M \times \mathbb{R}, \bar{g})$ by an action functional of the form

$$S(\bar{g}) = \int_{V} \left(\alpha R_{\bar{g}}^{2} + \beta \langle \operatorname{Ric}_{\bar{g}}, \operatorname{Ric}_{\bar{g}} \rangle \right) dV_{\bar{g}}, \tag{1}$$

where α and β are free parameters in this variational setting. Let us highlight that the study of these kinds of higher-order gravitational action functional is well-motivated within contemporary theoretical physics as they appear in connection with effective field theories of gravity [6, 16, 39], as well as in the context of inflationary cosmology [44] and certain approaches to quantum gravity, such as conformal gravity [31, 32].

In order to make sense of the above functional, we can assume that the class of metrics considered above are such that $R_{\bar{g}}^2$ and $\langle \operatorname{Ric}_{\bar{g}}, \operatorname{Ric}_{\bar{g}} \rangle$ are integrable. Then, the functional $\bar{g} \mapsto S(\bar{g})$ is well-defined and we have an L^2 -gradient for this functional, given by a divergence-free tensor field $A_{\bar{g}} \in \Gamma(T_2^0 V)$, which is explicitly given by

$$A_{\bar{g}} = \beta \Box_{\bar{g}} \operatorname{Ric}_{\bar{g}} + (\frac{1}{2}\beta + 2\alpha) \Box_{\bar{g}} R \, \bar{g} - (2\alpha + \beta) \bar{\nabla}^2 R_{\bar{g}} - 2\beta \operatorname{Ric}_{\bar{g}} \operatorname{Riem}_{\bar{g}} + 2\alpha R_{\bar{g}} \operatorname{Ric}_{\bar{g}} - \frac{1}{2} \alpha R_{\bar{g}}^2 \bar{g} - \frac{1}{2} \beta \langle \operatorname{Ric}_{\bar{g}}, \operatorname{Ric}_{\bar{g}} \rangle_{\bar{g}} \bar{g},$$

$$(2)$$

^{*}Federal University of Ceará, Mathematics Department, Fortaleza, Ceará, Brazil

[†]Institut Mathématique de Jussieu, Université de Paris, Bâtiment Sophie Germain, Case 7052, 75205 Paris Cédex 13, France & DMA, Ecole normale supérieure, CNRS, PSL Research University, 75005 Paris.

where above we denoted $\operatorname{Ric}_{\bar{g}}\operatorname{Riem}_{\bar{g}_{ij}} \doteq \operatorname{Ric}_{\bar{g}}^{kl}\operatorname{Riem}_{\bar{g}_{kijl}}$. In a parallel situation to what is well-known in the context of general relativity (GR), we have shown that there is a canonical notion of energy, which we denote by $\mathcal{E}_{\alpha,\beta}(\bar{g})$, associated to asymptotically Euclidean (AE) solutions of the space-time field equations $A_{\bar{g}} = 0$ which arise as perturbations of solutions \bar{g}_0 which possess a time-like Killing field. Although the analysis of such an energy could be quite involved in general, we are able to identify some particular choices of the parameters α and β for which its analysis is tractable. More important, we establish positivity and rigidity results results for the energy in those cases. These results are intimately connected with the existence of metrics with positive constant Q-curvature

Our aim here is to analyse the particular case of stationary solutions of the fourth-order field equations parameterized by $2\alpha + \beta = 0$. Recall that globally hyperbolic stationary space-times are manifolds of the form $V = M \times \mathbb{R}$ endowed with a Lorentzian metric that can be written as

$$\bar{g} = -N^2 dt^2 + \tilde{g},\tag{3}$$

where $N: M \mapsto \mathbb{R}^+$ is the *lapse* function and $\tilde{g} \in \Gamma(T_2^0 M)$ restricts to a Riemannian metric g on each t = constant hypersurface. In this setting, the appropriate notion of energy associated to the action S and the corresponding field equations $A_{\bar{q}} = 0$ becomes

$$\mathcal{E}_{\alpha}(g) \doteq -\alpha \lim_{r \to \infty} \int_{S_r^{n-1}} \left(\partial_j \partial_i \partial_i g_{aa} - \partial_j \partial_u \partial_i g_{ui} \right) \nu^j d\omega_r.$$
(4)

In this expression, we are assuming that (M, g) is an AE manifold. This means that for a given compact set K the asymptotic region $M \setminus K$ is diffeomorphic to $\mathbb{R}^n \setminus \overline{B_1(0)}$ and it is foliated by topological (n-1)-dimensional spheres S_r^{n-1} , whose Euclidean volume element is denoted in the expression above by $d\omega_r$. There, ν stands for the Euclidean unit normal field to these spheres.

The nature of (4) as a conserved quantity in the context of higher-order gravitational theories make $\mathcal{E}_{\alpha}(g)$ a very good fourth-order analogue to the Arnowitt, Deser and Misner (ADM) energy in the context of GR. Let us recall that the total energy of an isolated gravitational system in GR, whose initial data is modelled as an AE manifold, is given by (see [1])

$$E_{ADM}(g) = c(n) \lim_{r \to \infty} \int_{S_r^{n-1}} \left(\partial_i g_{ji} - \partial_j g_{ii} \right) \nu^j d\omega_r, \tag{5}$$

where c(n) stands for a dimensional constant (see [4] for the detailed analytical properties of (5)). This energy E_{ADM} has had a huge impact both within GR and in geometric analysis. Most notoriously, it was the work of R. Schoen in [40] that elucidated the role that the ADM energy plays in the final resolution of the Yamabe problem (see also [28] for a review on this topic). In particular, in order to solve the Yamabe problem in the positive Yamabe class, in dimensions 3,4 and 5 or in locally conformally flat manifolds, Schoen noticed that it was enough to prove that an appropriate constant appearing in the expansion of the Green function G_{L_g} associated to the conformal Laplacian L_g was non-negative and that the zero case implied rigidity with the round sphere. Furthermore, it was pointed out that this constant was precisely the ADM energy of the AE manifold obtained via an stereographic projection. Therefore, the proof of the positive energy theorem in GR actually underlies the proof of the Yamabe problem in these cases. This beautiful relation shows what a fundamental result the positive energy theorem actually is within geometric analysis, being a cornerstone in the resolution of the Yamabe problem. Since then, the ADM energy has influenced many other constructions within geometric analysis and mathematical GR, which are not necessarily concerned with the Yamabe problem. For instance, rigidity phenomena associated to positive scalar curvature [7, 9, 10], isoperimetric problems on AE manifolds [5, 17, 25], geometric foliations and center of mass constructions [17, 24, 25, 37] and even gluing constructions [10] (for nice reviews of many of these topics, see [8, 27]).

In view of the above paragraph, we consider that the analysis of the positive energy theorem associated to (4) stands as a highly well-motivated problem both for the development of a program devoted to the mathematical analysis of fourth order theories of gravity, as well as a tool which can potentially play a fundamental role in several fourth order geometric problems. Thus, the main objective of this paper is the establishment of such a positive energy theorem, and, afterwards, we will show how it underlies some fundamental problems in geometric analysis.

Since we will be concerned with positivity issues related to $\mathcal{E}_{\alpha}(g)$, we must actually fix the sign of α a priori. For reasons that will become apparent through this paper, we will fix $\alpha = -1$ and define $\mathcal{E}(g) \doteq \mathcal{E}_{-1}(g)$. Let us notice that before embarking on the proof of the positive energy theorem, we should first analyse under what conditions $\mathcal{E}(g)$ is actually well-defined and prove that it is actually an intrinsic geometric object within a suitable class of AE-manifolds. This will be done in Proposition 1, Proposition 2 and Theorem 1. Once this is done, we will first explore appropriate geometric conditions that could in principle provide a rigidity statement in the critical cases. Explicitly, such conditions will involve a Q-curvature condition of the form $Q_g \geq 0$ (see Proposition 3 and Theorem 4). Let us recall that given a Riemannian manifold (M^n, g) , $n \geq 3$, its Q-curvature is defined by¹

$$Q_g \doteq -\frac{1}{2(n-1)}\Delta_g R_g - \frac{2}{(n-2)^2} |\operatorname{Ric}_g|_g^2 + \frac{n^3 - 4n^2 + 16n - 16}{8(n-1)^2(n-2)^2} R_g^2.$$

In this context, we will establish the following theorem:

Theorem A (Positive Energy). Let (M^n, g) be an n-dimensional AE manifold, with $n \geq 3$, which satisfies the decaying conditions: (i) $g_{ij} - \delta_{ij} = O_4(r^{-\tau})$, with $\tau > \max\{0, \frac{n-4}{2}\}$, in some coordinate system associated to an structure of infinity; (ii) $Q_g \in L^1(M, dV_g)$, and such that Y([g]) > 0 and $Q_g \geq 0$. Then, the fourth order energy $\mathcal{E}(g)$ is non-negative and $\mathcal{E}(g) = 0$ if and only if (M, g) is isometric to (\mathbb{R}^n, δ) .

The proof of this statement will be the content of Theorem 2 (see also theorems 3 and 4). Let us notice that some hypotheses in the above theorem can be relaxed while keeping important results. In particular, the positivity and rigidity statements are somehow decoupled. This implies that under hypotheses (i)-(ii) and $R_g > 0$ at infinity, it follows that $\mathcal{E}(g) \geq 0$ (see Theorem 3), while under hypotheses (i)-(ii), $Q_g \geq 0$ and g Yamabe positive, if $\mathcal{E}(g) = 0$, then the rigidity statement follows (see Theorem 4). Furthermore, let us notice that the borderline case of n = 4 is special, since it follows that any 4-dimensional AE manifold satisfies the decaying conditions (i)-(ii) and, in fact, under these conditions $\mathcal{E}(g) = 0$. These last comments also apply to the case n = 3, which can be checked explicitly quite straightforwardly. This imposes restrictions on the positive curvature conditions that these manifolds can admit, providing us with the following corollaries.

¹See Appendix B for details on our conventions on Q-curvature.

Corollary. Any n-dimensional AE-Riemannian manifold (M^n, g) , with $n \in \{3, 4\}$, such that $Q_g \ge 0$ and Y([g]) > 0, is isometric to (\mathbb{R}^n, δ) .

Let us notice that Yamabe positive AE manifolds in low dimensions (specially three) are very natural objects for instance in GR, since maximal vacuum initial data for isolated systems belong to this class of manifolds. Therefore, the above corollary, in some sense, can be used to separate the trivial solutions via an appeal to their *Q*-curvature.

The following result, which in particular gives a conformally invariant statement, is also a direct consequence of the analysis related to the above positive energy theorem for $\mathcal{E}(g)$ (see Proposition 5 for more details).

Corollary. Let (M, g) be an asymptotically Euclidean four manifold with $\kappa_g = \int_M Q_g \, dv_g \ge 0$ and Y([g]) > 0, then (M, g) is conformal to the euclidean \mathbb{R}^4 .

After presenting the above results concerning the positive energy theorem associated to $\mathcal{E}(g)$, we will make contact with important problems in geometric analysis. In particular, our aim is to show that that there is a very clear parallel in the role played by the ADM positive energy theorems of Schoen-Yau [41, 42, 43] in the resolution of the Yamabe problem, to the role Theorem A plays in the resolution of the positive *Q*-curvature Yamabe-type problems which are known up to date. This problem concerns finding a conformal deformation of a closed Riemannian manifold (M, g) so that the resulting metric has constant *Q*-curvature. There has been great progress in this program in recent times and the analysis depends on whether $n \geq 5$ or n = 3, 4. In particular, for the most updated resolutions of this problem to our knowledge in dimension four see [15, 18, 11] and references therein, while for $n \geq 5$, with different degrees of generality, let us draw the reader's attention to [23, 20, 38]. Although we will not be concerned with the three dimensional case, let us point out that the problem has been addressed in this case in [21, 22].

Let us now focus on the results of [23]. There, the authors prove that given a closed manifold (M,g) of dimension $n \ge 5$ which is Yamabe non-negative and satisfies $Q_g \ge 0$ not identically zero,² there is a conformal deformation to constant positive Q-curvature. In particular, this is done in Theorem 4 therein where in the cases of dimensions n = 5, 6, 7 or $n \geq 5$ and locally conformally flat, there is a parallel to Schoen's resolution of the Yamabe problem. That is, the problem can be reduced to showing that a certain coefficient in the Green function expansion of the Paneitz operator near a pole is non-negative. In analogy to Schoen's ideas, this constant has been called the mass of the Paneitz operator (whenever defined) and its positivity has been analysed first by Humbert-Raulot in the locally conformally flat case [26], then by Gursky-Malchiodi who incorporated the cases $5 \leq n \leq 7$ without this last restriction, and finally by Hang-Yang [21], who weakened the hypotheses on the scalar curvature imposed in [20] and achieved the final form of this results which was applied in Theorem 4 of [23]. Along these lines, let us also draw the reader's attention to an unpublished work by B. Michel [34], who also analysed these kinds of positive mass theorems associated to the Green function of the Paneitz operator previously and obtained related results through similar methods to those of [26, 20, 21].

In Section 2.2 we will show that exactly in dimensions $5 \leq n \leq 7$ or if (M, g) has a point p around which it is conformally flat, then the mass of the Green function G_{P_g} associated to the Paneitz operator P_g is positively proportional to the energy $\mathcal{E}(\hat{g})$ of the asymptotically euclidean manifold $(\hat{M} = M \setminus \{p\}, \hat{g} = G_{P_g}^{\frac{4}{n-4}}g)$ obtained via a stereographic

²In this cases we will say that Q_g is **semi-positive**.

projection. Then, we show that the following Q-curvature positive mass theorem follows from Theorem A:

Theorem B (Q-curvature Positive Mass [20, 23, 26]). Let (M, g) be a closed n-dimensional Riemannian manifold, with $5 \le n \le 7$ or $n \ge 8$ and locally conformally flat around some point $p \in M$. If $Y([g]) \ge 0$ and (M, g) admits a conformal metric with semi-positive Q-curvature, then the mass of G_{P_g} at p is non-negative and vanishes if and only if (M, g)is conformal to the standard sphere.

The above theorem is exactly the positive mass theorem used by Hang-Yang in [23] to solve the Q-curvature prescription problem in dimensions $5 \le n \le 7$ or $n \ge 5$ and M locally conformally flat around a point. This highlights the potential parallel of energy $\mathcal{E}(g)$ in the analysis of fourth-order problems to the role of the ADM energy in classical second order geometric problems.

Finally, along the lines of the remarks of the previous paragraphs, we will analyse the critical four-dimensional case and provide an independent proof a Theorem B in [18] appealing to the techniques derived in this paper. Concretely, the following theorem follows from our analysis:

Theorem C (Gursky). Let (M^4, g) a 4-dimensional manifold with $Y([g]) \ge 0$, then $\kappa_q \le 16\pi^2$ with equality holding iff (M^4, g) is conformal to the standard sphere.

We would like to highlight that, in this case, our techniques make contact with the positive mass theorem of 2-dimensional manifolds, as presented in [27].

Finally, let us comment that, with the aim of delivering a self-contained presentation but trying to avoid a long introduction to preliminary results concerning analysis on AE-manifolds, *Q*-curvature analysis and constructions concerning conformal normal coordinates, which are things very well-known for experts in each of these fields, we have compiled the main results which will be used in this paper is the Appendices A,B and C, where detailed references can be found.

It has been brought to our attention by Marc Herzlich that one of his students, Benoît Michel, has previously analysed a similar notion of energy, see [34].

Acknowledgments: The authors would like to thank the CAPES-COFECUB, CAPES-PNPD and ANR (ANR- 18-CE40-002) for their financial support. Also, we would like thank the comments, suggestions and critics made by an anonymous referee to this paper, which have helped us improve the presentation and content of the paper.

2 Preliminaries

In this section we will collect some definitions and notational conventions which will be used along the paper.

Some notational conventions:

- (M^n, g) will denote an *n*-dimensional Riemannian manifold.
- Given a Riemannian manifold (M, g), we will denote by $\Delta_g u = g^{ij} \nabla_i \nabla_j u$ the negative Laplacian.

• Given a Riemannian manifold (M, g) we denote by ∇ its Riemannian connection and define the curvature tensor by

$$R(V,W)Z = \nabla_V \nabla_W Z - \nabla_W \nabla_V Z - \nabla_{[V,W]} Z, \quad V,W,Z \in \Gamma(TM),$$

and we label its components in some coordinate system $\{x^i\}_{i=1}^n$ via $R_{jkl}^i = dx^i (R(\partial_k, \partial_l)\partial_j)$. Then, the Ricci tensor is defined locally via the contraction $R_{jl} = R_{jkl}^k$.

- Given a tensor field T, indices of $\nabla^k T$ resulting from covariant differentiation will be separated by a comma. That is, if T_{ij} are the components of a 2-tensor, then we denote the components of $\nabla^2 T$ by $T_{ij,kl}$.
- Given $\Sigma \hookrightarrow M$ an embedded compact hypersuface in (M, g), we will denote by ν the outward pointing g-unit normal to Σ .
- Given $\Sigma \hookrightarrow \mathbb{R}^n$ an embedded compact hypersuface, we will denote by ν^e Euclidean outward pointing unit normal to Σ .
- ω_{n-1} will denote the volume of the unit sphere $\mathbb{S}^{n-1} \hookrightarrow \mathbb{R}^n$ and $d\omega_{n-1}$ is its canonical volume measure.
- Given a compact manifold (M, g), the conformal class of g is $\{ug ; u \in C^{\infty}(M, \mathbb{R}^*_+)\}$, denoted [g].
- Given a compact manifold (M, g), its Yamabe invariant is defined as

$$Y([g]) = \inf_{\tilde{g}} \frac{\int_M R_{\tilde{g}} dv_{\tilde{g}}}{\operatorname{vol}(M)^{\frac{n-2}{2}}}.$$

• Given a function $f \in C^k(\mathbb{R}^n)$, we say that $f = O_k(|x|^{\tau})$ for some $k \ge 0$ and $\tau \in \mathbb{R}$ if for any multi-index α with $|\alpha| \le k$, the functions $\partial^{\alpha} f = O(|x|^{\tau-|\alpha|})$ either as $|x| \to \infty$ or $|x| \to 0$, depending on the context.

Definition 1 (Weighted spaces). Given $x \in \mathbb{R}^n$, let us define $r(x) \doteq |x|$, $\sigma(x) \doteq (1 + r^2(x))^{\frac{1}{2}}$ and consider $\delta \in \mathbb{R}$. Then, we set

- $L^p_{\delta}(\mathbb{R}^n) = \{ u \in L^p_{loc} \mid \int_{\mathbb{R}^n} |u|^p \sigma^{-\delta p n} dx < +\infty \}, we equip this set with the norm <math>||u||^p_{p,\delta} = \int_{\mathbb{R}^n} |u|^p \sigma^{-\delta p n} dx.$
- $L^{\infty}_{\delta}(\mathbb{R}^n) = \left\{ u \in L^{\infty}_{loc} \mid \sup_{\mathbb{R}^n} |u|\sigma^{-\delta} < +\infty \right\}$, we equip this set with the norm $||u||_{\infty,\delta} = \sup_{\mathbb{R}^n} |u|\sigma^{-\delta}$.
- $W^{k,p}_{\delta}(\mathbb{R}^n) = \left\{ u \in W^{k,p}_{loc} \mid \sum_{j=0}^k \|d^j u\|_{p,\delta-j} < +\infty \right\}, \text{ we equip this set with the norm} \|u\|_{k,p,\delta} = \sum_{j=0}^k \|d^j u\|_{p,\delta-j}.$

In particular, by Sobolev embedding, when kp > n if $u \in W^{k,p}_{\delta}(\mathbb{R}^n)$ then $u = O(r^{\delta})$ at infinity. See [4] for more details about weighted spaces.

Then we introduce the notion of asymptotically flat manifolds.

Definition 2 (AE manifolds). A complete Riemannian manifold (M, g) with $g \in W_{loc}^{k,q}(M)$ for some $k \ge 1$ and q > n is said to be asymptotically Euclidean (with one end) if there exists a compact set $K \subset M$ and a diffeomorphism $\phi : M \setminus K \to \mathbb{R}^n \setminus \overline{B_1(0)}$ such that • $\phi_*(g)$ is a uniformly positive defined metric, i.e. there exists $\lambda > 1$ such that

$$\frac{1}{\lambda} |\xi|^2 \le g_{ij}(x) \xi^i \xi^j \le \lambda |\xi|^2 \, \forall x \in \mathbb{R}^n \setminus B_1 \, \forall \xi \in \mathbb{R}^n.$$

•

$$\phi_*(g)_{ij} - \delta_{ij} \in W^{k,q}_{-\tau}(\mathbb{R}^n \setminus B(0,1))$$

for some $\tau > 0$ called the decreasing rate.

Important remark: In this chart we defined σ and we remark that the definition of L^q_{δ} is independent of the chart. But $W^{k,q}_{\delta}$ depends on the chart ϕ , since the partial derivatives will depend on the choice of coordinates. It will be denoted $W^{k,q}_{\delta}(\phi)$. In the rest of the section, we will call such a chart **a structure at infinity**. Once the structure at infinity is chosen, we naturally extend the definition of $W^{k,q}_{\delta}(\mathbb{R}^n \setminus B(0,1))$ to $W^{k,q}_{\delta}(M)$ on the compact part, noticing that all choices of charts in the compact region define equivalent norms.

In the following definition, which will be used in the core of this paper, we will restrict to AE manifold which posses more regularity than that of the general definition given above.

Definition 3. We will say that a (smooth) AE manifold (M,g) is of order $\tau > 0$ with respect to some structure at infinity $\Phi : M \setminus K \mapsto \mathbb{R}^n \setminus \overline{B}$, if, in such coordinates, $\Phi_* g_{ij} - \delta_{ij} = O_4(|x|^{-\tau})$.

Of course we can define asymptotically flat structures with multiple ends. But since analysis phenomena are determined by the behaviour at infinity, it is very easy to isolate each end and to consider that there is only one.

3 The positive energy theorem

The main result of this section will be a positive energy theorem related to (4) with its corresponding rigidity statement. But, before this, we will begin by analysing geometric conditions under which (4) is well-defined. In the following proposition, we will establish such geometric criteria. Let us first set

$$\tau_n = \begin{cases} 0 & \text{if } n = 3, 4\\ \frac{n}{2} - 2 & \text{if } n \ge 5 \end{cases}$$

Proposition 1. Let (M^n, g) be an AE manifold of dimension $n \ge 3$ satisfying the following conditions

- i) There are end rectangular coordinates, given by a structure of infinity Φ , where $g_{ij} = \delta_{ij} + O_4(r^{-\tau})$, where $\tau > \tau_n$;
- ii) The Q-curvature of g is in $L^1(M, dV_q)$.

Then, given an exhaustion of M by compact sets Ω_k such that $S_k \doteq \Phi(\partial \Omega_k)$ are smooth connected (n-1)-dimensional manifolds without boundary in \mathbb{R}^n satisfying

$$R_{k} \doteq \inf\{|x| : x \in S_{k}\} \xrightarrow[k \to \infty]{} \infty,$$

$$R_{k}^{n-1} \operatorname{area}(S_{k}) \text{ is bounded as } k \to \infty,$$
(6)

the limit

$$\mathcal{E}^{(\Phi)}(g) = \lim_{k \to \infty} \int_{S_k} \left(\partial_j \partial_i \partial_i g_{aa} - \partial_j \partial_u \partial_i g_{ui} \right) \nu^j dS,\tag{7}$$

exists and is independent of the sequence of $\{S_k\}$ used to compute it.

Proof. From the decaying conditions and the familiar expression

$$R_g = \partial_i \partial_j g_{ij} - \partial_j \partial_j g_{ii} + O_2((g - \delta)\partial^2 g) + O_2((\partial g)^2),$$

= $\partial_i \partial_j g_{ij} - \partial_j \partial_j g_{ii} + O_2(r^{-2\tau-2})$

we get

$$\begin{split} \Delta_{g}R_{g} &= g^{ab}\nabla_{a}\nabla_{b}R_{g} = g^{ab}\left(\partial_{a}\partial_{b}R_{g} + \Gamma_{ab}^{c}\partial_{c}R_{g}\right), \\ &= g^{ab}\partial_{a}\partial_{b}\left(\partial_{i}\partial_{j}g_{ij} - \partial_{j}\partial_{j}g_{ii}\right) + g^{ab}\partial_{a}\partial_{b}\left(O_{2}(r^{-2\tau-2})\right) + g^{ab}\Gamma_{ab}^{c}\partial_{c}\left(\partial_{i}\partial_{j}g_{ij} - \partial_{j}\partial_{j}g_{jj}\right) \\ &+ g^{ab}\Gamma_{ab}^{c}\partial_{c}\left(O_{2}\left(r^{-2\tau-2}\right)\right), \\ &= g^{ab}\partial_{a}\partial_{b}\left(\partial_{i}\partial_{j}g_{ij} - \partial_{j}\partial_{j}g_{ii}\right) + \left(g^{ab} - \delta^{ab}\right)O(r^{-2\tau-4}) + O(r^{-2\tau-4}) \\ &+ O_{3}\left((g - \delta)\partial g\right))O_{1}(r^{-\tau-3}) + O_{3}(\partial g)O_{1}(r^{-\tau-3}) + O_{3}\left((g - \delta)\partial g\right))O_{1}(r^{-2\tau-3}) \\ &+ O_{3}(\partial g)O_{1}(r^{-2\tau-3}), \\ &= \partial_{a}\partial_{a}\left(\partial_{i}\partial_{j}g_{ij} - \partial_{j}\partial_{j}g_{ii}\right) + O(r^{-2\tau-4}). \end{split}$$

In particular, denoting by D_k the annular region between S_{k+1} and S_k , we get that

$$\int_{D_k} \partial_j \partial_j \left(\partial_u \partial_i g_{ui} - \partial_u \partial_u g_{ii} \right) dV_e = \int_{S_{k+1}} (\partial_j \partial_u \partial_i g_{ui} - \partial_j \partial_u \partial_u g_{ii}) \nu_j^e dS^{n-1} - \int_{S_k} (\partial_j \partial_u \partial_i g_{ui} - \partial_j \partial_u \partial_u g_{ii}) \nu_j^e dS^{n-1},$$

where, following our convention, ν^e denotes the outward pointing Euclidean unit normal to each hypersurface. Therefore, if the left-hand side is integrable over $\mathbb{R}^n \setminus K$, then the above boundary integrals form a Cauchy sequence as $r \to \infty$ and therefore (7) is well defined. But from the above computations, we see that this can be reduced to $\Delta_g R_g \in L^1(M)$ and $2(\tau + 2) > n$, that is $\tau > \tau_n$. In particular, notice that under our decaying conditions R_g , $\operatorname{Ric}_g = O(r^{-\tau-2})$ near infinity, implying that

$$Q_g = -\frac{1}{2(n-1)}\Delta_g R_g + O(r^{-2\tau-4}),$$

therefore the leading order is carried on the first term and $\Delta_g R_g \in L^1(M)$ can be replaced by $Q_g \in L^1(M)$.

Now, we intend to show that (4) is a geometric object, independent of the structure of infinity we use. With this in mind, let us start by rewriting (4) in more geometric fashion, which will be proved to be more useful for its analysis.

Proposition 2. Let (M, g) be an AE manifold which satisfies the decaying conditions i) and ii) of Proposition 1. Then, we can rewrite the energy (4) as

$$\mathcal{E}(g) = -\lim_{r \to \infty} \int_{S_1^{n-1}} \partial_r R_g r^{n-1} d\omega_{n-1}.$$
(8)

Proof. Let us denote by ν the outward g-unit normal to $S_r^{n-1} \hookrightarrow \mathbb{R}^n$, with r sufficiently large, and by ν^e the outward euclidean unit normal to the same sphere. Then, it follows that $|\nu^e - \nu|_g = O(r^{-\tau})$. Thus, we find that

$$g(\nabla R_g, \nu) = g(\nabla R_g, \nu^e) + g(\nabla R_g, \nu - \nu^e),$$

= $\partial_i R_g \nu_i^e + \underbrace{(g - \delta)_{ij} \nabla^i R_g(\nu^e)^j}_{O_1(r^{-2\tau - 3})} + \underbrace{\partial_i R_g(\nu - \nu^e)_i}_{O_1(r^{-2\tau - 3})} + \underbrace{(g - \delta)_{ij} \nabla^i R_g(\nu - \nu^e)^j}_{O_1(r^{-3\tau - 3})}.$

Also, we already know that $R_g = \partial_i \partial_j g_{ij} - \partial_j \partial_j g_{ii} + O_2(r^{-2\tau-2})$, which implies

$$g(\nabla R_g, \nu) = \partial_i \left(\partial_k \partial_j g_{kj} - \partial_j \partial_j g_{kk} \right) \frac{x^i}{r} + O_1(r^{-2\tau-3})$$

Thus,

$$\int_{S_1^{n-1}} g(\nabla R_g, \nu) r^{n-1} d\omega_{n-1} = \int_{S_1^{n-1}} \partial_i \left(\partial_k \partial_j g_{kj} - \partial_j \partial_j g_{kk} \right) \frac{x^i}{r} r^{n-1} d\omega_{n-1} + O_1(r^{n-2\tau-4}).$$

Now, from our decaying condition $\tau > \tau_n$, we see that $n - 2\tau - 4 < 0$, which implies that in the limit $r \to \infty$ the last term vanishes. Therefore, under these conditions

$$\lim_{r \to \infty} \int_{S_1^{n-1}} g(\nabla R_g, \nu) r^{n-1} d\omega_{n-1} = \lim_{r \to \infty} \int_{S_1^{n-1}} \partial_i \left(\partial_k \partial_j g_{kj} - \partial_j \partial_j g_{kk} \right) \nu_i^e r^{n-1} d\omega_{n-1}.$$

That is,

$$\mathcal{E}(g) = -\lim_{r \to \infty} \int_{S_1^{n-1}} g(\nabla R_g, \nu) r^{n-1} d\omega_{n-1},$$

$$= -\lim_{r \to \infty} \int_{S_1^{n-1}} \nabla R_g \cdot \nu^e r^{n-1} d\omega_{n-1},$$

$$= -\lim_{r \to \infty} \int_{S_1^{n-1}} \partial_r R_g r^{n-1} d\omega_{n-1}.$$
(9)

With the aid of the above two results, we can establish the following theorem, which establishes that $\mathcal{E}(g)$ is a geometric object within a suitable class of AE manifolds.

Theorem 1. Let (ϕ, x) and (ψ, y) be two structures of infinity for the AE manifold (M, g)which satisfies the conditions i) and ii) of the above Proposition with decay rates τ_1, τ_2 respectively, satisfying $\tau \doteq \min\{\tau_1, \tau_2\} > \tau_n$. Then, the energies $\mathcal{E}^{(\phi)}(g)$ and $\mathcal{E}^{(\psi)}(g)$ are well-defined and equal. Moreover, the following coordinate independent identity follows:

$$\mathcal{E}(g) = -\int_M \Delta_g R_g dV_g. \tag{10}$$

Proof. Appealing to Proposition 1, let us compute the energy $\mathcal{E}^{\phi}(g)$, associated to the structure of infinity given by (ϕ, x) using a sequence of spheres near infinity of radii $\{r_k\}_{k=1}^{\infty}$. From the above proposition, we have that

$$\mathcal{E}^{(\phi)}(g) = -\lim_{|x| \to \infty} \int_{S_1^{n-1}} \nabla R_g \lrcorner dV_e,$$

where $dV_e = r^{n-1}dr \wedge d\omega_{n-1}$ stands for the canonical Euclidean volume form. Appealing to the decaying conditions of g, we know that $\sqrt{\det(g)} = 1 + O(r^{-\tau})$ and therefore

$$\int_{S_1^{n-1}} \nabla R_g \lrcorner dV_g = \int_{S_1^{n-1}} \nabla R_g \lrcorner dV_e + O(r^{n-2\tau-4}).$$

Since $n - 2\tau - 4 < 0$ under our hypotheses, the above implies that

$$\mathcal{E}^{(\phi)}(g) = -\lim_{r \to \infty} \int_{S_r^{n-1}} \nabla R_g \, \lrcorner \, dV_g.$$

Finally, since $\operatorname{div}_g(\nabla R_g)dV_g = d(\nabla R_g \lrcorner dV_g)$, we find that

$$\mathcal{E}^{(\phi)}(g) = -\lim_{r \to \infty} \int_{D_r} \Delta_g R_g dV_g = -\int_M \Delta_g R_g dV_g,$$

where $D_r \hookrightarrow M$ is the inner region in M with $\partial D_r = S_r^{n-1}$. In particular, the right-hand side of the above expression is independent of the coordinates used near infinity. Thus we can drop the reference to (ϕ, x) and claim that for any structure of infinity of M were gsatisfies conditions (i)-(ii) of Proposition 1, it holds that

$$\mathcal{E}(g) = -\int_M \Delta_g R_g dV_g.$$

Remark 1. Let us highlight that the form for \mathcal{E} given by (10) allows us to make contact with the ideas of [35], where the author introduced a general method to produce asymptotic charges from a Riemannian functional on a complete manifold with a model asymptotic structure. In particular, (10) is an example of such a functional. Let us nevertheless stress that the definition of energy we are interested in is the one (a priori) defined by equation (4), which is the quantity appearing as a conserved charge for the fourth order gravitational theories described in the introduction. In this sense, the connection between (4) and (10) is only revealed after the work done above through Propositions 1, 2 and Theorem 1, after which the conditions necessary to achieve asymptotic invariance have already been shown explicitly. Such conditions correspond to a somewhat simpler and explicit formulation of the corresponding abstract conditions presented in [35], which must be checked in a case by case basis.

Besides establishing sufficient conditions for the energy to be a well-defined intrinsic geometric object, the above theorem provides us with an easy positive energy corollary.

Corollary 1. Let (M^n, g) be an n-dimensional AE manifold, with $n \ge 3$, which satisfies the decaying conditions i) and ii) of Proposition 1 and such that $\Delta_g R_g \le 0$. Then, the fourth order energy $\mathcal{E}(g)$ is non-negative and $\mathcal{E}(g) = 0$ if and only if (M, g) is scalar flat.

The positivity statement in the above corollary is self-evident. On the other hand, the rigidity statement follows from the injectivity of the Laplacian under our hypotheses (see (A.1)).

Let us notice that the kind of rigidity that we get from the above corollary is quite weak. In fact, although it is well-known that topological obstructions exist, the set of AE manifolds with zero scalar curvature is in general quite large (see, for instance, the discussion after Theorem 2). Therefore, in order to get a stronger rigidity statement, we will need to impose so other geometric condition. We will explore this in the next proposition and in the main theorem given below. The following proposition, which is an adaptation of a result in [20] will be used to insure that $R_g > 0$ as long as g is not flat.

Proposition 3. Let (M^n, g) be an AE Riemannian manifold with $n \ge 3$ satisfying (1) $Q_g \ge 0$ and (2) $R_g \ge 0$. Then, either $R_g > 0$ or g must be flat.

Proof. If $Q_g \ge 0$, then we find that

$$\Delta_g R_g - c_1(n) R_g^2 \le -c_2(n) |\operatorname{Ric}_g|_g^2 \le 0,$$

for some constants $c_1(n), c_2(n) > 0$. From the the condition $R_g \ge 0$ we can appeal to Lemma 4 in [33] to conclude that if R_g vanishes at a single point, then it must be identically zero. Thus, we already see that either $R_g > 0$ or $R_g \equiv 0$. In the second case, we find that $0 \le Q_g = -c_2(n) |\operatorname{Ric}_g|_g^2 \le 0$, and thus $\operatorname{Ric}_g \equiv 0$, which implies (from asymptotic flatness) that g is flat. Indeed, since $\operatorname{Ric}_g \ge 0$, then by Bishop-Gromov theorem insures that $r \mapsto \frac{vol(B_g(p,r))}{\omega_r r^n}$ is non-increasing, hence it must be constant by asymptotic flatness, finally the equality case of Bishop-Gromov theorem implies that g must be flat. \Box

We will now present the main result of this paper. We should highlight that, in particular, the proof of the rigidity statement in the following theorem follows ideas close to the proofs of the positive mass theorems associated to the Paneitz operator of [26, 20, 21].

Theorem 2. Let (M^n, g) be an n-dimensional AE manifold, with $n \ge 3$, which satisfies the decaying conditions i) and ii) of Proposition 1 and such that $Q_g, R_g \ge 0$. Then, the fourth order energy $\mathcal{E}(g)$ is non-negative and $\mathcal{E}(g) = 0$ if and only if $Q_g \equiv 0$ and (M, g)is isometric to (\mathbb{R}^n, \cdot) .

Proof. Let us first notice that we need only work with the case $Q_g \ge 0$ and $R_g > 0$, since in the remaining case the result follows from Proposition 3. Thus, in what follows we assume that $Q_g \ge 0$ and $R_g > 0$.

Proof of positivity

Let us start by noticing that under our hypotheses we have $R_g = O_2(r^{-\tau-2})$ with $\tau > \tau_n$. Then, the energy exists and is independent of the sequence of spheres used to compute it. Furthermore, these decaying conditions imply that (8) also holds. We can rewrite

$$\int_{S_1^{n-1}} \partial_r R_g r^{n-1} d\omega_{n-1} = \frac{\partial}{\partial r} \left(r^{n-1} \int_{S_1^{n-1}} R_g d\omega_{n-1} \right) - (n-1)r^{n-2} \int_{S_1^{n-1}} R_g d\omega_{n-1} d\omega$$

Define

$$h(r) \doteq r^{n-2} \int_{S_1^{n-1}} R_g d\omega_{n-1}.$$
 (11)

Then,

$$\int_{S_1^{n-1}} \partial_r R_g r^{n-1} d\omega_{n-1} = \frac{\partial}{\partial r} \left(rh(r) \right) - (n-1)h(r).$$
(12)

By assumption we have that h > 0 and from our decreasing assumption that there is some r_0 sufficiently large, so that $R_g(x) \leq Cr^{-\tau-2}(x)$ for all $x \in M$ such that $r(x) > r_0$. In particular, this last condition implies that $h(r) \leq C\omega_{n-1}r^{-\tau+n-4}$ for all $r > r_0$. Suppose that there is some $r^* > r_0$ such that (12) is positive at r^* . Then, by continuity, there is some interval $(r^* - \epsilon, r^* + \epsilon)$ so that

$$rh'(r) > (n-2)h(r),$$

which implies

$$\log\left(\frac{h(r)}{h(r^*)}\right) > (n-2)\log\left(\frac{r}{r^*}\right), \text{ for any } r \in (r^*, r^* + \epsilon).$$

That is $h(r) > \frac{h(r^*)}{r^{*n-2}} r^{n-2}$ for all $r \in (r^*, r^* + \epsilon)$. But notice that also $h(r) \le C\omega_{n-1}r^{-\tau+n-4}$ for all such r, which shows that

$$\frac{h(r^*)}{r^{*n-2}}r^{n-2} < h(r) \le C\omega_{n-1}r^{-\tau+n-4} \text{ for any } r \in (r^*, r^* + \epsilon).$$

Since r^* and C are fixed, the above inequalities are clearly falsified for large enough r, since r^{n-2} grows faster that $r^{n-4-\tau}$. This, in turn, implies that the interval $I = [r^*, r^* + \epsilon)$ where (12) is positive cannot be extended indefinitely, since that would force us into the above inequalities being valid for arbitrary large $r > r^*$, presenting a contradiction. Therefore, this implies that there must be some ϵ_{max} where (12) is non-positive at $r^* + \epsilon_{max}$. That is, there must be some $r_1 > r^*$ where

$$r_1^{n-1} \int_{S_1^{n-1}} \partial_r R_g(r_1, \theta) d\omega_{n-1}(\theta) = \left(\frac{\partial}{\partial r} (rh(r)) - (n-1)h(r)\right)|_{r=r_1} \le 0.$$
(13)

The idea is now to repeat this procedure so as to select a sequence of spheres $\{S_{r_i}^{n-1}\}_{i=1}^{\infty}$ along which (12) is manifestly non-positive. The fact that such radii can be chosen under our hypotheses is a consequence of the above argument. Then

$$\mathcal{E}(g)_j = -r_j^{n-1} \int_{S_{r_j}} \partial_r R_g d\omega_{n-1} \ge 0 \text{ for all } j,$$

which implies

$$\mathcal{E}(g) = -\lim_{j \to \infty} \int_{S_{r_j}} \partial_r R_g r^{n-1} d\omega_{n-1} \ge 0.$$
(14)

Proof of rigidity

Now, the idea is to analyze whether $\mathcal{E}(g) = 0$ implies g-flatness. First we look for a conformal metric $\tilde{g} = u^{\frac{4}{n-2}}g$ such that $R_{\tilde{g}} \equiv 0$, that is,

$$L_g(u) = \Delta_g u - c_n R_g u = 0, \tag{15}$$

where $c_n = \frac{n-2}{4(n-1)}$. Setting $u = 1 + \phi$, this is equivalent to solve

$$\Delta_g \phi - c_n R_g \phi = c_n R_g, \tag{16}$$

We know that $R_g = O_2(r^{-\tau-2})$, this implies that $R_g \in L^p_{-\delta-2}$ for any $\delta < \tau$ and any p. Thanks to corollary A.1, we know that $L_g : W^{2,p}_{\rho} \mapsto L^p_{\rho-2}$ is an isomorphism for $2 - n < \rho < 0$. Thus, choosing p > n/2, we find a (smooth) solution $\phi \in W^{2,p}_{-\delta}$ for

any $0 < \delta < \sigma \doteq \min\{\tau, n-2\}$.³. By our choice of p, ϕ goes to zero at infinity, then, the fact that $R_g > 0$ and the maximum principle, see lemma 2 and 4 [33], insure that $u = 1 + \phi > 0$.

I) $n \neq 4$

Then, let $\Phi \doteq u^{-\frac{n-4}{n-2}}$; rewrite $g = \Phi^{\frac{4}{n-4}}\tilde{g}$ and notice that transformation rule for the Paneitz operator gives us

$$\frac{n-4}{2}\Phi^{\frac{n+4}{n-4}}Q_g = P_{\tilde{g}}\Phi.$$

Also, since $R_{\tilde{g}} = 0$, we have that $Q_{\tilde{g}} = -\frac{2}{(n-2)^2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2$, and thus

$$P_{\tilde{g}}\Phi = \Delta_{\tilde{g}}^2\Phi + \operatorname{div}_{\tilde{g}}\left(4S_{\tilde{g}}(\nabla\Phi, \cdot)\right) - \frac{(n-4)}{(n-2)^2}|\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2\Phi,\tag{17}$$

where, $S_{\tilde{g}} = \frac{1}{n-2} \operatorname{Ric}_{\tilde{g}}$ since $R_{\tilde{g}} = 0$. Therefore, denoting by D_r the bounded region in M whose boundary is given by the sphere S_r in the end of M, we see that

$$\int_{D_r} \frac{n-4}{2} \Phi^{\frac{n+4}{n-4}} Q_g + \frac{(n-4)}{(n-2)^2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \Phi \, dv_{\tilde{g}} = \int_{S_r} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu}) d\omega_{\tilde{g}} + \frac{4}{n-2} \int_{S_r} \operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu}) d\omega_{\tilde{g}}$$
(18)

Let us now estimate the terms in the right-hand side.

Estimates on $\operatorname{Ric}_{\tilde{q}}(\nabla \Phi, \tilde{\nu})$

From the above analysis we find that $u = 1 + O_4(r^{-\delta})$ for any $\delta < \sigma = \min\{\tau, n-2\}$, and therefore

$$\tilde{g}_{ij} = (1 + O_4(r^{-\delta}))(\delta_{ij} + O_4(r^{-\tau})), = \delta_{ij} + O_4(r^{-\delta}).$$

Then, $\operatorname{Ric}_{\tilde{g}} = O_2(r^{-\delta-2})$ and also

$$\nabla \Phi = -\frac{n-4}{n-2}u^{-\frac{n-4}{n-2}-1}\nabla u = O_3(r^{-\delta-1}).$$

Which implies

$$\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu}) = \operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\nu^{e}) + \operatorname{Ric}_{\tilde{g}_{ij}}\nabla^{i}\Phi(\nu^{e}-\tilde{\nu})^{j} = O_{2}(r^{-2\delta-3}),$$
$$\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu})r^{n-1} = O_{2}(r^{n-2\delta-4}).$$

Since $\sigma > \frac{n}{2} - 2$, then, we can choose δ such that

$$|\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu})|r^{n-1}\to 0 \text{ as } r\to +\infty.$$

Estimates on $\Delta_{\tilde{q}} \Phi$

Let us rewrite

$$\tilde{\nabla}_i \Phi = -\frac{n-4}{n-2} u^{-2\frac{n-3}{n-2}} \tilde{\nabla}_i u,$$
$$\tilde{\nabla}_j \tilde{\nabla}_i \Phi = -\frac{n-4}{n-2} \left(u^{-2\frac{n-3}{n-2}} \tilde{\nabla}_j \tilde{\nabla}_i u - 2\frac{n-3}{n-2} u^{-\frac{3n-8}{n-2}} \tilde{\nabla}_j u \tilde{\nabla}_i u \right)$$

³In the case $\tau > n-2$, we can achieve $\delta = \sigma$.

from which we get

$$\Delta_{\tilde{g}}\Phi = -\frac{n-4}{n-2}u^{-2\frac{n-3}{n-2}}\Delta_{\tilde{g}}u + 2\frac{(n-4)(n-3)}{(n-2)^2}u^{-\frac{3n-8}{n-2}}|\tilde{\nabla}u|_{\tilde{g}}^2$$

Now, we can compute that

$$\Delta_{\tilde{g}}u = \tilde{g}^{ij}\tilde{\nabla}_{j}\nabla_{i}u = \tilde{g}^{ij}\left(\nabla_{j}\nabla_{i}u - (\Gamma_{ij}^{k}(\tilde{g}) - \Gamma_{ij}^{k}(g))\nabla_{k}u\right),$$

$$= \Delta_{g}u + (\tilde{g}^{ij} - g^{ij})\nabla_{j}\nabla_{i}u - \tilde{g}^{ij}(\Gamma_{ij}^{k}(\tilde{g}) - \Gamma_{ij}^{k}(g))\nabla_{k}u$$

and we can estimate $\tilde{g}^{ij} - g^{ij} = O_4(r^{-\delta}) - O_4(r^{-\tau}) = O_4(r^{-\delta})$ and $\Gamma_{ij}^k(\tilde{g}) - \Gamma_{ij}^k(g) = O_3(r^{-\delta-1}) - O_3(r^{-\tau-1}) = O_3(r^{-\delta-1})$, since $\sigma = \min\{\tau, n-2\}$, and also $\nabla u = O_3(r^{-\delta-1})$. Therefore

$$\tilde{g}^{ij}(\Gamma_{ij}^{k}(\tilde{g}) - \Gamma_{ij}^{k}(g))\nabla_{k}u = O_{3}(r^{-2\delta-2}), (\tilde{g}^{ij} - g^{ij})\nabla_{j}\nabla_{i}u = O_{2}(r^{-2\delta-2}).$$
(19)

On the other hand, from (15) we find that⁴

$$\Delta_g u = c_n R_g + c_n \underbrace{R_g}_{O_2(r^{-\delta-2})} \underbrace{\phi}_{O_4(r^{-\delta})} = c_n R_g + O_2(r^{-2\delta-2}).$$
(20)

Putting together (19)-(20), we finally find that

$$\Delta_{\tilde{g}}u = c_n R_g + O_2(r^{-2\delta-2}), \qquad (21)$$

which, in turn, implies

$$\Delta_{\tilde{g}} \Phi = -\frac{n-4}{4(n-1)} u^{-2\frac{n-3}{n-2}} R_g + O_2(r^{-2\delta-2}),$$

$$\nabla(\Delta_{\tilde{g}} \Phi) = -\frac{n-4}{4(n-1)} u^{-2\frac{n-3}{n-2}} \nabla R_g + O_1(r^{-2\delta-3}).$$
(22)

Therefore

$$\begin{split} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu}) &= \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\nu^{e}) + \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu}-\nu^{e}), \\ &= \nabla_{i}\Delta_{\tilde{g}}\Phi\frac{x^{i}}{r} + \underbrace{(\tilde{g}_{ij}-\delta_{ij})\nabla^{i}(\Delta_{\tilde{g}}\Phi)}_{O_{1}(r^{-2\delta-3})} \underbrace{\frac{x^{j}}{r}}_{O_{1}(r^{-2\delta-3})} + \underbrace{\tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu}-\nu^{e})}_{O_{1}(r^{-2\delta-3})}, \\ &= -\frac{n-4}{4(n-1)}u^{-2\frac{n-3}{n-2}}\nabla_{i}R_{g}\frac{x^{i}}{r} + O_{1}(r^{-2\delta-3}), \end{split}$$

implying

$$\tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu})r^{n-1} = -\frac{n-4}{4(n-1)}\underbrace{\underbrace{u^{-2\frac{n-3}{n-2}}}_{r\to\infty}\partial_r R_g r^{n-1} + \underbrace{O_1(r^{n-2\delta-4})}_{r\to\infty},$$

⁴Notice that, actually, $R_g = O_2(r^{-\tau-2})$ which is stronger than they the decay we are using, but for our purposes keeping just one weight parameter given by δ is enough to achieve the required estimates.

where we have used, as above, that $\delta > \frac{n}{2} - 2$, implying we can choose $\delta > 0$ such that $n - 2\delta - 4 < 0$. From all this and $d\omega_{\tilde{g}} = (1 + O(r^{-\delta}))d\omega_r$, we find that

$$\lim_{r \to \infty} \int_{S_r^{n-1}} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi, \tilde{\nu}) d\omega_r = -\frac{n-4}{4(n-1)} \lim_{r \to \infty} \left(\int_{S_1^{n-1}} \partial_r R_g r^{n-1} d\omega_{n-1} + \int_{S_1^{n-1}} \underbrace{\partial_r R_g O(r^{-\delta}) r^{n-1}}_{O(r^{n-2\delta-4})} d\omega_{n-1} + \underbrace{\int_{S_1^{n-1}} \underbrace{\partial_r R_g O(r^{-\delta}) r^{n-1}}_{O(r^{n-2\delta-4})} + \underbrace{\int_{S_1^{n-1}} \underbrace{\partial_r R_g O(r^{-\delta}) r^{n-1}}_{O(r^{n-2\delta-4})} + \underbrace{\int_{S_1^{n-1}} \underbrace{\partial_r R_g O(r^{-\delta}) r^{n-1}}_{O(r^{n-2\delta-4})} + \underbrace{\int_{S_1^{n-2}} \underbrace{\partial_r R_g O(r$$

Putting together the above analysis, from (18), we find that

$$\int_{M} \frac{n-4}{2} \Phi^{\frac{n+4}{n-4}} Q_g + \frac{(n-4)}{(n-2)^2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \Phi \, dv_{\tilde{g}} = \frac{n-4}{4(n-1)} \mathcal{E}(g).$$
(23)

Thus, if $\mathcal{E}(g) = 0$, then $Q_g \equiv 0$ and $\operatorname{Ric}_{\tilde{g}} \equiv 0$, which implies, through results such as those exposed in the proof of Proposition 3 that \tilde{g} is flat and $M \cong \mathbb{R}^n$. Therefore, since $\tilde{g} = u^{\frac{4}{n-2}}g$, then g is conformally-flat.

II)
$$n = 4$$

Now, fix $\Phi = -\ln(u)$ and rewrite $\tilde{g} = e^{-2\Phi}g$. Then, using the transformation rule (B.7) we find

$$0 \le Q_g = e^{-4\Phi} (P_{\tilde{g}} \Phi + Q_{\tilde{g}}).$$
(24)

Since $R_{\tilde{g}} = 0$, then $Q_{\tilde{g}} = -\frac{1}{2} |\text{Ric}_{\tilde{g}}|_{\tilde{g}}^2$, which implies

$$0 \le \int_{D_r} \left(Q_g e^{4\Phi} + \frac{1}{2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \right) dV_{\tilde{g}} = \int_{S_r} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi, \tilde{\nu}) d\omega_{\tilde{g}} + 2 \int_{S_r} \operatorname{Ric}_{\tilde{g}}(\nabla\Phi, \tilde{\nu}) d\omega_{\tilde{g}}$$
(25)

Estimates on $\operatorname{Ric}_{\tilde{q}}(\nabla \Phi, \tilde{\nu})$

We clearly have $u = 1 + O_4(r^{-\delta})$ for some $\delta < \sigma$. Then, from $\nabla \Phi = -\frac{1}{u} \nabla u = O_3(r^{-\delta-1})$, we find that

$$\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu})r^{3} = \underbrace{\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\nu_{e})}_{O_{3}(r^{-2\delta-3})}r^{3} + \underbrace{\operatorname{Ric}_{\tilde{g}}(\nabla\Phi,\tilde{\nu}-\nu_{e})}_{O_{3}(r^{-3\delta-3})}r^{3} = O_{3}(r^{-2\delta}).$$
(26)

Combining this with $d\omega_{\tilde{g}} = (1 + O(r^{-\delta}))d\omega_r$ we get

$$\int_{S_r} \operatorname{Ric}_{\tilde{g}}(\nabla \Phi, \tilde{\nu}) d\omega_{\tilde{g}} = o(1).$$
(27)

Estimates on $\tilde{g}(\nabla \Delta_{\tilde{q}} \Phi, \tilde{\nu})$

Straightforwardly we see that

$$\tilde{\nabla}_{j}\tilde{\nabla}_{i}\Phi = -u^{-1}\tilde{\nabla}_{j}\tilde{\nabla}_{i}u + u^{-2}\tilde{\nabla}_{j}u\tilde{\nabla}_{i}u = -u^{-1}\tilde{\nabla}_{j}\tilde{\nabla}_{i}u + O_{3}(r^{-2\delta-2}),$$

$$\tilde{\nabla}_{j}\tilde{\nabla}_{i}u = \nabla_{j}\nabla_{i}u - \Delta\Gamma_{ji}^{k}\nabla_{k}u = \nabla_{j}\nabla_{i}u + O_{3}(r^{-2\delta-2}),$$
(28)

where $\Delta \Gamma_{ji}^k = \Gamma_{ij}^k(\tilde{g}) - \Gamma_{ij}^k(g)$. Thus, we see that

$$\begin{split} \Delta_{\tilde{g}} \Phi &= -u^{-1} \Delta_g u - u^{-1} \underbrace{(\tilde{g}^{ij} - g^{ij}) \nabla_j \nabla_i u}_{O_3(r^{-2\delta - 2})} + O_3(r^{-2\delta - 2}) = -u^{-1} \Delta_g u + O_3(r^{-2\delta - 2}), \\ &= -c_n R_g + O_3(r^{-2\delta - 2}), \end{split}$$

thus,

$$\nabla \Delta_{\tilde{g}} \Phi = -c_n \nabla R_g + O_2(r^{-2\delta-3}),$$

$$\tilde{g}(\nabla \Delta_{\tilde{g}} \Phi, \tilde{\nu}) = -c_n \tilde{g}(\nabla R_g, \nu_e) - c_n \underbrace{\tilde{g}(\nabla R_g, \tilde{\nu} - \nu_e)}_{O_1(r^{-2\delta-3})} + O_2(r^{-2\delta-3})$$

implying

$$\begin{split} \int_{S_r} \tilde{g}(\nabla \Delta_{\tilde{g}} \Phi, \tilde{\nu}) d\omega_{\tilde{g}} &= -c_n \int_{S_r} \tilde{g}(\nabla R_g, \nu_e) d\omega_{\tilde{g}} + O(r^{-2\delta}), \\ &= -c_n \int_{S_r} \tilde{g}(\nabla R_g, \nu_e) d\omega_r - \int_{S_r} \underbrace{\tilde{g}(\nabla R_g, \nu_e)}_{O(r^{-\tau-3})} O(r^{-\delta}) d\omega_r + O(r^{-2\delta}), \\ &= -c_n \int_{S_r} \tilde{g}(\nabla R_g, \nu_e) d\omega_r + o(1) \\ &= -c_n \int_{S_r} g(\nabla R_g, \nu_e) d\omega_r - c_n \int_{S_r} \underbrace{(\underline{u}^2 - 1)}_{o(1)} \underbrace{g(\nabla R_g, \nu_e)}_{O(r^{-\delta-3})} d\omega_r + o(1) \\ &= -c_n \int_{S_r} g(\nabla R_g, \nu_e) d\omega_r + o(1). \end{split}$$

Therefore, we find that

$$\lim_{r \to \infty} \int_{S_r} \tilde{g}(\nabla \Delta_{\tilde{g}} \Phi, \tilde{\nu}) d\omega_{\tilde{g}} = c_n \mathcal{E}(g).$$
⁽²⁹⁾

Finally, putting together (25),(27) and (29), we find that

$$0 \le \int_M \left(Q_g e^{4\Phi} + \frac{1}{2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \right) dV_{\tilde{g}} = c_n \mathcal{E}(g), \tag{30}$$

which implies that if $\mathcal{E}(g) = 0$, then $\operatorname{Ric}_{\tilde{g}} \equiv 0$ and therefore g is conformally-flat with $Q_g \equiv 0$.

Finally, $g = v^{\frac{4}{n-4}}\delta$ (or $g = e^{2v}\delta$ if n = 4), with $\Delta^2 v = 0$ and $\lim_{r \to \infty} v = 1$ (or $\lim_{r \to \infty} v = 0$), hence, by maximum principle, we get that $v \equiv 1$ (or $v \equiv 0$) which achieves the proof of the theorem.

It is worth noticing that, looking more carefully to the above proof, we can weaken the hypotheses of the previous theorem and still get interesting results. Indeed, in the positivity part, we only use the fact that the scalar curvature is positive at infinity, while for the rigidity statement, we only use the fact that we can make the scalar curvature flat via a conformal transformation, which is ensured by assuming that Y([g]) > 0 by theorem 5.1 of [14], and that Q_g is non-negative. Furthermore, under this last condition, equations (23) and (30) also provide proofs of positivity. All these observations give us the following results.

Theorem 3. Let (M^n, g) be an n-dimensional AE manifold, with $n \ge 3$, which satisfies the decaying conditions i) and ii) of Proposition 1 and such that $R_g > 0$ on $M \setminus K$, for some compact set K then the fourth order energy $\mathcal{E}(g) \ge 0$. **Theorem 4.** Let (M^n, g) be an n-dimensional AE manifold, with $n \ge 3$, which satisfies the decaying conditions i) and ii) of Proposition 1 and such that $Q_g \ge 0$ and Y([g]) > 0, then $\mathcal{E}(g) \ge 0$ with equality holding if and only if (M, g) is isometric to (\mathbb{R}^n, \cdot) .

Let us now notice that the Q-curvature assumption in Theorem 4 cannot be weakened while keeping the rest of the hypotheses. This can be seen as follows: Consider (\mathbb{R}^n, δ) , with $n \geq 3$, and let $\{h_k\}_{k=1}^{\infty} \in C^{\infty}(U)$, be a sequence of smooth compactly supported symmetric second rank tensor fields, $\bar{U} \subset M$, with $||h_k|| < \epsilon$, for some fixed small $\epsilon > 0$, such that $h_k \xrightarrow[k\to\infty]{} 0$. Then, consider the sequence of metrics $g_k = \delta + h_k$, where the construction of the h_k can be made so that g_k are not conformally flat. Furthermore, this construction can be fit so that $\{g_k\}_{k=1}^{\infty}$ are all Yamabe positive.⁵ Therefore, there are conformal factors $\{u_k = 1 + \varphi_k\}$, with $\varphi_k \in H_{-\delta}^s$, with $\delta < n-2$ arbitrary, such that $\Delta_{g_k} u_k - R_{g_k} u_k = 0$.⁶ This means that $R_{u_k^{\frac{4}{n-2}}g_k} = 0$ and thus $Q_{u_k^{\frac{4}{n-2}}g_k} = -c_2(n)|\operatorname{Ric}_{u_k^{\frac{4}{n-2}}g_k}|_{u_k^{\frac{4}{n-2}}g_k}^2 < 0$. Furthermore, all the metrics $u_k^{\frac{4}{n-2}}g_k$ satisfy the decaying conditions of Proposition 1 and are, by construction, Yamabe positive. Nevertheless, from their asymptotics $u_k^{\frac{4}{n-2}}g_{k_{ij}} = (1 + \frac{a_{k,n}}{r^{n-2}})\delta_{ij} + o_k(r^{-(n-2)})$ we see that $\mathcal{E}(g_k) = 0$ for all k, although none of the g_k are conformally flat. Finally, by considering k sufficiently large, we can make $-\frac{1}{k} < Q_{u_k^{\frac{4}{n-2}}g_k} < 0$, which shows that there are Yamabe positive AE metrics satisfying conditions (i)-(ii) of Proposition 1 which have negative Q-curvature which is as small as we want, zero energy and are not conformally flat.

In dimensions n = 3, 4, under the above hypotheses, we fall into a curious situation, since any such *n*-dimensional AE-manifold has $\mathcal{E}(g) = 0$ as a consequence of the fall-off conditions, since $\nabla R_g = 0(r^{-\tau-3})$ decreases faster than the volume of the S_r^{n-1} .

Corollary 2. Any n-dimensional AE-Riemannian manifold (M^n, g) , with $n \in \{3, 4\}$, such that $Q_g \ge 0$ and Y([g]) > 0, is isometric to (\mathbb{R}^n, δ) .

Rigidity in critical cases

The aim of this section is to comment further on the especial case that occurs in dimension four. In particular, the fact that the energy $\mathcal{E}(g)$ is always zero has some topological consequences as a corollary. It is worth to contrast this with the second order case associated to the ADM energy. In this last case, the critical case is given in dimension two, where asymptotic flatness is too strong a condition in order to detect any meaningful information concerning the ADM energy (for details, see Chapter 3 in [27]). Let us start by briefly commenting on this case as a warm-up.

Let (M, g) be an asymptotically flat surface with $\tau > 0$ and non-negative integrable Gaussian curvature. Let us consider a chart a infinity, and recall the definition of the ADM energy

$$m(g) = \lim_{R \to +\infty} \int_{\partial B_R} (\partial_i g_{ij} - \partial_j g_{ii}) \nu^j \, ds,$$

$$u_k = 1 + \frac{a_k}{r^{n-2}} + o(r^{-(n-2)}),$$

where a_k are constants.

⁵For an explicit construction of this see Example 1 in [2]

⁶In fact, from Theorem 1.17 in [4], it follows that

where ν is the outer normal. Because of the decreasing assumptions, we easily see that the mass must vanish. Let us write the Gauss-Bonnet formula which gives

$$2\chi(M) = \int_{M \setminus B(0,R)^c} K \, d\sigma + \int_{\partial B(0,R)} k_g \, ds.$$

Here we write $\chi(M)$ instead of $\chi(M \setminus B(0, R)^c)$ since they are equal for R large. Then in polar coordinate we have,

$$\begin{aligned} k_g &= -\frac{1}{r^2} g\left(\nabla_{\partial_\theta} \partial_\theta, \partial_r \right) = -\frac{1}{r^2} \Gamma^r_{\theta\theta}(g) + O\left(\frac{1}{r^{1+\tau}}\right) = -\frac{1}{r^2} \Gamma^r_{\theta\theta}(\delta) + O\left(\frac{1}{r^{1+\tau}}\right) \\ &= \frac{1}{r} + O\left(\frac{1}{r^{1+\tau}}\right), \end{aligned}$$

hence passing to the limit we obtain that

$$2\pi\chi(M) - 2\pi = \int_M K \, d\sigma.$$

But we know that

$$\chi(M) = 2 - 2g - r,$$

where g is the genus of M and r its number of ends. Then we necessary get that g = 0, r = 1 and $K \equiv 0$. Hence we recover the following well-known proposition, see chapter 3 [27],

Proposition 4. Let (M, g) an asymptotically flat surface with non-negative Gauss curvature. Then, (M, g) is the Euclidean plane.

Let us turn to the 4-dimensional case. Let (M, g) be a 4-dimensional AE manifold with $\tau > 0$ with only one end such that $|W_g|^2$ is integrable. From Corollary 2, we know that the mass is necessarily vanishing

$$\mathcal{E}(g) = -\lim_{R \to \infty} \int_{S_R} \partial_r R_g r^3 d\omega_3 = 0.$$
(31)

Along the lines of the 2-dimensional case, let us try to get a similar proof of the rigidity result. First, we apply the Gauss-Bonnet-Chern formula with boundary (see [12] for instance)

$$32\pi^2 \chi(M) = \int_{K_R} |W|^2 + 16\sigma_2(S_g) \, dv + 8 \int_{\partial B(0,R)} B_g \, dv$$

where $K_R = \left(M \setminus \overline{B(0,R)}\right)^c$, $S_g = \frac{1}{2} \left(\operatorname{Ric} - \frac{1}{6}R_g g\right)$ is the Schouten tensor and $B_g = \frac{1}{2}R_g H - \operatorname{Ric}(\nu,\nu)H - R_{\gamma\alpha\beta\gamma}\Pi^{\alpha\beta} + \frac{1}{3}H^3 - H|\Pi|^2 + \frac{2}{3}tr(\Pi^3)$, where II and H are respectively the second form and the mean curvature of a hypersurface $\Sigma \hookrightarrow M$, $H = \operatorname{tr}_{\Sigma}\Pi$, taken with respect to the inner unit normal to Σ , the Greek indices are indices tangent to Σ , and Π^3 is defined in local coordinates by $\Pi_j^{3i} = \Pi_k^i \Pi_l^k \Pi_j^l$. Moreover, in dimension 4, we have

$$Q_g = -\frac{1}{6}\Delta_g R_g + 4\sigma_2(S_g),$$

and thanks to our decreasing assumption, we get

$$B_g = \frac{2}{r^3} + O\left(\frac{1}{r^{3+\tau}}\right)$$

which gives, using once more our decreasing assumption, that

$$32\pi^{2}(\chi(M) - 1) = \int_{K_{R}} |W|^{2} + 4Q_{g} + \frac{2}{3}\Delta_{g}R_{g} \, dv_{g} + O\left(\frac{1}{R^{\tau}}\right)$$
$$= \int_{K_{R}} |W|^{2} + 4Q_{g} \, dv_{g} + O\left(\frac{1}{R^{\tau}}\right)$$
(32)

Hence passing to the limit, we get

$$32\pi^2(\chi(M) - 1) = \int_M |W|^2 + 4Q_g \, dv_g$$

Let us set

$$\kappa_g = \int_M Q_g \, dv_g$$

One can use the above expression to recover Corollary 2 in the four dimensional case as consequence of the conformal invariance of κ_g . Also, we can easily deduce the following curvature-topology proposition. This proposition is not really new but we just would like to put them in perspective with our rigidity result

Proposition 5. Let (M,g) be an AE 4-manifold with $\kappa_g \geq 0$, then $\chi(M) \geq 1$ with equality if (M,g) is locally conformally flat with $\kappa_g = 0$. In particular, if $\kappa_g \geq 0$ and Y([g]) > 0 or the second betti number vanishes, then $\kappa_g = 0$ and (M,g) is conformal to the euclidean \mathbb{R}^4 .

3.1 The Q-curvature positive mass theorem

In this section we will make contact with a series of recent results associated with the positive mass theorem for the Paneitz operator, namely [20, 21, 26]. In particular, we will show that these results follow from Theorem 4.

Let us consider a closed manifold (M^n, g) with $n \ge 5$. If, the Panietz operator is coercive, that is to say

$$\inf_{\in H^2(M)\,;\,\|u\|_2=1}\int_M P_g(u)u\,dv_g>0,$$

then it admits a Green functions G_{P_g} . This is in particular guaranteed if $R_g \ge 0$ and Q_g is semi-positive, see proposition B of [20], or more generally by the fact that

$$Y_4([g]) = \inf_{u \in H^2(M) ; ||u||_{2^{\#}} = 1} \int_M P_g(u) u \, dv_g > 0,$$

where $2^{\#} = \frac{2n}{n-4}$. This last infimum has the advantage to be conformally invariant and it plays a similar role to the Yamabe invariant for the *Q*-curvature. Through this section, we will assume that $Y_4([g]) > 0$ and therefore G_{P_g} exists for every $g \in [g]$.

Nevertheless, contrary to the conformal Laplacian, nothing here guarantees that G_{P_g} is positive. This will be one of our assumptions, which is in particular satisfied if $Y([g]) \ge 0$ and $Q_{\tilde{g}}$ is semi-positive for some conformal metric \tilde{g} , due to lemma 3.2 in [23]. **Remark 2.** In fact Hang and Yang assume Y([g]) > 0, but if Y([g]) = 0 and there is some $\tilde{g} \in [g]$ with $Q_{\tilde{g}}$ semi-positive, then there exists $\bar{g} = u^{\frac{4}{n-2}}\tilde{g}$ such that $R_{\bar{g}} = 0$ which implies that $Q_{\bar{g}} \leq 0$ which contradicts the fact that

$$\int_M Q_{\tilde{g}} v \, dv_{\tilde{g}} = \int_M v^{\frac{n+4}{n-4}} Q_{\bar{g}} \, dv_{\tilde{g}},$$

with $v = u^{\frac{n-4}{n-2}}$. Hence the existence of a conformal metric with semi-positive Q-curvature forces the Yamabe invariant to be positive.

In fact, if $Y([g]) \ge 0$, thanks to, theorem 1.1 of [23], the existence of a positive G_{P_g} is equivalent to the existence of a conformal metric \tilde{g} with semi-positive Q-curvature.

Let now us focus on the expansion of the Green function around a singularity. If we assume that $5 \leq n \leq 7$ or g is locally conformally flat, then, in conformal normal coordinates $\{x^i\}$ for the conformal metric \tilde{g} , the Green function G_P of $P_{\tilde{g}}$ admits an expansion of the form (see Proposition 2.5 in [20])

$$G_P(p,x) = \frac{\gamma_n}{r^{n-4}} + \alpha + O_4(r), \qquad (33)$$

where $r(x) \doteq d_{\tilde{g}}(p, x)$, $\gamma_n \doteq \frac{1}{2(n-2)(n-4)\omega_{n-1}}$ and α is a constant **called the mass**.

Remark 3. As remarked in [23], under the condition $\text{Ker}(P_g) = 0$ (which is itself conformally invariant), the sign of the Green function is a conformal invariant since,

$$G_{P_{u^{\frac{4}{n-4}}g}}(p,q) = u(p)^{-1}u(q)^{-1}G_{P_g}(p,q).$$

Hence, the sign of the mass is also a conformal invariant.

This terminology arises in analogy to the Yamabe problem, where the Green function G_{L_g} of the conformal Laplacian is involved and, in important cases, it admits a similar expansion to (33). In the case of the conformal Laplacian, it was an observation of R. Schoen that the constant which appears in the place of α is precisely the ADM mass of the AE-manifold obtained by via the stereographic projection $(M \setminus \{p\}, G_L^{\frac{4}{n-2}}g)$. This showed that the resolution of the positive mass conjecture in general relativity would amount to completing the resolution of the Yamabe problem, which concerned the cases Y(g) > 0 and dimensions n = 3, 4, 5 or M locally conformally flat (see [40] and [28]).

The positive mass theorem presented in [20] as Theorem 2.9 states that under the conditions described above which provide us with the expansion (33), the mass satisfies $\alpha \geq 0$ with equality holding if and only if (M, g) is conformal to the round sphere. This came about as a generalisation of [26], where this result was proven in the conformally flat case, and, in turn, all this was generalised in [21], where the condition $R_g \geq 0$ was replaced by Y([g]) > 0. Furthermore, in Theorem 1.4 of [23], this positive mass theorem is used to solve the $Y_4([g]) > 0$ Q-curvature prescription problem in very much the same spirit as the usual positive mass theorem was used by Schoen in [40]. We intend to show that all these results are special cases of the positive energy theorem presented above.

Finally, let us notice that, for instance, in the Q-curvature prescription problem it is the sign of the mass that is actually important rather than its precise value. This becomes especially useful when it is combined with the additional observation that the sign of the mass is itself a conformal invariant, which was highlighted in Remark 3. Thus, being concerned with a conformal problem, given $(M^n, [g])$, we will always consider a choice of g satisfying the conformal normal coordinate properties (C.1), (C.3), (C.4). Notice that from the discussion presented in that appendix, this can always be achieved by first going to a related conformal metric.

In order to prepare for the main statements of this section, let us first establish a couple of preliminary results.

Proposition 6. Let (M^n, g) be a closed manifold satisfying $n \ge 5$ and whose Panietz operator admits a positive Green function G_P with an expansion as (33) around some point $p \in M$. Then, the manifold $(\hat{M} \doteq M \setminus \{p\}, \hat{g} \doteq G_P(p, \cdot)^{\frac{4}{n-4}}g)$ is an asymptotically flat manifold of order $\tau = 1$ if n = 5 and $\tau = 2$ if n > 5. Furthermore, either if $5 \le n \le 7$ or g is flat around p, then $\mathcal{E}(\hat{g}) = 8(n-1)(n-2)\omega_{n-1}\gamma_n\alpha$.

Proof. Let us fix the conformal normal coordinates $\{x^i\}$ where (33) holds. Let us appeal to the conformal normal coordinate construction of order $N \ge 4$ described Appendix B so that (C.3)-(C.4) hold, and start by considering the following expansion near p

$$g_{ij} = \delta_{ij} + \frac{1}{3}R_{iklj}(p)x^k x^l + \frac{1}{6}R_{iklj,a}(p)x^k x^l x^a + O(r^4),$$
(34)

Now, consider the inverted coordinates $z^i = \gamma_n^{\frac{2}{n-4}} \frac{x^i}{r^2}$ in a neighborhood of p and define $\rho^2 \doteq |z|^2$, so that $\rho^2 = \gamma_n^{\frac{4}{n-4}} r^{-2}$ and $x^i = \gamma_n^{\frac{2}{n-4}} \frac{z^i}{\rho^2}$. Then, it holds that $\gamma_n r^{-(n-4)} = \gamma_n^{-1} \rho^{n-4}$ and

$$\frac{\partial}{\partial z^{i}} = \gamma_{n}^{\frac{2}{n-4}} \rho^{-2} \left(\delta_{i}^{j} - 2\rho^{-2} z^{i} z^{j} \right) \frac{\partial}{\partial x^{j}}, \tag{35}$$

which, appealing to (33), implies

$$\hat{g}_{ij}(z) = \left(\gamma_n^{-1}\rho^{n-4} + \alpha + O_4(\rho^{-1})\right)^{\frac{4}{n-4}} g(\partial_{z^i}, \partial_{z^j}), \\ = \left(1 + \frac{\gamma_n \alpha}{\rho^{n-4}} + O_4(\rho^{3-n})\right)^{\frac{4}{n-4}} \left(\delta_k^i - 2\rho^{-2}z^i z^k\right) \left(\delta_l^j - 2\rho^{-2}z^j z^l\right) g_{kl}\left(\gamma_n^{\frac{2}{n-4}}\rho^{-2}z\right).$$

Invoking (34), this implies

$$\hat{g}_{ij}(z) = \left(1 + \frac{\gamma_n \alpha}{\rho^{n-4}} + O_4(\rho^{3-n})\right)^{\frac{4}{n-4}} h_{ij}(z),$$

where

$$h_{ij}(z) = \left(\delta_k^i \delta_l^j - 2\frac{z^i z^k}{\rho^2} \delta_l^j - 2\frac{z^j z^l}{\rho^2} \delta_k^i + 4\frac{z^i z^j z^k z^l}{\rho^4}\right) \left(\delta_{kl} + \frac{\gamma_n^{\frac{4}{n-4}}}{3} R_{kmnl}(p) \frac{z^m z^n}{\rho^4} + \frac{\gamma_n^{\frac{6}{n-4}}}{6} R_{kmnl,p}(p) \frac{z^m z^n z^p}{\rho^6} + O_4\left(\frac{1}{\rho^4}\right)\right),$$
$$= \delta_{ij} + H_{ij}^2(z) + H_{ij}^3(z) + O_4\left(\frac{1}{\rho^4}\right)$$

where

$$\begin{split} \gamma_n^{-\frac{4}{n-4}} H_{ij}^2 &= \frac{1}{3} R_{imnj}(p) \frac{z^m z^n}{\rho^4} - \frac{2}{3} \underbrace{R_{kmnj}(p) z^k z^m}_{=0} \frac{z^n z^i}{\rho^6} - \frac{2}{3} \underbrace{R_{imnl}(p) z^n z^l}_{=0} \frac{z^m z^j}{\rho^6} \\ &+ \frac{4}{3} \underbrace{R_{kmnl}(p) z^k z^m}_{=0} \frac{z^n z^l z^i z^j}{\rho^8}, \\ &= \frac{1}{3} R_{imnj}(p) \frac{z^m z^n}{\rho^4} \end{split}$$

and

$$\begin{split} \gamma_n^{-\frac{6}{n-4}} H_{ij}^3 &= \frac{1}{6} R_{imnj,a}(p) \frac{z^m z^n z^a}{\rho^6} - \frac{1}{3} \underbrace{R_{kmnj,a}(p) z^k z^m}_{=0} \frac{z^n z^i z^a}{\rho^8} - \frac{1}{3} \underbrace{R_{imnl,a}(p) z^n z^l}_{=0} \frac{z^m z^j z^a}{\rho^8} \\ &+ \frac{2}{3} \underbrace{R_{kmnl,a}(p) z^k z^m}_{=0} \frac{z^n z^l z^i z^j z^a}{\rho^{10}}, \\ &= \frac{1}{6} R_{imnj,a}(p) \frac{z^m z^n z^a}{\rho^6}. \end{split}$$

Putting together all the above, we find that

$$\hat{g}_{ij}(z) = \left(1 + \frac{4}{n-4} \frac{\gamma_n \alpha}{\rho^{n-4}}\right) \delta_{ij} + \frac{\gamma_n^{\frac{4}{n-4}}}{3} R_{iabj}(p) \frac{z^a z^b}{\rho^4} + \frac{\gamma_n^{\frac{6}{n-4}}}{6} R_{iabj,c}(p) \frac{z^a z^b z^c}{\rho^6} + O_4(\rho^{-(n-3)}) + O_4(\rho^{-4}),$$

which establishes the asymptotic flatness condition claimed in the proposition. Notice that if $5 \leq n \leq 7$, then, from Proposition 1, we know that $\mathcal{E}(\hat{g})$ is well-defined and the above expansion implies

$$\partial_a \hat{g}_{ij} = -4\gamma_n \alpha \frac{z^a}{\rho^{n-2}} \delta_{ij} + \partial_a H_{ij}^2 + \partial_a H_{ij}^3 + O_3 \left(\rho^{-(n-2)}\right), \\ \partial_{ba} \hat{g}_{ij} = -4\gamma_n \alpha \left(\frac{\delta_{ab}}{\rho^{n-2}} - (n-2)\frac{z^b z^a}{\rho^n}\right) \delta_{ij} + \partial_{ba} H_{ij}^2 + \partial_{ba} H_{ij}^3 + O_2 \left(\rho^{-(n-1)}\right).$$

Appealing to (C.3)-(C.4), it is not difficult to explicitly compute that $\partial_{aa}H_{ii}^2$ and $\partial_{aa}H_{ii}^3$ both vanish. This implies

$$\partial_{aa}\hat{g}_{ii} = -8n\gamma_n \alpha \rho^{-(n-2)} + O_2(\rho^{-(n-1)}).$$
(36)

Similarly, appealing to (C.3)-(C.4), it also holds that $\partial_{ij}H_{ij}^2 = 0$ and $\partial_{ij}H_{ij}^3 = 0$, and therefore we find that

$$\partial_{ij}\hat{g}_{ij} = -8\gamma_n \alpha \rho^{-(n-2)} + O_2(\rho^{-(n-1)}).$$
(37)

Thus, putting together (36)-(37), we see that

$$\left(\partial_{caa}\hat{g}_{ii}(z) - \partial_{cij}\hat{g}_{ij}\right)\frac{z^c}{\rho} = 8\gamma_n\alpha(n-1)(n-2)\rho^{-(n-1)} + O_1(\rho^{-n}),\tag{38}$$

implying that

$$\mathcal{E}(\hat{g}) = \lim_{\rho \to \infty} \int_{S_{\rho}} \left(\partial_{caa} \hat{g}_{ii}(z) - \partial_{cij} \hat{g}_{ij}(z) \right) \frac{z^c}{\rho} d\omega_{\rho} = \omega_{n-1} 8 \gamma_n \alpha (n-1)(n-2).$$
(39)

Finally, in g is flat near p, without restriction on the dimension, we know that the expansion (33) holds in rectangular coordinates around p, where $g_{ij}(x) = \delta_{ij}$. Thus, the same computations as above show that

$$\hat{g}_{ij}(z) = \left(1 + \frac{4}{n-4} \frac{\gamma_n \alpha}{\rho^{n-4}}\right) \delta_{ij} + O_4(\rho^{-(n-3)}).$$

In this case the order of decay is improved for $n \ge 7$ and in particular we know that $\mathcal{E}(\hat{g})$ is well-defined. From the above expression, the same computations that led us to (39) prove that $\mathcal{E}(\hat{g}) = 8\omega_{n-1}(n-1)(n-2)\gamma_n\alpha$.

In order to apply the positive energy theorem to establish that $\alpha \ge 0$, we first need to check that \hat{g} satisfies its hypotheses. With this in mind, consider the following proposition.

Proposition 7. Consider a closed Riemannian manifold (M^n, g) , with $n \geq 5$, which admits a conformal metric with positive Q-curvature such that $Y([g]) \geq 0$. Then, there exists a conformal metric \tilde{g} such that the asymptotically flat manifold $(\hat{M} = M \setminus \{p\}, \hat{g} = G_{P_{\tilde{g}}}^{\frac{4}{n-4}} \tilde{g})$ satisfies $Y([\hat{g}]) > 0$ and $Q_{\hat{g}} \equiv 0$.

Proof. Thanks to Remark 2, Y([g]) > 0 and $G_{P_{\tilde{g}}}$ exists and is positive for every $\tilde{g} \in [g]$. Then, thanks to proposition 6, we only have to prove that $Y([\hat{g}]) > 0$. Since Y([g]) > 0, let $G_{L_{\tilde{q}}}$ be the Green's function of the conformal Laplacian. We trivially get that $\bar{g} =$

$$\left(\frac{G_L^2}{G_P^{\frac{2(n-2)}{n-4}}}\right)^{\frac{2}{n-2}}\hat{g} \text{ is scalar flat.}$$

Equipped with the above two propositions, we can recover the following theorem, originally proved by Hang-Yang in [23], and which is an extension of results of Gursky-Malchiodi [20] and Humbert-Raulot [26].

Theorem 5. Let (M, g) be a closed n-dimensional Riemannian manifold, with $5 \le n \le 7$ or $n \ge 8$ and locally conformally flat around some point $p \in M$. If $Y([g]) \ge 0$ and (M, g)admits a conformal metric with semi-positive Q-curvature, then the mass of G_P at p is non-negative and vanishes if and only if (M, g) is conformal to the standard sphere.

It is important to note that here the hypothesis are conformally invariant. In dimension $n \ge 6$, thanks to Corollary 1.1 [19] and remark 2, they are equivalent to Y([g]) > 0 and $Y_4^*([g]) > 0$, where

$$Y_4^*([g]) = \frac{n-4}{2} \inf_{\tilde{g} \in [g], R_{\tilde{g}} > 0} \frac{\int_M Q_{\tilde{g}} \, dv_{\tilde{g}}}{\operatorname{Vol}_{\tilde{q}}(M)}$$

Of course $Y_4^*([g])$ is also conformally invariant.

Proof. Under these conditions, we know that the Green function G_P exists and is positive for every element in [g]. From Proposition 6 we also know that there is a conformal metric \tilde{g} such that the manifold $(\hat{M} = M \setminus \{p\}, \hat{g} = G_{P_{\tilde{g}}}^{\frac{4}{n-4}} \tilde{g})$ is AE, satisfies the decay assumptions of Theorem 4 and, furthermore the energy is positively proportional to the mass of $G_{P_{\hat{g}}}$. Finally, from Proposition 7, we know that $Y([\hat{g}]) > 0$ and $Q_{\hat{g}} \equiv 0$. Therefore, (\hat{M}, \hat{g}) satisfies all the hypotheses of Theorem 4, and thus the non-negativity follows directly from $\mathcal{E}(\hat{g})$ being positively proportional to α . Finally, if $\alpha = 0$, we find that \hat{M} is isometric to \mathbb{R}^n . Being M the one point compactification of \hat{M} , we see that $M \cong S^n$ and g is conformal to the round metric.

3.2 The 4-dimensional case

Finally, we would like to end by briefly discussing some peculiarities concerning the analysis of the four dimensional case. In particular, a large part of what we have done above for the Q-curvature positive mass theorem can be translated perfectly well to n = 4. Nevertheless, in view of Corollary 2 it should be clear that strong restrictions should appear in the treatment of this special case. Although the origins of some of these restriction may be evident to experts in Q-curvature analysis, the aim in what follows is to make explicit the subtle differences of this case and where the particularities arise. In this spirit, let us start by presenting the following lemma which combines results of [36]-[29].

Lemma 1. Let (M^4, g) be a closed Riemannian manifold satisfying $\text{Ker}(P_g) = \mathbb{R}$ and $\kappa_g > 0$. Then, given $p \in M$, there exists a Green's function $G_p \doteq G_{P_g}(p, \cdot) \in C^{\infty}(M \setminus \{p\})$ with a pole at p, unique up to an additive constant, which satisfies

$$P_g G_p + Q_g = \kappa_g \delta_p \tag{40}$$

as distributions. Furthermore, near p, in g-normal coordinates $\{x^i\}_{i=1}^4$ the following expansion holds

$$G_p = \frac{\kappa_g}{16\pi^2} \ln(r^{-2}) + \frac{\kappa_g}{16\pi^2} S_0 + \frac{\kappa_g}{16\pi^2} a_i x^i + \frac{\kappa_g}{16\pi^2} b_{ij} x^i x^j + o_4(r^2).$$
(41)

for some constants S_0, a_i, b_{ij} .

Proof. From Lemma 2.1 in [36] we know that if $\text{Ker}(P_g) = \mathbb{R}$, then the Paneitz operator P_g admits a Green function G_P , which is to say that for every $u \in C^4(M)$ it holds that

$$u(x) - \bar{u} = \int_{M} G_{P}(x, y) P_{g}(u(y)) dV_{g}(y), \qquad (42)$$

where $\bar{u} \doteq \operatorname{vol}_{g}(M)^{-1} \int_{M} u dV_{g}$. Also, since $\operatorname{Ker}(P_{g}) = \mathbb{R}$, we know that there is a smooth function U satisfying

$$P_g U = -(Q_g - \bar{Q}_g).$$

Define $G_p \doteq \kappa_g G_P + U$, so that

$$\langle P_g G_p, u \rangle = \kappa_g u(p) - \kappa_g \bar{u} + \langle P_g U, u \rangle, = \kappa_g u_p - \kappa_g \bar{u} - \langle Q_g, u \rangle + \langle \frac{\kappa_g}{\operatorname{vol}_g(M)}, u \rangle = \kappa_g u_p - \langle Q_g, u \rangle.$$

Thus, we see that

$$\langle P_g G_p + Q_g, u \rangle = \kappa_g u(p) \ \forall u \in C^{\infty}(M),$$
(43)

which is to say that $P_g G_p + Q_g = \kappa_g \delta_p$. The uniqueness claim follows since two different such functions G_p and \tilde{G}_p must satisfy $P_g(G_p - \tilde{G}_p) = 0$, implying $G_p = \tilde{G}_p + c$. If we rewrite (40) as $P_g\left(\frac{16\pi^2}{k_g}G_p\right) + \frac{16\pi^2}{k_g}Q_g = 16\pi^2\delta_p$ we can appeal quite straightforwardly to the computations of the Appendix in [29] to conclude that, in *g*-normal coordinates, the Green function has the following expansion near p:

$$\frac{16\pi^2}{\kappa_g}G_p = -2\ln(r) + S_0 + a_i x^i + b_{ij} x^i x^j + o_4(r^2).$$
(44)

which proves (41).

Let us now consider the same setting as in the above lemma and consider the inverted coordinates $z^i = \frac{x^i}{r^2}$ and define $(\hat{M} = M \setminus \{p\}, \hat{g} = e^{2G_p}g)$. Then,

$$Q_{\hat{g}}(q) = e^{-4G_p(q)} \left(P_g G_p(q) + Q_g(q) \right) = 0, \text{ for any } q \in \hat{M}.$$

and

$$\hat{g}_{ij}(z) = e^{\ln(\rho^{4}\frac{\kappa_{g}}{16\pi^{2}}) + \frac{\kappa_{g}}{8\pi^{2}}S_{0} + \frac{\kappa_{g}}{8\pi^{2}}a_{i}\frac{z^{i}}{\rho^{2}} + \frac{\kappa_{g}}{8\pi^{2}}b_{ij}\frac{z^{i}z^{j}}{\rho^{4}} + O_{4}(\rho^{-3})}g(\partial_{z^{i}},\partial_{z^{j}}),
= \rho^{4}\frac{\kappa_{g}}{16\pi^{2}}e^{\frac{\kappa_{g}}{8\pi^{2}}S_{0}}e^{\frac{\kappa_{g}}{8\pi^{2}}a_{i}\frac{z^{i}}{\rho^{2}} + \frac{\kappa_{g}}{8\pi^{2}}b_{ij}\frac{z^{i}z^{j}}{\rho^{4}} + O_{4}(\rho^{-3})}\rho^{-4}\left(\delta_{k}^{i} - 2\rho^{-2}z^{i}z^{k}\right)\left(\delta_{l}^{j} - 2\rho^{-2}z^{j}z^{l}\right)g_{kl}(\rho^{-2}z),
= \rho^{4\Delta\kappa_{g}}e^{\frac{\kappa_{g}}{8\pi^{2}}S_{0}}e^{\frac{\kappa_{g}}{8\pi^{2}}a_{i}\frac{z^{i}}{\rho^{2}} + \frac{\kappa_{g}}{8\pi^{2}}b_{ij}\frac{z^{i}z^{j}}{\rho^{4}} + O_{4}(\rho^{-3})}\left(\delta_{ij} + O_{4}(\rho^{-2})\right),
= \rho^{4\Delta\kappa_{g}}e^{\frac{\kappa_{g}}{8\pi^{2}}S_{0}}\left(1 + \frac{\kappa_{g}}{8\pi^{2}}a_{i}\frac{z^{i}}{\rho^{2}} + O_{4}(\rho^{-2})\right)\left(\delta_{ij} + O_{4}(\rho^{-2})\right),$$
(45)

where we have defined $\Delta \kappa_g \doteq \frac{\kappa_g}{16\pi^2} - 1$. Let us notice that the above inversion gives an AE gentric \hat{g} iff $\Delta \kappa_g = 0$, which is to say $\kappa_g = 16\pi^2$. In particular, in this case, via a coordinate change of the form $\bar{z}^i = e^{S_0} z^i$, we find that

$$\hat{g}_{ij}(\bar{z}) = \left(1 + 2e^{S_0}a_k\frac{\bar{z}^k}{\bar{\rho}^2} + O_4(\bar{\rho}^{-2})\right)\left(\delta_{ij} + O_4(\bar{\rho}^{-2})\right),\tag{46}$$

where $\bar{\rho} = |\bar{z}| = e^{S_0}\rho$. That is, $\rho^{-k} = e^{kS_0}\bar{\rho}^{-k}$. We see that (\hat{M}, \hat{g}) is an AE-manifold or order $\tau = 1$. Therefore, since by construction $Q_{\hat{g}} \equiv 0$, if Y(g) > 0, in view of Corollary 2, we must conclude that (M, g) is conformal to the round sphere. That is, the above computations together with Corollary 2 imply the following Corollary, which is not new, since it concerns the rigidity statement involved in Theorem B in [18].

Corollary 3. Let (M, g) be a closed 4-dimensional manifold which satisfies $\text{Ker}(P_g) = \mathbb{R}$ and $\kappa_g = 16\pi^2$. If Y(g) > 0, then (M, g) is conformal to the round sphere.

Finally, let us highlight that the techniques developed so far allow us to actually get more than the previous corollary. In fact, we can recover the full statement of Theorem B in [18] by an independent and simple proof.

Theorem 6 (Gursky). Let (M^4, g) a 4-dimensional manifold with $Y([g]) \ge 0$, then $\kappa_g \le 16\pi^2$ with equality holding iff (M^4, g) is conformal to the standard sphere.

Proof. Since we need only pay attention to the cases $\kappa_g > 0$, let us start by noticing that, due to Theorem A in [18], under our hypotheses $\operatorname{Ker}(P_g) = \mathbb{R}$. Furthermore, the case Y([g]) = 0 is also trivial, since in this case $\kappa_g \leq 0$, thus we will assume Y([g]) > 0. Since our hypotheses are conformally invariant, let us start assuming that g has been picked within [g] so as to be the metric of a conformal normal coordinate system. We can now appeal to Lemma 1 and the expansion (45) to construct the manifold $(\hat{M} = M \setminus \{p\}, \hat{g})$. Although in general this manifold will not be AE, it holds that $Q_{\hat{g}} \equiv 0$. Also, since Y([g]) > 0, we now that the Green function G_L of the conformal Laplacian (with a pole at $p \in M$) exists, it is positive and therefore we can construct the metric $\tilde{g} \doteq G_L^2 g$ on \hat{M} . In inverted normal coordinates around p, from [28], we know that

$$\tilde{g}(z)(\partial_{z^i},\partial_{z^j}) = \delta_{ij} + O_4(\rho^{-2}),$$

which shows that (\hat{M}, \tilde{g}) is AE of order $\tau = 2$. Furthermore, \hat{g} and \tilde{g} are related via

$$\hat{g} = e^{-\ln(G_L^2)} e^{2G_p} \tilde{g} = e^{2(G_p - \ln(G_L))} \tilde{g}$$

and therefore, from the conformal covariance associate to the Paneitz operator, we see that

$$0 = e^{4\Phi} Q_{\hat{g}} = P_{\tilde{g}} \Phi + Q_{\tilde{g}}, \tag{47}$$

where we have defined $\Phi \doteq G_P - \ln(G_L)$. Following the proof theorem 2, we know that $Q_{\tilde{g}} = -\frac{1}{2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2$, and therefore we find that, for ρ large enough,

$$\int_{D_{\rho}} \frac{1}{2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^{2} dV_{\tilde{g}} = \int_{S_{\rho}} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi, \tilde{\nu}) d\tilde{\omega} + 2 \int_{S_{\rho}} \operatorname{Ric}_{\tilde{g}}(\tilde{\nabla}\Phi, \tilde{\nu}) d\tilde{\omega}.$$
(48)

Now, the main difference with respect to theorem 2 is that we cannot estimate the derivatives of Φ in the same way, since it does not solve the same equation as in that proof. Nevertheless, in this case, we have an explicit expression for Φ , at least asymptotically. Thus, let us notice that for sufficiently large ρ the following holds

$$\Phi = \ln(\rho^{2\alpha}) + S_0 + O_4(\rho^{-1}) - \ln(\rho^2 + A + O_4(\rho^{-1})),$$

= $\ln(\rho^{2\alpha}) + S_0 - \ln(\rho^2) - \ln(1 + A\rho^{-2} + O_4(\rho^{-3})) + O_4(\rho^{-1}),$
= $\ln(\rho^{2(\alpha-1)}) + S_0 - \ln(1 + A\rho^{-2} + O_4(\rho^{-3})) + O_4(\rho^{-1}),$

where above we have defined $\alpha \doteq \frac{\kappa_g}{16\pi^2}$ and both A and S_0 are constants. From all this we can directly compute that

$$\tilde{\nabla}_{i}\Phi = 2(\alpha - 1)\frac{z^{i}}{\rho^{2}} + O_{3}(\rho^{-2}),$$
$$\Delta_{\tilde{g}}\Phi = \frac{4(\alpha - 1)}{\rho^{2}} + O_{2}(\rho^{-3}),$$
$$\tilde{\nabla}_{i}\Delta_{\tilde{g}}\Phi = -\frac{8(\alpha - 1)}{\rho^{3}}\frac{z^{i}}{\rho} + O_{2}(\rho^{-4}).$$

From the above expression, it follows that

$$\operatorname{Ric}_{\tilde{g}}(\tilde{\nabla}\Phi,\tilde{\nu}) = o(\rho^{-3}),$$
$$\tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi,\tilde{\nu}) = \frac{8(1-\alpha)}{\rho^3} + O_1(\rho^{-4})$$

implying that

$$\int_{D_{\rho}} \frac{1}{2} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 dV_{\tilde{g}} = 8\omega_3(1-\alpha) + o(1).$$
(49)

Finally, passing to the limit as r goes to infinity, we find that

$$\int_{\hat{M}} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 dV_{\tilde{g}} = 8\omega_3(1-\alpha) \ge 0.$$
(50)

This implies that $\alpha \leq 1$ and the equality case has already been dealt with in the previous corollary, which establishes the theorem.

3.3 The 3-dimensional case

In this section, we briefly explain how to recover the following theorem due to Hang and Yang [22].

Theorem 7 (Proposition 2.4 [22]). Assume the Yamabe invariant Y([g]) > 0, ker $P_g = 0$. If there is some $p \in M$ such that $G_p \doteq G_{P_g}(p, \cdot) < 0$ on $M \setminus \{p\}$, then $G_p(p) < 0$ except when (M, g) is conformally equivalent to the standard S^3 .

In this case the proof runs along the same lines as in the previous theorem. In particular, its hypotheses and conclusions are conformally invariant. Thus, we can assume that g is the metric of a conformal normal coordinate system. Then, let $\hat{M} = M \setminus \{p\}$ and define $\hat{g} \doteq G_p^{-4}g$ and $\tilde{g} = G_L^4 g$ on this manifold, where, as in the proof of the previous theorem, G_L denotes the Green function of the conformal Laplacian with a pole at p. Let us recall the following expansions, valid in conformal normal coordinates around p (see, [22] and [28])

$$G_p = A + O_4(r),$$

$$G_L = \frac{1}{r} + \alpha + O_4(r),$$

where A and α are constants. After going to inverted coordinates $z = \frac{x}{r}$, with $\rho = r^{-1}$, we find that

$$\widetilde{g}_{ij}(z) = \left(1 + \frac{\alpha}{\rho}\right)\delta_{ij} + O(\rho^{-2}),$$

and, clearly, on $\hat{g} = \Phi^{-4}\tilde{g}$, with $\Phi \doteq G_P G_L$ and $Q_{\hat{g}} \equiv 0$. Therefore, from the conformal covariance of the Paneitz operator, we find that $P_{\tilde{g}}\Phi \equiv 0$, which translates to

$$0 = \Delta_{\tilde{g}}^2 \Phi + 4 \operatorname{div}_{\tilde{g}} \left(\operatorname{Ric}_{\tilde{g}} (\nabla \Phi, \cdot) \right) + |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \Phi$$

That is

$$\int_{D_{\rho}} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \Phi dV_{\tilde{g}} = -\int_{S_{\rho}} \tilde{g}(\tilde{\nabla}\Delta_{\tilde{g}}\Phi, \tilde{\nu}) d\tilde{\omega}_{\rho} - 4 \int_{S_{\rho}} \operatorname{Ric}_{\tilde{g}}(\nabla\Phi, \tilde{\nu}) d\tilde{\omega}_{\rho}.$$
 (51)

As in the previous theorem, we now can compute explicitly the terms in the right-hand side. That is,

$$\Phi = \left(A + O_4(\rho^{-1})\right) \left(\rho + \alpha + O_4(\rho^{-1})\right) = A\rho + O_4(\rho^0)$$
$$\nabla_i \Phi = A \frac{z^i}{\rho} + O_3(\rho^{-1}),$$
$$\Delta_{\tilde{g}} \Phi = \frac{2A}{\rho} + O_2(\rho^{-2}),$$
$$\nabla_i \Delta_{\tilde{g}} \Phi = -2A \frac{z^i}{\rho^3} + O_1(\rho^{-3}),$$

which, together with $\operatorname{Ric}_{\tilde{q}} = O_2(\rho^{-3})$ implies that

$$\int_{D_{\rho}} |\operatorname{Ric}_{\tilde{g}}|_{\tilde{g}}^2 \Phi dV_{\tilde{g}} = 8\pi A + O(\rho^{-1}).$$

Passing to the limit and remembering that $\Phi < 0$, we find that $A \leq 0$ with equality holding iff $\operatorname{Ric}_{\tilde{q}} \equiv 0$. That is, if $\hat{M} \cong \mathbb{R}^3$ and $\tilde{g} = \delta$, which implies the final result.

A Appendix: Some analytic results concerning AE manifolds

In this appendix we will collect some facts results concerning AE manifolds which are used in the core of the paper. Most of these results are well-known for experts. We include them for the sake completeness and to deliver a self-contained presentation. For detailed proofs and discussions on these topics, we refer the reader to Bartnik [4], Lee-Parker [28].

Let us start with the following fundamental theorem regarding the properties of the Laplacian on AE manifolds.

Theorem A.1. Let (M, g) an asymptotically Euclidean manifold with a structure at infinity $\phi : M \setminus K \to \mathbb{R}^n \setminus B_1$ with decay rate τ . If $\delta \notin (\mathbb{Z} \setminus \{-1, \dots, 3-n\})$ and q > 1then

$$\Delta_g: W^{2,q}_{\delta}(\phi) \to L^q_{\delta-2}$$

is Fredholm. Moreover

$$\begin{cases} if \ \delta > 2 - n \ then \ \Delta_g \ is \ surjective, \\ if \ 2 - n < \delta < 0 \ then \ \Delta_g \ is \ bijective, \\ if \ \delta < 0 \ then \ \Delta_g \ is \ injective. \end{cases}$$
(A.1)

Remark A.1. The decay rates in the set $\mathbb{Z} \setminus \{-1, \dots, 3-n\}$ are called **exceptional** and we say that δ is **non-exceptional** if $\delta \notin \mathbb{Z} \setminus \{-1, \dots, 3-n\}$.

Let us now analyse the relation between different potential structures of infinity. In particular, the following theorem concerns the existence of harmonic coordinates.

Theorem A.2. Let (M,g) be an AE manifold, with $g \in W_{loc}^{k,q}$, $k \ge 1$, q > n, and $(\Phi, x) : M \setminus K \mapsto E_R \doteq \mathbb{R}^n \setminus \overline{B_R(0)}$ where $K \subset M$, $R \ge 1$ is a structure of infinity of order $\tau > 0$ with $1 - \tau$ non-exceptional, and fix $1 < \eta < 2$. There are functions $y^i \in W_{\eta}^{k,q}$,

 $i = 1, \dots, n$, such that $\Delta_g y^i = 0$ and $(x^i - y^i) \in W^{k,q}_{1-\tau^*}(E_R)$ for $\tau^* \doteq \min\{\tau, n-2\}$, which implies

$$|x^{i} - y^{i}| = o_{k}(r^{1-\tau^{*}}),$$

$$|g(\partial_{x^{i}}, \partial_{x^{j}}) - g(\partial_{y^{i}}, \partial_{y^{j}})| = o_{k}(r^{-\tau^{*}}).$$
(A.2)

Furthermore, the set of functions $\{1, y^i\}$ is a basis for $H_1 = \{u \in W_n^{k,q} : \Delta_g u = 0\}$.

Proof. Let us first extend the functions x^i smoothly to all of M. Then, near infinity, $\Delta_g x^i = g^{kl} \Gamma_{kl}^i \doteq \Gamma^i \in W_{1-\tau}^{k-1,q}$. Notice that if $1 - \tau > 2 - n$, then Theorem A.1 implies the existence of $v^i \in W_{1-\tau}^{k+1,q}$ solving $\Delta_g v^i = \Gamma^i$. Now, if $1 - \tau < 2 - n$ (the equality case is an exceptional case), we cannot, a priori, improve the decay given by r^{2-n} . This actually follows from Theorem 1.17 in [4]. Therefore, in any case, we know that there are solutions $v^i \in W_{1-\tau^*}^{k+1,q}$ to $\Delta_g v^i = \Gamma^i$, where $\tau^* = \min\{\tau, n-2\}$. Therefore, there is some $v^i \in W_{1-\tau^*}^{k+1,q}$ satisfying $\Delta_g(x^i - v^i) = 0$. Let us then define $y^i \doteq x^i - v^i$, which implies the first estimate in (A.2), and furthermore, since $\frac{\partial y^i}{\partial x^j} = \delta_j^i + o(r^{-\tau^*})$, we see that near infinity $\{y^i\}$ are coordinates which are asymptotically Cartesian. This, in turn, implies the second estimate in (A.2). The final claim concerning H_1 follows since $\{1, y^i\}$ spans an (n+1)-dimensional subspace of H_1 , but from Proposition 2.2 in [4] dim $(H_1) = n + 1$. □

Remark A.2. Let us highlight that the statement of the above theorem is slightly different than the well-known Theorem 3.1 in [4]. The difference relies in the fact that, in [4], it does not seem to be explicitly stated that, a priori, the decay rate r^{2-n} cannot be improved, regardless of how large τ may be. Nevertheless, as can be seen in the above proof, the above mild correction comes about from results contained in the same reference, as is also clear from a careful examination of the proof of Theorem 3.1 in [4]. Similar comments apply to the following theorem, which can be found in [4] as Corollary 3.2, and presents the relation between two different structures of infinity.

Theorem A.3. Let (M, g) an asymptotically flat manifold, $g \in W_{loc}^{k,q}$, $k \ge 1$ and q > n, with two structures at infinity $\phi, \psi : M \setminus K \to \mathbb{R}^n \setminus \overline{B_1(0)}$ with decay rates τ_{ϕ} and τ_{ψ} , where of each of these weights satisfies that $1 - \tau$ is non-exceptional. There exists $(O, a) \in O(n) \times \mathbb{R}^n$ such that

$$x^{i} - (O_{i}^{i}z^{j} + a^{i}) \in W^{k,q}_{1-\tau}(\mathbb{R}^{n})$$

which implies

$$|x^{i} - (O_{j}^{i}z^{j} + a^{i})| = o_{k}(r^{1-\tau}), \qquad (A.3)$$

where $\tau \doteq \min\{\tau_{\phi}, \tau_{\psi}, n-2\}, x = \phi^{-1} \text{ and } z = \psi^{-1}.$

Proof. Let y^i and w^i be the harmonic coordinates constructed in the previous theorem associated to ϕ and ψ respectively. Then, since H_1 is intrinsic to M and $\{1, y^i\}$ and $\{1, w^i\}$ are bases for this space, we get that

$$w^i = A^i_i y^j + a^i,$$

where we $A \in \operatorname{GL}(n, \mathbb{R})$ a priori, but actually, from the construction of the previous theorem, we know that $z^i = A^i_j x^j + a^i - A^i_j v^j + \overline{v}^i$, with $v^i \in W^{k,q}_{1-\tau^*_{\phi}}$ and $\overline{v}^i \in W^{k,q}_{1-\tau^*_{\phi}}$, where τ^* was defined in the previous theorem. Since these last two systems are Cartesian, then $A \in O(n)$ and we explicitly see that $z^i - A^i_j x^j - a^i \in W^{k,q}_{1-\tau}$.

The above estimate is the best possible and it is determined by the Ricci curvature as shown by the next theorem which corresponds to Proposition 3.3 in [4], see also [13].

Theorem A.4. Let (M, g) an asymptotically flat manifold with a structure at infinity $\phi: M \setminus K \to \mathbb{R}^n \setminus B(0, 1)$ with decay rate τ such that $(\phi_*g - \delta) \in W^{2,q}_{-\tau}(\mathbb{R}^n \setminus B(0, 1))$ for q > n and such that

$$Ric_g \in L^q_{-2-n}(M)$$
 for some $\eta > \tau$ and $\eta \notin (\mathbb{Z} \setminus \{-1, \cdots, 3-n\}).$

Then there exists a structure at infinity $\Theta : M \setminus K' \to \mathbb{R}^n \setminus B_1$ such that $(\Theta_* g - \delta) \in W^{2,q}_{-n}(\mathbb{R}^n \setminus B_1)$.

Finally, the following statement concerning the conformal Laplacian will be important in our analysis. It is a variant of theorem 9.2 of [28]

Corollary A.1. Let (M, g) an asymptotically flat manifold of order $\tau > 0$ and assume $R_g \ge 0$. If $2 - n < \delta < 0$ and q > 1, then

$$L_g = \Delta_g - c_n R_g : W^{2,q}_{\delta} \to L^q_{\delta-2}$$

is an isomorphism.

Proof. Since L_g is a compact perturbation of the Laplacian, it is also Fredholm, hence thanks to its self-adjointness it suffices to proof injectivity to get the result. Consider $u \in W^{2,q}_{\delta}$ such that $\Delta_g u = c_n R_g u$, and first notice that by elliptic regularity we can assume that $u \in W^{2,q}_{\delta}$ with q > n. Also, notice that $R_g u \in W^{2,q}_{\delta'-2}$ for any $\delta' > \delta - \tau$. Since $\tau > 0$, we can pick δ' satisfying max $\{\delta - \tau, 2 - n\} < \delta' < \delta$ and appeal to Theorem A.1 to obtain $u \in W^{2,q}_{\delta'}$ with $\delta' > 2 - n$ taken arbitrary close to 2 - n. Then, we can multiply $L_g(u) = 0$ by u and integrate by parts to obtain

$$\int_{M \setminus (\mathbb{R}^n \setminus B_R(0))} \left(|\nabla u|_g^2 + c_n R_g u^2 \right) dv_g = \int_{\partial B_R(0)} u \partial_\nu u \, d\sigma$$

Since $u = O_2(r^{\delta'})$ near infinity, we can estimate $|\nabla u|_g^2 = O(r^{2\delta'-2})$ and $R_g u^2 = O(r^{2\delta'-\tau-2})$ which implies these quantities are in $L^1(M)$ as long as chose $2 - n < \delta' < \frac{2-n}{2}$. Furthermore, under these condition one finds $u\partial_{\nu}u = O(r^{2\delta'-1})$ and therefore

$$\int_{M \setminus (\mathbb{R}^n \setminus B_R(0))} \left(|\nabla u|_g^2 + c_n R_g u^2 \right) dv_g = O(R^{n-2\delta'-2})$$

so that we can pass to the limit as $R \to +\infty$, which gives that $u \equiv 0$ since $R_g \ge 0$ and $u \to 0$ as $|x| \to \infty$, which concludes the proof.

B Appendix: Conventions on Q-curvature

Let us adopt the following general definition of Q-curvature for an arbitrary Riemannian manifold (M^n, g) with $n \ge 3$:

$$Q_g = -\Delta_g \sigma_1(S_g) + 4\sigma_2(S_g) + \frac{n-4}{2} \left(\sigma_1(S_g)\right)^2,$$
(B.1)

where $S_g \doteq \frac{1}{n-2} \left(\operatorname{Ric}_g - \frac{1}{2(n-1)} R_g g \right)$ stands for the Schouten tensor and $\sigma_k(S_g)$ stands for the *k*-th elementary symmetric function of the eigenvalues of S_g . In this context, the Paneitz operator is defined by

$$P_{g}u \doteq \Delta_{g}^{2}u + \operatorname{div}_{g}\left((4S_{g} - (n-2)\sigma_{1}(S_{g})g)(\nabla u, \cdot)\right) + \frac{n-4}{2}Q_{g}u,$$
(B.2)

for all $u \in C^{\infty}(M)$. In this context, the following relations hold:

$$\sigma_1(S_g) = \frac{R_g}{2(n-1)},$$

$$\sigma_2(S_g) = \frac{1}{2} \left(\frac{n^2 - n}{4(n-1)^2(n-2)^2} R_g^2 - \frac{|\operatorname{Ric}_g|_g^2}{(n-2)^2} \right),$$
(B.3)

which implies that

$$Q_g = -\frac{1}{2(n-1)} \Delta_g R_g - \frac{2}{(n-2)^2} |\operatorname{Ric}_g|_g^2 + \frac{n^3 - 4n^2 + 16n - 16}{8(n-1)^2(n-2)^2} R_g^2,$$

$$P_g u = \Delta^2 u + \operatorname{div}_g \left(\left(\frac{4}{n-2} \operatorname{Ric}_g - \frac{n^2 - 4n + 8}{2(n-1)(n-2)} R_g g \right) (\nabla u, \cdot) \right) + \frac{n-4}{2} Q_g u.$$
(B.4)

In particular, for $n \neq 4$, if $\bar{g} = u^{\frac{4}{n-4}}g$, then

$$Q_{\bar{g}} = \frac{2}{n-4} u^{-\frac{n+4}{n-4}} P_g u.$$
(B.5)

In the case of n = 4 we can apply the above definitions to get

$$Q_g = -\frac{1}{6}\Delta_g R_g - \frac{1}{2}|\operatorname{Ric}_g|_g^2 + \frac{1}{6}R_g^2,$$

$$P_g u = \Delta_g^2 u + \operatorname{div}_g \left(\left(2\operatorname{Ric}_g - \frac{2}{3}R_g g \right) (\nabla u, \cdot) \right),$$
(B.6)

and in this case, if $\bar{g} = e^{2u}g$, we have that [30]

$$Q_{\bar{g}} = e^{-4u} \left(P_g u + Q_g \right).$$
 (B.7)

Now, let us notice that it is also quite standard to *redefine* the Q-curvature in dimension 4 via (see [20, 29])

$$Q_g^{(4)} = \frac{1}{2}Q_g = -\frac{1}{12}\Delta_g R_g - \frac{1}{4}|\operatorname{Ric}_g|_g^2 + \frac{1}{12}R_g^2$$
(B.8)

and in this case, it follows that

$$2Q_{\bar{g}}^{(4)} = e^{-4u} \left(P_g u + 2Q_g^{(4)} \right).$$
(B.9)

We will not adopt this redefinitions and keep a unified notation via (B.1) along this paper.

C Appendix: Conformal normal coordinates

In order to deliver a presentation as self-cointained as possible, this section is meant to summarise some of the results concerning conformal normal coordinates presented in [28], which which are used in the core of this paper. The basic construction is given on a smooth Riemannian manifold (M^n, g) where we intend to expand g around a fixed point $p \in M$. In particular, we are interested in finding an element \tilde{g} within the conformal class [g] for which, in \tilde{g} -normal coordinates $\{x^i\}$, the following expansion holds around p:

$$\det(\tilde{g}) = 1 + O(r^N) \tag{C.1}$$

for any chosen $N \ge 2$, where r = |x| (see Theorem 5.1 in [28]). This type of expansion is achieved by first noticing that, for any Riemannian metric g, in g-normal coordinates $\{x^i\}_{i=1}^n$ around $p \in M$, the following holds

$$g_{ij}(x) = \delta_{ij} + \frac{1}{3}R_{iklj}x^kx^l + \frac{1}{6}R_{iklj,a}x^kx^lx^a + \left(\frac{1}{20}R_{iklj,ab} + \frac{2}{45}R_{iklc}R_{jabc}\right)x^kx^lx^ax^b + O(r^5)$$

where all the coefficients at evaluated at p. From this expression it is possible to compute det(g) around p for any such metric, so as to get

$$\det(g)(x) = 1 - \frac{1}{3}R_{ij}x^{i}x^{j} - \frac{1}{6}R_{ij,k}x^{i}x^{j}x^{k} - \left(\frac{1}{20}R_{ij,kl} + \frac{1}{90}R_{aijb}R_{aklb} - \frac{1}{18}R_{ij}R_{kl}\right)x^{i}x^{j}x^{k}x^{l} + O(r^{5}),$$
(C.2)

where again all the coefficients are evaluated at p. Assuming an expansion of the form $det(g) = 1 + O(r^N)$ with $N \ge 2$ (the case N = 2 is valid for any metric in its own normal coordinates) and appelaing to Theorem 5.2 in [28], the authors can find a homogeneous polynomial $f \in \mathcal{P}_N$ so that the expansion of $det(\tilde{g}) = 1 + O(r^{N+1})$ for $\tilde{g} = e^{2f}g$, which establishes an inductive proof (see the proof of Theorem 5.1). But notice that this implies that the symmetrization of the coefficients in (C.2) of order up to N must vanish for \tilde{g} . Thus, in the case we do this construction for $N \ge 4$, we see that this implies

$$R_{ij}(p) = 0, \tilde{R}_{ij,k} + \tilde{R}_{ki,j} + \tilde{R}_{jk,i}(p) = 0.$$
(C.3)

Putting together the second condition above with the contracted Bianchi identities, we also see that

$$\ddot{R}_{,k}(p) = 0. \tag{C.4}$$

References

- [1] Richard Arnowitt, Stanley Deser, and Charles W. Misner. Coordinate invariance and energy expressions in general relativity. *Phys. Rev.*, 122(3):9971006, 2006.
- [2] Rodrigo Avalos and Jorge H. Lira. Reduced thin-sandwich equations on manifolds euclidean at infinity and on closed manifolds: Existence and multiplicity. J. Math. Phy., 61:122501, 2020.

- [3] Rodrigo Avalos, Jorge H. Lira, and Nicolas Marque. Energy in Fourth Order Gravity. *arXiv e-prints*, page arXiv:2102.00545, January 2021.
- [4] Robert Bartnik. The mass of an asymptotically flat manifold. Comm. Pure Appl. Math., 39(5):661-693, 1986.
- [5] Simon Brendle and Michael. Eichmair. Large outlying stable constant mean curvature spheres in initial data sets. *Invent. Math.*, 107(3):663–682, 2014.
- [6] Cliff P. Burgess. Quantum gravity in everyday life: General relativity as an effective field theory. *Living Rev. Relativity*, 7:5, 2004.
- [7] Alessandro Carlotto. Rigidity of stable minimal hypersurfaces in asymptotically flat spaces. Calc. Var. Partial Differ. Equ., 55(3):1–20, 1988.
- [8] Alessandro Carlotto. Four Lectures on Asymptotically Flat Riemannian Manifolds. in: Einstein Equations: Physical and Mathematical Aspects of General Relativity. Eds: S. Cacciatori, Batu Güneysu and S. Pigola. Springer Nature Switzerland AG, 2019.
- [9] Alessandro Carlotto, Otis Chodosh, and Michael. Eichmair. Effective versions of the positive mass theorem. *Invent. Math.*, 206(3):1–20, 2016.
- [10] Alessandro Carlotto and Richard. Schoen. Localizing solutions of the einstein constraint equations. *Invent. Math.*, 205(3):1–20, 2016.
- [11] Sun-Yung Alice Chang and Paul C. Yang. Extremal metrics of zeta functional determinants on 4-manifolds. Ann. of Math., 142:171–212, 1995.
- [12] Szu-yu Sophie Chen. Conformal deformation on manifolds with boundary. Geom. Funct. Anal., 19(4):1029–1064, 2009.
- [13] Dennis M. DeTurck and Jerry L. Kazdan. Some regularity theorems in Riemannian geometry. Ann. Sci. École Norm. Sup. (4), 14(3):249–260, 1981.
- [14] James Dilts and David Maxwell. Yamabe classification and prescribed scalar curvature in the asymptotically Euclidean setting. Comm. Anal. Geom., 26(5):1127–1168, 2018.
- [15] Zindine Djadli and Andrea Malchiodi. Existence of conformal metrics with constant q-curvature. Ann. Math., 168:813–858, 2008.
- [16] John F. Donoghue. General relativity as an effective field theory: The leading quantum cerrections. *Phys. Rev. D*, 50(6):3874–3888, 1994.
- [17] Michael Eichmair and Jan Metzger. Large isoperimetric surfaces in initial data sets. J. Differ. Geom., 94(1):159–186, 2013.
- [18] Matthew J. Gursky. The principal eigenvalue of a conformally invariant differential operator, with an application to semilinear elliptic PDE. Comm. Math. Phys., 207(1):131–143, 1999.

- [19] Matthew J. Gursky, Fengbo Hang, and Yueh-Ju Lin. Riemannian manifolds with positive Yamabe invariant and Paneitz operator. Int. Math. Res. Not. IMRN, (5):1348– 1367, 2016.
- [20] Matthew J. Gursky and Andrea Malchiodi. A strong maximum principle for the Paneitz operator and a non-local flow for the Q-curvature. J. Eur. Math. Soc. (JEMS), 17(9):2137–2173, 2015.
- [21] Fengbo Hang and Paul C. Yang. Sign of Green's function of Paneitz operators and the Q curvature. Int. Math. Res. Not. IMRN, (19):9775–9791, 2015.
- [22] Fengbo Hang and Paul C. Yang. Q curvature on a class of 3-manifolds. Comm. Pure Appl. Math., 69(4):734–744, 2016.
- [23] Fengbo Hang and Paul C. Yang. Q-curvature on a class of manifolds with dimension at least 5. Comm. Pure Appl. Math., LXIX:1452–1491, 2016.
- [24] Luan. H. Huang. Foliations by stable spheres with constant mean curvature for isolated systems with general asymptotics. *Commun. Math. Phys.*, 300(2):331–373, 2010.
- [25] Gerhard Huisken and Shing T. Yau. Definition of center of mass for isolated physical systems and unique foliations by stable spheres with constant mean curvature. *Invent. Math.*, 124:281–311, 1996.
- [26] Emmanuel Humbert and Simon Raulot. Positive mass theorem for the Paneitz-Branson operator. Calc. Var. Partial Differential Equations, 36(4):525–531, 2009.
- [27] Dan A. Lee. *Geometric relativity*, volume 201 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2019.
- [28] John M. Lee and Thomas H. Parker. The Yamabe problem. Bull. Amer. Math. Soc. (N.S.), 17(1):37–91, 1987.
- [29] Jiayu Li, Yuxiang Li, and Pan Liu. The *Q*-curvature on a 4-dimensional Riemannian manifold (M, g) with $\int_M Q dV_g = 8\pi^2$. Adv. Math., 231(3-4):2194–2223, 2012.
- [30] Yueh-Ju Lin and Wei Yuan. Deformations of Q-curvature I. Calc. Var. Partial Differential Equations, 55(4):Art. 101, 29, 2016.
- [31] Juan Maldacena. Einstein gravity from conformal gravity. arXiv:1105.5632v2.
- [32] Philip D. Mannheim. Making the case for conformal gravity. Found Phys, 42:388–420, 1980.
- [33] David Maxwell. Solutions of the Einstein constraint equations with apparent horizon boundaries. *Comm. Math. Phys.*, 253(3):561–583, 2005.
- [34] Benoît Michel. Masse des opérateurs GJMS. *arXiv e-prints*, page arXiv:1012.4414, December 2010.
- [35] Benoît Michel. Geometric invariance of mass-like asymptotic invariants. Journal of Mathematical Physics, 52(5):052504, 2011.

- [36] Cheikh Birahim Ndiaye. Constant q-curvature metrics in arbitrary dimension. Journal of Functional Analysis, 251:1–58, 2007.
- [37] Christopher Nerz. Foliations by stable spheres with constant mean curvature for isolated systems without asymptotic symmetry. *Calc. Var. Partial Differ. Equ*, 54(2):1911–1946, 2015.
- [38] J. Qing and D. Raske. On positive solutions to semilinear conformally invariant equations on locally conformally flat manifolds. *Int. Math. Res. Not.*, 2006(9):94172– 94172, 2006.
- [39] Gerard 'T Hooft and Martinus Veltman. One loop divergencies in the theory of gravitation. Ann. Inst. Henri Poincare, A, 20:69–94, 1974.
- [40] Richard Schoen. Conformal deformation of a riemannian metric to constant scalar curvature. J. Diff. Geom, 20:479–595, 1984.
- [41] Richard Schoen and Shing T. Yau. On the proof of the positive mass conjecture in general relativity. *Comm. Math. Phys*, 65:45–76, 1979.
- [42] Richard Schoen and Shing T. Yau. Comm. Math. Phys, 79:231–260, 1981.
- [43] Richard Schoen and Shing T. Yau. Conformally flat manifolds, kleinian groups and scalar curvature. *Invent. math.*, 92:47–71, 1988.
- [44] Alexei Starobinsky. A new type of isotropic cosmological models without singularity. *Physics Letters B*, 91(1):99–102, 1980.