

On an isomorphism theorem for the Feichtinger's Segal algebra on locally compact groups^{*}

R. Lakshmi Lavanya

*Indian Institute of Science Education and Research (IISER) Tirupati,
Karakambadi Road, Mangalam,
Tirupati - 517507, India.*

Abstract

In this article we observe that a locally compact group G is completely determined by the algebraic properties of its Feichtinger's Segal algebra $S_0(G)$. Let G and H be locally compact groups. Then any linear (not necessarily continuous) bijection of $S_0(G)$ onto $S_0(H)$ which preserves the convolution and pointwise products is essentially a composition with a homeomorphic isomorphism of H onto G .

Keywords: Locally compact groups, Fourier algebra, Feichtinger's Segal algebra, Isomorphism theorem

2010 MSC: Primary 43A30, 46J35; Secondary 22D15, 47J25.

1. Introduction

In 1981, Feichtinger introduced and studied a new Segal algebra on locally compact Abelian groups([3]). This algebra, now called the Feichtinger algebra has been extensively studied over the years. A recent survey of this progress can be found in [5]. The genesis of the study of the Feichtinger's Segal algebra for locally compact groups came up almost a quarter century later([9]). In [9], the author introduces an operator space structure on Feichtinger's Segal algebra for locally compact groups. It was also observed that $S_0(G)$ for a locally compact group has many functorial properties very similar to that in the Abelian setting. Infact, some of these properties could be obtained even without exploiting the operator space structure of the Feichtinger's Segal algebra. One such example is the isomorphism theorem, viz., Theorem 2.1 proved in [9]. This result states that any bounded linear bijection which preserves the convolution and pointwise products on the Feichtinger's Segal algebra determines the underlying locally

^{*}Dedicated to the memory of Prof. Elias M. Stein

^{**}The author's research is supported by DST-SERB, Government of India, under the Early Career Research Award Scheme(No. ECR/2017/003366).

compact group.

The aim of this article is to observe that the algebraic properties of the Feichtinger's Segal algebra determine the underlying locally compact group. More precisely, a locally compact group is completely determined by any linear bijection (not necessarily continuous), which preserves the convolution and pointwise products on its Feichtinger's Segal algebra.

For more on isomorphism theorems, we refer to [6], [9], [10], [11] and the references therein.

2. Preliminaries and Notation

For a locally compact group G , let $\mathcal{C}_c(G)$ denote the space of all compactly supported continuous functions on G . As per the standard convention, we denote by $A(G)$ and $B(G)$, the Fourier and the Fourier-Stieltjes algebras on G , respectively. The Feichtinger's Segal algebra on G , denoted by $S_0(G)$, is defined as follows([9]):

Let \mathcal{A} be a regular Banach algebra with Gelfand spectrum X . An ideal \mathcal{I} of \mathcal{A} is said to have *compact support* if $\text{supp } \mathcal{I} = \overline{X \setminus h(\mathcal{I})}$ is compact in X . Here, $h(\mathcal{I})$ denotes the *hull* of \mathcal{I} , defined as $h(\mathcal{I}) = \{x \in X : u(x) = 0 \text{ for all } u \in \mathcal{I}\}$.

Let \mathcal{I} be a non-zero ideal in $A(G)$ which has compact support. Identify $l^1(G)$ with the closed linear span of the Dirac measures $\{\delta_s : s \in G\}$. Consider $l^1(G)$ with the maximal operator space structure.

Define

$$q_{\mathcal{I}} : l^1(G) \widehat{\otimes} \mathcal{I} \rightarrow A(G)$$

as

$$q_{\mathcal{I}}(x \otimes u) = \sum_{s \in G} \alpha_s s \cdot u,$$

where $x = \sum_{s \in G} \alpha_s \delta_s$ and $s \cdot u(t) = u(s^{-1}t)$, $t \in G$. Here, $\widehat{\otimes}$ denotes the operator projective tensor product. The *Feichtinger's Segal algebra* on G is defined as

$$S_0(G) = \text{ran}(q_{\mathcal{I}}) \cong l^1(G) \widehat{\otimes} \mathcal{I} / \ker(q_{\mathcal{I}}).$$

Equipped with the quotient operator space structure, $S_0(G)$ becomes an operator space.

For functions $u, v \in S_0(G)$, $u * v$ denotes the usual convolution product, given by

$$u * v(x) = \int_G u(y) v(y^{-1}x) dy.$$

Here, we use dy to denote a fixed left Haar measure on G .

In [9], it was illustrated that $S_0(G)$ satisfies several functorial properties analogous to its Abelian counterpart.

The following isomorphism theorem for S_0 was proved in [9].

Theorem 2.1 ([9]). *Let G and H be locally compact groups and $\Phi : S_0(G) \rightarrow S_0(H)$ be a bounded linear bijection which satisfies*

$$\Phi(uv) = \Phi u \Phi v \quad \text{and} \quad \Phi(u * v) = \Phi u * \Phi v$$

for every $u, v \in S_0(G)$. There is a homeomorphic isomorphism $\alpha : H \rightarrow G$ such that

$$\Phi u = u \circ \alpha$$

for each u in $S_0(G)$.

The aim of the present article is to observe that the conclusion of the above result holds good even if the map Φ is not apriori continuous (see Theorem 3.4). This leads us to the fact that the algebraic properties of the Feichtinger's Segal algebra solely determine the underlying locally compact group.

For further details and properties of $S_0(G)$, we refer to [9].

3. An Isomorphism theorem for $S_0(G)$

First we obtain the form of any bijection which is multiplicative on $S_0(G)$. Later we will obtain the isomorphism theorem by additionally assuming that this bijection preserves the convolution product too.

In this section, we use letters x, y, \dots to denote the elements of the respective groups, as indicated in the context.

Our preliminary result is the following, which exploits only the pointwise product structure of $S_0(G)$.

Theorem 3.1. *Let G and H be locally compact groups and $\Phi : S_0(G) \rightarrow S_0(H)$ be a bijection which satisfies the following conditions for every $u, v \in S_0(G)$:*

$$(1) \quad \Phi(u + \overline{v}) = \Phi(u) + \overline{\Phi(v)}$$

$$(2) \quad \Phi(uv) = \Phi u \Phi v$$

Then there exists a decomposition of H as $H = H_1 \cup H_2$ and a homeomorphism $\alpha : H \rightarrow G$ such that

$$\Phi u = \chi_{H_1} (u \circ \alpha) + \chi_{H_2} (\overline{u \circ \alpha}),$$

for each $u \in S_0(G)$.

PROOF. The proof consists of 7 steps. Some of these overlap with a characterisation of the Fourier transform on the Fourier twins on locally compact Abelian groups (as in Theorem 3.2 of [7]). For the sake of completeness, a proof of these ideas (steps 1-5) is included in the Appendix. Here, we only state these steps.

Step 1. If $u \in \mathcal{C}_c(G)$, then $\Phi u \in \mathcal{C}_c(H)$.

Step 2. For any $x_0 \in G$, there exists $y_0 \in H$ such that $\Phi u(y_0) = 0$ whenever $u(x_0) = 0$, for any $u \in S_0(G)$.

We say that a function $u : G \rightarrow \mathbb{C}$ satisfies condition (\star) if the following holds good:

$$(\star) \quad u(x_0) = 0 \text{ if and only if } \Phi u(y_0) = 0.$$

Step 3. Define a map $\beta : G \rightarrow H$ as follows: $\beta(x) = y$ if $\Phi u(y) = 0$ whenever $u(x) = 0$, for any $u \in S_0(G)$. Then the map β is well-defined.

Step 4. The map $\beta : G \rightarrow H$ is a bijection.

Step 5. The map β is a homeomorphism of G onto H .

Step 6. 'Extension' of the map Φ to scalars.

Illustration of Step 6. For $u, v \in S_0(G)$, and $c(\neq 0) \in \mathbb{C}$, we have

$$\Phi(cu)(x) \Phi v(x) = \Phi(cuv)(x) = \Phi(u)(x) \Phi(cv)(x), \quad x \in G.$$

Let $w \in S_0(G)$ be such that $\Phi w(x) \neq 0$ for any $x \in G$. Then we have

$$\begin{aligned} \Phi(cu)(x) &= \frac{\Phi(cw)(x)}{\Phi w(x)} \Phi u(x) \text{ for all } u \in S_0(G) \\ &= m(c, x) \Phi u(x) \text{ (say).} \end{aligned}$$

When $c = 0$, we have $m(c, x) \Phi u(x) = \Phi(cu)(x) = 0$, for all $x \in H$ and for all $u \in S_0(G)$. Hence it is reasonable to define $m(0, x) = 0$ for all $x \in H$.

(Here, we have used that $\Phi(0_G) = 0_H$, where 0_G and 0_H are the zero functions on the respective groups. This follows by condition(1) since $\Phi(0_G) = \Phi(0_G + 0_G) = \Phi(0_G) + \Phi(0_G)$.)

Thus $\Phi(cu)(x) = m(c, x) \Phi u(x)$, for all $x \in H$.

In particular, the function $m(\cdot, \cdot)$ is continuous in the second variable, as a function of $x \in H$.

Step 7. For fixed $y \in H$, the map $m(\cdot, y) : \mathbb{C} \rightarrow \mathbb{C}$ is an additive and multiplicative bijection, which maps \mathbb{R} onto \mathbb{R} , and hence we have either $m(c, y) = c$, for all $c \in \mathbb{C}$, or $m(c, y) = \bar{c}$, for all $c \in \mathbb{C}$.

Proof of Step 7. Choose $u \in S_0(G)$, with $u(x) \neq 0$ for any $x \in G$. Then by the condition (\star) , we get $\Phi u(\beta(x)) \neq 0$, for any $x \in G$.

Suppose $m(c, y) = m(d, y)$ for some $c, d \in \mathbb{C}$. Then

$$\Phi(cu)(y) = m(c, y) \Phi u(y) = m(d, y) \Phi u(y) = \Phi(du)(y), \quad y \in H.$$

As Φ is a bijection, this gives $cu = du$. Since u is never vanishing, we get that $c = d$.

By hypothesis(1), we have

$$\begin{aligned} m(c + \bar{d}, y) \Phi u(y) &= \Phi((c + \bar{d})u)(y) = \Phi(cu + \bar{d}u)(y) \\ &= \Phi(cu)(y) + \overline{\Phi(\bar{d}u)}(y) = (m(c, y) + \overline{m(d, y)}) \Phi u(y). \end{aligned}$$

Since Φu is never zero, we get $m(c + \bar{d}, y) = m(c, y) + \overline{m(d, y)}$. In particular, $m(d, y) \in \mathbb{R}$ whenever $d \in \mathbb{R}$.

Now, hypothesis(2) gives

$$m(cd, y) \Phi u(y) = \Phi(cdu)(y) = m(c, y) \Phi(du)(y) = m(c, y) m(d, y) \Phi u(y).$$

Again, since Φu is nowhere vanishing, we get $m(cd, y) = m(c, y) m(d, y)$ for all $c, d \in \mathbb{C}$.

Thus $m(\cdot, y)$ is a field automorphism of \mathbb{C} which preserves \mathbb{R} . So for fixed $y \in H$, we have either $m(c, y) = c$ for all $c \in \mathbb{C}$, or $m(c, y) = \bar{c}$ for all $c \in \mathbb{C}$. Let

$$\begin{aligned} H_1 &= \{y \in H : m(c, y) = c \text{ for all } c \in \mathbb{C}\}, \\ H_2 &= \{y \in H : m(c, y) = \bar{c} \text{ for all } c \in \mathbb{C}\}. \end{aligned}$$

Step 8. For $u \in S_0(G)$, and $x_0 \in G$, we have $\Phi u[\beta(x_0)] = m(u(x_0), \beta(x_0))$.

Proof of Step 8. As before, choose $u \in S_0(G)$ such that $u(x) \neq 0$ for any $x \in G$. Then $\Phi u(y) \neq 0$ for any $y \in H$. Let $v \in S_0(G)$.

Define

$$w(x) := v(x_0) u(x) - v(x) u(x), \quad x \in G.$$

Then $w \in S_0(G)$ and $w(x_0) = 0$. By Condition (\star) , we have $\Phi w[\beta(x_0)] = 0$. This gives

$$\begin{aligned} 0 = \Phi w[\beta(x_0)] &= \Phi[v(x_0) \cdot u - v \cdot u][\beta(x_0)] \\ &= m(v(x_0), \beta(x_0)) \Phi u[\beta(x_0)] - \Phi v[\beta(x_0)] \Phi u[\beta(x_0)]. \end{aligned}$$

Since Φu is never zero, this gives $\Phi v[\beta(x_0)] = m(v(x_0), \beta(x_0))$.

Since $\alpha = \beta^{-1}$, using Step 7, we get that

$$\text{either } \Phi v(x_0) = v[\alpha(x_0)] \text{ or } \Phi v(x_0) = \overline{v[\alpha(x_0)]}.$$

Combining this with the definition of the sets H_1 and H_2 gives the representation $\Phi v = \chi_{H_1} \alpha \circ v + \chi_{H_2} \overline{\alpha \circ v}$. \square

Corollary 3.2. *In addition to the hypothesis of Theorem 3.1, if H is connected, then the following conclusion holds:*

There exists a homeomorphism $\alpha : H \rightarrow G$ such that

$$\begin{aligned} &\text{either } \Phi u = u \circ \alpha \text{ for all } u \in S_0(G), \\ &\text{or } \Phi u = \overline{u \circ \alpha} \text{ for all } u \in S_0(G). \end{aligned}$$

PROOF. Since the map $m(\cdot, \cdot) : \mathbb{C} \times H \rightarrow \mathbb{C}$ is continuous in the second variable, we get by Step 7 of the previous theorem that either $m(c, y) = c$ for all $y \in H$, or $m(c, y) = \bar{c}$ for all $y \in H$. \square

As of now, we have not made use of the convolution product in $S_0(G)$. As mentioned in [9], it is reasonable to expect that any isomorphism theorem for the Feichtinger's Segal algebra involves both the convolution and the pointwise product. This leads us to the following result. The proof of this result is along the lines of the proof of Step 6 of Theorem 3.2 in [7]. We state it here for the sake of completeness.

Theorem 3.3. *In addition to the hypothesis of Theorem 3.1, assume that Φ satisfies*

$$\Phi(u * v) = \Phi u * \Phi v, \text{ for all } u, v \in S_0(G).$$

Then there exists a decomposition of H as $H = H_1 \cup H_2$ and a homeomorphic isomorphism $\alpha : H \rightarrow G$ such that

$$\Phi u = \chi_{H_1} (u \circ \alpha) + \chi_{H_2} (\overline{u \circ \alpha}),$$

for each $u \in S_0(G)$.

PROOF. In view of the proof of Theorem 3.1, we are left with proving that α is a homomorphism.

Claim: The map $(\alpha^{-1})^\beta : G \rightarrow H$ is a homomorphism.

Proof of Claim. Suppose $\beta(xy) \neq \beta(x)\beta(y)$ for some $x, y \in G$.

Then there exist disjoint neighbourhoods $V_{xy}, V_{x \diamond y}$ in H with $\beta(xy) \in V_{xy}$ and $\beta(x)\beta(y) \in V_{x \diamond y}$. By continuity of the map β , this gives rise to a neighbourhood $W_{xy,1}$ of xy in G with $\beta(W_{xy,1}) \subseteq V_{xy}$. By continuity of multiplication in G , we get neighbourhoods $W_{x,1}, W_{y,1}$ of x and y , respectively, in G , such that $W_{x,1}W_{y,1} \subseteq W_{xy,1}$. Thus

$$\beta(W_{x,1}W_{y,1}) \subseteq \beta(W_{xy,1}) \subseteq V_{xy}. \quad (3.1)$$

On the other hand, by continuity of multiplication in H , $\beta(x)\beta(y) \in V_{x \diamond y}$ gives neighbourhoods $V_{x,2}, V_{y,2}$ in H , such that

$$\beta(x) \in V_{x,2}, \beta(y) \in V_{y,2} \text{ and } V_{x,2} V_{y,2} \subseteq V_{x \diamond y}. \quad (3.2)$$

This implies there exist neighbourhoods $W_{x,2}, W_{y,2}$ of x and y , respectively, with $\beta(W_{x,2}) \subseteq V_{x,2}$ and $\beta(W_{y,2}) \subseteq V_{y,2}$.

Define $W_x = W_{x,1} \cap W_{x,2}$, $W_y = W_{y,1} \cap W_{y,2}$. Then

$$\begin{aligned} \beta(W_x) &\subseteq \beta(W_{x,2}) \subseteq V_{x,2} = V_x \text{ (say)} \\ \beta(W_y) &\subseteq \beta(W_{y,2}) \subseteq V_{y,2} = V_y \text{ (say)} \\ \beta(W_x W_y) &\subseteq \beta(W_{x,1} W_{y,1}) \subseteq \beta(W_{xy,1}) \subseteq V_{xy} \end{aligned} \quad (3.3)$$

Choose $u_x, u_y \in S_0(G)$ such that $\text{Supp } u_x \subseteq W_x, \text{Supp } u_y \subseteq W_y$ and $u_x * u_y \neq 0$. Let $v_x = \Phi u_x$ and $v_y = \Phi u_y$. Then $\Phi(u_x * u_y) = v_x * v_y \neq 0$.

We have

$$\begin{aligned} \text{Supp}(v_x * v_y) &= \text{Supp } \Phi(u_x * u_y) \subseteq (\text{Supp } \Phi u_x) (\text{Supp } \Phi u_y) \\ &= \beta(\text{Supp } u_x) \beta(\text{Supp } u_y) \subseteq \beta(W_x) \beta(W_y) \\ &\subseteq V_{x,2} V_{y,2} \subseteq V_{x \diamond y} \text{ (by (3.2))} \end{aligned} \quad (3.4)$$

But $\text{Supp } (u_x * u_y) \subseteq (\text{Supp } u_x) (\text{Supp } u_y) \subseteq W_x W_y$. By (3.3), this gives

$$\begin{aligned} \text{Supp}(v_x * v_y) &= \text{Supp}(\Phi u_x * \Phi u_y) = \text{Supp } \Phi(u_x * u_y) \\ &= \beta(\text{Supp}(u_x * u_y)) \subseteq \beta(W_x W_y) \subseteq V_{xy}. \end{aligned} \quad (3.5)$$

From (3.4) and (3.5), we get

$$\text{Supp}(v_x * v_y) \subseteq V_{x \diamond y} \cap V_{xy} = \emptyset.$$

This gives $v_x * v_y \equiv 0$, a contradiction. This proves the multiplicativity of the map β .

Thus the map β is a multiplicative homeomorphism of G onto H . As $\alpha = \beta^{-1}$, this proves our result. \square

We now state an isomorphism theorem for the Feichtinger's Segal algebra.

Theorem 3.4. *Let G and H be locally compact groups and $\Phi : S_0(G) \rightarrow S_0(H)$ be a linear bijection which satisfies the following conditions for every $u, v \in S_0(G)$:*

$$\Phi(uv) = \Phi u \Phi v \text{ and } \Phi(u * v) = \Phi u * \Phi v.$$

Then there exists an isomorphic homeomorphism $\alpha : G \rightarrow H$ such that

$$\Phi u = u \circ \alpha, \text{ for all } u \in S_0(G).$$

PROOF. We proceed as in Theorem 3.1, till Step 5.

Let $u \in S_0(G)$ be such that $u(x) \neq 0$ for any $x \in G$. Then $\Phi u(y) \neq 0$ for any $y \in H$ (by Condition (\star)). Let $v \in S_0(G)$.

Define $w(x) := v(x_0) u(x) - v(x) u(x)$, $x \in G$.

Then $w \in S_0(G)$ and $w(x_0) = 0$. By Condition (\star) , we get $\Phi w[\beta(x_0)] = 0$. Since Φ is linear, this gives

$$\begin{aligned} 0 = \Phi w[\beta(x_0)] &= \Phi[v(x_0) \cdot u - v \cdot u][\beta(x_0)] \\ &= v(x_0) \Phi u[\beta(x_0)] - \Phi v[\beta(x_0)] \Phi u[\beta(x_0)]. \end{aligned}$$

Since Φu is never zero, this gives $\Phi v[\beta(x_0)] = v(x_0)$, whenever $v(x_0) \neq 0$.

If $v(x_0) = 0$, then condition (\star) gives that $\Phi v(\beta(x_0)) = 0$. Since $\alpha = \beta^{-1}$, this proves that

$$\Phi v = v \circ \alpha \text{ for all } v \in S_0(G).$$

As in the proof of Theorem 3.3, we get that α is an isomorphism. \square

Remark 3.5. *As stated in [9], the above isomorphism theorems too, hold good for any regular Banach algebra of functions in G (whose spectrum apriori need not be G), which is contained in $C_0(G)$.*

Remark 3.6. *In particular, Theorem 3.4 holds good for the Rajchman algebra $B_0(G) = B(G) \cap C_0(G)$, defined and studied in [6].*

4. Appendix

Part of proof of Theorem 3.1: (as in [7])

For $x \in G$, let $C(x) = \{f \in S_0(G) : f(x) \neq 0\}$.

Step 1. If $u \in \mathcal{C}_c(G)$, then $\Phi u \in \mathcal{C}_c(H)$.

Proof of Step 1. If $v = 1$ on $\text{Supp } u$, we have $u \cdot v = u$. This gives $\Phi u = \Phi(u \cdot v) = \Phi u \cdot \Phi v$, and so $\Phi v = 1$ on the set $\{x : \Phi u(x) \neq 0\}$. By the continuity of Φv , we get $\Phi v = 1$ on $\text{Supp}(\Phi u)$.

For $u(\neq 0) \in \mathcal{C}_c(G)$, choose $v \in S_0(G)$ such that $v = 1$ on $\text{Supp } u$. Then $\Phi v = 1$ on $\text{Supp } \Phi u$. Since $\Phi v \in S_0(H) \subset C_0(H)$, we get $\text{Supp } \Phi u$ is compact.

Step 2. For any $x_0 \in G$, there exists $y_0 \in H$ such that $\Phi u \in C(y_0)$ whenever $u \in C(x_0)$.

Proof of Step 2. Let $E := \{u \in S_0(G) : u(x_0) \neq 0\}$. Fix a function $v \in \mathcal{C}_c(G)$ with $v(x_0) \neq 0$. By Step 1, we have $K := \text{Supp } \Phi v$ is compact.

For $u \in E$, define $K_u := K \cap \text{Supp } \Phi u$. For finitely many functions $u_0 := v, u_1, \dots, u_k \in E$, we have $\prod_{j=0}^k u_j \neq 0$, and so $\prod_{j=0}^k \Phi u_j = \Phi(\prod_{j=0}^k u_j) \neq 0$, which gives $\cap_{j=0}^k K_{u_j} \neq \emptyset$. This means, the collection $\{K_u : u \in E\}$ of closed subsets of K has finite intersection property. Since K is compact, this gives $\bigcap_{u \in E} K_u \neq \emptyset$. Let $y_0 \in \bigcap_{u \in E} K_u$.

Claim. For $u \in S_0(G)$, we have $u(x_0) = 0$ if and only if $\Phi u(y_0) = 0$.

Proof of Claim. We prove the claim in two separate cases.

Suppose $u(x_0) \neq 0$.

Then u never vanishes on a neighbourhood, say, V of x_0 . Let $v \in S_0(G)$ be such that $u \cdot v = 1$ on V . Choose $w \in S_0(G)$ such that $w = 1$ on a neighbourhood W of x_0 , and satisfies $W \subseteq \text{Supp } w \subseteq V$. Since $u \cdot v = 1$ on $\text{Supp } w$, by Step 1, we get $\Phi(u \cdot v) = \Phi u \cdot \Phi v = 1$ on $\text{Supp } \Phi w$. Since $h \in E$, by definition, $y_0 \in \text{Supp } \Phi w$. This implies $\Phi u(y_0) \neq 0$, and hence $\Phi u \in C(y_0)$.

Note that all our arguments till now can be applied to the map Φ^{-1} as well, and so we have proved that $u(x_0) \neq 0$ if and only if $\Phi u(y_0) \neq 0$.

A function $u : G \rightarrow \mathbb{C}$ is said to satisfy the condition (\star) if the following holds:

$$u(x_0) = 0 \text{ if and only if } \Phi u(y_0) = 0. \quad (\star)$$

By the above discussion, we have that all functions in $S_0(G)$ satisfy the Condition (\star) .

Step 3. Define a map $\beta : G \rightarrow H$ as follows: $\beta(x) = y$ if $\Phi u \in C(y)$ whenever $u \in C(x)$. Then the map β is well-defined.

Proof of Step 3. Suppose for some $x_0 \in G$, we have $\beta(x_0) = y_1$ and $\beta(x_0) = y_2$ with $y_1 \neq y_2$. Let V_1 and V_2 be disjoint open neighbourhoods in H , of y_1 and y_2 , respectively. There exists functions v_1 and v_2 in $S_0(H)$ which are supported in V_1 and V_2 , respectively, such that $v_1(y_1) \neq 0$ and $v_2(y_2) \neq 0$. Let $u_1, u_2 \in S_0(G)$ with $\Phi u_1 = v_1$ and $\Phi u_2 = v_2$. Then $0 \equiv v_1 \cdot v_2 = \Phi u_1 \cdot \Phi u_2 = \Phi(u_1 \cdot u_2)$ and so $u_1 \cdot u_2 \equiv 0$.

On the other hand, as $v_j(y_j) = \Phi u_j(y_j) \neq 0$ for $j = 1, 2$, we have by Condition (\star) that $u_j(x_0) \neq 0$ for $j = 1, 2$, which is in contradiction to $u_1 \cdot u_2 \equiv 0$.

Step 4. The map $\beta : G \rightarrow H$ is a bijection.

Proof of Step 4. The hypotheses of the theorem hold good for the map Φ^{-1} as well. Applying the preceding steps to the map Φ^{-1} gives rise to a well-defined function, say, $\alpha : H \rightarrow G$. Then $\alpha = \beta^{-1}$, proving that β is bijective.

Step 5. The map β is a homeomorphism of G onto H .

Proof of Step 5. First we prove that β is continuous. Suppose not. Then there exists $x \in G$, and a net (x_τ) in G with $x_\tau \rightarrow x$, but $\beta(x_\tau)$ does not converge to $\beta(x)$.

Let V be a neighbourhood of $\beta(x)$ in H , such that $\beta(x_\tau) \notin V$ for any τ . Let $w \in S_0(H)$ with $\text{Supp } w \subseteq V$, and $w[\beta(x)] = 1$. Let $v \in S_0(G)$ be such that $\Phi v = w$. Then $\beta(x_\tau) \notin \text{Supp } \Phi v$ for any τ , and so $x_\tau \notin \text{Supp } v$ for any τ . This gives $v(x_\tau) = 0$ for all τ , implying $v(x) = 0$, which is not possible by Condition (\star) , since $\Phi v[\beta(x)] = 1$.

We observe that the above argument holds good when the maps Φ and β are replaced with Φ^{-1} and β^{-1} respectively, yielding that the map $\beta : G \rightarrow H$ is a homeomorphism.

Since β is a homeomorphism, Condition (\star) gives:

$$\text{Supp } \Phi u = \beta(\text{Supp } u), \quad \text{for all } u \in S_0(G).$$

—

References

References

- [1] E.G. Effros, Z.-J. Ruan, *Operator Spaces*, London Math. Soc. Monogr. (N.S.), vol. **23**, Oxford Univ. Press, New York, 2000.
- [2] P. Eymard, L'algèbre de Fourier d'un groupe localement compact, Bull. Soc. Math. France **92** (1964), 181-236.
- [3] H. G. Feichtinger, *On a New Segal Algebra*, Monatsh. Math. **92**(1981), 269-289.
- [4] G.B. Folland, *A Course in Abstract Harmonic Analysis*, Studies in Advanced Mathematics, CRC Press, Boca Raton, FL, 1995.
- [5] M. S. Jakobsen, *On a (No Longer) New Segal Algebra: A Review of the Feichtinger Algebra*, J. Fourier Anal. Appl. **24** (2018), no. 6, 1579-1660.

- [6] E. Kaniuth, A.T. Lau and A. Ulger, *The Rajchman algebra $B_0(G)$ of a locally compact group G* , Bull. Sci. Math. **140** (2016), no. 3, 273-302.
- [7] R. Lakshmi Lavanya, *On the Fourier transform on function algebras on locally compact Abelian groups*, Monatsh. Math. (4) **184** (2017), 597-610.
- [8] H. Reiter and J. D. Stegeman, *Classical Harmonic Analysis and Locally Compact Groups*, LMS Monographs, New series **22**, Clarendon Press, Oxford, 2000.
- [9] N. Spronk, *Operator space structure on Feichtinger's Segal algebra*, J. Funct. Anal. **248** (2007), no. 1, 152-174.
- [10] M.E. Walter, *Group duality and isomorphisms of Fourier and Fourier-Stieltjes algebras from a W^* -algebra point of view*, Bull. Amer. Math. Soc. **76** (1970), 1321-1325.
- [11] J.G. Wendel, *Left centralizers and isomorphisms of group algebras*, Pacific J. Math. **2** (1952), 251-261.