LOCAL COMMUTANTS AND ULTRAINVARIANT SUBSPACES

JANKO BRAČIČ

ABSTRACT. For an operator A on a complex Banach space \mathfrak{X} and a closed subspace $\mathfrak{M} \subseteq \mathfrak{X}$, the local commutant of A at \mathfrak{M} is the set $\mathcal{C}(A; \mathfrak{M})$ of all operators T on \mathfrak{X} such that TAx = ATx for every $x \in \mathfrak{M}$. It is clear that $\mathcal{C}(A; \mathfrak{M})$ is a closed linear space of operators, however it is not an algebra, in general. For a given A, we show that $\mathcal{C}(A; \mathfrak{M})$ is an algebra if and only if the largest subspace \mathfrak{M}_A such that $\mathcal{C}(A; \mathfrak{M}) = \mathcal{C}(A; \mathfrak{M}_A)$ is invariant for every operator in $\mathcal{C}(A; \mathfrak{M})$. We say that these are ultrainvariant subspaces of A. For several types of operators we prove that there exist non-trivial ultrainvariant subspace is ultrainvariant. On the other hand, the lattice of all ultrainvariant subspaces of a non-zero nilpotent operator can be strictly smaller than the lattice of all hyperinvariant subspaces.

1. INTRODUCTION

Let \mathfrak{X} be a complex Banach space and let $B(\mathfrak{X})$ be the Banach algebra of all bounded linear operators on \mathfrak{X} . The dual space of \mathfrak{X} will be denoted by \mathfrak{X}^* . For $A \in B(\mathfrak{X})$ and a closed subspace $\mathfrak{M} \subseteq \mathfrak{X}$, the local commutant of A at \mathfrak{M} is $\mathfrak{C}(A; \mathfrak{M}) = \{T \in B(\mathfrak{X}); TAx = ATx \text{ for all } x \in \mathfrak{M}\}$. It is clear that (A)', the commutant of A, is a subset of $\mathfrak{C}(A; \mathfrak{M})$. Of course, $\mathfrak{C}(A; \mathfrak{X}) = (A)'$ and $\mathfrak{C}(A; \{0\}) = B(\mathfrak{X})$.

Local commutants were introduced by Larson [13, 6] more than twenty years ago, however there are only a few results related to them (see [4], for instance) although that there are some interesting open problems related to them. One among these problems is the question when is $\mathcal{C}(A; \mathcal{M})$ an algebra. Of course, for an arbitrary A and \mathcal{M} which is equal either to $\{0\}$ or \mathcal{X} the local commutant is an algebra. However, for A which is not a scalar multiple of $I_{\mathcal{X}}$, the identity operator on \mathcal{X} , there exists a closed subspace $\mathcal{M} \subseteq \mathcal{X}$ such that $\mathcal{C}(A; \mathcal{M})$ is not a subalgebra of $B(\mathcal{X})$ but just a linear subspace.

²⁰²⁰ Mathematics Subject Classification. Primary 47A15 47L05.

Key words and phrases. Commutant, local commutant, hyperinvariant subspace, ultrainvariant subspace.

Recall that a closed subspace $\mathcal{M} \subseteq \mathfrak{X}$ is said to be invariant for $A \in B(\mathfrak{X})$ if $A\mathcal{M} \subseteq \mathcal{M}$. It is well-known that the family of all closed subspaces which are invariant for A form a lattice; the lattice operations are the intersection and the closed linear span. As usual, we will denote this lattice by Lat(A). A sublattice of Lat(A) is $Lat_h(A)$, the lattice of all hyperinvariant subspaces of A: a subspace $\mathcal{M} \in \operatorname{Lat}(A)$ belongs to $\operatorname{Lat}_h(A)$ if and only if $T\mathcal{M} \subseteq \mathcal{M}$ for every $T \in (A)'$. It is obvious that the trivial lattice $\{\{0\}, \mathcal{X}\}$ is a sublattice of $\operatorname{Lat}_h(A)$ and therefore of $\operatorname{Lat}(A)$. In 1970s, Enflo showed that there exists a Banach space \mathfrak{X} and a bounded operator A on it such that Lat(A) is trivial. The result was published much later in [9] and meanwhile Read published his example of an operator without a non-trivial invariant subspace, see [17]. Nowadays several examples of Banach spaces (including l_1) with operators whose lattice of invariant subspaces is trivial are known. On the other hand, Lomonosov [15] proved that an operator has a non-trivial invariant subspace if it commutes with an operator which is not a scalar multiple of I_x and commutes with a non-zero compact operator. It follows that for some infinite dimensional Banach spaces \mathfrak{X} (see example given by Argyros and Haydon [2]) every operator in $B(\mathfrak{X})$ has a non-trivial invariant subspaces.

However for many Banach spaces (for instance, for the infinite dimensional separable Hilbert space) it is not known if Lat(A) is non-trivial for every operator A. Similarly, for more or less the same class of Banach spaces, it is not known if $Lat_h(A)$ can be trivial for an operator A which is not a scalar multiple of I_x . Work on these problems is still very active, see [20] and references therein. The reader is referred to [5] and [16] for more details about the problem of invariant subspaces.

It turns out that our problem about the algebraic structure of $\mathcal{C}(A; \mathcal{M})$ is connected with hyperinvariant subspaces of A. Our main observation is the following. Let $A \in B(\mathfrak{X})$ and let $\mathcal{M} \subseteq \mathfrak{X}$ be a closed subspace. Denote by $\mathcal{M}_A \subseteq \mathfrak{X}$ the set of all vectors $x \in \mathfrak{X}$ such that TAx = ATx for every $T \in \mathcal{C}(A; \mathcal{M})$. Then \mathcal{M}_A is a closed subspace of \mathfrak{X} such that $\mathcal{M} \subseteq \mathcal{M}_A$ and $\mathcal{C}(A; \mathcal{M}) = \mathcal{C}(A; \mathcal{M}_A)$. The local commutant $\mathcal{C}(A; \mathcal{M})$ is an algebra if and only if \mathcal{M}_A is invariant for every operator in $\mathcal{C}(A; \mathcal{M})$; we say that \mathcal{M}_A is an ultrainvariant subspace of A. Since $(A)' \subseteq \mathcal{C}(A; \mathcal{M})$ an ultrainvariant subspace is a hyperinvariant subspace, as well. The opposite is not true, in general.

Ultrainvariant subspaces are the main theme of the second part of this paper. Denote by $\operatorname{Lat}_u(A)$ the family of all ultrainvariant subspaces of A. It is a sublattice of $\operatorname{Lat}_h(A)$. For several types of operators A we show that $\operatorname{Lat}_u(A)$ is not trivial. There are classes of operators for which we are able to characterize the lattice of ultrainvariant subspaces completely. For instance, for algebraic operators, in particular, for operators on a finite dimensional Banach space, and for normal operators on a Hilbert space. If A is a unicellular operator on a Hilbert space, then $\operatorname{Lat}_u(A) = \operatorname{Lat}_u(A)$. This holds, for instance, for the Volterra operator and Donoghue operators.

Several questions about ultrainvariant subspaces remain open. For instance, we do not know if every compact operator has a non-trivial ultrainvariant subspace. Actually we believe that it is hard to find an operator which has a non-trivial hyperinvariant subspace but does not have a non-trivial ultrainvariant subspace.

In the end of this section, let us say a few words about the terminology. Sz.-Nagy and Foiaş initiated the study of hyperinvariant subspaces in their book Analyse harmonique des opérateurs de l'espace Hilbert however they called them ultrainvariant subspaces. In this paper we use their terminology (see Definition 4.1) for a special type of hyperinvariant subspaces. Another new term which we use is girder. The girder of $\mathcal{C}(A; \mathcal{M})$ is the largest subspace $\mathcal{M}_A \subseteq \mathfrak{X}$ such that $\mathcal{C}(A; \mathcal{M}) = \mathcal{C}(A; \mathcal{M}_A)$. In our opinion the term girder is suitable as \mathcal{M}_A "carries" the whole space $\mathcal{C}(A; \mathcal{M})$ in the sense that \mathcal{M}_A is the largest subspace whose vectors give the information whether T belongs to $\mathcal{C}(A; \mathcal{M})$ or not.

2. Local commutants

Although that we are mainly interested in local commutants we begin our study by considering a slightly more general objects. Let \mathfrak{X} and \mathfrak{Y} be complex Banach spaces and let $B(\mathfrak{X}, \mathfrak{Y})$ be the space of all bounded linear operators from \mathfrak{X} to \mathfrak{Y} . For $A \in B(\mathfrak{X})$ and $B \in B(\mathfrak{Y})$, let $\mathfrak{I}(A, B)$ be the set of all operators $S \in B(\mathfrak{X}, \mathfrak{Y})$ which intertwine A and B, that is, $\mathfrak{I}(A, B) = \{S \in B(\mathfrak{X}, \mathfrak{Y}); SA = BS\}$. Of course, if $\mathfrak{X} = \mathfrak{Y}$ and A = B, then $\mathfrak{I}(A, A) = (A)'$. Let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. The space of operators which intertwine A and B locally at \mathfrak{M} is $\mathfrak{I}(A, B; \mathfrak{M}) = \{S \in B(\mathfrak{X}, \mathfrak{Y}); SAx = BSx \text{ for every } x \in \mathfrak{M}\}$. In particular, $\mathfrak{I}(A, A; \mathfrak{M}) = \mathfrak{C}(A; \mathfrak{M})$ is the local commutant of A at \mathfrak{M} .

It is easily seen that $\mathcal{I}(A, B; \mathcal{M})$ is closed in the strong operator topology and that $\mathcal{I}(A, B; \mathcal{M}) = \mathcal{I}(A - \lambda I_x, B - \lambda I_y; \mathcal{M})$, for every $\lambda \in \mathbb{C}$. For arbitrary invertible operators $U \in B(\mathfrak{X})$ and $V \in B(\mathfrak{Y})$, we have

(1)
$$\mathfrak{I}(A,B;\mathfrak{M}) = V^{-1}\mathfrak{I}(UAU^{-1},VBV^{-1};U\mathfrak{M})U.$$

If $\mathcal{M}_1, \mathcal{M}_2$ are closed subspaces of \mathfrak{X} such that $\mathcal{M}_2 \subseteq \mathcal{M}_1$, then $\mathfrak{I}(A, B; \mathcal{M}_1) \subseteq \mathfrak{I}(A, B; \mathcal{M}_2)$. In particular, $\mathfrak{I}(A, B) = \mathfrak{I}(A, B; \mathfrak{X}) \subseteq \mathfrak{I}(A, B; \mathfrak{M}) \subseteq \mathfrak{I}(A, B; \{0\}) = B(\mathfrak{X}, \mathfrak{Y})$.

Let $\{\mathcal{M}_j; j \in J\}$ be a non-empty family of closed subspaces of \mathcal{X} . It is not hard to see that

$$\bigvee_{j \in J} \{ \mathfrak{I}(A, B; \mathcal{M}_j); \ j \in J \} \subseteq \mathfrak{I}(A, B; \bigcap_{j \in J} \mathcal{M}_j).$$

On the other hand,

(2)
$$\mathbb{J}(A, B; \bigvee_{j \in J} \mathfrak{M}_j) = \bigcap_{j \in J} \mathbb{J}(A, B; \mathfrak{M}_j).$$

Here we denoted by \bigvee the closed linear span of the involved sets.

To verify (2), note that $\mathcal{I}(A, B; \bigvee_{j \in J} \mathcal{M}_j) \subseteq \bigcap_{j \in J} \mathcal{I}(A, B; \mathcal{M}_j)$ holds since $\mathcal{M}_i \subseteq \bigvee_{j \in J} \mathcal{M}_j$ and therefore $\mathcal{I}(A, B; \bigvee_{j \in J} \mathcal{M}_j) \subseteq \mathcal{I}(A, B; \mathcal{M}_i)$, for every $i \in J$. For the opposite inclusion, observe that $BSx_j = SAx_j$ for every $x_j \in \mathcal{M}_j$ and every $j \in J$ if $S \in \bigcap_{j \in J} \mathcal{I}(A, B; \mathcal{M}_j)$. Hence $S \in \mathcal{I}(A, B; \bigvee_{j \in J} \mathcal{M}_j)$. (Here we used a simple fact that BSx = SAx holds for every $x \in \bigvee \mathcal{U}$ if it holds for every $x \in \mathcal{U} \subseteq \mathcal{X}$.)

Let $A \in B(\mathfrak{X})$, $B \in B(\mathfrak{Y})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a complemented closed subspace, that is, there exists a closed subspace $\mathfrak{N} \subseteq \mathfrak{X}$ such that $\mathfrak{X} = \mathfrak{M} \oplus \mathfrak{N}$. Let $\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$ be the operator matrix of A with respect to this decomposition. For arbitrary operators $S \in B(\mathfrak{X}, \mathfrak{Y})$ and $T \in B(\mathfrak{X})$, let $[S_1 S_2]$, respectively $\begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$, be their operator matrices with respect to the decomposition $\mathfrak{X} = \mathfrak{M} \oplus \mathfrak{N}$. These assumptions and notation will be used throughout this section.

A straightforward computation shows that $S \in \mathcal{I}(A, B)$ if and only if

(3)
$$BS_1 = S_1 A_{11} + S_2 A_{21}$$

and

(4)
$$BS_2 = S_1 A_{12} + S_2 A_{22}.$$

Similarly, $T \in (A)'$ if and only if

(5)
$$T_{11}A_{11} + T_{12}A_{21} = A_{11}T_{11} + A_{12}T_{21}, \quad T_{21}A_{11} + T_{22}A_{21} = A_{21}T_{11} + A_{22}T_{21}$$

and

(6)
$$T_{11}A_{12} + T_{12}A_{22} = A_{11}T_{12} + A_{12}T_{22}, \quad T_{21}A_{12} + T_{22}A_{22} = A_{21}T_{12} + A_{22}T_{22}.$$

Lemma 2.1. Operator $S \in B(\mathfrak{X}, \mathfrak{Y})$ is in $\mathfrak{I}(A, B; \mathfrak{M})$ if and only if (3) holds and $T \in B(\mathfrak{X})$ is in $\mathfrak{C}(A; \mathfrak{M})$ if and only if (5) holds.

Proof. Let $x \in \mathfrak{X}$ be arbitrary and let $x = x_1 \oplus x_2$, where $x_1 \in \mathfrak{M}$ and $x_2 \in \mathfrak{N}$. Then $SAx = (S_1A_{11} + S_2A_{21})x_1 + (S_1A_{12} + S_2A_{22})x_2$ and $BSx = BS_1x_1 + BS_2x_2$. Hence, if $S \in \mathfrak{I}(A, B; \mathfrak{M})$, that is, $BS(x_1 \oplus 0) = SA(x_1 \oplus 0)$ for every $x_1 \in \mathfrak{M}$, then $BS_1x_1 = (S_1A_{11} + S_2A_{21})x_1$ for every $x_1 \in \mathfrak{M}$ which gives (3). It is clear that the opposite implication holds, as well. The second part of the assertion follows by the first one if we put $B = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$, $S_1 = \begin{bmatrix} T_{11} \\ T_{21} \end{bmatrix}$ and $S_2 = \begin{bmatrix} T_{12} \\ T_{22} \end{bmatrix}$ in (3).

It is clear that $\mathcal{I}(A, B; \{0\}) = B(\mathfrak{X}, \mathfrak{Y})$ for arbitrary operators A and B. Let $\mathfrak{M} \neq \{0\}$. The following corollary characterizes those pairs of operators $A \in B(\mathfrak{X})$ and $B \in B(\mathfrak{Y})$ for which $\mathcal{I}(A, B; \mathfrak{M}) = B(\mathfrak{X}, \mathfrak{Y})$.

Proposition 2.2. Let $\mathfrak{M} \neq \{0\}$. Then $\mathfrak{I}(A, B; \mathfrak{M}) = B(\mathfrak{X}, \mathfrak{Y})$ if and only if $A_{11} = \lambda I_{\mathfrak{M}}$, $B = \lambda I_{\mathfrak{Y}}$, for some $\lambda \in \mathbb{C}$, and $A_{21} = 0$. In particular, $\mathfrak{C}(A; \mathfrak{M}) = B(\mathfrak{X})$ if and only if A is a scalar multiple of $I_{\mathfrak{X}}$.

Proof. Assume that $\mathcal{I}(A, B; \mathcal{M}) = B(\mathfrak{X}, \mathfrak{Y})$. Let $S = [S_1 0]$, where $S_1 \in B(\mathcal{M}, \mathfrak{Y})$ is arbitrary. By Lemma 2.1, $BS_1 = S_1A_{11}$. Let $y \in \mathcal{Y}$ and $0 \neq \xi \in \mathcal{M}^*$ be arbitrary. Then $y \otimes \xi \in B(\mathcal{M}, \mathfrak{Y})$ and therefore $B(y \otimes \xi) = (y \otimes \xi)A_{11}$. If $x \in \mathcal{M}$ is such that $\langle x, \xi \rangle = 1$, then $By = B(y \otimes \xi)x = (y \otimes \xi)A_{11}x = \langle A_{11}x, \xi \rangle y$. Hence, $B \in B(\mathfrak{Y})$ is an operator such that every $0 \neq y \in \mathcal{Y}$ is its eigenvector. It follows that there exists $\lambda \in \mathbb{C}$ such that $B = \lambda I_{\mathfrak{Y}}$. Now we have $\lambda S_1 = S_1A_{11}$ for every $S_1 \in B(\mathcal{M}, \mathfrak{Y})$ which gives $A_{11} = \lambda I_{\mathfrak{M}}$. Since every $S = [S_1 S_2] \in B(\mathfrak{X}, \mathfrak{Y})$ satisfies (3) we have $S_2A_{21} = 0$ for every $S_2 \in B(\mathcal{N}, \mathfrak{Y})$. Hence, $A_{21} = 0$. The opposite implication is simple: if $A_{11} = \lambda I_{\mathfrak{M}}$, $B = \lambda I_{\mathfrak{Y}}$ and $A_{21} = 0$, then (3) holds for every $S = [S_1 S_2]$.

It follows from the first part of this corollary that in the case when $\mathfrak{X} = \mathfrak{Y}$ and A = Bwe have $\mathfrak{C}(A; \mathfrak{M}) = B(\mathfrak{X})$ if and only if $A = \lambda I_{\mathfrak{X}}$ for some $\lambda \in \mathbb{C}$.

Equality $\mathcal{I}(A, B; \mathcal{M}) = \mathcal{I}(A, B)$ holds if and only if every pair of operators $S_1 \in B(\mathcal{M}, \mathcal{Y})$ and $S_2 \in B(\mathcal{N}, \mathcal{Y})$ which satisfies (3) satisfies (4), as well. In the following proposition we determine those pairs $A \in B(\mathcal{X})$ and $B \in B(\mathcal{Y})$ for which $\mathcal{I}(A, B; \mathcal{M}) = \mathcal{I}(A, B)$ holds for a given subspace $\mathcal{M} \in \text{Lat}(A), \ \mathcal{M} \neq \mathcal{X}$. We will use the following notation. For $S \in B(\mathcal{X}, \mathcal{Y})$, the kernel of S is denoted by ker(S) and the range (image) of S by im(S).

If \mathcal{Z} is a non-empty set of vectors in \mathcal{X} , then by \mathcal{Z}^{\perp} we denote its annihilator in \mathcal{X}^* , that is, $\mathcal{Z}^{\perp} = \{\xi \in \mathcal{X}^*; \langle x, \xi \rangle = 0 \text{ for all } x \in \mathcal{Z} \}.$

Proposition 2.3. Let $\mathfrak{M} \neq \mathfrak{X}$. Then $\mathfrak{I}(A, B; \mathfrak{M}) = \mathfrak{I}(A, B)$ and $\mathfrak{M} \in \operatorname{Lat}(A)$ if and only if there exists $\lambda \in \mathbb{C}$ such that $A = \begin{bmatrix} A_{11} & A_{12} \\ 0 & \lambda I_{\mathfrak{N}} \end{bmatrix}$, $B = \lambda I_{\mathfrak{Y}}$ and $\overline{\operatorname{im}(A_{12})} \subseteq \overline{\operatorname{im}(\lambda I_{\mathfrak{M}} - A_{11})}$. In this case, $\mathfrak{I}(A, B; \mathfrak{M}) = \{ [S_1 S_2] \in B(\mathfrak{X}, \mathfrak{Y}); \overline{\operatorname{im}(\lambda I_{\mathfrak{M}} - A_{11})} \subseteq \operatorname{ker}(S_1), S_2 \in B(\mathfrak{N}, \mathfrak{Y}) \}$.

Proof. Assume that $\mathcal{I}(A, B; \mathcal{M}) = \mathcal{I}(A, B)$, where $\mathcal{X} \neq \mathcal{M} \in \text{Lat}(A)$. Then the operator matrix of A with respect to the decomposition $\mathcal{X} = \mathcal{M} \oplus \mathcal{N}$ is $A = \begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$. Hence, by Lemma 2.1, an operator $S = [S_1 S_2] \in B(\mathcal{X}, \mathcal{Y})$ intertwines A and B at \mathcal{M} if and only if it satisfies (3) which in our case means

$$BS_1 = S_1 A_{11}$$

It follows that $[0 S_2] \in \mathcal{I}(A, B; \mathcal{M})$, and therefore $[0 S_2] \in \mathcal{I}(A, B)$, for every $S_2 \in B(\mathcal{N}, \mathcal{Y})$. Thus, $[0 S_2]$ satisfies (4), that is, $BS_2 = S_2A_{22}$ holds for every $S_2 \in B(\mathcal{N}, \mathcal{Y})$. By Proposition 2.2, there exists $\lambda \in \mathbb{C}$ such that $A_{22} = \lambda I_{\mathcal{N}}$ and $B = \lambda I_{\mathcal{Y}}$. Now (7) reads as

(8)
$$S_1(\lambda I_{\mathcal{M}} - A_{11}) = 0.$$

Hence, if $[S_1 S_2]$ is an operator in $\mathcal{I}(A, \lambda I_y; \mathcal{M})$, then $\overline{\mathrm{im}(\lambda I_M - A_{11})} \subseteq \ker(S_1)$ and S_2 is arbitrary. Note that $[S_1 S_2]$ is in $\mathcal{I}(A, \lambda I_y)$ if and only if it satisfies (4) which reads as

(9)
$$S_1 A_{11} = 0$$

in our case. Since $\mathcal{I}(A, \lambda I_{y}; \mathcal{M}) = \mathcal{I}(A, \lambda I_{y})$ every S_{1} which satisfies (8) has to satisfy (9), as well. Suppose towards a contradiction that there exists $x = A_{12}u \in \operatorname{im}(A_{12})$ such that $x \notin \operatorname{im}(\lambda I_{\mathcal{M}} - A_{11})$. Let $\xi \in \mathcal{M}^{*}$ be such that $\xi \in \operatorname{im}(\lambda I_{\mathcal{M}} - A_{11})^{\perp}$ and $\langle x, \xi \rangle = 1$. For any $0 \neq y \in \mathcal{Y}$, the rank one operator $y \otimes \xi \in B(\mathcal{M}, \mathcal{Y})$ satisfies $(y \otimes \xi)(\lambda I_{\mathcal{M}} - A_{11}) = 0$, which means $y \otimes \xi \in \mathcal{I}(A, \lambda I_{y}; \mathcal{M})$, however $(y \otimes \xi)A_{12} \neq 0$ since $(y \otimes \xi)A_{12}u = \langle x, \xi \rangle y \neq 0$. This proves that $\operatorname{im}(A_{12}) \subseteq \operatorname{im}(\lambda I_{\mathcal{M}} - A_{11})$.

The opposite implication is easily checked and one gets that $\mathcal{I}(A, B; \mathcal{M}) = \{[S_1 S_2] \in B(\mathfrak{X}, \mathfrak{Y}); \overline{\mathrm{im}(\lambda I_{\mathfrak{M}} - A_{11})} \subseteq \mathrm{ker}(S_1), S_2 \in B(\mathfrak{N}, \mathfrak{Y})\}.$

Corollary 2.4. If $\mathfrak{M} \in \operatorname{Lat}(A)$, $\mathfrak{M} \neq \mathfrak{X}$, and A is not a scalar multiple of $I_{\mathfrak{X}}$, then $(A)' \subsetneq \mathfrak{C}(A; \mathfrak{M})$.

Theorem 2.5. Let $A \in B(\mathfrak{X})$ and assume that A is not a scalar multiple of $I_{\mathfrak{X}}$. Let $\mathfrak{M} \in \operatorname{Lat}(A), \{0\} \neq \mathfrak{M} \neq \mathfrak{X}$. The local commutant $\mathfrak{C}(A; \mathfrak{M})$ is an algebra if and only if $\mathfrak{C}(A; \mathfrak{M}) = \{T \in B(\mathfrak{X}); T_{11} \in (A_{11})' \text{ and } T_{21} = 0\}.$

Proof. We begin the proof with the following simple observation.

Claim. Let $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3$ and \mathfrak{X}_4 be Banach spaces. If $T \in B(\mathfrak{X}_1, \mathfrak{X}_2)$ and $R \in B(\mathfrak{X}_3, \mathfrak{X}_4)$ are such that $R(u \otimes \xi)T = 0$ for all $u \in \mathfrak{X}_3$ and $\xi \in \mathfrak{X}_2^*$, then either R = 0 or T = 0.

Indeed! Assume that $T \neq 0$. Then there exists $x \in \mathfrak{X}_1$ such that $Tx \neq 0$. Let $\xi \in \mathfrak{X}_2^*$ be such that $\langle Tx, \xi \rangle = 1$. For an arbitrary $u \in \mathfrak{X}_3$ we have, by the assumption, $R(u \otimes \xi)T = 0$ and therefore $R(u \otimes \xi)Tx = Ru = 0$. This shows that R = 0.

Assume that $\mathcal{C}(A; \mathcal{M})$ is an algebra. Since $\mathcal{M} \in \text{Lat}(A)$ the operator matrix of A is of the form $\begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$. Hence, by Lemma 2.1, an operator $T = \begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$ is in $\mathcal{C}(A; \mathcal{M})$ if and only if

(10)
$$T_{11}A_{11} - A_{11}T_{11} = A_{12}T_{21}, \quad T_{21}A_{11} - A_{22}T_{21} = 0$$

and $T_{12} \in B(\mathcal{N}, \mathcal{M}), T_{22} \in B(\mathcal{N})$ are arbitrary. In particular, the projection P onto \mathcal{M} along \mathcal{N} is in $\mathcal{C}(A; \mathcal{M})$. Since $\mathcal{C}(A; \mathcal{M})$ is assumed to be an algebra we see that operators PTP and $(I_x - P)TP$ are in $\mathcal{C}(A; \mathcal{M})$ for every $T \in \mathcal{C}(A; \mathcal{M})$. Note that the operator matrices of PTP and $(I_x - P)TP$ are $\begin{bmatrix} T_{11} & 0\\ 0 & 0 \end{bmatrix}$ and $\begin{bmatrix} 0 & 0\\ T_{21} & 0 \end{bmatrix}$, respectively. Hence, if $T \in \mathcal{C}(A; \mathcal{M})$, then the entries of its operator matrix have to satisfy

(11)
$$T_{11}A_{11} - A_{11}T_{11} = 0, \quad A_{12}T_{21} = 0, \quad T_{21}A_{11} - A_{22}T_{21} = 0.$$

Suppose that $A_{12} \neq 0$. Let $u \in \mathbb{N}$ and $\xi \in \mathbb{N}^*$ be arbitrary. Then U whose operator matrix is $\begin{bmatrix} 0 & 0\\ 0 & u \otimes \xi \end{bmatrix}$ satisfies (10) and therefore $U \in \mathcal{C}(A; \mathcal{M})$ which gives $U(I_x - P)TP \in \mathcal{C}(A; \mathcal{M})$. The operator matrix of $U(I_x - P)TP$ is $\begin{bmatrix} 0 & 0\\ (u \otimes \xi)T_{21} & 0 \end{bmatrix}$ and the entries of this matrix satisfy (11); in particular, $A_{12}(u \otimes \xi)T_{21} = 0$. Since $u \otimes \xi \in B(\mathcal{N})$ is an arbitrary operator of rank at most 1 and $A_{12} \neq 0$ we conclude, by Claim, that $T_{21} = 0$.

Now we may assume that $A_{12} = 0$. If $A_{11} = \lambda I_{\mathcal{M}}$ and $A_{22} = \mu I_{\mathcal{N}}$, then $\lambda \neq \mu$ as A is not a scalar multiple of I_x . It follows, by the third equality in (11), that $T_{21} = 0$. Thus, it remains to consider the case when either A_{11} is not a scalar multiple of $I_{\mathcal{M}}$ or A_{22} is not a scalar multiple of $I_{\mathcal{N}}$. Suppose that this holds. Let $Q \in B(\mathcal{N}, \mathcal{M})$ be arbitrary and let $V \in B(\mathfrak{X})$ be the operator whose operator matrix is $\begin{bmatrix} 0 & Q \\ 0 & 0 \end{bmatrix}$. Then $V \in \mathbb{C}(A; \mathcal{M})$ and therefore $V(I_x - P(TP \in \mathbb{C}(A; \mathcal{M}))$ for every $T \in \mathbb{C}(A; \mathcal{M})$.

operator matrix of $V(I_x - P(TP \in C(A; \mathcal{M}) \text{ is } \begin{bmatrix} QT_{21} & 0 \\ 0 & 0 \end{bmatrix}$. Hence, $QT_{21} \in (A_{11})'$. Since T_{21} satisfies the third equality in (11) we have $A_{11}QT_{21} = QT_{21}A_{11} = QA_{22}T_{21}$ which gives $(A_{11}Q - QA_{22})T_{21} = 0$. We have supposed that either A_{11} is not a scalar multiple of $I_{\mathcal{M}}$ or A_{22} is not a scalar multiple of $I_{\mathcal{N}}$. Hence, by Proposition 2.2, there exists $Q_0 \in B(\mathcal{N}, \mathcal{M})$ such that $R = A_{11}Q_0 - Q_0A_{22} \neq 0$. Thus, $RT_{21} = 0$ for every $T \in C(A; \mathcal{M})$. Let T be fixed now. As in the previous paragraph, let $u \in \mathcal{N}$ and $\xi \in \mathcal{N}^*$ be arbitrary and let $U \in C(A; \mathcal{M})$ be the operator whose operator matrix is $\begin{bmatrix} 0 & 0 \\ 0 & u \otimes \xi \end{bmatrix}$. Then $U(I_x - P)TP \in$ $C(A; \mathcal{M})$ and therefore $R(u \otimes \xi)T_{21} = 0$. Since $u \otimes \xi \in B(\mathcal{N})$ is an arbitrary operator of rank at most 1 and $R \neq 0$ we conclude, by Claim, that $T_{21} = 0$. This proves that $C(A; \mathcal{M}) = \{T \in B(\mathcal{X}); T_{11} \in (A_{11})'$ and $T_{21} = 0\}$ if $C(A; \mathcal{M})$ is assumed to be an algebra. The opposite implication is obvious.

In the proof of Theorem 2.5 we have seen that the projection P which maps onto \mathcal{M} along \mathcal{N} is in $\mathcal{C}(A; \mathcal{M})$ if $\mathcal{M} \in \text{Lat}(A)$. It is not hard to see that the opposite implication holds as well.

Corollary 2.6. Projection P is in $\mathcal{C}(A; \mathcal{M})$ if and only if \mathcal{M} is invariant for A.

Let $\sigma(T)$ denote the spectrum of $T \in B(\mathfrak{X})$. It follows from a well-known theorem proved by Rosenblum that $\mathcal{I}(A, B) = \{0\}$ if $\sigma(A) \cap \sigma(B) = \emptyset$, see [18, Corollary 3.3] and [10, Theorem I.4.1]. Note that the condition $\sigma(A) \cap \sigma(B) = \emptyset$ is not necessary for $\mathcal{I}(A, B) = \{0\}$. For instance, let B = 0 and let $A \in B(\mathfrak{X})$ be such that $\overline{\mathrm{im}(A)} = \mathfrak{X}$ and ker $(A) \neq \{0\}$. Then $0 \in \sigma(A) \cap \sigma(B)$, however $\mathcal{I}(A, 0) = \{0\}$ because SA = 0 holds for $S \in B(\mathfrak{X}, \mathfrak{Y})$ if and only if S = 0 as $\overline{\mathrm{im}(A)} = \mathfrak{X}$. On the other hand, note that $\mathcal{I}(0, A) \neq \{0\}$. Indeed, if $S \in B(\mathfrak{Y}, \mathfrak{X})$ is such that $\mathrm{im}(S) \subseteq \mathrm{ker}(A)$, then AS = 0.

Corollary 2.7. If $\mathcal{M} \in \text{Lat}(A)$ and $\sigma(A_{11}) \cap \sigma(A_{22}) = \emptyset$, then $\mathcal{C}(A; \mathcal{M})$ is an algebra.

Proof. By Lemma 2.1, $T \in \mathcal{C}(A; \mathcal{M})$ if and only if (5) holds. Since $A_{21} = 0$ the equalities in (5) simplify to (10). Because of $\sigma(A_{11}) \cap \sigma(A_{22}) = \emptyset$, only $T_{21} = 0$ satisfies the second equation in (10). Thus, $\mathcal{C}(A; \mathcal{M}) = \{T \in B(\mathcal{X}); T_{11} \in (A_{11})' \text{ and } T_{21} = 0\}$ and therefore it is an algebra, by Theorem 2.5.

In the following example we will see that $\mathcal{C}(Q; \mathcal{L})$ can be an algebra for $Q \in B(\mathfrak{X})$ and a closed subspace $\mathcal{L} \subseteq \mathfrak{X}$ which is not invariant for Q. **Example 2.8.** Let $\mathfrak{X}_1, \mathfrak{X}_2, \mathfrak{X}_3$ be closed subspaces such that $\mathfrak{X}_1, \mathfrak{X}_2$ are non-trivial. Let $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2 \oplus \mathfrak{X}_3$ and suppose that there exists a surjective operator in $B(\mathfrak{X}_1, \mathfrak{X}_2)$. Let $Q \in B(\mathfrak{X})$ be the projection onto \mathfrak{X}_1 along $\mathfrak{X}_2 \oplus \mathfrak{X}_3$. Choose and fix a surjective operator $U \in B(\mathfrak{X}_1, \mathfrak{X}_2)$. Let $\mathcal{L} \subseteq \mathfrak{X}$ be the subspace of all vectors $x \in \mathfrak{X}$ which are of the form $x = x_1 \oplus Ux_1 \oplus x_3$, where $x_1 \in \mathfrak{X}_1$ and $x_3 \in \mathfrak{X}_3$ are arbitrary. Denote by $P_j : \mathfrak{X} \to \mathfrak{X}_j$ (j = 1, 2, 3) the operators given by $P_j(x_1 \oplus x_2 \oplus x_3) = x_j$. It is clear that $P_1\mathcal{L} = \mathfrak{X}_1$ and $P_3\mathcal{L} = \mathfrak{X}_3$. Since U is surjective we have $P_2\mathcal{L} = \mathfrak{X}_2$, as well. Note that $\mathcal{L} \notin \text{Lat}(Q)$. Namely, if $x = x_1 \oplus Ux_1 \oplus x_3 \in \mathcal{L}$ is such that $Ux_1 \neq 0$, then $Qx = x_1 \oplus 0 \oplus 0 \notin \mathcal{L}$.

Assume that $T \in \mathcal{C}(Q; \mathcal{L})$. Let $[T_{ij}]_{i,j=1}^3$ be its operator matrix with respect to the decomposition $\mathcal{X} = \mathcal{X}_1 \oplus \mathcal{X}_2 \oplus \mathcal{X}_3$. For an arbitrary $x = x_1 \oplus Ux_1 \oplus x_3 \in \mathcal{L}$, we have $TQx = T_{11}x_1 \oplus T_{21}x_1 \oplus T_{31}x_1$ and $QTx = (T_{11}x_1 + T_{12}Ux_1 + T_{13}x_3) \oplus 0 \oplus 0$. Since $x_1 \in \mathcal{X}_1$ and $x_3 \in \mathcal{X}_3$ are arbitrary and U is surjective it follows from TQx = QTx that $T_{21} = 0$, $T_{31} = 0$, $T_{12} = 0$ and $T_{13} = 0$. Hence $T \in (Q)'$, that is $\mathcal{C}(Q; \mathcal{L}) = (Q)'$; in particular, $\mathcal{C}(Q; \mathcal{L})$ is an algebra.

Note that this example shows that $\mathcal{C}(A; \mathcal{M})$ and (A)' can be equal for an operator which is not a scalar multiple of I_x and $\mathcal{M} \neq \mathfrak{X}$ which is not invariant for A (cf. Corollary 2.4). \Box

Although that the condition $\mathcal{M} \in \text{Lat}(A)$ does not need to be fulfilled for $\mathcal{C}(A; \mathcal{M})$ being an algebra the invariance is not far away as we shall see in the following section.

3. Multiplicative structure of the space of local intertwiners

Let $A \in B(\mathfrak{X})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. We know that $\mathcal{C}(A; \mathfrak{M})$ is not an algebra in general. However it has some non-trivial multiplicative structure. For instance, let $T \in (A)'$ be such that $\mathfrak{M} \in \operatorname{Lat}(T)$. Then ASTx = SATx = STAx for all $S \in \mathcal{C}(A; \mathfrak{M})$ and $x \in \mathfrak{M}$. Hence, $\mathcal{C}(A; \mathfrak{M})T \subseteq \mathcal{C}(A; \mathfrak{M})$. The following proposition is a generalization of this simple observation.

Proposition 3.1. Let $A \in B(\mathfrak{X})$, $B \in B(\mathfrak{Y})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. Inclusion $\mathfrak{I}(A, B; \overline{T\mathfrak{M}})T \subseteq \mathfrak{I}(A, B; \mathfrak{M})$ holds for every $T \in \mathfrak{C}(A; \mathfrak{M})$. If T is invertible, then $\mathfrak{I}(A, B; \overline{T\mathfrak{M}})T = \mathfrak{I}(A, B; \mathfrak{M})$.

Proof. Let $T \in \mathcal{C}(A; \mathcal{M})$. For arbitrary $S \in \mathcal{I}(A, B; \overline{T\mathcal{M}})$ and $x \in \mathcal{M}$, we have STAx = SATx = BSTx and therefore $ST \in \mathcal{I}(A, B; \mathcal{M})$.

Assume now that $T \in \mathcal{C}(A; \mathcal{M})$ is invertible. Let $S \in \mathcal{I}(A, B; \mathcal{M})$ be arbitrary. Then for every $x \in \mathcal{M}$ we have $ST^{-1}ATx = SAx = BSx = BST^{-1}Tx$ which implies that $ST^{-1} \in \mathcal{I}(A, B; \overline{T\mathcal{M}})$, that is, $S \in \mathcal{I}(A, B; \overline{T\mathcal{M}})T$.

For a non-empty set of operators $S \subseteq B(\mathfrak{X}, \mathfrak{Y})$ and a closed subspace $\mathcal{M} \subseteq \mathfrak{X}$, let $S\mathcal{M}$ denote the closed linear span in \mathfrak{Y} of all subspaces $S\mathcal{M}$, where $S \in S$, that is, $S\mathcal{M} = \bigvee_{S \in S} S\mathcal{M}$.

Theorem 3.2. Let $A \in B(\mathfrak{X})$, $B \in B(\mathfrak{Y})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. Then $\mathcal{C}(B; \mathfrak{I}(A, B; \mathfrak{M})\mathfrak{M})$ is a subalgebra of $B(\mathfrak{Y})$ and it is the largest subalgebra of $B(\mathfrak{Y})$ such that $\mathfrak{I}(A, B; \mathfrak{M})$ is a left module over it.

Proof. Let us show that $\mathcal{C}(B; \mathcal{I}(A, B; \mathcal{M})\mathcal{M}) \cdot \mathcal{I}(A, B; \mathcal{M}) \subseteq \mathcal{I}(A, B; \mathcal{M})$. Assume that $T \in \mathcal{C}(B; \mathcal{I}(A, B; \mathcal{M})\mathcal{M})$ and $S \in \mathcal{I}(A, B; \mathcal{M})$. For every $x \in \mathcal{M}$, we have BSx = SAx and therefore BTSx = TBSx = TSAx because $Sx \in \mathcal{I}(A, B; \mathcal{M})\mathcal{M}$. We conclude that $TS \in \mathcal{I}(A, B; \mathcal{M})$.

Let $T_1, T_2 \in \mathcal{C}(B; \mathfrak{I}(A, B; \mathfrak{M})\mathfrak{M})$. If $S \in \mathfrak{I}(A, B; \mathfrak{M})$, then $T_2S \in \mathfrak{I}(A, B; \mathfrak{M})$, by the inclusion from the previous paragraph, and, by the same reason, $T_1T_2S \in \mathfrak{I}(A, B; \mathfrak{M})$. Hence, for every $S \in \mathfrak{I}(A, B; \mathfrak{M})$ and every $x \in \mathfrak{M}$, we have $BT_1T_2Sx = T_1T_2SAx = T_1T_2BSx$. Because of linearity and countinuity we may conclude that T_1T_2 commutes with B on the subspace $\mathfrak{I}(A, B; \mathfrak{M})\mathfrak{M}$ of \mathfrak{Y} , that is, $T_1T_2 \in \mathcal{C}(B; \mathfrak{I}(A, B; \mathfrak{M})\mathfrak{M})$. Thus, $\mathcal{C}(B; \mathfrak{I}(A, B; \mathfrak{M})\mathfrak{M})$ is a subalgebra of $\mathcal{B}(\mathfrak{Y})$ and $\mathfrak{I}(A, B; \mathfrak{M})$ is a left module over it.

Assume that $T \in B(\mathcal{Y})$ is such that $T\mathcal{I}(A, B; \mathcal{M}) \subseteq \mathcal{I}(A, B; \mathcal{M})$. Let $S \in \mathcal{I}(A, B; \mathcal{M})$ and $x \in \mathcal{M}$ be arbitrary. Then BTSx = TSAx = TBSx which implies $T \in \mathcal{C}(B; \mathcal{I}(A, B; \mathcal{M})\mathcal{M})$. This shows that $\mathcal{C}(B; \mathcal{I}(A, B; \mathcal{M})\mathcal{M})$ is the largest subalgebra of $B(\mathcal{Y})$ such that $\mathcal{I}(A, B; \mathcal{M})$ is a left module over it.

Corollary 3.3. Let $A \in B(\mathfrak{X})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. Then $\mathfrak{C}(A; \mathfrak{C}(A; \mathfrak{M})\mathfrak{M})$ is the largest algebra contained in $\mathfrak{C}(A; \mathfrak{M})$. If $\mathfrak{M} \in \mathrm{Lat}(\mathfrak{C}(A; \mathfrak{M}))$, then $\mathfrak{C}(A; \mathfrak{M})$ is an algebra.

Proof. By Theorem 3.2, $\mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})$ is the largest algebra contained in $\mathcal{C}(A; \mathcal{M})$. If $\mathcal{M} \in \operatorname{Lat}(\mathcal{C}(A; \mathcal{M}))$, then $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathcal{M}$ and therefore $\mathcal{C}(A; \mathcal{M})$ is an algebra.

Now we consider the right module structure of $\mathcal{J}(A, B; \mathcal{M})$. Let $A \in B(\mathfrak{X}), B \in B(\mathfrak{Y})$ and let $\mathcal{S} \subseteq B(\mathfrak{X}, \mathfrak{Y})$ be an arbitrary set of operators. It is clear that every $S \in \mathcal{S}$

intertwines A and B at vector 0, that is, $S \subseteq \mathfrak{I}(A, B; \{0\})$. Hence, the family \mathcal{F} of all closed subspaces $\mathcal{M} \subseteq \mathfrak{X}$ such that $S \subseteq \mathfrak{I}(A, B; \mathcal{M})$ is not empty. Let $\mathcal{M}_S = \bigvee_{\mathcal{M} \in \mathcal{F}} \mathcal{M}$. By (2), $S \subseteq \bigcap_{\mathcal{M} \in \mathcal{F}} \mathfrak{I}(A, B; \mathcal{M}) = \mathfrak{I}(A, B; \mathcal{M}_S)$, that is, $\mathcal{M}_S \in \mathcal{F}$. Of course, \mathcal{M}_S is the largest subspace in \mathcal{F} . In particular, there exists the largest subspace $\mathcal{M}_{\mathfrak{I}(A,B;\mathcal{M})}$ of \mathfrak{X} on which all operators from $\mathfrak{I}(A, B; \mathcal{M})$ intertwine A and B. We will simplify the notation to $\mathcal{M}_{A,B}$ instead of $\mathcal{M}_{\mathfrak{I}(A,B;\mathcal{M})}$ (and \mathcal{M}_A instead of $\mathcal{M}_{A,A}$ if $\mathfrak{X} = \mathcal{Y}$ and A = B) and we will call this subspace a girder, more precisely, $\mathcal{M}_{A,B}$ is the girder of $\mathfrak{I}(A, B; \mathcal{M})$ (and \mathcal{M}_A is the girder of $\mathfrak{C}(A; \mathcal{M})$) induced by \mathcal{M} . Of course, \mathfrak{X} is the girder of $\mathfrak{I}(A, B)$. On the other hand, $\{0\}$ is not necessary a girder — think about $I_{\mathfrak{X}}$. However, by Proposition 2.2, $\{0\}$ is the girder of $\mathfrak{C}(A; \{0\}) = B(\mathfrak{X})$ if A is not a scalar multiple of $I_{\mathfrak{X}}$.

It is clear that $\bigcap_{S \in \mathfrak{I}(A,B;\mathfrak{M})} \ker(BS-SA)$ is a closed subspace of \mathfrak{X} such that BSx = SAxfor every x in this subspace. Hence $\bigcap_{S \in \mathfrak{I}(A,B;\mathfrak{M})} \ker(BS-SA) \subseteq \mathfrak{M}_{A,B}$. On the other hand, if $x \in \mathfrak{M}_{A,B}$, then BSx = SAx for every $S \in \mathfrak{I}(A,B;\mathfrak{M})$, by the definition of $\mathfrak{M}_{A,B}$. Hence,

(12)
$$\mathfrak{M}_{A,B} = \bigcap_{S \in \mathfrak{I}(A,B;\mathfrak{M})} \ker(SA - BS).$$

It follows that

(13)
$$\mathbb{J}(A, B; \mathcal{M}_{A,B}) = \mathbb{J}(A, B; \mathcal{M}) \quad \text{and} \quad (\mathcal{M}_{A,B})_{A,B} = \mathcal{M}_{A,B}.$$

Indeed, since $\mathcal{M} \subseteq \mathcal{M}_{A,B}$ we have $\mathcal{I}(A, B; \mathcal{M}_{A,B}) \subseteq \mathcal{I}(A, B; \mathcal{M})$. On the other hand, the opposite inclusion follows by (12). Thus, $\mathcal{I}(A, B; \mathcal{M}_{A,B}) = \mathcal{I}(A, B; \mathcal{M})$ and therefore $(\mathcal{M}_{A,B})_{A,B} = \mathcal{M}_{A,B}$, by (12).

Proposition 3.4. Let $A \in B(\mathfrak{X})$, $B \in B(\mathfrak{Y})$ and let $\mathfrak{K}, \mathfrak{M} \subseteq \mathfrak{X}$ be closed subspaces. If $\mathfrak{M} \subseteq \mathfrak{K}$, then $\mathfrak{M}_{A,B} \subseteq \mathfrak{K}_{A,B}$. In particular, if $\mathfrak{M} \subseteq \mathfrak{K} \subseteq \mathfrak{M}_{A,B}$, then $\mathfrak{M}_{A,B} = \mathfrak{K}_{A,B}$ and therefore $\mathfrak{I}(A, B; \mathfrak{M}_{A,B}) = \mathfrak{I}(A, B; \mathfrak{K})$.

Proof. Assume that $\mathcal{M} \subseteq \mathcal{K}$. Then $\mathcal{I}(A, B; \mathcal{M}) \supseteq \mathcal{I}(A, B; \mathcal{K})$. Hence, if $x \in \mathcal{M}_{A,B}$, then SAx = BSx for every $S \in \mathcal{I}(A, B; \mathcal{K})$ and therefore $x \in \mathcal{K}_{A,B}$.

If $\mathfrak{M} \subseteq \mathfrak{K} \subseteq \mathfrak{M}_{A,B}$, then $\mathfrak{I}(A, B; \mathfrak{M}) \supseteq \mathfrak{I}(A, B; \mathfrak{K}) \supseteq \mathfrak{I}(A, B; \mathfrak{M}_{A,B}) = \mathfrak{I}(A, B; \mathfrak{M})$, that is, $\mathfrak{I}(A, B; \mathfrak{M}_{A,B}) = \mathfrak{I}(A, B; \mathfrak{K})$. It follows from this and the definition of $\mathfrak{K}_{A,B}$ that $\mathfrak{K}_{A,B} = \mathfrak{M}_{A,B}$.

For a closed subspace $\mathcal{M} \subseteq \mathcal{X}$, we denote by $\operatorname{Alg}(\mathcal{M})$ the algebra of all operators $T \in B(\mathcal{X})$ such that $\mathcal{M} \in \operatorname{Lat}(T)$.

Theorem 3.5. Let $A \in B(\mathfrak{X})$, $B \in B(\mathfrak{Y})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. Then $\mathcal{C}(A; \mathfrak{M}_{A,B}) \cap \operatorname{Alg}(\mathfrak{M}_{A,B})$ is a subalgebra of $B(\mathfrak{X})$ and $\mathfrak{I}(A, B; \mathfrak{M})$ is a right module over it.

If $\mathfrak{X} = \mathfrak{Y}$ and A = B, then $\mathfrak{C}(A; \mathfrak{M}_A) \cap \operatorname{Alg}(\mathfrak{M}_A)$ is the largest subalgebra of $B(\mathfrak{X})$ such that $\mathfrak{C}(A; \mathfrak{M})$ is a right module over it.

Proof. Recall that $\mathcal{M} \subseteq \mathcal{M}_{A,B}$, by the definition of $\mathcal{M}_{A,B}$, and $\mathcal{I}(A, B; \mathcal{M}) = \mathcal{I}(A, B; \mathcal{M}_{A,B})$, by (13). Let $S \in \mathcal{I}(A, B; \mathcal{M})$ and $T \in \mathcal{C}(A; \mathcal{M}_{A,B}) \cap \operatorname{Alg}(\mathcal{M}_{A,B})$ be arbitrary. If $x \in \mathcal{M}$, then STAx = SATx = BSTx. The first equality holds because of $T \in \mathcal{C}(A; \mathcal{M}_{A,B})$ and the second because of $Tx \in \mathcal{M}_{A,B}$ and $S \in \mathcal{I}(A, B; \mathcal{M}_{A,B})$. We have proved that

$$\mathbb{J}(A, B; \mathfrak{M}) \cdot \left(\mathbb{C}(A; \mathcal{M}_{A, B}) \cap \mathrm{Alg}(\mathcal{M}_{A, B}) \right) \subseteq \mathbb{J}(A, B; \mathfrak{M})$$

If $T_1, T_2 \in \mathcal{C}(A; \mathcal{M}_{A,B}) \cap \operatorname{Alg}(\mathcal{M}_{A,B})$, then, of course, $T_1T_2 \in \operatorname{Alg}(\mathcal{M}_{A,B})$. Let $x \in \mathcal{M}_{A,B}$ be arbitrary. Since $T_2 \in \mathcal{C}(A; \mathcal{M}_{A,B})$ we have $T_1T_2Ax = T_1AT_2x$. Because of $T_2x \in \mathcal{M}_{A,B}$ we have $AT_1T_2x = T_1AT_2x$, as well. Hence, $T_1T_2Ax = AT_1T_2x$ for every $x \in \mathcal{M}_{A,B}$, that is, $T_1T_2 \in \mathcal{C}(A; \mathcal{M}_{A,B})$. We have proved that $\mathcal{C}(A; \mathcal{M}_{A,B}) \cap \operatorname{Alg}(\mathcal{M}_{A,B})$ is a subalgebra of $B(\mathfrak{X})$ and $\mathcal{I}(A, B; \mathfrak{M})$ is a right module over it.

Assume now that $\mathfrak{X} = \mathfrak{Y}$ and A = B. By the first part of the proof we already know that $\mathfrak{C}(A; \mathfrak{M}_A) \cap \operatorname{Alg}(\mathfrak{M}_A)$ is a subalgebra of $B(\mathfrak{X})$ and $\mathfrak{C}(A; \mathfrak{M}_A)$ is a right module over it. Let $T \in B(\mathfrak{X})$ be an arbitrary operator such that $\mathfrak{C}(A; \mathfrak{M}_A)T \subseteq \mathfrak{C}(A; \mathfrak{M}_A)$. Since $(A)' \subseteq \mathfrak{C}(A; \mathfrak{M}_A)$ and (A)' contains the identity operator I we see that $T \in \mathfrak{C}(A; \mathfrak{M}_A)$. Let $x \in \mathfrak{M}_A$ and $S \in \mathfrak{C}(A; \mathfrak{M}_A)$ be arbitrary. Then SATx = STAx. Because of $ST \in \mathfrak{C}(A; \mathfrak{M}_A)$ we also have ASTx = STAx. Hence (SA - AS)Tx = 0 for all $S \in \mathfrak{C}(A; \mathfrak{M}_A)$ which means, see (12), that $Tx \in \mathfrak{M}_A$. This proves that $T \in \operatorname{Alg}(\mathfrak{M}_A)$. Thus, $\mathfrak{C}(A; \mathfrak{M}_A) \cap \operatorname{Alg}(\mathfrak{M}_A)$ is the largest subalgebra of $B(\mathfrak{X})$ such that $\mathfrak{C}(A; \mathfrak{M})$ is a right module over it. \Box

In the following theorem we list equivalent conditions for $\mathcal{C}(A; \mathcal{M})$ being an algebra.

Theorem 3.6. Let $A \in B(\mathfrak{X})$, $A \neq \lambda I_{\mathfrak{X}}$ for every $\lambda \in \mathbb{C}$, and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. The following assertions are equivalent:

(i) $\mathcal{C}(A; \mathcal{M})$ is an algebra; (ii) $\mathcal{C}(A; \mathcal{M})\mathcal{M} \subseteq \mathcal{M}_A$; (iii) $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathcal{M}_A$; (iv) $\mathcal{M}_A \in \operatorname{Lat}(\mathcal{C}(A; \mathcal{M}))$. *Proof.* (*i*) ⇒ (*ii*). Let $S, T \in C(A; M)$ be arbitrary. Since $ST \in C(A; M)$ we have (SA - AS)Tx = STAx - STAx = 0 for every $x \in M$. Hence, $Tx \in \ker(SA - AS)$. Since $S \in C(A; M)$ is arbitrary we have $Tx \in M_A$ which means that $TM \subseteq M_A$. Now we may conclude that $C(A; M)M \subseteq M_A$.

 $(ii) \Rightarrow (iii)$. If $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathfrak{X}$, then $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathcal{M}_A$, of course. Assume therefore that $\mathcal{C}(A; \mathcal{M})\mathcal{M} \neq \mathfrak{X}$. Then there exist $z \in \mathfrak{X} \setminus \mathcal{C}(A; \mathcal{M})\mathcal{M}$ and $\xi \in (\mathcal{C}(A; \mathcal{M})\mathcal{M})^{\perp}$ such that $\langle z, \xi \rangle = 1$. Since A is not a scalar multiple of I_x there exists $0 \neq y \in \mathfrak{X}$ which is not an eigenvector of A. Since $Ax \in \mathcal{C}(A; \mathcal{M})\mathcal{M}$ whenever $x \in \mathcal{C}(A; \mathcal{M})\mathcal{M}$ we have $A(y \otimes \xi)x = 0 = (y \otimes \xi)Ax$ for every $x \in \mathcal{C}(A; \mathcal{M})\mathcal{M}$. Hence, $y \otimes \xi \in \mathcal{C}(A; \mathcal{M})$. It follows from $A(y \otimes \xi)z = Ay$ and $(y \otimes \xi)Az = \langle Az, \xi \rangle y$ and the assumption that y is not an eigenvector of A that $A(y \otimes \xi)z \neq (y \otimes \xi)Az$ which implies $z \notin \mathcal{M}_A$.

 $(iii) \Rightarrow (i)$. Let $S, T \in C(A; \mathcal{M})$ be arbitrary. Then, for every $x \in \mathcal{M}$, we have (STA - AST)x = SATx - SATx = 0 since it follows by the assumption that $Tx \in \mathcal{M}_A$ and we already know, see (13), that $C(A; \mathcal{M}) = C(A; \mathcal{M}_A)$. Hence, $ST \in C(A; \mathcal{M})$.

 $(i) \Rightarrow (iv)$. Since $\mathcal{C}(A; \mathcal{M}) = \mathcal{C}(A; \mathcal{M}_A)$ we have $\mathcal{C}(A; \mathcal{M}_A) \mathcal{M}_A \subseteq (\mathcal{M}_A)_A = \mathcal{M}_A$, by the equivalence of assertions (i) and (ii). Hence, $\mathcal{M}_A \in \text{Lat}(\mathcal{C}(A; \mathcal{M}))$.

 $(iv) \Rightarrow (i)$. If $\mathcal{M}_A \in \text{Lat}(\mathcal{C}(A; \mathcal{M}))$, then $\mathcal{C}(A; \mathcal{M})\mathcal{M} \subseteq \mathcal{C}(A; \mathcal{M})\mathcal{M}_A \subseteq \mathcal{M}_A$ and therefore $\mathcal{C}(A; \mathcal{M})$ is an algebra, by equivalence of assertions (i) and (ii).

4. Ultrainvariant subspaces

We begin this section with a definition.

Definition 4.1. A closed subspace $\mathcal{M} \subseteq \mathcal{X}$ is *ultrainvariant* for $A \in B(\mathcal{X})$ if it is invariant for every operator in $\mathcal{C}(A; \mathcal{M})$.

Hence, every ultrainvariant subspace of A is a hyperinvariant subspace, as well. We will see that the opposite implication does not hold in general. For instance, if $A \neq 0$ is a nilpotent operator, then $\overline{\operatorname{im}(A^k)}$ is a hyperinvariant subspace of A for every $k \in \mathbb{N}$, however it is not necessary an ultrainvariant subspace (see Theorem 5.11).

Proposition 4.2. Let $A \in B(\mathfrak{X})$, $A \neq \lambda I_{\mathfrak{X}}$ for every $\lambda \in \mathbb{C}$. A closed subspace $\mathfrak{M} \subseteq \mathfrak{X}$ is ultrainvariant for A if and only if $\mathfrak{C}(A; \mathfrak{M})$ is an algebra and \mathfrak{M} is its girder.

Proof. Assume that \mathcal{M} is an ultrainvariant subspace of A. Let $T_1, T_2 \in \mathcal{C}(A; \mathcal{M})$ be arbitrary. Then $T_1T_2Ax = T_1AT_2x = AT_1T_2x$ for every $x \in \mathcal{M}$ because $T_2x \in \mathcal{M}$. Thus,

 $\mathcal{C}(A; \mathcal{M})$ is an algebra. By Theorem 3.6, $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathcal{M}_A$. However, $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathcal{M}$ as \mathcal{M} is invariant for every operator in $\mathcal{C}(A; \mathcal{M})$, that is, $\mathcal{M} = \mathcal{M}_A$. The opposite implication follows by Theorem 3.6.

Subspaces $\{0\}$ and \mathcal{X} are trivial ultrainvariant subspaces of every $A \in B(\mathcal{X})$. If A is a scalar multiple of I_x , then it is obvious that A has only trivial ultrainvariant subspaces as it has only trivial hyperinvariant subspaces.

In this section we will prove a few general results about ultrainvariant subspaces. Examples of ultrainvariant subspaces for some classes of operators will be given in sections that follow.

Proposition 4.3. Let $A \in B(\mathfrak{X})$ and let $U \in B(\mathfrak{X})$ be an invertible operator. A closed subspace $\mathcal{M} \subseteq \mathfrak{X}$ is an ultrainvariant subspace of A if and only if $U\mathcal{M}$ is an ultrainvariant subspace of UAU^{-1} .

Proof. Assume that \mathcal{M} is ultrainvariant. By (1), every operator in $\mathcal{C}(UAU^{-1}; U\mathcal{M})$ is of the form UTU^{-1} for some $T \in \mathcal{C}(A; \mathcal{M})$. Hence, for every $x \in \mathcal{M}$, we have $(UTU^{-1})(Ux) = UTx \in U\mathcal{M}$, that is, $U\mathcal{M}$ is invariant for every operator from $\mathcal{C}(UAU^{-1}; U\mathcal{M})$ and is therefore ultrainvariant. It is clear that the opposite implication holds, as well. \Box

Recall that a closed subspace $\mathfrak{X}_1 \subseteq \mathfrak{X}$ is reducing for $A \in B(\mathfrak{X})$ if $\mathfrak{X}_1 \in \operatorname{Lat}(A)$ and there exists $\mathfrak{X}_2 \in \operatorname{Lat}(A)$ such that $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2$. In this case we will say that $(\mathfrak{X}_1, \mathfrak{X}_2)$ is a reducing pair for A. Note that $A = A_1 \oplus A_2$ with respect to the decomposition $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2$. Let $P_j : \mathfrak{X} \to \mathfrak{X}_j$ (j = 1, 2) be given by $P_1(x_1 \oplus x_2) = x_1$ and $P_2(x_1 \oplus x_2) = x_2$.

Proposition 4.4. Let $A \in B(\mathfrak{X})$ and let $(\mathfrak{X}_1, \mathfrak{X}_2)$ be a reducing pair for A. If \mathfrak{M} is an ultrainvariant subspace of A, then $\mathfrak{M}_j = P_j \mathfrak{M}$ is an ultrainvariant subspace of A_j (j = 1, 2).

Proof. Assume that $\mathcal{M} = \mathcal{M}_1 \oplus \mathcal{M}_2$ is ultrainvariant. Let $T_1 \in \mathcal{C}(A_1; \mathcal{M}_1)$ be arbitrary and let $T \in B(\mathfrak{X})$ be such that $T = T_1 \oplus 0$ with respect to the decomposition $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2$. For an arbitrary $x = x_1 \oplus x_2 \in \mathcal{M}$, we have $TAx = T_1A_1x_1 \oplus 0 = A_1T_1x_1 \oplus 0 = ATx$ which means that $T \in \mathcal{C}(A; \mathcal{M})$. Since \mathcal{M} is ultrainvariant it is invariant for T. In particular, for every $x_1 \in \mathcal{M}_1$ we have $T(x_1 \oplus 0) = T_1x_1 \oplus 0 \in \mathcal{M}_1 \oplus \mathcal{M}_2$, that is, \mathcal{M}_1 is invariant for $T_1 \in \mathcal{C}(A_1; \mathcal{M}_1)$. We have seen that \mathcal{M}_1 is an ultrainvariant subspace of A_1 and a similar proof shows that \mathcal{M}_2 is an ultrainvariant subspace of A_2 . Let $n \geq 2$ be an integer. For every $j \in \{1, \ldots, n\}$, let \mathfrak{X}_j be a Banach space and $A_j \in B(\mathfrak{X}_j)$. Denote $\mathfrak{X} = \mathfrak{X}_1 \oplus \cdots \oplus \mathfrak{X}_n$ and $A = A_1 \oplus \cdots \oplus A_n$. Let $\mathfrak{M}_j \subseteq \mathfrak{X}_j$ be a complemented closed subspace and let $\mathfrak{N}_j \subseteq \mathfrak{X}_j$ be a closed subspace such that $\mathfrak{X}_j = \mathfrak{M}_j \oplus \mathfrak{N}_j$. Let $\begin{bmatrix} A_{11}^{(j)} & A_{12}^{(j)} \\ A_{21}^{(j)} & A_{22}^{(j)} \end{bmatrix}$ be the operator matrix of A_j with respect to the decomposition $\mathfrak{X}_j = \mathfrak{M}_j \oplus \mathfrak{N}_j$.

Proposition 4.5. If, for every $j \in \{1, ..., n\}$, \mathcal{M}_j is an ultrainvariant subspace of A_j and $\sigma(A_{11}^{(i)}) \cap \sigma(A_j) = \emptyset$ whenever $i \neq j$, then $\mathcal{M} := \mathcal{M}_1 \oplus \cdots \oplus \mathcal{M}_n$ is an ultrainvariant subspace of A.

Proof. It is enough to consider the case n = 2. We will identify \mathfrak{X} with $\mathfrak{M}_1 \oplus \mathfrak{N}_1 \oplus \mathfrak{M}_2 \oplus \mathfrak{N}_2$. Then $\mathfrak{M} = \mathfrak{M}_1 \oplus \{0\} \oplus \mathfrak{M}_2 \oplus \{0\}$. With respect to the decomposition $\mathfrak{X} = \mathfrak{M}_1 \oplus \mathfrak{N}_1 \oplus \mathfrak{M}_2 \oplus \mathfrak{N}_2$, operator A has operator matrix

$$\begin{bmatrix} A_{11}^{(1)} & A_{12}^{(1)} & 0 & 0 \\ 0 & A_{22}^{(1)} & 0 & 0 \\ 0 & 0 & A_{11}^{(2)} & A_{12}^{(2)} \\ 0 & 0 & 0 & A_{22}^{(2)} \end{bmatrix}.$$

Let $T \in \mathcal{C}(A; \mathcal{M})$ be arbitrary and let $[T_{ij}]_{i,j=1}^4$ be its operator matrix with respect to the decomposition $\mathcal{X} = \mathcal{M}_1 \oplus \mathcal{N}_1 \oplus \mathcal{M}_2 \oplus \mathcal{N}_2$. Hence, TAx = ATx for every $x = x_1 \oplus 0 \oplus x_2 \oplus 0 \in \mathcal{M}$. If $x = x_1 \oplus 0 \oplus 0 \oplus 0$, where $x_1 \in \mathcal{M}_1$ is arbitrary, then it follows from TAx = ATx that

(14)

$$T_{11}A_{11}^{(1)} = A_{11}^{(1)}T_{11} + A_{12}^{(1)}T_{21}$$

$$T_{21}A_{11}^{(1)} = A_{22}^{(1)}T_{21}$$

$$T_{31}A_{11}^{(1)} = A_{11}^{(2)}T_{31} + A_{12}^{(2)}T_{41}$$

$$T_{41}A_{11}^{(1)} = A_{22}^{(2)}T_{41}.$$

Let $W_{11} \in B(\mathfrak{X}_1)$ be the operator whose operator matrix with respect to the decomposition $\mathfrak{X}_1 = \mathfrak{M}_1 \oplus \mathfrak{N}_1$ is $\begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$. The first two equalities in (14) give $W_{11} \in \mathcal{C}(A_1; \mathfrak{M}_1)$. Since \mathfrak{M}_1 is an ultrainvariant subspace of A_1 it is invariant for W_{11} and therefore $T_{21} = 0$. Hence, by the first equality in (14), $T_{11} \in (A_{11}^{(1)})'$. Denote by W_{21} the operator in $B(\mathfrak{X}_1, \mathfrak{X}_2)$ whose operator matrix with respect to the decompositions $\mathfrak{X}_1 = \mathfrak{M}_1 \oplus \mathfrak{N}_1$ and $\mathfrak{X}_2 = \mathfrak{M}_2 \oplus \mathfrak{N}_2$ is $\begin{bmatrix} T_{31} & T_{32} \\ T_{41} & T_{42} \end{bmatrix}$. It follows from the last two equalities in (14) that column $\begin{bmatrix} T_{31} \\ T_{41} \end{bmatrix}$ intertwines $A_{11}^{(1)}$ and A_2 . Since, by the assumption, $\sigma(A_{11}^{(1)}) \cap \sigma(A_2) = \emptyset$ we conclude that $T_{31} = 0$ and $T_{41} = 0$. Let now $x_2 \in \mathcal{M}_2$ be arbitrary and $x = 0 \oplus 0 \oplus x_2 \oplus 0$. A similar reasoning as in the previous paragraph shows that TAx = ATx if and only if $T_{13} = 0$, $T_{23} = 0$, $T_{43} = 0$ and $T_{33} \in (A_{11}^{(2)})'$. Hence, the operator matrix of T with respect to the decomposition $\mathcal{X} = \mathcal{M}_1 \oplus \mathcal{N}_1 \oplus \mathcal{M}_2 \oplus \mathcal{N}_2$ is

$$\begin{bmatrix} T_{11} & T_{12} & 0 & T_{14} \\ 0 & T_{22} & 0 & T_{24} \\ 0 & T_{32} & T_{33} & T_{34} \\ 0 & T_{42} & 0 & T_{44} \end{bmatrix}$$

It is not hard to check now that $\mathcal{M} = \mathcal{M}_1 \oplus \{0\} \oplus \mathcal{M}_2 \oplus \{0\}$ is invariant for T. Since $T \in \mathcal{C}(A; \mathcal{M})$ was arbitrary we conclude that \mathcal{M} is an ultrainvariant subspace of A. \Box

Proposition 4.6. Let $A \in B(\mathfrak{X})$ and let $\mathfrak{M} \subseteq \mathfrak{X}$ be a closed subspace. Then

$$\mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})\mathcal{C}(A; \mathcal{M})\mathcal{M}$$

is the smallest ultrainvariant subspace of A which contains \mathcal{M} .

Proof. By Corollary 3.3, $\mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})$ is the largest algebra which is contained in $\mathcal{C}(A; \mathcal{M})$. It follows, by Theorem 3.6, that $(\mathcal{C}(A; \mathcal{M})\mathcal{M})_A = \mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})\mathcal{C}(A; \mathcal{M})\mathcal{M}$ and $(\mathcal{C}(A; \mathcal{M})\mathcal{M})_A$ is an ultrainvariant subspace of A. It is clear that $\mathcal{M} \subseteq (\mathcal{C}(A; \mathcal{M})\mathcal{M})_A$.

Suppose that $\mathcal{K} \subseteq \mathcal{X}$ is an ultrainvariant subspace of A such that $\mathcal{M} \subseteq \mathcal{K}$. Then $\mathcal{C}(A; \mathcal{K})$ is an algebra and $\mathcal{C}(A; \mathcal{M}) \supseteq \mathcal{C}(A; \mathcal{K})$. Since $\mathcal{C}(A; (\mathcal{C}(A; \mathcal{M})\mathcal{M})_A) = \mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})$ is the largest algebra contained in $\mathcal{C}(A; \mathcal{M})$ we have $\mathcal{C}(A; \mathcal{K}) \subseteq \mathcal{C}(A; \mathcal{C}(A; \mathcal{M})\mathcal{M})$. Hence, if $T \in \mathcal{C}(A; \mathcal{K})$, then it commutes with A at every vector from $\mathcal{C}(A; \mathcal{M})\mathcal{M}$ which means that $\mathcal{C}(A; \mathcal{M})\mathcal{M} \subseteq \mathcal{K}_A = \mathcal{K}$ and therefore $(\mathcal{C}(A; \mathcal{M})\mathcal{M})_A \subseteq \mathcal{K}$.

For $A \in B(\mathfrak{X})$, let $\operatorname{Lat}_u(A)$ denote the family of all ultrainvariant subspaces of A.

Proposition 4.7. Let $A \in B(\mathfrak{X})$ and let $\{\mathfrak{M}_j; j \in J\}$ be an arbitrary family of ultrainvariant subspaces of A. Then $\bigvee_{j \in J} \mathfrak{M}_j$ and $\bigcap_{j \in J} \mathfrak{M}_j$ are ultrainvariant subspaces of A. Hence, $\operatorname{Lat}_u(A)$ is a sublattice of $\operatorname{Lat}_h(A)$.

Proof. Denote $\mathcal{K} = \bigvee_{j \in J} \mathcal{M}_j$. By (2), we have $\mathcal{C}(A; \mathcal{K}) = \bigcap_{j \in J} \mathcal{C}(A; \mathcal{M}_j)$. Hence, $\mathcal{M}_j \ (j \in J)$ is invariant for an arbitrary $T \in \mathcal{C}(A; \mathcal{K})$. Assume that $x \in \mathcal{K}$ is such that $x = x_1 + \cdots + x_k$, where $x_i \in \mathcal{M}_{j_i} \ (i = 1, \dots, k)$. Then $Tx = Tx_1 + \cdots + Tx_k \in \mathcal{M}_{j_1} + \cdots + \mathcal{M}_{j_k} \subseteq \mathcal{M}$. It follows that $T\mathcal{K} \subseteq \mathcal{K}$. Since $T \in \mathcal{C}(A; \mathcal{K})$ is arbitrary we conclude that \mathcal{K} is ultrainvariant.

Denote $\mathcal{L} = \bigcap_{j \in J} \mathcal{M}_j$ and let $\mathcal{L}' \subseteq \mathcal{X}$ be the smallest ultrainvariant subspace of A such that $\mathcal{L} \subseteq \mathcal{L}'$ (see Proposition 4.6). Since, for every $j \in J$, $\mathcal{L} \subseteq \mathcal{M}_j$ and \mathcal{M}_j is ultrainvariant

we have $\mathcal{L}' \subseteq \mathcal{M}_j$. Hence $\mathcal{L}' \subseteq \bigcap_{j \in J} \mathcal{M}_j = \mathcal{L}$, that is, $\mathcal{L} = \mathcal{L}'$, i.e., \mathcal{L} is an ultrainvariant subspace of A.

The following example shows that even when $\mathcal{M} \in \text{Lat}(A)$ is the girder of $\mathcal{C}(A; \mathcal{M})$ it is not necessary an ultrainvariant subspace of A, that is, $\mathcal{C}(A; \mathcal{M})$ is not necessary an algebra.

Example 4.8. Let $\mathcal{X} = \mathcal{X}_1 \oplus \mathcal{X}_2 \oplus \mathcal{X}_3$, where \mathcal{X}_j (j = 1, 2, 3) are non-trivial subspaces of \mathcal{X} . Let A be the projection on $\mathcal{X}_1 \oplus \mathcal{X}_2$ along \mathcal{X}_3 and let $\mathcal{M} = \mathcal{X}_1 \oplus \{0\} \oplus \mathcal{X}_3$, that is, in \mathcal{M} are vectors of the form $x_1 \oplus 0 \oplus x_3$, where $x_1 \in \mathcal{X}_1$ and $x_3 \in \mathcal{X}_3$ are arbitrary. Hence, \mathcal{M} is a non-trivial subspace of \mathcal{X} and $\mathcal{M} \in \text{Lat}(A)$.

Let $T \in \mathcal{C}(A; \mathcal{M})$ be arbitrary and let $[T_{ij}]_{i,j=1}^3$ be its operator matrix with respect to the decomposition $\mathcal{X} = \mathcal{X}_1 \oplus \mathcal{X}_2 \oplus \mathcal{X}_3$. For every $x = x_1 \oplus 0 \oplus x_3 \in \mathcal{M}$, we have $TAx = T_{11}x_1 \oplus T_{21}x_1 \oplus T_{31}x_1$ and $ATx = (T_{11}x_1 + T_{13}x_3) \oplus (T_{21}x_1 + T_{23}x_3) \oplus 0$. It follows from TAx = ATx that $T_{11}x_1 = T_{11}x_1 + T_{13}x_3$, $T_{21}x_1 = T_{21}x_1 + T_{23}x_3$ and $T_{31}x_1 = 0$ for all $x_1 \in \mathcal{X}_1$ and $x_3 \in \mathcal{X}_3$. Thus, $T_{13} = 0$, $T_{23} = 0$ and $T_{31} = 0$. An operator T is in $\mathcal{C}(A; \mathcal{M})$ if and only if its operator matrix with respect to the decomposition $\mathcal{X} = \mathcal{X}_1 \oplus \mathcal{X}_2 \oplus \mathcal{X}_3$ is of the form $\begin{bmatrix} T_{11} & T_{12} & 0 \\ 0 & T_{32} & T_{33} \end{bmatrix}$, where $T_{ij} \in B(\mathcal{X}_j, \mathcal{X}_i)$ are arbitrary. It follows that $\mathcal{C}(A; \mathcal{M})$ is not an algebra.

To determine \mathcal{M}_A , let $x = x_1 \oplus x_2 \oplus x_3 \in \mathfrak{X}$ be such that TAx = ATx for every $T \in \mathcal{C}(A; \mathcal{M})$. Since $TAx = (T_{11}x_1 + T_{12}x_2) \oplus (T_{21}x_1 + T_{22}x_2) \oplus T_{32}x_2$ and $ATx = (T_{11}x_1 + T_{12}x_2) \oplus (T_{21}x_1 + T_{22}x_2) \oplus (T_{21}x_1 + T_{22}x_2) \oplus 0$ we see that $T_{32}x_2 = 0$ for every $T_{32} \in B(\mathfrak{X}_2, \mathfrak{X}_3)$. As the involved subspaces are non-trivial we conclude that $x_2 = 0$. This proves that $\mathcal{M}_A = \mathcal{M}$. However, \mathcal{M} is not an ultrainvariant subspace of A because $\mathcal{C}(A; \mathcal{M})$ is not an algebra. Moreover, $\mathcal{C}(A; \mathcal{M})\mathcal{M} = \mathfrak{X}$. Namely, let $x = x_1 \oplus x_2 \oplus x_3$ be an arbitrary vector from \mathfrak{X} . Let $y = y_1 \oplus 0 \oplus y_3 \in \mathcal{M}$ be such that $y_1 \neq 0$ and $y_3 \neq 0$. Then there exist $T_{11} \in B(\mathfrak{X}_1), T_{21} \in B(\mathfrak{X}_1, \mathfrak{X}_2)$ and $T_{33} \in B(\mathfrak{X}_3)$ such that $T_{11}y_1 = x_1, T_{21}y_1 = x_2$ and $T_{33}y_3 = x_3$. It is clear that T whose operator matrix with respect to the decomposition $\mathfrak{X} = \mathfrak{X}_1 \oplus \mathfrak{X}_2 \oplus \mathfrak{X}_3$ is $\begin{bmatrix} T_{11} & 0 & 0 \\ T_{21} & 0 & 0 \\ 0 & 0 & T_{33} \end{bmatrix}$ is in $\mathcal{C}(A; \mathcal{M})$ and Ty = x.

5. Operators with non-trivial ultrainvariant subspaces

Let $A \in B(\mathfrak{X})$ and assume that $\mathfrak{M} \in \operatorname{Lat}(A)$ is complemented, say $\mathfrak{X} = \mathfrak{M} \oplus \mathfrak{N}$. Let $\begin{bmatrix} A_{11} & A_{12} \\ 0 & A_{22} \end{bmatrix}$ be the operator matrix of A with respect this decomposition. Next proposition is basically a reformulation of Corollary 2.7.

Proposition 5.1. If $\sigma(A_{11}) \cap \sigma(A_{22}) = \emptyset$, then \mathfrak{M} is an ultrainvariant subspace of A.

Proof. If \mathfrak{M} is trivial, then there is nothing to prove. Assume therefore that $\{0\} \neq \mathfrak{M} \neq \mathfrak{X}$. By Corollary 2.7, $\mathfrak{C}(A; \mathfrak{M})$ is an algebra. Hence, if $\begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$ is the operator matrix of $T \in \mathfrak{C}(A; \mathfrak{M})$ with respect to the decomposition $\mathfrak{X} = \mathfrak{M} \oplus \mathfrak{N}$, then $T_{11} \in (A_{11})'$ and $T_{21} = 0$ by Theorem 2.5. It follows that \mathfrak{M} is invariant for every $T \in \mathfrak{C}(A; \mathfrak{M})$. \Box

Recall that the ascent $\alpha(A)$ of $A \in B(\mathfrak{X})$ is the smallest positive integer k such that $\ker(A^k) = \ker(A^{k+1})$. If there is no positive integer with this property, then $\alpha(A) = \infty$. The descent $\delta(A)$ of A is the smallest positive integer k such that $\operatorname{im}(A^k) = \operatorname{im}(A^{k+1})$ and $\delta(A) = \infty$ if there is no k with this property. Assume that A has finite ascent and descent. Then $\alpha(A) = n = \delta(A)$ for a positive integer n and, by [1, Lemma 2.21], $\operatorname{im}(A^n)$ is a closed complement of $\ker(A^n)$, that is $\mathfrak{X} = \ker(A^n) \oplus \operatorname{im}(A^n)$. Hence, $(\ker(A^n), \operatorname{im}(A^n))$ is a reducing pair of A and therefore $A = A_1 \oplus A_2$, where $A_1 \in B(\ker(A^n))$ and $A_2 \in B(\operatorname{im}(A^n))$. By [1, Theorem 2.23], A_1 is nilpotent and A_2 is invertible which implies $\sigma(A_1) \cap \sigma(A_2) = \emptyset$. It is clear now that the following corollary is a simple consequence of Proposition 5.1

Corollary 5.2. If $A \in B(\mathfrak{X})$ has finite ascent and descent n, then ker (A^n) and im (A^n) are ultrainvariant subspaces of A.

It is not hard to see that for every $\lambda \in \mathbb{C}$ the kernel $\ker(A - \lambda I_x)$ is an ultrainvariant subspace of $A \in B(\mathfrak{X})$. Namely, if $T \in \mathcal{C}(A; \ker(A - \lambda I_x))$, then $ATx = TAx = \lambda Tx$ for every $x \in \ker(A - \lambda I_x)$, that is, $\ker(A - \lambda I_x)$ is invariant for every $T \in \mathcal{C}(A; \ker(A - \lambda I_x))$. Hence, if A is not a scalar multiple of I_x and the point spectrum $\sigma_p(A)$ is not empty, then A has a non-trivial ultrainvariant subspace. More can be said. We need the following lemma.

Lemma 5.3. Let $A \in B(\mathfrak{X})$ and let $\mathcal{A} \subseteq B(\mathfrak{X})$ be the strongly closed subalgebra generated by A and $I_{\mathfrak{X}}$. If $\mathfrak{M} \in \operatorname{Lat}(A)$, then $\mathfrak{C}(A; \mathfrak{M}) \subseteq \mathfrak{C}(B; \mathfrak{M})$ for every $B \in \mathcal{A}$.

Proof. Assume first that $B = A - \lambda I_x$, where $\lambda \in \mathbb{C}$ is such that B is an invertible operator. Hence, $B\mathcal{M} = \mathcal{M}$. It is clear that $C(A; \mathcal{M}) = \mathcal{C}(B; \mathcal{M})$. Let us check that $\mathcal{C}(B; \mathcal{M}) \subseteq \mathcal{C}(B^k; \mathcal{M})$ for every integer $k \geq 0$. If k = 0 or k = 1, then the assertion is trivial. Since B is invertible and $B\mathcal{M} = \mathcal{M}$ we have $\mathcal{C}(B; \mathcal{M}) = \mathcal{C}(B; \mathcal{M})B$, by Proposition 3.1. Hence, if $T \in \mathcal{C}(B; \mathcal{M})$, then $TB \in \mathcal{C}(B; \mathcal{M})$ and, by induction, $TB^j \in \mathcal{C}(B; \mathcal{M})$ for every $j \ge 2$. Let $k \ge 2$ and assume that $TB^j x = B^j Tx$ holds for every $0 \le j \le k - 1$ and every $x \in \mathcal{M}$. Then $TB^k x = B^{k-1}TBx$ and TBx = BTx for every $x \in \mathcal{M}$. It follows that $TB^k x = B^k Tx$ for every $x \in \mathcal{M}$. Since $A = B + \lambda I_x$ we conclude that

$$TA^{k}x = T\sum_{j=0}^{k} \binom{k}{j}\lambda^{j}B^{k-j}x = \sum_{j=0}^{k} \binom{k}{j}\lambda^{j}B^{k-j}Tx = A^{k}Tx \quad \text{for every } x \in \mathcal{M}.$$

It follows that Tp(A)x = p(A)Tx for every $x \in \mathcal{M}$ and every polynomial $p \in \mathbb{C}[z]$. Let $B \in \mathcal{A}$ be arbitrary. Let $T \in \mathbb{C}(A; \mathcal{M})$ and $x \in \mathcal{M}$. For every $\varepsilon > 0$, there exists a polynomial p such that $||Bx - p(A)x|| < \varepsilon$ and $||BTx - p(A)Tx|| < \varepsilon$. Hence $||TBx - BTx|| \leq ||T|| ||Bx - p(A)x|| + ||BTx - p(A)Tx|| < \varepsilon(||T|| + 1)$, which gives TBx = BTx. \Box

Proposition 5.4. Let $A \in B(\mathfrak{X})$ and let $\mathcal{A} \subseteq B(\mathfrak{X})$ be the strongly closed subalgebra generated by A and $I_{\mathfrak{X}}$. If $B \in \mathcal{A}$ and $\mathfrak{M} \in \operatorname{Lat}(A)$ is an ultrainvariant subspace of B, then \mathfrak{M} is an ultrainvariant subspace of A. In particular, every kernel ker $(B - \lambda I_{\mathfrak{X}})$ $(\lambda \in \mathbb{C})$ is an ultrainvariant subspace of A.

Proof. By Lemma 5.3, $\mathcal{C}(A; \mathcal{M})\mathcal{M} \subseteq \mathcal{C}(B; \mathcal{M})\mathcal{M} = \mathcal{M}$. Thus, \mathcal{M} is invariant for every operator from $\mathcal{C}(A; \mathcal{M})$, that is, it is an ultrainvariant subspace of A.

It follows from the following proposition that a reducing subspace does not need to be ultrainvariant.

Proposition 5.5. Let $A \in B(\mathfrak{X})$. Assume that $(\mathfrak{M}, \mathfrak{N})$ is a pair of reducing subspaces for A. Let A be equal to $A_{\mathfrak{M}} \oplus A_{\mathfrak{N}}$ with respect to the decomposition $\mathfrak{X} = \mathfrak{M} \oplus \mathfrak{N}$. Then \mathfrak{M} is an ultrainvariant subspace of A if and only if $\mathfrak{I}(A_{\mathfrak{M}}, A_{\mathfrak{N}}) = \{0\}$.

Proof. Suppose that \mathcal{M} is ultrainvariant and let $S \in \mathcal{I}(A_{\mathcal{M}}, A_{\mathcal{N}})$ be arbitrary. Then $T(x \oplus y) = 0 \oplus Sx$, where $x \in \mathcal{M}, y \in \mathcal{N}$ are arbitrary, defines a bounded linear operator on \mathcal{X} . Since $AT(x \oplus 0) = 0 \oplus A_{\mathcal{N}}Sx = 0 \oplus SA_{\mathcal{M}}x = TA(x \oplus 0)$ for all $x \in \mathcal{M}$ we have $T \in \mathcal{C}(A; \mathcal{M})$. It follows from $T\mathcal{M} \subseteq \mathcal{M}$ that Sx = 0 for every $x \in \mathcal{M}$, that is, S = 0.

Assume now that $\mathcal{J}(A_{\mathfrak{M}}, A_{\mathfrak{N}}) = \{0\}$. Let $T \in \mathcal{C}(A; \mathcal{M})$ be arbitrary and let $\begin{bmatrix} T_{11} & T_{12} \\ T_{21} & T_{22} \end{bmatrix}$ be its operator matrix with respect to the decomposition $\mathfrak{X} = \mathcal{M} \oplus \mathfrak{N}$. It follows from $TA(x \oplus 0) = AT(x \oplus 0)$, where $x \in \mathcal{M}$ is arbitrary, that $T_{21} \in \mathcal{J}(A_{\mathfrak{M}}, A_{\mathfrak{N}})$, that is, $T_{21} = 0$. Hence, \mathcal{M} is an ultrainvariant subspace of A.

Let $A \in B(\mathfrak{X})$. A subset $\emptyset \subsetneq \sigma \subsetneq \sigma(A)$ is an isolated part of $\sigma(A)$ if both σ and $\sigma(A) \setminus \sigma$ are closed sets. It is well-known, see [10, §I.2], that for an isolated part σ of $\sigma(A)$ there

exists an idempotent $P_{\sigma} \in B(\mathfrak{X})$, the Riesz projection of A corresponding to σ , such that $\mathfrak{M} = \operatorname{im}(P_{\sigma})$ and $\mathfrak{N} = \operatorname{im}(I - P_{\sigma})$ are in $\operatorname{Lat}(A)$ and $\sigma(A|_{\mathfrak{M}}) = \sigma$, $\sigma(A|_{\mathfrak{N}}) = \sigma(A) \setminus \sigma$. Hence, the following result is an immediate consequence of Proposition 5.5.

Corollary 5.6. Let $A \in B(\mathfrak{X})$. If σ is an isolated part of $\sigma(A)$, then $im(P_{\sigma})$ is an ultrainvariant subspace of A.

Corollary 5.6 is a special case of the next result for which we need some notions from the local spectral theory. Let $A \in B(\mathfrak{X})$ and $x \in \mathfrak{X}$. The local resolvent set $\rho_A(x)$ of Aat x is the union of all open subsets $U \subseteq \mathbb{C}$ for which there exists an analytic function $f: U \to \mathfrak{X}$ such that $(A - \lambda I_x)f(\lambda) = x$ for all $\lambda \in U$. The local spectrum of A at xis then defined as $\sigma_A(x) = \mathbb{C} \setminus \rho_A(x)$. It is obvious that $\sigma_A(x)$ is a closed subset of the spectrum $\sigma(A)$. For an arbitrary $F \subseteq \mathbb{C}$, the local spectral subspace of A corresponding to F is $\mathfrak{X}_A(F) = \{x \in \mathfrak{X}; \sigma_A(x) \subseteq F\}$. By [14, Proposition 1.2.16 (a)], $\overline{\mathfrak{X}_A(F)}$ is a hyperinvariant subspace of A.

Proposition 5.7. $\overline{\mathfrak{X}_A(F)}$ is an ultrainvariant subspace of A.

Proof. Let $T \in \mathcal{C}(A; \overline{\mathcal{X}_A(F)})$ be arbitrary. Let $x \in \mathcal{X}_A(F)$ and let $\lambda_0 \in \rho_A(x)$. Then there is an open neighborhood $U \subseteq \rho_A(x)$ of λ_0 and an analytic function $f: U \to \mathcal{X}$ such that $(A - \lambda I_x)f(\lambda) = x$ for all $\lambda \in U$. By [14, Lemma 1.2.14], $\sigma_A(f(\lambda)) = \sigma_A(x)$ for every $\lambda \in U$. Hence, $f(\lambda) \in \mathcal{X}_A(F)$ for all $\lambda \in U$. It follows that $(A - \lambda I_x)Tf(\lambda) =$ $T(A - \lambda I_x)f(\lambda) = Tx$ for all $\lambda \in U$. Since $\lambda \mapsto Tf(\lambda)$ is an analytic function on U we conclude that $\lambda_0 \in \sigma_A(Tx)$ and therefore $\sigma_A(Tx) \subseteq \sigma_A(x) \subseteq F$. Thus, $\overline{\mathcal{X}_A(F)}$ is invariant for T.

Recall that $A \in B(\mathfrak{X})$ is a decomposable operator if for every open cover $\mathbb{C} = U \cup V$ there exist spaces $\mathcal{M}, \mathcal{N} \in \text{Lat}(A)$ such that $\mathfrak{X} = \mathcal{M} + \mathcal{N}$ and $\sigma(A|_{\mathcal{M}}) \subseteq U, \sigma(A|_{\mathcal{N}}) \subseteq V$ (see Definition 1.1.1 in [14]). If A is decomposable, then the local spectral subspace $\mathfrak{X}_A(F)$ is closed whenever F is a closed subset of \mathbb{C} (see Definition 1.2.18 and Theorem 1.2.29 in [14]) and $\sigma(A) = \bigcup \{\sigma_A(x); x \in \mathfrak{X}\}$ (see Proposition 1.3.2 in [14]).

Corollary 5.8. If $A \in B(\mathfrak{X})$ is a decomposable operator such that the spectrum $\sigma(A)$ contains at least two points, then A has a proper non-trivial ultrainvariant subspace.

Proof. Let $\lambda_1 \neq \lambda_2$ be points in $\sigma(A)$ and let $\mathbb{C} = U \cup V$ be an open cover such that $\lambda_1 \in U$ and $\lambda_2 \notin \overline{U}$. Then $\mathfrak{X}_A(\overline{U})$ and $\mathfrak{X}_A(\overline{V})$ are non-trivial and proper closed subspaces of \mathfrak{X} .

Indeed, by [14, Theorem 1.2.23], $\mathfrak{X} = \mathfrak{X}_A(\overline{U}) + \mathfrak{X}_A(\overline{V})$. Since $\sigma_A(x) \subseteq \overline{U} \cap \sigma(A) \subsetneq \sigma(A)$ and $\sigma(A) = \bigcup \{\sigma_A(x); x \in \mathfrak{X}\}$ there are vectors which are not in $\mathfrak{X}_A(\overline{U})$. Similarly, there are vectors which are not in $\mathfrak{X}_A(\overline{V})$. Hence, $\mathfrak{X}_A(\overline{U})$ and $\mathfrak{X}_A(\overline{V})$ are proper and non-trivial. By Proposition 5.7, these subspaces are ultrainvariant.

Assume that $A \in B(\mathfrak{X})$ is a nilpotent operator of order $n \geq 2$, that is, $A^n = 0$ and $A^{n-1} \neq 0$. It is well-known that

$$\{0\} = \ker(A^0) \subseteq \ker(A) \subseteq \ker(A^2) \subseteq \dots \subseteq \ker(A^{n-1}) \subseteq \ker(A^n) = \mathcal{X}$$

and

$$\{0\} = \operatorname{im}(A^n) \subseteq \overline{\operatorname{im}(A^{n-1})} \subseteq \overline{\operatorname{im}(A^{n-2})} \subseteq \dots \subseteq \overline{\operatorname{im}(A)} \subseteq \operatorname{im}(A^0) = \mathfrak{X}$$

are two chains of hyperinvariant subspaces of A. By [3, Remarque p. 317], subspaces $\overline{\operatorname{im}(A^{n-1})}$ and $\operatorname{ker}(N^{n-1})$ are the smallest, respectively the largest, non-trivial hyperinvariant subspaces of A. For every $j = 0, 1, \ldots, n-1$, we have $\overline{\operatorname{im}(A^{n-j})} \subsetneq \operatorname{im}(A^{n-j-1})$ and $\operatorname{ker}(A^j) \subsetneq \operatorname{ker}(A^{j+1})$. Of course, $\overline{\operatorname{im}(A^{n-j})} \subseteq \operatorname{ker}(A^j)$. Later we will need the following simple facts.

Lemma 5.9. (i) For every j = 1, ..., n - 1, there exists $u \in im(A^{n-j})$ such that $u \notin ker(A^{j-1})$.

(ii) Let $e \in \mathfrak{X}$ be such that $A^k e \neq 0$ for an integer $k \in \{0, 1, \dots, n-1\}$. Then $A^k e \notin \bigvee \{e, Ae, \dots, A^{k-1}e\}$, that is, $e, Ae, \dots, A^k e$ are linearly independent.

Proof. (i) If $im(A^{n-j}) \subseteq ker(A^{j-1})$, then we would have $A^{j-1}(N^{n-j}x) = 0$ for every $x \in \mathfrak{X}$, that is, it would be $A^{n-1} = 0$.

(ii) Towards a contradiction assume that there exist numbers $\alpha_0, \ldots, \alpha_{k-1}$ such that

$$A^k e = \alpha_0 e + \alpha_1 A e + \dots + \alpha_{k-1} A^{k-1} e.$$

Since $A^k e \neq 0$ there exists the smallest index $0 \leq j \leq k-1$ such that $\alpha_j \neq 0$. Hence,

$$A^k e = \alpha_j A^j e + \alpha_{j+1} A^{j+1} e + \dots + \alpha_{k-1} A^{k-1} e.$$

Let m be the integer such that $A^m e \neq 0$ and $A^{m+1}e = 0$. Multiply the above equality by A^{m-j} . Since j < k we have

$$0 = A^{m-j+k}e = \alpha_j A^m e + \alpha_{j+1} A^{m+1} + \dots + \alpha_{k-1} A^{m-j+k-1}e = \alpha_j A^m e.$$

It follows that $\alpha_j = 0$ as $A^m e \neq 0$. This is a contradiction.

Lemma 5.10. Let $j \in \{1, \ldots, n\}$. If $e \in \ker(A^j)$ and $\xi \in \mathfrak{X}^*$, then

$$(e \otimes \xi)A^{j-1} + A(e \otimes \xi)A^{j-2} + \dots + A^{j-1}(e \otimes \xi) \in \mathcal{C}(A; \overline{\mathrm{im}(A^{n-j})}).$$

Proof. It is not hard to check that $(e \otimes \xi)A^{j-1} + A(e \otimes \xi)A^{j-2} + \dots + A^{j-1}(e \otimes \xi)$ and A commute locally at every vector $A^{n-j}x \in im(A^{n-j})$, where $x \in \mathfrak{X}$ is arbitrary. \Box

Theorem 5.11. Let $A \in B(\mathfrak{X})$ be a nilpotent operator of order $n \geq 2$. A closed subspace $\mathcal{M} \subseteq \mathfrak{X}$ is ultrainvariant for A if and only if $\mathcal{M} = \ker(A^j)$ for some $j \in \{0, 1, ..., n\}$.

Proof. By Proposition 5.4, every kernel $\ker(A^j)$ (j = 0, 1, ..., n) is an ultrainvariant subspace of A. For the opposite implication, suppose that \mathcal{M} is an ultrainvariant subspace for A. Then \mathcal{M} is hyperinvariant for A and therefore, by [3, Lemma 5], there exists a unique $j \in \{0, 1, ..., n\}$ such that $\overline{\operatorname{im}(A^{n-j})} \subseteq \mathcal{M} \subseteq \ker(A^j)$.

We claim that $\mathcal{C}(A, \overline{\mathrm{im}(A^{n-j})}) \overline{\mathrm{im}(A^{n-j})} = \ker(A^j)$, for every $j \in \{0, 1, \ldots, n\}$. If j = 0or j = n, then the assertion is trivial. Assume therefore that $1 \leq j \leq n-1$. Let $T \in \mathcal{C}(A, \overline{\mathrm{im}(A^{n-j})})$ and $x \in \mathrm{im}(A^{n-j})$ be arbitrary. Let $z \in \mathcal{X}$ be such that $x = A^{n-j}z$. By Lemma 5.3, $A^jTx = TA^jx = TA^nz = 0$. This proves that $\mathcal{C}(A, \overline{\mathrm{im}(A^{n-j})}) \overline{\mathrm{im}(A^{n-j})} \subseteq \ker(A^j)$. For the opposite inclusion, assume that $x \in \ker(A^j)$. Let $e \in \mathrm{im}(A^{n-j})$ be such that $A^{j-1}e \neq 0$ (it exists by Lemma 5.9 (i)). Since, by Lemma 5.9 (ii), vectors $e, Ae, \ldots, A^{j-1}e$ are linearly independent there exists $\xi \in \mathfrak{X}^*$ such that $\langle A^{j-1}e, \xi \rangle = 1$ and $\langle A^i e, \xi \rangle = 0$ for $i = 0, \ldots, j - 2$. Denote $T = (x \otimes \xi)A^{j-1} + A(x \otimes \xi)A^{j-2} + \cdots + A^{j-1}(x \otimes \xi)$. By Lemma 5.10, $T \in \mathcal{C}(A, \overline{\mathrm{im}(A^{n-j})})$. Since Te = x the equality $\mathcal{C}(A, \overline{\mathrm{im}(A^{n-j})}) \overline{\mathrm{im}(A^{n-j})} = \ker(A^j)$ is proved. By Proposition 4.6,

$$\mathcal{C}(A; \mathcal{C}(A, \overline{\operatorname{im}(A^{n-j})})\overline{\operatorname{im}(A^{n-j})}) \mathcal{C}(A, \overline{\operatorname{im}(A^{n-j})})\overline{\operatorname{im}(A^{n-j})} = \mathcal{C}(A; \ker(A^j)) \ker(A^j) = \ker(A^j)$$

is the smallest ultrainvariant subspace of A which contains $\overline{\operatorname{im}(A^{n-j})}$. Hence, $\mathcal{M} = \ker(A^j)$.
$$\square$$

An operator $A \in B(\mathfrak{X})$ is algebraic if there exists a non-zero polynomial p such that p(A) = 0. It is not hard to see that for an algebraic operator A there exists a unique monic polynomial q_A of the minimal degree, called the minimal polynomial of A, such that $q_A(A) = 0$. Every operator on a finite dimensional Banach space is algebraic. Let $q_A(z) = (z - \lambda_1)^{n_1} \cdots (z - \lambda_k)^{n_k}$ with $\lambda_1, \ldots, \lambda_k$ distinct. For every $j = 1, \ldots, k$, let $\mathfrak{X}_j = \ker((A - \lambda_j I_x)^{n_i})$. Then $\mathfrak{X} = \mathfrak{X}_1 \oplus \cdots \oplus \mathfrak{X}_k$ and with respect to this decomposition $A = (\lambda_1 I_{\mathfrak{X}_1} + A_1) \oplus \cdots \oplus (\lambda_k I_{\mathfrak{X}_k} + A_k)$, where $A_j \in B(\mathfrak{X}_j)$ is a nilpotent operator of order

 n_j (if $n_j = 1$, then $A_j = 0$). The reader is referred to [19, §5.9] for details about algebraic operators on a complex Banach space.

Corollary 5.12. Let $A \in B(\mathfrak{X})$ be algebraic and let $A = (\lambda_1 I_{\mathfrak{X}_1} + A_1) \oplus \cdots \oplus (\lambda_k I_{\mathfrak{X}_k} + A_k)$ be its decomposition as described above. Then a closed subspace $\mathcal{M} \subseteq \mathfrak{X}$ is an ultrainvariant subspace of A if and only if $\mathcal{M} = \ker(A_1^{m_1}) \oplus \cdots \oplus \ker(A_k^{m_k})$ for some $0 \leq m_j \leq n_j$ $(j = 1, \dots, k)$.

Proof. Assume that \mathcal{M} is an ultrainvariant subspace of A. Let $\mathcal{M} = \mathcal{M}_1 \oplus \cdots \oplus \mathcal{M}_k$, where $\mathcal{M}_j \subseteq \mathcal{X}_j$ $(j = 1, \ldots, k)$. Since every subspace \mathcal{X}_j is reducing subspace of A, it follows, by Proposition 4.4, that \mathcal{M}_j is an ultrainvariant subspace of $\lambda_1 I_{\mathcal{X}_j} + A_j$. By Theorem 5.11, there exists $m_j \in \{0, 1, \ldots, n_j\}$ such that $\mathcal{M}_j = \ker(A_j^{m_j})$. The opposite implication follows by Proposition 4.5 because $\sigma(\lambda_i I_{\mathcal{X}_i} + A_i) \cap \sigma(\lambda_j I_{\mathcal{X}_j} + A_j) = \emptyset$ whenever $i \neq j$. \Box

6. Ultrainvariant subspaces of operators on a Hilbert space

In this section, \mathcal{H} is the infinite dimensional separable complex Hilbert space. The inner product of vectors $x, y \in \mathcal{H}$ is denoted by (x, y). For a closed subspace $\mathcal{M} \subseteq \mathcal{H}$, let \mathcal{M}^{\perp} be its orthogonal complement.

Our first result in this section is the complete description of the lattice of ultrainvariant subspaces of a normal operator. It turns out that $\operatorname{Lat}_u(A) = \operatorname{Lat}_h(A)$ whenever A is normal.

Let A be a normal operator on \mathcal{H} and let E be its spectral measure. For a Borel subset $\sigma \subseteq \sigma(A)$, operator $E(\sigma) \in B(\mathcal{H})$ is an orthogonal projection. Subspace $\mathcal{M} = \operatorname{im}(E(\sigma))$ is reducing for A and $\mathcal{M}^{\perp} = \operatorname{im}(E(\sigma(A) \setminus \sigma))$ because $E(\sigma(A) \setminus \sigma) = I - E(\sigma)$ (see Section XXXI.7 in [11] for details).

Theorem 6.1. Let $A \in B(\mathcal{H})$ be a normal operator and let E be its spectral measure. A subspace $\mathcal{M} \subseteq \mathcal{H}$ is an ultrainvariant subspace of A if and only if \mathcal{M} is a hyperinvariant subspace of A, that is, if and only if there exists a Borel subset $\sigma \subseteq \sigma(A)$ such that $\mathcal{M} = \operatorname{im}(E(\sigma))$.

Proof. By [7, Theorem 3], $\mathcal{M} \subseteq \mathcal{H}$ is a hyperinvariant subspace of A if and only if there exists a Borel subset $\sigma \subseteq \sigma(A)$ such that $\mathcal{M} = \operatorname{im}(E(\sigma))$. Hence, it remains to prove that for every Borel subset $\sigma \subseteq \sigma(A)$ the subspace $\mathcal{M} = \operatorname{im}(E(\sigma))$ is ultrainvariant. We can write $A = A_{\mathcal{M}} \oplus A_{\mathcal{M}^{\perp}}$ with respect to the decomposition $\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^{\perp}$. We have

to see that $\mathcal{I}(A_{\mathfrak{M}}, A_{\mathfrak{M}^{\perp}}) = \{0\}$ because then, by Proposition 5.5, \mathcal{M} is an ultrainvariant subspace of A. There are a few ways how to see that $\mathcal{I}(A_{\mathfrak{M}}, A_{\mathfrak{M}^{\perp}}) = \{0\}$. We will show it as follows. Assume that $S \in \mathcal{I}(A_{\mathfrak{M}}, A_{\mathfrak{M}^{\perp}})$. Then operator $T \in B(\mathcal{H})$ whose operator matrix with respect to the decomposition $\mathcal{H} = \mathcal{M} \oplus \mathcal{M}^{\perp}$ is $\begin{bmatrix} 0 & 0 \\ S & 0 \end{bmatrix}$ commutes with A. It follows, by The Fuglede-Putnam Theorem, that T commutes with $E(\sigma)$ (see Theorems XXXI.7.4 and XXXI.7.9 in [11], for instance). It is easy to see now that $TE(\sigma) = E(\sigma)T$ gives S = 0.

Let $1 \leq p < \infty$ and let (X, μ) be a measure space such that $L^p(X, \mu)$ is separable. For $\phi \in L^{\infty}(X, \mu)$, let M_{ϕ} be the multiplication by ϕ in $L^p(X, \mu)$. By [12, Theorem 1], a closed subspace $\mathcal{M} \subseteq L^p(X, \mu)$ is hyperinvariant for M_{ϕ} if and only if $\mathcal{M} = M_{\chi_{\sigma} \circ \phi}(L^p(X, \mu))$ for some Borel set $\sigma \subseteq X$. Here χ_{σ} denotes the characteristic function of σ and $M_{\chi_{\sigma} \circ \phi}$ is the multiplication operator induced by $\chi_{\sigma} \circ \phi$. Note that $M_{\chi_{\sigma} \circ \phi}$ is an idempotent. Hence $M_{\chi_{\sigma} \circ \phi}(L^p(X, \mu))$ is complemented and its complement $M_{\chi_{X\setminus \sigma} \circ \phi}(L^p(X, \mu))$ is a hyperinvariant subspace of M_{ϕ} . A similar reasoning as in the proof of Theorem 6.1 gives that every hyperinvariant subspace of M_{ϕ} is an ultrainvariant subspace.

For a closed subspace $\mathcal{M} \subseteq \mathcal{H}$, let $P_{\mathcal{M}}$ be the orthogonal projection onto \mathcal{M} . Let $A \in B(\mathcal{H})$. A distance on $\operatorname{Lat}(A)$ is the mapping θ which is given by $\theta(\mathcal{M}, \mathcal{N}) = ||P_{\mathcal{M}} - P_{\mathcal{N}}||$ $(\mathcal{M}, \mathcal{N} \in \operatorname{Lat}(A))$. A subspace $\mathcal{M} \in \operatorname{Lat}(A)$ is said to be inaccessible if the only continuous mapping $f : [0, 1] \to \operatorname{Lat}(A)$ with $f(0) = \mathcal{M}$ is the constant map $f(t) = \mathcal{M}$ ($t \in [0, 1]$). A simple modification of the proof of [7, Theorem 1] gives the following result.

Theorem 6.2. Let $A \in B(\mathcal{H})$. If $\mathcal{M} \in Lat(A)$ is inaccessible, then it is an ultrainvariant subspace of A.

Proof. Let $0 \neq T \in \mathcal{C}(A; \mathcal{M})$ be an arbitrary operator. Denote $\Lambda = \{\lambda \in \mathbb{R}; 0 \leq \lambda < \|T\|^{-1}\}$. Then for each $\lambda \in \Lambda$, the operator $I_{\mathcal{H}} - \lambda T$ is invertible and therefore $\mathcal{M}_{\lambda} = (I_{\mathcal{H}} - \lambda T)\mathcal{M}$ is a closed subspace of \mathcal{H} . If $y \in \mathcal{M}_{\lambda}$, then there exists $x \in \mathcal{M}$ such that $y = (I_{\mathcal{H}} - \lambda T)x$. Hence, $Ay = A(I_{\mathcal{H}} - \lambda T)x = (I_{\mathcal{H}} - \lambda T)Ax \in \mathcal{M}_{\lambda}$ because $Ax \in \mathcal{M}$. This shows that $\mathcal{M}_{\lambda} \in \text{Lat}(A)$ for every $\lambda \in \Lambda$. The map $\lambda \mapsto \mathcal{M}_{\lambda}$ is continuous (see the proof of Theorem 1 in [7]). Since $\mathcal{M} = \mathcal{M}_0$ and \mathcal{M} is inaccessible we have $\mathcal{M} = \mathcal{M}_{\lambda}$ for each $\lambda \in \Lambda$. Thus, $\mathcal{M} = (I_{\mathcal{H}} - \lambda T)\mathcal{M}$ for a non-zero λ which implies that \mathcal{M} is invariant for T, that is, it is an ultrainvariant subspace of A.

It is clear that $\mathcal{M} \in \text{Lat}(A)$ is inaccessible if it is an isolated point.

Corollary 6.3. Let $A \in B(\mathcal{H})$. If $\mathcal{M} \in Lat(A)$ is such that $P_{\mathcal{M}}$ commutes with every $P_{\mathcal{N}}$ $(\mathcal{N} \in Lat(A))$, then \mathcal{M} is an ultrainvariant subspace of A.

Proof. Assume that $\mathcal{M} \in \text{Lat}(A)$ is such that $P_{\mathcal{M}}P_{\mathcal{N}} = P_{\mathcal{N}}P_{\mathcal{M}}$ for every $\mathcal{N} \in \text{Lat}(A)$. Then, for every $\mathcal{N} \in \text{Lat}(A)$, $P_{\mathcal{M}}P_{\mathcal{N}}$ is an orthogonal projection such that $P_{\mathcal{M}}P_{\mathcal{N}} \leq P_{\mathcal{M}}$ and $P_{\mathcal{M}}P_{\mathcal{N}} \leq P_{\mathcal{N}}$. Suppose that $\mathcal{N} \neq \mathcal{M}$. Then either $P_{\mathcal{M}}P_{\mathcal{N}} < P_{\mathcal{M}}$ or $P_{\mathcal{M}}P_{\mathcal{N}} < P_{\mathcal{N}}$. If the former holds, then there exists $x \in \text{im}(P_{\mathcal{M}})$, ||x|| = 1, such that $P_{\mathcal{M}}P_{\mathcal{N}}x = 0$. Since $P_{\mathcal{M}}x = x$ it follows that $P_{\mathcal{N}}x = P_{\mathcal{N}}P_{\mathcal{M}}x = 0$. Thus, $||P_{\mathcal{M}} - P_{\mathcal{N}}|| \geq ||P_{\mathcal{M}}x - P_{\mathcal{N}}x|| = ||x|| = 1$ which gives $||P_{\mathcal{M}} - P_{\mathcal{N}}|| = 1$. We get the same conclusion if $P_{\mathcal{M}}P_{\mathcal{N}} < P_{\mathcal{N}}$ holds. We conclude that \mathcal{M} is an isolated point of Lat(A). Hence, it is inaccessible and therefore it is an ultrainvariant subspace of A, by Theorem 6.3. □

Recall that $A \in B(\mathcal{H})$ is unicellular if Lat(A) is totally ordered, that is, for any two subspaces $\mathcal{M}, \mathcal{N} \in Lat(A)$, either $\mathcal{M} \subseteq \mathcal{N}$ or $\mathcal{M} \supseteq \mathcal{N}$.

Corollary 6.4. If $A \in B(\mathcal{H})$ is unicellular, then every $\mathcal{M} \in Lat(A)$ is an ultrainvariant subspace of A.

Proof. Assume that $A \in B(\mathcal{H})$ is unicellular and let $\mathcal{M} \in Lat(A)$ be arbitrary. If $\mathcal{N} \in Lat(A)$, then either $\mathcal{M} \subseteq \mathcal{N}$ or $\mathcal{M} \supseteq \mathcal{N}$ which implies that $P_{\mathcal{M}}$ and $P_{\mathcal{N}}$ commute. By Corollary 6.3, \mathcal{M} is an ultrainvariant subspace of A.

An example of a unicellular operator is the Volterra operator V on $L^2[0,1]$. By [8, Example 3] (see also [16, Theorem 4.14]), Lat(V) consists of subspaces $\mathcal{M}_{\alpha} = \{f \in L^2[0,1]; f = 0 \text{ a.e. on } [0,\alpha]\}$, where $\alpha \in [0,1]$. Hence, by Corollary 6.4, every invariant subspace of V is actually an ultrainvariant subspace.

Let $(e_n)_{n=0}^{\infty}$ be an orthonormal basis of \mathcal{H} and let $w = (w_n)_{n=1}^{\infty}$ be a bounded sequence of complex numbers. Then w determines an operator $W_w \in \mathcal{H}$ by $W_w e_0 = 0$ and $W_w e_n = w_n e_{n-1}$ for $n \in \mathbb{N}$ (hence, W_w is the weighted backward shift determined by w). For each $n \geq 0$, denote $\mathcal{M}_n = \bigvee \{e_0, \ldots, e_n\}$.

Proposition 6.5. Every $\mathcal{M}_n \subseteq \mathcal{H}$ $(n \ge 0)$ is an ultrainvariant subspace of W_w .

Proof. Since $\mathcal{M}_0 = \ker(W_w)$ and kernels are ultrainvariant (see the paragraph before Lemma 5.3) we may assume that $n \ge 1$. Observe that $W_w x \in \mathcal{M}_k$ for some $x \in \mathcal{H}$ and $k \ge 0$ if and only if $x \in \mathcal{M}_{k+1}$. Indeed, if $x = \sum_{j=0}^{\infty} (x, e_j)e_j$ and $(x, e_j) \ne 0$ for some j > k+1, then $W_w x = \sum_{j=1}^{\infty} (x, e_j)w_j e_{j-1} \notin \mathcal{M}_k$.

Let $T \in \mathcal{C}(W_w; \mathcal{M}_n)$ be arbitrary. Then $W_w Te_0 = TW_w e_0 = 0$ and therefore $Te_0 \in \mathcal{M}_0$. Suppose that for j < n we have $Te_j \in \mathcal{M}_j$. Then $W_w Te_{j+1} = TW_w e_{j+1} = w_j Te_j \in \mathcal{M}_j$ which gives $Te_{j+1} \in \mathcal{M}_{j+1}$. We conclude that \mathcal{M}_n is invariant for every $T \in \mathcal{C}(W_w; \mathcal{M}_n)$, that is, \mathcal{M}_n is ultrainvariant.

If w is a sequence of non-zero complex numbers such that $|w_1| > |w_2| > \cdots$ and $\sum_{n=1}^{\infty} |w_n|^2 < \infty$, then W_w is called the Donoghue operator with weight sequence w. By [16, Theorem 4.12] (a special case was considered by Donoghue [8, Example 1]), a closed subspace $\mathcal{M} \subseteq \mathcal{H}$ is a non-trivial invariant subspace of the Donoghue operator W_w if and only if $\mathcal{M} = \mathcal{M}_n$ for some $n \ge 0$. Hence the Donoghue operator is unicellular and therefore every \mathcal{M}_n is an ultrainvariant subspace of W_w .

Acknowledgments. The author was supported by the Slovenian Research Agency through the research program P2-0268.

References

- Y. A. Abramovich, C. D. Aliprantis, An invitation to operator theory, Graduate studies in mathematics, 50, AMS, Providence, RI, 2002.
- [2] S. A. Agryros, R. G. Haydon, A hereditarily indecomposable \mathcal{L}_{∞} -space that solves the scalar-pluscompact problem, Acta Math. 54 (2011), 1–54.
- [3] M. Barraa, Sous-espaces hyperinvariants d'un operateur nilpotent sur un espace de Banach, J. Operator Theory 21 (1989), 315–321.
- [4] J. Bračič, V. Müller, M. Zajac, Reflexivity and hyperreflexivity of the space of locally intertwining operators, J. Operator Theory 63:1 (2010), 101–114.
- [5] I. Chalendar, J. R. Partington, Modern Approaches to the Invariant Subspace Problem, Cambridge University Press, 2011.
- [6] X. Dai, D. R. Larson, Wandering Vectors for Unitary Systems and Orthogonal Wavelets, Mem. Amer. Math. Soc. 134 (1998), No. 640.
- [7] R. G. Douglas, C. Pearcy, On a topology for invariant subspaces, J. Funct. Anal. 2 (1968), 323–341.
- [8] W. F. Donoghue, Jr., The lattice of invariant subspaces of a completely continuous quasinilpotent transformation, Pacific J. Math. 7 (1957), 1031–1035.
- [9] P. Enflo, On the invariant subspace problem for Banach spaces, Acta Math. 158 (1987), 213–313.
- [10] I. Gohberg, S. Goldberg, M. A. Kaashoek, *Classes of linear operators, Vol. I*, Birkhäuser Verlag, 1990.
- [11] I. Gohberg, S. Goldberg, M. A. Kaashoek, Classes of linear operators, Vol. II, Birkhäuser Verlag, 1993.

- [12] S. Z. Huang, Each hyperinvariant subspace for a multiplication operator is spectral, Proc. Amer. Math. Soc. 106 (1989), 1057–1061.
- [13] D. R. Larson, Von Neumann algebras and wavelets, Operator algebras and applications (Samos, 1996), 267–312, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., 495, Kluwer Acad. Publ., Dordrecht, 1997.
- [14] K. B Laursen, M. M. Neumann, An introduction to local spectral theory, Clarendon Press, Oxford, 2000.
- [15] V. I. Lomonosov, Invariant subspaces of the family of operators that commute with a completely continuous operator. (Russian), Funkcional. Anal. i Priložen. 7 (1973), 55–56.
- [16] H. Radjavi, P. Rosenthal, Invariant Subspaces, Springer 1973.
- [17] C. J. Read, A solution to the invariant subspace problem, Bull. Lond. Math. Soc. 16 (1984), 337-401.
- [18] M. Rosenblum, On the operator equation BX XA = Q, Duke Math, J. 23 (1956), 263–269.
- [19] A. E. Taylor, Introduction to Functional Analysis, John Wiley & Sons, Inc., 1958.
- [20] A. Tcaciuc, On quasinilpotent operators and the invariant subspace problem, J. Math. Anal. Appl. 477 (2019), 187–195.

FACULTY OF NATURAL SCIENCES AND ENGINEERING, UNIVERSITY OF LJUBLJANA, AŠKERČEVA C. 12, SI-1000 LJUBLJANA, SLOVENIA

Email address: janko.bracic@ntf.uni-lj.si