

# Generalized atomic subspaces for operators in Hilbert spaces

**Prasenjit Ghosh**

Department of Pure Mathematics, University of Calcutta,  
35, Ballygunge Circular Road, Kolkata, 700019, West Bengal, India  
e-mail: prasenjitpuremath@gmail.com

**T. K. Samanta**

Department of Mathematics, Uluberia College,  
Uluberia, Howrah, 711315, West Bengal, India  
e-mail: mumpu\_tapas5@yahoo.co.in

## Abstract

*We introduce the notion of a  $g$ -atomic subspace for a bounded linear operator and construct several useful resolutions of the identity operator on a Hilbert space using the theory of  $g$ -fusion frames. Also we shall describe the concept of frame operator for a pair of  $g$ -fusion Bessel sequences and some of their properties.*

**Keywords:** *Frame, atomic subspace,  $g$ -fusion frame,  $K$ - $g$ -fusion frame.*

**2010 Mathematics Subject Classification:** *Primary 42C15; Secondary 46C07.*

## 1 Introduction

Frames for Hilbert spaces were first introduced by Duffin and Schaeffer [6] in 1952 to study some fundamental problems in non-harmonic Fourier series. Later on, after some decades, frame theory was popularized by Daubechies, Grossman, Meyer [4]. At present, frame theory has been widely used in signal and image processing, filter bank theory, coding and communications, system modeling and so on. Several generalizations of frames namely,  $K$ -frames,  $g$ -frames, fusion frames etc. have been introduced in recent times.

$K$ -frames were introduced by L. Gavruta [8] to study the atomic system with respect to a bounded linear operator. Using frame theory techniques, the author also studied the atomic decompositions for operators on reproducing kernel Hilbert spaces [9]. Sun [15] introduced a  $g$ -frame and a  $g$ -Riesz basis in complex Hilbert spaces and discussed several properties of them. Huang [11] began to study  $K$ - $g$ -frame by combining  $K$ -frame and  $g$ -frame. P. Casazza [2] was first to introduce the notion of fusion frames or frames of subspaces and gave various ways to obtain a resolution of the identity operator from a fusion frame. The concept of an atomic subspace with respect to a bounded linear operator were introduced by A. Bhandari and S. Mukherjee [1]. Construction of  $K$ - $g$ -fusion frames and their dual were presented by Sadri and Rahimi [14] to generalize the theory of  $K$ -frame, fusion frame

and  $g$ -frame. P. Ghosh and T. K. Samanta [10] studied the stability of dual  $g$ -fusion frames in Hilbert spaces.

In this paper, we present some useful results about resolution of the identity operator on a Hilbert space using the theory of  $g$ -fusion frames. We give the notion of  $g$ -atomic subspace with respect to a bounded linear operator. The frame operator for a pair of  $g$ -fusion Bessel sequences are discussed and some properties are going to be established.

The paper is organized as follows; in Section 2, we briefly recall the basic definitions and results. Various ways of obtaining resolution of the identity operator on a Hilbert space in  $g$ -fusion frame are studied in Section 3.  $g$ -atomic subspaces are introduced and discussed in Section 4. In Section 5, frame operator for a pair of  $g$ -fusion Bessel sequences are given and establish various properties.

Throughout this paper,  $H$  is considered to be a separable Hilbert space with associated inner product  $\langle \cdot, \cdot \rangle$  and  $\{H_j\}_{j \in J}$  are the collection of Hilbert spaces, where  $J$  is subset of integers  $\mathbb{Z}$ .  $I_H$  is the identity operator on  $H$ .  $\mathcal{B}(H_1, H_2)$  is a collection of all bounded linear operators from  $H_1$  to  $H_2$ . In particular  $\mathcal{B}(H)$  denote the space of all bounded linear operators on  $H$ . For  $T \in \mathcal{B}(H)$ , we denote  $\mathcal{N}(T)$  and  $\mathcal{R}(T)$  for null space and range of  $T$ , respectively. Also,  $P_V \in \mathcal{B}(H)$  is the orthonormal projection onto a closed subspace  $V \subset H$ . Define the space

$$l^2\left(\{H_j\}_{j \in J}\right) = \left\{ \{f_j\}_{j \in J} : f_j \in H_j, \sum_{j \in J} \|f_j\|^2 < \infty \right\}$$

with inner product is given by  $\langle \{f_j\}_{j \in J}, \{g_j\}_{j \in J} \rangle = \sum_{j \in J} \langle f_j, g_j \rangle_{H_j}$ . Clearly  $l^2\left(\{H_j\}_{j \in J}\right)$  is a Hilbert space with the pointwise operations [14].

## 2 Preliminaries

**Theorem 2.1.** (*Douglas' factorization theorem*) [5] Let  $U, V \in \mathcal{B}(H)$ . Then the following conditions are equivalent:

- (1)  $\mathcal{R}(U) \subseteq \mathcal{R}(V)$ .
- (2)  $UU^* \leq \lambda^2 VV^*$  for some  $\lambda > 0$ .
- (3)  $U = VW$  for some bounded linear operator  $W$  on  $H$ .

**Theorem 2.2.** [12] The set  $\mathcal{S}(H)$  of all self-adjoint operators on  $H$  is a partially ordered set with respect to the partial order  $\leq$  which is defined as for  $T, S \in \mathcal{S}(H)$

$$T \leq S \Leftrightarrow \langle Tf, f \rangle \leq \langle Sf, f \rangle \quad \forall f \in H.$$

**Theorem 2.3.** [7] Let  $V \subset H$  be a closed subspace and  $T \in \mathcal{B}(H)$ . Then  $P_V T^* = P_V T^* P_{\overline{TV}}$ . If  $T$  is an unitary operator (i.e  $T^*T = I_H$ ), then  $P_{\overline{TV}} T = T P_V$ .

**Definition 2.4.** [3] A sequence  $\{f_j\}_{j \in J}$  of elements in  $H$  is a frame for  $H$  if there exist constants  $A, B > 0$  such that

$$A \|f\|^2 \leq \sum_{j \in J} |\langle f, f_j \rangle|^2 \leq B \|f\|^2 \quad \forall f \in H.$$

The constants  $A$  and  $B$  are called frame bounds.

**Definition 2.5.** [2] Let  $\{W_j\}_{j \in J}$  be a collection of closed subspaces of  $H$  and  $\{v_j\}_{j \in J}$  be a collection of positive weights. A family of weighted closed subspaces  $\{(W_j, v_j) : j \in J\}$  is called a fusion frame for  $H$  if there exist constants  $0 < A \leq B < \infty$  such that

$$A \|f\|^2 \leq \sum_{j \in J} v_j^2 \|P_{W_j}(f)\|^2 \leq B \|f\|^2 \quad \forall f \in H.$$

The constants  $A, B$  are called fusion frame bounds. If  $A = B$  then the fusion frame is called a tight fusion frame, if  $A = B = 1$  then it is called a Parseval fusion frame.

**Definition 2.6.** [1] Let  $\{W_j\}_{j \in J}$  be a family of closed subspaces of  $H$  and  $\{v_j\}_{j \in J}$  be a family of positive weights and  $K \in \mathcal{B}(H)$ . Then  $\{(W_j, v_j) : j \in J\}$  is said to be an atomic subspace of  $H$  with respect to  $K$  if following conditions hold:

(I)  $\sum_{j \in J} v_j f_j$  is convergent for all  $\{f_j\}_{j \in J} \in \left( \sum_{j \in J} \oplus W_j \right)_{l_2}$ .

(II) For every  $f \in H$ , there exists  $\{f_j\}_{j \in J} \in \left( \sum_{j \in J} \oplus W_j \right)_{l_2}$  such that

$$K(f) = \sum_{j \in J} v_j f_j \quad \text{and} \quad \|\{f_j\}\|_{\left( \sum_{j \in J} \oplus W_j \right)_{l_2}} \leq C \|f\|_H$$

for some  $C > 0$ , where

$$\left( \sum_{j \in J} \oplus W_j \right)_{l_2} = \left\{ \{f_j\}_{j \in J} : f_j \in W_j, \sum_{j \in J} \|f_j\|^2 < \infty \right\}$$

with inner product is given by  $\langle \{f_j\}_{j \in J}, \{g_j\}_{j \in J} \rangle = \sum_{j \in J} \langle f_j, g_j \rangle_H$ .

**Definition 2.7.** [15] A sequence  $\{\Lambda_j \in \mathcal{B}(H, H_j) : j \in J\}$  is called a generalized frame or g-frame for  $H$  with respect to  $\{H_j\}_{j \in J}$  if there are two positive constants  $A$  and  $B$  such that

$$A \|f\|^2 \leq \sum_{j \in J} \|\Lambda_j f\|^2 \leq B \|f\|^2 \quad \forall f \in H.$$

The constants  $A$  and  $B$  are called the lower and upper frame bounds, respectively.

**Definition 2.8.** [13, 14] Let  $\{W_j\}_{j \in J}$  be a collection of closed subspaces of  $H$  and  $\{v_j\}_{j \in J}$  be a collection of positive weights and let  $\Lambda_j \in \mathcal{B}(H, H_j)$  for each  $j \in J$ . Then the family  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  is called a generalized fusion frame or a  $g$ -fusion frame for  $H$  respect to  $\{H_j\}_{j \in J}$  if there exist constants  $0 < A \leq B < \infty$  such that

$$A \|f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2 \quad \forall f \in H. \quad (1)$$

The constants  $A$  and  $B$  are called the lower and upper bounds of  $g$ -fusion frame, respectively. If  $A = B$  then  $\Lambda$  is called tight  $g$ -fusion frame and if  $A = B = 1$  then we say  $\Lambda$  is a Parseval  $g$ -fusion frame. If  $\Lambda$  satisfies only the condition

$$\sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2 \quad \forall f \in H$$

then it is called a  $g$ -fusion Bessel sequence with bound  $B$  in  $H$ .

**Definition 2.9.** [14] Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion Bessel sequence in  $H$  with a bound  $B$ . The synthesis operator  $T_\Lambda$  of  $\Lambda$  is defined as

$$T_\Lambda : l^2(\{H_j\}_{j \in J}) \rightarrow H,$$

$$T_\Lambda \left( \{f_j\}_{j \in J} \right) = \sum_{j \in J} v_j P_{W_j} \Lambda_j^* f_j \quad \forall \{f_j\}_{j \in J} \in l^2(\{H_j\}_{j \in J})$$

and the analysis operator is given by

$$T_\Lambda^* : H \rightarrow l^2(\{H_j\}_{j \in J}), \quad T_\Lambda^*(f) = \{v_j \Lambda_j P_{W_j}(f)\}_{j \in J} \quad \forall f \in H.$$

The  $g$ -fusion frame operator  $S_\Lambda : H \rightarrow H$  is defined as follows:

$$S_\Lambda(f) = T_\Lambda T_\Lambda^*(f) = \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* \Lambda_j P_{W_j}(f)$$

and it can be easily verify that

$$\langle S_\Lambda(f), f \rangle = \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \quad \forall f \in H.$$

Furthermore, if  $\Lambda$  is a  $g$ -fusion frame with bounds  $A$  and  $B$  then from (1),

$$\langle Af, f \rangle \leq \langle S_\Lambda(f), f \rangle \leq \langle Bf, f \rangle \quad \forall f \in H.$$

The operator  $S_\Lambda$  is bounded, self-adjoint, positive and invertible. Now, according to the Theorem (2.2), we can write,  $AI_H \leq S_\Lambda \leq BI_H$  and this gives

$$B^{-1}I_H \leq S_\Lambda^{-1} \leq A^{-1}I_H.$$

**Definition 2.10.** [14] Let  $\{W_j\}_{j \in J}$  be a collection of closed subspaces of  $H$  and  $\{v_j\}_{j \in J}$  be a collection of positive weights and let  $\Lambda_j \in \mathcal{B}(H, H_j)$  for each  $j \in J$  and  $K \in \mathcal{B}(H)$ . Then the family  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  is called a  $K$ - $g$ -fusion frame for  $H$  if there exist constants  $0 < A \leq B < \infty$  such that

$$A \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2 \quad \forall f \in H. \quad (2)$$

**Theorem 2.11.** [14] Let  $\Lambda$  be a  $g$ -fusion Bessel sequence in  $H$ . Then  $\Lambda$  is a  $K$ - $g$ -fusion frame for  $H$  if and only if there exists  $A > 0$  such that  $S_\Lambda \geq AKK^*$ .

**Definition 2.12.** [2] A family of bounded operators  $\{T_j\}_{j \in J}$  on  $H$  is called a resolution of identity operator on  $H$  if for all  $f \in H$ , we have  $f = \sum_{j \in J} T_j(f)$ , provided the series converges unconditionally for all  $f \in H$ .

### 3 Resolution of the identity operator in $g$ -fusion frame

In this section, we present several useful results of resolution of the identity operator on a Hilbert space using the theory of  $g$ -fusion frames.

**Theorem 3.1.** Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H$  with frame bounds  $C, D$  and  $S_\Lambda$  be its associated  $g$ -fusion frame operator. Then the family  $\left\{v_j^2 P_{W_j} \Lambda_j^* T_j\right\}_{j \in J}$  is resolution of the identity operator on  $H$ , where  $T_j = \Lambda_j P_{W_j} S_\Lambda^{-1}$ ,  $j \in J$ . Furthermore, for all  $f \in H$ , we have

$$\frac{C}{D^2} \|f\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \leq \frac{D}{C^2} \|f\|^2.$$

*Proof.* For any  $f \in H$ , we have the reconstruction formula for  $g$ -fusion frame:

$$f = S_\Lambda S_\Lambda^{-1}(f) = \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} S_\Lambda^{-1}(f) = \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* T_j(f).$$

Thus,  $\left\{v_j^2 P_{W_j} \Lambda_j^* T_j\right\}_{j \in J}$  is a resolution of the identity operator on  $H$ . Since  $\Lambda$  is a  $g$ -fusion frame with bounds  $C$  and  $D$ , for each  $f \in H$ , we have

$$\begin{aligned} \sum_{j \in J} v_j^2 \|T_j(f)\|^2 &= \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j} S_\Lambda^{-1}(f)\|^2 \\ &\leq D \|S_\Lambda^{-1}(f)\|^2 \leq D \|S_\Lambda^{-1}\|^2 \|f\|^2 \\ &\leq \frac{D}{C^2} \|f\|^2 \text{ [since } D^{-1} I_H \leq S_\Lambda^{-1} \leq C^{-1} I_H \text{]}. \end{aligned}$$

On the other hand,

$$\sum_{j \in J} v_j^2 \|T_j(f)\|^2 = \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j} S_\Lambda^{-1}(f)\|^2$$

$$\geq C \|S_{\Lambda}^{-1}(f)\|^2 \geq \frac{C}{D^2} \|f\|^2.$$

Therefore,

$$\frac{C}{D^2} \|f\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \leq \frac{D}{C^2} \|f\|^2 \quad \forall f \in H.$$

□

**Theorem 3.2.** Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H$  with frame bounds  $C, D$  and let  $T_j : H \rightarrow H_j$  be a bounded operator such that  $\left\{v_j^2 P_{W_j} \Lambda_j^* T_j\right\}_{j \in J}$  is a resolution of the identity operator on  $H$ . Then

$$\frac{1}{D} \left\| \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* T_j(f) \right\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \quad \forall f \in H.$$

*Proof.* Assume  $I \subset J$  with  $|I| < \infty$ . If our inequality holds for all finite subsets then it would hold for all subset. Let  $f \in H$  and set  $g = \sum_{j \in I} v_j^2 P_{W_j} \Lambda_j^* T_j(f)$ . Then

$$\begin{aligned} \|g\|^4 &= (\langle g, g \rangle)^2 = \left( \left\langle g, \sum_{j \in I} v_j^2 P_{W_j} \Lambda_j^* T_j(f) \right\rangle \right)^2 \\ &= \left( \sum_{j \in I} v_j \langle \Lambda_j P_{W_j}(g), v_j T_j(f) \rangle \right)^2 \\ &\leq \left( \sum_{j \in I} v_j \| \Lambda_j P_{W_j}(g) \| \| v_j T_j(f) \| \right)^2 \\ &\leq \left( \sum_{j \in I} v_j^2 \| \Lambda_j P_{W_j}(g) \|^2 \right) \left( \sum_{j \in I} \| v_j T_j(f) \|^2 \right) \\ &\leq D \|g\|^2 \sum_{j \in I} \| v_j T_j(f) \|^2 \quad [\text{since } \Lambda \text{ is a } g\text{-fusion frame}]. \\ &\Rightarrow \frac{1}{D} \|g\|^2 \leq \sum_{j \in I} \| v_j T_j(f) \|^2 \\ &\Rightarrow \frac{1}{D} \left\| \sum_{j \in I} v_j^2 P_{W_j} \Lambda_j^* T_j(f) \right\|^2 \leq \sum_{j \in I} v_j^2 \|T_j(f)\|^2 \quad \forall f \in H. \end{aligned}$$

Since the inequality holds for any finite subset  $I \subset J$ , we have

$$\frac{1}{D} \left\| \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* T_j(f) \right\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \quad \forall f \in H.$$

This completes the proof.  $\square$

**Theorem 3.3.** *Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H$  with frame bounds  $C, D$  and let  $T_j : H \rightarrow H_j$  be a bounded operator such that  $\{v_j^2 P_{W_j} \Lambda_j^* T_j\}_{j \in J}$  is a resolution of the identity operator on  $H$ . If  $T_j^* \Lambda_j P_{W_j} = T_j$ , then*

$$\frac{1}{D} \|f\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \leq D E \|f\|^2 \quad \forall f \in H,$$

where  $E = \sup_j \|T_j\|^2 < \infty$ .

*Proof.* Since  $\{v_j^2 P_{W_j} \Lambda_j^* T_j\}_{j \in J}$  is a resolution of the identity on  $H$ ,

$$f = \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* T_j(f), \quad f \in H.$$

Now, for each  $f \in H$ , using Theorem (3.2), we get

$$\begin{aligned} \frac{1}{D} \|f\|^2 &= \frac{1}{D} \left\| \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* T_j(f) \right\|^2 \leq \sum_{j \in J} v_j^2 \|T_j(f)\|^2 \\ &= \sum_{j \in J} v_j^2 \|T_j^* \Lambda_j P_{W_j}(f)\|^2 \quad [\text{since } T_j^* \Lambda_j P_{W_j} = T_j] \\ &\leq \sum_{j \in J} v_j^2 \|T_j\|^2 \|\Lambda_j P_{W_j}(f)\|^2 \\ &\leq E \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \quad [\text{using } E = \sup_j \|T_j\|^2] \\ &\leq D E \|f\|^2 \quad [\text{since } \Lambda \text{ is a } g\text{-fusion frame}]. \end{aligned}$$

This completes the proof.  $\square$

**Theorem 3.4.** *Let  $\{W_j\}_{j \in J}$  be a family of closed subspaces of  $H$  and  $\{v_j\}_{j \in J}$  be a family of bounded weights and let  $\Lambda_j \in \mathcal{B}(H, H_j)$ ,  $j \in J$ . Then  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  is a  $g$ -fusion frame for  $H$  if the following conditions are hold:*

(I) *For all  $f \in H$ , there exists  $A > 0$  such that*

$$\sum_{j \in J} \|\Lambda_j P_{W_j}(f)\|^2 \leq \frac{1}{A} \|f\|^2.$$

(II)  $\left\{ v_j P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} \right\}_{j \in J}$  is a resolution of the identity operator on  $H$ .

*Proof.* Since  $\left\{ v_j P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} \right\}_{j \in J}$  is a resolution of the identity operator on  $H$ , for  $f \in H$ , we have

$$f = \sum_{j \in J} v_j P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} (f).$$

By Cauchy-Schwarz inequality, we have

$$\begin{aligned} \|f\|^4 &= (\langle f, f \rangle)^2 = \left( \left\langle \sum_{j \in J} v_j P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} (f), f \right\rangle \right)^2 \\ &= \left( \sum_{j \in J} v_j \langle \Lambda_j P_{W_j} (f), \Lambda_j P_{W_j} (f) \rangle \right)^2 = \left( \sum_{j \in J} v_j \| \Lambda_j P_{W_j} (f) \|^2 \right)^2 \\ &\leq \left( \sum_{j \in J} \| \Lambda_j P_{W_j} (f) \|^2 \right) \left( \sum_{j \in J} v_j^2 \| \Lambda_j P_{W_j} (f) \|^2 \right) \\ &\leq \frac{1}{A} \|f\|^2 \sum_{j \in J} v_j^2 \| \Lambda_j P_{W_j} (f) \|^2 \quad [\text{using given condition (I)}] \\ &\Rightarrow A \|f\|^2 \leq \sum_{j \in J} v_j^2 \| \Lambda_j P_{W_j} (f) \|^2. \end{aligned}$$

On the other hand,

$$\begin{aligned} \sum_{j \in J} v_j^2 \| \Lambda_j P_{W_j} (f) \|^2 &\leq B \sum_{j \in J} \| \Lambda_j P_{W_j} (f) \|^2 \quad [\text{where } B = \sup_{j \in J} \{ v_j^2 \}] \\ &\leq \frac{B}{A} \|f\|^2 \quad [\text{using given condition (I)}] \end{aligned}$$

and hence  $\Lambda$  is a  $g$ -fusion frame.  $\square$

## 4 $g$ -atomic subspace

In this section, we define generalized atomic subspace or  $g$ -atomic subspace of a Hilbert space with respect to a bounded linear operator.

**Definition 4.1.** Let  $K \in \mathcal{B}(H)$  and  $\{W_j\}_{j \in J}$  be a collection of closed subspaces of  $H$ , let  $\{v_j\}_{j \in J}$  be a collection of positive weights and  $\Lambda_j \in \mathcal{B}(H, H_j)$  for each  $j \in J$ . Then the family  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  is said to be a generalized atomic subspace or  $g$ -atomic subspace of  $H$  with respect to  $K$  if the following statements hold:



(I)  $\Lambda$  is a  $g$ -fusion Bessel sequence in  $H$ .

(II) For every  $f \in H$ , there exists  $\{f_j\}_{j \in J} \in l^2(\{H_j\}_{j \in J})$  such that

$$K(f) = \sum_{j \in J} v_j P_{W_j} \Lambda_j^* f_j \quad \text{and} \quad \left\| \{f_j\}_{j \in J} \right\|_{l^2(\{H_j\}_{j \in J})} \leq C \|f\|_H$$

for some  $C > 0$ .

**Theorem 4.2.** Let  $K \in \mathcal{B}(H)$  and  $\{W_j\}_{j \in J}$  be a collection of closed subspaces of  $H$ , let  $\{v_j\}_{j \in J}$  be a collection of positive weights and  $\Lambda_j \in \mathcal{B}(H, H_j)$  for each  $j \in J$ . Then the following statements are equivalent:

(I)  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .

(II)  $\Lambda$  is a  $K$ - $g$ -fusion frame for  $H$ .

*Proof.* (I)  $\Rightarrow$  (II) Suppose  $\Lambda$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ . Then  $\Lambda$  is a  $g$ -fusion Bessel sequence, So, there exists  $B > 0$  such that

$$\sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j}(f) \right\|^2 \leq B \|f\|^2 \quad \forall f \in H.$$

Now, for any  $f \in H$ , we have

$$\|K^* f\| = \sup_{\|g\|=1} |\langle K^* f, g \rangle| = \sup_{\|g\|=1} |\langle f, K g \rangle|,$$

by definition (4.1), for  $g \in H$ , there exists  $\{f_j\}_{j \in J} \in l^2(\{H_j\}_{j \in J})$  such that

$$K(g) = \sum_{j \in J} v_j P_{W_j} \Lambda_j^* f_j \quad \text{and} \quad \left\| \{f_j\}_{j \in J} \right\|_{l^2(\{H_j\}_{j \in J})} \leq C \|g\|_H$$

for some  $C > 0$ . Thus

$$\begin{aligned} \|K^* f\| &= \sup_{\|g\|=1} \left| \left\langle f, \sum_{j \in J} v_j P_{W_j} \Lambda_j^* f_j \right\rangle \right| = \sup_{\|g\|=1} \left| \sum_{j \in J} v_j \langle \Lambda_j P_{W_j}(f), f_j \rangle \right| \\ &\leq \sup_{\|g\|=1} \left( \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j}(f) \right\|^2 \right)^{\frac{1}{2}} \left( \sum_{j \in J} \|f_j\|^2 \right)^{\frac{1}{2}} \\ &\leq C \sup_{\|g\|=1} \left( \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j}(f) \right\|^2 \right)^{\frac{1}{2}} \|g\| \\ &\Rightarrow \frac{1}{C^2} \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j}(f) \right\|^2. \end{aligned}$$

Therefore,  $\Lambda$  is a  $K$ - $g$ -fusion frame for  $H$  with bounds  $\frac{1}{C^2}$  and  $B$ .

(II)  $\Rightarrow$  (I) Suppose that  $\Lambda$  is a  $K$ - $g$ -fusion frame with the corresponding synthesis operator  $T_\Lambda$ . Then obviously  $\Lambda$  is a  $g$ -fusion Bessel sequence in  $H$ . Now, for each  $f \in H$ ,

$$A \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 = \|T_\Lambda^* f\|^2$$

gives  $AKK^* \leq T_\Lambda T_\Lambda^*$  and by Theorem (2.1),  $\exists L \in \mathcal{B}(H, l^2(\{H_j\}_{j \in J}))$  such that  $K = T_\Lambda L$ . Define  $L(f) = \{f_j\}_{j \in J}$ , for every  $f \in H$ . Then, for each  $f \in H$ , we have

$$K(f) = T_\Lambda L(f) = T_\Lambda \left( \{f_j\}_{j \in J} \right) = \sum_{j \in J} v_j P_{W_j} \Lambda_j^* f_j \text{ and}$$

$$\left\| \{f_j\}_{j \in J} \right\|_{l^2(\{H_j\}_{j \in J})} = \|L(f)\|_{l^2(\{H_j\}_{j \in J})} \leq C \|f\|,$$

where  $C = \|L\|$ . Hence,  $\Lambda$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .  $\square$

**Theorem 4.3.** Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H$ . Then  $\Lambda$  is a  $g$ -atomic subspace of  $H$  with respect to its  $g$ -fusion frame operator  $S_\Lambda$ .

*Proof.* Since  $\Lambda$  is a  $g$ -fusion frame in  $H$ , there exist  $A, B > 0$  such that

$$A \|f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2 \quad \forall f \in H.$$

Since  $\mathcal{R}(T_\Lambda) = H = \mathcal{R}(S_\Lambda)$ , by Theorem (2.1), there exists some  $\alpha > 0$  such that  $\alpha S_\Lambda S_\Lambda^* \leq T_\Lambda T_\Lambda^*$  and therefore for each  $f \in H$ , we have

$$\alpha \|S_\Lambda^* f\|^2 \leq \|T_\Lambda^* f\|^2 = \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B \|f\|^2.$$

Thus,  $\Lambda$  is a  $S_\Lambda$ - $g$ -fusion frame and hence by Theorem (4.2),  $\Lambda$  is a  $g$ -atomic subspace of  $H$  with respect to  $S_\Lambda$ .  $\square$

**Theorem 4.4.** Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  and  $\Gamma = \{(W_j, \Gamma_j, v_j)\}_{j \in J}$  be two  $g$ -atomic subspaces of  $H$  with respect to  $K \in \mathcal{B}(H)$  with the corresponding synthesis operators  $T_\Lambda$  and  $T_\Gamma$ , respectively. If  $T_\Lambda T_\Gamma^* = \theta_H$  ( $\theta_H$  is a null operator on  $H$ ) and  $U, V \in \mathcal{B}(H)$  such that  $U + V$  is invertible operator on  $H$  with  $K(U + V) = (U + V)K$ , then

$$\left\{ \left( (U + V)W_j, (\Lambda_j + \Gamma_j) P_{W_j} (U + V)^*, v_j \right) \right\}_{j \in J}$$

is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .

*Proof.* Since  $\Lambda$  and  $\Gamma$  are *g*-atomic subspaces with respect to  $K$ , by Theorem (4.2), they are  $K$ -*g*-fusion frames for  $H$ . So, for each  $f \in H$ , there exist positive constants  $(A_1, B_1)$  and  $(A_2, B_2)$  such that

$$A_1 \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \leq B_1 \|f\|^2 \text{ and}$$

$$A_2 \|K^* f\|^2 \leq \sum_{j \in J} v_j^2 \|\Gamma_j P_{W_j}(f)\|^2 \leq B_2 \|f\|^2.$$

Since  $T_\Lambda T_\Gamma^* = \theta_H$ , for any  $f \in H$ , we have

$$T_\Lambda \{v_j \Gamma_j P_{W_j}(f)\}_{j \in J} = \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* \Gamma_j P_{W_j}(f) = 0. \quad (3)$$

Also,  $U + V$  is invertible, so

$$\begin{aligned} \|K^* f\|^2 &= \left\| \left( (U + V)^{-1} \right)^* (U + V)^* K^* f \right\|^2 \\ &\leq \left\| (U + V)^{-1} \right\|^2 \| (U + V)^* K^* f \|^2. \end{aligned} \quad (4)$$

Now, for any  $f \in H$ , we have

$$\begin{aligned} &\sum_{j \in J} v_j^2 \left\| (\Lambda_j + \Gamma_j) P_{W_j} (U + V)^* P_{(U+V)W_j}(f) \right\|^2 \\ &= \sum_{j \in J} v_j^2 \left\| (\Lambda_j + \Gamma_j) P_{W_j} (U + V)^* (f) \right\|^2 \text{ [ using Theorem (2.3) ]} \\ &= \sum_{j \in J} v_j^2 \langle (\Lambda_j + \Gamma_j) P_{W_j} (T^* f), (\Lambda_j + \Gamma_j) P_{W_j} (T^* f) \rangle \text{ [ taking } T = U + V \text{ ]} \\ &= \sum_{j \in J} v_j^2 \left( \left\| \Lambda_j P_{W_j} (T^* f) \right\|^2 + \left\| \Gamma_j P_{W_j} (T^* f) \right\|^2 + 2 \operatorname{Re} \langle T P_{W_j} \Lambda_j^* \Gamma_j P_{W_j} (T^* f), f \rangle \right) \\ &= \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (T^* f) \right\|^2 + \sum_{j \in J} v_j^2 \left\| \Gamma_j P_{W_j} (T^* f) \right\|^2 \text{ [ using (3) ]} \\ &\leq B_1 \|T^* f\|^2 + B_2 \|T^* f\|^2 \text{ [ since } \Lambda, \Gamma \text{ are } K\text{-}g\text{-fusion frames ]} \\ &= (B_1 + B_2) \|(U + V)^* f\|^2 \text{ [ since } T = U + V \text{ ]} \\ &\leq (B_1 + B_2) \|U + V\|^2 \|f\|^2 \text{ [ as } U + V \text{ is bounded ].} \end{aligned}$$

On the other hand,

$$\begin{aligned} &\sum_{j \in J} v_j^2 \left\| (\Lambda_j + \Gamma_j) P_{W_j} (U + V)^* P_{(U+V)W_j}(f) \right\|^2 \\ &= \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (U + V)^* f \right\|^2 + \sum_{j \in J} v_j^2 \left\| \Gamma_j P_{W_j} (U + V)^* f \right\|^2 \end{aligned}$$

$$\begin{aligned}
&\geq \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (U + V)^* f \right\|^2 \\
&\geq A_1 \left\| K^* (U + V)^* f \right\|^2 \quad [\text{since } \Lambda \text{ is } K\text{-}g\text{-fusion frame}] \\
&= A_1 \left\| (U + V)^* K^* f \right\|^2 \quad [\text{using } K(U + V) = (U + V)K] \\
&\geq A_1 \left\| (U + V)^{-1} \right\|^{-2} \left\| K^* f \right\|^2 \quad [\text{using (4)}].
\end{aligned}$$

Therefore,  $\left\{ ((U + V)W_j, (\Lambda_j + \Gamma_j) P_{W_j} (U + V)^*, v_j) \right\}_{j \in J}$  is a  $K$ - $g$ -fusion frame and by Theorem (4.2), it is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .  $\square$

**Corollary 4.5.** *Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  and  $\Gamma = \{(W_j, \Gamma_j, v_j)\}_{j \in J}$  be two  $g$ -atomic subspaces of  $H$  with respect to  $K \in \mathcal{B}(H)$  with the corresponding synthesis operators  $T_\Lambda$  and  $T_\Gamma$ . If  $T_\Lambda T_\Gamma^* = \theta_H$  and  $U \in \mathcal{B}(H)$  is an invertible operator with  $KU = UK$ , then  $\{(UW_j, (\Lambda_j + \Gamma_j) P_{W_j} U^*, v_j)\}_{j \in J}$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .*

*Proof.* The proof of this Corollary directly follows from the Theorem (4.4), by putting  $V = \theta_H$ .  $\square$

**Theorem 4.6.** *Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -atomic subspace for  $K \in \mathcal{B}(H)$  and  $S_\Lambda$  be the frame operator of  $\Lambda$ . If  $U \in \mathcal{B}(H)$  be a positive and invertible operator on  $H$ , then  $\Lambda' = \{((I_H + U)W_j, \Lambda_j P_{W_j} (I_H + U)^*, v_j)\}_{j \in J}$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ . Moreover, for any natural number  $n$ ,  $\Lambda'' = \{((I_H + U^n)W_j, \Lambda_j P_{W_j} (I_H + U^n)^*, v_j)\}_{j \in J}$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .*

*Proof.* Since  $\Lambda$  is a  $g$ -atomic subspace with respect to  $K$ , by Theorem (4.2), it is a  $K$ - $g$ -fusion frame for  $H$ . Then, according to the Theorem (2.11), there exists  $A > 0$  such that  $S_\Lambda \geq AKK^*$ . Now, for each  $f \in H$ , we have

$$\begin{aligned}
&\sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (I_H + U)^* P_{(I_H + U)W_j} (f) \right\|^2 \\
&= \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (I_H + U)^* (f) \right\|^2 \quad [\text{using Theorem (2.3)}] \\
&\leq B \left\| (I_H + U)^* (f) \right\|^2 \quad [\text{since } \Lambda \text{ is a } K\text{-}g\text{-fusion frame}] \\
&\leq B \left\| I_H + U \right\|^2 \left\| f \right\|^2 \quad [\text{since } (I_H + U) \in \mathcal{B}(H)].
\end{aligned}$$

Thus  $\Lambda'$  is a  $g$ -fusion Bessel sequence in  $H$ . Also, for each  $f \in H$ , we have

$$\begin{aligned}
&\sum_{j \in J} v_j^2 P_{(I_H + U)W_j} (\Lambda_j P_{W_j} (I_H + U)^*)^* \Lambda_j P_{W_j} (I_H + U)^* P_{(I_H + U)W_j} (f) \\
&= \sum_{j \in J} v_j^2 P_{(I_H + U)W_j} (I_H + U) P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} (I_H + U)^* P_{(I_H + U)W_j} (f)
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{j \in J} v_j^2 \left( P_{W_j} (I_H + U)^* P_{(I_H + U)W_j} \right)^* \Lambda_j^* \Lambda_j \left( P_{W_j} (I_H + U)^* P_{(I_H + U)W_j} (f) \right) \\
 &= \sum_{j \in J} v_j^2 \left( P_{W_j} (I_H + U)^* \right)^* \Lambda_j^* \Lambda_j P_{W_j} (I_H + U)^* (f) \quad [\text{using Theorem (2.3)}] \\
 &= \sum_{j \in J} v_j^2 (I_H + U) P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} (I_H + U)^* (f) \\
 &= (I_H + U) \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* \Lambda_j P_{W_j} (I_H + U)^* (f) = (I_H + U) S_\Lambda (I_H + U)^* (f).
 \end{aligned}$$

This shows that the frame operator of  $\Lambda'$  is  $(I_H + U) S_\Lambda (I_H + U)^*$ . Now,

$$(I_H + U) S_\Lambda (I_H + U)^* \geq S_\Lambda \geq AKK^* \quad [\text{since } U, S_\Lambda \text{ are positive}].$$

Then by Theorem (2.11), we can conclude that  $\Lambda'$  is a  $K$ - $g$ -fusion frame and therefore by Theorem (4.2),  $\Lambda'$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ . According to the preceding procedure, for any natural number  $n$ , the frame operator of  $\Lambda''$  is  $(I_H + U^n) S_\Lambda (I_H + U^n)^*$  and similarly it can be shown that  $\Lambda''$  is a  $g$ -atomic subspace of  $H$  with respect to  $K$ .  $\square$

## 5 Frame operator for a pair of $g$ -fusion Bessel sequences

In this section, we shall discuss about the frame operator for a pair of  $g$ -fusion Bessel sequences and established some properties relative to frame operator. At the end of this section, we shall construct a new  $g$ -fusion frame for the Hilbert space  $H \oplus X$ , using the  $g$ -fusion frames of the Hilbert spaces  $H$  and  $X$ .

**Definition 5.1.** Let  $\Lambda = \{(W_j, \Lambda_j, w_j)\}_{j \in J}$  and  $\Gamma = \{(V_j, \Gamma_j, v_j)\}_{j \in J}$  be two  $g$ -fusion Bessel sequences in  $H$  with bounds  $D_1$  and  $D_2$ . Then the operator  $S_{\Gamma\Lambda} : H \rightarrow H$ , defined by

$$S_{\Gamma\Lambda}(f) = \sum_{j \in J} v_j w_j P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f) \quad \forall f \in H,$$

is called the frame operator for the pair of  $g$ -fusion Bessel sequences  $\Lambda$  and  $\Gamma$ .

**Theorem 5.2.** The frame operator  $S_{\Gamma\Lambda}$  for the pair of  $g$ -fusion Bessel sequences  $\Lambda$  and  $\Gamma$  is bounded and  $S_{\Gamma\Lambda}^* = S_{\Lambda\Gamma}$ .

*Proof.* For each  $f, g \in H$ , we have

$$\begin{aligned}
 \langle S_{\Gamma\Lambda}(f), g \rangle &= \left\langle \sum_{j \in J} v_j w_j P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f), g \right\rangle \\
 &= \sum_{j \in J} v_j w_j \langle \Lambda_j P_{W_j}(f), \Gamma_j P_{V_j}(g) \rangle. \quad (5)
 \end{aligned}$$

By the Cauchy-Schwarz inequality, we obtain  $|\langle S_{\Gamma\Lambda}(f), g \rangle|$

$$\begin{aligned} &\leq \left( \sum_{j \in J} v_j^2 \|\Gamma_j P_{V_j}(g)\|^2 \right)^{\frac{1}{2}} \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \quad (6) \\ &\leq \sqrt{D_2} \|g\| \sqrt{D_1} \|f\|. \end{aligned}$$

This shows that  $S_{\Gamma\Lambda}$  is a bounded operator with  $\|S_{\Gamma\Lambda}\| \leq \sqrt{D_1 D_2}$ . Now,

$$\begin{aligned} \|S_{\Gamma\Lambda} f\| &= \sup \{ |\langle S_{\Gamma\Lambda}(f), g \rangle| : \|g\| = 1 \} \\ &\leq \sup \left\{ \sqrt{D_2} \|g\| \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} : \|g\| = 1 \right\} \quad [\text{using (6)}] \\ &\leq \sqrt{D_2} \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \quad (7) \end{aligned}$$

and similarly it can be shown that

$$\|S_{\Gamma\Lambda}^* g\| \leq \sqrt{D_1} \left( \sum_{j \in J} v_j^2 \|\Gamma_j P_{V_j}(g)\|^2 \right)^{\frac{1}{2}}. \quad (8)$$

Also, for each  $f, g \in H$ , we have

$$\begin{aligned} \langle S_{\Gamma\Lambda}(f), g \rangle &= \left\langle \sum_{j \in J} v_j w_j P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f), g \right\rangle \\ &= \sum_{j \in J} v_j w_j \langle f, P_{W_j} \Lambda_j^* \Gamma_j P_{V_j}(g) \rangle \\ &= \left\langle f, \sum_{j \in J} w_j v_j P_{W_j} \Lambda_j^* \Gamma_j P_{V_j}(g) \right\rangle = \langle f, S_{\Lambda\Gamma}(g) \rangle \end{aligned}$$

and hence  $S_{\Gamma\Lambda}^* = S_{\Lambda\Gamma}$ .  $\square$

**Theorem 5.3.** *Let  $S_{\Gamma\Lambda}$  be the frame operator for a pair of  $g$ -fusion Bessel sequences  $\Lambda$  and  $\Gamma$  with bounds  $D_1$  and  $D_2$ , respectively. Then the following statements are equivalent:*

- (I)  $S_{\Gamma\Lambda}$  is bounded below.
- (II) There exists  $K \in \mathcal{B}(H)$  such that  $\{T_j\}_{j \in J}$  is a resolution of the identity operator on  $H$ , where  $T_j = v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}$ ,  $j \in J$ .

If one of the given conditions hold, then  $\Lambda$  is a *g*-fusion frame.

*Proof.* (I)  $\Rightarrow$  (II) Suppose that  $S_{\Gamma\Lambda}$  is bounded below. Then for each  $f \in H$ , there exists  $A > 0$  such that

$$\|f\|^2 \leq A \|S_{\Gamma\Lambda} f\|^2 \Rightarrow \langle I_H f, f \rangle \leq A \langle S_{\Gamma\Lambda}^* S_{\Gamma\Lambda} f, f \rangle \Rightarrow I_H^* I_H \leq A S_{\Gamma\Lambda}^* S_{\Gamma\Lambda}.$$

So, by Theorem (2.1), there exists  $K \in \mathcal{B}(H)$  such that  $K S_{\Gamma\Lambda} = I_H$ . Therefore, for each  $f \in H$ , we have

$$\begin{aligned} f &= K S_{\Gamma\Lambda}(f) = K \left( \sum_{j \in J} v_j w_j P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f) \right) \\ &= \sum_{j \in J} v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f) = \sum_{j \in J} T_j(f) \end{aligned}$$

and hence  $\{T_j\}_{j \in J}$  is a resolution of the identity operator on  $H$ , where  $T_j = v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}$ .

(II)  $\Rightarrow$  (I) Since  $\{T_j\}_{j \in J}$  is a resolution of the identity operator on  $H$ , for any  $f \in H$ , we have

$$\begin{aligned} f &= \sum_{j \in J} T_j(f) = \sum_{j \in J} v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f) \\ &= K \left( \sum_{j \in J} v_j w_j P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f) \right) = K S_{\Gamma\Lambda}(f). \end{aligned}$$

Thus,  $I_H = K S_{\Gamma\Lambda}$ . So, by Theorem (2.1), there exists some  $\alpha > 0$  such that  $I_H I_H^* \leq \alpha S_{\Gamma\Lambda} S_{\Gamma\Lambda}^*$  and hence  $S_{\Gamma\Lambda}$  is bounded below.

Last part: First we suppose that  $S_{\Gamma\Lambda}$  is bounded below. Then for all  $f \in H$ , there exists  $M > 0$  such that  $\|S_{\Gamma\Lambda} f\| \geq M \|f\|$  and this implies that

$$\begin{aligned} M^2 \|f\|^2 &\leq \|S_{\Gamma\Lambda} f\|^2 \leq D_2 \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \quad [\text{using (7)}] \\ &\Rightarrow \frac{M^2}{D_2} \|f\|^2 \leq \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2. \end{aligned}$$

Hence,  $\Lambda$  is a *g*-fusion frame for  $H$  with bounds  $\frac{M^2}{D_2}$  and  $D_1$ .

Next, we suppose that the given condition (II) holds. Then for any  $f \in H$ , we have

$$f = \sum_{j \in J} v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f), \quad K \in \mathcal{B}(H).$$

By Cauchy-Schwarz inequality, for each  $f \in H$ , we have

$$\begin{aligned}
\|f\|^2 &= \langle f, f \rangle = \left\langle \sum_{j \in J} v_j w_j K P_{V_j} \Gamma_j^* \Lambda_j P_{W_j}(f), f \right\rangle \\
&= \sum_{j \in J} v_j w_j \langle \Lambda_j P_{W_j}(f), \Gamma_j P_{V_j}(K^* f) \rangle \\
&\leq \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \left( \sum_{j \in J} v_j^2 \|\Gamma_j P_{V_j}(K^* f)\|^2 \right)^{\frac{1}{2}} \\
&\leq \sqrt{D_2} \|K^* f\| \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \\
&\leq \sqrt{D_2} \|K\| \|f\| \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \\
&\Rightarrow \frac{1}{D_2 \|K\|^2} \|f\|^2 \leq \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2.
\end{aligned}$$

Therefore, in this case  $\Lambda$  is also a  $g$ -fusion frame for  $H$ .  $\square$

**Theorem 5.4.** *Let  $S_{\Gamma\Lambda}$  be the frame operator for a pair of  $g$ -fusion Bessel sequences  $\Lambda$  and  $\Gamma$  with bounds  $D_1$  and  $D_2$ , respectively. Suppose  $\lambda_1 < 1$ ,  $\lambda_2 > -1$  such that for each  $f \in H$ ,*

$$\|f - S_{\Gamma\Lambda} f\| \leq \lambda_1 \|f\| + \lambda_2 \|S_{\Gamma\Lambda} f\|.$$

*Then  $\Lambda$  is a  $g$ -fusion frame for  $H$ .*

*Proof.* For each  $f \in H$ , we have

$$\begin{aligned}
\|f\| - \|S_{\Gamma\Lambda} f\| &\leq \|f - S_{\Gamma\Lambda} f\| \leq \lambda_1 \|f\| + \lambda_2 \|S_{\Gamma\Lambda} f\| \\
\Rightarrow (1 - \lambda_1) \|f\| &\leq (1 + \lambda_2) \|S_{\Gamma\Lambda} f\| \\
\Rightarrow \left( \frac{1 - \lambda_1}{1 + \lambda_2} \right) \|f\| &\leq \sqrt{D_2} \left( \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2 \right)^{\frac{1}{2}} \quad [\text{using (7)}] \\
\Rightarrow \frac{1}{D_2} \left( \frac{1 - \lambda_1}{1 + \lambda_2} \right)^2 \|f\|^2 &\leq \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2. \quad (9)
\end{aligned}$$

Thus,  $\Lambda$  is a  $g$ -fusion frame for  $H$  with bounds  $\frac{1}{D_2} \left( \frac{1 - \lambda_1}{1 + \lambda_2} \right)^2$  and  $D_1$ .  $\square$



**Theorem 5.5.** *Let  $S_{\Gamma\Lambda}$  be the frame operator for a pair of  $g$ -fusion Bessel sequences  $\Lambda$  and  $\Gamma$  of bounds  $D_1$  and  $D_2$ , respectively. Assume  $\lambda \in [0, 1)$  such that*

$$\|f - S_{\Gamma\Lambda} f\| \leq \lambda \|f\| \quad \forall f \in H.$$

Then  $\Lambda$  and  $\Gamma$  are  $g$ -fusion frames for  $H$ .

*Proof.* By putting  $\lambda_1 = \lambda$  and  $\lambda_2 = 0$  in (9), we get

$$\frac{(1 - \lambda)^2}{D_2} \|f\|^2 \leq \sum_{j \in J} w_j^2 \|\Lambda_j P_{W_j}(f)\|^2$$

and therefore  $\Lambda$  is a  $g$ -fusion frame. Now, for each  $f \in H$ , we have

$$\begin{aligned} \|f - S_{\Gamma\Lambda}^* f\| &= \|(I_H - S_{\Gamma\Lambda})^* f\| \leq \|(I_H - S_{\Gamma\Lambda})\| \|f\| \leq \lambda \|f\| \\ \Rightarrow (1 - \lambda) \|f\| &\leq \|S_{\Gamma\Lambda}^* f\| \leq \sqrt{D_1} \left( \sum_{j \in J} v_j^2 \|\Gamma_j P_{V_j}(f)\|^2 \right)^{\frac{1}{2}} \quad [\text{using (8)}] \\ \Rightarrow \sum_{j \in J} v_j^2 \|\Gamma_j P_{V_j}(f)\|^2 &\geq \frac{(1 - \lambda)^2}{D_1} \|f\|^2 \quad \forall f \in H. \end{aligned}$$

Hence,  $\Gamma$  is a  $g$ -fusion frame with bounds  $\frac{(1 - \lambda)^2}{D_1}$  and  $D_2$ . □

**Definition 5.6.** *Let  $H$  and  $X$  be two Hilbert spaces. Define*

$$H \oplus X = \{(f, g) : f \in H, g \in X\}. \text{ Then } H \oplus X \text{ forms a}$$

*Hilbert space with respect to point-wise operations and inner product defined by*

$$\langle (f, g), (f', g') \rangle = \langle f, f' \rangle_H + \langle g, g' \rangle_X \quad \forall f, f' \in H \ \& \ \forall g, g' \in X.$$

*Now, if  $U \in \mathcal{B}(H, Z)$ ,  $V \in \mathcal{B}(X, Y)$  then for all  $f \in H, g \in X$ , we define*

$$U \oplus V \in \mathcal{B}(H \oplus X, Z \oplus Y) \text{ by } (U \oplus V)(f, g) = (Uf, Vg),$$

*and  $(U \oplus V)^* = U^* \oplus V^*$ , where  $Z, Y$  are Hilbert spaces and also we define  $P_{M \oplus N}(f, g) = (P_M f, P_N g)$ , where  $P_M, P_N$  and  $P_{M \oplus N}$  are the orthonormal projection onto the closed subspaces  $M \subset H, N \subset X$  and  $M \oplus N \subset H \oplus X$ , respectively.*

From here we assume that for each  $j \in J$ ,  $W_j \oplus V_j$  are the closed subspaces of  $H \oplus X$  and  $\Gamma_j \in \mathcal{B}(X, X_j)$ , where  $\{X_j\}_{j \in J}$  are the collection of Hilbert spaces and  $\Lambda_j \oplus \Gamma_j \in \mathcal{B}(H \oplus X, H_j \oplus X_j)$ .

**Theorem 5.7.** Let  $\Lambda = \{(W_j, \Lambda_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H$  with bounds  $A, B$  and  $\Gamma = \{(V_j, \Gamma_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $X$  with bounds  $C, D$ . Then  $\Lambda \oplus \Gamma = \{(W_j \oplus V_j, \Lambda_j \oplus \Gamma_j, v_j)\}_{j \in J}$  is a  $g$ -fusion frame for  $H \oplus X$  with bounds  $\min\{A, C\}, \max\{B, D\}$ . Furthermore, if  $S_\Lambda, S_\Gamma$  and  $S_{\Lambda \oplus \Gamma}$  are  $g$ -fusion frame operators for  $\Lambda, \Gamma$  and  $\Lambda \oplus \Gamma$  respectively then we have  $S_{\Lambda \oplus \Gamma} = S_\Lambda \oplus S_\Gamma$ .

*Proof.* Let  $(f, g) \in H \oplus X$  be an arbitrary element. Then

$$\begin{aligned}
& \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \right\|^2 \\
&= \sum_{j \in J} v_j^2 \left\langle (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g), (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \right\rangle \\
&= \sum_{j \in J} v_j^2 \left\langle \Lambda_j \oplus \Gamma_j (P_{W_j} (f), P_{V_j} (g)), \Lambda_j \oplus \Gamma_j (P_{W_j} (f), P_{V_j} (g)) \right\rangle \\
&= \sum_{j \in J} v_j^2 \left\langle (\Lambda_j P_{W_j} (f), \Gamma_j P_{V_j} (g)), (\Lambda_j P_{W_j} (f), \Gamma_j P_{V_j} (g)) \right\rangle \\
&= \sum_{j \in J} v_j^2 \left( \left\langle \Lambda_j P_{W_j} (f), \Lambda_j P_{W_j} (f) \right\rangle_H + \left\langle \Gamma_j P_{V_j} (g), \Gamma_j P_{V_j} (g) \right\rangle_X \right) \\
&= \sum_{j \in J} v_j^2 \left( \left\| \Lambda_j P_{W_j} (f) \right\|_H^2 + \left\| \Gamma_j P_{V_j} (g) \right\|_X^2 \right) \\
&= \sum_{j \in J} v_j^2 \left\| \Lambda_j P_{W_j} (f) \right\|_H^2 + \sum_{j \in J} v_j^2 \left\| \Gamma_j P_{V_j} (g) \right\|_X^2 \\
&\leq B \|f\|_H^2 + D \|g\|_X^2 \quad [\text{since } \Lambda, \Gamma \text{ are } g\text{-fusion frames}] \\
&\leq \max\{B, D\} (\|f\|_H^2 + \|g\|_X^2) = \max\{B, D\} \|(f, g)\|^2.
\end{aligned}$$

Similarly, it can be shown that

$$\min\{A, C\} \|(f, g)\|^2 \leq \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \right\|^2.$$

Therefore, for all  $(f, g) \in H \oplus X$ , we have

$$A_1 \|(f, g)\|^2 \leq \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \right\|^2 \leq B_1 \|(f, g)\|^2$$

and hence  $\Lambda \oplus \Gamma$  is a  $g$ -fusion frame for  $H \oplus X$  with bounds  $A_1 = \min\{A, C\}$  and  $B_1 = \max\{B, D\}$ . Furthermore, for  $(f, g) \in H \oplus X$ , we have

$$\begin{aligned}
S_{\Lambda \oplus \Gamma} (f, g) &= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \\
&= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) (P_{W_j} (f), P_{V_j} (g))
\end{aligned}$$

$$\begin{aligned}
&= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j P_{W_j}(f), \Gamma_j P_{V_j}(g)) \\
&= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j^* \oplus \Gamma_j^*) (\Lambda_j P_{W_j}(f), \Gamma_j P_{V_j}(g)) \\
&= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j^* \Lambda_j P_{W_j}(f), \Gamma_j^* \Gamma_j P_{V_j}(g)) \\
&= \sum_{j \in J} v_j^2 (P_{W_j} \Lambda_j^* \Lambda_j P_{W_j}(f), P_{V_j} \Gamma_j^* \Gamma_j P_{V_j}(g)) \\
&= \left( \sum_{j \in J} v_j^2 P_{W_j} \Lambda_j^* \Lambda_j P_{W_j}(f), \sum_{j \in J} v_j^2 P_{V_j} \Gamma_j^* \Gamma_j P_{V_j}(g) \right) \\
&= (S_\Lambda(f), S_\Gamma(g)) = (S_\Lambda \oplus S_\Gamma)(f, g) \quad \forall (f, g) \in H \oplus X.
\end{aligned}$$

Hence,  $S_{\Lambda \oplus \Gamma} = S_\Lambda \oplus S_\Gamma$ . This completes the proof.  $\square$

**Theorem 5.8.** Let  $\Lambda \oplus \Gamma = \{(W_j \oplus V_j, \Lambda_j \oplus \Gamma_j, v_j)\}_{j \in J}$  be a g-fusion frame for  $H \oplus X$  with frame operator  $S_{\Lambda \oplus \Gamma}$ . Then

$$\Delta' = \left\{ \left( S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(W_j \oplus V_j), (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}, v_j \right) \right\}_{j \in J}$$

is a Parseval g-fusion frame for  $H \oplus X$ .

*Proof.* Since  $S_{\Lambda \oplus \Gamma}$  is a positive operator, there exists a unique positive square root  $S_{\Lambda \oplus \Gamma}^{\frac{1}{2}}$  (or  $S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}$ ) and they commute with  $S_{\Lambda \oplus \Gamma}$  and  $S_{\Lambda \oplus \Gamma}^{-1}$ . Therefore, each

$$\begin{aligned}
(f, g) \in H \oplus X, \text{ can be written as } (f, g) &= S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}} S_{\Lambda \oplus \Gamma} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(f, g) \\
&= \sum_{j \in J} v_j^2 S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}} P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(f, g).
\end{aligned}$$

Now, for each  $(f, g) \in H \oplus X$ , we have

$$\begin{aligned}
&\|(f, g)\|^2 = \langle (f, g), (f, g) \rangle \\
&= \left\langle \sum_{j \in J} v_j^2 S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}} P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(f, g), (f, g) \right\rangle \\
&= \sum_{j \in J} v_j^2 \left\langle (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(f, g), (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}}(f, g) \right\rangle
\end{aligned}$$

$$\begin{aligned}
&= \sum_{j \in J} v_j^2 \left\| \left( \Lambda_j \oplus \Gamma_j \right) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}} (f, g) \right\|^2 \\
&= \sum_{j \in J} v_j^2 \left\| \left( \Lambda_j \oplus \Gamma_j \right) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-\frac{1}{2}} P \left( \begin{array}{c} -\frac{1}{2} \\ S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j) \end{array} \right) (f, g) \right\|^2 \quad [\text{by Theorem (2.3)}].
\end{aligned}$$

This shows that  $\Delta'$  is a Parseval  $g$ -fusion frame for  $H \oplus X$ .  $\square$

**Theorem 5.9.** *Let  $\Lambda \oplus \Gamma = \{(W_j \oplus V_j, \Lambda_j \oplus \Gamma_j, v_j)\}_{j \in J}$  be a  $g$ -fusion frame for  $H \oplus X$  with bounds  $A_1, B_1$  and  $S_{\Lambda \oplus \Gamma}$  be the corresponding frame operator. Then*

$$\Delta = \left\{ \left( S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j), (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1}, v_j \right) \right\}_{j \in J}$$

is a  $g$ -fusion frame for  $H \oplus X$  with frame operator  $S_{\Lambda \oplus \Gamma}^{-1}$ .

*Proof.* For any  $(f, g) \in H \oplus X$ , we have

$$\begin{aligned}
(f, g) &= S_{\Lambda \oplus \Gamma} S_{\Lambda \oplus \Gamma}^{-1} (f, g) \\
&= \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} (f, g). \quad (10)
\end{aligned}$$

By Theorem (2.3), for any  $(f, g) \in H \oplus X$ , we have

$$\begin{aligned}
&\sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)} (f, g) \right\|^2 \\
&= \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} (f, g) \right\|^2 \quad (11) \\
&\leq B_1 \left\| S_{\Lambda \oplus \Gamma}^{-1} \right\|^2 \|(f, g)\|^2 \quad [\text{since } \Lambda \oplus \Gamma \text{ is } g\text{-fusion frame}].
\end{aligned}$$

On the other hand, using (10), we get

$$\begin{aligned}
&\|(f, g)\|^4 = |\langle (f, g), (f, g) \rangle|^2 \\
&= \left| \left\langle \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} (f, g), (f, g) \right\rangle \right|^2 \\
&= \left| \sum_{j \in J} v_j^2 \langle (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} (f, g), (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} (f, g) \rangle \right|^2
\end{aligned}$$

$$\begin{aligned}
 &\leq \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1}(f, g) \right\|^2 \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j}(f, g) \right\|^2 \\
 &\leq \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1}(f, g) \right\|^2 B_1 \|(f, g)\|^2 \text{ [as } \Lambda \oplus \Gamma \text{ is } g\text{-fusion frame]} \\
 &= B_1 \|(f, g)\|^2 \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)}(f, g) \right\|^2 \text{ [from (11)].}
 \end{aligned}$$

Therefore,

$$B_1^{-1} \|(f, g)\|^2 \leq \sum_{j \in J} v_j^2 \left\| (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)}(f, g) \right\|^2.$$

Hence,  $\Delta$  is a *g*-fusion frame for  $H \oplus X$ . Let  $S_\Delta$  be the *g*-fusion frame operator for  $\Delta$  and take  $\Delta_j = \Lambda_j \oplus \Gamma_j$ . Now, for each  $(f, g) \in H \oplus X$ ,  $S_\Delta(f, g)$

$$\begin{aligned}
 &= \sum_{j \in J} v_j^2 P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)} (\Delta_j P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1})^* (\Delta_j P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1}) P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)}(f, g) \\
 &= \sum_{j \in J} v_j^2 \left( P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)} \right)^* \Delta_j^* \Delta_j \left( P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} P_{S_{\Lambda \oplus \Gamma}^{-1}(W_j \oplus V_j)} \right) (f, g) \\
 &= \sum_{j \in J} v_j^2 \left( P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} \right)^* \Delta_j^* \Delta_j \left( P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} \right) (f, g) \text{ [ using Theorem (2.3) ]} \\
 &= \sum_{j \in J} v_j^2 S_{\Lambda \oplus \Gamma}^{-1} P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) \left( P_{W_j \oplus V_j} S_{\Lambda \oplus \Gamma}^{-1} \right) (f, g) \\
 &= S_{\Lambda \oplus \Gamma}^{-1} \left( \sum_{j \in J} v_j^2 P_{W_j \oplus V_j} (\Lambda_j \oplus \Gamma_j)^* (\Lambda_j \oplus \Gamma_j) P_{W_j \oplus V_j} \left( S_{\Lambda \oplus \Gamma}^{-1}(f, g) \right) \right) \\
 &= S_{\Lambda \oplus \Gamma}^{-1} S_{\Lambda \oplus \Gamma} \left( S_{\Lambda \oplus \Gamma}^{-1}(f, g) \right) \text{ [ by definition of } S_{\Lambda \oplus \Gamma} \text{].} \\
 &= S_{\Lambda \oplus \Gamma}^{-1}(f, g).
 \end{aligned}$$

Thus,  $S_\Delta = S_{\Lambda \oplus \Gamma}^{-1}$ . This completes the proof.  $\square$

**Note 5.10.** Form Theorem (5.9), we can conclude that if  $\Lambda \oplus \Gamma$  is a *g*-fusion frame for  $H \oplus K$ , then  $\Delta$  is also a *g*-fusion frame for  $H \oplus K$ . The *g*-fusion frame  $\Delta$  is called the canonical dual *g*-fusion frame of  $\Lambda \oplus \Gamma$ .

## References

- [1] A. Bhandari, A. Mukherjee, *Atomic subspaces for operators*, Submitted, arXiv: 1705. 06042.
- [2] P. Casazza, G. Kutyniok, *Frames of subspaces*, Cotemporary Math, AMS 345 (2004), 87-114.

- [3] O. Christensen, *An introduction to frames and Riesz bases*, Birkhauser (2008).
- [4] I. Daubechies, A. Grossmann, Y. Mayer, *Painless nonorthogonal expansions*, Journal of Mathematical Physics 27 (5) (1986) 1271-1283.
- [5] R. G. Douglas, *On majorization, factorization, and range inclusion of operators on Hilbert space*. Proc. Am. Math. Soc. 17, 413-415 (1966).
- [6] R. J. Duffin, A. C. Schaeffer, *A class of nonharmonic Fourier series*, Trans. Amer. Math. Soc., 72, (1952), 341-366.
- [7] P. Gavruta, *On the duality of fusion frames*, J. Math. Anal. Appl. 333 (2007) 871-879.
- [8] L. Gavruta, *Frames for operator*, Appl. Comput. Harmon. Anal. 32 (1), 139-144 (2012).
- [9] L. Gavruta, *Atomic decompositions for operators in reproducing kernel Hilbert spaces*, Math. Reports 17(67), 3(2015), 303-314.
- [10] P. Ghosh, T. K. Samanta, *Stability of dual  $g$ -fusion frame in Hilbert spaces*, Method of Functional Analysis and Topology, Vol. 26, no. 3, pp. 227-240.
- [11] D. L. Hua, Y. D. Huang,  *$K$ - $g$ -frames and stability of  $K$ - $g$ -frames in Hilbert spaces*, J. Korean Math. Soc. 53(6), 1331-1345.
- [12] Pawan K. Jain, Om P. Ahuja, *Functional Analysis*, New Age International Publisher, 1995.
- [13] V. Sadri, Gh. Rahimlou, R. Ahmadi and R. Zarghami Farfar, *Generalized Fusion Frames in Hilbert Spaces*, Submitted, arXiv: 1806.03598v1 [math.FA] 10 Jun 2018.
- [14] V. Sadri, Gh. Rahimlou, R. Ahmadi, and R. Zarghami Farfar, *Constructions of  $K$ - $g$  fusion frames and their duals in Hilbert spaces*, Bull. Transilvania Un. Brasov, 13(62), No. 1, (2020), 17-32.
- [15] W. Sun,  *$G$ -frames and  $G$ -Riesz bases*, Journal of Mathematical Analysis and Applications 322 (1) (2006), 437-452.