

Additive Average Schwarz Method for Elliptic Mortar Finite Element Problems with Highly Heterogeneous Coefficients

Ali Khademi · Leszek Marcinkowski ·
Sanjib Kumar Acharya · Talal Rahman

Received: date / Accepted: date

Abstract In this paper, we extend the additive average Schwarz method to solve second order elliptic boundary value problems with heterogeneous coefficients inside the subdomains and across their interfaces by the mortar technique, where the mortar finite element discretization is on nonmatching meshes. In this two-level method, we enrich the coarse space in two different ways, i.e., by adding eigenfunctions of two variants of the generalized eigenvalue problems. We prove that the condition numbers of the systems of algebraic equations resulting from the extended additive average Schwarz method, corresponding to both coarse spaces, are of the order $O(H/h)$ and independent of jumps in the coefficients, where H and h are the mesh parameters.

Keywords Domain decomposition method · Additive Schwarz method · Generalized eigenvalue problem · Mortar finite elements

Ali Khademi
Department of Computer science, Electrical engineering and Mathematical sciences,
Western Norway University of Applied Sciences, P.O. Box 7030, Bergen, Norway
E-mail: ali.khademi@hvl.no, akhademi.math@gmail.com

Leszek Marcinkowski
Department of Mathematics, Warsaw University, Poland
E-mail: lmarcin@mimuw.edu.pl

Sanjib Kumar Acharya
Institute of Chemical Technology Mumbai, Indian Oil Campus, Odisha, Bhubaneswar, India
E-mail: acharya.k.sanjib@gmail.com

Talal Rahman
Department of Computer science, Electrical engineering and Mathematical sciences,
Western Norway University of Applied Sciences, P.O. Box 7030, Bergen, Norway
E-mail: talal.rahman@hvl.no

1 Introduction

Domain decomposition methods are efficient and powerful iterative methods to solve large algebraic systems arising from a finite element discretization of elliptic boundary value problems [32,35,37]. They can also be regarded as a procedure of producing preconditioners for other iterative methods, such as the conjugate gradient method, for achieving fast convergence. In both approaches, to solve an original problem defined on a bounded Lipschitz domain $\bar{\Omega} = \cup_{i=1}^N \bar{\Omega}_i$, it is equivalent to solve many subproblems defined locally on the subdomains Ω_i in parallel. To obtain fast convergence, Dryja and Widlund [12], and Matsonki and Nepomnyaschikh [27] proposed to add one global problem and introduced the additive Schwarz methods to solve the global and local problems in parallel.

Corresponding to the global and local problems, different coarse and fine spaces can be constructed, for instance, see [5,7,11,29]. It is worth mentioning that coarse spaces are more important than fine spaces since they have a key role in the central of scalability for domain decomposition methods. Therefore, in [7] one of the simplest and efficient ways to construct coarse space, called the average coarse space, was proposed. Consequently, the two-level additive average Schwarz method was introduced and developed to solve many kinds of elliptic problems with the continuous and discontinuous coefficients, see [8,14,17,24]. Hence, an extension of the additive average Schwarz method to solve elliptic model problems arising from many applications such as composite materials with highly heterogeneous coefficients is of particular interest in this paper because it has a high-level performance.

In general, in terms of distributions of the coefficients, we can classify elliptic problems with the heterogeneous coefficients into two classes, i.e., elliptic problems with jumps in the coefficients only inside the subdomains and ones with jumps in the coefficients both inside the subdomains and on the subdomain interfaces. For the first class, where the heterogeneous coefficients are piecewise constants for subdomains $\Omega_i, i = 1, \dots, N$, the additive Schwarz method was developed and analyzed in [7,9,30,36] and references therein. For the second class with the large variation in the heterogeneous coefficients, the classical coarse spaces lead the condition numbers of the preconditioned systems to blow up. Consequently, the convergence rates of iterative methods will deteriorate. To alleviate this difficulty, the coarse spaces can be enriched by combining their structure with spectral spaces.

The idea of the coarse spectral space was introduced in [6] and extended as the spectral algebraic multigrid method in [15]. The spectral construction of this new space is achieved by solving the generalized eigenvalue problems locally. Due to its crucial role in preventing the impact of large jumps in the coefficients on bounds of the condition numbers of preconditioners, the new coarse space has received considerable attention. As a result, it is extensively developed for overlapping Schwarz methods [16,18,19,34], nonoverlapping additive Schwarz method [26], balancing domain decomposition methods [20,

21,25,33], and nonlinear domain decomposition [22], where all nodes on the subdomains interfaces for all references are matching grids. Hence, in this paper, we focus on the nonmatching grids, which are common due to heterogeneous materials in real-life problems. In practice, one may use different triangulations for polygonal subdomains independent of other triangulated subdomains. More precisely, there exist situations, where two subdomains with a common interface have fine and coarse (or fine but with different mesh sizes compared to other) triangulations. Therefore, the nonmatching grids on the subdomains interfaces are unavoidable and cause consistency errors in numerical methods, which can be handled by using the mortar techniques [1,2,3,4,31]. Hence, the additive Schwarz method for the mortar finite element method was introduced [8], and modified [28] with Crouzeix-Raviart mortar finite elements.

The main aim of this paper is to enrich the coarse space used in the classical nonoverlapping additive average Schwarz method by using the idea of coarse spectral space for elliptic problems with highly heterogeneous coefficients inside the subdomains and across their interfaces where the mortar finite element discretization is on the nonmatching meshes. To achieve this, our new coarse space consists of two subspaces. The first subspace is the common coarse space in the classical additive average Schwarz method used in [8], i.e., for the fixed i , is the range of a linear operator defined on Ω_i such that it is either the nodal values of a function $u \in V_h$ inside Ω_i or the average of nodal values of the function u on the mortar and nonmortar sides of Ω_i , where V_h is a finite space of P1 conforming elements defined on a fine triangulation of Ω and vanishing on $\partial\Omega$. The second subspace has a particular spectral structure. To obtain the basis functions for this subspace, we solve the generalized eigenvalue problems restricted to each subdomain as in [26].

To define the proper generalized eigenvalues problems, we require determining minimum values of the coefficients over each subdomain's triangulation to estimate the condition number of additive average Schwarz preconditioners independent of the large eigenvalues caused by the large jumps in the coefficients. Motivated by the ideas from [26], we consider two different types of the generalized eigenvalue problems based on either minimum of the coefficients over the whole subdomain or just minimum of ones over the layer connected to the boundary of the subdomain with one vertex or with one edge of the triangles inside of the subdomain. Solving these generalized eigenvalue problems lead to finding orthogonal basis functions enriching the coarse spaces. With these new coarse spaces, we prove that the condition numbers of the produced preconditioners are of the order $O(H/h)$ and independent of the number of subdomains. Due to the definition of the second type layer, it has faster performance than the first one through the implementations with numerical-software packages. See also Section 6 for numerical results.

The outline of this paper is as follows. In Section 2, after introducing a discrete problem, we define the mortar condition and the space of basis

functions satisfying that condition. Furthermore, several figures related to those functions are also given there. Section 3 is devoted to introducing the additive average Schwarz method, where the average interpolation operator has two different types. This operator consists of the natural extension of the standard average interpolation operator for mortar case and orthogonal operators defined in the next section. In Section 4, the generalized eigenvalue problems in terms of how the minimum values of jumps in the coefficients over the subdomains' triangulations can be defined are introduced. In Section 5, following the standard additive Schwarz framework [32], we drive an optimal estimate of the condition number of the produced preconditioners with the aid of removing bad eigenvalues, which are influenced by the large jumps in the coefficients. Finally, to verify our theoretical results' validity, in Section 6 some numerical experiments are reported.

2 Discrete problem

Let $\Omega \subset \mathbb{R}^2$ be a bounded Lipschitz domain with a nonoverlapping partition $\{\Omega_i\}_{i=1}^N$ of polygonal subdomains such that $\overline{\Omega} = \cup_{i=1}^N \overline{\Omega}_i$. We consider an elliptic model problem defined on Ω : Find $u_* \in H_0^1(\Omega)$ such that

$$a(u_*, v) = f(v), \quad v \in H_0^1(\Omega), \quad (1)$$

where

$$a(u, v) = \sum_{i=1}^N a_i(u, v) = \sum_{i=1}^N (\alpha_i \nabla u, \nabla v)_{L^2(\Omega_i)}$$

and

$$f(v) = \int_{\Omega} f v \, dx = \sum_{i=1}^N \int_{\Omega_i} f v \, dx.$$

Here, $f \in L^2(\Omega)$, $\alpha \in L^\infty(\Omega)$ and $\alpha_i(x)$ is the restriction of $\alpha(x)$ over Ω_i . Further, we assume that there exists a positive constant α_0 such that $\alpha(x) > \alpha_0$.

The partition $\{\Omega_i\}_{i=1}^N$ forms a coarse triangulation of Ω with the mesh parameter $H = \max\{H_i, i = 1, \dots, N\}$, where H_i are diameters of Ω_i . We assume this partition to be geometrically conforming, i.e., $\partial\Omega_i \cap \partial\Omega_j$ ($i \neq j$) is a vertex or a whole edge of both subdomains Ω_i and Ω_j or is empty. Further, we denote the set of all vertices of such coarse triangulation, except those belonging to $\partial\Omega$, by \mathcal{N}_H . We also denote the triangulation of the subdomain Ω_i by \mathcal{T}_i which consists of triangles satisfying the shape regular property [10] inside of Ω_i , and quasi-uniform triangles touching $\partial\Omega_i$ with the mesh size h_i . Further, we denote the set of all internal nodes of the fine triangulation \mathcal{T}_i by \mathcal{I}_i .

We define the product space X_h on the computational domain Ω by

$$X_h(\Omega) = X_1(\Omega_1) \times \dots \times X_N(\Omega_N),$$




Fig. 1 The mortar and nonmortar sides of subdomains Ω_i and Ω_j which are denoted by $\gamma_{m(i)}$ and $\delta_{m(j)}$, respectively.

where $X_i(\Omega_i), i = 1, \dots, N$ are the finite element spaces of the continuous piecewise linear functions defined on \mathcal{T}_i and vanishing on $\partial\Omega \cap \partial\Omega_i$. We denote all nodal points on $\bigcup_{i=1}^N \mathcal{T}_i$ except those on $\partial\Omega$ by \mathcal{N}_Ω . Further, we denote the set of basis functions associated with the set of nodal points \mathcal{N}_Ω by $\{\phi_l\}_{l \in \mathcal{N}_\Omega}$.

Due to independent triangulations inside of each subdomain, on each side of the interface $\bar{\Gamma}_{ij} = \bar{\Omega}_i \cap \bar{\Omega}_j$ we may have different discretization (cf. Figure 1). We select one side of Γ_{ij} as the mortar side, and the other side as the nonmortar side denoted by $\gamma_{m(i)}$ and $\delta_{m(j)}$, respectively. It is obvious that $\Gamma_{ij} = \gamma_{m(i)} = \delta_{m(j)}$. Further, we denote the nodes on the mortar and nonmortar sides by $m_0, m_1, \dots, m_{n_m+1}$ and $s_0, s_1, \dots, s_{n_s+1}$, respectively.

Let $W^{h_i}(\Gamma_{ij})$ and $W^{h_j}(\Gamma_{ij})$ denote the restrictions of $X_i(\Omega_i)$ and $X_j(\Omega_j)$ onto Γ_{ij} , respectively. Now, the nonmatching grids on the subdomain interfaces impose discontinuities for the functions belong to X_h . Therefore, we need to define a weak continuity condition. To this end, we first define the projection $\Pi_m(u_i, \text{Tr } v_j) : L^2(\delta_{m(j)}) \rightarrow W^{h_j}(\delta_{m(j)})$ by

$$\int_{\delta_{m(j)}} \left(u_i|_{\gamma_{m(i)}} - \Pi_m(u_i, \text{Tr } v_j) \right) \psi \, dx = 0, \quad \gamma_{m(i)} = \Gamma_{ij} = \delta_{m(j)}, \quad (2)$$

for all functions $\psi \in M^{h_j}(\delta_{m(j)}) = \text{span}\{\varphi_l\}_{l=0}^{n_s+1}$ and

$$\Pi_m(u_i, \text{Tr } v_j)|_{\partial\delta_{m(j)}} = v_j|_{\partial\delta_{m(j)}},$$

where u_i , $\text{Tr } v_j$, and $M^{h_j}(\delta_{m(j)})$ are the restriction of u into each subdomains $\Omega_i, i = 1, \dots, N$, the trace of v_j , and a subspace of $W^{h_j}(\delta_{m(j)})$ with constant values on the elements touching $\partial\delta_{m(j)}$, more precisely, at two end points of $\delta_{m(j)}$, respectively. Now, we say a function $u_h = \{u_i\}_{i=1}^N \in X_h$ satisfies the mortar condition on $\delta_{m(j)}$, if (2) holds. We denote the mortar space by V_h in terms of the mortar condition, i.e.,

$$V_h = \{u_h, v_h \in X_h \mid \Pi_m(u_h, \text{Tr } v_h) = 0, \forall \Gamma_{ij}, i, j = 1, \dots, N\}.$$




Fig. 2 The node x_k is represented by the black and thick dot, where the value of the basis function $\phi_k(x)$ associated with x_k is 1. Figures (I)-(IV) illustrate the different positions of x_k .

To see the structure of the basis functions spanning the mortar space V_h , let $\phi_k^{(i)}(x)$ be a basis function defined on Ω_i and define $V_h = \text{span}\{\phi_k\}$, where $\phi_k(x)$ is a basis function associated with a node x_k . Due to different positions of x_k , $\phi_k(x)$ takes different forms as follows (cf. Figure 2).

1. $x_k \in \mathcal{V}_i$:

$$\phi_k(x) = \phi_k^{(i)}(x).$$

2. $x_k \in \{m_1, \dots, m_{n_m}\}$:

$$\phi_k(x) = \begin{cases} \phi_k^{(i)}(x), & \text{on } \overline{\Omega}_i, \\ \Pi_m(\phi_k^{(i)}(x), 0)(x), & \text{on } \overline{\delta}_{m(j)}, \text{ where } \gamma_{m(i)} = \delta_{m(j)}, \\ 0, & \text{otherwise.} \end{cases}$$

3. $x_k \in \mathcal{N}_H$:

(a) x_k is a point of intersection between two mortar sides $\gamma_{m(i)}$ and $\gamma_{n(i)}$:

$$\phi_k(x) = \begin{cases} \phi_k^{(i)}(x), & \text{on } \bar{\gamma}_{m(i)} \text{ and } \bar{\gamma}_{n(i)}, \\ \Pi_m(\phi_k^{(i)}, 0)(x), & \text{on } \bar{\delta}_{m(j)}, \\ \Pi_n(\phi_k^{(i)}, 0)(x), & \text{on } \bar{\delta}_{n(j)}, \\ 0, & \text{otherwise.} \end{cases}$$

(b) x_k is a point of intersection between mortar side $\gamma_{m(i)}$ and nonmortar side $\delta_{n(i)}$:

$$\phi_k(x) = \begin{cases} \phi_k^{(i)}(x), & \text{on } \bar{\gamma}_{m(i)}, \\ \Pi_m(\phi_k^{(i)}, 0)(x), & \text{on } \bar{\delta}_{m(j)}, \\ \Pi_n(0, \text{Tr } \phi_k^{(i)})(x), & \text{on } \bar{\delta}_{n(i)}, \\ 0, & \text{otherwise.} \end{cases}$$

(c) x_k is a point of intersection between two nonmortar sides $\delta_{m(i)}$ and $\delta_{n(i)}$:

$$\phi_k(x) = \begin{cases} \phi_k^{(i)}(x), & \text{on } \bar{\gamma}_{m(j)} \text{ and } \bar{\gamma}_{n(j)}, \\ \Pi_m(0, \text{Tr } \phi_k^{(i)})(x), & \text{on } \bar{\delta}_{m(i)}, \\ \Pi_n(0, \text{Tr } \phi_k^{(i)})(x), & \text{on } \bar{\delta}_{n(i)}, \\ 0, & \text{otherwise.} \end{cases}$$

We now express the main problem as in the following form: Find $u_h^* = \{u_i^h\}_{i=1}^N \in V_h$ such that

$$a_h(u_h^*, v_h) = f(v_h) \quad \forall v_h \in V_h. \quad (3)$$

In what follows, we consider the following matrix representation of the linear systems arising from the discretization of (3).

$$\mathbf{A}\mathbf{v} = \mathbf{f},$$

where \mathbf{v} is the vector of all unknown coefficients defined on $\bar{\Omega}$. Further, we consider the submatrices $\mathbf{A}_{\Omega_i} = \mathbf{R}_i \mathbf{A} \mathbf{R}_i^T$, where $\mathbf{R}_i, i = 1, \dots, N$ are the restriction matrices such that $\mathbf{v}_i = \mathbf{R}_i \mathbf{v}$ are the vectors of coefficients defined on $\Omega_i \setminus \partial\Omega_i$.

Employing the mortar condition, we can compute some coefficients of \mathbf{v} in terms of other ones. More precisely, where $\Gamma_{ij} = \gamma_{m(i)} = \delta_{m(j)}$ assume $\nu_s = (v_h(s_i))_{i=1}^{n_s}, \nu_c = (v_h(s_0), v_h(s_{n_s+1}))^T, \nu_m = (v_h(m_i))_{i=0}^{n_m+1}$ and consider the following matrix representations as in [13].

$$\begin{aligned} \mathbf{M}_\gamma &:= ((\varphi_{s_i}, \phi_{m_j})_{L^2(\gamma)})_{i=1; j=0}^{n_s \quad n_m+1} \in \mathbb{R}^{n_s \times (n_m+2)}, \\ \mathbf{S}_\gamma &:= ((\varphi_{s_i}, \phi_{s_j})_{L^2(\gamma)})_{i=1; j=1}^{n_s \quad n_s} \in \mathbb{R}^{n_s \times n_s}, \\ \mathbf{C}_\gamma &:= ((\varphi_{s_i}, \phi_{m_j})_{L^2(\gamma)})_{i=1; j=0, n_s+1}^{n_s} \in \mathbb{R}^{n_s \times 2}. \end{aligned}$$

Hence

$$\nu_s = \mathbf{S}_\gamma^{-1}(\mathbf{M}_\gamma \nu_m - \mathbf{C}_\gamma \nu_c).$$

Consequently, in the next sections, we will focus only on all mortar, corner, and interior nodes.

3 The additive average Schwarz method

Let $V^{(i)}$, $i = 1, \dots, N$ be decompositions of V_h that are restrictions of V_h to Ω_i with zero on $\partial\Omega_i$ and on the subdomains Ω_j , $j \neq i$. Further, let $V_0^{(type)}$ stands for two different types of the coarse spaces, distinguished by two notations I and II, such that

$$V_h = V_0^{(type)} + \sum_{i=1}^N V^{(i)}, \quad type \in \{I, II\},$$

where

$$V_0^{(type)} = I_0 V_h + \sum_{i=1}^N W_i^{type}, \quad type \in \{I, II\}.$$

We define the operator I_0 and the spaces W_i^{type} , $type \in \{I, II\}$ in the next section. Here, we use the exact bilinear form for all our subproblems. Thus, we have

$$a(u^h, v^h) = a_i(u_i^h, v_i^h), \quad i = 1, \dots, N,$$

where $u^h = \{u_i^h\}_{i=1}^N \in V^{(i)}$ and $v^h = \{v_i^h\}_{i=1}^N \in V^{(i)}$. We define the projection-like operators $T^{(i)} : V_h \rightarrow V^{(i)}$ $i = 1, \dots, N$ and $T_0^{(type)} : V_h \rightarrow V_0^{(type)}$ such that for $u^h \in V_h$, $T^{(i)}u^h$ and $T_0^{(type)}u^h$ are the solutions of

$$a(T^{(i)}u^h, v^h) = a(u^h, v^h), \quad v^h \in V^{(i)}, \quad i = 1, \dots, N$$

and

$$a(T_0^{(type)}u^h, v^h) = a(u^h, v^h), \quad v^h \in V_0^{(type)}, \quad type \in \{I, II\}.$$

Now, the additive Schwarz operator $T^{type} : V_h \rightarrow V_h$ is

$$T^{type} = T_0^{(type)} + \sum_{i=1}^N T^{(i)}, \quad type \in \{I, II\}$$

and the problem (3) can be written as follow.

$$T^{type}u_h^* = g^{type}, \quad type \in \{I, II\},$$

where $g^{type} = g_0^{(type)} + \sum_{i=1}^N g^{(i)}$ with $g_0^{(type)} = T_0^{(type)}u_h^*$ and $g^{(i)} = T^{(i)}u_h^*$.

4 Enrichment of the coarse space for the mortar discretization

In this section, we design two different coarse spaces for the additive average Schwarz method. To this end, we first denote the sets of nodal nodes of all mortar and nonmortar sides, and also all interior nodes of all subdomains Ω_i , $i = 1, \dots, N$ by \mathcal{N}_m , \mathcal{N}_s , and \mathcal{N}_i , respectively. Now, the average interpolation operator $I_0 : V_h \rightarrow V_h$ for the mortar discretization as in [8] has the following structure.

$$I_0 u^h(x) = \begin{cases} u_i^h(x) & x \in \mathcal{N}_m \cup \mathcal{N}_s, \\ \bar{u}_i^h & x \in \mathcal{N}_i, \end{cases} \quad i = 1, \dots, N,$$

where \bar{u}_i^h is the average value of u_i^h over Ω_i , i.e.,

$$\bar{u}_i^h = \frac{1}{\mu_{(\delta, \gamma)}^i} \left(\sum_{\gamma_{m(i)} \subset \partial \Omega_i} \bar{u}_{\gamma_{m(i)}}^h + \sum_{\delta_{m(i)} \subset \partial \Omega_i, \gamma_{m(j)} = \delta_{m(i)}} \bar{u}_{\gamma_{m(j)}}^h \right), \quad (4)$$

where

$$\bar{u}_{\gamma_{m(i)}}^h = \frac{1}{|\gamma_{m(i)}|} \int_{\gamma_{m(i)}} u_i^h ds$$

and $\mu_{(\delta, \gamma)}^i$ is the number of all mortar and nonmortar sides of Ω_i , and $|\gamma_{m(i)}|$ is the length of $\gamma_{m(i)}$. To express the interpolation operator I_0 in terms of the matrix form denoted by \mathbf{R}_0 , let \mathcal{N}_c be a set of all nodal nodes at the end of all nonmortar sides of Ω and let also $\mathbf{I}_{(c)} \in \mathbb{R}^{N_c \times N_c}$ and $\mathbf{I}_{(m)} \in \mathbb{R}^{N_m \times N_m}$ be identity matrices, where $N_c = \dim(\mathcal{N}_c)$ and $N_m = \dim(\mathcal{N}_m)$. Further, consider $\mathbf{H} = \text{diag}(\mathbf{H}_1, \dots, \mathbf{H}_N) \in \mathbb{R}^{(N_c + N_m) \times N_i}$, where $\mathbf{H}_i = (\bar{u}_i^h)_{l=1}^{4+n_m} n_i$, $N_i = \dim(\mathcal{N}_i)$, and $n_i = \dim(\mathcal{N}_i)$. Hence

$$\mathbf{R}_0 = \left[\text{diag}(\mathbf{I}_{(c)}, \mathbf{I}_{(m)}) \middle| \mathbf{H} \right] \in \mathbb{R}^{(N_c + N_m) \times N_\Omega}, \quad N_\Omega = N_c + N_m + N_i.$$




Fig. 3 The boundary layer $\Omega_i^B \subset \Omega_i$ is highlighted by the colorful triangles.

To define two different coarse spaces, we first introduce the boundary layers Ω_i^B , $i = 1, \dots, N$, where $\Omega_i^B \subset \Omega_i$ is the sum of all triangles such as $\tau \in \mathcal{T}_i$ such that $\partial \tau \cap \partial \Omega_i \neq \emptyset$ (cf. Figure 3). We then define the following local

minimums of the coefficients over the subdomains Ω_i :

$$\underline{\alpha}_i := \min_{x \in \Omega_i} \alpha(x), \quad \underline{\alpha}_{i,B} := \min_{x \in \Omega_i^B} \alpha(x), \quad i = 1, \dots, N.$$

We now proceed to construct the second part of $V_0^{(type)}$, i.e., $W_i^{type}, type \in \{I, II\}, i = 1, \dots, N$ which are the spaces of adaptively chosen eigenfunctions of specially constructed generalized eigenvalue problems defined locally in each subdomain and extended by zero to the rest of the domain. Hence, the generalized eigenvalue problems is to find all eigen pairs $(\lambda_k^{i,type}, \psi_k^{i,type}) \in (\mathbb{R}_+, V^{(i)})$ such that

$$\mathbf{A}_{\Omega_i} \mathbf{x} = \lambda^{type} (\mathbf{R}_i \mathbf{B}^{type} \mathbf{R}_i^T) \mathbf{x}, \quad i = 1, \dots, N, \quad (5)$$

where $\mathbf{B}^{type}(\cdot, \cdot), type \in \{I, II\}$ are the matrix representations of the following bilinear forms.

$$b^I(u, v) := \sum_{i=1}^N \int_{\Omega_i} \underline{\alpha}_i \nabla u \cdot \nabla v \, dx, \quad u, v \in V_h,$$

$$b^{II}(u, v) := \sum_{i=1}^N \left(\int_{\Omega_i^B} \underline{\alpha}_{i,B} \nabla u \cdot \nabla v \, dx + \int_{\Omega_i \setminus \Omega_i^B} \alpha \nabla u \cdot \nabla v \, dx \right), \quad u, v \in V_h.$$

We also denote all eigenvalues of (5) by

$$\mathbf{D}^{type} = \text{diag}(\mathbf{D}_1^{type}, \mathbf{D}_2^{type}, \dots, \mathbf{D}_N^{type}) \in \mathbb{R}^{N_i \times N_i}, \quad type \in \{I, II\},$$

where

$$\mathbf{D}_i^{type} = \text{diag}(\lambda_1^{i,type}, \lambda_2^{i,type}, \dots, \lambda_{n_i}^{i,type}) \in \mathbb{R}^{n_i \times n_i}, \quad i = 1, \dots, N,$$

such that

$$\lambda_1^{i,type} \geq \lambda_2^{i,type} \geq \dots \geq \lambda_{n_i}^{i,type} > 0.$$

We define

$$W_i^{type} := \text{span}\{\psi_k^{i,type}\}_{k=1}^{n_i}, \quad type \in \{I, II\}, i = 1, \dots, N$$

as the space of the eigenfunctions associated with the eigenvalues $\lambda_j^{i,type}$. We also correspond to these spectral spaces the matrix forms denoted by $\mathbf{W}_i^{type}, i = 1, \dots, N$ such that

$$\mathbf{W}^{type} = \text{diag}(\mathbf{W}_1^{type}, \mathbf{W}_2^{type}, \dots, \mathbf{W}_N^{type}), \quad type \in \{I, II\}.$$

Further, we use the notation \mathbf{R}_0^{type} as the matrix representation of the operator $T_0^{(type)}$ which has the following structure.

$$\mathbf{R}_0^{type} = \left[\begin{array}{c|c} \mathbf{R}_0 & O \\ \hline O & \mathbf{W}^{type} \end{array} \right], \quad \mathbf{R}_0^{type} \in \mathbb{R}^{N_\Omega \times (N_\Omega + N_i)}, \quad type \in \{I, II\}.$$

4.1 The additive average Schwarz preconditioner

The main aim of this section is to introduce new enrichment preconditioners denoted by \mathbf{B}_E^{type} , $type \in \{I, II\}$. The idea is based on expressing these preconditioners in terms of a combination of \mathbf{B}_C^{type} , $type \in \{I, II\}$ and $\sum_{i=1}^N \mathbf{R}_i^T \mathbf{A}_{\Omega_i}^{-1} \mathbf{R}_i$, where $\mathbf{B}_C^{type} = (\mathbf{R}_0^{type})^T (\mathbf{R}_0^{type} \mathbf{A}_N (\mathbf{R}_0^{type})^T)^{-1} \mathbf{R}_0^{type}$ and

$$\mathbf{A}_N = \left[\begin{array}{c|c} \mathbf{A}_N^{(11)} & \mathbf{A}_N^{(12)} \\ \hline \mathbf{A}_N^{(21)} & \mathbf{A}_N^{(22)} \end{array} \right], \quad \mathbf{A}_N \in \mathbb{R}^{(N_\Omega + N_i) \times (N_\Omega + N_i)}$$

such that

$\mathbf{A}_N^{(11)}$:= square submatrix of \mathbf{A} whose rows and columns are corresponding to the nodes belong to $\mathcal{N}_c \cup \mathcal{N}_m \cup \mathcal{N}_i$,

$\mathbf{A}_N^{(12)}$:= rectangular submatrix of \mathbf{A} whose rows are corresponding to all nodes belong to $\mathcal{N}_c \cup \mathcal{N}_m \cup \mathcal{N}_i$, meanwhile whose columns are corresponding to the nodes belong to \mathcal{N}_i ,

$$\mathbf{A}_N^{(21)} = (\mathbf{A}_N^{(12)})^T,$$

$\mathbf{A}_N^{(22)}$:= square submatrix of \mathbf{A} whose rows and columns are corresponding to the nodes belong to \mathcal{N}_i . Indeed, $\mathbf{A}_N^{(22)} = \text{diag}(\mathbf{A}_{\Omega_1}, \mathbf{A}_{\Omega_2}, \dots, \mathbf{A}_{\Omega_N})$.

Hence

$$\mathbf{R}_0^{type} \mathbf{A}_N (\mathbf{R}_0^{type})^T = \left[\begin{array}{c|c} \mathbf{R}_0 \mathbf{A}_N^{(11)} \mathbf{R}_0^T & \mathbf{R}_0 \mathbf{A}_N^{(12)} (\mathbf{W}^{type})^T \\ \hline (\mathbf{R}_0 \mathbf{A}_N^{(12)} (\mathbf{W}^{type})^T)^T & \mathbf{W}^{type} \mathbf{A}_N^{(22)} (\mathbf{W}^{type})^T \end{array} \right].$$

Since $\mathbf{W}^{type} \mathbf{A}_N^{(22)} (\mathbf{W}^{type})^T = \mathbf{D}^{type}$, $type \in \{I, II\}$ and it is the nonsingular matrix, we can easily compute \mathbf{B}_C^{type} . To this end and for the sake of simplicity, we first assume

$$\begin{aligned} \mathbf{G}^{type} &= \mathbf{R}_0 \mathbf{A}_N^{(12)} (\mathbf{W}^{type})^T, \\ \mathbf{S}^{type} &= \mathbf{R}_0 \mathbf{A}_N^{(11)} \mathbf{R}_0^T - \mathbf{G}^{type} (\mathbf{D}^{type})^{-1} (\mathbf{G}^{type})^T. \end{aligned}$$

Hence, the inverse of 2×2 block matrix $\mathbf{R}_0^{type} \mathbf{A}_N (\mathbf{R}_0^{type})^T$ implies

$$\begin{aligned} \mathbf{B}_{C(11)}^{type} &= \mathbf{R}_0^T (\mathbf{S}^{type})^{-1} \mathbf{R}_0, \\ \mathbf{B}_{C(12)}^{type} &= -\mathbf{R}_0^T (\mathbf{S}^{type})^{-1} \mathbf{G}^{type} (\mathbf{D}^{type})^{-1} \mathbf{W}^{type}, \\ \mathbf{B}_{C(21)}^{type} &= -(\mathbf{B}_{C(12)}^{type})^T, \\ \mathbf{B}_{C(22)}^{type} &= (\mathbf{W}^{type})^T \left((\mathbf{D}^{type})^{-1} + (\mathbf{G}^{type} (\mathbf{D}^{type})^{-1})^T (\mathbf{S}^{type})^{-1} \mathbf{G}^{type} (\mathbf{D}^{type})^{-1} \right) \mathbf{W}^{type}. \end{aligned}$$

Due to different sizes of the submatrices of \mathbf{B}_C^{type} , we can use the restriction matrices $\mathbf{R}_i, i = 1, \dots, N$ to obtain new matrices not only with identical sizes, but also with their previous performances. Let $\mathbf{R}_r = \text{diag}(\mathbf{R}_1, \mathbf{R}_2, \dots, \mathbf{R}_N)$, $\mathbf{B}_0^{type} = \mathbf{B}_{C(11)}^{type} + \mathbf{R}_r^T \mathbf{B}_{C(21)}^{type}$ and $\mathbf{B}_{00}^{type} = \mathbf{B}_{C(12)}^{type} \mathbf{R}_r + \mathbf{R}_r^T \mathbf{B}_{C(22)}^{type} \mathbf{R}_r$. Hence, $\mathbf{B}_E^{type}, type \in \{I, II\}$ as the enrichment preconditioners can be written explicitly in the following matrix form.

$$\begin{aligned} \mathbf{B}_E^{type} &= \mathbf{B}_0^{type} + \mathbf{B}_{00}^{type} + \sum_{i=1}^N \mathbf{R}_i^T \mathbf{A}_{\Omega_i}^{-1} \mathbf{R}_i \\ &= \mathbf{B}_0^{type} + \left(\mathbf{R}_r^T - \mathbf{B}_0^{type} \mathbf{A}_N^{(12)} \right) (\mathbf{W}^{type})^T (\mathbf{D}^{type})^{-1} \mathbf{W}^{type} \mathbf{R}_r + \sum_{i=1}^N \mathbf{R}_i^T \mathbf{A}_{\Omega_i}^{-1} \mathbf{R}_i. \end{aligned}$$

To estimate an upper bound for the condition number of \mathbf{B}_E^{type} in the next section, we define the $b_i^{type}(\cdot, \cdot)$ -orthogonal projection operator $\pi_i^{type} : V^{(i)} \rightarrow V^{(i)}$ as

$$\pi_i^{type} v^h = \sum_{k=1}^{N_i} b_i^{type}(v^h, \psi_k^{i,type}) \psi_k^{i,type}, \quad v^h \in V_h, \quad type \in \{I, II\}.$$

For any $u^h \in V_h$, we consider the function $w^h = u^h - I_0 u^h \in V_h$ with zero value both on $\partial\Omega_i$ and on the rest of the domain Ω . In addition, we define $I_0^{type} : V_h \rightarrow V_0^{(type)}$ as

$$I_0^{type} u^h = I_0 u^h + \sum_{i=1}^N \pi_i^{type} w^h, \quad u^h \in V_h, \quad type \in \{I, II\}.$$

5 On the estimation of the condition number bound

This section is devoted to obtaining condition number estimates of the additive average Schwarz preconditioners, which are defined in the previous section for $type \in \{I, II\}$. The proof is based on the standard additive Schwarz framework, where three assumptions to be held, see [32, p. 155]. Here, we need only to show that there exists the stable splitting for all $u \in V_h$ as in Theorem 1. Note that the other assumptions hold since there is no overlapping, and we also use the exact bilinear forms. For simplicity, throughout this section, we use C as a positive constant, which is independent of the mesh sizes. We add a notation or an index to C if we want to emphasize a special constant. Furthermore, we use the notation \lesssim to remove any constants except the mesh sizes in the inequalities.

Let M_i^{type} be a given number such that $0 \leq M_i^{type} < n_i$ and $\lambda_{M_i^{type}+1}^{i,type} < \lambda_{M_i^{type}}^{i,type}$. We define

$$\widetilde{W}_i^{type} := \text{span}\{\psi_k^{i,type}\}_{k=1}^{M_i^{type}}, \quad type \in \{I, II\}$$

and

$$\tilde{V}_0^{(type)} = I_0 V_h + \sum_{i=1}^N \tilde{W}_i^{type}, \quad type \in \{I, II\}.$$

For the analysis of the additive average Schwarz method, we define $\tilde{I}_0^{type} : V_h \rightarrow \tilde{V}_0^{(type)}$ as

$$\tilde{I}_0^{type} u^h = I_0 u^h + \sum_{i=1}^N \tilde{\pi}_i^{type} w^h, \quad u^h \in V_h, \quad type \in \{I, II\},$$

where

$$\tilde{\pi}_i^{type} v^h = \sum_{k=1}^{M_i^{type}} b_i^{type}(v^h, \psi_k^{i,type}) \psi_k^{i,type}, \quad type \in \{I, II\}.$$

To prove our main result, we need the following lemma.

Lemma 1. For all $u \in V^{(i)}, i = 1, \dots, N$,

$$\begin{aligned} |u - \tilde{\pi}_i^I u|_{H^1(\Omega_i), \alpha}^2 &\leq C \lambda_{M_i^I+1}^{i,I} \|\underline{\alpha}_i^{1/2} \nabla u\|_{L^2(\Omega_i)}^2, \\ |u - \tilde{\pi}_i^{II} u|_{H^1(\Omega_i), \alpha}^2 &\leq C \lambda_{M_i^{II}+1}^{i,II} \left(\|\underline{\alpha}_{iB}^{1/2} \nabla u\|_{L^2(\Omega_i^B)}^2 + \|\alpha_i^{1/2} \nabla u\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 \right), \end{aligned}$$

where $|\cdot|_{H^1(\Omega_i), \alpha}^2 = a_i(\cdot, \cdot)$.

Proof. We first express any $u \in V^{(i)}$ uniquely in terms of the eigenfunctions, i.e., $u = \sum_{k=1}^{n_i} b_i^{type}(u, \psi_k^{i,type}) \psi_k^{i,type}$. Hence

$$u - \tilde{\pi}_i^{type} u = \sum_{k=M_i^{type}+1}^{n_i} b_i^{type}(u, \psi_k^{i,type}) \psi_k^{i,type}, \quad type \in \{I, II\}$$

and we have

$$|u - \tilde{\pi}_i^{type} u|_{H^1(\Omega_i), \alpha}^2 \leq \sum_{k=M_i^{type}+1}^{n_i} \left(b_i^{type}(u, \psi_k^{i,type}) \right)^2 |\psi_k^{i,type}|_{H^1(\Omega_i), \alpha}^2.$$

Furthermore, $(\mathbf{W}_i^{type})^T \mathbf{A}_{\Omega_i} \mathbf{W}_i^{type} = \mathbf{D}_i^{type}$ is equivalent to

$$|\nabla \psi_k^{i,type}|_{H^1(\Omega_i), \alpha}^2 = \lambda_k^{i,type}, \quad k = 1, \dots, n_i \quad i = 1, \dots, N, \quad type \in \{I, II\}.$$

Therefore

$$|u - \tilde{\pi}_i^{type} u|_{H^1(\Omega_i), \alpha}^2 \leq \sum_{k=M_i^{type}+1}^{n_i} \left(b_i^{type}(u, \psi_k^{i,type}) \right)^2 \lambda_k^{i,type}, \quad type \in \{I, II\}. \quad (6)$$

Using the Schwarz inequality for $type = I$, yields

$$|u - \tilde{\pi}_i^I u|_{H^1(\Omega_i), \alpha}^2 \leq \|\underline{\alpha}_i^{1/2} \nabla u\|_{L^2(\Omega_i)}^2 \sum_{k=M_i^I+1}^{n_i} \|\underline{\alpha}_i^{1/2} \nabla \psi_k^{i,I}\|_{L^2(\Omega_i)}^2 \lambda_k^{i,I}.$$

Since $(\mathbf{W}_i^{type})^T (\mathbf{R}_i \mathbf{B}^{type} \mathbf{R}_i^T) \mathbf{W}_i^{type} = \mathbf{I}$, $type \in \{I, II\}$ and $i = 1, \dots, N$ we have equivalently

$$\|\underline{\alpha}_i^{1/2} \nabla \psi_k^{i,I}\|_{L^2(\Omega_i)}^2 = 1, \quad \|\underline{\alpha}_{iB}^{1/2} \nabla \psi_k^{i,II}\|_{L^2(\Omega_i^B)}^2 + \|\alpha_i^{1/2} \nabla \psi_k^{i,II}\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 = 1. \quad (7)$$

Then

$$|u - \tilde{\pi}_i^I u|_{H^1(\Omega_i), \alpha}^2 \leq C \lambda_{M_i^I+1}^{i,I} \|\alpha_i^{1/2} \nabla u\|_{L^2(\Omega_i)}^2.$$

For $type = \text{II}$, from (6) we get

$$\begin{aligned} |u - \tilde{\pi}_i^{\text{II}} u|_{H^1(\Omega_i), \alpha}^2 &\leq C \sum_{k=M_i^{\text{II}}+1}^{n_i} \left(\|\alpha_i^{1/2} \nabla u\|_{L^2(\Omega_i^B)}^2 \|\alpha_i^{1/2} \nabla \psi_k^{i,\text{II}}\|_{L^2(\Omega_i^B)}^2 \right. \\ &\quad \left. + \|\alpha_i^{1/2} \nabla u\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 \|\alpha_i^{1/2} \nabla \psi_k^{i,\text{II}}\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 \right) \lambda_k^{i,\text{II}}. \end{aligned}$$

To complete the proof, it is sufficient to use (7). \square

Theorem 1. For all $u^h \in V_h$ the following results hold:

$$a(\tilde{I}_0^{type} u^h, \tilde{I}_0^{type} u^h) \lesssim \max_i \lambda_{M_i^{type}+1}^{i,type} \frac{H}{h} a(u^h, u^h), \quad type \in \{\text{I}, \text{II}\}, \quad i = 1, \dots, N.$$

Proof. We first consider the following splitting.

$$u^h = \tilde{I}_0^{type} u^h + \sum_{i=1}^N u^i,$$

where $u^i = (0, \dots, 0, w_i, 0, \dots, 0) \in V^{(i)}$ and $w_i = (u^h - \tilde{I}_0^{type} u^h)|_{\Omega_i}$. Consequently, we get

$$a(\tilde{I}_0^{type} u^h, \tilde{I}_0^{type} u^h) \preceq a(u^h, u^h) + a(u^h - \tilde{I}_0^{type} u^h, u^h - \tilde{I}_0^{type} u^h),$$

where

$$\begin{aligned} a(u^h - \tilde{I}_0^{type} u^h, u^h - \tilde{I}_0^{type} u^h) &= \sum_{i=1}^N a(u^i, u^i) = \sum_{i=1}^N a_i(u^h - \tilde{I}_0^{type} u^h, u^h - \tilde{I}_0^{type} u^h) \\ &= \sum_{i=1}^N a_i(w - \tilde{\pi}_i^{type} w, w - \tilde{\pi}_i^{type} w), \quad type \in \{\text{I}, \text{II}\}. \end{aligned}$$

We first use Lemma 1 for the $type = \text{I}$. Then, we have

$$\begin{aligned} a_i(w - \tilde{\pi}_i^I w, w - \tilde{\pi}_i^I w) &\leq C \lambda_{M_i^I+1}^{i,I} \alpha_i \|\nabla(u^h - I_0 u^h)\|_{L^2(\Omega_i)}^2 \\ &\leq C \lambda_{M_i^I+1}^{i,I} \alpha_i \left(\frac{H}{h} \|\nabla u^h\|_{L^2(\Omega_i)}^2 + \|\nabla I_0 u^h\|_{L^2(\Omega_i)}^2 \right). \end{aligned}$$

Further, from [8, pp. 8-9] we get similarly

$$\alpha_i \|\nabla I_0 u^h\|_{L^2(\Omega_i)}^2 \leq C \frac{H}{h} a_i(u^h, u^h). \quad (8)$$

For $type = \text{I}$, the proof is completed by summing (8) over $i = 1, \dots, N$. We now proceed to prove a similar result for $type = \text{II}$. Due to definition of $I_0 u_i^h$, we use this fact that $\nabla I_0 u_i^h$ is equal to zero on each triangle $\tau \notin \Omega_i^B$, $i = 1, \dots, N$. Hence, we use Lemma 1 to get the first inequality as follow.

$$\begin{aligned}
& a_i(w - \tilde{\pi}_i^\Pi w, w - \tilde{\pi}_i^\Pi w) \\
& \leq C\lambda_{M_i^\Pi+1}^{i,\Pi} \left(\|\alpha_i^{1/2} \nabla(u^h - I_0 u^h)\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 + \underline{\alpha}_{i,B} \|\nabla(u^h - I_0 u^h)\|_{L^2(\Omega_i^B)}^2 \right) \\
& \leq C\lambda_{M_i^\Pi+1}^{i,\Pi} \left(\|\alpha_i^{1/2} \nabla u^h\|_{L^2(\Omega_i \setminus \Omega_i^B)}^2 + \underline{\alpha}_{i,B} \|\nabla(u^h - I_0 u^h)\|_{L^2(\Omega_i^B)}^2 \right) \\
& \leq C\lambda_{M_i^\Pi+1}^{i,\Pi} \left(\|\alpha_i^{1/2} \nabla u^h\|_{L^2(\Omega_i)}^2 + \underline{\alpha}_{i,B} \left(\|\nabla u^h\|_{L^2(\Omega_i^B)}^2 + \|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 \right) \right) \\
& \leq C\lambda_{M_i^\Pi+1}^{i,\Pi} \left(a_i(u^h, u^h) + \|\alpha_i^{1/2} \nabla u^h\|_{L^2(\Omega_i)}^2 + \underline{\alpha}_{i,B} \|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 \right) \\
& \leq C\lambda_{M_i^\Pi+1}^{i,\Pi} \left(a_i(u^h, u^h) + \underline{\alpha}_{i,B} \|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 \right).
\end{aligned}$$

Using the inverse inequality and the definition of the operator I_0 , implies

$$\begin{aligned}
\|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 &= \sum_{\tau \in \overline{\Omega}_i^B} \|\nabla I_0 u^h\|_{L^2(\tau)}^2 \\
&= \sum_{\tau \in \overline{\Omega}_i^B} \|\nabla(I_0 u^h - \bar{u}^h)\|_{L^2(\tau)}^2 \\
&\leq C \sum_{\tau \in \overline{\Omega}_i^B} h_\tau^{-2} \|I_0 u^h - \bar{u}^h\|_{L^2(\tau)}^2 \\
&\leq C \sum_{x \in \partial\Omega_{ih}} (u_i^h(x) - \bar{u}_i^h)^2.
\end{aligned} \tag{9}$$

In what follow, we need to use the following Poincaré inequality [36].

$$\|g\|_{L^2(\Omega_i)}^2 \leq C_1 \|g\|_{H^1(\Omega_i)}^2 + C_2 \left(\int_{\Omega_i} g \, dx \right)^2, \quad g \in H^1(\Omega_i), \tag{10}$$

where C_1 and C_2 are two positive constants independent of the mesh size of Ω . Consider the function $g = u_i^h - c$, where $c = \frac{1}{\text{meas}(\Omega_i)} \int_{\Omega_i} u_i^h \, dx$. Now, (10) implies

$$\|u_i^h - c\|_{L^2(\Omega_i)}^2 \leq C_1 \|u_i^h\|_{H^1(\Omega_i)}^2. \tag{11}$$

Furthermore, we use (4) and this property that all boundary elements of subdomains $\Omega_i, i = 1, \dots, N$ are quasi-uniform which implies that the number of all nodes belonging to $\partial\Omega_i$, denoted by M_i , is of the order H_i/h_i , to get

$$\begin{aligned}
M_i \left(\overline{u_i^h - c} \right)^2 &\leq C M_i \frac{1}{\mu_{(\delta, \gamma)}^i} \sum_{\Gamma \subset \partial\Omega_i} \frac{1}{\Gamma} \|u_i^h - c\|_{L^2(\Gamma)}^2 \\
&\leq C h_i^{-1} \|u_i^h - c\|_{L^2(\partial\Omega_i)}^2,
\end{aligned} \tag{12}$$

where Γ is $\gamma_{m(i)}$ and $\delta_{m(i)}$.

Now, we use g and (12) to estimate an upper bound for (9) as follows.

$$\begin{aligned} \|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 &\leq C \sum_{x \in \partial\Omega_{ih}} (g(x) - \bar{g})^2 \\ &\leq Ch_i^{-1} \|g\|_{L^2(\partial\Omega_i)}^2 \\ &\leq Ch_i^{-1} H_i \|\hat{g}\|_{L^2(\partial\hat{\Omega})}^2, \end{aligned}$$

where $\hat{\Omega}$ is the reference element of unit diameter. From this inequality we deduce

$$\begin{aligned} \|\nabla I_0 u^h\|_{L^2(\Omega_i^B)}^2 &\leq Ch_i^{-1} H_i \left(\|\hat{g}\|_{H^1(\hat{\Omega})}^2 + \|\hat{g}\|_{L^2(\hat{\Omega})}^2 \right) \\ &\leq Ch_i^{-1} \left(H_i \|g\|_{H^1(\Omega_i)}^2 + H_i^{-1} \|g\|_{L^2(\Omega_i)}^2 \right) \\ &\leq C \frac{H_i}{h_i} a_i(u^h, u^h), \end{aligned}$$

by the trace inequality, Theorem 3.1.2 in [10] and (11). To complete the proof, it suffices to take a summation over $i = 1, \dots, N$. \square

To estimate upper bounds of the condition numbers of $\mathbf{B}_E^{type} \mathbf{A}$, $type \in \{I, II\}$ it suffices to use Lemma 3 in [32, pp. 156-158] and Theorem 1.

Theorem 2. *The condition numbers of the enriched additive average Schwarz preconditioners are bounded by*

$$\kappa(\mathbf{B}_E^{type} \mathbf{A}) \leq C \left(\frac{H}{h} \right) \max_i \lambda_{M_i^{type}+1}^{i,type}, \quad type \in \{I, II\}, \quad i = 1, \dots, N,$$

where C is a positive constant.

6 Numerical results

In this section, numerical results confirm the additive average Schwarz method's validity and efficiency with adaptive enrichment, where the jumps in the coefficient $\alpha(x)$ in (1) are very large and even change rapidly. Those jumps might be occurred inside of the subdomains, or even across the subdomain boundaries. To have a complicated distribution of jumps, we use the following pattern [26], depicted in Figure 4, which is a periodic pattern when the number of subdomains are increased (cf. Figure 5). We also consider background channels, crossing channels and corner channels denoted by α_b , α_i and α_c , respectively. For different values of α_b , α_c and α_i , our test problem has the right-hand side function $f(x, y) = 2\pi^2 \sin(\pi x) \sin(\pi y)$ defined in $\Omega = [0, 1] \times [0, 1]$.

All presented numerical results are based on the nonmatching triangulations across the subdomains interfaces, where $H \in \{1/6, 1/9\}$, $h \in \{1/36, 1/54\}$, and $h^* \in \{1/54, 1/81\}$ (cf. Figure 5). We use the similar




Fig. 4 The decomposition of $\Omega = [0, 1] \times [0, 1]$ into 3×3 subdomains (numbered from 1 to 9) and the locations of $\alpha(x)$ consisting of α_b , α_c and α_i in the white, red and green areas, respectively. Further, all mortar sides are denoted by the thick and black line segments.




Fig. 5 The distribution of all jumps in $\alpha(x)$ following the extended pattern in Figure 4, where $\Omega = [0, 1] \times [0, 1]$ is divided into 6×6 subdomains. The nonmatching discretization parameters are $H = 1/6$, $h = 1/36$ and $h^* = 1/54$. Further, all mortar sides are the coarse mortar.

locations for α_c and α_i as the periodic patterns and the similar mortar sides, as depicted in Figure 4, for $N \geq 9$. For instance, see Figure 5, where $N = 36$. Further, in what follows, we use two terms, i.e., coarse mortar and fine mortar to distinguish between the coarse and fine triangulations connected

to the mortar sides. To withdraw numerical results, we use the additive average Schwarz method to produce the enrichment preconditioners. We estimate the condition number of such preconditioners for $type \in \{I, II\}$. Moreover, the iteration numbers in all tables come from the preconditioned conjugate gradient method, with the tolerance $5e - 6$, based on the produced preconditioners.

ADD		Different Values for α_b , α_c and α_i			
		1, 1e3, 1e4		1, 1e4, 1e6	
		Coarse Mortar	Fine Mortar	Coarse Mortar	Fine Mortar
Number of Subdomains	6×6	6.07e1 (52)	4.25e1 (45)	5.99e1 (56)	4.24e1 (49)
	9×9	7.99e1 (55)	5.56e1 (50)	7.69e1 (58)	5.51e1 (52)

Table 1 The condition number of $\mathbf{B}_E^{type} \mathbf{A}$ and the number of iterations of the preconditioned conjugate gradient method (in parentheses) for $type = II$ with different values for α_b , α_c , and α_i . For 6×6 and 9×9 subdomains, h and h^* belong to $\{1/36, 1/54\}$ and $\{1/54, 1/81\}$, respectively. In addition, selecting the number of eigenfunctions for each subdomain to construct the enrichment coarse space is based on the adaptive enrichment, where the threshold is 50.

In Table 1, we set different values for α_b , α_c , and α_i for different number of subdomains, for instance, see Figure 5 including 6×6 subdomains and the coarse mortar case. We implement the additive average Schwarz method with the adaptive enrichment coarse space $type = II$, where the threshold is 50, i.e., all eigenfunctions associated with all eigenvalues larger than 50 are considered. Note that the condition number of the non-enrichment preconditioner is very large. For instance, in the case 6×6 subdomains and coarse mortar, it is $1.32e7$. We first observe that the condition numbers of the enrichment preconditioners are proportional to the ratio H/h and independent of the number of subdomains. We also observe that if we consider the same number of eigenfunctions for each subdomain based on a proper threshold, consequently, the condition number estimates are independent of values of α_b , α_c and α_i .

Table 2 demonstrates only the condition number estimates and iteration numbers of the preconditioned conjugate gradient method (in parentheses) in the cases $type = II$, coarse mortar, 6×6 and 9×9 subdomains, where $\alpha_b = 1$, $\alpha_c = 1e4$, and $\alpha_i = 1e6$. For the other cases and also different values of jumps in the coefficient $\alpha(x)$, we have similar results. Also, the construction of the enrichment coarse space is relied on the fixed number of eigenfunctions for each subdomain varying from 0 to 7.

It is conclusive that both approaches to enrich the standard coarse space, i.e., using the given threshold and imposing the fixed numbers of

ADD		Number of Eigenfunctions in Each Subdomain							
		0	1	2	3	4	5	6	7
Number of Subdomains	6×6	1.32e7 (1756)	3.25e6 (1168)	2.65e5 (584)	4.10e4 (295)	4.12e3 (112)	5.7e1 (50)	5.6e1 (48)	5.58e1 (47)
	9×9	2.29e7 (4981)	7.54e6 (3889)	1.41e6 (2368)	3.82e5 (1180)	1.23e5 (353)	6.21e3 (118)	5.80e2 (63)	7.38e1 (50)

Table 2 The implementation of the additive average Schwarz method, $type = II$, with the fixed number of eigenfunctions for each subdomain to estimate the condition number of $\mathbf{B}_E^{type} \mathbf{A}$ and the number of iterations of the preconditioned conjugate gradient method (in parentheses) for 6×6 ($h = 1/36$, $h^* = 1/54$) and 9×9 ($h = 1/54$, $h^* = 1/81$) subdomains and the coarse mortar case. Further, the distribution of jumps in $\alpha(x)$ are $\alpha_b = 1$, $\alpha_c = 1e4$ and $\alpha_i = 1e6$.

ADD				Different Values for α_b , α_c and α_i			
				1, 1e3, 1e4		1, 1e4, 1e6	
				Coarse Mortar	Fine Mortar	Coarse Mortar	Fine Mortar
Number of Subdomains	6×6	$type$	I	625	525	635	534
			II	75	80	75	80
	9×9	$type$	I	1988	1680	1992	1681
			II	240	233	240	233

Table 3 The total numbers of the eigenfunctions associated with the eigenvalues greater than 50 to enrich the coarse spaces used in the additive average Schwarz method for both types I and II.

eigenfunctions for each subdomain, lead to similar results. This fact can be viewed by comparing, for instance, the third column of Table 1 with the last column of Table 2.

Table 3 gives the total number of required eigenfunctions for the adaptive enrichment coarse spaces in the cases 6×6 and 9×9 subdomains, $type = I$ and II and different values for $\alpha(x)$. As we can see from this table, solving the second type of the generalized eigenvalue problem (5) is more efficient than solving the first type. To analyze the distribution of eigenfunctions, Figure 6 contains the polar histograms for both coarse and fine mortars in the case 6×6 subdomains, and $type \in \{I, II\}$. It clearly shows that we need only consider a few eigenfunctions for each subdomain in the case $type = II$ compared to $type = I$.





(a) Fine mortar and $type = I$.(b) Fine mortar and $type = II$.(c) Coarse mortar and $type = I$.(d) Coarse mortar and $type = II$.

Fig. 6 The histograms of the number of eigenfunctions associated with the eigenvalues greater than 50 corresponding to the partition of $\Omega = [0, 1] \times [0, 1]$ into 6×6 subdomains, where $\alpha_b = 1$, $\alpha_c = 1e4$ and $\alpha_i = 1e6$. The number of subdomains are ordered anticlockwise around the polar histograms from 1 to 36. For $type = I$ and II , the number of eigenfunctions are grouped into the ranges 0 to 70 and 0 to 10, respectively. The largest numbers of the eigenfunctions in (a) – (d) are 39, 6, 63 and 5, respectively.

7 Conclusion

In this paper, we have employed the additive average Schwarz method with the enrichment coarse spaces to solve second order elliptic boundary problem coupled with very large jumps in the function $\alpha(x)$, where the finite element discretization of that problem has been based on the nonmatching triangulations across the subdomains interfaces. We have proved that the condition number estimates for the produced preconditioners by the additive average Schwarz methods for $type \in \{I, II\}$ are proportional to the ratio H/h and independent of the number of subdomains. Besides, we have compared the numerical results to conclude that, in practice, $type = II$ has much better performance than $type = I$.

Acknowledgements

The work of Leszek Marcinkowski was partially supported by Polish Scientific Grant: National Science Center: 2016/21/B/ST1/00350.

References

1. T. Arbogast, L. C. Cowsar, M. F. Wheeler and I. Yotov, Mixed finite element methods on nonmatching multiblock grids, *SIAM J. Numer. Anal.* 37(4), 1295–1315, (2000)
2. T. Arbogast, G. Pencheva, M. F. Wheeler and I. Yotov, A multiscale mortar mixed finite element method, *Multiscale Model. Simul.* 6(1), 319–346, (2007)
3. T. Arbogast and H. Xiao, A multiscale mortar mixed space based on homogenization for heterogeneous elliptic problems, *SIAM J. Numer. Anal.* 51(1), 377–399, (2013)
4. C. Bernardi, Y. Maday and F. Rapetti, Basics and some application of mortar element method, *GAMM-Mitt.* 28, 97–123, (2005)
5. S. C. Brenner, Two-level additive Schwarz preconditioners for nonconforming finite element methods, *Math. Comp.* 65, 897–921, (1996)
6. M. Brezina, C. Heberton, J. Mandel and P. Vaněk, An iterative method with convergence rate chosen a priori, Tech. Report 140, Center for Computational Mathematics, University of Colorado Denver, Denver, (1999)
7. P. E. Bjørstad, M. Dryja and E. Vainikko, Parallel implementation of a Schwarz domain decomposition algorithms, in *Applied Parallel Computing in Industrial Problems and Optimization* (J. Wasniewski, J. Dongara, K. Madsen, and D. Olsem, eds.) Lecture Notes in Computer Science, Springer, 1184, 141–157, (1996)
8. P. E. Bjørstad, M. Dryja and T. Rahman, Additive Schwarz methods for elliptic mortar finite element problems, *Numer. Math.* 95, 427–457, (2003)
9. T. F. Chan and T. P. Mathew, Domain decomposition algorithms, *Acta Numerica.* 3, 61–143, (1994)
10. P. G. Ciarlet, *The Finite Element Method for Elliptic Problems*, North-Holland, Amsterdam, (1978)
11. L. C. Cowsar, Domain decomposition methods for nonconforming finite elements spaces of Lagrange-type, *Proceedings of the Sixth Copper Mountain Conference on Multigrid Methods*, NASA Conference Publication 3224, 93–109, (1993)
12. M. Dryja and O. B. Widlund, An additive variant of the Schwarz alternating method for the case of many subregions, Technical Report 339, also Ultracomputer Note 131, Department of Computer Science, Courant Institute, New York University, (1987)
13. M. Dryja, A. Gantner, O. B. Widlund and B. I. Wohlmuth, Multilevel additive Schwarz preconditioner for nonconforming mortar finite element methods, *J. Numer. Math.* 12, 23–38, (2004)

14. M. Dryja and M. Sarkis, Additive average Schwarz methods for discretization of elliptic problems with highly discontinuous coefficients, *Comput. Methods Appl. Math.* 10, 164–176, (2010)
15. Y. Efendiev, J. Galvis and P. S. Vassilevski, Spectral element agglomerate algebraic multigrid methods for elliptic problems with high contrast coefficients. In: Y. Huang, R. Kornhuber, O. B. Widlund, J. Xu (eds.) *Domain Decomposition Methods in Science and Engineering XIX*, Lecture Notes in Computational Science and Engineering, vol. 78, pp. 407–414. Springer, Berlin (2011)
16. E. Eikeland, L. Marcinkowski and T. Rahman, Overlapping Schwarz methods with adaptive coarse spaces for multiscale problems in 3D, *Numer. Math.* 142, 103–128, (2019)
17. X. Feng and T. Rahman, An additive average Schwarz method for the plate bending problem, *J. Numer. Math.* 10, 109–125, (2002)
18. M. J. Gander, A. Loneland and T. Rahman, Analysis of a new harmonically enriched multiscale coarse space for domain decomposition methods, arXiv preprint arXiv:1512.05285 (2015)
19. A. Heinlein, A. Klawonn, J. Knepper and O. Rheinbach, Adaptive GDSW coarse spaces for overlapping Schwarz methods in three dimensions, *SIAM J. Sci. Comput.* 41(5), A3045–A3072, (2019)
20. H. H. Kim and E. T. Chung, A BDDC algorithm with enriched coarse spaces for two-dimensional elliptic problems with oscillatory and high contrast coefficients, *Multiscale Model. Simul.* 13, 571–593, (2015)
21. H. H. Kim, E. Chung and J. Wang, BDDC and FETI-DP preconditioners with adaptive coarse spaces for three-dimensional elliptic problems with oscillatory and high contrast coefficients, *J. Comput. Phys.* 349, 191–214, (2017)
22. A. Klawonn, M. Lanser, P. Radtke and O. Rheinbach, On an adaptive coarse space and on nonlinear domain decomposition. In: J. Erhel, M. J. Gander, L. Halpern, G. Pichot, T. Sassi, O. B. Widlund (eds.) *Domain Decomposition Methods in Science and Engineering XXI*, Lecture Notes in Computational Science and Engineering, vol. 98, pp. 71–83. Springer, Switzerland (2014)
23. A. Klawonn, P. Radtke and O. Rheinbach, FETI-DP methods with an adaptive coarse space, *SIAM J. Numer. Anal.* 53(1), 297–320, (2015)
24. A. Loneland, L. Marcinkowski and T. Rahman, Additive average Schwarz method for a Crouzeix-Raviart finite volume element discretization of elliptic problems with heterogeneous coefficients, *Numer. Math.* 134, 91–118, (2016)
25. J. Mandel, B. Sousedk and J. Šístek, Adaptive BDDC in three dimensions, *Math. Comput. Simulation.* 82, 1812–1831, (2012)
26. L. Marcinkowski and T. Rahman, Additive average Schwarz with adaptive coarse spaces: scalable algorithms for multiscale problems, *Electron. Trans. Numer. Anal.* 49, 28–40., (2018)
27. A. M. Matsokin and S. V. Nepomnyaschikh, A Schwarz alternating method in a subspace. *Sov. Math.* 29(10), 78–84, (1985)
28. T. Rahman, X. Xu and R. Hoppe, Additive Schwarz methods for the Crouzeix-Raviart mortar finite element for elliptic problems with discontinuous coefficients, *Numer. Math.* 101(3), 551–572, (2005)
29. M. Sarkis, Two-level Schwarz methods for nonconforming finite elements and discontinuous coefficients. Proceedings of the Sixth Copper Mountain Conference on Multigrid Methods, NASA Conference Publication 3224, 543–565, (1993)
30. M. Sarkis, Nonstandard coarse spaces and Schwarz methods for elliptic problems with discontinuous coefficients using non-conforming elements, *Numer. Math.* 77(3), 383–406, (1997)
31. P. Seshaiyer and M. Suri, Uniform hp convergence results for the mortar finite element method, *Math. Comp.* 69, 521–546, (2000)
32. B. F. Smith, P. E. Bjørstad and W. D. Gropp, *Domain Decomposition: Parallel Multilevel Methods for Elliptic Partial Differential Equations*, Cambridge University Press, (1996).
33. N. Spillane and D. J. Rixen, Automatic spectral coarse spaces for robust finite element tearing and interconnecting and balanced domain decomposition algorithms, *Int. J. Numer. Methods. Eng.* 95(11), 953–990, (2013)

-
34. N. Spillane, V. Dolean, P. Hauret, F. Nataf, C. Pechstein and R. Scheichl, Abstract robust coarse spaces for systems of PDEs via generalized eigenproblems in the overlaps, *Numer. Math.* 126(4), 741–770, (2014)
 35. P. Le Tallec, Y. H. De Roeck and M. Vidrascu, Domain decomposition methods for large linearly elliptic three-dimensional problems, *J. Comput. Appl. Math.* 34, 93–117, (1991)
 36. A. Toselli and O. B. Widlund, Domain decomposition methods-algorithms and theory, volume 34 of Springer Series in Computational Mathematics, Springer-Verlag, Berlin, (2005)
 37. J. Xu and J. Zou, Some nonoverlapping domain decomposition methods, *SIAM Rev.* 40, 857–914, (1998)