## STABILITY IMPLIES ROBUST CONVERGENCE OF A CLASS OF PRECONDITIONED PARALLEL-IN-TIME ITERATIVE ALGORITHMS

SHU-LIN WU\*, TAO ZHOU<sup>†</sup>, AND ZHI ZHOU<sup>‡</sup>

Abstract. Solving evolutionary equations in a parallel-in-time manner is an attractive topic and many algorithms are proposed in recent two decades. The algorithm based on the block  $\alpha$ -circulant preconditioning technique has shown promising advantages, especially for wave propagation problems. By fast Fourier transform for factorizing the involved circulant matrices, the preconditioned iteration can be computed efficiently via the so-called diagonalization technique, which yields a direct parallel implementation across all time levels. In recent years, considerable efforts have been devoted to exploring the convergence of the preconditioned iteration by studying the spectral radius of the iteration matrix, and this leads to many case-by-case studies depending on the used time-integrator. In this paper, we propose a unified convergence analysis for the algorithm applied to u' + Au = f, where  $\sigma(A) \subset \mathbb{C}^+$  with  $\sigma(A)$  being the spectrum of  $A \in \mathbb{C}^{m \times m}$ . For any one-step method (such as the Runge-Kutta methods) with stability function  $\mathcal{R}(z)$ , we prove that the decay rate of the global error is bounded by  $\alpha/(1 - \alpha)$ , provided the method is stable, i.e.,  $\max_{\lambda \in \sigma(A)} |\mathcal{R}(\Delta t\lambda)| \leq 1$ . For any linear multistep method, such a bound becomes  $c\alpha/(1 - c\alpha)$ , where  $c \geq 1$  is a constant specified by the multistep method itself. Our proof only relies on the stability of the time-integrator and the estimate is independent of the step size  $\Delta t$  and the spectrum  $\sigma(A)$ .

Key words. time-parallel algorithm,  $\alpha$ -circulant preconditioner, convergence analysis, stability

AMS subject classifications. 65M55, 65M12, 65M15, 65Y05

**1. Introduction.** We are interested in solving the following evolutionary equation parallel-in-time (PinT):

$$y' + Ay = g, \ A \in \mathbb{C}^{m \times m},\tag{1.1}$$

with initial value  $y(0) = y_0$ . The spectrum of the matrix A is supposed to lay on the right hand-side of the complex plane, i.e.,

$$\sigma(A) \subseteq \mathbb{C}^+ = \{ z = \delta + i\omega : \delta \ge 0, \omega \in \mathbb{R} \}.$$

Note that the system (1.1) is widely used as an spatially semi-discrete scheme of many evolutionary partial differential equations, e.g.:

- Diffusion equation:  $u_t \Delta u = f$ , where  $A = -\Delta_h \approx \Delta$  denotes the discrete Laplacian, and we have  $\sigma(A) \subseteq \mathbb{R}^+$ .
- Acoustic wave equation:  $u_{tt} \Delta u = f$ . In this case, we can make an order reduction to yield the first-order system (1.1) with

$$y = \begin{bmatrix} u \\ v \end{bmatrix}, \ A = \begin{bmatrix} & -I_x \\ -\Delta_h \end{bmatrix}, \ g = \begin{bmatrix} 0 \\ f \end{bmatrix},$$
(1.2)

Then we have  $\sigma(A) \subseteq i\mathbb{R}^+$ .

• Schrodinger equation:  $iu_t - \Delta u + V(x)u = f$  with a positive potential term V(x) > 0. Then we have  $\sigma(A) \subseteq i\mathbb{R}^+$ .

In the recent two decades, the research toward reducing the computation time via time parallelization is a hot topic (see [10] for a comprehensive review). In this paper,

<sup>\*</sup>School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China. E-mail: wushulin84@hotmail.com

<sup>&</sup>lt;sup>†</sup>LSEC, Institute of Computational Mathematics and Scientific/Engineering Computing, AMSS, Chinese Academy of Sciences, Beijing, 100190, China. E-mail: tzhou@lsec.cc.ac.cn

<sup>&</sup>lt;sup>‡</sup>Department of Applied Mathematics, The Hong Kong Polytechnic University, Kowloon, Hong Kong. E-mail: zhizhou@polyu.edu.hk

we will focus on the approach that uses the diagonalization technique proposed by Maday and Ronquist in [29]. In that pioneer work, they formulated the time-stepping system into a space-time *all-at-once* system. Then they diagonalize the time stepping matrix and solve all time steps in parallel. Such a technique was first used in [15] for wave equations. The concrete algorithm lies in using the geometrically increasing time step-sizes  $\{\Delta t_n = \mu^{n-1}\Delta t_1\}_{n=1}^{N_t}$  with some parameter  $\mu > 1$ , in order to make the time discretization matrix diagonalizable. The algorithm is directly parallel, but the roundoff error arising from the diagonalization procedure increases dramatically for large  $N_t$ . In order to balance the roundoff error and the discretization error, the quantity  $N_t$  can not be too large (in practice, people choose  $N_t = 20 \sim 25$ ).

To overcome the restriction on  $N_t$ , the diagonalization technique was used in an iterative fashion [19, 21, 23, 25, 26, 28], that will be investigated in this paper. The idea is briefly explained as follows. First, we partition the time domain (0, T) by  $N_t$  equally spaced time points  $\{t_n\}_{n=0}^{N_t}$  with step size  $\Delta t = T/N_t$ . Then we apply a time-integrator to (1.1) (such as Runge-Kutta (RK) methods)

$$y_{n+1} + \mathcal{R}(\Delta t A)y_n = \eta_n, \quad n = 0, 1, \dots, N_t - 1,$$
 (1.3)

with a given initial value  $y_0$ , where  $\eta_n$  is a known quantity specified by the source term at  $t_n$ . The increment matrix  $\mathcal{R}(\Delta tA)$  is deduced from the stability function of the timeintegrator. Instead of solving the  $N_t$  difference equations (1.3) level-by-level, we form these equations into an *all-at-once* system

$$\mathcal{K}\boldsymbol{u} = \boldsymbol{b},\tag{1.4}$$

where 
$$\boldsymbol{u} = (y_1^{\top}, y_2^{\top}, \dots, y_{N_t}^{\top})^{\top}, \boldsymbol{b} = ((\eta_0 - \mathcal{R}(\Delta t A)y_0)^{\top}, \eta_1^{\top}, \dots, \eta_{N_t}^{\top})^{\top}$$
 and  
 $\mathcal{K} = I_t \otimes I_x + B \otimes \mathcal{R}(\Delta t A).$ 
(1.5)

Here and hereafter  $I_t \in \mathbb{R}^{N_t \times N_t}$  and  $I_x \in \mathbb{R}^{m \times m}$  denotes the identity matrices in time and space respectively and the Toeplitz matrix B is

$$B = \begin{bmatrix} 0 & & & \\ 1 & 0 & & \\ & \ddots & \ddots & \\ & & 1 & 0 \end{bmatrix} \in \mathbb{R}^{N_t \times N_t}.$$
 (1.6)

To solve the all-at-once system (1.4), we propose a block  $\alpha$ -circulant preconditioner:

$$\mathcal{P}_{\alpha} = I_t \otimes I_x + C(\alpha) \otimes \mathcal{R}(\Delta tA), \tag{1.7}$$

where  $C(\alpha)$  denotes the following  $\alpha$ -circulant matrix

$$C(\alpha) = \begin{bmatrix} 0 & & & \alpha \\ 1 & 0 & & \\ & \ddots & \ddots & \\ & & 1 & 0 \end{bmatrix} \in \mathbb{R}^{N_t \times N_t},$$
 (1.8)

and apply the preconditioned iteration:

$$\mathcal{P}_{\alpha}\Delta \boldsymbol{u}^{k} = \boldsymbol{r}^{k}, \boldsymbol{u}^{k+1} = \boldsymbol{u}^{k} + \Delta \boldsymbol{u}^{k}, \ \boldsymbol{r}^{k} := \boldsymbol{b} - \mathcal{K}\boldsymbol{u}^{k}, \ k = 0, 1, \dots$$
(1.9)

Thanks to the property of the  $\alpha$ -circulant matrices (see, e.g., [1, Lemma 2.1]), we can make a block Fourier diagonalization for the preconditioner  $\mathcal{P}_{\alpha}$  and this allows to compute  $\mathcal{P}_{\alpha}^{-1}(\mathcal{K}\boldsymbol{u}^{k}-\boldsymbol{b})$  in (1.9) highly parallel for all time levels [19,21,23,25,26,28].

In this paper, we aim to answer the question: under what conditions the iterative algorithm (1.9) converges rapidly and robustly? In existing references, people answered this question by examining the radius of the spectrum of the iteration matrix  $\mathcal{I} - \mathcal{P}_{\alpha}^{-1}\mathcal{K}$ , see e.g., [19, 25, 26, 28, 37] for parabolic problems and [16, 23] for the wave equations. For some special time-integrators (e.g., the implicit Euler method [25, 26], the implicit leap-frog method [23], and the two-stage singly diagonal implicit RK method [37]) it was shown

$$\rho(\mathcal{I} - \mathcal{P}_{\alpha}^{-1}\mathcal{K}) \le \frac{\alpha}{1 - \alpha}, \ \alpha \in (0, 1).$$
(1.10)

The analysis in these works is rather technical and heavily depends on the special property of the time-integrator, e.g., the sparseity, Toeplitz structure and diagonal dominance of the time-discretization matrix.

Instead of exploring the spectrum of the iterative matrix, we shall examine the error of the preconditioned iteration directly, and prove that for *any* one-step time stepping method (1.3) the error  $err^k = u^k - u$  of the algorithm (1.9) satisfies (Theorem 2.1)

$$\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty} \leq \frac{\alpha}{1-\alpha} \|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty},$$
(1.11)

where P is the eigenvector matrix of A, provided the method solving (1.1) is stable, i.e.,  $\max_{\lambda \in \sigma(A)} |\mathcal{R}(\Delta t\lambda)| \leq 1$ . If  $\alpha \in (0, 1/2)$ , the preconditioned iteration (1.9) converges linearly with the convergence factor  $\alpha/(1-\alpha)$ . The result (1.11) well explains the robust convergence observed in [16, 17, 25, 26], because all the time-integrators used there are unconditionally stable.

Similarly, for any stable linear multistep method, we can also develop a preconditioned iterative solver, and show that the iteration converges linearly (for suitable parameter  $\alpha$ ) and satisfies (Theorem 3.1)

$$\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty} \le \frac{c\alpha}{1-c\alpha} \|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty},$$
(1.12)

where  $c \ge 1$  is a generic constant only depending on the stability property of the multistep method. For parabolic equations and subdiffusion models with memory effects, such an iterative algorithm for r-step backward differentiation formula up to r = 6 has been analyzed in a most recent work [35]. The current work provides a more systematic argument showing the relation between stability of time-integrators and the convergence of the iterative solver (1.9).

These estimates imply that the convergence of the iterative algorithm (1.9) is very fast when we choose a very small  $\alpha$ . However, in practice, the parameter  $\alpha$  can not be arbitrarily small, because the roundoff error arising from the block diagonalization of  $\mathcal{P}_{\alpha}$ will increase dramatically when  $\alpha \to 0$ . Such a roundoff error is explained as follows. For any diagonalizable square matrix Q with eigenvalue matrix D and eigenvector matrix V, due to the floating point operations Q can not precisely equal to  $VDV^{-1}$  and it only holds  $Q \approx VDV^{-1}$ . In our case, for the  $\alpha$ -circulant matrix  $C(\alpha)$  in (1.8), the difference between  $C(\alpha)$  and  $VDV^{-1}$  continuously increase as  $\alpha$  decreases. See [16,35] for some detailed studies on the roundoff error of the diagonalization procedure, where the authors proved

roundoff error = 
$$O(\epsilon \alpha^{-2})$$
, (1.13)

where  $\epsilon$  is the machine precision ( $\epsilon = 2.2204 \times 10^{-16}$  for a 32-bit computer).

The rest of this paper is organized as follows. In Section 2, we prove (1.11) for the one-step time-integrators. The proof lies in representing the preconditioned iteration (1.9) as a special difference equations with head-tail coupled condition and relays on the stability

function only. We also consider the linear multistep methods in Section 3, for which we prove (1.12) and the stability of the numerical method plays a central role in the proof as well. We conclude this paper in Section 4 by listing some important issues that need to be addressed in the future.

2. Convergence of iteration (1.9) for one-step methods. In this section, we prove the convergence of the iterative solver (1.9). To this end, we assume that the one-step time-integrator (1.3) is stable in the following sense.

ASSUMPTION 1. The matrix A in the linear system (1.1) is diagonalizable as  $A = PD_AP^{-1}$  with  $\sigma(A) \subset \mathbb{C}^+$ . For such a system, it holds

$$|\mathcal{R}(\Delta t\lambda)| \le 1 \qquad \forall \ \lambda \in \sigma(A), \tag{2.1}$$

where  $\mathcal{R}(\cdot)$  denotes the stability function of the one-step method (1.3).

Then we are ready to state our main theorem.

THEOREM 2.1. Under Assumption 1, the error at the k-th iteration in (1.9), denoted by  $err^{k} = u^{k} - u$ , satisfies

$$\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty} \le \frac{\alpha}{1-\alpha} \|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty} \qquad \forall \ k \ge 1.$$
(2.2)

Therefore, the iteration (1.9) converges linearly if  $\alpha \in (0, 1/2)$ .

*Proof.* From (1.9), we observe that the error  $err^k$  satisfies

$$err^{k+1} = err^k - (\mathcal{P}_{\alpha}^{-1}\mathcal{K})err^k, \ k = 0, 1, \dots$$

Now we define

$$\boldsymbol{\zeta}^k = ((\boldsymbol{\xi}_1^k)^\top, (\boldsymbol{\xi}_2^k)^\top, \dots, (\boldsymbol{\xi}_m^k)^\top)^\top = (I_t \otimes P)\boldsymbol{err}^k.$$

Let z be an arbitrary eigenvalue of  $\Delta t A$  and  $\boldsymbol{\xi}^k \in \mathbb{R}^{N_t}$  be the corresponding subvector of  $\boldsymbol{\zeta}^k$ . Then, it is clear that

$$\boldsymbol{\xi}^{k+1} = \boldsymbol{\xi}^k - (P_{\alpha}^{-1}(z)K(z))\boldsymbol{\xi}^k, \ k = 0, 1, \dots$$
(2.3)

with

$$K(z) = I_t + \mathcal{R}(z)C_t, \quad P_\alpha(z) = I_t + \mathcal{R}(z)C(\alpha).$$

By applying  $\mathcal{P}_{\alpha}$  on (2.3), we derive that for  $k \geq 1$ 

$$P_{\alpha}(z)\boldsymbol{\xi}^{k+1} = (P_{\alpha}(z) - K(z))\boldsymbol{\xi}^{k}.$$
(2.4)

Let

$$oldsymbol{\xi}^k = (\xi_1^k, \xi_2^k, \dots, \xi_{N_t}^k)^ op$$

Then the relation (2.4) implies

$$(I_t + \mathcal{R}(z)C_t) \boldsymbol{\xi}^{k+1} = \alpha \begin{bmatrix} \mathcal{R}(z)(\boldsymbol{\xi}_{N_t}^{k+1} - \boldsymbol{\xi}_{N_t}^k) \\ 0 \\ \vdots \\ 0 \end{bmatrix}.$$

Therefore, (2.3) is equivalent to the following difference equations with a head-tail coupled condition

$$\begin{cases} \xi_{n+1}^{k+1} + \mathcal{R}(z)\xi_{n}^{k+1} = 0, \ n = 0, 1, \dots, N_t - 1, \\ \xi_0^{k+1} = \alpha(\xi_{N_t}^{k+1} - \xi_{N_t}^k). \end{cases}$$
(2.5)

By Assumption 1, we know that the time-integrator is stable, i.e.,  $|\mathcal{R}(z)| \leq 1$ . This together with the first relation in (2.5) immediately implies for all  $k \geq 0$ 

$$\left|\xi_{n+1}^{k+1}\right| = |\mathcal{R}(z)| \left|\xi_{n}^{k+1}\right| \le \left|\xi_{n}^{k+1}\right|.$$

Hence, the the second relation in (2.5) leads to

$$\left|\xi_{n}^{k+1}\right| \leq \left|\xi_{0}^{k+1}\right| = \alpha \left|\xi_{N_{t}}^{k+1} - \xi_{N_{t}}^{k}\right| \leq \alpha \left|\xi_{N_{t}}^{k+1}\right| + \alpha \left|\xi_{N_{t}}^{k}\right| \leq \alpha \left|\xi_{n}^{k+1}\right| + \alpha \left|\xi_{n}^{k}\right|,$$

where  $n = 0, 1, \ldots, N_t$ . That is,

$$\left|\xi_{n}^{k+1}\right| \leq \frac{\alpha}{1-\alpha} \left|\xi_{n}^{k}\right|.$$

To summarize, it holds

$$\|\boldsymbol{\xi}^{k+1}\|_{\infty} \le \frac{\alpha}{1-\alpha} \|\boldsymbol{\xi}^k\|_{\infty}.$$
(2.6)

For  $\boldsymbol{\zeta}^k = ((\boldsymbol{\xi}_1^k)^{\top}, (\boldsymbol{\xi}_2^k)^{\top}, \dots, (\boldsymbol{\xi}_m^k)^{\top})^{\top}$  with any index k, it is clear that

$$\|\boldsymbol{\zeta}^k\|_{\infty} = \max_{j=1,2,\dots,m} \|\boldsymbol{\xi}_j^k\|_{\infty}.$$

Since  $\boldsymbol{\xi}^k$  is an arbitrary subvector of  $\boldsymbol{\zeta}^k$ , it follows from (2.6) that  $\|\boldsymbol{\zeta}^{k+1}\|_{\infty} \leq \frac{\alpha}{1-\alpha} \|\boldsymbol{\zeta}^k\|_{\infty}$ . This, together with  $\boldsymbol{\zeta}^k = (I_t \otimes P) \boldsymbol{err}^k$ , gives the desired result.  $\Box$ 

REMARK 2.1. Theorem 2.1 reveals the relation between the stability of the time stepping scheme (1.3) and the convergence of preconditioned iterative solver (1.9). The proof only utilizes the stability of the time stepping scheme instead of any matrix structure of the all-at-once system explicitly, and can be extended to other numerical schemes, e.g. linear multistep methods, which will be introduced in the next section.

To test the sharpness of the estimate (2.2), we consider the following advection-diffusion equation

$$\begin{cases} u_t - \nu u_{xx} + u_x = 0, & (x, t) \in (-\frac{1}{2}, \frac{1}{2}) \times (0, T), \\ u(x, 0) = \sin(2\pi x), & x \in (-\frac{1}{2}, \frac{1}{2}), \end{cases}$$
(2.7)

with the periodic boundary condition, where  $\nu > 0$  is a constant and for  $\nu$  small the solution u has obvious characteristic of wave propagation. We use the centered finite difference formula to discretize the spatial derivates

$$u_{xx}(x_i, t) \approx \frac{u_{i+1}(t) - 2u_i(t) + u_{i-1}(t)}{\Delta x^2}$$
 and  $u_x(x_i, t) \approx \frac{u_{i+1}(t) - u_{i-1}(t)}{2\Delta x}$ 

with  $i = 1, 2, ..., N_x - 1$  and  $\Delta x = \frac{1}{N_x}$ . This leads to ODE system (1.1) with

$$A = \frac{\nu}{\Delta x^2} \begin{bmatrix} 2 & -1 & & -1 \\ -1 & 2 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & -1 & 2 & -1 \\ -1 & & & -1 & 2 \end{bmatrix} + \frac{1}{2\Delta x} \begin{bmatrix} 0 & -1 & & 1 \\ 1 & 0 & -1 & & \\ & \ddots & \ddots & \ddots & \\ & & 1 & 0 & -1 \\ -1 & & & 1 & 0 \end{bmatrix}.$$
 (2.8)

For time-discretization, we consider the following two-stage SDIRK method

$$\begin{array}{c|cc} \gamma & \gamma & 0\\ \underline{\gamma + \tilde{\gamma}} & \tilde{\gamma} & \gamma\\ \hline b & 1 - b \end{array},$$
(2.9)

where  $\gamma > 0$  and  $b \neq 1$ . From [22, Chapter IV.6], the method is of order 2 if

$$\gamma b + (\tilde{\gamma} + \gamma)(1 - b) = \frac{1}{2} \left( \Rightarrow \tilde{\gamma} = \frac{\frac{1}{2} - \gamma b}{1 - b} - \gamma \right).$$
(2.10a)

The stability function is

$$\mathcal{R}(z) = \frac{(2\gamma^2 - 4\gamma + 1)z^2 - (2 - 4\gamma)z + 2}{2(\gamma z + 1)^2}.$$
(2.10b)

It is easy to verify that  $|\mathcal{R}(z)| \leq 1 (\forall z \in \mathbb{C}^+)$ , i.e., the SDIRK method (2.9) is unconditionally stable, if and only if  $\gamma \geq \frac{1}{4}$ . Otherwise the method is only conditionally stable. Here, we consider two SDIRK methods

which correspond to  $\gamma = 0.2$  and  $\gamma = \frac{3+\sqrt{3}}{6}$  respectively (for both methods  $b = \frac{1}{2}$ ). Let  $\Delta x = 0.01$  and  $\Delta t = 0.02$ . Then, for two values of  $\nu$ :  $\nu = 10^{-3}$  and  $\nu = 2 \times 10^{-4}$ , we show in Figure 2.1 the spectrum  $\sigma(\Delta tA)$  and the stability region of the SDIRK method with  $\gamma = 0.2$ , i.e., the second method in (2.11). We see that for  $\nu = 10^{-3}$  this SDIRK method is stable, while it is not for  $\nu = 2 \times 10^{-4}$ . For  $\nu = 2 \times 10^{-4}$ , the second SDIRK method in (2.11) is suitable because it is unconditionally stable. By using the two methods in (2.11) for  $\nu = 10^{-3}$  and  $\nu = 2 \times 10^{-4}$  respectively, we show in Figure 2.2 the measured error  $\|\mathbf{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\mathbf{err}^k\|_{\infty}$ , together with the upper bound predicted by  $\frac{\alpha}{1-\alpha}$  (dotted line). We see that  $\|\mathbf{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\mathbf{err}^k\|_{\infty}$  decay with a very similar rate and that the estimate (2.2) predicts such a rate very well. Both  $\|\mathbf{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\mathbf{err}^k\|_{\infty}$  do not smoothly decay to the machine precision 2.2204 \times 10^{-16} and at the last few iterations they stagnant at the level  $O(10^{-13})$  for  $\alpha = 0.1$  and  $O(10^{-12})$  for  $\alpha = 0.01$ . Such a stagnation is due to the roundoff error arising from the block diagonalization of the preconditioner  $\mathcal{P}_{\alpha}$ .



FIG. 2.1. The stability region (shadow region) of the second SDIRK method in (2.11) (i.e.,  $\gamma = 0.2$ ) and the spectrum  $\sigma(\Delta tA)$  with A being the matrix given by (2.8) and  $\Delta t = 0.02$ .

We note that the advection-dominated diffusion equation is a well-known difficult problem for PinT computation. In particular, for equation (2.7) with  $\nu = 10^{-3}$  and  $\nu = 2 \times 10^{-4}$ 



FIG. 2.2. The measured error  $\|\boldsymbol{err}^{k}\|_{\infty}$  and  $\|(I_t \otimes P)\boldsymbol{err}^{k}\|_{\infty}$  together with the upper bound by  $\frac{\alpha}{1-\alpha}$ . Left:  $\nu = 10^{-3}$  and we use the SDIRK method in (2.11) with  $\gamma = 0.2$ . Right:  $\nu = 2 \times 10^{-4}$  and we use the SDIRK method in (2.11) with  $\gamma = \frac{3+\sqrt{3}}{6}$ . Here, T = 10 and  $\Delta t = 0.02$ .

the convergence rate of the mainstream PinT algorithms *parareal* and MGRiT is rather disappointing and the convergence rate continuously deteriorates as we refine  $\Delta x$  (see [17, Section 1] for more details).

**3.** Convergence of iteration (1.9) for linear multistep methods. In this section, we consider the linear multistep methods. For parabolic equations, the parallel algorithms for BDF methods (with initial corrections) up to order six has been proposed and analyzed in [35], and has been generalized to the diffusion models with historical memory effects. In the present work, we propose a systematic approach, showing the convergence of the preconditioned iteration, which works for all stable linear multistep methods.

We consider the r-step method  $(r \ge 1)$  with time levels  $t_n = n\Delta t$   $(n = 1, ..., N_t)$ : for given  $y_0, y_1, \ldots, y_{r-1}$ , we find  $y_{n+r}$  such that

$$\sum_{j=0}^{r} a_j y_{n+r-j} = \Delta t \sum_{j=0}^{r} b_j (-Ay_{n+r-j} + g_{n+r-j}), \quad n = 0, 1, \dots, N_t - r.$$
(3.1)

Without loss of generality, we assume that  $a_0 = 1$ . Then we define the characteristic polynomials for  $z \in \mathbb{C}^+$ :

$$p(s;z) = \sum_{j=0}^{r} a_j s^{r-j} + z b_j s^{r-j}.$$
(3.2)

We assume that the r-step method is stable (Assumption 2). In principle, for any given angle  $\theta \in (0, \pi/2)$ , there exist  $A(\theta)$ -stable r-step methods of order r for every  $r \ge 1$ .

ASSUMPTION 2. The matrix A in the linear system (1.1) is diagonalizable as  $A = PD_AP^{-1}$  with  $\sigma(A) \subset \mathbb{C}^+$ . Then for  $z = \Delta t\lambda$  with  $\lambda \in \sigma(A)$ , the characteristic polynomial satisfies the following root condition:

$$p(s;z) = 0 \implies \begin{cases} either \ |s| < 1, \\ or \ |s| = 1 \text{ and it is a root of multiplicity 1.} \end{cases}$$
(3.3)

Assumption 2 immediately implies a stability estimate of the following discrete initial value problem: for given  $\xi_0, \xi_1, \ldots, \xi_{r-1}$ , find  $\xi_{r+n}$  such that

$$\sum_{j=0}^{r} a_j \xi_{n+r-j} = \sum_{j=0}^{r} b_j (-z\xi_{n+r-j} + \Delta t f_{n+r-j}), \quad n = 0, 1, \dots, N_t - r,$$

where  $z = \lambda \Delta t$  and  $\lambda \in \sigma(A)$ . Then Assumption 2 leads to the following stability result [18, Theorem 6.3.2]

$$|\xi_n| \le c \Big( \max_{0 \le j \le r-1} |\xi_j| + \Delta t \sum_{j=r}^n |f_j| \Big),$$

where the constant c is independent of n,  $\Delta t$  and  $N_t$ .

Next, we describe the parallel-in-time solver of the time-stepping scheme (3.1). Similar to the one-step methods, we can also form (3.1) into an *all-at-once* system (1.4) with

$$\mathcal{K} = B_1 \otimes I_x + \Delta t B_2 \otimes A, \tag{3.4}$$

where  $B_1, B_2 \in \mathbb{R}^{N_t \times N_t}$  are Toeplitz matrices

$$B_{1} = \begin{bmatrix} a_{0} & & & & & \\ a_{1} & a_{0} & & & & \\ \vdots & \ddots & \ddots & & & \\ a_{r} & & \ddots & \ddots & & \\ & \ddots & & a_{1} & a_{0} \\ & & & a_{r} & \dots & a_{1} & a_{0} \end{bmatrix}, B_{2} = \begin{bmatrix} b_{0} & & & & & \\ b_{1} & b_{0} & & & \\ \vdots & \ddots & \ddots & & \\ b_{r} & & \ddots & \ddots & \\ & \ddots & & b_{1} & b_{0} \\ & & & b_{r} & \dots & b_{1} & b_{0} \end{bmatrix}.$$

Again, we solve the all-at-once system (1.4) by the preconditioned iteration (1.9) with preconditioner

$$\mathcal{P}_{\alpha} = C_1(\alpha) \otimes I_x + C_2(\alpha) \otimes A, \tag{3.5}$$

where  $C_1(\alpha)$  and  $C_2(\alpha)$  are  $\alpha$ -circulant matrices

$$C_{1}(\alpha) = \begin{bmatrix} a_{0} & \alpha a_{r} & \dots & \alpha a_{2} & \alpha a_{1} \\ a_{1} & a_{0} & & & \alpha a_{2} \\ \vdots & \ddots & \ddots & & \ddots & \vdots \\ a_{r} & & \ddots & \ddots & & \alpha a_{r} \\ & \ddots & & a_{1} & a_{0} \\ & & & a_{r} & \dots & a_{1} & a_{0} \end{bmatrix}, C_{2}(\alpha) = \begin{bmatrix} b_{0} & \alpha b_{r} & \dots & \alpha b_{2} & \alpha b_{1} \\ b_{1} & b_{0} & & & \alpha b_{2} \\ \vdots & \ddots & \ddots & & \ddots & \vdots \\ b_{r} & & \ddots & \ddots & & \ddots & \vdots \\ b_{r} & & \ddots & \ddots & & \alpha b_{r} \\ & \ddots & & b_{1} & b_{0} \\ & & & b_{r} & \dots & b_{1} & b_{0} \end{bmatrix}$$

Then we are ready to show the convergence of the preconditioned iteration (1.9) for the linear multistep methods.

THEOREM 3.1. Under Assumption 2, the error at the k-th iteration in (1.9), denoted by  $err^k = u^k - u$ , satisfies

$$\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty} \leq \frac{c\alpha}{1-c\alpha}\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty},$$
(3.6)

where c > 0 denotes a generic constant independent of  $\Delta t$  and  $N_t$ . Therefore, the iteration (1.9) converges linearly if  $\alpha \in (0, 1/2c)$ .

*Proof.* From (1.9), the error  $err^k$  satisfies

$$err^{k+1} = err^k - (\mathcal{P}_{\alpha}^{-1}\mathcal{K})err^k \qquad \forall \ k \ge 0.$$

We apply  $I_t \otimes P$  on both sides of the above equation and define

$$oldsymbol{\zeta}^k = ((oldsymbol{\xi}_1^k)^ op, (oldsymbol{\xi}_2^k)^ op, \dots, (oldsymbol{\xi}_m^k)^ op)^ op = (I_t\otimes P)oldsymbol{err}^k.$$

Let z be an arbitrary eigenvalue of  $\Delta tA$ ,  $\boldsymbol{\xi}^k \in \mathbb{R}^{N_t}$  be the corresponding subvector of  $\boldsymbol{\zeta}^k$ . Define

$$K(z) = C_1 + zC_2, \ P_{\alpha}(z) = C_1(\alpha) + zC_2(\alpha)$$

Then, it is clear that

$$\boldsymbol{\xi}^{k+1} = \boldsymbol{\xi}^k - (P_{\alpha}^{-1}(z)K(z))\boldsymbol{\xi}^k, \ \forall k \ge 0.$$
(3.7)

By applying  $P_{\alpha}$  on both sides of (3.7), we derive

$$P_{\alpha}(z)\boldsymbol{\xi}^{k+1} = (P_{\alpha}(z) - K(z))\boldsymbol{\xi}^{k}.$$
(3.8)

Let  $\boldsymbol{\xi}^k = (\xi_r^k, \xi_{r+1}^k, \dots, \xi_{N_t}^k)^{\top}$ . Then a routine calculation yields

$$(C_1 + zC_2) \boldsymbol{\xi}^{k+1} = \alpha \begin{bmatrix} \sum_{j=1}^r (a_j + zb_j)(\xi_{N_t - j + 1}^{k+1} - \xi_{N_t - j + 1}^k) \\ \sum_{j=2}^r (a_j + zb_j)(\xi_{N_t - j + 2}^{k+1} - \xi_{N_t - j + 2}^k) \\ \vdots \\ (a_r + zb_r)(\xi_{N_t}^{k+1} - \xi_{N_t}^k) \\ \vdots \\ 0 \end{bmatrix}.$$

Since  $P_{\alpha}(z) - K(z) = C_1 + zC_2$ , (3.8) is equivalent to the following difference equations

$$\begin{cases} \sum_{j=0}^{r} a_{j} \xi_{n-j}^{k+1} = -z \sum_{j=0}^{r} b_{j} \xi_{n-j}^{k+1}, & n = r, r+1, \dots, N_{t} - r, \\ \xi_{r-n}^{k+1} = \alpha(\xi_{N_{t}+1-n}^{k+1} - \xi_{N_{t}+1-n}^{k}), & n = 1, \dots, r. \end{cases}$$
(3.9)

By Assumption 2, we have the stability estimate

$$\max_{0 \le n \le N_t} |\xi_n^{k+1}| \le c_0 \max_{0 \le n \le r-1} |\xi_n^{k+1}| \le c_0 \alpha \max_{0 \le n \le r-1} |\xi_{N_t-n}^{k+1} - \xi_{N_t-n}^k|,$$
(3.10)

with a generic constant  $c_0 > 1$ . This implies

$$\max_{0 \le n \le r-1} |\xi_{N_t - n}^{k+1}| \le c_0 \alpha \Big( \max_{0 \le n \le r-1} |\xi_{N_t - n}^{k+1}| + \max_{0 \le n \le r-1} |\xi_{N_t - n}^{k}| \Big).$$

Rearranging terms in the above inequality, we arrive at

$$\max_{0 \le n \le r-1} |\xi_{N_t - n}^{k+1}| \le \frac{c_0 \alpha}{1 - c_0 \alpha} \max_{0 \le n \le r-1} |\xi_{N_t - n}^k|.$$

Then, applying the stability estimate (3.10) again leads to

$$\max_{r \le n \le N_t - r} |\xi_n^{k+1}| \le c_0 \alpha \Big( \max_{0 \le n \le r-1} |\xi_{N_t - n}^{k+1}| + \max_{0 \le n \le r-1} |\xi_{N_t - n}^k| \Big) 
\le c_0 \alpha \Big( c_0 \max_{r \le n \le N_t - r} |\xi_n^{k+1}| + c_0 \max_{r \le n \le N_t - r} |\xi_n^k| \Big).$$
(3.11)

We arrange the inequality and obtain

$$\max_{r \le n \le N_t - r} |\xi_n^{k+1}| \le \frac{c_0^2 \alpha}{1 - c_0^2 \alpha} \max_{r \le n \le N_t - r} |\xi_n^k|.$$

Letting  $c = c_0^2$ , we derive

$$\max_{r \le n \le N_t} |\xi_n^{k+1}| \le \frac{c\alpha}{1 - c\alpha} \max_{r \le n \le N_t} |\xi_n^k|,$$

which implies that

$$\|\boldsymbol{\xi}^{k+1}\|_{\infty} \le \frac{c\alpha}{1-c\alpha} \|\boldsymbol{\xi}^{k}\|_{\infty}.$$
(3.12)

Now, substituting  $\boldsymbol{\zeta}^k = (I_t \otimes P) \boldsymbol{err}^k$  into (3.12) gives the desired result (3.6).  $\Box$ 

REMARK 3.1. For the multistep method, we note that the convergence factor involves a generic constant  $c \geq 1$ . It depends on the stability of the time stepping scheme, but it is always independent of the step size  $\Delta t$ , the time level n and the total number of steps  $N_t$ .

To complete this section, we test two four-step multistep methods, the four-step BDF method (BDF4) [35] and the modified Adams-Moulton method (AM) [3]:

(BDF4) 
$$y_{n+1} - \frac{48}{25}y_n + \frac{36}{25}y_{n-1} - \frac{16}{25}y_{n-2} + \frac{3}{25}y_{n-3} + \frac{12\Delta tA}{25}y_{n+1} = 0,$$
  
(AM)  $y_{n+1} - y_n + \Delta tA\left(\frac{2}{3}y_{n+1} + \frac{5}{12}y_{n-1} - \frac{1}{12}y_{n-3}\right) = 0.$  (3.13)

We apply the above multistep methods to the semi-discrete system of the advection-diffusion equation (2.7) with T = 8,  $\nu = 10^{-3}$  and  $\Delta t = \Delta x = \frac{1}{128}$ .

In Figure 3.1, we plot the stability regions of the above two multistep methods (shadow regions) together with  $\sigma(\Delta tA)$ , the spectrum of  $\Delta tA$ . We see that both methods are stable for the semi-discrete system.

In Figure 3.2, we plot the measured error  $\|\boldsymbol{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty}$ , together with the upper bound predicted by  $\frac{\alpha}{1-\alpha}$  (dotted line). We see that for AM scheme,  $\|\boldsymbol{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty}$  decay with a very similar rate close to  $\alpha/(1-\alpha)$ . While for BDF4, as we see in Figure 3.2 on the right, the decay rate is not uniform (see the error of the first 6 iterations in Figure 3.3). In particular, for  $\alpha = 0.1$  we have

$$\frac{\|(I_t \otimes P)\boldsymbol{err}^2\|_{\infty}}{\|(I_t \otimes P)\boldsymbol{err}^1\|_{\infty}} = 0.54, \ \frac{\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty}}{\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty}} \approx 0.11 \approx \frac{\alpha}{1-\alpha} \quad (\text{for } k \ge 1),$$

and for  $\alpha = 0.01$  we have

$$\frac{\|(I_t \otimes P)\boldsymbol{err}^2\|_{\infty}}{\|(I_t \otimes P)\boldsymbol{err}^1\|_{\infty}} = 0.05, \ \frac{\|(I_t \otimes P)\boldsymbol{err}^{k+1}\|_{\infty}}{\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty}} \approx 0.01 \approx \frac{\alpha}{1-\alpha} \quad (\text{for } k \ge 1)$$

This contrasts sharp with the one-step methods, where the convergence factor is alway bounded (from above) by  $\alpha/(1-\alpha)$  as suggested in Theorem 2.1. However, the numerical results suggest that the convergence rate will approaches  $\alpha/(1-\alpha)$  from the second iteration. This interesting phenomenon warrants further investigation.

Similar to Figure 2.2,  $\|\boldsymbol{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\boldsymbol{err}^k\|_{\infty}$  do not decay to the machine precision and at the last few iterations the error stagnants at certain level, due to the roundoff error arising in the block diagonalization of the preconditioner  $\mathcal{P}_{\alpha}$ .

4. Conclusion and discussion. We gave a uniformed proof for a class of PinT algorithm, which is based on formulating the time discretization into an all-at-once system  $\mathcal{K} \boldsymbol{u} = \boldsymbol{b}$  and then solving this system iteratively by using a block  $\alpha$ -circulant matrix  $\mathcal{P}_{\alpha}$  as the preconditioner. Existing work focus on examining the spectrum of the iterative matrix  $\mathcal{I} - \mathcal{P}_{\alpha}^{-1}\mathcal{K}$  and this leads to many case-by-case studies depending on the underlying time-integrator. In this short paper, we proved that the error of the algorithm decays with a rate only depending on the parameter  $\alpha$  for all *stable* one-step methods. We also considered the linear multistep methods, for which we proved similar result provided the method is stable as well. The error decaying rate is slightly different from that of the one-step methods and in practical computation we indeed observed such a difference. The proof given in this



FIG. 3.1. The stability region (shadow region) of the BDF4 method (left) and the AM method (right) in (3.13) and the spectrum  $\sigma(\Delta tA)$  with A being the matrix given by (2.8) and  $\Delta t = \Delta x = \frac{1}{128}$  and  $\nu = 10^{-3}$ .



FIG. 3.2. Applying the preconditioned iteration method (1.9) to the advection-diffusion equation (2.7), the measured error  $\|\mathbf{err}^k\|_{\infty}$  and  $\|(I_t \otimes P)\mathbf{err}^k\|_{\infty}$  together with the upper bound by  $\frac{\alpha}{1-\alpha}$ . Left: the 4-step Adams method in (3.13). Right: the 4-step BDF method in (3.13). Here, T = 8,  $\Delta t = \Delta x = \frac{1}{128}$  and  $\nu = 10^{-3}$ .



FIG. 3.3. For Figure 3.2 on the right, the local details for the first 6 iterations. Left:  $\alpha = 0.1$ . Right:  $\alpha = 0.01$ .

paper is completely different from existing work. It lies in representing the preconditioned iteration as a special difference equations with head-tail coupled condition and relays on the stability function only.

An important work that we do not concern is how to get the increment  $\Delta u^k = \mathcal{P}_{\alpha}^{-1} r^k$ in (1.9). For one-step methods, we can factorize  $\mathcal{P}_{\alpha}$  in (1.7) as

$$\mathcal{P}_{\alpha} = (V \otimes I_x)(I_t \otimes I_x + D \otimes \mathcal{R}(\Delta tA))(V^{-1} \otimes I_x),$$

where V and D are the eigenvector and eigenvalue matrices of the circulant matrix C, i.e.,  $C = VDV^{-1}$ . Then, we can compute  $\Delta u^k$  via the following steps

$$\begin{cases} \boldsymbol{p} = (V^{-1} \otimes I_x) \boldsymbol{r}^k, & \text{Step-(a)} \\ (I_x + \lambda_n \mathcal{R}(\Delta tA)) q_n = p_n, \ n = 1, 2, \dots, N_t, & \text{Step-(b)} \\ \Delta \boldsymbol{u}^k = (V \otimes I_x) \boldsymbol{q}, & \text{Step-(c)} \end{cases}$$
(4.1)

where  $\boldsymbol{p} = (p_1^{\top}, \ldots, p_{N_t}^{\top})^{\top}$  and  $\boldsymbol{q} = (q_1^{\top}, \ldots, q_{N_t}^{\top})^{\top}$ . For Step-(a) and Step-(c), we only need to do matrix-vector multiplications and the major computation burden is Step-(b). These are completely decoupled linear systems and can be solved in parallel, but there are still many concrete issues that need to be explored.

The first issue is about the general implicit RK method of stage s specified by the Butcher tableau

$$\begin{array}{c|c} c & \Theta \\ \hline & b^{\top} \end{array},$$

for which  $\mathcal{R}(\Delta tA) = I_x - b^\top \otimes (\Delta tA)(I_s \otimes I_x + \Theta \otimes (\Delta tA))^{-1}(\mathbf{1} \otimes I_x)$ , where  $I_s \in \mathbb{R}^{s \times s}$ is an identity matrix and  $\mathbf{1} = (1, 1, \dots, 1)^\top \in \mathbb{R}^s$ . Then how to efficiently solve the linear system in Step-(b) is an very important topic. Some preliminary experiences are as follows. For the two-stage RK methods, by an *elimination* operation for the stage variables it can be shown that such a linear system is equivalent to the following

$$(a_0 I_x + a_1 (\Delta tA) + a_2 (\Delta tA)^2) \tilde{p} = \tilde{q}, \qquad (4.2)$$

where  $a_0$ ,  $a_1$ ,  $a_2$  depend on  $\lambda_n$  and the RK method. The quadratic term  $(\Delta t A)^2$  will be a serious problem for both the memory storage and the computation cost. The approach for this issue is to notice that equation (4.2) can be represented as

where  $p_{\dagger}$  is an auxiliary variable and  $\mu \neq 0$  is a free parameter. Then, we can factorize  $W_s$  as  $W_s = V_s D_s V_s^{-1}$  and solve (4.3) by using the diagonalization technique again. In this way, we only need to solve two independent linear systems of the form

$$(\eta I_x + \Delta t A)v = b, \tag{4.4}$$

where  $\eta = a_0/D_s(1,1)$  (or  $\eta = a_0/D_s(2,2)$ ). (Here we assume  $D_s(1,1) \neq 0$ , since otherwise it is trivial to treat the linear system.) The above idea was used in [36] for the two-stage Lobatto IIIC method and in [37] for the two-stage SDIRK method. Similar to (4.2), for a general *s*-stage RK method we believe that the linear system in Step-(b) of (4.1) can be equivalently represented as

$$(a_0 I_x + a_1 (\Delta tA) + \dots + a_s (\Delta tA)^s) \tilde{p} = \tilde{q}.$$

However, it is unclear how to further represent it as (4.3) with a suitable matrix  $W_s$ . Such a  $W_s$  should be diagonalizable with small condition number of the eigenvector matrix  $V_s$ .

The second issue is about the computation of the complex-shift problem (4.4). It seems this is a very fundamental problem arising in various fields, such as the Laplace inversion approach, preconditioning technique for the Helmholtz equations and the diagonalization technique here. For ODEs with first-order temporal derivative, in most cases we found that the quantity  $\eta$  arising from the diagonalization technique has non-negative real part, i.e.,  $\Re(\eta) \ge 0$ , and thus it is not difficult to solve (4.4). This issue was carefully addressed for the multigrid method in [27, 31]. For second-order ODEs, it happens  $\Re(\eta) < 0$  for some  $\lambda_n$  in (4.1) and therefor (4.4) is a Helmholtz-like problem, a typical hard problem in numerics when  $|\Re(\eta)|$  is large and the ratio  $r := |\Im(\eta)/\Re(\eta)| \ll 1$ . Cocquet and Gander made an interesting analysis for the multigrid method in [4] and the main conclusion is that the multigrid method always converges if  $|\Im(\eta)| = \mathcal{O}(|\Re(\eta)|)$ . However, numerical results indicate that the multigrid method could converge arbitrarily slow and the convergence rate actually heavily depends on the ratio r. A local Fourier analysis will reveal how the convergence rate depends on r and this is one of our ongoing work. Besides the multigrid method, there are also other work concerning the complex-shift problem (4.4) and to name a few we mention the preconditioned GMRES method [14], the domain decomposition method [20] and the approach based on representing the real and imaginary parts separately, leading to a real symmetric linear system, which can be solved by an Uzawa-type method (see [34, Section 5] for this approach).

## REFERENCES

- D. BINI, G. LATOUCHE, AND B. MEINI, Numerical methods for structured Markov chains, Oxford University Press: New York, 2005.
- [2] F. CHEN, J. S. HESTHAVEN, AND X. ZHU, On the use of reduced basis methods to accelerate and stabilize the Parareal method, in: Quarteroni, A., Rozza, G. (eds.) Reduced Order Methods for Modeling and Computational Reduction, MS&A-Modeling, Simulation and Applications, vol. 9, pp. 187-214. Springer, Berlin, 2014.
- [3] W. CHEN, M. GUNZBURGER, D. SUN AND X. WANG, An efficient and long-time accurate third-order algorithm for the Stokes-Darcy system, Numer. Math., 134 (2016), pp. 857–879.
- [4] P.-H. COCQUET AND M. J. GANDER, How large a shift is needed in the shifted Helmholtz preconditioner for its effective inversion by multigrid?, SIAM J. Sci. Comput., 39 (2017), pp. A438–A478.
- [5] X. DAI AND Y. MADAY, Stable parareal in time method for first- and second-order hyperbolic systems, SIAM J. Sci. Comput., 35 (2013), pp. A52-A78.
- [6] A. EGHBAL, A. G. GERBER, AND E. AUBANEL, Acceleration of unsteady hydrodynamic simulations using the Parareal algorithm, J. Comput. Sci., 19 (2017), pp. 57-76.
- [7] M. EMMETT AND M. L. MINION, Toward an efficient parallel in time method for partial differential equations, Comm. App. Math. Comp. Sci., 7 (2012), pp. 105-132.
- [8] C. FARHAT, J. CORTIAL, C. DASTILLUNG, AND H. BAVESTRELLO, Time-parallel implicit integrators for the near-real-time prediction of linear structural dynamic responses, Int. J. Numer. Methods Eng., 67 (2006), pp. 697-724.
- [9] R. D. FALGOUT, S. FRIEDHOFF, T.KOLEV, S. P. MACLACHLAN, AND J. B. SCHRODER, Parallel time integration with multigrid, SIAM J. Sci. Comput., 36 (2014), pp. C635-C661.
- [10] M. J. Gander. 50 years of time parallel time integration. In Multiple shooting and time domain decomposition methods, volume 9 of Contrib. Math. Comput. Sci., pages 69–113. Springer, Cham, 2015.
- [11] M. J. GANDER AND M. PETCU, Analysis of a modified parareal algorithm for second-order ordinary differential equations, AIP Conference Proceedings, 936(1) (2007), pp. 233-236.
- [12] M. J. GANDER AND M. PETCU, Analysis of a Krylov subspace enhanced parareal algorithm for linear problems, ESAIM: Proceedings, 25 (2008), pp. 114-129.
- [13] M. J. GANDER AND S. VANDEWALLE, Analysis of the parareal time-parallel time-integration method, SIAM J. Sci. Comput., 29 (2007), pp. 556-578.
- [14] M. J. GANDER, I. GRAHAM, AND E. SPENCE, Applying GMRES to the Helmholtz equation with shifted

Laplacian preconditioning: what is the largest shift for which wavenumber- independent convergence is guaranteed?, Numer. Math., 131 (2015), pp. 567–614.

- [15] M. J. GANDER, L. HALPERN, J. RANNOU, AND J. RYAN, A direct time parallel solver by diagonalization for the wave equation, SIAM J. Sci. Comput., 41 (2019), pp. A220–A245.
- [16] M. J. GANDER AND S. L. WU, Convergence analysis of a periodic-like waveform relaxation method for initial-value problems via the diagonalization technique, Numer. Math., 143 (2019), pp. 489–527.
- [17] M. J. GANDER AND S. L. WU, A diagonalization-based parareal algorithm for dissipative and wave propagation problems, SIAM J. Numer. Anal., 58 (2020), pp. 2981–3009.
- [18] W. GAUTSCHI, Numerical analysis: An introduction, Second Edition, Birkhäuser Boston, Inc., Boston, MA, 2012.
- [19] A. GODDARD AND A. WATHEN, A note on parallel preconditioning for all-at-once evolutionary PDEs, ETNA-Electron. Trans. Numer. Anal., 51 (2019), pp. 135-150
- [20] I. GRAHAM, E. SPENCE, AND E. VAINIKKO, Domain decomposition preconditioning for high frequency Helmholtz problems with absorption, Math. Comp., 86 (2017), pp. 2089–2127.
- [21] X. M. GU, Y. L. ZHAO, X. L. ZHAO, AND B. CARPENTIERI, AND Y. Y. HUANG, A note on parallel preconditioning for the all-at-once solution of Riesz fractional diffusion equations, arXiv:2003.07020.
- [22] E. HAIRER AND G. WANNER, Solving ordinary differential equations II: stiff and differential-algebraic problems, Springer, Berlin, 2000.
- [23] J. LIU AND S. L. WU, A fast block α-circulant preconditioner for all-at-once systems from wave equations. SIAM J. Mat. Anal. Appl, 41 (2020), pp. 1912–1943.
- [24] J. L. LIONS, Y. MADAY, AND G. TURINICI, A "parareal" in time discretization of PDEs, C. R. Acad. Sci. Paris Ser. I Math., 332 (2001), pp. 661-668.
- [25] X.-L. LIN AND M. NG, An all-at-once preconditioner for evolutionary partial differential equations, arXiv:2002.01108.
- [26] X. LIN, M. NG, AND H. SUN, Efficient preconditioner of one-sided space fractional diffusion equation, BIT Numer. Math., 58 (2018), pp. 729-748.
- [27] S. P. MACLACHLAN AND C. W. OOSTERLEE, Algebraic multigrid solvers for complex-valuedmatrices, SIAM J. Sci. Comput., 30 (2008), pp. 1548-1571.
- [28] E. MCDONALD, J. PESTANA, AND A. WATHEN, Preconditioning and iterative solution of all-at-once systems for evolutionary partial differential equations, SIAM J. Sci. Comput., 40 (2018), pp. A1012–A1033.
- [29] Y. MADAY AND E. M. RONQUIST, Parallelization in time through tensor-product space?time solvers, Comptes Rendus Mathematique, 346 (2008), pp. 113?118.
- [30] H. NGUYEN AND R. TSAI, A stable parareal-like method for the second order wave equation, J. Comput. Phys., 405 (2020), pp. 109156.
- [31] Y. NOTAY, An aggregation-based algebraic multigrid method, Electron. Trans. Numer. Anal., 37 (2010), pp. 123-146.
- [32] D. RUPRECHT AND R. KRAUSE, Explicit parallel-in-time integration of a linear acoustic-advection system, Comput. Fluids, 59 (2012), pp. 72-83.
- [33] D. RUPRECHT, Wave propagation characteristics of Parareal, Comput. Visual Sci., 19 (2018), pp. 1-17.
- [34] M. STOLL, One-shot solution of a time-dependent time-periodic PDE-constrained optimization problem, IMA J. Numer. Anal., 34 (2014), pp. 1554–1577.
- [35] S. WU AND Z. ZHOU. Parallel-in-time high-order BDF schemes for diffusion and subdiffusion equations, Preprint, arXiv: 2007.13125.
- [36] S. L. WU AND T. ZHOU. Acceleration of the two-level MGRIT algorithm via the diagonalization technique, SIAM J. Sci. Comput., Vol. 41 (2019), pp. A3421–A3448.
- [37] S. L. WU AND T. ZHOU. Parallel implementation for the two-stage SDIRK methods via diagonalization, J. Comput. Phys., 428 (2021), pp. 110076.