

A priori and a posteriori error analysis of the Crouzeix-Raviart and Morley FEM with original and modified right-hand sides

Carsten Carstensen* and Neela Nataraj†

December 24, 2021

Abstract

This article on nonconforming schemes for m harmonic problems simultaneously treats the Crouzeix-Raviart ($m = 1$) and the Morley finite elements ($m = 2$) for the original and for modified right-hand side F in the dual space $V^* := H^{-m}(\Omega)$ to the energy space $V := H_0^m(\Omega)$. The smoother $J : V_{\text{nc}} \rightarrow V$ in this paper is a companion operator, that is a linear and bounded right-inverse to the nonconforming interpolation operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$, and modifies the discrete right-hand side $F_h := F \circ J \in V_{\text{nc}}^*$. The best-approximation property of the modified scheme from Veerer et al. (2018) is recovered and complemented with an analysis of the convergence rates in weaker Sobolev norms. Examples with oscillating data show that the original method may fail to enjoy the best-approximation property but can also be better than the modified scheme. The a posteriori analysis of this paper concerns data oscillations of various types in a class of right-hand sides $F \in V^*$. The reliable error estimates involve explicit constants and can be recommended for explicit error control of the piecewise energy norm. The efficiency follows solely up to data oscillations and examples illustrate this can be problematic.

Keywords: nonconforming schemes, Crouzeix-Raviart, Morley, biharmonic problem, medius error analysis, best-approximation, a priori and a posteriori analysis

AMS Classification: 65N30, 65N12, 65N50

1 Introduction

The beginning of the nonconforming finite element methodology, its motivating examples in fluid mechanics, and its connections with mixed formulations are outlined in [7] together with the relevant literature that (partly) led to the Strang lemmas (also named after Strang-Fix and Strang-Berger etc.) in finite element textbooks. This article aims at a simple a priori

*Department of Mathematics, Humboldt-Universität zu Berlin, 10099 Berlin, Germany. Distinguished Visiting Professor, Department of Mathematics, Indian institute of Technology Bombay, Powai, Mumbai-400076, India. cc@math.hu-berlin.de

†Department of Mathematics, Indian Institute of Technology Bombay, Powai, Mumbai 400076, India. neela@math.iitb.ac.in

and a posteriori error analysis for m harmonic problems, that involve the energy space $V := H_0^m(\Omega)$ endowed with the energy scalar product $a : V \times V \rightarrow \mathbb{R}$. The emphasis is on the lowest-order nonconforming schemes with the piecewise polynomial, nonconforming discrete space $V_{\text{nc}} \subset P_m(\mathcal{T})$, that simultaneously describes the Crouzeix-Raviart (CR) for $m = 1$ and Morley finite element methods (FEM) for $m = 2$. Those nonconforming FEMs allow for an interpolation operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ with the best-approximation property and interpolation error estimates with explicit (and small) constants (called κ_m throughout) from [16, 17] for $m = 1$ and [12] for $m = 2$. The word Strang lemma is no longer used in this paper and a *companion operator* $J : V_{\text{nc}} \rightarrow V$ circumvents the medius analysis from [38].




Figure 1: Crouzeix-Raviart FEM (left) and Morley FEM (right)

A (*conforming*) *companion operator* $J : V_{\text{nc}} \rightarrow V$ is defined in this paper as a right-inverse of the interpolation operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ with additional benefits such as the L^2 orthogonality $v_{\text{nc}} - Jv_{\text{nc}} \perp P_m(\mathcal{T})$ for all $v_{\text{nc}} \in V_{\text{nc}}$. The extra L^2 orthogonality is *not* needed for the sole best-approximation [51–53], but it enables a duality argument for error estimates in weaker Sobolev norms [15].

The paper departs with a revisit of the medius analysis [38], the best-approximation results from [51–53] and recovers the same best-approximation constant C_{qo} for a modified right-hand side F_h in Section 3. It thereafter contributes a comprehensive analysis of convergence rates in Sobolev norms weaker and stronger than the energy norm in V in Section 4.

The precise convergence rate $\min\{2\sigma, m + \sigma - s\}$ is guaranteed for quasi-uniform meshes for the error $u - Ju_{\text{nc}}$ of the post-processed approximation Ju_{nc} in the norm of $H^s(\Omega)$ for $-m \leq s < m + 1/2$ with the positive index $\sigma \leq 1$ in terms of elliptic regularity. This holds under the generic assumption that the convergence rate in the energy norm is exactly $\sigma < m$ and so excludes untypical smooth solutions. The only former result is the seminal contribution [39] with a proof that the convergence rate in the L^2 error in nonconforming schemes is no better than 2σ not only for one model scenario but for many other nonconforming schemes.

The paper will focus on shape-regular triangulations of the planar bounded polygonal Lipschitz domain $\Omega \subset \mathbb{R}^2$ into triangles for simplicity. The abstract analysis holds verbatim for any space dimensions once the companion operator J is designed. This started in 2D for $m = 1$ with [15] for the Crouzeix-Raviart finite element method (CRFEM) and is almost immediately generalized to any space dimension. For $m = 2$ and the Morley FEM it started in the PhD thesis [34] and details can be found in [35, 53] in 2D, while the companion operator J for generalization of the Morley FEM from [44] is designed in

[25] in 3D. The most refined analysis in 2D with explicit constants and the connection to a posteriori error estimates are summarized in [18, 26]. The paper will focus on lowest-order schemes because, first, the interpolation operators are less clear for higher-order CRFEM in 2D and 3D [27], hence, second, conforming companions are unknown. In fact, there are two aspects that make the nonconforming FEM attractive in comparison to conforming FEM: The interpolation operator is local (simplex-oriented) and defined for any function in the energy space. Those advantages are possible for simplicial meshes in any space dimension, but *not* available for all higher polynomial degrees.

The modified nonconforming FEM specifies the discrete right-hand side $F_h := F \circ J \in V_{\text{nc}}^*$ with the companion operator $J : V_{\text{nc}} \rightarrow V$ as a smoother. Thereby the modified version allows for general data $F \in V^*$. Modifications of the right-hand side F_h of this type had been introduced in [2, 8], before [51–53] systematically studied the resulting best-approximation property and established the name smoother. The modified FEM is a Petrov-Galerkin scheme discussed in Section 3 and hence quasi-optimally convergent (once stable). If the source term $F \in V^* \setminus L^2(\Omega)$ is *not* a Lebesgue function, the application of the original discrete problem is less obvious. This paper studies a class of right-hand sides F in V^* that includes point forces for $m = 2$ in the model example to define F_h (without any smoother) adapted to the data. This provokes a comparison of the nonconforming FEM with the two different discrete right-hand sides with advantages for both sides. The model example is rich enough to allow for a *counterexample for the best-approximation* of the original method. The latter holds up to the data oscillations only; cf. [24] for $m = 1$ and $F \in L^2(\Omega)$. Conversely, another example illustrates that the modified nonconforming method (with the discrete right-hand side $F \circ J$) can be *worse* than the original discretization.

The a posteriori error control for the (original) nonconforming FEM and $F \in L^2(\Omega)$ started with [31] and continued with [9–11, 14, 19, 20, 50] for $m = 1$ and [5, 13, 40, 41] for $m = 2$. The initial work was based on a Helmholtz decomposition of the piecewise energy norm of the nonconforming error. The Helmholtz decompositions become Hodge decompositions for multiply-connected domains and are not all known for 3D. Their discrete counterparts from [3] for $m = 1$ and [14] for $m = 2$ are not known for all domains and space dimensions. That motivates the use of other techniques and this paper suggests a paradigm shift: Utilize the companion operator J for a split in the conforming error $u - Ju_{\text{nc}}$ and in the nonconforming error $u_{\text{nc}} - Ju_{\text{nc}}$. This approach leads to a posteriori error estimates even if a Helmholtz decomposition is unknown and without a piecewise integration by parts. As a consequence, the error estimators do *not* contain weighted jumps of the normal components of derivatives of the discrete solution u_{nc} across interior edges. On the other hand, the efficiency requires some other additional benefit of the companion operator, which actually motivated its first design in a posteriori error control. Rate-optimal adaptive nonconforming FEM are analyzed in [4, 14, 15, 22, 25, 47] and the references therein. Amongst all second-order schemes, the Morley FEM appears to be the most simple and method of choice for fourth-order problems [21]. Hence the generalization of the methodology in this article for $m \geq 3$ with nonconforming finite elements from [55] appears appealing.

Outline of the structure. The remaining parts of this paper are organised as follows. In Section 2 nonconforming schemes, interpolation and companion operators, and examples

are presented. The best-approximation results available in literature are reviewed and an abstract best-approximation result is established for data $F \in H^{-m}(\Omega)$ in Section 3. Section 4 presents lower-order error estimates and precise convergence rates under further hypotheses. Section 5 discusses examples of data in V^* , an alternate choice of the right-hand side functional that leads to a modified scheme and a comparison of the original and the modified scheme. This is followed by an a posteriori analysis in Section 6.

General notation. Standard notation of Lebesgue and Sobolev spaces, their norms, and L^2 scalar products applies throughout the paper such as the abbreviation $\|\bullet\|$ for $\|\bullet\|_{L^2(\Omega)}$. For a shape-regular triangulation \mathcal{T} of Ω into triangles, the product space $\prod_{T \in \mathcal{T}} H^s(T)$ denoted as $H^s(\mathcal{T})$ is defined as $\{v_{\text{pw}} \in L^2(\Omega) : \forall T \in \mathcal{T}, v_{\text{pw}}|_T \in H^s(T)\}$ and is equipped with the Euclid norm of those contributions $\|\bullet\|_{H^s(T)}$ for all $T \in \mathcal{T}$. The triple norm $\|\bullet\| := |\bullet|_{H^m(\Omega)}$ is the energy norm and $\|\bullet\|_{\text{pw}} := |\bullet|_{H^m(\mathcal{T})} := \|D_{\text{pw}}^m \bullet\|$ is its piecewise version with the piecewise partial derivatives D_{pw}^m of order $m \in \mathbb{N}$. For non-integer real $t = \ell + s$ with $\ell \in \mathbb{N}_0$ and $0 < s < 1$, the 2D Sobolev-Slobodeckii semi-norm [37] of $f \in H^t(\Omega)$ reads

$$|f|_{H^t(\Omega)} := \left(\sum_{|\beta|=\ell} \int_{\Omega} \int_{\Omega} \frac{|\partial^{\beta} f(x) - \partial^{\beta} f(y)|^2}{|x-y|^{2+2s}} dx dy \right)^{1/2}. \quad (1.1)$$

The vector space of piecewise polynomials of at most degree $k \in \mathbb{N}$ is denoted by $P_k(\mathcal{T})$, $S^k(\mathcal{T}) := P_k(\mathcal{T}) \cap C^0(\Omega) \subset H^1(\Omega)$, and $S_0^k(\mathcal{T}) := P_k(\mathcal{T}) \cap C^0(\Omega) \subset H_0^1(\Omega)$ includes the homogeneous boundary conditions. For a Lebesgue-measurable set $\omega \subseteq \mathbb{R}^n$ and a Lebesgue-measurable function $v : \omega \rightarrow X$ with values in a finite-dimensional real Banach space X , the integral with respect to the d -dimensional Lebesgue measure is denoted by $\int_{\omega} v dx := 1/|\omega| \int_{\omega} v dx$. The integral over a $(n-1)$ -dimensional hypersurface Γ with respect to the $(n-1)$ -dimensional Hausdorff measure reads as $\int_{\Gamma} v ds$. The notation $A \lesssim B$ abbreviates $A \leq CB$ for some positive generic constant C , which depends on the solution u ; $A \approx B$ abbreviates $A \lesssim B \lesssim A$. The set of all 2×2 real symmetric matrices is denoted by \mathbb{S} . The symmetric part of $A \in \mathbb{R}^{2 \times 2}$ is denoted by $\text{sym}(A)$. That is, $\text{sym}(A) = 1/2(A + A^T)$. For any differentiable scalar function v resp. vector field $\Phi = (\phi_1, \phi_2)^T$,

$$\text{Curl } v = \begin{pmatrix} -\frac{\partial v}{\partial x_2} \\ \frac{\partial v}{\partial x_1} \end{pmatrix} \quad \text{resp.} \quad \text{Curl } \Phi := \text{Curl} \begin{pmatrix} \phi_1 \\ \phi_2 \end{pmatrix} = \begin{pmatrix} -\frac{\partial \phi_1}{\partial x_2} & \frac{\partial \phi_1}{\partial x_1} \\ -\frac{\partial \phi_2}{\partial x_2} & \frac{\partial \phi_2}{\partial x_1} \end{pmatrix}.$$

The symbol $D^1 v := \nabla v$ (resp. $\nabla_{\text{pw}} v$) denotes the gradient (resp. piecewise gradient) of a scalar function v and $\sigma := D^2 u$ (resp. $D_{\text{pw}}^2 u$) denotes the Hessian (piecewise Hessian) of v with the four entries σ_{11} , σ_{12} , σ_{21} , and σ_{22} in the form of a 2×2 matrix.

2 Nonconforming schemes

2.1 Continuous model problem

Suppose $u \in V := H_0^m(\Omega)$ solves the m -harmonic equation $(-1)^m \Delta^m u = F$ for a given right-hand side $F \in V^* \equiv H^{-m}(\Omega)$ in a planar bounded Lipschitz domain $\Omega \subset \mathbb{R}^2$ with

polygonal boundary $\partial\Omega$, $m \in \mathbb{N}$. The weak form of this equation reads

$$a(u, v) = F(v) \quad \text{for all } v \in V \quad (2.1)$$

with $a(v, w) := \int_{\Omega} D^m v : D^m w \, dx$ for all $v, w \in V$. The examples in this paper are merely for $m = 1, 2$, but the abstract framework can be written for any $m \in \mathbb{N}$. It is well known that (2.1) has a unique solution u and elliptic regularity [1, 6, 36, 46] holds in the sense that $F \in H^{-s}(\Omega)$ implies $u \in H^{2m-s}(\Omega)$ for all $m - \sigma \leq s \leq m$ with the positive index of elliptic regularity σ . The latter is well established for $m = 1, 2$ and an overall hypothesis in this paper establishes for other m . The lowest-order nonconforming finite element schemes suggest linear convergence rates in the energy norm only $\|\bullet\| := a(\bullet, \bullet)^{1/2} \approx \|\bullet\|_V$ and so it is sufficient to assume throughout this paper that the quantity $\sigma \leq 1$ is the minimum of the index of elliptic regularity and one, whence $0 < \sigma \leq 1$ depends exclusively on Ω and m . The index of elliptic regularity $\sigma = 1$ follows for a convex polygonal domain Ω while $1/2 < \sigma < 1$ characterizes a nonconvex polygonal domain Ω [1, 6, 36, 46].

2.2 Nonconforming discretisation

Suppose that $V_{\text{nc}} \subset P_m(\mathcal{T})$ is a nonconforming FE space based on a shape-regular triangulation \mathcal{T} of Ω into triangles. Let the semi-scalar product a_{pw} be defined by the piecewise differential operator D_{pw}^m (the gradient for $m = 1$ and the Hessian for $m = 2$)

$$a_{\text{pw}}(v_{\text{pw}}, w_{\text{pw}}) := \sum_{T \in \mathcal{T}} \int_T D^m v_{\text{pw}} : D^m w_{\text{pw}} \, dx \quad \text{for all } v_{\text{pw}}, w_{\text{pw}} \in H^m(\mathcal{T}),$$

that induces a piecewise H^m seminorm $\|\bullet\|_{\text{pw}} = a_{\text{pw}}(\bullet, \bullet)^{1/2}$ that is also a norm in V_{nc} . Then $(V_{\text{nc}}, a_{\text{pw}})$ is a (finite-dimensional) Hilbert space so that, given any $F_h \in V_{\text{nc}}^*$, there exists a unique discrete solution $u_{\text{nc}} \in V_{\text{nc}}$ to

$$a_{\text{pw}}(u_{\text{nc}}, v_{\text{nc}}) = F_h(v_{\text{nc}}) \quad \text{for all } v_{\text{nc}} \in V_{\text{nc}}. \quad (2.2)$$

The nonconforming FEM is accompanied with an interpolation operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ and a best-approximation property in $P_m(\mathcal{T})$, i.e.,

$$a_{\text{pw}}(v - I_{\text{nc}}v, w_m) = 0 \quad \text{for all } v \in V \text{ and all } w_m \in P_m(\mathcal{T}). \quad (2.3)$$

Suppose that there exists a universal constant $\kappa_m > 0$ with

$$\|h_{\mathcal{T}}^{-m}(v - I_{\text{nc}}v)\| \leq \kappa_m \|v - I_{\text{nc}}v\|_{\text{pw}} \quad \text{for all } v \in V \quad (2.4)$$

for the L^2 norm $\|\bullet\| := \|\bullet\|_{L^2(\Omega)}$ and the piecewise H^m seminorm $\|\bullet\|_{\text{pw}}$ for the piecewise constant mesh-size $h_{\mathcal{T}} \in P_0(\mathcal{T})$ with $h_{\mathcal{T}}|_T := h_T = \text{diam}(T)$.

Remark 2.1 (Pythagoras). The semi-inner product a_{pw} in $H^m(\mathcal{T}) := \{v_{\text{pw}} \in L^2(\Omega) : \forall T \in \mathcal{T}, v_{\text{pw}}|_T \in H^m(T)\}$ allows for the concept of orthogonality: Two piecewise Sobolev functions $v_{\text{pw}}, w_{\text{pw}} \in H^m(\mathcal{T})$ are orthogonal if $a_{\text{pw}}(v_{\text{pw}}, w_{\text{pw}}) = 0$ holds and then the Pythagoras theorem

$$\|v_{\text{pw}} + w_{\text{pw}}\|_{\text{pw}}^2 = \|v_{\text{pw}}\|_{\text{pw}}^2 + \|w_{\text{pw}}\|_{\text{pw}}^2$$

follows. An example is the orthogonality of $v - I_{\text{nc}}v$ and w_{nc} from (2.3) that guarantees in particular

$$\|v - w_{\text{nc}}\|_{\text{pw}}^2 = \|v - I_{\text{nc}}v\|_{\text{pw}}^2 + \|w_{\text{nc}} - I_{\text{nc}}v\|_{\text{pw}}^2 \quad (2.5)$$

for all $v \in V$ and $w_{\text{nc}} \in V_{\text{nc}}$.

Remark 2.2 ($V + V_{\text{nc}}$ is a Hilbert space with inner product a_{pw}). Recall the assumption that $\|\bullet\|_{\text{pw}}$ is a norm in V_{nc} and that some $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ satisfies (2.3)-(2.4). Then $\|\bullet\|_{\text{pw}}$ is a norm in $V + V_{\text{nc}}$. Since $\|\bullet\|_{\text{pw}}$ is always a seminorm, the definiteness has to be clarified: Suppose that $\|v + v_{\text{nc}}\|_{\text{pw}} = 0$ for some $v \in V$ and $v_{\text{nc}} \in V_{\text{nc}}$. This and $w_{\text{nc}} := -v_{\text{nc}}$ imply in (2.5) that $\|v - I_{\text{nc}}v\|_{\text{pw}} = 0 = \|v_{\text{nc}} + I_{\text{nc}}v\|_{\text{pw}}$. Since $\|\bullet\|_{\text{pw}}$ is a norm in V_{nc} , it follows $v_{\text{nc}} = -I_{\text{nc}}v$. This shows in (2.4) that $v + v_{\text{nc}} = 0$ a.e. as claimed.

Consequently, $V + V_{\text{nc}}$ is a pre-Hilbert space with inner product a_{pw} with a complete subspace V and a finite-dimensional subspace V_{nc} . The completeness of $V + V_{\text{nc}}$ follows from this by standard arguments with the closed subspace $W := V \cap V_{\text{nc}}$ and its orthogonal complement W^\perp in $V + V_{\text{nc}}$: Let Q and R denote the orthogonal projection onto W and $V \cap W^\perp$, respectively. Given any $v + v_{\text{nc}} \in W^\perp$ for $v \in V$ and $v_{\text{nc}} \in V_{\text{nc}}$, the consequence $v + v_{\text{nc}} = (1 - Q)v + (1 - Q)v_{\text{nc}}$ leads to $W^\perp = (V \cap W^\perp) \oplus (V_{\text{nc}} \cap W^\perp)$. The orthogonal split $v + v_{\text{nc}} = (1 - Q)v + R(1 - Q)v_{\text{nc}} + (1 - R)(1 - Q)v_{\text{nc}}$ leads to $W^\perp = (V \cap W^\perp) \oplus (1 - R)(V_{\text{nc}} \cap W^\perp)$. The space $V + V_{\text{nc}}$ is therefore the orthogonal sum of the complete subspaces W , $V \cap W^\perp$, and $(1 - R)(V_{\text{nc}} \cap W^\perp)$ and so complete.

Remark 2.3 (uniqueness and extension of $I_{\text{nc}} : V + V_{\text{nc}} \rightarrow V_{\text{nc}}$). Since $(V + V_{\text{nc}}, a_{\text{pw}})$ is a Hilbert space, (2.3) shows that $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ maps any v to its (unique) best-approximation $I_{\text{nc}}v$ in V_{nc} . In other words, the operator I_{nc} in (2.3)-(2.4) is unique and equal to the restriction to V of the best-approximation $I_{\text{nc}} : V + V_{\text{nc}} \rightarrow V_{\text{nc}}$ that is the orthogonal projection onto V_{nc} in the Hilbert space $(V + V_{\text{nc}}, a_{\text{pw}})$. Throughout this paper, the symbol I_{nc} denotes this orthogonal projection in the extended space $I_{\text{nc}} : V + V_{\text{nc}} \rightarrow V_{\text{nc}}$ as well as its restriction to arguments in V . Notice that $I_{\text{nc}} = \text{id}$ in V_{nc} and $I_{\text{nc}}(v + w_{\text{nc}}) = I_{\text{nc}}v + w_{\text{nc}}$ for all $v \in V$ and $w_{\text{nc}} \in V_{\text{nc}}$.

2.3 Examples for $m = 1$ and $m = 2$

For $m = 1$ (resp. $m = 2$), the nonconforming discretization models the Crouzeix-Raviart (resp. Morley) FEM. Besides the best approximation property (2.3), these examples allow for the L^2 orthogonality

$$D^m(v - I_{\text{nc}}v) \perp P_0(\mathcal{T}; \mathbb{R}^{2m}). \quad (2.6)$$

2.3.1 Crouzeix-Raviart FEM for the harmonic equation

The set of vertices \mathcal{V} (resp. edges \mathcal{E}) in the shape-regular triangulation \mathcal{T} of the bounded polygonal Lipschitz domain Ω into triangles is divided into the interior vertices $\mathcal{V}(\Omega)$ (resp. edges $\mathcal{E}(\Omega)$) and the vertices $\mathcal{V}(\partial\Omega)$ (resp. edges $\mathcal{E}(\partial\Omega)$) on the boundary $\partial\Omega$. Let Π_k denote the L^2 projection onto the space of piecewise polynomials $P_k(\mathcal{T})$ of total degree at most $k \in \mathbb{N}_0$. The mesh-size $h_{\mathcal{T}} \in P_0(\mathcal{T})$ is defined by $h_{\mathcal{T}}|_T := h_T := |T|^{1/2} \approx \text{diam}(T)$ in any triangle $T \in \mathcal{T}$ of area $|T|$; $|E|$ denotes the length of an edge $E \in \mathcal{E}$.

The CR finite element space $V_{\text{nc}} := \text{CR}_0^1(\mathcal{T})$ for $m = 1$ reads [29]

$$\begin{aligned} \text{CR}^1(\mathcal{T}) &:= \{v_{\text{CR}} \in P_1(\mathcal{T}) \mid v_{\text{CR}} \text{ is continuous at all midpoints of interior edges of } \mathcal{T}\}, \\ \text{CR}_0^1(\mathcal{T}) &:= \{v_{\text{CR}} \in \text{CR}^1(\mathcal{T}) \mid v_{\text{CR}} \text{ vanishes at all midpoints of boundary edges of } \mathcal{T}\}. \end{aligned}$$

The CR interpolation operator $I_{\text{nc}} := I_{\text{CR}} : V \rightarrow \text{CR}_0^1(\mathcal{T})$ is defined by

$$(I_{\text{CR}}v)(\text{mid}(E)) = \int_E v \, ds \text{ for all } E \in \mathcal{E} \text{ for all } v \in V.$$

It satisfies (2.3)-(2.5) and (2.6) with $\kappa_1 = \sqrt{j_{1,1}^{-2} + 1/48}$ for the first positive root $j_{1,1}$ of the Bessel function of the first kind [12, 16], that does not even depend on the shape of the triangles.

For a simply-connected domain Ω , the *discrete Helmholtz decomposition* [3] shows

$$P_0(\mathcal{T}; \mathbb{R}^2) = \nabla_{\text{pw}} \text{CR}_0^1(\mathcal{T}) \oplus \text{Curl}(S^1(\mathcal{T})/\mathbb{R})$$

and, as $h \rightarrow 0$, leads to the Helmholtz decomposition

$$L^2(\Omega; \mathbb{R}^2) = \nabla H_0^1(\Omega) \oplus \text{Curl}(H^1(\Omega)/\mathbb{R}).$$

2.3.2 Morley FEM for the biharmonic equation

The nonconforming Morley finite element space $V_{\text{nc}} = \text{M}(\mathcal{T})$ for $m = 2$ reads [45]

$$\begin{aligned} \text{M}'(\mathcal{T}) &:= \left\{ v_{\text{M}} \in P_2(\mathcal{T}) \left| \begin{array}{l} v_{\text{M}} \text{ is continuous at the vertices and its normal derivatives} \\ v_E \cdot \nabla_{\text{pw}} v_{\text{M}} \text{ are continuous at the midpoints of interior edges} \end{array} \right. \right\}, \\ \text{M}(\mathcal{T}) &:= \left\{ v_{\text{M}} \in \text{M}'(\mathcal{T}) \left| \begin{array}{l} v_{\text{M}} \text{ vanishes at the vertices of } \partial\Omega \text{ and its normal derivatives} \\ v_E \cdot \nabla v_{\text{M}} \text{ vanish at the midpoints of boundary edges} \end{array} \right. \right\}. \end{aligned}$$

For any $v \in V$, the Morley interpolation operator $I_{\text{nc}}v := I_{\text{M}}v \in \text{M}(\mathcal{T})$ is defined by the degrees of freedom

$$(I_{\text{M}}v)(z) = v(z) \text{ for any } z \in \mathcal{V}(\Omega) \text{ and } \int_E \frac{\partial I_{\text{M}}v}{\partial \nu_E} \, ds = \int_E \frac{\partial v}{\partial \nu_E} \, ds \text{ for any } E \in \mathcal{E}.$$

The Morley interpolation operator $I_{\text{M}} : V \rightarrow \text{M}(\mathcal{T})$ satisfies (2.3)-(2.5) and (2.6) [14, 41] with $\kappa_2 = 0.25745784465$ [12] that does not even depend on the shape of the triangles.

For a simply-connected domain Ω , the *discrete Helmholtz decomposition* [14] reads

$$P_0(\mathcal{T}; \mathbb{S}) = D_{\text{pw}}^2 \text{M}(\mathcal{T}) \oplus \text{sym Curl}(S^1(\mathcal{T})^2/\mathbb{R}^2)$$

and, as $h \rightarrow 0$, leads to the Helmholtz decomposition

$$L^2(\Omega; \mathbb{R}^2) = D^2 H_0^2(\Omega) \oplus \text{sym Curl}(H^1(\Omega; \mathbb{R}^2)/\mathbb{R}^2).$$

2.4 Companion operator

The interpolation operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ allows for a right-inverse J with benefits [26]. Recall the conforming finite element subspace V_c of V and suppose that a linear operator $J : V_{\text{nc}} \rightarrow V$ and a constant Λ_J exist with

$$I_{\text{nc}}J = \text{id in } V_{\text{nc}}, \text{ i.e., } I_{\text{nc}}Jv_{\text{nc}} = v_{\text{nc}}, \quad (2.7)$$

$$\|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J \|v_{\text{nc}} - v\|_{\text{pw}}, \quad (2.8)$$

$$v_{\text{nc}} - Jv_{\text{nc}} \perp P_m(\mathcal{T}) \quad \text{in } L^2(\Omega) \quad (2.9)$$

for all $v_{\text{nc}} \in V_{\text{nc}}$ and for all $v \in V$.

The companion operator J in this paper (a) allows for an elegant proof (in Subsection 3.2), (b) provides an appropriate right-hand side $F_h := F(J\bullet) \in V_{\text{nc}}^*$ in (2.1) for general $F \in V^*$ with best-approximation (in Subsection 3.3), and (c) controls the inconsistency term $\min_{v \in V} \|u_{\text{nc}} - v\|_{\text{pw}}$ in the a posteriori error control (in Section 6).

Example 2.1 (companion operators). For any $v_{\text{CR}} \in CR_0^1(\mathcal{T})$, there exists $Jv_{\text{CR}} \in (S_0^1(\mathcal{T}) + P_4(\mathcal{T})) \cap V$ [15] that satisfies (2.7)-(2.9). For any $v_{\text{M}} \in M(\mathcal{T})$, there exists a $Jv_{\text{M}} \in (HCT(\mathcal{T}) + P_8(\mathcal{T})) \cap V$ (with the HCT finite element space $HCT(\mathcal{T})$) that satisfies (2.7)-(2.9) [26, 34, 35].

Remark 2.4 (extra orthogonality in (2.9)). The L^2 orthogonality in (2.9) allows a direct proof of Theorem 3.1 that circumvents the a posteriori error analysis of the consistency term as part of the medius analysis [38]. It also allows control over dual norm estimates of the form $\|v_{\text{M}} - Jv_{\text{M}}\|_{H^{-s}(\Omega)} \lesssim \|h_{\mathcal{T}}^s(v_{\text{M}} - Jv_{\text{M}})\|$ for $0 \leq s \leq 2$. This is critical e.g. in eigenvalue analysis or for problems with low-order terms [22, 26]. The proof of the best-approximation of Theorem 3.2 for the modified scheme (3.9), however, does not require the L^2 orthogonality in (2.9).

3 Best-Approximation

The first subsection is a reformulation of the medius analysis in the seminal work [38].

3.1 First error analysis for $F \in L^2(\Omega)$

Suppose $F \in L^2(\Omega)$ and let u (resp. u_{nc}) denote the exact (resp. discrete) solution to (2.1) (resp. (2.2)). The analysis is first illustrated for data $F \equiv f \in L^2(\Omega)$ such that $F_h(v_{\text{nc}}) = F(v_{\text{nc}}) := \int_{\Omega} f v_{\text{nc}} \, dx$ for all $v_{\text{nc}} \in V_{\text{nc}}$ in (2.2).

The error $e := u - u_{\text{nc}} \in V + V_{\text{nc}}$ and its interpolation $e_{\text{nc}} := I_{\text{nc}}u - u_{\text{nc}} \in V_{\text{nc}}$ are naively split into

$$\|e\|_{\text{pw}}^2 = a(u, u - Ju_{\text{nc}}) - a_{\text{pw}}(u_{\text{nc}}, e_{\text{nc}}) + a_{\text{pw}}(u - I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}}). \quad (3.1)$$

(The identity (3.1) follows from elementary algebra in the Hilbert space $V + V_{\text{nc}}$ and $a_{\text{pw}}(u_{\text{nc}}, u - I_{\text{nc}}u) = 0 = a_{\text{pw}}(I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}})$ from (2.3) and (2.7).) The first and second term on the right-hand side of (3.1) allow for (2.1) and (2.2), respectively, and their sum

is equal to $F(u - Ju_{\text{nc}} - e_{\text{nc}})$ (recall $F_h \equiv F$). This and the last term in (3.1) allow Cauchy inequalities. This, (2.7), and (2.8) lead in (3.1) to

$$\|e\|_{\text{pw}}^2 \leq \|h_{\mathcal{T}}^m f\| \|h_{\mathcal{T}}^{-m}(1 - I_{\text{nc}})(u - Ju_{\text{nc}})\| + \Lambda_J \|u - I_{\text{nc}}u\|_{\text{pw}} \|e\|_{\text{pw}}. \quad (3.2)$$

The interpolation estimate in (2.4) is followed by a triangle inequality, (2.8) with $v = u \in V$ and (2.5) to derive

$$\|h_{\mathcal{T}}^{-m}(1 - I_{\text{nc}})(u - Ju_{\text{nc}})\| \leq \kappa_m \|u - I_{\text{nc}}u\|_{\text{pw}} + \kappa_m \Lambda_J \|e\|_{\text{pw}} \leq \kappa_m (1 + \Lambda_J) \|e\|_{\text{pw}}.$$

The combination with (3.2) shows best-approximation up to a data approximation term

$$\|u - u_{\text{nc}}\|_{\text{pw}} \leq \max\{\Lambda_J, \kappa_m(1 + \Lambda_J)\} (\|u - I_{\text{nc}}u\|_{\text{pw}} + \|h_{\mathcal{T}}^m f\|). \quad (3.3)$$

Some local efficiency is enfolded to verify the a posteriori error estimate $\|h_{\mathcal{T}}^m f\| \lesssim \text{osc}_m(f, \mathcal{T}) + \|u - I_{\text{nc}}u\|_{H^m(\mathcal{T})}$ in the examples of [38]. This is a standard argument in residual-based a posteriori error control and links (3.3) to the alternate version in (3.4).

3.2 Alternative proof for $F \in L^2(\Omega)$

The medius analysis in [38] does *not* employ (2.9) to derive the slightly improved version

$$\|u - u_{\text{nc}}\|_{\text{pw}} \leq C_1 (\|u - I_{\text{nc}}u\|_{\text{pw}} + \text{osc}_m(f, \mathcal{T})) \quad (3.4)$$

in terms of the data-oscillations

$$\text{osc}_m(f, \mathcal{T}) := \|h_{\mathcal{T}}^m(f - \Pi_m f)\| \quad (3.5)$$

for $f \in L^2(\Omega)$ and its L^2 projection $\Pi_m f$ onto $P_m(\mathcal{T})$. The beneficial property (2.9) allows for a direct proof and this is detailed next.

The key observation is that the split (3.1) leads to $F((1 - I_{\text{nc}})(u - Ju_{\text{nc}}))$ and the contribution $F(u - I_{\text{nc}}u)$ (that stems from $a(u, u) - a_{\text{pw}}(u_{\text{nc}}, I_{\text{nc}}u)$ by (2.1)-(2.2)) does *not* allow for (2.9). The alternative split

$$\|e\|_{\text{pw}}^2 = a(u, Je_{\text{nc}}) - a_{\text{pw}}(u_{\text{nc}}, e_{\text{nc}}) + a_{\text{pw}}(u, e - Je_{\text{nc}}) \quad (3.6)$$

(this follows with the arguments in (3.1)) does allow for (2.9). Because of (2.1)-(2.2), the sum of the first two terms on the right-hand side of (3.6) is equal to

$$F((J - 1)e_{\text{nc}}) = \int_{\Omega} (f - \Pi_m f)((J - 1)e_{\text{nc}}) \, dx \leq \text{osc}_m(f, \mathcal{T}) \|h_{\mathcal{T}}^{-m}(1 - J)e_{\text{nc}}\|$$

with (2.9) followed by a weighted Cauchy inequality in $L^2(\Omega)$ in the end. Since $(1 - J)e_{\text{nc}} = (1 - I_{\text{nc}})(1 - J)e_{\text{nc}}$ from (2.7), (2.4) implies

$$\kappa_m^{-1} \|h_{\mathcal{T}}^{-m}(1 - J)e_{\text{nc}}\| \leq \|(1 - J)e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J \|e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J \|e\|_{\text{pw}} \quad (3.7)$$

with $v = 0$ in (2.8) for the second last and the Pythagoras theorem (e.g. (2.5) with $v = u$ and $w_{\text{nc}} = u_{\text{nc}}$) in the last step.

The last term of (3.6) is recast with $I_{\text{nc}}(e - Je_{\text{nc}}) = 0$ from (2.7) and $a_{\text{pw}}(I_{\text{nc}}u, e - Je_{\text{nc}}) = 0$ from (2.3) as

$$\begin{aligned} a_{\text{pw}}(u - I_{\text{nc}}u, e - Je_{\text{nc}}) &= \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + a_{\text{pw}}(u - I_{\text{nc}}u, (1 - J)e_{\text{nc}}) \\ &\leq (1 + \Lambda_J) \|u - I_{\text{nc}}u\|_{\text{pw}} \|e\|_{\text{pw}} \end{aligned} \quad (3.8)$$

with a Cauchy inequality and $\|(1 - J)e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J \|e\|_{\text{pw}}$ from (3.7) and (2.5) in the last step. The summary of the inequalities in (3.6) concludes the proof of Theorem 3.1 (with suboptimal constants as outlined in Remark 3.8 below).

Theorem 3.1 (best-approximation up to data approximation). *The assumptions (2.1)-(2.9) imply (3.4) for $C_1 := \max\{\kappa_m \Lambda_J, 1 + \Lambda_J\}$.*

3.3 Best-approximation for the modified scheme

The discrete scheme (2.2) requires a discrete right-hand side F_h for a general $F \in H^{-m}(\Omega)$. The evaluation of $F_h := F \circ J$ is feasible with $F_h(v_{\text{nc}}) := F(Jv_{\text{nc}})$ for all $v_{\text{nc}} \in V_{\text{nc}}$ and the (*modified*) *nonconforming scheme* seeks the solution $u_{\text{nc}} \in V_{\text{nc}}$ to

$$a_{\text{pw}}(u_{\text{nc}}, v_{\text{nc}}) = F(Jv_{\text{nc}}) \quad \text{for all } v_{\text{nc}} \in V_{\text{nc}}. \quad (3.9)$$

Let Λ_0 denote the norm of $1 - J$ of the right-inverse $J \in L(V_{\text{nc}}; V)$ regarded as a linear map between V_{nc} and V ,

$$\Lambda_0 := \sup_{v_{\text{nc}} \in V_{\text{nc}} \setminus \{0\}} \|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} / \|v_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J. \quad (3.10)$$

Theorem 3.2 (best-approximation). *(a) Under the assumptions (2.3)-(2.4), (2.7)-(2.8), the exact solution u to (2.1) and the discrete solution u_{nc} to (3.9) satisfy*

$$\|u - u_{\text{nc}}\|_{\text{pw}} \leq \sqrt{1 + \Lambda_0^2} \|u - I_{\text{nc}}u\|_{\text{pw}}.$$

(b) The best-approximation constant $\sqrt{1 + \Lambda_0^2} = C_{\text{qo}}$ is attained in the sense that there exist some $F = a(u, \bullet) \in V^$ and $u \in V$ with $\|u - u_{\text{nc}}\|_{\text{pw}} = C_{\text{qo}} \|u - I_{\text{nc}}u\|_{\text{pw}} > 0$ for the discrete solution $u_{\text{nc}} \in V_{\text{nc}}$ to (3.9).*

Proof of (a). Recall the error $e := u - u_{\text{nc}} \in V + V_{\text{nc}}$ and its interpolation $e_{\text{nc}} := I_{\text{nc}}u - u_{\text{nc}} = I_{\text{nc}}e \in V_{\text{nc}}$. The key identity

$$a(u, Je_{\text{nc}}) = F(Je_{\text{nc}}) = a_{\text{pw}}(u_{\text{nc}}, e_{\text{nc}}) \quad (3.11)$$

follows directly from (2.1) (resp. (3.9)) with the test function $Je_{\text{nc}} \in V$ (resp. e_{nc}). This proves in (3.6) that

$$\|e\|_{\text{pw}}^2 = a_{\text{pw}}(u, e - Je_{\text{nc}}) = a_{\text{pw}}(u - I_{\text{nc}}u, e - Je_{\text{nc}}) \quad (3.12)$$

with $a_{\text{pw}}(I_{\text{nc}}u, e - Je_{\text{nc}}) = 0$ in the last step. The term $a_{\text{pw}}(u - I_{\text{nc}}u, e - Je_{\text{nc}})$ is estimated in (3.8). It is true that u_{nc} denotes another discrete solution in (3.7)-(3.8), but the related

arguments remain valid verbatim in this proof as well and lead to the upper bound $(1 + \Lambda_J)\|e\|_{\text{pw}}\|u - I_{\text{nc}}u\|_{\text{pw}}$. This concludes an easy proof of the best-approximation result

$$\|u - u_{\text{nc}}\|_{\text{pw}} \leq (1 + \Lambda_J)\|u - I_{\text{nc}}u\|_{\text{pw}}.$$

The proof of the optimal best-approximation constant in Theorem 3.2 revisits (3.12) and the Pythagoras theorem $\|e\|_{\text{pw}}^2 = \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + \|e_{\text{nc}}\|_{\text{pw}}^2$ (e.g. (2.5) with $v = e$ and $w_{\text{nc}} = 0$). The combination with (3.12) proves

$$\|e_{\text{nc}}\|_{\text{pw}}^2 = a_{\text{pw}}(u - I_{\text{nc}}u, e_{\text{nc}} - Je_{\text{nc}}).$$

The Cauchy inequality and the continuity constant Λ_0 show that

$$\|e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_0\|u - I_{\text{nc}}u\|_{\text{pw}}$$

(after a division of $\|e_{\text{nc}}\|_{\text{pw}}$, if positive). This and the above Pythagoras theorem result in

$$\|e\|_{\text{pw}}^2 = \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + \|e_{\text{nc}}\|_{\text{pw}}^2 \leq (1 + \Lambda_0^2)\|u - I_{\text{nc}}u\|_{\text{pw}}^2.$$

This concludes the proof under conditions weaker than (2.8) and with a better constant.

Proof of (b). The second assertion on the attainment is immediate for $\Lambda_0 = 0$; for $V_{\text{nc}} \subset V$; see Remark 3.1 for a discussion of this case. Hence suppose $\Lambda_0 > 0$ in the sequel.

Note that Λ_0^2 from (3.10) is an eigenvalue and there exists $v_{\text{nc}} \in V_{\text{nc}}$ with $\|v_{\text{nc}}\|_{\text{pw}} = 1$ and

$$a_{\text{pw}}(v_{\text{nc}} - Jv_{\text{nc}}, w_{\text{nc}} - Jw_{\text{nc}}) = \Lambda_0^2 a_{\text{pw}}(v_{\text{nc}}, w_{\text{nc}}) \text{ for all } w_{\text{nc}} \in V_{\text{nc}}. \quad (3.13)$$

For $w_{\text{nc}} = v_{\text{nc}}$, (3.13) shows $\|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} = \Lambda_0$. Since $a_{\text{pw}}(v_{\text{nc}} - Jv_{\text{nc}}, w_{\text{nc}}) = 0$, (3.13) implies

$$a_{\text{pw}}(Jv_{\text{nc}}, Jw_{\text{nc}}) = (\Lambda_0^2 + 1)a_{\text{pw}}(v_{\text{nc}}, w_{\text{nc}}) \text{ for all } w_{\text{nc}} \in V_{\text{nc}}. \quad (3.14)$$

Define $F := -a(Jv_{\text{nc}}, \bullet)$; so that (2.1) has the exact solution $u = -Jv_{\text{nc}} \in V$ and the discrete problem for $u_{\text{nc}} \in V_{\text{nc}}$ reads

$$a_{\text{pw}}(u_{\text{nc}}, w_{\text{nc}}) = F(Jw_{\text{nc}}) = -a(Jv_{\text{nc}}, Jw_{\text{nc}}) \text{ for all } w_{\text{nc}} \in V_{\text{nc}}. \quad (3.15)$$

The combination of (3.14)-(3.15) shows $u_{\text{nc}} = -(1 + \Lambda_0^2)v_{\text{nc}}$. Consequently, $u - u_{\text{nc}} = v_{\text{nc}} - Jv_{\text{nc}} + \Lambda_0^2 v_{\text{nc}}$ and $v_{\text{nc}} = -I_{\text{nc}}u$. This and the orthogonality $a(v_{\text{nc}} - Jv_{\text{nc}}, v_{\text{nc}}) = 0$ lead to

$$\|u - u_{\text{nc}}\|_{\text{pw}}^2 = \|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}}^2 + \Lambda_0^4 \|v_{\text{nc}}\|_{\text{pw}}^2 = \Lambda_0^2(1 + \Lambda_0^2) > 0$$

with $\|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} = \Lambda_0$ and $\|v_{\text{nc}}\|_{\text{pw}} = 1$ in the last step. Since $\|u - I_{\text{nc}}u\|_{\text{pw}} = \|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} = \Lambda_0$ from the design of $u = -Jv_{\text{nc}}$ with $I_{\text{nc}}u = -v_{\text{nc}}$, this proves $\|u - u_{\text{nc}}\|_{\text{pw}} = \sqrt{1 + \Lambda_0^2}\|u - I_{\text{nc}}u\|_{\text{pw}}$. \square

Remark 3.1 ($V_{\text{nc}} \not\subset V$). The case $\Lambda_0 = 0$ corresponds to the optimal case $\|J\| = 1$ and implies that J is the identity in $V_{\text{nc}} \subset V$. This is reserved for the conforming Ritz-Galerkin scheme and excluded throughout this paper.

Remark 3.2 ($V_{\text{nc}} \neq P_m(\mathcal{T})$). It is clear that $V_{\text{nc}} \subset P_m(\mathcal{T})$ and $V_{\text{nc}} = P_m(\mathcal{T})$ is excluded in the following sense. If a linear operator $I_{\text{nc}} : V \rightarrow V_{\text{nc}}$ exists with (2.3) for $V_{\text{nc}} = P_m(\mathcal{T})$ and dimension $n = 2$, $m = 1$ or $m = 2$, then there exists no right-inverse J of I_{nc} . The proof argues by contradiction, so let $J : P_m(\mathcal{T}) \rightarrow V$ be a right-inverse of I_{nc} and start with the observation that (2.3) implies $D_{\text{pw}}^m I_{\text{nc}} v = \Pi_0 D^m v$ for all $v = J v_{\text{nc}} \in V$, $v_{\text{nc}} \in P_m(\mathcal{T})$. Since $I_{\text{nc}} v = v_{\text{nc}}$ (from $I_{\text{nc}} J v_{\text{nc}} = v_{\text{nc}}$), this means

$$D_{\text{pw}}^m v_{\text{nc}} = \Pi_0 D^m J v_{\text{nc}} \text{ for all } v_{\text{nc}} \in P_m(\mathcal{T}). \quad (3.16)$$

For a contradiction, let $\varphi_c \in S_0^1(\mathcal{T}) \setminus P_0(\Omega)$ (so that $\|\varphi_c\| \neq 0$) for $m = 1$ and let $\Psi_c \in S^1(\mathcal{T}; \mathbb{R}^2) \setminus P_0(\Omega; \mathbb{R}^2)$ (so that $\|\Psi_c\| \neq 0$) and then define $v_m \in P_m(\mathcal{T})$ for $x \in T \in \mathcal{T}$ by

$$\begin{aligned} v_1(x) &:= (\text{Curl } \varphi_c) \cdot (x - \text{mid}(T)) \\ \text{resp. } v_2(x) &:= \frac{1}{2}(x - \text{mid}(T)) \cdot (\text{sym}(\text{Curl } \Psi_c))(x - \text{mid}(T)). \end{aligned}$$

The point is that $\nabla_{\text{pw}} v_1 = \text{Curl } \varphi_c \in L^2(\Omega; \mathbb{R}^2)$ resp. $D_{\text{pw}}^2 v_2 = \text{sym}(\text{Curl } \Psi_c) \in L^2(\Omega; \mathbb{S})$ is L^2 perpendicular to ∇v resp. $D^2 v$ for $v := J v_m \in V$. Hence

$$0 = \int_{\Omega} D_{\text{pw}}^m v_m : D^m v \, dx = \int_{\Omega} D_{\text{pw}}^m v_m : D^m J v_m \, dx = \|v_m\|_{\text{pw}}^2$$

with (3.16) for $v_m \in P_m(\mathcal{T})$ in the last step. This is a contradiction to the nontrivial choice of φ_c resp. Ψ_c with $\|v_m\|_{\text{pw}} > 0$.

3.4 Interpretation as a Petrov-Galerkin scheme

This section continues with another interpretation of the modified nonconforming finite element method as a Petrov-Galerkin scheme in the Hilbert space $(\widehat{V}, a_{\text{pw}})$ for $\widehat{V} = V + V_{\text{nc}}$. The best-approximation operator $I_{\text{nc}} : \widehat{V} \rightarrow V_{\text{nc}}$ in the Hilbert space $(\widehat{V}, a_{\text{pw}})$ is characterized by the orthogonality $a_{\text{pw}}(u_{\text{nc}}, v - I_{\text{nc}} v) = 0$ for all $v \in V$. Since J is a right-inverse, this and $v = J v_{\text{nc}}$ for $v_{\text{nc}} \in V_{\text{nc}}$ imply $a_{\text{pw}}(u_{\text{nc}}, J v_{\text{nc}} - v_{\text{nc}}) = 0$. Suppose $u \in V$ defines $F := a(u, \bullet) \in V^*$ (or vice versa owing to the Riesz isomorphism), this and $a_{\text{pw}}(u_{\text{nc}}, v_{\text{nc}}) = a_{\text{pw}}(u_{\text{nc}}, J v_{\text{nc}})$ lead in (3.9) to

$$a_{\text{pw}}(u - u_{\text{nc}}, J v_{\text{nc}}) = 0 \quad \text{for all } v_{\text{nc}} \in V_{\text{nc}} \quad (3.17)$$

(and this is equivalent to (3.9) for $F = a(u, \bullet)$). This defines a *Petrov-Galerkin operator* $u \mapsto u_{\text{nc}}$ that is naturally extended to $\widehat{V} := V + V_{\text{nc}}$ with (3.17) as follows. Any $\widehat{u} \in \widehat{V}$ solves (3.9) with $F(J v_{\text{nc}})$ replaced by $a_{\text{pw}}(\widehat{u}, J v_{\text{nc}})$ for all $v_{\text{nc}} \in V_{\text{nc}}$ with the solution $P \widehat{u} := u_{\text{nc}} \in V_{\text{nc}}$. Therefore $P : \widehat{V} \rightarrow \widehat{V}$ is characterized by

$$a_{\text{pw}}(\widehat{v} - P \widehat{v}, J v_{\text{nc}}) = 0 \quad \text{for all } v_{\text{nc}} \in V_{\text{nc}} \text{ and for all } \widehat{v} \in \widehat{V}. \quad (3.18)$$

This characterizes a Petrov-Galerkin scheme in the Hilbert space $(\widehat{V}, a_{\text{pw}})$ with the trial space V_{nc} and the test space $V_{\widehat{c}} := J V_{\text{nc}}$. The Petrov-Galerkin operator $P \in L(\widehat{V})$ is regarded as a map from \widehat{V} into \widehat{V} . It is well defined, linear, bounded and idempotent. (In particular, any $u \in V$ is approximated by $u_{\text{nc}} = P u$.)

Based on the interpretation as a Petrov-Galerkin scheme, several conclusions may be drawn in the subsequent subsection.

3.5 Comments on the best-approximation constant C_{qo}

The subsection gives certain characterizations of the best-approximation constant C_{qo} in (3.19) and shows in particular, that the multiplicative constant $C_{\text{qo}} = \sqrt{1 + \Lambda_0^2}$ in Theorem 3.2 is optimal in the resulting estimate

$$\|u - u_{\text{nc}}\|_{\text{pw}} \leq C_{\text{qo}} \|u - I_{\text{nc}}u\|_{\text{pw}} \quad (3.19)$$

for the exact solution $u \in V$ to (2.1) and the discrete solution $u_{\text{nc}} \in V_{\text{nc}}$ to (3.9) and an arbitrary right-hand side $F \in V^*$. To be more precise, adopt the notation of a Petrov-Galerkin operator $P : V \rightarrow V_{\text{nc}}$ from (3.18) and specify

$$C_{\text{qo}} := \sup_{v \in V \setminus V_{\text{nc}}} \|v - Pv\|_{\text{pw}} / \|v - I_{\text{nc}}v\|_{\text{pw}} \quad (3.20)$$

with a supremum over all $v \in V$ with $v \neq I_{\text{nc}}v$ (for this leads to an undefined quotient $0/0$ that can be excluded) in the definition of the best multiplicative constant (3.20) for (3.19). To avoid a pathological situation, suppose throughout this paper that $V \setminus V_{\text{nc}}$ is non-empty and that $V_{\text{nc}} \neq \{0\}$ is nontrivial (as well as V). (The overall intention is that the nonconforming finite element space $V_{\text{nc}} \not\subset V$ is *not* conforming, but the case $V_{\text{nc}} \subset V$ is actually included in the analysis of this section.) Let $\|J\| := \|J\|_{L(V_{\text{nc}}; V)}$ abbreviate the operator norm of the right-inverse $J : V_{\text{nc}} \rightarrow V$.

Theorem 3.3 (characterization of best-approximation constant). $C_{\text{qo}} = \sqrt{1 + \Lambda_0^2} = \|J\|$.

Proof. *Step 1* recalls $\|1 - P\|_{L(\widehat{V})} = C_{\text{qo}}$. This is well known in the mathematical foundations of Petrov-Galerkin schemes and the short proof is given for completeness. Any $\widehat{v} \in \widehat{V} = V + V_{\text{nc}}$ is of the form $\widehat{v} = v + v_{\text{nc}} = w + w_{\text{nc}}$ for $w := (1 - I_{\text{nc}})v \in (1 - I_{\text{nc}})V = V_{\text{nc}}^\perp$ and $w_{\text{nc}} = I_{\text{nc}}v + v_{\text{nc}} \in V_{\text{nc}}$. Since $P\widehat{v} = Pw + w_{\text{nc}}$ and $\widehat{v} - P\widehat{v} = w - Pw$, the orthogonal split $\widehat{v} = w + w_{\text{nc}}$ shows

$$\|1 - P\|_{L(\widehat{V})} = \sup_{w, w_{\text{nc}}} \|w - Pw\|_{\text{pw}} (\|w\|_{\text{pw}}^2 + \|w_{\text{nc}}\|_{\text{pw}}^2)^{-1/2} = \sup_{w \in V_{\text{nc}}^\perp \setminus \{0\}} \|w - Pw\|_{\text{pw}} / \|w\|_{\text{pw}}$$

with the abbreviation $\sup_{w, w_{\text{nc}}}$ for the supremum over all $(w, w_{\text{nc}}) \in (V_{\text{nc}}^\perp \times V_{\text{nc}}) \setminus \{0\}$. This and the re-substitution $w = v - I_{\text{nc}}v$ and $w - Pw = v - Pv$ for $v \in V$ in the second equality conclude the proof,

$$\|1 - P\|_{L(\widehat{V})} = \|1 - P\|_{L(V_{\text{nc}}^\perp; \widehat{V})} = \sup_{v \in V \setminus V_{\text{nc}}} \|v - Pv\|_{\text{pw}} / \|v - I_{\text{nc}}v\|_{\text{pw}} = C_{\text{qo}}. \quad \square$$

Step 2 establishes $\|P\|_{L(\widehat{V})} = \|P\|_{L(V_{\text{c}}; V_{\text{nc}})}$ for the conforming companion finite element space $V_{\text{c}} := JV_{\text{nc}}$. Since $\|P\|_{L(V_{\text{c}}; V_{\text{nc}})} \leq \|P\|_{L(\widehat{V})}$ is obvious (for $V_{\text{c}} \subset \widehat{V}$), the proof of the converse inequality concerns an arbitrary $\widehat{v} \in \widehat{V}$ and its orthogonal projection $v_{\text{c}} \in V_{\text{c}}$ onto the (finite-dimensional, whence closed) subspace $V_{\text{c}} := JV_{\text{nc}}$ in the Hilbert space $(\widehat{V}, a_{\text{pw}})$ with orthogonality written as \perp . In particular, $v_{\text{c}} \in V_{\text{c}}$ and $\widehat{v} - v_{\text{c}} \perp V_{\text{c}}$. Recall from (3.18) that $P\widehat{v}$ is characterized by $P\widehat{v} \in V_{\text{nc}}$ and $\widehat{v} - P\widehat{v} \perp V_{\text{c}}$. The two orthogonalities lead to $v_{\text{c}} - P\widehat{v} \perp V_{\text{c}}$, whence $P\widehat{v} = Pv_{\text{c}}$. This and $\|v_{\text{c}}\|_{\text{pw}} \leq \|\widehat{v}\|_{\text{pw}}$ imply

$$\|P\widehat{v}\|_{\text{pw}} = \|Pv_{\text{c}}\|_{\text{pw}} \leq \|P\|_{L(V_{\text{c}}; V_{\text{nc}})} \|v_{\text{c}}\|_{\text{pw}} \leq \|P\|_{L(V_{\text{c}}; V_{\text{nc}})} \|\widehat{v}\|_{\text{pw}}. \quad \square$$

Step 3 computes $\|P\|_{L(V_c; V_{nc})} = \|J\|$ with linear algebra. Given any basis (ψ_1, \dots, ψ_N) of V_{nc} , $(J\psi_1, \dots, J\psi_N)$ forms a basis of V_c (for $v_{nc} \in V_{nc}$ with $Jv_{nc} = 0$ implies $v_{nc} = I_{nc}Jv_{nc} = 0$). The stiffness matrices $A, B \in \mathbb{R}^{N \times N}$ are defined by

$$A_{jk} := a_{pw}(\psi_j, \psi_k) \quad \text{and} \quad B_{jk} := a_{pw}(J\psi_j, J\psi_k) \quad \text{for all } j, k = 1, \dots, N.$$

Then A and B are SPD and characterize the Petrov-Galerin map $P : V_c \rightarrow V_{nc}$ in coefficient vectors $x, y \in \mathbb{R}^N$ for $v_{nc} = \sum_{j=1}^N x_j \psi_j$ and $v_c = \sum_{j=1}^N y_j J\psi_j$ by $v_{nc} = Pv_c$ is equivalent to $Ax = By$. The norm $\|P\|_{L(V_c; V_{nc})}$ of $P : V_c \rightarrow V_{nc}$ involves the norms with the squares $\|Pv_c\|_{pw}^2 = \|v_{nc}\|_{pw}^2 = x \cdot Ax$ and $\|v_c\|_{pw}^2 = y \cdot By$ and lead, for $x, y \in \mathbb{R}^N \setminus \{0\}$ with $Ax = By$, to $y \cdot By = x \cdot AB^{-1}Ax$. This and the Rayleigh-Ritz min-max principle show

$$\|P\|_{L(V_c; V_{nc})}^2 = \sup_{v_c \in V_c \setminus \{0\}} \frac{\|Pv_c\|_{pw}^2}{\|v_c\|_{pw}^2} = \sup_{x \in \mathbb{R}^N \setminus \{0\}} \frac{x \cdot Ax}{x \cdot AB^{-1}Ax} = \lambda$$

for the maximal eigenvalue of an eigenpair $(\lambda, x) \in \mathbb{R}_+ \times \mathbb{R}^N \setminus \{0\}$ of the generalized eigenvalue problem $Ax = \lambda AB^{-1}Ax$. Since the latter is equivalent to $Bx = \lambda Ax$, the maximal eigenvalue is

$$\lambda = \sup_{x \in \mathbb{R}^N \setminus \{0\}} \frac{x \cdot Bx}{x \cdot Ax} = \sup_{v_{nc} \in V_{nc} \setminus \{0\}} \frac{\|Jv_{nc}\|_{pw}^2}{\|v_{nc}\|_{pw}^2} = \|J\|^2. \quad \square$$

Step 4 verifies $\|J\|^2 = 1 + \Lambda_0^2$. Since J is a right-inverse to I_{nc} , the Pythagoras theorem $\|Jv_{nc}\|_{pw}^2 = \|v_{nc} - Jv_{nc}\|_{pw}^2 + \|v_{nc}\|_{pw}^2$ concludes the proof by the definition of $\|J\|$ and Λ_0 :

$$\|J\|^2 = \sup_{v_{nc} \in V_{nc} \setminus \{0\}} \frac{\|v_{nc} - Jv_{nc}\|_{pw}^2 + \|v_{nc}\|_{pw}^2}{\|v_{nc}\|_{pw}^2} = 1 + \sup_{v_{nc} \in V_{nc} \setminus \{0\}} \frac{\|v_{nc} - Jv_{nc}\|_{pw}^2}{\|v_{nc}\|_{pw}^2} = 1 + \Lambda_0^2. \quad \square$$

Step 5 concludes the proof with Kato's lemma. Any oblique projection (that is a bounded linear map $P \in L(\widehat{V})$ in a Hilbert space \widehat{V} with $P^2 = P$) satisfies $\|1 - P\|_{L(\widehat{V})} = \|P\|_{L(\widehat{V})}$ provided $0 \neq P \neq 1$. This is known as Kato's lemma [42] with many very different proofs in [48]. The summary of all steps concludes the proof. \square

Remark 3.3. Veesper et al. [53] provide the bound $C_{q0} = \|J\|$, while Theorem 3.2 gives $C_{q0} = \sqrt{1 + \Lambda_0^2}$ and both results are optimal.

Remark 3.4 (J vs V_c). The Petrov-Galerkin scheme and its operator P require the trial space V_{nc} and the test space V_c , but no knowledge about J . However, $V_c := J(V_{nc})$ depends very much on the right-inverse. In fact, given any $V_c \subset V$ of dimension $\dim V_c = \dim V_{nc} \in \mathbb{N}$, there is at most one right inverse: If I_{nc} is not injective as a linear map from V_c into V_{nc} , or equivalently (for $\dim V_c = \dim V_{nc}$) it is not surjective, then there exists no right-inverse $J : V_{nc} \rightarrow V$ with $V_c = J(V_{nc})$. But otherwise J exists and is uniquely determined by V_c . (If $J : V_{nc} \rightarrow V_c$ is a linear map and $I_{nc}J$ is identity in V_{nc} then J is injective and hence bijective for $\dim V_c = \dim V_{nc}$. Then J is the inverse of I_{nc} , when I_{nc} is regarded as a bijection of V_c onto V_{nc} .)

Remark 3.5 ($\|J\| > 1$). Theorem 3.3 asserts $\|J\| \geq 1$ and the optimal case $\|J\| = 1$ for the conforming Ritz-Galerkin scheme is discussed in Remark 3.1. For the truly nonconforming situation $V_{nc} \not\subset V$, the quality of the scheme is controlled by $C_{q0} = \|J\| > 1$.

Remark 3.6 (another characterization of Λ_0). Theorem 3.3 also reveals the identity $\Lambda_0 = \|P\|_{L(V_{\text{nc}}^\perp; \widehat{V})}$ the orthogonal complement of V_{nc} in \widehat{V} . This identity is behind the refined proof of Theorem 3.2. The proof of it considers $Pw \perp w \in V_{\text{nc}}^\perp$ and the Pythagoras theorem $\|w - Pw\|_{\text{pw}}^2 = \|Pw\|_{\text{pw}}^2 + \|w\|_{\text{pw}}^2$ for the first equation in

$$\|1 - P\|_{L(V_{\text{nc}}^\perp; \widehat{V})}^2 = \sup_{w \in V_{\text{nc}}^\perp \setminus \{0\}} (\|Pw\|_{\text{pw}}^2 + \|w\|_{\text{pw}}^2) \|w\|_{\text{pw}}^{-2} = 1 + \|P\|_{L(V_{\text{nc}}^\perp; \widehat{V})}^2$$

Theorem 3.3 and the last line in Step 1 of its proof show that this is equal to $1 + \Lambda_0^2$. \square

Remark 3.7 (post-processing). The error $u - Ju_{\text{nc}}$ of the post-processed discrete solution can be measured in all Sobolev scales $H^s(\Omega)$ for $-m \leq s \leq m$ and we start with $s = m$ and the affirmative estimate

$$\left| \|u - u_{\text{nc}}\|_{\text{pw}} - \|u - Ju_{\text{nc}}\| \right| \leq \Lambda_J \|u - u_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J C_{\text{qo}} \|u - I_{\text{nc}}u\|_{\text{pw}}.$$

(This follows from a reverse triangle inequality and (2.8) plus Theorem 3.2.) This estimate compares with errors in the continuous and piecewise energy norms up to a best-approximation term on the right-hand side. In particular, the post-processing is quasi-optimal. However, it is *not* quasi-optimal in comparison with the best-approximation in $V_c := JV_{\text{nc}}$: Otherwise, with the Petrov-Galerkin operator $P : V \rightarrow V_{\text{nc}}$ from (3.18), any inequality $\|v - JPv\|_{\text{pw}} \lesssim \|v - v_c\|_{\text{pw}}$ for all $v \in V$ and $v_c \in V_c$ implies, in particular, that $v_c = JPv_c$ for all $v = v_c \in V_c$. Since J is a right-inverse, the last identity shows that $I_{\text{nc}}v_c = Pv_c$ for all $v_c \in V_c$. But this implies $\|P\|_{L(V_c; V_{\text{nc}})} \leq 1$. Step 3 in the proof of Theorem 3.3 leads to $\|J\| = 1$. This is characterized in Remark 3.1 as the conforming case $V_{\text{nc}} = V_c \subset V$.

Remark 3.8 (sharper constants in Theorem 3.1). The constant C_1 in (3.4) is not much larger than $\sqrt{1 + \Lambda_0^2}$. Follow the proof until (3.7) and replace Λ_J by Λ_0 and ignore the last estimate. Utilize Λ_0 for $\|(1 - J)e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_0 \|e_{\text{nc}}\|_{\text{pw}}$ in the equality (3.8). This leads to an estimate for the right-hand side of (3.6) that includes the additive term $\|u - I_{\text{nc}}u\|_{\text{pw}}^2$ (still from (3.8)) that arises also in the Pythagoras theorem $\|e\|_{\text{pw}}^2 = \|e_{\text{nc}}\|_{\text{pw}}^2 + \|u - I_{\text{nc}}u\|_{\text{pw}}^2$ on the left-hand side. This proves (after a division of $\|e_{\text{nc}}\|_{\text{pw}}$ if positive)

$$\|e_{\text{nc}}\|_{\text{pw}} \leq \Lambda_0 (\kappa_m \text{osc}_m(f, \mathcal{T}) + \|u - I_{\text{nc}}u\|_{\text{pw}}).$$

This identity and the Pythagoras theorem again conclude the proof of

$$\|e_{\text{nc}}\|_{\text{pw}}^2 \leq (1 + \Lambda_0^2) \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + 2\Lambda_0^2 \kappa_m \text{osc}_m(f, \mathcal{T}) \|u - I_{\text{nc}}u\|_{\text{pw}} + \kappa_m^2 \text{osc}_m(f, \mathcal{T})^2.$$

The right-hand side is equal to $(\|u - I_{\text{nc}}u\|_{\text{pw}}, \text{osc}_m(f, \mathcal{T})) \cdot A (\|u - I_{\text{nc}}u\|_{\text{pw}}, \text{osc}_m(f, \mathcal{T}))$ with the symmetric 2×2 matrix A that has entries $A_{11} = 1 + \Lambda_0^2$, $A_{12} = \Lambda_0^2 \kappa_m$, and $A_{22} = \kappa_m^2$. The eigenvalues of A can be computed and the bigger one leads to a bound (3.4) with

$$2(1 + \Lambda_0^2) < 2C_1^2 = 1 + \kappa_m^2 + \Lambda_0^2 + \sqrt{(1 - \kappa_m^2 + \Lambda_0^2)^2 + 4\kappa_m^2 \Lambda_0^4}.$$

Since κ_m is small, the constant C_1 is not significantly larger than C_{qo} in (3.19). \square

4 Weaker and piecewise Sobolev norm estimates

4.1 Lower-order error estimates

This section lists further conditions sufficient for a lower-order a priori error estimate for the discrete problem (2.2) under the hypotheses (2.3)-(2.4) and (2.7)-(2.8).

The duality argument relies on the elliptic regularity and the approximation properties of the dual and discrete problem with constants $0 < \sigma \leq 1$ and $0 < C_{\text{reg}}(s), C_{\text{int}}(s) < \infty$ such that for any $G \in H^{-s}(\Omega)$ with $m - \sigma \leq s \leq m$, the weak solution $z \in V = H_0^m(\Omega)$ to the PDE $(-1)^m \Delta^m z = G$ satisfies $z \in V \cap H^{2m-s}(\Omega)$ and

$$\|z\|_{H^{2m-s}(\Omega)} \leq C_{\text{reg}}(s) \|G\|_{H^{-s}(\Omega)}, \quad (4.1)$$

$$\|z - I_{\text{nc}}z\|_{\text{pw}} \leq C_{\text{int}}(s) h_{\text{max}}^{m-s} \|z\|_{H^{2m-s}(\Omega)}. \quad (4.2)$$

Since in general $u_{\text{nc}} \in V_{\text{nc}}$ may not belong to $H^{m-s}(\Omega)$, the post-processing Ju_{nc} is considered in the lower-order a priori error estimate.

Theorem 4.1 (lower-order error estimates). *Suppose the hypotheses of Theorem 3.2, adopt the notation (4.1)-(4.2) for some s with $m - \sigma \leq s \leq m$. Given any $F \in H^{-s}(\Omega)$, the solutions $u \in V$ to (2.1) and $u_{\text{nc}} \in V_{\text{nc}}$ to (2.2) satisfy (a) $\|u - Ju_{\text{nc}}\|_{H^s(\Omega)} \leq C_2(s) h_{\text{max}}^{m-s} \|u - u_{\text{nc}}\|_{\text{pw}}$ and (b) $\|u - u_{\text{nc}}\|_{H^s(\mathcal{T})} \leq C_3(s) h_{\text{max}}^{m-s} \|u - u_{\text{nc}}\|_{\text{pw}}$.*

Proof of (a). The duality argument is applied for the exact solution $u \in V$ to (2.1) and the post-processing $v := Ju_{\text{nc}} \in V$ for the discrete solution $u_{\text{nc}} \in V_{\text{nc}}$ to (2.2). The duality of $H_0^s(\Omega)$ and $H^{-s}(\Omega)$ reveals

$$\|u - v\|_{H^s(\Omega)} = \sup_{0 \neq G \in H^{-s}(\Omega)} \frac{G(u - v)}{\|G\|_{H^{-s}(\Omega)}} = G(u - v)$$

and the supremum is attained for some $G \in H^{-s}(\Omega) \subset V^*$ with norm $\|G\|_{H^{-s}(\Omega)} = 1$ (owing to a corollary of the Hahn-Banach theorem). The functional has a unique Riesz representation $z \in V$ in the Hilbert space (V, a) , $a(\bullet, z) = G \in V^*$; i.e., $z \in V$ the weak solution to the PDE $(-1)^m \Delta^m z = G$. The elliptic regularity (4.1) leads to $z \in V \cap H^{2m-s}(\Omega)$ and (4.1)-(4.2); in particular,

$$\|u - v\|_{H^s(\Omega)} = a(u - v, z) \quad \text{and} \quad \|z\|_{H^{2m-s}(\Omega)} \leq C_{\text{reg}}(s). \quad (4.3)$$

Let $v := Ju_{\text{nc}} \in V$ and $\zeta := JI_{\text{nc}}z \in V$, where u_{nc} (resp. z) denote the discrete (resp. dual) solution. The test function $\zeta = JI_{\text{nc}}z \in V$ in (2.1) and the test function $I_{\text{nc}}z$ in (2.2) lead to

$$a_{\text{pw}}(u_{\text{nc}}, I_{\text{nc}}z) = F(\zeta) = a(u, \zeta).$$

This and elementary algebra with $a_{\text{pw}}((1 - J)u_{\text{nc}}, I_{\text{nc}}z) = 0 = a_{\text{pw}}(u_{\text{nc}}, (1 - J)I_{\text{nc}}z)$ (from (2.3) and (2.7)), result in

$$a(u - v, z) = a(u - v, z - \zeta) + a_{\text{pw}}(u_{\text{nc}} - v, \zeta - I_{\text{nc}}z). \quad (4.4)$$

From (2.8) deduce that

$$\|v - u_{\text{nc}}\|_{\text{pw}} \leq \Lambda_J \|u - u_{\text{nc}}\|_{\text{pw}} \quad \text{and} \quad \|\zeta - I_{\text{nc}}z\|_{\text{pw}} \leq \Lambda_J \|z - I_{\text{nc}}z\|_{\text{pw}}. \quad (4.5)$$

The combination of (4.5) and triangle inequality leads to

$$\|u - v\| \leq \|u - u_{\text{nc}}\|_{\text{pw}} + \|v - u_{\text{nc}}\|_{\text{pw}} \leq (1 + \Lambda_J) \|u - u_{\text{nc}}\|_{\text{pw}}. \quad (4.6)$$

Similar arguments for $(z, \zeta, I_{\text{nc}}z)$ replacing (u, v, u_{nc}) reveal

$$\|z - \zeta\| \leq (1 + \Lambda_J) \|z - I_{\text{nc}}z\|_{\text{pw}}. \quad (4.7)$$

A Cauchy inequality for the terms on the right-hand side of (4.4) followed by an application of (4.5)-(4.7) implies that $a(u - v, z)$ is bounded by $((1 + \Lambda_J)^2 + \Lambda_J^2) \|u - u_{\text{nc}}\|_{\text{pw}} \|z - I_{\text{nc}}z\|_{\text{pw}}$. A substitution to (4.3) leads to

$$\|u - v\|_{H^s(\Omega)} \leq ((1 + \Lambda_J)^2 + \Lambda_J^2) \|u - u_{\text{nc}}\|_{\text{pw}} \|z - I_{\text{nc}}z\|_{\text{pw}}.$$

This, (4.2), and (4.3b) lead to $C_2(s) := C_{\text{int}}(s)C_{\text{reg}}(s)((1 + \Lambda_J)^2 + \Lambda_J^2)$. \square

Proof of (b). The Sobolev-Slobodeckii semi-norm over Ω involves double integrals over $\Omega \times \Omega$ and so is larger than or equal to the sum of the integrals over $T \times T$ for all the triangles $T \in \mathcal{T}$, i.e., $\sum_{T \in \mathcal{T}} |\bullet|_{H^s(T)}^2 \leq |\bullet|_{H^s(\Omega)}^2$ for any $m - 1 < s < m$. Hence Theorem 4.1.a implies for all s with $m - \sigma \leq s \leq m$ and a constant $C_2(s)$ that

$$\|u - Ju_{\text{nc}}\|_{H^s(\mathcal{T})} \leq C_2(s) h_{\text{max}}^{m-s} \|u - u_{\text{nc}}\|_{\text{pw}}. \quad (4.8)$$

The equivalence of the Sobolev-Slobodeckii norm and the norm by interpolation of Sobolev spaces [43, Remark 9.1], for instance for a fixed reference triangle $T = T_{\text{ref}}$ with $C_4(s) = C_4(s, T_{\text{ref}})$, provides for $w := (u_{\text{nc}} - Ju_{\text{nc}})|_T \in H^m(T)$ the estimate

$$\|w\|_{H^s(T)} \leq C_4(s) \|w\|_{H^{m-1}(T)}^{m-s} \|w\|_{H^m(T)}^{s-m+1} \quad \text{for } m - 1 < s < m. \quad (4.9)$$

A straightforward transformation of Sobolev norms [28, Theorem 3.1.2] shows (4.9) for any triangle $T \in \mathcal{T}$ with $C_4(s) = C_4(s, T) = \kappa^{1+s} C_4(s, T_{\text{ref}})$ for the condition number $\kappa = \sigma_1/\sigma_2$ of the affine transformation $a + Bx$ of T_{ref} to T with the 2×2 matrix B and its positive singular values $\sigma_2 \leq \sigma_1$. A more detailed analysis [23] reveals that $C_4(s)$ exclusively depends on s (but exploits singularities as s approaches the end-points $m - 1$ and m). The estimate (4.9) shows the first inequality in

$$C_4(s)^{-2} \|w\|_{H^s(T)}^2 \leq \|w\|_{H^{m-1}(T)}^{2(m-s)} \|w\|_{H^m(T)}^{2(s-m+1)} \leq \|w\|_{H^{m-1}(T)}^2 + \|w\|_{H^{m-1}(T)}^{2(m-s)} |w|_{H^m(T)}^{2(s-m+1)}$$

with the subadditivity $(a + b)^p \leq a^p + b^p$ for $a, b \geq 0$ and $0 < p = s - m + 1 < 1$ (e.g. from the concavity of $x \mapsto x^p$ for non-negative x) in the last step. An elementary estimate is followed by the Young's inequality $ab \leq a^p/p + b^q/q$ for $p = (m - s)^{-1}$, $q = (s - m + 1)^{-1}$, $a = \|h_T^{-1}w\|_{H^1(T)}^{2(m-s)}$, and $b = |w|_{H^2(T)}^{2(s-m+1)}$ to prove

$$h_{\text{max}}^{2(s-m)} \|w\|_{H^{m-1}(T)}^{2(m-s)} |w|_{H^m(T)}^{2(s-m+1)} \leq \|h_T^{-1}w\|_{H^{m-1}(T)}^{2(m-s)} |w|_{H^m(T)}^{2(s-m+1)} \leq \|h_T^{-1}w\|_{H^{m-1}(T)}^2 + |w|_{H^m(T)}^2.$$

This and the trivial estimate $h_{\text{max}}^{2(s-m+1)} \leq \text{diam}(\Omega)^{2(s-m+1)}$ lead to $C_5(s) = C_4(s)^2(1 + \text{diam}(\Omega)^{2(s-m+1)})$ in

$$\|w\|_{H^s(T)}^2 \leq C_5(s) h_{\text{max}}^{2(m-s)} \left(\|h_T^{-1}w\|_{H^{m-1}(T)}^2 + |w|_{H^m(T)}^2 \right).$$




Figure 2: Rates of convergence t of $\|u - Ju_{\text{nc}}\|_{H^s(\Omega)}$ as $h_{\text{max}} \rightarrow 0$ for $-m \leq s < m + 1/2$.

The sum over all those contributions over $T \in \mathcal{T}$ reads

$$\|u_{\text{nc}} - Ju_{\text{nc}}\|_{H^s(\mathcal{T})} \leq \sqrt{C_5(s)} h_{\text{max}}^{m-s} \left(\|h_T^{-1}(u_{\text{nc}} - Ju_{\text{nc}})\|_{H^{m-1}(\mathcal{T})} + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}} \right).$$

An inverse estimate shows $\|h_T^{-1}(u_{\text{nc}} - Ju_{\text{nc}})\|_{H^{m-1}(T)} \lesssim h_T^{-m} \|u_{\text{nc}} - Ju_{\text{nc}}\|_{L^2(T)}$ for all $T \in \mathcal{T}$ with a constant in \lesssim that depends only on the shape-regularity. This, (2.4), a triangle inequality, (4.8), and (2.8) conclude the proof of (b). \square

4.2 Precise convergence rates for quasi-uniform meshes

This subsection comments on the sharpness of the a priori estimates of Theorem 4.1 and discusses conditions sufficient for the convergence rate $t := \min\{2\sigma, m + \sigma - s\}$ of the error $u - Ju_{\text{nc}} \in V = H_0^m(\Omega)$ in the norm of $H^s(\Omega)$ for $-m \leq s \leq m$ as $h_{\text{max}} \rightarrow 0$. Under the generic assumption $\|u - u_{\text{nc}}\|_{\text{pw}} \approx h_{\text{max}}^\sigma$, the estimate

$$\|u - Ju_{\text{nc}}\|_{H^s(\Omega)} \approx h_{\text{max}}^t \text{ for } -m \leq s < m + 1/2 \quad (4.10)$$

holds for a class of quasi-uniform shape-regular triangulations \mathcal{T} with sufficiently small maximal mesh-size $h_{\text{max}} \ll 1$. Figure 2 depicts the convergence rate t in (4.10) as $h_{\text{max}} \rightarrow 0$.

Theorem 4.2. *Given $F \in H_0^{\max\{0, -s\}}(\Omega)$ for $-m \leq s \leq m + 1/2$, let $u \in V$ solve (2.1) and let $u_{\text{nc}} \in V_{\text{nc}}$ solve (3.9). Suppose $\sigma < m$, $t := \min\{2\sigma, m + \sigma - s\}$, and that for a sequence of quasi-uniform shape-regular triangulations*

$$h_{\text{max}}^\sigma \lesssim \|u - u_{\text{nc}}\|_{\text{pw}} \quad (4.11)$$

holds. Then $\|u - Ju_{\text{nc}}\|_{H^s(\Omega)} \approx h_{\text{max}}^t$ as $h_{\text{max}} \rightarrow 0$.

The proof of the theorem for $-m \leq s \leq m - \sigma$ rests on the subsequent lemma.

Lemma 4.3. *For $F \in H_0^{\max\{0, -s\}}(\Omega)$ and $-m \leq s \leq m - \sigma$, the exact solution $u \in V$ to (2.1) and the discrete solution $u_{\text{nc}} \in V_{\text{nc}}$ to (3.9) satisfy*

$$\|u - u_{\text{nc}}\|_{\text{pw}}^2 \leq 2\kappa_m h_{\text{max}}^m (1 + \Lambda_J) \|F\| \|u - u_{\text{nc}}\|_{\text{pw}} + \|F\|_{H^{-s}(\Omega)} \|u - Ju_{\text{nc}}\|_{H^s(\Omega)}.$$

Proof. The weak solution u to $(-1)^m \Delta^m u = F$ satisfies $a(u, u) = F(u)$. The discrete solution u_{nc} to (3.9) satisfies $a_{\text{pw}}(u_{\text{nc}}, 2I_{\text{nc}}u - u_{\text{nc}}) = F(2JI_{\text{nc}}u - Ju_{\text{nc}})$. Consequently,

$$\|u - u_{\text{nc}}\|_{\text{pw}}^2 = F(Ju_{\text{nc}} - u) + 2F(u - JI_{\text{nc}}u).$$

The first term on the right-hand side of the above expression is $\leq \|F\|_{H^{-s}(\Omega)} \|u - Ju_{\text{nc}}\|_{H^s(\Omega)}$ while the second term, for $F \in L^2(\Omega)$, is twice

$$\begin{aligned} F(u - JI_{\text{nc}}u) &\leq \kappa_m h_{\text{max}}^m \|F\| \|u - JI_{\text{nc}}u\|_{\text{pw}} \\ &\leq \kappa_m (1 + \Lambda_J) h_{\text{max}}^m \|F\| \|u - I_{\text{nc}}u\|_{\text{pw}} \end{aligned}$$

with (2.4) in the first step and a triangle inequality plus (2.8) in the last step. A combination of the estimates concludes the proof. \square

Proof of Theorem 4.2 for $-m \leq s \leq m - \sigma$. Suppose that Ω is a (possibly nonconvex) polygon with elliptic regularity index $1/2 < \sigma \leq 1$ and \mathcal{T} is quasi-uniform with mesh-size h_{max} . The estimate $\|u - u_{\text{nc}}\|_{\text{pw}} \lesssim h_{\text{max}}^\sigma$ follows from Theorem 3.2 and the interpolation approximation estimate $\|u - I_{\text{nc}}u\|_{\text{pw}} \lesssim h_{\text{max}}^\sigma$ for $u \in H^{m+\sigma}(\Omega)$. The additional assumption (4.11) is that the convergence rate is not better than σ . This and Lemma 4.3 imply

$$C_6 h_{\text{max}}^{2\sigma} \leq \|u - Ju_{\text{nc}}\|_{H^s(\Omega)} + h_{\text{max}}^{m+\sigma}$$

for a positive constant $C_6 \approx 1$. In case $m = 1 = \sigma$, this estimate is inconclusive. But otherwise $2\sigma < m + \sigma$ (e.g. for $m = 2$ or $\sigma < 1$) and $h_{\text{max}}^{2\sigma} (C_6 - h_{\text{max}}^{m-\sigma}) \leq \|u - Ju_{\text{nc}}\|_{H^s(\Omega)}$ shows, at least for sufficiently small $h_{\text{max}} \ll 1$, that $h_{\text{max}}^{2\sigma} \lesssim \|u - Ju_{\text{nc}}\|_{H^s(\Omega)}$. The Sobolev embedding for $s \leq m - \sigma$ shows $\|u - Ju_{\text{nc}}\|_{H^s(\Omega)} \lesssim \|u - Ju_{\text{nc}}\|_{H^{m-\sigma}(\Omega)}$. Theorem 4.1 and $\|u - u_{\text{nc}}\|_{\text{pw}} \lesssim h_{\text{max}}^\sigma$ imply $\|u - Ju_{\text{nc}}\|_{H^{m-\sigma}(\Omega)} \lesssim h_{\text{max}}^{2\sigma}$. The combination of all those estimates concludes the proof for $-m \leq s \leq m - \sigma$. \square

Suppose \mathcal{T} is quasi-uniform and $0 < \varepsilon < 1/2$ throughout the remaining parts of this subsection. A quasi-uniform mesh allows inverse estimates even in the Sobolev-Slobodeckii semi-norm. Throughout the remaining part of this subsection, the generic factors hidden in \lesssim may also depend on ε .

Lemma 4.4 (inverse estimate). *It holds $\varepsilon^{1/2} \|u - Ju_{\text{nc}}\|_{H^{m+\varepsilon}(\Omega)} \lesssim h_{\text{max}}^{\sigma-\varepsilon} |u|_{H^{m+\sigma}(\Omega)}$.*

Proof. The best-approximation property leads to $\|u - Ju_{\text{nc}}\|_{H^m(\Omega)} \lesssim h_{\text{max}}^\sigma |u|_{H^{m+\sigma}(\Omega)}$ and controls the low-order contributions. Hence the proof focusses on the Sobolev-Slobodeckii semi-norm of each partial-derivative ∂^α of order $|\alpha| = m$. Let $f := \partial^\alpha u$, $g := \partial^\alpha Ju_{\text{nc}}$. The triangle inequality motivates the split

$$|f - g|_{H^\varepsilon(\Omega)} \leq |f - \Pi_0 f|_{H^\varepsilon(\Omega)} + |\Pi_0 f - g|_{H^\varepsilon(\Omega)}.$$

The first term is controlled by a Poincare inequality for the Sobolev-Slobodeckii semi-norm

$$\|f - \Pi_0 f\|_{H^\varepsilon(\Omega)} \lesssim \varepsilon^{-1/2} h_{\text{max}}^{s-\varepsilon} |f|_{H^s(\Omega)}.$$

(This is a standard estimate for the reference triangle e.g. by interpolation and a scaling argument completes the proof with a generic constant that may depend on ε . The claimed multiplicative constant $\approx \varepsilon^{-1/2}$ follows from a refined estimation [23].) The second term

is an inverse estimate for $Ju_{\text{nc}} \in V$ that is a piecewise polynomial (e.g. with respect to a HCT refinement of \mathcal{T}) with mesh-sizes equivalent to $h_{\mathcal{T}}$. Since $0 < \varepsilon < 1/2$, any piecewise polynomial $p_r \in P_r(\mathcal{T})$ of a degree at most $r \in \mathbb{N}_0$ belongs to $H^\varepsilon(\Omega)$ and the inverse estimate $\|p_r\|_{H^\varepsilon(\Omega)} \lesssim \varepsilon^{-1/2} \|h_{\mathcal{T}}^{-\varepsilon} p_r\|_{L^2(\Omega)}$ holds [30] for a quasi-uniform mesh (the multiplicative constant $\approx \varepsilon^{-1/2}$ follows from a refined estimation [23]). Since this is available for the above piecewise polynomial $\Pi_0 f - g$, it implies

$$\varepsilon^{1/2} \|\Pi_0 f - g\|_{H^\varepsilon(\Omega)} \lesssim \|h_{\mathcal{T}}^{-\varepsilon} (\Pi_0 f - g)\| \lesssim h_{\min}^{-\varepsilon} \|Ju_{\text{nc}} - I_{\text{nc}} u\|_{\text{pw}}$$

for the minimal mesh-size $h_{\min} := \min_{T \in \mathcal{T}} h_T$ in \mathcal{T} and $h_{\min} \approx h_{\max}$ for a quasi-uniform triangulation. A triangle inequality with (2.5), (2.8), and Theorem 3.2 lead to

$$\begin{aligned} \|I_{\text{nc}} u - Ju_{\text{nc}}\|_{\text{pw}} &\leq \|I_{\text{nc}} u - u_{\text{nc}}\|_{\text{pw}} + \|u_{\text{nc}} - Ju_{\text{nc}}\| \\ &\leq C_{\text{qo}}(1 + \Lambda_J) \|u - I_{\text{nc}} u\|_{\text{pw}} \lesssim h_{\max}^\sigma |u|_{H^{m+\sigma}(\Omega)}. \end{aligned}$$

The combination of the above estimates for each derivative $f = \partial^\alpha u$ of order $m = |\alpha|$ shows $\varepsilon^{1/2} \|u - Ju_{\text{nc}}\|_{H^{m+\varepsilon}(\Omega)} \lesssim h_{\max}^{\sigma-\varepsilon} |u|_{H^{m+\sigma}(\Omega)}$. This concludes the proof for a quasi-uniform triangulation. \square

Lemma 4.5 (interpolation). *If $0 < s < m$ and $0 < \varepsilon < 1/2$, then*

$$\|u - Ju_{\text{nc}}\|_{H^m(\Omega)}^{s+\varepsilon} \lesssim h_{\max}^{s(\sigma-\varepsilon)} |u|_{H^{m+\sigma}(\Omega)}^s \|u - Ju_{\text{nc}}\|_{H^{m-s}(\Omega)}^\varepsilon.$$

Proof. The interpolation [49] of $H^m(\Omega)$ between $H^{m-s}(\Omega)$ and $H^{m+\varepsilon}(\Omega)$ for $e := u - Ju_{\text{nc}} \in V \cap H^{m+\varepsilon}(\Omega)$ leads to $\|e\|_{H^m(\Omega)} \lesssim \|e\|_{H^{m-s}(\Omega)}^{s/(s+\varepsilon)} \|e\|_{H^{m+\varepsilon}(\Omega)}^{\varepsilon/(s+\varepsilon)}$. Lemma 4.4 reveals $\|e\|_{H^{m+\varepsilon}(\Omega)} \lesssim h_{\max}^{\sigma-\varepsilon} |u|_{H^{m+\sigma}(\Omega)}$ and concludes the proof. \square

Proof of Theorem 4.2 for $m - \sigma \leq s \leq m$. The upper bound for the error $e := u - Ju_{\text{nc}} \in V$ in the norm of $H^s(\Omega)$ follows from Theorem 4.1. The assumption (4.11) and Lemma 4.5 reveal $h_{\max}^{\sigma+m-s} \lesssim |u|_{H^{m+\sigma}(\Omega)}^{s/(\sigma+\varepsilon)} \|e\|_{H^s(\Omega)}$. This concludes the proof for $m - \sigma \leq s \leq m$. \square

Proof of Theorem 4.2 for $m \leq s < m + 1/2$. The upper bounds for quasi-uniform meshes follow from Lemma 4.4. The lower bounds are derived by interpolation for $e := u - Ju_{\text{nc}} \in V \cap H^{m+\varepsilon}(\Omega)$. For the convex coefficients $\theta = \varepsilon/(\sigma + \varepsilon)$ and $1 - \theta = \sigma/(\sigma + \varepsilon)$ in $m = \theta(m - \sigma) + (1 - \theta)(m + \varepsilon)$, an interpolation [49] shows

$$\|e\| \leq C(\varepsilon) \|e\|_{H^{m-\sigma}(\Omega)}^\theta \|e\|_{H^{m+\varepsilon}(\Omega)}^{1-\theta}$$

for some constant $C(\varepsilon)$ that exclusively depends on ε and Ω . This and Theorem 4.1 result in $\|e\|^{\sigma+\varepsilon} \lesssim h_{\max}^{2\sigma} \varepsilon \|e\|_{H^{m+\varepsilon}(\Omega)}^\sigma$. Recall (4.11) and derive the asserted lower bound $h_{\max}^{\sigma-\varepsilon} \lesssim \|e\|_{H^{m+\varepsilon}(\Omega)}$. This concludes the proof. \square

5 Comments and examples of data in $H^{-m}(\Omega)$

5.1 The space $H^{-m}(\Omega)$

The space $H^{-m}(\Omega)$ is the dual to $H_0^m(\Omega)$ and is a real Hilbert space. It is known that for any functional $F \in H^{-m}(\Omega)$, there exist $f_\alpha \in L^2(\Omega)$ for all multi-indices α with $|\alpha| \leq m$ such that

$$F(v) = \sum_{|\alpha| \leq m} \int_{\Omega} f_\alpha \partial_\alpha v \, dx \quad \text{for all } v \in H_0^m(\Omega) \quad (5.1)$$

(with multi-index notation $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_n) \in \mathbb{N}_0^n$ and $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n$ for the partial derivatives $\partial_\alpha v := \partial^{|\alpha|} v / (\partial^{\alpha_1} x_1 \cdots \partial^{\alpha_n} x_n)$ etc). The family $(f_\alpha : |\alpha| \leq m)$ of $L^2(\Omega)$ functions with (5.1) is *not* unique. For instance, given any $F \in L^2(\Omega)$ one may solve $(-1)^m \Delta^m u = F$ for a weak solution $u \in H_0^m(\Omega)$ and consider $G := D^m u \in L^2(\Omega; \mathbb{R}^{(n^m)})$ with

$$F(v) = a(u, v) = \int_{\Omega} G : D^m v \, dx \text{ for all } v \in H_0^m(\Omega).$$

$$\text{Hence } f_\alpha := \begin{cases} F & \text{if } \alpha = (0, 0, \dots, 0) \\ 0 & \text{if } 1 \leq |\alpha| \leq m \end{cases} \quad \text{and } f_\alpha := \begin{cases} 0 & \text{if } |\alpha| \leq m-1 \\ G_\alpha & \text{if } |\alpha| = m \end{cases} \quad \text{with } G_\alpha = \partial_\alpha u$$

define two sets of data $(f_\alpha : |\alpha| \leq m)$ on the right-hand side of (5.1) for the same functional F . It is also known [49, Lemma 15.9, p.78] or [32, Thm.1. in 5.9, p.253] for $m = 1$ that $\|F\|_{H^{-m}(\Omega)}$ is the infimum of $\sqrt{\sum_{|\alpha| \leq m} \|f_\alpha\|_{L^2(\Omega)}^2}$ for all $(f_\alpha : |\alpha| \leq m)$ with (5.1).

It cannot be overemphasized that the preprocessing of the data set $(f_\alpha : |\alpha| \leq m)$ in (5.1) is a subtle issue that essentially depends on the way the right-hand side F is provided. For instance, point forces may apply, a right-hand side $f \in L^1(\Omega)$ for $m \geq 2$ and line integrals may be modeled for $m \geq 1$. The main idea will be explained for a model example with the right-hand side $F \in H^{-m}(\Omega)$, $\Omega \subset \mathbb{R}^2$ in (5.1) of the form

$$F(v) := \int_{\Omega} G : D^m v \, dx + \int_{\Omega} g v \, dx \quad \text{for all } v \in V = H_0^m(\Omega) \quad (5.2)$$

for given $G \in L^2(\Omega; \mathbb{R}^{2^m})$ and $g \in L^2(\Omega)$. Consider the vector space $H(\text{div}^m, \Omega; \mathbb{R}^{2^m})$ of all L^2 functions $Q \in L^2(\Omega; \mathbb{R}^{2^m})$ such that the m -th distributional divergence $\text{div}^m Q \in L^2(\Omega)$ is a Lebesgue function: $q \in L^2(\Omega)$ is called the m -th divergence of some $Q \in L^2(\Omega; \mathbb{R}^{2^m})$ if, for all $v \in D(\Omega)$, $\int_{\Omega} Q : D^m v \, dx = (-1)^m \int_{\Omega} q v \, dx$. This holds for all $v \in H_0^m(\Omega)$ as well (by density of $D(\Omega)$ in $H_0^m(\Omega)$) and one writes $q = (-1)^m \text{div}^m Q$. Hence any Q and any $v \in H_0^m(\Omega)$ satisfy

$$F(v) = \int_{\Omega} (G - Q) : D^m v \, dx + \int_{\Omega} (g + \text{div}^m Q) v \, dx \quad (5.3)$$

In other words, the data G and g of F in (5.2) can be modified to $G' := G - Q$ and $g' := g + \text{div}^m Q$ without changing the action of F on the continuous level. This data preprocessing may reduce the data oscillations as they arise e.g. in the a posteriori error control.

5.2 Point forces ($m = 2$)

The mechanical model often involves point loads. The Dirac delta functional $\delta_z \in H^{-2}(\Omega)$ is the evaluation of the continuous test function $u \in V = H_0^2(\Omega) \hookrightarrow C(\overline{\Omega})$ at a fixed point $z \in \overline{\Omega}$, called atom of the delta functional. Unlike for $m = 1$, $\delta_z : V \rightarrow \mathbb{R}$ is linear and bounded for $m \geq 2$. It is well known that (if $\langle \bullet, \bullet \rangle$ abbreviates the duality brackets in $H^{-2}(\Omega) \times H^2(\Omega)$ that extends the L^2 scalar product)

$$\langle \delta_z, v \rangle = v(z) = \frac{1}{2\pi} \int_{\Omega} \Delta v(x) \log |x - z| \, dx \quad \text{for all } v \in V = H_0^2(\Omega).$$

(cf. e.g. [33, Ex.1, page 82] or [32, Thm 1, page 23] for further details and proofs.) Hence δ_z is represented by $G(x) := \frac{1}{2\pi} \log |x - z|_{1_{2 \times 2}}$ for $x \in \Omega$ with the 2×2 unit matrix $1_{2 \times 2}$ and $g \equiv 0$. Since $G \in L^2(\Omega; \mathbb{S})$, the results of this survey apply to this case as well, but the practical computation may avoid integration with the weakly singular function $\log |x - z|$ and just apply δ_z on the point evaluation. The point force at $a \in A \subset \mathcal{V}(\Omega)$ does not contribute to the upper bound below in Proposition 5.1 when located at an interior vertex of the triangulation. The model example will suppose this for simplicity and because it can easily be arranged by appropriate triangulations in practice at least for a small number of point forces.

Before the next subsection continues with the corresponding model example, the effect of a general location shall at least be illuminated.

Example 5.1 (point force not located at a vertex). Suppose that there is a point force at $z \in \Omega$ that is not correlated to the triangulation \mathcal{T} . The original Morley FEM has to model δ_z also in case $z \in \partial T_+ \cap \partial T_- = E$ is shared by two triangles T_+ and T_- . The test function $v_M \in M(\mathcal{T})$ is (in general) discontinuous at z and

$$F_h := \mu \delta_{T_+,z} + (1 - \mu) \delta_{T_-,z} \quad (5.4)$$

applies with the piecewise point evaluation $\langle \delta_{T_{\pm},z}, v_{pw} \rangle_{H^2(\mathcal{T})} := (v_{pw}|_{T_{\pm}})(z)$ for all $v_{pw} \in H^2(\mathcal{T})$ and a convex coefficient $0 \leq \mu \leq 1$; $\mu = 1/2$ appears natural. The choice of μ does not affect the modified Morley FEM, but it will do so for the original scheme in general. The a priori error analysis of the best-approximation for the modified Morley FEM remains valid and the Proposition 5.1 is easily modified. A standard Bramble-Hilbert argument shows for $w \in H^2(T)$, $T \in \mathcal{T}$, that $w := v_M - Jv_M$ satisfies

$$\|w\|_{L^\infty(T)} \leq C_7 h_T |w|_{H^2(T)}$$

(In fact, the P_1 FEM interpolation error analysis shows in the above displayed inequality for a left-hand side $|w|_{H^1(T)}$ based on $w \in H^2(T)$ with $w(z) = 0$ for all vertices $z \in \mathcal{V}(T)$. The H^1 seminorm and the L^∞ norm do not scale in 2D [28] and this leads to the assertion.) It follows that the absolute value of

$$F_h(v_M) - F(Jv_M) = \mu (w|_{T_+})(z) + (1 - \mu) (w|_{T_-})(z)$$

is smaller than or equal to $\mu \|w\|_{L^\infty(T_+)} + (1 - \mu) \|w\|_{L^\infty(T_-)} \leq C_7 (\mu h_{T_+} + (1 - \mu) h_{T_-}) \|w\|_{pw}$. This leads to the additional data-oscillation term $\mu h_{T_+} + (1 - \mu) h_{T_-}$.

5.3 Model example

The smoother J in the right-hand side of the discrete problem (3.9) leads to an a priori best-approximation result for general data $F \in H^{-m}(\Omega)$. The a posteriori error control requires data evaluation in explicit residual-based a posteriori error estimates. Throughout this section the particular format of the right-hand side $F \in H^{-m}(\Omega)$ in (2.1) is assumed of the form

$$F(v) := \int_{\Omega} G : D^m v \, dx + \int_{\Omega} g v \, dx + \sum_{a \in A} \beta(a) v(a) \quad \text{for all } v \in V \quad (5.5)$$

for given $G \in L^2(\Omega; \mathbb{R}^{(n^m)})$, $g \in L^2(\Omega)$, A is a finite set of interior vertices, and $a \in A \subset \mathcal{V}(\Omega)$ is the position and $\beta(a)$ the strength of the point force. The *natural right-hand side* $F_h := \widehat{F}|_{V_{\text{nc}}} \in V_{\text{nc}}^*$ in the discrete problem for (5.5) reads, for all $\widehat{v} = v + v_{\text{nc}} \in \widehat{V} \equiv V + V_{\text{nc}}$,

$$\widehat{F}(\widehat{v}) := \int_{\Omega} G : D_{\text{pw}}^m \widehat{v} \, dx + \int_{\Omega} g \widehat{v} \, dx + \sum_{a \in A} \beta(a) \widehat{v}(a). \quad (5.6)$$

5.4 Comparison

The two schemes (3.9) and (2.2) for $F_h = \widehat{F}|_{V_{\text{nc}}}$ from (5.6) are in fact different. While the modified scheme (3.9) with right-hand side $F \circ J$ enjoys a best-approximation property, the original version for the right-hand side $\widehat{F}|_{V_{\text{nc}}}$ from (5.6) does so only up to data-oscillations.

Proposition 5.1. *Given F and \widehat{F} in (5.5)-(5.6), let $u_{\text{nc}}^{\text{org}} \in V_{\text{nc}}$ solve (2.2) and let $u_{\text{nc}}^{\text{mod}}$ solve (3.9). Then $\Lambda_0^{-1} \|u_{\text{nc}}^{\text{org}} - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}} \leq \|G - \Pi_0 G\| + \kappa_m \text{osc}_m(g, \mathcal{T})$.*

Proof. Since $\|u_{\text{nc}}^{\text{org}} - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}}^2 = \widehat{F}((1 - J)(u_{\text{nc}}^{\text{org}} - u_{\text{nc}}^{\text{mod}})) \leq \|\widehat{F} \circ (1 - J)\|_{V_{\text{nc}}^*} \|u_{\text{nc}}^{\text{org}} - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}}$, it remains to analyse $\|\widehat{F} \circ (1 - J)\|_{V_{\text{nc}}^*}$. For any $v_{\text{nc}} \in V_{\text{nc}}$ with $\|v_{\text{nc}}\|_{\text{pw}} = 1$, the definition (5.6) shows

$$\begin{aligned} \widehat{F} \circ (1 - J)(v_{\text{nc}}) &= \int_{\Omega} G : D_{\text{pw}}^2(v_{\text{nc}} - Jv_{\text{nc}}) \, dx + \int_{\Omega} g(v_{\text{nc}} - Jv_{\text{nc}}) \, dx \\ &= \int_{\Omega} (G - \Pi_0 G) : D_{\text{pw}}^2(v_{\text{nc}} - Jv_{\text{nc}}) \, dx + \int_{\Omega} (g - \Pi_m g)(v_{\text{nc}} - Jv_{\text{nc}}) \, dx \end{aligned}$$

with the assumptions that $\Pi_0 D^2 Jv_{\text{nc}} = D_{\text{pw}}^2 v_{\text{nc}}$ and $\Pi_m Jv_{\text{nc}} = v_{\text{nc}}$ for $m = 1$ and $m = 2$ in the last step that holds for CR and Morley FEM (see Sections 2.3.1 and 2.3.2). This and $\|h_{\mathcal{T}}^{-m}(v_{\text{nc}} - Jv_{\text{nc}})\| \leq \kappa_m \|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}}$ from (2.4) and (2.7) prove

$$\widehat{F} \circ (1 - J)(v_{\text{nc}}) \leq (\|G - \Pi_0 G\| + \kappa_m \text{osc}_m(g, \mathcal{T})) \|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}}.$$

The definition of Λ_0 in (3.10) shows $\|v_{\text{nc}} - Jv_{\text{nc}}\|_{\text{pw}} \leq \Lambda_0 \|v_{\text{nc}}\|_{\text{pw}} = \Lambda_0$. The combination of the preceding estimates proves $\widehat{F} \circ (1 - J)(v_{\text{nc}}) \leq \Lambda_0 (\|G - \Pi_0 G\| + \kappa_m \text{osc}_m(g, \mathcal{T}))$. This concludes the proof. \square

The data-oscillation term $\text{osc}_m(g, \mathcal{T}) = o(h_{\max})$ can be of higher-order if $g \in H^1(\mathcal{T})$ is piecewise smooth for $m \geq 2$, while this is less clear for the data-oscillation $\|G - \Pi_0 G\|$.

The subsequent example shows that (a) the original scheme can be optimal while the modified one is not, (b) the bound in the Proposition 5.1 is sharp in general (at least if $g \equiv 0$). The methodology is similar to the analysis in Theorem 3.3; in fact (c) is another proof of the optimality of $C_{\text{qo}} = 1 + \Lambda_0^2$ (recall $\Lambda_0 > 0$ throughout the paper).

Example 5.2. There exist $G := D^m Jz_{\text{nc}} \in D^m V_{\text{c}}$ and $g := 0$ in (5.5) for some $z_{\text{nc}} \in V_{\text{nc}}$ such that the exact solution $u \in V$ to (2.1), the discrete solution $u_{\text{nc}}^{\text{org}}$ to the original scheme (2.2) and the discrete solution $u_{\text{nc}}^{\text{mod}} \in V_{\text{nc}}$ to (3.9) with F and $F_h = \widehat{F}|_{V_h}$ from (5.5)-(5.6), satisfy (a) $u_{\text{nc}}^{\text{org}} = I_{\text{nc}} u$, (b) $\|u_{\text{nc}}^{\text{org}} - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}} = \Lambda_0^2$, and (c) $\|u - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}}^2 = \|u - u_{\text{nc}}^{\text{org}}\|_{\text{pw}}^2 + \Lambda_0^4 = \Lambda_0^2(1 + \Lambda_0^2)$.

Proof of (a). Given V_{nc} and J from Subsections 2.2 and 2.4, there exists $z_{\text{nc}} \in V_{\text{nc}}$ with $\|z_{\text{nc}}\|_{\text{pw}} = 1$ and $\|z_{\text{nc}} - Jz_{\text{nc}}\|_{\text{pw}} = \Lambda_0 \|z_{\text{nc}}\|_{\text{pw}} = \Lambda_0$. Let $G := D^m Jz_{\text{nc}}$ and recall $\Pi_0 G = D_{\text{pw}}^m I_{\text{nc}} Jz_{\text{nc}} = D_{\text{pw}}^m z_{\text{nc}}$ with $\|\Pi_0 G\| = \|z_{\text{nc}}\|_{\text{pw}} = 1$ and $\|G - \Pi_0 G\| = \|z_{\text{nc}} - Jz_{\text{nc}}\|_{\text{pw}} = \Lambda_0$. For all $v_{\text{nc}} \in V_{\text{nc}}$, (2.3) shows

$$a_{\text{pw}}(z_{\text{nc}}, v_{\text{nc}}) = a_{\text{pw}}(Jz_{\text{nc}}, v_{\text{nc}}) = \int_{\Omega} G : D_{\text{pw}}^m v_{\text{nc}} \, dx$$

with $G := D^m Jz_{\text{nc}}$ in the last step. Hence $u_{\text{nc}}^{\text{org}} = z_{\text{nc}}$. Since $g = 0$, for $u = Jz_{\text{nc}}$, (5.5) shows

$$a(u, v) = a(Jz_{\text{nc}}, v) = \int_{\Omega} G : D^m v \, dx \text{ for all } v \in V.$$

This and $I_{\text{nc}} u = I_{\text{nc}} Jz_{\text{nc}} = z_{\text{nc}} = u_{\text{nc}}^{\text{org}}$ proves the equality in (a).

Proof of (b). Recall

$$\begin{aligned} \Lambda_0^2 &= \|z_{\text{nc}} - Jz_{\text{nc}}\|_{\text{pw}}^2 = a_{\text{pw}}(Jz_{\text{nc}} - z_{\text{nc}}, Jz_{\text{nc}}) \\ &= a(u, u) - a_{\text{pw}}(z_{\text{nc}}, z_{\text{nc}}) = \int_{\Omega} G : D^m Jz_{\text{nc}} \, dx - \int_{\Omega} G : D_{\text{pw}}^m z_{\text{nc}} \, dx \\ &= a_{\text{pw}}(u_{\text{nc}}^{\text{mod}} - u_{\text{nc}}^{\text{org}}, z_{\text{nc}}). \end{aligned}$$

A Cauchy inequality in the semi-scalar inner product a_{pw} with $\|z_{\text{nc}}\|_{\text{pw}} = 1$ leads to

$$\Lambda_0^2 = a_{\text{pw}}(u_{\text{nc}}^{\text{mod}} - u_{\text{nc}}^{\text{org}}, z_{\text{nc}}) \leq \|u_{\text{nc}}^{\text{mod}} - u_{\text{nc}}^{\text{org}}\|_{\text{pw}}.$$

For $g = 0$ and $\|G - \Pi_0 G\| = \Lambda_0$ from above, the Proposition 5.1 shows $\|u_{\text{nc}}^{\text{mod}} - u_{\text{nc}}^{\text{org}}\|_{\text{pw}} \leq \Lambda_0^2$. This concludes the proof of (b). \square

Proof of (c). The assertion (c) follows from the above and the Pythagoras theorem

$$\|u - u_{\text{nc}}^{\text{mod}}\|_{\text{pw}}^2 = \|Jz_{\text{nc}} - z_{\text{nc}}\|_{\text{pw}}^2 + \|u_{\text{nc}}^{\text{mod}} - u_{\text{nc}}^{\text{org}}\|_{\text{pw}}^2. \quad \square$$

5.5 Counterexamples for best-approximation

The best-approximation property for the original nonconforming FEM with the natural right-hand side fails in general: The data oscillation terms cannot be removed. Suppose Ω a simply-connected domain throughout this subsection.

5.5.1 Counterexample to best-approximation of CRFEM for $m = 1$

Suppose $|\mathcal{T}| \geq 2$ so that there exists some $\beta_{\text{CR}} \in \text{CR}^1(\mathcal{T}) \setminus S^1(\mathcal{T})$. Since the quotient space $S^1(\mathcal{T})/\mathbb{R}$ is a Hilbert space with scalar product $a_{\text{pw}}(\bullet, \bullet)$, the linear functional $v_{\text{c}} \mapsto a_{\text{pw}}(\beta_{\text{CR}}, v_{\text{c}}) \in (S^1(\mathcal{T}) \setminus \mathbb{R})^*$ has a Riesz representation β_{c} so that $b_{\text{CR}} := \beta_{\text{CR}} - \beta_{\text{c}} \in \text{CR}^1(\mathcal{T}) \setminus S^1(\mathcal{T})$ satisfies $a_{\text{pw}}(b_{\text{CR}}, v_{\text{c}}) = 0$ for all $v_{\text{c}} \in S^1(\mathcal{T})$. Without loss of generality, suppose $\|b_{\text{CR}}\|_{\text{pw}} = 1$. Since $\text{Curl}_{\text{pw}} b_{\text{CR}}$ is a rotated gradient $\nabla_{\text{pw}} b_{\text{CR}}$ etc. in 2D, the L^2 orthogonality $\text{Curl}_{\text{pw}} b_{\text{CR}} \perp \text{Curl } S^1(\mathcal{T})$ in $L^2(\Omega; \mathbb{R}^2)$ follows. The companion operators in this paper obey homogeneous boundary conditions but are naturally extended to functions without any restriction. This leads to some $b := Jb_{\text{CR}} \in H^1(\Omega)$ with $b_{\text{CR}} := I_{\text{CR}} b$. Given those functions, define the data $G := \text{Curl } b \perp \nabla H_0^1(\Omega)$ and

$g \equiv 0$ in (5.5)-(5.6) for F and \widehat{F} . Then the exact solution $u \equiv 0$ and hence the best approximation provides $u_{\text{CR}}^{\text{mod}} \equiv 0$ as well. The investigation of the original CRFEM solution $u_{\text{nc}}^{\text{org}} \in \text{CR}_0^1(\mathcal{T})$ to (2.2) recalls the discrete Helmholtz decomposition from Subsection 2.3.1: $P_0(\mathcal{T}; \mathbb{R}^2) = \nabla_{\text{pw}} \text{CR}_0^1(\mathcal{T}) \oplus \text{Curl}(S^1(\mathcal{T}) \setminus \mathbb{R})$. The L^2 orthogonality of $\Pi_0 G = \Pi_0 \text{Curl } b = \text{Curl}_{\text{pw}} I_{\text{CR}} b = \text{Curl}_{\text{pw}} b_{\text{CR}} \perp \text{Curl } S^1(\mathcal{T})$ in $L^2(\Omega; \mathbb{R}^2)$ therefore proves $\Pi_0 G = \nabla_{\text{pw}} u_{\text{CR}}$ for some $u_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$. In fact, u_{CR} solves the original CRFEM because of

$$a_{\text{pw}}(u_{\text{CR}}, v_{\text{CR}}) = \int_{\Omega} \Pi_0 G \cdot \nabla_{\text{pw}} v_{\text{CR}} \, dx = \int_{\Omega} G \cdot \nabla_{\text{pw}} v_{\text{CR}} \, dx$$

for all $v_{\text{CR}} \in \text{CR}_0^1(\mathcal{T})$. That is, $\|u_{\text{CR}}^{\text{org}}\|_{\text{pw}} = \|u_{\text{CR}}\|_{\text{pw}} = \|\Pi_0 G\| = \|I_{\text{CR}} b\|_{\text{pw}} = \|b_{\text{CR}}\|_{\text{pw}} = 1$. This contradicts the best-approximation property.

5.5.2 Counterexample to best-approximation of Morley FEM for $m = 2$

Suppose $|\mathcal{T}| \geq 2$ and recall $S^1(\mathcal{T}) \subsetneq \text{CR}_0^1(\mathcal{T})$ for the existence of $\beta_{\text{CR}} \in \text{CR}^1(\mathcal{T}; \mathbb{R}^2) \setminus S^1(\mathcal{T}; \mathbb{R}^2)$. The linear Green strain $\varepsilon := \text{sym } D$ from linear elasticity allows for a Korn inequality with the rigid body motions

$$\text{RH}(\Omega) := \{\gamma \in P_1(\Omega; \mathbb{R}^2) : D\gamma + D\gamma^T = 0\}$$

(a linear subspace of dimension 3 of the form $\gamma(x) = (a, b) + c(x_2, -x_1)$ for all $x \in \Omega$). The semi-scalar product $(\varepsilon(\bullet), \varepsilon(\bullet))_{L^2(\Omega)}$ in the vector space $S^1(\mathcal{T}; \mathbb{R}^2)/\text{RH}(\Omega)$ is a scalar product (from the Korn inequality). The Riesz representation $\beta_c \in S^1(\mathcal{T}; \mathbb{R}^2)/\text{RH}(\Omega)$ of the functional $v_c \mapsto (\varepsilon_{\text{pw}}(\beta_{\text{CR}}), \varepsilon(v_c))$ in $(S^1(\mathcal{T}; \mathbb{R}^2)/\text{RH}(\Omega))^*$ provides $b_{\text{CR}} := \beta_{\text{CR}} - \beta_c \in \text{CR}^1(\mathcal{T}; \mathbb{R}^2) \setminus S^1(\mathcal{T}; \mathbb{R}^2)$ with $\varepsilon_{\text{pw}}(b_{\text{CR}}) \perp \varepsilon(S^1(\mathcal{T}; \mathbb{R}^2))$ in $L^2(\Omega; \mathbb{S})$. Without loss of generality suppose $\|\varepsilon(b_{\text{CR}})\| = 1$. Recall the companion operator $J : \text{CR}^1(\mathcal{T}) \rightarrow H^1(\Omega)$ from the previous example and apply $Jb_{\text{CR}} = (Jb_{\text{CR}}(1), Jb_{\text{CR}}(2)) \in H^1(\Omega; \mathbb{R}^2)$ componentwise to $b_{\text{CR}} \in \text{CR}^1(\mathcal{T}; \mathbb{R}^2)$. Then let $G := \text{sym Curl}(Jb_{\text{CR}}(2), -Jb_{\text{CR}}(1))$ and $g \equiv 0$. Since $G \perp D^2 H_0^2(\Omega)$, the solutions $u = 0 = u_{\text{M}}^{\text{mod}}$ coincide. On the other hand, for all $\phi, \psi \in H^1(T; \mathbb{R}^2)$,

$$(\text{sym Curl}(\phi_2, -\phi_1)) : (\text{sym Curl}(\psi_2, -\psi_1)) = \varepsilon(\phi) : \varepsilon(\psi) \text{ a.e. in } T.$$

This relation shows

$$\begin{aligned} \Pi_0 G &= \text{sym Curl}_{\text{pw}} I_{\text{CR}}(Jb_{\text{CR}}(2), -Jb_{\text{CR}}(1)) = \text{sym Curl}_{\text{pw}}(b_{\text{CR}}(2), -b_{\text{CR}}(1)) \\ &\perp \text{sym Curl } S^1(\mathcal{T}; \mathbb{R}^2) \text{ in } L^2(\Omega; \mathbb{S}). \end{aligned}$$

The discrete Helmholtz decomposition $P_0(\mathcal{T}; \mathbb{S}) = D_{\text{pw}}^2 \mathbf{M}(\mathcal{T}) \oplus \text{sym Curl } S^1(\mathcal{T}; \mathbb{R}^2/\mathbb{R}^2)$ from Subsection 2.3.2 implies $\Pi_0 G = D_{\text{pw}}^2 u_{\text{M}}$ for some $u_{\text{M}} \in \mathbf{M}(\mathcal{T})$. Then $u_{\text{M}} = u_{\text{M}}^{\text{org}}$ and $\|u_{\text{M}}^{\text{org}}\|_{\text{pw}} = \|\Pi_0 G\| = \|b_{\text{CR}}\|_{L^2(T)} = 1$. This contradicts the best-approximation property.

6 A posteriori analysis

Unlike conforming schemes where the guaranteed error bound involves constants from approximation in the error estimates of a quasi-interpolation operator, the constants in the

nonconforming schemes appear small and are related to the simplicial element domain (and not the shape of nodal patches). Hence the resulting a posteriori error estimators can indeed be employed directly in the a posteriori error control.

6.1 Reliable error estimates

The *natural right-hand side* in the discrete problem is $F_h := \widehat{F}|_{V_{\text{nc}}} \in V_{\text{nc}}^*$ with \widehat{F} from (5.6).

Theorem 6.1 (a posteriori estimate for $F_h = \widehat{F}|_{V_{\text{nc}}}$). *Suppose the L^2 orthogonality in (2.6). Let $u \in V$ solve (2.1) and let $u_{\text{nc}} \in V_{\text{nc}}$ solve (2.2) with right-hand side $F_h := \widehat{F}|_{V_{\text{nc}}}$ from (5.6). Then*

$$\begin{aligned} (a) \quad & \|u - Ju_{\text{nc}}\|^2 + \|I_{\text{nc}}u - u_{\text{nc}}\|_{\text{pw}}^2 \leq (\|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\| + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}})^2, \\ (b) \quad & \|u - u_{\text{nc}}\|_{\text{pw}}^2 + \|I_{\text{nc}}u - u_{\text{nc}}\|_{\text{pw}}^2 \leq (\|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\| + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}})^2 \\ & \quad + 2\widehat{F}(u_{\text{nc}} - Ju_{\text{nc}}). \end{aligned}$$

The former analysis in [5, 13, 41] is based on Helmholtz decompositions and worked out for simply-connected domains. This is problematic and the three-dimensional application in [41] requires some 3D Helmholtz decompositions for a domain that is simply-connected and has a boundary that is connected and a discrete counterpart is unknown. The present analysis is rather simple and can immediately be extended to any bounded polyhedral Lipschitz domain with the help of the companion operator J in 3D from [25].

Proof of (a). For $e := u - Ju_{\text{nc}}$, elementary algebra and (2.1)-(2.3) lead to

$$\begin{aligned} \|e\|^2 &= a(u, e) + a_{\text{pw}}(u_{\text{nc}} - Ju_{\text{nc}}, e) - a_{\text{pw}}(u_{\text{nc}}, I_{\text{nc}}e) \\ &= \widehat{F}(e - I_{\text{nc}}e) + a_{\text{pw}}(u_{\text{nc}} - Ju_{\text{nc}}, e - I_{\text{nc}}e). \end{aligned} \quad (6.1)$$

This, the definition of \widehat{F} in (5.6), (2.6), and a Cauchy inequality show

$$\begin{aligned} \|e\|^2 &\leq \int_{\Omega} (G - \Pi_0 G) : D^m(e - I_{\text{nc}}e) \, dx + \int_{\Omega} g(e - I_{\text{nc}}e) \, dx \\ &\quad + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}} \|e - I_{\text{nc}}e\|_{\text{pw}}. \end{aligned}$$

This, (weighted) Cauchy inequalities and $\|h_{\mathcal{T}}^{-m}(e - I_{\text{nc}}e)\| \leq \kappa_m \|e - I_{\text{nc}}e\|_{\text{pw}}$ from (2.4) imply

$$\|e\|^2 \leq (\|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\| + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}}) \|e - I_{\text{nc}}e\|_{\text{pw}}.$$

The Pythagoras theorem (2.5) shows $\|e\|^2 = \|e - I_{\text{nc}}e\|_{\text{pw}}^2 + \|I_{\text{nc}}e\|_{\text{pw}}^2$. This and Young's inequality conclude the proof. \square

Proof of (b). Elementary algebra and $a_{\text{pw}}(u_{\text{nc}}, u - I_{\text{nc}}u) = 0 = a_{\text{pw}}(I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}})$ from (2.3) result in

$$\|u - u_{\text{nc}}\|_{\text{pw}}^2 = a_{\text{pw}}(u - I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}}) + a_{\text{pw}}(u, u - Ju_{\text{nc}}) - a_{\text{pw}}(u_{\text{nc}}, I_{\text{nc}}u - u_{\text{nc}}).$$

This, the Pythagoras theorem (2.5), a Cauchy inequality, the continuous problem (2.1) with the test function $u - Ju_{\text{nc}}$, and (2.2) with the test function $I_{\text{nc}}u - u_{\text{nc}}$ lead to

$$\begin{aligned} \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + \|I_{\text{nc}}u - u_{\text{nc}}\|_{\text{pw}}^2 &\leq \|u - I_{\text{nc}}u\|_{\text{pw}} \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}} \\ &\quad + \widehat{F}(u - I_{\text{nc}}u) + \widehat{F}(u_{\text{nc}} - Ju_{\text{nc}}). \end{aligned} \quad (6.2)$$

The definition of \widehat{F} in (5.6), (2.6), (weighted) Cauchy inequalities, and (2.4) show

$$\begin{aligned} \widehat{F}(u - I_{\text{nc}}u) &= \int_{\Omega} (G - \Pi_0 G) : D^m(u - I_{\text{nc}}u) \, dx + \int_{\Omega} g(u - I_{\text{nc}}u) \, dx \\ &\leq (\|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\|) \|u - I_{\text{nc}}u\|_{\text{pw}}. \end{aligned}$$

The last two displayed estimates and Young's inequality conclude the proof. \square

Remark 6.1. The term $\widehat{F}(u_{\text{nc}} - Ju_{\text{nc}})$ can be computed and so (b) is an a posteriori error estimate. Moreover, (2.9) and (weighted) Cauchy inequalities show

$$\widehat{F}(u_{\text{nc}} - Ju_{\text{nc}}) \leq (\|G - \Pi_0 G\| + \kappa_m \text{osc}_m(g, \mathcal{T})) \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}}.$$

The combination with (6.2) leads to (a) up to the extra factor 2.

The *modified choice of the right-hand side* reads $F_h = F \circ J$ for the right-inverse J of I_{nc} from Section 2.4. The discrete problem for this choice of F_h from (3.9) leads to a (new) a posteriori estimate in Theorem 6.2.

Theorem 6.2 (a posteriori estimate for $F_h = F \circ J$). *Suppose (5.5) and the L^2 orthogonality in (2.6). Let $u \in V$ solve (2.1) and let $u_{\text{nc}} \in V_{\text{nc}}$ solve (3.9) with right-hand side $F_h := F \circ J$. Then*

$$\begin{aligned} (a) \quad \|u - Ju_{\text{nc}}\| &\leq \sqrt{1 + \Lambda_0^2} \|G - \Pi_0 G\| + \sqrt{(\kappa_m \|h_{\mathcal{T}}^m g\| + \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}})^2 + \kappa_m^2 \Lambda_0^2 \text{osc}_m^2(g, \mathcal{T})}, \\ (b) \quad \|u - u_{\text{nc}}\|_{\text{pw}} &\leq \sqrt{2(\|u_{\text{nc}} - Ju_{\text{nc}}\|^2 + \text{apx}(F)^2)} \end{aligned}$$

with $\text{apx}(F) := (1 + \Lambda_J) \|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\| + \kappa_m \Lambda_J \text{osc}_m(g, \mathcal{T})$ and Λ_0 (resp. Λ_J) from (3.10) (resp. (2.8)).

Proof of (a). Elementary algebra with $e := u - Ju_{\text{nc}} \in V$, (2.1), $a_{\text{pw}}(u_{\text{nc}}, e - I_{\text{nc}}e) = 0$ from (2.3), $a_{\text{pw}}(u_{\text{nc}} - Ju_{\text{nc}}, I_{\text{nc}}e) = 0$ from (2.7), and (3.9) lead to

$$\begin{aligned} \|e\|^2 &= a(u, e) + a_{\text{pw}}(u_{\text{nc}} - Ju_{\text{nc}}, e) - a_{\text{pw}}(u_{\text{nc}}, e) \\ &= F(e - JI_{\text{nc}}e) + a_{\text{pw}}(u_{\text{nc}} - Ju_{\text{nc}}, e - I_{\text{nc}}e). \end{aligned} \quad (6.3)$$

For any $v \in V$, (2.9) and (2.6) imply $\int_{\Omega} \Pi_0 G : D^m(v - JI_{\text{nc}}v) \, dx = 0$. This, the definition (5.5), and (2.9) result in

$$\begin{aligned} F(v - JI_{\text{nc}}v) &= \int_{\Omega} (G - \Pi_0 G) : D^m(v - JI_{\text{nc}}v) \, dx + \int_{\Omega} g(v - I_{\text{nc}}v) \, dx \\ &\quad + \int_{\Omega} (g - \Pi_0 g)((1 - J)I_{\text{nc}}v) \, dx \\ &\leq \|G - \Pi_0 G\| \|v - JI_{\text{nc}}v\|_{\text{pw}} + \|h_{\mathcal{T}}^m g\| \|h_{\mathcal{T}}^{-m}(v - I_{\text{nc}}v)\| \\ &\quad + \text{osc}_m(g, \mathcal{T}) \|h_{\mathcal{T}}^{-m}(1 - J)I_{\text{nc}}v\| \end{aligned} \quad (6.4)$$

with (weighted) Cauchy inequalities in the last step. For $\Lambda_0 = \|1 - J\|_{L(V_{\text{nc}}, \widehat{V})}$ from (3.10), a triangle inequality and $\|e\|_{\text{pw}}^2 = \|e - I_{\text{nc}}e\|_{\text{pw}}^2 + \|I_{\text{nc}}e\|_{\text{pw}}^2$ from (2.5) prove

$$\|e - JI_{\text{nc}}e\|_{\text{pw}} \leq \|e - I_{\text{nc}}e\|_{\text{pw}} + \|(1 - J)I_{\text{nc}}e\|_{\text{pw}} \leq \sqrt{1 + \Lambda_0^2} \|e\|_{\text{pw}}.$$

Recall from (2.4) and (2.5) that

$$\|h_{\mathcal{T}}^{-m}(e - I_{\text{nc}}e)\| \leq \kappa_m \|e - I_{\text{nc}}e\|_{\text{pw}} \text{ and } \|h_{\mathcal{T}}^{-m}(1 - J)I_{\text{nc}}e\| \leq \kappa_m \Lambda_0 \|I_{\text{nc}}e\|_{\text{pw}}.$$

Substitute the last three displayed inequalities in (6.4) with $v := e$ and then substitute the resulting estimate in (6.3). Apply a Cauchy inequality for the second term in the right-hand side of (6.3) and use (2.5) to conclude the proof of (a). \square

Proof of (b). Elementary algebra with $a_{\text{pw}}(u_{\text{nc}}, u - I_{\text{nc}}u) = 0 = a_{\text{pw}}(I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}})$ from (2.3) and the continuous problem (2.1) with test function $u - Ju_{\text{nc}}$ show

$$\|u - u_{\text{nc}}\|_{\text{pw}}^2 = F(u - Ju_{\text{nc}}) + a_{\text{pw}}(u - I_{\text{nc}}u, Ju_{\text{nc}} - u_{\text{nc}}) - a_{\text{pw}}(u_{\text{nc}}, I_{\text{nc}}u - u_{\text{nc}}).$$

The Pythagoras theorem (2.5), a Cauchy inequality, and (3.9) reveal in the last identity that

$$\|u - I_{\text{nc}}u\|_{\text{pw}}^2 + \|I_{\text{nc}}u - u_{\text{nc}}\|_{\text{pw}}^2 \leq \|u - I_{\text{nc}}u\|_{\text{pw}} \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}} + F(u - JI_{\text{nc}}u). \quad (6.5)$$

The choice of $v := u$ in (6.4), a triangle inequality, (2.4), and twice (2.8) for $\|I_{\text{nc}}u - JI_{\text{nc}}u\|_{\text{pw}} \leq \Lambda_J \|u - I_{\text{nc}}u\|_{\text{pw}}$ lead to

$$\begin{aligned} F(u - JI_{\text{nc}}u) &\leq \|u - I_{\text{nc}}u\|_{\text{pw}} ((1 + \Lambda_J) \|G - \Pi_0 G\| + \kappa_m \|h_{\mathcal{T}}^m g\| \\ &\quad + \kappa_m \Lambda_J \text{osc}_m(g, \mathcal{T})) = \text{apx}(F) \|u - I_{\text{nc}}u\|_{\text{pw}}. \end{aligned} \quad (6.6)$$

A substitution of (6.6) in (6.5) and Young's inequality show

$$\frac{1}{2} \|u - I_{\text{nc}}u\|_{\text{pw}}^2 + \|I_{\text{nc}}u - u_{\text{nc}}\|_{\text{pw}}^2 \leq \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}}^2 + \text{apx}(F)^2.$$

This concludes the proof. \square

Remark 6.2 (generalization). The model scenario of the right-hand side in (5.5) solely coincides with the lowest and highest order terms in (5.1) because oscillations occur in those terms because of (2.3) and (2.9). The generalization of Theorems 6.1-6.2 to the general situation (5.1) is straightforward and the extra terms read $\sum_{1 \leq |\alpha| \leq m-1} \|h_{\mathcal{T}}^{m-|\alpha|} f_{\alpha}\|_{L^2(\Omega)}$.

Remark 6.3 (a posteriori estimates for (5.3)). For the right-hand side in (5.3), the discretisation will change because \widehat{F} changes to

$$\widehat{F}(\widehat{v}) := \int_{\Omega} (G - Q) : D_{\text{pw}}^m \widehat{v} \, dx + \int_{\Omega} (g + \text{div}^m Q) \widehat{v} \, dx \text{ for all } \widehat{v} = v + v_{\text{nc}} \in \widehat{V} \equiv V + V_{\text{nc}}.$$

It may be seen as an advantage of the modified scheme (3.9) that u_{nc} is not affected by a change of (G, g) to $(G - Q, g + \text{div}^m Q)$ for any Q . The a posteriori error control involves the terms $\|G - \Pi_0 G\| + \|h_{\mathcal{T}}^m g\|$ and this motivates the minimisation

$$\min_{Q \in H(\text{div}^m, Q)} (\|(1 - \Pi_0)(G - Q)\|^m + \kappa_m \|h_{\mathcal{T}}^m (g + \text{div}^m Q)\|). \quad (6.7)$$

The modified data $(G', g') = (G - Q, g + \operatorname{div}^m Q)$ for a minimizer Q leads to optimal a posteriori error control in Theorem 6.2 when the upper bound is simplified to

$$\frac{1}{2} \|u - u_{\text{nc}}\|_{\text{pw}}^2 \leq \|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}}^2 + 2(1 + \Lambda_J)^2 (\|G' - \Pi_0 G'\| + \kappa_m \|h_{\mathcal{T}}^m g'\|)^2. \quad (6.8)$$

The situation is less clear in Theorem 6.1 because u_{nc} may be affected by a change of \widehat{F} and this may affect $\|u_{\text{nc}} - Ju_{\text{nc}}\|_{\text{pw}}$. However, it is unrealistic to solve the minimisation problem (6.7) exactly for general G and g . Nevertheless, a preprocessing of the data G and g can reduce the data approximation terms in (guaranteed) upper error bounds. In return, the data selected can be poor as well and ruin the efficiency of the reliable error estimates in Theorems 6.1 and 6.2.

6.2 Efficiency up to data-oscillations

The efficiency of the term $\|h_{\mathcal{T}}^m g\|_{L^2(\Omega)}$ up to data-oscillations is established next.

Theorem 6.3 (efficiency up to data-oscillations). *Let u solve (2.1) with right-hand side (5.5) for $G, g \in L^2(\Omega)$. Then $\|h_{\mathcal{T}}^m g\|_{L^2(\Omega)} \lesssim \|u - I_{\text{nc}} u\|_{\text{pw}} + \operatorname{osc}_m(g, \mathcal{T}) + \|G - \Pi_0 G\|$.*

Proof. The efficiency holds in a local form for each triangle $T \in \mathcal{T}$. Let $\varphi_z \in S^1(\mathcal{T})$ be the P_1 nodal basis function associated to a vertex $z \in \mathcal{V}(T) = \{a, b, c\}$ of the triangle T , i.e., $\varphi_z|_T$ is a barycentric coordinate of z in T and let $b_T := 27\varphi_a\varphi_b\varphi_c \in P_3(T) \cap H_0^1(T)$ be the cubic-bubble function. Then $b_T^m \in H_0^m(T) \subset V$ satisfies $0 \leq b_T^m \leq 1$ and inverse estimates $\|b_T^m\| \lesssim h_T^{-m} \|b_T\|$. Abbreviate $g_m := \Pi_m g|_T \in P_m(T)$ and observe

$$\|h_T^m g\|_{L^2(T)} \leq \|h_T^m g_m\|_{L^2(T)} + \operatorname{osc}_m(g, T) \quad (6.9)$$

from the definition of $\operatorname{osc}_m(g, T)$ and a triangle inequality. An inverse estimate $\|g_m\|_{L^2(T)} \leq C_8 \|b_T^{m/2} g_m\|_{L^2(T)}$ follows from the equivalence of the norms $\|b_T \bullet\|_{L^2(T)} \approx \|\bullet\|_{L^2(T)}$ in $P_m(T)$. (A scaling argument reveals that the equivalence constant does not depend on the size or shape of the triangles and exclusively depends on m .) The inverse estimate shows

$$\begin{aligned} C_8^{-2} h_T^{2m} \|g_m\|_{L^2(T)}^2 &\leq h_T^{2m} \|b_T^{m/2} g_m\|_{L^2(T)}^2 = h_T^{2m} \int_T b_T^m g_m (g_m - g) \, dx + h_T^{2m} \int_T b_T^m g_m g \, dx \\ &\leq \operatorname{osc}_m(g, \mathcal{T}) h_T^m \|b_T^m g_m\|_{L^2(T)} + h_T^m \int_T g v_T \, dx \end{aligned} \quad (6.10)$$

with a Cauchy inequality and the abbreviation $v_T := h_T^m b_T^m g_m \in P_{4m}(T) \cap H_0^2(T) \subset V$ in the last step. Since $a(u, v_T) = \int_T g v_T \, dx + \int_T G : D^m v_T \, dx$ (from (2.1) for the test function $v_T \in V$), the crucial term in (6.10) is equal to

$$h_T^m \int_T g v_T \, dx = h_T^m \int_T (D^2 u - G) : D^m v_T \, dx. \quad (6.11)$$

The bubble-function methodology due to [54] observes at this point that $\Pi_0 D^m v_T = 0$ from an integration by parts with $v_T \in H_0^m(T)$, whence $D^m v_T = 0$ on ∂T , and so $\int_T D^m v_T \, dx = \int_{\partial T} (D^{m-1} v_T) v_T \, ds = 0$. Consequently,

$$\begin{aligned} \int_T (D^m u - G) : D^m v_T \, dx &= \int_T D^m v_T : (1 - \Pi_0)(D^m u - G) \, dx \\ &\leq (\|u - I_{\text{nc}} u\|_{H^2(T)} + \|G - \Pi_0 G\|_{L^2(T)}) \|D^m v_T\|_{L^2(T)}. \end{aligned}$$

A standard inverse estimate $\|D^m v_T\|_{L^2(T)} \lesssim h_T^{-m} \|v_T\|_{L^2(T)}$ for the polynomial $v_T \in P_{4m}(T)$ depends on m and on the shape-regularity of the triangle T . Note that $h_T^{-m} \|v_T\|_{L^2(T)} = \|b_T^m g_m\|_{L^2(T)} \leq \|g_m\|_{L^2(T)}$ follows from the definition of $v_T = h_T^m b_T^m g_m$ and $0 \leq b_T \leq 1$. The aforementioned estimates lead in (6.11) to

$$h_T^m \int_T g v_T \, dx \leq C_9 (|u - I_{\text{nc}} u|_{H^2(T)} + \|G - \Pi_0 G\|_{L^2(T)}) h_T^m \|g_m\|_{L^2(T)}$$

for some generic constant $C_9 \approx 1$ that depends on the shape of T and m . The combination of the previous estimate with (6.10) proves

$$h_T^{2m} \|g_m\|_{L^2(T)}^2 \leq C_8 h_T^m \|g_m\|_{L^2(T)} (\text{osc}_m(g, T) + C_9 (|u - I_{\text{nc}} u|_{H^2(T)} + \|G - \Pi_0 G\|_{L^2(T)})),$$

whence $h_T^m \|g_m\|_{L^2(T)} \lesssim |u - I_{\text{nc}} u|_{H^2(T)} + \|G - \Pi_0 G\|_{L^2(T)} + \text{osc}_m(g, T)$. This and (6.9) lead to the asserted local efficiency $\|h_T^m g\|_{L^2(T)}^2 \lesssim |u - I_{\text{nc}} u|_{H^2(T)}^2 + \|G - \Pi_0 G\|_{L^2(T)}^2 + \text{osc}_m(g, T)^2$ for each $T \in \mathcal{T}$. The sum over all $T \in \mathcal{T}$ concludes the proof. \square

6.3 Example with dominating data oscillations

The data-oscillation term $\|G - \Pi_0 G\|$ dominates the error zero and makes the error estimates useless in the following alarming example. Suppose $m = 2$ (the situation is simpler for $m = 1$) and choose some function $z_c \in S^1(\mathcal{T}; \mathbb{R}^2)$. Given z_c , consider any $z \in H^1(\Omega; \mathbb{R}^2)$ with $\int_E z \, ds = \int_E z_c \, ds$ for all edges $E \in \mathcal{E}$. (This can be done, e.g. by volume bubble functions added to z_c or other functions that do not change the above edge integrals.) Let $g \equiv 0$ and $G := \text{sym Curl } z \in L^2(\Omega; \mathbb{S})$ in (5.5). Since $\Pi_0 \text{Curl } z = \text{Curl}_{\text{pw}} I_{\text{CR}} z = \text{Curl } z_c$, it follows that $\int_\Omega G : D^2 v \, dx = 0 = \int_\Omega G : D_{\text{pw}}^2 v_M \, dx$ for all $v \in V$ and $v_M \in \mathbf{M}(\mathcal{T})$. The first identity follows immediately from an integration by parts and the second from a piecewise integration by parts; for instance, any $v_M \in \mathbf{M}(\mathcal{T})$ satisfies

$$\begin{aligned} \int_\Omega G : D_{\text{pw}}^2 v_M \, dx &= \int_\Omega \text{sym Curl } z : D_{\text{pw}}^2 v_M \, dx = \int_\Omega \text{Curl } z : D_{\text{pw}}^2 v_M \, dx \\ &= \int_\Omega \text{Curl } z_c : D_{\text{pw}}^2 v_M \, dx = \sum_{E \in \mathcal{E}} \int_E v_E \cdot \langle \text{Curl } z_c \rangle_E [\nabla_{\text{pw}} v_M]_E \, ds \\ &= \sum_{E \in \mathcal{E}} \left(\int_E [\nabla_{\text{pw}} v_M]_E \, ds \right) \cdot \langle \text{Curl } z_c \rangle_E v_E \, ds = 0 \end{aligned}$$

(because $[\text{Curl } z_c]_E \times v_E = 0$ and $\langle \text{Curl } z_c \rangle_E v_E = 0$ on E while $\int_E [\nabla_{\text{pw}} v_M]_E \, ds = 0$). This latter orthogonality property implies $u = 0 = u_M$ for the scheme (2.2) as well as for (3.9). On the other hand, $\|G - \Pi_0 G\| = \|\text{sym Curl } (z - z_c)\|$. If $\|G - \Pi_0 G\|_{L^2(T)} = 0$ for $T \in \mathcal{T}$, then $w = (w_1, w_2) := z - z_c \in H^1(T; \mathbb{R}^2)$ satisfies $\text{sym Curl } w = 0$ a.e. in T . This is equivalent to $w_{1,2} = 0 = w_{2,1}$ and $w_{1,1} = w_{2,2}$. Hence (w_2, w_1) is divergence-free and there exists some $\beta \in H^2(T)$ with $(w_2, w_1) = \nabla \beta = (\partial \beta / \partial x_1, \partial \beta / \partial x_2)$. Consequently, β satisfies $\partial^2 \beta / \partial x_j^2 = 0$ for $j = 1, 2$. Since β is harmonic, it is smooth in T . Without loss of generality, assume $0 \in \text{int}(T)$. For $x = (x_1, x_2) \in T$, an integration along the line $\text{conv}\{0, x_1\} \times \{x_2\}$ shows $\beta(x) = \beta(0, x_2) + \int_0^{x_1} \beta_{,1}(\xi, x_2) \, d\xi$ (with the abbreviation $\beta_{,j} := \partial \beta / \partial x_j$, $\beta_{,ij} := \partial^2 \beta / \partial x_i \partial x_j$, $j = 1, 2$ for the partial derivatives). Since $\beta_{,11} = 0$,

$\beta_{,1}(\xi, x_2) = \beta_{,1}(0, x_2)$ for a.e. $\xi \in \text{conv}\{0, x_1\}$. Hence $\beta(x) = \beta(0, x_2) + x_1\beta_{,1}(0, x_2)$. The same argument applies for an integration along $\{x_1\} \times \text{conv}\{0, x_2\}$ and leads to $\beta(x) = \beta(x_1, 0) + x_2\beta_{,2}(x_1, 0)$. Recall that β is smooth in $\text{int}(T)$ and consider derivatives of the two identities to verify $\beta_{,12}(x) = \beta_{,12}(0, x_2) = \beta_{,12}(x_1, 0)$ for all $x \in \text{int}(T)$. The latter identity shows first for $(0, x_2) \in T$ that $\beta_{,12}(0, x_2) = \beta_{,12}(0, 0)$ and second for all $x \in \text{int}(T)$ that $\beta_{,12}(x) = \beta_{,12}(0, 0)$. Consequently, the Hessian $D^2\beta$ of β is constant as $\beta \in P_2(T)$. Recall that $\int_E w \, ds = 0 = \int_E \nabla\beta \, ds$ for all edges $E \in \mathcal{E}(T)$ of T . Therefore the affine vector field $\nabla\beta \in P_1(T; \mathbb{R}^2)$ vanishes at all the edges, midpoints and so everywhere in T . Consequently, β is a constant and $w := (w_1, w_2) = (\beta_{,2}, \beta_{,1}) = 0$ vanishes a.e. in T . This shows that any choice of $z \in H^1(\Omega; \mathbb{R}^2)$ with $\|z - z_c\|_{L^2(\Omega)} > 0$ leads to $\|G - \Pi_0 G\| > 0$. In other words, the error vanishes while the data approximation does not.

Acknowledgements

The research of the first author has been supported by the Deutsche Forschungsgemeinschaft in the Priority Program 1748 under the project "foundation and application of generalized mixed FEM towards nonlinear problems in solid mechanics" (CA 151/22-2). The finalization of this paper has been supported by SPARC project (id 235) entitled *the mathematics and computation of plates*.

References

- [1] S. Agmon, *Lectures on elliptic boundary value problems*, AMS Chelsea Publishing, Providence, RI, 2010.
- [2] D. N. Arnold and F. Brezzi, *Mixed and nonconforming finite element methods: implementation, postprocessing and error estimates*, RAIRO Modél. Math. Anal. Numér. **19** (1985), no. 1, 7–32.
- [3] D. N. Arnold and R. S. Falk, *A uniformly accurate finite element method for the Reissner–Mindlin plate*, SIAM J. Numer. Anal. **26** (1989), no. 6, 1276–1290.
- [4] R. Becker, S. Mao, and Z. Shi, *A convergent nonconforming adaptive finite element method with quasi-optimal complexity*, SIAM J. Numer. Anal. **47** (2010), no. 6, 4639–4659.
- [5] L. Beirão da Veiga, J. Niiranen, and R. Stenberg, *A posteriori error estimates for the Morley plate bending element*, Numer. Math. **106** (2007), no. 2, 165–179.
- [6] H. Blum and R. Rannacher, *On the boundary value problem of the biharmonic operator on domains with angular corners*, Math. Methods Appl. Sci. **2** (1980), no. 4, 556–581.
- [7] S. C. Brenner, *Forty years of the Crouzeix–Raviart element*, Numer. Meth. for PDEs **31** (2015), 367–396.

- [8] S. C. Brenner and L.-Y. Sung, *C^0 interior penalty methods for fourth order elliptic boundary value problems on polygonal domains*, J. Sci. Comput. **22/23** (2005), 83–118.
- [9] C. Carstensen, *Lectures on adaptive mixed finite element methods*, Mixed finite element technologies, CISM Courses and Lectures, vol. 509, Springer Wien New York, Vienna, 2009, pp. 1–56.
- [10] C. Carstensen, S. Bartels, and S. Jansche, *A posteriori error estimates for nonconforming finite element methods*, Numer. Math. **92** (2002), no. 2, 233–256.
- [11] C. Carstensen, M. Eigel, R. H. W. Hoppe, and C. Löbhard, *A review of unified a posteriori finite element error control*, Numer. Math. Theory Methods Appl. **5** (2012), no. 4, 509–558.
- [12] C. Carstensen and D. Gallistl, *Guaranteed lower eigenvalue bounds for the biharmonic equation*, Numer. Math. **126** (2014), no. 1, 33–51.
- [13] C. Carstensen, D. Gallistl, and J. Hu, *A posteriori error estimates for nonconforming finite element methods for fourth-order problems on rectangles*, Numer. Math. **124** (2013), no. 2, 309–335.
- [14] ———, *A discrete Helmholtz decomposition with Morley finite element functions and the optimality of adaptive finite element schemes*, Comput. Math. Appl. **68** (2014), no. 12, part B, 2167–2181.
- [15] C. Carstensen, D. Gallistl, and M. Schedensack, *Adaptive nonconforming Crouzeix-Raviart FEM for eigenvalue problems*, Math. Comp. **84** (2015), 1061–1087.
- [16] C. Carstensen and J. Gedicke, *Guaranteed lower bounds for eigenvalues*, Math. Comput. **83** (2014), no. 290, 2605–2629.
- [17] C. Carstensen, J. Gedicke, and D. Rim, *Explicit error estimates for Courant, Crouzeix-Raviart and Raviart-Thomas finite element methods*, J. Comput. Math. **30** (2012), no. 4, 337–353.
- [18] C. Carstensen and F. Hellwig, *Constants in discrete Poincaré and Friedrichs inequalities and discrete quasi-interpolation*, CMAM **18** (2017), no. 3, 433–450.
- [19] C. Carstensen and J. Hu, *A unifying theory of a posteriori error control for nonconforming finite element methods*, Numer. Math. **107** (2007), no. 3, 473–502.
- [20] C. Carstensen, J. Hu, and A. Orlando, *Framework for the a posteriori error analysis of nonconforming finite element*, SIAM J. Numer. Anal. **45** (2007), no. 1, 68–82.
- [21] C. Carstensen, G. Mallik, and N. Nataraj, *Nonconforming finite element discretization for semilinear problems with trilinear nonlinearity*, IMA J. Numer. Anal., arXiv:1708.07627 (2021), no. 41, 164–205.
- [22] C. Carstensen and N. Nataraj, *Adaptive Morley FEM for the von Kármán equations with optimal convergence rates*, SIAM J. Numer. Anal. (Accepted for publication) (2020), arXiv:1908.08013.

- [23] C. Carstensen and N. Nataraj, *Mathematics and computation of plates*, Under preparation, 2020.
- [24] C. Carstensen, D. Peterseim, and M. Schedensack, *Comparison results of finite element methods for the Poisson model problem*, SIAM J. Numer. Anal. **50** (2012), no. 6, 2803–2823.
- [25] C. Carstensen and S. Puttkammer, *Direct guaranteed lower eigenvalue bounds with optimal a priori convergence rates for the bi-Laplacian*, (2021) Under preparation.
- [26] ———, *How to prove the discrete reliability for nonconforming finite element methods*, J. Comput. Math **38** (2020), no. 1, 142–175.
- [27] P. Ciarlet, C. F. Dunkl, and S. A. Sauter, *A family of Crouzeix-Raviart finite elements in 3D*, Anal. Appl. (Singap.) **16** (2018), no. 5, 649–691.
- [28] P. G. Ciarlet, *The finite element method for elliptic problems*, North-Holland, Amsterdam, 1978.
- [29] M. Crouzeix and P.-A. Raviart, *Conforming and nonconforming finite element methods for solving the stationary Stokes equations. I*, Rev. Française Automat. Informat. Recherche Opérationnelle Sér. Rouge **7** (1973), no. R-3, 33–75.
- [30] W. Dahmen, B. Faermann, I. G. Graham, W. Hackbusch, and S. A. Sauter, *Inverse inequalities on non-quasi-uniform meshes and application to the mortar element method*, Math. Comp. **73** (2004), no. 247, 1107–1138.
- [31] E. Dari, R. Duran, C. Padra, and V. Vampa, *A posteriori error estimators for non-conforming finite element methods*, RAIRO Modél. Math. Anal. Numér. **30** (1996), no. 4, 385–400. MR 1399496
- [32] L. C. Evans, *Partial differential equations*, vol. 19, American Mathematical Society, 1998.
- [33] G. B. Folland, *Introduction to partial differential equations*, second ed., Princeton University Press, Princeton, NJ, 1995.
- [34] D. Gallistl, *Adaptive finite element computation of eigenvalues*, Ph.D. thesis, Humboldt-Universität zu Berlin, Mathematisch-Naturwissenschaftliche Fakultät, 2014.
- [35] ———, *Morley finite element method for the eigenvalues of the biharmonic operator*, IMA J. Numer. Anal. **35** (2015), no. 4, 1779–1811.
- [36] D. Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition.
- [37] P. Grisvard, *Singularities in boundary value problems*, vol. RMA 22, Masson & Springer-Verlag, 1992.

- [38] T. Gudi, *A new error analysis for discontinuous finite element methods for linear elliptic problems*, Math. Comp. **79** (2010), no. 272, 2169–2189.
- [39] J. Hu and Z. Shi, *The best L_2 norm error estimate of lower order finite element methods for the fourth order problem*, J. of Comp. Math. **30** (2012), no. 5, 449–460.
- [40] J. Hu, Z. Shi, and J. Xu, *Convergence and optimality of the adaptive Morley element method*, Numer. Math. **121** (2012), no. 4, 731–752.
- [41] J. Hu and Z. C. Shi, *A new a posteriori error estimate for the Morley element*, Numer. Math. **112** (2009), no. 1, 25–40.
- [42] T. Kato, *Estimation of iterated matrices, with application to the von Neumann condition*, Num. Math. **2** (1960), 22–29.
- [43] J.-L. Lions and E. Magenes, *Non-homogeneous boundary value problems and applications. Vol. I*, Springer-Verlag, New York-Heidelberg, 1972, Translated from the French by P. Kenneth, Die Grundlehren der mathematischen Wissenschaften, Band 181.
- [44] W. Ming and J. Xu, *The Morley element for fourth order elliptic equations in any dimensions*, Numer. Math. **103** (2006), 155–169.
- [45] L. S. D. Morley, *The triangular equilibrium element in the solution of plate bending problems*, Aero. Quart. **19** (1968), 149–169.
- [46] J. Nečas, *Les méthodes directes en théorie des équations elliptiques*, Masson et Cie, Éditeurs, Paris; Academia, Éditeurs, Prague, 1967.
- [47] H. Rabus, *A natural adaptive nonconforming FEM of quasi-optimal complexity*, Comput. Methods Appl. Math. **10** (2010), no. 3, 315–325.
- [48] D. B. Szyld, *The many proofs of an identity on the norm of oblique projections*, Numer. Algorithms **42** (2006), 309–323.
- [49] L. Tartar, *An introduction to Sobolev spaces and interpolation spaces*, Springer, Berlin; Heidelberg, 2010.
- [50] R. Vanselow, *New results concerning the DWR method for some nonconforming FEM*, Appl. Math. **57** (2012), no. 6, 551–568.
- [51] A. Veeseer and P. Zanotti, *Quasi-optimal nonconforming methods for symmetric elliptic problems. I—Abstract theory*, SIAM J. Numer. Anal. **56** (2018), no. 3, 1621–1642.
- [52] ———, *Quasi-optimal nonconforming methods for symmetric elliptic problems. III—Discontinuous Galerkin and other interior penalty methods*, SIAM J. Numer. Anal. **56** (2018), no. 5, 2871–2894.
- [53] ———, *Quasi-optimal nonconforming methods for symmetric elliptic problems. II—Overconsistency and classical nonconforming elements*, SIAM J. Numer. Anal. **57** (2019), no. 1, 266–292.

- [54] R. Verfürth, *A posteriori error estimation techniques for finite element methods*, Numerical Mathematics and Scientific Computation, Oxford University Press, Oxford, 2013.
- [55] M. Wang and J. Xu, *Minimal finite element spaces for $2m$ -th-order partial differential equations in R^n* , Math. Comp. **82** (2013), no. 281, 25–43.