Debiased Kernel Methods

Rahul Singh MIT Economics rahul.singh@mit.edu

Abstract

I propose a practical procedure based on bias correction and sample splitting to calculate confidence intervals for functionals of generic kernel methods, i.e. non-parametric estimators learned in a reproducing kernel Hilbert space (RKHS). For example, an analyst may desire confidence intervals for functionals of kernel ridge regression. I propose a bias correction that mirrors kernel ridge regression. The framework encompasses (i) evaluations over discrete domains, (ii) derivatives over continuous domains, (iii) treatment effects of discrete treatments, and (iv) incremental treatment effects of continuous treatments. For the target quantity, whether it is (i)-(iv), I prove \sqrt{n} consistency, Gaussian approximation, and semiparametric efficiency by finite sample arguments. I show that the classic assumptions of RKHS learning theory also imply inference.

1 Introduction

The reproducing kernel Hilbert space (RKHS) is a popular nonparametric setting in machine learning. For example, kernel ridge regression is a classic algorithm for prediction, with impressive attributes. Computationally, it has a closed form solution; empirically, it adapts well to local nonlinearity while preserving smoothness; statistically, it is minimax optimal with respect to mean square error [16].

A recent literature proposes kernel methods for causal learning problems, such as nonparametric instrumental variable regression [79] and nonparametric treatment effects [81, 78], that share the impressive attributes of kernel ridge regression. The motivation for these causal kernel methods is policy evaluation in social science and epidemiology. In social science and epidemiology, however, confidence intervals are essential.

The goal of this project is to provide practical confidence intervals for functionals of kernel ridge regression *under the classic assumptions of RKHS learning theory*. I provide conceptual, algorithmic, and statistical contributions to RKHS theory.

Conceptual. The generality of the framework is threefold. First, I unify (i) evaluations over discrete domains, (ii) derivatives over continuous domains, (iii) treatment effects of discrete treatments, and (iv) incremental treatment effects of continuous treatments into one general inference problem. Second, I allow for any nonparametric kernel method with a known learning rate. Third, I allow for the covariates to be discrete or continuous, and low, high, or infinite dimensional. As such, the main results apply to kernel methods trained on texts, graphs, or images.

Algorithmic. I provide a general purpose procedure to automatically debias (i)-(iv) for semiparametric inference. The debiasing is novel, and I show that is has a one line, closed form solution. Its only hyperparameters are kernel hyperparameters and ridge regression penalties, just like kernel ridge regression. The former have well known heuristics, and the latter are easily tuned by cross validation.

Statistical. For the target quantity, whether it is (i)-(iv), I prove pointwise \sqrt{n} consistency, Gaussian approximation, and semiparametric efficiency by finite sample arguments. The analysis explicitly

accounts for each source of error in any finite sample size. Formally, I show that the classic assumptions of RKHS learning theory also imply inference.

The structure of the paper is as follows. Section 1.1 describes related work. Section 2 articulates the general inference problem. Section 3 proposes the general purpose debiasing procedure and furnishes confidence intervals. Sections 4 and 5 present the main theoretical contributions. Section 6 concludes.

1.1 Related work

Functionals. By focusing on functionals, i.e. scalar summaries, of nonparametric quantities, this paper continues the tradition of classic semiparametric statistical theory [50, 33, 36, 63, 48, 69, 95, 13, 53, 67, 96, 14, 55, 1, 56, 2, 90, 49, 3]. Whereas classic semiparametric theory studies functionals of densities or regressions over low dimensional domains, I study functionals of generic kernel methods over domains that may include discrete or continuous treatments and low, high, or infinite dimensional covariates. In classic semiparametric theory, an object called the Riesz representer of the functional appears in asymptotic variance calculations [53, 37]. For exactly the same reasons, it appears in the practical procedure for confidence intervals that I propose.

Debiasing. In asymptotic inference, the Riesz representer is inevitable. Motivated by this insight, a growing literature directly incorporates the Riesz representer into estimation, which amounts to debiasing known estimators [33, 14, 97, 7, 8, 9, 10, 42, 43, 44, 91, 62, 21, 58, 64, 39, 40, 41, 15, 99, 100]. Doubly robust estimating equations serve this purpose [68, 67, 94, 93, 51, 89]. A geometric perspective on debiasing emphasizes Neyman orthogonality: by debiasing, the learning problem for the functional of interest becomes orthogonal to the learning problem for the underlying nonparametric object [59, 60, 96, 65, 98, 9, 10, 20, 6, 19, 31]. In this work, I debias kernel methods using doubly robust estimating equations. The learning problem for (i)-(iv) becomes orthogonal to the learning problem for the underlying kernel method.

Sample splitting. With debiasing alone, a key challenge remains: for inference, the function class in which the nonparametric quantity is learned must be Donsker. However, popular nonparametric settings in machine learning such as the RKHS are not Donsker. A solution to this challenging issue is to combine debiasing with sample splitting [12, 75, 48, 96, 65]. The debiased machine learning (DML) and targeted maximum likelihood literatures are responsible for this insight [74, 72, 73, 94, 98, 93, 27, 92, 20, 46, 19, 45]. In particular, DML delivers simple sufficient conditions for inference on functionals in terms of learning rates of the underlying regression and the Riesz representer [20, 19]. The analysis of Section 4 is written at this level of generality. Whereas the existing DML theory is usually asymptotic, I provide the first finite sample analysis with black box machine learning.

Automatic debiasing. Since standard statistical learning theory provides rates for nonparametric regression, what remains is estimation and analysis that provides sufficiently fast rates for the Riesz representer. However, the Riesz representer may be a complex object. Even for simple functionals such as policy effects, its closed form involves ratios of densities, which would imply slow rates. A crucial insight is that the Riesz representer is directly identified from data; with some creativity, it can be estimated and analyzed directly, without estimating or analyzing its components [66, 57, 4, 22, 24, 34, 35, 80, 71, 23]. For the RKHS context, I propose a novel estimator of the Riesz representer that generalizes kernel ridge regression. I derive its fast rate in Section 5.

Companion work on adversarial debiasing. In companion work [23], we develop an automatic debiasing procedure that generalizes adversarial estimation. At a high level, both projects are concerned with inference for functionals over machine learning function spaces that are not Donsker. An advantage of the analysis in [23] is that its critical radius arguments apply to a broader class of function spaces: not only the RKHS, but also sparse linear and neural network spaces. The goal of this project is to guarantee inference under the RKHS learning theory assumptions that appear in [16, 79, 81, 78]. Computationally, the kernel ridge Riesz representer involves much simpler operations of kernel matrices than the adversarial Riesz representer.

Kernel methods. Recent developments in the kernel methods literature motivate this project. In particular, a new toolkit addresses causal learning problems such as instrumental variable regression [79, 52, 28], treatment effects under selection on observables [61, 81], and treatment effects with negative controls [78]. Confidence intervals are essential for practical use in policy evaluation. As

explained above, a key contribution is fast rate analysis for a new Riesz representer estimator. My analysis directly builds on the seminal work of [16], which provides minimax optimal rates for kernel ridge regression. As such, classic learning theory assumptions imply the learning rate for the Riesz representer and hence inference for the functional.

2 Framework

2.1 General inference problem

The general inference problem is to find a confidence interval for $\theta_0 \in \mathbb{R}$ where

$$\theta_0 = \mathbb{E}[m(W, \gamma_0)], \quad \gamma_0 \in \mathbb{L}_2$$

 $m: \mathcal{W} \times \mathbb{L}_2 \to \mathbb{R}$ is an abstract formula. $W \in \mathcal{W}$ is a concatenation of random variables in the model excluding the outcome $Y \in \mathcal{Y} \subset \mathbb{R}$. \mathbb{L}_2 is the space of functions of the form $\gamma: \mathcal{W} \to \mathbb{R}$ that are square integrable with respect to measure \mathbb{P} .

I consider functionals of nonparametric regression. These statistical quantities have causal interpretation under the assumption of unconfoundedness, also called selection on observables. The simplest example is evaluation over a discrete domain.

Example 2.1 (Evaluation of nonparametric regression). W are covariates, and $\gamma_0(w) = \mathbb{E}[Y|W=w]$. Then evaluation at location w is $\theta_0 = \gamma_0(w)$. I require that W are discrete.

As we will see, a virtue of working in the RKHS is the ability to handle functionals that involve differentiation.

Example 2.2 (Average partial derivative of nonparametric regression). W are covariates, and $\gamma_0(w) = \mathbb{E}[Y|W=w]$. Then average partial derivative with respect to W_j is $\theta_0 = \mathbb{E}[\nabla_j \gamma_0(W)]$ where $\nabla_j = \frac{\partial}{\partial w_j}$. I require that W_j is continuous.

Next, I turn to classic treatment effects in causal inference: average treatment effect (ATE), average treatment on the treated (ATT), and conditional average treatment effect (CATE). I generalize each of these classic effects to have discrete treatment values rather than binary treatment values. The expressions follow from [70, 81].

Example 2.3 (Discrete treatment effects). W = (D, X) are treatment and covariates, and $\gamma_0(d, x) = \mathbb{E}[Y|D = d, X = x]$. For heterogeneous effects, I replace X with (V, X) where V is the low dimensional subcovariate of interest. I require that (D, V) are discrete.

- 1. ATE. The mean potential outcome of treatment value d is given by $\theta_0 = \mathbb{E}[\gamma_0(d, X)]$.
- 2. ATT. The mean potential outcome of treatment value d' for the subpopulation who received treatment value d is given by $\beta_0 = \frac{\theta_0}{\mathbb{P}(d)}$ where $\theta_0 = \mathbb{E}[\gamma_0(d',X)\mathbb{1}_{D=d}]$.
- 3. CATE. The mean potential outcome of treatment value d for the subpopulation with subcovariate value v is given by $\beta_0 = \frac{\theta_0}{\mathbb{P}(v)}$ where $\theta_0 = \mathbb{E}[\gamma_0(d,v,X)\mathbb{1}_{V=v}]$.

In a couple instances, the quantity of interest is $\beta_0 \in \mathbb{R}$, which is a ratio of $\theta_0 \in \mathbb{R}$ and a marginal probability. By standard arguments, inference for θ_0 implies inference for β_0 by delta method.

In the examples so far, I have restricted treatment to be discrete. I now allow treatment to be continuous and consider incremental average treatment effect (ATE- ∇) and incremental conditional average treatment effect (CATE- ∇).

These effects can be viewed as relaxations of average derivatives of potential outcomes. For example, ATE- ∇ is a relaxation of $\mathbb{E}[\nabla Y^{(d)}]$ where $\nabla = \frac{\partial}{\partial d}$. Specifically, ATE- ∇ is $\mathbb{E}[\nabla Y^{(D)}]$, sometimes called the joint average treatment effect. The expressions follow from [38, 81].

Example 2.4 (Incremental treatment effects). W = (D, X) are treatment and covariates, and $\gamma_0(d, x) = \mathbb{E}[Y|D = d, X = x]$. I now allow D to be continuous but maintain that V is discrete.

1. ATE- ∇ . The mean incremental treatment effect is given by $\mathbb{E}[\nabla \gamma_0(D,X)]$.

2. CATE- ∇ . The mean incremental treatment effect for the subpopulation with subcovariate value v is given by $\beta_0 = \frac{\theta_0}{\mathbb{P}(v)}$ where $\theta_0 = \mathbb{E}[\nabla \gamma_0(D, V, X) \mathbb{1}_{V=v}]$

Note that this relaxation is not possible for ATT, because I require the conditioning variable that defines the subpopulation to be discrete.

2.2 Structure on the functional

To prove validity of the confidence interval, I require that the formula m is mean square continuous.

Assumption 2.1 (Mean square continuity). There exists $\bar{L}^{(m)} < \infty$ s.t.

$$\mathbb{E}[m(W,\gamma)]^2 \le \bar{L}^{(m)} \cdot \mathbb{E}[\gamma(W)]^2$$

This condition will be key in Section 4, where I reduce the problem of inference for θ_0 into the problem of learning $(\gamma_0, \alpha_0^{\min})$, where α_0^{\min} is introduced below. It is a powerful condition, yet it is easily satisfied in all of the discrete examples above.

Proposition 2.1 (Verifying continuity for discrete examples). For Examples 2.1 and 2.3, Assumption 2.1 holds for all $\gamma \in \mathbb{L}_2$ if the corresponding propensity scores are bounded away from zero.

For the continuous examples, which involve derivatives, some additional care is needed. It turns out that Assumption 2.1 can be verified for all $\gamma \in \mathbb{H}_2^s \subset \mathbb{L}_2$, which I now define. As far as I know, the insight that derivative functionals may be mean square continuous is original. This possibility is assumed away in [54, Assumption 6] and [18, Assumption 3.9].

Definition 2.1 (Sobolev space). For simplicity, let $W \subset \mathbb{R}^d$. $\mathbb{H}_2^s \subset \mathbb{L}_2$ is a Sobolev space if it consists of functions of the form $\gamma: W \to \mathbb{R}$ such that their s-order derivatives are square integrable.

Proposition 2.2 (Verifying continuity for continuous examples). For Examples 2.2 and 2.4, Assumption 2.1 holds for all $\gamma \in \mathbb{H}_2^s$ if $s > \frac{d}{2} + 1$.

The restriction that $\gamma_0 \in \Gamma \subset \mathbb{L}_2$, where Γ is some convex function space, is called a restricted model in semiparametric statistical theory. In RKHS learning theory, mean square rates are adaptive to the smoothness of γ_0 , encoded by $\gamma_0 \in \Gamma$.

Proposition 2.3 (Riesz representation; Lemma S3.1 of [22]). *Suppose Assumption 2.1 holds. Further suppose* $\gamma_0 \in \Gamma$. *Then there exists a Riesz representer* $\alpha_0 \in \mathbb{L}_2$ *s.t.*

$$\mathbb{E}[m(W,\gamma)] = \mathbb{E}[\alpha_0(W)\gamma(W)], \quad \forall \gamma \in \Gamma$$

Moreover, there exists a unique minimal Riesz representer $\alpha_0^{\min} \in closure(span(\Gamma))$ that satisfies this equation.

Riesz representation delivers a doubly robust formulation of the target $\theta_0 \in \mathbb{R}$. I view $(\gamma_0, \alpha_0^{\min})$ as nuisance parameters that I must learn in order to learn and infer θ_0 .

$$0 = \mathbb{E}[\psi_0(W)], \quad \psi_0(w) = \psi(w, \theta_0, \gamma_0, \alpha_0^{\min}), \quad \psi(w, \theta, \gamma, \alpha) = m(w, \gamma) + \alpha(w)[y - \gamma(w)] - \theta$$

The term $\alpha(w)[y-\gamma(w)]$ is the product of the Riesz representer and the regression residual. It serves as a bias correction for the term $m(w,\gamma)$. This formulation is doubly robust in the sense that the moment function for θ_0 remains valid if either γ_0 or α_0 is correct, i.e.

$$0 = \mathbb{E}[\psi(W, \theta_0, \gamma_0, \alpha)] \quad \forall \alpha \in \mathbb{L}_2, \quad 0 = \mathbb{E}[\psi(W, \theta_0, \gamma, \alpha_0)] \quad \forall \gamma \in \mathbb{L}_2$$

While any Riesz representer α_0 will suffice for valid learning and inference, the minimal Riesz representer α_0^{\min} confers semiparametric efficiency [22, Theorem 4.2].

¹As a preview, I will take $\Gamma = \mathcal{H}^c$, which is a smoother RKHS than \mathcal{H} . Conveniently, $closure(span(\mathcal{H}^c)) = \mathcal{H}^c$, and $\mathcal{H}^1 = \mathcal{H}$. For the continuous examples, I will choose $\mathcal{H} = \mathbb{H}_2^s$ and hence $\Gamma = [\mathbb{H}_3^s]^c$.

2.3 Reproducing kernel Hilbert space

Kernel ridge regression is an algorithm that conducts estimation of γ_0 in a reproducing kernel Hilbert space (RKHS) $\mathcal{H} \subset \mathbb{L}_2$. \mathcal{H} consists of functions of the form $\gamma: \mathcal{W} \to \mathbb{R}$. I denote its kernel by $k: \mathcal{W} \times \mathcal{W} \to \mathbb{R}$, and its feature map $\phi: \mathcal{W} \to \mathcal{H}$, $w \mapsto k(w, \cdot)$.

To gain familiarity with \mathcal{H} , I present its series interpretation. This exposition sets up notation that I will use in Sections 3 and 5. In preparation, define the convolution operator $L_k: \mathbb{L}_2 \to \mathbb{L}_2$, $f \mapsto \int k(\cdot,w)f(w)d\mathbb{P}(w)$. L_k is a self adjoint, positive, compact operator, so by the spectral theorem we can denote its countable eigenvalues and eigenfunctions by $\{\lambda_j\}$ and $\{\varphi_j\}$, respectively: $L_k f = \sum_{j=1}^\infty \lambda_j \langle \varphi_j, f \rangle_2 \cdot \varphi_j$.

With this notation, I express \mathbb{L}_2 and the RKHS \mathcal{H} in terms of the series $\{\varphi_j\}$. If $f \in \mathbb{L}_2$, then f can be uniquely expressed as $f = \sum_{j=1}^{\infty} a_j \varphi_j$ and the partial sums $\sum_{j=1}^{J} a_j \varphi_j$ converge to f in \mathbb{L}_2 . Indeed,

$$\mathbb{L}_2 = \left\{ f = \sum_{j=1}^{\infty} a_j \varphi_j : \sum_{j=1}^{\infty} a_j^2 < \infty \right\}, \quad \langle f, g \rangle_2 = \sum_{j=1}^{\infty} a_j b_j$$

for $f=\sum_{j=1}^\infty a_j\varphi_j$ and $g=\sum_{j=1}^\infty b_j\varphi_j$. By [26, Theorem 4], the RKHS $\mathcal H$ corresponding to the kernel k can be explicitly represented as

$$\mathcal{H} = \left\{ f = \sum_{j=1}^{\infty} a_j \varphi_j : \sum_{j=1}^{\infty} \frac{a_j^2}{\lambda_j} < \infty \right\}, \quad \langle f, g \rangle_{\mathcal{H}} = \sum_{j=1}^{\infty} \frac{a_j b_j}{\lambda_j}$$

Let us interpret this result, known as Picard's criterion [5]. Recall that $\{\lambda_j\}$ is a weakly decreasing sequence. The RKHS \mathcal{H} is the subset of \mathbb{L}_2 for which higher order terms in the series $\{\varphi_j\}$ have a smaller contribution. Finally, I express the feature map, kernel, and evaluation in terms of the series $\{\varphi_j\}$ appealing to Mercer's theorem.

$$\phi(w) = \{\sqrt{\lambda_j}\varphi_j(w)\}_{j=1}^{\infty}, \quad k(w, w') = \langle \phi(w), \phi(w') \rangle_{\mathcal{H}}, \quad \gamma(w) = \langle \gamma, \phi(w) \rangle_{\mathcal{H}}$$

With this series notation, we are now prepared to express a critical insight that first appears in [23, Section 5.3]: causal inference introduces counterfactual features, which must appear in learning and inference. Since each $\varphi_j \in \mathbb{L}_2$, $m(w, \varphi_j)$ is well defined. I therefore write the counterfactual feature map, kernel, and evaluation as

$$\phi^{(m)}(w) = \{\sqrt{\lambda_j} m(w, \varphi_j)\}_{j=1}^{\infty}, \ k^{(m)}(w, w') = \langle \phi^{(m)}(w), \phi^{(m)}(w') \rangle_{\mathcal{H}}, \ m(w, \gamma) = \langle \gamma, \phi^{(m)}(w) \rangle_{\mathcal{H}}$$

In the simple example of ATE at d=1, the formula is $m(w,\gamma)=\gamma(1,x)$. Then

$$\phi^{(m)}(W_i) = \{\sqrt{\lambda_i}\varphi_i(1, X_i)\}_{i=1}^{\infty} = \phi(1, X_i)$$

We see that the counterfactual feature map applied to observation W_i replaces the observed treatment value D_i with the counterfactual value d=1.

3 Algorithm

3.1 Confidence interval

The goal of this paper is general purpose learning and inference for θ_0 , where $\theta_0 \in \mathbb{R}$ may be an (i) evaluation over discrete domain, (ii) average partial derivative over a continuous domain, (iii) treatment effect of discrete treatment, or (iv) incremental treatment effect of continuous treatment. In Section 2, I demonstrated that any such θ_0 is a mean square continuous functional of $\gamma_0 \in \Gamma$ and hence has a unique minimal representer $\alpha_0^{\min} \in \Gamma$. In this section, I describe a meta algorithm to turn kernel estimators $\hat{\gamma}$ of γ_0 and $\hat{\alpha}$ for α_0^{\min} into an estimator $\hat{\theta}$ of θ_0 such that $\hat{\theta}$ has a valid and practical confidence interval. This meta algorithm is precisely debiased machine learning (DML) [20, 19].

Recall that $\hat{\gamma}$ may be any one of a number of different kernel methods such as kernel ridge regression. To preserve this generality, I do not instantiate a choice of $\hat{\gamma}$; I treat it as a black box. In subsequent

analysis, I will only require that $\hat{\gamma}$ converges to γ_0 in mean square error. This mean square rate is guaranteed by existing learning theory. As such, the inferential theory builds directly on the learning theory.

The target estimator $\hat{\theta}$ as well as its confidence interval will depend on nuisance estimators $\hat{\gamma}$ and $\hat{\alpha}$. For now, I refrain from instantiating the estimator $\hat{\alpha}$ for α_0^{\min} . As we will see in subsequent analysis, the general theory only requires that $\hat{\alpha}$ converges to α_0^{\min} in mean square error. This property can be verified by extending classic learning theory arguments, which I reserve for Section 5.

Algorithm 3.1 (Target and confidence interval; eq. 2.2 and 2.3 of [20]). *Partition the sample into folds* $\{I_{\ell}\}_{\ell=1:L}$. *Denote by* I_{ℓ}^{c} *observations not in fold* I_{ℓ}

- 1. For each fold ℓ , estimate $\hat{\gamma}_{\ell}$ and $\hat{\alpha}_{\ell}$ from observations in I_{ℓ}^{c}
- 2. Estimate $\hat{\theta}$ as

$$\hat{\theta} = \frac{1}{n} \sum_{\ell=1}^{L} \sum_{i \in I_{\ell}} \{ m(W_i, \hat{\gamma}_{\ell}) + \hat{\alpha}_{\ell}(W_i) [Y_i - \hat{\gamma}_{\ell}(W_i)] \}$$

3. Estimate its $(1-a) \cdot 100\%$ confidence interval as

$$\hat{\theta} \pm c_a \frac{\hat{\sigma}}{\sqrt{n}}, \quad \hat{\sigma}^2 = \frac{1}{n} \sum_{\ell=1}^{L} \sum_{i \in I_{\ell}} \{ m(W_i, \hat{\gamma}_{\ell}) + \hat{\alpha}_{\ell}(W_i) [Y_i - \hat{\gamma}_{\ell}(W_i)] - \hat{\theta} \}^2$$

where c_a is the $1-\frac{a}{2}$ quantile of the standard Gaussian.

3.2 Bias correction

Algorithm 3.1 is a meta algorithm that takes as inputs the sample, the algorithm $\hat{\gamma}$ for γ_0 , and the algorithm $\hat{\alpha}$ for α_0^{\min} . A broad literature already proposes and analyzes estimators $\hat{\gamma}$. I now propose an estimator $\hat{\alpha}$ that extends kernel ridge regression. As such, existing learning guarantees for kernel ridge regression will extend to $\hat{\alpha}$.

I motivate the new estimator by the following property.

Proposition 3.1 (Riesz representer loss). Suppose the kernels k and $k^{(m)}$ are bounded. Then

$$\alpha_0^{\min} \in \underset{\alpha \in \mathcal{H}}{\operatorname{argmin}} \mathcal{L}(\alpha), \quad \mathcal{L}(\alpha) = -2\langle \alpha, \mu^{(m)} \rangle_{\mathcal{H}} + \langle \alpha, T\alpha \rangle_{\mathcal{H}}$$

where $T = \mathbb{E}[\phi(W) \otimes \phi(W)]$ is the uncentered covariance operator and $\mu^{(m)} = \mathbb{E}[\phi^{(m)}(W)]$ is the counterfactual kernel mean embedding.

The quantities (μ,T) generalize the sufficient statistics (\tilde{M},\tilde{G}) in automatic debiased machine learning (Auto-DML) [22, 24]. In Auto-DML, which uses p explicit basis functions, $\tilde{M} \in \mathbb{R}^p$ is the moment vector and $\tilde{G} \in \mathbb{R}^{p \times p}$ is the covariance matrix. In my approach, which uses the countable spectrum of the kernel k as implicit basis functions, $\mu^{(m)} \in \mathcal{H}$ is the counterfactual kernel mean embedding [83] and $T: \mathcal{H} \to \mathcal{H}$ is the covariance operator [32].

Next, I define the regularized Riesz representer in this context. The regularization is a ridge penalty with regularization parameter λ .

Definition 3.1 (Kernel ridge Riesz representer: Population).

$$\alpha_{\lambda} = \underset{\alpha \in \mathcal{H}}{\operatorname{argmin}} \mathcal{L}_{\lambda}(\alpha), \quad \mathcal{L}_{\lambda}(\alpha) = \mathcal{L}(\alpha) + \lambda \|\alpha\|_{\mathcal{H}}^{2}$$

I define the estimator as the empirical analogue. To lighten notation, I abstract from sample splitting and consider the setting where n observations are used.

Definition 3.2 (Kernel ridge Riesz representer: Sample).

$$\hat{\alpha} = \operatorname*{argmin}_{\alpha \in \mathcal{H}} \mathcal{L}_{\lambda}^{n}(\alpha), \quad \mathcal{L}_{\lambda}^{n}(\alpha) = -2\langle \alpha, \hat{\mu}^{(m)} \rangle_{\mathcal{H}} + \langle \alpha, \hat{T}\alpha \rangle_{\mathcal{H}} + \lambda \|\alpha\|_{\mathcal{H}}^{2}$$

where
$$\hat{T} = \frac{1}{n} \sum_{i=1}^{n} [\phi(W_i) \otimes \phi(W_i)]$$
 and $\hat{\mu}^{(m)} = \frac{1}{n} \sum_{i=1}^{n} \phi^{(m)}(W_i)$.

The generalized kernel ridge loss in Definition 3.2 is original. It differs from Dantzig [22], Lasso [25], and adversarial [23] losses previously considered.

As a kernel estimator, $\hat{\alpha}$ can be implemented by simple matrix operations. Kernel methods typically involve matrix operations of the matrix $K^{(1)} \in \mathbb{R}^{n \times n}$ with (i,j)-th entry $k(W_i,W_j)$; the kernel matrix collects inner product of features $\{\phi(W_i)\}$. The proposed kernel method involves an extended kernel matrix $K \in \mathbb{R}^{2n \times 2n}$, which includes inner products with counterfactual features $\{\phi^{(m)}(W_i)\}$.

Wherever possible, I preserve the notation of companion work on adversarial estimation [23]. Define the operator $\Phi: \mathcal{H} \to \mathbb{R}^n$ with i-th row $\langle \phi(W_i), \cdot \rangle_{\mathcal{H}}$ and likewise the operator $\Phi^{(m)}: \mathcal{H} \to \mathbb{R}^n$ with i-th row $\langle \phi^{(m)}(W_i), \cdot \rangle_{\mathcal{H}}$. Finally define the concatenated operator $\Psi: \mathcal{H} \to \mathbb{R}^{2n}$ that is constructed by concatenating Φ and $\Phi^{(m)}$. By construction, $K = \Psi \Psi^*$. For intuition, it is helpful to conceptualize these operators as matrices. Informally,

$$\Psi = \begin{bmatrix} \Phi \\ \Phi^{(m)} \end{bmatrix}, \quad K = \begin{bmatrix} K^{(1)} & K^{(2)} \\ K^{(3)} & K^{(4)} \end{bmatrix} = \begin{bmatrix} \Phi \Phi^* & \Phi(\Phi^{(m)})^* \\ \Phi^{(m)} \Phi^* & \Phi^{(m)} (\Phi^{(m)})^* \end{bmatrix}$$

Note that $\{K^{(j)}\}_{j\in[4]}\in\mathbb{R}^{n\times n}$ and hence $K\in\mathbb{R}^{2n\times 2n}$ can be computed from data, though they depend on the choice of formula m.

Proposition 3.2 (Extended kernel matrix: Computation; Proposition 9 of [23]). *In the case of ATE with* $m(W, \gamma) = \gamma(d, X)$,

$$K_{ij}^{(1)} = k(D_i, D_j)k(X_i, X_j), \quad K_{ij}^{(2)} = K_{ji}^{(3)} = k(D_i, d)k(X_i, X_j), \quad K_{ij}^{(4)} = k(d, d)k(X_i, X_j)$$

With this additional notation, I prove that $\hat{\alpha}$ has a closed form solution and derive that solution. My argument is an extension of the classic representer theorem for kernel methods [47, 76].

Proposition 3.3 (Generalized representer theorem). There exists some $\rho \in \mathbb{R}^{2n}$ s.t. $\hat{\alpha} = \Psi^* \rho$.

I use this generalized representation theorem to derive the closed form of $\hat{\alpha}$ in terms of matrix operations.

Algorithm 3.2 (Kernel ridge Riesz representer: Computation). Given observations $\{W_i\}$, formula m, and evaluation location w

- 1. Calculate $K^{(j)} \in \mathbb{R}^{n \times n}$ as defined above
- 2. Calculate $\Omega^{2n\times 2n}$ and $u,v\in\mathbb{R}^{2n}$ by

$$\Omega = \begin{bmatrix} K^{(1)}K^{(1)} & K^{(1)}K^{(3)} \\ K^{(3)}K^{(1)} & K^{(3)}K^{(3)} \end{bmatrix}, \ v = \begin{bmatrix} K^{(2)} \\ K^{(4)} \end{bmatrix} \mathbbm{1}_n, \ u_i = \begin{cases} k(W_i,w) & \text{if } i \in \{1,...,n\} \\ \tilde{k}(W_i,w) & \text{if } i \in \{n+1,...,2n\} \end{cases}$$

where $\mathbb{1}_n \in \mathbb{R}^n$ is a vector of ones, $k(w, w') = \langle \phi(w), \phi(w') \rangle_{\mathcal{H}}$, and $\tilde{k}(w, w') = \langle \phi^{(m)}(w), \phi(w') \rangle_{\mathcal{H}}$

3. Set
$$\hat{\alpha}(w) = v^{\top} (\Omega + n\lambda K)^{-1} u$$

I give a theoretical rate of regularization λ that balances bias and variance in Theorem 5.2 below. I give a practical procedure based on leave-one-out cross validation to empirically balance bias and variance in Appendix E. Conveniently, the cross validation procedure has a closed form solution.

For comparison, if $\hat{\gamma}$ is kernel ridge regression, then $\hat{\gamma}(w) = v^{\top} (K^{(1)} + n\lambda I)^{-1} u$ where $v_i = Y_i$ and $u_i = k(W_i, w)$. Algorithm 3.2 involves only one inversion of one matrix in $\mathbb{R}^{2n \times 2n}$. Its simplicity confers numerical stability and robustness to tuning. By contrast, [23, Corollary 14] involves several inversions (and inversions of products of inversions) of matrices in $\mathbb{R}^{2n \times 2n}$.

4 Validity of confidence interval

I write this section at a high level of generality so that it can be used in future work. In particular, I abstract from the RKHS framework. I assume high level conditions and consider black box estimators $(\hat{\gamma}, \hat{\alpha})$. I prove by finite sample arguments that $\hat{\theta}$ defined by Algorithm 3.1 is consistent, and

its confidence interval is valid and semiparametrically efficient. In Section 5, I will verify the high level conditions in the RKHS framework.

Towards this end, define $\sigma^2 = \mathbb{E}[\psi_0(W)]^2$, $\eta^3 = \mathbb{E}|\psi_0(W)|^3$, and $\zeta^4 = \mathbb{E}[\psi_0(W)]^4$. Write the Berry Esseen constant as $c^{BE} = 0.4748$ [77]. The result will be in terms of abstract mean square rates on the nuisances.

Definition 4.1 (Mean square error). Write the conditional mean square error of $(\hat{\gamma}_{\ell}, \hat{\alpha}_{\ell})$ trained on I_{ℓ}^{c} as

$$\mathcal{R}(\hat{\gamma}_{\ell}) = \mathbb{E}[\{\hat{\gamma}_{\ell}(W) - \gamma_0(W)\}^2 | I_{\ell}^c], \quad \mathcal{R}(\hat{\alpha}_{\ell}) = \mathbb{E}[\{\hat{\alpha}_{\ell}(W) - \alpha_0(W)\}^2 | I_{\ell}^c]$$

We are now ready to state the first main result, which is a finite sample Gaussian approximation.

Theorem 4.1 (Gaussian approximation). Suppose

$$\mathbb{E}[Y - \gamma_0(W)]^2 \le \bar{\sigma}^2, \quad \mathbb{E}[m(W, \gamma)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma(W)]^2, \quad \|\alpha_0\|_{\infty} \le \bar{\alpha}$$

Then w.p. $1 - \epsilon$,

$$\sup_{z \in \mathbb{R}} \left| \mathbb{P} \left(\frac{\sqrt{n}}{\sigma} (\hat{\theta} - \theta_0) \le z \right) - \Phi(z) \right| \le c^{BE} \left(\frac{\eta}{\sigma} \right)^3 n^{-\frac{1}{2}} + \frac{\Delta}{\sqrt{2\pi}} + \epsilon$$

where $\Phi(z)$ is the standard Gaussian c.d.f. and

$$\Delta = \frac{3L}{\epsilon \cdot \sigma} \left\{ (\sqrt{\bar{L}^{(m)}} + \bar{\alpha}) \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} + \bar{\sigma} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} + \sqrt{n} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} \right\}$$

Theorem 4.1 is a finite sample Gaussian approximation for DML with black box machine learning. It is a finite sample refinement of the asymptotic black box result in [20, Lemma 15]. It is a black box generalization of the main result in [22, Theorem 4.1], which provides a finite sample analysis specific to the Dantzig selector. By Theorem 4.1, the neighborhood of Gaussian approximation scales as $\frac{\sigma}{\sqrt{n}}$. It is therefore helpful to bound the magnitude of the asymptotic variance σ^2 .

Proposition 4.1 (Variance bound). Suppose the conditions of Theorem 4.1 hold. Then

$$\sigma^2 \le 3(\bar{L}^{(m)}\mathbb{E}[\gamma_0(W)]^2 + \bar{\alpha}^2\bar{\sigma}^2 + \theta_0^2)$$

Observe that finite sample Gaussian approximation is in terms of the true asymptotic variance σ^2 . I now provide a guarantee for its estimator $\hat{\sigma}^2$.

Theorem 4.2 (Variance estimation). Suppose

$$\mathbb{E}[Y - \gamma_0(W)]^2 \le \bar{\sigma}^2, \quad \mathbb{E}[m(W, \gamma)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma(W)]^2, \quad \|\hat{\alpha}_\ell\|_{\infty} \le \bar{\alpha}'$$

Then w.p. $1 - \epsilon'$,

$$|\hat{\sigma}^2 - \sigma^2| \le \Delta' + 2\sqrt{\Delta'}(\sqrt{\Delta''} + \sigma) + \Delta''$$

where

$$\Delta' = \frac{24L}{\epsilon'} \{ [\hat{\theta} - \theta_0]^2 + [\bar{L}^{(m)} + (\bar{\alpha}')^2] \mathcal{R}(\hat{\gamma}_\ell) + \bar{\sigma}^2 \mathcal{R}(\hat{\alpha}_\ell) \}, \quad \Delta'' = \sqrt{\frac{2}{\epsilon'}} \zeta^2 n^{-\frac{1}{2}}$$

Theorem 4.2 is a both a black box generalization and a finite sample refinement of the asymptotic theorem in [24] specific to the Lasso. Theorems 4.1 and 4.2 immediately imply sufficient conditions for validity of the proposed confidence interval.

Corollary 4.1 (Confidence interval). Suppose the assumptions of Theorems 4.1 and 4.2 hold as well as the regularity condition on moments $\left\{ \left(\frac{\eta}{\sigma} \right)^3 + \zeta^2 \right\} n^{-\frac{1}{2}} \to 0$. Assume

$$(\sqrt{\bar{L}^{(m)}} + \bar{\alpha} + \bar{\alpha}')\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \xrightarrow{p} 0, \quad \bar{\sigma}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} \xrightarrow{p} 0, \quad \sqrt{n}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \xrightarrow{p} 0$$

Then the estimator $\hat{\theta}$ given in Algorithm 3.1 is consistent, i.e. $\hat{\theta} \stackrel{p}{\to} \theta_0$, and the confidence interval given in Algorithm 3.1 includes θ_0 with probability approaching the nominal level, i.e. $\lim_{n\to\infty} \mathbb{P}\left(\theta_0 \in \left[\hat{\theta} \pm c_a \frac{\hat{\sigma}}{\sqrt{n}}\right]\right) = 1 - a$.

Corollary 4.1 summarizes simple sufficient conditions for inference in terms of learning rates. Verifying these conditions is the goal of the next section.

5 Validity of bias correction

5.1 Goals

To employ Corollary 4.1, I must verify several conditions. In terms of the general inference problem, I must show

$$\mathbb{E}[Y - \gamma_0(W)]^2 \le \bar{\sigma}^2, \quad \mathbb{E}[m(W, \gamma)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma(W)]^2, \quad \|\alpha_0\|_{\infty} \le \bar{\alpha}, \quad \|\hat{\alpha}_{\ell}\|_{\infty} \le \bar{\alpha}'$$

Proposition 2.1 already previewed some of the relevant analysis. In what follows, I articulate the mild requirements for these conditions to hold. The same mild requirements will also furnish learning rates to satisfy

$$(\sqrt{\bar{L}^{(m)}} + \bar{\alpha} + \bar{\alpha}')\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \xrightarrow{p} 0, \quad \bar{\sigma}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} \xrightarrow{p} 0, \quad \sqrt{n}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \xrightarrow{p} 0$$

Note that the product condition $\sqrt{n}\sqrt{\mathcal{R}(\hat{\gamma}_\ell)}\sqrt{\mathcal{R}(\hat{\gamma}_\ell)}\overset{p}{\to} 0$ allows a tradeoff: one of the learning rates may be slow, as long as the other is sufficiently fast to compensate. While proving these results, I will abstract from sample splitting in order to improve readability. In other words, I will prove results for $(\hat{\gamma}, \hat{\alpha})$ trained on the full sample rather than I_ℓ^e .

5.2 Assumptions

I place standard, weak assumptions on the supports of (Y, W).

Assumption 5.1 (Original spaces). Assume

- 1. $W \in \mathcal{W}$ is a Polish space, i.e. separable and completely metrizable topological spaces
- 2. $Y \in \mathcal{Y} \subset \mathbb{R}$

A Polish space may be low, high, or infinite dimensional. Random variables with support in a Polish space may be discrete or continuous. Next, I place standard, weak assumptions on the RKHS \mathcal{H} .

Assumption 5.2 (RKHS regularity). *Assume*

- 1. k and $k^{(m)}$ are bounded. Formally, $\sup_{w \in \mathcal{W}} \|\phi(w)\|_{\mathcal{H}} \leq \sqrt{\kappa}$ and $\sup_{w \in \mathcal{W}} \|\phi^{(m)}(w)\|_{\mathcal{H}} \leq \sqrt{\kappa}$
- 2. $\phi(w)$ and $\phi^{(m)}(w)$ are measurable
- 3. $k^{(m)}$ is characteristic in its components that vary

Commonly used kernels are bounded. Boundedness implies Bochner integrability, which permits the exchange of expectation and inner product. Measurability is a similarly weak condition. The characteristic property ensures injectivity of the counterfactual kernel mean embedding $\mu^{(m)}$ with respect to its random components, and hence uniqueness of the RKHS representation [85, 86, 84].

Proposition 2.1 verifies $\mathbb{E}[m(W,\gamma)]^2 \leq \bar{L}^{(m)}\mathbb{E}[\gamma(W)]^2$ for a broad class of important examples. I now verify $\mathbb{E}[Y-\gamma_0(W)]^2 \leq \bar{\sigma}^2$, $\|\alpha_0\|_{\infty} \leq \bar{\alpha}$, and $\|\hat{\alpha}\|_{\infty} \leq \bar{\alpha}'$ under Assumptions 5.1 and 5.2.

Proposition 5.1 (Bounded noise and bounded Riesz representer). Suppose $\gamma_0 \in \mathcal{H}$ and Assumptions 5.1 and 5.2 hold. Then

$$\bar{\sigma}^2 = 2(\mathbb{E}[Y]^2 + \kappa \|\gamma_0\|_{\mathcal{H}}^2), \quad \bar{\alpha} = \sqrt{\kappa} \|\alpha_0^{\min}\|_{\mathcal{H}}, \quad \bar{\alpha}' = \frac{\kappa \|\alpha_0^{\min}\|_{\mathcal{H}}}{\sqrt{\lambda}}$$

The next two assumptions are the crux of the RKHS learning paradigm [16, 82]. The former is an assumption about the smoothness of γ_0 , parametrized by a scalar $c \in [1, 2]$. The latter is an assumption about the effective dimension of the RKHS, parametrized by a scalar $b \in (1, \infty]$.

Recall the spectral decomposition in Section 2. I define

$$\mathcal{H}^c = \left\{ f = \sum_{j=1}^{\infty} a_j \varphi_j : \sum_{j=1}^{\infty} \frac{a_j^2}{\lambda_j^c} < \infty \right\} \subset \mathcal{H}, \quad c \in [1, 2]$$

which is interpretable as an even smoother RKHS. We are now ready to state the smoothness assumption.

Assumption 5.3 (Smoothness). Assume $\gamma_0 \in \mathcal{H}^c$ for some $c \in [1, 2]$.

In other words, γ_0 is a particularly smooth element of \mathcal{H} ; it is well approximated by the leading terms in the series $\{\varphi_j\}$. This condition appears under the name of the *source* condition in both statistical learning [82, 16] and econometrics [17]. An abstract, equivalent version of this assumption is helpful for analysis.

Proposition 5.2 (Consequences of smoothness). The following statements are equivalent: (i) $\gamma_0 \in \mathcal{H}^c$; (ii) there exists some witness function $f \in \mathcal{H}$ s.t. $\gamma_0 = T^{\frac{c-1}{2}}f$; (iii) there exists some witness function $\tilde{f} \in \mathbb{L}_2$ s.t. $\gamma_0 = L_k^{\frac{c}{2}}\tilde{f}$. Moreover, $\gamma_0 \in \mathcal{H}^c$ implies $\alpha_0^{\min} \in \mathcal{H}^c$.

The effective dimension condition is also in terms of the spectrum.

Assumption 5.4 (Effective dimension). *If the RKHS is finite dimensional, write* $J \leq \beta < \infty$ *and* $b = \infty$. *If the RKHS is infinite dimensional, the eigenvalues* $\{\lambda_j\}$ *decay at a polynomial rate:* $\iota \leq j^b \cdot \lambda_j \leq \beta$ *for* $j \geq 1$, *where* $\iota, \beta > 0$ *and* $b \in (1, \infty)$.

Note that $\{\lambda_j\}$ are also the eigenvalues of the covariance operator T.

Proposition 5.3 (Consequences of effective dimension). If Assumption 5.4 holds then $\lambda_j = \Theta(j^{-b})$. Moreover, $L_k(\cdot) = \sum_{j=1}^{\infty} \lambda_j \langle \varphi_j, \cdot \rangle_2 \cdot \varphi_j$ if and only if $T(\cdot) = \sum_{j=1}^{\infty} \lambda_j \langle e_j, \cdot \rangle_2 \cdot e_j$ where $e_j = \sqrt{\lambda_j} \varphi_j$.

The spectrum of a random vector's covariance characterizes its dimension. In this sense, the assumption on spectral decay quantifies effective dimension. Following [16], I denote by $\mathcal{P}(b,c)$ the class of distributions that satisfy Assumptions 5.3 and 5.4.

Next I relate these assumptions to the Sobolev space introduced in Definition 2.1.

Proposition 5.4 (Sobolev space as RKHS [29, 11, 30]). A Sobolev space \mathbb{H}_2^s with $s > \frac{d}{2}$ is an RKHS. Suppose $\mathcal{H} = \mathbb{H}_2^s$ is chosen as the RKHS for estimation. If $\gamma_0 \in \mathbb{H}_2^{s_0}$ then $c = \frac{s_0}{s}$. Moreover, $b = \frac{2s}{d}$.

An RKHS is a generalization of a Sobolev space, and its assumptions generalize those typically made for Sobolev spaces.

In the RKHS setting, the approximation properties of γ_0 immediately imply approximation properties of α_0^{\min} . The fact that $\gamma_0 \in \mathcal{H}^c$ implies $\alpha_0^{\min} \in \mathcal{H}^c$ follows from Proposition 2.3. Likewise the fact that γ_0 and α_0^{\min} are governed by the same covariance operator and hence share the same effective dimension follows from Proposition 2.3. This symmetry does *not* hold for Dantzig or Lasso approaches to debiasing, where the analyst must place separate approximation assumptions for γ_0 and α_0^{\min} . The favorable geometry of the RKHS sets it apart from other approaches, and it allows us to use only learning theory assumptions to prove inference.

5.3 Guarantees

Assumptions 5.1, 5.2, 5.3, and 5.4 are classic assumptions in RKHS learning theory. I quote a classic result on the mean square rate for kernel ridge regression. This result also requires a weak condition on the noise

Assumption 5.5 (Noise). $Y = \gamma_0(W) + \varepsilon$ where ε is bounded, Gaussian, or subGaussian.

Theorem 5.1 (Kernel ridge regression: MSE; Theorems 1 and 2 of [16]). Suppose Assumptions 5.1, 5.2, 5.3, 5.4, and 5.5 hold. Calibrate the ridge regularization sequence s.t.

$$\lambda = \begin{cases} n^{-\frac{1}{2}} & \text{if} \quad b = \infty \\ n^{-\frac{b}{bc+1}} & \text{if} \quad b \in (1, \infty), \ c \in (1, 2] \\ \ln^{\frac{b}{b+1}}(n) \cdot n^{-\frac{b}{b+1}} & \text{if} \quad b \in (1, \infty), \ c = 1 \end{cases}$$

Then kernel ridge regression $\hat{\gamma}$ trained on $\{W_i\}_{i\in[n]}$ satisfies

$$\lim_{\tau \to \infty} \lim_{n \to \infty} \sup_{\rho \in \mathcal{P}(b,c)} \mathbb{P}_{\{W_i\} \sim \rho^n} (\mathcal{R}(\hat{\gamma}) > \tau \cdot r_n) = 0$$

where $\mathcal{R}(\hat{\gamma}) = \mathbb{E}[\{\hat{\gamma}(W) - \gamma_0(W)\}^2 | \{W_i\}_{i \in [n]}]$ and

$$r_n = \begin{cases} n^{-1} & \text{if } b = \infty \\ n^{-\frac{bc}{bc+1}} & \text{if } b \in (1, \infty), c \in (1, 2] \\ \ln^{\frac{b}{b+1}}(n) \cdot n^{-\frac{b}{b+1}} & \text{if } b \in (1, \infty), c = 1 \end{cases}$$

Moreover, the rate is optimal when $b \in (1, \infty)$ and $c \in (1, 2]$. It is optimal up to a logarithmic factor when $b \in (1, \infty)$ and c = 1.

Recall that $b=\infty$ is the finite dimensional regime; $b\in(1,\infty),\ c\in(1,2]$ is the infinite dimensional regime with polynomial spectral decay and additional smoothness; $b\in(1,\infty),\ c=1$ is the infinite dimensional regime with polynomial spectral decay and no additional smoothness.

Remarkably, an almost identical result is possible for the kernel ridge Riesz representer. This is the second main result of the paper. The result is almost for free in the sense that it only requires replacing Assumption 5.5 with Assumptions 2.1. Indeed, the core techniques are the same.

Theorem 5.2 (Kernel ridge Riesz representer: MSE). Suppose Assumption 2.1 as well Assumptions 5.1, 5.2, 5.3, and 5.4 hold. Calibrate the ridge regularization sequence λ as in Theorem 5.1. Then the kernel ridge Riesz representer $\hat{\alpha}$ trained on $\{W_i\}_{i\in[n]}$ satisfies

$$\lim_{\tau \to \infty} \lim_{n \to \infty} \sup_{\rho \in \mathcal{P}(b,c)} \mathbb{P}_{\{W_i\} \sim \rho^n} (\mathcal{R}(\hat{\alpha}) > \tau \cdot r_n) = 0$$

where
$$\mathcal{R}(\hat{\alpha}) = \mathbb{E}[\{\hat{\alpha}(W) - \alpha_0^{\min}(W)\}^2 | \{W_i\}_{i \in [n]}]$$
 and r_n is as in Theorem 5.1

Theorem 5.2 demonstrates that an extremely simple kernel ridge Riesz representer attains the optimal rate in many cases. The ridge regularization sequence for $\hat{\alpha}$ is the same sequence that balances bias and variance for $\hat{\gamma}$ in Theorem 5.1. Therefore the same tuning procedure for $\hat{\gamma}$ also applies to $\hat{\alpha}$. This symmetry does *not* hold for Dantzig and Lasso approaches to debiasing, where the analyst must tune $\hat{\gamma}$ and $\hat{\alpha}$ separately. Again, the favorable geometry of the RKHS sets it apart from other approaches.

This fast rate aligns with the fast rate of [23, Corollary 7] for an adversarial estimator of the Riesz representer. The rate in [23, Corollary 7] is in terms of a critical radius that adapts to the empirical eigenvalues of the kernel matrix, while the rate in Theorem 5.2 adapts to the smoothness c and effective dimension b defined above. Both the empirical eigenvalues and the effective dimension reflect the kernel's spectral characteristics; both Theorem 5.2 and [23, Corollary 7] are adaptive to the same underlying phenomena.

Corollary 5.1 (Confidence interval). Suppose the assumptions of Theorems 5.1 and 5.2 hold as well as the regularity condition on moments $\left\{ \left(\frac{\eta}{\sigma} \right)^3 + \zeta^2 \right\} n^{-\frac{1}{2}} \to 0$. Then for any $c \in (1,2]$ and $b \in (1,\infty]$, the estimator $\hat{\theta}$ is consistent, i.e. $\hat{\theta} \stackrel{p}{\to} \theta_0$, and the confidence interval includes θ_0 with probability approaching the nominal level, i.e. $\lim_{n \to \infty} \mathbb{P}\left(\theta_0 \in \left[\hat{\theta} \pm c_a \frac{\hat{\sigma}}{\sqrt{n}}\right]\right) = 1 - a$.

Recall that in Assumption 5.3, c=1 means γ_0 is correctly specified as an element in \mathcal{H} , while c>1 means γ_0 is in the interior of \mathcal{H} . As long as γ_0 is in the interior of the RKHS, rather than at the boundary, we can construct valid confidence intervals by kernel matrix operations. Meanwhile, the RKHS may be finite or infinite dimensional, as long as the spectrum decays at some polynomial rate $\lambda_j = \Theta(j^{-b})$ with b>1.

5.4 Trimming

Examining the proof of Corollary 5.1, the only condition that fails to hold when c=1 is $\bar{\alpha}'\sqrt{\mathcal{R}(\hat{\gamma})} \stackrel{p}{\to} 0$. Specifically, the bound on $\bar{\alpha}'$ given in Proposition 5.1 diverges too quickly. A solution is to trim the kernel ridge Riesz representer using a known bound on the minimal Riesz representer α_0^{\min} . In particular, consider the following estimator.

Algorithm 5.1 (Kernel ridge Riesz representer: Trimming). *Given* $\hat{\alpha}$ *calculated from Algorithm 3.2, set*

$$\tilde{\alpha}(w) = \begin{cases} -\bar{\alpha} & \text{if} \quad \hat{\alpha}(w) < -\bar{\alpha} \\ \hat{\alpha}(w) & \text{if} \quad \hat{\alpha}(w) \in [-\bar{\alpha}, \bar{\alpha}] \\ \bar{\alpha} & \text{if} \quad \hat{\alpha}(w) > \bar{\alpha} \end{cases}$$

Denote by $(\tilde{\theta}, \tilde{\sigma}^2)$ the estimators for (θ_0, σ^2) calculated from Algorithm 3.1 using $(\hat{\gamma}, \tilde{\alpha})$ rather than $(\hat{\gamma}, \hat{\alpha})$.

Corollary 5.2 (Confidence interval with trimming). Suppose the assumptions of Theorems 5.1 and 5.2 hold as well as the regularity condition on moments $\left\{ \left(\frac{\eta}{\sigma} \right)^3 + \zeta^2 \right\} n^{-\frac{1}{2}} \to 0$. Then for any $c \in [1,2]$ and $b \in (1,\infty]$, the estimator $\tilde{\theta}$ is consistent, i.e. $\tilde{\theta} \stackrel{p}{\to} \theta_0$, and the confidence interval includes θ_0 with probability approaching the nominal level, i.e. $\lim_{n\to\infty} \mathbb{P}\left(\theta_0 \in \left[\tilde{\theta} \pm c_a \frac{\tilde{\sigma}}{\sqrt{n}}\right]\right) = 1-a$.

Now the case c=1 is allowed. Since c=1 simply means $\gamma_0 \in \mathcal{H}$, the inference result allows us to actually relax Assumption 5.3. In other words, smoothness beyond correct specification hastens learning but is irrelevant to inference. Meanwhile, the RKHS may be finite or infinite dimensional, as long as the spectrum decays at some polynomial rate $\lambda_j = \Theta(j^{-b})$ with b>1.

6 Conclusion

For any kernel method with a mean square learning rate, I propose a practical procedure based on bias correction and sample splitting to calculate confidence intervals for its functionals. The inferential results are almost free in the sense that classic learning theory assumptions, together with a mean square continuity condition, are enough to justify the procedure. By providing relatively simple confidence intervals for functionals of kernel ridge regression, I facilitate their use in social science and epidemiology.

Acknowledgments and disclosure of funding

I am grateful to Alberto Abadie, Anish Agarwal, Victor Chernozhukov, Arthur Gretton, Anna Mikusheva, Whitney Newey, Devavrat Shah, Vasilis Syrgkanis, Suhas Vijaykumar, and Yinchu Zhu for helpful discussions.

References

- [1] Chunrong Ai and Xiaohong Chen. Efficient estimation of models with conditional moment restrictions containing unknown functions. *Econometrica*, 71(6):1795–1843, 2003.
- [2] Chunrong Ai and Xiaohong Chen. Estimation of possibly misspecified semiparametric conditional moment restriction models with different conditioning variables. *Journal of Econometrics*, 141(1):5–43, 2007.
- [3] Chunrong Ai and Xiaohong Chen. The semiparametric efficiency bound for models of sequential moment restrictions containing unknown functions. *Journal of Econometrics*, 170(2):442–457, 2012.
- [4] Susan Athey, Guido W Imbens, and Stefan Wager. Approximate residual balancing: Debiased inference of average treatment effects in high dimensions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 80(4):597–623, 2018.
- [5] Frank Bauer, Sergei Pereverzev, and Lorenzo Rosasco. On regularization algorithms in learning theory. *Journal of Complexity*, 23(1):52–72, 2007.
- [6] Alexandre Belloni, Victor Chernozhukov, Ivan Fernández-Val, and Christian Hansen. Program evaluation and causal inference with high-dimensional data. *Econometrica*, 85(1):233–298, 2017.
- [7] Alexandre Belloni, Victor Chernozhukov, and Christian Hansen. Inference for high-dimensional sparse econometric models. *arXiv:1201.0220*, 2011.
- [8] Alexandre Belloni, Victor Chernozhukov, and Christian Hansen. Inference on treatment effects after selection among high-dimensional controls. *The Review of Economic Studies*, 81(2):608–650, 2014.
- [9] Alexandre Belloni, Victor Chernozhukov, and Kengo Kato. Uniform post-selection inference for least absolute deviation regression and other Z-estimation problems. *Biometrika*, 102(1):77–94, 2014.

- [10] Alexandre Belloni, Victor Chernozhukov, and Lie Wang. Pivotal estimation via square-root lasso in nonparametric regression. *The Annals of Statistics*, 42(2):757–788, 2014.
- [11] Alain Berlinet and Christine Thomas-Agnan. *Reproducing Kernel Hilbert Spaces in Probability and Statistics*. Springer Science & Business Media, 2011.
- [12] Peter J Bickel. On adaptive estimation. *The Annals of Statistics*, pages 647–671, 1982.
- [13] Peter J Bickel, Chris AJ Klaassen, Peter J Bickel, Ya'acov Ritov, J Klaassen, Jon A Wellner, and YA'Acov Ritov. Efficient and Adaptive Estimation for Semiparametric Models, volume 4. Johns Hopkins University Press Baltimore, 1993.
- [14] Peter J Bickel and Yaacov Ritov. Estimating integrated squared density derivatives: Sharp best order of convergence estimates. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 381–393, 1988.
- [15] Jelena Bradic and Mladen Kolar. Uniform inference for high-dimensional quantile regression: Linear functionals and regression rank scores. *arXiv:1702.06209*, 2017.
- [16] Andrea Caponnetto and Ernesto De Vito. Optimal rates for the regularized least-squares algorithm. *Foundations of Computational Mathematics*, 7(3):331–368, 2007.
- [17] Marine Carrasco, Jean-Pierre Florens, and Eric Renault. Linear inverse problems in structural econometrics estimation based on spectral decomposition and regularization. *Handbook of Econometrics*, 6:5633–5751, 2007.
- [18] Xiaohong Chen. Large sample sieve estimation of semi-nonparametric models. *Handbook of Econometrics*, 6:5549–5632, 2007.
- [19] Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, and James Robins. Double/debiased machine learning for treatment and structural parameters: Double/debiased machine learning. *The Econometrics Journal*, 21(1), 2018.
- [20] Victor Chernozhukov, Juan Carlos Escanciano, Hidehiko Ichimura, Whitney K Newey, and James M Robins. Locally robust semiparametric estimation. *arXiv:1608.00033*, 2016.
- [21] Victor Chernozhukov, Christian Hansen, and Martin Spindler. Valid post-selection and post-regularization inference: An elementary, general approach. *Annual Review of Economics*, 7(1):649–688, 2015.
- [22] Victor Chernozhukov, Whitney Newey, and Rahul Singh. De-biased machine learning of global and local parameters using regularized Riesz representers. *arXiv:1802.08667*, 2018.
- [23] Victor Chernozhukov, Whitney Newey, Rahul Singh, and Vasilis Syrgkanis. Adversarial estimation of Riesz representers. arXiv:2101.00009, 2020.
- [24] Victor Chernozhukov, Whitney K Newey, and Rahul Singh. Automatic debiased machine learning of causal and structural effects. *arXiv:1809.05224*, 2018.
- [25] Victor Chernozhukov, Whitney K Newey, and Rahul Singh. Automatic debiased machine learning of causal and structural effects, 2020.
- [26] Felipe Cucker and Steve Smale. On the mathematical foundations of learning. *Bulletin of the American Mathematical Society*, 39(1):1–49, 2002.
- [27] Iván Díaz and Mark J van der Laan. Targeted data adaptive estimation of the causal dose–response curve. *Journal of Causal Inference*, 1(2):171–192, 2013.
- [28] Nishanth Dikkala, Greg Lewis, Lester Mackey, and Vasilis Syrgkanis. Minimax estimation of conditional moment models. *arXiv*:2006.07201, 2020.
- [29] David E Edmunds and Hans Triebel. Function Spaces, Entropy Numbers and Differential Operators. Cambridge University Press, 2008.
- [30] Simon Fischer and Ingo Steinwart. Sobolev norm learning rates for regularized least-squares algorithm. *arXiv:1702.07254*, 2017.
- [31] Dylan J Foster and Vasilis Syrgkanis. Orthogonal statistical learning. *arXiv:1901.09036*, 2019.
- [32] Kenji Fukumizu, Francis R Bach, and Michael I Jordan. Dimensionality reduction for supervised learning with reproducing kernel hilbert spaces. *Journal of Machine Learning Research*, 5(Jan):73–99, 2004.

- [33] Rafail Z Hasminskii and Ildar A Ibragimov. On the nonparametric estimation of functionals. In *Proceedings of the Second Prague Symposium on Asymptotic Statistics*, 1979.
- [34] David A Hirshberg and Stefan Wager. Debiased inference of average partial effects in single-index models. *arXiv*:1811.02547, 2018.
- [35] David A Hirshberg and Stefan Wager. Augmented minimax linear estimation. arXiv:1712.00038v5, 2019.
- [36] I Ibragimov and R Has'minskii. Statistical estimation, vol. 16 of. *Applications of Mathematics*, 1981.
- [37] Hidehiko Ichimura and Whitney K Newey. The influence function of semiparametric estimators. arXiv:1508.01378, 2015.
- [38] Guido W Imbens and Whitney K Newey. Identification and estimation of triangular simultaneous equations models without additivity. *Econometrica*, 77(5):1481–1512, 2009.
- [39] Jana Jankova and Sara Van De Geer. Confidence intervals for high-dimensional inverse covariance estimation. *Electronic Journal of Statistics*, 9(1):1205–1229, 2015.
- [40] Jana Jankova and Sara Van De Geer. Confidence regions for high-dimensional generalized linear models under sparsity. *arXiv:1610.01353*, 2016.
- [41] Jana Jankova and Sara Van De Geer. Semiparametric efficiency bounds for high-dimensional models. *The Annals of Statistics*, 46(5):2336–2359, 2018.
- [42] Adel Javanmard and Andrea Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *The Journal of Machine Learning Research*, 15(1):2869–2909, 2014.
- [43] Adel Javanmard and Andrea Montanari. Hypothesis testing in high-dimensional regression under the Gaussian random design model: Asymptotic theory. *IEEE Transactions on Information Theory*, 60(10):6522–6554, 2014.
- [44] Adel Javanmard and Andrea Montanari. Debiasing the lasso: Optimal sample size for Gaussian designs. *The Annals of Statistics*, 46(6A):2593–2622, 2018.
- [45] Edward H Kennedy. Optimal doubly robust estimation of heterogeneous causal effects. *arXiv:2004.14497*, 2020.
- [46] Edward H Kennedy, Zongming Ma, Matthew D McHugh, and Dylan S Small. Nonparametric methods for doubly robust estimation of continuous treatment effects. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 79(4):1229, 2017.
- [47] George Kimeldorf and Grace Wahba. Some results on Tchebycheffian spline functions. *Journal of Mathematical Analysis and Applications*, 33(1):82–95, 1971.
- [48] Chris AJ Klaassen. Consistent estimation of the influence function of locally asymptotically linear estimators. *The Annals of Statistics*, pages 1548–1562, 1987.
- [49] Michael R Kosorok. *Introduction to Empirical Processes and Semiparametric Inference*. Springer Science & Business Media, 2007.
- [50] B Ya Levit. On the efficiency of a class of non-parametric estimates. *Theory of Probability & Its Applications*, 20(4):723–740, 1976.
- [51] Alexander R Luedtke and Mark J Van Der Laan. Statistical inference for the mean outcome under a possibly non-unique optimal treatment strategy. *Annals of Statistics*, 44(2):713, 2016.
- [52] Krikamol Muandet, Arash Mehrjou, Si Kai Lee, and Anant Raj. Dual iv: A single stage instrumental variable regression. *arXiv:1910.12358*, 2019.
- [53] Whitney K Newey. The asymptotic variance of semiparametric estimators. *Econometrica*, pages 1349–1382, 1994.
- [54] Whitney K Newey. Convergence rates and asymptotic normality for series estimators. *Journal of Econometrics*, 79(1):147–168, 1997.
- [55] Whitney K Newey, Fushing Hsieh, and James M Robins. Undersmoothing and bias corrected functional estimation. Technical report, MIT Department of Economics, 1998.
- [56] Whitney K Newey, Fushing Hsieh, and James M Robins. Twicing kernels and a small bias property of semiparametric estimators. *Econometrica*, 72(3):947–962, 2004.

- [57] Whitney K Newey and James R Robins. Cross-fitting and fast remainder rates for semiparametric estimation. arXiv:1801.09138, 2018.
- [58] Matey Neykov, Yang Ning, Jun S Liu, and Han Liu. A unified theory of confidence regions and testing for high-dimensional estimating equations. *Statistical Science*, 33(3):427–443, 2018.
- [59] Jerzy Neyman. Optimal asymptotic tests of composite statistical hypotheses. In *Probability and Statistics*, page 416–444. Wiley, 1959.
- [60] Jerzy Neyman. C (α) tests and their use. Sankhyā: The Indian Journal of Statistics, Series A, pages 1–21, 1979.
- [61] Xinkun Nie and Stefan Wager. Quasi-oracle estimation of heterogeneous treatment effects. *arXiv:1712.04912*, 2017.
- [62] Yang Ning and Han Liu. A general theory of hypothesis tests and confidence regions for sparse high dimensional models. *The Annals of Statistics*, 45(1):158–195, 2017.
- [63] Johann Pfanzagl. Lecture notes in statistics. *Contributions to a General Asymptotic Statistical Theory*, 13, 1982.
- [64] Zhao Ren, Tingni Sun, Cun-Hui Zhang, and Harrison H Zhou. Asymptotic normality and optimalities in estimation of large Gaussian graphical models. *The Annals of Statistics*, 43(3):991–1026, 2015.
- [65] James Robins, Lingling Li, Eric Tchetgen, Aad van der Vaart, et al. Higher order influence functions and minimax estimation of nonlinear functionals. In *Probability and Statistics: Essays in Honor of David A. Freedman*, pages 335–421. Institute of Mathematical Statistics, 2008.
- [66] James Robins, Mariela Sued, Quanhong Lei-Gomez, and Andrea Rotnitzky. Comment on "performance of double-robust estimators when inverse probability weights are highly variable". Statistical Science, 22(4):544–559, 2007.
- [67] James M Robins and Andrea Rotnitzky. Semiparametric efficiency in multivariate regression models with missing data. *Journal of the American Statistical Association*, 90(429):122–129, 1995
- [68] James M Robins, Andrea Rotnitzky, and Lue Ping Zhao. Analysis of semiparametric regression models for repeated outcomes in the presence of missing data. *Journal of the American Statistical Association*, 90(429):106–121, 1995.
- [69] Peter M Robinson. Root-n-consistent semiparametric regression. *Econometrica*, pages 931–954, 1988.
- [70] Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983.
- [71] Dominik Rothenhäusler and Bin Yu. Incremental causal effects. arXiv:1907.13258, 2019.
- [72] Dan Rubin and Mark J van der Laan. A general imputation methodology for nonparametric regression with censored data. Technical report, UC Berkeley Division of Biostatistics, 2005.
- [73] Daniel Rubin and Mark J van der Laan. Extending marginal structural models through local, penalized, and additive learning. Technical report, UC Berkeley Division of Biostatistics, 2006.
- [74] Daniel O Scharfstein, Andrea Rotnitzky, and James M Robins. Adjusting for nonignorable drop-out using semiparametric nonresponse models. *Journal of the American Statistical As*sociation, 94(448):1096–1120, 1999.
- [75] Anton Schick. On asymptotically efficient estimation in semiparametric models. *The Annals of Statistics*, 14(3):1139–1151, 1986.
- [76] Bernhard Schölkopf, Ralf Herbrich, and Alex J Smola. A generalized representer theorem. In *International conference on computational learning theory*, pages 416–426. Springer, 2001.
- [77] Irina Shevtsova. On the absolute constants in the Berry-Esseen type inequalities for identically distributed summands. *arXiv:1111.6554*, 2011.
- [78] Rahul Singh. Kernel methods for unobserved confounding: Negative controls, proxies, and instruments. *arXiv*:2012.10315, 2020.

- [79] Rahul Singh, Maneesh Sahani, and Arthur Gretton. Kernel instrumental variable regression. In *Advances in Neural Information Processing Systems*, pages 4595–4607, 2019.
- [80] Rahul Singh and Liyang Sun. De-biased machine learning in intrumental variable models for treatment effects. *arXiv*:1909.05244, 2019.
- [81] Rahul Singh, Liyuan Xu, and Arthur Gretton. Kernel methods for policy evaluation: Treatment effects, mediation analysis, and off-policy planning. *arXiv:2010.04855*, 2020.
- [82] Steve Smale and Ding-Xuan Zhou. Learning theory estimates via integral operators and their approximations. *Constructive Approximation*, 26(2):153–172, 2007.
- [83] Alex Smola, Arthur Gretton, Le Song, and Bernhard Schölkopf. A Hilbert space embedding for distributions. In *International Conference on Algorithmic Learning Theory*, pages 13–31, 2007.
- [84] Bharath Sriperumbudur. On the optimal estimation of probability measures in weak and strong topologies. *Bernoulli*, 22(3):1839–1893, 2016.
- [85] Bharath Sriperumbudur, Kenji Fukumizu, and Gert Lanckriet. On the relation between universality, characteristic kernels and RKHS embedding of measures. In *International Conference on Artificial Intelligence and Statistics*, pages 773–780, 2010.
- [86] Bharath K Sriperumbudur, Kenji Fukumizu, and Gert RG Lanckriet. Universality, characteristic kernels and RKHS embedding of measures. *Journal of Machine Learning Research*, 12(7), 2011.
- [87] Ingo Steinwart and Andreas Christmann. Support Vector Machines. Springer Science & Business Media, 2008.
- [88] Dougal J Sutherland. Fixing an error in Caponnetto and de Vito (2007). arXiv:1702.02982, 2017.
- [89] B Toth and MJ van der Laan. TMLE for marginal structural models based on an instrument. Technical report, UC Berkeley Division of Biostatistics, 2016.
- [90] Anastasios Tsiatis. *Semiparametric Theory and Missing Data*. Springer Science & Business Media, 2007.
- [91] Sara Van de Geer, Peter Bühlmann, Ya'acov Ritov, and Ruben Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202, 2014.
- [92] Mark J van der Laan and Alexander R Luedtke. Targeted learning of an optimal dynamic treatment, and statistical inference for its mean outcome. Technical report, UC Berkeley Division of Biostatistics, 2014.
- [93] Mark J Van der Laan and Sherri Rose. *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer Science & Business Media, 2011.
- [94] Mark J Van Der Laan and Daniel Rubin. Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1), 2006.
- [95] Aad Van Der Vaart et al. On differentiable functionals. The Annals of Statistics, 19(1):178–204, 1991.
- [96] Aad W Van der Vaart. Asymptotic Statistics, volume 3. Cambridge University Press, 2000.
- [97] Cun-Hui Zhang and Stephanie S Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B* (Statistical Methodology), 76(1):217–242, 2014.
- [98] Wenjing Zheng and Mark J Van Der Laan. Asymptotic theory for cross-validated targeted maximum likelihood estimation. 2010.
- [99] Yinchu Zhu and Jelena Bradic. Breaking the curse of dimensionality in regression. *arXiv:1708.00430.*, 2017.
- [100] Yinchu Zhu and Jelena Bradic. Linear hypothesis testing in dense high-dimensional linear models. *Journal of the American Statistical Association*, 113(524):1583–1600, 2018.

A Framework

Proof of Proposition 2.1. I consider each functional of kernel ridge regression.

1. Evaluation.

$$\mathbb{E}[\gamma(w)]^2 = \{\gamma(w)\}^2 = \left\{\mathbb{E}\left[\gamma(W)\frac{1\{W=w\}}{\mathbb{P}(w)}\right]\right\}^2 \leq \left\{\frac{\mathbb{E}[\gamma(W)]}{\mathbb{P}(w)}\right\}^2 \leq \frac{\mathbb{E}[\gamma(W)]^2}{\mathbb{P}(w)^2}$$
 i.e. $\bar{L}^{(m)} = \mathbb{P}(w)^{-2}$

2. Treatment effects. For ATE,

$$\mathbb{E}[\gamma(d,X)]^2 = \mathbb{E}\left[\frac{1\{D=d\}}{\mathbb{P}(d|X)}\gamma(D,X)^2\right] \le \frac{\mathbb{E}\left[\gamma(D,X)\right]^2}{\inf_x \mathbb{P}(d|x)}$$

i.e. $\bar{L}^{(m)} = \{\inf_x \mathbb{P}(d|x)\}^{-1}$. ATT and CATE are similar.

Proof of Proposition 2.2. I prove the result for Example 2.2. The argument for Example 2.4 is similar. I proceed in steps

1. Sobolev and RKHS preliminaries.

By [11, Theorem 132], \mathbb{H}_2^s is an RKHS iff $s>\frac{d}{2}$. By [11, Theorem 133], the r-order derivative operator is continuous and linear iff $s>\frac{d}{2}+r$.

In an RKHS, $\gamma(w) = \langle \gamma, \phi(w) \rangle_{\mathcal{H}}$ where $\phi: w \mapsto k(w,\cdot)$. Moreover, if the derivative $\nabla_w \nabla_{w'} k(w,w')$ exists and is continuous then by [87, Lemma 4.34] $\nabla \gamma(w) = \langle \gamma, \nabla \phi(w) \rangle_{\mathcal{H}}$.

Let $\mathcal{H}=\mathbb{H}_2^s$ and set r=1. \mathcal{H} is an RKHS and the derivative operator is bounded if $s>\frac{d}{2}+1$. I wish to verify mean square continuity of the derivative functional for this particular function space.

2. Main argument. To begin, write

$$\mathbb{E}[\gamma(W)]^2 = \mathbb{E}\langle\gamma,\phi(W)\rangle_{\mathcal{H}}^2 = \mathbb{E}\langle\gamma,[\phi(W)\otimes\phi(W)]\gamma\rangle_{\mathcal{H}} = \langle\gamma,T\gamma\rangle_{\mathcal{H}}$$
 where $T:=\mathbb{E}[\phi(W)\otimes\phi(W)].$

Since ∇ is a continuous linear operator over \mathcal{H} , its adjoint ∇^* exists and both are bounded with the same operator norm. Write

$$\nabla \gamma(w) = \langle \gamma, \nabla \phi(w) \rangle_{\mathcal{H}} = \langle \nabla^* \gamma, \phi(w) \rangle_{\mathcal{H}}$$

Hence

$$\mathbb{E}[\nabla \gamma(W)]^2 = \mathbb{E}\langle \nabla^* \gamma, \phi(W) \rangle_{\mathcal{H}}^2 = \mathbb{E}\langle \nabla^* \gamma, [\phi(W) \otimes \phi(W)] \nabla^* \gamma \rangle_{\mathcal{H}} = \langle \nabla^* \gamma, T \nabla^* \gamma \rangle_{\mathcal{H}}$$

Next observe

$$\langle \nabla^* \gamma, T \nabla^* \gamma \rangle_{\mathcal{H}} = \langle \gamma, \nabla T \nabla^* \gamma \rangle_{\mathcal{H}} = Tr(\nabla T \nabla^* [\gamma \otimes \gamma]) \leq \|\nabla\|_{op} Tr(T \nabla^* [\gamma \otimes \gamma])$$

Moreover

$$Tr(T\nabla^*[\gamma \otimes \gamma]) = Tr(\nabla^*[\gamma \otimes \gamma]T) \leq \|\nabla\|_{op}Tr([\gamma \otimes \gamma]T) = \|\nabla\|_{op}\langle \gamma, T\gamma \rangle_{\mathcal{H}}$$

In summary

$$\mathbb{E}[\nabla \gamma(W)]^2 \leq \|\nabla\|_{on}^2 \langle \gamma, T\gamma \rangle_{\mathcal{H}} = \|\nabla\|_{on}^2 \mathbb{E}[\gamma(W)]^2$$

i.e.
$$\bar{L}^{(m)} = \|\nabla\|_{op}^2$$
.

Proof of Proposition 2.3. First, I show that the functional $\gamma \mapsto \mathbb{E}[m(W,\gamma)]$ is bounded as a functional over \mathbb{L}_2 . Write

$$\{|\mathbb{E}[m(W,\gamma)]|\}^2 \leq \{\mathbb{E}|m(W,\gamma)|\}^2 \leq \mathbb{E}[m(W,\gamma)]^2 \leq \bar{L}^{(m)}\mathbb{E}[\gamma(W)]^2$$

Therefore

$$\frac{\|\mathbb{E}[m(W,\gamma)]\|}{\|\gamma\|_2} \le \frac{\sqrt{\bar{L}^{(m)}} \|\gamma\|_2}{\|\gamma\|_2} \le \sqrt{\bar{L}^{(m)}}$$

Hence Riesz representation theorem w.r.t \mathbb{L}_2 implies existence of the RR $\alpha_0 \in \mathbb{L}_2$. [22, Lemma S3.1] guarantees existence of the unique minimal Riesz representer $\alpha_0^{\min} \in closure(span(\Gamma))$. \square

B Algorithm

Proof of Proposition 3.1. Recall that $\gamma_0 \in \Gamma$ implies that the minimal representer $\alpha_0^{\min} \in \Gamma \subset \mathcal{H}$ by Proposition 2.3. Trivially,

$$\alpha_0^{\min} \in \operatorname*{argmin}_{\alpha \in \mathcal{H}} \mathcal{R}(\alpha), \quad \mathcal{R}(\alpha) = \mathbb{E}[\alpha_0^{\min}(W) - \alpha(W)]^2$$

Note that

$$\mathcal{R}(\alpha) = \mathbb{E}[\alpha_0^{\min}(W) - \alpha(W)]^2$$

$$= \mathbb{E}[\alpha_0^{\min}(W)]^2 - 2\mathbb{E}[\alpha_0(W)\alpha(W)] + \mathbb{E}[\alpha(W)]^2$$

$$= C - 2\mathbb{E}[m(W, \alpha)] + \mathbb{E}[\alpha(W)]^2$$

Boundedness of the kernel implies Bochner integrability of $\phi(\cdot)$ and $\phi^{(m)}(\cdot)$, which then allows us to exchange expectation and inner product. By the RKHS representation property, linearity of m, and Bochner integrability of $\phi(\cdot)$ and $\phi^{(m)}(\cdot)$

$$\mathbb{E}[m(W,\alpha)] = \mathbb{E}[\langle \alpha, \phi^{(m)}(w) \rangle_{\mathcal{H}}] = \langle \alpha, \mu^{(m)} \rangle_{\mathcal{H}}, \quad \mu^{(m)} = \mathbb{E}[\phi^{(m)}(W)]$$

and

$$\mathbb{E}[\alpha(W)]^2 = \mathbb{E}[\langle \alpha, \phi(W) \rangle_{\mathcal{H}}^2] = \langle \alpha, T\alpha \rangle_{\mathcal{H}}, \quad T = \mathbb{E}[\phi(W) \otimes \phi(W)]$$

Proof of Proposition 3.2. Write

$$K_{ij}^{(1)} = \langle \phi(W_i), \phi(W_j) \rangle_{\mathcal{H}} = \langle \phi(D_i) \otimes \phi(X_i), \phi(D_j) \otimes \phi(X_j) \rangle_{\mathcal{H}} = k(D_i, D_j) k(X_i, X_j)$$

$$K_{ij}^{(2)} = \langle \phi(W_i), \phi^{(m)}(W_j) \rangle_{\mathcal{H}} = \langle \phi(D_i) \otimes \phi(X_i), \phi(d) \otimes \phi(X_j) \rangle_{\mathcal{H}} = k(D_i, d) k(X_i, X_j)$$

$$K_{ij}^{(4)} = \langle \phi^{(m)}(W_i), \phi^{(m)}(W_j) \rangle_{\mathcal{H}} = \langle \phi(d) \otimes \phi(X_i), \phi(d) \otimes \phi(X_j) \rangle_{\mathcal{H}} = k(d, d) k(X_i, X_j)$$

Proof of Proposition 3.3. Note that

$$\hat{\mu}^{(m)} = \frac{1}{n} \sum_{i=1}^{n} \phi^{(m)}(W_i)$$

and

$$\hat{T}\alpha = \frac{1}{n} \sum_{i=1}^{n} [\phi(W_i) \otimes \phi(W_i)]\alpha = \frac{1}{n} \sum_{i=1}^{n} \langle \alpha, \phi(W_i) \rangle_{\mathcal{H}} \phi(W_i)$$

Write the objective as

$$\mathcal{L}_{\lambda}^{n}(\alpha) = \frac{1}{n} \sum_{i=1}^{n} \left\{ -2\langle \alpha, \phi^{(m)}(W_{i}) \rangle_{\mathcal{H}} + \langle \alpha, \phi(W_{i}) \rangle_{\mathcal{H}}^{2} \right\} + \lambda \|\alpha\|_{\mathcal{H}}^{2}$$

Recall that for an RKHS, evaluation is a continuous functional represented as the inner product with the feature map. Due to the ridge penalty, the stated objective is coercive and strongly convex w.r.t α . Hence it has a unique minimizer $\hat{\alpha}$ that obtains the minimum.

Write $\hat{\alpha} = \hat{\alpha}_n + \hat{\alpha}_n^{\perp}$ where $\hat{\alpha}_n \in row(\Psi)$ and $\hat{\alpha}_n^{\perp} \in null(\Psi)$. Substituting this decomposition of $\hat{\alpha}$ into the objective, we see that

$$\mathcal{L}_{\lambda}^{n}(\hat{\alpha}) = \mathcal{L}_{\lambda}^{n}(\hat{\alpha}_{n}) + \lambda \|\hat{\alpha}_{n}^{\perp}\|_{\mathcal{H}}^{2}$$

Therefore

$$\mathcal{L}_{\lambda}^{n}(\hat{\alpha}) \geq \mathcal{L}_{\lambda}^{n}(\hat{\alpha}_{n})$$

Since $\hat{\alpha}$ is the unique minimizer, $\hat{\alpha} = \hat{\alpha}_n$.

Derivation of Algorithm 3.2. By Proposition 3.3, $\hat{\alpha} = \Psi^* \rho$. Substituting this expression into $\mathcal{L}^n_{\lambda}(\alpha)$ gives $\mathcal{L}^n_{\lambda}(\rho)$.

$$\mathcal{L}_{\lambda}^{n}(\rho) = -2\rho^{\top}\Psi\frac{1}{n}\sum_{i=1}^{n}\phi^{(m)}(W_{i}) + \rho^{\top}\Psi\hat{T}\Psi^{*}\rho + \lambda\rho^{\top}\Psi\Psi^{*}\rho$$
$$= -\frac{2}{n}\rho^{\top}\sum_{i=1}^{n}\Psi\phi^{(m)}(W_{i}) + \frac{1}{n}\rho^{\top}\Psi\Phi^{*}\Phi\Psi^{*}\rho + \lambda\rho^{\top}\Psi\Psi^{*}\rho$$
$$= -\frac{2}{n}\rho^{\top}v + \frac{1}{n}\rho^{\top}\Omega\rho + \lambda\rho^{\top}K\rho$$

where

$$\Omega = \Psi \Phi^* \Phi \Psi^*, \quad v = \sum_{i=1}^n \Psi \phi^{(m)}(W_i)$$

Differentiating, the FOC gives

$$0 = -\frac{2}{n}v + \frac{2}{n}\Omega\rho + 2\lambda K\rho \iff \rho = (\Omega + n\lambda K)^{-1}v$$

To evaluate the estimator at a test location w involves computing

$$\hat{\alpha}(w) = \langle \hat{\alpha}, \phi(w) \rangle_{\mathcal{H}} = \rho^{\top} \Psi \phi(w) = v^{\top} (\Omega + n\lambda K)^{-1} \Psi \phi(w)$$

C Validity of confidence interval

C.1 Gateaux differentiation

For readability, I introduce the following notation for Gateaux differentiation.

Definition C.1 (Gateaux derivative). Let u(w), v(w) be functions and let $\tau, \zeta \in \mathbb{R}$ be scalars. The Gateaux derivative of $\psi(w, \theta, \gamma, \alpha)$ with respect to its argument γ in the direction u is

$$[\partial_{\gamma}\psi(w,\theta,\gamma,\alpha)](u) = \frac{\partial}{\partial \tau}\psi(w,\theta,\gamma+\tau u,\alpha)\bigg|_{\tau=0}$$

The cross derivative of $\psi(w, \theta, \gamma, \alpha)$ with respect to its argument (γ, α) in the directions (u, v) is

$$\left[\partial_{\gamma,\alpha}^2 \psi(w,\theta,\gamma,\alpha)\right](u,v) = \frac{\partial^2}{\partial \tau \partial \zeta} \psi(w,\theta,\gamma+\tau u,\alpha+\zeta v) \bigg|_{\tau=0,\zeta=0}$$

Proposition C.1 (Calculation of derivatives).

$$[\partial_{\gamma}\psi(w,\theta,\gamma,\alpha)](u) = m(w,u) - \alpha(w)u(w)$$
$$[\partial_{\alpha}\psi(w,\theta,\gamma,\alpha)](v) = v(w)[y - \gamma(w)]$$
$$[\partial_{\gamma,\alpha}^{2}\psi(w,\theta,\gamma,\alpha)](u,v) = -v(w)u(w)$$

Proof. For the first result, write

$$\psi(w, \theta, \gamma + \tau u, \alpha) = m(w, \gamma) + \tau m(w, u) + \alpha(w)[y - \gamma(w) - \tau u(w)] - \theta$$

For the second result, write

$$\psi(w, \theta, \gamma, \alpha + \zeta v) = m(w, \gamma) + \alpha(w)[y - \gamma(w)] + \zeta v(w)[y - \gamma(w)] - \theta$$

For the final result, write

$$\psi(w, \theta, \gamma + \tau u, \alpha + \zeta v)$$

$$= m(w, \gamma) + \tau m(w, u) + \alpha(w)[y - \gamma(w) - \tau u(w)] + \zeta v(w)[y - \gamma(w) - \tau u(w)] - \theta$$

Finally, take scalar derivatives with respect to (τ, ζ) .

By using the doubly robust moment function, we have the following helpful property.

Proposition C.2 (Mean zero derivatives). For any (u, v),

$$\mathbb{E}[\partial_{\gamma}\psi_0(W)](u) = 0, \quad \mathbb{E}[\partial_{\alpha}\psi_0(W)](v) = 0$$

Proof. For the first result, write

$$\mathbb{E}[\partial_{\gamma}\psi_0(W)](u) = \mathbb{E}[m(W, u) - \alpha_0(W)u(W)]$$

Then appeal to the definition of the Riesz representer. For the second result, write

$$\mathbb{E}[\partial_{\alpha}\psi_0(W)](v) = \mathbb{E}[v(W)[y - \gamma_0(W)]]$$

In the case of nonparametric regression, $\gamma_0(w) = \mathbb{E}[Y|W=w]$ and we appeal to law of iterated expectations.

C.2 Taylor expansion

Train $(\hat{\gamma}_{\ell}, \hat{\alpha}_{\ell})$ on observations in I_{ℓ}^c . Let $m = |I_{\ell}| = \frac{n}{L}$ be the number of observations in I_{ℓ} . Denote by $\mathbb{E}_{\ell}[\cdot]$ the average over observations in I_{ℓ} .

Definition C.2 (Foldwise target and oracle).

$$\hat{\theta}_{\ell} = \mathbb{E}_{\ell}[m(W, \hat{\gamma}_{\ell}) + \hat{\alpha}_{\ell}(W)\{Y - \hat{\gamma}_{\ell}(W)\}]$$

$$\bar{\theta}_{\ell} = \mathbb{E}_{\ell}[m(W, \gamma_{0}) + \alpha_{0}(W)\{Y - \gamma_{0}(W)\}]$$

Proposition C.3 (Taylor expansion). Let $u = \hat{\gamma}_{\ell} - \gamma_0$ and $v = \hat{\alpha}_{\ell} - \alpha_0$. Then $\sqrt{m}(\hat{\theta}_{\ell} - \bar{\theta}_{\ell}) = \sum_{i=1}^{3} \Delta_{j\ell}$ where

$$\Delta_{1\ell} = \sqrt{m} \mathbb{E}_{\ell}[m(W, u) - \alpha_0(W)u(W)]$$

$$\Delta_{2\ell} = \sqrt{m} \mathbb{E}_{\ell}[v(W)\{Y - \gamma_0(W)\}]$$

$$\Delta_{3\ell} = \sqrt{m} \mathbb{E}_{\ell}[-u(W)v(W)]$$

Proof. An exact Taylor expansion gives

$$\psi(w,\theta_0,\hat{\gamma}_\ell,\hat{\alpha}_\ell) - \psi_0(w) = [\partial_\gamma \psi_0(w)](u) + [\partial_\alpha \psi_0(w)](v) + [\partial_{\gamma,\alpha}^2 \psi_0(w)](u,v)$$

Averaging over observations in I_ℓ

$$\hat{\theta}_{\ell} - \bar{\theta}_{\ell} = \mathbb{E}_{\ell}[\psi(W, \theta_0, \hat{\gamma}_{\ell}, \hat{\alpha}_{\ell})] - \mathbb{E}_{\ell}[\psi_0(W)]$$

$$= \mathbb{E}_{\ell}[\partial_{\gamma}\psi_0(W)](u) + \mathbb{E}_{\ell}[\partial_{\alpha}\psi_0(W)](v) + \mathbb{E}_{\ell}[\partial_{\gamma,\alpha}^2\psi_0(W)](u, v)$$

Finally appeal to Proposition C.1.

C.3 Residuals

Definition C.3 (Mean square error). Write the conditional mean square error of $(\hat{\gamma}_{\ell}, \hat{\alpha}_{\ell})$ trained on I_{ℓ}^{c} as

$$\mathcal{R}(\hat{\gamma}_{\ell}) = \mathbb{E}[\{\hat{\gamma}_{\ell}(W) - \gamma_0(W)\}^2 | I_{\ell}^c]$$

$$\mathcal{R}(\hat{\alpha}_{\ell}) = \mathbb{E}[\{\hat{\alpha}_{\ell}(W) - \alpha_0(W)\}^2 | I_{\ell}^c]$$

Proposition C.4 (Residuals). Suppose

$$\mathbb{E}[Y-\gamma_0(W)]^2 \leq \bar{\sigma}^2, \quad \mathbb{E}[m(W,\gamma)]^2 \leq \bar{L}^{(m)}\mathbb{E}[\gamma(W)]^2, \quad \|\alpha_0\|_{\infty} \leq \bar{\alpha}$$
 Then w.p. $1-\frac{\epsilon}{L}$,

$$|\Delta_{1\ell}| \le t_1 = \sqrt{\frac{6L}{\epsilon}} \sqrt{\bar{L}^{(m)} + \bar{\alpha}^2} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}$$
$$|\Delta_{2\ell}| \le t_2 = \sqrt{\frac{3L}{\epsilon}} \bar{\sigma} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$
$$|\Delta_{3\ell}| \le t_3 = \frac{3\sqrt{L}}{\epsilon} \sqrt{n} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$

Proof. I proceed in steps

1. Markov inequality

$$\mathbb{P}(|\Delta_{1\ell}| > t_1) \le \frac{\mathbb{E}[\Delta_{1\ell}^2]}{t_1^2}$$

$$\mathbb{P}(|\Delta_{2\ell}| > t_2) \le \frac{\mathbb{E}[\Delta_{2\ell}^2]}{t_2^2}$$

$$\mathbb{P}(|\Delta_{3\ell}| > t_3) \le \frac{\mathbb{E}|\Delta_{3\ell}|}{t_3}$$

2. Law of iterated expectations

$$\begin{split} \mathbb{E}[\Delta_{1\ell}^2] &= \mathbb{E}[\mathbb{E}[\Delta_{1\ell}^2|I_\ell^c]] \\ \mathbb{E}[\Delta_{2\ell}^2] &= \mathbb{E}[\mathbb{E}[\Delta_{2\ell}^2|I_\ell^c]] \\ \mathbb{E}|\Delta_{3\ell}| &= \mathbb{E}[\mathbb{E}||\Delta_{3\ell}||I_\ell^c]] \end{split}$$

3. Bounding conditional moments

Conditional on I_{ℓ}^c , (u,v) are nonrandom. Moreover, observations within fold I_{ℓ} are independent. Hence

$$\mathbb{E}[\Delta_{1\ell}^{2}|I_{\ell}^{c}] = \mathbb{E}[\{m(W, u) - \alpha_{0}(W)u(W)\}^{2}|I_{\ell}^{c}]$$

$$\leq 2\mathbb{E}[m(W, u)^{2}|I_{\ell}^{c}] + 2\mathbb{E}[\{\alpha_{0}(W)u(W)\}^{2}|I_{\ell}^{c}]$$

$$\leq 2(\bar{L}^{(m)} + \bar{\alpha}^{2})\mathcal{R}(\hat{\gamma}_{\ell})$$

Similarly

$$\begin{split} \mathbb{E}[\Delta_{2\ell}^{2}|I_{\ell}^{c}] &= \mathbb{E}[\{v(W)[Y - \gamma_{0}(W)]\}^{2}|I_{\ell}^{c}] \\ &= \mathbb{E}[v(W)^{2}\mathbb{E}[\{Y - \gamma_{0}(W)\}^{2}|W, I_{\ell}^{c}]|I_{\ell}^{c}] \\ &\leq \bar{\sigma}^{2}\mathcal{R}(\hat{\alpha}_{\ell}) \end{split}$$

Finally

$$\mathbb{E}[|\Delta_{3\ell}||I_{\ell}^{c}] = \sqrt{m}\mathbb{E}[|-u(W)v(W)||I_{\ell}^{c}]$$

$$\leq \sqrt{m}\mathbb{E}[u(W)^{2}|I_{\ell}^{c}]\mathbb{E}[v(W)^{2}|I_{\ell}^{c}]$$

$$= \sqrt{m}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$

4. Collecting results

$$\mathbb{P}(|\Delta_{1\ell}| > t_1) \le \frac{2(\bar{L}^{(m)} + \bar{\alpha}^2)\mathcal{R}(\hat{\gamma}_{\ell})}{t_1^2} = \frac{\epsilon}{3L}$$

$$\mathbb{P}(|\Delta_{2\ell}| > t_2) \le \frac{\bar{\sigma}^2 \mathcal{R}(\hat{\alpha}_{\ell})}{t_2^2} = \frac{\epsilon}{3L}$$

$$\mathbb{P}(|\Delta_{3\ell}| > t_3) \le \frac{\sqrt{m}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}}{t_2} = \frac{\epsilon}{3L}$$

Therefore w.p. $1 - \frac{\epsilon}{L}$, the following inequalities hold

$$|\Delta_{1\ell}| \le t_1 = \sqrt{\frac{6L}{\epsilon}} \sqrt{\bar{L}^{(m)} + \bar{\alpha}^2} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}$$
$$|\Delta_{2\ell}| \le t_2 = \sqrt{\frac{3L}{\epsilon}} \bar{\sigma} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$
$$|\Delta_{3\ell}| \le t_3 = \frac{3L}{\epsilon} \sqrt{m} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$

Finally recall $m = \frac{n}{L}$

C.4 Main argument

Definition C.4 (Overall target and oracle).

$$\hat{\theta} = \frac{1}{L} \sum_{\ell=1}^{L} \hat{\theta}_{\ell}, \quad \bar{\theta} = \frac{1}{L} \sum_{\ell=1}^{L} \bar{\theta}_{\ell}$$

Proposition C.5 (Oracle approximation). *Suppose the conditions of Proposition C.4 hold. Then w.p.* $1 - \epsilon$

$$\frac{\sqrt{n}}{\sigma}|\hat{\theta} - \bar{\theta}| \le \Delta = \frac{3L}{\epsilon \cdot \sigma} \left\{ (\sqrt{\bar{L}^{(m)}} + \bar{\alpha})\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} + \bar{\sigma}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} + \sqrt{n}\sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}\sqrt{\mathcal{R}(\hat{\alpha}_{\ell})} \right\}$$

Proof. I proceed in steps

1. Decomposition

Write

$$\sqrt{n}(\hat{\theta} - \bar{\theta}) = \frac{\sqrt{n}}{\sqrt{m}} \frac{1}{L} \sum_{\ell=1}^{L} \sqrt{m}(\hat{\theta}_{\ell} - \bar{\theta}_{\ell})$$
$$= \sqrt{L} \frac{1}{L} \sum_{\ell=1}^{L} \sum_{j=1}^{3} \Delta_{j\ell}$$

2. Union bound

Define the events

$$\mathcal{E}_{\ell} = \{ \forall j \in [3], \ |\Delta_{j\ell}| \le t_j \}, \quad \mathcal{E} = \cap_{\ell=1}^L \mathcal{E}_{\ell}, \quad \mathcal{E}^c = \cup_{\ell=1}^L \mathcal{E}^c_{\ell}$$

Hence by the union bound and Proposition C.4,

$$\mathbb{P}(\mathcal{E}^c) \le \sum_{\ell=1}^{L} \mathbb{P}(\mathcal{E}^c_{\ell}) \le L \frac{\epsilon}{L} = \epsilon$$

3. Collecting results

Therefore w.p. $1 - \epsilon$,

$$\sqrt{n}|\hat{\theta} - \bar{\theta}| \le \sqrt{L} \frac{1}{L} \sum_{\ell=1}^{L} \sum_{j=1}^{3} |\Delta_{jk}|$$

$$\le \sqrt{L} \frac{1}{L} \sum_{\ell=1}^{L} \sum_{j=1}^{3} t_j$$

$$= \sqrt{L} \sum_{j=1}^{3} t_j$$

Finally I simplify $\{t_j\}$. For a,b>0, $\sqrt{a+b}\leq \sqrt{a}+\sqrt{b}$. Moreover, $3\geq \sqrt{6},\sqrt{3}$. Finally, for $\epsilon\leq 1,\,\frac{1}{\sqrt{\epsilon}}\leq \frac{1}{\epsilon}$. In summary

$$t_1 \le \frac{3\sqrt{L}}{\epsilon} (\sqrt{\bar{L}^{(m)}} + \bar{\alpha}) \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})}$$
$$t_2 \le \frac{3\sqrt{L}}{\epsilon} \bar{\sigma} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$
$$t_3 \le \frac{3\sqrt{L}}{\epsilon} \sqrt{n} \sqrt{\mathcal{R}(\hat{\gamma}_{\ell})} \sqrt{\mathcal{R}(\hat{\alpha}_{\ell})}$$

Proof of Theorem 4.1. By Berry Esssen theorem, as in [22, Proof of Theorem 4.1, Step 3]

$$\mathbb{P}\left(\frac{\sqrt{n}}{\sigma}(\hat{\theta}-\theta_0) \leq z\right) - \Phi(z) \leq c^{BE} \left(\frac{\eta}{\sigma}\right)^3 n^{-\frac{1}{2}} + \frac{\Delta}{\sqrt{2\pi}} + \epsilon^{BE} \left(\frac{\eta}{\sigma}\right)^3 n^{-\frac{1}{2}} + \frac{\Delta}{\sqrt{2\pi}} + \frac{\Delta}$$

and

$$\mathbb{P}\left(\frac{\sqrt{n}}{\sigma}(\hat{\theta}-\theta_0) \leq z\right) - \Phi(z) \geq c^{BE} \left(\frac{\eta}{\sigma}\right)^3 n^{-\frac{1}{2}} - \frac{\Delta}{\sqrt{2\pi}} - \epsilon$$

where $\Phi(z)$ is the standard Gaussian c.d.f. and Δ is defined in Proposition C.5.

C.5 Variance estimation

Proof of Proposition 4.1. Write

$$\begin{split} \sigma^2 &= \mathbb{E}[m(W, \gamma_0) + \alpha_0^{\min}(W) \{Y - \gamma_0(W)\} - \theta_0]^2 \\ &\leq 3 \left(\mathbb{E}[m(W, \gamma_0)]^2 + \mathbb{E}[\alpha_0^{\min}(W) \{Y - \gamma_0(W)\}]^2 + \theta_0^2 \right) \end{split}$$

Then note

$$\mathbb{E}[m(W, \gamma_0)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma_0(W)]^2$$

and

$$\mathbb{E}[\alpha_0^{\min}(W)\{Y - \gamma_0(W)\}]^2 \le \bar{\alpha}^2 \mathbb{E}[Y - \gamma_0(W)]^2 \le \bar{\alpha}^2 \bar{\sigma}^2$$

Definition C.5 (Shorter notation). For $i \in I_{\ell}$, define

$$\psi_i = \psi(W_i, \theta_0, \gamma_0, \alpha_0)$$
$$\hat{\psi}_i = \psi(W_i, \hat{\theta}, \hat{\gamma}_\ell, \hat{\alpha}_\ell)$$

Proposition C.6 (Foldwise second moment).

$$\mathbb{E}_{\ell}[\hat{\psi}_i - \psi_i]^2 \le 4 \left\{ [\hat{\theta} - \theta_0]^2 + \sum_{j=4}^6 \Delta_{j\ell} \right\}$$

where

$$\Delta_{4\ell} = \mathbb{E}_{\ell}[m(W, u)]^2$$

$$\Delta_{5\ell} = \mathbb{E}_{\ell}[\hat{\alpha}_{\ell}(W)u(W)]^2$$

$$\Delta_{6\ell} = \mathbb{E}_{\ell}[v(W)\{Y - \gamma_0(W)\}]^2$$

Proof. Write

$$\psi_{i} - \hat{\psi}_{i} = m(W_{i}, \hat{\gamma}_{\ell}) + \hat{\alpha}_{\ell}(W_{i})[Y_{i} - \hat{\gamma}_{\ell}(W_{i})] - \hat{\theta} - \{m(W_{i}, \gamma_{0}) + \alpha_{0}(W_{i})[Y_{i} - \gamma_{0}(W_{i})] - \theta_{0}\}$$

$$\pm \hat{\alpha}_{\ell}[Y - \gamma_{0}(W_{i})]$$

$$= [\theta_{0} - \hat{\theta}] + m(W_{i}, u) - \hat{\alpha}_{\ell}(W_{i})u(W_{i}) + v(W_{i})[Y - \gamma_{0}(W_{i})]$$

Hence

$$[\psi_i - \hat{\psi}_i]^2 \le 4 \left\{ [\theta_0 - \hat{\theta}]^2 + m(W_i, u)^2 + [\hat{\alpha}_\ell(W_i)u(W_i)]^2 + [v(W_i)\{Y - \gamma_0(W_i)\}]^2 \right\}$$

Finally take $\mathbb{E}_{\ell}[\cdot]$ of both sides.

Proposition C.7 (Residuals). Suppose

$$\mathbb{E}[Y - \gamma_0(W)]^2 \le \bar{\sigma}^2, \quad \mathbb{E}[m(W, \gamma)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma(W)]^2, \quad \|\hat{\alpha}_\ell\|_{\infty} \le \bar{\alpha}'$$

Then w.p. $1 - \frac{\epsilon'}{2L}$,

$$\Delta_{4\ell} \le t_4 = \frac{6L}{\epsilon'} \bar{L}^{(m)} \mathcal{R}(\hat{\gamma}_{\ell})$$
$$\Delta_{5\ell} \le t_5 = \frac{6L}{\epsilon'} (\bar{\alpha}')^2 \mathcal{R}(\hat{\gamma}_{\ell})$$
$$\Delta_{6\ell} \le t_6 = \frac{6L}{\epsilon'} \bar{\sigma}^2 \mathcal{R}(\hat{\alpha}_{\ell})$$

Proof. I proceed in steps analogous to Proposition C.4.

1. Markov inequality

$$\mathbb{P}(|\Delta_{4\ell}| > t_4) \le \frac{\mathbb{E}[\Delta_{4\ell}]}{t_4}$$

$$\mathbb{P}(|\Delta_{5\ell}| > t_5) \le \frac{\mathbb{E}[\Delta_{5\ell}]}{t_5}$$

$$\mathbb{P}(|\Delta_{6\ell}| > t_6) \le \frac{\mathbb{E}[\Delta_{6\ell}]}{t_6}$$

2. Law of iterated expectations

$$\mathbb{E}[\Delta_{4\ell}] = \mathbb{E}[\mathbb{E}[\Delta_{4\ell}|I_{\ell}^c]]$$

$$\mathbb{E}[\Delta_{5\ell}] = \mathbb{E}[\mathbb{E}[\Delta_{5\ell}|I_{\ell}^c]]$$

$$\mathbb{E}[\Delta_{6\ell}] = \mathbb{E}[\mathbb{E}[\Delta_{6\ell}|I_{\ell}^c]]$$

3. Bounding conditional moments

Conditional on $I_\ell^c,\,(u,v)$ are nonrandom. Moreover, observations within fold I_ℓ are independent. Hence

$$\mathbb{E}[\Delta_{4\ell}|I_{\ell}^c] = \mathbb{E}[\{m(W,u)\}^2|I_{\ell}^c] \le \bar{L}^{(m)}\mathcal{R}(\hat{\gamma}_{\ell})$$

Similarly

$$\mathbb{E}[\Delta_{5\ell}|I_{\ell}^c] = \mathbb{E}[\{\hat{\alpha}_{\ell}(W)u(W)\}^2|I_{\ell}^c] \le (\bar{\alpha}')^2 \mathcal{R}(\hat{\gamma}_{\ell})$$

Finally

$$\mathbb{E}[\Delta_{6\ell}|I_{\ell}^{c}] = \mathbb{E}[\{v(W)[Y - \gamma_{0}(W)]\}^{2}|I_{\ell}^{c}]$$

$$= \mathbb{E}[v(W)^{2}\mathbb{E}[\{Y - \gamma_{0}(W)\}^{2}|W, I_{\ell}^{c}]|I_{\ell}^{c}]$$

$$\leq \bar{\sigma}^{2}\mathcal{R}(\hat{\alpha}_{\ell})$$

4. Collecting results

$$\mathbb{P}(|\Delta_{4\ell}| > t_4) \le \frac{\bar{L}^{(m)}\mathcal{R}(\hat{\gamma}_{\ell})}{t_4} = \frac{\epsilon'}{6L}$$

$$\mathbb{P}(|\Delta_{5\ell}| > t_5) \le \frac{(\bar{\alpha}')^2\mathcal{R}(\hat{\gamma}_{\ell})}{t_5} = \frac{\epsilon'}{6L}$$

$$\mathbb{P}(|\Delta_{6\ell}| > t_6) \le \frac{\bar{\sigma}^2\mathcal{R}(\hat{\alpha}_{\ell})}{t_6} = \frac{\epsilon'}{6L}$$

Therefore w.p. $1 - \frac{\epsilon'}{2L}$, the following inequalities hold

$$|\Delta_{4\ell}| \le t_4 = \frac{6L}{\epsilon'} \bar{L}^{(m)} \mathcal{R}(\hat{\gamma}_{\ell})$$
$$|\Delta_{5\ell}| \le t_5 = \frac{6L}{\epsilon'} (\bar{\alpha}')^2 \mathcal{R}(\hat{\gamma}_{\ell})$$
$$|\Delta_{6\ell}| \le t_6 = \frac{6L}{\epsilon'} \bar{\sigma}^2 \mathcal{R}(\hat{\alpha}_{\ell})$$

Proposition C.8 (Oracle approximation). Suppose the conditions of Proposition C.7 hold. Then w.p. $1 - \frac{\epsilon'}{2}$

$$\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2} \leq \Delta' = 4\{\hat{\theta} - \theta_{0}\}^{2} + \frac{24L}{\epsilon'}\{[\bar{L}^{(m)} + (\bar{\alpha}')^{2}]\mathcal{R}(\hat{\gamma}_{\ell}) + \bar{\sigma}^{2}\mathcal{R}(\hat{\alpha}_{\ell})\}$$

Proof. I proceed in steps analogous to Proposition C.5

1. Decomposition

By Proposition C.6

$$\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2} = \frac{1}{L} \sum_{\ell=1}^{L} \mathbb{E}_{\ell}[\hat{\psi}_{i} - \psi_{i}]^{2}$$

$$\leq 4[\hat{\theta} - \theta_{0}]^{2} + \frac{4}{L} \sum_{\ell=1}^{L} \sum_{j=4}^{6} \Delta_{j\ell}$$

2. Union bound

Define the events

$$\mathcal{E}'_{\ell} = \{ \forall j \in \{4, 5, 6\}, \ |\Delta_{j\ell}| \le t_j \}, \quad \mathcal{E}' = \cap_{\ell=1}^L \mathcal{E}'_{\ell}, \quad (\mathcal{E}')^c = \cup_{\ell=1}^L (\mathcal{E}'_{\ell})^c$$

Hence by the union bound and Proposition C.7,

$$\mathbb{P}((\mathcal{E}')^c) \le \sum_{\ell=1}^L \mathbb{P}((\mathcal{E}'_\ell)^c) \le L \frac{\epsilon'}{2L} = \frac{\epsilon'}{2}$$

3. Collecting results

Therefore w.p. $1 - \frac{\epsilon'}{2}$,

$$\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2} \leq 4[\hat{\theta} - \theta_{0}]^{2} + \frac{4}{L} \sum_{\ell=1}^{L} \sum_{j=4}^{6} |\Delta_{j\ell}|$$

$$\leq 4[\hat{\theta} - \theta_{0}]^{2} + \frac{4}{L} \sum_{\ell=1}^{L} \sum_{j=4}^{6} t_{j}$$

$$= 4[\hat{\theta} - \theta_{0}]^{2} + 4 \sum_{j=4}^{6} t_{j}$$

Proposition C.9 (Markov inequality). Suppose $\mathbb{E}[\psi_0(W)]^4 < \infty$. Then w.p. $1 - \frac{\epsilon'}{2}$

$$|\mathbb{E}_n[\psi_i]^2 - \sigma^2| \le \Delta'' = \sqrt{\frac{2}{\epsilon'}} \frac{\sqrt{\mathbb{E}[\psi_0(W)]^4}}{\sqrt{n}}$$

Proof. Let

$$Z_i = \psi_i^2, \quad \bar{Z} = \mathbb{E}_n[Z_i], \quad \mathbb{E}[Z] = \sigma^2$$

By Markov inequality

$$\mathbb{P}(|\bar{Z} - \mathbb{E}[Z]| > t) \le \frac{\mathbb{V}[\bar{Z}]}{t^2} = \frac{\epsilon'}{2}$$

Note that

$$\mathbb{V}[\bar{Z}] = \frac{\mathbb{V}(Z)}{n} = \frac{\mathbb{E}[\psi_0(W)^2]^2}{n} = \frac{\mathbb{E}[\psi_0(W)]^4}{n}$$

Solving,

$$\frac{\mathbb{E}[\psi_0(W)]^4}{nt^2} = \frac{\epsilon'}{2} \iff t = \sqrt{\frac{2}{\epsilon'}} \frac{\sqrt{\mathbb{E}[\psi_0(W)]^4}}{\sqrt{n}}$$

Proof of Theorem 4.2. I proceed in steps.

1. Decomposition of variance estimator

Write

$$\hat{\sigma}^2 = \mathbb{E}_n[\hat{\psi}_i]^2 = \mathbb{E}_n[\{\hat{\psi}_i - \psi_i\} + \psi_i]^2 = \mathbb{E}_n[\hat{\psi}_i - \psi_i]^2 + 2\mathbb{E}_n[\{\hat{\psi}_i - \psi_i\}\psi_i] + \mathbb{E}_n[\psi_i]^2$$

Hence

$$\hat{\sigma}^2 - \mathbb{E}_n[\psi_i]^2 = \mathbb{E}_n[\hat{\psi}_i - \psi_i]^2 + 2\mathbb{E}_n[\{\hat{\psi}_i - \psi_i\}\psi_i]$$

2. Decomposition of difference

Next write

$$\hat{\sigma}^2 - \sigma^2 = {\{\hat{\sigma}^2 - \mathbb{E}_n[\psi_i]^2\}} + {\{\mathbb{E}_n[\psi_i]^2 - \sigma^2\}}$$

Focusing on the former term

$$\hat{\sigma}^2 - \mathbb{E}_n[\psi_i]^2 = \mathbb{E}_n[\hat{\psi}_i - \psi_i]^2 + 2\mathbb{E}_n[\{\hat{\psi}_i - \psi_i\}\psi_i]$$

Moreover

$$\mathbb{E}_{n}[\{\hat{\psi}_{i} - \psi_{i}\}\psi_{i}] \leq \sqrt{\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2}} \sqrt{\mathbb{E}_{n}[\psi_{i}]^{2}}
\leq \sqrt{\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2}} \sqrt{|\mathbb{E}_{n}[\psi_{i}]^{2} - \sigma^{2}| + \sigma^{2}}
\leq \sqrt{\mathbb{E}_{n}[\hat{\psi}_{i} - \psi_{i}]^{2}} \left(\sqrt{|\mathbb{E}_{n}[\psi_{i}]^{2} - \sigma^{2}|} + \sigma\right)$$

3. High probability events

From the previous step, we see that to control $|\hat{\sigma}^2 - \sigma^2|$, it is sufficient to control two expressions: $\mathbb{E}_n[\hat{\psi}_i - \psi_i]^2$ and $|\mathbb{E}_n[\psi_i]^2 - \sigma^2|$. These are controlled in Propositions C.8 and C.9, respectively. Therefore w.p. $1 - \epsilon'$,

$$|\hat{\sigma}^2 - \sigma^2| \le \left\{ \Delta' + 2\sqrt{\Delta'}(\sqrt{\Delta''} + \sigma) \right\} + \Delta''$$

C.6 Corollary

Proof of Corollary 4.1. Immediately from Δ in Theorem 4.1, $\hat{\theta} \stackrel{p}{\rightarrow} \theta_0$ and

$$\lim_{n \to \infty} \mathbb{P}\left(\theta_0 \in \left[\hat{\theta} \pm \frac{\sigma}{\sqrt{n}}\right]\right) = 1 - a$$

For the desired result, it is sufficient that $\hat{\sigma}^2 \stackrel{p}{\to} \sigma^2$, which follows from Δ' and Δ'' in Theorem 4.2.

D Validity of bias correction

D.1 Preliminaries

Proposition D.1 (First order conditions). The first order conditions give

$$T\alpha_0^{\min} = \mu^{(m)}, \quad \alpha_\lambda = (T+\lambda)^{-1}\mu^{(m)}, \quad \hat{\alpha} = (\hat{T}+\lambda)^{-1}\hat{\mu}^{(m)}$$

Proof. Take derivatives of \mathcal{L} , \mathcal{L}_{λ} , and $\mathcal{L}_{\lambda}^{n}$

Proposition D.2 (Relation of norms).

$$\mathcal{R}(\alpha) = \mathbb{E}[\alpha(W) - \alpha_0^{\min}(W)]^2 = \|T^{\frac{1}{2}}(\alpha - \alpha_0^{\min})\|_{\mathcal{H}}^2$$

Proof. [16, Proposition 1.iii], using the fact that mean square error equals excess risk.

Proposition D.3 (Concentration; Proposition 2 of [16]). Let $(\Omega, \mathcal{F}, \rho)$ be a probability space. Let ξ be a random variable on Ω taking value in a real separable Hilbert space K. Assume there exists (a,b) > 0 s.t.

$$\|\xi(\omega)\|_{\mathcal{K}} \le \frac{a}{2}, \quad \mathbb{E}\|\xi\|_{\mathcal{K}}^2 \le b^2$$

Then for all $n \in \mathbb{N}$ and $\eta \in (0, 1)$,

$$\mathbb{P}_{\{\omega_i\} \sim \rho^n} \left(\left\| \frac{1}{n} \sum_{i=1}^n \xi(\omega_i) - \mathbb{E}[\xi] \right\|_{\mathcal{K}} \le 2 \ln(2/\eta) \left[\frac{a}{n} + \frac{b}{\sqrt{n}} \right] \right) \ge \frac{1}{\eta}$$

D.2 Learning theory quantities

Proof of Proposition 5.1. Write

$$\mathbb{E}[Y - \gamma_0(W)]^2 \le 2\mathbb{E}[Y]^2 + 2\mathbb{E}[\gamma_0(W)]^2$$

Since $\gamma_0 \in \mathcal{H}$,

$$|\gamma_0(w)| = |\langle \gamma_0, \phi(w) \rangle_{\mathcal{H}}| \le \sqrt{\kappa} ||\gamma_0||_{\mathcal{H}}$$

 $\gamma \in \mathcal{H}$ implies $\alpha_0^{\min} \in \mathcal{H}$, so

$$|\alpha_0^{\min}(w)| = |\langle \alpha_0^{\min}, \phi(w) \rangle_{\mathcal{H}}| \le \sqrt{\kappa} ||\alpha_0^{\min}||_{\mathcal{H}}$$

and likewise

$$|\hat{\alpha}(w)| = |\langle \hat{\alpha}, \phi(w) \rangle_{\mathcal{H}}| \le \sqrt{\kappa} ||\hat{\alpha}||_{\mathcal{H}}$$

Finally, I bound $\|\hat{\alpha}\|_{\mathcal{H}}$. By identical arguments to Proposition 3.1,

$$\hat{\alpha} = \operatorname*{argmin}_{\alpha \in \mathcal{H}} \mathcal{R}^{n}_{\lambda}(\alpha), \quad \mathcal{R}^{n}_{\lambda}(\alpha) = \mathbb{E}_{n}[\alpha^{\min}_{0}(W) - \alpha(W)]^{2} + \lambda \|\alpha\|_{\mathcal{H}}^{2}$$

Hence

$$\lambda \|\hat{\alpha}\|_{\mathcal{H}}^2 \leq \mathcal{R}_{\lambda}^n(\hat{\alpha}) \leq \mathcal{R}_{\lambda}^n(0) = \mathbb{E}_n[\alpha_0^{\min}(W)]^2 \leq \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2$$

Therefore

$$\|\hat{\alpha}\|_{\mathcal{H}} \leq \frac{\sqrt{\kappa} \|\alpha_0^{\min}\|_{\mathcal{H}}}{\sqrt{\lambda}}, \quad |\hat{\alpha}(w)| \leq \frac{\kappa \|\alpha_0^{\min}\|_{\mathcal{H}}}{\sqrt{\lambda}}$$

Proof of Proposition 5.2. For the first result, write $f = \sum_{k=1}^{\infty} a_k \varphi_k$ and appeal to [16, Remark 2]. The second result follows from Proposition 2.3.

Proof of Proposition 5.3. The first result is immediate. The second result follows from [16, Remark 2]. \Box

Definition D.1 (Learning theory quantities). *Define*

$$\mathcal{A}(\lambda) = \mathcal{R}(\alpha_{\lambda})$$

$$\mathcal{B}(\lambda) = \|\alpha_{\lambda} - \alpha_{0}^{\min}\|_{\mathcal{H}}^{2}$$

$$\mathcal{N}(\lambda) = Tr((T+\lambda)^{-1}T)$$

$$\mathcal{N}^{(m)}(\lambda) = Tr((T+\lambda)^{-1}T^{(m)})$$

where $T^{(m)} := \mathbb{E}[\phi^{(m)} \otimes \phi^{(m)}]$ is the counterfactual covariance operator. $\mathcal{A}(\lambda)$, $\mathcal{B}(\lambda)$, and $\mathcal{N}(\lambda)$ are standard in the kernel methods literature; $\mathcal{N}^{(m)}(\lambda)$ is new.

Proposition D.4 (Bounds on standard learning theory quantities). *Under Assumption 5.3*

$$\mathcal{A}(\lambda) \leq \lambda^{c} R$$
, $\mathcal{B}(\lambda) \leq \lambda^{c-1} R$, $R = \|T^{\frac{1-c}{2}} \gamma_0\|_{\mathcal{H}}^2$

Under Assumption 5.4

$$\mathcal{N}(\lambda) = Tr((T+\lambda)^{-1}T) \le \begin{cases} \beta^{\frac{1}{b}} \frac{\pi/b}{\sin(\pi/b)} \lambda^{-\frac{1}{b}} & \text{if } b < \infty \\ J & \text{if } b = \infty \end{cases}$$

Proof. See [16, Proposition 3] for $\mathcal{A}(\lambda)$ and $\mathcal{B}(\lambda)$. See [88, Section 1] for $\mathcal{N}(\lambda)$.

Proposition D.5 (Bound on new learning theory quantity). Suppose the kernels k and $k^{(m)}$ are bounded. Under Assumption 2.1,

$$\mathcal{N}^{(m)}(\lambda) \leq \bar{L}^{(m)} \cdot \mathcal{N}(\lambda)$$

Proof. Mean square continuity implies that there exists $\bar{L}^{(m)} < \infty$ s.t. $\forall \gamma \in \mathcal{H}$

$$\mathbb{E}[m(W,\gamma)]^2 \le \bar{L}^{(m)} \mathbb{E}[\gamma(W)]^2$$

Focusing on the LHS, boundedness of the kernel implies Bochner integrability hence

$$\mathbb{E}[m(W,\gamma)]^2 = \mathbb{E}\left[\langle \gamma, \phi^{(m)}(W)\rangle_{\mathcal{H}}^2\right] = \mathbb{E}\left[\langle \gamma, [\phi^{(m)}(W)\otimes\phi^{(m)}(W)]\gamma\rangle_{\mathcal{H}}\right] = \langle \gamma, T^{(m)}\gamma\rangle_{\mathcal{H}}$$

Likewise for the RHS

$$\mathbb{E}[\gamma(W)]^2 = \mathbb{E}\left[\langle \gamma, \phi(W) \rangle_{\mathcal{H}}^2\right] = \mathbb{E}\left[\langle \gamma, [\phi(W) \otimes \phi(W)] \gamma \rangle_{\mathcal{H}}\right] = \langle \gamma, T\gamma \rangle_{\mathcal{H}}$$

In summary, $\forall \gamma \in \mathcal{H}$

$$\langle \gamma, T^{(m)} \gamma \rangle_{\mathcal{H}} \leq \bar{L}^{(m)} \langle \gamma, T \gamma \rangle_{\mathcal{H}}$$

i.e. $\bar{L}^{(m)}T - T^{(m)} \succeq 0$ in the sense of Loewner ordering. Therefore

$$\bar{L}^{(m)}\mathcal{N}(\lambda) - \mathcal{N}^{(m)}(\lambda) = \bar{L}^{(m)}Tr((T+\lambda)^{-1}T) - Tr((T+\lambda)^{-1}T^{(m)})$$
$$= Tr((T+\lambda)^{-1}\{\bar{L}^{(m)}T - T^{(m)}\})$$
$$\geq 0$$

D.3 High probability events

In preparation for the main argument, I prove high probability events by concentration, appealing to Proposition D.3.

Definition D.2 (High probability events). *Define*

$$\mathcal{E}_{1} = \left\{ \| (T+\lambda)^{-1} (\hat{T} - T) \|_{\mathcal{L}_{2}(\mathcal{H}, \mathcal{H})} \leq 2 \ln(6/\eta) \left[\frac{2\kappa}{\lambda n} + \sqrt{\frac{\kappa \mathcal{N}(\lambda)}{\lambda n}} \right] \right\}$$

$$\mathcal{E}_{2} = \left\{ \| (T-\hat{T}) (\alpha_{\lambda} - \alpha_{0}^{\min}) \|_{\mathcal{H}} \leq 2 \ln(6/\eta) \left[\frac{2\kappa \sqrt{\mathcal{B}(\lambda)}}{n} + \sqrt{\frac{\kappa \mathcal{A}(\lambda)}{n}} \right] \right\}$$

$$\mathcal{E}_{3} = \left\{ \| (T+\lambda)^{-\frac{1}{2}} (\hat{\mu}^{(m)} - \hat{T}\alpha_{0}^{\min}) \|_{\mathcal{H}} \leq 2 \ln(6/\eta) \left[\frac{1}{n} \sqrt{\Gamma^{2} \frac{\kappa'}{\lambda}} + \sqrt{\frac{\Sigma^{2} \mathcal{N}(\lambda)}{n}} \right] \right\}$$

where

$$\Gamma = 2(1 + \sqrt{\kappa} \|\alpha_0^{\min}\|_{\mathcal{H}}), \quad \Sigma^2 = 2(\bar{L}^{(m)} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2), \quad \kappa' = \max\{\kappa, \kappa^{(m)}\}$$

I design \mathcal{E}_3 to resemble [16, eq. 48].

Proposition D.6 (High probability events). Suppose Assumptions 2.1 as well Assumptions 5.1 and 5.2 hold. Then

$$\mathbb{P}(\mathcal{E}_j^c) \le \frac{\eta}{3}, \quad j \in \{1, 2, 3\}$$

Proof. The argument for \mathcal{E}_1 immediately precedes [16, eq. 41]. The argument for \mathcal{E}_2 is identical to [16, eq. 43]. I verify \mathcal{E}_3 appealing to Proposition D.3. Define

$$\xi_i = (T+\lambda)^{-\frac{1}{2}} \{ \phi^{(m)}(w_i) - [\phi(w_i) \otimes \phi(w_i)] \alpha_0^{\min} \}$$

Note that

$$\mathbb{E}[\xi_i] = (T+\lambda)^{-\frac{1}{2}} \{ \mu^{(m)} - T\alpha_0^{\min} \} = 0$$

Towards concentration, I analyze $\|\xi\|_{\mathcal{H}}$ and $\mathbb{E}\|\xi\|_{\mathcal{H}}^2$

1. Bound on $\|\xi\|_{\mathcal{H}}$

Write

$$\|\xi_i\|_{\mathcal{H}} \leq \|(T+\lambda)^{-\frac{1}{2}}\|_{op}\|\phi^{(m)}(w_i) - [\phi(w_i)\otimes\phi(w_i)]\alpha_0^{\min}\|_{\mathcal{H}}$$

$$\leq \frac{1}{\sqrt{\lambda}}(\sqrt{\kappa^{(m)}} + \kappa\|\alpha_0^{\min}\|_{\mathcal{H}})$$

$$= \frac{a}{2}$$

2. Bound on $\mathbb{E}\|\xi\|_{\mathcal{H}}^2$

Write

$$\xi_i = A - B$$
, $A = (T + \lambda)^{-\frac{1}{2}} \phi^{(m)}(w_i)$, $B = (T + \lambda)^{-\frac{1}{2}} \alpha_0^{\min}(w_i) \phi(w_i)$

Hence

$$\|\xi_i\|_{\mathcal{H}}^2 = \|A - B\|_{\mathcal{H}}^2 \le 2\|A\|_{\mathcal{H}}^2 + 2\|B\|_{\mathcal{H}}^2$$

Focusing on the former term

$$||A||_{\mathcal{H}}^{2} = \langle (T+\lambda)^{-1}\phi^{(m)}(w_{i}), \phi^{(m)}(w_{i})\rangle_{\mathcal{H}}$$

= $Tr((T+\lambda)^{-1}[\phi^{(m)}(w_{i})\otimes\phi^{(m)}(w_{i})])$

Therefore by Proposition D.5

$$\mathbb{E}||A||_{\mathcal{H}}^2 = \mathcal{N}^{(m)}(\lambda) \le \bar{L}^{(m)}\mathcal{N}(\lambda)$$

Focusing on the latter term

$$||B||_{\mathcal{H}}^2 = \langle (T+\lambda)^{-1} \alpha_0^{\min}(w_i) \phi(w_i), \alpha_0^{\min}(w_i) \phi(w_i) \rangle_{\mathcal{H}}$$

$$< \kappa ||\alpha_0^{\min}||_{\mathcal{H}}^2 Tr((T+\lambda)^{-1} [\phi(w_i) \otimes \phi(w_i)])$$

Therefore

$$\mathbb{E}\|B\|_{\mathcal{H}}^2 \le \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2 \mathcal{N}(\lambda)$$

In summary

$$\mathbb{E}\|\xi\|_{\mathcal{H}}^2 \leq 2(\bar{L}^{(m)} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2) \cdot \mathcal{N}(\lambda) = b^2$$

3. Concentration

I have shown

$$a = \frac{2}{\sqrt{\lambda}} (\sqrt{\kappa^{(m)}} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}), \quad b = \sqrt{2(\bar{L}^{(m)} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2)} \sqrt{\mathcal{N}(\lambda)}$$

Therefore w.p. $1 - \eta/3$

$$\begin{split} & \left\| (T+\lambda)^{-\frac{1}{2}} (\hat{\mu}^{(m)} - \hat{T}\alpha_0^{\min}) \right\|_{\mathcal{H}} \\ & \leq 2 \ln(6/\eta) \left(\frac{a}{n} + \frac{b}{\sqrt{n}} \right) \\ & = 2 \ln(6/\eta) \left(\frac{2(\sqrt{\kappa^{(m)}} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}})}{n\sqrt{\lambda}} + \frac{\sqrt{2(\bar{L}^{(m)} + \kappa \|\alpha_0^{\min}\|_{\mathcal{H}}^2)} \sqrt{\mathcal{N}(\lambda)}}{\sqrt{n}} \right) \\ & \leq 2 \ln(6/\eta) \left(\frac{1}{n} \sqrt{\Gamma^2 \frac{\kappa'}{\lambda}} + \sqrt{\frac{\Sigma^2 \mathcal{N}(\lambda)}{n}} \right) \end{split}$$

D.4 Main argument

Proposition D.7 (Abstract rate). Suppose Assumptions 2.1 as well Assumptions 5.1 and 5.2 hold. If

$$C_{\eta} = 96 \ln^2(6/\eta), \quad n \ge \frac{2C_{\eta}\kappa'\mathcal{N}(\lambda)}{\lambda}, \quad \lambda \le ||T||_{op}$$

then w.p. $1 - \eta$

$$\mathcal{R}(\hat{\alpha}) \leq C_{\eta} \left(\mathcal{A}(\lambda) + \frac{\kappa^2 \mathcal{B}(\lambda)}{n^2 \lambda} + \frac{\kappa \mathcal{A}(\lambda)}{n \lambda} + \frac{\kappa' \Gamma^2}{n \lambda} + \frac{\Sigma^2 \mathcal{N}(\lambda)}{n} \right)$$

Proof. I proceed in steps, following the proof structure of [16, Theorem 4].

1. Decomposition

By [16, eq. 36],

$$\mathcal{R}(\hat{\alpha}) \le 3 \left\{ \mathcal{A}(\lambda) + \mathcal{S}_1(\lambda, \{w_i\}) + \mathcal{S}_2(\lambda, \{w_i\}) \right\}$$

where

$$S_{1}(\lambda, \{w_{i}\}) = \left\| T^{\frac{1}{2}}(\hat{T} + \lambda)^{-1}(\hat{\mu}^{(m)} - \hat{T}\alpha_{0}^{\min}) \right\|_{\mathcal{H}}^{2}$$

$$S_{2}(\lambda, \{w_{i}\}) = \left\| T^{\frac{1}{2}}(\hat{T} + \lambda)^{-1}(T - \hat{T})(\alpha_{\lambda} - \alpha_{0}^{\min}) \right\|_{\mathcal{H}}^{2}$$

2. Bound on $S_2(\lambda, \{w_i\})$

Write

$$S_2(\lambda, \{w_i\}) \le \|T^{\frac{1}{2}}(\hat{T} + \lambda)^{-1}\|_{op}^2 \|(T - \hat{T})(\alpha_\lambda - \alpha_0^{\min})\|_{\mathcal{H}}^2$$

(a) Bound on $\left\|T^{\frac{1}{2}}(\hat{T}+\lambda)^{-1}\right\|_{op}$ By [16, eq. 39], under \mathcal{E}_1

$$\left\| T^{\frac{1}{2}} (\hat{T} + \lambda)^{-1} \right\|_{op} \le \frac{1}{\sqrt{\lambda}}$$

(b) Bound on $\left\| (T - \hat{T})(\alpha_{\lambda} - \alpha_0^{\min}) \right\|_{\mathcal{H}}$ Under \mathcal{E}_2

$$\left\| (T - \hat{T})(\alpha_{\lambda} - \alpha_0^{\min}) \right\|_{\mathcal{H}} \le 2\ln(6/\eta) \left(\frac{2\kappa\sqrt{\mathcal{B}(\lambda)}}{n} + \sqrt{\frac{\kappa\mathcal{A}(\lambda)}{n}} \right)$$

In summary, under \mathcal{E}_1 and \mathcal{E}_2

$$S_{2}(\lambda, \{w_{i}\}) \leq \left\| T^{\frac{1}{2}}(\hat{T} + \lambda)^{-1} \right\|_{op}^{2} \left\| (T - \hat{T})(\alpha_{\lambda} - \alpha_{0}^{\min}) \right\|_{\mathcal{H}}^{2}$$

$$\leq \frac{1}{\lambda} \cdot \left\{ 2\ln(6/\eta) \right\}^{2} \cdot 2 \left(\left\{ \frac{2\kappa\sqrt{\mathcal{B}(\lambda)}}{n} \right\}^{2} + \left\{ \sqrt{\frac{\kappa\mathcal{A}(\lambda)}{n}} \right\}^{2} \right)$$

$$= 8\ln^{2}(6/\eta) \left(\frac{4\kappa^{2}\mathcal{B}(\lambda)}{n^{2}\lambda} + \frac{\kappa\mathcal{A}(\lambda)}{n\lambda} \right)$$

3. Bound on $S_1(\lambda, \{w_i\})$

Write

$$S_1(\lambda, \{w_i\}) \le \left\| T^{\frac{1}{2}} (\hat{T} + \lambda)^{-1} (T + \lambda)^{\frac{1}{2}} \right\|_{op}^2 \left\| (T + \lambda)^{-\frac{1}{2}} (\hat{\mu}^{(m)} - \hat{T} \alpha_0^{\min}) \right\|_{\mathcal{H}}^2$$

(a) Bound on
$$\|T^{\frac{1}{2}}(\hat{T}+\lambda)^{-1}(T+\lambda)^{\frac{1}{2}}\|_{op}$$

By [16, eq. 47], under \mathcal{E}_1
 $\|T^{\frac{1}{2}}(\hat{T}+\lambda)^{-1}(T+\lambda)^{\frac{1}{2}}\|_{op} \le 2$

$$\left\| T^{\frac{1}{2}} (\hat{T} + \lambda)^{-1} (T + \lambda)^{\frac{1}{2}} \right\|_{op} \le 2$$

(b) Bound on $\|(T+\lambda)^{-\frac{1}{2}}(\hat{\mu}^{(m)}-\hat{T}\alpha_0^{\min})\|_{\mathcal{H}}$ Under \mathcal{E}_3

$$\|(T+\lambda)^{-\frac{1}{2}}(\hat{\mu}^{(m)} - \hat{T}\alpha_0^{\min})\|_{\mathcal{H}} \le 2\ln(6/\eta) \left[\frac{1}{n} \sqrt{\Gamma^2 \frac{\kappa'}{\lambda}} + \sqrt{\frac{\Sigma^2 \mathcal{N}(\lambda)}{n}} \right]$$

Therefore by [16, eq. 49], under \mathcal{E}_1 and \mathcal{E}_3

$$S_1(\lambda, \{w_i\}) \le 32 \ln^2(6/\eta) \left(\frac{\kappa' \Gamma^2}{n^2 \lambda} + \frac{\Sigma^2 \mathcal{N}(\lambda)}{n}\right)$$

Proof of Theorem 5.2. Finally, I combine the abstract rate in Proposition D.7 with the bounds of Proposition D.4, following [88, Section 2]. The absolute constant depends only on $(R, \kappa, \kappa^{(m)}, \Gamma, \Sigma, \beta, b, c)$. Note that (Γ, Σ) introduce dependence on $(\|\alpha_0^{\min}\|_{\mathcal{H}}, \bar{L}^{(m)})$.

Proof of Corollary 5.1. I verify the conditions of Corollary 4.1 appealing to Theorems 5.1 and 5.2. In particular, by the symmetry of Theorems 5.1 and 5.2, I must show

$$(\sqrt{\bar{L}^{(m)}} + \bar{\alpha} + \bar{\alpha}' + \bar{\sigma})\sqrt{r_n} \to 0, \quad \sqrt{n} \cdot r_n \to 0$$

By Proposition 2.1, $\sqrt{\bar{L}^{(m)}} < \infty$ is a constant. By Proposition 5.1, $\bar{\alpha} < \infty$ and $\bar{\sigma} < \infty$ are also constants. However, $\bar{\alpha}'$ scales as $\lambda^{-\frac{1}{2}}$. Therefore the remaining conditions to verify are

$$\lambda^{-\frac{1}{2}}\sqrt{r_n} \to 0, \quad \sqrt{n} \cdot r_n \to 0$$

I consider each case

- 1. $b = \infty$.
 - (a) $n^{\frac{1}{4}} \cdot n^{-\frac{1}{2}} \to 0$
 - (b) $\sqrt{n} \cdot n^{-1} \to 0$
- 2. $b \in (1, \infty), c \in (1, 2]$.
 - (a) $n^{\frac{1}{2}\frac{b}{bc+1}} \cdot n^{-\frac{1}{2}\frac{bc}{bc+1}} \to 0 \iff c > 1$
 - (b) $\sqrt{n} \cdot n^{-\frac{bc}{bc+1}} \to 0 \iff bc > 1$
- 3. $b \in (1, \infty)$, c = 1.
 - (a) $\ln^{-\frac{1}{2}\frac{b}{b+1}}(n)n^{\frac{1}{2}\frac{b}{b+1}} \cdot \ln^{\frac{1}{2}\frac{b}{b+1}}(n)n^{-\frac{1}{2}\frac{b}{b+1}} = 1$
 - (b) $\sqrt{n} \cdot \ln^{\frac{b}{b+1}}(n) \cdot n^{-\frac{b}{b+1}} \to 0 \iff b > 1$

Proof of Corollary 5.2. Since $\|\alpha_0^{\min}\|_{\infty} \leq \bar{\alpha}$, the trimming can only possibly improve mean square error, and the rates in Theorem 5.2 continue to hold for the trimmed estimator $\tilde{\alpha}$ defined in Algorithm 5.1.

As in Corollary 5.1, I must show

$$(\sqrt{\bar{L}^{(m)}} + \bar{\alpha} + \bar{\alpha}' + \bar{\sigma})\sqrt{r_n} \to 0, \quad \sqrt{n} \cdot r_n \to 0$$

By Proposition 2.1, $\sqrt{\bar{L}^{(m)}} < \infty$ is a constant. By Proposition 5.1, $\bar{\alpha} < \infty$ and $\bar{\sigma} < \infty$ are also constants. Moreover, trimming implies that $\bar{\alpha}' = \bar{\alpha} < 0$ is a constant. Therefore the remaining condition to verify is

$$\sqrt{n} \cdot r_n \to 0$$

In the proof of Corollary 5.1, I verify this condition for across regimes.

E Tuning

E.1 Ridge penalty

I present two tuning procedures based on cross validation for the ridge penalty λ of the kernel ridge Riesz representer estimated in Algorithm 3.2. Each procedure finds a cross validated value $\lambda^* \in \Lambda$, where $\Lambda \subset \mathbb{R}$ is a grid of possible values.

The first approach exploits the sample splitting in Algorithm 3.1. Recall that $\hat{\alpha}_{\ell}$ is the kernel ridge Riesz representer estimated using observations in I_{ℓ}^c . By construction, $\hat{\alpha}_{\ell}$ depends on λ via Algorithm 3.2. The cross validation loss arises from the definition of the empirical loss $\mathcal{L}_{\lambda}^n(\alpha)$, dropping the ridge term and setting $\alpha = \hat{\alpha}_{\ell}$ for each observation $i \in I_{\ell}$.

Algorithm E.1 (Foldwise cross validation). *Partition the sample into folds* $\{I_{\ell}\}_{\ell=1:L}$. *Denote by* I_{ℓ}^{c} *observations not in fold* I_{ℓ} .

- 1. For each possible value $\lambda \in \Lambda$ and for each fold ℓ , estimate $\hat{\alpha}_{\ell}$ from observations in I_{ℓ}^{c} according to Algorithm 3.2
- 2. Set λ^* as

$$\lambda^* \in \operatorname*{argmin}_{\lambda \in \Lambda} \mathcal{L}(\lambda), \quad \mathcal{L}(\lambda) = \frac{1}{n} \sum_{\ell=1}^{L} \sum_{i \in I_{\ell}} \left\{ -2m(W_i, \hat{\alpha}_{\ell}) + \hat{\alpha}_{\ell}(W_i)^2 \right\}$$

The second approach exploits the theoretical symmetry between $\hat{\gamma}$ and $\hat{\alpha}$. Theorems 5.1 and 5.2 demonstrate that they require the same regularization sequence that balances bias and variance. The leave-one-out cross validation procedure for $\hat{\gamma}$ has a closed form solution derived in [81, Appendix G]. The tuned λ^* for $\hat{\gamma}$ can then be used with $\hat{\alpha}$.

Algorithm E.2 (Leave-one-out cross validation). Construct the matrices

$$H_{\lambda} := I - K_{WW}(K_{WW} + n\lambda)^{-1} \in \mathbb{R}^{n \times n}, \quad \tilde{H}_{\lambda} := diag(H_{\lambda}) \in \mathbb{R}^{n \times n}$$

where \tilde{H}_{λ} has the same diagonal entries as H_{λ} and off diagonal entries of 0. Then set

$$\lambda^* = \underset{\lambda \in \Lambda}{\operatorname{argmin}} \frac{1}{n} \|\tilde{H}_{\lambda}^{-1} H_{\lambda} Y\|_2^2, \quad \Lambda \subset \mathbb{R}$$

E.2 Kernel

In principle, I could instead use cross validation to tune kernel hyperparameters in the same way that I use cross validation to tune ridge penalties. However, given the choice of product kernel, this approach becomes impractical in high dimensions. Therefore I rely on heuristics to set kernel hyperparameters.