WEIERSTRASS MODELS

YING ZONG

1. Introduction

This paper draws inspiration from lagrangian fibrations on irreducible symplectic manifolds in general, and more specifically, from the question whether all these fibrations have smooth base. Recall that lagrangian fibrations on irreducible symplectic manifolds are equidimensional, at least in the framework of projective varieties, see Matsushita [6]. And when such a base S is also assumed to be smooth, S is then a projective space by Hwang [5].

In this direction, we proved the following theorem 1 in [9]. The theorem 2 is a further effort in studying a certain class of these fibrations.

Theorem 1. — Let S be a locally noetherian normal algebraic space of residue characteristic zero with regular formal fibers. Assume that there is a morphism $f: X \to S$, which is locally of finite type, equidimensional and admits étale local sections around every point of S, where X is a regular algebraic space. Then S is regular and f is flat.

Corollary. — If a lagrangian fibration $f: X \to S$ on an irreducible symplectic manifold X has local holomorphic sections around a point s of S, and if X and S are both projective varieties, then S is smooth over \mathbb{C} at s.

For then f admits formal sections around s, and one can apply Theorem 1.

Definition. — Keeping the hypotheses of Theorem 1, we say that f is a Weierstrass model on X if furthermore:

- a) f is locally for the étale topology of S projective.
- b) For every geometric codimension ≤ 1 point s of S, the fiber $f^{-1}(s)$ is irreducible.
- c) Each connected component S_0 of S has a geometric point s_0 such that $f^{-1}(s_0)$ admits a $k(s_0)$ -abelian variety structure.

The definition is motivated by elliptic K3, whose study often benefits from their contraction to plane models, the Weierstrass models.

It may happen that such a contraction is trivial, a phenomenon that is generalized here. The following theorem says that Weierstrass models are, to a large extent, group theoretical, being compactification of torsors in a very tight form. We propose a possible application below.

Note that in the condition c) if $S_{(s_0)}$ denotes the strict localization of S at s_0 , then $f \times_S S_{(s_0)}$ is smooth, and $X \times_S S_{(s_0)}$ admits an abelian scheme structure over $S_{(s_0)}$ (cf. [8] 7.2), and hence in particular, for every geometric generic point t of $S_{(s_0)}$, $f^{-1}(t)$ has a k(t)-abelian variety structure.

It follows that, if f is a Weierstrass model on X, then $f \times_S S'$ is a Weierstrass model on $X \times_S S'$ for every regular morphism $S' \to S$.

Theorem 2. — Let $f: X \to S$ be a Weierstrass model on X. Then there exist:

- a) An S-smooth group algebraic space A with connected fibers, which is Néron at every codimension 1 point of S.
 - b) An A-torsor Q on S for the étale topology.
 - c) An open S-immersion $j: Q \to X$.

Such (A, Q, j) is unique up to unique isomorphisms and of formation compatible with every base change by a regular morphism $S' \to S$.

We call Q the *identity component*, and A the *albanese*, of the Weierstrass model. As in the case of plane models, A in general has fibers with non-trivial unipotent part.

The proof, unlike that of Theorem 1, does not need assume S has excellent local rings, and is insensitive to characteristics.

Proposition 3. — Let $f: X \to S$ be a Weierstrass model on X, with S of finite type over a field k. Let s be a point of S. Assume that, in a neighborhood of $f^{-1}(s)$, f is lagrangian with respect to a symplectic structure Ψ on X/k.

Then A_s is of abelian rank at least tr.deg. k(s)/k.

This estimate says, for example, that, for s of codimension 1 in S, A_s , if it degenerates, is extension of an abelian variety B by a form of \mathbf{G}_m or \mathbf{G}_a . In the \mathbf{G}_m case, if \overline{s} denotes a geometric point over s, there is a \mathbf{P}^1 -bundle P over $B_{\overline{s}}$ with two sections $0, \infty : B_{\overline{s}} \to P$ such that $f^{-1}(\overline{s})$ is obtained from P by pinching 0 and ∞ ; this $f^{-1}(\overline{s})$ is one particular instance of a minimal model of L^* we classified in [10].

Keeping the hypotheses of Theorem 2, assume now that X is an irreducible symplectic manifold with its symplectic structure Ψ and hyperkahler metric ω_q (cf. [2]). Then $H^0(\mathcal{O}_S) = H^0(f_*\mathcal{O}_X) = H^0(\mathcal{O}_X) = \mathbf{C}$, S is proper smooth over \mathbb{C} , and f is a lagrangian fibration, at least when S is projective over \mathbb{C} ([6] Theorem 1). Notice that all terms of the exact sequence

$$0 \to f^*\Omega^1_S|Q \to \Omega^1_Q \to \Omega^1_{Q/S} \to 0$$

have natural action by A. So by descent one has the exact sequence

$$0 \to \Omega_S^1 \to \omega_Q^1 \to \omega_{Q/S}^1 \to 0$$

with ω_Q^1 and $\omega_{Q/S}^1$ consisting of the A-invariant forms of Ω_Q^1 and $\Omega_{Q/S}^1$, respectively. Also, $\Psi|Q$ by descent induces a symplectic form ψ on ω_Q^1 , relative to which Ω_S^1 is lagrangian.

We now define a hermitian metric on S: for each local holomorphic 1-form α on S, let it be considered as an A-invariant form on Q and then extended uniquely to X ($\operatorname{codim}(X-Q,X)\geq 2$), say α' ; let $|\alpha'|_g$ denote its norm relative to the hyperkahler metric ω_g on X. Clearly, the fiberwise integral

$$\alpha \mapsto |\alpha|^2 := \int_{X/S} |\alpha'|_g^2 \,\omega_g^n$$

where n is the relative dimension of X/S, induces a hermitian metric on S, which I speculate is Kahler-Einstein given the symmetry it enjoys.

2. Proof of Theorem 2

Recall that Néron models are of formation compatible with all base change by regular morphisms of spectra of discrete valuation rings [3] 7.2/1. The question is thus local for the étale topology of S. We may assume f projective, and that S is a scheme, quasi-compact, irreducible and with generic point t.

I) Case where S = t:

By assumptions of a Weierstrass model, $X \times_t \overline{t}$ admits a \overline{t} -abelian variety structure, for \overline{t} a geometric point of t = S. The assertion then follows by [8] 7.2: X = Q admits a unique torsor structure under its albanese A, dual abelian variety of $\operatorname{Pic}_{X/t}^o$.

II) Case where S is the spectrum of a discrete valuation ring:

Let s be the closed point of S, and let U consist of all those points of X where f is smooth. Since we assume that f admits étale local sections around s, U_s is non-empty and hence dense in the geometrically irreducible closed fiber X_s .

We have shown that the generic fiber $U_t = X_t$ is a torsor under its albanese A_t . It follows that U is weak S-Néron [3] 3.5/1 by definition,

and, as U_s is irreducible, is also minimal, loc.cit. 4.3/2, and finally, by loc.cit. 6.5 Corollary 3, that U is S-dense open in the S-Néron model Q of $U_t = X_t$, and that Q is a torsor under the Néron model A of A_t .

The fiber Q_s , thus A_s as well, is geometrically irreducible, since U_s is geometrically irreducible and dense in Q_s .

To finish, it suffices to show that U = Q.

Let L be an ample invertible module on X; such L exists by the hypothesis that X be projective over S. Notice that Q-U (resp. X-U) is of codimension ≥ 2 in Q (resp. X). As Q is regular, Q is parafactorial along Q-U, so L|U uniquely extends to an invertible module N on Q. It remains only to see that N is ample on Q, because

$$X = \operatorname{Proj} \sum H^0(L^{\otimes n}) = \operatorname{Proj} \sum H^0(L^{\otimes n}|U) = \operatorname{Proj} \sum H^0(N^{\otimes n}).$$

That N is ample follows immediately from Raynaud's theory of cube [7] VI 2.1; it is enough to find an integer n > 0 and a section $\sigma \in H^0(N^{\otimes n})$ such that

$$W := \{ y \in Q, \sigma(y) \neq 0 \}$$

is contained in U, affine, with $W_s \neq \emptyset$. Such a section σ exists, since

$$H^0(N^{\otimes n}) = H^0(L^{\otimes n}|U) = H^0(L^{\otimes n})$$

and L is ample.

- III) General case:
- III.1) Uniqueness and existence of A:
- i) If A exists, it is unique up to unique isomorphisms:

This follows by [4] Note (1) 3: The functor $A \mapsto A_t$ is fully faithful for those A that is Néron at all codimension 1 points of S.

ii) In order that the S-smooth group algebraic space A exists, it suffices that it exists over an S-algebraic space S' that is smooth surjective over S:

For, by i) and I), the asserted S'-smooth group algebraic space A' will admit a descent datum relative to $S' \to S$. Such a descent is effective by Artin [1] 6.3.

iii) Let S' be the open sub-scheme of X consisting of all points where f is smooth. Then $S' \to S$ is surjective:

Because S is regular and we assume that f admits étale local sections around every point of S, cf. [3] 3.1/3.

iv) For the existence of A, one can assume that f admits an S-section, say $e: S \to X$:

One takes the base change $S' \to S$, with S' as in iii), and applies ii).

v) Let $e: S \to X$ be as in iv), let U consist of all points of X where f is smooth, and let V consist of all those points x of U such that x and e(f(x)) lie in the same connected component of $U_{f(x)}$. Then V is an open sub-scheme of U and has geometrically irreducible fibers over S:

That V is open in U follows from EGA IV 15.6.5. Its S-fibers are by definition connected, thus geometrically connected, since the S-section $e:S\to X$ factors through V, and finally geometrically irreducible, since $V\to S$ is smooth.

vi) Let V be as in v). There exists on V an S-birational group law with zero section $e: S \to V$. Such a structure is unique up to unique isomorphisms:

Notice that $V_t = X_t$, which is, in a unique way, an abelian variety A_t with identity e(t). And, we have shown above that, when localized at any codimension 1 point s of S, V = U coincides with the Néron model of A_t over $\operatorname{Spec}(\mathcal{O}_{S,s})$. The sought-after S-birational composition law m on V therefore, by the technique of "passage à la limite projective", exists over an open sub-scheme S' of S, with S - S' of codimension ≥ 2 in S.

Now [7] IX 1.1 implies that the domain of definition of m is an open neighborhood of $(e \times_S e)(S)$, thus m exists throughout; it verifies the axioms of commutativity, associativity, etc, because it does over t.

viii) There exists an open neighborhood V' of e(S) in V such that (V', m|V', e) admits an S-dense open immersion into an S-group algebraic space A, compatible with the group laws.

One applies SGA 3 XVIII 3.2+3.5+3.7 to the birational group law of vii).

ix) The S-group algebraic space A of viii) is a scheme, separated, smooth, of finite type, and with connected fibers over S:

That A is a scheme (resp. S-separated, resp. of finite type over S) follows by [3] 6.2 b), (resp. a), resp. b)), because V' is quasi-projective over S. And, A is S-smooth with connected fibers, because V' is S-smooth with geometrically connected fibers and the morphism $V' \times_S V' \to A$, induced by the group law of A, is faithfully flat, which one can verify fiber by fiber.

x) One has V = A:

Let L be an S-ample invertible module on X, which exists since f is by assumption projective. The restriction L|V' extends uniquely to an invertible module N on A. Indeed, as A, being smooth over S, is regular and $\operatorname{codim}(A-V',A) \geq 2$, one has that A is parafactorial along

A-V'. Observe also that X-V' is of codimension ≥ 2 in X. Now, N is S-ample on A, which one proves by localizing at each point s of S and argues similarly as in II). This concludes the proof, because

$$X = \operatorname{Proj} \sum f_* L^{\otimes n} = \operatorname{Proj} \sum (f|V')_* (L^{\otimes n}|V') = \operatorname{Proj} \sum a_* N^{\otimes n},$$

where $a:A\to S$ denotes the structural morphism of A.

III.2) When f admits an S-section e, the (A,Q,j) exists as desired and remains so after every base change by a regular morphism :

One takes $j: Q \to X$ to be the inclusion of A = V in X.

III.3) The (A, Q, j) in III.2) is independent of the choice of S-sections of f, when such sections exist:

Let e_1, e_2 be two S-sections of f, and let (A_i, Q_i, j_i) denote the triple constructed relative to e_i , i = 1, 2. Identify Q_i with its image in X, and write $Q = Q_1 \cap Q_2$.

Observe that $Q_i - Q$ is of codimension ≥ 2 in Q_i . Indeed, Q_1 and Q_2 coincide over every codimension ≤ 1 point of S, by I)+II). It follows that, this intersection Q, when viewed as a birational correspondence between Q_1 and Q_2 , is, according to Weil [3] 4.4/1, in fact biregular. So the claim $Q_1 = Q_2$ is justified.

Write + for the group law of A = Q with origin e_1 . There exists a unique element $\alpha_{12} \in A(S)$ satisfying

$$\alpha_{12} + e_1 = e_2$$
.

III.4) Completion of the proof:

Clearly, varying the sections of f over all S-smooth schemes S', we have produced, through the system of α_{12} , a unique descent datum gluing the locally constructed (A, Q, j). This finishes the proof.

3. Proof of Proposition 3

The field k necessarily is of characteristic 0, and X and S are smooth over $\operatorname{Spec}(k)$.

Restricting to a neighborhood of $f^{-1}(s)$, we write $L: T_{X/k} \simeq \Omega^1_{X/k}$ for the isomorphism induced by the symplectic structure Ψ .

Let the closed image of Q_s in X_s be W, and let $Z \to W$ be a resolution of singularities, centered in $W - Q_s$, so that Q_s admits naturally an open dense immersion φ into Z.

Consider the composition

$$T_{Z/k} \to T_{X/k} \otimes \mathcal{O}_Z \stackrel{L \otimes 1}{\to} \Omega^1_{X/k} \otimes \mathcal{O}_Z \to \Omega^1_{Z/k} \to \Omega^1_{Z/s}$$

where

$$T_{Z/k} \to T_{X/k} \otimes \mathcal{O}_Z$$

 $\Omega^1_{X/k} \otimes \mathcal{O}_Z \to \Omega^1_{Z/k} \to \Omega^1_{Z/s}$

are induced by the morphism $Z/s \to X/k$.

Call this composition $T_{Z/k} \to \Omega^1_{Z/s}$, β ; we show below that it is of kernel $T_{Z/s}$. Note first that all terms of the exact sequence

$$0 \to T_{Z/s} \to T_{Z/k} \to T_{s/k} \otimes \mathcal{O}_Z \to 0$$

are locally free over \mathcal{O}_Z .

a) One has $\beta | T_{Z/s} = 0$, hence β induces a morphism $\overline{\beta} : T_{s/k} \otimes \mathcal{O}_Z \to \Omega^1_{Z/s}$:

For, $\beta|T_{Z/s}$, a section over Z of $\mathcal{H}om(T_{Z/s},\Omega^1_{Z/s})\simeq (\Omega^1_{Z/s})^{\otimes 2}$, vanishes on Q_s by assumption that f is lagrangian.

b) The morphism $\overline{\beta}$ is injective:

It is injective over Q_s , because f is lagrangian, and hence is injective throughout, since $T_{s/k} \otimes \mathcal{O}_Z$ is locally free over \mathcal{O}_Z .

c) One has $\dim_{k(s)} H^0(Z, \Omega^1_{Z/s}) \ge \dim_{k(s)} T_{s/k} = \text{tr.deg. } k(s)/k$:

This follows by b).

d) Reduction to a lemma:

The restriction $\varphi^*: H^0(Z,\Omega^1_{Z/s}) \to H^0(Q_s,\Omega^1_{Q_s/s})$ is injective. To finish, one combines c) with the lemma below applied to the open immersion $\varphi_{\overline{s}}: Q_{\overline{s}} \hookrightarrow Z_{\overline{s}}$, where \overline{s} is a geometric point over s.

Lemma. — Let k be a field of characteristic 0. Let A be a connected k-algebraic group, with generic point η , and with Chevalley decomposition

$$1 \to L \to A \to B \to 1$$

so that B is an abelian variety and L affine connected. Let φ be a k-rational map of A to a proper k-algebraic space Z.

Then

Image
$$\varphi^*: H^0(Z, \Omega^1_{Z/k}) \to \Omega^1_{\eta/k}$$

is contained in

Image
$$H^0(B, \Omega^1_{B/k}) \to H^0(A, \Omega^1_{A/k}) \to \Omega^1_{n/k}$$

In particular, the abelian rank of A verifies:

$$\dim(B) \ge \operatorname{rank}_k(\varphi^*)$$

Proof. Recall that every algebraic group over a field of characteristic 0 is smooth (Cartier). In particular, A and L are smooth over k, the projection $p:A\to B$ is smooth and faithfully flat, $p^*:H^0(B,\Omega^1_{B/k})\to\Omega^1_{\eta/k}$ is injective and of rank dim(B), and the lemma amounts to asserting the existence of a unique k-linear map

$$\rho: H^0(Z, \Omega^1_{Z/k}) \to H^0(B, \Omega^1_{B/k})$$

such that

$$\varphi^* = p^* \circ \rho.$$

Under this form, we may and will assume $k = \mathbf{C}$.

Write U for the domain of definition of φ , and consider φ as a morphism of U to Z. One has $\operatorname{codim}(A - U, A) \geq 2$, since A, being k-smooth, is normal, and Z/k is proper.

a) One may assume φ is an open schematically dense immersion : Decompose φ as

$$U \stackrel{\varphi'}{\hookrightarrow} Z' \stackrel{f}{\rightarrow} Z$$

with f proper, and φ' an open schematically dense immersion. Clearly, the lemma holds for (φ, Z) if it holds for (φ', Z') . It suffices to consider the latter.

b) One may assume furthermore Z smooth over k:

Assume by a) that φ is an open schematically dense immersion. Let $f: Z' \to Z$ be a resolution of singularities, centered in $Z - \varphi(U)$, so that φ factors as

$$U \stackrel{\varphi'}{\hookrightarrow} Z' \stackrel{f}{\rightarrow} Z$$

with φ' an open schematically dense immersion. As in a), it is sufficient to study (φ', Z') .

c) The compostion

$$H^0(Z,\Omega^1_{Z/k}) \stackrel{\varphi^*}{\to} H^0(U,\Omega^1_{A/k}) \to H^0(U,\Omega^1_{A/B})$$

is zero:

This is obvious if $p:A\to B$ is an isomorphism, i.e., if L=1. Assume L non-trivial.

By structure of affine algebraic groups, it is enough to show that, for each $b \in B(k)$ and each curve C in $p^{-1}(b) \simeq L$ of the form \mathbf{G}_a or \mathbf{G}_m , the composition

$$(\varphi|C)^*: H^0(Z,\Omega^1_{Z/k}) \xrightarrow{\varphi^*} H^0(U,\Omega^1_{A/k}) \to H^0(U,\Omega^1_{A/B}) \to H^0(C\cap U,\Omega^1_{C/k})$$
 is zero.

Now Z/k being proper, if $C \cap U \neq \emptyset$, $\varphi|C : C \cap U \to Z$ extends to a morphism of \mathbf{P}^1 to Z, but $H^0(\mathbf{P}^1, \Omega^1_{\mathbf{P}^1/k}) = 0$, hence c).

d) Let (φ, Z) be as in a), b). For each $\alpha \in H^0(Z, \Omega^1_{Z/k})$, one has

$$\varphi^*\alpha = \sum a_i \ p^*e^i$$

where $\{e^i\}$ is a fixed basis of $H^0(B,\Omega^1_{B/k})$, and $a_i \in H^0(U,\mathcal{O}_A)$:

This follows from c) and the exact sequence

$$0 \to p^* \Omega^1_{B/k} \to \Omega^1_{A/k} \to \Omega^1_{A/B} \to 0$$

e) All coefficients $a_i \in H^0(U, \mathcal{O}_A)$ in d) belong to k:

As $H^0(U, \mathcal{O}_A) = H^0(A, \mathcal{O}_A)$, the claim e) is clear if B = 1 or if $p: A \to B$ is an isomorphism. Assume $n = \dim(B) > 0$ and L non-trivial. Choose a volume form

$$\Phi \in H^0(B,\Omega^n_{B/k})$$

and let $e_j \in H^0(B, \Omega^{n-1}_{B/k})$ be such that

$$e^i \wedge e_j = \delta^i_j \Phi$$

 $\forall i, j \in \{1, \cdots, n\}.$

One deduces from the expression of $\varphi^*\alpha$ in d) that

$$\varphi^* \alpha \wedge p^* e_i = a_i \ p^* \Phi$$

and by differentiation that

$$d_{U/k}(a_j) \wedge p^* \Phi = 0$$

Indeed,

$$d_{Z/k}(\alpha) = 0, \ d_{B/k}(\Phi) = 0, \ d_{B/k}(e_i) = 0$$

by the E_1 -degeneration of Hodge-De Rham spectral sequences.

Now, $d_{U/k}(a_j) \wedge p^*\Phi = 0$ is equivalent to $d_{U/B}(a_j) = 0$. In fact, in terms of the Koszul filtration

$$K^i := p^* \Omega^i_{B/k} \wedge \Omega^{*-i}_{A/k}$$

on $\Omega_{A/k}^*$, one has $d_{U/k}(a_j) \wedge p^*\Phi \in H^0(U, K^n\Omega_{A/k}^{n+1})$, which, under the isomorphism

$$K^n = K^n/K^{n+1} \simeq \Omega^{*-n}_{A/B} \otimes \Omega^n_{B/k}$$

is mapped to $d_{U/B}(a_j) \otimes \Phi$.

And, $d_{U/B}(a_i) = 0$ implies that, when viewed as an analytic function,

$$a_j \in \text{Image } H^0(V^{an}, \mathcal{O}_B^{an}) \to H^0(U^{an}, \mathcal{O}_A^{an})$$

where V = p(U), since $p: U \to V$ is smooth surjective with geometrically connected fibers.

Observe that U intersects with all codimension 1 fibers of p. One has hence $\operatorname{codim}(B-V,B) \geq 2$, $H^0(V^{an},\mathcal{O}_B^{an}) = H^0(B^{an},\mathcal{O}_B^{an}) = k$. This concludes that $a_j \in k$.

f) Completion of the proof:

By e), all $a_i \in k$. The map

$$\rho: \alpha \mapsto \sum a_i e^i$$

is as desired.

References

- M. Artin. Versal deformations and algebraic stacks. Invent. math. 27, 165–189, 1974.
- 2. A. Beauville. Variétès kahleriennes dont la première classe de Chern est nulle. JDG 18, 755-782, 1983.
- 3. S. Bosch, W. Lütkebohmert, M. Raynaud. *Néron models*. Ergebnisse der Mathematik und ihrer Grenzgebiete, 1990.
- 4. P. Deligne. Le lemme de Gabber. Astérisque 127, 1985.
- 5. J. Hwang. Base manifolds for fibrations of projective irreducible symplectic manifolds. Invent. math. 174, 625-644, 2008.
- 6. D. Matsushita. Equidimensionality of lagrangian fibrations on holomorphic symplectic manifolds. Mathematical Research Letters 7, 389-391, 2000.
- M. Raynaud. Faisceaux amples sur les schémas en groupes et les espaces homogènes. LNM 119, 1970.
- 8. Y. Zong. Basic finite étale equivalence relations. arxiv.org/abs/1512.00097.
- 9. Y. Zong. Equidimensionality and regularity. arxiv.org/abs/1802.03846.
- 10. Y. Zong. Minimal models of L^* . arxiv.org/abs/1903.09740.

Email address: zongying.upenn@gmail.com