

FROBENIUS GROUPS WITH PERFECT ORDER CLASSES

JAMES MCCARRON

ABSTRACT. The purpose of this paper is to investigate the finite Frobenius groups with “perfect order classes”; that is, those for which the number of elements of each order is a divisor of the order of the group. If a finite Frobenius group has perfect order classes then so too does its Frobenius complement, the Frobenius kernel is a homocyclic group of odd prime power order, and the Frobenius complement acts regularly on the elements of prime order in the Frobenius kernel. The converse is also true.

Combined with elementary number-theoretic arguments, we use this to provide characterisations of several important classes of Frobenius groups. The insoluble Frobenius groups with perfect order classes are fully characterised. These turn out to be the perfect Frobenius groups whose Frobenius kernel is a homocyclic 11-group of rank 2.

We also determine precisely which nilpotent Frobenius complements have perfect order classes, from which it follows that a Frobenius group with nilpotent complement has perfect order classes only if the Frobenius complement is a cyclic $\{2, 3\}$ -group of even order.

Those Frobenius groups for which the Frobenius complement is a biprimary group are also described fully, and we show that no soluble Frobenius group whose Frobenius complement is a $\{2, 3, 5\}$ -group with order divisible by 30 has perfect order classes.

CONTENTS

1. Introduction	2
2. Preliminaries	4
2.1. A Bit of Number Theory	4
2.2. General Remarks on Groups	5
2.3. Frobenius Groups	7
2.4. Groups with Perfect Order Classes	7
2.5. Supporting Computations	9
3. The Structure of Order Classes in Frobenius Groups	9
4. Insoluble Frobenius Groups	12

Date: July 13, 2023.

5. Soluble Frobenius Groups	17
5.1. Frobenius Groups with Cyclic Complement	18
5.2. Frobenius Groups with Nilpotent Complement	21
5.3. Frobenius Groups with Non-Nilpotent Biprimary Complement	26
5.4. Frobenius Groups with Soluble $\{2, 3, 5\}$ -Complement That Do Not Exist	32
6. Concluding Remarks	41
References	41

1. INTRODUCTION

A finite group (and all our groups are supposed to be finite) is said to have **perfect order classes** if the number of elements of any given order either is zero or is a divisor of the order of the group. The non-cyclic group of order 4 does not have perfect order classes since the number of elements of order 2 is equal to 3, which is not a divisor of its order 4. However, the cyclic group of order 4 does have perfect order classes, because it has 2 elements of order 4 and one element each of orders 1 and 2.

Groups with perfect order classes (often, “subsets”) were introduced by Finch and Jones in [7], and have subsequently been studied by a number of authors. See, for example, [6, 8, 9, 12, 21, 22, 24].

The object of this paper is to investigate Frobenius groups with perfect order classes. Our first main result reduces the study of Frobenius groups with perfect order classes to that of Frobenius complements with perfect order classes.

Theorem A. *Let $G = K \rtimes_f H$ be a Frobenius group with Frobenius kernel K and Frobenius complement H . Then G has perfect order classes if, and only if, the following are true:*

- (a) H has perfect order classes;
- (b) K is a homocyclic p -group for an odd prime number p ; and,
- (c) $|H| = p^r - 1$, where r is the rank of K .

Condition (c) of this theorem is equivalent to the assertion that the Frobenius complement H acts regularly on the elements of order p in G , and it follows that they comprise a single conjugacy class in G . We have chosen the arithmetic formulation for this condition because it is most directly applicable in subsequent results.

This is advantageous because a great deal is known about the structure of Frobenius complements. For example, we are able to give a

complete and precise description of the insoluble Frobenius groups with perfect order classes.

Theorem B. *An insoluble Frobenius group has perfect order classes if, and only if, it is perfect and its Frobenius kernel is a homocyclic 11-group of rank 2.*

We remark that a Frobenius group is perfect if, and only if, its Frobenius complement is isomorphic to the binary icosahedral group of order 120.

We now turn to the richer class of soluble Frobenius groups, for which we are able to describe several important cases. We are able to completely characterise Frobenius groups with perfect order classes whose Frobenius complement is nilpotent. To state our result for this case, we note that a **Pierpont prime** [1, 18] is a prime number p of the form $p = 1 + 2^\alpha 3^\beta$, for some integers $\alpha, \beta \geq 0$. (If $\beta = 0$, then p is a Fermat prime, provided that $\alpha > 0$. But see [5].) At the time of writing it remains an open problem whether there exist infinitely many Pierpont primes [25].

Theorem C. *Let $G = K \rtimes_f H$ be a Frobenius group with nilpotent Frobenius complement H . Then G has perfect order classes if, and only if, one of the following is true.*

- (1) *K is a cyclic p -group, for some Pierpont prime $p = 1 + |H| > 2$, and H is a cyclic $\{2, 3\}$ -group of even order; or*
- (2) *G is isomorphic to one of the following groups:*
 - (a) $(C_{3^k} \times C_{3^k}) \rtimes_f C_8$;
 - (b) $(C_{5^k} \times C_{5^k}) \rtimes_f C_{24}$;
 - (c) $(C_{7^k} \times C_{7^k}) \rtimes_f C_{48}$; or,
 - (d) $(C_{17^k} \times C_{17^k}) \rtimes_f C_{288}$,*for some positive integer k .*

Observe that, in all cases, the Frobenius complement is cyclic.

It seems an interesting feature of our investigations that there are Frobenius complements with perfect order classes that nevertheless do not “fit” within any Frobenius group with perfect order classes. We completely characterise the non-cyclic nilpotent Frobenius complements with perfect order classes (there are two infinite series), but show that none occurs in a Frobenius group with perfect order classes. We also exhibit some soluble but non-nilpotent examples.

For the case in which the Frobenius complement is soluble but not nilpotent we are able to characterise the Frobenius groups with perfect order classes among those whose Frobenius complement is biprimary (that is, whose order has just two prime divisors).

Theorem D. *Let $G = K \rtimes_f H$ be a Frobenius group and assume that the Frobenius complement H is a non-nilpotent $\{2, q\}$ -group, where q is an odd prime. Then G has perfect order classes if, and only if, q is equal either to 3 or to 5 and G has one of the following forms:*

- (1) $G \simeq C_{5^k}^2 \rtimes_f \text{SL}(2, 3)$;
- (2) $G \simeq C_{5^k}^2 \rtimes_f (C_3 \rtimes C_8)$;
- (3) $G \simeq C_{7^k}^2 \rtimes_f (C_3 \rtimes C_{16})$;
- (4) $G \simeq C_{17^k}^2 \rtimes_f (C_9 \rtimes C_{32})$; or,
- (5) $G \simeq C_{3^k}^4 \rtimes_f (C_5 \rtimes C_{16})$.

We saw that an insoluble Frobenius group with perfect order classes is perfect with Frobenius complement isomorphic to the $\{2, 3, 5\}$ -group $\text{SL}(2, 5)$. It turns out that there are no Frobenius groups with perfect order classes in which the Frobenius complement is a soluble $\{2, 3, 5\}$ -group.

Theorem E. *A soluble Frobenius group whose Frobenius complement is a $\{2, 3, 5\}$ -group with order divisible by 30 does not have perfect order classes.*

Our results suffice to explain all the Frobenius groups with perfect order classes of order at most 15000 in the library of Frobenius groups up to that order in the computer algebra system MAPLE.

As is common in studying groups with perfect order classes, a considerable amount of number theory (all of an elementary nature) makes its way into our arguments.

In Section 2 we review some necessary background. Our first main result, Theorem A, is proved in Section 3. The proof of Theorem B occupies Section 4, and Theorems C, D and E are proved in Section 5. In Section 6 we suggest some problems for future work.

2. PRELIMINARIES

In this section, we describe our notation and terminology, as well as some required background from number theory and required results on groups with perfect order classes.

2.1. A Bit of Number Theory. Throughout the paper, a **prime** means a positive rational prime. By a **prime power** we mean a non-negative integer power of a prime. A prime power different from unity is said to be **well-defined**, and a prime power that is composite is said to be **proper**. If π is a set of primes, then a positive integer is a **π -number** if every one of its prime divisors belongs to π . For instance, 12 is a $\{2, 3, 61\}$ -number. The greatest common divisor of integers a

and b is denoted by $\gcd(a, b)$. We write $\phi(n)$ for Euler's function of the positive integer n . Recall that, for a well-defined prime power p^k , we have

$$\phi(p^k) = p^{k-1}(p - 1),$$

and that for relatively prime a and b , we have $\phi(ab) = \phi(a)\phi(b)$.

We shall make extensive use of the following result [2–4, 27].

2.1. Theorem (Zsigmondy's Theorem). *Let a and b be coprime positive integers such that $a > b$, let n be an integer greater than 1, and let $\epsilon = \pm 1$. If $\epsilon = 1$, assume that $(a, b, n) \neq (2, 1, 3)$ and, if $\epsilon = -1$, assume that $(a, b, n) \neq (2, 1, 6)$ and that, if $n = 2$, then $a + b$ is not a power of 2. Then $a^n + \epsilon b^n$ has a prime divisor q such that q does not divide $a^k + \epsilon b^k$, for any positive integer k less than n .*

The prime q in the statement of Zsigmondy's theorem is called a **primitive prime divisor** of $a^n + \epsilon b^n$.

Most often, Zsigmondy's theorem will be applied in the case for which $b = 1$ and $\epsilon = -1$. We can then interpret Zsigmondy's theorem as an assertion that, apart from the indicated exceptions, any positive integer a can be taken to have any given order modulo a suitable prime. Since, by Euler's theorem, we also have $a^{q-1} \equiv 1 \pmod{q}$, it follows that $q - 1$ is a multiple of n and, in particular, that $q > n$.

We shall also make frequent use of several well-known results on consecutive prime powers. The first says that 8 and 9 are the only consecutive proper prime powers. (This is much more elementary than the now proved [16] Catalan conjecture.) See, for example, [19, Chapter 7].

2.2. Lemma. *Let p and q be prime numbers. If a and b are integers greater than 1 such that $p^a - q^b = 1$, then $(p, q, a, b) = (3, 2, 2, 3)$.*

We shall also need to allow improper prime powers in certain cases.

2.3. Lemma. *If a and b are non-negative integers such that $2^a - 3^b = 1$, then $(a, b) \in \{(1, 0), (2, 1)\}$.*

2.4. Lemma. *If a and b are non-negative integers such that $3^b - 2^a = 1$, then $(a, b) \in \{(1, 1), (3, 2)\}$.*

2.2. General Remarks on Groups. All groups are supposed to be finite, and will generally be written multiplicatively. We use 1 to denote the identity element of any group, as well as to denote the trivial subgroup of a group.

If g is an element of a group G , then g^G denotes the conjugacy class of g in G , and $|g|$ is the order of g . In general, $|S|$ denotes the cardinality of a set S .

For a positive integer n , the cyclic group of order n is written C_n , while S_n , A_n and D_n denote, respectively, the symmetric group, the alternating group, and the dihedral group of degree n . Note that with this convention, we have $|D_n| = 2n$.

If $n \geq 3$, then Q_n denotes the generalised quaternion group of order 2^n , so that Q_3 is the ordinary quaternion group of order 8. Recall that Q_n has a presentation of the form

$$Q_n = \langle x, y \mid x^{2^{n-1}} = 1, y^2 = x^{2^{n-2}}, xy = x^{-1} \rangle.$$

It has an unique maximal cyclic subgroup $\langle x \rangle$ of index 2. Every element of Q_n belonging to the non-trivial coset of $\langle x \rangle$ in Q_n has order 4, and $\langle x \rangle$ has just two elements of order 4, for a total of $2^{n-1} + 2$ elements of order 4. Elements of every other order belong to the maximal cyclic subgroup $\langle x \rangle$, so we have the following result.

2.5. Lemma. *Let Q_n be a generalised quaternion group of order 2^n , where $n \geq 3$, and for each positive integer d , let $f_d(Q_n)$ denote the number of elements of order d in Q_n . Then*

$$f_{2^k}(Q_n) = \begin{cases} 1, & \text{for } k = 0, 1 \\ 2^{n-1} + 2, & \text{for } k = 2 \\ 2^{k-1}, & \text{for } 3 \leq k < n. \end{cases}$$

If G is a group, and r is a non-negative integer, then G^r denotes the direct product of r copies of G , with the convention that G^0 is a trivial group. In particular, if n is also a positive integer, then C_n^r denotes the **homocyclic group** of rank r and exponent n . (In the sequel, n will usually be a power of a prime.) We remark that $A \times B$ is the direct product of groups A and B , while $A \rtimes B$ is a semidirect product of A and B , where A is normal in $A \rtimes B$, and A and B have trivial intersection.

For homocyclic p -groups, we use frequently the following result counting elements of a given order.

2.6. Lemma. [8, Lemma 1] *Let p be a prime, and let r and k be positive integers. For each integer i with $1 \leq i \leq k$, the number of elements of order p^i in a homocyclic p -group of rank r and exponent p^k is given by*

$$f_{p^i}(C_{p^k}^r) = p^{r(i-1)}(p^r - 1).$$

If p is a prime, G is a p -group and n is a non-negative integer, then $\Omega_{(n)}(G) = \{g \in G : g^{p^n} = 1\}$, and $\Omega_n(G) = \langle \Omega_{(n)}(G) \rangle$. If G is abelian, then $\Omega_n(G) = \Omega_{(n)}(G)$.

The derived subgroup of a group G is denoted $[G, G]$, and the derived quotient, or **abelianisation** $G/[G, G]$ of G is written G^{ab} .

A group G is a **Z-group** if every Sylow subgroup of G is cyclic. It is well-known that every Z-group is metacyclic and has a presentation of the form

$$G = \langle x, y \mid x^\alpha, y^\beta, x^y = x^\gamma \rangle,$$

where the positive integers α, β and γ satisfy $\gcd(\alpha, \beta) = 1 = \gcd(\alpha, \gamma - 1)$ and $\gamma^\beta \equiv 1 \pmod{\alpha}$. Moreover, $|G| = \alpha\beta$ and $\langle x \rangle$ is the derived subgroup of G . If σ is the order of γ in the group $(\mathbb{Z}/\alpha\mathbb{Z})^\times$ of units of the ring $\mathbb{Z}/\alpha\mathbb{Z}$, then the centre of G is the subgroup $\langle y^\sigma \rangle$.

2.3. Frobenius Groups. For background on the general theory of Frobenius groups the reader may consult [11, 17, 20, 26]. Proofs of everything not proved here may be found in those cited references.

We recall that a group G is said to be a **Frobenius group** if it has a proper non-trivial subgroup H , called a **Frobenius complement**, such that $H \cap H^g = 1$, for all $g \in G \setminus H$. (We also say that H is **malnormal** in G .) Then Frobenius' theorem asserts that the set

$$K = \{1\} \cup \bigcup_{g \in G} G \setminus H^g$$

is a normal (indeed, characteristic) subgroup of G , called the **Frobenius kernel** of G . It follows that H acts faithfully and without fixed points on K . We say that the action of H on K is “Frobenius”. Then G is the semidirect product $K \rtimes H$, which we write as $K \rtimes_f H$ to indicate that the action of H on K is Frobenius. The Frobenius kernel of G is uniquely determined as the Fitting subgroup, and a Frobenius complement is uniquely determined up to conjugacy.

The Frobenius kernel and complement have relatively prime orders and, in fact, $|H|$ is a divisor of $|K| - 1$. Every normal subgroup of G either is contained in K or contains it. If N is a non-trivial normal subgroup of G contained in K , then the subgroup NH is a Frobenius group and, if N is a proper subgroup of K , then the quotient group G/N is a Frobenius group with Frobenius kernel isomorphic to K/N and Frobenius complement isomorphic to H . The Sylow subgroups of every Frobenius complement are either cyclic or generalised quaternion groups. If p and q are (not necessarily distinct) primes, then every subgroup of H with order pq is cyclic. No subgroup of a Frobenius complement is a Frobenius group. The Frobenius kernel K is necessarily a nilpotent group and, if the Frobenius complement H has even order, then K is, in fact, abelian.

2.4. Groups with Perfect Order Classes. This sub-section provides some background on groups with perfect order classes, beginning with a precise definition.

Given a finite group G , we define an equivalence relation on G by identifying elements which have the same order. The equivalence class of an element g in G with respect to this equivalence relation is called the **order class** of g , and is denoted by $g^{[G]}$. So we have

$$g^{[G]} = \{x \in G : |x| = |g|\}.$$

The cardinality of $g^{[G]}$ is denoted by $f_g(G)$. We say that G has **perfect order classes** if $f_g(G)$ is a divisor of the order of G , for all elements g in G .

For convenience, given a positive integer n , we also denote by $f_n(G)$ the cardinality of the order class of elements of order n . Note that $f_1(G) = 1$, and this is a divisor of the order of any group, so we usually do not mention the order class of 1.

It is clear that $g^G \subseteq g^{[G]}$, but in general the inclusion is proper. However, each order class is a disjoint union of one or more conjugacy classes.

2.7. Example. From Lemma 2.5 it follows that no generalised quaternion group Q_n has perfect order classes, since the number of elements of order 4 has an odd divisor.

We shall use some basic results on order classes and their cardinalities.

2.8. Lemma. [15] *If H is a normal Hall subgroup of a group G , then H contains the complete order class of each of its members.*

2.9. Lemma. [15] *Let A and B be groups whose orders are relatively prime. If a and b are positive integers such that a divides $|A|$ and b divides $|B|$, then*

$$f_{ab}(A \times B) = f_a(A)f_b(B).$$

The cyclic groups with perfect order classes have been completely characterised by Das [6]. This result follows from the elementary result that a positive integer $n > 1$ is a multiple of $\phi(n)$ precisely when it is an even $\{2, 3\}$ -number.

2.10. Proposition. [6, Proposition 2.1] *A cyclic group has perfect order classes if, and only if, it is trivial or its order is an even $\{2, 3\}$ -number.*

The next result is a key result in the study of groups with perfect order classes.

2.11. Proposition. [6, Proposition 2.2] *If G is a finite group, and g is an element of G , then $f_g(G)$ is divisible by $\phi(|g|)$.*

2.12. Corollary. [6, Corollary 2.3] *If G is a finite group with perfect order classes then, for each prime divisor p of the order of G , the order of G is divisible by $p - 1$. In particular, every non-trivial finite group with perfect order classes has even order.*

It is also a consequence of Corollary 2.12 that the smallest odd prime divisor of a group with perfect order classes that is not a 2-group is a Fermat prime.

We shall need the another result, originally due to Das [6], for which we provide a very short proof.

2.13. Proposition. [6, Proposition 2.4] *A finite 2-group has perfect order classes if, and only if, it is cyclic.*

Proof. Every group of even order has an odd number of involutions so, if a finite 2-group is to have perfect order classes, it must have a unique involution, which implies that it is either cyclic or a generalised quaternion group. But the number of elements of order 4 in a generalised quaternion group has an odd prime divisor, so a generalised quaternion group cannot have perfect order classes. \square

2.5. Supporting Computations. Some of our arguments require the use of computer calculations to verify the existence or non-existence of groups having particular properties. All computations were performed using the computer algebra systems GAP[10], MAGMA[13] and MAPLE[14] and, when feasible, were replicated in two or more of these systems.

3. THE STRUCTURE OF ORDER CLASSES IN FROBENIUS GROUPS

To prove our main theorem, we must first paint a clear picture of the structure of order classes in Frobenius groups.

3.1. Lemma. *Let $G = K \rtimes_f H$ be a Frobenius group with Frobenius kernel K and Frobenius complement H .*

- (1) *If $g \in K$, then $g^{[G]} = g^{[K]}$.*
- (2) *If $g \in G \setminus K$, then $g^{[G]}$ is the disjoint union*

$$g^{[G]} = \bigcup_{x \in K} (h^x)^{[H^x]},$$

where $h \in H \cap g^G$.

Proof. Let g be an element of K . Since K is a normal Hall subgroup of G , we have from Lemma 2.8 that $g^{[G]} \subseteq K$. Therefore,

$$g^{[K]} = K \cap g^{[G]} = g^{[G]}.$$

To prove the second assertion, fix an element g of $G \setminus K$. Since G is a Frobenius group, $G \setminus K$ is the disjoint union

$$G \setminus K = \bigcup_{x \in K} H^x \setminus 1.$$

Consequently, $H \cap g^G$ is non-empty, so we may choose $h \in H \cap g^G$. Then $|g| = |h| = |h^x|$, so $g^{[G]} = (h^x)^{[G]}$, for any $x \in K$. Keeping in mind that $1 \notin g^{[G]}$, we therefore have for each $x \in K$ that

$$g^{[G]} \cap (H^x \setminus 1) = (h^x)^{[G]} \cap (H^x \setminus 1) = (h^x)^{[G]} \cap H^x = (h^x)^{[H^x]}.$$

Since $K \cap g^{[G]} = \emptyset$, that is, $g^{[G]} \subseteq G \setminus K$, we have

$$g^{[G]} = g^{[G]} \cap \left(\bigcup_{x \in K} H^x \setminus 1 \right) = \bigcup_{x \in K} (h^x)^{[H^x]},$$

as claimed. \square

3.2. Corollary. *Let $G = K \rtimes_f H$ be a Frobenius group as in Lemma 3.1.*

- (1) *If $g \in K$, then $f_g(G) = f_g(K)$.*
- (2) *If $g \in G \setminus K$, then $f_g(G) = |K| f_h(H)$, where $h \in H \cap g^G$.*

With these preparations in hand, we are now ready to prove our main result, stated as Theorem A in the Introduction.

3.3. Theorem. *Let $G = K \rtimes_f H$ be a Frobenius group with Frobenius kernel K and Frobenius complement H . Then G has perfect order classes if, and only if, the following are true:*

- (a) *H has perfect order classes;*
- (b) *K is a homocyclic p -group for an odd prime number p ; and,*
- (c) *$|H| = p^r - 1$, where r is the rank of K .*

Proof. Suppose that H has perfect order classes and order equal to $p^r - 1$, where K is a homocyclic p -group of rank r . The set $G \setminus 1$ is the disjoint union of $K \setminus 1$ and the sets $H^x \setminus 1$, for $x \in K \setminus 1$. If d is the order of some non-trivial element of H , then $f_d(H)$ divides $|H|$ because H is assumed to have perfect order classes. Therefore, by Corollary 3.2, we have that $f_d(G) = |K| f_d(H)$ divides the order $|G| = |K| |H|$ of G . Now, K is homocyclic of rank r , so we can write

$$K \simeq C_{p^e}^r,$$

where p^e is the exponent of K . Then every member of K has order a divisor of p^e and, for each integer i with $1 \leq i \leq e$, we have from Lemma 2.6 and condition (c) that

$$f_{p^i}(G) = f_{p^i}(K) = p^{r(i-1)}(p^r - 1) = p^{r(i-1)} |H|,$$

which is clearly a divisor of the order of G . Therefore, any Frobenius group satisfying the conditions (a)-(c) has perfect order classes.

For the converse, suppose that G has perfect order classes. To show that H has perfect order classes, let m be the order of a non-trivial element of H . Since G has perfect order classes, it follows that $f_m(G)$ divides $|G|$, so we can write

$$|G| = sf_m(G),$$

for some positive integer s . Now H has exactly $|K|$ distinct conjugates H^g in G , for which we clearly have $f_m(H^g) = f_m(H)$. Moreover, these conjugates have pairwise trivial intersection, so that

$$f_m(G) = |K| f_m(H).$$

Therefore,

$$|K| |H| = |G| = sf_m(G) = s |K| f_m(H);$$

whence,

$$|H| = sf_m(H).$$

Therefore, $f_m(H)$ divides $|H|$ and, since m was the order of an arbitrary member of H , it follows that H has perfect order classes.

Since H has perfect order classes, and must be non-trivial, Corollary 2.12 tells us that the order of H is even. Therefore, the Frobenius kernel K is an abelian group. Suppose that the order of K is divisible by distinct (odd) primes p and q . Then K has elements of orders p , q and pq , every member of G with order equal to any of these numbers belongs to K and, since K is abelian, we have $f_{pq}(G) = f_p(G)f_q(G)$. Because K is abelian, we have also that $|x^G| = |H|$, for each non-trivial element x in K . Since each non-trivial order class in K is the disjoint union of conjugacy G -classes, each of length $|H|$, it follows that $|H|$ divides $f_x(G)$, for each $x \in K \setminus 1$. Therefore, $|H|$ divides both $f_p(G)$ and $f_q(G)$, so $|H|^2$ divides $f_{pq}(G)$. Since $f_{pq}(G)$ divides $|G|$, this is impossible. Therefore, K is a p -group, for some (odd) prime number p .

If K has rank r , then

$$f_p(G) = f_p(K) = |\Omega_1(K)| = p^r - 1.$$

Since this must be a divisor of $|G|$, and since $\gcd(p, p^r - 1) = 1$, it follows that $p^r - 1$ divides $|H|$. On the other hand, $\Omega_1(K)H$ is a Frobenius subgroup of G , since $\Omega_1(K)$ is characteristic in K , hence, normal in G . Therefore, the order of H divides $p^r - 1 = |\Omega_1(K)| - 1$, and we conclude that $|H| = p^r - 1$, as claimed.

Finally, suppose that K is not homocyclic, and write $K = I \times J$, where I is the homocyclic component of K with exponent equal to the

exponent p^e of K , and J has exponent p^d with $d < e$. Then $\Omega_d(K)$ is again a characteristic subgroup of K , and is proper since $d < e$, so $G/\Omega_d(K)$ is a Frobenius group with Frobenius complement isomorphic to H and Frobenius kernel isomorphic to $K/\Omega_d(K)$. But $K/\Omega_d(K)$ has rank s less than the rank r of K , and $|H| = p^r - 1$ must be a divisor of $p^s - 1$, which is impossible. Therefore, K must be homocyclic.

This completes the proof. \square

4. INSOLUBLE FROBENIUS GROUPS

In this section, we give a precise description of the Frobenius groups with perfect order classes that are insoluble.

Let us begin with an example to show that insoluble Frobenius groups with perfect order classes exist.

4.1. Example. The smallest insoluble Frobenius group G has order 14520, and has a Frobenius complement isomorphic to $\mathrm{SL}(2, 5)$, while the Frobenius kernel is the elementary abelian group C_{11}^2 . It is a perfect group ($G = [G, G]$), and has the order class profile

d	1	2	3	4	5	6	10	11
$f_d(G)$	1	121	2420	3630	2904	2420	2904	120

A simple numerical check shows that G has perfect order classes. Alternatively, one can check that $\mathrm{SL}(2, 5)$ has perfect order classes, and that $|\mathrm{SL}(2, 5)| = 120 = 11^2 - 1$, while the rank of the Frobenius kernel is 2, so the fact that G has perfect order classes follows from Theorem 3.3.

We are going to show that this example is essentially typical of insoluble Frobenius groups with perfect order classes. To this end, we shall need to solve a particular Diophantine equation involving prime powers. The purely number-theoretic proof of the following result we defer until the end of this section.

4.2. Proposition. *Let p and q be prime numbers, both greater than 5, and let r and m be positive integers with $r > 1$. Suppose that $q - 1$ is a divisor of 240, but that q does not divide 240. Then the equation*

$$p^r - 1 = 240q^m$$

has exactly two solutions:

$$(p, r, q, m) \in \{(11, 4, 61, 1), (41, 2, 7, 1)\}.$$

4.3. Lemma. *An insoluble Frobenius group with perfect order classes is perfect.*

Proof. Let G be a Frobenius group with perfect order classes and assume that G is insoluble but not perfect. Then by Theorem 3.3, the Frobenius kernel K of G is a homocyclic p -group of the form $C_{p^k}^r$, for some odd prime p , and positive integers k and r , a Frobenius complement H of G has perfect order classes, and we have

$$|H| = p^r - 1.$$

Since H is not abelian, it follows that the rank r of K is greater than 1 since, otherwise, H could not be isomorphic to a subgroup of $\text{Aut } K$.

Since H is insoluble, there is a subgroup L , of index at most 2 in H , such that $L = S \times M$, where S is isomorphic to $\text{SL}(2, 5)$, and M is a Z -group whose order is coprime to 30. Then we have

$$|H| = 120 |M| [H : L].$$

Suppose first that the subgroup M is trivial. Then $[H : L] = 2$, since H is not perfect, so $p^r = 241$, whence $p = 241$ and $r = 1$, contrary to the fact that $r > 1$. Therefore, it must be that M is non-trivial.

Let q be a prime divisor of the order of M . Since $\gcd(|M|, 30) = 1$, hence, $q > 5$. Then $L = S \times M$ has an element of order $5q$, and every element of H with order $5q$ belongs to L , since each member of $H \setminus L$ has even order if it is non-empty. Then

$$f_{5q}(H) = f_{5q}(L) = f_5(S)f_q(M) = 24(q - 1).$$

Since $q - 1$ is an even number, it follows that $f_{5q}(H)$ is divisible by 16. Thus, $[H : L] = 2$ because, otherwise, the order of H would not be divisible by 16.

Now suppose that s is another prime divisor of the order of M . Then $s > 5$ is also odd and, if M has an element of order qs , then every member of H whose order is equal to qs belongs to L , and so $f_{5qs}(H) = f_{5qs}(L)$ is a divisor of the order of H . But

$$f_{5qs}(H) = f_5(S)f_{qs}(M) = 24(q - 1)(s - 1),$$

which is divisible by 32 since q and s are odd. This is not possible, since $f_{5qs}(H)$ must divide the order of H . In particular, if M is cyclic, then M is a q -group, for some prime q with $q > 5$.

If M is not cyclic, then M is a Z -group and we can write

$$M = \langle x, y \mid x^\alpha, y^\beta, x^y = x^\gamma \rangle,$$

where α , β and γ are positive integers such that $\gcd(\alpha, \beta) = 1 = \gcd(\alpha, \gamma - 1)$. If either of α and β is not a prime power, then M has an element whose order is a product of two distinct primes, which we have just argued is not possible. Therefore, α and β are powers of different primes. We claim that the action of $\langle y \rangle$ on $\langle x \rangle$ is fixed-point

free. For, if not, then some non-trivial power of x commutes with some non-trivial power of y , and so their product has a power whose order is a product of two distinct primes. Therefore, M is a Frobenius group, with Frobenius kernel $\langle x \rangle$ and Frobenius complement $\langle y \rangle$. But M is a subgroup of the Frobenius complement H of G , which cannot contain a Frobenius subgroup. Therefore M must be a cyclic q -group. Let $|M| = q^m$, where m is a positive integer, so that

$$|H| = 240q^m.$$

We have the relation

$$p^r - 1 = 240q^m,$$

where $q - 1$ is a divisor of 240, and Proposition 4.2 implies that the only solutions are

$$(p, r, q, m) \in \{(11, 4, 61, 1), (41, 2, 7, 1)\}.$$

In the first case, $L \simeq \text{SL}(2, 5) \times C_{61}$ (and $K \simeq C_{11}^4$), whilst in the second case, we have $L \simeq \text{SL}(2, 5) \times C_7$ (and $K \simeq C_{41}^2$).

If $L \simeq \text{SL}(2, 5) \times C_7$, then L has an element of order 35 and every element of H with order equal to 35 belongs to L . Hence,

$$f_{35}(H) = f_{35}(L) = f_5(S)f_7(M) = 24 \cdot 6 = 2^4 \cdot 3^2,$$

which cannot divide $|H| = 2^4 \cdot 3 \cdot 5 \cdot 7$.

If $L \simeq \text{SL}(2, 5) \times C_{61}$, then L has an element of order 305 and, as in the previous case, we have

$$f_{305}(H) = f_5(S)f_{61}(M) = 24 \cdot 60 = 2^5 \cdot 3^2 \cdot 5$$

which, again, cannot divide $|H| = 2^4 \cdot 3 \cdot 5 \cdot 61$.

This final contradiction forces us to abandon our hypothesis that G is not perfect, completing the proof. \square

If $G = K \rtimes_f H$ is a Frobenius group with Frobenius kernel K and Frobenius complement H , then $[G, G] = K[H, H]$. Consequently, G is perfect if, and only if, H is perfect, and a well-known result of Zassenhaus states that a Frobenius complement is perfect if, and only if, it is isomorphic to $\text{SL}(2, 5)$. It follows that the only special linear groups that can occur as Frobenius complements are $\text{SL}(2, 5)$ and $\text{SL}(2, 3)$. In fact, $\text{SL}(2, 2) \simeq S_3$ is itself a Frobenius group, so it cannot be a Frobenius complement. And, for prime powers $q > 5$, the group $\text{SL}(2, q)$ is perfect, as is the simple group $\text{SL}(2, 4) \simeq A_5$.

Our next result provides a complete characterisation of the insoluble Frobenius groups with perfect order classes.

4.4. Theorem. *Let $G = K \rtimes_f H$ be an insoluble Frobenius group, with Frobenius kernel K and Frobenius complement H . Then G has perfect order classes if, and only if, $K \simeq C_{11^k}^2$, for some positive integer k , and $H \simeq \text{SL}(2, 5)$.*

Proof. According to Lemma 4.3, an insoluble Frobenius group with perfect order classes must be perfect. Let $G = K \rtimes_f H$ be a perfect Frobenius group with perfect order classes. Then $H \simeq \text{SL}(2, 5)$ and K is a homocyclic p -group of the form $C_{p^k}^r$, for an odd prime p and positive integers k and r . We also have $p^r - 1 = |H| = 120$, whence, $p^r = 121 = 11^2$, so $p = 11$ and $r = 2$. Therefore, $K \simeq C_{11^k}^2$, as claimed.

For the converse, assume that $H \simeq \text{SL}(2, 5)$ and that K is a homocyclic 11-group of rank 2, so that $K \simeq C_{11^k}^2$, for some positive integer k . Then $11^2 - 1 = |H|$, so G has perfect order classes, by Theorem 3.3. \square

4.5. Example. There are perfect Frobenius groups, $C_{19}^2 \rtimes_f \text{SL}(2, 5)$ of order 43320, $C_{29}^2 \rtimes_f \text{SL}(2, 5)$ of order 100920, and, $C_{59}^2 \rtimes_f \text{SL}(2, 5)$ of order 417720, but none of these has perfect order classes.

Let us now turn to our proof of Proposition 4.2.

Proof of Proposition 4.2. We are solving the equation

$$p^r - 1 = 240q^m,$$

where r is an integer greater than 1, and p and q are odd primes such that $q - 1$ divides 240, but q does not divide 240.

The prime numbers q such that $q - 1$ is a divisor of 240, but q does not divide 240 are

$$q \in \{7, 11, 13, 17, 31, 41, 61, 241\}.$$

That the stated solutions are valid is verified by direct numerical calculation:

$$11^4 = 1 + 240 \cdot 61^1, 41^2 = 1 + 240 \cdot 7^1.$$

We show that these are the only solutions. The method of solution is to factor the expression $p^r - 1$, and then to determine how the prime divisors of $240q^m$ may be allocated to each factor. To this end, we consider two primary cases, according to whether r is even or odd. In each case, there are a number of sub-cases,

First suppose that r is even, say $r = 2s$, where s is a positive integer. Then $p^r - 1$ is a difference of squares, so we have

$$2^4 \cdot 3 \cdot 5 \cdot q^m = p^{2s} - 1 = (p^s - 1)(p^s + 1).$$

Since the even integers $p^s - 1$ and $p^s + 1$ differ by 2, exactly one of them is divisible by 4, hence, it is divisible by 8, while the other is

twice an odd divisor of $240q^m$. These odd divisors are 3, 5, 15 as well as multiples of those three numbers by powers of q . Furthermore, $\gcd(p^s - 1, p^s + 1) = 2$, so that they have no odd prime divisors in common. In particular, each of 3, 5 and q divides exactly one of $p^s - 1$ and $p^s + 1$. Therefore, if q divides one of these two factors, then q^m divides that same factor.

Suppose that $p^s + 1$ is exactly divisible by 2, so that $p^s - 1$ is divisible by 8, and $p^s + 1$ is twice an odd divisor of $240q^m$. We consider the possibilities in turn; two of these lead to the expected solutions, while the others result in absurdities.

If $p^s + 1$ is divisible by q , then it is divisible by q^m . If $p^s + 1 = 2 \cdot q^m$, then $p^s - 1 = 2^3 \cdot 3 \cdot 5 = 120$, so $p^s = 121 = 11^2$, whence, $p = 11$ and $s = 2$. (Therefore, $r = 2s = 4$.) Then $2q^m = p^s + 1 = 122$, so that $q^m = 61$, yielding $q = 61$ and $m = 1$. This produces one of our expected solutions $11^4 = 1 + 240 \cdot 61$.

If $p^s + 1 = 2 \cdot 3 \cdot q^m$, then $p^s - 1 = 8 \cdot 5 = 40$, so that $p^s = 41$ and hence $p = 41$ and $s = 1$ (so $r = 2$). Then $p^s + 1 = 42$, so $q^m = 7$, giving $q = 7$ and $m = 1$. This produces our second expected solution $41^2 = 1 + 240 \cdot 7$.

The remaining sub-cases in which $p^s + 1 \in \{6, 10, 30, 10q^m, 30q^m\}$ all lead quickly either to arithmetic absurdities or to a contradiction of our stated hypotheses.

We must also consider all the possibilities for which $p^s - 1$ is equal to twice an odd divisor of $240q^m$. However, each of these also result in contradictions by elementary considerations.

This concludes our analysis for the case in which r is even, so assume now that r is odd. We examine a number of sub-cases, each leading to a contradiction.

We no longer have the convenient factorisation afforded by a difference of squares, but we can pull out a factor of $p - 1$ and write

$$2^4 \cdot 3 \cdot 5 \cdot q^m = p^r - 1 = (p - 1)(1 + p + \cdots + p^{r-1}).$$

Since r is odd, hence, $1 + p + \cdots + p^{r-1}$ is odd, so $16 = 2^4 \mid p - 1$. Therefore, $p - 1$ is a divisor of $240q^m$ divisible by 16. We cannot have either $p - 1 = 16 \cdot 3$ or $p - 1 = 16 \cdot 5$ since, in either case, p is not a prime.

Suppose that $p - 1 = 16$, so that $p = 17$, and

$$1 + p + \cdots + p^{r-1} = 15q^m.$$

Reducing this equation modulo 15, we obtain

$$1 + 2 + \cdots + 2^{r-1} \equiv 0 \pmod{15}.$$

Noting that $1 + 2 + 4 + 8 = 15$, and taking four terms at a time on the left hand side of this equation, it follows that r is divisible by 4, contrary to our assumption that r is odd.

If $p - 1 = 240$, then $p = 241$ and we have

$$241^r - 1 = 240q^m,$$

so that $241^r \equiv 1 \pmod{240q^m}$. In particular, $241^r \equiv 1 \pmod{q}$. For $q \in \{7, 11, 13, 17, 31, 41\}$, we have that 241 has even order modulo q , which implies that r is even. Thus, the only possibility for q is $q = 61$, since 241 has order equal to 5 modulo 61, which entails that r is a multiple of 5. Write $r = 5s$. If $s = 1$ so that $r = 5$, then we have

$$241^r - 1 = 241^5 - 1 = 812990017200 = 2^4 \cdot 3 \cdot 5^2 \cdot \mathbf{61} \cdot 11106421.$$

Hence, $(241^5 - 1)/240 = 5 \cdot 61 \cdot 11106421$, which is not a power of 61, so $s > 1$ and $r > 5$. But notice that $241^5 - 1$ is divisible by 61, as well as by 2, 3 and 5. Zsigmondy's theorem then implies that $241^r - 1$ has a prime divisor that does not divide $241^5 - 1$, since $r > 5$. This is impossible, so we conclude that $p \neq 241$.

If q^m divides $p - 1$ then, since 16 also divides $p - 1$, it follows that $1 + p + \cdots + p^{r-1}$ is a divisor of 15, which is clearly impossible for any prime $p > 5$.

We are left with the case in which q is a divisor of both $p - 1$ and $1 + p + \cdots + p^{r-1}$. We show that this case also cannot occur. Now, 3 must divide one of $p - 1$ and $p + 1$, so 3 is a divisor of $p^2 - 1$. By Euler's Theorem, we also have $p^4 \equiv 1 \pmod{5}$, so that 5 is a divisor of $p^4 - 1$. Since $p \equiv 1 \pmod{q}$ and q divides $1 + p + \cdots + p^{r-1}$, we have

$$0 \equiv 1 + p + \cdots + p^{r-1} \equiv r \pmod{q};$$

whence q divides r and, in particular, $r \geq q > 5$. Therefore, $p^r - 1$ has a primitive prime divisor s , for which $s \mid p^r - 1$ but $s \nmid p^i - 1$, for $1 \leq i < r$. But we have just seen that each of 2, 3, 5 and q occurs as a divisor of $p^i - 1$, for some $i < r$. Therefore, q cannot divide both $p - 1$ and $1 + p + \cdots + p^{r-1}$.

This shows that $p^r - 1 = 240 \cdot q^m$ has only the two solutions claimed. \square

5. SOLUBLE FROBENIUS GROUPS

We now turn to our study of the much richer class of soluble Frobenius groups. We are able to characterise those with perfect order classes in a number of important cases. Recall that the Sylow subgroups of a Frobenius complement are either cyclic or generalised quaternion. Therefore, a Frobenius complement is abelian if, and only if, it is cyclic.

We therefore consider first the Frobenius groups with cyclic Frobenius complement and provide a complete characterisation of those that have perfect order classes. Next, we consider those Frobenius groups whose Frobenius complement is nilpotent, and show that no new examples of groups with perfect order classes arise other than those with cyclic complement. Next, we turn to Frobenius groups whose complements have order divisible by just one odd prime; these are necessarily soluble, by Burnside's theorem, and we can assume that the Frobenius complement is not nilpotent, in light of previous results. Finally, since we have seen that a Frobenius complement in an insoluble Frobenius group with perfect order classes is isomorphic to the $\{2, 3, 5\}$ -group $\text{SL}(2, 5)$, we show that there are no soluble examples whose Frobenius complement is a $\{2, 3, 5\}$ -group.

5.1. Frobenius Groups with Cyclic Complement. We now examine Frobenius groups whose Frobenius complement is cyclic. (Recall that a Frobenius complement is abelian if, and only if, it is cyclic.)

We shall need the following simple lemma stating that the only pairs of consecutive $\{2, 3\}$ -numbers are $(1, 2)$, $(2, 3)$, $(3, 4)$ and $(8, 9)$. The reader should have no difficulty supplying a proof using Lemma 2.3 and Lemma 2.4.

5.1. Lemma. *If x, y, u, v are non-negative integers such that $2^x 3^y - 2^u 3^v = 1$, then*

$$(x, y, u, v) \in \{(1, 0, 0, 0), (0, 1, 1, 0), (2, 0, 0, 1), (0, 2, 3, 0)\}.$$

5.2. Corollary. *Let p be a positive integer such that both $p - 1$ and $p + 1$ are $\{2, 3\}$ -numbers. Then $p \in \{2, 3, 5, 7, 17\}$.*

Proof. Since $p + 1$ and $p - 1$ are both $\{2, 3\}$ -numbers, we can write $p + 1 = 2^a 3^b$ and $p - 1 = 2^c 3^d$, for suitable non-negative integers a, b, c and d . Then $2^a 3^b - 1 = p = 2^c 3^d + 1$, so we have

$$2^c 3^d = 2^a 3^b - 2.$$

First suppose that $a = 0$, so we have $2^c 3^d = 3^b - 2$, or $3^b = 2^c 3^d + 2$. If $c > 0$, this yields $3^b = 2(2^{c-1} 3^d + 1)$, which implies that 3^b is even, so it must be that $c = 0$ also. This yields $3^d = 3^b - 2$ or $3^b - 3^d = 2$, which clearly forces $b = 1$ and $d = 0$. Thus, we have $(a, b, c, d) = (0, 1, 0, 0)$, from which we obtain $p = 2$.

Now assume that $a > 0$. Then

$$2^c 3^d = 2(2^{a-1} 3^b - 1).$$

If $c = 0$, this again implies that 3^d is even, so c must also be positive. Then $2^{c-1}3^d = 2^{a-1}3^b - 1$, or

$$2^{a-1}3^b - 2^{c-1}3^d = 1.$$

From Lemma 5.1 we have

$$(a, b, c, d) \in \{(1, 1, 2, 0), (1, 2, 4, 0), (2, 0, 1, 0), (3, 0, 1, 1)\}.$$

These solutions yield (using either $p = 1 + 2^c 3^d$ or $p = 2^a 3^b - 1$) $p \in \{5, 17, 3, 7\}$. This completes the proof. \square

We now are ready to analyse the Frobenius groups with perfect order classes and cyclic Frobenius complement.

5.3. Proposition. *Let $G = K \rtimes_f H$ be a Frobenius group with cyclic Frobenius complement H . Then G has perfect order classes if, and only if, one of the following is true:*

- (1) *K is a cyclic p -group, for some Pierpont prime $p = 1 + |H| > 2$, and H is a $\{2, 3\}$ -group of even order; or*
- (2) *G is isomorphic to one of the following groups:*
 - (a) $(C_{3^k} \times C_{3^k}) \rtimes_f C_8$;
 - (b) $(C_{5^k} \times C_{5^k}) \rtimes_f C_{24}$;
 - (c) $(C_{7^k} \times C_{7^k}) \rtimes_f C_{48}$;
 - (d) $(C_{17^k} \times C_{17^k}) \rtimes_f C_{288}$,*for some positive integer k .*

Note that in the cyclic case, the possibility that the Frobenius complement is a 2-group is not excluded.

Proof. Suppose that G has perfect order classes. Then H has perfect order classes, and there is an odd prime p such that K is a homocyclic group of the form $C_{p^k}^r$, for some positive integers k and r , and we have $|H| = p^r - 1$. Since H is cyclic with perfect order classes, it follows from Proposition 2.10 that H is a $\{2, 3\}$ -group of even order, so we may write

$$|H| = 2^\alpha 3^\beta,$$

where α is a positive integer and β is a non-negative integer.

If $r = 1$, then K is cyclic and $p = 1 + |H|$ is a Pierpont prime, so assume that $r > 1$.

If r is odd, then $r \geq 3$. If H is a 2-group, whence $\beta = 0$, then $p^r - 1$ is a power of 2. If $p^r - 1 = 2$, then $p^r = 3$, so $p = 3$ and $r = 1$, contrary to the assumption that $r > 1$. Therefore, $p^r - 1$ must be a proper power of 2, say, $p^r - 1 = 2^m$, where the integer m is greater than 1. According to Lemma 2.2, this implies that $p = 3$, $r = 2$ and $m = 3$, this time contradicting the assumption that r is odd. Thus, in this

case, H cannot be a 2-group, and 3 is a divisor of $|H| = p^r - 1$. Since p and the order of H must be relatively prime, we see that $p > 3$, and hence, $3 \mid p^2 - 1$. Certainly, 2 divides $p - 1$ (and so also $p^2 - 1$). Since $r \geq 3$, Zsigmondy's theorem implies that $p^r - 1$ has a prime divisor q such that q does not divide $p^i - 1$, for any integer i with $1 \leq i < r$. In particular, q does not divide $p^2 - 1$, so $q > 3$. This is in contradiction to the hypothesis that $p^r - 1 = |H|$ is a $\{2, 3\}$ -number. We conclude, therefore, that r is odd only if $r = 1$.

Suppose now that $r = 2s$ is even, where s is a positive integer. If H is a 2-group, then $|H| = 2^\alpha$, and we have

$$2^\alpha = p^{2s} - 1 = (p^s - 1)(p^s + 1) = (p^2 - 1)(1 + p^2 + p^4 + \cdots + p^{2(s-1)}).$$

This implies that $p^2 - 1$ and $p^s - 1$ are powers of 2. Since $p^2 - 1 \neq 2$, for any odd prime p , therefore, $p^2 - 1$ is a proper power of 2, so $p^2 - 1 = 2^3$, and $p = 3$, thanks to Lemma 2.2. Since $p^s - 1 = 3^s - 1$ is also a power of 2, we must have $s = 1$ and $r = 2$. Then, from $|H| = p^r - 1$, we see that $|H| = 8$. Therefore,

$$G \simeq C_{3^k}^2 \rtimes_f C_8,$$

giving the first of our four cases with a Frobenius kernel of rank 2.

If H is not a 2-group, then $\beta > 0$ and $p > 3$. Then, from

$$2^\alpha 3^\beta = p^{2s} - 1 = (p^s - 1)(p^s + 1),$$

we see that both $p^s - 1$ and $p^s + 1$ are $\{2, 3\}$ -numbers, so from Corollary 5.2, we have $p^s \in \{2, 3, 5, 7, 17\}$, whence $s = 1$ and $p \in \{5, 7, 17\}$.

For $p = 5$, we have $H \simeq C_{24}$, and so

$$G \simeq C_{5^k}^2 \rtimes_f C_{24}.$$

For $p = 7$, we have $H \simeq C_{48}$, giving

$$G \simeq C_{7^k}^2 \rtimes_f C_{48}.$$

Finally, for $p = 17$ we have $H \simeq C_{288}$, so that

$$G \simeq C_{17^k}^2 \rtimes_f C_{288}.$$

For the converse, we consider each case in turn, beginning with the case of a cyclic Frobenius kernel. Suppose then that K is a cyclic p -group, where $p = 1 + |H|$ is a Pierpont prime and $p > 2$. Then $|H| \geq 2$ is a $\{2, 3\}$ -number and, since $|H| + 1$ is an odd prime, it follows that $|H|$ is even. Therefore, H has perfect order classes by Proposition 2.10. Since K is cyclic, it has rank equal to 1, so the relation $|H| = p - 1$ is satisfied. Therefore, by Theorem 3.3, G has perfect order classes.

For the rank 2 cases, we have $G \simeq C_{p^k}^2 \rtimes_f C_n$, where

$$(p, n) \in \{(3, 8), (5, 24), (7, 48), (17, 288)\},$$

and k is an arbitrary positive integer. In each case C_n is a $\{2, 3\}$ -group of even order and, as such, has perfect order classes, according to Proposition 2.10. The Frobenius kernels $C_{p^k}^2$ are homocyclic, and it is easily verified numerically that, in each case, we have $n = p^2 - 1$. Now invoking Theorem 3.3 suffices to complete the proof. \square

The following is a slight generalisation of [24, Theorem 2.1], which asserts that a dihedral group has perfect order class if, and only if, its degree is a power of 3.

5.4. Corollary. *A Frobenius group whose Frobenius complement is a group of order 2 has perfect order classes if, and only if, it is a dihedral group of degree equal to a power of 3.*

5.5. Example. Let $G = K \rtimes_f H$ be a Frobenius group with Frobenius complement H a cyclic group of order 4. Then G has perfect order class if, and only if, K is a non-trivial cyclic 5-group.

5.6. Example. If p is a prime number, then $\text{Hol}(C_p)$ has perfect order classes if, and only if, p is an odd Pierpont prime.

5.2. Frobenius Groups with Nilpotent Complement. In this section we aim to show that a nilpotent Frobenius complement in a Frobenius group with perfect order classes is necessarily cyclic. Recall that a Sylow 2-subgroup of a Frobenius complement is either cyclic or generalised quaternion, and that the odd order Sylow subgroups of any Frobenius complement are cyclic. Because a nilpotent Frobenius complement is the direct product of its Sylow subgroups, it can be non-abelian only if it is a direct product of a generalised quaternion group and a cyclic group of odd order.

Let us begin with an example demonstrating the existence of Frobenius complements with perfect order classes that are nilpotent and non-abelian.

5.7. Example. Let $H \simeq Q_4 \times C_{5^k}$, for a positive integer k , where Q_4 is a generalised quaternion group of order 16. Then H is a non-abelian nilpotent Frobenius complement with perfect order classes. To see this, we tabulate the number of elements of each possible order m in H , where i is any integer with $1 \leq i \leq k$.

m	1	2	4	8	5^i	$2 \cdot 5^i$	$4 \cdot 5^i$	$8 \cdot 5^i$
$f_m(H)$	1	1	10	4	$4 \cdot 5^i$	$4 \cdot 5^{i-1}$	$2 \cdot 5^i$	$16 \cdot 5^i$

In each case, we see that $f_m(H)$ is a divisor of the order $|H| = 16 \cdot 5^k$.

Our next result shows that this example is one of just two families of non-abelian nilpotent Frobenius complements that have perfect order classes.

5.8. Theorem. *Let H be a non-abelian nilpotent Frobenius complement. Then H has perfect order classes if, and only if, H is isomorphic either to $Q_3 \times C_{3^k}$ or to $Q_4 \times C_{5^k}$, for some positive integer k .*

Proof. Assume that H has perfect order classes; hence, the order of H is even. Since H is nilpotent, it is the direct product of its Sylow subgroups. And, since H is a Frobenius complement, its Sylow subgroups are cyclic or generalised quaternion. Because H is non-abelian, its Sylow 2-subgroup Q must be a generalised quaternion group, say $Q \simeq Q_n$, where $n \geq 3$. Therefore,

$$H = Q \times T,$$

where T is a cyclic group of odd order (the direct product of the odd order Sylow subgroups of H). Since generalised quaternion groups do not have perfect order classes, it follows that the subgroup T must be non-trivial. Therefore

$$|H| = 2^n |T| > 2^n.$$

Every element of order 4 in H belongs to Q , so we have

$$f_4(H) = f_4(Q) = 2^{n-1} + 2 = 2(2^{n-2} + 1).$$

Since $n \geq 3$, hence, $2^{n-2} + 1$ is an odd number greater than 1. And, since $f_4(H)$ divides the order of H , it follows that $2^{n-2} + 1$ is a divisor of the order of T .

We claim that $2^{n-2} + 1$ is square-free. To see this, suppose that there is a prime q such that q^2 divides $2^{n-2} + 1$, so that q^2 is also a divisor of $|T|$. Let q^m be the largest power of q dividing the order of T , where $m \geq 2$ is an integer. Since T is a cyclic group, therefore, T has $\phi(q^m) = q^{m-1}(q-1)$ elements of order q^m , and we have

$$f_{4q^m}(H) = f_4(Q)f_{q^m}(T) = 2(2^{n-2} + 1)q^{m-1}(q-1).$$

Since q^2 divides $2^{n-2} + 1$, it follows that q^{m+1} divides $f_{4q^m}(H)$, which implies that q^{m+1} divides $|T|$, in contradiction to the choice of m . Consequently, $2^{n-2} + 1$ must be square-free.

Next, let p , q and r be different prime divisors of the order of T . Then the order of H is divisible by

$$f_{2^{n-1}pqr}(H) = f_{2^{n-1}}(Q_n)f_{pqr}(T) = 2^{n-2}(p-1)(q-1)(r-1).$$

Since p , q and r are odd, we see that $f_{2^{n-1}pqr}(H)$ is divisible by 2^{n+1} , which cannot divide $|H| = 2^n |T|$. Now, if $2^{n-2} + 1$ had three distinct prime divisors, then so too would $|T|$, so it must be that $2^{n-2} + 1$ is a prime or a product of two distinct primes.

Suppose that

$$2^{n-2} + 1 = pq,$$

where p and q are (odd) primes with $p < q$. Let a and b be positive integers such that $|T| = p^a q^b$, so that

$$|H| = 2^n p^a q^b.$$

Then

$$f_{2^{n-1}pq}(H) = f_{2^{n-1}}(Q)f_{pq}(T) = 2^{n-2}(p-1)(q-1),$$

and so we must have

$$2^{n-2}(p-1)(q-1) \mid 2^n p^a q^b.$$

From this we see that the even numbers $p-1$ and $q-1$ must be exactly divisible by 2 (that is, that 4 divides neither). Because $p-1 < p < q$, it follows that $p-1$ can only be equal to 2, so $p = 3$. We therefore have

$$2^{n-1}(q-1) \mid 2^n 3^a q^b,$$

so $q-1 = 2 \cdot 3^i$, for some non-negative integer i , and $q = 1 + 2 \cdot 3^i$ is a Pierpont prime. Now,

$$2^{n-2} + 1 = 3q = 3(1 + 2 \cdot 3^i) = 3 + 2 \cdot 3^{i+1};$$

whence, $2 \cdot 3^{i+1} = 2^{n-2} - 2 = 2(2^{n-3} - 1)$, so $3^{i+1} = 2^{n-3} - 1$, or

$$2^{n-3} - 3^{i+1} = 1.$$

By Lemma 2.3, we have

$$(n, i) \in \{(4, -1), (5, 0)\}.$$

Since $i = -1$ is absurd, it must be that $i = 0$. But then we are led to the further absurdity $q = 3 > p = 3$. We are forced to conclude, therefore, that $2^{n-2} + 1$ is a prime.

Let $p = 2^{n-2} + 1$. Now, T has $\phi(p) = p - 1 = 2^{n-2}$ elements of order p , and

$$f_{2^{n-1}p}(H) = f_{2^{n-1}}(Q)f_p(T) = 2^{n-2} \cdot 2^{n-2} = 2^{2n-4}.$$

Since this quantity must divide $|H| = 2^n |T|$, therefore, $2n - 4 \leq n$, so $n \leq 4$. But also, n is greater than or equal to 3 so, in fact, $n = 3$ or $n = 4$. If $n = 3$, then $p = 2^{n-2} + 1 = 3$; and, if $n = 4$, then $p = 2^{n-2} + 1 = 5$.

We know that the order of T cannot be divisible by three primes, but we have yet to rule out the possibility that $|T|$ has a prime divisor other than p . In the case that $p = 3$ and $n = 3$, this follows from [15] since, in this case, H is a Hamiltonian group with perfect order classes, so T must be a non-trivial cyclic 3-group. So suppose that $p = 5$ and $n = 4$, and also that T has a prime divisor q other than 5. Then

$$f_{40q}(H) = f_8(Q)f_{5q}(T) = 4 \cdot 4 \cdot (q-1),$$

which, since q is odd, is divisible by 32. It follows that $f_{40q}(H)$ cannot divide $|H| = 16|T|$ in this case, so T must be a cyclic 5-group.

Thus, in summary, H is isomorphic either to $Q_3 \times C_{3^k}$ or to $Q_4 \times C_{5^k}$, for some positive integer k .

Conversely, we saw in Example 5.7 that $Q_4 \times C_{5^k}$ has perfect order classes. That $Q_3 \times C_{3^k}$ has perfect order classes follows from [15], or by the method of Example 5.7. \square

Now that we have a classification of the non-abelian nilpotent complements with perfect order classes, we shall see next that none of them “fits” into a Frobenius group that has perfect order classes.

5.9. Theorem. *A nilpotent Frobenius complement in a Frobenius group with perfect order classes is cyclic.*

Proof. Let $G = K \rtimes_f H$ be a Frobenius group with perfect order classes, and assume, for an eventual contradiction, that the Frobenius complement H is a non-abelian nilpotent group. By Theorem 3.3, H has perfect order classes, K is a homocyclic group of the form $C_{p^k}^r$, for an odd prime p and positive integers k and r , and we have $|H| = p^r - 1$. By Theorem 5.8, H is isomorphic to $Q_n \times C_{q^t}$, for some positive integer t , where $(n, q) \in \{(3, 3), (4, 5)\}$. Thus,

$$p^r - 1 = 2^n q^t.$$

Note that r must be greater than 1 since H is not abelian.

Suppose first that r is odd, so that $r \geq 3$. If $q = 3$, then $p > 3$, and so 3 divides $p^2 - 1$, and 2 is of course also a divisor of $p^2 - 1$. By Zsigmondy’s theorem, $p^r - 1$ has a primitive prime divisor other than 2 and 3, in contradiction to the equality $p^r - 1 = 2^4 \cdot 3^t$. It must be then that $q = 5$ and $n = 4$, in consequence of which we have either that $p = 3$ or that $p > 5$. If $p = 3$, then

$$2^4 \cdot 5^t = 3^r - 1 = 2(1 + 3 + 3^2 + \cdots + 3^{r-1}).$$

Since r is odd, therefore, $1 + 3 + \cdots + 3^{r-1}$ is odd, which implies the absurdity $2 = 16$. Therefore, it must be that p is greater than 5.

Now if $r = 3$, then

$$2^4 \cdot 5^t = p^3 - 1 = (p - 1)(p^2 + p + 1)$$

and, since $p^2 + p + 1$ is odd, it follows that 16 divides $p - 1$, and $p^2 + p + 1 = 5^i$, for some integer i satisfying $0 \leq i \leq t$. It is clear that i must be positive, however, so we have

$$p(p + 1) = 5^i - 1 = 4(1 + 5 + \cdots + 5^{i-1}).$$

Thus, 4 divides $p+1$. But this means that 4 is a divisor of both $p-1$ and $p+1$, which is impossible. Thus, r cannot be equal to 3 and we must have $r \geq 5$. But 5 divides $p^4 - 1$, by Euler's theorem and 2 certainly is a divisor of $p-1$, so Zsigmondy's theorem implies that $p^r - 1$ has a prime divisor other than 2 and 5, contradicting $p^r - 1 = 2^4 \cdot 5^t$. We conclude, therefore, that r must be even.

Write $r = 2s$, where s is a positive integer. Then

$$2^n q^t = p^{2s} - 1 = (p^s - 1)(p^s + 1),$$

from which it follows that both $p^s - 1$ and $p^s + 1$ are $\{2, q\}$ -numbers.

If $q = 3$ and $n = 3$ then, by Corollary 5.2, we have $p^s \in \{2, 3, 5, 7, 17\}$, whence, $s = 1$ and $r = 2$, and in fact $p \in \{5, 7, 17\}$, since p is odd and $q = 3$ forces $p > 3$. If $p = 7$, then $p^2 - 1 = 48 = 16 \cdot 3$, which cannot be equal to $8 \cdot 3^t$, for any t . If $p = 17$, then $p^2 - 1 = 288 = 2^5 \cdot 3^2 \neq 8 \cdot 3^t$, regardless of the value of t . It must therefore be that $p = 5$, so that

$$8 \cdot 3^t = 5^2 - 1 = 24,$$

and hence, $t = 1$. Then $H \simeq Q_3 \times C_3$ and $K \simeq C_{5^k} \times C_{5^k}$, for some positive integer k . Presuming the existence of such a group, we infer the existence of its Frobenius subgroup $\Omega_1(K)H$ of order $25 \cdot 24 = 600$ and with Frobenius complement $H \simeq Q_3 \times C_3$. (Note that we do not assert that $\Omega_1(K)H$ has perfect order classes.) However, consulting the library of Frobenius groups in MAPLE, we see that none of the Frobenius groups of order 600 has a Frobenius complement isomorphic to $Q_3 \times C_3$. (They are isomorphic to $C_3 \rtimes C_8$, C_{24} or $SL(2, 3)$.) We conclude, therefore, that no such group G exists.

Now suppose that $q = 5$ and $n = 4$, so $H \simeq Q_4 \times C_{5^t}$, and we have $p^r - 1 = 16 \cdot 5^t$. Suppose that $p > 3$, so that 3 divides $p^2 - 1$. Since r is even, hence, $p^2 - 1$ divides $p^r - 1$, so 3 also divides $p^r - 1$. Therefore, $p^r - 1$ cannot be a $\{2, 5\}$ -number, so we must have $p = 3$, and hence,

$$2^4 \cdot 5^t = 3^r - 1.$$

Since $3^r \equiv 1 \pmod{5}$, it follows that r is a multiple of 4, so we can write $r = 4s$, for a positive integer s . Suppose that $s > 1$, so that $r > 4$. Since 2 and 5 both divide $3^4 - 1 = 80$, Zsigmondy's theorem implies that $3^r - 1$ has a prime divisor other than 2 and 5, in contradiction to $3^r - 1 = 2^4 \cdot 5^t$. Thus, r must be equal to 4, and we have

$$2^4 \cdot 5^t = 3^4 - 1 = 80 = 2^4 \cdot 5,$$

whence $t = 1$ and $H \simeq Q_4 \times C_5$, and also $K \simeq C_{3^k}^4$, for some positive integer k . Now $\Omega_1(K)H$ is a Frobenius subgroup of G of order 6480, since $\Omega_1(K) \simeq C_3^4$. However, from the library of Frobenius groups in MAPLE, we see that no Frobenius group of order 6480 has Frobenius

complement isomorphic to $Q_4 \times C_5$. It follows, therefore, that G does not exist.

Having thus exhausted every possibility, the proof is complete. \square

Proof of Theorem C. It follows from Theorem 5.9 that the Frobenius groups with a nilpotent Frobenius complement and perfect order classes are exactly those described in Proposition 5.3. \square

We now have a complete picture of the Frobenius groups with perfect order classes whose Frobenius complement is nilpotent. In the following sections, we begin to study soluble Frobenius groups with perfect order classes whose Frobenius complement is not nilpotent.

5.3. Frobenius Groups with Non-Nilpotent Biprimary Complement. For Frobenius groups whose Frobenius complement is not nilpotent, we begin by considering cases for which the order of the Frobenius complement has just one odd prime divisor. By Burnside's theorem, any such group is soluble.

We begin by characterising Frobenius groups with perfect order classes, assuming that the Frobenius complement is a non-nilpotent $\{2, 3\}$ -group or a non-nilpotent $\{2, 5\}$ -group.

5.10. Theorem. *Let $G = K \rtimes_f H$ be a Frobenius group and assume that the Frobenius complement H is a non-nilpotent $\{2, 3\}$ -group. Then G has perfect order classes if, and only if, G has one of the following forms:*

- (1) $G \simeq C_{5^k}^2 \rtimes_f \text{SL}(2, 3)$;
- (2) $G \simeq C_{5^k}^2 \rtimes_f \langle a, b \mid a^3, b^8, a^b = a^{-1} \rangle$;
- (3) $G \simeq C_{7^k}^2 \rtimes_f \langle a, b \mid a^3, b^{16}, a^b = a^{-1} \rangle$; or
- (4) $G \simeq C_{17^k}^2 \rtimes_f \langle a, b \mid a^9, b^{32}, a^b = a^{-1} \rangle$,

for some positive integer k .

Proof. Suppose that G has perfect order classes. Then H also has perfect order classes, $K \simeq C_{p^k}^r$ is a homocyclic p -group for some odd prime p , and positive integers k and r , and $|H| = p^r - 1$. Since H is non-abelian, it follows that the rank r of K is greater than 1. It must also be that the order of H is divisible by 3 since, otherwise, H would be a 2-group, and we have assumed that H is not nilpotent.

First assume that r is odd, so that $r \geq 3$. Since $p > 3$, it follows that $p^2 - 1$ is divisible by 3 and, of course, $p - 1$ is divisible by 2. By Zsigmondy's theorem, $p^r - 1$ has a prime divisor q such that q does not divide $p^i - 1$, for $1 \leq i < r$. In particular, $q \neq 2$ and $q \neq 3$. But this contradicts $p^r - 1 = |H|$ being a $\{2, 3\}$ -number. Therefore, r must be

even and we can write $r = 2s$, for some positive integer s . Then we have

$$|H| = p^{2s} - 1 = (p^s - 1)(p^s + 1);$$

whence, both $p^s - 1$ and $p^s + 1$ are $\{2, 3\}$ -numbers. Therefore, by Corollary 5.2, we have $p^s \in \{2, 3, 5, 7, 17\}$, whence $s = 1$ (and $r = 2$), and since p is greater than 3 we have $p \in \{5, 7, 17\}$. If $p = 5$, then $|H| = p^2 - 1 = 24$; if $p = 7$, then $|H| = 48$; and, if $p = 17$, then $|H| = 288$.

Consulting the database of small groups in MAPLE, we find that the non-nilpotent groups of order 24 with perfect order classes are $\text{SL}(2, 3)$, $\langle a, b \mid a^3, b^8, a^b = a^{-1} \rangle \simeq C_3 \rtimes C_8$, and $\langle a, b, c \mid a^4, b^2, c^3, c^a = c^2, [a, b], [b, c] \rangle \simeq C_2 \times (C_3 \rtimes C_4)$. The last of these three contains a subgroup of the form $C_2 \times C_2$, so it cannot be a Frobenius complement. The first two do occur as Frobenius complements.

The non-nilpotent groups of order 48 with perfect order classes are

$$\langle a, b \mid a^3, b^{16}, a^b = a^{-1} \rangle \simeq C_3 \rtimes C_{16},$$

$$\langle a, b, c \mid a^8, b^2, c^3, [a, b], c^a = c^2, [b, c] \rangle \simeq (C_3 \rtimes C_8) \times C_2,$$

$$\langle a, b, c \mid a^8, b^2, c^3, c^a = c^2, [a, b], [b, c], aba^3b, a^{-1}ba^2ba^{-1} \rangle \simeq (C_3 \rtimes C_8) \rtimes C_2,$$

as well as $C_2 \times \text{SL}(2, 3)$. The last three of these contain a subgroup of the form $C_2 \times C_2$, so cannot be Frobenius complements. Therefore, the only choice for a group of order 48 among them is the first, with structure $C_3 \rtimes C_{16}$.

Similarly, among the 26 non-nilpotent groups of order 288 with perfect order classes only one group,

$$\langle a, b \mid a^9, b^{32}, a^b = a^{-1} \rangle \simeq C_9 \rtimes C_{32}$$

can be a Frobenius complement. For instance, there are two groups

$$G_{188} = \langle a, b, c \mid a^3, b^3, [a, b], c^{32}, a^c = ab^{-1}, b^c = (ba)^{-1} \rangle \simeq (C_3 \times C_3) \rtimes C_{32}$$

and

$$G_{373} = \langle a, b, c \mid a^3, b^3, [a, b], c^{32}, [a, c], b^c = ab^{-1} \rangle \simeq (C_3 \times C_3) \rtimes C_{32}$$

which cannot be Frobenius complements because they have subgroups of the form $C_3 \times C_3$. The remaining 23 groups all have a subgroup of the form $C_2 \times C_2$.

For the converse, assume that H has perfect order classes, and that $K \simeq C_{p^k}$, where $(p, |H|) \in \{(5, 24), (7, 48), (17, 288)\}$. (Of course, H must be among those groups identified above.) Since H has perfect order classes, we need only check that the number of elements whose

order is a divisor of $|K| = p^{2k}$ divides the order of G . For $1 \leq i \leq k$, we have

$$f_{p^i}(G) = f_{p^i}(K) = p^{2(i-1)}(p-1).$$

If $p = 5$, then $p - 1 = 4$, which divides 24. If $p = 7$, then $p - 1 = 6$, which divides 48. If $p = 17$, then $p - 1 = 16$, which divides 288. Finally, $|H| = p^2 - 1$ in each case. Therefore, G has perfect order classes. \square

5.11. Theorem. *Let G be a Frobenius group whose Frobenius complement H is a non-nilpotent $\{2, 5\}$ -group. Then G has perfect order classes if, and only if, the Frobenius kernel is a homocyclic 3-group of rank 4, and $H \simeq \langle a, b \mid a^5, b^{16}, a^b = a^4 \rangle$.*

Proof. One checks directly that the group $\langle a, b \mid a^5, b^{16}, a^b = a^{-1} \rangle$ is a non-nilpotent group of order $80 = 3^4 - 1$ with perfect order classes. Therefore, if H is isomorphic to this group, and if the Frobenius kernel of G is a homocyclic 3-group of rank 4, then G has perfect order classes by Theorem 3.3.

To prove the converse, assume that G has perfect order classes. Then by Theorem 3.3, the Frobenius kernel K of G is a homocyclic p -group of the form $C_{p^k}^r$, for some odd prime p and suitable positive integers k and r , H has perfect order classes, and we have $|H| = p^r - 1$. Since H is assumed not nilpotent, both 2 and 5 divide the order of H , and the rank r of K must be greater than 1. Since the orders of H and K are relatively prime, therefore, p cannot be equal to 5.

Suppose first that r is odd, so in fact, r must be greater than or equal to 3. If $r = 3$, then $p^3 - 1 = |H|$ is a $\{2, 5\}$ -number divisible by 10, so $p^3 \equiv 1 \pmod{5}$. Together with $p^4 \equiv 1 \pmod{5}$, from Euler's theorem, this implies that $p \equiv 1 \pmod{5}$, that is, that 5 divides $p - 1$. But 2 also divides $p - 1$, so Zsigmondy's theorem implies that there is a prime divisor q of $p^3 - 1$ that is not a divisor of $p - 1$ (or of $p^2 - 1$). In particular, q cannot be either of 2 and 5, in contradiction to the assumption that $p^3 - 1 = |H|$ is a $\{2, 5\}$ -number. Therefore, r cannot be equal to 3, and we must have $r \geq 5$. But, from the divisibility of $p^4 - 1$ by 5, another application of Zsigmondy's theorem shows that r cannot be odd.

So, r is even. If p is not equal to 3, then p must be greater than 5 and 3 divides $p^2 - 1$ which, in turn, divides $p^r - 1$. Since $|H| = p^r - 1$, this contradicts the assumption that H is a $\{2, 5\}$ -group. Consequently, it must be that p is equal to 3.

From $3^r \equiv 1 \pmod{5}$ it follows that r is a multiple of 4, and so we can write $r = 4s$, for some positive integer s . If $s = 1$ so that $r = 4$, then we have

$$|H| = 3^4 - 1 = 80 = 2^4 \cdot 5.$$

If $s > 1$, so that $r \geq 8$, then we again apply Zsigmondy's theorem to obtain a prime divisor q of $p^r - 1$ such that q does not divide $p^i - 1$, for $1 \leq i < r$. In particular, q does not divide $3^4 - 1 = 80$, so q cannot be either of 2 and 5 and this contradicts that assumed fact that $3^r - 1$ must be a $\{2, 5\}$ -number. Therefore, the only possibility is that the rank r of K is equal to 4, giving $K \simeq C_{3^k}^4$, for a suitable positive integer k , and also that H is a group of order 80.

Now, consulting the library of small groups in MAPLE we find that, among the groups of order 80, just three have perfect order classes, and one of those, $Q_4 \times C_5$, is nilpotent, leaving the two metacyclic groups

$$A = \langle a, b \mid a^5, b^{16}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^5, b^{16}, a^b = a^4 \rangle.$$

Both A and B are indeed Frobenius complements, but only B occurs as a Frobenius complement in a Frobenius group with Frobenius kernel isomorphic to $C_{3^k}^4$. To see this, suppose that there is a Frobenius group $F \simeq C_{3^k}^4 \rtimes_f A$. Then F has a Frobenius subgroup isomorphic to $C_3^4 \rtimes_f A$ of order 6480. Checking the MAPLE library of Frobenius groups, we find that there are three Frobenius groups of order 6480 of which two have a Frobenius complement of order 80. One of those complements is the cyclic group of that order, while the other is our group B . Therefore, we must have $H \simeq B$. \square

5.12. *Remark.* The proof of Theorem 5.11 furnishes us with an example

$$A = \langle a, b \mid a^5, b^{16}, a^b = a^2 \rangle,$$

of a soluble but non-nilpotent Frobenius complement with perfect order classes that is not a Frobenius complement in a Frobenius group with perfect order classes.

Having dealt with Frobenius complements that are either $\{2, 3\}$ -groups or $\{2, 5\}$ -groups, let us now show that there are no other $\{2, q\}$ -groups to be considered.

5.13. **Lemma.** *Let q be an odd prime, and let x, y, u and v be non-negative integers such that*

$$2^x q^y - 2^u q^v = 1.$$

(1) *If $q = 2^p - 1$ is a Mersenne prime, where p is a prime, then*

$$(x, y, u, v) \in \{(1, 0, 0, 0), (p, 0, 0, 1)\}.$$

(2) *If $q = 2^{2^t} + 1$ is a Fermat prime, for a non-negative integer t , then*

$$(x, y, u, v) \in \{(1, 0, 0, 0), (0, 1, 1, 0), (0, 1, 2^t, 0)\}.$$

(3) *Otherwise,*

$$(x, y, u, v) \in \{(1, 0, 0, 0), (0, 1, 1, 0), (0, 2, 3, 0)\}.$$

In all cases, if $(x, y, u, v) \in \{(0, 1, 1, 0), (0, 2, 3, 0)\}$, then $q = 3$.

Proof. Since $2^x q^y$ and $2^u q^v$ are consecutive integers, one must be even and the other odd, so exactly one of x and u is equal to zero.

Suppose first that $x = 0$ and $u > 0$, so we have

$$1 = q^y - 2^u q^v.$$

If $y \leq v$, then $1 = q^y(1 - 2^u q^{v-y})$, so $y = 0$ and $1 = 1 - 2^u q^v$. But this implies the impossibility $2^u q^v = 0$, so we must have $v < y$. Then

$$1 = q^v(q^{y-v} - 2^u);$$

whence, $v = 0$ and

$$q^y = 2^u + 1.$$

If $u = 0$, then $q^y = 2$, contradicting the assumption that q is odd. If $u = 1$, then $q^y = 3$, so $q = 3$ and $y = 1$. In this case, we have

$$(x, y, u, v) = (0, 1, 1, 0), \quad q = 3.$$

Assume now that $u > 1$. If $y = 0$, we arrive at the impossible $2^u + 1 = 1$. If $y = 1$, then $q = 2^u + 1$ is a Fermat prime, so there is a non-negative integer t for which $u = 2^t$. Then

$$(x, y, u, v) = (0, 1, 2^t, 0), \quad q = 2^{2^t} + 1.$$

Otherwise, $y > 1$ and, since $u > 1$ also, the only solution is $q = 3$, $u = 3$ and $y = 2$, so we have

$$(x, y, u, v) = (0, 2, 3, 0), \quad q = 3.$$

Now suppose that $u = 0$ and $x > 0$, so we have

$$1 = 2^x q^y - q^v.$$

If $v \leq y$, then

$$1 = q^v(2^x q^{y-v} - 1);$$

whence, $v = 0$ and $1 = 2^x q^y - 1$, or $2^x q^y = 2$. Then $x = 1$ and $y = 0$ and we have

$$(x, y, u, v) = (1, 0, 0, 0).$$

If $y < v$, then

$$1 = q^y(2^x - q^{v-y});$$

whence, $y = 0$ and

$$q^v = 2^x - 1.$$

It is clear that x cannot be zero and, if $x = 1$, then $q^v = 1$, so $v = 0$. Thus,

$$(x, y, u, v) = (1, 0, 0, 0)$$

in this case. Otherwise, we have $x > 1$, in which case it is clear that $v > 0$. If $v = 1$, then $q = 2^x - 1$ is a Mersenne prime, so x is a prime p and we have

$$(x, y, u, v) = (p, 0, 0, 1), \quad q = 2^p - 1.$$

Otherwise, we have $v > 1$ and, since $x > 1$, there is no solution in this case. \square

5.14. Proposition. *Let q be an odd prime, and let n be a positive integer such that $n^2 - 1$ is divisible by 2 and q , but by no other primes. Then one of the following is true.*

- (1) *If $q = 2^p - 1$ is a Mersenne prime, where p is a prime, then $n = 2^{p+1} - 1$; and, if n is a prime power, then $(q, n) = (3, 7)$.*
- (2) *If $q = 2^{2^t} + 1$ is a Fermat prime, then $n = 2^{1+2^t} + 1$; and, if n is a prime power, then $(q, n) \in \{(3, 5), (5, 9)\}$.*
- (3) *Otherwise, $(q, n) \in \{(3, 5), (3, 17)\}$.*

In particular, if n is a prime power, then $q \in \{3, 5\}$.

Proof. Note that $n > 1$ since 0 is divisible by every prime, and n must be odd, since $n^2 - 1$ is even. Since $n^2 - 1 = (n + 1)(n - 1)$ and both $n + 1$ and $n - 1$ are even positive integers, we can write

$$n + 1 = 2^a q^b \text{ and } n - 1 = 2^c q^d,$$

where a and c are positive integers, and b and d are non-negative integers. (In fact, one of a and c must be equal to 1, and one of b and d must be zero.) Since a and c are positive, we have

$$2 = (n + 1) - (n - 1) = 2^a q^b - 2^c q^d,$$

and hence,

$$2^{a-1} q^b - 2^{c-1} q^d = 1,$$

with all of $a - 1$, b , $c - 1$ and d non-negative. We now consider the cases from Lemma 5.13.

If $(a - 1, b, c - 1, d) = (1, 0, 0, 0)$, then $n = 3$. But $3^2 - 1 = 8$ is not divisible by any odd prime, so this case does not occur. If $(a - 1, b, c - 1, d) = (0, 1, 1, 0)$, then $q = 3$ and $n = 5$. If $(a - 1, b, c - 1, d) = (0, 2, 3, 0)$, then $n = 2q^2 - 1 = 17$, from which we obtain $q = 3$.

If $q = 2^p - 1$ is a Mersenne prime, and $(a - 1, b, c - 1, d) = (p, 0, 0, 1)$, then $n = 2^{p+1} - 1$. If n is prime, then $p + 1$ is prime, so $p = 2$, $q = 3$ and $n = 7$. In this case, n cannot be a proper power.

If $q = 2^{2^t} + 1$ is a Fermat prime and $(a - 1, b, c - 1, d) = (0, 1, 2^t, 0)$, then $n = 2^{1+2^t} + 1$. If n is a prime, it is a Fermat prime, so $1 + 2^t$ is a power of 2, say, $1 + 2^t = 2^i$. Then $t = 0$ and $i = 1$, so $q = 3$ and $n = 5$. If n is a proper prime power, then $n - 1 = 2^{1+2^t}$, so $n = 9$ and $2^{1+2^t} = 8$, which implies that $3 = 1 + 2^t$, so $t = 1$ and $q = 5$. \square

5.15. Theorem. *Let $G = K \rtimes_f H$ be a Frobenius group with perfect order classes, and suppose that H is a non-nilpotent $\{2, q\}$ -group, where q is an odd prime. Then $q = 3$ or $q = 5$ and the rank of K is either 2 or 4.*

Proof. Since H is a non-nilpotent $\{2, q\}$ -group, we can write $|H| = 2^a q^b$, where a and b are positive integers. By Theorem 3.3, H has perfect order classes, $K \simeq C_{p^k}^r$, for some odd prime p and positive integers k and r , and we have

$$p^r - 1 = |H| = 2^a q^b.$$

Since H has perfect order classes, it follows that $q - 1$ divides $|H| = 2^a q^b$ and, since $\gcd(q, q - 1) = 1$, hence, $q - 1$ is a divisor of 2^a , which means that it is a power of 2, say $q - 1 = 2^\tau$, where τ is a positive integer. Then $q = 2^\tau + 1$ is a Fermat prime, so $\tau = 2^t$ is a power of 2, for some non-negative integer t .

By Euler's theorem, we have $p^{q-1} \equiv 1 \pmod{q}$, so q divides $p^{q-1} - 1$. Since q also divides $p^r - 1$, it follows that q is a divisor of

$$\gcd(p^r - 1, p^{q-1} - 1) = p^d - 1,$$

where $d = \gcd(r, q - 1)$. If $d < r$ then, since 2 divides $p - 1$ and q divides $p^d - 1$, Zsigmondy's theorem implies that $p^r - 1$ has a prime divisor other than 2 and q , contrary to $p^r - 1 = 2^a q^b$. Therefore, $r = d = \gcd(r, q - 1)$, so r divides

$$q - 1 = (2^{2^t} + 1) - 1 = 2^{2^t}.$$

Thus, r is a power of 2. Since H is not abelian, hence, $r > 1$ and, in particular, r is even. Applying Proposition 5.14, we see that $q \in \{3, 5\}$ and, since $r > 1$ divides $q - 1$, it follows that $r = 2$ or $r = 4$. \square

5.4. Frobenius Groups with Soluble $\{2, 3, 5\}$ -Complement That Do Not Exist. We saw in Theorem 4.4 that a Frobenius complement of an insoluble Frobenius group with perfect order classes is isomorphic to the $\{2, 3, 5\}$ -group $\text{SL}(2, 5)$. This leads us to ask whether there are soluble Frobenius groups with perfect order classes whose Frobenius complement is a $\{2, 3, 5\}$ -group. We conclude by showing that there are no such soluble examples.

We begin, as usual, with some number theoretic preliminaries.

5.16. Lemma. *Suppose that x, y, u and v are non-negative integers such that*

$$2^x \cdot 3^y - 2^u \cdot 5^v = 1.$$

Then

$$(x, y, u, v) \in \{(0, 4, 4, 1), (1, 1, 0, 1), (1, 0, 0, 0), (0, 1, 1, 0), (0, 2, 3, 0)\}.$$

Proof. First assume that $x \leq u$. Then

$$1 = 2^x \cdot 3^y - 2^u \cdot 5^v = 2^x(3^y - 2^{u-x} \cdot 5^v),$$

so we must have $2^x = 1 = 3^y - 2^{u-x} \cdot 5^v$, whence $x = 0$ and

$$2^u \cdot 5^v = 3^y - 1.$$

If $v = 0$ then this becomes $2^u = 3^y - 1$, from which we conclude, with the help of Lemma 2.4, either that $y = u = 1$ or that $y = 2$ and $u = 3$. In this way we obtain the solutions

$$(x, y, u, v) \in \{(0, 1, 1, 0), (0, 2, 3, 0)\}.$$

Otherwise, v is positive, so $3^y \equiv 1 \pmod{5}$, from which it follows that y is a multiple of 4. Write $y = 4t$, where t is a positive integer. Then

$$2^u \cdot 5^v = 3^y - 1 = (3^{2t})^2 - 1 = (3^{2t} - 1)(3^{2t} + 1).$$

Since $\gcd(3^{2t} - 1, 3^{2t} + 1) = 2$, hence, $3^{2t} - 1$ and $3^{2t} + 1$ have no common odd prime divisors. Thus, one of $3^{2t} - 1$ and $3^{2t} + 1$ must be a power of 2, while the other is exactly divisible by 2. Now, $3^{2t} + 1 = 2^{u-1}$ implies that $2t = 1$, which is not possible. Hence, $3^{2t} - 1 = 2^{u-1}$, from which we get $2t = 2$ and $u - 1 = 3$. Therefore, $t = 1$ and $y = 4$, while $u = 4$. This gives, from $3^y - 2^u \cdot 5^v = 1$ that

$$1 = 3^y - 2^u \cdot 5^v = 3^4 - 2^4 \cdot 5^v = 81 - 16 \cdot 5^v;$$

whence, $16 \cdot 5^v = 81 - 1 = 80$, so $v = 1$. Thus, we obtain the solution

$$(x, y, u, v) = (0, 4, 4, 1).$$

Now suppose that $u < x$. Then

$$1 = 2^x \cdot 3^y - 2^u \cdot 5^v = 2^u(2^{x-u} \cdot 3^y - 5^v),$$

so that $u = 0$ and $1 = 2^x \cdot 3^y - 5^v$, or

$$2^x \cdot 3^y = 5^v + 1.$$

If $v = 0$ then $x = 1$ and $y = 0$, so we get the solution

$$(x, y, u, v) = (1, 0, 0, 0).$$

If $v = 1$, then $2^x \cdot 3^y = 6$, so we must have $x = y = 1$, and this case produces the solution

$$(x, y, u, v) = (1, 1, 0, 1).$$

If $v > 1$, then by Zsigmondy's theorem, $5^v + 1$ has a prime divisor q such that q does not divide 6, Therefore, $5^v + 1$ cannot have the form $2^x \cdot 3^y$ for any non-negative integers x and y , so there are no further solutions for $v > 1$. \square

5.17. Lemma. *Suppose that x, y, u and v are non-negative integers such that*

$$2^u \cdot 5^v - 2^x \cdot 3^y = 1.$$

Then

$$(x, y, u, v) \in \{(1, 0, 0, 0), (0, 0, 1, 0), (0, 1, 2, 0), (0, 2, 1, 1), (3, 1, 0, 2)\}.$$

Proof. Assume first that $x \leq u$, so that

$$1 = 2^u \cdot 5^v - 2^x \cdot 3^y = 2^x(2^{u-x} \cdot 5^v - 3^y),$$

whence, $x = 0$ and $2^u \cdot 5^v - 3^y = 1$, or

$$2^u \cdot 5^v = 3^y + 1.$$

If $y = 0$, then $2^u \cdot 5^v = 2$, so $u = 1$ and $v = 0$, producing the solution

$$(x, y, u, v) = (0, 0, 1, 0).$$

If $y = 1$, then $2^u \cdot 5^v = 4$, which implies that $u = 2$ and $v = 0$, resulting in the solution

$$(x, y, u, v) = (0, 1, 2, 0).$$

For $y = 2$, we have $2^u \cdot 5^v = 10$, from which we see that $u = 1 = v$, and this yields the solution

$$(x, y, u, v) = (0, 2, 1, 1).$$

If $y > 2$, then $3^y + 1$ has a prime divisor q such that q does not divide $3^2 + 1 = 10$, thanks to Zsigmondy's theorem. Since $q \notin \{2, 5\}$, it follows that there are no further solutions for $y > 2$.

Now suppose that $u < x$. Then

$$1 = 2^u \cdot 5^v - 2^x \cdot 3^y = 2^u(5^v - 2^{x-u} \cdot 3^y),$$

whence, $u = 0$ and $1 = 5^v - 2^x \cdot 3^y$, or

$$2^x \cdot 3^y = 5^v + 1.$$

If $y = 0$, then this becomes $2^x = 5^v + 1$ which has only the trivial solution $x = 1$ and $v = 0$, so we get

$$(x, y, u, v) = (1, 0, 0, 0).$$

Otherwise, y is positive, and we have $5^v \equiv 1 \pmod{3}$, so v is even and we can write $v = 2s$, for a suitable positive integer s . Then

$$2^x \cdot 3^y = 5^{2s} - 1 = (5^s - 1)(5^s + 1).$$

Since $\gcd(5^s - 1, 5^s + 1) = 2$, one of $5^s - 1$ and $5^s + 1$ is a power of 2 while the other is exactly divisible by 2. If $5^s + 1$ is a power of 2, then $s = 0 = v$, which results in the impossibility $2^{u-1} \cdot 3^y = 5^0 - 1 = 0$. Thus, it must be that $5^s - 1$ is a power of 2, which can only be 4, so $s = 1$ and $v = 2$. Then $2^x \cdot 3^y = 5^2 - 1 = 24$, so $x = 3$ and $y = 1$. We therefore have our final solution

$$(x, y, u, v) = (3, 1, 0, 2)$$

and, with that, the proof is complete. \square

We now use the preceding two lemmas to prove the following result.

5.18. Lemma. *If n is a positive integer such that $n^2 - 1$ is divisible by 2, 3 and 5, and by no other prime, then*

$$n \in \{11, 19, 31, 49, 161\}.$$

Proof. It is clear that n must be odd and greater than unity.

Since $n^2 - 1 = (n - 1)(n + 1)$ and $\gcd(n - 1, n + 1) = 2$, it follows either that one of $n - 1$ and $n + 1$ is a power of 2, while the other is divisible by 2, 3 and 5 and by no other prime and is furthermore indivisible by 4, or, that one of $n - 1$ and $n + 1$ is a $\{2, 3\}$ -number while the other is a $\{2, 5\}$ -number, neither being a prime power. Thus, there are four cases to consider.

Case 1. Suppose that

$$n - 1 = 2^u \quad \text{and} \quad n + 1 = 2 \cdot 3^s \cdot 5^t,$$

where s, t and u are positive integers. Then

$$2 = (n + 1) - (n - 1) = 2 \cdot 3^s \cdot 5^t - 2^u = 2(3^s \cdot 5^t - 2^{u-1});$$

whence, $3^s \cdot 5^t - 2^{u-1} = 1$, or

$$3^s \cdot 5^t = 2^{u-1} + 1.$$

Because s and t are positive, we have $2^{u-1} + 1 \geq 15$, so that $u - 1 \geq 4$. By Zsigmondy's theorem, $2^{u-1} + 1$ has a prime divisor q such that q does not divide $2^i + 1$, for $1 \leq i < u - 1$. In particular, q divides neither $3 = 2^1 + 1$ nor $5 = 2^2 + 1$. This means that $2^{u-1} + 1$ cannot be equal to $3^s \cdot 5^t$, for any choice of s and t , so this case does not occur.

Case 2. Suppose that

$$n + 1 = 2^u \quad \text{and} \quad n - 1 = 2 \cdot 3^s \cdot 5^t,$$

for some positive integers s, t and u . Then $n + 1 \geq 32$, so $u \geq 5$, and we have

$$2 = (n + 1) - (n - 1) = 2^u - 2 \cdot 3^s \cdot 5^t = 2(2^{u-1} - 3^s \cdot 5^t);$$

whence, $2^{u-1} - 3^s \cdot 5^t = 1$, or

$$3^s \cdot 5^t = 2^{u-1} - 1.$$

If $u = 5$, then $2^{u-1} - 1 = 15$, so $s = t = 1$ and $n = 31$.

If $u = 6$, then $2^{u-1} - 1 = 31$ which is not a $\{3, 5\}$ -number. There are no solutions s and t for $u = 7$ either since, in that case, $2^{u-1} - 1 = 63 = 3^2 \cdot 7$. If $u > 7$, then $u - 1 > 6$, so Zsigmondy's theorem tells us that $2^{u-1} - 1$ cannot be a $\{3, 5\}$ -number. Thus, this case yields only the solution $n = 31$.

Case 3. Suppose that

$$n - 1 = 2^a \cdot 3^b \text{ and } n + 1 = 2^c \cdot 5^d,$$

for some positive integers a, b, c and d . Then

$$2 = (n + 1) - (n - 1) = 2^c \cdot 5^d - 2^a \cdot 3^b;$$

whence,

$$2^{c-1} \cdot 5^d - 2^{a-1} \cdot 3^b = 1.$$

Then by Lemma 5.17 we have

$$(a, b, c, d) \in \{(2, 0, 1, 0), (1, 0, 2, 0), (1, 1, 3, 0), (1, 2, 2, 1), (4, 1, 1, 2)\}.$$

However, because a, b, c and d are all positive we have, in fact, that

$$(a, b, c, d) \in \{(1, 2, 2, 1), (4, 1, 1, 2)\}.$$

Using either $n = 2^a \cdot 3^b + 1$ or $n = 2^c \cdot 5^d - 1$, we obtain

$$n \in \{19, 49\}.$$

Case 4. Suppose that a, b, c and d are positive integers for which

$$n + 1 = 2^a \cdot 3^b \text{ and } n - 1 = 2^c \cdot 5^d.$$

Then

$$2 = (n + 1) - (n - 1) = 2^a \cdot 3^b - 2^c \cdot 5^d = 2(2^{a-1} \cdot 3^b - 2^{c-1} \cdot 5^d);$$

from which we obtain

$$2^{a-1} \cdot 3^b - 2^{c-1} \cdot 5^d = 1.$$

From Lemma 5.16 we have that

$$(a, b, c, d) \in \{(1, 4, 5, 1), (2, 1, 1, 1)\},$$

taking account of the fact that $a, b, c, d > 0$. Again using either $n = 2^a \cdot 3^b + 1$ or $n = 2^c \cdot 5^d - 1$, we obtain

$$n \in \{161, 11\}.$$

This completes the proof. □

Our next lemma will help us prove the three that follow by constraining the possibilities to be considered.

5.19. Lemma. *Let p be a prime number greater than 3. Let H be a \mathbb{Z} -subgroup of $\mathrm{GL}(2, p)$ of order $|H| = p^2 - 1$, and with presentation $\langle x, y \mid x^\alpha, y^\beta, x^y = x^\gamma \rangle$, where $\gcd(\alpha, \beta(\gamma - 1)) = 1$ and $\gamma^\beta \equiv 1 \pmod{\alpha}$. Then the index $[H : H \cap \mathrm{SL}(2, p)]$ is an even divisor of $\gcd(p - 1, \beta)$.*

Proof. For notational simplicity, let $G = \mathrm{GL}(2, p)$ and $S = \mathrm{SL}(2, p)$. Notice that $|S| = p|H|$ so the Sylow 2-subgroups of H and S have the same order. Since a Sylow 2-subgroup of H is cyclic, while a Sylow 2-subgroup of S is generalised quaternion, it follows that the index $[H : H \cap S]$ must be even.

Since

$$H/(H \cap S) \simeq HS/S \leq G/S \simeq C_{p-1},$$

hence, $[H : H \cap S]$ is a divisor of $p - 1$. Furthermore, since $H/(H \cap S)$ is cyclic, it follows that the derived subgroup $[H, H] = \langle x \rangle$ is contained in $H \cap S$, so $[H : H \cap S]$ divides β . Therefore, $[H : H \cap S]$ is an even divisor of $\gcd(p - 1, \beta)$, as advertised. \square

Next, we need to establish that certain potential Frobenius complements do not occur together with specific Frobenius kernels. To this end, we shall need the following result of Dickson listing the subgroups of two-dimensional special linear groups. (We state here only the part we shall need, for subgroups with order prime to the characteristic; see [23, Theorem 6.17, p. 404] for the full statement.)

5.20. Lemma (Dickson's Theorem). [23, Theorem 6.17] *Let q be a well-defined power of an odd prime p , and let H be a subgroup of $\mathrm{SL}(2, q)$ such that p does not divide the order of H . Then one of the following occurs:*

- (1) H is cyclic;
- (2) H is isomorphic to the dicyclic group $L_n = \langle x, y \mid x^n = y^2, x^y = x^{-1} \rangle$, for some n ;
- (3) $H \simeq \mathrm{SL}(2, 3)$;
- (4) $H \simeq \mathrm{SL}(2, 5)$;
- (5) $H \simeq \widehat{S}_4$, a central extension of the symmetric group S_4 .

We note that the dicyclic group L_n of order $4n$ in the statement of Dickson's theorem is a \mathbb{Z} -group only for odd positive integers n ; for even n , the Sylow 2-subgroup of L_n is a generalised quaternion group. Thus, apart from the cyclic subgroups, the dicyclic groups L_n for odd n are the only \mathbb{Z} -subgroups of $\mathrm{SL}(2, q)$.

5.21. Lemma. *No subgroup of $\text{GL}(2, 11)$ is isomorphic to either of the groups*

$$A = \langle a, b \mid a^5, b^{24}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{15}, b^8, a^b = a^2 \rangle.$$

Proof. Let $G = \text{GL}(2, 11)$ and $S = \text{SL}(2, 11)$. Note that both A and B are Z-groups of order $120 = 11^2 - 1$. Suppose that H is a subgroup of $\text{GL}(2, 11)$ isomorphic to either of A and B . Then Lemma 5.19 implies that the index $[H : H \cap S]$ is an even divisor of $\gcd(10, b)$, where $b = 24$ if $H \simeq A$ and $b = 8$ if $H \simeq B$.

If $H \simeq A$, then $[H : H \cap S]$ is an even divisor of $\gcd(10, 24) = 2$, hence $[H : H \cap S] = 2$, and we have

$$H \cap S = \langle a, b^2 \rangle \simeq \langle x, y \mid u^5, v^{12}, u^v = u^4 \rangle.$$

Since $H \cap S$ is not cyclic, it can only be isomorphic to the dicyclic group $L_{15} = \langle x, y \mid x^{15} = y^2, x^y = y^{-1} \rangle$. But, $L_{15}^{\text{ab}} \simeq C_4$, while $(H \cap S)^{\text{ab}} \simeq C_{12}$, so $H \simeq A$ cannot be a subgroup of $\text{GL}(2, 11)$.

If $H \simeq B$, then $[H : H \cap S]$ is an even divisor of $\gcd(10, 8) = 2$, so $[H : H \cap S] = 2$, and we have

$$H \cap S = \langle a, b^2 \rangle \simeq \langle s, t \mid s^{15}, t^4, s^t = s^4 \rangle.$$

As before, the only subgroup of S that $H \cap S$ might be isomorphic to is L_{15} . But it is easy to see that the centre of $H \cap S$ has order 6, while the centre of any dicyclic group has order 2. \square

5.22. Lemma. *No subgroup of $\text{GL}(2, 19)$ is isomorphic to either of the groups*

$$A = \langle a, b \mid a^5, b^{72}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{45}, b^8, a^b = a^8 \rangle.$$

Proof. Let $G = \text{GL}(2, 19)$ and $S = \text{SL}(2, 19)$, and suppose that H is a subgroup of G isomorphic to either of A and B . Then H is a Z-group of order $360 = 19^2 - 1$. Again, Lemma 5.19 implies that the index $[H : H \cap S]$ is an even divisor of $\gcd(18, b)$, where $b = 72$ if $H \simeq A$ and $b = 8$ if $H \simeq B$.

If indeed $H \simeq B$, then $[H : H \cap S] = 2$, and we have

$$H \cap S = \langle a, b^2 \rangle \simeq \langle s, t \mid s^{45}, t^4, s^t = s^{19} \rangle.$$

Since $H \cap S$ is not cyclic, and has order 180, therefore, the only subgroup of S that might be isomorphic to H is the dicyclic group L_{45} . But the centre of $H \cap S$ has order 18, while $|Z(L_{45})| = 2$. Thus, B cannot be isomorphic to a subgroup of $\text{GL}(2, 19)$.

If $H \simeq A$, we cannot apply the same reasoning, since $H \cap S$ contains a subgroup isomorphic to L_5 , which does, in fact, occur in $S = \text{SL}(2, 19)$ (in two conjugacy classes). However, in this case, we can check directly

using GAP or MAGMA that A is not isomorphic to a subgroup of $\text{GL}(2, 19)$. \square

5.23. Lemma. *No subgroup of $\text{GL}(2, 31)$ is isomorphic to either of the groups*

$$A = \langle a, b \mid a^5, b^{192}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{15}, b^{64}, a^b = a^2 \rangle.$$

Proof. Let $G = \text{GL}(2, 31)$ and $S = \text{SL}(2, 31)$, and suppose that H is a subgroup of G isomorphic to one of A and B . Then H is a Z-group and $|H| = 960 = 31^2 - 1$, so we can apply Lemma 5.19 as in the previous two lemmas.

If $H \simeq A$, then $[H : H \cap S] \in \{2, 6\}$, so $|H \cap S| \in \{480, 160\}$. Thus, $H \cap S$ must be isomorphic either to L_{120} or to L_{40} , but neither of these dicyclic groups is a Z-group. If $H \simeq B$, then $[H : H \cap S] = 2$ and we would again have $H \cap S \simeq L_{120}$, which is not a Z-group. \square

With these preparations in hand, we are now ready to analyse the Frobenius groups with a soluble Frobenius complement that is a $\{2, 3, 5\}$ -group.

5.24. Theorem. *There is no Frobenius group with perfect order classes in which a Frobenius complement is a soluble, non-nilpotent $\{2, 3, 5\}$ -group that is neither a $\{2, 3\}$ -group nor a $\{2, 5\}$ -group.*

Proof. Let G be a Frobenius group with perfect order classes, and assume that a Frobenius complement H for G is soluble, not nilpotent and is a $\{2, 3, 5\}$ -group, but neither a $\{2, 3\}$ -group nor a $\{2, 5\}$ -group. According to Theorem 3.3, there is an odd prime number p and positive integers k and r such that, if K is the Frobenius kernel for G , then $K \simeq C_{p^k}^r$. Furthermore, H has perfect order classes, and we have

$$|H| = p^r - 1.$$

Now, r cannot be equal to 1, since H is not abelian. Since $p > 5$, it follows that 2 and 3 divide $p^2 - 1$, and 5 divides $p^4 - 1$, by Euler's theorem. Therefore, if $r > 4$, then Zsigmondy's theorem implies that $p^r - 1$ has a prime divisor q other than 2, 3 and 5, contrary to $p^r - 1 = |H|$. So, r must satisfy $2 \leq r \leq 4$.

Suppose that $r = 3$, so that 5 is a divisor of $p^3 - 1$. Then 5 divides

$$(p^4 - 1) - (p^3 - 1) = p^4 - p^3 = p^3(p - 1).$$

Since $\gcd(p, 5) = 1$, we conclude that 5 divides $p - 1$. Another application of Zsigmondy's theorem implies that $p^3 - 1$ has a prime divisor other than 2, 3 and 5, so r cannot be equal to 3.

Suppose that $r = 2$, so that $p^2 - 1 = |H|$ is divisible by 2, 3 and 5 and by no other prime. Then $p \in \{11, 19, 31\}$, by Lemma 5.18, since 49 and 161 are composite. We must then have (respectively),

$$|H| \in \{120, 360, 960\}.$$

If $r = 4$, then $p^4 - 1 = (p^2)^2 - 1 = |H|$ is a $\{2, 3, 5\}$ -number, so $p^2 = 49$, and $p = 7$, since 49 is the only square of a prime from Lemma 5.18. In this case, we have

$$|H| = 2400.$$

In summary, we have shown so far that G has one of the structures:

- $G \simeq C_{11^k}^2 \rtimes_f H$, where $|H| = 120$;
- $G \simeq C_{19^k}^2 \rtimes_f H$, where $|H| = 360$;
- $G \simeq C_{31^k}^2 \rtimes_f H$, where $|H| = 960$; or,
- $G \simeq C_{7^k}^4 \rtimes_f H$, where $|H| = 2400$,

for some positive integer k , where H is a Frobenius complement of the indicated order with perfect order classes. We now proceed to argue that no such groups can exist.

Using MAPLE we identify the groups of each of these orders with perfect order classes and their structure. Using our lemmas above, as well as computations in GAP and MAGMA, we determine whether each possibility occurs.

There are five soluble groups of order 120 with perfect order classes that are not nilpotent. Three of them contain subgroups isomorphic to $C_2 \times C_2$ and therefore cannot be Frobenius complements. The remaining two are

$$A = \langle a, b \mid a^5, b^{24}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{15}, b^8, a^b = a^2 \rangle.$$

(Since 2 has order 4 in both $(\mathbb{Z}/5\mathbb{Z})^\times$ and $(\mathbb{Z}/15\mathbb{Z})^\times$, it follows that both A and B are Frobenius complements.) By Lemma 5.21, $\text{GL}(2, 11)$ contains no subgroup isomorphic either to A or to B , so neither can be a Frobenius complement in a Frobenius group with Frobenius kernel isomorphic to $C_{11^k}^2$, for any positive integer k .

There are four soluble, non-nilpotent groups of order 360 with perfect order classes. Two of these contain a copy of $C_2 \times C_2$ and so cannot be Frobenius complements. The other two are

$$A = \langle a, b \mid a^5, b^{72}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{45}, b^8, a^b = a^8 \rangle.$$

Again, $\text{GL}(2, 19)$ contains no subgroup isomorphic to either of A and B , by Lemma 5.22, so neither can be a Frobenius complement in a Frobenius group with Frobenius kernel of the form $C_{19^k}^2$. (Again, however, both A and B are Frobenius complements.)

The soluble, non-nilpotent groups of order 960 with perfect order classes that do not contain a copy of $C_2 \times C_2$ are

$$A = \langle a, b \mid a^5, b^{192}, a^b = a^2 \rangle \text{ and } B = \langle a, b \mid a^{15}, b^{64}, a^b = a^2 \rangle.$$

Once again, this time by Lemma 5.23, neither appears as a subgroup of $GL(2, 31)$ up to isomorphism, so neither can be a Frobenius complement in a Frobenius group with Frobenius kernel isomorphic to $C_{31^k} \times C_{31^k}$.

Among the soluble, non-nilpotent groups of order 2400 with perfect order classes, all but two contain a copy of either $C_2 \times C_2$ or $C_5 \times C_5$, leaving the groups

$$A = \langle a, b \mid a^{25}, b^{96}, a^b = a^7 \rangle \text{ and } B = \langle a, b \mid a^{75}, b^{32}, a^b = a^{32} \rangle.$$

However, a MAGMA computation shows that $GL(4, 7)$ does not contain a subgroup isomorphic to either A or B , so this case does not occur either.

□

6. CONCLUDING REMARKS

Our results all describe Frobenius groups $G = K \rtimes_f H$ with perfect order classes, which entails that the Frobenius complement H itself has perfect order classes. However, along the way we have seen that there are examples of a Frobenius complement H , such as $Q \times C_3$, with perfect order classes that do not, however, occur as complements of Frobenius groups with perfect order classes. This suggests an independent characterisation of Frobenius complements with perfect order classes. While our results do fully classify the nilpotent Frobenius complements with perfect order classes, a complete description for those that are either insoluble or soluble but not nilpotent remains open.

6.1. Problem. Describe all Frobenius complements with perfect order classes, whether or not they occur as complements of a Frobenius group with perfect order classes.

REFERENCES

- [1] OEIS, *A005109 Class 1— (or Pierpont) primes: primes of the form $2^t \cdot 3^u + 1$* , <https://oeis.org/A005109>. ↑3
- [2] A. S. Bang, *Taltheoretiske Undersøgelser*, Tidsskrift for Matematik **4** (1886), 70–80. ↑5
- [3] ———, *Taltheoretiske Undersøgelser*, Tidsskrift for Matematik **4** (1886), 130–137. ↑5
- [4] Geo. D. Birkhoff and H. S. Vandiver, *On the integral divisors of $a^n - b^n$* , Annals Math. **5** (1904), no. 4, 173–180. ↑5

- [5] Kent D. Boklan and John H. Conway, *Expect at most one billionth of a new Fermat Prime!*, Math. Intelligencer **39** (2017), no. 1, 3–5. ↑3
- [6] Ashish Kumar Das, *On Finite Groups Having Perfect Order Subsets*, Int. J. Algebra **3** (2009), no. 13, 629–637. ↑2, 8, 9
- [7] Carrie E. Finch and Lenny Jones, *A Curious Connection Between Fermat Numbers and Finite Groups*, Amer. Math. Monthly **109** (2002), no. 6, 517–524. ↑2
- [8] ———, *Non-Abelian groups with perfect order subsets*, The JP Journal of Algebra and Number Theory **3** (2003), no. 1, 13–26. ↑2, 6
- [9] Kevin Ford and Sergei Konyagin and Florian Luca, *On groups with Perfect Order subsets*, Moscow J. Combinatorics and Number Theory **2** (2012), no. 4, 297–312. ↑2
- [10] The GAP Group, *GAP – Groups, Algorithms, and Programming, Version 4.11.0*, 2020. ↑9
- [11] Daniel Gorenstein, *Finite Groups*, Harper’s series in modern mathematics, Harper & Row, New York, Evanston, and London, 1967. ↑7
- [12] Lenny Jones and Kelly Toppin, *On three questions concerning groups with perfect order subsets*, Involve **4** (2011), no. 3, 251–261. ↑2
- [13] Wieb and Cannon Bosma John and Playoust, *The Magma algebra system. I. The user language*, J. Symbolic Comput. **24** (1997), no. 3-4, 235–265. Computational algebra and number theory (London, 1993). ↑9
- [14] L. Bernardin and P. Chin and P. DeMarco and K. O. Geddes and D. E. G. Hare and K. M. Heal and G. Labahn and J. P. May and J. McCarron and M. B. Monagan and D. Ohashi and S. M. Vorkoetter, *Maple Programming Manual*, Maplesoft, 1996–2021. ↑9
- [15] James McCarron, *Hamiltonian groups with perfect order classes*, Math. Proc. Royal Irish Acad. **121A** (2021), no. 1, 1–8, DOI <https://doi.org/10.3318/pria.2021.121.01>. ↑8, 23, 24
- [16] P. Mihăilescu, *Primary cyclotomic units and a proof of Catalans conjecture*, Journal für die reine und angewandte Mathematik **2004** (2004), no. 572, 167–195. ↑5
- [17] D. S. Passman, *Permutation Groups*, Dover Publications, Inc., 2012. ↑7
- [18] James Pierpont, *On an undemonstrated theorem of the Disquisitiones Arithmeticae*, Bull. Amer. Math. Soc. **2** (1895), no. 3, 77–83. ↑3
- [19] Paulo Ribenboim, *My Numbers, My Friends*, Springer Verlag, 2000. ↑5
- [20] D.J.S. Robinson, *A Course in the Theory of Groups*, Graduate Texts in Mathematics, vol. 80, Springer-Verlag, New York, 1993. ↑7
- [21] R. Shen, *A note on finite groups having perfect order subsets*, Int. J. Algebra **13** (2010), no. 4, 643–646. ↑2
- [22] R. Shen and W. Shi and J. Shi, *POS-Groups with some cyclic Sylow subgroups*, Bull. Iran. Math. Soc. **39** (2013), no. 5, 941–957. ↑2
- [23] M. Suzuki, *Group Theory I*, Grundlehren der mathematischen Wissenschaften, vol. 247, Springer Berlin Heidelberg, 2014. ↑37
- [24] Nguyen Trong Tuan and Bui Xuan Hai, *On perfect order subsets in finite groups*, Int. J. Algebra **4** (2010), no. 21, 1021–1029. ↑2, 21
- [25] Wikipedia, *Pierpont-prime*, https://en.wikipedia.org/wiki/Pierpont_prime. ↑3

- [26] Joseph A. Wolf, *Spaces of Constant Curvature*, Sixth Edition, AMS Chelsea Publishing, AMS Chelsea Publishing, 2011. ↑7
- [27] K. Zsigmondy, *Zur Theorie der Potenzreste*, J. Monatshefte für Mathematik **3** (1892), no. 1, 265–284. ↑5

MAPLESOFT, 615 KUMPF DRIVE,, WATERLOO, ON, CANADA, N2V 1K8
Email address: james@maplesoft.com