

Monogeneity in totally real extensions of imaginary quadratic fields with an application to simplest quartic fields

István Gaál

University of Debrecen, Mathematical Institute

H-4002 Debrecen Pf.400., Hungary,

e-mail: gaal.istvan@unideb.hu,

October 10, 2021

Dedicated to Professor Michael Pohst on his 75th birthday.

Mathematics Subject Classification: Primary 11R04; 11R16. Secondary 11Y50.

Key words and phrases: monogeneity; power integral basis; totally real fields; simplest quartic fields; calculating the solutions

Abstract

We describe an efficient algorithm to calculate generators of power integral bases in composites of totally real fields with imaginary quadratic fields. We show that the calculation can be reduced to solving index form equations in the original totally real fields.

We illustrate our method by investigating monogeneity in the infinite parametric family of imaginary quadratic extensions of the simplest quartic fields.

1 Introduction

There is an extensive literature of monogeneity of number fields and power integral bases, see [14], [5]. A number field K of degree n is *monogenic* if its ring of integers \mathbb{Z}_K is a simple ring extension of \mathbb{Z} , that is there exists $\alpha \in \mathbb{Z}_K$ with $\mathbb{Z}_K = \mathbb{Z}[\alpha]$. In this case $(1, \alpha, \dots, \alpha^{n-1})$ is an integral basis of K , called *power integral basis*. We also call α the *generator* of this power integral basis. The algebraic integer α generates a power integral basis if and only if its *index*

$$I(\alpha) = \sqrt{\left| \frac{D(\alpha)}{D_K} \right|}$$

is equal to 1, where $D(\alpha)$ is the discriminant of α .

The calculation of generators of power integral bases can be reduced to certain diophantine equations, called *index form equations*, cf. [5].

There exist general algorithms for solving index form equations in cubic, quartic, quintic, sextic fields, however the general algorithms for quintic and sextic fields are already quite tedious, see [5]. Therefore it is worthy to develop efficient methods for the resolution of special types of higher degree number fields, cf. [10], [5].

In some cases we considered monogeneity in composites of fields, see [4], [11]. In these cases the index form factorizes that makes the resolution of the index form equation easier.

On the other hand, considering totally real relative Thue equations over imaginary quadratic fields, it has turned out, that the relative Thue equations can be reduced to absolute Thue equations (over \mathbb{Z}), cf. [8]. A similar idea was used in [3], [7].

In the present paper we study composites $K = LM$ of a totally real fields L and imaginary quadratic fields M . We show that in this case the resolution of the index form equation in K can be reduced to solving the index form equation in L . If L is of degree n , then the index form equation of L is of degree $n(n-1)/2$ in $n-1$ variables, but the index form equation in K is of degree $2n(2n-1)/2$ in $2n-1$ variables. Therefore our statement simplifies a lot the calculation of generators of power integral bases in fields of type K of degree $2n$.

Surprisingly the proofs of our statements are quite simple. However, they provide a powerful tool. Our Theorem 2 immediately implies some results of [4] (see our remarks after the proof of Theorem 2).

The strength and usefulness of our results is also demonstrated in the application given in Section 4, where we consider composites of the so called simplest quartic fields and imaginary quadratic fields.

2 Composites of totally real fields and imaginary quadratic fields

Let $f(x) \in \mathbb{Z}[x]$ be a monic irreducible polynomial of degree n having all roots $\xi = \xi^{(1)}, \dots, \xi^{(n)}$ in \mathbb{R} . The field $L = \mathbb{Q}(\xi)$ is then totally real. Assume that L has integral basis $(\ell_1 = 1, \ell_2, \dots, \ell_n)$ and discriminant D_L . We shall denote by $\gamma^{(j)}$ the conjugates of any $\gamma \in L$ ($j =$

$1, \dots, n$). The index form corresponding to the basis $(\ell_1 = 1, \ell_2, \dots, \ell_n)$ is defined by

$$I_L(x_2, \dots, x_n) = \frac{1}{\sqrt{|D_L|}} \prod_{1 \leq j_1 < j_2 \leq n} \left(x_2 \left(\ell_2^{(j_1)} - \ell_2^{(j_2)} \right) + \dots + x_n \left(\ell_n^{(j_1)} - \ell_n^{(j_2)} \right) \right).$$

As it is known, $I_L(x_2, \dots, x_n) \in \mathbb{Z}[x_2, \dots, x_n]$ and for any $x_1, x_2, \dots, x_n \in \mathbb{Z}$ the algebraic integer element

$$\alpha = x_1 + x_2 \ell_2 + \dots + x_n \ell_n$$

generates a power integral basis $(1, \alpha, \dots, \alpha^{n-1})$ if and only if

$$I_L(x_2, \dots, x_n) = \pm 1.$$

Let $0 < d \in \mathbb{Z}$ be a square-free integer, set

$$\omega = \begin{cases} i\sqrt{d} & \text{if } -d \equiv 2, 3 \pmod{4}, \\ \frac{1 + i\sqrt{d}}{2} & \text{if } -d \equiv 1 \pmod{4}. \end{cases}$$

The conjugates of ω are $\omega^{(1)} = \omega, \omega^{(2)} = \bar{\omega}$ (the complex conjugate of ω). Let $M = \mathbb{Q}(i\sqrt{d})$ with discriminant D_M .

Our purpose is to investigate the composite field $K = L \cdot M$ of degree 2ℓ and discriminant D_K . We assume $(D_L, D_M) = 1$. Then $(1, \ell_2, \dots, \ell_n, \omega, \omega \ell_2, \dots, \omega \ell_n)$ is an integral basis of K and $D_K = D_M^n \cdot D_L^2$ (cf. [14]).

In the ring of integers \mathbb{Z}_K of K any element α can be represented as

$$\alpha = x_1 + x_2 \ell_2 + \dots + x_n \ell_n + y_1 \omega + y_2 \omega \ell_2 + \dots + y_n \omega \ell_n = X_1 + X_2 \ell_2 + \dots + X_n \ell_n, \quad (1)$$

with $x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{Z}$ and $X_j = x_j + \omega y_j$ are elements in the ring \mathbb{Z}_M of algebraic integers of M .

We shall use the following consequence of Theorem 1 of [4]:

Lemma 1 *If α of (1) generates a power integral basis in K then*

$$N_{M/\mathbb{Q}}(I_L(X_2, \dots, X_n)) = \pm 1 \quad (2)$$

and

$$N_{L/\mathbb{Q}}(y_1 + y_2 \ell_2 + \dots + y_n \ell_n) = \pm 1. \quad (3)$$

The conjugates of α of (1) are obtained obviously as

$$\alpha^{(j,k)} = x_1 + x_2 \ell_2^{(j)} + \dots + x_n \ell_n^{(j)} + y_1 \omega^{(k)} + y_2 \omega^{(k)} \ell_2^{(j)} + \dots + y_n \omega^{(k)} \ell_n^{(j)},$$

for $j = 1, \dots, n, k = 1, 2$. Note that in this case the index form has three factors, two of them are the polynomials on the left hand sides of equations (2), (3). The third factor is

$$F(x_2, \dots, x_n, y_1, y_2, \dots, y_n) = \prod_{1 \leq j_1, j_2 \leq n, j_1 \neq j_2} (\alpha^{(j_1, 1)} - \alpha^{(j_2, 2)}), \quad (4)$$

(cf. [4], [5]). This is also a polynomial with coefficients in \mathbb{Z} . Equations (2), (3) together with

$$F(x_2, \dots, x_n, y_1, y_2, \dots, y_n) = \pm 1 \quad (5)$$

are already equivalent to α generating a power integral basis.

In our main result we reduce the relative index form equation (2) to absolute equations, inequalities. This makes the resolution of (2) much easier.

Theorem 2 *Assume α of (1) generates a power integral basis in K .*

If $-d \equiv 2, 3 \pmod{4}$, then

$$|I_L(x_2, \dots, x_n)| \leq 1 \quad (6)$$

and

$$|I_L(y_2, \dots, y_n)| \leq \frac{1}{(\sqrt{d})^{n(n-1)/2}}. \quad (7)$$

If $-d \equiv 1 \pmod{4}$, then

$$|I_L(2x_2 + y_2, \dots, 2x_n + y_n)| \leq 2^{n(n-1)/2} \quad (8)$$

and

$$|I_L(y_2, \dots, y_n)| \leq \left(\frac{2}{\sqrt{d}}\right)^{n(n-1)/2}. \quad (9)$$

Proof of Theorem 2. According to the arguments in the proof of Theorem 1 of [4] we have

$$|N_{M/Q}(I_L(X_2, \dots, X_n))| = \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} |\alpha^{(j_1, k)} - \alpha^{(j_2, k)}| \right). \quad (10)$$

If $-d \equiv 2, 3 \pmod{4}$, then

$$\operatorname{Re}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)}) = x_2(\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + x_n(\ell_n^{(j_1)} - \ell_n^{(j_2)})$$

and

$$\operatorname{Im}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)}) = \sqrt{d} \cdot \left(y_2(\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + y_n(\ell_n^{(j_1)} - \ell_n^{(j_2)}) \right).$$

We have

$$\begin{aligned} |I_L(x_2, \dots, x_n)|^2 &= |N_{M/Q}(I_L(x_2, \dots, x_n))| \\ &= \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} \left| x_2(\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + x_n(\ell_n^{(j_1)} - \ell_n^{(j_2)}) \right| \right) \\ &= \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} |\operatorname{Re}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)})| \right) \\ &\leq \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} |\alpha^{(j_1, k)} - \alpha^{(j_2, k)}| \right) \\ &= |N_{M/Q}(I_L(X_2, \dots, X_n))| \end{aligned}$$

whence by (2) we obtain (6). Similarly,

$$\begin{aligned}
(\sqrt{d})^{n(n-1)} |I_L(y_2, \dots, y_n)|^2 &= (\sqrt{d})^{n(n-1)} |N_{M/Q}(I_L(y_2, \dots, y_n))| \\
&= \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} \sqrt{d} \left| y_2(\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + y_n(\ell_n^{(j_1)} - \ell_n^{(j_2)}) \right| \right) \\
&= \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} |\operatorname{Im}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)})| \right) \\
&\leq \prod_{k=1}^2 \left(\frac{1}{|\sqrt{D_L}|} \prod_{1 \leq j_1 < j_2 \leq n} |\alpha^{(j_1, k)} - \alpha^{(j_2, k)}| \right) \\
&= |N_{M/Q}(I_L(X_2, \dots, X_n))|,
\end{aligned}$$

whence by (2) we obtain (7).

If $-d \equiv 1 \pmod{4}$, then

$$\operatorname{Re}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)}) = \frac{2x_2 + y_2}{2} \cdot (\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + \frac{2x_n + y_n}{2} \cdot (\ell_n^{(j_1)} - \ell_n^{(j_2)})$$

and

$$\operatorname{Im}(\alpha^{(j_1, k)} - \alpha^{(j_2, k)}) = \sqrt{d} \cdot \left(\frac{y_2}{2} \cdot (\ell_2^{(j_1)} - \ell_2^{(j_2)}) + \dots + \frac{y_n}{2} \cdot (\ell_n^{(j_1)} - \ell_n^{(j_2)}) \right).$$

The above arguments lead us to

$$\left| I_L \left(\frac{2x_2 + y_2}{2}, \dots, \frac{2x_n + y_n}{2} \right) \right| \leq 1$$

and

$$\left| I_L \left(\frac{y_2}{2}, \dots, \frac{y_n}{2} \right) \right| \leq \frac{1}{(\sqrt{d})^{n(n-1)/2}},$$

whence we obtain (8) and (9) accordingly. \square

Remark. Our Theorem 2 immediately implies the result of [4] on monogeneity of composites of totally real cyclic fields of prime degree and imaginary quadratic fields, as well as on monogeneity of composites of Lehmer's quintics and imaginary quadratic fields. Actually, our result yields, that the above statements of [4] are valid for any totally real number field.

3 Applying Theorem 2

Let $-d \equiv 2, 3 \pmod{4}$.

If $-d = -1$, then left hand side of both (6) and (7) is ± 1 . Concerning equation (6) this yields that either $I(x_2, \dots, x_n) = 0$, that is $x_2 = \dots = x_n = 0$ or $I(x_2, \dots, x_n) = \pm 1$, that is $\beta = x_2 \ell_2 + \dots + x_n \ell_n$ generates a power integral basis in L . Similary, either $y_2 = \dots = y_n = 0$, or $\gamma = y_2 \ell_2 + \dots + y_n \ell_n$ generates a power integral basis in L . For given y_2, \dots, y_n we calculate

y_1 from (3). We test all these $x_2, \dots, x_n, y_1, y_2, \dots, y_n$ in (5). Note that $x_2 = \dots = x_n = 0$, and simultaneously $y_2 = \dots = y_n = 0$ is not possible, since ω does not generate K .

Otherwise, if $-d \neq -1$, (7) gives $y_2 = \dots = y_n = 0$ and (3) gives $y_1 = \pm 1$. The values of x_2, \dots, x_n are obtained from (6). If $I_L(x_2, \dots, x_n) = 0$ then $x_2 = \dots = x_n = 0$ and $\alpha = \pm \omega$ which is again impossible, since ω does not generate K . Hence we have to take those x_2, \dots, x_n for which $I_L(x_2, \dots, x_n) = \pm 1$, that is $\beta = x_2 \ell_2 + \dots + x_n \ell_n$ generates a power integral basis in L . We test all these $x_2, \dots, x_n, y_1, y_2, \dots, y_n$ in (5).

Let now $-d \equiv 1 \pmod{4}$.

If $-d = -3$, then the left hand side of (8) is $2^{n(n-1)/2}$, the left hand side of (9) is $(2/\sqrt{3})^{n(n-1)/2}$. Hence we have to determine the elements $\beta = z_2 \ell_2 + \dots + z_n \ell_n$ having index $\leq 2^{n(n-1)/2}$ and to select from those the elements $\gamma = y_2 \ell_2 + \dots + y_n \ell_n$ having index $\leq (2/\sqrt{3})^{n(n-1)/2}$. We test if there exist $x_i \in \mathbb{Z}$ with $x_i = (z_i - y_i)/2$. For given y_2, \dots, y_n we calculate y_1 from (3). We test all possible $x_2, \dots, x_n, y_1, y_2, \dots, y_n$ in (5).

Otherwise, if $-d \neq -3$, equation (9) has left hand side < 1 , therefore $y_2 = \dots = y_n = 0$ and (3) gives $y_1 = \pm 1$. (8) implies $|I_L(x_2, \dots, x_n)| \leq 1$. $I_L(x_2, \dots, x_n) = 0$ is not possible again, $I_L(x_2, \dots, x_n) = \pm 1$ yields that $\beta = x_2 \ell_2 + \dots + x_n \ell_n$ generates a power integral basis in L . We test all these $x_2, \dots, x_n, y_1, y_2, \dots, y_n$ in (5).

This means that for calculating generators of power integral basis in K we only need the generators of power integral bases in L and, in case $-d = -3$, elements of small indices of L . This is easy to calculate in lower degree (cubic, quartic) fields, therefore we obtain an efficient method to calculate generators of power integral bases in sextic, octic fields, that are composites of cubic, quartic fields and imaginary quadratic fields.

As a consequence of the above arguments we have

Theorem 3 *If $-d \neq -1, -3$ then all generators of power integral bases of K are of the form*

$$\alpha = x + \beta \pm \omega$$

where β generates a power integral basis in L and $x \in \mathbb{Z}$ is arbitrary.

Remark. Our Theorem 3 implies Theorem 2 of [4] on generators of power integral bases in composites of totally real cyclic fields of prime degree and imaginary quadratic fields.

4 Composites of the simplest quartic fields and imaginary quadratic fields

In this section we shall give an application to an infinite parametric family of number fields, that shows the strength of our method.

Let $a > 0$ be an integer with $a \neq 3$. Let ξ be a root of

$$f(x) = x^4 - ax^3 - 6x^2 + ax + 1.$$

The parametric family of fields $L = \mathbb{Q}(\xi)$ is called *simplest quartic fields*, see M. N. Gras [12]. (She showed that for $a > 0, a \neq 3$ the polynomial $f(x)$ is irreducible.) In the following we shall also assume that $a^2 + 16$ is not divisible by an odd square. This condition was needed by H. K. Kim and J. H. Lee [13] to determine an integral basis of L . (M.N.Grass [12] showed that $a^2 + 16$ represents infinitely many square free integers.) Using the discriminant

$$D(f) = 4(a^2 + 16)^3$$

of the polynomial we obtain the discriminant of L . $v_2(x)$ denotes the exponent of 2 in the prime power decomposition of the integer x .

Lemma 4 *Under the above assumptions an integral basis and the discriminant of L is given by*

$$\begin{aligned} & \left(1, \xi, \xi^2, \frac{1+\xi^3}{2}\right), & D_L &= (a^2 + 16)^3, & \text{if } v_2(t) &= 0, \\ & \left(1, \xi, \frac{1+\xi^2}{2}, \frac{\xi+\xi^3}{2}\right), & D_L &= \frac{(a^2+16)^3}{4}, & \text{if } v_2(t) &= 1, \\ & \left(1, \xi, \frac{1+\xi^2}{2}, \frac{1+\xi+\xi^2+\xi^3}{4}\right), & D_L &= \frac{(a^2+16)^3}{16}, & \text{if } v_2(t) &= 2, \\ & \left(1, \xi, \frac{1+2\xi-\xi^2}{4}, \frac{1+\xi+\xi^2+\xi^3}{4}\right), & D_L &= \frac{(a^2+16)^3}{64}, & \text{if } v_2(t) &\geq 3. \end{aligned}$$

P.Olajos [15] determined all generators of power integral bases (up to sign and translation by elements of \mathbb{Z}).

Lemma 5 *Under the above assumptions power integral bases exist only for $a = 2$ and $a = 4$. All generators of power integral bases are given by*

- $a = 2$, $\alpha = x\xi + y\frac{1+\xi^2}{2} + z\frac{\xi+\xi^3}{2}$ where
 $(x, y, z) = (4, 2, -1), (-13, -9, 4), (-2, 1, 0), (1, 1, 0),$
 $(-8, -3, 2), (-12, -4, 3), (0, -4, 1), (6, 5, -2), (-1, 1, 0), (0, 1, 0)$
- $a = 4$, $\alpha = x\xi + y\frac{1+\xi^2}{2} + z\frac{1+\xi+\xi^2+\xi^3}{4}$ where
 $(x, y, z) = (3, 2, -1), (-2, -2, 1), (4, 8, -3), (-6, -7, 3),$
 $(0, 3, -1), (1, 3, -1).$

Note also that I. Gaál and G. Petrányi [9] calculated the minimal indices and all elements of minimal index for all parameters a .

Our purpose is to study composites of simplest quartic fields with imaginary quadratic fields. Let d be a squarefree positive integer, $M = \mathbb{Q}(i\sqrt{d})$. Set

$$\omega = i\sqrt{d} \text{ if } -d \equiv 2, 3 \pmod{4} \text{ and } \omega = \frac{1+i\sqrt{d}}{4} \text{ if } -d \equiv 1 \pmod{4}.$$

We assume that $(D_M, D_L) = 1$. If $v_2(a) \geq 1$ then D_L is even, hence in that case we must have $-d \equiv 1 \pmod{4}$.

We consider monogeneity of the composite field $K = LM$ of degree 8. This is an infinite parametric family of octic fields, depending on the parameters d, a .

We show:

Theorem 6 *Under the above conditions for $d \neq 3$ the field K is not monogenic.*

Proof of Theorem 6

For a given parameter a denote by $(1, \ell_2, \ell_3, \ell_4)$ an integral basis of L . Denote by $I_L(x_2, x_3, x_4)$ the index form corresponding to this integral basis. The condition $(D_L, D_M) = 1$ implies that an integral basis of K is given by $(1, \ell_2, \ell_3, \ell_4, \omega, \omega\ell_2, \omega\ell_3, \omega\ell_4)$. Represent any $\alpha \in \mathbb{Z}_K$ in the form

$$\alpha = x_1 + x_2\ell_2 + x_3\ell_3 + x_4\ell_4 + y_1\omega + y_2\omega\ell_2 + y_3\omega\ell_3 + y_4\omega\ell_4$$

with integer coefficients x_i, y_i ($1 \leq i \leq 4$).

I. Assume $d \neq 1$. By Theorem 2 we have (in both cases $-d \equiv 2, 3 \pmod{4}$ and $-d \equiv 1 \pmod{4}$) $I_L(y_2, y_3, y_4) = 0$ implying $y_2 = y_3 = y_4 = 0$. Theorem 1, equation (3) implies $y_1 = \pm 1$. Further, by Theorem 2 $|I_L(x_2, x_3, x_4)| \leq 1$. $I_L(x_2, x_3, x_4) = 0$ would imply $x_2 = x_3 = x_4 = 0$ which is impossible, while ω does not generate K . Therefore $I_L(x_2, x_3, x_4) = \pm 1$ which means that $\beta = x_2\ell_2 + x_3\ell_3 + x_4\ell_4$ generates a power integral basis in L . By Lemma 5 this is only possible for $a = 2, 4$ in which cases all possible x_2, x_3, x_4 are known.

Consider now the third factor (5) of the index form equation of K . In both cases $-d \equiv 2, 3 \pmod{4}$ and $-d \equiv 1 \pmod{4}$, both for $a = 2$ and $a = 4$ we substitute the possible x_2, x_3, x_4 and $y_2 = y_3 = y_4 = 0, y_1 = \pm 1$ into $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4)$. In all cases we obtain a polynomial of degree 6 in d with positive coefficients, implying that $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4) = \pm 1$ is not possible.

II. Let now $d = 1$. By Theorem 2 we have

$$|I_L(x_2, x_3, x_4)| \leq 1, \quad |I_L(y_2, y_3, y_4)| \leq 1.$$

a.) $I_L(x_2, x_3, x_4) = I_L(y_2, y_3, y_4) = 0$ implies $x_2 = x_3 = x_4 = 0$ and $y_2 = y_3 = y_4 = 0$, this is not possible, while ω does not generate K . Hence we must have either $I_L(x_2, x_3, x_4) = \pm 1$ or $I_L(y_2, y_3, y_4) = \pm 1$ (or both), hence there must be a power integral basis in L , whence only $a = 2$ and $a = 4$ is possible. In the following we restrict ourselves to $a = 2$ and $a = 4$.

b.) Assume $I_L(x_2, x_3, x_4) = \pm 1$ and $I_L(y_2, y_3, y_4) = 0$. The possible x_2, x_3, x_4 are listed in Lemma 5, $y_2 = y_3 = y_4 = 0, y_1 = \pm 1$. We substitute the possible values of the variables into $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4)$ and find that in all cases it takes huge values, not ± 1 .

c.) Assume $I_L(x_2, x_3, x_4) = 0$ and $I_L(y_2, y_3, y_4) = \pm 1$. Then $x_2 = x_3 = x_4 = 0$ and the possible y_2, y_3, y_4 are listed in Lemma 5. We substitute the possible values $x_2 = x_3 = x_4 = 0$ and y_2, y_3, y_4 into $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4)$ and obtain a polynomial of degree 12 in y_1 . Solving $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4) = \pm 1$ in y_1 we never get integer solutions for y_1 .

d.) If $I_L(x_2, x_3, x_4) = \pm 1$ and $I_L(y_2, y_3, y_4) = \pm 1$, then x_2, x_3, x_4 and y_2, y_3, y_4 may run independently through the possible triplets listed in Lemma 5. In all cases we substitute x_2, x_3, x_4

and y_2, y_3, y_4 into $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4)$ and obtain a polynomial of degree 12 in y_1 . Solving $F(x_2, x_3, x_4, y_1, y_2, y_3, y_4) = \pm 1$ in y_1 we never get integer solutions for y_1 . \square

Remark. The case $a = 3$ is not covered by Theorem 6. In this case we have

$$|I_L(2x_2 + y_2, 2x_3 + y_3, 2x_4 + y_4)| \leq 64$$

and

$$|I_L(y_2, y_3, y_4)| \leq 2.$$

According to [9] we have minimal index 2 in L for infinitely many parameters a . Especially the elements of index ≤ 64 seems very difficult to determine. This could be the subject of a further research.

Remark. Note that in [11] we obtained conditions on the monogeneity of composites of fields, among others of simplest quartic fields and quadratic fields. We did not assume that the discriminants are relative prime and involved also real quadratic fields. However we only obtained certain divisibility conditions on the parameters as a consequence of monogeneity.

5 One more example

We also provide a positive example to show that such composite fields may happen to be monogenic. Consider the totally real quartic field L generated by a root ξ of the polynomial $f(x) = x^4 - 4x^2 - x + 1$. In this field $(1, \xi, \xi^2, \xi^3)$ is an integral basis, $D_L = 1957$. Let $M = \mathbb{Q}(i)$ with $D_M = -4$, coprime to D_L . The composite field $K = LM$ can be generated e.g. by $\alpha = i\xi$ having minimal polynomial $g(x) = x^8 + 8x^6 + 18x^4 + 9x^2 + 1$. In K the element α generates a power integral basis.

6 Computational aspects

All calculations connected with the above examples were performed in Maple [2]. Our procedures were executed on an average laptop running under Windows. The CPU time took all together some seconds.

References

- [1] M. A. Bennett and A. Ghadermarzi, *Extremal families of cubic Thue equations*, J. Théor. Nombres Bordx. **27**(2015), 389-403.
- [2] B. W. Char, K. O. Geddes, G. H. Gonnet, M. B. Monagan, S. M. Watt (eds.) MAPLE, Reference Manual, Watcom Publications, Waterloo, Canada, 1988.
- [3] I. Gaál, *Computing elements of given index in totally complex cyclic sextic fields*, J. Symbolic Comput. **20**(1995), No. 1, 61-69.

- [4] I. Gaál, *Power integral bases in composites of number fields*, *Canad. Math. Bulletin*, **41**(1998), 158–165.
- [5] I. Gaál, *Diophantine equations and power integral bases. Theory and algorithms*. 2nd edition, Birkhäuser, Boston, 2019.
- [6] I. Gaál, *Calculating "small" solutions of inhomogeneous relative Thue inequalities*, submitted.
- [7] I. Gaál, *Monogeneity in totally complex sextic fields, revisited*, *J. Pure and Applied Math.*, **47**(2020), no. 1., 87-98.
- [8] I. Gaál, B. Jadrijević and L. Remete, *Totally real Thue inequalities over imaginary quadratic fields*, *Glasnik Matematički*, **53**(2018), No. 2, 229–238.
- [9] I. Gaál and G. Petrányi, *Calculating all elements of minimal index in the infinite parametric family of simplest quartic fields*, *Czech. Math. J.* **64**(2014), No. 2, 465–475.
- [10] I. Gaál and L. Remete, *Power integral bases in a family of sextic fields with quadratic subfields*, *Tatra Mt. Math. Publ.* **64**(2015), 59-66.
- [11] I. Gaál and L. Remete, *Integral bases and monogeneity of composite fields*, *Exp. Math.* **28**(2019), No. 2, 209-222.
- [12] M. N.Gras, *Table numérique du nombre de classes et des unités des extensions cycliques réelles de degré 4 de \mathbb{Q}* , *Publ. Math. Fac. Sci. Besancon, Théor. Nombres, Année 1977-1978*, Fasc. 2, 1978. (In French.)
- [13] H. K. Kim and J. H. Lee, *Evaluation of the Dedekind zeta function at $s = -1$ of the simplest quartic fields*, *Trends in Math., New Ser., Inf. Center for Math. Sci.*, **11**(2009), 63-79.
- [14] W. Narkiewicz, *Elementary and Analytic Theory of Algebraic Numbers*, Third Edition, Springer, 2004.
- [15] P. Olajos, *Power integral bases in the family of simplest quartic fields*, *Exp. Math.* **14**(2005), No. 2, 129-132 (2005).