

THE FINITE DUALS OF AFFINE PRIME REGULAR HOPF ALGEBRAS OF GK-DIMENSION ONE

KANGQIAO LI AND GONGXIANG LIU

ABSTRACT. This paper is an attempt to construct a special kind of Hopf pairing $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$. Specifically, H^\bullet and H should be both affine, noetherian and of the same GK-dimension. In addition, some properties of them would be dual to each other. We test the ideas in two steps for all the affine prime regular Hopf algebras H of GK-dimension one: 1) We compute the finite duals H° of them, which are given by generators and relations; 2) the Hopf pairings desired are determined by choosing certain Hopf subalgebras H^\bullet of H° , where $\langle -, - \rangle$ becomes the evaluation.

1. INTRODUCTION

Throughout this paper, \mathbb{k} is assumed to be an algebraically closed field of characteristic 0, and all matrices, vector spaces, algebras and Hopf algebras are assumed to be over \mathbb{k} .

Roughly speaking, the concept of Hopf algebras is still valid under taking the linear duals. This elementary point of view suggests to us that structures of the dual are effective when we deal with certain problems on a Hopf algebra. The most basic definition of the dual for a Hopf algebra H would be the *finite dual* H° described by Heyneman and Sweedler [11], which is also a Hopf algebra over the same base field. Of course, there are a number of relationships between H and H° , and this motivates the authors to consider the structure of duals for varieties of Hopf algebras.

Of course it is known that when H is a finite-dimensional Hopf algebra, its dual Hopf algebra H^* has the same dimension as well as similar or dual properties with H . Indeed, the structures of H are determined conversely by those of H^* . However for an infinite-dimensional Hopf algebra H , its finite dual H° might be much too large (or small) to be studied easily. This motivates us to consider some kinds of dual Hopf algebras H^\bullet for certain (infinite-dimensional) Hopf algebras H , where the original evaluation is replaced by Hopf pairings $H^\bullet \otimes H \rightarrow \mathbb{k}$. Our primary goal is to formulate such Hopf pairings for noetherian affine Hopf algebras H of finite GK-dimensions, such that:

Key words and phrases. GK-dimension one, Hopf algebra, Finite dual, Prime, Regular, Hopf pairing.

2020 *Mathematics Subject Classification.* 16E05, 16T05 (primary), 16P40, 16S34 (secondary).

[†]The first author was supported by Zhejiang Provincial Natural Science Foundation of China under No.LQ23A010003. The second author was supported by NSFC 12271243.

- (HP1) The pairing is non-degenerate;
- (HP2) H^\bullet and H are both affine and noetherian;
- (HP3) $\text{GKdim}(H^\bullet) = \text{GKdim}(H)$.

Meanwhile, we wish that some properties of H^\bullet and H would be dual to each other (such as (HP4) in Section 6.2), which should be noted in different situations. Finally, H^\bullet would be better to exist uniquely for any given H under our assumptions. In this paper, we try to establish the desired Hopf pairings for affine prime regular Hopf algebras H of GK-dimension one. For this purpose, the finite dual H° of each H is determined, and afterwards H^\bullet would be chosen as a suitable Hopf subalgebra of H° .

However, it is not easy to determine the structure of H° for a Hopf algebra H in general, especially for infinite-dimensional ones. Of course, the most direct way to get H° is by definition. For this, recall that H° is the Hopf algebra generated by $f \in H^*$ which vanish on an ideal $I \subseteq H$ of finite codimension. This means that we need a description of *all* finite codimensional ideals which is impossible in general. To the authors' knowledge, there are two other ways to get H° if H is good enough. One is applying the well-known Cartier-Konstant-Milnor-Moore's Theorem ([22]) if H happens to be commutative. The related idea and method were generalized further (see [23, Chapter 9], [7]). Another one is applying representation theoretical way if the representation category $\text{Rep-}H$ of finite dimensional modules happens to be very nice (see [30]).

In the literature, significant progress has been made in classifying infinite dimensional noetherian Hopf algebras of low Gelfand-Kirillov dimensions (GK-dimension for short). The program of classifying Hopf algebras of GK-dimension one was initiated by Lu, Wu and Zhang [20]. Then the second author found a new class of examples about prime regular Hopf algebras of GK-dimension one [18]. Brown and Zhang [5] made further efforts in this direction and classified all prime regular Hopf algebras of GK-dimension one under an extra hypothesis. In 2016, Wu, Ding and the second author [35] removed this hypothesis and gave a complete classification of prime regular Hopf algebras of GK-dimension one, in which some non-pointed Hopf algebras were found. Recently, the second author classified prime Hopf algebras of GK-dimension one satisfying two certain conditions [19]. For Hopf algebras H of GK-dimension two, Goodearl and Zhang [10] gave a classification of the case when H is a domain satisfying $\text{Ext}_H^1(\mathbb{k}, \mathbb{k}) \neq 0$. For those with vanishing Ext-groups, some interesting examples were constructed by Wang, Zhang and Zhuang [32]. The progress can also be found in a survey by Brown and Zhang [6]. As for Hopf algebras of GK-dimensions three and four, all connected ones were classified by Zhuang [36] and Wang, Zhang and Zhuang [33] respectively.

Our main interest in this paper is to deal with the affine prime Hopf algebras of GK-dimension one, whose detailed structures are recalled in Section 2.1. An interesting fact is that all affine prime regular Hopf algebras are commutative-by-finite [3], that is, a finite module over a normal commutative Hopf subalgebra. This suggests that we have a chance to get the finite duals of affine prime regular Hopf algebras of GK-dimension one explicitly. Actually, Ge and the second author use this observation together with some other techniques to determine the finite dual of the infinite

dihedral group algebra $\mathbb{k}\mathbb{D}_\infty$ recently [9]. In addition, after finishing this paper we realized that A. Jahn and M. Couto already considered this question along this line in their theses [12, 8].

This paper is to develop the idea and techniques in [9] further and as the main result we get the finite duals of all the affine prime regular Hopf algebras of GK-dimension one in terms of generators and relations. Now let us recall in our notations that there are five kinds of such Hopf algebras listed in [35]:

$$\mathbb{k}[x], \mathbb{k}[x^{\pm 1}], \mathbb{k}\mathbb{D}_\infty, T_\infty(n, v, \xi), B(n, \omega, \gamma), D(m, d, \xi).$$

Since the finite duals of $\mathbb{k}[x]$, $\mathbb{k}[x, x^{-1}]$ and $\mathbb{k}\mathbb{D}_\infty$ are known, it remains for us to determine the latter three kinds of Hopf algebras. Our main result is:

Theorem 1.1. (1) *As a Hopf algebra, $T_\infty(n, v, \xi)^\circ$ is isomorphic to the Hopf algebra $T_{\infty^\circ}(n, v, \xi)$ constructed in Subsection 3.1,*

(2) *As a Hopf algebra, $B(n, \omega, \gamma)^\circ$ is isomorphic to the Hopf algebra $B_\circ(n, \omega, \gamma)$ constructed in Subsection 4.1,*

(3) *As a Hopf algebra, $D(m, d, \xi)^\circ$ is isomorphic to the Hopf algebra $D_\circ(m, d, \xi)$ constructed in Subsection 5.1.*

These three claims are presented respectively as Theorems 3.4, 4.4, and 5.7. In brief, our computation for finite dual H° starts with properties of certain polynomial or Laurent polynomial subalgebra P , by which we figure cofinite left ideals of H . Then we construct some key elements in H° and determine their relations under Hopf operations. Finally it is proved that these elements generate H° with desired relations.

Using our descriptions of finite duals, we find that for an affine prime regular Hopf algebra H one can always get a Hopf algebra H^\bullet such that there is a non-degenerate Hopf pairing (see Majid [21] for the definition) between H and H^\bullet . The definition can also be found as Definition 6.1 in this paper. Some properties of H^\bullet are listed in the following.

Proposition 1.2. (1) *All Hopf algebras $(\mathbb{k}\mathbb{D}_\infty)^\bullet$, $T_\infty(n, v, \xi)^\bullet$, $B(n, \omega, \gamma)^\bullet$ and $D(m, d, \xi)^\bullet$ have GK-dimension one and are minimal under inclusion relation;*

(2) *As algebras, $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is regular while $T_\infty(n, v, \xi)^\bullet$, $B(n, \omega, \gamma)^\bullet$ and $D(m, d, \xi)^\bullet$ are not for $n, m \geq 2$.*

Proposition 1.2 is a direct consequence of Propositions 6.4 and 6.5. Through the pairing we constructed and this proposition, we get two observations: 1) For each affine prime regular Hopf algebra H of GK-dimension one, we get a quantum group in the sense of Takeuchi [31] naturally (see Remark 6.8 for details); 2) We give a counterexample to an infinite-dimensional analogous version of the semisimplicity result by Larson and Radford [15] which states that for a finite-dimensional Hopf algebra H , H is semisimple if and only if H^* is so.

After we finished the present paper, we found the nice preprint [4] written by K.A. Brown, M. Couto and A. Jahn. They refined the results in [12, 8] and determined

the structures of the finite dual of affine commutative-by-finite Hopf algebras. As the Hopf algebras of GK-dimension one considered in this paper are affine commutative-by-finite, the conclusions gotten in [4] can apply to our case directly. We remark that a number of formulas obtained in Section 3, 4 and 5 in our paper were essentially the same as those in [4, Section 7.5]. As pointed out in [4], the results in [4] and the present paper are largely complementary. Indeed, we can use our results to answer a question appeared in [4, Remarks 7.1(3)] (see Remark 5.1 in detail).

The organization of this paper is as follows: We gather in Section 2 necessary concepts and techniques, including the classification result for H , some equivalent definitions of H° for computation, and invertible matrices used for later proofs. Sections 3, 4 and 5 are devoted to giving the structures for $T_\infty(n, v, \xi)^\circ$, $B(n, \omega, \gamma)^\circ$ and $D(m, d, \xi)^\circ$ respectively. Finally in Section 6, constructions and related properties of desired non-degenerate Hopf pairings on $\mathbb{k}\mathbb{D}_\infty$, $T_\infty(n, v, \xi)$, $B(n, \omega, \gamma)$ and $D(m, d, \xi)$ are studied. We pose a question at the last section too.

1.1. Notation. We end this section by introducing some notations used in this paper. Let $\lfloor \frac{\cdot}{\cdot} \rfloor$ stand for the floor function, that is, for any natural numbers m, n , $\lfloor \frac{m}{n} \rfloor$ denotes the biggest integer which is not bigger than $\frac{m}{n}$.

Moreover, let n be a positive integer. We usually denote condition $0 \leq i \leq n-1$ simply by “ $i \in n$ ”. This can be referred to some convention in set theory that $n = \{0, 1, \dots, n-1\}$ as a set. Moreover, the cartesian product of $\{0, 1, \dots, m-1\}$ and $\{0, 1, \dots, n-1\}$ are denoted by $m \times n$, which is always ordered lexicographically.

Another notation which will be used is that: For an algebra H , we write $I \triangleleft_l H$ if I is a left ideal of H , and denote the principal left ideal generated by $h \in H$ by (h) .

2. PRELIMINARIES

In this section, we recall the classification result on affine prime regular Hopf algebras of GK-dimension one and give necessary tools and results for the determination of their finite duals. About general background knowledge, the reader is referred to [23, 29] for Hopf algebras and [20, 5] for exposition about noetherian Hopf algebras.

2.1. The Classification Result. At first, let us recall the classification of affine prime regular Hopf algebras of GK-dimension one given in [35, Theorem 8.3].

Lemma 2.1. *Any prime regular Hopf algebra of GK-dimension one must be isomorphic to one of the following:*

- (1) *Connected algebraic groups of dimension one: $\mathbb{k}[x]$ and $\mathbb{k}[x^{\pm 1}]$;*
- (2) *Infinite dihedral group algebra $\mathbb{k}\mathbb{D}_\infty$;*
- (3) *Infinite dimensional Taft algebras $T_\infty(n, v, \xi)$, where n, v are integers satisfying $0 \leq v \leq n-1$, and ξ is a primitive n th root of 1;*
- (4) *Generalized Liu’s algebras $B(n, \omega, \gamma)$, where n, ω are positive integers and γ is a primitive n th root of 1;*

- (5) The Hopf algebras $D(m, d, \xi)$, where m, d are positive integers satisfying $(1 + m)d$ is even and ξ is a primitive $2m$ th root of 1.

Detailed structures of $T_\infty(n, v, \xi)$, $B(n, \omega, \gamma)$ and $D(m, d, \xi)$ are recalled as follows:

Definition 2.2. Let n be a positive integer, $0 \leq v \leq n - 1$, and ξ be a primitive n th root of 1. As an algebra, $T_\infty(n, v, \xi)$ is generated by g and x with relations

$$g^n = 1, \quad xg = \xi gx.$$

Then $T_\infty(n, v, \xi)$ becomes a Hopf algebra with comultiplication, counit and antipode given by

$$\begin{aligned} \Delta(g) &= g \otimes g, \quad \Delta(x) = 1 \otimes x + x \otimes g^v, \quad \varepsilon(g) = 1, \quad \varepsilon(x) = 0, \\ S(g) &= g^{n-1}, \quad S(x) = -\xi^{-v} g^{n-v} x. \end{aligned}$$

Definition 2.3. Let n and ω be positive integers, and γ be a primitive n th root of 1. As an algebra, $B(n, \omega, \gamma)$ is generated by $x^{\pm 1}$, g and y with relations

$$\begin{cases} xx^{-1} = x^{-1}x = 1, & xg = gx, \quad xy = yx, \\ yg = \gamma gy, \\ y^n = 1 - x^\omega = 1 - g^n. \end{cases}$$

Then $B(n, \omega, \gamma)$ becomes a Hopf algebra with comultiplication, counit and antipode given by

$$\begin{aligned} \Delta(x) &= x \otimes x, \quad \Delta(g) = g \otimes g, \quad \Delta(y) = 1 \otimes y + y \otimes g, \\ \varepsilon(x) &= \varepsilon(g) = 1, \quad \varepsilon(y) = 0, \\ S(x) &= x^{-1}, \quad S(g) = g^{-1}, \quad S(y) = -\gamma^{-1} g^{-1} y. \end{aligned}$$

Definition 2.4. Let m, d be positive integers such that $(1 + m)d$ is even and ξ a primitive $2m$ th root of unity. Define

$$\omega := md, \quad \gamma := \xi^2.$$

As an algebra, $D(m, d, \xi)$ is generated by $x^{\pm 1}$, g , y and u_0, u_1, \dots, u_{m-1} with relations

$$\begin{aligned} &xx^{-1} = x^{-1}x = 1, \quad gx = xg, \quad yx = xy, \\ &yg = \gamma gy, \quad y^m = 1 - x^\omega = 1 - g^m, \\ &u_i x = x^{-1} u_i, \quad y u_i = \phi_i u_{i+1} = \xi x^d u_i y, \quad u_i g = \gamma^i x^{-2d} g u_i, \\ u_i u_j &= \begin{cases} (-1)^{-j} \xi^{-j} \gamma^{\frac{j(j+1)}{2}} \frac{1}{m} x^{-\frac{1+m}{2}d} \phi_i \phi_{i+1} \cdots \phi_{m-2-j} y^{i+j} g & (i+j \leq m-2) \\ (-1)^{-j} \xi^{-j} \gamma^{\frac{j(j+1)}{2}} \frac{1}{m} x^{-\frac{1+m}{2}d} y^{i+j} g & (i+j = m-1) \\ (-1)^{-j} \xi^{-j} \gamma^{\frac{j(j+1)}{2}} \frac{1}{m} x^{-\frac{1+m}{2}d} \phi_i \cdots \phi_{m-1} \phi_0 \cdots \phi_{m-2-j} y^{i+j-m} g & (i+j \geq m) \end{cases} \end{aligned}$$

where $\phi_i := 1 - \gamma^{-i-1} x^d$ and $i, j \in m$.

Then $D(m, d, \xi)$ becomes a Hopf algebra with comultiplication, counit and the antipode given by

$$\begin{aligned}\Delta(x) &= x \otimes x, \quad \Delta(g) = g \otimes g, \quad \Delta(y) = y \otimes g + 1 \otimes y, \\ \Delta(u_i) &= \sum_{j=0}^{m-1} \gamma^{j(i-j)} u_j \otimes x^{-jd} g^j u_{i-j}; \\ \varepsilon(x) &= \varepsilon(g) = \varepsilon(u_0) = 1, \quad \varepsilon(y) = \varepsilon(u_l) = 0, \\ S(x) &= x^{-1}, \quad S(g) = g^{-1}, \quad S(y) = -yg^{-1} = -\gamma^{-1}g^{-1}y, \\ S(u_i) &= (-1)^i \xi^{-i} \gamma^{-\frac{i(i+1)}{2}} x^{id + \frac{3}{2}(1-m)d} g^{m-i-1} u_i,\end{aligned}$$

for $i \in m$ and $1 \leq l \leq m-1$.

2.2. Finite Dual for Hopf algebras. In this subsection, we recall the concept of finite dual of a Hopf algebra, and list some lemmas in order to compute finite duals for the cases we are concerned with. The definition of finite dual is well-known and can be found in [29, Chapter 6] or [23, Chapter 9], for example.

Definition 2.5. Let H be a Hopf algebra over \mathbb{k} and denote its dual space by H^* . The finite dual of H is defined as

$$H^\circ := \{f \in H^* \mid f \text{ vanishes on a cofinite ideal } I \triangleleft H\}.$$

It is well-known that H° has a Hopf algebra structure naturally [23, Theorem 9.1.3]. We want to simplify H° a little. It is not hard to see that H° has the following alternative description: Let \mathcal{I} be the family of all the cofinite left ideals of H . Then by [23, Lemma 9.1.1], we have $H^\circ = \sum_{I \in \mathcal{I}} I^\perp$, where $I^\perp = \{f \in H^* \mid f(I) = 0\}$. Therefore, we have

Lemma 2.6. Let \mathcal{I}_0 be a subset of \mathcal{I} . Suppose for any $I \in \mathcal{I}$, there exist some $I_1, I_2, \dots, I_m \in \mathcal{I}_0$ such that $\bigcap_{i=1}^m I_i \subseteq I$. Then $H^\circ = \sum_{I \in \mathcal{I}_0} I^\perp$.

Proof. The condition implies that for any $I \in \mathcal{I}$, there exist some $I_1, I_2, \dots, I_m \in \mathcal{I}_0$ such that $\sum_{i=1}^m I_i^\perp = (\bigcap_{i=1}^m I_i)^\perp \supseteq I^\perp$. Thus

$$H^\circ = \sum_{I \in \mathcal{I}} I^\perp \subseteq \sum_{I \in \mathcal{I}_0} I^\perp \subseteq H^\circ,$$

and the claim is then verified. \square

Now we turn to a common feature of Hopf algebras in our cases. Specifically:

- For the remaining of this subsection, we assume that H is a Hopf algebra which is a finitely generated right module over some subalgebra P , such that P is a polynomial algebra $\mathbb{k}[x]$ or a Laurent polynomial algebra $\mathbb{k}[x^{\pm 1}]$.

We would note in subsequent sections that this assumption holds for all affine prime regular Hopf algebras of GK-dimension one. This assumption helps us to figure all the cofinite left ideals of H :

Lemma 2.7. For any left ideal $I \triangleleft_l H$, I is cofinite in H if and only if $I \cap P \neq 0$.

Proof. Note that whenever P is $\mathbb{k}[x]$ or $\mathbb{k}[x^{\pm 1}]$, $I \cap P$ is an ideal of P and thus a principal ideal generated by some polynomial $q(x) \in P$. It follows that

$$I \cap P \neq 0 \iff q(x) \neq 0 \iff I \cap P \text{ is cofinite in } P.$$

If I is cofinite in H , then the following relations of \mathbb{k} -vector spaces

$$H/I \supseteq (P + I)/I \cong P/(I \cap P)$$

implies that $P/(I \cap P)$ must be finite-dimensional. In other words, $I \cap P \neq 0$ holds.

Conversely if $I \cap P \neq 0$, then clearly $P/(I \cap P)$ is finite-dimensional, and so is $(P + I)/I$. However, the assumption above this lemma implies that $H = \sum_{i=1}^m h_i P$ for some $h_1, h_2, \dots, h_m \in H$. As I is a left ideal of H , we can know that $(P + I)/I$ is a finite-dimensional subspace of the quotient left H -module H/I , and

$$H/I = \sum_{i=1}^m h_i (P + I/I)$$

holds. It follows that H/I is also finite-dimensional. \square

Combining Lemmas 2.6 and 2.7, in order to compute H° , it is sufficient to find enough appropriate polynomials $p(x)$'s, such that each $I \in \mathcal{I}$ must contain some principal left ideal $(p(x))$. Concerning our cases of $T_\infty(n, v, \xi)$, $B(n, \omega, \gamma)$ and $D(m, d, \xi)$, we can choose such $p(x)$'s respectively as follows.

Lemma 2.8. *Let H be an example from any of the above three classes of Hopf algebras. Let $I \in \mathcal{I}$, and m, n be positive integers.*

- (1) *There is a polynomial $p(x) \in P = \mathbb{k}[x]$ with form:*

$$p(x) = \prod_{\alpha=1}^N (x^m - \lambda_\alpha)^{nr_\alpha} \quad (2.1)$$

for some $N \geq 1$, $r_\alpha \in \mathbb{N}$ and distinct $\lambda_\alpha \in \mathbb{k}$, such that I contains the cofinite left ideal generated by $p(x)$ in H ;

- (2) *There is a polynomial $p(x) \in P = \mathbb{k}[x^{\pm 1}]$ with form:*

$$p(x) = \prod_{\alpha=1}^N (x^n - \lambda_\alpha)^{r_\alpha} \quad (2.2)$$

for some $N \geq 1$, $r_\alpha \in \mathbb{N}$ and distinct $\lambda_\alpha \in \mathbb{k}^$, such that I contains the cofinite left ideal generated by $p(x)$ in H ;*

- (3) *There is a polynomial $p(x) \in P = \mathbb{k}[x^{\pm 1}]$ with form:*

$$p(x) = (x^m - 1)^{r'} (x^m + 1)^{r''} \prod_{\alpha=1}^N (x^m - \lambda_\alpha)^{r_\alpha} (x^m - \lambda_\alpha^{-1})^{r_\alpha} \quad (2.3)$$

for some $N \geq 1$, $r', r'', r_\alpha \in \mathbb{N}$, and $\lambda_\alpha \in \mathbb{k}^ \setminus \{\pm 1\}$ which are distinct and not inverses of each other, such that I contains the cofinite left ideal generated by $p(x)$ in H .*

Proof. We prove (2) for example, since (1) and (3) hold according to the same reason. It can be known from the proof of Lemma 2.7 that $I \cap P$ is a principal ideal of P generated by some non-zero polynomial $q(x)$. Without loss of generality, assume that

$$q(x) = (x - \mu_1)(x - \mu_2) \cdots (x - \mu_M)$$

for some $M \geq 1$ and $\mu_1, \mu_2, \dots, \mu_M \in \mathbb{k}^*$. Now define a polynomial

$$p(x) = (x^n - \mu_1^n)(x^n - \mu_2^n) \cdots (x^n - \mu_M^n),$$

which gives the desired form (2.2). Since $q(x) \mid p(x)$, I contains the left ideal of H generated by $p(x)$ in H . \square

This lemma suggests us to consider the following subfamilies $\mathcal{I}_0 \subseteq \mathcal{I}$ respectively.

Corollary 2.9. *Let m, n be positive integers.*

(1) *For $P = \mathbb{k}[x]$, denote*

$$\mathcal{I}_0 = \{((x^m - \lambda)^{nr}) \triangleleft_l H \mid \lambda \in \mathbb{k}, r \in \mathbb{N}\}.$$

If \mathcal{I}_0 satisfies the condition in Lemma 2.6, then $H^\circ = \sum_{I \in \mathcal{I}_0} (H/I)^$;*

(2) *For $P = \mathbb{k}[x^{\pm 1}]$, denote*

$$\mathcal{I}_0 = \{((x^n - \lambda)^r) \triangleleft_l H \mid \lambda \in \mathbb{k}^*, r \in \mathbb{N}\}.$$

If \mathcal{I}_0 satisfies the condition in Lemma 2.6, then $H^\circ = \sum_{I \in \mathcal{I}_0} (H/I)^$;*

(3) *For $P = \mathbb{k}[x^{\pm 1}]$, denote*

$$\mathcal{I}_0 = \{((x^m - 1)^r), ((x^m + 1)^r), ((x^m - \lambda)^r(x^m - \lambda^{-1})^r) \triangleleft_l H \mid \lambda \in \mathbb{k}^* \setminus \{\pm 1\}, r \in \mathbb{N}\}.$$

If \mathcal{I}_0 satisfies the condition in Lemma 2.6, then $H^\circ = \sum_{I \in \mathcal{I}_0} (H/I)^$.*

We end this subsection by providing the following lemma on the intersection of principal left ideals in \mathcal{I} :

Lemma 2.10. *Let N be a positive integer, and let P be $\mathbb{k}[x]$ or $\mathbb{k}[x^{\pm 1}]$. Suppose that $q_1(x), q_2(x), \dots, q_N(x) \in P$ are coprime with each other. Then*

$$\bigcap_{\alpha=1}^N (q_\alpha(x)) = \left(\prod_{\alpha=1}^N q_\alpha(x) \right)$$

as left ideals in H .

Proof. It is sufficient to prove the case when $N = 2$. First there exist $p_1(x), p_2(x) \in P$ such that

$$q_1(x)p_1(x) + q_2(x)p_2(x) = 1.$$

Now for any $h \in (q_1(x)) \cap (q_2(x))$, we have $h = h_1 q_1(x) = h_2 q_2(x)$ for some $h_1, h_2 \in H$. Then

$$\begin{aligned} h_2 &= h_2(q_1(x)p_1(x) + q_2(x)p_2(x)) = h_2 q_1(x)p_1(x) + h_1 q_1(x)p_2(x) \\ &= (h_2 p_1(x) + h_1 p_2(x))q_1(x), \end{aligned}$$

and thus $h = h_2 q_2(x) = (h_2 p_1(x) + h_1 p_2(x))q_1(x)q_2(x) \in (q_1(x)q_2(x))$. \square

2.3. Invertibility of Matrices. The following primary fact on the Kronecker products is required, which is well-known in matrix theory.

Lemma 2.11. (1) *Let Γ_1 and Γ_2 be finite totally ordered sets, and their Cartesian product $\Gamma_1 \times \Gamma_2$ be ordered lexicographically. Then for any two square matrices*

$$A = (a_{i,i'} \mid i, i' \in \Gamma_1) \quad \text{and} \quad B = (b_{j,j'} \mid j, j' \in \Gamma_2)$$

over \mathbb{k} , A and B are both invertible if and only if their Kronecker product

$$(a_{i,i'} b_{j,j'} \mid (i, j), (i', j') \in \Gamma_1 \times \Gamma_2)$$

is invertible;

(2) *Let n_1, n_2, \dots, n_m be positive integers. Then the square matrices*

$$A_k = (a_{i_k, i'_k}^{(k)} \mid i_k, i'_k \in n_k) \quad (1 \leq k \leq m)$$

over \mathbb{k} are all invertible, if and only if their Kronecker product

$$(a_{i_1, i'_1}^{(1)} a_{i_2, i'_2}^{(2)} \cdots a_{i_m, i'_m}^{(m)} \mid (i_1, i_2, \dots, i_m), (i'_1, i'_2, \dots, i'_m) \in n_1 \times n_2 \times \cdots \times n_m)$$

is invertible.

A generalized version of above lemma is also required, and we provide a proof for convenience.

Lemma 2.12. *Let Γ_1 and Γ_2 be finite totally ordered sets, and their Cartesian product $\Gamma_1 \times \Gamma_2$ be ordered lexicographically. Suppose $A := (a_{i,i'} \mid i, i' \in \Gamma_1)$ is invertible. Then the following matrix*

$$C := (a_{i,i'} b_{i',j,j'} \mid (i, j), (i', j') \in \Gamma_1 \times \Gamma_2)$$

is invertible, if and only if $B_{i'} := (b_{i',j,j'} \mid j, j' \in \Gamma_2)$ is invertible for each $i' \in \Gamma_1$.

Proof. For clarity, denote the cardinalities of sets by $m := |\Gamma_1|$ and $n := |\Gamma_2|$. Clearly, C can be divided into blocks of order m :

$$C = (a_{i,i'} Y_{i'} \mid i, i' \in \Gamma_1).$$

Denote $A^{-1} := (\tilde{a}_{i,i'} \mid i, i' \in \Gamma_1)$, then Kronecker product of A^{-1} and I_n (the identity matrix of order n) can be written as a block matrix

$$(\tilde{a}_{i,i'} I_n \mid i, i' \in \Gamma_1), \tag{2.4}$$

which is invertible according to Lemma 2.11. One can directly verify that the product of (2.4) and C is equal to

$$\left(\sum_{k \in \Gamma_1} \tilde{a}_{i,k} a_{k,i'} B_{i'} \mid i, i' \in \Gamma_1 \right) = (\delta_{i,i'} B_{i'} \mid i, i' \in \Gamma_1),$$

and the claim is followed. \square

Within the work of [9], they proved the invertibility of a certain form of matrices. This plays a crucial role on the process of constructing $(\mathbb{k}\mathbb{D}_\infty)^\circ$. Our construction for $D(m, d, \xi)^\circ$ in this paper would also rely on this fact. However, we provide here another form of that result for simplicity.

Lemma 2.13. ([9, Corollary 2]) *Let r be a positive integer, and $\lambda \in \mathbb{k}^* \setminus \{\pm 1\}$. The $2r \times 2r$ matrix*

$$\left(\lambda^{(-1)^e(2s'+e')} (2s' + e')^s \mid (e, s), (e', s') \in 2 \times r \right) \quad (2.5)$$

is invertible.

Proof. Denote $M := 2r - 1$. It follows from [9, Corollary 2] that the $2r \times 2r$ matrix

$$\begin{pmatrix} 1 & 0 & \cdots & 0 & 1 & 0 & \cdots & 0 \\ \lambda & \lambda & \cdots & \lambda & \lambda^{-1} & \lambda^{-1} & \cdots & \lambda^{-1} \\ \lambda^2 & 2\lambda^2 & \cdots & 2^{r-1}\lambda^2 & \lambda^{-2} & 2\lambda^{-2} & \cdots & 2^{r-1}\lambda^{-2} \\ \lambda^3 & 3\lambda^3 & \cdots & 3^{r-1}\lambda^3 & \lambda^{-3} & 3\lambda^{-3} & \cdots & 3^{r-1}\lambda^{-3} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda^M & M\lambda^M & \cdots & M^{r-1}\lambda^M & \lambda^{-M} & M\lambda^{-M} & \cdots & M^{r-1}\lambda^{-M} \end{pmatrix} \quad (2.6)$$

is invertible, which can be obtained from the transpose of (2.5) through elementary operations of type 2. Thus (2.5) is also invertible. \square

This result has a slightly generalized version, which is required in this paper as well:

Corollary 2.14. *Let r be a positive integer, and $\lambda \in \mathbb{k}^* \setminus \{\pm 1\}$. Then:*

(1) *The $2r \times 2r$ matrix*

$$\left(\lambda^{(-1)^e(2s'+e')} (a + 2s' + e')^s \mid (e, s), (e', s') \in 2 \times r \right) \quad (2.7)$$

is invertible for any $a \in \mathbb{k}$;

(2) *The $2r \times 2r$ matrix*

$$\left(\lambda^{(-1)^e(b+2s'+e')} (a + 2s' + e')^s \mid (e, s), (e', s') \in 2 \times r \right) \quad (2.8)$$

is invertible for any $a \in \mathbb{k}$ and $b \in \mathbb{Q}$.

Proof. (1) Denote $M := 2r - 1$. Firstly, (2.6) can be obtained from the following matrix through elementary column operations of type 3:

$$\begin{pmatrix} 1 & a & \cdots & a^{r-1} & 1 & a & \cdots & a^{r-1} \\ \lambda & (a+1)\lambda & \cdots & (a+1)^{r-1}\lambda & \lambda^{-1} & (a+1)\lambda^{-1} & \cdots & (a+1)^{r-1}\lambda^{-1} \\ \lambda^2 & (a+2)\lambda^2 & \cdots & (a+2)^{r-1}\lambda^2 & \lambda^{-2} & (a+2)\lambda^{-2} & \cdots & (a+2)^{r-1}\lambda^{-2} \\ \lambda^3 & (a+3)\lambda^3 & \cdots & (a+3)^{r-1}\lambda^3 & \lambda^{-3} & (a+3)\lambda^{-3} & \cdots & (a+3)^{r-1}\lambda^{-3} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda^M & (a+M)\lambda^M & \cdots & (a+M)^{r-1}\lambda^M & \lambda^{-M} & (a+M)\lambda^{-M} & \cdots & (a+M)^{r-1}\lambda^{-M} \end{pmatrix}. \quad (2.9)$$

The process is by deleting powers of λ or λ^{-1} with coefficients containing a , from the r th column to the 2nd column, and the $2r$ th column to the $(r+1)$ th column. Thus (2.9) is also invertible.

However, (2.7) can be obtained from the transpose of (2.9) through elementary operations of type 2. This implies the invertibility of (2.7).

(2) The similar argument follows that (2.8) can be obtained from the transpose of the following matrix through elementary column operations:

$$\begin{pmatrix} \lambda^b & a\lambda^b & \cdots & a^{r-1}\lambda^b & \lambda^{-b} & a\lambda^{-b} & \cdots & a^{r-1}\lambda^{-b} \\ \lambda^{b+1} & (a+1)\lambda^{b+1} & \cdots & (a+1)^{r-1}\lambda^{b+1} & \lambda^{-(b+1)} & (a+1)\lambda^{-(b+1)} & \cdots & (a+1)^{r-1}\lambda^{-(b+1)} \\ \lambda^{b+2} & (a+2)\lambda^{b+2} & \cdots & (a+2)^{r-1}\lambda^{b+2} & \lambda^{-(b+2)} & (a+2)\lambda^{-(b+2)} & \cdots & (a+2)^{r-1}\lambda^{-(b+2)} \\ \lambda^{b+3} & (a+3)\lambda^{b+3} & \cdots & (a+3)^{r-1}\lambda^{b+3} & \lambda^{-(b+3)} & (a+3)\lambda^{-(b+3)} & \cdots & (a+3)^{r-1}\lambda^{-(b+3)} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \lambda^{b+M} & (a+M)\lambda^{b+M} & \cdots & (a+M)^{r-1}\lambda^{b+M} & \lambda^{-(b+M)} & (a+M)\lambda^{-(b+M)} & \cdots & (a+M)^{r-1}\lambda^{-(b+M)} \end{pmatrix}$$

and this matrix has the same determinant as (2.9). \square

2.4. Combinatorial Notions. We end this section by listing two other concepts for later use. The one is the well-known *quantum binomial coefficients* for a parameter $q \in \mathbb{k}^*$, which is defined as

$$\binom{l}{k}_q := \frac{l!_q}{k!_q(l-k)!_q}$$

for integers $l \geq k \geq 0$, where $l!_q := 1_q 2_q \cdots l_q$ and $l_q := 1 + q + \cdots + q^{l-1}$.

The other one is a part of the *second Stirling number* (see (13.13) in [16]), written as

$$\sum_{t=0}^r \binom{r}{t} (-1)^{r-t} t^s, \quad (2.10)$$

which is in fact zero if $0 \leq s < r$.

3. THE FINITE DUAL OF INFINITE-DIMENSIONAL TAFT ALGEBRA $T_\infty(n, v, \xi)$

This section is devoted to providing the structure of $T_\infty(n, v, \xi)^\circ$, which is isomorphic to the Hopf algebra $T_{\infty^\circ}(n, v, \xi)$ constructed by generators and relations in Subsection 3.1. For the purpose, we choose certain elements in $T_\infty(n, v, \xi)^\circ$ and study their properties in Subsection 3.2. The desired isomorphism is shown in Subsection 3.3 finally.

3.1. The Hopf Algebra $T_{\infty^\circ}(n, v, \xi)$. Let n be a positive integer, $0 \leq v \leq n-1$ and ξ be a primitive n th root of 1. For simplicity we denote $m := \frac{n}{\gcd(n, v)}$. The Hopf algebra $T_{\infty^\circ}(n, v, \xi)$ is constructed as follows. As an algebra, it is generated by Ψ_λ ($\lambda \in \mathbb{k}$), Ω , F_1 , F_2 with relations

$$\begin{aligned} \Psi_{\lambda_1} \Psi_{\lambda_2} &= \Psi_{\lambda_1 + \lambda_2}, \quad \Psi_0 = 1, \quad \Omega^n = 1, \quad F_1^m = 0, \\ \Omega \Psi_\lambda &= \Psi_\lambda \Omega, \quad F_2 \Omega = \Omega F_2, \quad F_1 \Omega = \xi^v \Omega F_1, \\ F_2 \Psi_\lambda &= \Psi_\lambda F_2, \quad F_1 \Psi_\lambda = \Psi_\lambda F_1, \quad F_1 F_2 = F_2 F_1 \end{aligned}$$

for all $\lambda, \lambda_1, \lambda_2 \in \mathbb{k}$. The comultiplication, counit and antipode are given by

$$\begin{aligned} \Delta(\Omega) &= \Omega \otimes \Omega, \quad \Delta(F_1) = 1 \otimes F_1 + F_1 \otimes \Omega, \\ \Delta(F_2) &= 1 \otimes F_2 + F_2 \otimes \Omega^m + \sum_{k=1}^{m-1} F_1^{[k]} \otimes \Omega^k F_1^{[m-k]}, \end{aligned}$$

$$\Delta(\Psi_\lambda) = \sum_{c=0}^{(n/m)-1} (\Psi_{\lambda\xi^{mc}} \otimes \Psi_{\lambda\varsigma_c})(1 \otimes 1 + \lambda \sum_{k=1}^{m-1} F_1^{[k]} \otimes \Omega^k F_1^{[m-k]}),$$

$$\varepsilon(\Omega) = \varepsilon(\Psi_\lambda) = 1, \quad \varepsilon(F_1) = \varepsilon(F_2) = 0,$$

$$S(\Omega) = \Omega^{n-1}, \quad S(F_1) = -\xi^{-v}\Omega^{n-1}F_1, \quad S(F_2) = -F_2, \quad S(\Psi_\lambda) = \sum_{c=0}^{(n/m)-1} \Psi_{-\lambda\xi^{-mc}\varsigma_c},$$

for $\lambda \in \mathbb{k}$, where $F_1^{[k]} := \frac{1}{k!\xi^v} F_1^k$ for $1 \leq k \leq m-1$, and

$$\varsigma_c := \frac{m}{n}(1 + \xi^{-mc}\Omega^m + \xi^{-2mc}\Omega^{2m} + \dots + \xi^{-(n-m)c}\Omega^{n-m}) \quad (c \in n/m).$$

The fact that $T_{\infty^\circ}(n, v, \xi)$ is a Hopf algebra would be stated and proved in Subsection 3.3 as Theorem 3.4(1). Here we note that

$$\{\Psi_\lambda \Omega^j F_2^s F_1^l \mid \lambda \in \mathbb{k}, j \in n, s \in \mathbb{N}, l \in m\}$$

is a linear basis, due to an application of the Diamond Lemma [2].

3.2. Certain Elements in $T_{\infty}(n, v, \xi)^\circ$. Let n be a positive integer, $0 \leq v \leq n-1$, and ξ be a primitive n th root of 1. According to the definition of $T_{\infty}(n, v, \xi)$ recalled in Definition 2.2, we know that it has a linear basis $\{g^j x^l \mid j \in n, l \in \mathbb{N}\}$. Denote $m := \frac{n}{\gcd(n, v)}$, and then ξ^v is a primitive m th root of 1. We define following elements in $T_{\infty}(n, v, \xi)^*$:

$$\psi_\lambda : g^j x^l \mapsto \sum_{u=0}^{\infty} \delta_{l, um} \lambda^u, \quad \omega : g^j x^l \mapsto \delta_{l, 0} \xi^j, \quad E_1 : g^j x^l \mapsto \delta_{l, 1}, \quad E_2 : g^j x^l \mapsto \delta_{l, m} \quad (3.1)$$

for any $j \in n, l \in \mathbb{N}$ and $\lambda \in \mathbb{k}$. We remark that these definitions make sense for all $j \in \mathbb{Z}$ as well due to a direct computation.

One can verify that ψ_λ vanishes on the principal left ideal $(x^m - \lambda)$ for any $\lambda \in \mathbb{k}$; ω vanishes on (x) ; E_2 vanishes on (x^{m+1}) ; E_1 vanishes on (x^2) . Clearly, these left ideals are cofinite in $T_{\infty}(n, v, \xi)$, and hence all the elements in (3.1) lie in $T_{\infty}(n, v, \xi)^\circ$.

Lemma 3.1. *Following equations hold in $T_{\infty}(n, v, \xi)^\circ$:*

$$\begin{aligned} \psi_{\lambda_1} \psi_{\lambda_2} &= \psi_{\lambda_1 + \lambda_2}, \quad \psi_0 = 1, \quad \omega^n = 1, \quad E_1^m = 0, \\ \omega \psi_\lambda &= \psi_\lambda \omega, \quad E_2 \omega = \omega E_2, \quad E_1 \omega = \xi^v \omega E_1, \\ E_2 \psi_\lambda &= \psi_\lambda E_2, \quad E_1 \psi_\lambda = \psi_\lambda E_1, \quad E_1 E_2 = E_2 E_1 \end{aligned}$$

for all $\lambda, \lambda_1, \lambda_2 \in \mathbb{k}$.

Proof. Note that

$$\Delta(g^j x^l) = (g \otimes g)^j (1 \otimes x + x \otimes g^v)^l = \sum_{k=0}^l \binom{l}{k}_{\xi^v} g^j x^k \otimes g^{j+kv} x^{l-k}.$$

We prove the lemma by checking values at the basis elements $g^j x^l$ ($j \in n$, $l \in \mathbb{N}$):

$$\begin{aligned}
\langle \psi_{\lambda_1} \psi_{\lambda_2}, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l,um} \sum_{i=0}^u \binom{um}{im}_{\xi^v} \langle \psi_{\lambda_1}, g^j x^{im} \rangle \langle \psi_{\lambda_2}, g^{j+imv} x^{(u-i)m} \rangle \\
&= \sum_{u=0}^{\infty} \delta_{l,um} \sum_{i=0}^u \binom{u}{i} \lambda_1^i \lambda_2^{u-i} = \langle \psi_{\lambda_1+\lambda_2}, g^j x^l \rangle, \\
\langle \psi_0, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l,um} 0^u = \delta_{l,0} = \langle \varepsilon, g^j x^l \rangle, \\
\langle \omega \psi_{\lambda}, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l,um} \langle \omega, g^j \rangle \langle \psi_{\lambda}, g^j x^{um} \rangle \\
&= \sum_{u=0}^{\infty} \delta_{l,um} \langle \psi_{\lambda}, g^j x^{um} \rangle \langle \omega, g^{j+umv} \rangle = \langle \psi_{\lambda} \omega, g^j x^l \rangle, \\
\langle E_2 \omega, g^j x^l \rangle &= \delta_{l,m} \langle E_2, g^j x^m \rangle \langle \omega, g^{j+mv} \rangle = \delta_{l,m} \langle \omega, g^j \rangle \langle E_2, g^j x^m \rangle = \langle \omega E_2, g^j x^l \rangle, \\
\langle E_1 \omega, g^j x^l \rangle &= \delta_{l,1} \langle E_1, g^j x \rangle \langle \omega, g^{j+v} \rangle = \delta_{l,1} \xi^v \langle \omega, g^j \rangle \langle E_1, g^j x \rangle = \langle \xi^v \omega E_1, g^j x^l \rangle, \\
\langle E_2 \psi_{\lambda}, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l,(u+1)m} \binom{(u+1)m}{m}_{\xi^v} \langle E_2, g^j x^m \rangle \langle \psi_{\lambda}, g^{j+mv} x^{um} \rangle \\
&= \sum_{u=0}^{\infty} \delta_{l,(u+1)m} \binom{(u+1)m}{um}_{\xi^v} \langle \psi_{\lambda}, g^j x^{um} \rangle \langle E_2, g^{j+umv} x^m \rangle = \langle \psi_{\lambda} E_2, g^j x^l \rangle, \\
\langle E_1 \psi_{\lambda}, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l,um+1} \binom{um+1}{1}_{\xi^v} \langle E_1, g^j x \rangle \langle \psi_{\lambda}, g^{j+v} x^{um} \rangle \\
&= \sum_{u=0}^{\infty} \delta_{l,um+1} \binom{um+1}{um}_{\xi^v} \langle \psi_{\lambda}, g^j x^{um} \rangle \langle E_1, g^{j+umv} x \rangle = \langle \psi_{\lambda} E_1, g^j x^l \rangle.
\end{aligned}$$

Also, we find by induction on $k \in n$ and $k' \in m$ that

$$\begin{aligned}
\langle \omega^k, g^j x^l \rangle &= \delta_{l,0} \langle \omega^k, g^j \rangle = \delta_{l,0} \langle \omega, g^j \rangle^k = \delta_{l,0} \xi^{jk}, \\
\langle E_1^{k'}, g^j x^l \rangle &= \delta_{l,k'} \binom{k'}{1}_{\xi^v} \langle E_1, g^j x \rangle \langle E_1^{k'-1}, g^{j+v} x^{k'-1} \rangle \\
&= \delta_{l,k'} \frac{k'!_{\xi^v}}{(k'-1)!_{\xi^v}} \langle E_1^{k'-1}, g^{j+v} x^{k'-1} \rangle \\
&= \delta_{l,k'} \frac{k'!_{\xi^v}}{(k'-1)!_{\xi^v}} \frac{(k'-1)!_{\xi^v}}{(k'-2)!_{\xi^v}} \langle E_1^{k'-2}, g^{j+2v} x^{k'-2} \rangle = \dots = \delta_{l,k'} k'!_{\xi^v}.
\end{aligned}$$

Thus $\omega^n = 1$ and $E_1^m = 0$ hold. \square

Furthermore if denote $E_1^{[k']} := \frac{1}{k'!_{\xi^v}} E_1^{k'}$, we verify that for any $k' \in m$, $k, j \in n$ and $l \in \mathbb{N}$,

$$\langle \omega^k E_1^{[k]}, g^j x^l \rangle = \delta_{l,k'} \langle \omega^k, g^j \rangle \langle E_1^{[k]}, g^j x^{k'} \rangle = \delta_{l,k'} \xi^{jk},$$

$$\begin{aligned}
\langle \psi_\lambda \omega^k E_1^{[k]}, g^j x^l \rangle &= \sum_{u=0}^{\infty} \delta_{l, um+k'} \binom{um+k'}{um}_{\xi^v} \langle \psi_\lambda, g^j x^{um} \rangle \langle \omega^k E_1^{[k]}, g^{j+umv} x^{k'} \rangle \\
&= \sum_{u=0}^{\infty} \delta_{l, um+k'} \lambda^u \xi^{jk},
\end{aligned}$$

If we denote $E_2^{[s]} := \frac{1}{s!} E_2^s$ for $s \geq 1$, followings can be obtained by induction:

$$\begin{aligned}
\langle E_2^s, g^j x^l \rangle &= \delta_{l, sm} \binom{sm}{m}_{\xi^v} \langle E_2, g^j x^m \rangle \langle E_2^{s-1}, g^{j+mv} x^{(s-1)m} \rangle \\
&= \delta_{l, sm} s \langle E_2^{s-1}, g^{j+mv} x^{(s-1)m} \rangle \\
&= \delta_{l, sm} s \binom{(s-1)m}{m}_{\xi^v} \langle E_2, g^{j+mv} x^m \rangle \langle E_2^{s-2}, g^{j+2mv} x^{(s-2)m} \rangle \\
&= \delta_{l, sm} s(s-1) \langle E_2^{s-2}, g^{j+2mv} x^{(s-2)m} \rangle = \dots = \delta_{l, sm} s!, \\
\langle E_2^{[s]}, g^j x^l \rangle &= \delta_{l, sn}, \\
\langle E_2^{[s]} E_1^{[k]}, g^j x^l \rangle &= \delta_{l, sm+k'} \binom{sm+k'}{sm}_{\xi^v} \langle E_2^{[s]}, g^j x^{sm} \rangle \langle E_1^{[k]}, g^{j+smv} x^{k'} \rangle = \delta_{l, sm+k'}, \\
\langle \omega^k E_2^{[s]} E_1^{[k]}, g^j x^l \rangle &= \delta_{l, sm+k'} \langle \omega^k, g^j \rangle \langle E_2^{[s]} E_1^{[k]}, g^j x^{sm+k'} \rangle = \delta_{l, sm+k'} \xi^{jk}.
\end{aligned}$$

Lemma 3.2. *Following equations hold in $T_\infty(n, v, \xi)^\circ$:*

$$\Delta(\omega) = \omega \otimes \omega, \quad \Delta(E_1) = 1 \otimes E_1 + E_1 \otimes \omega,$$

$$\Delta(E_2) = 1 \otimes E_2 + E_2 \otimes \omega^m + \sum_{k=1}^{m-1} E_1^{[k]} \otimes \omega^k E_1^{[m-k]},$$

$$\Delta(\psi_\lambda) = \sum_{c=0}^{(n/m)-1} (\psi_\lambda \xi^{mc} \otimes \psi_\lambda \sigma_c) (1 \otimes 1 + \lambda \sum_{k=1}^{m-1} E_1^{[k]} \otimes \omega^k E_1^{[m-k]}),$$

$$\varepsilon(\omega) = \varepsilon(\psi_\lambda) = 1, \quad \varepsilon(E_1) = \varepsilon(E_2) = 0,$$

$$S(\omega) = \omega^{n-1}, \quad S(E_1) = -\xi^{-v} \omega^{n-1} E_1, \quad S(E_2) = -E_2, \quad S(\psi_\lambda) = \sum_{c=0}^{(n/m)-1} \psi_{-\lambda \xi^{-mc} \sigma_c},$$

for $\lambda \in \mathbb{k}$, where $E_1^{[k]} := \frac{1}{k! \xi^v} E_1^k$ for $1 \leq k \leq m-1$, and

$$\sigma_c := \frac{m}{n} (1 + \xi^{-mc} \omega^m + \xi^{-2mc} \omega^{2m} + \dots + \xi^{-(n-m)c} \omega^{n-m}) \quad (c \in n/m).$$

Proof. Note that $(g^j x^l)(g^{j'} x^{l'}) = \xi^{j'l} g^{j+j'} x^{l+l'}$ ($j, j' \in n, l, l' \in \mathbb{N}$). We also prove the lemma by checking values at each $g^j x^l \otimes g^{j'} x^{l'}$.

$$\begin{aligned}
\langle \omega, \xi^{j'l} g^{j+j'} x^{l+l'} \rangle &= \delta_{l+l', 0} \xi^{j'l+j+j'} = \delta_{l, 0} \xi^j \delta_{l', 0} \xi^{j'} = \langle \omega \otimes \omega, g^j x^l \otimes g^{j'} x^{l'} \rangle, \\
\langle E_1, \xi^{j'l} g^{j+j'} x^{l+l'} \rangle &= \delta_{l+l', 1} \xi^{j'l} = \delta_{l, 0} \delta_{l', 1} + \delta_{l, 1} \delta_{l', 0} \xi^{j'} \\
&= \langle 1 \otimes E_1 + E_1 \otimes \omega, g^j x^l \otimes g^{j'} x^{l'} \rangle,
\end{aligned}$$

$$\begin{aligned}
\langle E_2, \xi^{j'l} g^{j+j'} x^{l+l'} \rangle &= \delta_{l+l',m} \xi^{j'l} = \delta_{l,0} \delta_{l',m} + \delta_{l,m} \delta_{l',0} \xi^{j'm} + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \xi^{j'k} \\
&= \langle 1 \otimes E_2 + E_2 \otimes \omega^m + \sum_{k=1}^{m-1} E_1^{[k]} \otimes \omega^k E_1^{[m-k]}, g^j x^l \otimes g^{j'} x^{l'} \rangle.
\end{aligned}$$

It is not hard to find that for each $c \in n/m$, $j \in n$ and $l \in \mathbb{N}$:

$$\langle \psi_\lambda \sigma_c, g^j x^l \rangle = \begin{cases} \sum_{u=0}^{\infty} \delta_{l,um} \lambda^u, & \text{if } j \equiv c \pmod{n/m}; \\ 0, & \text{otherwise.} \end{cases}$$

Now compute that

$$\begin{aligned}
\langle \psi_\lambda, \xi^{j'l} g^{j+j'} x^{l+l'} \rangle &= \sum_{u=0}^{\infty} \delta_{l+l',um} \lambda^u \xi^{j'l} \\
&= \sum_{u,u'=0}^{\infty} (\delta_{l,um} \delta_{l',u'm} \lambda^{u+u'} \xi^{j'um} + \sum_{k=1}^{m-1} \delta_{l,um+k} \delta_{l',u'm+(m-k)} \lambda^{u+u'+1} \xi^{j'(um+k)}) \\
&= \sum_{u=0}^{\infty} \delta_{l,um} (\lambda \xi^{j'm})^u \sum_{u'=0}^{\infty} \delta_{l',u'm} \lambda^{u'} + \lambda \sum_{k=1}^{m-1} \sum_{u=0}^{\infty} \delta_{l,um+k} (\lambda \xi^{j'm})^u \sum_{u'=0}^{\infty} \delta_{l',u'm+(m-k)} \lambda^{u'} \xi^{j'k} \\
&= \langle \psi_{\lambda \xi^{j'm}} \otimes \psi_\lambda + \lambda \sum_{k=1}^{m-1} \psi_{\lambda \xi^{j'm}} E_1^{[k]} \otimes \psi_\lambda \omega^k E_1^{[m-k]}, g^j x^l \otimes g^{j'} x^{l'} \rangle \\
&= \langle \sum_{c=0}^{(n/m)-1} (\psi_{\lambda \xi^{mc}} \otimes \psi_\lambda \sigma_c + \lambda \sum_{k=1}^{m-1} \psi_{\lambda \xi^{mc}} E_1^{[k]} \otimes \psi_\lambda \sigma_c \omega^k E_1^{[m-k]}), g^j x^l \otimes g^{j'} x^{l'} \rangle \\
&= \langle \sum_{c=0}^{(n/m)-1} (\psi_{\lambda \xi^{mc}} \otimes \psi_\lambda \sigma_c) (1 \otimes 1 + \lambda \sum_{k=1}^{m-1} E_1^{[k]} \otimes \omega^k E_1^{[m-k]}), g^j x^l \otimes g^{j'} x^{l'} \rangle.
\end{aligned}$$

The expressions of the counit and the antipode are clear. \square

3.3. The Generation Problem.

Proposition 3.3. *As an algebra, $T_\infty(n, v, \xi)^\circ$ is generated by ψ_λ , ω , E_2 and E_1 for $\lambda \in \mathbb{k}$.*

Proof. Clearly $T_\infty(n, v, \xi)$ has a polynomial subalgebra $P = \mathbb{k}[x]$. Choose

$$\mathcal{I}_0 = \{((x^m - \lambda)^{nr}) \triangleleft_l T_\infty(n, v, \xi) \mid \lambda \in \mathbb{k}, r \in \mathbb{N}\}.$$

First for any cofinite left ideal I of $T_\infty(n, v, \xi)$, according to Lemma 2.8(1), there must be some non-zero polynomial

$$p(x) = \prod_{\alpha=1}^N (x^m - \lambda_\alpha)^{nr_\alpha}$$

for some $N \geq 1$, $r_\alpha \in \mathbb{N}$ and distinct $\lambda_\alpha \in \mathbb{k}$, which generates a cofinite left ideal contained in I . It can be known by Lemma 2.10 that

$$\bigcap_{\alpha=1}^N ((x^m - \lambda_\alpha)^{nr_\alpha}) = (p(x)) \subseteq I$$

as left ideals of $T_\infty(n, v, \xi)$, and thus Lemma 2.6 or Corollary 2.9(1) can be applied to obtain that $T_\infty(n, v, \xi)^\circ = \sum_{I \in \mathcal{I}_0} (T_\infty(n, v, \xi)/I)^*$.

Now we try to prove that for each $I \in \mathcal{I}_0$, the subspace $(T_\infty(n, v, \xi)/I)^*$ can be spanned by some products of $\psi_\lambda, \omega, E_2, E_1$ ($\lambda \in \mathbb{k}$). Specifically, suppose $I = ((x^m - \lambda)^{nr})$ for some $r \in \mathbb{N}$ and $\lambda \in \mathbb{k}$, and we aim to show that

$$\{\psi_\lambda \omega^j E_2^s E_1^l \mid j \in n, s \in nr, l \in m\} \quad (3.2)$$

is a linear basis of n^2mr -dimensional space $(T_\infty(n, v, \xi)/I)^*$.

Evidently $I = \mathbb{k}\{g^j x^l (x^m - \lambda)^{nr} \mid j \in n, l \in \mathbb{N}\}$. Also, $T_\infty(n, v, \xi)/I$ has a linear basis

$$\{g^j x^l (x^m - \lambda)^s \mid j \in n, l \in m, s \in nr\},$$

and hence $\dim((T_\infty(n, v, \xi)/I)^*) = \dim(T_\infty(n, v, \xi)/I) = n^2mr$. Next we show that all elements in (3.2) vanish on I . Recall that

$$E_2^{[s]} E_1^{[l]} = \frac{1}{s! \cdot l!_{\xi^v}} E_2^s E_1^l : g^{j'} x^{l'} \mapsto \delta_{l', sm+l} \quad (l \in m, s, l' \in \mathbb{N}, j' \in n).$$

Therefore, for any $j, j' \in n, s \in nr, l \in m$ and $l' \in \mathbb{N}$, we have

$$\begin{aligned} & \langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, g^{j'} x^{l'} (x^m - \lambda)^{nr} \rangle \\ &= \langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, \sum_{t=0}^{nr} \binom{nr}{t} g^{j'} x^{l'+tm} (-\lambda)^{nr-t} \rangle \\ &= \sum_{t=0}^{nr} \binom{nr}{t} (-\lambda)^{nr-t} \langle \psi_\lambda \otimes \omega^j E_2^{[s]} E_1^{[l]}, \Delta(g^{j'} x^{l'+tm}) \rangle \\ &= \sum_{t=s}^{nr} \binom{nr}{t} (-\lambda)^{nr-t} \delta_{l', l} \binom{l+tm}{(t-s)m}_{\xi^v} \langle \psi_\lambda, g^{j'} x^{(t-s)m} \rangle \langle \omega^j E_2^{[s]} E_1^{[l]}, g^{j'+(t-s)mv} x^{l+sm} \rangle \\ &= \sum_{t=s}^{nr} \binom{nr}{t} (-\lambda)^{nr-t} \delta_{l', l} \binom{t}{t-s} \lambda^{t-s} \xi^{(j'+(t-s)mv)j} \\ &= \lambda^{nr-s} \delta_{l', l} \xi^{j'j} \sum_{t=s}^{nr} \binom{nr}{t} (-1)^{nr-t} \binom{t}{s} \\ &= \lambda^{nr-s} \delta_{l', l} \xi^{j'j} \sum_{t=s}^{nr} \binom{nr-s}{t-s} (-1)^{nr-t} = 0. \end{aligned}$$

In other words, $\langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, I \rangle = 0$ for any $j \in n, s \in nr$ and $l \in m$, which follows that the elements in (3.2) belong to $I^\perp = (T_\infty(n, v, \xi)/I)^*$.

Finally we prove that the elements in (3.2) are linearly independent. Choose a lexicographically ordered set of elements $\{g^{j'} x^{l'+s'm} \in T_\infty(n, v, \xi) \mid (j', s', l') \in n \times nr \times m\}$. Our goal is to show that the $n^2 mr \times n^2 mr$ square matrix

$$A := \left(\langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, g^{j'} x^{l'+s'm} \rangle \mid (j, s, l), (j', s', l') \in n \times nr \times m \right)$$

is invertible, which would imply the linear independence of (3.2). We try to compute

$$\langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, g^{j'} x^{l'+s'm} \rangle = \langle \psi_\lambda \otimes \omega^j E_2^{[s]} E_1^{[l]}, \Delta(g^{j'} x^{l'+s'm}) \rangle.$$

One finds that $\langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, g^{j'} x^{l'+s'm} \rangle = 0$ if $s > s'$. Otherwise when $s \leq s'$,

$$\begin{aligned} & \langle \psi_\lambda \omega^j E_2^{[s]} E_1^{[l]}, g^{j'} x^{l'+s'm} \rangle \\ &= \delta_{l', l} \binom{l + s'm}{(s' - s)m}_{\xi^v} \langle \psi_\lambda, g^{j'} x^{(s' - s)m} \rangle \langle \omega^j E_2^{[s]} E_1^{[l]}, g^{j' + (s' - s)mv} x^{l + sm} \rangle \\ &= \delta_{l', l} \binom{s'}{s} \lambda^{s' - s} \xi^{j' j}. \end{aligned}$$

The computation follows that the matrix A is the Kronecker product of following three invertible matrices:

- A Vandermonde matrix $(\xi^{j' j} \mid j, j' \in n)$,
- An $nr \times nr$ upper triangular matrix with entries $\binom{s'}{s} \lambda^{s' - s}$ for $s \leq s'$, and
- An identity matrix matrix $(\delta_{l', l} \mid l, l' \in m)$.

Thus A is invertible by Lemma 2.11, and hence (3.2) is a basis of $(T_\infty(n, v, \xi)/I)^*$. We conclude that

$$T_\infty(n, v, \xi)^\circ = \sum_{I \in \mathcal{I}_0} (T_\infty(n, v, \xi)/I)^* = \mathbb{k}\{\psi_\lambda \omega^j E_2^s E_1^l \mid \lambda \in \mathbb{k}, j \in n, s \in \mathbb{N}, l \in m\}.$$

□

Theorem 3.4. (1) $T_{\infty^\circ}(n, v, \xi)$ constructed in Subsection 3.1 is a Hopf algebra;
 (2) As a Hopf algebra, $T_{\infty^\circ}(n, v, \xi)^\circ$ is isomorphic to $T_{\infty^\circ}(n, v, \xi)$.

Proof. Consider the following map

$$\Theta : T_{\infty^\circ}(n, v, \xi) \rightarrow T_\infty(n, v, \xi)^\circ, \quad \Psi_\lambda \mapsto \psi_\lambda, \quad \Omega \mapsto \omega, \quad F_1 \mapsto E_1, \quad F_2 \mapsto E_2,$$

where $\lambda \in \mathbb{k}$. This is an epimorphism of algebras by Lemma 3.1 and Proposition 3.3. Furthermore, Θ would become an isomorphism of Hopf algebras with desired coalgebra structure and antipode, as long as it is injective (since $T_{\infty^\circ}(n, v, \xi)^\circ$ is in fact a Hopf algebra).

In order to show that Θ is injective, it is enough to show the linear independence of

$$\{\psi_\lambda \omega^j E_2^s E_1^l \mid \lambda \in \mathbb{k}, j \in n, s \in \mathbb{N}, l \in m\}$$

in $T_\infty(n, v, \xi)^\circ$. By the linear independence of elements in (3.2), we only need to show that any finite sum of form

$$\sum_{\alpha=1}^N (T_\infty(n, v, \xi) / ((x^m - \lambda_\alpha)^{nr_\alpha}))^* = \sum_{\alpha=1}^N ((x^m - \lambda_\alpha)^{nr_\alpha})^\perp \quad (3.3)$$

is direct as long as λ_α 's are distinct from each other. But this is clear:

$$\begin{aligned} & ((x^m - \lambda_{\alpha'})^{nr})^\perp \cap \left[\sum_{\alpha \neq \alpha'} ((x^m - \lambda_\alpha)^{nr})^\perp \right] = ((x^m - \lambda_{\alpha'})^{nr})^\perp \cap \left[\bigcap_{\alpha \neq \alpha'} ((x^m - \lambda_\alpha)^{nr})^\perp \right]^\perp \\ &= ((x^m - \lambda_{\alpha'})^{nr})^\perp \cap \left(\prod_{\alpha \neq \alpha'} (x^m - \lambda_\alpha)^{nr} \right)^\perp = [((x^m - \lambda_{\alpha'})^{nr}) + \left(\prod_{\alpha \neq \alpha'} (x^m - \lambda_\alpha)^{nr} \right)]^\perp \\ &= T_\infty(n, v, \xi)^\perp = 0. \end{aligned}$$

□

Remark 3.5. As a special case, the connected algebraic groups of dimension one $H_1 = \mathbb{k}[x]$ is equal to $T_\infty(1, 0, 1)$. Therefore, Theorem 3.4 gives H_1° , which is well-known (see [23, Example 9.1.7] for example).

4. THE FINITE DUAL OF GENERALIZED LIU'S ALGEBRA $B(n, \omega, \gamma)$

The concept and definition of a generalized Liu's algebra $B(n, \omega, \gamma)$ is given in [5, Section 3.4]. In this section, we use the same procedure as Section 3 to determine the structure of $B(n, \omega, \gamma)^\circ$.

4.1. The Hopf Algebra $B_\circ(n, \omega, \gamma)$. Let n and ω be positive integers, and γ be a primitive n th root of 1. For convenience, $\lambda^{\frac{1}{m}}$ always denotes an m th root of $\lambda \in \mathbb{k}^*$ for a positive integer m throughout the paper.

The Hopf algebra $B_\circ(n, \omega, \gamma)$ is constructed as follows. As an algebra, it is generated by

$$\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, F_1, F_2 \quad (\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*)$$

(where the parameters of Ψ are meant to range through $\{(\alpha, \beta) \in \mathbb{k}^* \times \mathbb{k}^* \mid \alpha^\omega = \beta^n\}$) with relations

$$\Psi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} \Psi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}} = \Psi_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{n}} \lambda_2^{\frac{1}{n}})}, \quad \Psi_{1,1} = 1, \quad F_1^n = 0,$$

$$F_2 \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} = \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} F_2, \quad F_1 \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} = \lambda^{\frac{1}{n}} \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} F_1, \quad F_1 F_2 = F_2 F_1 + \frac{1}{n} F_1$$

for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}, \lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}} \in \mathbb{k}^*$. The comultiplication, counit and antipode are given by

$$\Delta(F_1) = 1 \otimes F_1 + F_1 \otimes \Psi_{1,\gamma}, \quad \Delta(F_2) = 1 \otimes F_2 + F_2 \otimes 1 - \sum_{k=1}^{n-1} F_1^{[k]} \otimes \Psi_{1,\gamma}^k F_1^{[n-k]},$$

$$\begin{aligned}\Delta(\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) &= (\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \otimes \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}})(1 \otimes 1 + (1 - \lambda) \sum_{k=1}^{n-1} F_1^{[k]} \otimes \Psi_{1, \gamma}^k F_1^{[n-k]}), \\ \varepsilon(\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) &= 1, \quad \varepsilon(F_1) = \varepsilon(F_2) = 0, \\ S(F_1) &= -\gamma^{n-1} \Psi_{1, \gamma}^{n-1} F_1, \quad S(F_2) = -F_2, \quad S(\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) = \Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}\end{aligned}$$

for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$, where $F_1^{[k]} := \frac{1}{k! \gamma} F_1^k$ for $1 \leq k \leq n-1$.

The fact that $B_\circ(n, \omega, \gamma)$ is a Hopf algebra would be stated and proved in Subsection 4.3 as Theorem 4.4(1). Let η be a primitive ω th root of 1. Here we note that

$\{\Psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \Psi_{\eta, 1}^i \Psi_{1, \gamma}^j F_2^s F_1^l \mid \lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \text{ are fixed roots of } \lambda \in \mathbb{k}^*, i \in \omega, j \in n, s \in \mathbb{N}, l \in n\}$ is a linear basis, due to an application of the Diamond Lemma [2].

4.2. Certain Elements in $B(n, \omega, \gamma)^\circ$. Let n and ω be positive integers, and γ be a primitive n th root of 1. According to the definition of $B(n, \omega, \gamma)$ recalled in Definition 2.3, we know that it has a linear basis $\{x^i g^j y^l \mid i \in \omega, j \in \mathbb{Z}, l \in n\}$. Define following elements in $B(n, \omega, \gamma)^*$:

$$\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} : x^i g^j y^l \mapsto \delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}}, \quad E_1 : x^i g^j y^l \mapsto \delta_{l,1}, \quad E_2 : x^i g^j y^l \mapsto \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{n} \right) \quad (4.1)$$

for any $i \in \omega, j \in \mathbb{Z}, l \in n$ and $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$. We remark that these definitions make sense for all $i \in \mathbb{Z}$ as well.

One can verify that $\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}$ vanishes on the principal left ideal $(g^n - \lambda)$ for any $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$; E_2 vanishes on $((g^n - 1)^2)$; E_1 vanishes on $(g^n - 1)$. Therefore, these elements lie in $B(n, \omega, \gamma)^\circ$.

Lemma 4.1. *Following equations hold in $B(n, \omega, \gamma)^\circ$:*

$$\begin{aligned}\psi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} \psi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}} &= \psi_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{n}} \lambda_2^{\frac{1}{n}})}, \quad \psi_{1,1} = 1, \quad E_1^n = 0, \\ E_2 \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} &= \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} E_2, \quad E_1 \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} = \lambda^{\frac{1}{n}} \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} E_1, \quad E_1 E_2 = E_2 E_1 + \frac{1}{n} E_1\end{aligned}$$

for all $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}, \lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}} \in \mathbb{k}^*$.

Proof. Note that

$$\Delta(x^i g^j y^l) = (x \otimes x)^i (g \otimes g)^j (1 \otimes y + y \otimes g)^l = \sum_{k=0}^l \binom{l}{k}_\gamma x^i g^j y^k \otimes x^i g^{j+k} y^{l-k}.$$

We prove the lemma through checking their values on the basis $x^i g^j y^l$ ($i \in \omega, j \in \mathbb{Z}, l \in n$):

$$\begin{aligned}\langle \psi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} \psi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}}, x^i g^j y^l \rangle &= \delta_{l,0} \langle \psi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}}, x^i g^j \rangle \langle \psi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}}, x^i g^j \rangle = \delta_{l,0} \lambda_1^{\frac{i}{\omega}} \lambda_2^{\frac{i}{\omega}} \lambda_1^{\frac{j}{n}} \lambda_2^{\frac{j}{n}} \\ &= \langle \psi_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}} \lambda_2^{\frac{1}{n}}}, x^i g^j y^l \rangle,\end{aligned}$$

$$\begin{aligned}
\langle \psi_{1,1}, x^i g^j y^l \rangle &= \delta_{l,0} 1^i 1^j = \langle \varepsilon, x^i g^j y^l \rangle, \\
\langle E_2 \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j y^l \rangle &= \delta_{l,0} \langle E_2, x^i g^j \rangle \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j \rangle = \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{n} \right) \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}} \\
&= \delta_{l,0} \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j \rangle \langle E_2, x^i g^j \rangle = \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, E_2, x^i g^j y^l \rangle, \\
\langle E_1 \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j y^l \rangle &= \delta_{l,1} \langle E_1, x^i g^j y \rangle \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^{j+1} \rangle = \delta_{l,1} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j+1}{n}} \\
&= \lambda^{\frac{1}{n}} \delta_{l,1} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}} = \lambda^{\frac{1}{n}} \delta_{l,1} \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j \rangle \langle E_1, x^i g^j y \rangle \\
&= \langle \lambda^{\frac{1}{n}} \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, E_1, x^i g^j y^l \rangle, \\
\langle E_1 E_2, x^i g^j y^l \rangle &= \delta_{l,1} \langle E_1, x^i g^j y \rangle \langle E_2, x^i g^{j+1} \rangle = \delta_{l,1} \left(\frac{i}{\omega} + \frac{j+1}{n} \right) \\
&= \delta_{l,1} \left(\frac{i}{\omega} + \frac{j}{n} \right) + \delta_{l,1} \frac{1}{n} \\
&= \delta_{l,1} \langle E_2, x^i g^j \rangle \langle E_1, x^i g^j y \rangle + \delta_{l,1} \frac{1}{n} \langle E_1, x^i g^j y \rangle \\
&= \langle E_2 E_1 + \frac{1}{n} E_1, x^i g^j y^l \rangle.
\end{aligned}$$

Similar to the case in Lemma 3.1, it can be found by induction on $k \in n$ that

$$\langle E_1^k, x^i g^j y^l \rangle = \delta_{l,k} k!_{\gamma},$$

and thus $E_1^n = 0$. \square

Furthermore if denote $E_1^{[k]} := \frac{1}{k!_{\gamma}} E_1^k$, we can also verify that for any $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$ and $k \in n$,

$$\langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, E_1^{[k]}, x^i g^j y^l \rangle = \delta_{l,k} \langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, x^i g^j \rangle \langle E_1^{[k]}, x^i g^j y^k \rangle = \delta_{l,k} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}}. \quad (4.2)$$

Lemma 4.2. *Following equations hold in $B(n, \omega, \gamma)^{\circ}$:*

$$\Delta(E_1) = 1 \otimes E_1 + E_1 \otimes \psi_{1,\gamma}, \quad \Delta(E_2) = 1 \otimes E_2 + E_2 \otimes 1 - \sum_{k=1}^{n-1} E_1^{[k]} \otimes \psi_{1,\gamma}^k E_1^{[n-k]},$$

$$\Delta(\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) = (\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \otimes \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}})(1 \otimes 1 + (1 - \lambda) \sum_{k=1}^{n-1} E_1^{[k]} \otimes \psi_{1,\gamma}^k E_1^{[n-k]}),$$

$$\varepsilon(\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) = 1, \quad \varepsilon(E_1) = \varepsilon(E_2) = 0,$$

$$S(E_1) = -\gamma^{n-1} \psi_{1,\gamma}^{n-1} E_1, \quad S(E_2) = -E_2, \quad S(\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}) = \psi_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-1}{n}}}$$

for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$, where $E_1^{[k]} := \frac{1}{k!_{\gamma}} E_1^k$ for $1 \leq k \leq n-1$.

Proof. Note that $(x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) = \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'}$ ($i, i' \in \omega, j, j' \in \mathbb{Z}, l, l' \in n$). We also prove the lemma through checking their values on the basis $x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'}$.

$$\begin{aligned}
&\langle E_1, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\
&= \delta_{l+l',1} \gamma^{j'l} \langle E_1, x^{i+i'} g^{j+j'} y \rangle + \delta_{l+l',n+1} \gamma^{j'l} \langle E_1, x^{i+i'} g^{j+j'} y^{n+1} \rangle
\end{aligned}$$

$$\begin{aligned}
&= \delta_{l+l',1} \gamma^{j'l} = \delta_{l,0} \delta_{l',1} + \delta_{l,1} \delta_{l',0} \gamma^{j'} \\
&= \langle 1 \otimes E_1 + E_1 \otimes \psi_{1,\gamma}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle, \\
&\quad \langle E_2, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\
&= \gamma^{j'l} (\delta_{l+l',0} \langle E_2, x^{i+i'} g^{j+j'} \rangle + \delta_{l+l',n} \langle E_2, x^{i+i'} g^{j+j'} - x^{i+i'} g^{j+j'+n} \rangle) \\
&= \delta_{l,0} \delta_{l',0} \left(\frac{i+i'}{\omega} + \frac{j+j'}{n} \right) \\
&\quad + \delta_{l+l',n} \gamma^{j'l} \left(\frac{i+i'}{\omega} + \frac{j+j'}{n} - \frac{i+i'}{\omega} - \frac{j+j'+n}{n} \right) \\
&= \delta_{l,0} \delta_{l',0} \left(\frac{i+i'}{\omega} + \frac{j+j'}{n} \right) + \sum_{k=1}^{n-1} \delta_{l,k} \delta_{l',n-k} \gamma^{j'k} (-1) \\
&= \delta_{l,0} \delta_{l',0} \left(\frac{i'}{\omega} + \frac{j'}{n} \right) + \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{n} \right) \delta_{l',0} - \sum_{k=1}^{n-1} \delta_{l,k} \delta_{l',n-k} \gamma^{j'k} \\
&= \langle 1 \otimes E_2 + E_2 \otimes 1 - \sum_{k=1}^{n-1} E_1^{[k]} \otimes \psi_{1,\gamma}^k E_1^{[n-k]}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle,
\end{aligned}$$

and

$$\begin{aligned}
&\langle \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\
&= \delta_{l+l',0} \langle \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}, x^{i+i'} g^{j+j'} \rangle + \delta_{l+l',n} \gamma^{j'l} \langle \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}, x^{i+i'} g^{j+j'} - x^{i+i'} g^{j+j'+n} \rangle \\
&= \delta_{l+l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{j+j'}{n}} + \delta_{l+l',n} \gamma^{j'l} \left(\lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{j+j'}{n}} - \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{j+j'+n}{n}} \right) \\
&= \delta_{l,0} \delta_{l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{j+j'}{n}} + \delta_{l+l',n} \gamma^{j'l} (1 - \lambda) \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{j+j'}{n}} \\
&= (\delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}}) (\delta_{l',0} \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{n}}) + (1 - \lambda) \sum_{k=1}^{n-1} (\delta_{l,k} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{n}}) (\delta_{l',n-k} \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{n}} \gamma^{j'k}) \\
&= \langle \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} \otimes \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} + (1 - \lambda) \sum_{k=1}^{n-1} \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} E_1^{[k]} \otimes \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} \psi_{1,\gamma}^k E_1^{[n-k]}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle.
\end{aligned}$$

The expressions of the counit and the antipode are clear. \square

4.3. The Generation Problem.

Proposition 4.3. *As an algebra, $B(n, \omega, \gamma)^\circ$ is generated by $\psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}$, E_1 and E_2 for $\lambda \frac{1}{\omega}, \lambda \frac{1}{n} \in \mathbb{k}^*$.*

Proof. Clearly $B(n, \omega, \gamma)$ has a Laurent polynomial subalgebra $P = \mathbb{k}[g^{\pm 1}]$. Choose

$$\mathcal{I}_0 = \{((g^n - \lambda)^r) \triangleleft_l B(n, \omega, \gamma) \mid \lambda \in \mathbb{k}^*, r \in \mathbb{N}\}.$$

At first for any cofinite left ideal I of $B(n, \omega, \gamma)$, according to Lemma 2.8(2), there must be some non-zero polynomial

$$p(g) = \prod_{\alpha=1}^N (g^n - \lambda_\alpha)^{r_\alpha}$$

for some $N \geq 1$, $r_\alpha \in \mathbb{N}$ and distinct $\lambda_\alpha \in \mathbb{k}^*$, which generates a cofinite left ideal contained in I . It can be known by Lemma 2.10 that

$$\bigcap_{\alpha=1}^N ((g^n - \lambda_\alpha)^{r_\alpha}) = (p(g)) \subseteq I$$

as left ideals of $B(n, \omega, \gamma)$, and thus Lemma 2.6 or Corollary 2.9(2) can be applied to obtain that $B(n, \omega, \gamma)^\circ = \sum_{I \in \mathcal{I}_0} (B(n, \omega, \gamma)/I)^*$.

Now we try to prove that for each $I \in \mathcal{I}_0$, the subspace $(B(n, \omega, \gamma)/I)^*$ can be spanned by some products of $\psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}}, E_2, E_1$ ($\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}} \in \mathbb{k}^*$).

For the purpose, suppose $I = ((g^n - \lambda)^r)$ for some $r \in \mathbb{N}$ and $\lambda \in \mathbb{k}^*$. Let η be a primitive ω th root of 1, and $\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}$ be fixed roots of λ respectively. We aim to show that

$$\{\psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}} \psi_{\eta, 1}^i \psi_{1, \gamma}^j E_2^s E_1^l \mid i \in \omega, j \in n, s \in r, l \in n\} \quad (4.3)$$

is a linear basis of $\omega n^2 r$ -dimensional space $(B(n, \omega, \gamma)/I)^*$.

Evidently $I = \mathbb{k}\{x^i g^j (g^n - \lambda)^r y^l \mid i \in \omega, j \in \mathbb{Z}, l \in n\}$, since g^n is central. Also, $B(n, \omega, \gamma)/I$ has a linear basis

$$\{x^i g^j (g^n - \lambda)^s y^l \mid i \in \omega, j \in n, s \in r, l \in n\},$$

and hence $\dim((B(n, \omega, \gamma)/I)^*) = \dim(B(n, \omega, \gamma)/I) = \omega n^2 r$. Next we show that all elements in (4.3) vanish on I . Recall that

$$E_1^{[l]} = \frac{1}{l!_\gamma} E_1^l : x^{i'} g^{j'} y^{l'} \mapsto \delta_{l', l} \quad (i', j' \in \mathbb{Z}, l, l' \in n).$$

Therefore, for any $s \in r, l, l' \in n, i' \in \omega$ and $j' \in \mathbb{Z}$, and for arbitrary roots $\lambda_{\frac{1}{\omega}}$ and $\lambda_{\frac{1}{n}}$ of λ , we have

$$\begin{aligned} & \langle \psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}} E_2^s E_1^{[l]}, x^{i'} g^{j'} (g^n - \lambda)^r y^{l'} \rangle \\ &= \langle \psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}} E_2^s E_1^{[l]}, \sum_{t=0}^r \binom{r}{t} x^{i'} g^{j'+nt} (-\lambda)^{r-t} y^{l'} \rangle \\ &= \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \langle \psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}} E_2^s \otimes E_1^{[l]}, \Delta(x^{i'} g^{j'+nt} y^{l'}) \rangle \\ &= \delta_{l', l} \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \langle \psi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{n}}} E_2^s, x^{i'} g^{j'+nt} \rangle \langle E_1^{[l]}, x^{i'} g^{j'+nt} y^{l'} \rangle \\ &= \delta_{l', l} \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \lambda_{\frac{1}{\omega}}^{i'} \lambda_{\frac{1}{n}}^{j'+nt} \left(\frac{i'}{\omega} + \frac{j' + nt}{n} \right)^s \end{aligned} \quad (4.4)$$

$$\begin{aligned}
&= \delta_{l',l} \lambda^r \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{n}} \sum_{t=0}^r \binom{r}{t} (-1)^{r-t} \sum_{u=0}^s \binom{s}{u} \left(\frac{i'}{\omega} + \frac{j'}{n} \right)^u t^{s-u} \\
&= \delta_{l',l} \lambda^r \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{n}} \sum_{u=0}^s \binom{s}{u} \left(\frac{i'}{\omega} + \frac{j'}{n} \right)^u \sum_{t=0}^r \binom{r}{t} (-1)^{r-t} t^{s-u} \\
&= 0,
\end{aligned}$$

since $\sum_{t=0}^r \binom{r}{t} (-1)^{r-t} t^{s-u} = 0$ when $s-u < r$ is a part of the second Stirling number (2.10).

In other words, $\langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} E_2^s E_1^{[l]}, I \rangle = 0$ for any $s \in r$, $l \in n$, and arbitrary roots $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}$ of λ . It follows that the elements in (4.3) belong to $I^\perp = (B(n, \omega, \gamma)/I)^*$, as $\lambda^{\frac{1}{\omega}} \eta^i$ and $\lambda^{\frac{1}{n}} \gamma^j$ are still roots of λ .

Finally we prove that (4.3) are linearly independent. Choose a lexicographically ordered set of elements $\{x^{i'} g^{j'+s'n} y^{l'} \in B(n, \omega, \gamma) \mid (i', j', s', l') \in \omega \times n \times r \times n\}$. Our goal is to show that the $n^2 r \times n^2 r$ square matrix

$$A := \left(\langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \psi_{\eta, 1}^i \psi_{1, \gamma}^j E_2^s E_1^{[l]}, x^{i'} g^{j'+s'n} y^{l'} \rangle \mid (i, j, s, l), (i', j', s', l') \in \omega \times n \times r \times n \right)$$

is invertible for fixed $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}$, which would imply the linear independence of (4.3). We try to compute

$$\begin{aligned}
\langle \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \psi_{\eta, 1}^i \psi_{1, \gamma}^j E_2^s E_1^{[l]}, x^{i'} g^{j'+s'n} y^{l'} \rangle &= \langle \psi_{\lambda^{\frac{1}{\omega}} \eta^i, \lambda^{\frac{1}{n}} \gamma^j} E_2^s \otimes E_1^{[l]}, \Delta(x^{i'} g^{j'+s'n} y^{l'}) \rangle \\
&= \delta_{l',l} \langle \psi_{\lambda^{\frac{1}{\omega}} \eta^i, \lambda^{\frac{1}{n}} \gamma^j} E_2^s, x^{i'} g^{j'+s'n} \rangle \langle E_1^{[l]}, x^{i'} g^{j'+s'n} y^{l'} \rangle \\
&= \delta_{l',l} (\lambda^{\frac{1}{\omega}} \eta^i)^{i'} (\lambda^{\frac{1}{n}} \gamma^j)^{j'+s'n} \left(\frac{i'}{\omega} + \frac{j'+s'n}{n} \right)^s \\
&= \delta_{l',l} \lambda^{\frac{i'}{\omega} + \frac{j'+s'n}{n}} \eta^{i i'} \gamma^{j j'} \left(\frac{i'}{\omega} + \frac{j'}{n} + s' \right)^s,
\end{aligned}$$

and observe that $\det(A)$ equals to

$$(\lambda^{\frac{1}{\omega}})^{nr} (\lambda^{\frac{1}{n}})^{\omega r} \lambda^{\omega n \frac{(r-1)r}{2}} \det \left(\delta_{l',l} \eta^{i i'} \gamma^{j j'} \left(\frac{i'}{\omega} + \frac{j'}{n} + s' \right)^s \right).$$

However, there are following two facts:

- The matrix $(\delta_{l',l} \eta^{i i'} \gamma^{j j'} \mid (i, j, l), (i', j', l') \in \omega \times n \times n)$ is invertible by Lemma 2.11, since it is the Kronecker product of three invertible matrices;
- For each $(i', j') \in \omega \times n$, the matrix $((\frac{i'}{\omega} + \frac{j'}{n} + s')^s \mid s, s' \in r)$ is an invertible Vandermonde matrix.

Thus $(\delta_{l',l} \eta^{i i'} \gamma^{j j'} (\frac{i'}{\omega} + \frac{j'}{n} + s')^s \mid (i, j, s, l), (i', j', s', l') \in \omega \times n \times r \times n)$ is invertible according to Lemma 2.12, which follows that A is invertible. As a consequence, (4.3) is a basis of $(B(n, \omega, \gamma)/I)^*$.

Therefore,

$$B(n, \omega, \gamma)^\circ = \sum_{I \in \mathcal{I}_0} (B(n, \omega, \gamma)/I)^* = \mathbb{k} \{ \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} E_2^s E_1^l \mid \lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*, s \in \mathbb{N}, l \in n \}.$$

□

Theorem 4.4. (1) $B_\circ(n, \omega, \gamma)$ constructed in Subsection 4.1 is a Hopf algebra;
 (2) As a Hopf algebra, $B(n, \omega, \gamma)^\circ$ is isomorphic to $B_\circ(n, \omega, \gamma)$.

Proof. Consider the following map

$$\Theta : B_\circ(n, \omega, \gamma) \rightarrow B(n, \omega, \gamma)^\circ, \quad \Psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} \mapsto \psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}, \quad F_1 \mapsto E_1, \quad F_2 \mapsto E_2,$$

where $\lambda \frac{1}{\omega}, \lambda \frac{1}{n} \in \mathbb{k}^*$. This is an epimorphism of algebras by Lemma 4.1 and Proposition 4.3. Furthermore, Θ would become an isomorphism of Hopf algebras with desired coalgebra structure and antipode, as long as it is injective (since $B(n, \omega, \gamma)^\circ$ is in fact a Hopf algebra).

In order to show that Θ is injective, we aim to show the linear independence of

$$\{\psi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}} \psi_{\eta, 1}^i \psi_{1, \gamma}^j E_2^s E_1^l \mid \lambda \in \mathbb{k}^*, i \in \omega, j \in n, s \in \mathbb{N}, l \in n\}$$

in $B(n, \omega, \gamma)^\circ$, where $\lambda \frac{1}{\omega}$ and $\lambda \frac{1}{n}$ are fixed roots of each $\lambda \in \mathbb{k}^*$. By linear independence of elements in (4.3), we only need to show that any finite sum of form

$$\sum_{\alpha=1}^N (B(n, \omega, \gamma) / ((g^n - \lambda_\alpha)^r))^* = \sum_{\alpha=1}^N ((g^n - \lambda_\alpha)^r)^\perp \quad (4.5)$$

is direct, as long as λ_α 's are distinct from each other. But this is clear:

$$\begin{aligned} & ((g^n - \lambda_{\alpha'})^r)^\perp \cap \left[\sum_{\alpha \neq \alpha'} ((g^n - \lambda_\alpha)^r)^\perp \right] \\ &= ((g^n - \lambda_{\alpha'})^r)^\perp \cap \left[\bigcap_{\alpha \neq \alpha'} ((g^n - \lambda_\alpha)^r)^\perp \right] \\ &= ((g^n - \lambda_{\alpha'})^r)^\perp \cap \left(\prod_{\alpha \neq \alpha'} (g^n - \lambda_\alpha)^r \right)^\perp = [((g^n - \lambda_{\alpha'})^r) + \left(\prod_{\alpha \neq \alpha'} (g^n - \lambda_\alpha)^r \right)]^\perp \\ &= B(n, \omega, \gamma)^\perp = 0. \end{aligned} \quad (4.6)$$

□

Remark 4.5. As a special case, the connected algebraic groups of dimension one $H_2 = \mathbb{k}[g^{\pm 1}]$ is equal to $B(1, 1, 1)$. Therefore, Theorem 4.4 gives H_2° which is well-known too (see [23, Example 9.2.7] for example).

Remark 4.6. Let us contrast the structure of $B(n, \omega, \gamma)^\circ$ with the Hopf algebra introduced in [1]: Choose a set $I := \{(\alpha, \beta) \in \mathbb{k}^* \times \mathbb{k}^* \mid \alpha^\omega = \beta^n\}$. Then a particular case of the Hopf algebra

$$\mathfrak{D} := \mathfrak{D}(n, \gamma, (\lambda \frac{1}{n})_{(\lambda \frac{1}{\omega}, \lambda \frac{1}{n}) \in I}, 1)$$

defined in [1, Theorem 3.4] is generated by $u, x, a_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}^{\pm 1}$ ($\lambda \frac{1}{\omega}, \lambda \frac{1}{n} \in \mathbb{k}^*$) with relations

$$u^n = 1, \quad x^n = 0, \quad a_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}^{\pm 1} a_{\lambda \frac{1}{\omega}, \lambda \frac{1}{n}}^{\mp 1} = 1, \quad ux = \gamma xu,$$

$$\begin{aligned} ua_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} &= a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} u, \quad a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} x = \lambda^{\frac{1}{n}} x a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, \\ a_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} a_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}} &= a_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}} a_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} \end{aligned}$$

as an algebra for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}, \lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}} \in \mathbb{k}^*$. The comultiplication, counit and antipode on \mathfrak{D} is defined by

$$\begin{aligned} \Delta(u) &= u \otimes u, \quad \Delta(x) = u \otimes x + x \otimes 1, \\ \Delta(a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\pm 1}) &= (1 \otimes 1 + (1 - \lambda^{\pm 1}) \sum_{k=1}^{n-1} \frac{1}{k! \gamma (n-k)! \gamma} x^{n-k} u^k \otimes x^k) (a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\pm 1} \otimes a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\pm 1}) \\ \varepsilon(u) &= 1, \quad \varepsilon(x) = 0, \quad \varepsilon(a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\pm 1}) = 1, \\ S(u) &= u^{n-1}, \quad S(x) = -u^{n-1}x, \quad S(a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\pm 1}) = a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}^{\mp 1} \end{aligned}$$

for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$. Denote by $\overline{\mathfrak{D}}$ the quotient Hopf algebra of \mathfrak{D} modulo the ideal generated by

$$a_{1,1} - 1, \quad u - a_{1,\gamma}, \quad a_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}} a_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}}} - a_{\lambda_1^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{\omega}}} a_{\lambda_1^{\frac{1}{n}}, \lambda_2^{\frac{1}{n}}} \quad (\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{n}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{n}} \in \mathbb{k}^*).$$

It can be found that $\overline{\mathfrak{D}}^{\text{op cop}}$ is isomorphic to the Hopf subalgebra of $B(n, \omega, \gamma)^\circ$ generated by $\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}$, E_1 ($\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*$) via the isomorphism

$$\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}} \mapsto a_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}}}, \quad x \mapsto E_1 \quad (\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{n}} \in \mathbb{k}^*).$$

5. THE FINITE DUAL OF $D(m, d, \xi)$

We apply the same way used Sections 3 and 4 to give structure of $D(m, d, \xi)^\circ$ in this section.

5.1. The Hopf Algebra $D_\circ(m, d, \xi)$. Let m and d be positive integers such that $(1+m)d$ is even, and ξ be a primitive $2m$ th root of 1. Define

$$\omega := md, \quad \gamma := \xi^2.$$

The Hopf algebra $D_\circ(m, d, \xi)$ is constructed as follows. As an algebra, it is generated by

$$Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, F_1, F_2, \quad (\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*)$$

(where the parameters of Z and X are meant to range through $\{(\alpha, \beta) \in \mathbb{k}^* \times \mathbb{k}^* \mid \alpha^\omega = \beta^m\}$) with relations

$$\begin{aligned} Z_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} Z_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} &= Z_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}})}, \quad X_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} X_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = X_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}})}, \\ Z_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} X_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} &= X_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} Z_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = 0, \quad Z_{1,1} + X_{1,1} = 1, \quad F_1^m = \frac{1}{(1-\gamma)^m} X_{1,1}, \\ F_2 Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} &= Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} F_2, \quad F_1 Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \lambda^{\frac{1}{m}} Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} F_1, \end{aligned}$$

$$F_2 X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} F_2, \quad F_1 X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} F_1,$$

$$F_1 F_2 = F_2 F_1 + \frac{1}{m} Z_{1,1} F_1$$

for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}, \lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}} \in \mathbb{k}^*$. Denote $F_1^{[k]} := \frac{1}{k!_\gamma} F_1^k$ for $1 \leq k \leq m-1$. The comultiplication is given by:

$$\Delta(F_1) = 1 \otimes F_1 + F_1 \otimes (Z_{1,\gamma} + \xi X_{1,\gamma}),$$

$$\Delta(F_2) = (Z_{1,1} - X_{1,1}) \otimes F_2 + F_2 \otimes 1 - \sum_{k=1}^{m-1} (Z_{1,1} - X_{1,1}) F_1^{[k]} \otimes (Z_{1,\gamma} + \xi X_{1,\gamma})^{-m+k} F_1^{[m-k]},$$

$$\begin{aligned} \Delta(Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} + (1 - \lambda) \sum_{k=1}^{m-1} Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} F_1^{[k]} \otimes Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} Z_{1,\gamma}^k F_1^{[m-k]} \\ &\quad + \lambda^{\frac{(1-m)d/2}{\omega}} (1 - \lambda) (\theta_0^{-1} X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes X_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}}) \\ &\quad + \sum_{k=1}^{m-1} \theta_{m-k}^{-1} X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} F_1^{[k]} \otimes X_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} \xi^k X_{1,\gamma}^k F_1^{[m-k]} \\ \Delta(X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} - \theta_0 \sum_{k=1}^{m-1} \theta_1 \cdots \theta_{k-1} Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \xi^k X_{1,\gamma}^k F_1^{[m-k]} \\ &\quad + X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes Z_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} \\ &\quad - \theta_0 \sum_{k=1}^{m-1} \lambda^{\frac{-(m-k)d}{\omega}} \theta_1 \cdots \theta_{m-k-1} X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes Z_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} Z_{1,\gamma}^k F_1^{[m-k]}. \end{aligned}$$

where $\lambda^{\frac{1}{\omega}} \in \mathbb{k}^*$ and $\theta_0 = \frac{1-\lambda^{\frac{d}{\omega}}}{1/m}$, $\theta_k = \frac{1-\gamma^k \lambda^{\frac{d}{\omega}}}{1-\gamma^k}$ ($1 \leq k \leq m-1$). Note that $\theta_0 \theta_1 \cdots \theta_{m-1} = 1 - \lambda$ holds (see [13, Proposition IV.2.7]) and thus $(1 - \lambda) \theta_k^{-1}$ for $0 \leq k \leq m-1$ is well-defined. The coproducts on $Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$ and $X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$ for arbitrary $\lambda^{\frac{1}{\omega}}$ and $\lambda^{\frac{1}{m}}$ are given by

$$\Delta(Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = \Delta(Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) \Delta(Z_{1,\gamma} + \xi X_{1,\gamma})^k,$$

$$\Delta(X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = \xi^{-k} \Delta(X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) \Delta(Z_{1,\gamma} + \xi X_{1,\gamma})^k,$$

where k is a non-negative integer such that $\lambda^{\frac{1}{m}} = \lambda^{\frac{d}{\omega}} \gamma^k$ and $Z_{1,\gamma} + \xi X_{1,\gamma}$ is defined to be a group-like element.

The counit is given by

$$\varepsilon(Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = 1, \quad \varepsilon(X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = 0, \quad \varepsilon(F_1) = \varepsilon(F_2) = 0.$$

The antipode is given by

$$S(F_1) = -\gamma^{-1} (Z_{1,\gamma^{-1}} + \xi^{-1} X_{1,\gamma^{-1}}) F_1,$$

$$S(F_2) = -Z_{1,1} F_2 + X_{1,1} F_2 + \frac{1-m}{2m} X_{1,1},$$

$$S(Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = Z_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-1}{m}}},$$

$$S(X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) = \lambda^{\frac{(1-m)d/2}{\omega}} \gamma^{-k} X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}} \gamma^{-k}} \quad \text{when } \lambda^{\frac{1}{m}} = \lambda^{\frac{d}{\omega}} \gamma^k.$$

The fact that $D_{\circ}(m, d, \xi)$ is a Hopf algebra would be stated and proved in Subsection 5.3 as Theorem 5.7(1). Let η be an primitive ω th root of 1. Here we note that

$$\{Z_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} Z_{\eta, 1}^i Z_{1, \gamma}^j F_2^s F_1^l, X_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} X_{\eta, 1}^i X_{1, \gamma}^j F_2^s F_1^l \mid \lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \text{ are fixed roots of } \lambda \in \mathbb{k}^*, i \in \omega, j \in m, s \in \mathbb{N}, l \in m\}$$

is a linear basis, due to an application of the Diamond Lemma [2].

Remark 5.1. It is asked in [4, Remarks 7.1(3)] whether the cocycle σ non-trivial in the formula of [4, Proposition 7.2(V)]:

$$D(m, d, \xi)^{\circ} = (\mathbb{k}C_2 \#_{\sigma} T_f(m, 1, \xi^2)) \# (\mathbb{k}[f] \otimes \mathbb{k}(\mathbb{k}^{\times})).$$

Our results can be used to find that σ is trivial in fact.

5.2. Certain Elements in $D(m, d, \xi)^{\circ}$. Let m, d be positive integers such that $(1 + m)d$ is even and ξ a primitive $2m$ th root of unity. Define

$$\omega := md, \quad \gamma := \xi^2.$$

According to the definition of $D(m, d, \xi)$ recalled in Definition 2.4, we know that it has a linear basis $\{x^i g^j y^l, x^i g^j u_l \mid i \in \omega, j \in \mathbb{Z}, l \in m\}$ by [34, Lemma 3.3] and [35, Equation 4.7]. Define following elements in $D(m, d, \xi)^*$:

$$\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} : \begin{cases} x^i g^j y^l \mapsto \delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \\ x^i g^j u_l \mapsto 0 \end{cases}, \quad \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} : \begin{cases} x^i g^j y^l \mapsto 0 \\ x^i g^j u_l \mapsto \delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \end{cases},$$

$$E_1 : \begin{cases} x^i g^j y^l \mapsto \delta_{l,1} \\ x^i g^j u_l \mapsto \frac{\xi}{1-\gamma^{-1}} \delta_{l,1} \end{cases}, \quad E_2 : \begin{cases} x^i g^j y^l \mapsto \delta_{l,0} (\frac{i}{\omega} + \frac{j}{m}) \\ x^i g^j u_l \mapsto \delta_{l,0} (\frac{i}{\omega} + \frac{j}{m}) \end{cases}, \quad (5.1)$$

for any $i \in \omega, j \in \mathbb{Z}, l \in m$ and $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*$. We remark that these definitions make sense for all $i \in \mathbb{Z}$ as well, due to direct computations.

One can verify that $\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$ and $\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$ both vanish on the principal left ideal $((g^m - \lambda)(g^m - \lambda^{-1}))$ for any $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*$; E_2 vanishes on $((g^m - 1)^2)$; E_1 vanishes on $(g^m - 1)$. Therefore, these elements belong to $D(m, d, \xi)^{\circ}$.

Lemma 5.2. *Following equations hold in $D(m, d, \xi)^{\circ}$:*

$$\zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = \zeta_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}})}, \quad \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = \chi_{(\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}), (\lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}})},$$

$$\zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = 0, \quad \zeta_{1,1} + \chi_{1,1} = 1, \quad E_1^m = \frac{1}{(1-\gamma)^m} \chi_{1,1},$$

$$E_2 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2, \quad E_1 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \lambda^{\frac{1}{m}} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1,$$

$$E_2 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2, \quad E_1 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} = \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1,$$

$$E_1 E_2 = E_2 E_1 + \frac{1}{m} \zeta_{1,1} E_1$$

for all $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}, \lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}, \lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}} \in \mathbb{k}^*$.

Proof. Note that

$$\begin{aligned} \Delta(x^i g^j y^l) &= \sum_{k=0}^l \binom{l}{k}_{\gamma} x^i g^j y^k \otimes x^i g^{j+k} y^{l-k}, \\ \Delta(x^i g^j u_l) &= \sum_{k=0}^{m-1} \gamma^{k(l-k)} x^i g^j u_k \otimes x^{i-kd} g^{j+k} u_{l-k}. \end{aligned}$$

We prove the lemma through checking their values on the basis $x^i g^j y^l, x^i g^j u_l$ ($i \in \omega, j \in \mathbb{Z}, l \in m$):

$$\begin{aligned} \langle \zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle &= \delta_{l,0} \langle \zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}}, x^i g^j \rangle \langle \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j \rangle = \delta_{l,0} \lambda_1^{\frac{i}{\omega}} \lambda_2^{\frac{i}{\omega}} \lambda_1^{\frac{j}{m}} \lambda_2^{\frac{j}{m}} \\ &= \langle \zeta_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle, \\ \langle \zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j u_l \rangle &= 0 = \langle \zeta_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}}}, x^i g^j u_l \rangle, \\ \langle \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle &= 0 = \langle \chi_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle, \\ \langle \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j u_l \rangle &= \delta_{l,0} \langle \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}}, x^i g^j u_0 \rangle \langle \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j u_0 \rangle = \delta_{l,0} \lambda_1^{\frac{i}{\omega}} \lambda_2^{\frac{i}{\omega}} \lambda_1^{\frac{j}{m}} \lambda_2^{\frac{j}{m}} \\ &= \langle \zeta_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}}}, x^i g^j u_l \rangle, \\ \langle \zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle &= \langle \zeta_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \chi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j u_0 \rangle = 0, \\ \langle \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j y^l \rangle &= \langle \chi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \zeta_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}}, x^i g^j u_0 \rangle = 0, \\ \langle \zeta_{1,1} + \chi_{1,1}, x^i g^j y^l \rangle &= \delta_{l,0} 1^i 1^j = \langle \varepsilon, x^i g^j y^l \rangle, \\ \langle \zeta_{1,1} + \chi_{1,1}, x^i g^j u_l \rangle &= \delta_{l,0} 1^i 1^j = \langle \varepsilon, x^i g^j u_l \rangle, \\ \langle E_2 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j y^l \rangle &= \delta_{l,0} \langle E_2, x^i g^j \rangle \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j \rangle = \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m} \right) \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \\ &= \delta_{l,0} \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j \rangle \langle E_2, x^i g^j \rangle = \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, E_2, x^i g^j y^l \rangle, \\ \langle E_2 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j u_l \rangle &= 0 = \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, E_2, x^i g^j u_l \rangle, \\ \langle E_2 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j y^l \rangle &= 0 = \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, E_2, x^i g^j y^l \rangle, \\ \langle E_2 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j u_l \rangle &= \delta_{l,0} \langle E_2, x^i g^j u_0 \rangle \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j u_0 \rangle = \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m} \right) \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \\ &= \delta_{l,0} \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j u_0 \rangle \langle E_2, x^i g^j u_0 \rangle = \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, E_2, x^i g^j u_l \rangle, \\ \langle E_1 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^j y^l \rangle &= \delta_{l,1} \langle E_1, x^i g^j y \rangle \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^i g^{j+1} \rangle = \delta_{l,1} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j+1}{m}} \end{aligned}$$

$$\begin{aligned}
&= \lambda^{\frac{1}{m}} \delta_{l,1} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} = \lambda^{\frac{1}{m}} \delta_{l,1} \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^i g^j \rangle \langle E_1, x^i g^j y \rangle \\
&= \langle \lambda^{\frac{1}{m}} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1, x^i g^j y^l \rangle, \\
\langle E_1 \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^i g^j u_l \rangle &= 0 = \langle \lambda^{\frac{1}{m}} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1, x^i g^j u_l \rangle, \\
\langle E_1 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^i g^j y^l \rangle &= 0 = \langle \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1, x^i g^j y^l \rangle, \\
\langle E_1 \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^i g^j u_l \rangle &= \delta_{l,1} \langle E_1, x^i g^j u_1 \rangle \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^{i-d} g^{j+1} u_0 \rangle \\
&= \delta_{l,1} \frac{\xi}{1 - \gamma^{-1}} \lambda^{\frac{i-d}{\omega}} \lambda^{\frac{j+1}{m}} \\
&= \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} \delta_{l,1} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \frac{\xi}{1 - \gamma^{-1}} \\
&= \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} \delta_{l,1} \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} , x^i g^j u_0 \rangle \langle E_1, x^i g^j u_1 \rangle \\
&= \langle \lambda^{\frac{-d}{\omega}} \lambda^{\frac{1}{m}} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_1, x^i g^j u_l \rangle, \\
\langle E_1 E_2, x^i g^j y^l \rangle &= \delta_{l,1} \langle E_1, x^i g^j y \rangle \langle E_2, x^i g^{j+1} \rangle = \delta_{l,1} \left(\frac{i}{\omega} + \frac{j+1}{m} \right) \\
&= \delta_{l,1} \left(\frac{i}{\omega} + \frac{j}{m} \right) + \delta_{l,1} \frac{1}{m} \\
&= \delta_{l,1} \langle E_2, x^i g^j \rangle \langle E_1, x^i g^j y \rangle + \delta_{l,1} \frac{1}{m} \langle E_1, x^i g^j y \rangle \\
&= \langle E_2 E_1 + \frac{1}{m} E_1, x^i g^j y^l \rangle = \langle E_2 E_1 + \frac{1}{m} \zeta_{1,1} E_1, x^i g^j y^l \rangle, \\
\langle E_1 E_2, x^i g^j u_l \rangle &= \delta_{l,1} \langle E_1, x^i g^j u_1 \rangle \langle E_2, x^{i-d} g^{j+1} u_0 \rangle \\
&= \delta_{l,1} \frac{\xi}{1 - \gamma^{-1}} \left(\frac{i-d}{\omega} + \frac{j+1}{m} \right) \\
&= \delta_{l,1} \left(\frac{i}{\omega} + \frac{j}{m} \right) \frac{\xi}{1 - \gamma^{-1}} = \delta_{l,1} \langle E_2, x^i g^j u_0 \rangle \langle E_1, x^i g^j u_1 \rangle \\
&= \langle E_2 E_1, x^i g^j u_l \rangle = \langle E_2 E_1 + \frac{1}{m} \zeta_{1,1} E_1, x^i g^j u_l \rangle,
\end{aligned}$$

Similarly to the case in Lemmas 3.1 and 4.1, it can be found by induction on $k \in n$ that $\langle E_1^k, x^i g^j y^l \rangle = \delta_{l,k} k!_\gamma$ and

$$\begin{aligned}
\langle E_1^k, x^i g^j u_l \rangle &= \delta_{l,k} \gamma^{k-1} \langle E_1, x^i g^j u_1 \rangle \langle E_1^{k-1}, x^{i-d} g^{j+1} u_{k-1} \rangle \\
&= \delta_{l,k} \gamma^{k-1} \frac{\xi}{1 - \gamma^{-1}} \langle E_1^{k-1}, x^{i-d} g^{j+1} u_{k-1} \rangle \\
&= \delta_{l,k} \gamma^{(k-1)+(k-2)} \left(\frac{\xi}{1 - \gamma^{-1}} \right)^2 \langle E_1^{k-2}, x^{i-2d} g^{j+2} u_{k-2} \rangle = \dots \\
&= \delta_{l,k} \gamma^{(k-1)+(k-2)+\dots+1} \left(\frac{\xi}{1 - \gamma^{-1}} \right)^{k-1} \langle E_1, x^{i-(k-1)d} g^{j+(k-1)} u_1 \rangle \\
&= \delta_{l,k} \gamma^{(k-1)+(k-2)+\dots+1} \left(\frac{\xi}{1 - \gamma^{-1}} \right)^k = \delta_{l,k} \frac{\xi^{k^2}}{(1 - \gamma^{-1})^k}.
\end{aligned}$$

Thus $E_1^m = \frac{\xi^{m^2}}{(1-\gamma^{-1})^m} \chi_{1,1} = \frac{(-1)^m}{(1-\gamma^{-1})^m} \chi_{1,1} = \frac{1}{(1-\gamma)^m} \chi_{1,1}$. \square

It can be found in the proof above that

$$E_1^{[k]} := \frac{1}{k!_\gamma} E_1^k : \begin{cases} x^i g^j y^l \mapsto \delta_{l,k} \\ x^i g^j u_l \mapsto \frac{1}{k!_\gamma} \frac{\xi^{k^2}}{(1-\gamma^{-1})^k} \delta_{l,k} = \frac{\xi^k}{(1-\gamma^{-1})(1-\gamma^{-2})\dots(1-\gamma^{-k})} \delta_{l,k} \end{cases}.$$

Similarly to Equation (4.2), we can also verify that for any $\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{m}} \in \mathbb{k}^*$ and $k \in m$,

$$\begin{aligned} \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{m}}} E_1^{[k]} &: \begin{cases} x^i g^j y^l \mapsto \delta_{l,k} \lambda_{\frac{1}{\omega}}^{\frac{i}{\omega}} \lambda_{\frac{1}{m}}^{\frac{j}{m}} \\ x^i g^j u_l \mapsto 0 \end{cases}, \\ \chi_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{1}{m}}} E_1^{[k]} &: \begin{cases} x^i g^j y^l \mapsto 0 \\ x^i g^j u_l \mapsto \frac{\xi^k}{(1-\gamma^{-1})(1-\gamma^{-2})\dots(1-\gamma^{-k})} \delta_{l,k} \lambda_{\frac{1}{\omega}}^{\frac{i}{\omega}} \lambda_{\frac{1}{m}}^{\frac{j}{m}} \end{cases}. \end{aligned}$$

Lemma 5.3. *Following equations hold in $D(m, d, \xi)^\circ$:*

$$\Delta(E_1) = 1 \otimes E_1 + E_1 \otimes (\zeta_{1,\gamma} + \xi \chi_{1,\gamma}),$$

$$\Delta(E_2) = (\zeta_{1,1} - \chi_{1,1}) \otimes E_2 + E_2 \otimes 1 - \sum_{k=1}^{m-1} (\zeta_{1,1} - \chi_{1,1}) E_1^{[k]} \otimes (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^{-m+k} E_1^{[m-k]},$$

$$\varepsilon(E_1) = \varepsilon(E_2) = 0,$$

where $E_1^{[k]} := \frac{1}{k!_\gamma} E_1^k$ for $1 \leq k \leq m-1$. We remark that $\zeta_{1,1} - \chi_{1,1} = (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^m$.

Proof. Note that for each $i, i' \in \omega$, $j, j' \in \mathbb{Z}$ and $l, l' \in m$,

$$\begin{aligned} (x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) &= \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'}, \\ (x^i g^j u_l)(x^{i'} g^{j'} u_{l'}) &= \gamma^{j'l} x^{i-i'-2dj'} g^{j+j'} u_l u_{l'}, \\ (x^i g^j y^l)(x^{i'} g^{j'} u_{l'}) &= \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \dots \phi_{l'+l-1} g^{j+j'} u_{l+l'}, \\ (x^i g^j u_l)(x^{i'} g^{j'} y^{l'}) &= \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \dots \phi_{l+l'-1} g^{j+j'} u_{l+l'}. \end{aligned}$$

We also prove the lemma by through checking their values on the basis.

$$\begin{aligned} &\langle E_1, (x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) \rangle \\ &= \langle E_1, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\ &= \delta_{l+l',1} \gamma^{j'l} \langle E_1, x^{i+i'} g^{j+j'} y \rangle + \delta_{l+l',m+1} \gamma^{j'l} \langle E_1, x^{i+i'} g^{j+j'} y^{m+1} \rangle \\ &= \delta_{l+l',1} \gamma^{j'l} = \delta_{l,0} \delta_{l',1} + \delta_{l,1} \delta_{l',0} \gamma^{j'}, \\ &= \langle \zeta_{1,1} \otimes \zeta_{1,1} E_1 + \zeta_{1,1} E_1 \otimes \zeta_{1,\gamma}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle, \\ &\langle E_1, (x^i g^j u_l)(x^{i'} g^{j'} u_{l'}) \rangle \\ &= \langle E_1, \gamma^{j'l} x^{i-i'-2dj'} g^{j+j'} u_l u_{l'} \rangle = \delta_{l+l',1} \gamma^{j'l} \langle E_1, x^{i-i'-2dj'} g^{j+j'} u_l u_{l'} \rangle \\ &= \delta_{l,0} \delta_{l',1} \langle E_1, x^{i-i'-2dj'} g^{j+j'} u_0 u_1 \rangle + \delta_{l,1} \delta_{l',0} \gamma^{j'} \langle E_1, x^{i-i'-2dj'} g^{j+j'} u_1 u_0 \rangle \\ &= \delta_{l,0} \delta_{l',1} \langle E_1, u_0 u_1 \rangle + \delta_{l,1} \delta_{l',0} \gamma^{j'} \langle E_1, u_1 u_0 \rangle \\ &= \delta_{l,0} \delta_{l',1} \langle E_1, \frac{-\xi}{m} \phi_0 \phi_1 \dots \phi_{m-3} y g \rangle + \delta_{l,1} \delta_{l',0} \gamma^{j'} \langle E_1, \frac{1}{m} \phi_1 \phi_2 \dots \phi_{m-2} y g \rangle \end{aligned}$$

$$\begin{aligned}
&= \delta_{l,0}\delta_{l',1}\frac{-\xi\gamma}{m}(1-\gamma^{-1})(1-\gamma^{-2})\cdots(1-\gamma^{-(m-2)}) \\
&\quad + \delta_{l,1}\delta_{l',0}\gamma^{j'}\frac{\gamma}{m}(1-\gamma^{-2})(1-\gamma^{-3})\cdots(1-\gamma^{-(m-1)}) \\
&= \delta_{l,0}\delta_{l',1}\frac{-\xi\gamma}{1-\gamma} + \delta_{l,1}\delta_{l',0}\gamma^{j'}\frac{\gamma}{1-\gamma^{-1}} \\
&= \delta_{l,0}(\frac{\xi}{1-\gamma^{-1}}\delta_{l',1}) + \xi(\frac{\xi}{1-\gamma^{-1}}\delta_{l,1})(\delta_{l',0}\gamma^{j'}) \\
&= \langle \chi_{1,1} \otimes \chi_{1,1}E_1 + \chi_{1,1}E_1 \otimes \xi\chi_{1,\gamma}, x^i g^j u_l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
&\quad \langle E_1, (x^i g^j y^l)(x^{i'} g^{j'} u_{l'}) \rangle \\
&= \langle E_1, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{l'+l-1} g^{j+j'} u_{l+l'} \rangle \\
&= \delta_{l,0}\delta_{l',1} \langle E_1, x^{i+i'} g^{j+j'} u_1 \rangle + \delta_{l,1}\delta_{l',0}\gamma^{j'} \langle E_1, x^{i+i'} \phi_0 g^{j+j'} u_1 \rangle \\
&= \delta_{l,0}\delta_{l',1} \langle E_1, u_1 \rangle + \delta_{l,1}\delta_{l',0}\gamma^{j'} \langle E_1, \phi_0 u_1 \rangle \\
&= \delta_{l,0}\delta_{l',1} \frac{\xi}{1-\gamma^{-1}} + \delta_{l,1}\delta_{l',0}\gamma^{j'}(1-\gamma^{-1}) \frac{\xi}{1-\gamma^{-1}} \\
&= \delta_{l,0}(\frac{\xi}{1-\gamma^{-1}}\delta_{l',1}) + \xi\delta_{l,1}(\delta_{l',0}\gamma^{j'}) \\
&= \langle \zeta_{1,1} \otimes \chi_{1,1}E_1 + \zeta_{1,1}E_1 \otimes \xi\chi_{1,\gamma}, x^i g^j y^l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
&\quad \langle E_1, (x^i g^j u_l)(x^{i'} g^{j'} y^{l'}) \rangle \\
&= \langle E_1, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{l+l'-1} g^{j+j'} u_{l+l'} \rangle \\
&= \delta_{l,0}\delta_{l',1} \xi^{-1} \langle E_1, x^{i-i'-2dj'-d} \phi_0 g^{j+j'} u_1 \rangle + \delta_{l,1}\delta_{l',0}\gamma^{j'} \langle E_1, x^{i-i'-2dj'} g^{j+j'} u_1 \rangle \\
&= \delta_{l,0}\delta_{l',1} \xi^{-1} \langle E_1, \phi_0 u_1 \rangle + \delta_{l,1}\delta_{l',0}\gamma^{j'} \langle E_1, u_1 \rangle \\
&= \delta_{l,0}\delta_{l',1} \xi^{-1}(1-\gamma^{-1}) \frac{\xi}{1-\gamma^{-1}} + \delta_{l,1}\delta_{l',0}\gamma^{j'} \frac{\xi}{1-\gamma^{-1}} \\
&= \delta_{l,0}\delta_{l',1} + (\frac{\xi}{1-\gamma^{-1}}\delta_{l,1})(\delta_{l',0}\gamma^{j'}) \\
&= \langle \chi_{1,1} \otimes \zeta_{1,1}E_1 + \chi_{1,1}E_1 \otimes \zeta_{1,\gamma}, x^i g^j u_l \otimes x^{i'} g^{j'} y^{l'} \rangle.
\end{aligned}$$

It can be concluded that $\Delta(E_1) = 1 \otimes E_1 + E_1 \otimes (\zeta_{1,\gamma} + \xi\chi_{1,\gamma})$.

$$\begin{aligned}
&\langle E_2, (x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) \rangle \\
&= \langle E_2, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\
&= \gamma^{j'l} (\delta_{l+l',0} \langle E_2, x^{i+i'} g^{j+j'} \rangle + \delta_{l+l',m} \langle E_2, x^{i+i'} g^{j+j'} - x^{i+i'} g^{j+j'+m} \rangle) \\
&= \delta_{l,0}\delta_{l',0} (\frac{i+i'}{\omega} + \frac{j+j'}{m}) \\
&\quad + \delta_{l+l',m} \gamma^{j'l} (\frac{i+i'}{\omega} + \frac{j+j'}{m} - \frac{i+i'}{\omega} - \frac{j+j'+m}{m}) \\
&= \delta_{l,0}\delta_{l',0} (\frac{i+i'}{\omega} + \frac{j+j'}{m}) + \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k} \gamma^{j'k} (-1)
\end{aligned}$$

$$\begin{aligned}
&= \delta_{l,0}\delta_{l',0}\left(\frac{i'}{\omega} + \frac{j'}{m}\right) + \delta_{l,0}\left(\frac{i}{\omega} + \frac{j}{m}\right)\delta_{l',0} - \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k}\gamma^{j'k} \\
&= \langle \zeta_{1,1} \otimes \zeta_{1,1}E_2 + \zeta_{1,1}E_2 \otimes \zeta_{1,1} - \sum_{k=1}^{m-1} \zeta_{1,1}E_1^{[k]} \otimes \zeta_{1,\gamma}^k E_1^{[m-k]}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle, \\
&\quad \langle E_2, (x^i g^j u_l)(x^{i'} g^{j'} u_{l'}) \rangle \\
&= \langle E_2, \gamma^{j'l} x^{i-i'-2dj'} g^{j+j'} u_l u_{l'} \rangle \\
&= \delta_{l,0}\delta_{l',0} \langle E_2, \frac{1}{m} x^{i-i'-2dj' - \frac{(1+m)}{2}d} \phi_0 \phi_1 \cdots \phi_{m-2} g^{j+j'+1} \rangle \\
&\quad + \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k} \gamma^{j'k} \\
&\quad \quad \langle E_2, (-1)^{-m+k} \xi^{(m-k)^2} \frac{1}{m} x^{i-i'-2dj' - \frac{(1+m)}{2}d} \phi_k \cdots \phi_{m-1} \phi_0 \cdots \phi_{k-2} g^{j+j'+1} \rangle \\
&= \delta_{l,0}\delta_{l',0} \frac{1}{m} \left[\left(\frac{i-i'-2dj'}{\omega} - \frac{(1+m)}{2\omega}d + \frac{j+j'+1}{m} \right) m + \frac{m-1}{2} \right] \\
&\quad + \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k} \gamma^{j'k} (-1)^{-m+k} \xi^{(m-k)^2} \frac{1}{m} \\
&\quad \quad \left[-\frac{1}{m} (1 - \gamma^{-k-1}) \cdots (1 - \gamma^{-m+1}) (1 - \gamma^{-1}) \cdots (1 - \gamma^{-k+1}) \right] \\
&= \delta_{l,0}\delta_{l',0} \left(\frac{i-i'}{\omega} + \frac{j-j'}{m} \right) \\
&\quad + \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k} \gamma^{j'k} (-1)^{-m+k} \xi^{(m-k)^2} \frac{1}{m} \left(-\frac{1}{m} \frac{m}{1 - \gamma^{-k}} \right) \\
&= -\delta_{l,0}\delta_{l',0} \left(\frac{i'}{\omega} + \frac{j'}{m} \right) + \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m} \right) \delta_{l',0} \\
&\quad - \sum_{k=1}^{m-1} \frac{\xi^k}{(1 - \gamma^{-1}) \cdots (1 - \gamma^{-k})} \delta_{l,k} \xi^k \frac{\xi^{m-k}}{(1 - \gamma^{-1}) \cdots (1 - \gamma^{-(m-k)})} \delta_{l',m-k} \gamma^{j'k} \\
&= \langle -\chi_{1,1} \otimes \chi_{1,1}E_2 + \chi_{1,1}E_2 \otimes \chi_{1,1} - \sum_{k=1}^{m-1} \chi_{1,1}E_1^{[k]} \otimes \xi^k \chi_{1,\gamma}^k E_1^{[m-k]}, x^i g^j u_l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
&\quad \langle E_2, (x^i g^j y^l)(x^{i'} g^{j'} u_{l'}) \rangle \\
&= \langle E_2, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{l'+l-1} g^{j+j'} u_{l+l'} \rangle \\
&= \delta_{l,0}\delta_{l',0} \langle E_2, x^{i+i'} g^{j+j'} u_0 \rangle \\
&\quad + \sum_{k=1}^{m-1} \delta_{l,k}\delta_{l',m-k} \gamma^{j'k} \langle E_2, x^{i+i'} \phi_{m-k} \phi_{m-k+1} \cdots \phi_{m-1} g^{j+j'} u_0 \rangle \\
&= \delta_{l,0}\delta_{l',0} \left(\frac{i+i'}{\omega} + \frac{j+j'}{m} \right)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} \left(-\frac{1}{m}\right) (1 - \gamma^{k-1})(1 - \gamma^{k-2}) \cdots (1 - \gamma) \\
= & \delta_{l,0} \delta_{l',0} \left(\frac{i+i'}{\omega} + \frac{j+j'}{m}\right) \\
& - \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} \frac{1}{m} \frac{m}{(1 - \gamma^{-1})(1 - \gamma^{-2}) \cdots (1 - \gamma^{-(m-k)})} \\
= & \delta_{l,0} \delta_{l',0} \left(\frac{i'}{\omega} + \frac{j'}{m}\right) + \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m}\right) \delta_{l',0} \\
& + \sum_{k=1}^{m-1} \xi^k \delta_{l,k} \left[\frac{\xi^{m-k}}{(1 - \gamma^{-1})(1 - \gamma^{-2}) \cdots (1 - \gamma^{-(m-k)})} \delta_{l',m-k} \gamma^{j'k} \right] \\
= & \langle \zeta_{1,1} \otimes \chi_{1,1} E_2 + \zeta_{1,1} E_2 \otimes \chi_{1,1} + \sum_{k=1}^{m-1} \zeta_{1,1} E_1^{[k]} \otimes \xi^k \chi_{1,\gamma} E_1^{[m-k]}, x^i g^j y^l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
& \langle E_2, (x^i g^j u_l)(x^{i'} g^{j'} y^{l'}) \rangle \\
= & \langle E_2, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{l+l'-1} g^{j+j'} u_{l+l'} \rangle \\
= & \delta_{l,0} \delta_{l',0} \langle E_2, x^{i-i'-2dj'} g^{j+j'} u_0 \rangle \\
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \xi^{-(m-k)} \gamma^{j'k} \langle E_2, x^{i-i'-2dj'-d(m-k)} \phi_k \phi_{k+1} \cdots \phi_{m-1} g^{j+j'} u_0 \rangle \\
= & \delta_{l,0} \delta_{l',0} \left(\frac{i-i'-2dj'}{\omega} + \frac{j+j'}{m}\right) \\
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \xi^{-(m-k)} \gamma^{j'k} \left(-\frac{1}{m}\right) (1 - \gamma^{m-k-1})(1 - \gamma^{m-k-2}) \cdots (1 - \gamma) \\
= & \delta_{l,0} \delta_{l',0} \left(\frac{i-i'}{\omega} + \frac{j-j'}{m}\right) \\
& - \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \xi^{-(m-k)} \gamma^{j'k} \frac{1}{m} \frac{m}{(1 - \gamma^{-1})(1 - \gamma^{-2}) \cdots (1 - \gamma^{-k})} \\
= & -\delta_{l,0} \delta_{l',0} \left(\frac{i'}{\omega} + \frac{j'}{m}\right) + \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m}\right) \delta_{l',0} \\
& + \sum_{k=1}^{m-1} \left[\frac{\xi^k}{(1 - \gamma^{-1})(1 - \gamma^{-2}) \cdots (1 - \gamma^{-k})} \delta_{l,k} \right] (\delta_{l',m-k} \gamma^{j'k}) \\
= & \langle -\chi_{1,1} \otimes \zeta_{1,1} E_2 + \chi_{1,1} E_2 \otimes \zeta_{1,1} + \sum_{k=1}^{m-1} \chi_{1,1} E_1^{[k]} \otimes \zeta_{1,\gamma} E_1^{[m-k]}, x^i g^j u_l \otimes x^{i'} g^{j'} y^{l'} \rangle.
\end{aligned}$$

It can be concluded that

$$\Delta(E_2) = (\zeta_{1,1} - \chi_{1,1}) \otimes E_2 + E_2 \otimes 1 - \sum_{k=1}^{m-1} (\zeta_{1,1} - \chi_{1,1}) E_1^{[k]} \otimes (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^{-m+k} E_1^{[m-k]}.$$

The counit is clear. □

Lemma 5.4. *Following equations hold in $D(m, d, \xi)^\circ$:*

$$\Delta(\zeta_{1,\gamma} + \xi\chi_{1,\gamma}) = (\zeta_{1,\gamma} + \xi\chi_{1,\gamma}) \otimes (\zeta_{1,\gamma} + \xi\chi_{1,\gamma}), \quad \varepsilon(\zeta_{1,\gamma} + \xi\chi_{1,\gamma}) = 1,$$

and

$$\begin{aligned} \Delta(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} + (1 - \lambda) \sum_{k=1}^{m-1} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \zeta_{1,\gamma}^k E_1^{[m-k]} \\ &\quad + \lambda^{\frac{(1-m)d/2}{\omega}} (1 - \lambda) (\theta_0^{-1} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \\ &\quad + \sum_{k=1}^{m-1} \theta_{m-k}^{-1} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \xi^k \chi_{1,\gamma}^k E_1^{[m-k]}) \\ \Delta(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} - \theta_0 \sum_{k=1}^{m-1} \theta_1 \cdots \theta_{k-1} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \xi^k \chi_{1,\gamma}^k E_1^{[m-k]} \\ &\quad + \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \\ &\quad - \theta_0 \sum_{k=1}^{m-1} \lambda^{\frac{-(m-k)d}{\omega}} \theta_1 \cdots \theta_{m-k-1} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \zeta_{1,\gamma}^k E_1^{[m-k]}, \\ \varepsilon(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= 1, \quad \varepsilon(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) = 0, \end{aligned}$$

where $\lambda^{\frac{1}{\omega}} \in \mathbb{k}^*$, $E_1^{[k]} := \frac{1}{k!} E_1^k$, and $\theta_0 = \frac{1-\lambda^{\frac{d}{\omega}}}{1/m}$, $\theta_k = \frac{1-\gamma^k \lambda^{\frac{d}{\omega}}}{1-\gamma^k}$ ($1 \leq k \leq m-1$).

Proof. We also prove the lemma through checking their values on the basis.

$$\begin{aligned} &\langle \zeta_{1,\gamma} + \xi\chi_{1,\gamma}, (x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) \rangle \\ &= \langle \zeta_{1,\gamma}, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\ &= \delta_{l+l',0} \langle \zeta_{1,\gamma}, x^{i+i'} g^{j+j'} \rangle + \delta_{l+l',m} \langle \zeta_{1,\gamma}, \gamma^{j'l} x^{i+i'} g^{j+j'} - x^{i+i'} g^{j+j'+m} \rangle \\ &= \delta_{l,0} \delta_{l',0} \gamma^{j+j'} = (\delta_{l,0} \gamma^j)(\delta_{l',0} \gamma^{j'}) \\ &= \langle \zeta_{1,\gamma} \otimes \zeta_{1,\gamma}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle, \\ &\quad \langle \zeta_{1,\gamma} + \xi\chi_{1,\gamma}, (x^i g^j u_l)(x^{i'} g^{j'} u_{l'}) \rangle \\ &= \langle \zeta_{1,\gamma}, \gamma^{j'l} x^{i-i'-2dj'} g^{j+j'} u_l u_{l'} \rangle \\ &= \delta_{l,0} \delta_{l',0} \langle \zeta_{1,\gamma}, \frac{1}{m} x^{i-i'-2dj'-\frac{(1+m)}{2}d} \phi_0 \phi_1 \cdots \phi_{m-2} g^{j+j'+1} \rangle \\ &\quad + \delta_{l+l',m} \gamma^{j'l} \langle \zeta_{1,\gamma}, (-1)^{-l'} \xi^{l'^2} \frac{1}{m} x^{i-i'-2dj'-\frac{(1+m)}{2}d} \phi_l \cdots \phi_{m-1} \phi_0 \cdots \phi_{l-2} g^{j+j'+1} \rangle \\ &= \delta_{l,0} \delta_{l',0} \frac{1}{m} \langle \zeta_{1,\gamma}, x^{i-i'-2dj'-\frac{(1+m)}{2}d} \phi_0 \phi_1 \cdots \phi_{m-2} g^{j+j'+1} \rangle \\ &= \delta_{l,0} \delta_{l',0} \gamma^{j+j'+1} = (\xi \delta_{l,0} \gamma^j)(\xi \delta_{l',0} \gamma^{j'}) \\ &= \langle \xi\chi_{1,\gamma} \otimes \xi\chi_{1,\gamma}, x^i g^j u_l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\ &\quad \langle \zeta_{1,\gamma} + \xi\chi_{1,\gamma}, (x^i g^j y^l)(x^{i'} g^{j'} u_{l'}) \rangle \end{aligned}$$

$$\begin{aligned}
&= \langle \xi \chi_{1,\gamma}, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{l'+l-1} g^{j+j'} u_{l+l'} \rangle \\
&= \delta_{l,0} \delta_{l',0} \xi \langle \chi_{1,\gamma}, x^{i+i'} g^{j+j'} u_0 \rangle + \delta_{l+l',m} \xi \langle \chi_{1,\gamma}, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{m-1} g^{j+j'} u_m \rangle \\
&= \delta_{l,0} \delta_{l',0} \xi \gamma^{j+j'} = (\delta_{l,0} \gamma^j) (\xi \delta_{l',0} \gamma^{j'}) \\
&= \langle \zeta_{1,\gamma} \otimes \xi \chi_{1,\gamma}, x^i g^j y^l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
&\quad \langle \zeta_{1,\gamma} + \xi \chi_{1,\gamma}, (x^i g^j u_l) (x^{i'} g^{j'} y^{l'}) \rangle \\
&= \langle \xi \chi_{1,\gamma}, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{l+l'-1} g^{j+j'} u_{l+l'} \rangle \\
&= \delta_{l,0} \delta_{l',0} \xi \langle \chi_{1,\gamma}, x^{i-i'-2dj'} g^{j+j'} u_0 \rangle \\
&\quad + \delta_{l+l',m} \xi \langle \chi_{1,\gamma}, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{m-1} g^{j+j'} u_m \rangle \\
&= \delta_{l,0} \delta_{l',0} \xi \gamma^{j+j'} = (\xi \delta_{l,0} \gamma^j) (\delta_{l',0} \gamma^{j'}) \\
&= \langle \xi \chi_{1,\gamma} \otimes \zeta_{1,\gamma}, x^i g^j u_l \otimes x^{i'} g^{j'} y^{l'} \rangle.
\end{aligned}$$

It can be concluded that

$$\Delta(\zeta_{1,\gamma} + \xi \chi_{1,\gamma}) = (\zeta_{1,\gamma} + \xi \chi_{1,\gamma}) \otimes (\zeta_{1,\gamma} + \xi \chi_{1,\gamma}).$$

Now we compute that for any $\lambda_{\frac{1}{\omega}} \in \mathbb{k}^*$,

$$\begin{aligned}
&\langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, (x^i g^j y^l) (x^{i'} g^{j'} y^{l'}) \rangle \\
&= \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, \gamma^{j'l} x^{i+i'} g^{j+j'} y^{l+l'} \rangle \\
&= \delta_{l,0} \delta_{l',0} \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, x^{i+i'} g^{j+j'} \rangle + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, \gamma^{j'k} (x^{i+i'} g^{j+j'} - x^{i+i'} g^{j+j'+m}) \rangle \\
&= \delta_{l,0} \delta_{l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} (\lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} - \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j'+m)d}{\omega}}) \\
&= \delta_{l,0} \delta_{l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} + (1-\lambda) \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} \\
&= (\delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{id}{\omega}}) (\delta_{l',0} \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'd}{\omega}}) + (1-\lambda) \sum_{k=1}^{m-1} (\delta_{l,k} \lambda^{\frac{i}{\omega}} \lambda^{\frac{id}{\omega}}) (\delta_{l',m-k} \gamma^{j'k} \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'd}{\omega}}) \\
&= \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}} \otimes \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}} + (1-\lambda) \sum_{k=1}^{m-1} \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}} E_1^{[k]} \otimes \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}} \zeta_{1,\gamma}^k E_1^{[m-k]}, x^i g^j y^l \otimes x^{i'} g^{j'} y^{l'} \rangle, \\
&\quad \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, (x^i g^j u_l) (x^{i'} g^{j'} u_{l'}) \rangle \\
&= \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, \gamma^{j'l} x^{i-i'-2dj'} g^{j+j'} u_l u_{l'} \rangle \\
&= \delta_{l,0} \delta_{l',0} \langle \zeta_{\lambda_{\frac{1}{\omega}}, \lambda_{\frac{d}{\omega}}}, \frac{1}{m} x^{i-i'-2dj'-\frac{(1+m)}{2}d} \phi_0 \phi_1 \cdots \phi_{m-2} g^{j+j'+1} \rangle \\
&\quad + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k}
\end{aligned}$$

$$\begin{aligned}
& \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}}, (-1)^{-(m-k)} \xi^{(m-k)^2} \frac{1}{m} x^{i-i'-2dj' - \frac{(1+m)d}{2}} \phi_k \cdots \phi_{m-1} \phi_0 \cdots \phi_{k-2} g^{j+j'+1} \rangle \\
= & \delta_{l,0} \delta_{l',0} \frac{1}{m} \lambda^{\frac{i-i'-2dj'}{\omega} - \frac{(1+m)d}{2\omega}} (1 - \gamma^{-1} \lambda^{\frac{d}{\omega}}) (1 - \gamma^{-2} \lambda^{\frac{d}{\omega}}) \cdots (1 - \gamma^{-(m-1)} \lambda^{\frac{d}{\omega}}) \lambda^{\frac{(j+j'+1)d}{\omega}} \\
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} (-1)^{-(m-k)} \xi^{(m-k)^2} \frac{1}{m} \lambda^{\frac{i-i'-2dj'}{\omega} - \frac{(1+m)d}{2\omega}} \\
& (1 - \gamma^{-(k+1)} \lambda^{\frac{d}{\omega}}) \cdots (1 - \gamma^{-m} \lambda^{\frac{d}{\omega}}) (1 - \gamma^{-1} \lambda^{\frac{d}{\omega}}) \cdots (1 - \gamma^{-(k-1)} \lambda^{\frac{d}{\omega}}) \lambda^{\frac{(j+j'+1)d}{\omega}} \\
= & \delta_{l,0} \delta_{l',0} \frac{1}{m} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \lambda^{\frac{(1-m)d/2}{\omega}} \frac{1 - \lambda}{1 - \lambda^{\frac{d}{\omega}}} \\
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} (-1)^{-(m-k)} \xi^{(m-k)^2} \frac{1}{m} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \lambda^{\frac{(1-m)d/2}{\omega}} \frac{1 - \lambda}{1 - \gamma^{-k} \lambda^{\frac{d}{\omega}}} \\
= & (1 - \lambda) \lambda^{\frac{(1-m)d/2}{\omega}} \left[\frac{1/m}{1 - \lambda^{\frac{d}{\omega}}} \delta_{l,0} \delta_{l',0} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \right. \\
& + \sum_{k=1}^{m-1} \frac{1 - \gamma^{-k}}{1 - \gamma^{-k} \lambda^{\frac{d}{\omega}}} \frac{\xi^k}{(1 - \gamma^{-1}) \cdots (1 - \gamma^{-k})} \delta_{l,k} \\
& \left. \frac{\xi^{m-k}}{(1 - \gamma^{-1}) \cdots (1 - \gamma^{-(m-k)})} \delta_{l',m-k} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \xi^k \gamma^{j'k} \right] \\
= & \langle \lambda^{\frac{(1-m)d/2}{\omega}} (1 - \lambda) (\theta_0^{-1} \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \otimes \chi_{\lambda \frac{-1}{\omega}, \lambda \frac{-d}{\omega}} \\
& + \sum_{k=1}^{m-1} \theta_{m-k}^{-1} \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} E_1^{[k]} \otimes \chi_{\lambda \frac{-1}{\omega}, \lambda \frac{-d}{\omega}} \xi^k \chi_{1,\gamma}^k E_1^{[m-k]}), \\
& x^i g^j u_l \otimes x^{i'} g^{j'} u_{l'} \rangle, \\
& \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j y^l) (x^{i'} g^{j'} u_{l'}) \rangle \\
= & \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{l'+l-1} g^{j+j'} u_{l+l'} \rangle = 0, \\
& \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j u_l) (x^{i'} g^{j'} y^{l'}) \rangle \\
= & \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{l+l'-1} g^{j+j'} u_{l+l'} \rangle = 0.
\end{aligned}$$

It can be concluded that

$$\begin{aligned}
\Delta(\zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}) &= \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \otimes \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} + (1 - \lambda) \sum_{k=1}^{m-1} \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} E_1^{[k]} \otimes \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \zeta_{1,\gamma}^k E_1^{[m-k]} \\
&+ \lambda^{\frac{(1-m)d/2}{\omega}} (1 - \lambda) (\theta_0^{-1} \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \otimes \chi_{\lambda \frac{-1}{\omega}, \lambda \frac{-d}{\omega}} \\
&+ \sum_{k=1}^{m-1} \theta_{m-k}^{-1} \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} E_1^{[k]} \otimes \chi_{\lambda \frac{-1}{\omega}, \lambda \frac{-d}{\omega}} \xi^k \chi_{1,\gamma}^k E_1^{[m-k]}).
\end{aligned}$$

On the other hand,

$$\begin{aligned}
& \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j y^l)(x^{i'} g^{j'} y^{l'}) \rangle = 0, \\
& \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j u_l)(x^{i'} g^{j'} u_{l'}) \rangle = 0, \\
& \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j y^l)(x^{i'} g^{j'} u_{l'}) \rangle \\
= & \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \gamma^{j'l} x^{i+i'} \phi_{l'} \phi_{l'+1} \cdots \phi_{l'+l-1} g^{j+j'} u_{l+l'} \rangle \\
= & \delta_{l,0} \delta_{l',0} \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, x^{i+i'} g^{j+j'} u_0 \rangle \\
& + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \gamma^{j'k} x^{i+i'} \phi_{m-k} \phi_{m-k+1} \cdots \phi_{m-1} g^{j+j'} u_0 \rangle \\
= & \delta_{l,0} \delta_{l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} + \sum_{k=1}^{m-1} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} \lambda^{\frac{i+i'}{\omega}} (1 - \gamma^{k-1} \lambda^{\frac{d}{\omega}}) (1 - \gamma^{k-2} \lambda^{\frac{d}{\omega}}) \cdots (1 - \lambda^{\frac{d}{\omega}}) \lambda^{\frac{(j+j')d}{m}} \\
= & \delta_{l,0} \delta_{l',0} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} \\
& - \sum_{k=1}^{m-1} (1 - \lambda^{\frac{d}{\omega}}) (1 - \gamma \lambda^{\frac{d}{\omega}}) \cdots (1 - \gamma^{k-1} \lambda^{\frac{d}{\omega}}) \frac{m}{(1 - \gamma) \cdots (1 - \gamma^{k-1})} \\
& \delta_{l,k} \frac{\xi^{m-k}}{(1 - \gamma^{-1})(1 - \gamma^{-2}) \cdots (1 - \gamma^{-(m-k)})} \delta_{l',m-k} \xi^k \gamma^{j'k} \lambda^{\frac{i+i'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} \\
= & \langle \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \otimes \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} - \theta_0 \sum_{k=1}^{m-1} \theta_1 \cdots \theta_{k-1} \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} E_1^{[k]} \otimes \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}} \xi^k \chi_{1,\gamma} E_1^{[m-k]}, \\
& x^i g^j y^l \otimes x^{i'} g^{j'} u_{l'}, \\
& \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, (x^i g^j u_l)(x^{i'} g^{j'} y^{l'}) \rangle \\
= & \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \xi^{-l'} \gamma^{j'l} x^{i-i'-2dj'-dl'} \phi_l \phi_{l+1} \cdots \phi_{l+l'-1} g^{j+j'} u_{l+l'} \rangle \\
= & \delta_{l,0} \delta_{l',0} \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, x^{i-i'-2dj'} g^{j+j'} u_0 \rangle \\
& + \sum_{k=1}^{m-k} \delta_{l,k} \delta_{l',m-k} \langle \chi_{\lambda \frac{1}{\omega}, \lambda \frac{d}{\omega}}, \xi^{-(m-k)} \gamma^{j'k} x^{i-i'-2dj'-d(m-k)} \phi_k \phi_{k+1} \cdots \phi_{m-1} g^{j+j'} u_0 \rangle \\
= & \delta_{l,0} \delta_{l',0} \lambda^{\frac{i-i'-2dj'}{\omega}} \lambda^{\frac{(j+j')d}{\omega}} + \sum_{k=1}^{m-k} \delta_{l,k} \delta_{l',m-k} \xi^{-(m-k)} \gamma^{j'k} \lambda^{\frac{i-i'-2dj'-d(m-k)}{\omega}} \\
& (1 - \gamma^{m-k-1} \lambda^{\frac{d}{\omega}}) (1 - \gamma^{m-k-2} \lambda^{\frac{d}{\omega}}) \cdots (1 - \lambda^{\frac{d}{\omega}}) \lambda^{\frac{(j+j')d}{\omega}} \\
= & \delta_{l,0} \delta_{l',0} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} + \sum_{k=1}^{m-k} \delta_{l,k} \delta_{l',m-k} \xi^{-(m-k)} \gamma^{j'k} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \lambda^{\frac{-(m-k)d}{\omega}} \\
& (1 - \gamma^{m-k-1} \lambda^{\frac{d}{\omega}}) (1 - \gamma^{m-k-2} \lambda^{\frac{d}{\omega}}) \cdots (1 - \lambda^{\frac{d}{\omega}})
\end{aligned}$$

$$\begin{aligned}
&= \delta_{l,0} \delta_{l',0} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \\
&\quad - \sum_{k=1}^{m-k} \lambda^{\frac{-(m-k)d}{\omega}} (1 - \lambda^{\frac{d}{\omega}}) (1 - \gamma \lambda^{\frac{d}{\omega}}) \cdots (1 - \gamma^{m-k-1} \lambda^{\frac{d}{\omega}}) \frac{m}{(1 - \gamma) \cdots (1 - \gamma^{m-k-1})} \\
&\quad \frac{\xi^k}{(1 - \gamma^{-1}) \cdots (1 - \gamma^{-k})} \delta_{l,k} \delta_{l',m-k} \gamma^{j'k} \lambda^{\frac{i-i'}{\omega}} \lambda^{\frac{(j-j')d}{\omega}} \\
&= \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} - \theta_0 \sum_{k=1}^{m-1} \lambda^{\frac{-(m-k)d}{\omega}} \theta_1 \cdots \theta_{m-k-1} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} \zeta_{1,\gamma}^k E_1^{[m-k]}, \\
&\quad x^i g^j u_l \otimes x^{i'} g^{j'} y^{l'} \rangle,
\end{aligned}$$

It can be concluded that

$$\begin{aligned}
\Delta(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}}) &= \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} - \theta_0 \sum_{k=1}^{m-1} \theta_1 \cdots \theta_{k-1} \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \xi^k \chi_{1,\gamma}^k E_1^{[m-k]} \\
&\quad + \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} \otimes \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} \\
&\quad - \theta_0 \sum_{k=1}^{m-1} \lambda^{\frac{-(m-k)d}{\omega}} \theta_1 \cdots \theta_{m-k-1} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}}} E_1^{[k]} \otimes \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-d}{\omega}}} \zeta_{1,\gamma}^k E_1^{[m-k]}.
\end{aligned}$$

The counit is clear. \square

Lemma 5.5. *Following equations hold in $D(m, d, \xi)^\circ$:*

$$\begin{aligned}
S(\zeta_{1,\gamma} + \xi \chi_{1,\gamma}) &= \zeta_{1,\gamma^{-1}} + \xi^{-1} \chi_{1,\gamma^{-1}}, \\
S(E_1) &= -\gamma^{-1} (\zeta_{1,\gamma^{-1}} + \xi^{-1} \chi_{1,\gamma^{-1}}) E_1, \\
S(E_2) &= -\zeta_{1,1} E_2 + \chi_{1,1} E_2 + \frac{1-m}{2m} \chi_{1,1}, \\
S(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) &= \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-1}{m}}}, \\
S(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) &= \lambda^{\frac{(1-m)d/2}{\omega}} \gamma^{-k} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}} \gamma^{-k}} \text{ when } \lambda^{\frac{1}{m}} = \lambda^{\frac{d}{\omega}} \gamma^k.
\end{aligned}$$

Proof. We also prove the lemma through checking their values on the basis.

$$\begin{aligned}
\langle S(E_1), x^i g^j y^l \rangle &= \delta_{l,1} \langle E_1, -\gamma^{-j-1} x^{-i} g^{-j-1} y \rangle \\
&= -\delta_{l,1} \gamma^{-j-1} = \langle -\gamma^{-1} \zeta_{1,\gamma^{-1}} E_1, x^i g^j y^l \rangle, \\
\langle S(E_1), x^i g^j u_l \rangle &= \delta_{l,1} \langle E_1, -\xi^{-1} \gamma^{-j-1} x^{i+d+\frac{3}{2}(1-m)d+2dj} g^{m-2-j} u_1 \rangle \\
&= -\xi^{-1} \gamma^{-j-1} \frac{\xi}{1 - \gamma^{-1}} \delta_{l,1} = \langle -\gamma^{-1} \xi^{-1} \chi_{1,\gamma^{-1}} E_1, x^i g^j u_l \rangle.
\end{aligned}$$

It can be concluded that

$$S(E_1) = -\gamma^{-1} (\zeta_{1,\gamma^{-1}} + \xi^{-1} \chi_{1,\gamma^{-1}}) E_1.$$

Also,

$$\begin{aligned}
\langle S(E_2), x^i g^j y^l \rangle &= \delta_{l,0} \langle E_2, x^{-i} g^{-j} \rangle \\
&= -\delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m} \right) = \langle -\zeta_{1,1} E_2, x^i g^j y^l \rangle, \\
\langle S(E_2), x^i g^j u_l \rangle &= \delta_{l,0} \langle E_2, x^{i+2dj+\frac{3}{2}(1-m)d} g^{m-1-j} u_0 \rangle \\
&= \delta_{l,0} \left(\frac{i+2dj}{\omega} + \frac{3(1-m)d/2}{\omega} + \frac{m-1-j}{m} \right) \\
&= \delta_{l,0} \left(\frac{i}{\omega} + \frac{j}{m} + \frac{1-m}{2m} \right) = \langle \chi_{1,1} E_2 + \frac{1-m}{2m} \chi_{1,1}, x^i g^j u_l \rangle.
\end{aligned}$$

It can be concluded that

$$S(E_2) = -\zeta_{1,1} E_2 + \chi_{1,1} E_2 + \frac{1-m}{2m} \chi_{1,1}.$$

Moreover,

$$\begin{aligned}
\langle S(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}), x^i g^j y^l \rangle &= \delta_{l,0} \langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^{-i} g^{-j} \rangle = \delta_{l,0} \lambda^{\frac{-i}{\omega}} \lambda^{\frac{-j}{m}} \\
&= \langle \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-1}{m}}}, x^i g^j y^l \rangle, \\
\langle S(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}), x^i g^j u_l \rangle &= 0.
\end{aligned}$$

When $\lambda^{\frac{1}{m}} = \lambda^{\frac{d}{\omega}} \gamma^k$ holds where k is a non-negative integer,

$$\begin{aligned}
\langle S(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}), x^i g^j y^l \rangle &= 0, \\
\langle S(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}), x^i g^j u_l \rangle &= \delta_{l,0} \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, x^{i+2dj+\frac{3}{2}(1-m)d} g^{m-1-j} u_0 \rangle \\
&= \delta_{l,0} \lambda^{\frac{i+2dj}{\omega}} \lambda^{\frac{3(1-m)d/2}{\omega}} \lambda^{\frac{m-1-j}{m}} \\
&= \delta_{l,0} \lambda^{\frac{i+2dj}{\omega}} \lambda^{\frac{3(1-m)d/2}{\omega}} (\lambda^{\frac{d}{\omega}} \gamma^k)^{m-1-j} \\
&= \lambda^{\frac{(1-m)d/2}{\omega}} \gamma^{-k} \delta_{l,0} \lambda^{\frac{i}{\omega}} (\lambda^{\frac{d}{\omega}} \gamma^{-k})^j \\
&= \langle \lambda^{\frac{(1-m)d/2}{\omega}} \gamma^{-k} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}} \gamma^{-k}}, x^i g^j u_l \rangle.
\end{aligned}$$

It can be concluded that

$$\begin{aligned}
S(\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) &= \zeta_{\lambda^{\frac{-1}{\omega}}, \lambda^{\frac{-1}{m}}}, \\
S(\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}) &= \lambda^{\frac{(1-m)d/2}{\omega}} \gamma^{-k} \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{d}{\omega}} \gamma^{-k}} \text{ when } \lambda^{\frac{1}{m}} = \lambda^{\frac{d}{\omega}} \gamma^k.
\end{aligned}$$

□

5.3. The Generation Problem.

Proposition 5.6. *As an algebra, $D(m, d, \xi)^\circ$ is generated by $\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$, $\chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}$, E_1 and E_2 for $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*$.*

Proof. Clearly $D(m, d, \xi)$ has a Laurent polynomial subalgebra $P = \mathbb{k}[g^{\pm 1}]$. Choose $\mathcal{I}_0 = \{((g^m - 1)^r), ((g^m + 1)^r), ((g^m - \lambda)^r (g^m - \lambda^{-1})^r) \triangleleft_l D(m, d, \xi) \mid \lambda \in \mathbb{k}^* \setminus \{\pm 1\}, r \in \mathbb{N}\}$.

At first for any cofinite left ideal I of $D(m, d, \xi)$, according to Lemma 2.8(3), there must be some non-zero polynomial

$$p(g) = (g^m - 1)^{r'} (g^m + 1)^{r''} \prod_{\alpha=1}^N (g^m - \lambda_\alpha)^{r_\alpha} (g^m - \lambda_\alpha^{-1})^{r_\alpha}$$

for some $N \geq 1$, $r', r'', r_\alpha \in \mathbb{N}$ and distinct $\lambda_\alpha \in \mathbb{k}^* \setminus \{\pm 1\}$ which are not inverses of each other, such that $p(g)$ generates a cofinite left ideal contained in I . It can be known by Lemma 2.10 that

$$((g^m - 1)^{r'}) \cap ((g^m + 1)^{r''}) \cap \left[\bigcap_{\alpha=1}^N ((g^m - \lambda_\alpha)^{r_\alpha} (g^m - \lambda_\alpha^{-1})^{r_\alpha}) \right] = (p(g)) \subseteq I$$

as left ideals of $D(m, d, \xi)$, and thus Lemma 2.6 or Corollary 2.9(3) can be applied to obtain that $D(m, d, \xi)^\circ = \sum_{I \in \mathcal{I}_0} (D(m, d, \xi)/I)^*$.

Now we try to prove that for each $I \in \mathcal{I}_0$, the subspace $(D(m, d, \xi)/I)^*$ can be spanned by some products of $\zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}, E_2, E_1$ ($\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*$). For simplicity, we always denote

$$\psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} := \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} + \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} : \begin{cases} x^i g^j y^l \mapsto \delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \\ x^i g^j u_l \mapsto \delta_{l,0} \lambda^{\frac{i}{\omega}} \lambda^{\frac{j}{m}} \end{cases},$$

where $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}} \in \mathbb{k}^*$. Note that $\psi_{\lambda_1^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}}} \psi_{\lambda_2^{\frac{1}{\omega}}, \lambda_2^{\frac{1}{m}}} = \psi_{\lambda_1^{\frac{1}{\omega}} \lambda_2^{\frac{1}{\omega}}, \lambda_1^{\frac{1}{m}} \lambda_2^{\frac{1}{m}}}$ always holds. Moreover, let η be an primitive ω th root of 1.

We continue to show the spanning by a discussion on the form of $I \in \mathcal{I}_0$:

Case 1. Suppose $I = ((g^m - \lambda)^r (g^m - \lambda^{-1})^r)$ for some $r \in \mathbb{N}$ and $\lambda \in \mathbb{k}^* \setminus \{\pm 1\}$. Let $\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}$ be some fixed roots of λ respectively. In this case we aim to show that

$$\left\{ \psi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}^{(-1)^e} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l \mid e \in 2, i \in \omega, k \in 2m, s \in r, l \in m \right\} \quad (5.2)$$

is a linear basis of $4\omega m^2 r$ -dimensional space $(D(m, d, \xi)/I)^*$. We remark that (5.2) is linearly equivalent to

$$\left\{ \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}^{(-1)^e} \zeta_{\eta,1}^i \zeta_{1,\gamma}^j E_2^s E_1^l, \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}}^{(-1)^e} \chi_{\eta,1}^i \chi_{1,\gamma}^j E_2^s E_1^l \mid e \in 2, i \in \omega, j \in m, s \in r, l \in m \right\}. \quad (5.3)$$

Evidently

$$I = \mathbb{k}\{x^i g^j (g^m - \lambda)^r (g^m - \lambda^{-1})^r y^l, x^i g^j (g^m - \lambda)^r (g^m - \lambda^{-1})^r u_l \mid i \in \omega, j \in \mathbb{Z}, l \in m\},$$

since $y g^m = g^m y$ and $u_i g^m = g^{-m} u_i$ for each $i \in m$. Also, $D(m, d, \xi)/I$ has a linear basis

$$\{x^i g^j (g^m - \lambda)^s (g^m - \lambda^{-1})^s y^l, x^i g^j (g^m - \lambda)^s (g^m - \lambda^{-1})^s u_l \mid i \in \omega, j \in 2m, s \in r, l \in m\},$$

and hence $\dim((D(m, d, \xi)/I)^*) = \dim(D(m, d, \xi)/I) = 4\omega m^2 r$. Next we show that all elements in (5.3) vanish on I . We make computations for any $s \in r$, $l, l' \in m$ and $j' \in \mathbb{Z}$, especially for arbitrary roots $\lambda^{\frac{1}{\omega}}$ and $\lambda^{\frac{1}{m}}$ of λ : Clearly

$$\langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s E_1^l, x^{i'} g^{j'} (g^m - \lambda)^r u_{l'} \rangle = 0, \quad (5.4)$$

$$\langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s E_1^l, x^{i'} g^{j'} (g^m - \lambda)^r y^{l'} \rangle = 0, \quad (5.5)$$

and

$$\langle \zeta_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s E_1^l, x^{i'} g^{j'} (g^m - \lambda)^r y^{l'} \rangle = 0 \quad (5.6)$$

due to the same reason as Equation (4.4). Similarly,

$$\begin{aligned} & \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s E_1^l, x^{i'} g^{j'} (g^m - \lambda)^r u_{l'} \rangle \\ &= \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s E_1^l, \sum_{t=0}^r \binom{r}{t} x^{i'} g^{j'+mt} (-\lambda)^{r-t} u_{l'} \rangle \\ &= \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s \otimes E_1^l, \Delta(x^{i'} g^{j'+nt} u_{l'}) \rangle \\ &= \delta_{l',l} \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \langle \chi_{\lambda^{\frac{1}{\omega}}, \lambda^{\frac{1}{m}}} E_2^s, x^{i'} g^{j'+mt} u_0 \rangle \langle E_1^l, x^{i'} g^{j'+nt} u_l \rangle \\ &= \delta_{l',l} \sum_{t=0}^r \binom{r}{t} (-\lambda)^{r-t} \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'+mt}{m}} \left(\frac{i'}{\omega} + \frac{j'+mt}{m} \right)^s \frac{\xi^{l^2}}{(1-\gamma^{-1})^l} \\ &= \delta_{l',l} \frac{\xi^{l^2}}{(1-\gamma^{-1})^l} \lambda^r \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{m}} \sum_{t=0}^r \binom{r}{t} (-1)^{r-t} \sum_{u=0}^s \binom{s}{u} \left(\frac{i'}{\omega} + \frac{j'}{m} \right)^u t^{s-u} \\ &= \delta_{l',l} \frac{\xi^{l^2}}{(1-\gamma^{-1})^l} \lambda^r \lambda^{\frac{i'}{\omega}} \lambda^{\frac{j'}{m}} \sum_{u=0}^s \binom{s}{u} \left(\frac{i'}{\omega} + \frac{j'}{m} \right)^u \sum_{t=0}^r \binom{r}{t} (-1)^{r-t} t^{s-u} \\ &= 0, \end{aligned} \quad (5.7)$$

since $\sum_{t=0}^r \binom{r}{t} (-1)^{r-t} t^{s-u} = 0$ when $s-u < r$ is a part of the second Stirling number (2.10).

Of course, these equations still hold if we substitute $\lambda^{\frac{1}{\omega}}$ and $\lambda^{\frac{1}{m}}$ by $\lambda^{\frac{-1}{\omega}}$ and $\lambda^{\frac{-1}{m}}$ respectively. As a conclusion, all elements in (5.3) as well as (5.2) belong to $I^\perp = (D(m, d, \xi)/I)^*$.

Finally we prove that (5.2) are linearly independent. Choose a lexicographically ordered set of elements $\{h_{e', i', k', s', l'} \in D(m, d, \xi) \mid (e', i', k', s', l') \in 2 \times \omega \times 2m \times r \times m\}$, where

$$h_{e', i', k', s', l'} := \begin{cases} \frac{1}{l'! \gamma} x^{i'} g^{\frac{k'}{2} + 2s'm + e'm} y^{l'}, & \text{when } 2 \mid k'; \\ \frac{(1-\gamma^{-1})^{l'}}{\xi^{l'^2}} x^{i'} g^{\frac{k'-1}{2} + 2s'm + e'm} u_{l'}, & \text{when } 2 \nmid k'. \end{cases}$$

Our goal is to show that the $4\omega m^2 r \times 4\omega m^2 r$ square matrix

$$A := \left(\langle \psi_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l, h_{e',i',k',s',l'} \rangle \right. \\ \left. \mid (e, i, k, s, l), (e', i', k', s', l') \in 2 \times \omega \times 2m \times r \times m \right)$$

is invertible for fixed $\lambda \frac{1}{\omega}, \lambda \frac{1}{m}$, which would imply the linear independence of (5.2) as well as (5.3). We make following computations: When $2 \mid k'$,

$$\begin{aligned} & \langle \psi_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l, h_{e',i',k',s',l'} \rangle \\ &= \langle \zeta_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} \zeta_{\eta,1}^i \zeta_{1,\gamma}^k E_2^s E_1^l, \frac{1}{l'!_\gamma} x^{i'} g^{\frac{k'}{2} + 2s'm + e'm} y^{l'} \rangle \\ &= \delta_{l',l} \langle \zeta_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} \zeta_{\eta,1}^i \zeta_{1,\gamma}^k E_2^s, x^{i'} g^{\frac{k'}{2} + 2s'm + e'm} \rangle \frac{1}{l'!_\gamma} \langle E_1^l, x^{i'} g^{\frac{k'}{2} + 2s'm + e'm} y^{l'} \rangle \\ &= \delta_{l',l} \lambda^{(-1)^e (\frac{i'}{\omega} + \frac{k'}{2m} + 2s' + e')} \eta^{ii'} \xi^{kk'} \left(\frac{i'}{\omega} + \frac{k'}{2m} + 2s' + e' \right)^s; \end{aligned}$$

When $2 \nmid k'$,

$$\begin{aligned} & \langle \psi_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l, h_{e',i',k',s',l'} \rangle \\ &= \langle \chi_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} \chi_{\eta,1}^i (\xi \chi_{1,\gamma})^k E_2^s E_1^l, \frac{(1 - \gamma^{-1})^{l'}}{\xi^{l'^2}} x^{i'} g^{\frac{k'-1}{2} + 2s'm + e'm} u_{l'} \rangle \\ &= \delta_{l',l} \langle \chi_{\lambda \frac{(-1)^e}{\omega}, \lambda \frac{(-1)^e}{m}} \xi^k E_2^s, x^{i'} g^{\frac{k'-1}{2} + 2s'm + e'm} u_0 \rangle \frac{(1 - \gamma^{-1})^l}{\xi^{l^2}} \langle E_1^l, x^{i'} g^{\frac{k'-1}{2} + 2s'm + e'm} u_l \rangle \\ &= \delta_{l',l} \lambda^{(-1)^e (\frac{i'}{\omega} + \frac{k'-1}{2m} + 2s' + e')} \eta^{ii'} \xi^{kk'} \left(\frac{i'}{\omega} + \frac{k'-1}{2m} + 2s' + e' \right)^s. \end{aligned}$$

One concludes that

$$A := \left(\delta_{l',l} \lambda^{(-1)^e (\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s' + e')} \eta^{ii'} \xi^{kk'} \left(\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s' + e' \right)^s \right. \\ \left. \mid (e, i, k, s, l), (e', i', k', s', l') \in 2 \times \omega \times 2m \times r \times m \right).$$

However, there are following two facts:

- The matrix $(\delta_{l',l} \eta^{ii'} \xi^{kk'} \mid (i, k, l), (i', k', l') \in \omega \times 2m \times m)$ is invertible by Lemma 2.11, since it is the Kronecker product of three invertible matrices;
- For each $(i', k') \in \omega \times 2m$, the matrix

$$\left(\lambda^{(-1)^e (\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s' + e')} \left(\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s' + e' \right)^s \mid (e, s), (e', s') \in 2 \times r \right)$$

is invertible by Corollary 2.14(2).

Thus A is invertible according to Lemma 2.12, and (5.2) or (5.3) is a basis of $(D(m, d, \xi)/I)^*$.

Case 2. Suppose $I = ((g^m - 1)^r)$ for some $r \in \mathbb{N}$. In this case we aim to show that

$$\{(\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l \mid i \in \omega, k \in 2m, s \in r, l \in m\} \quad (5.8)$$

is a linear basis of $2\omega m^2 r$ -dimensional space $(D(m, d, \xi)/I)^*$. We remark that (5.8) is linearly equivalent to

$$\{\zeta_{\eta,1}^i \zeta_{1,\gamma}^j E_2^s E_1^l, \chi_{\eta,1}^i \chi_{1,\gamma}^j E_2^s E_1^l \mid i \in \omega, j \in m, s \in r, l \in m\}. \quad (5.9)$$

Evidently

$$I = \mathbb{k}\{x^i g^j (g^m - 1)^r y^l, x^i g^j (g^m - 1)^r u_l \mid i \in \omega, j \in \mathbb{Z}, l \in m\},$$

since $yg^m = g^m y$ and $u_i g^m = g^{-m} u_i$ for each $i \in m$. Also, $D(m, d, \xi)/I$ has a linear basis

$$\{x^i g^j (g^m - 1)^s y^l, x^i g^j (g^m - 1)^s u_l \mid i \in \omega, j \in m, s \in r, l \in m\},$$

and hence $\dim((D(m, d, \xi)/I)^*) = \dim(D(m, d, \xi)/I) = 2\omega m^2 r$. Similar computations to Equations (5.4) to (5.7) follows that all elements in (5.9) vanish on I , and hence all elements in (5.9) as well as (5.8) belong to $I^\perp = (D(m, d, \xi)/I)^*$.

Finally we prove that (5.8) are linearly independent. Our goal is to show that the $2\omega m^2 r \times 2\omega m^2 r$ square matrix

$$\begin{aligned} A &:= (\langle (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l, h_{0,i',k',s',l'} \rangle \\ &\quad \mid (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times r \times m) \\ &= \left(\delta_{l',l} \eta^{ii'} \xi^{kk'} \left(\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s' \right)^s \mid (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times r \times m \right) \end{aligned}$$

is invertible. This is true due to the same argument in the proof of Proposition 4.3 that A is invertible there.

Case 3. Suppose $I = ((g^m + 1)^r)$ for some $r \in \mathbb{N}$. Let $(-1)^{\frac{1}{\omega}}, (-1)^{\frac{1}{m}}$ be some fixed roots of -1 respectively. In this case we aim to show that

$$\{\psi_{(-1)^{\frac{1}{\omega}}, (-1)^{\frac{1}{m}}} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l \mid i \in \omega, k \in 2m, s \in r, l \in m\} \quad (5.10)$$

is a linear basis of $2\omega m^2 r$ -dimensional space $(D(m, d, \xi)/I)^*$. We remark that (5.10) is linearly equivalent to

$$\{\zeta_{(-1)^{\frac{1}{\omega}}, (-1)^{\frac{1}{m}}} \zeta_{\eta,1}^i \zeta_{1,\gamma}^j E_2^s E_1^l, \chi_{(-1)^{\frac{1}{\omega}}, (-1)^{\frac{1}{m}}} \chi_{\eta,1}^i \chi_{1,\gamma}^j E_2^s E_1^l \mid i \in \omega, j \in m, s \in r, l \in m\}. \quad (5.11)$$

Evidently

$$I = \mathbb{k}\{x^i g^j (g^m + 1)^r y^l, x^i g^j (g^m + 1)^r u_l \mid i \in \omega, j \in \mathbb{Z}, l \in m\},$$

since $yg^m = g^m y$ and $u_i g^m = g^{-m} u_i$ for each $i \in m$. Also, $D(m, d, \xi)/I$ has a linear basis

$$\{x^i g^j (g^m + 1)^s y^l, x^i g^j (g^m + 1)^s u_l \mid i \in \omega, j \in m, s \in r, l \in m\},$$

and hence $\dim((D(m, d, \xi)/I)^*) = \dim(D(m, d, \xi)/I) = 2\omega m^2 r$. Similar computations to Equations (5.4) to (5.7) follows that all elements in (5.11) vanish on I , and hence all elements in (5.11) as well as (5.10) belong to $I^\perp = (D(m, d, \xi)/I)^*$.

Finally we prove that (5.10) are linearly independent. Our goal is to show that the $2\omega m^2 r \times 2\omega m^2 r$ square matrix

$$A := (\langle \psi_{(-1)^{\frac{1}{\omega}}, (-1)^{\frac{1}{m}}} (\zeta_{\eta,1} + \chi_{\eta,1})^i (\zeta_{1,\gamma} + \xi \chi_{1,\gamma})^k E_2^s E_1^l, h_{0,i',k',s',l'} \rangle$$

$$\begin{aligned}
& | (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times r \times m) \\
= & ((-1)^{\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s'} \delta_{l', l} \eta^{ii'} \xi^{kk'} (\frac{i'}{\omega} + \lfloor \frac{k'}{2} \rfloor \frac{1}{m} + 2s')^s \\
& | (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times r \times m)
\end{aligned}$$

is invertible. This is true due to the same argument in the proof of Proposition 4.3 that A is invertible there.

We summarize three cases above in a result that

$$\begin{aligned}
D(m, d, \xi)^\circ &= \sum_{I \in \mathcal{I}_0} (D(m, d, \xi)/I)^* \\
&= \mathbb{k}\{\zeta_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} E_2^s E_1^l, \chi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} E_2^s E_1^l \mid \lambda \frac{1}{\omega}, \lambda \frac{1}{m} \in \mathbb{k}^*, s \in \mathbb{N}, l \in m\}.
\end{aligned}$$

□

Theorem 5.7. (1) $D_\circ(m, d, \xi)$ constructed in Subsection 5.1 is a Hopf algebra;
(2) As a Hopf algebra, $D(m, d, \xi)^\circ$ is isomorphic to $D_\circ(m, d, \xi)$.

Proof. Consider the following map

$$\begin{aligned}
\Theta : D_\circ(m, d, \xi) &\rightarrow D(m, d, \xi)^\circ, \\
Z_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} &\mapsto \zeta_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}}, X_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} \mapsto \chi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}}, F_1 \mapsto E_1, F_2 \mapsto E_2,
\end{aligned}$$

where $\lambda \frac{1}{\omega}, \lambda \frac{1}{m} \in \mathbb{k}^*$. This is a epimorphism of algebras by Lemma 5.2 and Proposition 5.6. Furthermore, Θ would become an isomorphism of Hopf algebras with desired coalgebra structure and antipode, as long as it is injective (since $D(m, d, \xi)^\circ$ is in fact a Hopf algebra).

In order to show that Θ is injective, we aim to show the linear independence of

$$\{\zeta_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} \zeta_{\eta, 1}^i \zeta_{1, \gamma}^j E_2^s E_1^l, \chi_{\lambda \frac{1}{\omega}, \lambda \frac{1}{m}} \chi_{\eta, 1}^i \chi_{1, \gamma}^j E_2^s E_1^l \mid \lambda \in \mathbb{k}^*, i \in \omega, j \in m, s \in \mathbb{N}, l \in m\}$$

in $D(m, d, \xi)^\circ$, where $\lambda \frac{1}{\omega}$ and $\lambda \frac{1}{m}$ are fixed roots of each $\lambda \in \mathbb{k}^*$. By linear independence of elements in (5.3), (5.9) and (5.11), we only need to show that

This is due to the fact that any finite sum of form

$$\begin{aligned}
& (D(m, d, \xi)/((g^m - 1)^r))^* + (D(m, d, \xi)/((g^m + 1)^r))^* \\
& + \sum_{\alpha=1}^N (D(m, d, \xi)/((g^m - \lambda_\alpha)^r (g^m - \lambda_\alpha^{-1})^r))^* \\
= & ((g^m - 1)^r)^\perp + ((g^m + 1)^r)^\perp + \sum_{\alpha=1}^N ((g^m - \lambda_\alpha)^r (g^m - \lambda_\alpha^{-1})^r)^\perp
\end{aligned}$$

is direct, as long as λ_α 's are distinct and not inverses of each other. The reason is the same as Equations (4.6). □

Remark 5.8. The infinite dihedral group \mathbb{D}_∞ is generated by g and x with relations

$$x^2 = 1, \quad xgx = g^{-1},$$

and the finite dual of $\mathbb{k}\mathbb{D}_\infty$ is established by [9]. Note that $\mathbb{k}\mathbb{D}_\infty = D(1, 1, -1)$ and thus we cover the presentation of $(\mathbb{k}\mathbb{D}_\infty)^\circ$. For our convenience, we use our notion to write $(\mathbb{k}\mathbb{D}_\infty)^\circ$ out: It is generated by

$$\zeta_\lambda : g^j x^k \mapsto \delta_{k,0} \lambda^j, \quad \chi_\lambda : g^j x^k \mapsto \delta_{k,1} \lambda^j, \quad E_2 : g^j x^k \mapsto j \quad (j \in \mathbb{Z}, k \in 2)$$

for $\lambda \in \mathbb{k}^*$, with relations

$$\begin{aligned} \zeta_{\lambda_1} \zeta_{\lambda_2} &= \zeta_{\lambda_1 \lambda_2}, \quad \chi_{\lambda_1} \chi_{\lambda_2} = \chi_{\lambda_1 \lambda_2}, \quad \zeta_{\lambda_1} \chi_{\lambda_2} = \chi_{\lambda_1} \zeta_{\lambda_2} = 0, \quad \chi_1 + \zeta_1 = 1, \\ E_2 \zeta_\lambda &= \zeta_\lambda E_2, \quad E_2 \chi_\lambda = \chi_\lambda E_2, \end{aligned}$$

and

$$\begin{aligned} \Delta(\zeta_\lambda) &= \zeta_\lambda \otimes \zeta_\lambda + \chi_\lambda \otimes \chi_{\lambda^{-1}}, \quad \Delta(\chi_\lambda) = \zeta_\lambda \otimes \chi_\lambda + \chi_\lambda \otimes \zeta_{\lambda^{-1}}, \\ \Delta(E_2) &= (\zeta_1 - \chi_1) \otimes E_2 + E_2 \otimes 1, \\ \varepsilon(\zeta_\lambda) &= 1, \quad \varepsilon(\chi_\lambda) = \varepsilon(E_2) = 0, \\ S(\zeta_\lambda) &= \zeta_{\lambda^{-1}}, \quad S(\chi_\lambda) = \chi_{\lambda^{-1}}, \quad S(E_2) = -(\zeta_1 - \chi_1)E_2. \end{aligned}$$

This presentation is equivalent to that in [9, Section 3.1] and the equivalence is given through:

$$\zeta_\lambda \mapsto \frac{1}{2}(\phi_\lambda + \psi_\lambda), \quad \chi_\lambda \mapsto \frac{1}{2}(\phi_\lambda - \psi_\lambda) \quad (\lambda \in \mathbb{k}^*), \quad E_2 \mapsto F,$$

where we used notions of [9] freely.

6. HOPF PAIRING AND CONSEQUENCES

With the help of Sections 3, 4 and 5, we show that there always is a non-degenerate Hopf pairing on each affine prime regular Hopf algebra of GK-dimension one. Using such Hopf pairing, we attempt to consider an infinite-dimensional version of some conclusions which are valid in the finite-dimensional situation.

6.1. Non-Degenerate Hopf Pairing. As mentioned in the introduction, the notion of pairing of a bialgebra or a Hopf algebra was given by Majid [21]:

Definition 6.1. *Let H and H^\bullet be Hopf algebras. A linear map $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$ is called a Hopf pairing (on H), if*

$$\begin{aligned} \text{(i)} \quad \langle ff', h \rangle &= \sum \langle f, h_{(1)} \rangle \langle f', h_{(2)} \rangle, & \text{(ii)} \quad \langle f, hh' \rangle &= \sum \langle f_{(1)}, h \rangle \langle f_{(2)}, h' \rangle, \\ \text{(iii)} \quad \langle 1, h \rangle &= \varepsilon(h), & \text{(iv)} \quad \langle f, 1 \rangle &= \varepsilon(f), \\ \text{(v)} \quad \langle f, S(h) \rangle &= \langle S(f), h \rangle \end{aligned}$$

hold for all $f, f' \in H^\bullet$ and $h, h' \in H$. Moreover, it is said to be non-degenerate, if for any $f \in H^\bullet$ and any $h \in H$,

$$\langle f, H \rangle = 0 \text{ implies } f = 0, \text{ and } \langle H^\bullet, h \rangle = 0 \text{ implies } h = 0.$$

Clearly, the definition follows that there are linear maps

$$\alpha : H^\bullet \rightarrow H^*, \quad f \mapsto \langle f, - \rangle \quad \text{and} \quad \beta : H \rightarrow H^{\bullet*}, \quad h \mapsto \langle -, h \rangle.$$

Furthermore, we know by (ii) in Definition 6.1 that for any $f \in H^\bullet$,

$$M^*(\alpha(f)) = \sum \alpha(f_{(1)}) \otimes \alpha(f_{(2)}) \in H^* \otimes H^*,$$

where M denotes the multiplication on H , and hence the image of α is in fact contained in H° by [23, Lemma 9.1.1].

As a conclusion, Definition 6.1 follows that there are two maps of Hopf algebras

$$\alpha : H^\bullet \rightarrow H^\circ, f \mapsto \langle f, - \rangle \quad \text{and} \quad \beta : H \rightarrow H^{\bullet\circ}, h \mapsto \langle -, h \rangle,$$

which are both injective if and only if the Hopf pairing $\langle -, - \rangle$ is non-degenerate. Thus we might be able to construct non-degenerate Hopf pairing on H , as long as the structure of finite dual H° is determined.

As examples, non-degenerate Hopf pairing on $\mathbb{k}\mathbb{D}_\infty$, $T_\infty(n, v, \xi)$, $B(n, \omega, \gamma)$ and $D(m, d, \xi)$ are constructed as follows, respectively.

Proposition 6.2. *For each affine prime regular Hopf algebra H of GK-dimension one, we can construct a Hopf algebra H^\bullet and a non-degenerate Hopf pairing $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$ as follows: Specifically, keeping the notions used in Sections 3, 4 and 5, we have*

- (1) *The evaluation $\langle -, - \rangle : (\mathbb{k}\mathbb{D}_\infty)^\bullet \otimes \mathbb{k}\mathbb{D}_\infty \rightarrow \mathbb{k}$ is a non-degenerate Hopf pairing, where*

$$\begin{aligned} (\mathbb{k}\mathbb{D}_\infty)^\bullet &= \mathbb{k}\{\zeta_1 E_2^s, \chi_1 E_2^s \mid s \in \mathbb{N}\} \\ &= \mathbb{k}\{(\zeta_1 - \chi_1)^k E_2^s \mid k \in \mathbb{N}, s \in \mathbb{N}\} \subseteq (\mathbb{k}\mathbb{D}_\infty)^\circ. \end{aligned}$$

- (2) *The evaluation $\langle -, - \rangle : T_\infty(n, v, \xi)^\bullet \otimes T_\infty(n, v, \xi) \rightarrow \mathbb{k}$ is a non-degenerate Hopf pairing, where*

$$\begin{aligned} T_\infty(n, v, \xi)^\bullet &= \mathbb{k}\{\omega^j E_2^s E_1^l \mid j \in n, s \in \mathbb{N}, l \in m\} \\ &= \mathbb{k}\{\omega^j E_2^{[s]} E_1^{[l]} \mid j \in n, s \in \mathbb{N}, l \in m\} \subseteq T_\infty(n, v, \xi)^\circ, \end{aligned}$$

$$\text{and } m = \frac{n}{\gcd(n, v)}, E_2^{[s]} E_1^{[l]} = \frac{1}{s! \cdot l! \xi^v} E_2^s E_1^l.$$

- (3) *The evaluation $\langle -, - \rangle : B(n, \omega, \gamma)^\bullet \otimes B(n, \omega, \gamma) \rightarrow \mathbb{k}$ is a non-degenerate Hopf pairing, where*

$$\begin{aligned} B(n, \omega, \gamma)^\bullet &= \mathbb{k}\{\psi_{1, \gamma}^j E_2^s E_1^l \mid j \in n, s \in \mathbb{N}, l \in n\} \\ &= \mathbb{k}\{\psi_{1, \gamma}^j E_2^s E_1^{[l]} \mid j \in n, s \in \mathbb{N}, l \in n\} \subseteq B(n, \omega, \gamma)^\circ, \end{aligned}$$

$$\text{and } E_1^{[l]} = \frac{1}{l! \gamma} E_1^l.$$

- (4) *The evaluation $\langle -, - \rangle : D(m, d, \xi)^\bullet \otimes D(m, d, \xi) \rightarrow \mathbb{k}$ is a non-degenerate Hopf pairing, where*

$$\begin{aligned} D(m, d, \xi)^\bullet &= \mathbb{k}\{\zeta_{1, \gamma}^j E_2^s E_1^l, \chi_{1, \gamma}^j E_2^s E_1^l \mid i \in \omega, j \in m, s \in \mathbb{N}, l \in m\} \\ &= \mathbb{k}\{(\zeta_{1, \gamma} + \xi \chi_{1, \gamma})^k E_2^s E_1^l \mid k \in 2m, s \in \mathbb{N}, l \in m\} \subseteq D(m, d, \xi)^\circ. \end{aligned}$$

Proof. It is not hard to see that all H^\bullet given above are Hopf subalgebras of the corresponding H° and the evaluation gives a Hopf pairing. Thus the remaining task is to show that the evaluation is non-degenerate. We only prove the non-degeneracy in (4) since the others can be proved similarly.

Clearly, the Hopf pairing $\langle -, - \rangle : D(m, d, \xi)^\bullet \otimes D(m, d, \xi) \rightarrow \mathbb{k}$ gives two Hopf algebra maps

$$\alpha : D(m, d, \xi)^\bullet \rightarrow D(m, d, \xi)^\circ \text{ and } \beta : D(m, d, \xi) \rightarrow D(m, d, \xi)^{\bullet\circ},$$

and the map α is just an inclusion. Therefore, to show that $\langle -, - \rangle$ is non-degenerate, we only need to prove the injectivity of β . Recall that in Section 5 that $D(m, d, \xi) = \mathbb{k}\{x^i g^j y^l, x^i g^j u_l \mid i \in \omega, j \in \mathbb{Z}, l \in m\}$. Also,

$$\begin{aligned} \langle (\zeta_{1,\gamma} + \xi\chi_{1,\gamma})^{k'} E_2^{s'} E_1^{l'}, x^i g^j y^l \rangle &= l!_\gamma \cdot \delta_{l,l'} \xi^{2jk'} \left(\frac{i}{\omega} + \frac{j}{m}\right)^{s'}, \\ \langle (\zeta_{1,\gamma} + \xi\chi_{1,\gamma})^{k'} E_2^{s'} E_1^{l'}, x^i g^j u_l \rangle &= \frac{\xi^{l^2}}{(1-\gamma^{-1})^l} \cdot \delta_{l,l'} \xi^{(2j+1)k'} \left(\frac{i}{\omega} + \frac{j}{m}\right)^{s'} \end{aligned}$$

hold for any $k' \in 2m$, $s' \in \mathbb{N}$, $l' \in m$ and $i \in \omega$, $j \in \mathbb{Z}$, $l \in m$. This means that we only need to show the linear independence of

$$\{\beta(x^i g^j y^l), \beta(x^i g^j u_l) \mid i \in \omega, j \in \mathbb{Z}, l \in m\} \subseteq D(m, d, \xi)^{\bullet\circ}.$$

Choose a lexicographically ordered set of elements for any positive integer N : $\{h_{i,k,s,l} \in D(m, d, \xi) \mid (i, k, s, l) \in \omega \times 2m \times \{-N, \dots, N\} \times m\}$, where

$$h_{i,k,s,l} := \begin{cases} \frac{1}{l!_\gamma} x^i g^{\frac{k}{2}+sm} y^l, & \text{when } 2 \mid k; \\ \frac{(1-\gamma^{-1})^l}{\xi^{l^2}} x^i g^{\frac{k-1}{2}+sm} u_l, & \text{when } 2 \nmid k. \end{cases}$$

The desired linear independence would be implied by the invertibility of the following $2\omega m^2(2N+1) \times 2\omega m^2(2N+1)$ matrix

$$\begin{aligned} A : &= (\langle \beta(h_{i,k,s,l}), (\zeta_{1,\gamma} + \xi\chi_{1,\gamma})^{k'} E_2^{i'+s'\omega+N\omega} E_1^{l'} \rangle \\ &\quad \mid (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times \{-N, \dots, N\} \times m) \end{aligned}$$

for any positive integer N . However,

$$\begin{aligned} A &= (\langle (\zeta_{1,\gamma} + \xi\chi_{1,\gamma})^{k'} E_2^{i'+s'\omega+N\omega} E_1^{l'}, h_{i,k,s,l} \rangle \\ &\quad \mid (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times \{-N, \dots, N\} \times m) \\ &= (\delta_{l,l'} \xi^{kk'} \left(\frac{i}{\omega} + \lfloor \frac{k}{2} \rfloor \frac{1}{m} + s\right)^{i'+s'\omega+N\omega} \mid (i, k, s, l), (i', k', s', l') \in \omega \times 2m \times \{-N, \dots, N\} \times m) \end{aligned}$$

is similar to the matrix

$$\begin{aligned} &(\delta_{l,l'} \xi^{kk'} \left(\lfloor \frac{k}{2} \rfloor \frac{1}{m} + \frac{s\omega+i}{\omega}\right)^{s'\omega+i'+N\omega} \mid (k, l, s, i), (k', l', s', i') \in 2m \times m \times \{-N, \dots, N\} \times \omega) \\ &= (\delta_{l,l'} \xi^{kk'} \left(\lfloor \frac{k}{2} \rfloor \frac{1}{m} + \frac{t}{\omega}\right)^{t'+N\omega} \mid (k, l, t), (k', l', t') \in 2m \times m \times \{-N\omega, \dots, (N+1)\omega-1\}), \end{aligned}$$

which is invertible according to Lemma 2.12, because of the invertibility of

$$(\delta_{l,l'} \xi^{kk'} \mid (k, l), (k', l') \in 2m \times m)$$

and

$$\left(\left(\lfloor \frac{k}{2} \rfloor \frac{1}{m} + \frac{t}{\omega}\right)^{t'+N\omega} \mid t, t' \in \{-N\omega, \dots, (N+1)\omega-1\}\right)$$

for each $k \in 2m$. □

For a general infinite-dimensional Hopf algebra H , there may be *no* H^\bullet such that there is a non-degenerate Hopf pairing between H and H^\bullet (see [23, Example 9.1.5(1)] for example). In fact, the existence of non-degenerate Hopf pairings over a Hopf algebra H is equivalent to require that H is residually finite-dimensional. Recall that an algebra H is said to be *residually finite-dimensional*, if there is a family $\{\pi_i\}$ of finite-dimensional \mathbb{k} -representations of H such that $\bigcap_i \text{Ker}(\pi_i) = 0$. See [23, Definition 9.2.8] for example.

Lemma 6.3. *Let H be a Hopf algebra over an arbitrary field \mathbb{k} . Then there is a non-degenerate Hopf pairing $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$, if and only if H is residually finite-dimensional.*

Proof. This is followed by [23, Proposition 9.2.10]. □

However, our constructions for H^\bullet in Proposition 6.2 satisfy further conditions, which might be crucial in a sense. Before that, we recall in [26, Definition 15.4.4] that a non-cosemisimple pointed Hopf algebra H is called *minimal-pointed*, if each of its proper Hopf subalgebras must be cosemisimple.

Proposition 6.4. *Suppose that H is any of $\mathbb{k}\mathbb{D}_\infty$, $T_\infty(n, v, \xi)$, $B(\omega, n, \gamma)$ and $D(m, d, \xi)$, and H^\bullet denotes the Hopf algebra constructed in Proposition 6.2. Then H^\bullet is a minimal (under inclusion) Hopf subalgebra of H° such that the evaluation $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$ is a non-degenerate Hopf pairing.*

Proof. Since H is infinite-dimensional, it is sufficient to show that H^\bullet is the only infinite-dimensional Hopf subalgebra of itself.

(1) For the case when $H = \mathbb{k}\mathbb{D}_\infty$, we find that $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is a minimal-pointed Hopf algebra (see [26, Proposition 15.4.6] and [25, Proposition 2]) with the finite-dimensional coradical. Thus $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is the only infinite-dimensional Hopf subalgebra as desired.

(2) For the case when $H = T_\infty(n, v, \xi)$, note that $K := \mathbb{k}\{\omega^j E_1^l \mid j, l \in n\}$ is a finite-dimensional normal Hopf subalgebra of $H^\bullet = T_\infty(n, v, \xi)^\bullet$. The quotient Hopf algebra $H^\bullet / K^+ H^\bullet = \mathbb{k}[\overline{E_2}]$ is the polynomial Hopf algebra, which is minimal-pointed too. Therefore, any infinite-dimensional Hopf subalgebra of $T_\infty(n, v, \xi)^\bullet$ must contain the element E_2 . By the coproduct of E_2 , it follows that $T_\infty(n, v, \xi)^\bullet$ is a minimal infinite-dimensional Hopf algebra.

(3) The case when $H = B(n, \omega, \gamma)$ is completely similar to the case $T_\infty(n, v, \xi)$, due to the facts that $K := \mathbb{k}\{\psi_{1,\gamma}^j E_1^l \mid j, l \in n\}$ is a finite-dimensional normal Hopf subalgebra, and the quotient $H^\bullet / K^+ H^\bullet$ is the polynomial Hopf algebra.

(4) The case when $H = D(m, d, \xi)$ is similar, too. Note that the finite-dimensional normal Hopf subalgebra of $D(m, d, \xi)^\bullet$ is chosen as a generalized Taft algebra $\mathbb{k}\{(\zeta_{1,\gamma} + \xi\chi_{1,\gamma})^k E_1^l \mid k \in 2m, l \in m\}$, and the responding quotient is also the polynomial Hopf algebra. □

6.2. Properties of H^\bullet and Consequences. For possible future applications, we might consider Hopf pairings $H^\bullet \otimes H \rightarrow \mathbb{k}$ with additional requirements (including the non-degeneracy), which are listed as follows:

- (HP1) The pairing is non-degenerate;
- (HP2) H^\bullet and H are both affine and noetherian;
- (HP3) $\text{GKdim}(H^\bullet) = \text{GKdim}(H)$.

Furthermore, we might also wish H^\bullet would have certain properties dual to some of H , for examples in this paper:

- (HP4) H^\bullet is indecomposable as a coalgebra, when H is prime as an algebra.

These considerations motivate us to discuss some properties of H^\bullet in the following. Recall that a coalgebra is indecomposable if and only if it is *link-indecomposable* (see [24, Corollary 2.2]). Besides, a ring is called *regular* if it has finite global dimension.

Proposition 6.5. *For affine prime regular Hopf algebras H of GK-dimension one, consider Hopf algebras H^\bullet constructed in Proposition 6.2. We have*

- (1) *All the Hopf algebras H^\bullet have GK-dimension one.*
- (2) *All the Hopf algebras H^\bullet are pointed and (link-)indecomposable as coalgebras.*
- (3) *All the Hopf algebras H^\bullet are noetherian.*
- (4) *The Hopf algebra $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is regular while $T_\infty(n, v, \xi)^\bullet$, $B(n, \omega, \gamma)^\bullet$ and $D(m, d, \xi)^\bullet$ are not when $n, m \geq 2$.*

Proof. (1) and (2) are clear by our constructions for H^\bullet .

(3) This is because every H^\bullet constructed is a finitely generated module (on both sides) over its noetherian subalgebra $\mathbb{k}[E_2]$.

(4) The regularity of $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is a direct consequence of the Hilbert Theorem on Syzygies (see [27, Theorem 8.36] e.g.). Recall that as an algebra, $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is generated by $\zeta_1 - \chi_1$ and E_2 with relations

$$(\zeta_1 - \chi_1)^2 = 1, \quad E_2(\zeta_1 - \chi_1) = (\zeta_1 - \chi_1)E_2.$$

Clearly, $(\mathbb{k}\mathbb{D}_\infty)^\bullet$ is the polynomial algebra over the subalgebra

$$\mathbb{k}\{1, \zeta_1 - \chi_1\} \cong \mathbb{k}\mathbb{Z}_2$$

with the indeterminate E_2 commuting with all the coefficients. However, $\mathbb{k}\mathbb{Z}_2$ is semisimple and hence regular. Thus $\text{gldim}((\mathbb{k}\mathbb{D}_\infty)^\bullet) = \text{gldim}(\mathbb{k}\mathbb{Z}_2) + 1 < \infty$, where gldim denotes the global dimension.

Next we show that $B(n, \omega, \gamma)^\bullet$ is not regular when $n \geq 2$ and the non-regularity for $T_\infty(n, v, \xi)^\bullet$ can be proved similarly. Recall that as an algebra, $B(n, \omega, \gamma)^\bullet$ is generated by $\psi_{1,\gamma}$, E_2 and E_1 with relations

$$\psi_{1,\gamma}^n = 1, \quad E_1^n = 0, \quad E_2\psi_{1,\gamma} = \psi_{1,\gamma}E_2, \quad E_1\psi_{1,\gamma} = \gamma\psi_{1,\gamma}E_1, \quad E_1E_2 = E_2E_1 + \frac{1}{n}E_1.$$

Note in Proposition 6.2(3) that

$$B(n, \omega, \gamma)^\bullet = \mathbb{k}\{\psi_{1,\gamma}^j E_2^s E_1^l \mid j \in n, s \in \mathbb{N}, l \in n\}$$

$$= \mathbb{k}\{E_2^s E_1^l \psi_{1,\gamma}^j \mid s \in \mathbb{N}, l \in n, j \in n\}.$$

Since $B(n, \omega, \gamma)^\bullet$ is a free $\mathbb{k}\langle E_1 \rangle$ -module by Poincaré-Birkhoff-Witt theorem, if $B(n, \omega, \gamma)^\bullet$ has finite global dimension, then the projective dimension $\text{prdim}_{\mathbb{k}\langle E_1 \rangle}(\mathbb{k}) < \infty$. However, this is not possible since $\mathbb{k}\langle E_1 \rangle$ is a finite-dimensional commutative local, but not a field.

Finally, let's show the non-regularity of $D(m, d, \xi)^\bullet$. By Proposition 6.2(4), we find that

$$\begin{aligned} D(m, d, \xi)^\bullet &= \mathbb{k}\{\zeta_{1,\gamma}^j E_2^s E_1^l, \chi_{1,\gamma}^j E_2^s E_1^l \mid j \in m, s \in \mathbb{N}, l \in m\} \\ &= \mathbb{k}\{\zeta_{1,\gamma}^j E_2^s E_1^l \mid j \in m, s \in \mathbb{N}, l \in m\} \\ &\quad \oplus \mathbb{k}\{\chi_{1,\gamma}^j E_2^s E_1^l \mid j \in m, s \in \mathbb{N}, l \in m\} \end{aligned}$$

is a direct sum of ideals. The former ideal is isomorphic to $B(m, \omega, \gamma)^\bullet$. and thus it is not regular as a ring when $m \geq 2$. Therefore,

$$\text{gldim}(D(m, d, \xi)^\bullet) \geq \text{gldim}(B(m, \omega, \gamma)^\bullet) = \infty.$$

□

Thus concluding Propositions 6.2 and 6.5, we find that

Corollary 6.6. *For affine prime regular Hopf algebras H of GK-dimension one, the Hopf pairings $\langle -, - \rangle : H^\bullet \otimes H \rightarrow \mathbb{k}$ constructed in Proposition 6.2 satisfy the requirements (HP1) to (HP4).*

Remark 6.7. *Due to Proposition 6.4, if one verifies that H^\bullet is exactly the (link-)indecomposable component of H° containing the unit element, then H^\bullet would be the unique Hopf subalgebra of H° such that (HP1) to (HP4) hold.*

Remark 6.8. Takeuchi [31] defined a quantum group G to be a triple $(A, U, \langle -, - \rangle)$ where $\langle -, - \rangle : A \otimes U \rightarrow \mathbb{k}$ is a Hopf pairing. By Proposition 6.2, for each affine prime regular Hopf algebra H of GK-dimension one, one can get a quantum group in Takeuchi's sense naturally.

A classical result by Larson and Radford [15, Theorem 3.3] states that a finite-dimensional Hopf algebra H in characteristic 0 is semisimple, if and only if H^* is semisimple. Note that the regularity can be regarded as an infinite-dimensional analogue for the semisimplicity. However according to Proposition 6.5(3), it is not true that H is regular if and only if H^\bullet is regular for a non-degenerate Hopf pairing $H^\bullet \otimes H \rightarrow \mathbb{k}$ (or a quantum group). This might be a version negating the semisimplicity result by Larson and Radford in infinite-dimensional cases.

Question 6.9. For a general infinite-dimensional Hopf algebra H which is residually finite-dimensional, natural questions are: When does a minimal Hopf algebra H^\bullet forming a non-degenerate Hopf pairing over H exist? When are such minimal Hopf algebras H^\bullet unique and of the same GK-dimensions with H ?

ACKNOWLEDGEMENTS

The authors would like to thank Doctor Ruipeng Zhu for discussions on additional techniques to estimate the regularity for Hopf algebras appeared, as well as the condition for the existence of non-degenerate Hopf pairings.

REFERENCES

- [1] N. Andruskiewitsch, J. Cuadra and P. Etingof, “On two finiteness conditions for Hopf algebras with nonzero integral”, *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* **14**:2 (2015), 401-440.
- [2] G. M. Bergman, “The diamond lemma for ring theory”, *Adv. Math.* **29**:2 (1978), 178-218.
- [3] K. A. Brown and M. Couto, “Affine commutative-by-finite Hopf algebras”, *J. Algebra* **573**:1 (2021), 56-94.
- [4] K. A. Brown, M. Couto, and A. Jahn, “The finite dual of commutative-by-finite Hopf algebras”, preprint, arXiv:2105.13874
- [5] K. A. Brown and J. J. Zhang, “Prime regular Hopf algebras of GK-dimension one”, *Proc. Lond. Math. Soc.* **101**:1 (2010), 260-302.
- [6] K. A. Brown and J. J. Zhang, “Survey on Hopf algebras of GK-dimension 1 and 2”, Hopf algebras, tensor categories and related topics, 43-62, *Contemp. Math.*, **771**, Amer. Math. Soc., 2021.
- [7] W. Chin and I. M. Musson, “Hopf algebra duality, injective modules and quantum groups”, *Comm. Algebra* **22**:12 (1994), 4661-4692.
- [8] M. Couto, “Commutative-by-finite Hopf algebras and their finite dual”, thesis, University of Glasgow (2019). <http://theses.gla.ac.uk/74418/>
- [9] F. Ge and G. Liu, “A combinatorial identity and the finite dual of infinite dihedral group algebra”, *Mathematika* **67** (2021), 498-513.
- [10] K. R. Goodearl and J. J. Zhang, “Noetherian Hopf algebra domains of Gelfand-Kirillov dimension two”. *J. Algebra* **324**:11 (2010), 3131-3168.
- [11] R. G. Heyneman and M. E. Sweedler, “Affine Hopf algebras. I”, *J. Algebra* **13** (1969), 192-241.
- [12] A. Jahn, “The finite dual of crossed products”, thesis, University of Glasgow (2015). <http://theses.gla.ac.uk/6158/>
- [13] C. Kassel, *Quantum groups*, Grad. Texts Math. **155**, Springer-Verlag, 1995.
- [14] T. Y. Lam, *Lectures on modules and rings*, Grad. Texts Math. **189**, Springer-Verlag, 1999.
- [15] R. G. Larson and D. E. Radford, “Finite-dimensional cosemisimple Hopf algebras in characteristic 0 are semisimple”, *J. Algebra* **117**:2 (1988), 267-289.
- [16] J. H. van Lint and R. M. Wilson, *A course in combinatorics*, second edition. Cambridge Univ. Press, 2001.
- [17] G. Liu, “A note on the global dimension of smash products”, *Comm. Algebra* **33**:8 (2005), 2625-2627.
- [18] G. Liu, “On Noetherian affine prime regular Hopf algebras of Gelfand-Kirillov dimension 1”, *Proc. Amer. Math. Soc.* **137**:3 (2009), 777-785.
- [19] G. Liu, “A classification result on prime Hopf algebras of GK-dimension one”, *J. Algebra* **547** (2020), 579-667.
- [20] D.-M. Lu, Q.-S. Wu and J. J. Zhang, “Homological integral of Hopf algebras”, *Trans. Amer. Math. Soc.* **359**:10 (2007), 4945-4975.
- [21] S. Majid, “Quasitriangular Hopf algebras and Yang-Baxter equations”, *Internat. J. Modern Phys. A* **5**:1 (1990), 1-91.
- [22] J. W. Milnor and J. C. Moore, “On the structure of Hopf algebras”, *Ann. of Math.* **81** (1965), 211-264.
- [23] S. Montgomery, *Hopf algebras and their actions on rings*, CBMS Regional Conference Series in Mathematics **82**, American Mathematical Society, Providence, RI, 1993.
- [24] S. Montgomery, “Indecomposable coalgebras, simple comodules, and pointed Hopf algebras”, *Proc. Amer. Math. Soc.* **123**:8 (1995), 2343-2351.

- [25] D. E. Radford, “Finite-dimensional simple-pointed Hopf algebras”, *J. Algebra* **211**:2 (1999), 686-710.
- [26] D. E. Radford, *Hopf algebras*, Series on Knots and Everything, **49**, World Scientific Publishing Co. Pte. Ltd., 2012.
- [27] J. J. Rotman, *An introduction to homological algebra*, second edition. Universitext. Springer, 2009.
- [28] L. W. Small, J. T. Stafford, and R. B. Warfield, “Affine algebras of Gel’fand-Kirillov dimension one are PI”, *Math. Proc. Cambridge Philos. Soc.* **97**:3 (1985), 407-414.
- [29] M. E. Sweedler, *Hopf Algebras*, Mathematics Lecture Note Series W. A. Benjamin, Inc., 1969.
- [30] M. Takeuchi, “Hopf algebra techniques applied to the quantum group $U_q(sl(2))$ ”, Deformation theory and quantum groups with applications to mathematical physics (Amherst, MA, 1990), 309-323, *Contemp. Math.* **134** (1992), Amer. Math. Soc., Providence, RI.
- [31] M. Takeuchi, “Some topics on $GL_q(n)$ ”, *J. Algebra* **147**:2 (1992), 379-410.
- [32] D.-G. Wang, J. J. Zhang and G.-B. Zhuang, “Hopf algebras of GK-dimension two with vanishing Ext-group”, *J. Algebra* **388** (2013), 219-247.
- [33] D.-G. Wang, J. J. Zhang and G.-B. Zhuang, “Connected Hopf algebras of Gelfand-Kirillov dimension four”, *Trans. Amer. Math. Soc.* **367**:8 (2015), 5597-5632.
- [34] J. Wu, “Note on the coradical filtration of $D(m, d, \xi)$ ”, *Comm. Algebra* **44**:11 (2016), 4844-4850.
- [35] J. Wu, G. Liu, and N. Ding, “Classification of affine prime regular Hopf algebras of GK-dimension one”, *Adv. Math.* **296** (2016), 1-54.
- [36] G.-B. Zhuang, “Properties of pointed and connected Hopf algebras of finite Gelfand-Kirillov dimension”, *J. Lond. Math. Soc.* (2) **87**:3 (2013), 877-898.

SCHOOL OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU 311121, CHINA

Email address: kqli@hznu.edu.cn

DEPARTMENT OF MATHEMATICS, NANJING UNIVERSITY, NANJING 210093, CHINA

Email address: gxliu@nju.edu.cn