Bayesian Filtering for Nonlinear Stochastic Systems Using Holonomic Gradient Method with Integral Transform

Tomoyuki Iori and Toshiyuki Ohtsuka

Abstract— This paper proposes a symbolic-numeric Bayesian filtering method for a class of discrete-time nonlinear stochastic systems to achieve high accuracy with a relatively small online computational cost. The proposed method is based on the holonomic gradient method (HGM), which is a symbolicnumeric method to evaluate integrals efficiently depending on several parameters. By approximating the posterior probability density function (PDF) of the state as a Gaussian PDF, the update process of its mean and variance can be formulated as evaluations of several integrals that exactly take into account the nonlinearity of the system dynamics. An integral transform is used to evaluate these integrals more efficiently using the HGM compared to our previous method. Further, a numerical example is provided to demonstrate the efficiency of the proposed method over other existing methods.

I. INTRODUCTION

In most engineering applications, knowledge of the state of a dynamical system is required for monitoring or control via feedback. The optimal filtering theory was developed to estimate the current state of a system using the history of its outputs that are observable but contaminated by random noise. Following the success of the Kalman filter (KF) [1], [2] for linear systems with Gaussian noise, optimal filtering theory has been extended to applications involving nonlinear systems as well as non-Gaussian noise [3]–[7].

Bayesian filtering is a general framework of optimal filtering. Although numerous problems can be formulated as the Bayesian filtering problems, they are difficult to implement as algorithms on computers, particularly for the real-time applications in systems and control. The Monte-Carlo scheme, which was first introduced for the particle filter (PF) [8], [9] and subsequently used in other filtering algorithms, such as [7], is a powerful tool to compute integrations accompanied by statistical operations, such as marginalizations and expectations in nonlinear Bayesian filtering problems. Moreover, the nonlinearity of the system dynamics can be exactly accounted for by updating the particles according to the dynamics. To reap these benefits of the Monte-Carlo scheme, however, the number of particles must be sufficiently large, which requires heavy computational resources and may be unacceptable for applications with short sampling intervals. Another approach involves extending the KF to nonlinear cases such as the extended KF (EKF) [10], unscented KF (UKF) [11], cubature KF (CKF) [12], and Gauss-Hermite KF (GHKF) [13]. In all these methods, the posterior probability density function (PDF) is assumed to be or approximated as a Gaussian PDF. Under this Gaussian approximation, the marginalizations and expectations of the nonlinear functions, which are required to update the posterior PDF, can be efficiently computed using the Gauss quadrature rules. However, in exchange for efficiency, the Gauss quadrature rules cannot provide exact values of the integrals, which implies that the nonlinearity of the system dynamics can only be considered in the approximate sense. Thus, there is still room to extend the boundary of the trade-off between the exact consideration of nonlinearity and reduction of computational cost.

In nonlinear cases, the integrals in the posterior PDF are ones of general nonlinear functions that depend on parameters such as the observed output, which are both analytically and numerically intractable to evaluate. In our previous work [14], a symbolic-numeric method called the holonomic gradient method (HGM) [15] was used to evaluate such integrals efficiently. By utilizing the symbolic computation in terms of differential operators, we can perform integrations offline while considering the nonlinearity of the system exactly. The state estimation is then reduced to solving a set of initial-value problems (IVPs) of nonlinear ordinary differential equations (ODEs) online, which can be efficiently accomplished with numerical integration methods such as the fourth order Runge-Kutta (RK4) or Adams-Bashforth-Moulton predictor-corrector (ADM4) methods. However, our previous method requires solving an IVP for each component of the mean and variance of the current posterior PDF in the one-step estimation. This incurs a high computational cost for the online computation and is unacceptable for short sampling intervals.

In this work, we introduce an integral transform to overcome the computational limitations of our previous method. We can efficiently derive all the components of the mean and variance from the integral transform and its derivatives using the HGM because the integral transform is similar to the moment generating function. Hence, the number of IVPs solved in the one-step estimation can be decreased, which reduces the online computational cost.

Notations: For the field of real numbers **R** and a vector of indeterminates $X = [X_1 \cdots X_n]^{\top}$, $\mathbf{R}(X)$ denotes the field of rational functions in the components of X over **R**. $\partial_X := [\partial_{X_1} \cdots \partial_{X_n}]^{\top}$ denotes a vector of differential operators, where $\partial_{X_i} = \partial/\partial X_i$. We abbreviate ∂_{X_i} by ∂_i if X is clearly specified according to the context. For a multi-index vector $d = [d_1 \cdots d_n]^{\top} \in \mathbf{Z}_{\geq 0}^n$, X^d and ∂^d denote $X_1^{d_1} \cdots X_n^{d_n}$ and $\partial_1^{d_1} \cdots \partial_n^{d_n}$, respectively. The

This work was partly supported by JSPS KAKENHI Grant Numbers JP18J22093 and JP15H02257.

T. Iori and T. Ohtsuka are with the Department of Systems Science, Graduate School of Informatics, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto 606–8501, Japan iori@sys.i.kyoto-u.ac.jp, ohtsuka@i.kyoto-u.ac.jp

symbol $\mathcal{R}_n := \mathbf{R}(X)\langle \partial \rangle$ denotes the noncommutative ring of differential operators with coefficients in $\mathbf{R}(X)$. The subscript n of \mathcal{R}_n is omitted if it is clear from the context. We denote the action of an element $l \in \mathcal{R}_n$ on a sufficiently smooth function $\alpha(X) = \alpha(X_1, \ldots, X_n)$ as $l \bullet \alpha(X)$; for instance, $\partial_i \bullet \alpha(X) = \partial \alpha / \partial X_i(X)$. The left ideal generated by a finite set of differential operators $\{l_1, \ldots, l_s\} \subset \mathcal{R}_n$ is defined as a set of differential operators $\{l_1, \ldots, l_s\} \subset \mathcal{R}_n$ is defined as a set of differential operators $\{l_1, \ldots, l_n\} := \{a_1 \cdot l_1 + \cdots + a_s \cdot l_s \mid a_1, \ldots, a_s \in \mathcal{R}_n\}$. We omit the adjective "left" and simply refer to them as ideals because all the ideals in this work are left ideals. For an ideal $I = \langle l_1, \ldots, l_s \rangle$, the set $\{l_1, \ldots, l_s\}$ is called a basis of I. We say that a differential operator $l \in \mathcal{R}$ annihilates a function α if $l \bullet \alpha = 0$. Similarly, we say that an ideal $I \subset \mathcal{R}$ annihilates α if $l \bullet \alpha = 0$ for all $l \in I$. We denote the set of all positive definite $n \times n$ matrices by PD(n).

II. MATHEMATICAL PRELIMINARIES

In this work, we consider the Bayesian filtering problem for a class of nonlinear functions called *holonomic functions* and use the *HGM* [15] to evaluate some integrals of the holonomic functions efficiently. Before describing the problem setting, we briefly introduce the notion of holonomic functions and the holonomic gradient method.

A. Holonomic functions

A holonomic function can be defined by a set of differential operators called a zero-dimensional ideal in \mathcal{R}_n .

Definition 1: [16] Let I be an ideal in $\mathcal{R}_n = \mathbf{R}(X)\langle \partial \rangle$. We call I a zero-dimensional ideal in \mathcal{R}_n if the quotient ring \mathcal{R}_n/I is a finite-dimensional vector space over $\mathbf{R}(X)$.

Definition 2: [16] An analytic function is called a *holonomic function* if there exists a zero-dimensional ideal annihilating the function.

Remark 1: Most nonlinear functions encountered in systems theory problems are holonomic functions. For example, polynomials, rational functions, exponentials, logarithms, trigonometric functions, as well as their sums and products are holonomic [17].

Holonomic functions have two remarkable properties: closure property and finiteness of the partial derivatives.

Theorem 1 (Closure property): [17] For holonomic functions $\alpha(X)$ and $\beta(X)$ in $X = [X_1 \cdots X_n]^{\top}$, the following properties hold.

- 1) The product $\alpha \cdot \beta$ is also a holonomic function.
- 2) Assume that $\alpha = \alpha(X_1, \ldots, X_n)$ is infinitely differentiable on \mathbf{R}^n and is rapidly decreasing with respect to X_n , meaning that $\lim_{X_n\to\infty} X_n^s \partial_n^t \alpha(X) = 0$ ($s, t \in \mathbb{Z}_{\geq 0}$) for any $[X_1 \cdots X_{n-1}]^\top \in \mathbf{R}^{n-1}$. Then, the integral $\int_{-\infty}^{\infty} \alpha(X) dX_n$ is also a holonomic function of $[X_1 \cdots X_{n-1}]^\top$.

By virtue of the closure property, when a given function is described as a product of two nonlinear functions, we can verify that the product is holonomic by confirming that each of the two nonlinear functions is holonomic; the same is true for an integral of a nonlinear function. Moreover, if we have bases of zero-dimensional ideals for the two nonlinear functions, a basis of a zero-dimensional ideal for the product can be computed symbolically using multiplications in terms of differential operators, and the same is true for an integral (see [17] for details).

On the other hand, the finiteness of the partial derivatives is stated as the following theorem.

Theorem 2 (Finiteness of the partial detivatives): [16] For a holonomic function $\alpha(X)$ in $X = [X_1 \dots X_n]^{\top}$, there exist a finite number of its partial derivatives $\partial^{d_1}\alpha(X), \dots, \partial^{d_{q-1}}\alpha(X)$ such that a *q*-dimensional vectorvalued function $Q(X) := [\alpha(X) \ \partial^{d_1}\alpha(X) \ \cdots \ \partial^{d_{q-1}}\alpha(X)]^{\top}$ satisfies the following PDEs:

$$\partial_{X_i}Q(X) = A_{X_i}(X)Q(X) \qquad (i = 1, \dots, n), \tag{1}$$

where each $A_{X_i}(X) \in \mathbf{R}(X)^{q \times q}$ is a matrix-valued function whose components are all rational functions of X.

The set of PDEs (1) is called the *Pfaffian system* for the holonomic function $\alpha(X)$. Theorem 2 shows that each partial derivative of Q(X) and consequently that of $\alpha(X)$ as the first component of Q(X) can be expressed as a linear combination of the components of Q(X) with coefficients in $\mathbf{R}(X)$; in other words, all the partial derivatives of $\alpha(X)$ are elements of a finite-dimensional vector space spanned by the components of Q(X) over $\mathbf{R}(X)$.

Let *I* be a zero-dimensional ideal annihilating a holonomic function $\alpha(X)$. If we have a basis of *I*, we can find the differential operators $\partial^{d_1}, \ldots, \partial^{d_{q-1}}$ in Theorem 2 and symbolically compute the Pfaffian system, that is, the matrices $A_{X_i}(X)$ $(i = 1, \ldots, n)$ in (1) [16].

B. Holonomic gradient method

The HGM [15] is an approach for evaluating a holonomic function at a given point using a basis of a zero-dimensional ideal annihilating it. In particular, the HGM is suitable for a holonomic function such that i) its explicit expression consists of relatively simple nonlinear functions using integrations and multiplications, and ii) its evaluation via the explicit expression is time-consuming and thus unacceptable.

Consider the evaluation of a holonomic function $\alpha(X)$, which is prescribed by such a complex expression, at a specific point \hat{X} . The HGM consists of the following three steps: a) computation of the Pfaffian system (1) from the explicit expression of $\alpha(X)$, b) computation of an initial vector $Q(X_{\text{init}})$ at an initial point X_{init} , and c) integration of the Pfaffian system along a path from X_{init} to \hat{X} .

a) Computation of the Pfaffian system: A basis of a zero-dimensional ideal annihilating $\alpha(X)$ is needed to compute the Pfaffian system for $\alpha(X)$. Although the expression of $\alpha(X)$ is complex, we can symbolically compute a basis of a zero-dimensional ideal for $\alpha(X)$ based on those for all the simple components comprising $\alpha(X)$ (see, e.g., [14], [16], [17]). This computation requires the notion of the *holonomic ideals* in Weyl algebra, which are roughly the counterparts of the zero-dimensional ideals in \mathcal{R} [16].

b) Computation of the initial vector: In computing the Pfaffian system, we obtain a finite set of differential operators $\{\partial^{d_1}, \ldots, \partial^{d_{q-1}}\}$. By applying these differential operators to the explicit expression of $\alpha(X)$, we can obtain a complex

but explicit expression for $Q(X) = [\alpha \ \partial^{d_1} \alpha \ \cdots \ \partial^{d_{q-1}} \alpha]^\top$. By evaluating the expression of Q at a fixed point X_{init} , a vector $Q(X_{\text{init}})$ can be numerically computed given sufficient computational resources.

c) Integration of the Pfaffian system along an integration path: For a specific point \hat{X} , we can set an integration path $X: [0,1] \ni s \mapsto X(s) \in \mathbf{R}^n$ such that $X(0) = X_{\text{init}}$ and $X(1) = \hat{X}$. The vector $Q(\hat{X})$ and first component $[Q(\hat{X})]_1 = \alpha(\hat{X})$ can then be computed by solving the following IVP:

$$\frac{dQ}{ds} = \frac{\partial Q}{\partial X}\frac{dX}{ds} = \sum_{i=1}^{n} A_{X_i}(X(s))Q(X(s))\frac{dX_i}{ds},$$
$$Q(X(0)) = Q(X_{\text{init}}). \quad (2)$$

Note that certain components of the coefficient matrices $A_{X_i}(X(s))$ may have denominators that vanish at certain points $\tilde{s} \in [0, 1]$ and that the integration fails at such instances. The zero set of the least common multiple of all the denominators in $A_{X_i}(X)$ (i = 1, ..., n) is called the *singular locus* of the Pfaffian system, which should be avoided when choosing the initial point and integration path.

The advantage of the HGM for state estimation is that we can perform steps a) and b) *offline*, that is, before commencing the estimation process. Therefore, the online process is reduced to solving the ODE (2), which is easier than evaluating the original expression of $\alpha(X)$ when it is too complex for numerical evaluation.

III. PROBLEM SETTING

In the Bayesian filtering approach for discrete-time nonlinear systems, the posterior PDF of the state x_k at time step k conditional on the output history $y_{[0:k]}$ from time steps 0 to k is recursively updated via the following equations [18].

$$p(x_{k} | y_{[0:k]}) = \frac{p(x_{k}, y_{k} | y_{[0:k-1]})}{p(y_{k} | y_{[0:k-1]})},$$

$$p(x_{k}, y_{k} | y_{[0:k-1]}) = p(y_{k} | x_{k})$$

$$\times \int_{\mathbf{R}^{n}} p(x_{k} | x_{k-1})p(x_{k-1} | y_{[0:k-1]})dx_{k-1},$$

$$p(y_{k} | y_{[0:k-1]}) = \int_{\mathbf{R}^{n}} p(x_{k}, y_{k} | y_{[0:k-1]})dx_{k}.$$
(3)

We assume that the system dynamics and observation process are defined by the following state and observation equations.

$$x_k = f(x_{k-1}, u_k) + w_k,$$
(4)

$$y_k = h(x_k) + v_k, \tag{5}$$

where w_k and v_k denote the system and observation noises that are assumed to be independent and identically distributed with the PDFs $p_w(w_k)$ and $p_v(v_k)$, respectively. Using the rule of transformation of PDFs [10], the conditional PDFs $p(x_k | x_{k-1})$ and $p(y_k | x_k)$ in (3) can be rewritten using functions f, h, p_w , and p_v as follows:

$$p(x_k \mid x_{k-1}, u_k) = p_w(x_k - f(x_{k-1}, u_k)),$$
(6)

$$p(y_k \mid x_k) = p_v(y_k - h(x_k)),$$
(7)

where the former PDF is conditional on not only x_{k-1} but also u_k owing to (4). By substituting (6) and (7) into (3), we can describe the update law of Bayesian filtering using f, h, p_w , and p_v .

Since Bayesian filtering is a recursive algorithm, we can focus on the one-step update of the posterior PDF and omit the subscript *k*. Moreover, the past history of outputs $y_{[0:k-1]}$ has nothing to do with the update at time step *k*; it only appears in the previous posterior PDF $p(x_{k-1} | y_{[0:k-1]})$, so we also omit the past history $y_{[0:k-1]}$. Thus, (3) can be summarized as

$$p(x \mid y, u) = \frac{p_{xy}(x, y \mid u)}{\int_{\mathbf{R}^n} p_{xy}(x, y \mid u) dx},$$
(8)

where x, y, and u denote the current state, output, and input, respectively; x^- denotes the previous state, and $p_{xy}(x, y \mid u)$ denotes the joint PDF of x and y defined as

$$p_{xy}(x, y \mid u) \coloneqq p(y \mid x) \int_{\mathbf{R}^n} p(x \mid x^-, u) p(x^-) dx^-.$$
 (9)

The update law (8) can be viewed as a *functional* that maps the previous posterior PDF $p(x^-)$ to the current one p(x | y, u). In the linear case with Gaussian noise, this functional can be reduced to simple arithmetic operations of the mean and variance of the previous posterior PDF to yield the prediction and update steps of the Kalman filter. However, in the nonlinear cases, the Gaussian property of the posterior PDF can no longer be guaranteed, which makes it extremely difficult to compute the functional. Therefore, as the first step to overcome this difficulty, we approximate the posterior PDF using a Gaussian.

For the following discussion, \tilde{p} denotes an approximation of the PDF p with the same stochastic variables. Suppose we have an approximating Gaussian $\tilde{p}(x^- | \mu^-, \Sigma^-)$ of the previous posterior PDF $p(x^-)$ with mean μ^- and variance Σ^- . By replacing $p(x^-)$ with $\tilde{p}(x^- | \mu^-, \Sigma^-)$ in (9), $p_{xy}(x, y | u)$ can be approximated as $\tilde{p}_{xy}(x, y | u, \mu^-, \Sigma^-)$, and p(x | y, u) can be approximated as

$$\tilde{p}(x \mid y, u, \mu^{-}, \Sigma^{-}) = \frac{\tilde{p}_{xy}(x, y \mid u, \mu^{-}, \Sigma^{-})}{\int_{\mathbf{R}^{n}} \tilde{p}_{xy}(x, y \mid u, \mu^{-}, \Sigma^{-}) dx}.$$
 (10)

Note that the approximating PDF on the left-hand side is *not* Gaussian since we consider the nonlinearities of functions f and h without any approximations. Hence, we further approximate $\tilde{p}(x \mid y, u, \mu^-, \Sigma^-)$ with a Gaussian $\tilde{p}(x \mid \mu, \Sigma)$ of the same mean and variance as those of $\tilde{p}(x \mid y, u, \mu^-, \Sigma^-)$, that is, $\tilde{p}(x \mid y, u, \mu^-, \Sigma^-) \approx \tilde{p}(x \mid \mu, \Sigma)$ where

$$\mu \coloneqq E_{\tilde{p}(x|y,u,\mu^{-},\Sigma^{-})}[x]$$
$$= \frac{\int_{\mathbf{R}^{n}} x \cdot \tilde{p}_{xy}(x,y \mid u,\mu^{-},\Sigma^{-})dx}{\int_{\mathbf{R}^{n}} \tilde{p}_{xy}(x,y \mid u,\mu^{-},\Sigma^{-})dx}$$
(11)

and

$$\Sigma := E_{\tilde{p}(x|y,u,\mu^{-},\Sigma^{-})}[xx^{\top}] - \mu\mu^{\top} = \frac{\int_{\mathbf{R}^{n}} xx^{\top} \cdot \tilde{p}_{xy}(x,y \mid u,\mu^{-},\Sigma^{-})dx}{\int_{\mathbf{R}^{n}} \tilde{p}_{xy}(x,y \mid u,\mu^{-},\Sigma^{-})dx} - \mu\mu^{\top}.$$
(12)

Definitions (11) and (12) provide a finite number of functions that map the parameters y, u, μ^- , and Σ^- to the current estimates μ and Σ . Hence, by evaluating the right-hand sides of (11) and (12) recursively, we can perform Bayesian filtering while exactly considering the nonlinearities of fand h. For simplicity, we define $\Phi[h(x)](y, u, \mu^-, \Sigma^-)$ for a scalar-, vector-, or matrix-valued function h(x) as

$$\Phi\left[h(x)\right]\left(y,u,\mu^{-},\Sigma^{-}\right) \coloneqq \int_{\mathbf{R}^{n}} h(x) \cdot \tilde{p}_{xy}(x,y \mid u,\mu^{-},\Sigma^{-}) dx,$$

which is an integral over x depending on the parameters y, u, μ^- , and Σ^- . The computations of (11) and (12) are then reduced to evaluations of three integrals, namely $\Phi[1]$, $\Phi[x]$, and $\Phi[xx^{\top}]$, because μ and Σ can be written as $\mu = \Phi[x]/\Phi[1]$ and $\Sigma = \Phi[xx^{\top}]/\Phi[1] - \mu\mu^{\top}$, respectively. We utilize the HGM to evaluate these integrals depending on the parameters. Hence, we make the following assumption.

Assumption 1: The conditional PDFs (6) and (7) are holonomic functions.

Under this assumption, $\tilde{p}_{xy}(x, y \mid u, \mu^-, \Sigma^-)$ is holonomic by the closure property (Theorem 1) because the Gaussian $\tilde{p}(x^- \mid \mu^-, \Sigma^-)$ is also holonomic.

IV. Evaluation of Mean and Variance via Integral Transform

The authors propose a method to evaluate all three integrals $\Phi[1]$, $\Phi[x]$, and $\Phi[xx^{\top}]$ directly using the HGM [14]. Although the previous method showed better performance than other existing methods for a numerical example, the IVP (2) has to be solved for each component of the three integrals. If we can decrease the number of IVPs, then it reduces the online computational cost. Hereafter, $z \in \mathbf{R}^N$ denotes a vector consisting of all the independent components of $y \in \mathbf{R}^r$, $u \in \mathbf{R}^m$, $\mu^- \in \mathbf{R}^n$, and $\Sigma^- \in PD(n)$, where N = r + m + n + n(n + 1)/2. With this notation, for example, $\tilde{p}_{xy}(x, z)$ denotes $\tilde{p}_{xy}(x, y \mid u, \mu^-, \Sigma^-)$ as a function of x, y, u, μ^- , and Σ^- .

In this work, we use an integral transform of \tilde{p}_{xy} to reduce the online computational cost. Consider an integral transform $\mathcal{T}[\tilde{p}_{xy}]$ that is defined as follows.

$$\mathcal{T}[\tilde{p}_{xy}](\xi, z) \coloneqq \int_{\mathbf{R}^n} \exp(\xi^\top x) \tilde{p}_{xy}(x, z) dx.$$
(13)

Note that although this definition is similar to that of the moment generating function, it is different because \tilde{p}_{xy} is the PDF of the *x* and *y* pair rather than *x*. We assume that there exists a compact subset $\Xi \subset \mathbb{R}^n$ including the origin such that for all $\xi \in \Xi$, (13) can be defined and smooth.

The integrals $\Phi[1]$, $\Phi[x]$, and $\Phi[xx^{\top}]$ are then obtained from the derivatives of $\mathcal{T}[\tilde{p}_{xy}]$ at $\xi = 0$ as follows:

$$\mathcal{T}[\tilde{p}_{xy}](0,z) = \int_{\mathbf{R}^n} \tilde{p}_{xy}(x,z)dx = \Phi[1](z),$$
$$\partial_{\xi_i} \mathcal{T}[\tilde{p}_{xy}](0,z) = \int_{\mathbf{R}^n} \left[\partial_{\xi_i} \exp(\xi^\top x) \tilde{p}_{xy}(x,z) \right] \Big|_{\xi=0} dx$$
$$= \Phi[x_i](z), \tag{14}$$

$$\partial_{\xi_i} \partial_{\xi_j} \mathcal{T}[\tilde{p}_{xy}](0,z) = \int_{\mathbf{R}^n} \left[\partial_{\xi_i} \partial_{\xi_j} \exp(\xi^\top x) \tilde{p}_{xy}(x,z) \right] \Big|_{\xi=0} dx$$
$$= \Phi\left[x_i x_j \right] (z). \tag{15}$$

Here, the approximating PDF \tilde{p}_{xy} is a holonomic function under Assumption 1, and the kernel $\exp(\xi^{\top}x)$ is also holonomic. This implies by the closure property that $\mathcal{T}[\tilde{p}_{xy}]$ is also a holonomic function (Theorem 1). Hence, there exist a finite set of differential operators $\{\partial^{d_1}, \ldots, \partial^{d_{q-1}}\}$ such that the following vector-valued function satisfies the Pfaffian system (1): $Q(\xi, z) \coloneqq [1 \ \partial^{d_0} \cdots \ \partial^{d_{q-1}}]^{\top} \bullet \mathcal{T}[\tilde{p}_{xy}]$, where • represents all the actions of all components on the scalarvalued function $\mathcal{T}[\tilde{p}_{xy}]$. Using the Pfaffian system, we can explicitly express $\mathcal{T}[\tilde{p}_{xy}]$ and its derivatives (14) as well as (15) as a linear combination of the components of Q with coefficients in $\mathbf{R}(\xi, z)$.

First, $\mathcal{T}[\tilde{p}_{xy}](\xi, z)$ itself is the first component of $Q(\xi, z)$, thus satisfying the following:

$$\Phi[1] = \mathcal{T}[\tilde{p}_{xy}] = C^{(0)}Q, \tag{16}$$

where $C^{(0)}$ is a constant coefficient vector $[1 \ 0 \ \cdots \ 0] \in \mathbf{R}^q$. Next, the first derivatives $\partial_{\xi_i} \mathcal{T}[\tilde{p}_{xy}]$ (i = 1, ..., n) are obtained using the first components of the left-hand sides of (1). Hence, the following identities hold.

$$\Phi[x_i] = \partial_{\xi_i} \mathcal{T}[\tilde{p}_{xy}] = C_i^{(1)} \mathcal{Q} \quad (i = 1, \dots, n),$$
(17)

where the coefficient vector $C_i^{(1)}(\xi, z)$ is the first row vector of $A_{\xi_i}(\xi, z) \in \mathbf{R}(\xi, z)^{q \times q}$ in (1). Finally, the second derivatives $\partial_{\xi_i} \partial_{\xi_j} \mathcal{T}[\tilde{p}_{xy}]$ (i, j = 1, ..., n) are obtained by differentiating both sides of (1); $\partial_{\xi_i} \bullet \partial_{\xi_j} Q = \partial_{\xi_i} \bullet (A_{\xi_j} Q) = \partial_{\xi_i} A_{\xi_j} Q + A_{\xi_j} \partial_{\xi_i} Q = (\partial_{\xi_i} A_{\xi_j} + A_{\xi_j} A_{\xi_i})Q$, and hence

$$\Phi[x_i x_j] = \partial_{\xi_i} \partial_{\xi_j} \mathcal{T}[\tilde{p}_{xy}] = C_{ij}^{(2)} Q \quad (i, j \in \{1, \dots, n\}),$$
(18)

where the coefficient vector $C_{ij}^{(2)}(\xi, z)$ is the first row vector of $\partial_{\xi_i} A_{\xi_i} + A_{\xi_i} A_{\xi_i}$.

Note that A_{ξ_i} (i = 1, ..., n) can be computed symbolically and offline from a basis of a zero-dimensional ideal annihilating $\mathcal{T}[\tilde{p}_{xy}]$, along with $C_i^{(1)}(\xi, z)$ (i = 1, ..., n) and $C_{ij}^{(2)}(\xi, z)$ (i, j = 1, ..., n). Therefore, if we can efficiently evaluate $Q(0, \hat{z})$ for a given \hat{z} , we can readily compute the values of $\Phi[1]$, $\Phi[x]$, and $\Phi[xx^{\top}]$ at $z = \hat{z}$ from (16)–(18). The computation of $Q(0, \hat{z})$ can be accomplished by numerically integrating (2) using numerical integration methods, such as the RK4 method. Therefore, the current mean μ and variance Σ for a given data \hat{z} can be computed from (11) and (12) as

$$\mu = \frac{\Phi[x](\hat{z})}{\Phi[1](\hat{z})}, \quad \Sigma = \frac{\Phi[xx^{\top}](\hat{z})}{\Phi[1](\hat{z})} - \mu\mu^{\top}.$$
 (19)

Now, we summarize the entire online process of the proposed method as Algorithm 1. Note that ξ is always zero during the numerical integration at line 4 of Algorithm 1. Therefore, we can substitute $\xi = 0$ into A_{λ} ($\lambda \in \{\xi_1, \ldots, \xi_n, z_1, \ldots, z_N\}$ in advance. Furthermore, we can omit evaluation of A_{ξ_i} ($i = 1, \ldots, n$) because if ξ is constant, the derivatives $\partial Q/\partial \xi_i = A_{\xi_i}Q$ ($i = 1, \ldots, n$) are not

required in the numerical integration.

V. NUMERICAL EXAMPLE

This section presents a numerical example to show the efficiency of the proposed method. In the following demonstration, we use Risa/Asir [19] for the symbolic computation, Maple for the offline numerical computation, and Python for the online numerical computation on a PC (Intel(R) Core(TM) i7-1065G7 CPU @ 1.30 GHz; RAM: 16 GB).

A. Problem setting and results of offline computation

Consider the nonlinear stochastic one-dimensional system considered in Section V of our previous work [14], where *u* is given as the function $\cos(0.6k)$ with time step *k*. First, we need to compute a basis of a holonomic ideal annihilating $\mathcal{T}[\tilde{p}_{xy}]$. From the definition, $\mathcal{T}[\tilde{p}_{xy}]$ consists of four simple components: $\tilde{p}(x^- | \mu^-, \Sigma^-)$, (6), (7), and the kernel exp (ξx) of the integral transform. Therefore, a basis of the holonomic ideal for $\mathcal{T}[\tilde{p}_{xy}]$ can be computed from bases of holonomic ideals annihilating these components using integrations and multiplications in terms of differential operators.

The one-dimensional Gaussian $\tilde{p}(x^- | \mu^-, \Sigma^-)$ is annihilated by the following three differential operators:

$$\Sigma^{-}\partial_{x^{-}} + x^{-} - \mu^{-}, \quad \Sigma^{-}\partial_{\mu^{-}} - x^{-} + \mu^{-},$$

$$2(\Sigma^{-})^{2}\partial_{\Sigma^{-}} + \Sigma^{-} - (x^{-} - \mu^{-})^{2},$$

$$(20)$$

which construct a basis of a holonomic ideal for $\tilde{p}(x^- | \mu^-, \Sigma^-)$. The conditional PDF (6) is obtained by substituting the state equation into the PDF of $\mathcal{N}(0, 1)$ and is annihilated by the following three differential operators:

$$\partial_x + x - (4/5)x^- - u, \quad \partial_u - x + (4/5)x^- + u, - (5/4)\partial_{x^-} + x - (4/5)x^- - u,$$
(21)

which construct a basis of a holonomic ideal for (6) in this example. The other conditional PDF (7) is also obtained by substituting the observation equation into the PDF of $\mathcal{N}(0, 1)$ and is annihilated by

$$\partial_y + y - \frac{2x}{1+x^2}, \quad -\frac{(1+x^2)^2}{2(1-x^2)}\partial_x + y - \frac{2x}{1+x^2}.$$
 (22)

After cancellation of the denominators, the set of differential operators becomes a basis of a holonomic ideal for (7).

Algorithm 1 One-step Estimation via HGM using (13)

Input: Explicit expressions of $C^{(0)}$, $C_i^{(1)}(\xi, z)$ (i = 1, ..., n), $C_{ij}^{(2)}(\xi, z)$ $(i, j \in \{1, ..., n\})$, $A_\lambda(\xi, z)$ $(\lambda \in \{\xi_1, ..., \xi_n, z_1, ..., z_N\})$, initial point z_{init} and vector $Q(0, z_{\text{init}})$, as well as given data \hat{z}

Output: Failure or estimates μ and Σ

- 1: Define z(s) such that $z(0) = z_{init}$ and $z(1) = \hat{z}$
- 2: if $\exists \tilde{s} \in [0, 1]$ such that $z(\tilde{s})$ lies in singular locus then
- 3: **return** algorithm has failed
- 4: Solve IVP (2) numerically using initial vector Q(0, z_{init}) and obtain Q(0, ẑ)
- 5: return (19) obtained from (16)–(18)

Finally, a basis of a holonomic ideal for $\exp(\xi x)$ can be obtained as

$$\partial_{\xi} - x, \quad \partial_x - \xi.$$
 (23)

From all bases (20)–(23), a basis \mathcal{B} of a holonomic ideal annihilating the integral transform $\mathcal{T}[\tilde{p}_{xy}](\xi, z)$, where $z = [y \ u \ \mu^- \Sigma^-]^\top \in \mathbf{R}^3 \times \mathbf{R}_+$, can be obtained via multiplication and integration in terms of differential operators [17]. In this example, \mathcal{B} consists of seven differential operators.

Next, we compute the Pfaffian system (1) for $\mathcal{T}[\tilde{p}_{xy}]$. It is known that any basis of a holonomic ideal generates a zero-dimensional ideal in \mathcal{R} [16]. Hence, \mathcal{B} also generates a zero-dimensional ideal over $\mathcal{R} = \mathbf{R}(\xi, z) \langle \partial_{\xi}, \partial_{z} \rangle$. From \mathcal{B} , we derive $\{\partial_{z_1}, \partial_{z_2}, \partial_{z_4}, \partial_{z_1}^2, \partial_{z_2}\partial_{z_1}, \partial_{z_4}\partial_{z_2}\}$ as the finite set of differential operators in Theorem 2. Hence, Q becomes a seven-dimensional vector-valued function:

$$Q = \begin{bmatrix} 1 \ \partial_{z_1} \ \partial_{z_2} \ \partial_{z_4} \ \partial_{z_1}^2 \ \partial_{z_2} \partial_{z_1} \ \partial_{z_4} \partial_{z_2} \end{bmatrix}^\top \bullet \mathcal{T}[\tilde{p}_{xy}], \quad (24)$$

where • represents the actions of all the components on $\mathcal{T}[\tilde{p}_{xy}]$. All entries of the coefficient matrices $A_{\lambda} \in \mathbf{R}(\xi, z)^{7\times7}$ ($\lambda \in \{\xi, z_1, z_2, z_3, z_4\}$) as well as the coefficient vectors in (17) and (18) are explicitly computed from \mathcal{B} .

Finally, we have to compute the initial points z_{init} and corresponding initial vectors $Q(0, z_{\text{init}})$. The choice of the initial points depends on the singular locus of the Pfaffian system since it should not be intersected by the integration path; otherwise, Algorithm 1 fails (line: 3). In this case, however, the least common multiple of all the denominators in A_{λ} ($\lambda \in \{\xi, z_1, z_2, z_3, z_4\}$) is $16z_4 +$ 25, which must not be zero because $z_4 = \Sigma^- > 0$. Therefore, the singular locus is the empty set. For this example, we choose the following eight initial points: $Z_{\text{init}} := \{[z_1 \ z_2 \ z_3 \ z_4]^\top | z_1 = \pm 1, z_2 = \pm 1, z_3 = \pm 1, z_4 = 1\}$. The corresponding initial vectors can be computed numerically from the definition (24).

B. Settings for online computation and estimation results

In the online computations, the data z consisting of the input u, output y, and previous estimates μ^- and Σ^- are given at each time step. Let \hat{z} denote the given data at a time step. To avoid failure of the integration in Algorithm 1, an initial point and an integration path should be appropriately chosen. For this example, we fix the integration path to a line segment from an initial point z_{init} to a given data \hat{z} , that is, $z(s) := s\hat{z} + (1 - s)z_{\text{init}}$. The initial point is thus selected for each given data such that the line segment does not intersect the singular locus and is shortest among the prescribed set Z_{init} . In this example, the selection process can be simplified because the singular locus is empty; hence, it is enough to choose the initial point closest to \hat{z} . However, this selection process considering the singular locus is fully implemented for a fair comparison with our previous method, which uses the Pfaffian systems for $\Phi[1]$, $\Phi[x]$, and $\Phi[xx^{\top}]$ with a nonempty singular locus [14]. The numerical integration along the defined path is performed using the ABM4 method, with the first three steps initialized using the RK4 method. Under these settings, the proposed method is implemented.



Fig. 1: Comparison of NLLs for proposed method (solid, thick), previous method (solid, thin), PFs of 100 particles (dashed), UKF (dotted), and EKF (dash-dotted, almost greater than 1.2 after k = 4)



Fig. 2: Boxplots of computational times for one-step estimations of all methods

For comparison, we also implemented the EKF, UKF, PF, and our previous method. The number of particles in the PF was set to 100 so that its accuracy would be almost the same as that of the proposed method. Three hundreds of realizations are used in this comparison.

To compare the performances of all the methods, the negative log-likelihood (NLL) [20] was computed. Figure 1 shows the NLL of each method averaged over all realizations. As can be seen from the figure, only the PF of 100 particles shows comparable performance to the proposed and previous methods. The averaged NLL of the previous method is identical to that of the proposed method because both approaches compute the same estimates (19).

The computational times for the one-step estimations of all the methods are summarized as boxplots in Fig. 2. It can be seen that the proposed method is faster than the previous method as well as the PF with 100 particles, which show comparable performances for the NLL values. These results indicate that the proposed method is more efficient than the PF and our previous method.

VI. CONCLUSION

A symbolic-numeric method is proposed herein to perform Bayesian filtering for a class of nonlinear stochastic systems. The posterior PDF of the state is first approximated using a Gaussian PDF, and the mean and variance are then updated while considering the nonlinearity of the system exactly. This update process is formulated as evaluations of several integrals using the HGM. We also propose an integral transform to compute the mean and variance efficiently from the vector-valued function obtained using the HGM.

It is a known fact that the numerical integration in the third step of the HGM may sometimes diverge owing to errors in the initial vectors. Therefore, in future work, the selection of integration paths or initial points will be studied in depth to enable more stable numerical integrations, for example, by considering the special structure of the problems considered in this work. Another direction for future work is approximating the posterior PDF more accurately using a PDF with more parameters than the Gaussian, such as the skew Gaussian.

References

- R. E. Kalman, "A new approach to linear filtering and prediction problems," ASME Journal of Basic Engineering, vol. 82, pp. 35–45, 1960.
- [2] R. E. Kalman and R. S. Bucy, "New results in linear filtering and prediction theory," ASME Journal of Basic Engineering, vol. 83, no. 1, pp. 95–108, 1961.
- [3] I. Rusnak, "Maximum likelihood optimal estimator of nonautonomous nonlinear dynamic systems," in *Proceedings of European Control Conference*, pp. 909–914, 2015.
- [4] X. Luo, Y. Jiao, W. L. Chiou, and S. S. Yau, "A novel suboptimal method for solving polynomial filtering problems," *Automatica*, vol. 62, pp. 26–31, 2015.
- [5] B. Chen and G. Hu, "Nonlinear state estimation under bounded noises," *Automatica*, vol. 98, pp. 159–168, 2018.
- [6] N. Duong, J. L. Speyer, J. Yoneyama, and M. Idan, "Laplace estimator for linear scalar systems," in *Proceedings of the IEEE Conference on Decision and Control*, pp. 2283–2290, 2018.
- [7] K. Li, F. Pfaff, and U. D. Hanebeck, "Unscented dual quaternion particle filter for SE(3) estimation," *IEEE Control Systems Letters*, vol. 5, no. 2, pp. 647–652, 2021.
- [8] G. Kitagawa, "Monte Carlo filtering and smoothing method for non-Gaussian nonlinear state space model," *Institute of Statistical Mathematics Research Memorandum*, vol. 462, 1993.
- [9] N. J. Gordon, D. J. Salmond, and A. F. M. Smith, "Novel approach to nonlinear/non-Gaussian Bayesian state estimation," in *IEE Proceedings F - Radar and Signal Processing*, vol. 140, pp. 107–113, 1993.
- [10] A. H. Jazwinski Stochastic Processes and Filtering Theory, Elsevier Science, 1970.
- [11] S. J. Julier and J. K. Uhlmann, "New extension of the Kalman filter to nonlinear systems," in SPIE, vol. 3068, 1997.
- [12] I. Arasaratnam and S. Haykin, "Cubature Kalman filters," *IEEE Transactions on Automatic Control*, vol. 54, no. 6, pp. 1254–1269, 2009.
- [13] K. Ito and K. Xiong, "Gaussian filters for nonlinear filtering problems," *IEEE Transactions on Automatic Control*, vol. 45, no. 5, pp. 910–927, 2000.
- [14] T. Iori and T. Ohtsuka, "Symbolic-numeric computation of posterior mean and variance for a class of discrete-time nonlinear stochastic systems," in *Proceedings of the IEEE Conference on Decision and Control*, pp. 4814–4821, 2020.
- [15] H. Nakayama, K. Nishiyama, M. Noro, K. Ohara, T. Sei, N. Takayama, and A. Takemura, "Holonomic gradient descent and its application to the Fisher-Bingham integral," *Advances in Applied Mathematics*, vol. 47, no. 3, pp. 639–658, 2011.
- [16] T. Hibi ed. Gröbner Bases: Statistics and Software Systems, Springer Japan, 1st edition, 2013.
- [17] T. Oaku, Y. Shiraki, and N. Takayama, "Algebraic algorithms for D-modules and numerical analysis," *Computer Mathematics (Proceedings of ASCM 2003)*, vol. 10, pp. 23–39, 2003.
- [18] J. A. E. Bryson and Y.-C. Ho Applied Optimal Control, John Wiley & Sons, 1st edition, 1975.
- [19] M. Noro, N. Takayama, H. Nakayama, K. Nishiyama, and K. Ohara, "Risa/Asir: A computer algebra system," http://www.math. kobe-u.ac.jp/Asir/asir.html, 2020.
- [20] M. P. Deisenroth *Efficient Reinforcement Learning Using Gaussian Processes*, KIT Scientific Publishing, 2010.