Principal Radical Systems, Lefschetz Properties, and Perfection of Specht Ideals of Two-Rowed Partitions

Chris McDaniel and Junzo Watanabe

Abstract

We show that the Specht ideal of a two-rowed partition is perfect over an arbitrary field, provided that the characteristic is either zero or bounded below by the size of the second row of the partition, and we show this lower bound is tight. We also establish perfection and other properties of certain variants of Specht ideals, and find a surprising connection to the weak Lefschetz property. Our results in particular give a self-contained proof of Cohen-Macaulayness of certain *h*-equals sets, a result previously obtained by Etingof-Gorsky-Losev over the complex numbers using rational Cherednik algebras.

Keywords: Specht ideal, Cohen-Macaulay, weak Lefschetz

1 Introduction

Fix an integer n, let \mathbb{F} be any field, and let $R = \mathbb{F}[x_1, \dots, x_n]$ be the polynomial ring with its standard grading, and equipped with the usual action of the symmetric group \mathfrak{S}_n by permuting the variables. For any partition $\lambda \vdash n$ the Specht module $V(\lambda)$ over \mathbb{F} is the \mathbb{F} -vector space generated by the Specht polynomials of λ which are indexed by the set of tableaux T on the Young diagram of λ . If \mathbb{F} has characteristic zero, then the Specht modules form a complete list of irreducible \mathfrak{S}_n -representations, highlighting their importance in representation theory. In this paper we take the point of view of commutative algebra, and study the ideals generated by Specht modules called Specht ideals.

Specifically we show that for partitions with two parts $\lambda = (n - k, k)$ (or Young diagrams with two rows), the Specht ideal, which we denote by a(n, k, k) is radical and, if the characteristic of \mathbb{F} is zero or sufficiently large, perfect. Our results are stated in terms of commutative algebra, but they can be interpreted geometrically as follows:

Proposition 1.1. Fix a field \mathbb{F} with char(\mathbb{F}) = $p \ge 0$, and fix $m \ge 3$. For each $2 \le h \le m$, define $X_{m,h} \subset \mathbb{F}^m$ as the union of \mathfrak{S}_m translates of the linear subspace cut out by the h-1 linear equations $x_1 = \cdots = x_h$, i.e.

$$X_{m,h} = \bigcup_{\sigma \in \mathfrak{S}_m} \sigma.\{(x_1,\ldots,x_m) \in \mathbb{F}^m \mid x_1 = \cdots = x_h\}.$$

Assume that $h > \left\lfloor \frac{m+1}{2} \right\rfloor$.

1. If p = 0 or $p \ge m - h + 1$, then $X_{m,h}$ is Cohen-Macaulay¹.

¹meaning that its homogeneous coordinate ring is Cohen-Macaulay

2. If $0 , then <math>X_{m,m-p}$ is not Cohen-Macaulay.

Over the field $\mathbb{F} = \mathbb{C}$, Proposition 1.1(1.) was obtained by Etingof-Gorsky-Losev [2, Proposition 3.11], using deep results from the representation theory of rational Cherednik algebras. Our proof of Proposition 1.1 is more elementary in that it uses only basic results from commutative algebra, which we hope will appeal to those uninitiated with rational Cherednik algebras. We emphasize however that our elementary proof is by no means easy.

The study of Specht ideals seems to have been initiated by Yanagawa in his recent paper [15], although they have appeared implicitly in the earlier works of others [1, 2, 4]. As Yanagawa shows in his paper, the connection between Proposition 1.1 and Specht ideals is as follows: Setting m = n + 1 and k + 1 = m - h + 1, and if $I_{m,h} \subset R$ is the ideal cutting out the union of linear spaces $X_{m,h}$ then we have

$$I_{m,h} = \bigcap_{\sigma \in \mathfrak{S}_m} \sigma. (x_1 - x_2, \dots, x_1 - x_h) = \sqrt{\mathfrak{a}(n+1, k+1, k+1)}.$$
 (1)

In his paper [15], Yanagawa proves that two-rowed Specht ideals are radical by an ingenious but complicated argument. He then invokes the Etingof-Gorsky-Losev result [2, Proposition 3.11] to prove that the Specht ideal $\mathfrak{a}(n+1,k+1,k+1)$ (although he used different notation) is perfect if the field has characteristic zero. The present paper grew out of an attempt to understand and simplify Yanagawa's arguments and to find an elementary proof of the Etingof-Gorsky-Losev result in the two-rowed case. As the reader will surmise, the distinguishing feature of Specht ideals of two-rowed partitions is that their minimal generators are square-free, a fact which will be exploited throughout this paper.

Recall that an ideal $I \subset R$ in a Noetherian ring is called *perfect* if its grade is equal to its homological (or projective) dimension. In a polynomial ring R, a homogeneous ideal $I \subset R$ is perfect if and only if its quotient R/I is Cohen-Macaulay. The following is one of the main results of this paper, and is the algebraic analogue of Proposition 1.1.

Theorem 1.2. Let \mathbb{F} be any field of characteristic $p \ge 0$, and fix positive integers n, k satisfying $n \ge 2k + 1$.

- 1. If p = 0 or $p \ge k + 1$, then the Specht ideal a(n + 1, k + 1, k + 1) is perfect.
- 2. If $n \ge 2p + 1$, then the Specht ideal a(n + 1, p + 1, p + 1) is not perfect.

In his paper [15] Yanagawa has conjectured that the Specht ideal a(n + 1, k + 1, k + 1) is perfect in characteristic p if and only if p = 0 or $p \ge k + 1$. Theorem 1.2 proves one implication and part of the other one in Yanagawa's conjecture.

Our proof of Theorem 1.2 is inspired by the seminal paper of Hochster-Eagon [5], in which they proved perfection of generic determinantal ideals using what they termed a principal radical system. Our method, which might be more aptly described as a *deconstructed* principal radical system is based on the following elementary facts from commutative algebra:

Lemma 1.3. Let $I \subset R$ be a homogeneous ideal, and let $x \in R \setminus I$ be a homogeneous polynomial.

- 1. If (I:x) = I and I + (x) is radical, then I is also radical.
- 2. If (I:x) = I and I + (x) is perfect, then I is also perfect.

Lemma 1.4. Suppose that ideals $I, J \subset R$ are homogeneous ideals, both perfect of the same grade g and suppose that I + J has grade g + 1. Then $I \cap J$ is perfect if and only if I + J is perfect.

Actually we use a slight modification of Lemma 1.3, but we wrote it that way to emphasize the connection between radical and perfection. In fact, before we prove that our Specht ideals are perfect, we must first prove that they are radical. More specifically, the main challenge in applying Lemma 1.3 is finding the right principal ideal (x) with respect to I, and it is this issue that Lemma 1.4 addresses. However, in applying Lemma 1.4, we must know what amounts to a primary decomposition of our ideal I. Our method then consists of repeated application of Lemma 1.3 and Lemma 1.4 to obtain a sequence of (families of) ideals beginning with the Specht ideal, and ending with some ideal for which perfection is already known, e.g. by induction. A schematic of our argument with our sequence of ideals is shown below:

$$\alpha(n+1,k+1,k+1) \xrightarrow{\text{Lem. 1.3}} \alpha(n,k,k+1) \xrightarrow{\text{Lem. 1.4}} I(n,k) \xrightarrow{\text{Lem. 1.4}} J(n,k)$$


$$I(n-1,k-1)$$

We give a more detailed description of our method and these ideals listed above, together with the other results of this paper, below.

Taking $I = \alpha(n+1,k+1,k+1)$ and $x = x_{n+1}$ in Lemma 1.3, it is easy to show that (I:x) = I, and that $I + (x) = \alpha(n,k,k+1) + (x_{n+1})$ where $\alpha(n,k,k+1)$ is the ideal generated by square-free products of Specht polynomials of type $\lambda = (n-k,k)$ with a linear monomial in the variables x_1, \ldots, x_n . Generalizing, we introduce, for integers $0 \le k \le d \le n-k$, the *d-shifted Specht ideal* $\alpha(n,k,d)$ generated by square-free products of Specht polynomials of type $\lambda = (n-k,k)$ and square-free monomials of degree d-k.

These shifted Specht ideals interpolate between Specht ideals a(n, k, k) in case d = k, and square-free monomial ideals in case k = 0, where $a(n, 0, d) = (x_1, \dots, x_n)^{(d)}$ is the ideal generated by all square-free monomials of fixed degree d. Our first step in understanding these shifted Specht ideals is to find a minimal generating set. Just as minimal generators for the Specht ideal are indexed by standard Young tableaux on λ , we show that minimal generators for the shifted Specht ideal are indexed by standard Young tableaux on a shifted version of λ .

Theorem 1.5. A minimal generating set for the shifted Specht ideal a(n, k, d) is formed by the shifted Specht polynomials $F_T(d)$ indexed by standard tableaux T on the d-shifted shape of $\lambda = (n - k, k)$ obtained from Young diagram of λ by moving the last n - k - d boxes on the top row to the first n - k - d boxes on the bottom row, i.e.



The linear span of the shifted Specht polynomials $F_T(d)$, $T \in \text{Tab}(\lambda(d))$ forms an \mathfrak{S}_n -representation V(n,k,d) that we call a *shifted Specht module*, which is reducible in general. We prove that our d-shifted Specht ideals satisfy the following decomposition formula, which is crucial in our quest for perfection, and which holds if and only if our shifted Specht ideals are radical.

Theorem 1.6. For any integers k, d satisfying $1 \le k < d \le n - k$, we have

$$a(n, k, d) = a(n, k, d - 1) \cap (x_1, \dots, x_n)^{(d)}.$$
 (2)

Yanagawa [15] has proved Theorem 1.6 in the special case d = k + 1 using a clever argument, which is described in further detail below. As it turns out, his argument goes through verbatim to prove Theorem 1.6 in the general case, and is in fact simplified by Theorem 1.5. It follows directly from Theorem 1.6 that our shifted Specht ideals are radical.

Theorem 1.7. Fix integers k, d satisfying $0 \le k \le d \le n - k$. Then the shifted Specht ideal a(n, k, d) is radical.

Perfection of shifted Specht ideals is more difficult to prove, and in fact, most shifted Specht ideals are not perfect. Indeed Theorem 1.6 implies that the shifted Specht ideal a(n, k, d) does not have pure height and hence cannot be perfect if $d \neq k, k+1$. To show that the shifted Specht ideal $a(n, k, k+1) = a(n, k, k) \cap (x_1, \dots, x_n)^{(k+1)}$ is perfect, we appeal to Lemma 1.4 and introduce the *Specht-monomial ideal I* $(n, k) = a(n, k, k) + (x_1, \dots, x_n)^{(k+1)}$.

While the Specht-monomial ideal is not radical in general, it does satisfy a decomposition formula similar to (2), but it depends on the field characteristic. We were pleased to discover that this dependence on field characteristic is the same one imposed by the weak Lefschetz property of certain monomial complete intersection algebras. This is summarized in the following, which can be considered the other main result of this paper.

Theorem 1.8. Let \mathbb{F} be a field with $\operatorname{char}(\mathbb{F}) = p \ge 0$, and let n and k be positive integers satisfying $n \ge 2k + 1$. The following are equivalent:

- 1. p = 0 or $p \ge k + 1$.
- 2. The quadratic monomial complete intersection

$$C = \frac{\mathbb{F}[x_1, \dots, x_{2k}]}{(x_1^2, \dots, x_{2k}^2)}$$
(3)

has the weak Lefschetz property.

3. The Specht-monomial ideal I(n, k) satisfies decomposition

$$I(n,k) = I(n-1,k-1) \cap \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \right),$$

$$where \quad y_i = \begin{cases} x_n - x_i & \text{if } 1 \le i \le n-1 \\ x_n & \text{if } i = n \end{cases}$$
(4)

4. The Specht-monomial ideal I(n, k) is perfect.

We shall break Theorem 1.8 into three parts, one for each equivalence $(1.) \Leftrightarrow (a.)$ which shall refer to hereafter as Theorem 1.8(a.) for a=2,3,4. Theorem 1.8(2.) is due to Kustin-Vraciu [7], and we will not re-prove it here. Theorem 1.8(3.) and Lemma 1.4 lead to yet another family of ideals, which remain unnamed:

$$J(n,k) = I(n-1,k-1) + (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

Finally to prove Theorem 1.8(4.), and hence also Theorem 1.2(1.) we apply Lemma 1.3 to show that our unnamed ideal J(n, k) is perfect. Unfortunately, Theorem 1.2(2.) seems to be the best we could do with our method, although we conjecture that the full converse of Theorem 1.2(1.) holds.

Conjecture 1.9. If $p = \text{char}(\mathbb{F})$ and n, k are positive integers satisfying $0 and <math>n \ge 2k+1$, then the Specht ideal $\mathfrak{a}(n+1,k+1,k+1)$ is not perfect.

Theorem 1.2(2.) says Conjecture 1.9 holds for k = p, but it remains open for k > p. Together with Theorem 1.2, Conjecture 1.9 is equivalent to Yanagawa's conjecture [15, Conjecture 5.5].

Some further remarks on the connection to the weak Lefschetz property are in order here. Since (shifted) Specht polynomials are square-free they are identified with elements of the algebra $A = \mathbb{F}[x_1, \dots, x_n]/(x_1^2, \dots, x_n^2)$, which carries an \mathfrak{sl}_2 -representation in which the raising operator is multiplication by the sum of variables, and the lowering operator is the corresponding linear partial differential operator. Moreover surjectivity of this lowering operator in degree k is equivalent to equality of the kernel of that lowering operator with the Specht module V(n,k,k), which is in turn equivalent to the weak Lefschetz property of C in (3). Surjectivity of the lowering operator on A is key to proving decomposition (4), and in fact reveals a hidden property of the Specht-monomial ideal: in small positive characteristic, i.e. 0 the ideal <math>I(n,k) has an embedded prime divisor, a phenomenon which does not occur for the Specht ideals.

The most technically difficult parts of our arguments are the decompositions in Theorem 1.6 and Theorem 1.8(3.). The two proofs we give are strikingly similar, and are both based on that ingenious and clever argument of Yanagawa mentioned above. This argument, in general terms, runs as follows: To prove that an ideal I satisfies a decomposition formula of the form $I = I' \cap J$ where J is a monomial ideal, first show that the intersection $I' \cap J$ can be generated by products of minimal generators of I' with monomials (not necessarily from J), with the additional property that if a sum of such products is in $I' \cap J$ then each of the summands is also in $I' \cap J$ (perhaps we should call such I an I'-monomial ideal). Next fix a monomial m and split the minimal generators V(I') into two parts say $V(I') = V_m(I') \oplus V^m(I')$, determined by whether or not m appears in their monomial expansion or not. In our situation, one can show that if m is not in the support of $v \in V^m(I')$, then $m \cdot v \in I$, and, with more effort, one can also show that if $v \in V_m$ and $m \cdot v \in I' \cap J$ then $m \cdot v \in I$. We highlight this argument here because it seems important in the theory of two-rowed Specht ideals.

This paper is organized as follows. In Section 2 we define Specht polynomials, shifted Specht polynomials, and the modules and ideals they generate. We then prove Theorem 1.5 and compute the dimensions of our shifted Specht modules (Theorem 2.1 and Theorem 2.14). In Section 3 we draw out the connection between Lefschetz properties and Specht modules, decompose the shifted Specht module into its irreducible representations, and derive other useful consequences. In Section 4 we prove Theorem 1.7 and Theorem 1.6 (Theorem 4.4 and Theorem 4.3). In Section 5 we prove Theorem 1.8(4.) (Theorem 5.6), Theorem 1.2 (Theorem 5.7), and Theorem 1.8(3.) (Theorem 5.5), in that order.

Acknowledgments

We would like to thank Larry Smith and University of Göttingen for their hospitality during our visit in the spring semester of 2019, when our discussions of Specht ideals and principal radical systems first began. We would also like to thank Larry Smith for pointing out that the idea of the principal radical system was first used by Kiyoshi Oka. The first author thanks Endicott College for the time and resources necessary to work on this project. The second author thanks K. Yanagawa for the introduction to Specht ideals and for his insightful problem.

Contents

1	Intr	oduction	1
2	Shif	ted Specht Polynomials, Modules, and Ideals	6
	2.1	Basis of a Shifted Specht Module	8
	2.2	Dimension of a Shifted Specht Module	14
3	Lefs	chetz Propoerties	16
	3.1	Weak Lefschetz Property	18
	3.2	Strong Lefschetz Property	20
4	Rad	ical of Shifted Specht Ideals	21
	4.1	Proof of Theorem 1.7 or Theorem 4.4	23
	4.2	Proof of Theorem 1.6 or Theorem 4.3	23
5	Perf	ection of Specht and Specht-Monomial Ideals	25
	5.1	Proof of Theorem 1.8(4.) or Theorem 5.6	27
	5.2	Proof of Theorem 1.2 or Theorem 5.7	31
		Proof of Theorem 1.8(3.) or Theorem 5.5	

2 Shifted Specht Polynomials, Modules, and Ideals

Fix a positive integer n, a field \mathbb{F} , and let $R = \mathbb{F}[x_1, \dots, x_n]$ be the standard graded polynomial ring in n-variables, equipped with the usual action of \mathfrak{S}_n which permutes the variables. A partition of n, denoted $\lambda \vdash n$ is a sequence of non-increasing integers which sum to n, i.e. $\lambda = (\lambda_1, \dots, \lambda_r)$ where $\lambda_1 \geq \dots \geq \lambda_r$ and $\sum_{i=1}^r \lambda_i = n$. The Young diagram of λ is a left-justified array of boxes with r-rows and λ_i boxes in each row, and a filling of those boxes with distinct numbers $1, \dots n$ is called a tableau of shape λ ; the set of all tableaux of shape λ will be written as $\mathrm{Tab}(\lambda)$. To each tableau $T \in \mathrm{Tab}(\lambda)$ of shape λ we associate a polynomial $F_T \in R$ as follows.

First for any subset $S \subset \{1, ..., n\}$, define the S-Vandermonde polynomial

$$\Delta_S = \prod_{i < j \in S} (x_i - x_j)$$

with the convention that if |S| < 2 then $\Delta_S = 1$. Then for any tableau T of shape λ with columns $C_1, \ldots, C_{\lambda_1}$, define its Specht polynomial by

$$F_T = \prod_{i=1}^{\lambda_1} \Delta_{C_i}.$$

The \mathbb{F} -linear span of Specht polynomials over Tab(λ) is an \mathfrak{S}_n -representation called the Specht module, which we denote $V(\lambda)$, i.e.

$$V(\lambda) = \langle F_T \mid T \in \text{Tab}(\lambda) \rangle \operatorname{sp}_{\mathbb{R}}(F_T \mid T \in \text{Tab}(\lambda));$$

it is well known, in the case char(\mathbb{F}) = 0, that $V(\lambda)$ is irreducible, and conversely that every irreducible \mathfrak{S}_n -representation is isomorphic to $V(\lambda)$ for some $\lambda \vdash n$, e.g. [12]. The Specht ideal

of λ is the ideal in R generated by $V(\lambda)$, i.e.


$$\mathfrak{a}(\lambda) = V(\lambda) \cdot R = (F_T \mid T \in \text{Tab}(\lambda)).$$

In his paper [15], Yanagawa has formulated the following conjecture:

Conjecture (Yanagawa '19). Over any field \mathbb{F} , and for any partition λ , the Specht ideal $\mathfrak{a}(\lambda)$ is radical.

This conjecture has been proved for partitions of the form $\lambda = (n-k, 1, \dots, 1)$ by Yanagawa-Watanabe [14], and for partitions of the form $\lambda = (n-k, k)$ and $\lambda = (d, d, 1)$ by Yanagawa [15]. However, even in these simple cases, Yanagawa's proof that $\mathfrak{a}(\lambda)$ is radical is by no means easy. This paper grew out of an attempt to understand and perhaps simplify Yanagawa's proof in the case of two rowed partitions $\lambda = (n-k, k)$, which we discuss next.

Fix an integer k satisfying $1 \le k \le n - k$, and let $\lambda = (n - k, k)$ be the corresponding partition, which we regard as a left-justified two-rowed Young diagram with n - k boxes in the first row and k boxes in the second. For each integer d satisfying $1 \le k \le d \le n - k$, define the d-shifted shape $\lambda(d)$ to be the Young diagram obtained by moving the righter-most n - d - k boxes on the first row to the left of the first boxes in the second row:



Note that $\lambda(k)$ is λ rotated by 180°.

A tableau T on shape $\lambda(d)$ is a labeling of the boxes of the Young diagram of $\lambda(d)$ with the numbers $\{1, \ldots, n\}$; we say that T is standard if the rows are increasing from left to right, and the columns are increasing from top to bottom. The set of tableaux on $\lambda(d)$ we denote by Tab(n, k, d), and the set of standard tableaux by STab(n, k, d).

Given a tableau $T \in \text{Tab}(n, k, d)$ on the d-shifted shape $\lambda(d)$, such as

$$T = \frac{i_1 \cdots i_k i_{k+1} \cdots i_d}{i_{d+1} \cdots i_{n-k} j_1 \cdots j_k}$$
(5)

define the associated d-shifted Specht polynomial to be the homogeneous polynomial of degree d by

$$F_T(d) = (x_{i_1} - x_{i_1}) \cdots (x_{i_k} - x_{i_k}) \cdot x_{i_{k+1}} \cdots x_{i_d}$$

Note that if d = k, then we recover the usual Specht polynomial:

$$F_T(k) = (x_{i_1} - x_{j_1}) \cdots (x_{i_k} - x_{j_k});$$

we sometimes use the alternative notation F_T or F_T^k to remember the degree. We allow k = 0, and in this case a tableau $S \in \text{Tab}(n, 0, d)$ has the form

$$S = \frac{i_1 \cdots i_d}{|i_{d+1}| \cdots |i_{n-k}| j_1 \cdots j_k}$$

$$(6)$$

and its associated shifted Specht polynomial is the monomial

$$M_S = x_{i_1} \cdots x_{i_d},$$

which we sometimes write as M_S^d to remember degree.

The *d-shifted Specht module of* $\lambda = (n - k, k)$, denoted by V(n, k, d), is defined to be the \mathbb{F} -linear span of the *d*-shifted Specht polynomials, i.e.

$$V(n, k, d) = \langle F_T(d) \mid T \in \text{Tab}(n, k, d) \rangle$$
;

like the Specht module, it is also an \mathfrak{S}_n -representation, although it is not irreducible for d > k, cf. Section 3. The *d-shifted Specht ideal of* $\lambda = (n - k, k)$, denoted by $\mathfrak{a}(n, k, d)$, is the ideal in R generated by the shifted Specht module, i.e.

$$\mathfrak{a}(n,k,d) = V(n,k,d) \cdot R = (F_T(d) \mid T \in \mathsf{Tab}(n,k,d)).$$

The remainder of the section will be devoted to finding a basis for the shifted Specht module V(n, k, d) and to computing its dimension.

2.1 Basis of a Shifted Specht Module

Theorem 2.1. A basis for the d-shifted Specht module V(n,k,d), and hence a minimal generating set of the ideal $\alpha(n,k,d)$, are indexed by the standard tableaux on the d-shifted shape $\lambda(d)$, i.e.

$$\{F_T(d) \mid T \in \operatorname{STab}(\lambda(d))\}\$$
.

The proof of Theorem 2.1 comes in two steps, which we state as Lemmas.

Lemma 2.2. The set $\{F_T(d)|T\in STab(n,k,d)\}$ is linearly independent.

Proof. By induction on $n \ge 2$. The base case, where n = 2 and k = d = n - k = 1 is trivial. For the inductive step, we assume that for every choice of k' and d' satisfying $1 \le k' \le d' \le (n-1)-k'$, the set $\{F_{T'}(d')|T' \in \operatorname{STab}(n-1,k',d')\}$ is linearly independent. Fix any integers $1 \le k \le d \le n - k$ and suppose we have a dependence relation

$$\sum_{T \in \text{STab}(n,k,d)} c_T F_T(d) = 0. \tag{7}$$

Note that for every d-standard tableau $T \in \operatorname{STab}(n,k,d)$ as in (5), we must have either $n=i_d$ or $n=j_k$; let $I_d \subset \operatorname{STab}(n,k,d)$ denote the set of d-standard tableaux with $n=i_d$ and let $J_d \subset \operatorname{STab}(n,k,d)$ be the ones with $n=j_k$. Let $\pi\colon R\to S=\mathbb{F}[x_1,\ldots,x_{n-1}]$ be the projection sending x_n to 0, and note that for every $T\in I_d$ we have $\pi(F_T(d))=0$. Moreover for every $T\in J_d$ we have $\pi(F_T(d))=F_{T'}(d)$ where

$$T' = \frac{i_1 \cdots i_{k-1} i_k |i_{k+1} \cdots |i_d|}{i_{d+1} \cdots |i_{n-k}| j_1 \cdots j_{k-1}}$$

We see that $T' \in \operatorname{STab}(n-1,k-1,d)$, hence the induction hypothesis applies. Note that since the map $J_d \to \operatorname{STab}(n-1,k-1,d)$, sending $T \mapsto T'$ is one-to-one, and since

$$\pi \left(\sum_{T \in \operatorname{STab}(n,k,d) = I_d \sqcup J_d} c_T F_T(d) \right) = \sum_{T \in J_d} c_T F_{T'}(d) = 0,$$

we deduce, by the induction hypothesis, that $c_T = 0$ for all $T \in J_d$. Then our dependence relation (7) becomes

$$\sum_{T \in I_d} c_T F_T(d) = 0 = x_n \cdot \left(\sum_{T \in I_d} c_T F_{T''}(d) \right)$$

where

$$T^{\prime\prime} = \frac{\begin{vmatrix} i_1 & \cdots & i_k & |i_{k+1}| \cdots |i_{d-1}| \\ |i_{d+1}| & \cdots & |i_{n-k}| & |j_1| & \cdots & |j_k| \end{vmatrix}}{\begin{vmatrix} i_{d+1}| & \cdots & |i_{n-k}| & |j_1| & \cdots & |j_k| \end{vmatrix}}$$

In this case we see that $T'' \in \operatorname{STab}(n-1,k,d-1)$, and again our induction hypothesis applies. Since the map $I_d \to \operatorname{STab}(n-1,k,d-1)$ sending $T \mapsto T''$ is one-to-one, and since

$$\sum_{T \in I_d} c_T F_{T''}(d) = 0$$

the induction hypothesis implies that $c_T = 0$ for every $T \in I_d$ too, and therefore the dependence relation (7) must be trivial.

To see that the *d*-standard polynomials span V(n, k, d) is a little more work. We will follow the general method described in [12, Section 2.6]. For T as in (5) and any index $1 \le a \le n$, define *a-composition vector* to be the integer vector with n - k components defined by

$$\gamma^{a}(T) = (\gamma_{1}^{a}(T), \dots, \gamma_{n-k}^{a}(T)), \text{ where } \gamma_{b}^{a}(T) = \#\{c \in \text{col}_{b}(T) \mid c \leq a\}.$$

Note that for a two rowed partition $\lambda = (n - k, k)$ the *a*-composition vector has entries 0, 1, or 2. Define the *composition series* for T to be the *n*-tuple of composition vectors $\gamma(T) = (\gamma^1(T), \dots, \gamma^n(T))$; we regard $\gamma(T)$ as a matrix whose columns are the composition vectors of T. Given two vectors $v = (v_1, \dots, v_{n-k})$ and $w = (w_1, \dots, w_{n-k})$, we say that w dominates v, and write $v \triangleleft w$, if $v_1 + \dots + v_p \leq w_1 + \dots + w_p$ for all $1 \leq p \leq n - k$. Finally given two tableaux $T, T' \in \text{Tab}(\lambda(d))$, we say that T' dominates T, and write $T \triangleleft T'$, if every composition vector of T' dominates the corresponding composition vector of T, i.e.

$$T \triangleleft T' \Leftrightarrow \gamma^a(T) \triangleleft \gamma^a(T'), \forall 1 \leq a \leq n.$$

This composition-dominance order is a partial order on the set of tableaux $Tab(\lambda(d))$. Moreover it is clear that the largest tableau is the one that fills the columns in order from left to right. For example, if n = 5, k = 1, and d = 3 the largest tableau (with increasing columns) is the standard tableau

$$T = \begin{array}{c|c|c} 2 & 4 & 5 \\ \hline 1 & 3 & \end{array}$$

with composition series

$$\gamma(T) = \left(\begin{array}{ccccc} 1 & 1 & 1 & 1 & 1 \\ 0 & 1 & 2 & 2 & 2 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 \end{array}\right)$$

Note that if T and T' have the same columns (possibly in different orders) then they have the same composition series, and they also have the same shifted Specht polynomials (up to sign), i.e. $\gamma(T) = \gamma(T')$ and $F_T(d) = \pm F_{T'}(d)$. The following lemma is useful for telling when one tableau dominates another.

Lemma 2.3. If $1 \le a < b \le n$ and a appears in a column to the right of b, then

$$T \triangleleft (a,b) \cdot T$$

where $(a,b) \cdot T$ is the tableau obtained from T by transposing a and b.

Proof. Note that for $1 \le i \le a-1$ and for $b \le i \le n$ we have $\gamma^i(T) = \gamma^i((a,b) \cdot T)$. Assume then that $a \le i \le b-1$, and suppose that a and b belong to columns r and q in T, respectively. Then

$$\gamma^{i}((a,b) \cdot T) = \gamma^{i}(T)$$
 with r^{th} part decreased by 1 and q^{th} part increased by 1.

and since we are assuming that q < r, it follows that $\gamma^i(T) \triangleleft \gamma^i((a,b) \cdot T)$, and the result follows.

We are now in a position to show the standard shifted Specht polynomials span the shifted Specht module.

Lemma 2.4. The set $\{F_T(d)|T\in STab(n,k,d)\}\$ spans V(n,k,d).

Proof. We will show by downward induction on the composition-dominance order on $\text{Tab}(\lambda(d))$ that for every $T \in \text{Tab}(\lambda(d))$, $F_T(d)$ can be written as a linear combination of shifted Specht polynomials associated to d-standard tableaux. For the base case, note, as above, that the largest tableau of shifted shape $\lambda(d)$ is already standard. Inductively, fix a shifted tableau $T \in \text{Tab}(\lambda(d))$ as in (5), and assume that for every tableau T' that dominates T, $F_{T'}(d)$ can be written as a linear combination of shifted Specht polynomials corresponding to standard tableaux. We may assume that the columns of T are increasing. If T has no row descents, then T must be standard and we are done. Otherwise T has some row descent, say between the a^{th} and $a + 1^{st}$ column. There are several cases to consider and we claim that in all cases we can write $F_T(d)$ as a linear combination

$$F_T(d) = \sum_{T \triangleleft T'} c_{T'} F_{T'}(d).$$

Case 1: $1 \le a \le n - k - d - 1$. Set b = d + a; in this case, we can merely swap i_b and i_{b+1} without affecting $F_T(d)$; in other words setting $T' = (i_b, i_{b+1}) \cdot T$ we have

$$F_T(d) = F_{T'}(d)$$

and since $T \triangleleft (i_b, i_{b+1}) \cdot T$, it follows from our inductive hypothesis that $F_T(d)$ can be written as a linear combination of d-standard Specht polynomials on the shifted shape $\lambda(d)$.

Case 2: a = n - k - d. Then we have

$$T = \frac{i_1 \cdots i_k |i_{k+1}| \cdots |i_d|}{|i_{d+1}| \cdots |i_{n-k}| |j_1| \cdots |j_k|}$$

and we have the descent $i_{n-k} > j_1 > i_1$. Then by lemma 2.3 we see that

$$T \triangleleft T' = (i_{n-k}, j_1) \cdot T$$
 and $T \triangleleft T'' = (i_1, i_{n-k}) \cdot T$.

Moreover one can easily check that

$$F_T(d) = F_{T'}(d) - F_{T''}(d) = G\left((x_{i_1} - x_{i_{n-k}}) - (x_{j_1} - x_{i_{n-k}})\right)$$

and inductive hypothesis applies.

Case 3: $n - k - d + 1 \le a \le n - d - 1$. Set b = a - n + k + d so we have

Sub-Case 3a: $i_{b+1} < i_b < j_b$. Then

$$T \triangleleft T' = (i_{b+1}, j_b) \cdot T$$
 and $T \triangleleft T'' = (i_{b+1}, i_b) \cdot T$

and again one can easily check that

$$F_T(d) = F_{T'}(d) + F_{T''}(d) = G \cdot \left((x_{i_b} - x_{i_{b+1}})(x_{j_b} - x_{j_{b+1}}) + (x_{i_{b+1}} - x_{j_b})(x_{i_b} - x_{j_{b+1}}) \right).$$

and our inductive hypothesis applies.

Sub-Case 3b: $i_b < i_{b+1} < j_{b+1} < j_b$. Then

$$T \triangleleft T' = (i_{b+1}, j_b) \cdot T$$
 and $T \triangleleft T'' = (j_{b+1}, j_b) \cdot T$

and again one can easily check that

$$F_T(d) = F_{T'}(d) + F_{T''}(d) = G \cdot \left((x_{i_b} - x_{i_{b+1}})(x_{j_b} - x_{j_{b+1}}) + (x_{i_b} - x_{j_{b+1}})(x_{i_{b+1}} - x_{j_b}) \right)$$

and our inductive hypothesis applies.

Case 4: a = n - d. Then

$$T = \frac{i_1 \cdots i_k |i_{k+1}| \cdots |i_d|}{|i_{d+1}| \cdots |i_{n-k}| |j_1| \cdots |j_k|}$$

and we have the descent $j_k > i_k > i_{k+1}$. Then by Lemma 2.3 we have

$$T \triangleleft T' = (i_{k+1}, i_k) \cdot T$$
 and $T \triangleleft T'' = (i_{k+1}, i_k) \cdot T$.

and again one can check that

$$F_T(d) = F_{T'}(d) + F_{T''}(d) = G\left(x_{i_k}\left(x_{i_{k+1}} - x_{j_k}\right) + x_{j_k}\left(x_{i_k} - x_{i_{k+1}}\right)\right)$$

to which the inductive hypothesis once again implies.

Case 5: $n-d+1 \le a \le n-k-1$. In this case set b=a-(n-k-d) so that we have

$$T = \frac{\begin{vmatrix} i_1 & \cdots & i_k & i_{k+1} & \cdots & i_b & i_{b+1} & \cdots & i_d \end{vmatrix}}{\begin{vmatrix} i_{d+1} & \cdots & i_{n-k} & j_1 & \cdots & j_k \end{vmatrix}}$$

with the descent $i_b > i_{b+1}$. Hence by Lemma 2.3 we have

$$T \triangleleft T' = (i_{b+1}, i_b) \cdot T$$

and in this case we clearly have

$$F_T(d) = F_{T'}(d)$$

to which the induction hypothesis applies again.

Therefore in all cases we have shown that $F_T(d)$ is a linear combination of shifted Specht polynomials indexed by tableaux on the shifted shape $\lambda(d)$ which dominate T. Therefore by induction, the d-standard shifted Specht polynomials

$$\{F_T(d)|T\in \operatorname{STab}(\lambda(d))\}\$$

must span V(n, k, d).

Proof of Theorem 2.1. By Lemma 2.2, the polynomials in the set

2 | 4

$$\{F_T(d) \mid T \in \operatorname{STab}(\lambda(d))\}\$$

are linearly independent, and by Lemma 2.4 they generate the shifted Specht module V(n, k, d), and hence they must form a basis.

Example 2.5. Let n = 5, k = 1, and d = 3. There are 9 standard tableau of shifted shape $\lambda(3)$ where $\lambda = (4, 1)$:

Hence a minimal generating set for the ideal a(5, 1, 3), and a basis for the representation V(5, 1, 3), is given by

$$\begin{cases}
(x_2 - x_3)x_4x_5, & (x_2 - x_4)x_3x_5, & (x_2 - x_5)x_3x_4, \\
(x_1 - x_3)x_4x_5, & (x_1 - x_4)x_3x_5, & (x_1 - x_5)x_3x_4, \\
(x_1 - x_4)x_2x_5, & (x_1 - x_5)x_2x_4, & (x_1 - x_5)x_2x_3
\end{cases}.$$

The utility of Theorem 2.1 is that it can transform set maps on the set of standard tableau $STab(\lambda(d))$ to linear maps on the shifted Specht module V(n, k, d), and sometimes the shifted Specht ideal $\mathfrak{a}(n, k, d)$. The following useful corollaries illustrate this point.

First note every standard tableau $T \in STab(n, k, k)$ has n in its support; in fact it must have the form

$$T = \frac{i_1 \cdots i_{k-1} i_k}{|i_{k+1}| \cdots |i_{n-k}| j_1 \cdots j_{k-1}| n}$$

We get a standard tableau $T' \in STab(n-1, k-1, k)$ from T by deleting the box containing n, i.e.

$$T' = \frac{\begin{vmatrix} i_1 & \cdots & i_{k-1} & i_k \\ \vdots & \vdots & \vdots & \vdots \\ i_{k+1} & \cdots & i_{n-k} & j_1 & \cdots & j_{k-1} \end{vmatrix}}{\begin{vmatrix} i_k & \cdots & \vdots \\ \vdots & \vdots & \vdots \\ \vdots & \vdots & \vdots \\ \end{vmatrix}}$$

Since this map is obviously a bijection it extends to a linear isomorphism

$$V(n, k, k) \rightarrow V(n-1, k-1, k)$$
.

In fact it can be extended to a ring isomorphism of Specht ideals using a linear change of coordinates: Define the variables y_1, \ldots, y_n by the formula

$$y_i = \begin{cases} x_n - x_i & \text{if } 1 \le i \le n - 1 \\ x_n & \text{if } i = n \end{cases}$$

and let $\Phi: R \to R$ be the change of coordinates map $\Phi(x_i) = y_i$.

Corollary 2.6. Fix integers k, n satisfying $1 \le k \le n - k$. Then we have a ring isomorphism

$$\mathfrak{a}(n, k, k) \cong \Phi(\mathfrak{a}(n, k, k)) = \mathfrak{a}(n - 1, k - 1, k).$$

In fact, in adding the principal ideal (x_n) , this isomorphism becomes equality:

$$a(n, k, k) + (x_n) = a(n - 1, k - 1, k) + (x_n).$$

Proof. With $T \in STab(n, k, k)$ and $T' \in STab(n - 1, k - 1, k)$ as above, we compute

$$F_T = \prod_{t=1}^{k} (x_{i_t} - x_{j_t}) = \prod_{t=1}^{k-1} (x_{i_t} - x_{j_t}) \cdot (x_{i_k} - x_n) = \prod_{t=1}^{k-1} (y_{j_t} - y_{i_t}) \cdot (-y_{i_k}) = (-1)^k \cdot \Phi(F_{T'})$$

and hence by Theorem 2.1, Φ sends a the first statement follows. To see the second statement note that the ring automorphism $\Phi: R \to R$ becomes the negative identity map on the quotient $\bar{\Phi} = -I: R/(x_n) \to R/(x_n)$, and hence

$$a(n, k, k) + (x_n) = a(n - 1, k - 1, k) + (x_n)$$

which is the second statement.

We say that an index i, $1 \le i \le n$ is in the support of tableau $T \in \text{Tab}(n, k, d)$, and write $i \in \text{supp}(T)$, if i appears in or to the right of a column with more than one row, i.e. if

$$T = \frac{i_1 \cdots i_k |i_{k+1}| \cdots |i_d|}{|i_{d+1}| \cdots |i_{n-k}| |j_1| \cdots |j_k|}$$

then $\operatorname{supp}(T) = \{i_1, j_1, \dots, i_k, j_k, i_{k+1}, \dots, i_d\}$. Fix an integer m satisfying $0 \le m \le d$, and define the subset $\operatorname{STab}_m(n, k, d) \subset \operatorname{STab}(n, k, d)$ consisting of standard tableaux which contain $\{1, \dots, m\}$ in its support. A standard tableau $T \in \operatorname{STab}_m(n, k, d)$ necessarily has the form

$$T = \frac{1 \cdots m_{m+1} \cdots k_{k+1} \cdots d}{i_{d+1} \cdots i_{n-k} j_1 \cdots j_m j_{m+1} \cdots j_k}$$

and we define its image tableau as the one obtained by removing the boxes containing the numbers $1, \ldots, m$:

$$T' = \frac{\begin{vmatrix} m+1 & \cdots & k & k+1 & \cdots & d \\ |i_{d+1}| & \cdots & |i_{n-k}| & |j_1| & \cdots & |j_m| & |j_{m+1}| & \cdots & |j_k| \end{vmatrix}}$$

Note that $T' \in \operatorname{STab}([n]_m, k-m, d-m)$ where $[n]_m$ means the tableaux are filled with numbers $\{m+1,\ldots,n\}$, and we count k-m as zero if $k \leq m$. The map of sets $\operatorname{STab}_m(n,k,d) \ni T \mapsto T' \in \operatorname{STab}([n]_m, k-m, d-m)$ is evidently one-to-one and onto, and by Theorem 2.1, it induces an bijective linear map of Specht modules, which is the key to some of the main technical arguments in this paper.

Corollary 2.7. *The induced linear map*

$$V_m(n, k, d) \longrightarrow V([n]_m, k - m, d - m)$$

 $F_T(d) \longmapsto F_{T'}(d - m)$

is bijective.

2.2 Dimension of a Shifted Specht Module

We compute the dimension of the shifted Specht module V(n, k, d) by counting the standard tableaux on the shifted shape $\lambda(d)$. As a first step, we observe that a standard tableau T on $\lambda(d)$ is uniquely determined by its d-row, or its (n-d)-row, i.e. if

$$T = \frac{\begin{vmatrix} i_1 & \cdots & i_k & i_{k+1} & \cdots & i_d \\ i_{d+1} & \cdots & i_{n-k} & j_1 & \cdots & j_k \end{vmatrix}}{\begin{vmatrix} i_{d+1} & \cdots & i_{n-k} & j_1 & \cdots & j_k \end{vmatrix}}$$

its d-row is the sequence (i_1, \ldots, i_d) , and its (n-d)-row is $(i_{d+1}, \ldots, i_{n-k}, j_1, \ldots, j_k)$.

Lemma 2.8. A necessary and sufficient condition for an increasing sequence of d integers in [n], say $\{i_1 < \cdots < i_d\}$, to be a d-row of a standard tableau of shape $\lambda(d)$ is that for each index $1 \le t \le k$, we have

$$\#\{j \in [n] \setminus \{i_1, \dots, i_d\} \mid j > i_{k+1-t}\} \ge t.$$
 (8)

Proof. Assume that (8) holds. Write the compliment $[n] \setminus \{i_1, \ldots, i_d\}$ sequence in increasing order as $i_{d+1} < \cdots < i_{n-k} < j_1 < \cdots < j_k$. Then Condition (8) implies that $i_t < j_t$ for each $1 \le t \le k$, and hence the tableau with d-row (i_1, \ldots, i_d) must be standard. Conversely, suppose that T in (8) is standard. Then for each $1 \le t \le k$ we have $i_t < j_t < j_{t+1} < \cdots < j_k$, which implies Condition (8).

In fact, we can encode a tableau T, with increasing rows but not necessarily standard, on the shifted shape $\lambda(d)$ as a NE lattice path from (0,0) to (n-d,d). Then the standard tableaux will correspond to NE lattice paths which stay below the shifted diagonal line y = x + (d-k). Here are some more precise statements.

Definition 2.9. Define a NE lattice path from (a, b) to (a', b') to be a sequence of points on the \mathbb{Z}^2 lattice, $P_0 = (a, b), P_1, \dots, P_m = (a', b')$ where if $P_i = (a_i, b_i)$ then either $P_{i+1} = (a_i + 1, b_i)$ in which case the i^{th} step is E, or $P_{i+1} = (a_i, b_i + 1)$ in which case the i^{th} step is N.

The following fact is well known.

Fact 2.10. The number of NE lattice paths from (a,b) to (a',b') is the binomial coefficient


$$\underbrace{(a'+b')-(a+b)}^{m} b'-b$$

There is a map from the set of NE lattice paths from (0,0) to (n-d,d) to the set of tableaux of shifted shape $\lambda(d)$ with increasing rows: label the i^{th} step n-i+1, i.e.

$$\overline{P_i P_{i+1}} \mapsto n - i + 1 \tag{9}$$

and fill in the shifted diagram $\lambda(d)$ with these numbers from right to left with the N-steps in the k-row and the E-steps in the (n-k)-row.


Example 2.11. Let n = 11, k = 4, d = 6, so that n - d = 5. Then setting $\lambda = (7, 4)$, the NE lattice path from (0, 0) to (n - d, d) = (5, 6) given by



corresponds to the tableau of shifted shape given by

	1	4	6	7	8	11
2	3	5	9	10		

Note that the NE lattice path here is sub-shifted-diagonal, and that the Young tableau here is standard. A non-sub-shifted-diagonal NE lattice path is given by



and corresponds to the non-standard Young tableau

The theorem, evidenced by Example 2.11, is that the above map gives a bijection between sub-shifted-diagonal NE lattice paths and standard Young tableaux.

Lemma 2.12. There is a bijection between NE lattice paths from (0,0) to (n-d,d) which do not cross the shifted diagonal line y = x + (d-k) and standard tableaux of shifted shape $\lambda(d)$.

Proof. Translating condition (8) from Lemma 2.8 into the language of NE lattice paths, it says:

E-steps to the left of the
$$(d - k + t)^{th}$$
 N-step $\geq t$. (10)

Hence if the $(d-k+t+1)^{th}$ *N*-step is the p^{th} step in the path, then the point $P_p = (e_p, n_p)$ has coordinates that satisfy $n_p = (d-k+t)$ and $e_p \ge t$, and hence $n_p - e_p \le d-k$. This implies that the point P_p lies below the shifted diagonal line y = x + (d-k), as desired.

Lemma 2.12 translates our problem of counting standard tableaux of shifted shape $\lambda(d)$ into the problem of counting sub-shifted-diagonal lattice paths from (0,0) to (n-d,d). The advantage of counting lattice paths lies in the so-called reflection principal; a proof can be found in [3].

Lemma 2.13 (Reflection Principle). Let ℓ : y = x + c be any line with slope 1, and suppose that (a',b') and (a,b) are two lattice points that lie on the same side of ℓ . Then the number of NE lattice paths from (a',b') to (a,b) which touch or cross ℓ is equal to the total number of NE lattice paths from (a'',b'') to (a,b), where (a'',b'') is the reflection of (a',b') over ℓ .

We are now in a position to compute the dimension of the shifted Specht module V(n, k, d).

Theorem 2.14. The number of NE lattice paths from (0,0) to (n-d,d) which do not cross the shifted diagonal y = x + (d-k), and hence the number of standard tableaux on the shifted shape $\lambda(d)$, is equal to

$$\dim(V(n,k,d)) = \binom{n}{d} - \binom{n}{k-1}.$$

Proof. To count the number of shifted subdiagonal NE lattice paths from (0,0) to (n-d,d) is the same as to count the NE lattice paths which do not touch or cross the line y = x + (d-k) + 1. Let us instead count the number of NE lattice paths from (0,0) to (n-d,d) that do touch or cross the line y = x + (d-k) + 1, which, according to the Reflection Principle, is the same as the total number of NE lattice paths from (-(d-k)-1,(d-k)+1) to (n-d,d). According to Fact 2.10, this number is

$$\binom{(n-d+d)+(-(d-k)-1+(d-k)+1)}{d-(d-k)-1} = \binom{n}{k-1}.$$
 (11)

We can also count the total number of NE lattice paths from (0,0) to (n-d,d) as

$$\binom{(n-d+d)-(0+0)}{d-0} = \binom{n}{d}.$$
(12)

We therefore get the number of NE lattice paths from (0,0) to (n-d,d) which do not cross the line y = x + (d-k) by subtracting (11) from (12), which gives the desired formula.

3 Lefschetz Propoerties

Let $E \subset R$ be the graded vector subspace spanned by square-free monomials in the variables x_1, \ldots, x_n . It will be convenient to identify E with a quotient of R as well, specifically the Artinian monomial complete intersection

$$A = \frac{\mathbb{F}[x_1, \dots, x_n]}{(x_1^2, \dots, x_n^2)}.$$

Define the following linear operators on A: The *raising operator* is multiplication by the sum of variables:

$$L = \times (x_1 + \cdots + x_n) \colon A \to A[1],$$

the *lowering operator* is the partial derivative map corresponding to the sum of variables:

$$D = \frac{\partial}{\partial x_1} + \dots + \frac{\partial}{\partial x_n} \colon A \to A[-1],$$

and the semi-simple operator:

$$H: A \rightarrow A, \ H(a) = (n-2k) \cdot a, \ \forall \ a \in A_k.$$

Lemma 3.1. The operators $\{D, L, H\}$ forms an \mathfrak{sl}_2 -triple on $E \cong A$, i.e. they satisfy the commutator relations:

$$[D, L] = H, [H, D] = 2D, [H, L] = -2L.$$

Proof. Since D, L, and H are linear, it suffices to check the relations on monomials, and by symmetry it suffices to check only the single square-free monomial $\mu = x_1 \cdots x_i$. We have

$$D \circ L(m) = D\left(\sum_{j=i+1}^{n} x_{1} \cdots x_{i} x_{j}\right) = \sum_{j=i+1}^{n} D(x_{1} \cdots x_{i} \cdot x_{j}) = \sum_{j=i+1}^{n} \left(x_{1} \cdots x_{i} + \sum_{k=1}^{i} x_{1} \cdots \hat{x}_{k} \cdots x_{i} \cdot x_{j}\right)$$

$$= (n-i) \cdot m + \sum_{j=i+1}^{n} \sum_{k=1}^{i} x_{1} \cdots \hat{x}_{k} \cdots x_{i} x_{j}$$

$$(13)$$

$$L \circ D(m) = L\left(\sum_{k=1}^{i} x_{1} \cdots \hat{x}_{k} \cdots x_{i}\right) = \sum_{k=1}^{i} \left(L(x_{1} \cdots \hat{x}_{k} \cdots x_{i})\right) = \sum_{k=1}^{i} \left(x_{1} \cdots x_{i} + \sum_{j=i+1}^{n} x_{1} \cdots \hat{x}_{k} \cdots x_{i} \cdot x_{j}\right)$$

$$= i \cdot m + \sum_{k=1}^{i} \sum_{j=i+1}^{n} x_{1} \cdots \hat{x}_{k} \cdots x_{i} \cdot x_{j}.$$

$$(14)$$

Subracting (13) and (14) yields $[D, L](m) = D \circ L(m) - L \circ D(m) = (n-2i) \cdot m = H(m)$, and hence verifies the relation [D, L] = H. Verifications of the other two relations are straightforward and left to the reader.

For each $0 \le i \le n$, define the i^{th} -primitive subspace $P_i \subset A_i$ by

$$P_i = \ker(D) \cap A_i = \{\alpha \in A_i \mid D(\alpha) = 0\}.$$

It follows from Lemma 3.1 that for any positive integer m and for any $\alpha \in P_k$ we have

$$D(L^{m}(\alpha)) = m \cdot (n - 2k + 1 - m) \cdot L^{m-1}(\alpha).$$

Also note that for any Specht polynomial $F_T \in V(n, k, k) \subset A_k$, we have

$$D(F_T) = 0.$$

In particular, we have a chain of containments

$$V(n, k, k) \subseteq P_k \subseteq \ker(L^{n-2k+1}) \cap A_k$$
.

Lemma 3.2. Equality $V(n, k, k) = P_k$ holds if and only if the derivative map, i.e. the lowering operator

$$D: A_{k} \to A_{k-1}$$

is surjective.

Proof. Assume that $V(n, k, k) = P_k$. Then by Theorem 2.14 we have

$$\dim(P_k) = \binom{n}{k} - \binom{n}{k-1} = \dim(A_k) - \dim(A_{k-1}).$$

Let $I_{k-1} = D(A_k)$ be the image of the derivative map. By linear algebra $\dim(I_{k-1}) + \dim(P_k) = \dim(A_k)$ hence $\dim(I_{k-1}) - \dim(A_{k-1}) = 0$, hence the derivative map is surjective. Conversely, if $D: A_k \to A_{k-1}$ is surjective, then $\dim(I_{k-1}) = \dim(A_{k-1})$, hence $\dim(P_k) = \dim(A_k) - \dim(A_{k-1}) = \dim(V(n,k,k))$. Since $V(n,k,k) \subseteq P_k$, and they have the same dimension, this containment must be equality.

As we shall see, surjectivity of the derivative map in Lemma 3.2 is dictated by the weak Lefschetz property.

3.1 Weak Lefschetz Property

For an arbitrary graded Artinian algebra C = R/I, we say that C has the weak Lefschetz property if there is a linear form $\ell \in C_1$ such that the multiplication maps

$$\times \ell \colon C_{i-1} \to C_i \tag{15}$$

have maximum rank for every degree $i \ge 0$; in this case we call ℓ a weak Lefschetz element for C. If C has a unimodal Hilbert function with socle degree d, then $\ell \in C_1$ is Lefschetz if and only if the multiplication maps (15) are injective for $1 \le i \le \left\lfloor \frac{d+1}{2} \right\rfloor$ and surjective for $\left\lfloor \frac{d+3}{2} \right\rfloor \le i \le d$. If C is Gorenstein, then it suffices only to check that (15) is injective in degrees $1 \le i \le \left\lfloor \frac{d+1}{2} \right\rfloor$. In fact one can show that if C is Gorenstein with the standard grading and if the multiplication map (15) is injective for some i_0 , then it is injective for all $i \le i_0$. Moreover if the ideal I is generated by monomials then C is weak Lefschetz if and only if $x_1 + \cdots + x_n \in C_1$ is a weak Lefschetz element. For more details, especially regarding these last two facts, see [9, Propositions 2.1, 2.2] or [13, Proposition 2.5].

In our situation A is a standard graded Artinian Gorenstein algebra with unimodal Hilbert function and cut out by a monomial ideal. In fact, in our situation, the matrix for the multiplication map $L: A_{k-1} \to A_k$ in the monomial basis is the transpose of the derivative map $D: A_k \to A_{k-1}$. Therefore we see that A has the weak Lefschetz property if and only if the derivative maps

$$D: A_k \to A_{k-1}$$

are surjective for all $1 \le k \le \left\lfloor \frac{n+1}{2} \right\rfloor$. The following result is due to Kustin-Vraciu [7], and we refer the reader there for a proof. As usual, $p = \operatorname{char}(\mathbb{F}) \ge 0$.

Lemma 3.3. The monomial complete intersection A has the weak Lefschetz property if and only if p = 0 or $p \ge \lfloor \frac{n+3}{2} \rfloor$.

From Lemma 3.3 we derive the following useful result.

Lemma 3.4. Fix any integer k satisfying $2k \le n$. Then the following are equivalent:

- 1. p = 0 or $p \ge k + 1$.
- 2. The derivative maps $D: A_i \to A_{i-1}$ are surjective for all $1 \le i \le k$.
- 3. The derivative map $D: A_k \to A_{k-1}$ is surjective.

Proof. (1.) \Rightarrow (2.). Assume that p = 0 or $p \ge k + 1$. For each index j, $2k \le j \le n$, define the nested chain of monomial complete intersections

$$C_{2k} \subset \cdots \subset C_n$$
, where $C_j = \frac{\mathbb{F}[x_1, \dots, x_j]}{(x_1^2, \dots, x_j^2)}$

By Lemma 3.3, the first monomial complete intersection C_{2k} has the weak Lefschetz property, and in particular the derivative map

$$D_{C_{2k}}: (C_{2k})_i \to (C_{2k})_{i-1}$$

is surjective for all $1 \le i \le k$. Inductively for j > 2k, note that for each $0 \le i \le k$ we have direct sum decomposition of graded vector spaces

$$(C_j)_i \cong (C_{j-1})_i \oplus x_j \cdot (C_{j-1})_{j-1}$$

and in particular the derivative map in degree k decomposes into block triangular form

$$D_{C_{j}} \colon \left(C_{j}\right)_{i} \to \left(C_{j}\right)_{i-1} = \begin{pmatrix} D_{C_{j-1}} \colon \left(C_{j-1}\right)_{i} \to \left(C_{j-1}\right)_{i-1} & I \colon \left(C_{j-1}\right)_{i-1} \to \left(C_{j-1}\right)_{i-1} \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & & \\ & & & \\ & & & & \\$$

Since the maps

$$D_{C_{j-1}}: \left(C_{j-1}\right)_i \rightarrow \left(C_{j-1}\right)_{i-1}$$

is surjective for all $1 \le i \le k$, it follows that the maps

$$D_{C_j}: (C_j)_i \to (C_j)_{i-1}$$

is also surjective for all $1 \le i \le k$. In particular, this argument shows that the derivative maps $D: A_i \to A_{i-1}$ must be surjective for all $1 \le i \le k$ as well.

- $(2.) \Rightarrow (3.)$ Obvious.
- $(3.) \Rightarrow (1.)$ Assume that $0 . Then setting <math>i = p \le k$, we claim that $D: A_p: A_{p-1}$ cannot be surjective. Indeed, set $\alpha \in A_p$ to be the p^{th} -elementary symmetric polynomial in the first $2p-1 \le n-1$ variables:

$$\alpha = e_p(x_1, \ldots, x_{2p-1}) \in A_p.$$

Note first that over a field of characteristic p, we have

$$D(\alpha) = p \cdot e_{p-1}(x_1, \dots, x_{2p-1}) \equiv 0.$$

On the other hand, we have

$$\alpha(1,\ldots,1) = \binom{2p-1}{p} = \frac{(2p-1)\cdots(p+1)}{(p-1)!} \neq 0.$$

This shows that $\alpha \in P_p = \ker(D) \cap A_p$, but $\alpha \notin V(n, p, p)$ (since every polynomial in V(n, p, p) necessarily vanishes at any point in which at least p + 1-entries are equal). Therefore, by Lemma 3.2, $D: A_p \to A_{p-1}$ is not surjective, and hence by [9, Proposition 2.1], $D: A_k \to A_{k-1}$ cannot be surjective either.

We can also derive a useful result on specialization of Specht modules.

Lemma 3.5. Assume that p = 0 or $p \ge k + 1$, fix j satisfying $2k \le j \le n$, and set

$$B = \frac{\mathbb{F}[x_1, \dots, x_j]}{(x_1^2, \dots, x_j^2)}.$$

Then

$$V(j, k, k) = V(n, k, k) \cap B_k$$
.

Proof. Containment $V(j, k, k) \subseteq V(n, k, k) \cap B_k$ is clear. For the reverse containment, suppose that $\alpha \in V(n, k, k) \cap B_k$. The key observation here is that the restriction of the derivative map on A, call it D_A , to the subspace $B \subset A$ is the same as D_B . Since $\alpha \in V(n, k, k) \subseteq P_{A,k} = \ker(D_A) \cap A_k$, it follows that $D_A(\alpha) = 0$, and hence also $D_B(\alpha) = 0$. This means that $\alpha \in P_{B,k}$. From the proof of Lemma 3.4, we deduce that $D_B \colon B_k \to B_{k-1}$ is surjective, which by Lemma 3.2 implies that $\alpha \in V(j, k, k)$, as desired.

As we shall see, the weak Lefschetz property, or lack thereof, can be used to detect embedded primary components of our Specht-monomial ideals. Next we shall use the strong Lefschetz property to decompose our shifted Specht modules into irreducible \mathfrak{S}_n -representations.

3.2 Strong Lefschetz Property

The pair (A, L) is *strong Lefschetz* if the restriction of L^i to the graded components of A always has maximal rank, or equivalently if

$$L^{n-2k}: A_k \to A_{n-k}$$

are isomorphisms for all $0 \le k \le \left| \frac{n}{2} \right|$. One can show that we have containment

$$V(n, k, d) \subseteq \ker \left(L^{n-k-d+1}\right) \cap A_k.$$

Lemma 3.6. Let $p = \text{char}(\mathbb{F})$. The following are equivalent.

- 1. p = 0 or $p \ge n + 1$,
- 2. The pair (A, L) is strong Lefschetz.
- 3. For all integers $0 \le k \le d \le n k$, the shifted Specht modules $V(n, k, d) \subset A$ satisfy

$$V(n,k,d) = \bigoplus_{i=k}^{d} L^{d-i}(P_i) = \ker\left(L^{n-k-d+1}\right) \cap A_d.$$

Proof. (1.) \Rightarrow (2.). This is due to Ikeda; see [6, Proposition 3.66].

- (2.) \Rightarrow (1.). Note that if $p \le n$, then we must have $L^p = 0$, and (A, L) cannot be strong Lefschetz.
 - $(2.) \Rightarrow (3.)$. If (A, L) is strong Lefschetz then the map

$$L^{n-k-d+1}: A_d \to A_{n-k+1}$$

must have maximal rank, and hence we must have

$$\dim(\ker(L^{n-k-d+1})\cap A_d) = \dim(A_d) - \dim(A_{n-k+1}) = \binom{n}{d} - \binom{n}{k-1} = \dim(V(n,k,d)).$$

But since we already have containment $V(n, k, d) \subseteq \ker(L^{n-k-d+1}) \cap A_d$ it must be equality. It follows that $P_i = \ker(L^{n-2i+1}) \cap A_i$, and hence we have the containment

$$\bigoplus_{i=k}^{d} L^{d-i}(P_i) \subseteq \ker(L^{n-k-d+1}) \cap A_d.$$

Since (A, L) is strong Lefschetz it follows that for each i, $L^{d-i}(P_i) \cong P_i$, and hence a simple dimension count reveals this containment must also be equality.

 $(3.) \Rightarrow (2.)$. Assume (3.) holds, and assume that (2.) does not. Fix integer k satisfying $1 \le k \le \lfloor \frac{n}{2} \rfloor$ and suppose that $\alpha \in \ker(L^{n-2k}) \cap A_k$. Then certainly $\alpha \in \ker(A^{n-2k+1}) \cap A_k$ hence $\alpha \in P_k$ by (3.). But also according to (3.) we have

$$A_{n-k} = V(n, 0, n-k) = \bigoplus_{i=0}^{n-k} L^{n-k-i}(P_i).$$

On the other hand if $L^{n-2k}(\alpha) = 0$, then $\dim(L^{n-2k}(P_k)) < P_k$, and hence we must have

$$\dim(A_{n-k}) = \sum_{i=0}^{n-k} \dim(L^{n-k-i}(P_i)) < \sum_{i=0}^{n-k} P_i = \binom{n}{k}$$

which is a contradiction. Therefore (2.) must hold after all.

If any one of the conditions in Lemma 3.6 is satisfied, one can show that $L^{d-i}(P_i) \cong P_i \cong V(n,k,k)$, and hence in this case we get a decomposition of the shifted Specht module V(n,k,d) into irreducible \mathfrak{S}_n -representations.

Corollary 3.7. Let $p = \text{char}(\mathbb{F})$, and assume that p = 0 or $p \ge n+1$. Then the primitive decomposition of the shifted Specht module is a decomposition into irreducible \mathfrak{S}_n -representations:

$$V(n,k,d) \cong \bigoplus_{i=k}^{d} L^{d-i}(P_i) \cong \bigoplus_{i=k} V(n,i,i)[d-i].$$

Here a basis for the irreducible component $L^{d-i}(P_i)$ is

$$\{e_{d-i}(T^c)\cdot F_T\mid T\in \mathrm{Tab}(n,i,i)\}$$

where $e_{d-i}(T^c)$ is the $(d-i)^{th}$ elementary symmetric polynomial in the variables which are not in the support of T.

4 Radical of Shifted Specht Ideals

Our main idea is principal radical systems, based on the following basic facts from commutative algebra:

Lemma 4.1. Let $I \subset R$ be a homogeneous ideal and $x \in R \setminus I$ be any homogeneous polynomial satisfying $(I : x) = (I : x^2)$.

- 1. If (I:x) = I and if I + (x) is radical, then I is radical too.
- 2. If $(I:x) \neq I$ and if (I:x) and I+(x) are both radical, then I is radical too.

Proof. Note that $I \subset I + (x)$ and if I + (x) is radical we also have

$$\sqrt{I} \subset I + (x)$$
.

Hence for $g \in \sqrt{I}$ we can find $a \in I$ and $b \in R$ such that g = a + xb. Since $g \in \sqrt{I}$, there is some integer N for which $g^N \in I$, and we have $g^N = a' + x^N b^N$ for some other $a' \in I$. Therefore $x^N b^N \in I$ and hence $b^N \in (I : x^n) = (I : x)$ and therefore $b \in \sqrt{(I : x)}$. If $(I : x) \neq I$ and (I : x) is radical, then $xb \in I$ and hence $g \in I$, and we are done. If (I : x) = I, then $b \in \sqrt{I}$, and hence we can find $a_1 \in I$ and $b_1 \in R$ for which $b = a_1 + xb_1$. Looking back to g, we have $g = (a + xa_1) + x^2b_1$. Repeating this procedure a number m-times will yield $g = (a + xa_1 + x^2a_2 + \cdots + x^ma_m) + x^{m+1}b_m$, which implies that

$$g \in \bigcap_{m=1}^{\infty} I + (x^m).$$

Since x is homogeneous, it follows that $\bigcap_{m=1}^{\infty} I + (x^m) = I$, and the result follows.

An easy application of principal radical systems is the radical of monomial ideal $(x_1, \ldots, x_n)^{(d)}$.

Lemma 4.2. For each integer d satisfying $1 \le d \le n$ the monomial ideal

$$(x_1,\ldots,x_n)^{(d)}$$

is radical.

Proof. By induction on $n \ge 1$, the base case being trivial. For the inductive step, assume that $(x_1, \ldots, x_{n-1})^{(e)}$ is radical for all $1 \le e \le n-1$, and fix an integer d satisfying $1 \le d \le n$. Set $I = (x_1, \ldots, x_n)^{(d)}$ and $x = x_n$. If d = n then $I = (x_1, \ldots, x_n)$ is principal, and clearly radical, hence we may assume that $1 \le d \le n-1$. Then we have

$$(I:x) = (x_1, \dots, x_{n-1})^{(d-1)}, \text{ and } I + (x) = (x_1, \dots, x_{n-1})^{(d)} + (x_n)$$

which are both radical by the induction hypothesis. Also note that $(I:x^2)=(I:x)$ since I is generated by square-free monomials. It therefore follows from Lemma 4.1 that the ideal $I=(x_1,\ldots,x_n)^{(d)}$ is radical.

Applying principal radical systems to Specht ideals requires the following decomposition of shifted Specht ideals, and is key to the further results of this paper. This is Theorem 1.6 from the Introduction.

Theorem 4.3. For any integers k, d satisfying $0 \le k < d \le n - k$, we have

$$a(n, k, d) = a(n, k, d - 1) \cap (x_1, \dots, x_n)^{(d)}.$$
 (16)

Before embarking on the proof of Theorem 4.3, we will show how to use Theorem 4.3 and principal radical systems to show that shifted Specht ideals are radical. This is Theorem 1.7 from the Introduction.

Theorem 4.4. For any integers k, d satisfying $0 \le k \le d \le n - k$, the shifted Specht ideal a(n, k, d) is radical.

4.1 Proof of Theorem 1.7 or Theorem 4.4

Proof. Assuming that Theorem 4.3 holds, we prove Theorem 4.4 by induction on $n \ge 2$. For the base case n = 2, the only possibilities for integers k, d are k = 0 and d = 1, 2 and k = 1 = d. In the case k = 0 we have $\mathfrak{a}(2,0,d) = (x_1,x_2)^{(d)}$ which is radical by Lemma 4.5. In the case, k = d = 1 we have $\mathfrak{a}(2,1,1) = ((x_1 - x_2))$ is prime, and therefore radical. For the inductive step, assume that $\mathfrak{a}(n-1,j,e)$ for all integers satisfying $0 \le j \le e \le n-1-j$. Fix integers k,d satisfying $1 \le k = d \le n-k$. First we argue for the case d = k. Let $I = \mathfrak{a}(n,k,k)$ and $x = x_n$. Then by Corollary 2.6 we have

$$I + (x) = \alpha(n, k, k) + (x_n) = \alpha(n - 1, k - 1, k) + (x_n).$$

By Theorem 4.3 we have

$$a(n-1, k-1, k) = a(n-1, k-1, k-1) \cap (x_1, \dots, x_{n-1})^{(k)}.$$

By the induction hypothesis a(n-1, k-1, k-1) is radical, and by Lemma 4.2 $(x_1, \dots, x_{n-1})^{(k)}$ is radical. It follows that a(n-1, k-1, k) and hence also I + (x) is also radical.

Also using the change of coordinates map Φ : $x_i \mapsto y_i$ in Corollary 2.6 we find that

$$\Phi(I:x) = (\Phi(I):\Phi(x_n)) = (\mathfrak{a}(n-1,k-1,k):x_n) = \mathfrak{a}(n-1,k-1,k) = \Phi(I)$$

from which it follows that (I : x) = I. Therefore it follows from Lemma 4.1 that $I = \mathfrak{a}(n, k, k)$ itself must be radical.

For d > k we appeal again to Theorem 4.3:

$$a(n, k, d) = a(n, k, d - 1) \cap (x_1, \dots, x_n)^{(d)} = a(n, k, k) \cap (x_1, \dots, x_n)^{(d)}$$

Since a(n, k, k) is radical, and $(x_1, \dots, x_n)^{(d)}$ is radical, it follows that a(n, k, d) is radical too. \square

4.2 Proof of Theorem 1.6 or Theorem 4.3

The proof of Theorem 4.3 (or Theorem 1.6 from the Introduction) comes in three steps, each of which we state as a lemma. First some notation: For any exponent vector $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{N}^n$ we denote the associated monomial by $\mathbf{x}^{\mathbf{a}} = x_1^{a_1} \cdots x_n^{a_n}$, and its radical by $\sqrt{\mathbf{x}^{\mathbf{a}}} = \prod_{a_i > 0} x_i$.

Lemma 4.5. The ideal $\alpha(n,k,d-1) \cap (x_1,\ldots,x_n)^{(d)}$ is generated by products of monomials and polynomials in the shifted Specht module V(n,k,d-1). In fact if the sum of any monomials times forms in V(n,k,d-1) lies in the intersection $\alpha(n,k,d-1) \cap (x_1,\ldots,x_n)^{(d)}$, then so do each of its summands.

Proof. It is not difficult to see that every polynomial $P \in \mathfrak{a}(n, k, d-1)$ decomposes into a sum of terms of the form

$$P = \sum_{\mathbf{a} \in \mathbb{N}^n} \mathbf{x}^{\mathbf{a}} \cdot \nu_{\mathbf{a}}$$

where $v_{\mathbf{a}} \in V(n, k, d-1)$. We want to show that if $P \in (x_1, \dots, x_n)^{(d)}$, then each of its summands are too, i.e. $\mathbf{x}^{\mathbf{a}} \cdot v_{\mathbf{a}} \in (x_1, \dots, x_n)^{(d)}$ for all $\mathbf{a} \in \mathbb{N}^n$. Suppose by way of contradiction that for some $\mathbf{a} \in \mathbb{N}^n$ and some $v_{\mathbf{a}} \in V(n, k, d-1)$ that $\mathbf{x}^{\mathbf{a}} \cdot v_{\mathbf{a}} \notin (x_1, \dots, x_n)^{(d)}$. Since $(x_1, \dots, x_n)^{(d)}$ is a monomial ideal, there must be some monomial $\mathbf{x}^{\mathbf{b}}$ which appears in the monomial expansion of $\mathbf{x}^{\mathbf{a}} \cdot v_{\mathbf{a}}$ such that $\mathbf{x}^{\mathbf{b}} \notin (x_1, \dots, x_n)^{(d)}$. Define the *weight* of monomial $\mathbf{x}^{\mathbf{b}}$ as $\mathrm{wt}(\mathbf{x}^{\mathbf{b}}) = \#\{b_i > 0\}$. Since $(x_1, \dots, x_n)^{(d)}$ consists of all monomials of weight at least d, it follows that $\mathrm{wt}(\mathbf{x}^{\mathbf{b}}) \leq d-1$.

On the other hand, since v_a is a linear combination of shifted Specht polynomials of type $\lambda(d)$, it follows that every monomial in the monomial expansion of v_a also has weight d-1. This implies that $\operatorname{wt}(\mathbf{x}^{\mathbf{b}}) = d-1$, and therefore that

$$\frac{x^b}{\sqrt{x^b}} = x^a.$$

In particular, we see that the monomial $\mathbf{x}^{\mathbf{b}}$ is unique to the term $\mathbf{x}^{\mathbf{a}} \cdot \nu_{\mathbf{a}}$, and hence must occur with the same coefficient in the monomial expansion of $\mathbf{x}^{\mathbf{a}} \cdot \nu_{\mathbf{a}}$ as it does in the monomial expansion of $P = \sum_{\mathbf{a} \in \mathbb{N}^n} \mathbf{x}^{\mathbf{a}} \cdot \nu_{\mathbf{a}}$. Therefore $P \notin (x_1, \dots, x_n)^{(d)}$, as desired.

Lemma 4.5 tells us that it suffices to check equation (16) in Theorem 4.3 on products of monomials with V(n, k, d-1). So we want to show that for each $v \in V(n, k, d-1)$ and for each $\mathbf{a} \in \mathbb{N}^n$ the following implication holds:

$$\mathbf{x}^{\mathbf{a}} \cdot \mathbf{v} \in (x_1, \dots, x_n)^{(d)} \implies \mathbf{x}^{\mathbf{a}} \cdot \mathbf{v} \in \mathfrak{a}(n, k, d).$$

Note that since $(x_1, \ldots, x_n)^{(d)}$ is generated by square-free monomials, we have

$$\mathbf{x}^{\mathbf{a}} \cdot \mathbf{v} \in (x_1, \dots, x_n)^{(d)} \iff \sqrt{\mathbf{x}^{\mathbf{a}}} \cdot \mathbf{v} \in (x_1, \dots, x_n)^{(d)}$$

In particular, we may assume without loss of generality that our monomials $\mathbf{x}^{\mathbf{a}}$ are square-free. For any polynomial $F \in R$ define its *support* to be the set of square-free monomials which divide some non-zero monomial term of F. For example, given a tableau $T \in \text{Tab}(n, k, d)$, the support of the shifted Specht polynomial $F_T(d-1)$ is the set of square-free monomials indexed by subsets of numbers in the support of T, no two of which lie in the same column of T.

Lemma 4.6. For each $T \in \text{Tab}(n, k, d)$, if $\mathbf{x}^{\mathbf{a}} \notin \text{supp}(F_T(d-1))$ then $\mathbf{x}^{\mathbf{a}} \cdot F_T(d-1) \in \mathfrak{a}(n, k, d)$.

Proof. If there is a variable $x_i \in \operatorname{supp}(\mathbf{x}^{\mathbf{a}})$ which is not in $\operatorname{supp}(F_T(d-1))$, then certainly $\mathbf{x}^{\mathbf{a}} \cdot F_T(d-1) \in \mathfrak{a}(n,k,d)$. Otherwise, there must be two indices $i \neq j$ such that $x_i, x_j \in \operatorname{supp}(\mathbf{x}^{\mathbf{a}})$ and i,j in the same column of T. Choose any index $r \neq i,j$ such that $x_r \notin \operatorname{supp}(F_T(d-1))$, which exists since $0 \leq k \leq d-1 < d \leq n-k$, and let $(i,r),(j,r) \in \mathfrak{S}_n$ be the transpositions swapping i,r and j,r, respectively. Then we have

$$F_T(d-1) = F_{(i,r),T}(d-1) + F_{(i,r),T}(d-1)$$

and since $x_i \cdot F_{(i,r),T}(d-1), x_j \cdot F_{(j,r),T}(d-1) \in \mathfrak{a}(n,k,d)$, it follows that

$$\mathbf{x}^{\mathbf{a}} \cdot F_T(d-1) \in \mathfrak{a}(n,k,d)$$

as desired.

Finally we must show what happens with square-free monomials which do lie in the support of the shifted Specht polynomials. Here we use symmetry to make the further reduction that our square-free monomial is initial, i.e.

$$\mathbf{x}^{\mathbf{a}} = \mathbf{x}^{\mathbf{m}} = x_1 \cdots x_m$$
, for some $1 \le m \le d - 1$.

Setting $V^m(n,k,d-1) \subset V(n,k,d-1)$ to be the span of shifted Specht polynomials indexed by standard tableaux $T \in \operatorname{STab}(n,k,d-1)$ for which $\{1,\ldots,m\} \not\subset \operatorname{supp}(T)$. Then the shifted Specht module decomposes into a direct sum

$$V(n, k, d-1) = V_m(n, k, d-1) \oplus V^m(n, k, d-1),$$

and Lemma 4.6 says $\mathbf{x}^{\mathbf{m}} \cdot \mathbf{v} \in \mathfrak{a}(n, k, d)$ for every $\mathbf{v} \in V^m(n, k, d - 1)$. Recall the bijective linear map

$$\phi \colon V(n,k,d-1) \longrightarrow V([n]_m,k-m,d-m)$$

$$F_T(d) \longmapsto F_{T'}(d)$$

where

$$T = \frac{1 \cdots m |_{m+1} \cdots |_{i_k} |_{i_{k+1}} \cdots |_{i_{d-1}}}{|_{i_d} \cdots |_{i_{n-k}} |_{j_1} \cdots |_{j_m} |_{j_{m+1}} \cdots |_{j_k}}$$
(17)

and

$$T' = \frac{i_{m+1} \cdots i_k |i_{k+1}| \cdots |i_{d-1}|}{|i_d| \cdots |i_{n-k}| |j_1| \cdots |j_m| |j_{m+1}| \cdots |j_k|}$$
(18)

Lemma 4.7. Fix $v \in V_m(n, k, d-1)$. If $\mathbf{x}^{\mathbf{m}} \cdot v \in (x_1, \dots, x_n)^{(d)}$ then v = 0, and hence $\mathbf{x}^{\mathbf{m}} \cdot v \in \mathfrak{a}(n, k, d)$.

Proof. We observe that

$$\mathbf{x}^{\mathbf{m}} \cdot (\mathbf{v} - \mathbf{x}^{\mathbf{m}} \cdot \mathbf{v}') \in (x_1, \dots, x_n)^{(d)}$$

where $v' \in V([n]_m, k-m, d-1-m)$ is the image of v in the map above. Indeed note that for each standard tableau $T \in \operatorname{STab}_m(n,k,d-1)$ as in (17), the support of the difference $F_T(d-1) - \mathbf{x^m} \cdot F_{T'}(d-1-m)$ does not contain the monomial $\mathbf{x^m}$, hence the product $\mathbf{x^m} \cdot (F_T(d) - \mathbf{x^m} \cdot F_{T'}(d-1-m)) \in (x_1,\ldots,x_n)^{(d)}$. Hence if $\mathbf{x^m} \cdot v \in (x_1,\ldots,x_n)^{(d)}$, it follows that $(\mathbf{x^m})^2 \cdot v' \in (x_1,\ldots,x_n)^{(d)}$, and hence that $v' \in ((x_1,\ldots,x_n)^{(d)} : \mathbf{x^m}) = (x_{m+1},\ldots,x_n)^{(d-m)}$. For degree reasons this implies that v' = 0, and hence by Corollary 2.7, v = 0 as well.

Proof of Theorem 4.3. The containment $a(n,k,d) \subset a(n,k,d-1) \cap (x_1,\ldots,x_n)^{(d)}$ is clear. For the reverse containment, Lemma 4.5 implies that it suffices to check it on products of monomials with polynomials in V(n,k,d-1). Since $(x_1,\ldots,x_n)^{(d)}$ is generated by square-free monomials we may assume that our monomials are square-free, and by symmetry we may assume that our monomial has the form $\mathbf{x}^{\mathbf{m}} = x_1 \cdots x_m$ for some integer $1 \le m \le d$. Then as above we have

$$V(n, k, d - 1) = V_m(n, k, d - 1) \oplus V^m(n, k, d - 1)$$

and by Lemma 4.7, $\mathbf{x}^{\mathbf{m}} \cdot \mathbf{v} \in \mathfrak{a}(n,k,d)$ if $\mathbf{v} \in V_m(n,k,d-1)$. Furthermore, Lemma 4.6 implies that for $\mathbf{v} \in V^m(n,k,d-1)$ if $\mathbf{x}^{\mathbf{m}} \cdot \mathbf{v} \in (x_1,\ldots,x_n)^{(d)}$ then $\mathbf{x}^{\mathbf{m}} \cdot \mathbf{v} \in \mathfrak{a}(n,k,d)$, and the result follows.

5 Perfection of Specht and Specht-Monomial Ideals

Recall that an ideal $I \subset R$ in a commutative ring has projective or homological dimension s if a minimal resolution of R/I as an R-module has length s. Its grade is the length g of a maximal R-sequence contained in I, or equivalently the smallest integer g for which the Ext group $\operatorname{Ext}_R^g(R/I,R)$ is non-zero. We say that the ideal $I \subset R$ is *perfect* if its projective dimension is equal to its grade. In our case, where R is polynomial (hence Cohen-Macaulay), the grade of an ideal is equal to its height, and by the Auslander-Buchsbaum formula, I is perfect if and only if the quotient R/I is Cohen-Macaulay. For more details on these matters we refer the reader to [8].

As in Section 4, the main idea here is to use principal radical systems.

Lemma 5.1. Let $I \subset R$ be any homogeneous ideal and let $x \in R \setminus I$ be any homogeneous polynomial. Then

- 1. if (I:x) = I and if I + (x) is perfect, then I is perfect too.
- 2. if $(I:x) \neq I$ and if (I:x) and I+(x) are both perfect of the same grade g, then I is perfect of that same grade too.

Proof. Item (1.) is well known, and can be found in any commutative algebra text, e.g. [8, Theorem 17.3]. For (2.) we use the long exact sequence for Ext-modules associated with the short exact sequence of R-modules

$$0 \longrightarrow R/(I:x) \xrightarrow{\times x} R/I \longrightarrow R/I + (x) \longrightarrow 0.$$

As in Section 4, we give an easy application of principal radical systems.

Lemma 5.2. For every integer d satisfying $1 \le d \le n$ the monomial ideal

$$(x_1,\ldots,x_n)^{(d)}$$

is perfect.

Proof. By induction on n, where the base case n = 1 is trivial. For the induction step, assume that $(x_1, \ldots, x_{n-1})^{(e)}$ is perfect for every $1 \le e \le n-1$, and fix an integer d satisfying $1 \le d \le n$. Set $I = (x_1, \ldots, x_n)^{(d)}$ and $x = x_n$. Then we have

$$(I:x) = (x_1, \dots, x_{n-1})^{(d-1)}$$
 $I + (x) = (x_1, \dots, x_{n-1})^{(d)} + (x_n)$

which are both perfect by our induction hypothesis. It follows from Lemma 5.1 that $I = (x_1, \dots, x_n)^{(d)}$ is perfect too.²

It is trickier to apply Lemma 5.1 to shifted Specht ideals. For one thing, not all shifted Specht ideals are perfect. Indeed, Theorem 4.3 gives the decomposition

$$\mathfrak{a}(n,k,d) = \mathfrak{a}(n,k,k) \cap (x_1,\ldots,x_n)^{(d)}.$$

Individually the grades (=heights) of the ideals a(n, k, k) and $(x_1, \ldots, x_n)^{(d)}$ are g = n - k and g = n - d + 1, respectively. In particular we see that if d > k + 1, then the two ideals a(n, k, k) and $(x_1, \ldots, x_n)^{(d)}$ have mixed heights, and in particular the shifted Specht ideal a(n, k, d) cannot be perfect. For d = k + 1, we must show that the intersection is perfect:

$$\alpha(n,k,k+1) = \alpha(n,k,k) \cap (x_1,\ldots,x_n)^{(d)}.$$

To this end, we appeal to the following basic fact, which appears in the paper [5] by Hochster-Eagon, and we refer the reader there for its proof.

Lemma 5.3 (Proposition 18, [5]). Suppose that $I, J \subset R$ are two perfect ideals of the same grade g, and assume that I + J has grade g + 1. Then I + J is perfect if and only if $I \cap J$ is perfect.

²Alternatively, one could also appeal to Reisner's theorem [11] here, since $(x_1, \ldots, x_n)^{(d)}$ is the Stanley-Reisner ideal of the d-1 skeleton of an n-1 simplex.

In the spirit of Lemma 5.3 we study the following sum of ideals, which plays a key role in this paper.

Definition 5.4. Fix an integer k satisfying $1 \le k < k + 1 \le n - k$, and define the *Spechtmonomial ideal* for the pair (k, n) to be the sum of ideals

$$I(n,k) = a(n,k,k) + (x_1, \dots, x_n)^{(k+1)}.$$
 (19)

The next result is key to our results in this section, and it plays a similar role as Theorem 4.3 in Section 4. Note however that this result depends on the characteristic of \mathbb{F} . It is Theorem 1.8(3.) from the Introduction.

Theorem 5.5. Let $p = \text{char}(\mathbb{F}) \ge 0$ and fix positive integers n, k satisfying $n \ge 2k + 1$. Then the following conditions are equivalent.

- 1. p = 0 or $p \ge k + 1$.
- 2. The Specht monomial ideal I(n, k) satisfies

$$I(n,k) = I(n-1,k-1) \cap \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \right)$$

$$where \quad y_i = \begin{cases} x_n - x_i & \text{if } 1 \le i \le n-1 \\ x_n & \text{if } i = n \end{cases}$$
(20)

Before embarking on the proof of Theorem 5.5, we will use it in conjunction with principal radical systems to prove Theorems 1.8(4.) and also Theorem 1.2 from the Introduction.

Theorem 5.6. Let $p = \text{char}(\mathbb{F}) \ge 0$ and fix positive integers n, k satisfying $n \ge 2k + 1$. Then the following conditions are equivalent.

- 1. p = 0 or $p \ge k + 1$.
- 2. The Specht-monomial ideal I(n, k) is perfect.

Theorem 5.7. Let $p = \text{char}(\mathbb{F}) \ge 0$ and fix positive integers n, k satisfying $n \ge 2k + 1$.

- 1. If p = 0 or $p \ge k + 1$, then the Specht ideal a(n + 1, k + 1, k + 1) is perfect.
- 2. If $n \ge 2p + 1$, then the Specht ideal a(n + 1, p + 1, p + 1) is not perfect.

The proof for Theorem 5.6 occupies the next subsection, followed by the proof of Theorem 5.7, and we save the proof of Theorem 5.5 for the end.

5.1 Proof of Theorem 1.8(4.) or Theorem 5.6

We will focus on the implication (1.) \Rightarrow (2.) first. Assuming that p = 0 or $p \ge k + 1$, Theorem 5.5 says that for all $1 \le k < k + 1 \le n - k$ the Specht-monomial ideal satisfies (20):

$$I(n,k) = I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)} + (x_n^2)).$$

We need the following Lemma, which tells us how to go between x-variables and y-variables, and provides a direct link to Lefschetz properties from Section 3.

Lemma 5.8. Let $P^j(x_1, ..., x_{n-1})$ be any square-free polynomial of degree j in the variables $x_1, ..., x_{n-1}$. Then modulo the principal ideal (x_n^2) we have

$$P^{j}(x_{1},...,x_{n-1}) \equiv (-1)^{j} \left(P^{j}(y_{1},...,y_{n-1}) - x_{n} \cdot D \left(P^{j}(y_{1},...,y_{n-1}) \right) \right) \mod (x_{n}^{2})$$

where D is the linear partial differentiation operator $D = \frac{\partial}{\partial y_1} + \dots + \frac{\partial}{\partial y_{n-1}}$. In particular, if $\alpha = \alpha(x_1, \dots, x_{n-1}) \in V(n-1, k-1, k-1)$ is a linear combination of Specht polynomials then

$$\alpha(x_1, \dots, x_{n-1}) = (-1)^{k-1} \cdot \alpha(y_1, \dots, y_{n-1}).$$

Proof. By linearity of D it suffices to assume that P^j is a square-free monomial, and by symmetry we may assume is $P^j(x_1, \ldots, x_{n-1}) = x_1 \cdots x_j$. Then we have

$$P^{j}(x_{1},...,x_{n-1}) = x_{1} \cdots x_{j} = (x_{n} - y_{1}) \cdots (x_{n} - y_{j})$$

$$= x_{n}^{2} \cdot (\text{stuff}) + (-1)^{j-1} \cdot x_{n} \left(\sum_{i=1}^{j} y_{1} \cdots \hat{y}_{i} \cdots y_{j} \right) + (-1)^{j} \cdot y_{1} \cdots y_{j}$$
(where \hat{y} means omission)
$$= (-1)^{j} \left(y_{1} \cdots y_{j} - x_{n} D \left(y_{1} \cdots y_{j} \right) \right) + x_{n}^{2} \cdot (\text{stuff})$$

$$= (-1)^{j} \left(P^{j}(y_{1},...,y_{n-1}) - x_{n} D \left(P^{j}(y_{1},...,y_{n-1}) \right) \right) \quad \text{mod } (x_{n}^{2})$$

as claimed. The second statement follows from the first since $D(\alpha) = 0$.

Next, for each pair of positive integers n, k satisfying $n \ge 2k + 1$ let us form the new ideal

$$J(n,k) = I(n-1,k-1) + (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$
(21)

Lemma 5.9. Assume that p = 0 or $p \ge k + 1$. Then the ideal J(n, k) satisfies

$$J(n,k) + (x_n) = I(n-1,k-1) + (x_n), \text{ and } (J(n,k):x_n) = (x_1,\ldots,x_{n-1})^{(k-1)} + (x_n).$$

Proof. We have

$$J(n,k) + (x_n) = I(n-1,k-1) + (y_1, \dots, y_{n-1})^{(k)} + (x_n^2) + (x_n)$$

$$= \alpha(n-1,k-1,k-1) + (x_1, \dots, x_{n-1})^{(k)} + (y_1, \dots, y_{n-1})^{(k)} + (x_n)$$

$$= \alpha(n-1,k-1,k-1) + (x_1, \dots, x_{n-1})^{(k)} + (x_n)$$

$$= I(n-1,k-1) + (x_n).$$

For the other equality, note first that containment $(J(n,k):x_n)\supseteq (x_1,\ldots,x_{n-1})^{(k-1)}+(x_n)=(y_1,\ldots,y_{n-1})^{(k-1)}+(x_n)$ follows from Lemma 5.8. Indeed identifying the space of square-free polynomials with the monomial complete intersection $B=\mathbb{F}[x_1,\ldots,x_{n-1}]/(y_1^2,\ldots,y_{n-1}^2)$, Lemma 3.4 implies that the derivative map $D\colon B_k\to B_{k-1}$ is surjective, and hence for any square-free monomial of degree k-1 in variables y_1,\ldots,y_{n-1} , say $P=y_1\cdots y_{k-1}$, we know by Lemma 3.4 there is square-free polynomial of degree k for which

$$D(Q(y_1,\ldots,y_{n-1})) = P(y_1,\ldots,y_{n-1}) = y_1\cdots y_{k-1}.$$

Then Lemma 5.8 implies that

$$P \cdot x_n = y_1 \cdots y_{k-1} \cdot x_n = x_n \cdot D(Q(y_1, \dots, y_{n-1}))$$

= $Q(x_1, \dots, x_{n-1}) \pm Q(y_1, \dots, y_{n-1})$ mod (x_n^2)

Since $Q(x_1, \ldots, x_{n-1}) \in I(n-1, k-1)$ and $Q(y_1, \ldots, y_{n-1}) \in (y_1, \ldots, y_{n-1})^{(k)}$, it follows that $x_n \cdot P = x_n \cdot D(Q(y_1, \ldots, y_{n-1})) \in J(n, k)$, and hence $P = y_1 \cdots y_{k-1} \in (J(n, k) : x_n)$. For the reverse containment, fix $G \in (J(n, k) : x_n)$. Then for each $S \in STab(n-1, 0, k)$ there exists polynomials $d_S \in R$ for which

$$x_nG - \sum_{S \in \text{STab}(n-1,0,k)} d_S M_S \in I(n-1,k-1) + (x_n^2)$$

where $M_S = M_S(y_1, \dots, y_{n-1}) \in (y_1, \dots, y_{n-1})^{(k)}$ are square-free monomials of degree k in the y-variables. By Lemma 5.8 we have for each $S \in STab(n-1, 0, k)$

$$M_S(y_1, \dots, y_{n-1}) \equiv \pm (M_S(x_1, \dots, x_{n-1}) - x_n D(M_S(x_1, \dots, x_{n-1})))$$
 mod (x_n^2)

and since $M_S(x_1, ..., x_{n-1}) \in (x_1, ..., x_{n-1})^{(k)} \subset I(n-1, k-1)$, we see that

$$x_nG - \sum_{S \in \text{STab}(n-1,0,k)} d_S M_S(y_1, \dots, y_{n-1}) \equiv x_nG - \sum_{S \in \text{STab}(n-1,0,k)} d_S x_n D(M_S(y_1, \dots, y_{n-1}))$$

$$\equiv 0 \mod I(n-1,k-1) + (x_n^2)$$

Since x_n is a non-zero divisor for I(n-1, k-1), it follows that

$$G - \sum_{S \in \text{STab}(n-1,0,k)} d_S D(M_S(x_1,\ldots,x_{n-1})) \in I(n-1,k-1) + (x_n) \subset (x_1,\ldots,x_{n-1})^{(k-1)} + (x_n)$$

and since $D(M_S(x_1,...,x_{n-1})) \in (x_1,...,x_{n-1})^{(k-1)}$ for all $S \in STab(n-1,0,k)$, we see also that $G \in (x_1,...,x_{n-1})^{(k-1)} + (x_n)$, as desired.

We are now in a position to prove implication $(1.) \Rightarrow (2.)$ in Theorem 5.6.

Proof of (1.) ⇒ (2.) in Theorem 5.6. Assume that p = 0 or $p \ge k + 1$. We prove by induction on $n \ge 3$ for each integer k satisfying $1 \le k < k + 1 \le n - k$ then the ideal I(n, k) is perfect. The base case is n = 3 where the only possible k value is k = 1. Here the assumption on p is vacuous, and we have

$$I(3,1) = \mathfrak{a}(3,1,1) + (x_1, x_2, x_3)^{(2)} = (x_1 - x_3, x_2 - x_3, x_1 x_2, x_1 x_3, x_2 x_3).$$

Note that $\mathbb{F}[x_1, x_2, x_3]/I(3, 1) \cong \mathbb{F}[z]/(z^2)$ is Cohen-Macaulay, which implies that I(3, 1) is perfect of grade n - k + 1 = 3.

For the inductive step, assume that I(n-1, j) is perfect for all integers j satisfying $1 \le j < j + 1 \le n - 1 - j$. Fix k satisfying $1 \le k < k + 1 \le n - k$, and consider the Specht-monomial ideal

$$I(n,k) = \alpha(n,k,k) + (x_1,\ldots,x_n)^{(k+1)}$$
.

Consider the sum J(n, k) as in (21), i.e.

$$J(n,k) = I(n-1,k-1) + (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

Note that I(n-1,k-1) is perfect by the induction hypothesis. Also $(y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$ is perfect since $(y_1,\ldots,y_{n-1})^{(k)}$ is perfect (by Lemma 5.2), and x_n^2 is $(y_1,\ldots,y_{n-1})^{(k)}$ -regular. Therefore by Lemma 5.3, I(n,k) is perfect if and only if J(n,k) is perfect. But according to Lemma 5.9 $J(n,k)+(x_n)=I(n-1,k-1)+(x_n)$ and $(J(n,k):x_n)=(x_1,\ldots,x_{n-1})^{(k-1)}+(x_n)$ which are both perfect of the same grade g=n-k+2, and hence by Lemma 5.2, it follows that J(n,k) is also perfect (of grade g=n-k+2). Thus it follows that I(n,k) is perfect completing the induction step, and hence the proof.

Next we prove the reverse implication (2.) \Rightarrow (1.) in Theorem 5.6 which also requires a bit of a set up. First note that Theorem 5.5 implies that if p = 0 or $p \ge k + 1$, then I(n, k) has the following irredundant primary decomposition:

$$I(n,k) = \mathfrak{a}(n,k,k) + (x_1, \dots, x_n)^{(k+1)} = \bigcap_{\sigma \in \mathfrak{S}_n} \underbrace{\sigma. \left(x_1 - x_2, \dots, x_1 - x_{n-k+1}, x_1^2 \right)}_{O_{\sigma}}$$
(22)

Note that for $\sigma = e$, we have $Q_e = \mathfrak{a}(n-k+1,1,1) + (x_1,\ldots,x_{n-k+1})^{(2)}$; in particular, Q_e contains all quadratic forms in the variables x_1,\ldots,x_{n-k+1} .

Lemma 5.10. Over any field \mathbb{F} , and for any positive integers n, k satisfying $n \ge 2k+1$, if I(n, k) is perfect then I(n, k) must satisfy the decomposition (22).

Proof. Assume that I(n, k) is perfect, and consider the intersection of primary ideals as in (22), i.e.

$$I'(n,k) = \bigcap_{\sigma \in \mathfrak{S}_n} Q_{\sigma}$$

with minimal associated prime divisors given by

$$P_{\sigma} = \sqrt{Q_{\sigma}} = \sigma.(x_1, \ldots, x_{n-k+1}).$$

We would like to show that I(n, k) = I'(n, k). Note first that they have the same radical:

$$\sqrt{I'(n,k)} = \bigcap_{\sigma \in \mathfrak{S}_n} P_{\sigma} = (x_1, \dots, x_n)^{(k)} = \sqrt{I(n,k)}.$$
 (23)

For the last equality, the containment $I(n, k) \subseteq (x_1, \dots, x_n)^{(k)}$ is clear, and the other containment follows from the containment:

$$x_{i_1}^2 \cdots x_{i_k}^2 = x_{i_1} \cdots x_{i_k} \cdot F_T + \left(\text{stuff in } (x_1, \dots, x_n)^{(k+1)} \right) \in I(n, k)$$

where

$$T = \frac{\begin{vmatrix} i_1 & \cdots & i_k \\ i_{k+1} & \cdots & i_{n-k} & j_1 & \cdots & j_k \end{vmatrix}}{\begin{vmatrix} i_1 & \cdots & i_k \\ \vdots & \vdots & \vdots & \vdots \end{vmatrix}}$$

It follows that $\{P_{\sigma} \mid \sigma \in \mathfrak{S}_n\}$ is a complete list of minimal prime divisors of I(n,k), which implies equation (23).

Set I' = I'(n, k) and I = I(n, k). Suppose that a primary decomposition of I is given by $I = U_1 \cap \cdots \cap U_m$. Since I is perfect, all of its associated prime divisors must be minimal, and hence the primary components must be indexed by the symmetric group and we can write

$$I = \bigcap_{\sigma \in \mathfrak{S}_n} U_{\sigma}$$

where $\sqrt{U_{\sigma}} = P_{\sigma}$. Fix $\sigma \in \mathfrak{S}_n$, and set $U = U_{\sigma}$, $P = P_{\sigma}$, and $Q = Q_{\sigma}$. We want to show that $U_{\sigma} = Q_{\sigma}$. Let R_P be the polynomial ring $R = \mathbb{F}[x_1, \ldots, x_n]$ localized at the prime ideal P. By a theorem of Nagata [10, Theorem 8.7] we have

$$U = IR_P \cap R$$
 and $Q = JR_P \cap R$.

We will prove that in the local ring R_P , the ideals are equal $IR_P = JR_P$. Certainly because of the containment $I \subseteq J$, we have also $IR_P \subseteq JR_P$. In the other direction we observe that

 $JR_P = QR_P$. Without loss of generality we may assume that $\sigma = e$, and $Q = (x_1 - x_2, \dots, x_1 - x_{n-k+1}, x_1^2)$. For each pair $1 \le r < s \le n-k+1$ we may choose $1 \le i_1 < \dots < i_{k-1} \le n-k+1$ such that $r, s \notin \{i_1, \dots, i_{k-1}\}$ and setting $j_i = n-k+1+i$ define the tableau

$$T = \frac{i_1 \cdots i_{k-1} r}{i_{k+1} \cdots i_{n-k} j_1 \cdots j_{k-1} s}$$

Then in the local ring R_P the Specht polynomial $F_T \in I$ has the form $F_T = u \cdot (x_r - x_s)$ where $u \in R_P$ is a unit. Also consider the monomial

$$M_S = x_{n-k+2} \cdots x_n \cdot x_r \cdot x_s \in (x_1, \dots, x_n)^{(k)} \subset I.$$

Then in the local ring R_P it has the form $M = w \cdot x_r \cdot x_s$ where $w \in R_P$ is a unit. Therefore the generators of QR_P satisfy $(x_r - x_s) \in IR_P$ and $x_r \cdot x_s \in IR_P$ it follows that $QR_P \subset IR_P$ and hence that $JR_P \subset IR_P$, as desired.

In particular Lemma 5.10 implies that if I(n, k) is not perfect then it must have an embedded prime divisor. The upshot here is that embedded prime divisors seem to be an easier check for imperfection than, say, extraneous syzygies in a minimal free resolution. Note that Theorem 4.4 implies that the Specht ideals a(n + 1, k + 1, k + 1) never have embedded prime divisors in any characteristic, and this is one of the main reasons for the disparity between Theorems 5.6 and 5.7. We are now in a position to prove the remainder of Theorem 5.6.

Proof of (2.) \Rightarrow (1.) in Theorem 5.6. Assume that p satisfies 0 . We will show that <math>I(n, k) is not perfect by showing it does not satisfy Decomposition (22), or equivalently that

$$\underbrace{\alpha(n,k,k) + (x_1, \dots, x_n)^{(k+1)}}_{I(n,k)} \neq \underbrace{\alpha(n-1,k-1,k-1) + (x_1, \dots, x_{n-1})^{(k)}}_{I(n-1,k-1)} \cap \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \right). \tag{24}$$

We identify the space of square-free x-monomials with the monomial complete intersection $B = \mathbb{F}[x_1,\ldots,x_{n-1}]/(x_1^2,\ldots,x_{n-1}^2)$ and $D = D_B = \frac{\partial}{\partial x_1} + \cdots + \frac{\partial}{\partial x_{n-1}}$ the associated lowering operator. From our assumptions on p, Lemma 3.4 implies that the derivative map $D \colon B_k \to B_{k-1}$ is not surjective, and hence (by Lemma 3.2), $V(n-1,k,k) = \ker(D) \cap B_k$. In particular, there must exist a square-free polynomial of degree k, say $f = f(x_1,\ldots,x_{n-1}) \in (x_1,\ldots,x_{n-1})^{(k)}$ with the property that D(f) = 0 but $f \notin V(n-1,k,k)$. Since D(f) = 0, we deduce from Lemma 5.8 that

$$f(x_1, \dots, x_{n-1}) \equiv f(y_1, \dots, y_{n-1}) \equiv 0 \mod (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

Therefore, $f \in I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)}+(x_n^2))$. On the other hand, since $f \notin V(n-1,k,k)$ it follows from Lemma 3.5 that $f \notin V(n,k,k)$ either, and it follows that $f \notin I(n,k)$. This shows that Inequality (24) holds, and hence by Lemma 5.10, the Specht-monomial ideal I(n,k) is not perfect.

5.2 Proof of Theorem 1.2 or Theorem 5.7

Proof. First assume that p = 0 or $p \ge k + 1$. We prove by induction on $n \ge 3$ that for each integer k satisfying $1 \le k < k + 1 \le n - k$, and each i satisfying $1 \le i \le k$ the Specht ideal

a(n+1, i+1, i+1) is perfect. First recall that Corollary 2.6 says the change of coordinates map Φ gives a ring isomorphism

$$a(n+1, i+1, i+1) \cong a(n, i, i+1) = a(n, i, i) \cap (x_1, \dots, x_n)^{(i+1)}$$

where the second equality follows from Theorem 1.6 or Theorem 4.3.

For the base case n = 3 and the only possible k = 1 gives

$$\mathfrak{a}(4,2,2) \cong \mathfrak{a}(3,1,2) = \mathfrak{a}(3,1,1) \cap (x_1,x_2,x_3)^{(2)}.$$

Note that a(3, 1, 1) and $(x_1, x_2, x_3)^{(2)}$ are both perfect of grade g = 2 Also note that the Spechtmonomial ideal $I(3, 1) = a(3, 1, 1) + (x_1, x_2, x_3)^{(2)}$ has grade g + 1 = 3 and its quotient

$$\frac{\mathbb{F}[x_1, x_2, x_3]}{I(3, 1)} \cong \frac{\mathbb{F}[z]}{(z^2)}$$

is Cohen-Macaulay. Therefore I(3, 1) is perfect and hence by Lemma 5.3, so is $\mathfrak{a}(4, 2, 2)$. For the inductive step, assume that $\mathfrak{a}(n, i, i)$ is perfect for every $1 \le i \le k$. We have

$$\mathfrak{a}(n+1, i+1, i+1) \cong \mathfrak{a}(n, i, i) \cap (x_1, \dots, x_n)^{(i+1)}$$
.

Also note that since $1 \le i < i + 1 \le k + 1 \le n - k \le n - i$, Theorem 5.6 implies that the Specht monomial ideal

$$I(n, i) = \alpha(n, i, i) + (x_1, \dots, x_n)^{(i+1)}$$

is perfect of grade g+1=n-i+1. Since $\mathfrak{a}(n,i,i)$ and $(x_1,\ldots,x_n)^{(i+1)}$ are both perfect of grade g=n-i. Then it follows from Lemma 5.3 that $\mathfrak{a}(n+1,i+1,i+1)\cong\mathfrak{a}(n,i,i)\cap(x_1,\ldots,x_n)^{(i+1)}$ is also perfect. This proves implication (1.).

For (2.), assume that $n \ge 2p+1$. By (1.), we see that a(n, p, p) is perfect. Also $(x_1, \dots, x_n)^{(p)}$ is perfect by Lemma 5.2. Therefore by Lemma 5.3, it follows that the Specht ideal a(n+1, p+1, p+1) and the Specht-monomial ideal I(n, p) are perfect or not, alike. But Theorem 5.6 implies that the Specht-monomial ideal

$$I(n, p) = a(n, p, p) + (x_1, \dots, x_n)^{(p+1)}$$

is not perfect, and hence the Specht monomial ideal a(n+1, p+1, p+1) is not perfect either. \Box

As stated in the Introduction, we conjecture that Theorem 1.2(2.) can be improved.

Conjecture 5.11. If $p = \text{char}(\mathbb{F})$ and n, k are positive integers satisfying $0 and <math>n \ge 2k+1$, then the Specht ideal $\mathfrak{a}(n+1,k+1,k+1)$ is not perfect.

Example 5.12. Taking p=2, Theorem 1.2(2.) says that a(n+1,3,3) is not perfect for every $n \ge 5$, a result also obtained by Yanagawa [15, Theorem 5.3]. For example if n=5, then a(6,3,3) is not perfect, and one obstruction to perfection is the elementary symmetric polynomial $\alpha=e_2(x_1,\ldots,x_4)$ which lies in the intersection

$$e_2(x_1,\ldots,x_4) \in \mathfrak{a}(4,1,1) \cap \left((y_1,\ldots,y_4)^{(2)} + (x_5^2) \right) = \bigcap_{\sigma \in \mathfrak{S}_5} \sigma.(x_1-x_2,x_1-x_3,x_1-x_4,x_1^2)$$

but $e_2(x_1, ..., x_4) \notin I(5, 2)$. This indicates that the Specht-monomial ideal I(5, 2) must have an embedded prime divisor in characteristic p = 3, which does not appear in higher characteristics. Macaulay2 reveals that the primary decomposition of I(5, 2) over the field $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$ is

$$I(5,2) = \bigcap_{\sigma \in \mathfrak{S}_5} \sigma.(x_1 - x_2, x_1 - x_3, x_1 - x_4, x_1^2) \cap Q$$

where Q is primary and satisfies $(x_1^2, \ldots, x_5^2) \subseteq Q$; in particular the maximal ideal $\mathfrak{m} = (x_1, \ldots, x_5)$ is an associated prime divisor of the Specht-monomial ideal $I(5,2) = \mathfrak{a}(5,2,2) + (x_1, \ldots, x_5)^{(3)}$ in characteristic p = 2. Since $\mathfrak{a}(5,2,2)$ and $(x_1, \ldots, x_5)^{(3)}$ are both perfect of grade g = 3 in characteristic p = 2 (and all characteristics), we deduce that $\mathfrak{a}(6,3,3) \cong \mathfrak{a}(5,2,3) = \mathfrak{a}(5,2,2) \cap (x_1,\ldots,x_5)^{(3)}$ is also not perfect by Lemma 5.3.

Similarly, for p = 3, Theorem 1.2(2.) implies that a(n + 1, 4, 4) is not perfect for $n \ge 7$ in characteristic p = 3. For example if n = 7 then a(8, 4, 4) is not perfect, and, as in the previous case, one can see that the Specht-monomial ideal I(7, 3) has an extra primary component that contains all the squared variables. Conjecture 5.11 would imply that, for example, a(n + 1, 5, 5) is not perfect for all $n \ge 9$ in characteristic p = 3. Computations in Macaulay2 show that a(10, 5, 5) is indeed not perfect, supporting this claim.

5.3 Proof of Theorem 1.8(3.) or Theorem 5.5

The proof of Theorem 5.5 is surprisingly similar to that of Theorem 4.3, and it too comes by way of several lemmas. We want to show that for every $1 \le k < k + 1 \le n - k$ the Spechtmonomial ideal $I(n,k) = \mathfrak{a}(n,k,k) + (x_1,\ldots,x_n)^{(k+1)}$ satisfies

$$I(n,k) = I(n-1,k-1) \cap \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \right).$$

Of course there is an assumption about characteristic here, but we will not assume it yet, and try to point out exactly where we need it.

Lemma 5.13. *Assume that* $1 \le k < k + 1 \le n - k$. *Then*

$$I(n,k) \subseteq I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)} + (x_n^2)).$$

Proof. Certainly we have that $a(n,k,k) \subset a(n-1,k,k) \subset a(n-1,k-1,k-1)$, and also $(x_1,\ldots x_n)^{(k+1)} \subset (x_1,\ldots ,x_{n-1})^{(k)}$, hence $I(n,k) \subset I(n-1,k-1)$ (if k=1, we should regard I(n-1,0) as R). It remains to see why $I(n,k) \subset (y_1,\ldots ,y_{n-1})^{(k)} + (x_n^2)$. For $T \in STab(n,k,k)$ we have

$$F_T^k(x_1,\ldots,x_n)=(x_{i_1}-x_{j_1})\cdots(x_{i_k}-x_n)=(y_{j_1}-y_{i_1})\cdots(y_{j_{k-1}}-y_{i_{k-1}})\cdot(-y_i)\in(y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$$

For $S \in \text{STab}(n, 0, k+1)$, if $x_n \in \text{supp}(M_S^{k+1})$ then $M_S^{k+1} = x_n \cdot M_S^k$, for $S' \in \text{STab}(n-1, 0, k)$ and by Lemma 5.8 we have

$$M_{S'}^k(x_1,\ldots,x_{n-1})=M_{S'}^k(y_1,\ldots,y_{n-1})+x_n\cdot D(M_{S'}^k(y_1,\ldots,y_{n-1})) \mod (y_1,\ldots,y_{n-1})^{(k)}.$$

It follows that $M_S^{k+1} = x_n \cdot M_S^k$, $\in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. If $x_n \notin \text{supp}(M_S^{k+1})$, then it is obvious that $M_S^{k+1} \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. Hence $I(n, k) \subseteq I(n-1, k-1) \cap ((y_1, \dots, y_{n-1})^{(k)} + (x_n^2))$, as desired.

We have added a superscript to our notation for the shifted Specht polynomials to help the reader remember degrees. The other containment

$$I(n,k) \supseteq I(n-1,k-1) \cap \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \right)$$
 (25)

is harder to prove.

Note that the ideal I(n-1,k-1) is generated in degrees k-1 and k by the following subspaces of forms:

$$V = V(n-1, k-1, k-1) = \left\langle F_T^{k-1} \mid T \in STab(n-1, k-1, k-1) \right\rangle$$

$$U = U(n-1, k-1, k) = \left\langle x_n F_T^{k-1} \mid T \in STab(n-1, k-1, k-1) \right\rangle + \left\langle M_S^k \mid S \in STab(n-1, 0, k) \right\rangle$$

$$= x_n \cdot V(n-1, k-1, k-1) + V(n-1, 0, k)$$

We make some preliminary observations, but first some notation. Denote by $\mathbb{N}^n(m)$ the set of exponent vectors of degree, i.e. $\mathbf{a}=(a_1,\ldots,a_n)$ with $a_1+\cdots+a_n=m$. A monomial in the y-variables (resp. the x-variables) will be denoted by $\mathbf{y}^{\mathbf{a}}=y_1^{a_1}\cdots y_n^{a_n}$ (resp. $\mathbf{x}^{\mathbf{a}}=x_1^{a_1}\cdots x_n^{a_n}$), and its radical is the square-free monomial $\sqrt{\mathbf{y}^{\mathbf{a}}}=\prod_{a_i>0}y_i$. We also define the weight of a monomial to be wt($\mathbf{y}^{\mathbf{a}}$) = $\#\{a_i>0\}$, the number of non-zero entries in its exponent vector.

Lemma 5.14. With U and V as above, we have

- 1. $x_n \cdot V \subseteq U$
- 2. $x_n \cdot U \subseteq I(n,k)$, and
- 3. for every $P \in I(n-1, k-1)$, and for all exponent vectors $\mathbf{a} \in \mathbb{N}^{n-1}(m)$ and $\mathbf{b} \in \mathbb{N}^{n-1}(m-1)$, there exists elements $v_{\mathbf{a}} \in V$ and $\mu_{\mathbf{b}} \in U$ such that

$$P \equiv \sum_{\mathbf{a} \in \mathbb{N}^{n-1}(m)} \mathbf{y}^{\mathbf{a}} \nu_{\mathbf{a}} + \sum_{\mathbf{b} \in \mathbb{N}^{n-1}(m-1)} \mathbf{y}^{\mathbf{b}} \mu_{\mathbf{b}} \qquad mod \ I(n, k).$$
 (26)

Proof. (1.) is obvious from the definitions. For (2.), note that for $S \in \text{STab}(n-1,0,k-1)$, $x_n \cdot M_S^k \in V(n,0,k+1) \subset I(n,k)$. Also for $T \in \text{STab}(n-1,k-1,k-1)$ and for index $1 \le i \le n-1$ such that $i \notin \text{supp}(T)$, which exists because $n-1 \ge 2k > 2(k-1)$, we have

$$x_n^2 \cdot F_T^{k-1} = x_n(x_n - x_i) \cdot F_T^{k-1} + x_n x_i \cdot F_T^{k-1}$$

and since $(x_n - x_i) \cdot F_T^{k-1} \in V(n, k, k)$ and $x_n x_i \cdot F_T^{k-1} \in V(n, 0, k+1)$, it follows that $x_n^2 \cdot F_T^{k-1} \in I(n, k)$, and (2.) follows. Finally fix a homogeneous polynomial $P \in I(n-1, k-1)$. Then for each $T \in \operatorname{STab}(n-1, k-1, k-1)$ and each $S \in \operatorname{STab}(n-1, 0, k)$, there exist polynomials, which we may take in the y-variables, $g_T(y)$ of degree m and $h_S(y)$ of degree m-1 for which

$$P = \sum_{T \in \operatorname{STab}(n-1,k-1,k-1)} g_T(\mathbf{y}) \cdot F_T^{k-1} + \sum_{S \in \operatorname{STab}(n-1,0,k)} h_S(\mathbf{y}) \cdot M_S.$$

Writing $g_T(\mathbf{y}) = g_T^0(y_1, \dots, y_{n-1}) + x_n g_T^1(y_1, \dots, y_{n-1}) + x_n^2 g_T^2(\mathbf{y})$ and also $h_S(\mathbf{y}) = h_S^0(y_1, \dots, y_{n-1}) + x_n h_S^1(\mathbf{y})$, it follows from (2.) that $x_n^2 g_T^2(\mathbf{y}) \cdot F_T^{k-1} \in I(n,k)$, $x_n h_S^1(\mathbf{y}) \cdot M_S^k \in I(n,k)$, and also that $x_n g_T^1(y_1, \dots, y_{n-1}) \cdot F_T^{k-1} \in U$. Then taking monomial expansions, reversing orders of summations, and grouping like monomial terms, we get

$$P \equiv \sum_{T \in \text{STab}(n-1,k-1,k-1)} \sum_{\mathbf{a} \in \mathbb{N}^{n-1}(m)} c_T^0(\mathbf{a}) \mathbf{y}^{\mathbf{a}} \cdot F_T^{k-1}$$

$$+ \sum_{T \in \text{STab}(n-1,k-1,k-1)} \sum_{\mathbf{b} \in \mathbb{N}^n(m-1)} c_T^1(\mathbf{b}) \mathbf{y}^{\mathbf{b}} \cdot x_n F_T^{k-1} + \sum_{S \in \text{STab}(n-1,0,k)} \sum_{\mathbf{b} \in \mathbb{N}^n(m-1)} d_S^0(\mathbf{b}) \mathbf{y}^{\mathbf{b}} \cdot M_S$$

$$= \sum_{\mathbf{a} \in \mathbb{N}^n(m)} \mathbf{y}^{\mathbf{a}} \cdot \left(\sum_{T \in \text{STab}(n-1,k-1,k-1)} c_T^0(\mathbf{a}) F_T^{k-1} \right)$$

$$+ \sum_{\mathbf{b} \in \mathbb{N}^{n-1}(m-1)} \mathbf{y}^{\mathbf{b}} \cdot \left(\sum_{T \in \text{STab}(n-1,k-1,k-1)} c_T^1(\mathbf{b}) F_T^{k-1} + \sum_{S \in \text{STab}(n-1,0,k)} d_S^0(\mathbf{b}) M_S^k \right) \quad \text{mod } I(n,k)$$
and (3.) follows.

The following Lemma is analogous to Lemma 4.5 in Section 4.

Lemma 5.15. If $P \in I(n-1, k-1) \cap ((y_1, \dots, y_{n-1})^{(k)} + (x_n^2))$ is expressed as in (26), then each of its monomial summands must lie in $(y_1, ..., y_{n-1})^{(k)} + (x_n^2)$ is too, i.e. $\mathbf{y^a} \cdot \mu_{\mathbf{a}} \in (y_1, ..., y_{n-1})^{(k)} + (x_n^2)^{(k)}$ (x_n^2) and $\mathbf{y^b} \cdot v_b \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ for all $\mathbf{a} \in \mathbb{N}^{n-1}(m)$ and $\mathbf{b} \in \mathbb{N}^{n-1}(m-1)$.

Proof. By Lemma 5.14, we may write

$$P = \underbrace{\sum_{\mathbf{a} \in \mathbb{N}^{n-1}(m)} \mathbf{y}^{\mathbf{a}} \nu_{\mathbf{a}}}_{P_1} + \underbrace{\sum_{\mathbf{b} \in \mathbb{N}^{n-1}(m-1)} \mathbf{y}^{\mathbf{b}} \mu_{\mathbf{b}}}_{P_2} + P_3$$

for some $v_{\mathbf{a}} \in V$, some $\mu_{\mathbf{b}} \in U$, and some $P_3 \in I(n,k)$. Next note that if $P \in (y_1, \dots, y_{n-1})^{(k)} + ($ (x_n^2) then both $P_1 \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ and $P_2 \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. Indeed note that in their respective y-monomial expansions, those monomials in P_1 are all independent of y_n , whereas all monomials in the y-monomial expansion of P_2 are either divisible by $y_n = x_n$, or in $(y_1, \ldots, y_{n-1})^{(k)} + (x_n^2)$ already, by Lemma 5.8.

For the monomial products in P_1 we assume by way of contradiction that $\mathbf{y^a} \cdot \mathbf{v_a} \notin (y_1, \dots, y_{n-1})^{(k)} +$ (x_n^2) for some $\mathbf{a} \in \mathbb{N}^{n-1}$. Then since $(y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ is a monomial ideal, it follows that in the y-monomial expansion of $y^a v_a$, there must be some monomial say y^d which is not in $(y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$. Since $\mathbf{y}^{\mathbf{a}}v_{\mathbf{a}}$ is independent of x_n , so is $\mathbf{y}^{\mathbf{d}}$. Since $(y_1,\ldots,y_{n-1})^{(k)}$ consists of all y-monomials of weight at least k, it follows that $wt(y^d) \le k - 1$. On the other hand, every y-monomial in the monomial expansion of v_a has weight equal to k-1, hence it follows that $wt(v^d) = k - 1$. But then we can deduce, as in the proof of Lemma 4.5, that

$$\frac{y^d}{\sqrt{y^d}} = y^a$$

and hence the exponent vector \mathbf{d} uniquely determines the exponent vector \mathbf{a} . This implies that y^{d} occurs with the same coefficient in the monomial expansion of the term $y^{a}v_{a}$ as it does in the entire sum

$$P_1 = \sum_{\mathbf{a} \in \mathbb{N}^{n-1}(m)} \mathbf{y}^{\mathbf{a}} v_{\mathbf{a}},$$

contradicting the fact that $P_1 \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. The argument for P_2 is similar.

Lemma 5.15 says that to check the containment

$$I(n,k) \supseteq I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)} + (x_n^2))$$

it suffices to check on products of monomials and forms in either V or U, i.e. for any $\mathbf{a}, \mathbf{b} \in$ \mathbb{N}^{n-1} and for any $\nu \in V$ and any $\mu \in U$

$$\mathbf{y}^{\mathbf{a}}\nu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \Rightarrow \qquad \mathbf{y}^{\mathbf{a}}\nu \in I(n, k)$$

$$\mathbf{y}^{\mathbf{b}}\mu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \Rightarrow \qquad \mathbf{y}^{\mathbf{b}}\mu \in I(n, k)$$

$$(27)$$

$$\mathbf{y}^{\mathbf{b}}\mu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2) \Rightarrow \qquad \mathbf{y}^{\mathbf{b}}\mu \in I(n, k)$$
 (28)

The next lemma verifies implications (27) and (28) in the special case where $\mathbf{a}, \mathbf{b} = \mathbf{0}$. This seems to be where we need our assumptions on p.

Lemma 5.16. Fix $v \in V(n-1, k-1, k-1)$ and $\mu \in U(n-1, k-1, k)$. If $v \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ then v = 0. If p = 0 or $p \ge k + 1$ and if $\mu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ then $\mu \in \mathfrak{a}(n, k, k) \subset I(n, k)$.

Proof. The first statement for v is obvious for degree reasons. For the second statement, we can write $\mu = \mu(x_1, \dots, x_n) = x_n \alpha + \beta$ for polynomials $\alpha = \alpha(x_1, \dots, x_{n-1}) \in V_{\mathbf{x}}(n-1, k-1, k-1)$ and $\beta = \beta(x_1, \dots, x_{n-1}) \in V_{\mathbf{x}}(n-1, 0, k)$. Here it will be important to distinguish between polynomials in x-variables and those in the y-variables, hence we adopt the notation $V_{\mathbf{x}}(m, j, j)$ and $V_{\mathbf{y}}(m, j, j)$ to denote the \mathbb{F} -span of Specht polynomials in the x-variables and y-variables, respectively; note that if $m \le n-1$ these two subspaces coincide.

Consider inclusions of monomial complete intersections, $B \subset A$ defined in the y-variables by:

$$B = \frac{\mathbb{F}[y_1, \dots, y_{n-1}]}{(y_1^2, \dots, y_{n-1}^2)} \hookrightarrow A = \frac{\mathbb{F}[y_1, \dots, y_n]}{(y_1^2, \dots, y_n^2)}$$

and let L_B , D_B , H_B , and L_A , D_A , H_A be their respective raising, lowering, and semi-simple operators, respectively, as in Section 3. Then since p = 0 or $p \ge k + 1$, Lemma 3.4 implies that the lowering maps for B and A,

$$D_B: B_k \to B_{k-1}$$
, and $D_A: A_k \to A_{k-1}$

are both surjective, which by Lemma 3.2 is equivalent to their primitive subspaces $P_{B,k} = \ker(D_B) \cap B_k$ and $P_{A,k} = \ker(D_A) \cap A_k$ satisfying

$$P_{B,k} = V_{y}(n-1, k, k)$$
, and $P_{A,k} = V_{y}(n, k, k)$.

We will identify B (resp. A) with the subspace spanned by square-free monomials in variables y_1, \ldots, y_{n-1} (resp. y_1, \ldots, y_n).

Then if $\mu = x_n \alpha(x_1, \dots, x_{n-1}) + \beta(x_1, \dots, x_{n-1}) \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$, by Lemma 5.8 we have

$$\mu \equiv (-1)^{k-1} x_n \left(\alpha(y_1, \dots, y_{n-1}) + D_B(\beta(y_1, \dots, y_{n-1})) \right) \equiv 0 \mod (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$$
 (29)

(note that $D_B(\alpha(y_1,\ldots,y_{n-1}))=0$ and $\beta(y_1,\ldots,y_{n-1})\in (y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$ automatically). Dividing by x_n in (29) we see that

$$\alpha(y_1, \dots, y_{n-1}) + D_B(\beta(y_1, \dots, y_{n-1})) \in ((y_1, \dots, y_{n-1})^{(k)} + x_n^2) : x_n = (y_1, \dots, y_{n-1})^{(k)} + (x_n)$$

which, for degree reasons, implies that

$$\alpha(v_1,\ldots,v_{n-1}) + D_R(\beta(v_1,\ldots,v_{n-1})) = 0.$$

Therefore $\beta_y = \beta(y_1, \dots, y_{n-1}) \in B_k$ is a square-free polynomial in y_1, \dots, y_{n-1} such that $D_B(\beta_y) = -\alpha_y \in V(n-1, k-1, k-1) = P_{B,k-1}$. Applying the commutator relation

$$[D_B, L_B] = H_B$$

to $\alpha_{\mathbf{v}} = \alpha(y_1, \dots, y_{n-1})$, we find that

$$D_B \circ L_B(\alpha_y) = H(\alpha_y) = (n-1-2(k-1)) \cdot \alpha_y$$

which implies that

$$\mu_B(y_1, \dots, y_{n-1}) := L_B(\alpha_{\mathbf{v}}) + (n - 2k + 1) \cdot \beta_{\mathbf{v}} \in P_{B,k} = V_{\mathbf{v}}(n - 1, k, k).$$
 (30)

Note that Equation (30) also implies that $\mu_B := \mu_B(x_1, \dots, x_{n-1}) \in V_{\mathbf{x}}(n-1, k, k)$. We can also apply the commutator relations for those operators on A. Note that the restriction of $D_A = D_B + \partial/\partial y_n$ to B is D_B , and that by Lemma 3.5, we have

$$V_{\mathbf{v}}(n-1,k,k) = V_{\mathbf{v}}(n,k,k) \cap B_k$$

It follows that $D_A(\beta_y) = D_B(\beta_y) = -\alpha_y \in V_y(n-1, k-1, k-1) \subset V_y(n, k-1, k-1)$, and hence applying the commutator relation

$$[D_A, L_A] = H_A$$

to $D_A(\beta_{\mathbf{v}}) = -\alpha_{\mathbf{v}}$ we also find that

$$\mu_A(y_1, \dots, y_{n-1}, y_n) := L_A(\alpha_{\mathbf{v}}) + (n - 2k + 2) \cdot \beta_{\mathbf{v}} \in P_{A,k} = V_{\mathbf{v}}(n, k, k)$$
 (31)

and hence $\mu_A = \mu_A(x_1, \dots, x_n) \in V_{\mathbf{x}}(n, k, k)$. Noting that $L_A = L_B + x_n$ we see that adding μ to μ_B yields μ_A , i.e.

$$\underbrace{(x_n\alpha+\beta)}_{\mu}+\underbrace{((x_1+\cdots+x_{n-1})\cdot\alpha+(n-2k+1)\beta)}_{\mu_B}=\underbrace{(x_1+\cdots+x_n)\cdot\alpha+(n-2k+2)\beta}_{\mu_A}.$$

It follows from (30) and (31) that $\mu = x_n \cdot \alpha + \beta = \mu_A - \mu_B \in V_{\mathbf{x}}(n, k, k) = V(n, k, k) \subset \mathfrak{a}(n, k, k)$, as claimed.

Next we check containment (25) for monomials which are not contained in the support of ν and μ . The following Lemma is analogous to Lemma 4.6 in Section 4. Recall that the support of the (shifted) Specht polynomial for T is the set of square-free monomials indexed by subsets of numbers in the support of T, no two of which lie in the same column of T.

Lemma 5.17. Fix tableaux $T \in \text{STab}(n-1, k-1, k-1)$ and $S \in \text{STab}(n-1, 0, k)$, and exponent vectors $\mathbf{a}, \mathbf{b} \in \mathbb{N}^{n-1}$.

- 1. If $\sqrt{\mathbf{x}^{\mathbf{a}}} \notin \operatorname{supp}(F_T^{k-1})$ then $\mathbf{y}^{\mathbf{a}} \cdot F_T^{k-1} \in \mathfrak{a}(n, k, k) \subset I(n, k)$.
- 2. If $\sqrt{\mathbf{x}^{\mathbf{b}}} \notin \text{supp}(M_{\mathbf{x}}^{k})$ then $\mathbf{y}^{\mathbf{b}} \cdot M_{\mathbf{x}}^{k} \in (x_{1}, \dots, x_{n})^{(k+1)} \subset I(n, k)$.

Proof. For (1.) assume $\sqrt{\mathbf{x}^{\mathbf{a}}} \notin \operatorname{supp}(F_T^{k-1})$. If some variable $x_i \in \operatorname{supp}(\mathbf{y}^{\mathbf{a}})$ but $x_i \notin \operatorname{supp}(F_T^{k-1})$, then clearly

$$\mathbf{y^a} \cdot F_T^{k-1} = \mathbf{y^{a'}} \cdot y_i \cdot F_T^{k-1} = \mathbf{y^{a'}} \cdot (x_n - x_i) \cdot F_T^{k-1} \in V(n,k,k).$$

We may therefore assume that every variable of $\mathbf{x}^{\mathbf{a}}$ lies in the support of F_T^{k-1} . But since $\sqrt{\mathbf{x}^{\mathbf{a}}} \notin \operatorname{supp}(F_T^{k-1})$ there must be two distinct indices i, j for which $y_i, y_j \in \operatorname{supp}(\mathbf{y}^{\mathbf{a}})$ and i, j lie in the same column in T. Choose any index $r \neq i, j$ such that $x_r \notin \operatorname{supp}(F_T^{k-1})$ -presumably there is such an r since we are assuming that $k+1 \leq n-k$ -and let $(i, r), (j, r) \in \mathfrak{S}_n$ be the transpositions swapping i, r and j, r, respectively. Then we have

$$\mathbf{y}^{\mathbf{a}} \cdot F_T^{k-1} = \mathbf{y}^{\mathbf{a}} \cdot \left(F_{(i,r).T}^{k-1} + F_{(j,r).T}^{k-1} \right)$$

and since $x_i \notin \text{supp}(F_{(i,r),T}^{k-1})$, we must have

$$\mathbf{y}^{\mathbf{a}} \cdot F_{(i,r),T}^{k-1} = \mathbf{y}^{\mathbf{a}'} \cdot y_i \cdot F_{(i,r),T}^{k-1} = \mathbf{y}^{\mathbf{a}'} \cdot (x_n - x_i) \cdot F_{(i,r),T}^{k-1} \in \mathfrak{a}(n,k,k) \subset I(n,k)$$

and similarly for $F_{(j,r),T}^{k-1}$. This proves that $\mathbf{y^a} \cdot F_T^{k-1} \in I(n,k)$. Item (2.) is easier and left to the reader.

Finally, we need to check what containment (25) for monomials which are contained in the support of ν or μ . Since the ideal $(y_1, \ldots, y_{n-1})^{(k)} + (x_n^2)$ is generated by square-free monomials in (y_1, \ldots, y_{n-1}) and the monomial x_n^2 , it suffices to prove implications (27) and (28) for square free $\mathbf{y}^{\mathbf{a}}$. Also, by symmetry, it will suffice to assume that $\mathbf{y}^{\mathbf{a}} = \mathbf{y}^{\mathbf{m}} = y_1 \cdots y_m$. As in Corollary 2.7, we define the subset of standard tableau on a shape λ as those which contain the set of integers $\{1, \ldots, m\}$ in their support, denoted by $\mathrm{STab}_m(\lambda) \subset \mathrm{STab}(\lambda)$, and define the subspaces

$$V_{m} = V_{m}(n-1, k-1, k-1) = \left\langle F_{T}^{k-1} \mid T \in STab_{m}(n-1, k-1, k-1) \right\rangle$$

$$U_{m} = U_{m}(n-1, k-1, k) = \left\langle x_{n}F_{T}^{k-1} \mid T \in STab_{m}(n-1, k-1, k-1) \right\rangle + \left\langle M_{S}^{k} \mid S \in STab_{m}(n-1, 0, k) \right\rangle$$

We also define their complimentary subspaces

$$\begin{split} V^m = & V^m(n-1,k-1,k-1) = & \left\langle F_T^{k-1} \mid T \in \operatorname{STab}(n-1,k-1,k-1) \setminus \operatorname{STab}_m(n-1,k-1,k-1) \right\rangle \\ U^m = & U^m(n-1,k-1,k) = & \left\langle x_n F_T^{k-1} \mid T \in \operatorname{STab}(n-1,k-1,k-1) \setminus \operatorname{STab}_m(n-1,k-1,k-1) \right\rangle \\ & + \left\langle M_S^k \mid S \in \operatorname{STab}(n-1,0,k) \setminus \operatorname{STab}_m(n-1,0,k) \right\rangle. \end{split}$$

Then we have vector space decompositions $V = V_m \oplus V^m$ and $U = U_m + U^m$, and Lemma 5.17 implies that $\mathbf{y^m} \cdot \alpha \in I(n,k)$ for $\alpha \in V^m \sqcup U^m$. Hence it only remains to check $\alpha \in V_m \sqcup U_m$. As in Corollary 2.7 we have the following bijective map of vector spaces:

$$V_m(n-1,k-1,k-1) \longrightarrow V([n-1]_m,k-1-m,k-1-m)$$

$$F_T^{k-1} \longmapsto F_{T'}^{k-1-m}$$

where

$$T = \frac{1 \cdots m i_{m+1} \cdots i_k}{|i_{k+1}| \cdots |i_{n-k}| j_1 \cdots j_m j_{m+1} \cdots j_k}$$

$$(32)$$

and

$$T' = \frac{i_{m+1} \cdots i_k}{|i_{k+1}| \cdots |i_{n-k}| \ j_1 | \cdots |j_m| j_{m+1} \cdots |j_k|}$$
(33)

We also have the (possibly non-injective) linear map

$$U_m(n-1,k-1,k-1) \longrightarrow U([n-1]_m,k-1-m,k-m)$$

$$x_n F_T^{k-1} \longmapsto x_n F_{T'}^{k-1-m}$$

$$M_S^k \longmapsto M_{S'}^{k-m}$$

where $T \in \operatorname{STab}_m(n-1,k-1,k-1)$ and $T' \in \operatorname{STab}([n-1]_m,k-1-m,k-1-m)$ are the tableau in (32) and (33), respectively, and where where $S \in \operatorname{STab}(n-1,0,k)$ and $S' \operatorname{STab}([n-1]_m,0,k-m)$ are given by

$$S = \frac{1 \left| \cdots \right| m \left| i_{m+1} \cdots \right| i_k}{\left| i_{k+1} \right| \cdots \left| i_{n-k} \right|}$$
(34)

and

$$S' = \frac{|i_{m+1}| \cdots |i_k|}{|i_{k+1}| \cdots |i_{n-k}|}$$

$$(35)$$

The following two lemmas, will be useful.

Lemma 5.18. For any $\beta \in V([n-m], k-m, k-m)$ where $[n-m] = \{m+1, ..., n\}$, we have

$$(\mathbf{x}^{\mathbf{m}})^2 \cdot \beta \in I(n,k).$$

Proof. Fix a tableau $T' \in STab([n-m], k-m, k-m)$ as in (33) and let $T \in STab_m(n-1, k-1, k-1)$ be the corresponding tableau as in (32). Then it suffices to show that we have

$$(\mathbf{x}^{\mathbf{m}})^2 \cdot F_{T'}^{k-m} = x_1^2 \cdots x_m^2 \cdot F_{T'}^{k-m} \in I(n,k) = \mathfrak{a}(n,k,k) + (x_1,\ldots,x_n)^{(k+1)}$$

On the other hand we clearly have

$$x_1 \cdots x_m \cdot F_T^k = x_1^2 \cdots x_m^2 \cdot F_{T'}^{k-m} + (\text{stuff in } (x_1, \dots, x_n)^{(k+1)}) \in \mathfrak{a}(n, k, k)$$

and the result follows.

Lemma 5.19. Fix elements $v \in V_m(n-1,k-1,k-1)$ and $\mu \in U_m(n-1,k-1,k)$, and let $v' \in V([n-1]_m,k-1-m,k-1-m)$ and $\mu' \in U([n-1]_m,k-1-m,k-m)$ their respective images under the maps above. Then we have

- 1. $\mathbf{y^m} (\nu (-1)^m \mathbf{y^m} \nu') \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$
- 2. $\mathbf{v}^{\mathbf{m}} (\mu (-1)^m \mathbf{v}^{\mathbf{m}} \mu') \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$
- 3. $\mathbf{x}^{\mathbf{m}} (\mu \mathbf{x}^{\mathbf{m}} \mu') \in I(n, k)$, and
- 4. $\mathbf{y}^{\mathbf{m}} \cdot \mu (-1)^m \mathbf{x}^{\mathbf{m}} \cdot \mu \in I(n, k)$.

Proof. For (1.) note that it suffices to see that for $T \in STab_m(n-1,k-1,k-1)$ and $T' \in STab([n-1]_m,k-1-m,k-1-m)$ as in (32) and (33) we have

$$\mathbf{y}^{\mathbf{m}} \left(F_T^{k-1} - (-1)^m \mathbf{y}^{\mathbf{m}} \cdot F_{T'}^{k-1} \right) \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

By Lemma 5.8, we have $F_{T,\mathbf{y}}^{k-1} = F_T^{k-1}(y_1,\ldots,y_{n-1}) = (-1)^{k-1} \cdot F_T^{k-1}(x_1,\ldots,x_{n-1}) = F_T^{k-1}$ and $F_{T',\mathbf{y}}^{k-1-m} = (-1)^{k-1-m} \cdot F_{T'}^{k-1}$. Since the difference $F_{T,\mathbf{y}}^{k-1} - \mathbf{y^m} \cdot F_{T',\mathbf{y}}^{k-1}$ is a combination of y-monomials which do not contain $\mathbf{y^m}$ in their support, it follows that the product

$$\mathbf{y^m} \left(F_T^{k-1} - (-1)^m \mathbf{y}^m \cdot F_{T'}^{k-1} \right) \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

For (2.), write $\mu = x_n \cdot \alpha + \beta$ where $\alpha \in V_m(n-1,k-1,k-1)$ and $\beta \in V_m(n-1,0,k)$. By (1.) we know that

$$\mathbf{y}^{\mathbf{m}}(x_n \cdot \alpha - (-1)^m x_n \cdot \alpha') \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$$

hence it suffices to take $\mu = \beta = M_S^k \in V_m(n-1,0,k)$ and show that

$$\mathbf{y}^{\mathbf{m}} \left(M_S^k - (-1)^m \mathbf{y}^{\mathbf{m}} M_{S'}^{k-m} \right) \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$$

for $S \in \operatorname{STab}_m(n-1,0,k)$ and $S' \in \operatorname{STab}([n-1]_m,0,k-m)$ as in (34) and (35). In this case, we have

$$M_S^k = x_1 \cdots x_m \cdot M_{S'}^{k-m}$$

and Lemma 5.8 implies that the difference satisfies

$$\begin{aligned} M_{S}^{k} - (-1)^{m} \mathbf{y}^{\mathbf{m}} \cdot M_{S'}^{k-m} &= (-1)^{k} M_{S,\mathbf{y}}^{k} + (-1)^{k-1} x_{n} D(M_{S,\mathbf{y}}^{k}) \\ &- (-1)^{m} \mathbf{y}^{\mathbf{m}} \left((-1)^{k-m} M_{S',\mathbf{y}}^{k-m} + (-1)^{k-1-m} x_{n} D(M_{S',\mathbf{y}}^{k-m}) \right) & \mod(x_{n}^{2}) \\ &= (-1)^{k-1} x_{n} \cdot \left(D(M_{S,\mathbf{y}}^{k}) - \mathbf{y}^{\mathbf{m}} \cdot D(M_{S'}^{k-m}) \right) & \mod(x_{n}^{2}) \\ &= (-1)^{k-1} x_{n} \cdot \left(\sum_{i=1}^{m} y_{1} \cdots \hat{y_{i}} \cdots y_{m} \cdot M_{S',\mathbf{y}}^{k-m} \right) & \mod(x_{n}^{2}) \end{aligned}$$

Therefore it follows that the product satisfies

$$\mathbf{y}^{m} \left(M_{S}^{k} - (-1)^{m} \mathbf{y}^{\mathbf{m}} \cdot M_{S'}^{k-m} \right) \in (y_{1}, \dots, y_{n-1})^{(k)} + (x_{n}^{2}).$$

For (3.), we write $\mu = x_n \alpha + \beta$ where $\alpha \in V_m(n-1,k-1,k-1)$ and $\beta \in V_m(n-1,0,k)$, and also $\mu' = x_n \alpha' + \beta'$. Then we have

$$\mathbf{x}^{\mathbf{m}}(x_n\alpha - \mathbf{x}^{\mathbf{m}}x_n\alpha') \in (x_1, \dots, x_n)^{(k+1)} \subset I(n, k)$$

and also

$$\mathbf{x}^m (\beta - \mathbf{x}^m \beta') = 0 \in I(n, k)$$

and the result follows.

For (4.), note that

$$\mathbf{y^m} - (-1)^m \mathbf{x^m} = (x_n - x_1) \cdots (x_n - x_m) - (-1)^m x_1 \cdots x_m = x_n \text{ (stuff)}$$

and since $x_n \cdot \mu \in I(n, k)$ by Lemma 5.14, the result follows.

The following is an analogue of Lemma 4.7 in Section 4.

Lemma 5.20. If $\mathbf{y^m} \cdot \nu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ for some $\nu \in V_m$, then $\nu = 0$, and in particular, $\mathbf{y^m} \cdot \nu \in I(n, k)$. If p = 0 or $p \ge k + 1$, then if $\mathbf{y^m} \cdot \mu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$ for some $\mu \in U_m$, then $\mathbf{v^m} \cdot \mu \in I(n, k)$.

Proof. First assume that for some $v \in V_m$ we have $\mathbf{y^m} \cdot v \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. By Lemma 5.19(1.) we have

$$\mathbf{y^m} (\nu - \mathbf{y^m} \cdot \nu') \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$$

and it follows also that

$$(\mathbf{y^m})^2 \cdot \mathbf{v'} \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2).$$

Therefore we have

$$v' \in ((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) : (\mathbf{y^m})^2) = ((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) : \mathbf{y^m}) = (y_{m+1}, \dots, y_{n-1})^{k-m} + (x_n^2).$$

and hence that $v' \in V([n-1]_m, k-1-m, k-1-m) \cap ((y_{m+1}, \dots, y_{n-1})^{(k-m)} + (x_n^2)$. By Lemma 5.16, it follows that v' = 0, and therefore also v = 0, which proves the first statement.

Next, assume that p = 0 or $p \ge k + 1$, and assume that for some $\mu = \mu(x_1, \dots, x_{n-1}) \in U_m$ we have $\mathbf{y^m} \cdot \mu \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$. Then by Lemma 5.19(2.), we must also have $(\mathbf{y^m})^2 \mu' \in (y_1, \dots, y_{n-1})^{(k)} + (x_n^2)$, and hence also we must have

$$\mu' \in \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) : (\mathbf{y^m})^2 \right) = \left((y_1, \dots, y_{n-1})^{(k)} + (x_n^2) : \mathbf{y^m} \right) = (y_{m+1}, \dots, y_{n-1})^{k-m} + (x_n^2).$$

Therefore we have $\mu' \in U([n-1]_m, k-1-m, k-m) \cap (y_{m+1}, \dots, y_{n-1})^{k-m} + (x_n^2)$, and it follows from Lemma 5.16 that $\mu' \in V([n]_m, k, k)$. Therefore by Lemma 5.18, it follows that $(\mathbf{x}^{\mathbf{m}})^2 \cdot \mu' \in I(n, k)$. Then by Lemma 5.19(3.) we must also have $\mathbf{x}^m \cdot \mu \in I(n, k)$, and from Lemma 5.19(4.) it follows that $\mathbf{y}^{\mathbf{m}} \cdot \mu \in I(n, k)$, as desired.

Finally we are in a position to prove Theorem 5.5:

Proof of Theorem 5.5. Assume that p = 0 or $p \ge k + 1$. By Lemma 5.13, we have

$$I(n,k) \subseteq I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)} + (x_n^2)).$$

For the reverse containment, Lemma 5.15 implies we only have to check on products of monomials in (y_1,\ldots,y_{n-1}) and forms in the subspaces V=V(n-1,k-1,k-1) and U=U(n-1,k-1,k). Since $(y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$ is generated by square-free monomials in (y_1,\ldots,y_{n-1}) it follows that we may assume our monomials are square-free, and by symmetry we may assume that our monomial is $\mathbf{y^m}=y_1\cdots y_m$. Write $V=V_m\oplus V^m$ and $U=U_m+U^m$ as above. Then Lemma 5.17 implies that $\mathbf{y^m}\cdot\alpha\in I(n,k)$ for any $\alpha\in V^m\sqcup U^m$. Also if $\beta\in V_m\sqcup U_m$, then Lemma 5.20 implies that if $\mathbf{y^m}\cdot\beta\in (y_1,\ldots,y_{n-1})^{(k)}+(x_n^2)$, then $\mathbf{y^m}\cdot\beta\in I(n,k)$, and the result follows.

Conversely, assume that $0 . Then as in the proof of Theorem 5.6 and in particular (24), we have <math>I(n,k) \neq I(n-1,k-1) \cap ((y_1,\ldots,y_{n-1})^{(k)}+(x_n^2))$.

References

- [1] C.B. Zamaere, S. Griffeth, S.V. Sam, *Jack Polynomials as Fractional Quantum Hall States and the Betti Numbers of the (k + 1)-Equals Ideal.*, Commun. Math. Phys. 330 (2014), 415–434.
- [2] P. Etingof, E. Gorsky, I. Losev, *Representations of Cherednik algebras with minimal support and torus knots.*, Adv. Math. 277 (2015), 124–180.
- [3] W. Feller, An Introduction to Probability Theory and Its Applications, Volume 1, Third Ed. Wiley, 1968
- [4] R. Fröberg, B. Shapiro, On Vandermonde Varieties., Math. Scand., 119, (2016), 76–91.
- [5] M. Hochster, J. Eagon, Cohen-Macaulay Rings, Invariant Theory, and the Generic Perfection of Determinantal Loci., Am. J. Math., vol. 93, no. 4, 1971, 1020–1058.
- [6] T. Harima, T. Maeno, H. Morita, Y. Numata, A. Wachi, J. Watanabe, *The Lefschetz Properties*, Lecture Notes in Mathematics 2080. Springer, 2013.
- [7] A. Kustin, A. Vraciu, *The weak Lefschetz property for monomial complete intersections in positive characteristic.*, Trans. Amer. Math. Soc., 366, (2014), 4571–4601.
- [8] H. Matsumura, *Commutative Ring Theory*, Cambridge Studies in Advanced Mathematics, No. 8. Cambridge University Press, 2006.
- [9] J. Migliore, R. Miró-Roig, U. Nagel, *Monomial ideals, almost complete intersections and the weak Lefschetz property.*, Trans. Amer. Math. Soc., 363, (2011), 229–257.
- [10] M. Nagata, Local Rings, Interscience Publishers, New York, 1962.
- [11] G.A. Reisner, *Cohen-Macaulay Quotients of Polynomial Rings*, Adv. Math. 21 (1976), 30–49.
- [12] B. Sagan, The Symmetric Group Representations, Combinatorial Algorithms, and Symmetric Functions, Second Edition, Graduate Texts in Mathematics, No. 203. Springer-Verlag, New York-Heidelberg, 2001.

- [13] J. Watanabe, *The Dilworth number of Artinian rings and finite posets with rank function*, In: Commutative Algebra and Combinatorics, Kyoto, 1985,(eds. M. Nagata and H. Matsumura), Adv. Stud. Pure Math., 11, North-Holland, Amsterdam; Kinokuniya, Tokyo, 1987, 303-312.
- [14] J. Watanabe, K. Yanagawa, *Vandermonde Determinantal Ideals.*, Math. Scand., Vol. 125, no. 2, Oct. 2019, pp. 179-184.
- [15] K. Yanagawa, When is a Specht ideal Cohen–Macaulay?, J. Commut. Algebra To appear.