

# Periodic Trajectories in P-Time Event Graphs and the Non-Positive Circuit Weight Problem

Davide Zorzenon, Jan Komenda and Jörg Raisch

**Abstract**—P-time event graphs (P-TEGs) are specific timed discrete-event systems, in which the timing of events is constrained by intervals. An important problem is to check, for all natural numbers  $d$ , the existence of consistent  $d$ -periodic trajectories for a given P-TEG. Let us consider a different, seemingly unrelated problem in graph theory: given three arbitrary square matrices  $P, I$  and  $C$  with elements in  $\mathbb{R} \cup \{-\infty\}$ , find all real values of parameter  $\lambda$  such that the parametric directed graph having arcs weights of the form  $w((j,i)) = \max(P_{ij} + \lambda, I_{ij} - \lambda, C_{ij})$  (for all arcs  $(j,i)$ ) does not contain circuits with positive weight. In a related paper, we have proposed a strongly polynomial algorithm that solves the latter problem faster than other algorithms reported in literature. In the present paper, we show that the first problem can be formulated as an instance of the second; consequently, we prove that the same algorithm can be used to find  $d$ -periodic trajectories in P-TEGs faster than with previous approaches.

## I. INTRODUCTION

P-time event graphs (P-TEGs) constitute a specific class of timed discrete-event systems, namely event graphs in which the sojourn time of tokens in places is constrained by lower and upper bounds. Events that do not occur in the specified time window correspond, in the real system modeled by the P-TEG, to failure of meeting time specifications. P-TEGs have been applied for modeling and analyzing time schedules in manufacturing systems such as electroplating lines, baking processes, cluster tools [2], [17], [7], [13].

Timed event graphs are usually assumed to operate under the earliest firing rule [1]. This makes their temporal evolution deterministic. In contrast, in P-TEGs the time evolution of the marking is non-deterministic, since transitions are not required to fire as soon as they are enabled. In fact, following the earliest firing rule could lead to the violation of upper bound constraints. For this reason, it is fundamental to characterize the set of all admissible infinite trajectories for a P-TEG (*i.e.*, trajectories for which all the constraints are satisfied), and to verify that it is non-empty; in this case, the P-TEG is said to be *consistent*. In [19], we introduced a stronger property than consistency that is even more desirable for P-TEGs applications; a P-TEG is *boundedly consistent* if it admits at least one consistent trajectory such that the delay between the  $k^{\text{th}}$  firings of every pair of transitions

is bounded for all  $k$ . Moreover, we proved that a P-TEG is boundedly consistent if and only if there exists at least one 1-periodic trajectory that is consistent for the P-TEG. Since the problem of checking the existence of 1-periodic trajectories can be solved using linear programming [2], [6], bounded consistency can be checked in *weakly* polynomial time (on the contrary, no algorithm that checks standard consistency has been found yet). However, the use of linear programming for finding 1-periodic trajectories implies two issues that we will try to overcome in this paper. First, in weakly polynomial time algorithms, in contrast to *strongly* polynomial time algorithms, execution time grows with the size of their numerical inputs (expressed in terms of the number of bits needed to represent them); this leads to slower execution when “large” values of the inputs (in our case, the values of lower and upper bounds) are employed. Second, linear programming does not provide an explicit closed formula of the solution.

To overcome these issues, we focus on graph theory. Given a matrix  $A \in \mathbb{R}_{\max}^{n \times n}$  (where  $\mathbb{R}_{\max} = \mathbb{R} \cup \{-\infty\}$ ), the precedence graph  $\mathcal{G}(A)$  is a weighted directed graph whose arc weights are equal to the finite elements of matrix  $A$ . The number of nodes in  $\mathcal{G}(A)$  is  $n$  and the number of arcs, denoted by  $q$ , is equal to the number of finite entries in  $A$ . When elements of  $A$  are functions of real parameters, we say that  $\mathcal{G}(A)$  is a *parametric precedence graph*. In this context, the *Non-positive Circuit weight Problem* (NCP) consists in finding all the values of the parameters such that  $\mathcal{G}(A)$  does not contain circuits with positive weight. The NCP has been studied in the literature for the case when elements of  $A$  depend on a single parameter  $\lambda \in \mathbb{R}$  and the relation  $\lambda \mapsto A(\lambda)$  assumes some specific form [12], [18]. For instance, an algorithm by Levner and Kats [14] solves the NCP in strongly polynomial time  $\mathcal{O}(qn^2)$  when  $A_{ij}(\lambda) = X_{ij} + Y_{ij} \times \lambda$ , with  $X_{ij} \in \mathbb{R}_{\max}$  and  $Y_{ij} \in \{-1, 0, +1\}$ . In this paper we consider the NCP on a larger class of parametric precedence graphs, in which  $A(\lambda)$  has the form  $A_{ij}(\lambda) = \max(P_{ij} + \lambda, I_{ij} - \lambda, C_{ij})$ , with  $P_{ij}, I_{ij}, C_{ij} \in \mathbb{R}_{\max}$ . Using the max-plus algebra notation, matrix  $A$  can be expressed compactly as  $A(\lambda) = \lambda P \oplus \lambda^{-1} I \oplus C$ , where  $P, I, C$  are three square matrices in  $\mathbb{R}_{\max}$ . Since weights in  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$  depend proportionally, inversely and constantly on  $\lambda$  (in the max-plus sense), the NCP on this class of parametric precedence graphs is called *Proportional-Inverse-Constant-NCP* (PIC-NCP). It is easy to see that the PIC-NCP can always be transformed into an equivalent NCP that can be solved using the Levner-Kats algorithm [14]. However in order to do so, the worst-case complexity of the algorithm increases to  $\mathcal{O}(n^6)$  [20].

D. Zorzenon and J. Raisch are with Technische Universität Berlin, Control Systems Group, Einsteinufer 17, D-10587 Berlin, Germany, {zorzenon, raisch}@control.tu-berlin.de. J. Komenda is with Institute of Mathematics, Czech Academy of Sciences, Žitkova 22, 616 62 Brno, Czech Republic, komenda@ipm.cz. This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation), Projektnummer RA 516/14-1. Partially supported by the GACR grant 19-06175J, by MSMT INTER-EXCELLENCE project LTAUSA19098, and by RVO 67985840.

The main purpose of the present paper is to show that the problem of finding all periods of 1-periodic trajectories that are consistent for a P-TEG can be formulated as an instance of the PIC-NCP. In [20], we provided a closed formula for the solution set of the PIC-NCP that can be computed in  $\mathcal{O}(n^4)$ , improving Levner-Kats worst-case complexity. As a consequence, we prove that the existence of admissible 1-periodic trajectories, and thus bounded consistency, can also be checked in strongly polynomial time complexity  $\mathcal{O}(n^4)$  in P-TEGs with initially 0 or 1 token per place. Moreover (as recalled in Section IV), every P-TEG can be transformed into an equivalent one in which the number of tokens initially residing in every place is at most 1, by increasing the number of transitions to  $\bar{n}$  (evaluated in (4), at page 4), see e.g. [15]. This transformation allows to check the existence of 1-periodic trajectories in general P-TEGs in complexity  $\mathcal{O}(\bar{n}^4)$ , which is strongly polynomial in the number of the transitions of the transformed P-TEG, but pseudo-polynomial considering the whole transformation procedure. In addition, exploiting the connection between periodic trajectories and the NCP, we prove a surprising result regarding more general trajectories: the set of all periods of  $d$ -periodic trajectories that are consistent for a P-TEG is the same for all  $d \in \mathbb{N}$ .

The remainder of the paper is organized as follows. In Section II, algebraic preliminaries are recalled. Precedence graphs and the NCP are introduced in Section III. Section IV contains the formal definition of P-TEGs, and the connection between the PIC-NCP and the problem of checking the existence of consistent  $d$ -periodic trajectories for P-TEGs is presented in Section V. Finally, conclusions are given in Section VI.

Notation: the set of positive, respectively non-negative, integers is denoted by  $\mathbb{N}$ , respectively  $\mathbb{N}_0$ . The set of non-negative real numbers is denoted by  $\mathbb{R}_{\geq 0}$ . Moreover,  $\mathbb{R}_{\max} := \mathbb{R} \cup \{-\infty\}$  and  $\bar{\mathbb{R}}_{\max} := \mathbb{R}_{\max} \cup \{+\infty\}$ .

## II. ALGEBRAIC PRELIMINARIES

In this section, some preliminary concepts from the theory of idempotent semirings and semifields are summarized. For more details, we refer to [1], [10], [9].

### A. Idempotent semirings and semifields

A set  $\mathcal{D}$  endowed with two binary operations  $\oplus$  (addition) and  $\otimes$  (multiplication),  $(\mathcal{D}, \oplus, \otimes)$ , is an *idempotent semiring* (or *dioid*), if the following properties hold:  $\oplus$  is associative, commutative, idempotent ( $\forall a \in \mathcal{D}, a \oplus a = a$ ), and has a neutral element  $\varepsilon$ ;  $\otimes$  is associative, distributes over  $\oplus$ , has a neutral element  $e$ , and  $\varepsilon$  is absorbing for  $\otimes$  ( $\forall a \in \mathcal{D}, a \otimes \varepsilon = \varepsilon \otimes a = \varepsilon$ ). The canonical order relation  $\preceq$  on  $\mathcal{D}$  is defined by:  $\forall a, b \in \mathcal{D}, a \preceq b \Leftrightarrow a \oplus b = b$ . From this definition, it follows that  $\otimes$  is order preserving, i.e.,  $\forall a, b, c \in \mathcal{D}, a \preceq b \Rightarrow a \otimes c \preceq b \otimes c$  and  $c \otimes a \preceq c \otimes b$ . Furthermore, we write that  $a \prec b$  if  $a \preceq b$  and  $a \neq b$ . A *complete dioid* is a dioid that is closed for infinite sums and such that  $\otimes$  distributes over infinite sums. In a complete dioid,  $\top \in \mathcal{D}$  denotes the unique greatest element of  $\mathcal{D}$ , defined as  $\top = \bigoplus_{x \in \mathcal{D}} x$ . Moreover, the Kleene star of an element  $a \in \mathcal{D}$ , denoted  $a^*$ , is

defined by  $a^* = \bigoplus_{i \in \mathbb{N}_0} a^i$ , where  $a^0 = e$ ,  $a^{i+1} = a \otimes a^i$ , and the greatest lower bound  $\wedge$  is defined by  $a \wedge b = \bigoplus_{\mathcal{D}_{ab}} x$ , where  $\mathcal{D}_{ab} = \{x \in \mathcal{D} \mid x \preceq a \text{ and } x \preceq b\}$ . The greatest lower bound  $\wedge$  is commutative, associative, idempotent, and satisfies the following property:  $\forall a, b \in \mathcal{D}, a \preceq b \Leftrightarrow a \wedge b = a$ .

*Remark 1.* The following equivalence holds:  $\forall a, b, c \in \mathcal{D}, a \succeq b$  and  $a \succeq c \Leftrightarrow a \succeq b \oplus c$ . Indeed,  $(\Leftarrow)$  comes from  $b \oplus c \succeq b, b \oplus c \succeq c$ .  $(\Rightarrow)$  comes from:  $a \succeq b \Leftrightarrow a \oplus b = a, a \succeq c \Leftrightarrow a \oplus c = a$ ; therefore,  $a \oplus (b \oplus c) = (a \oplus b) \oplus c = a \oplus c = a$ , which is equivalent to  $a \succeq b \oplus c$ . Analogously, it is possible to show that:  $a \preceq b$  and  $a \preceq c \Leftrightarrow a \preceq b \wedge c$ .

We can extend operations  $\oplus$  and  $\otimes$  to matrices as follows:  $\forall A, B \in \mathcal{D}^{m \times n}, C \in \mathcal{D}^{n \times p}$

$$(A \oplus B)_{ij} = A_{ij} \oplus B_{ij}, \quad (A \otimes C)_{ij} = \bigoplus_{k=1}^n (A_{ik} \otimes C_{kj}).$$

Furthermore, multiplication between a scalar and a matrix is defined as:  $\forall \lambda \in \mathcal{D}, A \in \mathcal{D}^{m \times n}, (\lambda \otimes A)_{ij} = \lambda \otimes A_{ij}$ . If  $(\mathcal{D}, \oplus, \otimes)$  is a complete dioid, then  $(\mathcal{D}^{n \times n}, \oplus, \otimes)$ , where  $\oplus$  and  $\otimes$  are as defined above, is a complete dioid. In this case, operation  $\wedge$  is defined,  $\forall A, B \in \mathcal{D}^{n \times n}$ , as  $(A \wedge B)_{ij} = A_{ij} \wedge B_{ij}$ . The neutral elements for  $\oplus$  and  $\otimes$  in  $(\mathcal{D}^{n \times n}, \oplus, \otimes)$  are, respectively, matrices  $\mathcal{E}$  and  $E_{\otimes}$ , where  $\mathcal{E}_{ij} = \varepsilon \forall i, j$  and  $E_{\otimes ij} = e$  if  $i = j$ ,  $E_{\otimes ij} = \varepsilon$  if  $i \neq j$ . Furthermore,  $A \preceq B \Leftrightarrow A \oplus B = B \Leftrightarrow \forall i, j A_{ij} \preceq B_{ij}$ .

**Definition II.1** (From [3]). A binary operation  $\odot$  is a *dual multiplication* in the complete dioid  $(\mathcal{D}, \oplus, \otimes)$ , if it is associative, distributes over  $\wedge$ ,  $e$  is its neutral element, and  $\top$  is absorbing for  $\odot$ .

$\odot$  can be extended to matrices as:  $\forall A \in \mathcal{D}^{m \times n}, C \in \mathcal{D}^{n \times p}, \lambda \in \mathcal{D}$

$$(A \odot C)_{ij} = \bigwedge_{k=1}^n (A_{ik} \odot C_{kj}), \quad (\lambda \odot A)_{ij} = \lambda \odot A_{ij};$$

moreover, if  $\odot$  is a dual product in  $(\mathcal{D}, \oplus, \otimes)$ , then its extension to matrices is a dual product in  $(\mathcal{D}^{n \times n}, \oplus, \otimes)$ .

We say that a complete dioid  $(\mathcal{D}, \oplus, \otimes)$  is a *complete idempotent semifield* if,  $\forall a \in \mathcal{D} \setminus \{\varepsilon, \top\}$ , there exists the multiplicative inverse of  $a$ , indicated by  $a^{-1}$ , i.e.,  $a \otimes a^{-1} = a^{-1} \otimes a = e$ . For notational convenience we write  $\varepsilon^{-1} := \top$ , and  $\top^{-1} := \varepsilon$ . Note however that  $\top \otimes \varepsilon = \varepsilon \otimes \top = \varepsilon$ .

**Proposition II.1.** *Let  $(\mathcal{D}, \oplus, \otimes)$  be a complete idempotent semifield. Then operation  $\odot$ , defined as*

$$a \odot b = \begin{cases} a \otimes b & \text{if } a, b \in \mathcal{D} \setminus \{\top\} \\ \top & \text{if } a = \top \text{ or } b = \top \end{cases}, \quad (1)$$

*is a dual product for  $(\mathcal{D}, \oplus, \otimes)$ .*

*Proof.* The only non-trivial property to prove is distributivity of  $\odot$  over  $\wedge$ , i.e.,  $a \odot (b \wedge c) = (a \odot b) \wedge (a \odot c)$  and  $(b \wedge c) \odot a = (b \odot a) \wedge (c \odot a)$ , when  $a, b, c \in \mathcal{D} \setminus \{\varepsilon, \top\}$ ; note that this is a direct consequence of Lemma 4.36 of [1]. ■

In the following two propositions, we will let  $(\mathcal{D}, \oplus, \otimes)$  be a complete idempotent semifield, and  $\odot$  be defined as in (1).

**Proposition II.2.** For all  $A \in (\mathcal{D} \setminus \{\top\})^{n \times n}$ ,  $x \in (\mathcal{D} \setminus \{\top\})^n$ ,

$$A \otimes x \preceq x \Leftrightarrow x \preceq A^\# \odot x, \quad (2)$$

where  $(A^\#)_{ij} := (A_{ji})^{-1}$ .

*Proof.* Let us first show that the proposition holds in the scalar case ( $n = 1$ ), in which  $A^\# = A^{-1}$ . If  $A \in \mathcal{D} \setminus \{\varepsilon, \top\}$ , then, from the facts that  $\otimes$  is order preserving and  $A$  has a multiplicative inverse,  $A \otimes x \preceq x \Rightarrow A^{-1} \otimes A \otimes x = x \preceq A^{-1} \otimes x = A^{-1} \odot x$ , and  $x \preceq A^{-1} \odot x = A^{-1} \otimes x \Rightarrow A \otimes x \preceq A \otimes A^{-1} \otimes x = x$ . In case  $A = \varepsilon$ , the proof comes from the absorbing properties of  $\varepsilon$  over  $\otimes$  and of  $\top$  over  $\odot$ .

In the matrix case, the left-hand side inequality of (2) can be written as  $\forall i \in \{1, \dots, n\}$ ,  $\bigoplus_{j=1}^n A_{ij} \otimes x_j \preceq x_i$ , which, from Remark 1, is equivalent to,  $\forall i, j \in \{1, \dots, n\}$ ,  $A_{ij} \otimes x_j \preceq x_i$ . From the scalar case, the latter expression is equivalent to,  $\forall i, j \in \{1, \dots, n\}$ ,  $x_j \preceq (A_{ij})^{-1} \odot x_i$ , which, using again Remark 1, can be written as  $\forall j \in \{1, \dots, n\}$   $x_j \preceq \bigwedge_{i=1}^n (A_{ij})^{-1} \odot x_i$ ; since the latter inequality is an alternative expression for the right-hand side inequality of (2), the proof is completed. ■

**Proposition II.3.** For all  $A \in (\mathcal{D} \setminus \{\top\})^{n \times n}$ ,  $B \in (\mathcal{D} \setminus \{\varepsilon\})^{n \times n}$  and  $x \in (\mathcal{D} \setminus \{\top\})^n$ ,  $A \otimes x \preceq x \preceq B \odot x$  if and only if  $(A \oplus B^\#) \otimes x \preceq x$ .

*Proof.* The inequalities to the left of the ‘‘if and only if’’ are equivalent to  $A \otimes x \preceq x$  and  $x \preceq B \odot x$ . From Proposition II.2, the second inequality can be written as  $B^\# \otimes x \preceq x$ . Using the first statement of Remark 1, the two inequalities can be combined into:  $A \otimes x \oplus B^\# \otimes x = (A \oplus B^\#) \otimes x \preceq x$ . ■

In the following, we recall the definition of tensor (or Kronecker) product and one of its properties, which holds in *commutative dioids*, i.e., dioids in which the product  $\otimes$  is commutative. Let  $A \in \mathcal{D}^{m \times n}$  and  $B \in \mathcal{D}^{p \times q}$ ; the tensor product  $A \otimes^t B$  is defined as the block matrix

$$A \otimes^t B = \begin{bmatrix} A_{11} \otimes B & \cdots & A_{1n} \otimes B \\ \vdots & & \vdots \\ A_{m1} \otimes B & \cdots & A_{mn} \otimes B \end{bmatrix}.$$

**Proposition II.4** (Mixed product property). Let  $(\mathcal{D}, \oplus, \otimes)$  be a commutative dioid,  $A \in \mathcal{D}^{m \times n}$ ,  $B \in \mathcal{D}^{p \times q}$ ,  $C \in \mathcal{D}^{n \times k}$ ,  $D \in \mathcal{D}^{q \times r}$ . Then  $(A \otimes^t B) \otimes (C \otimes^t D) = (A \otimes C) \otimes^t (B \otimes D)$ .

*Proof.* The proof is analogous (after replacing  $+$  with  $\oplus$  and  $\times$  with  $\otimes$ ) to the one of Lemma 4.2.10 of [11]. ■

### B. Max-plus algebra

The max-plus algebra,  $(\mathbb{R}_{\max}, \oplus, \otimes)$ , is the complete idempotent semifield defined as the set  $\mathbb{R}_{\max}$  endowed with operations  $\max$ , indicated by  $\oplus$ , and  $+$ , indicated by  $\otimes$ . In  $(\mathbb{R}_{\max}, \oplus, \otimes)$ ,  $\varepsilon = -\infty$ ,  $e = 0$ ,  $\top = \infty$ ,  $\wedge$  coincides with the standard  $\min$ , and  $\preceq$  coincides with the standard  $\leq$ . Following Proposition II.1, the dual product  $\odot$  is defined such that it coincides with  $\otimes$ , except when one of its arguments is  $\top$ . Moreover, note that multiplication  $\otimes$  and dual multiplication  $\odot$  are commutative in  $(\mathbb{R}_{\max}, \oplus, \otimes)$ . Since  $(\mathbb{R}_{\max}, \oplus, \otimes)$  is a complete idempotent semifield,  $(\mathbb{R}_{\max}^{n \times n}, \oplus, \otimes)$  is a complete

dioid. If  $A \in \mathbb{R}_{\max}^{n \times n}$  then  $A^\#$  is known as Butkovic conjugate matrix [4] and coincides, in standard algebra, with  $-A^T$ . We will indicate the product between a scalar  $\lambda \in \mathbb{R}$  and a matrix  $A \in \mathbb{R}_{\max}^{m \times n}$ ,  $\lambda \otimes A = \lambda \odot A$ , simply by  $\lambda A$ .

## III. PRECEDENCE GRAPHS

A *directed graph* is a pair  $(N, E)$  where  $N$  is the set of nodes and  $E \subseteq N \times N$  is the set of arcs. Given two nodes  $i, j \in N$  of a directed graph, we say that  $j$  is a downstream node of  $i$  (and  $i$  is an upstream node of  $j$ ) if  $(i, j) \in E$ . A *weighted directed graph* is a 3-tuple  $(N, E, w)$ , where  $(N, E)$  is a directed graph, and  $w: E \rightarrow \mathbb{R}$  is a function that associates a weight  $w((j, i))$  to every arc  $(j, i)$  of the graph.

**Definition III.1** (Precedence graph). Let  $A \in \mathbb{R}_{\max}^{n \times n}$ . The *precedence graph* associated with  $A$  is the weighted directed graph  $\mathcal{G}(A) = (N, E, w)$ , where  $N = \{1, \dots, n\}$ ,  $E$  is defined such that there is an arc  $(j, i) \in E$  from node  $j$  to node  $i$  if and only if  $A_{ij} \neq -\infty$ , and  $w$  is such that  $w((j, i)) = A_{ij}$ , for every arc  $(j, i) \in E$ . If matrix  $A$  depends on some real parameters,  $A = A(\lambda_1, \dots, \lambda_p)$ ,  $\lambda_1, \dots, \lambda_p \in \mathbb{R}$ , we say that  $\mathcal{G}(A)$  is a *parametric precedence graph*.

A *path*  $\rho$  in  $\mathcal{G}(A) = (N, E, w)$  is a sequence of nodes  $(\rho_1, \rho_2, \dots, \rho_{r+1})$ ,  $r \geq 1$ , with arcs from node  $\rho_i$  to node  $\rho_{i+1}$ , such that  $\forall i \in \{1, \dots, r\}$ ,  $(\rho_i, \rho_{i+1}) \in E$ . The length of a path  $\rho$  is denoted by  $|\rho|_L = r$ , and its weight, denoted by  $|\rho|_W$ , is the max-plus product (standard sum) of the weights of its arcs:  $|\rho|_W = \bigotimes_{i=1}^r A_{\rho_i, \rho_{i+1}}$ . A path  $\rho$  is called *circuit* if its initial and final nodes coincide, i.e., if  $\rho_1 = \rho_{|\rho|_L+1}$ . We recall that, if  $A \in \mathbb{R}_{\max}^{n \times n}$ ,  $(A^r)_{ij}$  is equal to the maximum weight of all paths of  $\mathcal{G}(A)$  from node  $j$  to node  $i$  of length  $r$ . We indicate by  $\Gamma$  the set of all precedence graphs that do not contain circuits with positive weight. The following statement connects max-plus linear inequalities with graphs.

**Proposition III.1** (From [1], [8], [4]). Let  $A \in \mathbb{R}_{\max}^{n \times n}$ . Inequality  $A \otimes x \preceq x$  has at least one solution  $x \in \mathbb{R}^n$  if and only if  $\mathcal{G}(A) \in \Gamma$ . Moreover, if  $\mathcal{G}(A) \in \Gamma$ , then  $A^*$  can be computed in  $\mathcal{O}(n^3)$ , and  $\{x \in \mathbb{R}^n \mid A \otimes x \preceq x\} = \{A^* \otimes u \mid u \in \mathbb{R}^n\}$ .

### A. The non-positive circuit weight problem

Given a parametric precedence graph  $\mathcal{G}(A)$ ,  $A = A(\lambda_1, \dots, \lambda_p) \in \mathbb{R}_{\max}^{n \times n}$ , the *Non-positive Circuit weight Problem* (NCP) consists in finding all  $\lambda_1, \dots, \lambda_p \in \mathbb{R}$  such that  $\mathcal{G}(A) \in \Gamma$ . The solution set of the NCP is indicated by  $\Lambda_{\text{NCP}}(A) := \{(\lambda_1, \dots, \lambda_p) \in \mathbb{R}^p \mid \mathcal{G}(A(\lambda_1, \dots, \lambda_p)) \in \Gamma\}$ . Note that, from Proposition III.1,  $\Lambda_{\text{NCP}}(A)$  coincides with the set of  $\lambda_1, \dots, \lambda_p \in \mathbb{R}$  such that the following inequality admits at least one solution  $x \in \mathbb{R}^n$ :

$$A(\lambda_1, \dots, \lambda_p) \otimes x \preceq x. \quad (3)$$

In the following section, we will be particularly interested in a subclass of the NCP, which is described as follows. Let  $P, I, C$  denote three arbitrary  $n \times n$  matrices in  $\mathbb{R}_{\max}$  and let us consider the parametric precedence graph  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$ , in which the weights of the arcs depend on a single parameter  $\lambda \in \mathbb{R}$  in a proportional  $(\lambda P)$ , inverse  $(\lambda^{-1} I)$  and constant

(C) way, in the max-plus sense. The NCP on graph  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$  is then called *Proportional-Inverse-Constant-NCP* (PIC-NCP). In standard algebra, the weight of a generic arc  $(j, i)$  of  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$  has the form  $w((j, i)) = \max(P_{ij} + \lambda, I_{ij} - \lambda, C_{ij})$ .

#### IV. P-TIME EVENT GRAPHS

In this section, P-time event graphs (P-TEGs) are presented following [19].

##### A. General description of P-time event graphs

**Definition IV.1.** An *event graph* is a 4-tuple  $(\mathcal{P}, \mathcal{T}, E, m)$ , where  $(\mathcal{P} \cup \mathcal{T}, E)$  is a directed graph in which the set of nodes is partitioned into the set of places,  $\mathcal{P}$  (represented as circles), and the set of transitions,  $\mathcal{T}$  (represented as bars), the set of arcs  $E$  is such that  $E \subseteq (\mathcal{P} \times \mathcal{T}) \cup (\mathcal{T} \times \mathcal{P})$ , every place has exactly one upstream transition and one downstream transition<sup>1</sup>, and  $m : \mathcal{P} \rightarrow \mathbb{N}_0$  is a map such that  $m(p)$  is the initial marking of place  $p \in \mathcal{P}$ , *i.e.*, the initial number of tokens (represented as dots) in  $p$ .

In an event graph, a transition  $t \in \mathcal{T}$  is enabled if all its upstream places contain at least 1 token, or if there are no upstream places of  $t$ . When a transition is enabled, it can fire: its firing causes the marking of all its upstream places to decrease by 1, and the marking of all its downstream places to increase by 1.

**Definition IV.2** (From [5]). A *P-time event graph* (P-TEG) is a 5-tuple  $(\mathcal{P}, \mathcal{T}, E, m, \iota)$ , where  $(\mathcal{P}, \mathcal{T}, E, m)$  is an event graph, and  $\iota : \mathcal{P} \rightarrow \{[\tau^-, \tau^+] \mid \tau^- \in \mathbb{R}_{\geq 0}, \tau^+ \in (\mathbb{R}_{\geq 0} \cup \{+\infty\})\}$ ,  $\tau^- \leq \tau^+$  is a map that associates to every place  $p \in \mathcal{P}$  a time interval  $\iota(p) = [\tau_p^-, \tau_p^+]$ , where  $\tau_p^-$ , respectively  $\tau_p^+$ , represent minimal, respectively maximal, sojourn times of tokens in place  $p$ .

In addition to the rules stated for event graphs, the dynamics of P-TEGs must respect the following firing rule. After a token enters a place  $p$  at time  $\tau$ , it must remain there at least until time  $\tau + \tau_p^-$  and at most until time  $\tau + \tau_p^+$ . If a token does not leave place  $p$  before or at time  $\tau + \tau_p^+$ , then it is said to be dead. This situation can occur if, for instance, the downstream transition of  $p$  is not enabled in time due to a late arrival of tokens in other upstream places of the transition. We assume that the firing of a transition causes tokens from the upstream places to move instantly to the downstream places; therefore, no delay is associated with the firing of transitions. Note that, since firing of transitions can occur at any time instant that satisfies the bounds associated with every upstream place, the time evolution of the marking of a P-TEG is non-deterministic.

We recall from [15] that every P-TEG  $(\mathcal{P}, \mathcal{T}, E, m, \iota)$  in which there exist places with initial marking greater than 1 can be transformed, without modifying the (non-deterministic) behavior of the original P-TEG, into another one, in which the initial marking of all places is 0 or 1. If

<sup>1</sup>The notion of upstream and downstream places/transitions derives from the definition of upstream and downstream nodes in a directed graph.

$n = |\mathcal{T}|$  is the number of transitions in the original P-TEG, then the number of transitions in the transformed P-TEG is

$$\bar{n} = n + \sum_{p \in \mathcal{P}} \max(0, m(p) - 1) . \quad (4)$$

Hence, without loss of generality, in the remainder of the paper, only P-TEGs in which every place has zero or unitary initial marking are considered, *i.e.*,  $\forall p \in \mathcal{P}, m(p) \in \{0, 1\}$ .

##### B. Dynamics of a P-TEG in the max-plus algebra

In the following, we characterize the time evolution of the marking of a P-TEG as a dynamical system of inequalities in the max-plus algebra. Let  $(\mathcal{P}, \mathcal{T}, E, m, \iota)$  be a P-TEG with  $n$  transitions  $t_1, \dots, t_n \in \mathcal{T}$ . Let the four square matrices  $A^0, A^1 \in (\mathbb{R}_{\geq 0} \cup \{-\infty\})^{n \times n}$ ,  $B^0, B^1 \in (\mathbb{R}_{\geq 0} \cup \{+\infty\})^{n \times n}$  be defined as follows: if there exists a place  $p$  with  $m(p) \in \{0, 1\}$ , upstream transition  $t_j$  and downstream transition  $t_i$ , then  $A_{ij}^{m(p)}$  and  $B_{ij}^{m(p)}$  represent, respectively, the lower and upper bound of interval  $\iota(p)$ , *i.e.*,  $\iota(p) = [A_{ij}^{m(p)}, B_{ij}^{m(p)}]$ ; otherwise, if there is no place  $p$  with initial marking  $\mu \in \{0, 1\}$ , upstream transition  $t_j$  and downstream transition  $t_i$ , then  $A_{ij}^\mu = -\infty$ ,  $B_{ij}^\mu = +\infty$ . Let  $x : \mathbb{N}_0 \rightarrow \mathbb{R}^n$  be a dater function, *i.e.*,  $x_i(k)$  represents the time at which transition  $t_i$  fires for the  $(k+1)$ <sup>st</sup> time. It is natural to require that, for every transition, the  $(k+1)$ <sup>st</sup> firing cannot occur before the  $k$ <sup>th</sup> firing:  $\forall i \in \{1, \dots, n\}, \forall k \in \mathbb{N} x_i(k) \geq x_i(k-1)$ . Hence, we will consider only daters that are non-decreasing in  $k$ .

In order to respect the constraints given by the time bounds  $\iota(p) \forall p \in \mathcal{P}$ , the dater function must satisfy

$$\forall k \in \mathbb{N}_0 \quad \begin{cases} A^0 \otimes x(k) \leq x(k) & \leq B^0 \odot x(k) \\ A^1 \otimes x(k) \leq x(k+1) & \leq B^1 \odot x(k) \end{cases} \quad (5)$$

(see [16] for more details). A non-decreasing trajectory  $\{x(k)\}_{k \in \mathbb{N}_0}$  that satisfies (5) for every  $k \in \mathbb{N}_0$  is said to be *consistent*, or *admissible* for the P-TEG. Note that following a consistent trajectory  $\{x(k)\}_{k \in \mathbb{N}_0}$  guarantees that no token dies, since all time constraints associated with every place of the P-TEG are satisfied.

#### V. PERIODIC TRAJECTORIES AND THE NCP

In this section, we show that the problem of finding all  $\lambda \in \mathbb{R}_{\geq 0}$  such that a P-TEG admits 1-periodic (and, more generally,  $d$ -periodic) trajectories of period  $\lambda$  is equivalent to the PIC-NCP for a particular choice of matrices  $P, I, C$ .

Given a number  $d \in \mathbb{N}$ , we say that a trajectory  $\{x(k)\}_{k \in \mathbb{N}_0}$  is  $d$ -periodic with period  $\lambda \in \mathbb{R}_{\geq 0}$  if, in the max-plus algebra,  $\forall k \in \mathbb{N}_0, x(k+d) = \lambda^d x(k)$ . In standard algebra,  $d$ -periodic trajectories satisfy:  $\forall k \in \mathbb{N}_0, \forall i \in \{1, \dots, n\}, x_i(k+d) = d \times \lambda + x_i(k)$ . We indicate by  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$  the set of all  $\lambda \in \mathbb{R}_{\geq 0}$  for which the P-TEG characterized by matrices  $A^0, A^1, B^0, B^1$  admits a  $d$ -periodic trajectory of period  $\lambda$ , for some  $x(0), x(1), \dots, x(d-1) \in \mathbb{R}^n$ . In the remainder of the section, we consider a P-TEG characterized by matrices  $A^0, A^1, B^0, B^1$ , and we define:  $P := B^{1\sharp}, I := A^1 \oplus E_\otimes, C := A^0 \oplus B^{0\sharp}$ . In the following theorem, we show the connection between  $d$ -periodic trajectories and a variant of the NCP.

**Theorem V.1.** *The set  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$  coincides with  $\Lambda_{\text{NCP}}((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))$ , where  $T_d^P = T_d^P(\lambda)$ ,  $T_d^I = T_d^I(\lambda)$ ,  $T_d^C$  are  $d \times d$  matrices defined as*

$$T_d^P := \begin{bmatrix} -\infty & 0 & -\infty & \cdots & -\infty \\ -\infty & -\infty & 0 & \cdots & -\infty \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\infty & -\infty & -\infty & \cdots & 0 \\ \lambda^d & -\infty & -\infty & \cdots & -\infty \end{bmatrix}, T_d^I := \begin{bmatrix} -\infty & -\infty & \cdots & -\infty & \lambda^{-d} \\ 0 & -\infty & \cdots & -\infty & -\infty \\ -\infty & 0 & \cdots & -\infty & -\infty \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ -\infty & -\infty & \cdots & 0 & -\infty \end{bmatrix}$$

(where expressions  $\lambda^d$  and  $\lambda^{-d}$  are meant in the max-plus sense and correspond, in the standard algebra, respectively to  $d \times \lambda$  and  $-d \times \lambda$ ) and  $T_d^C := E_{\otimes}$ .

*Proof.* Let us rewrite (5) adding conditions  $\forall k \in \mathbb{N}_0$   $x(k+d) = \lambda^d x(k)$  ( $d$ -periodicity),  $x(k+1) \succeq x(k)$  (non-decreasingness), using Propositions II.2 and II.3, and the fact that  $\otimes$  is order preserving:

$$\forall k \in \mathbb{N}_0 \left\{ \begin{array}{l} (A^0 \oplus B^{0\sharp}) \otimes x(k) \preceq x(k) \\ A^1 \otimes x(k) \preceq x(k+1) \\ B^{1\sharp} \otimes x(k+1) \preceq x(k) \\ E_{\otimes} \otimes x(k) \preceq x(k+1) \\ \vdots \\ (A^0 \oplus B^{0\sharp}) \otimes x(k+d-1) \preceq x(k+d-1) \\ \lambda^{-d} A^1 \otimes x(k+d-1) \preceq x(k) \\ \lambda^d B^{1\sharp} \otimes x(k) \preceq x(k+d-1) \\ \lambda^{-d} E_{\otimes} \otimes x(k+d-1) \preceq x(k) \end{array} \right.$$

Using Remark 1, the distributive property of  $\otimes$  over  $\oplus$  and the definition of tensor product, the system above can be written as

$$\forall k \in \mathbb{N}_0 \quad ((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C)) \otimes \tilde{x}(k) \preceq \tilde{x}(k),$$

where  $\tilde{x}(k) := [x^T(k), x^T(k+1), \dots, x^T(k+d-1)]^T \in \mathbb{R}^{dn}$ . Note that, if  $\tilde{x}(0) = \tilde{x}_0$  satisfies the above inequality, from the fact that  $\lambda^d$  admits a multiplicative inverse  $\lambda^{-d}$ , then  $\forall k \in \mathbb{N}_0$ ,  $\tilde{x}(k+1)$  automatically satisfies it. Therefore, there exists an admissible  $d$ -periodic trajectory of period  $\lambda$  if and only if inequality

$$((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C)) \otimes \tilde{x}_0 \preceq \tilde{x}_0, \quad (6)$$

admits a solution  $\tilde{x}_0 \in \mathbb{R}^{dn}$ . Hence, from (3), finding all  $\lambda$ s for which there exists a  $d$ -periodic trajectory of period  $\lambda$  that is consistent for the P-TEG is equivalent to solving the NCP for the parametric precedence graph  $\mathcal{G}((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))$ . ■

*Remark 2.* Given a  $\lambda \in \Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$ , it is possible to characterize, in time  $\mathcal{O}((dn)^3)$ , the set of vectors  $\tilde{x}_0 = [x_0^T, \dots, x_{d-1}^T]^T$  such that the  $d$ -periodic trajectory  $x(0) = x_0, \dots, x(d-1) = x_{d-1}$ ,  $\forall k \in \mathbb{N}_0$ ,  $x(k+d) = \lambda^d x(k)$  is consistent for the P-TEG. Indeed, as discussed in the proof of Theorem V.1, this set coincides with the set of solutions  $\tilde{x}_0 \in \mathbb{R}^{dn}$  to (6), which, from Proposition III.1, is equivalent to  $\{((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))^* \otimes u \mid u \in \mathbb{R}^{dn}\}$ .

Since  $T_1^P = \lambda$ ,  $T_1^I = \lambda^{-1}$ ,  $T_1^C = 0$ , an immediate consequence of Theorem V.1 is that  $\Lambda_{\text{P-TEG}_1}(A^0, A^1, B^0, B^1)$  coincides with the solution set of the PIC-NCP on graph  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$ , i.e.,  $\Lambda_{\text{NCP}}(\lambda P \oplus \lambda^{-1} I \oplus C)$ . As the PIC-NCP can be solved in time complexity  $\mathcal{O}(n^4)$  [20], this

proves that we can compute the set of  $\lambda$ s for which there exists a 1-periodic trajectory with period  $\lambda$  that is consistent for a P-TEG in the same complexity.

On the other hand, since  $T_d^P \otimes^t P$ ,  $T_d^I \otimes^t I$ ,  $T_d^C \otimes^t C \in \mathbb{R}_{\max}^{dn \times dn}$ , it is natural to think that the complexity for computing  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$  would grow with  $d$ . However, the following result gives a surprisingly simple characterization of the admissible periods of  $d$ -periodic trajectories.

**Theorem V.2.** *For all  $d \in \mathbb{N}$ , the set  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$  coincides with  $\Lambda_{\text{P-TEG}_1}(A^0, A^1, B^0, B^1)$ .*

*Proof.* Since every 1-periodic trajectory is also  $d$ -periodic, it is obvious that  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1) \supseteq \Lambda_{\text{P-TEG}_1}(A^0, A^1, B^0, B^1)$ . Therefore, we only need to prove that  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1) \subseteq \Lambda_{\text{P-TEG}_1}(A^0, A^1, B^0, B^1)$ ; from Theorem V.1, this is equivalent to prove that  $\Lambda_{\text{NCP}}((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C)) \subseteq \Lambda_{\text{NCP}}(\lambda P \oplus \lambda^{-1} I \oplus C)$ . In order to do so, in the following we suppose that, for a given  $\lambda \in \mathbb{R}_{\geq 0}$ , there exists a circuit in  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$  from node  $i \in \{1, \dots, n\}$  of length  $h$  and positive weight; we will prove that, as a consequence, there exists a circuit in  $\mathcal{G}((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))$  from node  $i$  of length  $h \times d$  and positive weight. Algebraically, this is the same as proving that, if  $((\lambda P \oplus \lambda^{-1} I \oplus C)^h)_{ii} > 0$ , then  $((((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))^h)_{ii} > 0$ .

By expanding the  $h$ -th power of the trinomial  $\lambda P \oplus \lambda^{-1} I \oplus C$  we observe that, since  $\oplus$  is idempotent, there must exist a certain matrix  $M$  of the form  $M = \lambda^{h_p - h_l} X_1 \otimes \dots \otimes X_h$ , with  $X_1, \dots, X_h \in \{P, I, C\}$ , obtained by multiplying, in some order,  $h_p$  times matrix  $\lambda P$ ,  $h_l$  times matrix  $\lambda^{-1} I$  and  $h_c$  times matrix  $C$ , with  $h_p + h_l + h_c = h$ , such that  $((\lambda P \oplus \lambda^{-1} I \oplus C)^h)_{ii} = M_{ii} > 0$ .

By expanding the  $h$ -th power of the trinomial  $(T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C)$  we get, by idempotency,

$$\begin{aligned} & (((T_d^P \otimes^t P) \oplus (T_d^I \otimes^t I) \oplus (T_d^C \otimes^t C))^h)_{ii} \succeq \\ & \succeq (((T_d^{X_1} \otimes^t X_1) \otimes (T_d^{X_2} \otimes^t X_2) \otimes \dots \otimes (T_d^{X_h} \otimes^t X_h))^d)_{ii}. \end{aligned}$$

Using Proposition II.4 ( $h \times d$ ) - 1 times (after writing the  $d$ -th power as the product of  $d$  terms), the latter expression can be written as

$$((T_d^{X_1} \otimes \dots \otimes T_d^{X_h})^d \otimes (X_1 \otimes \dots \otimes X_h)^d)_{ii}. \quad (7)$$

It is easy to verify that the following properties hold on matrices  $T_d^P$  and  $T_d^I$ :  $(T_d^P)^d = \lambda^d E_{\otimes}$ ,  $(T_d^I)^d = \lambda^{-d} E_{\otimes}$ , and  $T_d^P \otimes T_d^I = T_d^I \otimes T_d^P = T_d^C = E_{\otimes}$ . Therefore,

$$\begin{aligned} (T_d^{X_1} \otimes \dots \otimes T_d^{X_h})^d &= \begin{cases} ((T_d^P)^{h_p - h_l})^d & \text{if } h_p \geq h_l \\ ((T_d^I)^{h_l - h_p})^d & \text{otherwise} \end{cases} \\ &= (\lambda^{h_p - h_l})^d E_{\otimes}, \end{aligned}$$

and (7) simplifies to

$$\begin{aligned} & ((\lambda^{h_p - h_l})^d E_{\otimes} \otimes (X_1 \otimes \dots \otimes X_h)^d)_{ii} = \\ & = ((\lambda^{h_p - h_l} X_1 \otimes \dots \otimes X_h)^d)_{ii} = (M^d)_{ii} \succeq M_{ii} > 0. \quad \blacksquare \end{aligned}$$




Fig. 1. Example of P-TEG

### A. Example

In the following, we illustrate the previous theorems by a simple example taken from [19]. Let us consider the P-TEG shown in Fig. 1, characterized by matrices

$$A^0 = \begin{bmatrix} -\infty & -\infty & -\infty \\ 2 & -\infty & -\infty \\ 6 & 0.5 & -\infty \end{bmatrix}, A^1 = \begin{bmatrix} -\infty & 0 & -\infty \\ -\infty & -\infty & 0.5 \\ -\infty & -\infty & 0 \end{bmatrix},$$

$$B^0 = \begin{bmatrix} +\infty & +\infty & +\infty \\ 3 & +\infty & +\infty \\ +\infty & +\infty & +\infty \end{bmatrix}, B^1 = \begin{bmatrix} +\infty & +\infty & +\infty \\ +\infty & +\infty & +\infty \\ +\infty & +\infty & 4 \end{bmatrix}.$$

Let  $P = B^{1\sharp}$ ,  $I = A^1 \oplus E_{\otimes}$ ,  $C = A^0 \oplus B^{0\sharp}$ . From Theorems V.1 and V.2, for all  $d \in \mathbb{N}$ ,  $\Lambda_{\text{P-TEG}_d}(A^0, A^1, B^0, B^1)$  coincides with the set of  $\lambda$ s for which the parametric precedence graph  $\mathcal{G}(\lambda P \oplus \lambda^{-1} I \oplus C)$ , shown in Fig. 2, does not contain circuits with positive weight. Using the algorithm presented in [20], we obtain:  $\Lambda_{\text{NCP}}(\lambda P \oplus \lambda^{-1} I \oplus C) = \{\lambda \mid 3.5 \leq \lambda \leq 4\}$ . For instance, let us consider  $\lambda = 4$  and  $d = 2$ ; from Remark 2, the 2-periodic trajectory  $x(0) = x_0 = [0, 2.5, 6]^T$ ,  $x(1) = x_1 = [3.5, 6.5, 10]^T$ ,  $\forall k \in \mathbb{N}_0$ ,  $x(k+2) = 4^2 x(k) = 8x(k)$  is consistent for the P-TEG, since, for  $u = [0, 0, 0, 0, 0, 0]^T$ , we have

$$T_2^P = \begin{bmatrix} -\infty & 0 \\ 8 & -\infty \end{bmatrix}, T_2^I = \begin{bmatrix} -\infty & -8 \\ 0 & -\infty \end{bmatrix}, T_2^C = \begin{bmatrix} 0 & -\infty \\ -\infty & 0 \end{bmatrix},$$

$$((T_2^P \otimes^t P) \oplus (T_2^I \otimes^t I) \oplus (T_2^C \otimes^t C))^* \otimes u =$$

$$= \begin{bmatrix} 0 & -3 & -6.5 & -4.5 & -7.5 & -10.5 \\ 2.5 & 0 & -3.5 & -1.5 & -4.5 & -7.5 \\ 6 & 3 & 0 & 2 & -1 & -4 \\ 3.5 & 0.5 & -2.5 & 0 & -3 & -6.5 \\ 6.5 & 3.5 & 0.5 & 2.5 & 0 & -3.5 \\ 10 & 7 & 4 & 6 & 3 & 0 \end{bmatrix} \otimes \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} x_0 \\ x_1 \end{bmatrix}.$$

## VI. CONCLUSIONS

The present paper shows the connection between two problems: the PIC-NCP, and the problem of finding all periods of the admissible  $d$ -periodic trajectories for a P-TEG, for all  $d \in \mathbb{N}$ . Indeed, based on their description in the max-plus algebra, we formulate the second problem as a particular instance of the first one. Moreover, we prove that the solution set of the second problem is invariant with respect to  $d$ . In a related paper [20], we have suggested an algorithm that solves the PIC-NCP in time complexity  $\mathcal{O}(n^4)$ , where  $n$  is the number of nodes in the graph; this proves that the existence of consistent  $d$ -periodic trajectories can be checked in strongly polynomial time  $\mathcal{O}(n^4)$ , where  $n$  is the number of transitions of a P-TEG with initially 0 or 1 token per place. Since every P-TEG can be transformed into




Fig. 2. Parametric precedence graph associated with the P-TEG of Fig. 1

one with at most 1 initial token per place, a consequence is that the same property can be checked in generic P-TEGs in time complexity  $\mathcal{O}(\bar{n}^4)$ , where  $\bar{n}$  is defined in (4).

## REFERENCES

- [1] François Baccelli, Guy Cohen, Geert Jan Olsder, and Jean-Pierre Quadrat. *Synchronization and linearity: an algebra for discrete event systems*. John Wiley & Sons Ltd, 1992.
- [2] Tassadit Becha, Redouane Kara, S Collart Dutilleul, and Jean Jacques Loiseau. Modelling, analysis and control of electroplating line modelled by P-time event graphs. *IFAC Proceedings Volumes*, 46(24):311–316, 2013.
- [3] Thomas Brunsch, Laurent Hardouin, Carlos Andrey Maia, and Jörg Raisch. Duality and interval analysis over idempotent semirings. *Linear algebra and its applications*, 437(10):2436–2454, 2012.
- [4] Peter Butkovič. *Max-linear systems: theory and algorithms*. Springer Science & Business Media, 2010.
- [5] Soizick Calvez, Pascal Ayyalinc, and Wael Khansa. P-time Petri nets for manufacturing systems with staying time constraints. *IFAC Proceedings Volumes*, 30(6):1487–1492, 1997.
- [6] Philippe Declerck. Extremum cycle times in time interval models. *IEEE Transactions on Automatic Control*, 63(6):1821–1827, 2017.
- [7] Philippe Declerck. Critical subsystems in time interval models. *Discrete Event Dynamical Systems*, 2020.
- [8] Tibor Gallai. Maximum-minimum Sätze über Graphen. *Acta Mathematica Hungarica*, 9(3-4):395–434, 1958.
- [9] Laurent Hardouin, Bertrand Cottenceau, Ying Shang, and Jörg Raisch. Control and state estimation for max-plus linear systems. *Foundations and Trends® in Systems and Control*, 6(1):1–116, 2018.
- [10] Bernd Heidergott, Geert Jan Olsder, and Jacob Van Der Woude. *Max Plus at work: modeling and analysis of synchronized systems: a course on Max-Plus algebra and its applications*. Princeton University Press, 2014.
- [11] Roger A Horn and Charles R Johnson. *Topics in matrix analysis*. Cambridge University Press, 1991.
- [12] Richard M Karp and James B Orlin. Parametric shortest path algorithms with an application to cyclic staffing. *Discrete Applied Mathematics*, 3(1):37–45, 1981.
- [13] Ja-Hee Kim and Tae-Eog Lee. Schedulability analysis of time-constrained cluster tools with bounded time variation by an extended Petri net. *IEEE Transactions on Automation Science and Engineering*, 5(3):490–503, 2008.
- [14] Eugene Levner and Vladimir Kats. A parametric critical path problem and an application for cyclic scheduling. *Discrete Applied Mathematics*, 87(1-3):149–158, 1998.
- [15] Pavel Špaček and Jan Komenda. Analysis of cycle time in interval P-time event graphs in dioid algebras. *IFAC-PapersOnLine*, 50(1):13461–13467, 2017.
- [16] Pavel Špaček, Jan Komenda, and Sébastien Lahaye. Analysis of P-time event graphs in (max,+) and (min,+) semirings. *International Journal of Systems Science*, 0(0):1–16, 2021.
- [17] Pavel Špaček, Marie-Ange Manier, and Abdellah El Moudni. Control of an electroplating line in the max and min algebras. *International Journal of Systems Science*, 30(7):759–778, 1999.
- [18] Neal E Young, Robert E Tarjant, and James B Orlin. Faster parametric shortest path and minimum-balance algorithms. *Networks*, 21(2):205–221, 1991.
- [19] Davide Zorzenon, Jan Komenda, and Jörg Raisch. Bounded consistency of P-time event graphs. In *2020 59th IEEE Conference on Decision and Control (CDC)*, pages 79–85, 2020.
- [20] Davide Zorzenon, Jan Komenda, and Jörg Raisch. The non-positive circuit weight problem in parametric graphs: a fast solution based on dioid theory. 2021. Published on arXiv, available at <https://arxiv.org/abs/2102.12264>.