

ON THE DESCRIPTIVE COMPLEXITY OF HOMOGENEOUS CONTINUA

PAWEŁ KRUPSKI

Dedicated to Professor Jerzy Mioduszewski on his 90th birthday

ABSTRACT. It is shown that the family of all homogeneous continua in the hyperspace of all subcontinua of any finite-dimensional Euclidean cube or the Hilbert cube is an analytic subspace of the hyperspace which contains a topological copy of the linear space $c_0 = \{(x_k) \in \mathbb{R}^\omega : \lim x_k = 0\}$ as a closed subset.

A continuum X is homogeneous if

- (h) for every two points $x, y \in X$ there is a homeomorphism

$$h : X \rightarrow X \text{ such that } h(x) = y.$$

Let $I = [0, 1]$ and $2 \leq n \leq \omega$. Given a Polish space (X, d) , let 2^X be the hyperspace of all nonempty compact subsets of X with the Hausdorff metric H_d and $C(X) \subset 2^X$ be its closed subspace of all subcontinua in X . It is known that 2^X is a Polish space and if X is compact then 2^X is compact. We use standard notation $\text{Lim } A_k = A$ if $H_d(A_k, A) \rightarrow 0$ in the metric space $(2^X, H_d)$. Denote by $\mathcal{H}(I^n)$ the subspace of $C(I^n)$ consisting of all homogeneous continua in I^n .

In 2003, D. P. Bellamy raised the problem of determining the descriptive complexity of $\mathcal{H}(I^n)$. For $n = 2$, it is now slightly better understood due to L. C. Hoehn and L. G. Oversteegen [5] who settled the long-standing conjecture that there are four topological types of homogeneous plane continua: a point, a circle, the pseudo-arc and the circle of pseudo-arcs. It is also known that the exact Borel class of the family of all simple closed curves in I^n is $F_{\sigma\delta}$ in $C(I^n)$ [6], the family of all pseudo-arcs in I^n is dense G_δ in $C(I^n)$ [1] and the family of all circles of pseudo-arcs in I^n is a Borel subset of $C(I^n)$ [11] (unfortunately, the actual Borel class of the latter family is unknown). So, we conclude that $\mathcal{H}(I^2)$ is Borel in $C(I^2)$.

In higher dimensions, there are much more types of homogeneous continua. The Borel complexity of some of them, the most popular ones, was studied in [8, 9] where some special characterizations of the families were employed. For example, the family of Menger universal curves is exactly $F_{\sigma\delta}$ in $C(I^n)$ for $n \geq 3$, the family of all solenoids is a Borel subset of $C(I^n)$ for $n \geq 3$ but we do not know its exact Borel class, similarly for the family of solenoids of pseudo-arcs or Menger curves of pseudo-arcs.

These particular results may suggest that $\mathcal{H}(I^n)$ is a Borel subset of $C(I^n)$. A naive direct analysis of formula (h) gives the projective class Π_2^1 for $\mathcal{H}(I^n)$. In this

Date: January 1, 2022.

2020 Mathematics Subject Classification. Primary 54H05; Secondary 54F16.

Key words and phrases. analytic set, homogeneous continuum, hyperspace.

note we reduce the complexity of $\mathcal{H}(I^n)$ by showing that it is a Σ_1^1 -subset of $C(I^n)$ which is not $G_{\delta\sigma}$.

Recall from [10] that, given compact metric spaces X, Y , the set $P(X, Y) \subset 2^{X \times Y}$ of partial continuous maps $f : Z \rightarrow Y$, $Z \in 2^X$ (the maps are identified with their graphs), topologized by the convergence

(*) $f_k \rightarrow f$ if and only if $\text{Lim}(\text{dom } f_k) = \text{dom } f$ and

$$\lim f_k(x_k) = f(x) \text{ whenever } x_k \in \text{dom } f_k, x \in \text{dom } f \text{ and } \lim x_k = x,$$

($\text{dom } g$ denotes the domain of a partial map $g \in P(X, Y)$), is a Polish space and the subspace $H_p(X, Y) \subset P(X, Y)$ consisting of all partial homeomorphisms is a G_δ subset of $P(X, Y)$, so it is also a Polish space.

Let $\text{Aut}_p(I^n)$ denote the subspace of $H_p(I^n, I^n)$ of all partial autohomeomorphisms $h : X \rightarrow X$ where $X \in C(I^n)$. Given any $X \in 2^{I^n}$, if $H(X)$ is the space of all autohomeomorphisms $h : X \rightarrow X$ with the uniform convergence, then $H(X)$ is a closed subspace of $\text{Aut}_p(I^n)$

By a straightforward verification, we get the following

Lemma 1. *$\text{Aut}_p(I^n)$ is a closed subset of $H_p(I^n, I^n)$, so it is a Polish space.*

Part (1) of the next lemma follows from the Effros' theorem [4]; part (2) is a consequence of part (1) and the fact that open maps on completely metrizable spaces are compact-covering [3, Problem 5.5.11].

Lemma 2. *If X is a homogeneous compactum, then*

- (1) *for each $a \in X$, the valuation map $\phi_a : H(X) \rightarrow X$, $\phi_a(h) = h(a)$, is an open surjection.*
- (2) *For every closed $K \subset X$, there exists a compact $C \subset H(X)$ such that $\phi_a(C) = K$.*

Let $\mathcal{F} = \{ (X, a) \in C(I^n) \times I^n : a \in X, X \in \mathcal{H}(I^n) \}$. In view of Lemma 2, the following equation is obvious but it serves as a key observation in our evaluation of the complexity of $\mathcal{H}(I^n)$.

$$(\mathcal{F}) \quad \mathcal{F} = \{ (X, a) \in C(I^n) \times I^n : a \in X, \exists C \in 2^{\text{Aut}_p(I^n)} C \subset H(X), \phi_a(C) = X \}.$$

Lemma 3. *The set $\{(C, X) \in 2^{\text{Aut}_p(I^n)} \times C(I^n) : C \subset H(X)\}$ is closed in $2^{\text{Aut}_p(I^n)} \times C(I^n)$.*

Proof. Let $C_k \subset H(X_k)$, $C_k \rightarrow C$ in $2^{\text{Aut}_p(I^n)}$ and $X_k \rightarrow X$ in $C(I^n)$. In order to show that $C \subset H(X)$, take $h \in C$. There exist $h_k \in C_k$ such that $h_k \rightarrow h$ in $\text{Aut}_p(I^n)$ which means, by (*), that $\text{dom } h = \text{Lim}(\text{dom } h_k) = \text{Lim } X_k = X$. Since h is assumed to be in $\text{Aut}_p(I^n)$, we get $h \in H(X)$. \square

Lemma 4. *The set*

$$\{(C, X, a) \in 2^{\text{Aut}_p(I^n)} \times C(I^n) \times I^n : a \in X, C \subset H(X), \phi_a(C) = X\}$$

is closed in $2^{\text{Aut}_p(I^n)} \times C(I^n) \times I^n$.

Proof. Suppose $a_k \in X_k$, $C_k \subset H(X_k)$, $\phi_{a_k}(C_k) = X_k$ and $a_k \rightarrow a$, $X_k \rightarrow X$, $C_k \rightarrow C$ in respective spaces. Then $a \in X$ and $C \subset H(X)$ by Lemma 3 which implies the inclusion $\phi_a(C) \subset X$.

In order to show the converse inclusion, let $x \in X$. There exist a sequence $x_k \in X_k$ converging to x and a sequence $h_k \in C_k$ such that $h_k(a_k) = x_k$. The set $C \cup \bigcup_{k \in \omega} C_k$ being compact, there is a convergent subsequence (h_{k_i}) . Denote by h its limit in $\text{Aut}_p(I^n)$. Then

$$\text{dom } h = \text{Lim}_i (\text{dom } h_{k_i}) = \text{Lim}_i X_{k_i} = X$$

and

$$\lim_i x_{k_i} = \lim_i h_{k_i}(a_{k_i}) = h(a).$$

It follows that $h \in C$ and $\phi_a(h) = x$. □

Lemma 5. *The set \mathcal{F} is Σ_1^1 in $C(I^n) \times I^n$.*

Proof. Refer to equation (F) and observe that \mathcal{F} is the projection of the closed subset $\{(C, X, a) : a \in X, C \subset H(X), \phi_a(C) = X\}$ of the Polish space $2^{\text{Aut}_p(I^n)} \times C(I^n) \times I^n$ into the $C(I^n) \times I^n$. □

Theorem 6. *The space $\mathcal{H}(I^n)$ is a Σ_1^1 -subset of the hyperspace $C(I^n)$ and $\mathcal{H}(I^n)$ contains a closed homeomorphic copy of $c_0 = \{(x_k) \in \mathbb{R}^\omega : \lim x_k = 0\}$. Consequently, $\mathcal{H}(I^n)$ is not $G_{\delta\sigma}$.*

Proof. $\mathcal{H}(I^n)$ is the projection of \mathcal{F} into $C(I^n)$, therefore it is Σ_1^1 , by Lemma 5. The second part of the theorem follows directly from the proof of an analogous fact in [6, Theorem 2 (2)] (see also [7]). For reader's convenience, we recall a (slightly modified) simple construction of an embedding $f : I^\omega \rightarrow C(I^2)$ such that $f(x)$ is a simple closed curve if $x = (x_k) \in \hat{c}_0 := \{(x_k) \in I^\omega : \lim x_k = 0\}$ and $f(x)$ is not homogeneous otherwise. Let A_k be the segment in I^2 from the point $(1/k, 0)$ to $(1/(k+1), x_k/2)$ and B_k be the segment from $(1/(k+1), x_k/2)$ to $(1/(k+1), 0)$. Define

$$f(x) = (\{0\} \times I) \cup (I \times \{1\}) \cup (\{1\} \times I) \cup \bigcup_k (A_k \cup B_k).$$

Now, $f(\hat{c}_0) = f(I^\omega) \cap \mathcal{H}(I^2)$, hence $f(\hat{c}_0)$ is a closed subset of $\mathcal{H}(I^2)$. Clearly, we can regard f as an embedding in $C(I^n)$ for $n \geq 2$, so that $f(\hat{c}_0) = f(I^\omega) \cap \mathcal{H}(I^n)$. It is also known that \hat{c}_0 is homeomorphic to c_0 because the spaces are both characterized as $F_{\sigma\delta}$ -absorbers (see [2]). Thus, c_0 embeds in $\mathcal{H}(I^n)$ as a closed subset. It follows that the space $\mathcal{H}(I^n)$ is not $G_{\delta\sigma}$ in $C(I^n)$, because otherwise its closed subset $f(\hat{c}_0)$ would be $G_{\delta\sigma}$ in $C(I^n)$, so an absolute $G_{\delta\sigma}$ -set which is not possible since c_0 is a classical example of an absolute $F_{\sigma\delta}$ -set which is not absolute $G_{\delta\sigma}$. □

Acknowledgments. The author thanks the referee for careful reading and corrections.

REFERENCES

1. R. H. Bing, Concerning hereditarily indecomposable continua, Pacific J. Math. 1 (1951), 43–51.
2. J. J. Dijkstra, J. van Mill and J. Mogilski, The space of infinite-dimensional compacta and other topological copies of $(l_f^2)^\omega$, Pacific J. Math. 152 (1992), 255–273.
3. R. Engelking, General topology, Heldermann, Berlin, 1989.
4. E. G. Effros, Transformation groups and C^* -algebras, Ann. of Math. (2) 81 (1965), 38–55.
5. L. C. Hoehn and L. G. Oversteegen, A complete classification of homogeneous plane continua. Acta Math. 216 (2016), no. 2, 177–216.

6. P. Krupski, Hyperspaces of various locally connected subcontinua, *Acta Univ. Carolin. Math. Phys.*, 40, no. 2 (1999), 79–83.
7. P. Krupski, Errata to: "Hyperspaces of various locally connected subcontinua". *Acta Univ. Carolin. Math. Phys.* 41 (2000), no. 1, 81.
8. P. Krupski, Hyperspaces of universal curves and 2-cells are true $F_{\sigma\delta}$ -sets, *Colloq. Math.* 91 (2002), 91–98.
9. P. Krupski, Families of continua with the property of Kelley, arc continua and curves of pseudo-arcs, *Houston J. Math.* 30, no. 2 (2004), 459–482.
10. K. Kuratowski, Sur l'espace des fonctions partielles, *Ann. Mat. Pura Appl.* (4) 40 (1955), 61–67.
11. C. Ryll-Nardzewski, On Borel measurability of orbits. *Fund. Math.* 56 (1964), 129–130.

Email address: `pawel.krupski@pwr.edu.pl`

FACULTY OF PURE AND APPLIED MATHEMATICS, WROCLAW UNIVERSITY OF SCIENCE AND TECHNOLOGY, WYBRZEŻE WYSPIAŃSKIEGO 27, 50-370 WROCLAW, POLAND.