

# Unitarity of Minkowski non-local theories made explicit

Alexey S. Koshelev<sup>1</sup> and Anna Tokareva<sup>2</sup>

<sup>1</sup>*Departamento de Física, Centro de Matemática e Aplicações (CMA-UBI),  
Universidade da Beira Interior, 6200 Covilhã, Portugal*

<sup>2</sup>*Department of Physics, University of Jyväskylä, P.O.Box 35 (YFL), FIN-40014, Finland &  
Institute for Nuclear Research of Russian Academy of Sciences, 117312 Moscow, Russia &  
Helsinki Institute of Physics (HIP), University of Helsinki, P.O. Box 64, 00014, Finland*

## Abstract

In this work we explicitly show that the perturbative unitarity of analytic infinite derivative (AID) scalar field theories can be achieved using a modified prescription for computing scattering amplitudes. The crux of the new prescription is the analytic continuation of a result obtained in the Euclidean signature to the Minkowski external momenta. We explicitly elaborate an example of a non-local  $\phi^4$  model for various infinite derivative operators. General UV properties of amplitudes in non-local theories are discussed.

## 1 Introduction

Higher-Derivative theories appear in different contexts [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21] from pure phenomenological models to fundamental theories with the notable example of string field theory [22, 23, 24, 25] which belongs to the particular interesting class of models featuring analytic infinite derivative (AID) operators entering Lagrangians and acting as form factors. Analyticity of these operators at low momenta guarantees a smooth local IR limit and prompts for the UV modification of respective theories.

A model example providing such a higher derivative modification can be readily written as follows:

$$S = \int d^4x \left[ \frac{1}{2} \phi(\square - m^2) f(\square)^{-1} \phi - \frac{\lambda}{4!} \phi^4 \right] \quad (1)$$

where  $\square$  is the d'Alembertian operator. This action is the subject of our consideration in this paper which is aimed at illustrating explicitly that the unitarity can be maintained in this class of models. Having the metric signature fixed as  $(-+++)$  the above Lagrangian describes a normal non-ghost field provided  $f = 1$  which is also not a tachyon as long as

$m^2 > 0$ . Presence of extra derivatives may generically spoil the model by the appearance of ghosts in the spectrum. A simple way to avoid this is to demand that  $f(\square)^{-1}$  is an exponent of an entire function. Then the system has no new degrees of freedom as there are no finite poles on the complex plane and consequently there are no finite poles of the propagator apart from the already existing one at the point  $m^2$ . This picture seems good especially given that higher-derivative factor can easily provide better convergence of loop integrals.

Here exactly a serious issue arises because an exponent of an entire function must have an essential singularity at complex infinity making the use of the Wick rotation unjustified [26]. Indeed, the latter requires that the propagator has a pole at the complex infinity, not an essential singularity and this cannot be achieved in our model.

The situation, however, has a resolution presented in the paper by Pius and Sen [27] for the class of AID theories originating from the string field theory. The resolution consists of certain modifications to the prescription of how the loop integrals are computed. Namely specially designed integration contours for integrals over the loop momenta are prescribed. It is interesting to note that related idea was presented many years before by Efimov [2]. To be phrased in short, the idea of Efimov is to perform all the inner loop integrals assuming the Euclidean signature of all the momenta. After a result is obtained one should continue analytically this result for the computed amplitude to Minkowski signature for all external momenta. This procedure was claimed to gain unitary scattering amplitudes.

In the present work, using the simplest example of the fish diagram in  $\phi^4$  theory we illustrate that the idea by Efimov in fact leads to the same definition of the amplitude as the method elaborated by Pius and Sen<sup>1</sup>. The latter provides rules how to account for the poles at the complex plane when integrating over loop momenta. These rules are actually the same as in local theories, since non-local factors do not lead to new poles. Let us also mention that the definition of amplitudes by formal application of the familiar Wick rotation (which is the same as doing the analytic continuation of the result upon completion of loop computations) yields a well-defined local limit. In this limit, the normal Wick rotation is recovered, as well as the unitarity of a local theory.

In our analysis we first we provide an explicit example of computations fixing  $f(\square) = \exp(\alpha\square)$  which already comprise a highly non-trivial set of formulae and then move on by showing that one can integrate out internal momenta at one-loop level for an arbitrary form-factor  $f(\square)$ . This leads us to an extensive discussion on UV properties of the amplitudes in non-local theories. We then end up by conclusion and outline.

## 2 Fate of the Wick rotation

In a local quantum field theory the amplitudes can be defined in Minkowski space-time within the familiar rule for the poles. Than it is convenient to do a Wick rotation and go to Euclidean momenta. In non-local theory, however, it is often problematic to define the Minkowski amplitude because the corresponding integrals may diverge. In this Section we

---

<sup>1</sup>See also [28, 29].

show that there is no feasible way to choose a special non-local form-factor which would allow for obtaining convergent Minkowski amplitudes.

Let us start with the simplest tadpole correction to the propagator depicted in Fig. 1.




Figure 1: Tadpole one-loop contribution to the propagator in  $\phi^4$  theory.

Formally, in Minkowski space we can write

$$\mathcal{A} = \int \frac{d^4 k f(k_0^2 - \vec{k}^2)}{k_0^2 - \vec{k}^2 - m^2} \quad (2)$$

Then, we can do the Wick rotation and obtain the Euclidean integral,

$$\mathcal{A}_E = -i \int \frac{d^4 k_E f(-k_{0E}^2 - \vec{k}^2)}{k_{0E}^2 + \vec{k}^2 + m^2} \quad (3)$$

where  $k_{0E} = -ik_0$ . This integral can be made manifestly convergent due to an appropriate choice of the function  $f(\square)$ . However, we can not conclude that  $\mathcal{A} = \mathcal{A}_E$  unless we check that the integral over an infinite arc

$$\mathcal{A}_c = \int_0^{\frac{\pi}{2}} \frac{d^3 k R e^{i\theta} d\theta f(R^2 e^{2i\theta} - \vec{k}^2)}{R^2 e^{2i\theta} - \vec{k}^2 - m^2} \quad (4)$$

vanishes in the limit  $R \rightarrow \infty$ . In this limit,

$$\begin{aligned} \mathcal{A}_c^\infty &= \int_0^{\frac{\pi}{2}} \frac{R e^{i\theta} d\theta f(R^2 e^{2i\theta})}{R^2 e^{2i\theta}} \\ &= \int_r^R \frac{i f(-z^2)}{z^2} + \int_r^R \frac{f(z^2)}{z^2} + \int_0^{\frac{\pi}{2}} f(0) \frac{e^{-i\theta}}{r} d\theta \\ &= \int_r^R \frac{f(0)(i+1) - f(z^2) - i f(-z^2)}{z^2} \end{aligned} \quad (5)$$

If  $f(\square) = 1$  than this integral is zero which makes the Wick rotation consistent in a local theory. However, in non-local models the situation is less obvious. If  $\mathcal{A}_c^\infty = \text{const} < \infty$  than  $\mathcal{A}_c = \int \mathcal{A}_c^\infty d^3 \vec{k}$  is divergent which means that the Minkowski amplitude is divergent too. In principle it is possible to choose  $f(\square)$  in such a way that the condition

$$\int \frac{(f(0)(i+1) - f(z^2) - i f(-z^2)) dz}{z^2} = 0 \quad (6)$$

is satisfied. However, when considering other diagrams, we obtain other conditions, finally an infinite set of similar conditions on  $f(\square)$  which a priori would not be resolved simultaneously. We thus conclude, that it is nearly impossible to define non-local amplitudes in Minkowski space in a standard way unless a mysterious combination of a form-factor and the theory potential can be found such that an infinite tower of conditions allowing the Wick rotation is resolved. Independently of the form of the non-local propagator, the familiar Wick rotation can not be done. Therefore, another way to define physical non-local scattering amplitudes is required.

### 3 Unitarity of the fish diagram

As we have shown just before, the definition of physical amplitudes in non-local theories is a tricky point which most likely cannot be resolved using standard local field theory prescriptions by just making an appropriate choice of the non-local propagator. We thus turn to the method elaborated in papers [2, 27].

We are going to study the procedure using the so-called fish diagram in non-local theory (1). This diagram depends on external momenta in a combination  $p = p_1 + p_2$  but choosing the center of mass reference frame we can reduce the number of variables to only one scalar  $p_0$ . The corresponding Euclidean matrix element would be defined as,

$$\mathcal{M}_E = -\frac{\lambda^2}{32\pi^4} I(p_E) \quad (7)$$

The physical matrix element is to be defined as an analytic continuation of  $\mathcal{M}_E(p_E)$  to Minkowski momenta  $p$ .




Figure 2: One-loop contribution to  $2 \rightarrow 2$  scattering in  $\phi^4$  theory. Here only  $s$ -channel contribution is shown,  $p = p_1 + p_2$ .

Here  $I(p_E)$  is defined as Euclidean integral

$$I(p_E) = \int d^4 k_E \frac{f(k_E^2) f((p_E - k_E)^2)}{(k_E^2 + m^2)((p_E - k_E)^2 + m^2)} \quad (8)$$

The integral can be rewritten as

$$I(p_E) = 2 \int \frac{f(k_E^2) f((p_E - k_E)^2) d^4 k_E}{p_{0E}(k_E^2 + m^2)(p_{0E} - 2k_{0E})} \quad (9)$$

This integral has a pole at  $k_{0E} = p_{0E}/2$ . This results in the non-zero imaginary part,

$$I(p_E) = -\pi i \int \frac{(f(p_{0E}^2/4 + \vec{k}^2))^2 d^3 \vec{k}}{p_{0E}(\vec{k}^2 + m^2 + p_{0E}^2/4)} + P.V. \left( 2 \int \frac{f(k_E^2) f((p_E - k_E)^2) d^4 k_E}{p_{0E}(k_E^2 + m^2)(p_{0E} - 2k_{0E})} \right) \quad (10)$$

where *P.V.* stands for the Principal Value definition of the integral. Let us now continue analytically the function of  $p_E$  to the physical momenta. Thus, we change  $p_{0E} \rightarrow -ip_0$ ,

$$I(p) = -\pi \int \frac{(f(\vec{k}^2 - p_0^2/4))^2 d^3 \vec{k}}{p_0(\vec{k}^2 + m^2 - p_0^2/4 - i\epsilon)} - 2 \int \frac{f(k^2) f((-ip_0 - k_0)^2 - \vec{k}^2) d^4 k}{p_0(k^2 + m^2)(p_0 - 2ik_0)} \quad (11)$$

This integral is real for  $p_0^2 < 4m^2$ . In the opposite case, the first part would have an imaginary part, physically corresponding to the possibility of a particle production. The corresponding expression is,

$$\text{Im } I(p) = -\pi^3 \frac{f(m^2)^2 \sqrt{p_0^2 - 4m^2}}{p_0} \theta(p_0^2 - 4m^2) \quad (12)$$

Then, taking into account that non-local function should be normalized in such a way that  $f(m^2) = 1$  to preserve the value of the residue of the propagator, we recover the familiar result for the local theory,

$$\text{Im } \mathcal{M} = \frac{\lambda^2}{32\pi} \frac{\sqrt{p_0^2 - 4m^2}}{p_0} \theta(p_0^2 - 4m^2). \quad (13)$$

It is thus obvious that one observes the canonical optical theorem statement which tells us that the imaginary part of some Feynman graph is equivalent to the product of expressions corresponding to parts in which a given graph would be cut.

We can demonstrate that the same result for the amplitude can be obtained if one take the formal definition of non-local amplitude in Minkowski space,

$$\mathcal{M} = -i \frac{\lambda^2}{32\pi^4} I(p) \quad (14)$$

Here

$$I(p) = \int d^4 k \frac{f(k^2) f((p - k)^2)}{(k^2 - m^2 + i\epsilon)((p - k)^2 - m^2 + i\epsilon)} \quad (15)$$

and this integral over Minkowski momenta is divergent due to the inevitable presence of the essential singularity of function  $f(\square)$  and consequently some growth direction at infinity as explained in the previous Section. However, applying formally the Wick rotation like in a local theory we arrive to the convergent integral. This integral can be written in the following form,

$$I(p) = -2 \int \frac{f(k^2) f((p_M - k_E)^2) d^4 k_E}{p_0(k_E^2 - m^2 + i\epsilon)(k_0 - p_0/2 + i\epsilon)} \quad (16)$$

where we have not changed  $k_0$  to  $ik_{0E}$  in one instance to account the pole on the real axis of  $k_0$  properly. Namely, we first have to transform

$$\frac{1}{k_0 - p_0/2 + i\epsilon} = -i\pi\delta(k_0 - p_0/2) + P.V. \left( \frac{1}{k_0 - p_0/2 + i\epsilon} \right) \quad (17)$$

and then change in the second term  $k_0 \rightarrow ik_{0E}$ . The other pole is not pinched by the contour. After that we arrive at

$$I(p) = \pi i \int \frac{d^3\vec{k}(f(p_0^2/4 - \vec{k}^2))^2}{p_0(\vec{k}^2 + m^2 - p_0^2/4 - i\epsilon)} + 2i \int \frac{f(k_E^2) f((p - k_E)^2) d^4k_E}{p_0(p_0 - 2ik_{0E})(k_E^2 + m^2)} \quad (18)$$

which differs from (11) by the factor  $-i$  and this is precisely compensated by  $i$  in the definition of amplitude (14).

Thus, the accurate definition of non-locals amplitudes through Euclidean integrals analytically continued to Minkowski external momenta is equivalent to the familiar one used for local theories. Even though the latter cannot be formalized through the Wick rotation in non-local theories because the Minkowski integral is divergent, this formal approach gives the same results. Further, as it was proven in [sen], the way of amplitudes definition through Euclidean integrals preserves the unitarity of non-local theories and the optical theorem. Also one easily see that the local limit  $f(\square) \rightarrow 1$  restores standard textbook answers.

## 4 Fish diagram 2-2 scattering

In this section we consider the model with non-local function of the simplest form allowing for the analytic computation of the amplitude,  $f(\square) = e^{\alpha\square}$ . The matrix element can be formally defined as

$$\mathcal{M} = -i \frac{\lambda^2}{32\pi^4} I(p) \quad (19)$$

where

$$I(p) = \int d^4k \frac{e^{\alpha((p/2+k)^2 + (p/2-k)^2)}}{((p/2+k)^2 - m^2)((p/2-k)^2 - m^2)} \quad (20)$$

As has been already discussed above, the Minkowski integral is divergent and the physical amplitude is defined in another way. We replace loop momenta by their Euclidean counterparts as we would do for the Wick rotation in a local theory and then obtain  $I(p)$  as a sum of the Euclidean integral and the pole part,

$$I(p) = I_{\mathcal{E}}(p) + I_{pole}(p) \quad (21)$$

The pole part here stands for the portion of the whole expression which corresponds to poles which appear at points  $k_0 = \pm p_0/2 \pm \sqrt{\vec{k}^2 + m^2}$  as long as these poles lie in the first and third quarter of the integration contour plane upon the prescription  $m^2 \rightarrow m^2 - i\epsilon$ .

Let us start with the computation of the Euclidean part,

$$\begin{aligned}
I_{\mathcal{E}} &= ie^{\alpha p^2/2} \int \frac{dk_0 d^3 k e^{-2\alpha(k_0^2 + \vec{k}^2)}}{(k_0^2 + \vec{k}^2 + m^2 - p^2/4 + ipk_0)(k_0^2 + \vec{k}^2 + m^2 - p^2/4 - ipk_0)} = \\
&= 2\pi i e^{\alpha p^2/2} \int \frac{dr d\theta r^3 \sin^2 \theta e^{-\alpha r^2}}{(r^2 + m^2 - p^2/4)^2 + p^2 r^2 \cos^2 \theta} = \\
&= \frac{4\pi^2 i e^{\alpha p^2/2}}{p^2} \int r dr e^{-2\alpha r^2} \left( \sqrt{\frac{(r^2 + m^2 - p^2/4)^2 + p^2 r^2}{(r^2 + m^2 - p^2/4)}} - 1 \right)
\end{aligned} \tag{22}$$

In the massless limit,

$$I_{\mathcal{E}} = \frac{4\pi^2 i e^{\alpha p^2/2}}{p^2} \left( \int_0^{p^2/4 - \epsilon_0} dz e^{-2\alpha z} \frac{z}{p^2/4 - z} + \int_{p^2/4 + \epsilon_0}^{\infty} dz e^{-2\alpha z} \frac{p^2/4}{z - p^2/4} \right) \tag{23}$$

Evaluating these integrals one obtains the contribution to  $\text{Re } \mathcal{M}$

$$I_{\mathcal{E}}^{Re} = -\frac{2\pi^2 i}{p^2 \alpha} \left( e^{\alpha p^2/2} - 1 - 2\alpha p^2 \text{Ei}(\alpha p^2/2) + \alpha p^2 (\gamma + \log \epsilon_0) \right) \tag{24}$$

Here  $\gamma$  is Euler constant,  $\text{Ei}(z)$  is the integral exponent and we have taken the mean value of the integrals performing the limit  $\epsilon_0 \rightarrow 0$ ). As we will see later, the divergence  $\log \epsilon_0$  will be exactly cancelled by the pole part of the amplitude in the massless limit.

The pole part is non-zero only when there are poles in I and III sectors of the complex plane, i.e. when  $k^2 + m^2 > p^2/4$ . Fig. 3 illustrates this situation. In this case, their impact to the amplitude is given by the sum of the relevant residues in (20)

$$I_{pole} = -2\pi i \int d^3 k \left( \text{Res}(k_0 = p_0/2 - \sqrt{m^2 + \vec{k}^2}) - \text{Res}(k_0 = -p_0/2 + \sqrt{m^2 + \vec{k}^2}) \right) \tag{25}$$

which readily simplifies to

$$I_{pole} = -2\pi i \int \frac{d^3 k e^{2\alpha m^2 + 2\alpha p(p/2 - \sqrt{\vec{k}^2 + m^2})}}{2p\sqrt{m^2 + \vec{k}^2}(p/2 - \sqrt{m^2 + \vec{k}^2})} \tag{26}$$

For  $m = 0$  we obtain

$$I_{pole} = -\frac{8\pi^2 i e^{\alpha p^2}}{p^2} \int_0^{p^2 - \epsilon_0} \frac{z dz e^{-4\alpha z}}{p^2/4 - z} \tag{27}$$

We notice that integration up to  $p^2$  follows from the fact that  $k^2 + m^2$  should be greater than  $p^2$  in order to have poles in the I and III quarters of the integration plane.

$$I_{pole} = \frac{2\pi^2 i}{\alpha p^2} \left( e^{\alpha p^2} - 1 - \alpha p^2 \text{Ei}(\alpha p^2) + \alpha p^2 (\gamma + \log \epsilon_0) \right) \tag{28}$$




Figure 3: Poles of the amplitude (20).

Summing up and adding the computed in Sec. 3 imaginary part contribution  $-\pi^3$  to  $I(p)$  we get the final answer

$$I(p) = -\pi^3 + \frac{2i\pi^2}{\alpha p^2} \left( e^{\alpha p^2} - \alpha p^2 \text{Ei}(\alpha p^2) - e^{\alpha p^2/2} + \frac{1}{2} \alpha p^2 \text{Ei}(\alpha p^2/2) \right) \quad (29)$$

The real part of amplitude (19) is plotted on Fig. 4 where presence of a strong coupling regime seems obvious. We will see in next Sections that this situations depends on the form-factor.

One can further observe using that for small arguments  $\text{Ei}(z) \approx \gamma + \log(z) + z$  that the local limit and the logarithmic singularity usually arising in the cut-off regularization restore as long as  $\alpha \rightarrow 0$ .

The total scattering amplitude including the crossed diagrams can be obtained as,

$$\mathcal{M}_{tot} = -i \frac{\lambda^2}{32\pi^4} \left( I(\sqrt{s}) + I(\sqrt{t}) + I(\sqrt{u}) \right) \quad (30)$$

where standard Mandelstam variables are used.

## 5 Computation for a generic propagator

The computation of the previous Section can be extended to a generic propagator, i.e. generic form-factor  $f(\square)$ . This can be done by utilizing the method described in [?] which reduces






Figure 4: Amplitude (real part) dependence on the external momentum. Here  $\alpha = 1$ ,  $m = 0$ ,  $\lambda = 0.1$ .

in our case to the following quite simple expression for the matrix element

$$\mathcal{M}(p) = \frac{-i\lambda^2}{2(2\pi)^4} \int d^4x \Delta(x)^2 e^{ipx} \quad (31)$$

where  $\Delta(x)$  is the propagator in the  $x$ -representation and the whole formula is in a sense the Fourier transform of the product of the propagators.

The propagator in  $x$ -space can be easily written as

$$\Delta(x) = i \int d^4k \frac{f(k^2)}{k^2} e^{ikx} \quad (32)$$

and in contrast with a local theory the latter integral can be straightforwardly made well defined at short distances, i.e. small  $x$ . First, we go to spherical coordinates and simplify the last expression as follows

$$\Delta(x) = 4\pi^3 i \int_0^\infty dk f(k^2) \frac{J_1(kx)}{x} \quad (33)$$

where  $J_1(z)$  is the Bessel function.

Then we get by direct substitution

$$\mathcal{M}(p) = -2\pi^2 \lambda^2 \int_0^\infty \frac{J_1(px) J_1(kx) J_1(qx)}{px^2} f(k^2) f(q^2) dk dq dx \quad (34)$$

This form allows computing and verifying the optical theorem statement at least numerically for various form-factors  $f(k^2)$ . As the first check one can find out that the results of the previous Section can be reproduced as long as  $f(k^2) = \exp(-\alpha k^2)$ . Further results for the time being can be obtained using numeric integration.

The latter form-factor is characterized by polynomial decay at infinity making the theory avoiding a possible strong coupling regime.

Generically one can try performing the integration over  $x$  in (34) analytically in terms of the fourth Appell function  $F_4(z)$  as follows

$$\int_0^\infty \frac{J_1(px)J_1(kx)J_1(qx)}{x^2} dx = \frac{kq}{4p} F_4(0, 1; 2; 2; k^2/p^2; q^2/p^2)$$

[30, 31]. Even though this looks like explicit analytic answer its use is not obviously well motivated. The reason is that the given answer is justified for  $k, q < p$  and thus requires analytic continuation to other values to cover the full domain of the integration. The corresponding procedure has to be studied in order to be applied for actual computations.

We present the result of the numerical computation for the two examples of non-local function. As the first example we take

$$f_1(k^2) = e^{-k^4} \quad (35)$$

being the simplest non-local function allowing for staying in perturbative regime<sup>2</sup>. Besides that we consider a more involved function which falls polynomially along the real axis while it is an exponent of entire function. This function was first suggested by Tomboulis in [8]

$$f_2(k^2) = e^{-\Gamma(0, k^4) - \gamma - \log k^4} \quad (36)$$

Here  $\Gamma(0, k^4)$  is an incomplete gamma function. For large real  $k$   $f_2(k^2) \propto 1/k^4$ . In Fig. 5 we plot the modulus of this function on the whole complex plane to make its growth directions visible.

Also we plot the four-dimensional Fourier images  $\Delta_1(x)$  and  $\Delta_2(x)$  for the functions  $f_1(k)$  and  $f_2(k)$ , respectively, in Fig. (5). The resulting amplitude (real part) numerically computed with the use of (31) is plotted in Fig. 5. One can see that the curves grow for small momenta representing the logarithmic divergence in the local limit and fall for momenta much higher than non-locality scale. The asymptotes of this falling depends on the UV behaviour of non-local propagator and differs for the two cases under consideration. We obtained the amplitude for  $f_1$  falling exponentially, as  $e^{-p^4}$  while the function  $f_2$  leads to power law suppression. We see that the model remains weakly coupled for both chosen form-factors in contrast with the  $e^{\alpha p^2}$  form-factor discussed in Section 4.

## 6 UV properties of the amplitudes in non-local theory

The UV behaviour of the scattering amplitudes is determined by the form of non-local propagator. As we obtained in Sec. 4, for the propagator  $e^{\square/\Lambda^2}/\square$  the amplitude is growing exponentially for positive  $p^2$ . Does it mean that this kind of model are not valid from the point of view of quantum field theory [32]? We cannot immediately conclude that such kind of propagators cannot be considered because around the non-locality scale the model undergoes strong coupling regime which means that one-loop approximation is not valid

---

<sup>2</sup>Hereafter in our numerical studies we set the non-local scale to be unity.




Figure 5: Absolute value of the function  $f_2(\text{Re } k, \text{Im } k)$ .

anymore. However, this do not immediately mean that the unitarity is broken because non-perturbative methods can lead to the result which is still consistent with unitarity. We leave this question for future study.

In this work we concentrate on the results which can be obtained perturbatively. If the propagator  $f(k^2)/k^2$  is falling for both signs of  $k^2$  than we expect the loop contributions to fall for all physical values of external momenta. However, the scattering amplitude would inevitably grow for some directions of  $s$  and  $t$  in the complex plane. Let us check the consistency of such a behaviour with the known results about the properties of the amplitudes. For example, the Martin-Froissart bound [33] tells us that the amplitude in the forward limit




Figure 6: Left plot represents the functions  $f_1(k)$ ,  $f_2(k)$  and right plot shows their Fourier images  $\Delta_{1,2}$ .




Figure 7: Scattering 2-2 one-loop amplitude as a function of the external momentum. Red curve is for form-factor  $f_1(k)$  and blue curve is for form-factor  $f_2(k)$ . Here we have taken  $\lambda = 0.1$ .

$t \rightarrow 0$  is bounded by

$$|\mathcal{M}(s, 0)| < Cs(\log s)^2 \quad (37)$$

on the  $s$  complex plane. Although this bound was obtained in local theories, the recent work [32] provides a formal proof of the similar bound in non-local theories,

$$|\mathcal{M}(s, 0)| \lesssim Cs^{1+2a}. \quad (38)$$

This bound was obtained within rather weak assumptions including the validity of partial wave expansion, unitarity, analyticity and the exponential boundedness  $|\mathcal{M}(s, t)| < s^N e^{a|s|^\alpha}$  for fixed  $t$ . Here  $\alpha$ ,  $N$ ,  $C$  are some positive constants. These assumptions are expected to be fulfilled in the models we consider in this work. The bound (38) actually means that the amplitudes in the forward limit are *polynomially* bounded while one would expect the exponential growth in some directions on the complex  $s$  plane.

Let us discuss this issue using the simple model with cubic interaction where the exponential behaviour shows up already at tree level.

$$S = \int d^4x \left[ \frac{1}{2} \phi(\square - m^2) f(\square)^{-1} \phi - \frac{\lambda}{3!} \phi^3 \right] \quad (39)$$

The tree level amplitude in this model has the form

$$\mathcal{M}(s, t) \sim \lambda^2 \left( \frac{f(s)}{s - m^2} + \frac{f(t)}{t - m^2} + \frac{f(u)}{u - m^2} \right) \quad (40)$$

Thus,  $f(s)$  must have a direction of exponential growth since an entire function which is polynomially bounded on the whole complex plane is nothing else than polynomial of finite

order. However, for all physical momenta the amplitude is small if we require  $f(q^2)$  to fall for real  $q^2$ . Thus, with this choice, the model stays in perturbative regime for physical values of momenta. However, if one computes the amplitude for complex values of momenta this could lead to breaking of the perturbation theory. The naive computation leading to the exponential growth is not valid anymore. Therefore the conclusion about the amplitude's exponential growth in the nonphysical region is not correct. Moreover, the analyticity arguments leading eventually to the bound (38) allows us to conclude that the full nonperturbative amplitude is bounded polynomially.

The bound (38) leads to another interesting consequences for the model (39). Under the condition of polynomial boundedness of the amplitude in forward limit we have the Cerulus-Martin lower bound on the amplitude falling [34],

$$|\mathcal{M}(s, t \rightarrow 0)| > C e^{-\gamma\sqrt{s}\log(s/s_0)} \quad (41)$$

Here  $C$ ,  $\gamma$ ,  $s_0$  are positive constants. This bound is obviously violated if we take  $f(k^2) \propto e^{-\alpha k^4}$ . The tree level amplitude is falling too fast while the model is in a perturbative regime so we can trust this computation for physical momenta. This contradiction means that something is wrong with the models with exponentially falling propagators. We leave for the future study finding which assumption is actually broken in a model with  $f(k^2) \propto e^{-\alpha k^4}$ . Notice that in the work [8] the choice of polynomially suppressed propagators was suggested motivated by different arguments such as of power counting for example. The propagator falling as  $1/k^n$  is still consistent with the discussed bounds.

Notice that disfavouring of the exponentially suppressed propagator is not applicable for models with contact  $\phi^4$  interactions since the leading term is given by a constant tree level contribution. This kind of suppression for those models is not in a contradiction with Martin-Cerulus bound [34].

## 7 Conclusion and Outlook

We have shown explicitly that the optical theorem statement holds in non-local higher derivative theories upon utilizing a modified computation prescription. The prescription consist of computing Feynman graphs assuming all the internal momenta are Euclidean and then continuing the final answer analytically to Minkowski signature for external momenta. Resulting expressions satisfy the optical theorem statement and obey the correct local theory limit. We were able to show that the one-loop correction to the vertex in a non-local  $\phi^4$  theory can be computed for arbitrary form-factors and have presented both analytic and numeric derivations of the result.

As an important observation we have found that since the Martin-Cerulus bound [34] must be satisfied in non-local models as in local ones, than the amplitudes should not fall too fast obeying bound (41). If such a non-local theory has a contact interaction than this bound can always be satisfied irrespectively of the decay rate of the propagator as the contact interaction term always dominates in the perturbative expansion. On the other hand, in the absence of the contact interaction (for instance computing corrections to 2-2 scattering in  $\phi^3$

theory) one must demand that propagator decays polynomially favoring this way the type of functions suggested in [8].

As the next step one would aim at demonstrating that the background field method provides consistent results and thus can be applied to AID models which is a plausible expectation. Also the strong coupling regime which may arise in such non-local field theories [20] due to the growth of the amplitudes for complex parameters is an important question for further investigations. Yet another ambitious problem is to study such theories in the large- $N$  approximations [35] and see how they are compared to their local counterparts.

## Note added

Upon completion of the manuscript we have met with great interest a related study arXiv:2103.00353 discussing the same problem of unitarity of non-local field theories.

## Acknowledgements

Authors would like to thank I. Aref'eva and M. Shaposhnikov for comments and questions on the manuscript. AK is supported by FCT Portugal investigator project IF/01607/2015. AT is supported by the Academy of Finland grant 318319. The part of the work performed by AT related to the numeric computations of the amplitudes is supported by the Russian Science Foundation grant 19-12-00393.

## References

- [1] M. Ostrogradsky, “Memoires sur les equations differentielles relatives au probleme des isoperimetres,” *Mem. Ac. St. Petersbourg* **VI** no. 4, (1850) 385.
- [2] G. V. Efimov, “Non-local quantum theory of the scalar field,” *Commun. Math. Phys.* **5** no. 1, (1967) 42–56.
- [3] S. Coleman, “Resonance poles and resonance multipoles,”.
- [4] K. S. Stelle, “Renormalization of Higher Derivative Quantum Gravity,” *Phys. Rev. D* **16** (1977) 953–969.
- [5] E. S. Fradkin and A. A. Tseytlin, “Renormalizable Asymptotically Free Quantum Theory of Gravity,” *Phys. Lett. B* **104** (1981) 377–381.
- [6] Y. Kuzmin, “THE CONVERGENT NONLOCAL GRAVITATION. (IN RUSSIAN),” *Sov. J. Nucl. Phys.* **50** (1989) 1011–1014.
- [7] N. Krasnikov, “NONLOCAL GAUGE THEORIES,” *Theor. Math. Phys.* **73** (1987) 1184–1190.

- [8] E. Tomboulis, “Superrenormalizable gauge and gravitational theories,” [arXiv:hep-th/9702146](#).
- [9] L. Brekke, P. G. Freund, M. Olson, and E. Witten, “Nonarchimedean String Dynamics,” *Nucl. Phys. B* **302** (1988) 365–402.
- [10] V. Vladimirov, I. Volovich, and E. Zelenov, *p-adic analysis and mathematical physics*, vol. 1. 1994.
- [11] B. Dragovich, A. Y. Khrennikov, S. Kozyrev, I. Volovich, and E. Zelenov, “p-Adic Mathematical Physics: The First 30 Years,” *Anal. Appl.* **9** (2017) 87–121, [arXiv:1705.04758 \[math-ph\]](#).
- [12] D. Anselmi, “Fakeons And Lee-Wick Models,” *JHEP* **02** (2018) 141, [arXiv:1801.00915 \[hep-th\]](#).
- [13] A. Nicolis, R. Rattazzi, and E. Trincherini, “The Galileon as a local modification of gravity,” *Phys. Rev. D* **79** (2009) 064036, [arXiv:0811.2197 \[hep-th\]](#).
- [14] G. W. Horndeski, “Second-order scalar-tensor field equations in a four-dimensional space,” *Int. J. Theor. Phys.* **10** (1974) 363–384.
- [15] T. Kobayashi, “Horndeski theory and beyond: a review,” *Rept. Prog. Phys.* **82** no. 8, (2019) 086901, [arXiv:1901.07183 \[gr-qc\]](#).
- [16] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, “Towards singularity and ghost free theories of gravity,” *Phys. Rev. Lett.* **108** (2012) 031101, [arXiv:1110.5249 \[gr-qc\]](#).
- [17] T. Biswas, A. S. Koshelev, and A. Mazumdar, “Consistent higher derivative gravitational theories with stable de Sitter and anti-de Sitter backgrounds,” *Phys. Rev. D* **95** no. 4, (2017) 043533, [arXiv:1606.01250 \[gr-qc\]](#).
- [18] L. Keltner and A. J. Tolley, “UV properties of Galileons: Spectral Densities,” [arXiv:1502.05706 \[hep-th\]](#).
- [19] A. S. Koshelev, K. S. Kumar, and A. A. Starobinsky, “Analytic infinite derivative gravity,  $R^2$ -like inflation, quantum gravity and CMB,” [arXiv:2005.09550 \[hep-th\]](#).
- [20] A. S. Koshelev and A. Tokareva, “Non-local self-healing of Higgs inflation,” *Phys. Rev. D* **102** (2020) 123518, [arXiv:2006.06641 \[hep-th\]](#).
- [21] J. Boos, V. P. Frolov, and A. Zelnikov, “Ghost-free modification of the Polyakov action and Hawking radiation,” *Phys. Rev. D* **100** no. 10, (2019) 104008, [arXiv:1909.01494 \[hep-th\]](#).
- [22] E. Witten, “Noncommutative Geometry and String Field Theory,” *Nucl. Phys. B* **268** (1986) 253–294.

- [23] E. Witten, “Interacting Field Theory of Open Superstrings,” *Nucl. Phys. B* **276** (1986) 291–324.
- [24] K. Ohmori, “A Review on tachyon condensation in open string field theories,” other thesis, 2, 2001.
- [25] I. Arefeva, D. Belov, A. Giryavets, A. Koshelev, and P. Medvedev, “Noncommutative field theories and (super)string field theories,” in *11th Jorge Andre Swieca Summer School on Particle and Fields*, pp. 1–163. 11, 2001. [arXiv:hep-th/0111208](#).
- [26] M. E. Peskin and D. V. Schroeder, *An Introduction to quantum field theory*. Addison-Wesley, Reading, USA, 1995.
- [27] R. Pius and A. Sen, “Cutkosky rules for superstring field theory,” *JHEP* **10** (2016) 024, [arXiv:1604.01783 \[hep-th\]](#). [Erratum: *JHEP* 09, 122 (2018)].
- [28] R. Pius and A. Sen, “Unitarity of the Box Diagram,” *JHEP* **11** (2018) 094, [arXiv:1805.00984 \[hep-th\]](#).
- [29] F. Brischese and L. Modesto, “Cutkosky rules and perturbative unitarity in Euclidean nonlocal quantum field theories,” *Phys. Rev. D* **99** no. 10, (2019) 104043, [arXiv:1803.08827 \[gr-qc\]](#).
- [30] I. S. Gradshteyn and I. M. Ryzhik, *Table of integrals, series, and products*. Elsevier/Academic Press, Amsterdam, Amsterdam, The Netherlands, 2007.
- [31] W. N. Bailey, “Some infinite integrals involving bessel functions,” *Proceedings of the London Mathematical Society* **s2-40** no. 1, 37–48.
- [32] J. Tokuda, “Extension of positivity bounds to non-local theories: IR obstructions to Lorentz invariant UV completions,” *JHEP* **05** (2019) 216, [arXiv:1902.10039 \[hep-th\]](#).
- [33] A. Martin, “Extension of the axiomatic analyticity domain of scattering amplitudes by unitarity. 1.,” *Nuovo Cim. A* **42** (1965) 930–953.
- [34] F. A. Cerulus and A. Martin, “A lower bound for large-angle elastic scattering at high energies,” *Phys. Lett.* **8** (1964) 80–82.
- [35] G. ’t Hooft, “Large N,” in *The Phenomenology of Large N(c) QCD*, pp. 3–18. 4, 2002. [arXiv:hep-th/0204069](#).