Parametric Complexity Bounds for Approximating PDEs with Neural Networks

Tanya Marwah, Zachary C. Lipton, Andrej Risteski Machine Learning Department, Carnegie Mellon University {tmarwah, zlipton, aristesk}@andrew.cmu.edu

Abstract

Recent empirical results show that deep networks can approximate solutions to high dimensional PDEs, seemingly escaping the curse of dimensionality. However many open questions remain regarding the theoretical basis for such approximations, including the number of parameters required. In this paper, we investigate the representational power of neural networks for approximating solutions to linear elliptic PDEs with Dirichlet Boundary conditions. We prove that when a PDE's coefficients are representable by small neural networks, the parameters required to approximate its solution scale polynomially with the input dimension d and are proportional to the parameter counts of the coefficient neural networks. Our proof is based on constructing a neural network which simulates gradient descent in an appropriate Hilbert space which converges to the solution of the PDE. Moreover, we bound the size of the neural network needed to represent each iterate in terms of the neural network representing the previous iterate, resulting in a final network whose parameters depend polynomially on d and does not depend on the volume of the domain.

1 Introduction

A partial differential equation (PDE) relates a multivariate function defined over some domain to its partial derivatives. Typically, one's goal is to solve for the (unknown) function, often subject to additional constraints, such as the function's value on the boundary of the domain. PDEs are ubiquitous in both the natural and social sciences, where they model such diverse processes as heat diffusion [Crank and Nicolson, 1947, Özişik et al., 2017], fluid dynamics [Anderson and Wendt, 1995, Temam, 2001], and financial markets [Black and Scholes, 1973, Ehrhardt and Mickens, 2008]. Because most PDEs of interest lack closed-form solutions, computational approximation methods remain a vital and active field of research [Ames, 2014]. For low-dimensional functions, dominant approaches include the finite differences and finite element methods [LeVeque, 2007], which operate on discretizations of the domain. After partitioning the domain into a *mesh*, these methods solve for the function value at the vertices of the mesh. However, such mesh-based methods scale exponentially with the input dimension, rendering them unsuitable for high-dimensional problems.

Following breakthroughs in deep learning for approximating high-dimensional functions in such diverse domains as computer vision [Krizhevsky et al., 2012, Radford et al., 2015] and natural language processing [Bahdanau et al., 2014, Devlin et al., 2018, Vaswani et al., 2017], a burgeoning line of research leverages neural networks to approximate solutions to PDEs. This line of work has produced promising empirical results for common PDEs such as the Hamilton-Jacobi-Bellman and

Black-Scholes equations [Han et al., 2018, Grohs et al., 2018, Sirignano and Spiliopoulos, 2018]. Because they do not explicitly discretize the domain, and given their empirical success on highdimensional problems, these methods *appear* not to suffer the curse of dimensionality. However, these methods are not well understood theoretically, leaving open questions about when they are applicable, what their performance depends on, and just how much many parameters are required to approximate the solution to a given PDE.

Over the past three years, several theoretical works have investigated questions of representational power under various assumptions. Khoo et al. [2017] prove a kind of universal approximation power of neural networks, albeit with size exponential in the input dimension. Exploring a variety of settings, Kutyniok et al. [2019], Grohs et al. [2018], and Jentzen et al. [2018], proved that the number of parameters required to approximate a solution to a PDE exhibits a less than exponential dependence on the input dimension for some special parabolic PDEs that admit straightforward analysis. Grohs and Herrmann [2020] consider elliptic PDEs with Dirichlet boundary conditions. However, their rate depends on the volume of the domain, and thus can have an implicit exponential dependence on dimension (e.g., consider a hypercube with side length greater than one).

In this paper, we focus on a linear elliptic PDEs with Dirichlet boundary conditions, which are prevalent in science and engineering (e.g., the Laplace and Poisson equations). Most commonly, linear elliptic PDEs define the steady state of processes like heat diffusion and fluid dynamics. We provide an answer to the following:

Question. How many parameters suffice to approximate solutions to a linear elliptic PDE up to a specified level of precision using a neural network?

We show that when the coefficients of the PDE are expressible as small neural networks (note that PDE coefficients are functions), the number of parameters required to express an approximate solution to the PDE is proportional to the number of parameters required to express the coefficients. Furthermore, we show that the number of parameters depends polynomially on the dimension and does not depend upon the volume of the domain.

2 Overview of Results

We will consider PDEs that take the following form:

Definition 1 (Linear Elliptic PDE [Evans, 1998]). A linear elliptic PDE is a partial differential equation of the form

$$\begin{cases} (Lu)(x) \equiv (-div(A\nabla u) + cu)(x) = f(x), \forall x \in \Omega, \\ u(x) = 0, \forall x \in \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^d$ is a bounded open set with a boundary $\partial \Omega$. Further, for all $x \in \Omega$, $A : \Omega \to \mathbb{R}^{d \times d}$ is a matrix-valued function, s.t. $A(x) \succ 0$, and $c : \Omega \to \mathbb{R}$, s.t. c(x) > 0.¹

We will refer to A and c as the coefficients of the PDE. The divergence form in Definition 1 is one of two canonical ways to define a linear elliptic PDE [Evans, 1998] and is convenient for several

¹Here div denotes the divergence operator, which given a vector field $F : \mathbb{R}^d \to \mathbb{R}^d$ outputs div $(F) = \nabla \cdot F = \sum_{i=1}^d \frac{\partial F_i}{\partial x_i}$

technical reasons (see Section 4). The boundary condition being such that the solution takes a constant value (here 0) is also without loss of generality.

We will be interested in the complexity of the solution of the PDE (in terms of number of parameters of a neural network that approximates it) when the coefficients A and c can be expressed as neural networks with small number of parameters. (For technical details, see Section 5.) Informally, our result states the following:

Theorem (Informal). If the coefficients A, c and the function f are approximable by neural networks with at most N parameters, the solution u^* to the PDE in Definition 1 is approximable by a neural network with O(poly(d)N) parameters.

The formal statement is provided in Section 5.

Our results provide supporting theory that can help to explain the empirical efficacy of neural networks in approximating solutions to high-dimensional PDEs with boundary conditions as shown in Sirignano and Spiliopoulos [2018] and Li et al. [2020a]. Our relies on simulating gradient descent in an appropriate Hilbert space using neural networks. Namely, we show that each iterate can be approximated by a small neural network slightly larger than the neural network approximating the previous iterate, as well as bound the total number of iterates.

Organization of the paper We introduce the required notation along with some mathematical preliminaries on PDEs in Section 4. The problem setting and formal statement of the main result are provided in Section 5. Finally, we provide the proof of the main result in Section 6.

3 Prior Work

Among the first papers to leverage neural networks to approximate solutions to PDEs with boundary conditions are Lagaris et al. [1998], Lagaris et al. [2000], and Malek and Beidokhti [2006]. However, these methods discretize the input space and thus are not suitable for high-dimensional input spaces. More recently, mesh-free neural network approaches have been proposed for highdimensional PDEs [Han et al., 2018, Raissi et al., 2017, 2019], achieving impressive empirical results in various applications. Sirignano and Spiliopoulos [2018] design a loss function that penalizes failure to satisfy the PDE, training their network on minibatches sampled uniformly from the input domain. They also provide a universal approximation result that shows that for sufficiently regularized PDEs, there exists a multilayer network that approximates its solution. However, they do not comment on the complexity of the neural network or how it scales with the input dimension. Khoo et al. [2017] also prove universal approximation power, albeit with networks of size exponential in the input dimension. Recent work from Grohs et al. [2018], Jentzen et al. [2018] provide a better than exponential dependence on the input dimension for some special parabolic PDEs, for which the simulating a PDE solver by a neural network is straightforward.

Several recent works [Bhattacharya et al., 2020, Kutyniok et al., 2019, Li et al., 2020b,a] present empirical results showing that a single neural network can solve for an entire family of PDEs. They approximate the map from the parameters of the PDE to the solution of the PDE using a neural network. This can potentially save practitioners the trouble of retraining a new network each time the coefficients of the PDE change. The only one of these papers to provide theoretical grounding for this approach is Kutyniok et al. [2019]. However, they assume the existence of a finite low-dimensional space with basis functions that can approximate this parametric map—and it is unclear when this would obtain. Our work proves the existence of such maps in Section 7, under the assumption that the family of PDEs has coefficients described by neural networks with a fixed architecture.

In the work most closely related to ours, Grohs and Herrmann [2020] provide approximation rates polynomial in the input dimension d for the Poisson equation (a special kind of linear elliptic PDE) with Dirichlet boundary conditions. They introduce a walk-on-the-sphere algorithm, which simulates a stochastic differential equation that can be used to solve a Poisson equation with Dirichlet boundary conditions (see, e.g. Oksendal [2013] (Theorem 9.13)). The rates provided in Grohs and Herrmann [2020] depend on the volume of the domain, and thus have an implicit dependence on the input dimension d. However, as they show, their construction avoids the curse of dimensionality for domains where the volume remains constant in dimensions, e.g., a hypercube with side length one. Our results consider the boundary condition for the PDE and is independent of the volume of the sphere. Further, we note that our results are defined for a more general linear elliptic PDE, of which the Poisson equation is a special case.

4 Notation and Definitions

We now introduce several key concepts from PDEs and our notation. For any open set $\Omega \subset \mathbb{R}^d$, we denote its boundary by $\partial\Omega$ and denote its closure by $\bar{\Omega} := \Omega \cup \partial\Omega$. By $C^0(\Omega)$, we denote the space of real-valued continuous functions defined over the domain Ω . Furthermore, for $k \in \mathbb{N}$, a function $g \in C^k(\Omega)$ if $g \in C^0(\Omega)$ and $\partial^{\alpha}g \in C^0(\Omega)$ for any multi-index α such that $|\alpha| \leq k$. Finally, a function $g \in C^{\infty}(\Omega)$ if $g \in C^k(\Omega)$ for all $k \in N$. Next, we define several function spaces that are essential to our analysis:

Definition 2. For any $k \in \mathbb{N} \cup \{\infty\}$, $C_0^k(\Omega) := \{g : g \in C^k(\Omega), \overline{supp(g)} \subseteq \Omega\}$.

Definition 3. For a domain Ω and a function $g: \Omega \to \mathbb{R}$, the function space $L^2(\Omega)$ is defined as:

$$L^{2}(\Omega) := \left\{ g : \|g\|_{L^{2}(\Omega)} < \infty \right\},$$

where $\|g\|_{L^2(\Omega)} = \left(\int_{\Omega} |g(x)|^2 dx\right)^{\frac{1}{2}}$. This function space is equipped with the following inner product:

$$\langle g,h\rangle_{L^2(\Omega)} = \int_{\Omega} g(x)h(x)dx.$$

Definition 4. For a domain Ω and a function $g : \Omega \to \mathbb{R}$, the function space $L^{\infty}(\Omega)$ is defined analogously, where $\|g\|_{L^{\infty}(\Omega)} = \inf\{c \ge 0 : |g(x)| \le c \text{ for almost all } x \in \Omega\}.$

Definition 5. For a domain Ω and $m \in \mathbb{N}$, we define the Hilbert space $H^m(\Omega)$ as

$$H^{m}(\Omega) := \{g : \Omega \to \mathbb{R} : \partial^{\alpha}g \in L^{2}(\Omega), \ \forall \alpha \ s.t. \ |\alpha| \le m\}$$

Furthermore, $H^m(\Omega)$ is equipped with the inner product, $\langle g,h\rangle_{H^m(\Omega)} = \sum_{|\alpha| \le m} \int_{\Omega} (\partial^{\alpha} g) (\partial^{\alpha} h) dx$ and the corresponding norm

$$\|g\|_{H^m(\Omega)} = \left(\sum_{|\alpha| \le m} \|\partial^{\alpha}g\|_{L^2(\Omega)}^2\right)^{\frac{1}{2}}.$$

Definition 6. The closure of $C_0^{\infty}(\Omega)$ in $H^m(\Omega)$ is denoted by $H_0^m(\Omega)$.

Informally, $H_0^m(\Omega)$ is the set of functions belonging to $H^m(\Omega)$ that can be approximated by a sequence of functions $\phi_n \in C_0^\infty(\Omega)$. This also implies that if a function $g \in H_0^m(\Omega)$, then g(x) = 0 for all $x \in \partial \Omega$. This space (particularly with m = 1) is often useful when analyzing elliptic PDEs with Dirichlet boundary conditions.

Definition 7 (Gateaux Derivative). Let $G : \mathcal{U} \to \mathcal{V}$ be a function between two topological spaces. For $x, h \in \mathcal{U}$ and $h \neq 0$, the Gateaux differential of G at x in the direction of h is defined as

$$F'[x](h) = \lim_{\epsilon \to 0} \frac{G(x+\epsilon h) - G(x)}{\epsilon}$$

Having introduced these preliminaries, we now introduce some important concepts and formulations for a linear PDE, which we will use extensively in developing our analysis.

Proposition 1. For the PDE in Definition 1, given $f \in L^2(\Omega)$ the following hold:

- 1. There exists a function $\bar{f} \in H_0^1(\Omega)$ such that for all $v \in H_0^1(\Omega)$, $\langle f, v \rangle_{L^2(\Omega)} = \langle \bar{f}, v \rangle_{L^2(\Omega)}$.
- 2. The solution u^* to the PDE is unique and obtained as

$$u^* = \operatorname*{argmin}_{u \in H_0^1(\Omega)} F[u], \text{ where } F[u] = \frac{1}{2} \|Lu - \bar{f}\|_{L^2(\Omega)}^2$$

- 3. For all $v \in H_0^1(\Omega)$, the Gateaux derivative of F[u] at a point $u \in H_0^1(\Omega)$ in the direction v is given by $F'[u](v) = \langle Lu \bar{f}, v \rangle_{L^2(\Omega)}$.
- 4. u^{\star} is the unique stationary point of F, that is $F'[u](v) = 0, \forall v \in H_0^1(\Omega)$ if and only if $u = u^{\star}$. Equivalently, for all $v \in H_0^1(\Omega)$ we have,

$$\langle Lu, v \rangle_{L^2(\Omega)} = \langle \bar{f}, v \rangle_{L^2(\Omega)} = \langle f, v \rangle_{L^2(\Omega)}, \tag{1}$$

if and only if $u = u^*$.

Proof. First, we prove point 1. For all $v \in H_0^1(\Omega)$, let $l(v) := \langle f, v \rangle_{L^2(\Omega)}$ be a linear operator defined on $H_0^1(\Omega)$. Note that from the Riesz representation theorem there exists a function $\bar{f} \in H_0^1(\Omega)$ such that $l(v) = \langle \bar{f}, v \rangle_{L^2(\Omega)}$. This completes the proof.

To get the result in point 3, we calculate the Gateaux derivative of F[u], *i.e.*, for a $v \in H_0^1(\Omega)$,

$$\begin{aligned} F'[u](v) &= \lim_{\epsilon \to 0} \frac{F[u+v] - F[u]}{\epsilon} \\ &= \lim_{\epsilon \to 0} \frac{1}{2} \frac{\|L(u+\epsilon v) - \bar{f}\|_{L^2(\Omega)}^2 - \|Lu - \bar{f}\|_{L^2(\Omega)}^2}{\epsilon} \\ &= \lim_{\epsilon \to 0} \frac{1}{2} \left(\frac{\langle L(u+\epsilon v) - \bar{f}, L(u+\epsilon v) - \bar{f} \rangle_{L^2(\Omega)}}{\epsilon} - \frac{\langle Lu - \bar{f}, Lu - \bar{f} \rangle}{\epsilon} \right) \\ &= \langle Lu - \bar{f}, Lv \rangle_{L^2(\Omega)} \end{aligned}$$

Using the fact that $L: H_0^1(\Omega) \to H_0^1(\Omega)$ and therefore $Lv \in H_0^1(\Omega)$ (see Lemma A.1), we have that for all $v \in H_0^1(\Omega)$, any stationary point u of F will statisfy

$$F'[u](v) = \langle Lu - \bar{f}, v \rangle_{L^2(\Omega)} = 0$$

$$\implies \langle Lu, v \rangle_{L^2(\Omega)} = \langle \bar{f}, v \rangle_{L^2(\Omega)}$$

$$\implies \langle Lu, v \rangle_{L^2(\Omega)} = \langle f, v \rangle_{L^2(\Omega)} \quad \text{using part (1).}$$

Note that (1) has a unique solution (referred to as u^*). The proof for this is standard (see Proposition A.2 in the Appendix). Therefore, since the stationary point of F[u] satisfies (1) and we know that the solution to the equation is unique, it follows that F[u] has a unique stationary point (u^*) .

The first part of the proposition states that we can find a function $\bar{f} \in H_0^1(\Omega)$ where $f_{|\partial\Omega} = 0$ such that the $L^2(\Omega)$ inner products are equal. This ensures that the minimization defined in Part 2 is expressed over functions that satisfy the boundary condition for our PDE (Definition 1)—i.e., that they lie in $H_0^1(\Omega)$ —and hence the solution $u^* \in H_0^1(\Omega)$ as well. Note that equation (1) is also referred to as the variational formulation for the PDE in Definition 1.

Definition 8. Given the operator $L : H_0^1(\Omega) \to H_0^1(\Omega)$, the tuples $(\lambda, \varphi)_{i=1}^{\infty}$, where $\lambda_i \in \mathbb{R}$ and $\varphi_i \in H_0^1(\Omega)$ are (eigenvalue, eigenfunction) pairs that satisfy the following:

$$L\varphi = \lambda\varphi \qquad \forall x \in \Omega.$$

Since $\varphi \in H^1_0(\Omega)$ we know that $\varphi_{|\partial\Omega} = 0$. The eigenvalue can be written as

$$\lambda_i = \inf_{u \in X_i} \frac{\langle Lu, u \rangle_{L^2(\Omega)}}{\|u\|_{L^2(\Omega)}^2} \tag{2}$$

where $X_i := span\{\varphi_1, \ldots, \varphi_i\}^{\perp} = \{u \in H_0^1(\Omega) : \langle u, \varphi_j \rangle_{L^2(\Omega)} = 0 \ \forall i \in 1, \ldots, i\} \text{ and } \lambda_1 \leq \lambda_2 \leq \cdots$.

5 Main Result

Before stating the results, we give the formal assumptions on the PDE we will be considering.

Assumptions:

- (i) **Smoothness**: We assume that the function $f \in L^2(\Omega)$. The coefficient $A \in \Omega \to \mathbb{R}^{d \times d}$ is a symmetric matrix valued function, i.e., $A = (a_{ij}(x))$ and $a_{ij}(x) \in L^{\infty}(\Omega)$ for all $i, j \in [d]$. The function $c \in L^{\infty}(\Omega)$ and there exists a constant ζ such that $c(x) \geq \zeta > 0$ for all $x \in \Omega$.
- (ii) **Ellipticity**: There exist constants $M \ge m > 0$ such that, for all $x \in \Omega$ and $\xi \in \mathbb{R}^d$,

$$m \|\xi\|^2 \le \sum_{i,j=1}^d a_{ij}(x)\xi_i\xi_j \le M \|\xi\|^2.$$

(iii) Neural network approximability: There exist neural networks ϕ_A , ϕ_c with $N_A, N_c \in \mathbb{N}$ number of parameters, that approximate the functions A and c, i.e., $||A - \phi_A||_{L^{\infty}(\Omega)} \leq \epsilon_A$ and $||c - \phi_c||_{L^{\infty}(\Omega)} \leq \epsilon_c$, for small $\epsilon_A, \epsilon_c \geq 0$. We also assume that there exists a neural network $\tilde{f} \in \text{span}\{\varphi_1, \dots, \varphi_k\}$ $(k \in \mathbb{N})$ with $N_{\tilde{f}}$ parameters such that $||f - \tilde{f}||_{L^2(\Omega)} \leq \epsilon_f$. Further, let Σ denote the set of all activation functions used in the networks \tilde{f}, ϕ_A and ϕ_c .

Intuitively, ellipticity of L in a functional equation Lu = f is analogous to positive definiteness of the matrix $Q \in \mathbb{R}^d$ in a linear equation Qx = p, where Q is a matrix and $x, p \in \mathbb{R}^d$ (see Section 6). Further, given $u \in H_0^1(\Omega)$, the function $Lu \in H_0^1(\Omega)$ as well, *i.e.*, $L : H_0^1(\Omega) \to H_0^1(\Omega)$ and is self-adjoint (see Proposition A.1 in the Appendix).

In (iii), we have assumed that the coefficients A and c, and the function f can be approximated by small neural networks. By small, we mean that they have a less than exponential dependence on the input dimension d. In fact, for many PDEs used in practice, approximating the coefficients using neural networks is straightforward. For example, in heat diffusion (whose equilibrium is defined by a linear elliptic PDE) A(x) defines the conductivity of the material at point x. If it is a constant, then it requires a neural network with at most d parameters to approximate.

Given that the operator L is in fact self-adjoint and elliptic, the eigenvalues are real and countable in number. Further, the eigenfunctions form an orthonormal basis of $H_0^1(\Omega)$ (see Evans [1998], Section 6.5). The assumption in (iii) that $\tilde{f} \in H_0^1(\Omega)$ lies in the span of the first k eigenfunctions of the operator L is intuitively a constraint on the smoothness of \tilde{f} . For instance, if $L = -\Delta$ (the Laplacian operator), the Dirichlet form satisfies $\frac{\langle Lu, u \rangle_{L^2(\Omega)}}{\|u\|_{L^2(\Omega)}^2} = \frac{\|\nabla u\|_{L^2(\Omega)}}{\|u\|_{L^2(\Omega)}}$, so eigenfunctions corresponding to larger k exhibit a higher degree of spikiness on average. The reader can also think of the eigenfunctions corresponding to larger k as Fourier basis functions corresponding to higher frequencies.

With this in mind, we can formally state our main result.

Theorem 1 (Main theorem). Consider a linear elliptic PDE satisfying Assumptions (i)-(iii), and let $u^* \in H_0^1(\Omega)$ denotes its unique solution. If there exists a neural network $u_0 \in H_0^1(\Omega)$ with N_0 parameters, such that $||u^* - u_0||_{L^2(\Omega)} \leq R$, for some $R < \infty$, then for every $\epsilon > 0$, there exists a neural network u_{ϵ} with size

$$O\left(d^{2\frac{\log\left(\frac{R}{\epsilon}\right)}{\log\kappa}}\left(N_{0}+N_{A}\right)+\frac{\log\left(\frac{R}{\epsilon}\right)}{\log\kappa}\left(N_{\tilde{f}}+N_{c}\right)\right)$$

such that $\|u^* - u_{\epsilon}\|_{L^2(\Omega)} \leq \epsilon$, and $\kappa = \frac{\lambda_k + \lambda_1}{\lambda_k - \lambda_1}$ Furthermore, the activation functions used in u_{ϵ} belong to the set $\Sigma \cup \{\rho\}$ where $\rho(y) = y^2$ for all $y \in \mathbb{R}$ is the square activation function.

The above theorem statement shows that given an initial neural network estimate $u_0 \in H_0^1(\Omega)$ containing N_0 parameters, we can recover a neural network that is ϵ close to the unique solution u^* . The number of parameters in u_{ϵ} depend on how close the initial estimate u_0 is to the solution u^* and N_0 . This results in a trade-off, wherein a better approximation may require larger number of parameters compared to a poorer approximation with fewer parameters.

For our proof we introduce an iterative sequence with roughly $O\left(\frac{\log(R/\epsilon)}{\log(\kappa)}\right)$ functions to reach an ϵ -approximation to the solution u^* . The parameter κ depends upon λ_k and λ_1 , and acts as analogous

to the condition number of a matrix². Therefore, if $\lambda_k \gg \lambda_1$ then the number of parameters in u_{ϵ} increases.

Corollary 1. If u_0 is initialized to be identically 0, then $u_0 \in H_0^1(\Omega)$ (as it satisfies the boundary condition), then the number of parameters required in u_{ϵ} is bounded by

$$O\left(d^{2\frac{\log\left(\frac{\|f\|_{L^{2}(\Omega)}}{\lambda_{1}\epsilon}\right)}{\log\kappa}}(N_{0}+N_{A})+\frac{\log\left(\frac{\|f\|_{L^{2}(\Omega)}}{\lambda_{1}\epsilon}\right)}{\log\kappa}(N_{\tilde{f}}+N_{c})\right)$$

Proof. Given that u_0 is identically 0, the value of R in Theorem 1 equals $||u^* - u_0||_{L^2(\Omega)} = ||u^*||_{L^2(\Omega)}$ Using the inequality in (3), we have,

$$\|u^{\star}\|_{L^{2}(\Omega)}^{2} \leq \frac{\langle Lu^{\star}, u^{\star} \rangle}{\lambda_{1}}$$

$$\leq \frac{1}{\lambda_{1}} \langle f, u^{\star} \rangle_{L^{2}(\Omega)} \quad (\text{from } (1))$$

$$\leq \frac{1}{\lambda_{1}} \|f\|_{L^{2}(\Omega)} \|u^{\star}\|_{L^{2}(\Omega)}$$

$$\implies \|u^{\star}\|_{L^{2}(\Omega)} \leq \frac{1}{\lambda_{1}} \|f\|_{L^{2}(\Omega)}$$

We make few remarks about the theorem statement:

Remark 1. Note that even though we have assumed that $u^* \in H^1_0(\Omega)$ is the unique solution of (1) from the boundary regularity condition, we have that $u^* \in H^2(\Omega)$ (Evans [1998], Theorem 4 in Chapter 6). This ensures that the solution u^* is twice differentiable as well.

Remark 2. To get a sense of the scale of λ_1 and λ_k , when $L = -\Delta$ (the Laplacian operator), the eigenvalue $\lambda_1 = \inf_{u \in H_0^1(\Omega)} \frac{\|\nabla u\|_{L^2(\Omega)}}{\|u\|_{L^2(\Omega)}} = \frac{1}{C_p}$, where C_p is the Poincaré constant (see Theorem A.1 in Appendix). For geometrically well-behaved sets Ω (e.g. convex sets with a strongly convex boundary, like a sphere), C_p is even dimension-independent. Further from the Weyl's law operator (Evans [1998], Section 6.5) we have

$$\lim_{k \to \infty} \frac{\lambda_k^{d/2}}{k} = \frac{(2\pi)^d}{\operatorname{vol}(\Omega)\alpha(d)}$$

where $\alpha(d)$ is the volume of a unit ball in d dimensions. So, if $vol(\Omega) \geq 1/\alpha(d)$, λ_k grows as $O(k^{2/d})$, which is a constant so long as $\log k \ll d$.

6 Proof of Main Result

At the outset, we provide some intuition behind the proof, via an analogy between a uniformly elliptic operator and a positive definite matrix in linear algebra. We can think of finding the solution to the equation Lu = f for an elliptic L as analogous to finding the solution to the linear system

²If $\lambda_k \gg \lambda_1$ then it implies that $\kappa \approx 1 + \lambda_1 / \lambda_k$ and hence $1/\log(\kappa) \approx \lambda_k / \lambda_1$.

of equations Qx = k, where Q is a $d \times d$ positive definite matrix, and x and k are d-dimensional vectors. One way to solve such a linear system is by minimizing the strongly convex function $||Qx - b||^2$ using gradient descent. Since the objective is strongly convex, after $O(\log(1/\epsilon))$ gradient steps, we reach an ϵ -optimal point in an l_2 sense.

Our proof uses a similar strategy. First, we show that for the operator L, we can define a sequence of functions that converge to an ϵ -optimal function approximation (in this case in the $L^2(\Omega)$ norm) after $O(\log(1/\epsilon)$ steps—similar to the rate of convergence for strongly convex functions. Next, we inductively show that each iterate in the sequence can be approximated by a small neural network. More precisely, we show that given a bound on the size of the *t*-th iterate u_t , we can, in turn, upper bound the size of the (t+1)-th iterate u_{t+1} because the update transforming u_t to u_{t+1} can be simulated by a small neural network. These iterations look roughly like like $u_{t+1} \leftarrow u_t - \eta(Lu_t - f)$, and use a "backpropagation" lemma (Lemma 5) which bounds the size of the derivative of a neural network.

6.1 Defining a Convergent Sequence

We begin by defining the convergence sequence. The rough idea is to perform gradient descent in $L^2(\Omega)$ on an appropriate loss, which in our case resembles $\frac{1}{2} \|Lu - f\|_{L^2(\Omega)}$ [Neuberger, 2009, Faragó and Karátson, 2001, 2002] as suggested by Propositon 1. We show that our sequence converges to the solution u^* in the $L^2(\Omega)$ norm. Towards that end, we first note that the operator L is bounded from above and "positive definite" within the span of the its first k eigenfunctions. Although this is a standard lemma, we include the proof here for completeness:

Lemma 1. Let $L: H_0^1(\Omega) \to H_0^1(\Omega)$ be an elliptic operator. Then for all $v \in H_0^1(\Omega)$ such that $v \in \operatorname{span}\{\varphi_1, \cdots, \varphi_k\}$

1. The inner product is bounded,

$$\lambda_1 \|v\|_{L^2(\Omega)} \le \langle Lv, v \rangle_{L^2(\Omega)} \le \lambda_k \|v\|_{L^2(\Omega)} \tag{3}$$

2. For every $u \in \text{span}\{\varphi_1, \cdots, \varphi_k\}$, we have

$$\left\| \left(I - \frac{2}{\lambda_k + \lambda_k} L \right) u \right\|_{L^2(\Omega)} \le \frac{\lambda_k - \lambda_1}{\lambda_k + \lambda_1} \| u \|_{L^2(\Omega)}$$
(4)

Proof. Since $v \in \text{span}\{\varphi_1, \cdots, \varphi_k\}$, we use the definition of eigenvalues in (2) to get,

$$\frac{\langle Lv, v \rangle_{L^2(\Omega)}}{\|v\|_{L^2(\Omega)}} \le \sup_{v} \frac{\langle Lv, v \rangle_{L^2(\Omega)}}{\|v\|_{L^2(\Omega)}} \le \lambda_k$$
$$\implies \langle Lv, v \rangle_{L^2(\Omega)} \le \lambda_k \|v\|_{L^2(\Omega)}^2$$

and similarly

$$\frac{\langle Lv, v \rangle_{L^2(\Omega)}}{\|v\|_{L^2(\Omega)}} \ge \inf_v \frac{\langle Lv, v \rangle_{L^2(\Omega)}}{\|v\|_{L^2(\Omega)}} \le \lambda_1$$
$$\implies \langle Lv, v \rangle_{L^2(\Omega)} \ge \lambda_1 \|v\|_{L^2(\Omega)}^2$$

In order to prove (2.) let us first denote $\tilde{L} := \left(I - \frac{2}{\lambda_k + \lambda_1}L\right)$. Note if φ is an eigenfunction of L with corresponding eigenvalue λ , it is also an eigenfunction of \tilde{L} with corresponding eigenvalue $\frac{\lambda_k + \lambda_1 - 2\lambda}{\lambda_k + \lambda_1}$.

Hence, writing $u \in \text{span}\{\varphi_1, \cdots, \varphi_k\}$ as $u = \sum_{i=1}^k d_i \varphi_i$, where $d_i = \langle u, \varphi_i \rangle$, we have

$$\|\tilde{L}u\|_{L^{2}(\Omega)}^{2} = \left\|\sum_{i=1}^{k} \frac{\lambda_{k} + \lambda_{1} - 2\lambda_{i}}{\lambda_{k} + \lambda_{1}} d_{i}\varphi_{i}\right\|_{L^{2}(\Omega)}^{2} \le \max_{i \in k} \left(\frac{\lambda_{k} + \lambda_{1} - 2\lambda_{i}}{\lambda_{k} + \lambda_{1}}\right)^{2} \left\|\sum_{i=1}^{k} d_{i}\varphi_{i}\right\|_{L^{2}(\Omega)}^{2}$$
(5)

By the orthogonality of $\{\varphi_i\}_{i=1}^k$, we have

$$\left\|\sum_{i=1}^{k} d_{i}\varphi_{i}\right\|_{L^{2}(\Omega)}^{2} = \sum_{i=1}^{k} d_{i}^{2} = \|u\|_{L^{2}(\Omega)}^{2}$$

Since $\lambda_1 \leq \lambda_2 \cdots \leq \lambda_k$, we have $\lambda_k + \lambda_1 - 2\lambda_i \geq \lambda_1 - \lambda_k$ and $\lambda_k + \lambda_1 - 2\lambda_i \leq \lambda_k - \lambda_1$, so $|\lambda_k + \lambda_1 - 2\lambda_i| \leq \lambda_k - \lambda_1$. This implies $\max_{i \in k} \left(\frac{\lambda_k + \lambda_1 - 2\lambda_i}{\lambda_k + \lambda_1}\right)^2 \leq \left(\frac{\lambda_1 - \lambda_k}{\lambda_1 + \lambda_k}\right)^2$. Plugging this back in (5), we get the claim we wanted. \Box

Before we begin with the proof of convergence, we stress upon a few details. From Proposition A.2, we know that u^* is unique and $u^* \in H^1_0(\Omega)$. However, we still need to ensure that the same holds for the ϵ -approximation we derive. This is because for any function $u \in H^1_0(\Omega)$, it implies that $u_{|\partial\Omega} = 0$ (this follows from the definition of $H^1_0(\Omega)$ space) and hence, the Dirichlet boundary condition will be satisfied.

One way to do so is to ensure that all the functions in our sequence belong to the $H_0^1(\Omega)$ space. Since our updates are of the form $u_{t+1} \leftarrow u_t - \eta(Lu - f)$, we first show that the function we can use the function \tilde{f} instead of f, since $\tilde{f} \in \text{span}\{\varphi_1, \dots, \varphi_k\}$ and therefore belongs to the $H_0^1(\Omega)$ space. Using Assumption (iii), given that $||f - \tilde{f}|| \leq \epsilon_f$ we show in the following lemma that the solution to a new PDE defined as $Lu = \tilde{f}$ is close to u^* .

Lemma 2. Assume that \tilde{u} is a solution to the PDE, $Lu = \tilde{f}$, where $\tilde{f} : H_0^1(\Omega) \to \mathbb{R}$ and $\tilde{f} \in \text{span}\{\varphi_1, \dots, \varphi_k\}$ such that, $\|\tilde{f} - f\|_{L^2(\Omega)} \leq \epsilon \cdot \lambda_1$ for an $\epsilon > 0$. Then, we have $\|u^* - \tilde{u}\|_{H_0^1(\Omega)} \leq \epsilon$.

Proof. The fact that \tilde{u} is a unique solution to $Lu = \tilde{f}$ depends upon the properties of the operator L that follow from the Assumptions in Section 5. The proof is deferred to the appendix (Proposition A.2).

Now we show that the solutions to the two PDEs are close. Using the result from Equation 3 and Theorem A.1 we have,

$$\begin{aligned} \|u^{\star} - \tilde{u}\|_{L^{2}(\Omega)}^{2} &\leq \frac{1}{\lambda_{1}} \langle L(u^{\star} - \tilde{u}), u^{\star} - \tilde{u} \rangle_{L^{2}(\Omega)} \\ &= \frac{1}{\lambda_{1}} \langle f - \tilde{f}, u^{\star} - \tilde{u} \rangle_{L^{2}(\Omega)} \\ &\leq \frac{1}{\lambda_{1}} \|f - \tilde{f}\|_{L^{2}(\Omega)} \|u^{\star} - \tilde{u}\|_{L^{2}(\Omega)} \\ &\implies \|u^{\star} - \tilde{u}\|_{L^{2}(\Omega)} \leq \frac{1}{\lambda_{1}} \|f - \tilde{f}\|_{L^{2}(\Omega)} \leq \epsilon \end{aligned}$$

We define the iterative convergent sequence in (6), where we use f instead of f, and the sequence therefore converges to \tilde{u} instead of u^* . Since, we know that the solution to $Lu = \tilde{f}$ should minimize the norm of $\|Lu - \tilde{f}\|_{L^2(\Omega)}$ here $Lu - \tilde{f}$ resembles the gradient. With appropriately chosen step size η (here $\eta = \frac{2}{\lambda_k + \lambda_1}$) we show that this sequence converges to an approximate solution u_{ϵ} in $O(\log(1/\epsilon))$ steps. Since the sequence is defined using functions that lie in span{ $\varphi_1, \dots, \varphi_k$ }, all the iterates and therefore the final function $u_{\epsilon} \in \text{span}{\{\varphi_1, \dots, \varphi_k\}}$ as well.

Finally, in the next lemma we define the iterates converging to \tilde{u} , as well as characterize the convergence rate.

Lemma 3 (Convergence of gradient descent in H_0^1). Let \tilde{u} denote the unique solution to the PDE $Lu = \tilde{f}$, where $\tilde{f} \in H_0^1(\Omega)$ defined in Lemma 2, and the operator L satisfies the conditions in Lemma 1. Then for any $u_0 \in H_0^1(\Omega)$ such that $u_0 \in \text{span}\{\varphi_1, \dots, \varphi_k\}$, we define the sequence

$$u_{t+1} \leftarrow u_t - \frac{2}{\lambda_1 + \lambda_k} (Lu_t - \tilde{f}) \quad (t \in \mathbb{N})$$
(6)

where for all $t \in \mathbb{N}$, $u_t \in H_0^1(\Omega)$. Then for any $\epsilon > 0$ we have $||u_T - \tilde{u}||_{L^2(\Omega)} \leq \epsilon$ after T iterations where,

$$T \ge \frac{\log\left(\frac{\|u_0 - \tilde{u}\|_{L^2(\Omega)}}{\epsilon}\right)}{\log\left(\frac{\lambda_k + \lambda_1}{\lambda_k - \lambda_1}\right)}$$

Proof. Given that $u_0 \in H_0^1(\Omega)$ and $u_0 \in \operatorname{span}\{\varphi_1, \cdots, \varphi_k\}$ the function $Lu_0 \in H_0^1(\Omega)$ and $Lu_0 \in \operatorname{span}\{\varphi_1, \cdots, \varphi_k\}$ as well (because $L : H_0^1(\Omega) \to H_0^1(\Omega)$).

As $\tilde{f} \in \text{span}\{\varphi_1, \cdots, \varphi_k\}$, all the iterates in the sequence will also belong to $H_0^1(\Omega)$ and will lie in the span $\{\varphi_1, \cdots, \varphi_k\}$.

Now at a step t the iteration looks like,

$$u_{t+1} = u_n - \frac{2}{\lambda_k + \lambda_1} \left(L u_t - \tilde{f} \right)$$
$$u_{t+1} - \tilde{u} = \left(I - \frac{2}{\lambda_k + \lambda_1} L \right) \left(u_t - \tilde{u} \right)$$

Using the result in (4) we have,

$$\|u_{t+1} - \tilde{u}\|_{L^{2}(\Omega)} \leq \left(\frac{\lambda_{k} - \lambda_{1}}{\lambda_{k} + \lambda_{1}}\right) \|u_{t} - \tilde{u}\|_{L^{2}(\Omega)}$$
$$\implies \|u_{t+1} - \tilde{u}\|_{L^{2}(\Omega)} \leq \left(\frac{\lambda_{k} - \lambda_{1}}{\lambda_{k} + \lambda_{1}}\right)^{t} \|u_{0} - \tilde{u}\|_{L^{2}(\Omega)}$$

Hence this implies that, $||u_T - \tilde{u}||_{L^2(\Omega)} \leq \epsilon$ when

$$T \ge \frac{\log\left(\frac{\|u_0 - \tilde{u}\|_{L^2(\Omega)}}{\epsilon}\right)}{\log\left(\frac{\lambda_k + \lambda_1}{\lambda_k - \lambda_1}\right)}$$

Using $\kappa := \frac{\lambda_k + \lambda_1}{\lambda_k - \lambda_1}$ we can rewrite the above as

$$T \ge \frac{\log\left(\frac{\|u_0 - \tilde{u}\|_{L^2(\Omega)}}{\epsilon}\right)}{\log(\kappa)}$$

Further, combining the results from Lemma 2 and Lemma 3, we have the following result

$$\begin{aligned} \|u^{\star} - u_T\|_{L^2(\Omega)} &\leq \|u^{\star} - \tilde{u}\|_{L^2(\Omega)} + \|\tilde{u} - u_T\|_{L^2(\Omega)} \\ &\leq \epsilon + \epsilon \leq 2\epsilon \end{aligned}$$

From Lemma 3, we show that the convergence rate depends upon how far the initial estimate u_0 is to the solution u^* . Also, if λ_1 and λ_k are close, then we get faster convergence.

6.2 Approximating iterates by neural networks

The convergence proof in Lemma 3 shows that there exists a sequence of functions (6) that converges to the solution u^* . We now show that if we start from a neural network u_0 with N parameters, we can bound the increase in the size of a network representing u_{t+1} as a function of the size of the network representing u_t . The crux of the proof will be the following three observations:

- 1. The iterates have the form $u_{t+1} = u_t \eta (Lu_t f)$, so the next iterate can be written in terms of three operations on u_t , a and f: taking derivatives, multiplication and addition.
- 2. If g is representable as a neural network with N parameters, the coordinates of the vector ∇g can be represented by a neural network with O(N) parameters. This is a classic result (Theorem 5), essentially following from the backpropagation algorithm.
- 3. The addition—or multiplication—of two functions representable as neural networks with sizes N_1, N_2 can be represented as neural networks with size O(N + M).

Using these facts, we can write down a recurrence upper bounding the size of the network needed to represent u_{t+1} in terms of the size needed to represent u_t .

Lemma 4 (Recursion Lemma). Consider the update equation defined in 6, where $A : \mathbb{R}^d \to \mathbb{R}^{d \times d}$ is a neural network with N_A parameters, and $\tilde{f} : \mathbb{R}^d \to \mathbb{R}$ is a network with $N_{\tilde{f}}$ parameters. If at step $t, u_t : \mathbb{R}^d \to \mathbb{R}$ is a neural network with N_t parameters, then u_{t+1} is a neural network with $O(d^2(N_A + N_t) + N_t + N_{\tilde{f}} + N_c)$ parameters.

Proof. Consider the update,

$$u_{t+1} \leftarrow u_t - \eta \left(Lu_t - \tilde{f} \right)$$
$$u_{t+1} \leftarrow u_t - \eta \left(\sum_{i=1}^d \partial_i \left(A \nabla u_t \right)_i + cu_t - \tilde{f} \right)$$
$$u_{t+1} \leftarrow u_t - \eta \left(\sum_{i,j=1}^d a_{ij} \partial_{ij} u_t + \sum_{j=1}^d \left(\sum_{i=1}^d \partial_i a_{ij} \right) \partial_j u_t + cu_t - \tilde{f} \right)$$

Using Lemma 5, $\partial_{ij}u_t$, ∂_ju_t and $\partial_i a_{ij}$ can be approximated in $O(N_t)$, $O(N_t)$ and $O(N_A)$ parameters. Further, $\partial_i a_{ij} \partial_j u$ and $a_{ij} \partial_{ij} u$ can be approximated using $O(N_A + N_t)$ parameters, and cu_t can be approximated by $O(N_t + N_c)$ parameters, from Lemma 6. Hence u_{t+1} can be approximated in $O(d^2(N_A + N_t) + N_{\tilde{f}} + N_c + N_t)$ parameters. Note that, throughout the entire proofs O hides independent constants. Concerning the recurrence given in (6), Lemma 4 shows that at each step, the complexity of the algorithm increases by the factor of d^2 . Combining the results of Lemma 3 and Lemma 4, the following gives us the final result for the total number of parameters required,

$$N_{t+1} \le d^2 N_{t-1} + d^2 N_A + N_t + N_{\tilde{f}} + N_c$$

$$\implies N_T \le d^{2T} N_0 + \frac{d^2 (d^T - 1)}{d^2 - 1} N_A + T(N_{\tilde{f}} + N_c)$$

Hence, the total number of parameters required for a neural network to approximate a solution to a PDE of the form in Definition 1

$$O\left(d^{2\frac{\log\left(\frac{\|u_{0}-\tilde{u}\|_{L^{2}(\Omega)}}{\epsilon}\right)}{\log\kappa}}(N_{0}+N_{A})+\frac{\log\left(\frac{\|u_{0}-\tilde{u}\|_{L^{2}(\Omega)}}{\epsilon}\right)}{\log\kappa}(N_{\tilde{f}}+N_{c})\right)$$

Here O hides the smaller factors of the dimension d and their coefficients.

Operations on Neural Network functionals

Lemma 5 (Backpropagation Rumelhart et al. [1986]). Consider neural network $g : \mathbb{R}^m \to \mathbb{R}$ with depth l and N parameters. There exists a neural of size O(l+N) that calculates the gradient $\frac{dg}{di}$ for all $i \in [m]$.

Lemma 6 (Addition and Multiplication). Given neural networks $g: \Omega \to \mathbb{R}$, $h: \Omega \to \mathbb{R}$, with N_g and N_h parameters respectively, the operations g(x) + h(x) and $g(x) \cdot h(x)$ can be represented by neural networks of size $O(N_g + N_h)$, and square activation functions.

Proof. For Addition, there exists a network h containing both networks f and g as subnetworks and an extra layer to compute the addition between their outputs. Hence, the total number of parameters in such a network will be $O(N_f + N_g)$.

For Multiplication, consider the operation $f(x) \cdot g(x) = \frac{1}{2} \left((f(x) + g(x))^2 - f(x)^2 - g(x)^2 \right)$. Then following the same argument as for addition of two networks, we can construct a network h containing both networks and square activation function.

While the representation result in Lemma 6 is shown using square activation, we refer to Yarotsky [2017] for approximation results with ReLU activation. The scaling with respect to the number of parameters in the network remains the same.

7 Applications to Learning Operators

A number of recent works attempt to simultaneously approximate the solutions for an entire family of PDEs by learning a *parametric map* that takes as inputs (some representation of) the coefficients of a PDE and returns its solution [Bhattacharya et al., 2020, Li et al., 2020b,a]. For example, given a set of observations that $\{a_j, u_j\}_{j=1}^N$, where each a_j denotes a coefficient of a PDE with corresponding solution u_j , they learn a neural network G such that for all j, $u_j = G(a_j)$. Our parametric results provide useful insights for why simultaneously solving an entire family of PDEs with a *single neural network* G is possible in the case of linear elliptic PDEs. Consider the case where the coefficients a_j in the family of PDEs are given by neural networks with a fixed architecture, but where each instance of a PDE is characterized by a different setting of the weights in the models representing the coefficients. Lemma 4 shows that each iteration of our sequence (6) constructs a new network containing both the current solution and the coefficient networks as subnetworks. We can view our approximation as not merely approximating the solution to a single PDE but to every PDE in the family, by treating the coefficient networks as placeholder architectures whose weights are provided as inputs. With this view, our construction provides a parametric map between the coefficients of an elliptic PDE in this family and its solution.

8 Conclusion and Future Work

We derive the complexity bounds for neural network approximation for linear elliptic PDEs with Dirichlet boundary conditions where the coefficients are neural networks with finite number of parameters. By simulating gradient descent in function spaces using neural networks, we construct a neural network that approximates the solution of a PDE. We show that the number of parameters in the neural network depends on the parameters required to represent the coefficients and has a poly(d) dependence on the dimension of the input space, therefore avoiding the curse of dimensionality.

The most immediate open questions are related to tightening our results: can the assumption that \tilde{f} lies within the span of the first k eigenfunctions be relaxed? It seems like some kind of "regularity" assumption is needed, but we might imagine that weaker conditions could suffice. The dependencies in the exponent of d on R and κ may also be improvable. Finally, the idea of simulating an iterative algorithm by a neural network to derive a representation-theoretic result seems applicable more broadly, and may be a fertile ground for further work, both theoretically and empirically (as it suggest a particular kind of weight tying.)

References

W. F. Ames. Numerical methods for partial differential equations. Academic press, 2014.

- J. D. Anderson and J. Wendt. Computational fluid dynamics, volume 206. Springer, 1995.
- D. Bahdanau, K. Cho, and Y. Bengio. Neural machine translation by jointly learning to align and translate. arXiv preprint arXiv:1409.0473, 2014.
- K. Bhattacharya, B. Hosseini, N. B. Kovachki, and A. M. Stuart. Model reduction and neural networks for parametric pdes. arXiv preprint arXiv:2005.03180, 2020.
- F. Black and M. Scholes. The pricing of options and corporate liabilities. Journal of political economy, 81(3):637–654, 1973.
- J. Crank and P. Nicolson. A practical method for numerical evaluation of solutions of partial differential equations of the heat-conduction type. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 43, pages 50–67. Cambridge University Press, 1947.
- J. Devlin, M.-W. Chang, K. Lee, and K. Toutanova. Bert: Pre-training of deep bidirectional transformers for language understanding. arXiv preprint arXiv:1810.04805, 2018.

- M. Ehrhardt and R. E. Mickens. A fast, stable and accurate numerical method for the black–scholes equation of american options. *International Journal of Theoretical and Applied Finance*, 11(05): 471–501, 2008.
- L. C. Evans. Partial Differential Equations. graduate studies in mathematics. american mathematical society, 1998. ISBN 9780821807729.
- I. Faragó and J. Karátson. The gradient-finite element method for elliptic problems. Computers & Mathematics with Applications, 42(8-9):1043–1053, 2001.
- I. Faragó and J. Karátson. Numerical solution of nonlinear elliptic problems via preconditioning operators: Theory and applications, volume 11. Nova Publishers, 2002.
- D. Gilbarg and N. S. Trudinger. Elliptic partial differential equations of second order. 2001.
- P. Grohs and L. Herrmann. Deep neural network approximation for high-dimensional elliptic pdes with boundary conditions. arXiv preprint arXiv:2007.05384, 2020.
- P. Grohs, F. Hornung, A. Jentzen, and P. Von Wurstemberger. A proof that artificial neural networks overcome the curse of dimensionality in the numerical approximation of black-scholes partial differential equations. arXiv preprint arXiv:1809.02362, 2018.
- J. Han, A. Jentzen, and E. Weinan. Solving high-dimensional partial differential equations using deep learning. *Proceedings of the National Academy of Sciences*, 115(34):8505–8510, 2018.
- A. Jentzen, D. Salimova, and T. Welti. A proof that deep artificial neural networks overcome the curse of dimensionality in the numerical approximation of kolmogorov partial differential equations with constant diffusion and nonlinear drift coefficients. arXiv preprint arXiv:1809.07321, 2018.
- Y. Khoo, J. Lu, and L. Ying. Solving parametric pde problems with artificial neural networks. arXiv preprint arXiv:1707.03351, 2017.
- A. Krizhevsky, I. Sutskever, and G. E. Hinton. Imagenet classification with deep convolutional neural networks. In F. Pereira, C. J. C. Burges, L. Bottou, and K. Q. Weinberger, editors, Advances in Neural Information Processing Systems, volume 25, pages 1097–1105. Curran Associates, Inc., 2012.
- G. Kutyniok, P. Petersen, M. Raslan, and R. Schneider. A theoretical analysis of deep neural networks and parametric pdes. arXiv preprint arXiv:1904.00377, 2019.
- I. E. Lagaris, A. Likas, and D. I. Fotiadis. Artificial neural networks for solving ordinary and partial differential equations. *IEEE transactions on neural networks*, 9(5):987–1000, 1998.
- I. E. Lagaris, A. C. Likas, and D. G. Papageorgiou. Neural-network methods for boundary value problems with irregular boundaries. *IEEE Transactions on Neural Networks*, 11(5):1041–1049, 2000.
- P. D. Lax and A. N. Milgram. Parabolic equations, volume 33 of annals of mathematics studies, 1954.

- R. J. LeVeque. Finite difference methods for ordinary and partial differential equations: steady-state and time-dependent problems. SIAM, 2007.
- Z. Li, N. Kovachki, K. Azizzadenesheli, B. Liu, K. Bhattacharya, A. Stuart, and A. Anandkumar. Fourier neural operator for parametric partial differential equations. arXiv preprint arXiv:2010.08895, 2020a.
- Z. Li, N. Kovachki, K. Azizzadenesheli, B. Liu, K. Bhattacharya, A. Stuart, and A. Anandkumar. Neural operator: Graph kernel network for partial differential equations. arXiv preprint arXiv:2003.03485, 2020b.
- A. Malek and R. S. Beidokhti. Numerical solution for high order differential equations using a hybrid neural network—optimization method. *Applied Mathematics and Computation*, 183(1): 260–271, 2006.
- J. Neuberger. Sobolev gradients and differential equations. Springer Science & Business Media, 2009.
- B. Oksendal. Stochastic differential equations: an introduction with applications. Springer Science & Business Media, 2013.
- M. N. Ozişik, H. R. Orlande, M. J. Colaço, and R. M. Cotta. *Finite difference methods in heat transfer.* CRC press, 2017.
- A. Radford, L. Metz, and S. Chintala. Unsupervised representation learning with deep convolutional generative adversarial networks. *arXiv preprint arXiv:1511.06434*, 2015.
- M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics informed deep learning (part i): Datadriven solutions of nonlinear partial differential equations. arXiv preprint arXiv:1711.10561, 2017.
- M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, 2019.
- D. E. Rumelhart, G. E. Hinton, and R. J. Williams. Learning representations by back-propagating errors. nature, 323(6088):533–536, 1986.
- J. Sirignano and K. Spiliopoulos. Dgm: A deep learning algorithm for solving partial differential equations. *Journal of computational physics*, 375:1339–1364, 2018.
- R. Temam. Navier-Stokes equations: theory and numerical analysis, volume 343. American Mathematical Soc., 2001.
- A. Vaswani, N. Shazeer, N. Parmar, J. Uszkoreit, L. Jones, A. N. Gomez, L. Kaiser, and I. Polosukhin. Attention is all you need. In *Advances in neural information processing systems*, pages 5998–6008, 2017.
- D. Yarotsky. Error bounds for approximations with deep relu networks. *Neural Networks*, 94: 103–114, 2017.

A Brief Overview of Partial Differential Equations

In this section, we introduce few key definitions and results from PDE literature. We note that that the results in this section are standard and can be found in classical references on PDEs. We note that the results in this section are standard and have been included in the Appendix for completeness. We refer the reader to classical texts on PDEs ((Evans, 1998, Gilbarg and Trudinger, 2001)) for more details.

We will use the following Poincaré inequality throughout our proofs.

Theorem A.1 (Poincaré's inequality). Given $\Omega \subset \mathbb{R}^d$, a bounded open subset, there exists a constant $C_p > 0$ such that for all $u \in H_0^1(\Omega)$

$$||u||_{L^2(\Omega)} \le C_p ||\nabla u||_{L^2(\Omega)}.$$

Corollary A.1. For the bounded open subset $\Omega \subset \mathbb{R}^d$, for all $u \in H_0^1(\Omega)$, we define the norm in the Hilbert space $H_0^1(\Omega)$ as

$$\|u\|_{H^1_0(\Omega)} = \|\nabla u\|_{L^2(\Omega)}.$$
(7)

Further, the norm in $H^1_0(\Omega)$ is equivalent to the norm $H^1(\Omega)$.

Proof. Note that for $u \in H_0^1(\Omega)$ we have,

$$\begin{split} \|u\|_{H^{1}(\Omega)} &= \|\nabla u\|_{L^{2}(\Omega)} + \|u\|_{L^{2}(\Omega)} \\ &\geq \|\nabla u\|_{L^{2}(\Omega)} \\ \implies \|u\|_{H^{1}(\Omega)} \geq \|u\|_{H^{1}_{0}(\Omega)}. \end{split}$$

Where we have used the definition of the norm in $H_0^1(\Omega)$ space.

Further, using the result in Theorem A.1 we have

$$\|u\|_{H^{1}(\Omega)}^{2} = \left(\|u\|_{L^{2}(\Omega)}^{2} + \|\nabla u\|_{L^{2}(\Omega)}^{2}\right) \le \left(C_{p}^{2} + 1\right)\|\nabla u\|_{H^{1}(\Omega)}^{2}$$

Therefore, combining the two inequalities we have

$$\|u\|_{H^{1}_{0}(\Omega)} \le \|u\|_{H^{1}(\Omega)} \le C_{h} \|u\|_{H^{1}_{0}(\Omega)}$$
(8)

where $C_h = (C_p^2 + 1)$. Hence we have that the norm in $H_0^1(\Omega)$ and $H^1(\Omega)$ spaces are equivalent. \Box

A.1 Proof for Existence and Uniqueness of the Solution

We begin by introducing a form for $\langle Lu, v \rangle_{L^2(\Omega)}$ that is more amenable for the existence and uniqueness results.

Proposition A.1. For all $u, v \in H_0^1(\Omega)$ the inner product $\langle Lu, v \rangle_{L^2(\Omega)}$ equals,

$$\langle Lu, v \rangle_{L^2(\Omega)} = \int_{\Omega} \left(A \nabla u \cdot \nabla v + c u v \right) dx$$

Proof. We will be using the following integration by parts formula,

$$\int_{\Omega} \frac{\partial u}{\partial x_i} dx = -\int_{\Omega} u \frac{\partial v}{\partial x_i} dx + \int_{\partial \Omega} u v n_i \partial \Gamma$$

Where n_i is a normal at the boundary and $\partial \Gamma$ is an infinitesimal element of the boundary.

Hence we have for all $u, v \in H_0^1(\Omega)$,

$$\begin{split} \langle Lu, v \rangle_{L^{2}(\Omega)} &= \int_{\Omega} -\left(\sum_{i=1}^{d} \left(\partial_{i} \left(A \nabla u\right)_{i}\right)\right) v + cuv \ dx \\ &= \int_{\Omega} A \nabla u \cdot \nabla v dx - \int_{\partial \Omega} \left(\sum_{i=1}^{d} (A \nabla u)_{i} n_{i}\right) v d\Gamma + \int_{\Omega} cuv dx \\ &= \int_{\Omega} A \nabla u \cdot \nabla v dx + \int_{\Omega} cuv dx \quad (\because v_{|\partial\Omega} = 0) \end{split}$$

In order to prove for the uniqueness of the solution, we first state the Lax-Milgram theorem ((Lax and Milgram, 1954)).

Theorem A.2 (Lax Milgram). Let \mathcal{H} be a Hilbert space with inner-product $(\cdot, \cdot) : \mathcal{H} \times \mathcal{H} \to \mathbb{R}$, and let $b : \mathcal{H} \times \mathcal{H} \to \mathbb{R}$ and $l : \mathcal{H} \to \mathbb{R}$ be the bilinear form and linear form, respectively. Assume that there exists constants $C_1, C_2, C_3 > 0$ such that for all $u, v \in \mathcal{H}$ we have,

$$|C_1||u||_{\mathcal{H}}^2 \le b(u,u), \quad |b(u,v)| \le C_2||u||_{\mathcal{H}}||v||_{\mathcal{H}}, \quad and \ |l(u)| \le C_3||u||_{\mathcal{H}}.$$

Then there exists a unique $u \in \mathcal{H}$ such that,

$$b(u, v) = l(v)$$
 for all $v \in \mathcal{H}$

Having stated the Lax-Milgram Theorem, we make the following proposition,

Proposition A.2. Given the assumptions (i)-(iii), solution to the variational formulation in Equation 1 exists and is unique.

Proof. Using the variational formulation defined in (1), we introduce the bilinear form $b(\cdot, \cdot)$: $H_0^1(\Omega) \times H_0^1(\Omega) \to \mathbb{R}$ where $b(u, v) := \langle Lu, v \rangle$. Hence, we prove the theorem by showing that the bilinear form b(u, v) satisfies the conditions in Theorem A.2.

We first show that for all $u, v \in H_0^1(\Omega)$ the following holds,

$$\begin{aligned} |b(u,v)| &= \left| \int_{\Omega} \left(A \nabla u \cdot \nabla v + c u v \right) dx \right| \\ &\leq \int_{\Omega} \left| \left(A \nabla u \cdot \nabla v + c u v \right) \right| dx \\ &\leq \int_{\Omega} \left| A \nabla u \cdot \nabla v \right| dx + \int_{\Omega} \left| c u v \right| dx \\ &\leq \|A\|_{L^{\infty}(\Omega)} \|\nabla u\|_{L^{2}(\Omega)} \|\nabla v\|_{L^{2}(\Omega)} + \|c\|_{L^{\infty}(\Omega)} \|u\|_{L^{2}(\Omega)} \|v\|_{L^{2}(\Omega)} \\ &\leq \Lambda \|\nabla u\|_{L^{2}(\Omega)} \|\nabla v\|_{L^{2}(\Omega)} + \|c\|_{L^{\infty}(\Omega)} \|u\|_{L^{2}(\Omega)} \|v\|_{L^{2}(\Omega)} \\ &\leq \max \left(\Lambda, C_{p} \|c\|_{L^{\infty}(\Omega)} \right) \|u\|_{H^{1}_{0}(\Omega)} \|v\|_{H^{1}_{0}(\Omega)} \end{aligned}$$

Now we show that the bilinear form a(u, u) is lower bounded.

$$\begin{split} b(v,v) &= \int_{\Omega} \left(A \nabla v \cdot \nabla v + c v^2 \right) dx \\ &\geq \lambda \int_{\Omega} \| \nabla v \|^2 dx + \eta \int_{\Omega} v^2 dx \\ &\geq \lambda \int_{\Omega} \| \nabla v \|^2 dx = \lambda \| \nabla v \|_{H^1_0(\Omega)} \end{split}$$

Finally, for $v \in H_0^1(\Omega)$

$$|(f,v)| = \left| \int_{\Omega} f v dx \right| \le \|f\|_{L^{2}(\Omega)} \|v\|_{L^{2}(\Omega)} \le C_{p} \|f\|_{L^{2}(\Omega)} \|v\|_{H^{1}_{0}(\Omega)}$$

Hence, we satisfy the assumptions in required in Theorem A.2 and therefore the variational problem defined in (1) has a unique solution.

A.2 Some Properties of the Operator L

Lemma A.1. Given the assumptions (i)-(iii), if $u \in H^1_0(\Omega)$ then we have the following,

- 1. If $u \in H_0^1(\Omega)$ then $Lu \in H_0^1(\Omega)$ as well.
- 2. Operator L is self-adjoint.

Proof. 1. We first show that for a function $u: \Omega \to \mathbb{R}$ such that $\overline{\operatorname{supp}(u)} \subsetneq \Omega$ then $\overline{\operatorname{supp}(Lu)} \subsetneq \Omega$.

Since $\overline{\operatorname{supp}(u)} \subsetneq \Omega$ there exists an $\epsilon > 0$, such that for all $x \in \Omega \setminus \overline{\operatorname{supp}(u)}$ we have $d(x, \partial \Omega) < \epsilon$. For an $\epsilon' < \epsilon$, we define a open set $S_{\epsilon'} := \{x : d(x, \partial \Omega) < \epsilon'; x \in \Omega\}$. Note that for an $x' \in S_{\epsilon'}$ we have that $x' \notin \overline{\operatorname{supp}(\Omega)}$ (this follows from the definition of the set $S_{\epsilon'}$). This implies that for all $x' \in S_{\epsilon'}$ we have u(x') = 0. Further, we have $\nabla u(x') = \lim_{\delta \to 0} \frac{u(x') - u(x' + \delta)}{\delta} = 0$ and similarly $\nabla^2 u(x') = 0$. Therefore, for all $x' \in S_{\epsilon'}$ we have (Lu)(x') = 0 and $\overline{\operatorname{supp}(Lu)} = \frac{\Omega \setminus S_{\epsilon'}}{\delta}$. With $\epsilon' \to \epsilon$ we have (Lu)(x') = 0 for all $x' \in S_{\epsilon}$ and $\overline{\operatorname{supp}(Lu)} \to \Omega \setminus S_{\epsilon}$ and hence $\overline{\operatorname{supp}(Lu)} \subsetneq \Omega$.

Now we show that if $u \in H_0^1(\Omega)$ then $Lu \in H_0^1(\Omega)$ as well. Recalling the definition of $H_0^1(\Omega)$ space, for all $u \in H_0^1(\Omega)$ there exists a sequence of functions $\phi_n \in C_0^\infty(\Omega)$ such that $\phi_n \to u$ as $n \to \infty$. Further, $\phi_n \in C_0^\infty(\Omega)$ implies that $L\phi_n \in C^\infty(\Omega)$ as well, since ϕ_n for all $n \in \mathbb{N}$ is infinitely differentiable. Moreover, the $\operatorname{supp}(\phi_n) \subsetneq \Omega$ (follows from the definition of $C_0^\infty(\Omega)$ space) from the earlier proof we have also have that $\operatorname{supp}(L\phi_n) \subsetneq \Omega$.

Therefore, for all n the sequence of functions $L\phi_n \in C_0^{\infty}(\Omega)$ and $L\phi_n \to Lu$ as $n \to \infty$, and $Lu \in H_0^1(\Omega)$.

2. To show that the operator $L: H_0^1(\Omega) \to H_0^1(\Omega)$ is self-adjoint, we show that for all $u, v \in H_0^1(\Omega)$ we have $\langle Lu, v \rangle = \langle u, Lv \rangle$.

From Proposition A.1, for functions $u,v\in H^1_0(\Omega)$ we have

$$\langle Lu, v \rangle_{L^{2}(\Omega)} = \int_{\Omega} A \nabla u \cdot \nabla v dx + \int_{\Omega} cuv dx$$
$$= \int_{\Omega} A \nabla v \cdot \nabla u dx + \int_{\Omega} cvu dx$$
$$= \langle Lv, u \rangle$$