

EXAMPLES OF PLANE RATIONAL CURVES WITH TWO GALOIS POINTS IN POSITIVE CHARACTERISTIC, II

SATORU FUKASAWA AND KATSUSHI WAKI

ABSTRACT. It is proved that there exist plane rational curves of degree twelve (resp. twenty-four) with two different outer Galois points such that the Galois group at one of two Galois points is an alternating group A_4 (resp. a symmetric group S_4) of degree four, under the assumption that the characteristic of the ground field is eleven (resp. is twenty-three). For an alternating group A_5 of degree five, a similar existence theorem is confirmed, over a field of characteristic 59, by GAP system.

1. INTRODUCTION

Let $C \subset \mathbb{P}^2$ be an irreducible plane curve over an algebraically closed field k of characteristic $p \geq 0$ with $k(C)$ as its function field. For a point $P \in \mathbb{P}^2$, if the function field extension $k(C)/\pi_P^*k(\mathbb{P}^1)$ induced by the projection π_P is Galois, then P is called a Galois point for C . This notion was introduced by Yoshihara ([5, 6]). Furthermore, if a Galois point P is contained in $\mathbb{P}^2 \setminus C$, then P is said to be outer. The associated Galois group at P is denoted by G_P .

It was difficult to prove the existence of plane curves $C \subset \mathbb{P}^2$ of degree $d = 12$ with two Galois points $P_1, P_2 \in \mathbb{P}^2 \setminus C$ such that G_{P_1} is isomorphic to an alternating group A_4 of degree four. For example, for plane rational curves C of degree d over a field k of characteristic $p = 0$, the case where $d = 12, 24$ or 60 is excluded in Yoshihara's paper [7]. In the present article, it is proved that there exist plane rational curves of this kind, which update the Table in [8], as follows.

Theorem 1. *Let $p = 11$. Then there exist plane rational curves $C \subset \mathbb{P}^2$ of degree $d = 12$ with different outer Galois points $P_1, P_2 \in \mathbb{P}^2 \setminus C$ such that the associated Galois groups G_{P_1}, G_{P_2} are in the following each cases:*

- (a) $G_{P_1} \cong A_4, G_{P_2} \cong \mathbb{Z}/12\mathbb{Z};$

2010 *Mathematics Subject Classification.* 14H50, 20G40.

Key words and phrases. Galois point, plane curve, Galois group, projective linear groups.

The first author was partially supported by JSPS KAKENHI Grant Number JP19K03438.

- (b) $G_{P_1} \cong A_4$, $G_{P_2} \cong D_{12}$, where D_{12} is the dihedral group of order 12;
- (c) $G_{P_1} \cong G_{P_2} \cong A_4$.

For a symmetric group S_4 of degree four, we have the following theorem.

Theorem 2. *Let $p = 23$. Then there exist plane rational curves $C \subset \mathbb{P}^2$ of degree $d = 24$ with different outer Galois points $P_1, P_2 \in \mathbb{P}^2 \setminus C$ such that the associated Galois groups G_{P_1}, G_{P_2} are in the following each cases:*

- (a) $G_{P_1} \cong S_4$, $G_{P_2} \cong \mathbb{Z}/24\mathbb{Z}$;
- (b) $G_{P_1} \cong S_4$, $G_{P_2} \cong D_{24}$;
- (c) $G_{P_1} \cong G_{P_2} \cong S_4$.

For an alternating group A_5 of degree five, we have the following theorem, which will be confirmed by GAP system [4].

Theorem 3. *Let $p = 59$. Then there exist plane rational curves $C \subset \mathbb{P}^2$ of degree $d = 60$ with different outer Galois points $P_1, P_2 \in \mathbb{P}^2 \setminus C$ such that the associated Galois groups G_{P_1}, G_{P_2} are in the following each cases:*

- (a) $G_{P_1} \cong A_5$, $G_{P_2} \cong \mathbb{Z}/60\mathbb{Z}$;
- (b) $G_{P_1} \cong A_5$, $G_{P_2} \cong D_{60}$;
- (c) $G_{P_1} \cong G_{P_2} \cong A_5$.

2. PROOF OF THEOREM 1

According to [1, Theorem 1 and Remark 1] and Lüroth's theorem, the following theorem holds (see also [2, Fact 2], [3, Problem 1]).

Fact 1. *Let $Q \in \mathbb{P}^1$ and let $G_1, G_2 \subset PGL(2, k)$ be different finite subgroups with $|G_1| = |G_2|$. If the two conditions*

- (a) $G_1 \cap G_2 = \{1\}$, and
- (b) *the orbits G_1Q, G_2Q have length $|G_1| = |G_2|$ and $G_1Q = G_2Q$*

are satisfied, then there exists a plane rational curve $C \subset \mathbb{P}^2$ of degree $|G_1| = |G_2|$ with two different outer Galois points $P_1, P_2 \in \mathbb{P}^2 \setminus C$ such that $G_{P_i} = G_i$ for $i = 1, 2$.

To prove Theorem 1 (a) (resp. Theorem 1 (b), Theorem 1 (c)), we have to prove the existence of a pair (G_1, G_2) of subgroups $G_1, G_2 \subset PGL(2, k)$ such that $G_1 \cong A_4$ and $G_2 \cong \mathbb{Z}/12\mathbb{Z}$ (resp. $G_1 \cong A_4$ and $G_2 \cong D_{12}$, $G_1 \cong G_2 \cong A_4$), and conditions (a) and (b) in Fact 1 are satisfied for (G_1, G_2) .

Let $\alpha = 2 \in \mathbb{F}_{11}$, which is a primitive element. Let $\sigma, \tau, \eta \in \text{Aut}(\mathbb{P}^1) \cong PGL(2, k)$ be represented by matrices

$$A_\sigma = \begin{pmatrix} 0 & \alpha \\ 1 & 0 \end{pmatrix}, \quad A_\tau = \begin{pmatrix} 1 & \alpha \\ -1 & -1 \end{pmatrix}, \quad A_\eta = \begin{pmatrix} \alpha & \alpha^4 \\ 1 & \alpha^2 \end{pmatrix}$$

respectively, that is, $\sigma(s, t) = (s, t)A_\sigma$, $\tau(s, t) = (s, t)A_\tau$, $\eta(s, t) = (s, t)A_\eta$. The following lemma can be confirmed by hand.

Lemma 1. *For matrices $A, B \in GL(2, k)$, if the classes of A, B are the same, then it is denoted by $A \sim B$. Then:*

- (a) σ, τ are of order two.
- (b) η is of order three.
- (c) $A_\sigma A_\tau \sim A_\tau A_\sigma$.
- (d) $A_\eta^{-1} A_\sigma A_\eta \sim A_\tau$.
- (e) $A_\eta^{-1} A_\tau A_\eta \sim A_\sigma A_\tau$.
- (f) The group $G_1 := \langle \sigma, \tau, \eta \rangle$ is isomorphic to A_4 .
- (g) The following three matrices represent all elements of $G_1 \cong A_4$ of order two:

$$\begin{pmatrix} 0 & 2 \\ 1 & 0 \end{pmatrix}, \quad \begin{pmatrix} 10 & 9 \\ 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 9 & 9 \\ 1 & 2 \end{pmatrix}.$$

- (h) The following eight matrices represent all elements of $G_1 \cong A_4$ of order three:

$$\begin{pmatrix} 2 & 5 \\ 1 & 4 \end{pmatrix}, \quad \begin{pmatrix} 7 & 5 \\ 1 & 9 \end{pmatrix}, \quad \begin{pmatrix} 4 & 1 \\ 1 & 6 \end{pmatrix}, \quad \begin{pmatrix} 3 & 4 \\ 1 & 10 \end{pmatrix},$$

$$\begin{pmatrix} 1 & 4 \\ 1 & 8 \end{pmatrix}, \quad \begin{pmatrix} 8 & 3 \\ 1 & 5 \end{pmatrix}, \quad \begin{pmatrix} 6 & 3 \\ 1 & 3 \end{pmatrix}, \quad \begin{pmatrix} 5 & 4 \\ 1 & 7 \end{pmatrix}.$$

- (i) The group G_1 acts on the set $\mathbb{P}^1(\mathbb{F}_{11})$ transitively, where $\mathbb{P}^1(\mathbb{F}_{11})$ is the set of all \mathbb{F}_{11} -rational points of \mathbb{P}^1 .

Let ξ be an automorphism of \mathbb{P}^1 represented by

$$A_\xi = \begin{pmatrix} \alpha & 1 \\ 1 & 0 \end{pmatrix},$$

and let $G_2 := \langle \xi \rangle$. It is not difficult to confirm the following lemma.

Lemma 2. *The order of ξ is twelve, and G_2 acts on $\mathbb{P}^1(\mathbb{F}_{11})$ transitively.*

For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 1 (a).

Proposition 1. *The pair (G_1, G_2) satisfies the following two conditions:*

- (a) $G_1 \cap G_2 = \{1\}$.
- (b) *Let $Q \in \mathbb{P}^1(\mathbb{F}_{11})$. Then $G_1Q = \mathbb{P}^1(\mathbb{F}_{11}) = G_2Q$.*

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_2) .

Proof. We prove assertion (a). Assume by contradiction that $G_1 \cap G_2 \neq \{1\}$. Since $G_1 \cap G_2$ is a subgroup of A_4 and of a cyclic group $\langle \xi \rangle \cong \mathbb{Z}/12\mathbb{Z}$, it follows that $\xi^6 \in G_1$ or $\xi^4 \in G_1$. Note that

$$A_\xi^6 \sim \begin{pmatrix} 1 & 1 \\ 1 & 10 \end{pmatrix}, \quad A_\xi^4 \sim \begin{pmatrix} 7 & 1 \\ 1 & 5 \end{pmatrix}.$$

According to Lemma 1 (g) (resp. Lemma 1 (h)), it follows that $\xi^6 \notin G_1$ (resp. that $\xi^4 \notin G_1$). This is a contradiction. \square

Proof of Theorem 1 (a). Proposition 1 and Fact 1 imply Theorem 1 (a). \square

Let $\sigma', \tau' \in \text{Aut}(\mathbb{P}^1)$ be represented by matrices

$$A_{\sigma'} = \begin{pmatrix} 0 & \alpha^3 \\ 1 & 0 \end{pmatrix}, \quad A_{\tau'} = \begin{pmatrix} \alpha^2 & 1 \\ \alpha^2 & \alpha^4 \end{pmatrix}$$

respectively. It is not difficult to confirm the following lemma.

Lemma 3. (a) σ' is of order two.

(b) τ' is of order six.

(c) $A_{\sigma'}^{-1} A_{\tau'} A_{\sigma'} \sim A_{\tau'}^{-1}$.

(d) The group $G_3 := \langle \sigma', \tau' \rangle$ is isomorphic to D_{12} .

(e) The group G_3 acts on $\mathbb{P}^1(\mathbb{F}_{11})$ transitively.

For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 1 (b).

Proposition 2. *The pair (G_1, G_3) satisfies the following two conditions:*

- (a) $G_1 \cap G_3 = \{1\}$.
- (b) *Let $Q \in \mathbb{P}^1(\mathbb{F}_{11})$. Then $G_1Q = \mathbb{P}^1(\mathbb{F}_{11}) = G_3Q$.*

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_3) .

Proof. We prove assertion (a). Assume by contradiction that $G_1 \cap G_3 \neq \{1\}$. Since $G_1 \cap G_3$ is a subgroup of A_4 , it follows that $G_1 \cap G_3$ contains an element of order two or three.

Assume that $G_1 \cap G_3$ contains an element of order three. Since $G_1 \cap G_3$ is a subgroup of $G_3 \cong D_{12}$, it follows that $\tau'^2 \in G_1$. Note that

$$A_{\tau'}^2 \sim \begin{pmatrix} 3 & 3 \\ 1 & 6 \end{pmatrix}$$

According to Lemma 1 (h), it follows that $\tau'^2 \notin G_1$. This is a contradiction.

Assume that $G_1 \cap G_3$ contains an element of order two. It follows that $\sigma \in G_1 \cap G_3$, $\tau \in G_1 \cap G_3$ or $\sigma\tau \in G_1 \cap G_3$. Assume that $\sigma \in G_1 \cap G_3$. Then $\sigma\tau'^3 = \tau'^3\sigma$. Note that

$$A_{\tau'}^3 \sim \begin{pmatrix} 4 & 3 \\ 1 & 7 \end{pmatrix}$$

It follows that

$$A_\sigma A_{\tau'}^3 \sim \begin{pmatrix} 6 & 9 \\ 1 & 9 \end{pmatrix}, \quad A_{\tau'}^3 A_\sigma \sim \begin{pmatrix} 2 & 9 \\ 1 & 5 \end{pmatrix}.$$

This implies that $\sigma\tau'^3 \neq \tau'^3\sigma$. This is a contradiction.

Assume that $\tau \in G_1 \cap G_3$. Then $\tau\tau'^3 = \tau'^3\tau$. It follows that

$$A_\tau A_{\tau'}^3 \sim \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}, \quad A_{\tau'}^3 A_\tau \sim \begin{pmatrix} 9 & 1 \\ 1 & 10 \end{pmatrix}.$$

This implies that $\tau\tau'^3 \neq \tau'^3\tau$. This is a contradiction.

Assume that $\sigma\tau \in G_1 \cap G_3$. Then $(\sigma\tau)\tau'^3 = \tau'^3(\sigma\tau)$. It follows that

$$A_{\sigma\tau} A_{\tau'}^3 \sim \begin{pmatrix} 2 & 4 \\ 1 & 1 \end{pmatrix}, \quad A_{\tau'}^3 A_{\sigma\tau} \sim \begin{pmatrix} 10 & 4 \\ 1 & 9 \end{pmatrix}.$$

This implies that $(\sigma\tau)\tau'^3 \neq \tau'^3(\sigma\tau)$. This is a contradiction. \square

Proof of Theorem 1 (b). Proposition 2 and Fact 1 imply Theorem 1 (b). \square

Let ι be an automorphism of \mathbb{P}^1 represented by

$$A_\iota = \begin{pmatrix} \alpha & 0 \\ 0 & 1 \end{pmatrix},$$

and let $G_4 := \iota G_1 \iota^{-1}$. For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 1 (c).

Proposition 3. *The pair (G_1, G_4) satisfies the following two conditions:*

- (a) $G_1 \cap G_4 = \{1\}$.
- (b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{11})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{11}) = G_4 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_4) .

Proof. We prove assertion (a). The following three matrices represent all elements of G_4 of order two:

$$\begin{pmatrix} 0 & 6 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 5 & 5 \\ 1 & 6 \end{pmatrix}, \begin{pmatrix} 10 & 5 \\ 1 & 1 \end{pmatrix}.$$

According to Lemma 1 (g), it follows that $G_1 \cap G_4$ does not contain an element of order two. The following eight matrices represent all elements of G_4 of order three:

$$\begin{pmatrix} 1 & 4 \\ 1 & 2 \end{pmatrix}, \begin{pmatrix} 9 & 4 \\ 1 & 10 \end{pmatrix}, \begin{pmatrix} 2 & 3 \\ 1 & 3 \end{pmatrix}, \begin{pmatrix} 7 & 1 \\ 1 & 5 \end{pmatrix}, \\ \begin{pmatrix} 6 & 2 \\ 1 & 4 \end{pmatrix}, \begin{pmatrix} 4 & 9 \\ 1 & 8 \end{pmatrix}, \begin{pmatrix} 3 & 9 \\ 1 & 7 \end{pmatrix}, \begin{pmatrix} 8 & 3 \\ 1 & 9 \end{pmatrix}.$$

According to Lemma 1 (h), it follows that $G_1 \cap G_4$ does not contain an element of order three. The claim follows. \square

Proof of Theorem 1 (c). Proposition 3 and Fact 1 imply Theorem 1 (c). \square

3. PROOF OF THEOREM 2

Let $\alpha = 5 \in \mathbb{F}_{23}$, which is a primitive element. Let $\sigma, \tau, \eta, \mu := \eta^2 \in \text{Aut}(\mathbb{P}^1) \cong \text{PGL}(2, k)$ be represented by matrices

$$A_\sigma = \begin{pmatrix} 0 & 1 \\ \alpha^7 & 0 \end{pmatrix}, A_\tau = \begin{pmatrix} \alpha^{12} & \alpha^7 \\ 1 & \alpha^3 \end{pmatrix}, A_\eta = \begin{pmatrix} 1 & \alpha^{10} \\ \alpha^6 & \alpha^{15} \end{pmatrix}, A_\mu = \begin{pmatrix} -1 & 1 \\ -\alpha^7 & 1 \end{pmatrix}$$

respectively. The following lemma can be confirmed by hand.

Lemma 4. (a) σ, μ are of order two.

(b) τ is of order three.

(c) η is of order four.

(d) $A_\sigma A_\mu \sim A_\mu A_\sigma$.

(e) $A_\tau^{-1} A_\sigma A_\tau \sim A_\mu, A_\tau^{-1} A_\mu A_\tau \sim A_\sigma A_\mu$.

(f) $A_\eta^{-1} A_\sigma A_\eta \sim A_\sigma A_\mu, A_\eta^{-1} A_\mu A_\eta \sim A_\mu$.

(g) $A_\eta^{-1} A_\tau A_\eta \sim A_\sigma A_\mu A_\tau^2$.

(h) $A_\eta^{-1} A_\tau^2 A_\eta \sim A_\mu A_\tau$.

(i) $\langle \sigma, \mu, \tau \rangle \cong A_4$, and $G_1 := \langle \sigma, \tau, \eta, \mu \rangle \cong S_4$.

(j) The group G_1 acts on the set $\mathbb{P}^1(\mathbb{F}_{23})$ transitively.

(k) *Let*

$$O_1 = \{(0 : 1), (1 : \alpha), (1 : \alpha^3), (1 : \alpha^6), (1 : \alpha^7), (1 : \alpha^{18})\},$$

$$O_2 = \{(1 : 0), (1 : \alpha^8), (1 : \alpha^9), (1 : \alpha^{12}), (1 : \alpha^{14}), (1 : \alpha^{19})\},$$

$$O_3 = \{(1 : 1), (1 : \alpha^2), (1 : \alpha^4), (1 : \alpha^{10}), (1 : \alpha^{17}), (1 : \alpha^{21})\},$$

$$O_4 = \{(1 : \alpha^5), (1 : \alpha^{11}), (1 : \alpha^{13}), (1 : \alpha^{15}), (1 : \alpha^{16}), (1 : \alpha^{20})\},$$

namely, $\mathbb{P}^1(\mathbb{F}_{23}) = O_1 \cup O_2 \cup O_3 \cup O_4$. Then it follows that

$$\sigma(O_1) = O_2, \sigma(O_2) = O_1, \sigma(O_3) = O_4, \sigma(O_4) = O_3,$$

$$\tau(O_1) = O_1, \tau(O_2) = O_3, \tau(O_3) = O_4, \tau(O_4) = O_2,$$

$$\eta(O_1) = O_2, \eta(O_2) = O_3, \eta(O_3) = O_4, \eta(O_4) = O_1.$$

In particular, G_1 acts on the set $\{O_1, O_2, O_3, O_4\}$ faithfully.

Let ξ be an automorphism of \mathbb{P}^1 represented by

$$A_\xi = \begin{pmatrix} 0 & -1 \\ -1 & 1 \end{pmatrix},$$

and let $G_2 := \langle \xi \rangle$. It is not difficult to confirm the following lemma.

Lemma 5. *The order of ξ is twenty-four, and G_2 acts on $\mathbb{P}^1(\mathbb{F}_{23})$ transitively.*

For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 2 (a).

Proposition 4. *The pair (G_1, G_2) satisfies the following two conditions:*

(a) $G_1 \cap G_2 = \{1\}$.

(b) *Let $Q \in \mathbb{P}^1(\mathbb{F}_{23})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{23}) = G_2 Q$.*

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_2) .

Proof. We prove assertion (a). Assume by contradiction that $G_1 \cap G_2 \neq \{1\}$. Since $G_1 \cap G_2$ is a subgroup of S_4 and of a cyclic group $\langle \xi \rangle \cong \mathbb{Z}/24\mathbb{Z}$, it follows that $\xi^{12} \in G_1$ or $\xi^8 \in G_1$. Note that

$$A_\xi^{12} \sim \begin{pmatrix} -1 & -3 \\ -3 & 1 \end{pmatrix}, \quad A_\xi^8 \sim \begin{pmatrix} 13 & 2 \\ 2 & 11 \end{pmatrix}.$$

It follows that

$$\xi^{12}(0 : 1) = (1 : \alpha^{17}) \in O_3 \quad \text{and} \quad \xi^8(0 : 1) = (1 : \alpha^7) \in O_1$$

for the point $(0 : 1) \in O_1$. According to Lemma 4 (k), it follows that $\xi^{12}(1 : \alpha) \in O_3$ and $\xi^8(1 : \alpha^3) \in O_1$, since $(1 : \alpha), (1 : \alpha^3) \in O_1$. However,

$$\xi^{12}(1 : \alpha) = (1 : \alpha^5) \in O_4, \quad \xi^8(1 : \alpha^3) = (1 : \alpha^{16}) \in O_4.$$

This is a contradiction. □

Proof of Theorem 2 (a). Proposition 4 and Fact 1 imply Theorem 2 (a). □

Let $\sigma', \tau' \in \text{Aut}(\mathbb{P}^1)$ be represented by matrices

$$A_{\sigma'} = \begin{pmatrix} 0 & \alpha^{10} \\ \alpha^9 & 0 \end{pmatrix}, \quad A_{\tau'} = \begin{pmatrix} \alpha^{15} & \alpha \\ -1 & \alpha^7 \end{pmatrix}$$

respectively. The following lemma can be confirmed by hand.

Lemma 6. (a) σ' is of order two.

(b) τ' is of order twelve.

(c) $A_{\sigma'}^{-1} A_{\tau'} A_{\sigma'} \sim A_{\tau'}^{-1}$.

(d) The group $G_3 := \langle \sigma', \tau' \rangle$ is isomorphic to D_{24} .

(e) The group G_3 acts on $\mathbb{P}^1(\mathbb{F}_{23})$ transitively.

(f) Let

$$\begin{aligned} T_1 &= \{(0 : 1), (1 : \alpha^{18}), (1 : \alpha^3), (1 : \alpha^{11}), (1 : \alpha^4), (1 : \alpha^9), \\ &\quad (1 : 1), (1 : \alpha^{21}), (1 : \alpha^{13}), (1 : \alpha^{16}), (1 : \alpha^{17}), (1 : \alpha^{15})\}, \\ T_2 &= \{(1 : 0), (1 : \alpha^8), (1 : \alpha^6), (1 : \alpha^7), (1 : \alpha^{10}), (1 : \alpha^2), \\ &\quad (1 : \alpha), (1 : \alpha^{14}), (1 : \alpha^{19}), (1 : \alpha^{12}), (1 : \alpha^{20}), (1 : \alpha^5)\}, \end{aligned}$$

namely, $\mathbb{P}^1(\mathbb{F}_{23}) = T_1 \cup T_2$. Then it follows that

$$\sigma'(T_1) = T_2, \sigma'(T_2) = T_1, \tau'(T_1) = T_1, \tau'(T_2) = T_2.$$

In particular, G_3 acts on the set $\{T_1, T_2\}$.

For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 2 (b).

Proposition 5. The pair (G_1, G_3) satisfies the following two conditions:

(a) $G_1 \cap G_3 = \{1\}$.

(b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{23})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{23}) = G_3 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_3) .

Proof. We prove assertion (a). It follows from Lemmas 4 (k) and 6 (f) that

$$\begin{aligned} O_1 \cap T_1 &= \{(0 : 1), (1 : \alpha^3), (1 : \alpha^{18})\}, O_1 \cap T_2 = \{(1 : \alpha), (1 : \alpha^6), (1 : \alpha^7)\}, \\ O_2 \cap T_1 &= \{(1 : \alpha^9)\}, O_2 \cap T_2 = \{(1 : 0), (1 : \alpha^8), (1 : \alpha^{12}), (1 : \alpha^{14}), (1 : \alpha^{19})\}, \\ O_3 \cap T_1 &= \{(1 : 1), (1 : \alpha^4), (1 : \alpha^{17}), (1 : \alpha^{21})\}, O_3 \cap T_2 = \{(1 : \alpha^2), (1 : \alpha^{10})\}, \\ O_4 \cap T_1 &= \{(1 : \alpha^5), (1 : \alpha^{20})\}, O_4 \cap T_2 = \{(1 : \alpha^{11}), (1 : \alpha^{13}), (1 : \alpha^{15}), (1 : \alpha^{16})\}. \end{aligned}$$

Note that if $O_i \cap T_j$ consists of a unique point, then $i = 2$ and $j = 1$. Let $\gamma \in G_1 \cap G_3$. Then $\gamma(1 : \alpha^9) = (1 : \alpha^9)$. Since G_3 acts on $\mathbb{P}(\mathbb{F}_{23})$ transitively, it follows that $\gamma = 1$. \square

Proof of Theorem 2 (b). Proposition 5 and Fact 1 imply Theorem 2 (b). \square

Let ι be an automorphism of \mathbb{P}^1 represented by

$$A_\iota = \begin{pmatrix} \alpha^7 & 0 \\ 0 & 1 \end{pmatrix},$$

and let $G_4 := \iota G_1 \iota^{-1}$. For the following proposition, we give the proof of assertion (a), which is the most complicated part of the proof of Theorem 2 (c).

Proposition 6. *The pair (G_1, G_4) satisfies the following two conditions:*

- (a) $G_1 \cap G_4 = \{1\}$.
- (b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{23})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{23}) = G_4 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_4) .

Proof. We prove assertion (a). It follows that

$$\begin{aligned} \iota(O_1) &= \{(0 : 1), (1 : \alpha^{16}), (1 : \alpha^{18}), (1 : \alpha^{21}), (1 : 1), (1 : \alpha^{11})\}, \\ \iota(O_2) &= \{(1 : 0), (1 : \alpha), (1 : \alpha^2), (1 : \alpha^5), (1 : \alpha^7), (1 : \alpha^{12})\}, \\ \iota(O_3) &= \{(1 : \alpha^{15}), (1 : \alpha^{17}), (1 : \alpha^{19}), (1 : \alpha^3), (1 : \alpha^{10}), (1 : \alpha^{14})\}, \\ \iota(O_4) &= \{(1 : \alpha^{20}), (1 : \alpha^4), (1 : \alpha^6), (1 : \alpha^8), (1 : \alpha^9), (1 : \alpha^{13})\}. \end{aligned}$$

Note that $O_i \cap \iota(O_j) = \emptyset$ implies $i = 2$ and $j = 1$. Assume by contradiction that there exists $\gamma \in G_1 \cap G_4 \setminus \{1\}$. According to Lemma 4 (k), it follows that $\gamma(\iota(O_1)) = \iota(O_1)$. Since $\iota(O_1) \subset O_1 \cup O_3 \cup O_4$, it follows that γ acts on $\{O_1, O_3, O_4\}$, that is, $\gamma(O_2) = O_2$.

Assume that γ is of order three. Then $\gamma(O_1) = O_3$ or $\gamma^2(O_1) = O_3$. We can assume that $\gamma(O_1) = O_3$. Then γ coincides with the automorphism $\sigma\tau^2\sigma \in G_1$, which is represented by the matrix

$$A_\sigma A_\tau^2 A_\sigma \sim \begin{pmatrix} 0 & 1 \\ 17 & 0 \end{pmatrix} \begin{pmatrix} -4 & -7 \\ 5 & 2 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 17 & 0 \end{pmatrix} \sim \begin{pmatrix} 11 & 5 \\ 1 & 1 \end{pmatrix}.$$

Then for points $(1 : 0), (1 : \alpha^2) \in \iota(O_2)$, $\gamma(1 : 0) = (1 : \alpha^{14}) \in \iota(O_3)$ and $\gamma(1 : \alpha^2) = (1 : \alpha^5) \in \iota(O_2)$. Since $\gamma \in G_4$, this is a contradiction.

Assume that γ is of order two. Then $\gamma(O_i) = O_i$ for some $i = 1, 3, 4$. If $i = 1$, then γ coincides the automorphism $\tau^{-1}\sigma\eta\tau \in G_1$, which is represented by

$$A_\tau A_\eta A_\sigma A_\tau^2 \sim \begin{pmatrix} -5 & -6 \\ 1 & 10 \end{pmatrix} \begin{pmatrix} 1 & 9 \\ 8 & -4 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 17 & 0 \end{pmatrix} \begin{pmatrix} -4 & -7 \\ 5 & 2 \end{pmatrix} \sim \begin{pmatrix} -10 & 1 \\ 6 & 10 \end{pmatrix}.$$

Then for points $(0 : 1), (1 : 1) \in \iota(O_1)$, $\gamma(0 : 1) = (1 : \alpha^7) \in \iota(O_2)$ and $\gamma(1 : 1) = (1 : \alpha^{16}) \in \iota(O_1)$. Since $\gamma \in G_4$, this is a contradiction. If $i = 4$, then γ coincides the automorphism $\gamma_0 := \eta^2\sigma\eta \in G_1$, which is represented by

$$A_\eta A_\sigma A_\eta^2 \sim \begin{pmatrix} 1 & 9 \\ 8 & -4 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 17 & 0 \end{pmatrix} \begin{pmatrix} 4 & -4 \\ -1 & -4 \end{pmatrix} \sim \begin{pmatrix} -10 & 5 \\ -4 & 10 \end{pmatrix}.$$

Then for points $(0 : 1), (1 : 1) \in \iota(O_1)$, $\gamma(0 : 1) = (1 : \alpha^{10}) \in \iota(O_3)$ and $\gamma(1 : 1) = (1 : \alpha^7) \in \iota(O_2)$. Since $\gamma \in G_4$, this is a contradiction. If $i = 3$, then γ coincides the automorphism $\tau\gamma_0\tau^2$, which is represented by

$$A_\tau^2 A_{\eta^2\sigma\eta} A_\tau \sim \begin{pmatrix} -4 & -7 \\ 5 & 2 \end{pmatrix} \begin{pmatrix} -10 & 5 \\ -4 & 10 \end{pmatrix} \begin{pmatrix} -5 & -6 \\ 1 & 10 \end{pmatrix} \sim \begin{pmatrix} 7 & 3 \\ -10 & -7 \end{pmatrix}.$$

Then for points $(0 : 1), (1 : 1) \in \iota(O_1)$, $\gamma(0 : 1) = (1 : \alpha^{16}) \in \iota(O_1)$ and $\gamma(1 : 1) = (1 : \alpha^{10}) \in \iota(O_3)$. Since $\gamma \in G_4$, this is a contradiction. \square

Proof of Theorem 2 (c). Proposition 6 and Fact 1 imply Theorem 2 (c). \square

4. PROOF OF THEOREM 3

All lemmas and propositions in this section can be confirmed by GAP system [4].

Let $\alpha = 2 \in \mathbb{F}_{59}$, which is a primitive element. Let $\sigma, \tau \in \text{Aut}(\mathbb{P}^1) \cong PGL(2, k)$ be represented by matrices

$$A_\sigma = \begin{pmatrix} -\alpha^{26} & 1 \\ \alpha^{27} & \alpha^{26} \end{pmatrix}, \quad A_\tau = \begin{pmatrix} 1 & \alpha \\ \alpha^6 & \alpha^{34} \end{pmatrix},$$

respectively.

- Lemma 7.** (a) σ is of order two.
 (b) τ is of order three.
 (c) The group $G_1 := \langle \sigma, \tau \rangle$ is isomorphic to A_5 .
 (d) The group G_1 acts on the set $\mathbb{P}^1(\mathbb{F}_{59})$ transitively.

Let ξ be an automorphism of \mathbb{P}^1 represented by

$$A_\xi = \begin{pmatrix} 1 & 1 \\ \alpha^{12} & 0 \end{pmatrix},$$

and let $G_2 := \langle \xi \rangle$.

Lemma 8. The order of ξ is sixty, and G_2 acts on $\mathbb{P}^1(\mathbb{F}_{59})$ transitively.

Proposition 7. The pair (G_1, G_2) satisfies the following two conditions:

- (a) $G_1 \cap G_2 = \{1\}$.
 (b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{59})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{59}) = G_2 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_2) .

Proof of Theorem 3 (a). Proposition 7 and Fact 1 imply Theorem 3 (a). \square

Let $\sigma', \tau' \in \text{Aut}(\mathbb{P}^1)$ be represented by matrices

$$A_{\sigma'} = \begin{pmatrix} 0 & \alpha^2 \\ \alpha^{-1} & 0 \end{pmatrix}, \quad A_{\tau'} = \begin{pmatrix} \alpha^2 & \alpha^3 \\ -1 & -1 \end{pmatrix}$$

respectively.

- Lemma 9.** (a) σ' is of order two.
 (b) τ' is of order 30.
 (c) $A_{\sigma'}^{-1} A_{\tau'} A_{\sigma'} \sim A_{\tau'}^{-1}$.
 (d) The group $G_3 := \langle \sigma', \tau' \rangle$ is isomorphic to D_{60} .
 (e) The group G_3 acts on $\mathbb{P}^1(\mathbb{F}_{59})$ transitively.

Proposition 8. The pair (G_1, G_3) satisfies the following two conditions:

- (a) $G_1 \cap G_3 = \{1\}$.
 (b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{59})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{59}) = G_3 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_3) .

Proof of Theorem 3 (b). Proposition 8 and Fact 1 imply Theorem 3 (b). \square

Let ι be an automorphism of \mathbb{P}^1 represented by

$$A_\iota = \begin{pmatrix} 1 & \alpha^{30} \\ 0 & -\alpha^{15} \end{pmatrix},$$

and let $G_4 := \iota G_1 \iota^{-1}$.

Proposition 9. *The pair (G_1, G_4) satisfies the following two conditions:*

- (a) $G_1 \cap G_4 = \{1\}$.
- (b) Let $Q \in \mathbb{P}^1(\mathbb{F}_{59})$. Then $G_1 Q = \mathbb{P}^1(\mathbb{F}_{59}) = G_4 Q$.

In particular, conditions (a) and (b) in Fact 1 are satisfied for the pair (G_1, G_4) .

Proof of Theorem 3 (c). Proposition 9 and Fact 1 imply Theorem 3 (c). □

REFERENCES

- [1] S. Fukasawa, A birational embedding of an algebraic curve into a projective plane with two Galois points, *J. Algebra* **511** (2018), 95–101.
- [2] S. Fukasawa and K. Higashine, A birational embedding with two Galois points for quotient curves, *J. Pure Appl. Algebra* **225** (2021), 106525, 10 pages.
- [3] S. Fukasawa and K. Waki, Examples of plane rational curves with two Galois points in positive characteristic, *Finite Fields and their Applications: Proceedings of the 14th International Conference on Finite Fields and their Applications, Vancouver, June 3-7, 2019*, pp.181–188, De Gruyter, 2020.
- [4] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.11.0; 2020. (<https://www.gap-system.org>)
- [5] K. Miura and H. Yoshihara, Field theory for function fields of plane quartic curves, *J. Algebra* **226** (2000), 283–294.
- [6] H. Yoshihara, Function field theory of plane curves by dual curves, *J. Algebra* **239** (2001), 340–355.
- [7] H. Yoshihara, Galois points for plane rational curves, *Far East J. Math.* **25** (2007), 273–284; Errata, *ibid.* **29** (2008), 209–212.
- [8] H. Yoshihara and S. Fukasawa, List of problems, available at: <http://hyoshihara.web.fc2.com/openquestion.html>

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, YAMAGATA UNIVERSITY,
KOJIRAKAWA-MACHI 1-4-12, YAMAGATA 990-8560, JAPAN

Email address: s.fukasawa@sci.kj.yamagata-u.ac.jp

Email address: waki@sci.kj.yamagata-u.ac.jp